

Number of partitions of n into parts not divisible by m

Damanvir Singh Binner

Department of Mathematics

Indian Institute of Science Education and Research (IISER)

Mohali, Punjab, India

damanvirbinner@iisermohali.ac.in

Abstract

In this note, we obtain a formula which leads to a practical and efficient method to calculate the number of partitions of n into parts not divisible by m for given natural numbers n and m . Our formula is a generalization of Euler's recurrence for integer partitions which can be viewed as the case $m = 1$ of our formula. Our approach primarily involves the principle of inclusion and exclusion. We also use our approach to obtain a natural combinatorial proof of a identity of Glaisher which generalizes a classical theorem of Euler.

1 Introduction

Euler [1, Corollary 1.2] proved that the number of partitions of n into distinct parts is equal to the number of partitions into distinct parts. Glaisher [4, 5] generalized Euler's result by proving that for any positive integers m and n , the number of partitions of n into parts not divisible by m is equal to the number of partitions of n with each part appearing less than m times. In this note, we obtain a formula that helps us to quickly calculate these numbers. Our formula generalizes Euler's recurrence relation for integer partitions [1, Corollary 1.8]. We use the principle of inclusion and exclusion (PIE) to study these objects, an approach used by the present author and Rattan in the author's PhD Thesis [2, Section 5.1] to obtain a natural combinatorial proof of Euler's recurrence. This approach was also used by the present author [3] to obtain a formula for the number of partitions of n with a given parity of the smallest part.

Note that the calculation of $p(n)$ is very easy compared to finding all the partitions of n because of the availability of formulae such as Hardy-Ramanujan-Rademacher formula and recurrences such as Euler's recurrence. Therefore, Theorem 1 below describes an efficient method to calculate the number of partitions of n into parts not divisible by m , as demonstrated by examples in Section 4.

2 Main Theorem

Theorem 1. Let $P_m(n)$ denote the number of partitions of n into parts not divisible by m . Then, $P_m(n)$ is given by the following formula.

$$P_m(n) = p(n) + \sum_{k \geq 1} (-1)^k \left(p \left(n - \frac{mk(3k-1)}{2} \right) + p \left(n - \frac{mk(3k+1)}{2} \right) \right).$$

Remark 2. For $m = 1$, $P_m(n) = 0$, and Theorem 1 immediately yields Euler's recurrence.

Proof of Theorem 1. We recall some notation defined in [2, Section 5.1] and define some new notation.

- $A_{j,k}(n)$ is the set of partitions of n having exactly k parts of size j ;
- $B_{j,k}(n)$ is the set of partitions of n having at least k parts of size j .
- $C_{j,k}(n)$ is the set of partitions of n having at most k parts of size j ;

The following properties of these sets are immediate.

1. $|B_{j,k}(n)| = p(n - jk)$.
2. If $j \neq j'$, then $|B_{j,k}(n) \cap B_{j',k'}(n)| = p(n - jk - j'k')$.
3. $C_{j,k}(n) = B_{j,k+1}^c(n)$, where the complementation is with respect to the set $Par(n)$, consisting of all partitions of n .
4. In particular, $A_{j,0}(n) = C_{j,0}(n) = B_{j,1}^c(n)$.

We also need the following notation.

- D denotes the set of nonempty distinct partitions.
- For $s \in \mathbb{N}$, T_s denotes the set of partitions into s distinct parts.
- For a partition π , $n(\pi)$ denotes the number of parts in π .

Then using PIE, we have

$$\begin{aligned}
P_m(n) &= |A_{m,0}(n) \cap A_{2m,0}(n) \cap A_{3m,0}(n) \cap \cdots| \\
&= |\cap_{i \geq 1} A_{im,0}(n)| \\
&= |\cup_{i \geq 1} B_{im,1}^{\complement}(n)| \\
&= \sum_{s \geq 0} (-1)^s \sum_{(i_1, i_2, \dots, i_s) \in T_s} |B_{i_1 m, 1}(n) \cap B_{i_2 m, 1}(n) \cap \cdots \cap B_{i_s m, 1}(n)| \\
&= \sum_{s \geq 0} (-1)^s \sum_{(i_1, i_2, \dots, i_s) \in T_s} p(n - i_1 m - i_2 m - \cdots - i_s m) \\
&= \sum_{s \geq 0} \sum_{\pi \in T_s} (-1)^s p(n - m|\pi|) \\
&= p(n) + \sum_{s \geq 1} \sum_{\pi \in T_s} (-1)^s p(n - m|\pi|) \\
&= p(n) + \sum_{\pi \in D} (-1)^{n(\pi)} p(n - m|\pi|),
\end{aligned}$$

which completes the proof by Euler's pentagonal number theorem [1, Theorem 1.6]. Note that here we are able to use PIE even though there are infinitely many sets because all except finitely many are empty. \square

3 A proof of Glaisher's identity

Let $Q_m(n)$ denote the number of partitions of n with each part appearing less than m times. Then using PIE, we have

$$\begin{aligned}
Q_m(n) &= |\cap_{i \geq 1} C_{i, m-1}(n)| \\
&= |\cup_{i \geq 1} B_{i, m}^{\complement}(n)| \\
&= \sum_{s \geq 0} (-1)^s \sum_{(i_1, i_2, \dots, i_s) \in T_s} |B_{i_1, m}(n) \cap B_{i_2, m}(n) \cap \cdots \cap B_{i_s, m}(n)| \\
&= \sum_{s \geq 0} (-1)^s \sum_{(i_1, i_2, \dots, i_s) \in T_s} p(n - i_1 m - i_2 m - \cdots - i_s m) \\
&= \sum_{s \geq 0} \sum_{\pi \in T_s} (-1)^s p(n - m|\pi|) \\
&= p(n) + \sum_{s \geq 1} \sum_{\pi \in T_s} (-1)^s p(n - m|\pi|) \\
&= p(n) + \sum_{\pi \in D} (-1)^{n(\pi)} p(n - m|\pi|),
\end{aligned}$$

completing the proof of Glaisher's generalization of Euler's identity.

4 Examples

For computational purposes, it is convenient to note a few terms and compare the pattern with the terms appearing in Euler's recurrence for partitions, which is given as

$$p(n) = p(n-1) + p(n-2) - p(n-5) - p(n-7) \\ + p(n-12) + p(n-15) - p(n-22) - p(n-26) + \dots$$

Then, by Theorem 1, we have the following formula for $P_m(n)$.

$$P_m(n) = p(n) - p(n-m) - p(n-2m) + p(n-5m) + p(n-7m) \\ - p(n-12m) - p(n-15m) + p(n-22m) + p(n-26m) - \dots$$

First suppose $n = 17$ and $m = 3$. Then, we have

$$P_3(17) = p(17) - p(14) - p(11) + p(2) \\ = 297 - 135 - 56 + 2 \\ = 108.$$

Thus, 108 out of 297 partitions of 17 have parts not divisible by 3 and the remaining 189 partitions have at least one part that is divisible by 3.

Next, suppose $n = 164$ and $m = 7$. Then, we have

$$P_7(164) = p(164) - p(157) - p(150) + p(129) + p(115) - p(80) - p(59) + p(10) \\ = 156919475295 - 80630964769 - 40853235313 \\ + 4835271870 + 1064144451 - 15796476 - 831820 + 42 \\ = 41318063280.$$

Thus, 41318063280 out of 156919475295 partitions of 164 have parts not divisible by 7 and the remaining 115601412015 partitions have at least one part that is divisible by 7.

References

- [1] G. E. Andrews, *The Theory of Partitions*, Cambridge Mathematical Library, Reprint of the 1976 original Cambridge University Press, Cambridge, 1998.
- [2] D. S. Binner, Combinatorial methods for integer partitions, *PhD Thesis*, Simon Fraser University, 2021, available online at <http://summit.sfu.ca/item/21451>.
- [3] D. S. Binner, Number of partitions of n with a given parity of the smallest part, 2022, arXiv preprint, available online at <https://arxiv.org/pdf/2204.07218.pdf>.
- [4] J. W. L. Glaisher, A theorem in partitions, *Messenger of Math.* **12**(1883), 158–170.
- [5] I. Konan, Weighted words at degree two, I: Bressoud's algorithm as an energy transfer, accepted for publication in *Ann. of IHP D*, available online at <https://arxiv.org/pdf/2001.10927.pdf> (2021).