

ON CHARACTERISTIC POWER SERIES OF DUAL SIGNED SELMER GROUPS

JISHNU RAY AND FLORIAN SPRUNG

ABSTRACT. We relate the cardinality of the p -primary part of the Bloch-Kato Selmer group over \mathbb{Q} attached to a modular form at a non-ordinary prime p to the constant term of the characteristic power series of the signed Selmer groups over the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} . This generalizes a result of Vigni and Longo in the ordinary case. In the case of elliptic curves, such results follow from earlier works by Greenberg, Kim, the second author, and Ahmed-Lim, covering both the ordinary and most of the supersingular case.

1. INTRODUCTION

The aim of this paper is to relate the size of a p -adic Selmer group attached to a modular form which is non-ordinary at p to the constant term of the characteristic power series of the cyclotomic deformation of this Selmer group. This has been done in the case of elliptic curves, in the ordinary and most of the supersingular (i.e. non-ordinary) case, and in the case of modular forms, but working with ordinary p . We review what is known in the next three subsections before describing our contribution.

1.1. The case of elliptic curves (ordinary at p). Let E be an elliptic curve defined over a number field F such that E has good ordinary reduction at all the primes of F above an odd prime p . Assume the p -primary Selmer group $\text{Sel}_p(E/F)$ is finite. Let F_∞ be the cyclotomic \mathbb{Z}_p -extension of F . Then by a control theorem, $\text{Sel}_p(E/F_\infty)$ is known to be cotorsion over the corresponding Iwasawa algebra of the Galois group $\Gamma_0 := \text{Gal}(F_\infty/F)$. Let $f_E(X)$ generate the characteristic ideal of the Pontryagin dual of $\text{Sel}_p(E/F_\infty)$, considered as a power series by identifying $1 + X$ with a topological generator of Γ_0 . In [Gre99, Thm. 4.1], Greenberg showed that $f_E(0) \neq 0$ and

$$(1.1) \quad f_E(0) \sim \frac{\#\text{Sel}_p(E/F) \cdot \prod_{v \text{ bad}} c_v(E) \cdot \prod_{v|p} \#(\tilde{E}_v(\mathbb{F}_v)_p)^2}{\#(E(F)_p)^2},$$

2020 *Mathematics Subject Classification*. Primary: 11R23, Secondary: 11R18, 11F11, 11F85.

Key words and phrases. Euler characteristic, non-ordinary, modular forms, signed Selmer groups, algebraic p -adic L -function.

where the symbol \sim means that the two quantities differ by a p -adic unit. Here $c_v(E)$ is the Tamagawa number of E at a prime v of bad reduction, \mathbb{F}_v is the residue field of F at v , $\tilde{E}_v(\mathbb{F}_v)_p$ is the p -torsion of the group of \mathbb{F}_v -rational points of the reduction \tilde{E}_v of E at v and $E(F)_p$ is the p -torsion subgroup of the Mordell-Weil group $E(F)$.

The product of quantities in (1.1) can be reinterpreted as an Euler characteristic, so that

$$f_E(0) \sim \chi(\Gamma_0, \text{Sel}_p(E/F_\infty));$$

see [Gre99, Lemma 4.2] both for the statement and the definition of $\chi(\Gamma_0, \text{Sel}_p(E/F_\infty))$.

1.2. The case of elliptic curves (supersingular and mixed reduction).

In the supersingular case, the classical Selmer group $\text{Sel}_p(E/F_\infty)$ is not cotorsion. However, there are remedies. Denote by S_p the primes of F above p . Suppose for the moment that for each $v \in S_p$, $a_v = 1 + p - \#\tilde{E}_v(\mathbb{F}_p) = 0$, i.e. we are in a subcase of supersingular reduction. Kobayashi constructed signed (plus/minus) Selmer groups $\text{Sel}_p^\pm(E/F_\infty)$, and proved that when $F = \mathbb{Q}$, they are cotorsion over the corresponding Iwasawa algebra [Kob03]. Suppose p splits completely in F and is totally ramified in F_∞ . The signed Selmer groups $\text{Sel}_p^\pm(E/F_\infty)$ are conjectured to be cotorsion over the Iwasawa algebra of Γ_0 . When this is the case, let f^\pm be generators of the characteristic power series of their Pontryagin duals.

When $\text{Sel}_p(E/F)$ is finite, one can use a control theorem to prove $\text{Sel}_p^\pm(E/F_\infty)$ are cotorsion. Kim then showed that

$$f^\pm(0) \sim \#\text{Sel}_p(E/F) \cdot \prod_\ell c_\ell.$$

Here ℓ runs over every prime and the Tamagawa factor $c_\ell = [E(\mathbb{Q}_\ell) : E^0(\mathbb{Q}_\ell)]$. See [Kim13, Thm. 1.2 or Corollary 3.15 and the explanations in its proof] for the above assertions.

If one drops the assumption $a_v = 0$, the second author has generalized Kobayashi's construction to construct a pair of chromatic Selmer groups Sel^\sharp and Sel^\flat [Spr12], and given an analogous formula of Kim's, all assuming that $F = \mathbb{Q}$, see [Spr16, Lemmas 4.4, 4.5, 4.8].

Returning to the case $a_v = 0$, in Kim's results, the local condition appearing in the definition of $\text{Sel}_p^+(E/F_\infty)$ is the same ("plus condition" of Kobayashi) for each v . Analogously, the local condition in $\text{Sel}_p^-(E/F_\infty)$ is the same "minus condition" for each v . However, one may consider *mixed* signed Selmer groups, where the local conditions may be different for different v . In this case, Kim's Euler characteristic formula has been generalized by Ahmed and Lim under certain additional hypotheses (see [AL20, Thm. 1.1]).

1.3. The case of modular forms (ordinary at p). Suppose $f(q) = \sum_{n \geq 1} a_n(f)q^n$ is a newform of even weight $k \geq 4$ and level $\Gamma_0(N)$. Let $\mathbb{Q}_f = \mathbb{Q}(a_n(f) \mid n \geq 1)$ be the Hecke field of f . Suppose $p \nmid 2N$ and fix a prime \mathfrak{p} of \mathbb{Q}_f above p . Assume that $a_p(f)$ is a \mathfrak{p} -adic unit (i.e. \mathfrak{p} is ordinary for f) and $a_p(f) \not\equiv 1 \pmod{\mathfrak{p}}$. Following the notations of Longo and Vigni, we write E for the completion of \mathbb{Q}_f at \mathfrak{p} and \mathcal{O}_E for the valuation ring of E . Let V be the self-dual twist of the representation $V_{f,\mathfrak{p}}$ of $G_{\mathbb{Q}}$ attached to f by Deligne; hence $V = V_{f,\mathfrak{p}}(k/2)$. Choose a $G_{\mathbb{Q}}$ -stable \mathcal{O}_E -lattice T of V and set $A_f = V/T$. Let Σ be a finite set of primes of \mathbb{Q} containing p , primes dividing the level N and the archimedean prime. Suppose that the Bloch-Kato Selmer group $\text{Sel}_{\text{BK}}(A_f/\mathbb{Q})$ is finite. Then Greenberg's Selmer group $\text{Sel}_{\text{Gr}}(A_f/K_{\infty})$ over the cyclotomic \mathbb{Z}_p -extension K_{∞} of \mathbb{Q} is cotorsion [LV21, Prop. 4.1]. The reader may consult [LV21, Sec. 3.4] for the definition of the Bloch-Kato and Greenberg Selmer groups. Let \mathcal{F} be the power series generating the Pontryagin dual of $\text{Sel}_{\text{Gr}}(A_f/K_{\infty})$. Vigni and Longo then showed the following Euler characteristic formula (see [LV21, Thm. 1.1]).

$$(1.2) \quad \#(\mathcal{O}_E/\mathcal{F}(0) \cdot \mathcal{O}_E) = \#\text{Sel}_{\text{BK}}(A_f/\mathbb{Q}) \cdot \prod_{v \in \Sigma, v \nmid p} c_v(A_f).$$

Here $c_v(A_f) := [H_{\text{ur}}^1(\mathbb{Q}_\ell, A_f) : H_f^1(\mathbb{Q}_\ell, A_f)]$ (see [LV21, Defn. 3.2]).

Vigni and Longo were actually able to deduce this Euler characteristic result for general ordinary representations satisfying additional hypotheses (see assumption 2.1 of [LV21]).

1.4. The case of modular forms (non-ordinary at p). This is the case missing from the literature and this is precisely what this article addresses. In this case, using the theory of Wach modules, Lei, Loeffler, and Zerbes constructed signed Selmer groups over K_{∞} (see [Lei11], [LLZ10], [LLZ11]). These signed Selmer groups are generally conjectured to be cotorsion and this is known to hold in a large number of cases. Many arithmetic properties of the signed Selmer groups have recently been proved by Hatley and Lei [HL19]. Our main goal in this article is to generalize (1.2) for the characteristic power series obtained from the signed Selmer groups over K_{∞} .

We work with infinitely many twists $A(s)$ of A_f , where s is an integer, and consider the signed Selmer groups $\text{Sel}_1(A(s)/K_{\infty})$ and $\text{Sel}_2(A(s)/K_{\infty})$ (see section 2.2 and the beginning of section 3 for the precise definitions). When they are cotorsion, let $f_{1,s}$ and $f_{2,s}$ generate their Pontryagin duals. Under some mild hypotheses made in [HL19] and some hypotheses on s , our main theorem is (see theorem 3.2)

$$\# \left(\frac{\mathcal{O}_E}{f_{i,s}(0)\mathcal{O}_E} \right) = \#\chi(\Gamma_0, \text{Sel}_i(A(s)/K_{\infty})) = \#\text{Sel}_{\text{BK}}(A(s)/\mathbb{Q}) \cdot \prod_{\ell \in \Sigma, \ell \nmid p} c_{\ell}(A(s)),$$

for $i = 1$ or $i = 2$, where $c_{\ell}(A(s)) := [H_{\text{ur}}^1(\mathbb{Q}_{\ell}, A(s)) : H_f^1(\mathbb{Q}_{\ell}, A(s))]$ is the p -part of the *Tamagawa number* of $A(s)$ at ℓ .

One comment about the hypotheses on s : The first equality holds for all but finitely many, while for the second, we need s to be one of $0, \dots, k - 2$.

The astute reader might have noticed that the Tamagawa numbers are over the primes not dividing p in both our formula and in that of Longo–Vigni’s, in contrast to the elliptic curve case. This is because only over those places the local conditions for the (signed) Selmer groups coincide with Greenberg’s local conditions (and with the usual Bloch-Kato local conditions, again because we are away from p , see section 2.3). The recent techniques of Vigni and Longo [LV21] developed to deal with primes not dividing p in the ordinary case are sufficiently general to come to the rescue in our setting, the non-ordinary case, as well.

So what about the prime p ? Here, the local condition, that for the signed Selmer group $\text{Sel}_i(A/K_\infty)$, is defined as the Tate-local orthogonal complement of the kernel of the signed Coleman map. The arithmetic properties of such local conditions at the prime p has been extensively studied by Hatley and Lei [HL19], and we develop them further to prove a “Mini-Control Lemma” (lemma 3.4), equating local cohomology groups of the form $H^1(\mathbb{Q}_p, -)$ to $H^1(K_{\infty,p}, -)^{\Gamma_0}$ sitting in an exact sequence.

We use this to perform a calculation of the Euler characteristic, and are then in a position to deduce our result relating the characteristic power series of the Pontryagin dual of the signed Selmer groups with that of the cardinality of Bloch-Kato Selmer group over \mathbb{Q} and the Tamagawa factors.

1.5. Outlook. Ponsinet obtained a similar formula for the constant terms of the characteristic power series for the dual signed Selmer groups of abelian varieties with supersingular reduction at p such that the Hodge-Tate weights lie in $[0, 1]$ (see [Pon20, Corollary 2.10]), under various other additional hypotheses (see [Pon20, Sections 1.2 and 1.3]). Proving such a result in a more general setting is currently underway.

ACKNOWLEDGEMENTS AND DEDICATION

We would like to thank Jeffrey Hatley, Antonio Lei and Meng-Fai Lim for answering many questions and for their comments on an earlier draft of this article. We also thank Matteo Longo and Stefano Vigni for their encouragement. The first author is supported by the Institute for Advancing Intelligence (IAI), TCG CREST. The second author is supported by a Simons grant and an NSF grant.

Sujatha started her work in Iwasawa theory by computing an Euler characteristic for abelian varieties (in a noncommutative setting) back in the year 1999 [CS99]. It is a pleasure to dedicate this paper to her; she introduced Iwasawa theory both to the first author of this article, and – through the two books co-authored with John Coates [CS06, CS10] – to the second author.

2. PRELIMINARIES

2.1. Notation. Several results in our paper rely on the results of [HL19]. In order to make it easier for the reader to jump between papers, we follow their notation (which in turn follows mostly that of [Gre99]) as much as possible. Thus, let p be a fixed odd prime. The letter ℓ also denotes a prime. Let $\mathbb{Q}_{p,n} = \mathbb{Q}_p(\mu_{p^n})$, $\mathbb{Q}_\infty = \mathbb{Q}(\mu_{p^\infty})$ and $\Gamma = \text{Gal}(\mathbb{Q}_\infty/\mathbb{Q}) \cong \text{Gal}(\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p) \cong \Gamma_0 \times \Delta$, where $\Gamma_0 \cong \mathbb{Z}_p$ and $\Delta \cong \mathbb{Z}/(p-1)\mathbb{Z}$. Let κ and ω be the restriction of the cyclotomic character χ to Γ_0 and Δ . Then $K_\infty = \mathbb{Q}_\infty^\Delta$ is the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} . In any subfield of \mathbb{Q}_∞ , we write p for its unique prime above p , since p is totally ramified in \mathbb{Q}_∞ .

Let $f = \sum a_n(f)q^n$ be a normalized new cuspidal eigenform of even weight $k \geq 2$, level N and nebentypus ε . We assume that $p \nmid N$, f is non-ordinary at p and $a_n(f)$ is defined over a totally real field for all n . Let E be a finite extension of \mathbb{Q}_p containing $a_n(f)$ for all n , ϖ a uniformizer of E , and V_f the E -linear Galois representation attached to f constructed by Deligne [Del71]. Let T_f be the canonical $G_{\mathbb{Q}}$ -stable \mathcal{O}_E -lattice in V_f defined by Kato [Kat04, Sec. 8.3]. Let $A_f = V_f/T_f(1)$. Note that V_f has Hodge-Tate weights $\{0, 1-k\}$, where the convention is that the Hodge-Tate weight of the cyclotomic character is 1. Let \mathcal{T}_f be the Tate twist $T_f(k-1)$ which has Hodge-Tate weights $\{0, k-1\}$.

We let $\Lambda := \mathcal{O}_E[[\Gamma_0]]$ and let $\Lambda(\Gamma) := \mathcal{O}_E[\Delta][[X]]$, the Iwasawa algebra of Γ . There are certain $\Lambda(\Gamma)$ -morphisms called Coleman maps for each $i = 1, 2$ constructed using the theory of Wach modules,

$$\text{Col}_{f,i} : H_{\text{Iw}}^1(\mathbb{Q}_p, \mathcal{T}_f) \rightarrow \Lambda(\Gamma)$$

(see e.g. [HL19, p. 1264 after eq. (2.1)]). Here, $H_{\text{Iw}}^1(\mathbb{Q}_p, \mathcal{T}_f) := \varprojlim H^1(\mathbb{Q}_{p,n}, \mathcal{T}_f)$. For any finite extension K/\mathbb{Q}_p contained in $\mathbb{Q}_{\infty,p}$, by Tate-duality, there exists the Tate pairing

$$(2.1) \quad H^1(K, \mathcal{T}_f) \times H^1(K, A_f) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p.$$

Upon taking direct and inverse limits, we obtain a pairing

$$H_{\text{Iw}}^1(\mathbb{Q}_p, \mathcal{T}_f) \times H^1(\mathbb{Q}_{\infty,p}, A_f) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p.$$

Let $H_i^1(\mathbb{Q}_{\infty,p}, A_f)$ be the orthogonal complement of $\ker \text{Col}_{f,i}$ under the above pairing. This is the local condition at p which will be used in defining the signed Selmer groups below. When v is a prime of \mathbb{Q}_∞ not above the prime p , the local condition is

$$(2.2) \quad H_i^1(\mathbb{Q}_{\infty,v}, A_f) := H_{\text{un}}^1(\mathbb{Q}_{\infty,v}, A_f),$$

which is the unramified subgroup of $H^1(\mathbb{Q}_{\infty,v}, A_f)$.

2.2. The signed Selmer groups. The signed Selmer groups over \mathbb{Q}_∞ are defined as

$$\mathrm{Sel}_i(A_f/\mathbb{Q}_\infty) = \ker \left(H^1(\mathbb{Q}_\infty, A_f) \rightarrow \prod_v H_{/i}^1(\mathbb{Q}_{\infty,v}, A_f) \right),$$

where v runs through all the places of \mathbb{Q}_∞ and $H_{/i}^1(\mathbb{Q}_{\infty,v}, A_f) := \frac{H^1(\mathbb{Q}_{\infty,v}, A_f)}{H_i^1(\mathbb{Q}_{\infty,v}, A_f)}$.

When K is a subfield of \mathbb{Q}_∞ and v is a prime of K not above p , we can define $H_i^1(K_v, A_f)$ in the same way as in (2.2). When $v \mid p$, $H_i^1(K_v, A_f)$ is defined as the orthogonal complement (under the Tate-pairing (2.1)) of the image of $\ker \mathrm{Col}_{f,i}$ under the natural map $H_{\mathrm{Iw}}^1(\mathbb{Q}_p, \mathcal{T}_f) \rightarrow H^1(K_v, \mathcal{T}_f)$.

Hence the signed Selmer groups are defined as

$$\mathrm{Sel}_i(A_f/K) := \ker \left(H^1(K, A_f) \rightarrow \prod_v H_{/i}^1(K_v, A_f) \right).$$

Given an integer s , let $A_{f,s} = A_f \otimes \chi^s$, and $\mathcal{T}_{f,s} = \mathcal{T}_f \otimes \chi^{-s}$. These twists commute with cohomology, so that one can define twisted the signed Selmer groups $\mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)$ and $\mathrm{Sel}_i(A_{f,s}/K)$ (cf. [HL19, p. 1268 before Remark 2.6]). For $i \in \{1, 2\}$, the signed Selmer groups over the cyclotomic \mathbb{Z}_p -extension K_∞ are then given by

$$\mathrm{Sel}_i(A_{f,s}/K_\infty) \cong \mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)^\Delta \cong \mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)^{\omega^{-s}} \otimes \kappa^s,$$

as $\Lambda = \mathcal{O}_E[[\Gamma_0]]$ -modules, see [HL19, Remark 2.6]. Here $\mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)^{\omega^{-s}}$ is the ω^{-s} -isotypic component of $\mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)$. If θ is a character of Δ , then the θ -isotypic component of $\mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)$ is $\mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)^\theta$ and we have the following isomorphisms of Λ -modules.

$$\mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)^\theta \cong \mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)^{\theta\omega^{-s}} \otimes \kappa^s \cong \mathrm{Sel}_i(A_{f,s}(\theta^{-1})/K_\infty)$$

(See again [HL19, a little further down in Rem. 2.6]). We record the following hypotheses also made in [HL19].

(irred): The $G_{\mathbb{Q}}$ -representation $T_f/\varpi T_f$ is irreducible.

(inv): For all $m \geq 0$, $A_f(m)^{G_{\mathbb{Q}_\infty.p}} = 0$.

(tor): For any character θ of Δ , the Selmer groups $\mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)^\theta$ are both cotorsion over Λ for $i = 1, 2$ ¹.

These hypotheses are known to hold in many cases. The following table gives conditions when they are satisfied with references.

¹In [HL19], the statement is about $\mathrm{Sel}_i(A_{f,s}/\mathbb{Q}_\infty)^\theta$ for any fixed integer s , which is equivalent to our hypothesis thanks to [HL19, Remark 2.6], noting that twisting by the cyclotomic character does not change Λ -cotorsion-ness of a module.

Hypothesis	Sufficient conditions	References
(irred)	(holds for almost all primes: $p > k$ and $a_q(f) \not\equiv 1 + q^{k-1} \pmod{\varpi}$ for some prime $q \equiv 1 \pmod{N}$)	[DFG04, Proof of Lem. 2.4], cf. [HL19, Lem. 7.1]
(inv)	$k \leq p$	[Lei11, Lem. 4.4]
(tor)	$a_p = 0$ or $k \geq 3$	[HL19, Start of Sec. 7, p. 1289]

TABLE 1.

The references when the hypothesis **(tor)** is known to hold are [Kob03, Thm. 7.3], [Lei11, Prop. 6.4], [LLZ10, Thm. 6.5], [Spr12, Thm. 7.14].

2.3. The Bloch-Kato Selmer groups. Let V be a \mathbb{Q}_p -vector space defined with a $G_{\mathbb{Q}}$ -action and T be a \mathbb{Z}_p -stable lattice of V . Let us suppose that $\ell \neq p$. Recall that for $* \in \{V, V/T\}$ the unramified local cohomology group is defined as

$$H_{\text{ur}}^1(\mathbb{Q}_{\ell}, *) := \ker(H^1(\mathbb{Q}_{\ell}, *) \rightarrow H^1(I_{\ell}, *)),$$

where I_{ℓ} is the inertia subgroup at ℓ .

Then the Bloch-Kato local conditions are defined as

$$H_f^1(\mathbb{Q}_{\ell}, V) = \begin{cases} H_{\text{ur}}^1(\mathbb{Q}_{\ell}, V) & \ell \neq p, \\ \ker(H^1(\mathbb{Q}_{\ell}, V) \rightarrow H^1(\mathbb{Q}_{\ell}, V \otimes \mathbb{B}_{\text{cris}})) & \ell = p, \end{cases}$$

where \mathbb{B}_{cris} is the Fontaine's crystalline period ring. We also define $H_f^1(\mathbb{Q}_{\ell}, T)$ (resp. $H_f^1(\mathbb{Q}_{\ell}, V/T)$) as the preimage (resp. image) of $H_f^1(\mathbb{Q}_{\ell}, V)$ under the natural inclusion map (resp. projection map). In the $\ell \neq p$ case, we have the following diagram (see e.g [LV19, Section 3.1])

$$(2.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_{\text{ur}}^1(\mathbb{Q}_{\ell}, V/T) & \longrightarrow & H^1(\mathbb{Q}_{\ell}, V/T) & \longrightarrow & H^1(I_{\ell}, V/T) \\ & & & & \uparrow & & \uparrow \\ 0 & \longrightarrow & H_{\text{ur}}^1(\mathbb{Q}_{\ell}, V) & \longrightarrow & H^1(\mathbb{Q}_{\ell}, V) & \longrightarrow & H^1(I_{\ell}, V), \end{array}$$

which shows that $H_f^1(\mathbb{Q}_{\ell}, V/T) \subset H_{\text{ur}}^1(\mathbb{Q}_{\ell}, V/T)$. The index

$$c_{\ell}(V/T) := [H_{\text{ur}}^1(\mathbb{Q}_{\ell}, V/T) : H_f^1(\mathbb{Q}_{\ell}, V/T)]$$

is finite (see [Rub14, Lemma 1.3.5] or [LV21, Lemma 3.1]) and is defined as the p -part of the Tamagawa number of V/T at ℓ .

Let Σ be a finite set of places of \mathbb{Q} containing the prime p , all the places that divide the level N , and the archimedean prime. Let \mathbb{Q}_{Σ} be the maximal extension

of \mathbb{Q} unramified outside Σ . The Bloch-Kato Selmer group is defined as

$$\mathrm{Sel}_{\mathrm{BK}}(A_{f,s}/\mathbb{Q}) := \ker \left(H^1(\mathbb{Q}_\Sigma/\mathbb{Q}, A_{f,s}) \rightarrow \prod_{\ell \in \Sigma} H_{/f}^1(\mathbb{Q}_\ell, A_{f,s}) \right),$$

where

$$H_{/f}^1(\mathbb{Q}_\ell, A_{f,s}) := \frac{H^1(\mathbb{Q}_\ell, A_{f,s})}{H_f^1(\mathbb{Q}_\ell, A_{f,s})}.$$

To conclude this section, we recall an important result of Greenberg.

Lemma 2.1 (Greenberg's Lemma). *Let M be a cofinitely generated, cotorsion module over the Iwasawa algebra $\mathcal{O}[[T]]$, where \mathcal{O} is the ring of integers of any finite degree extension of \mathbb{Q}_p . Let $f(T)$ be a generator of the characteristic ideal of the Pontryagin dual of M . Assume that M^Γ is finite. Then M_Γ is finite, $f(0) \neq 0$, and*

$$f(0) \sim |M^\Gamma|/|M_\Gamma| = \chi(M)$$

The quantity on the right is the *Euler characteristic* of M , which will be made precise in the main section.

Proof. When $\mathcal{O} = \mathbb{Z}_p$, this is [Gre89, Lemma 4.2]. To make [Gre99, Proof of Lemma 4.2] work for general \mathcal{O} , notice that all you have to do is replace the first appearance of ' \mathbb{Z}_p ' by ' \mathcal{O} .' \square

3. MAIN RESULT

Let s be any integer and θ be a character of Δ . The Pontryagin dual of $A_f(\kappa^s \theta^{-1})$ is $\mathcal{T}_f(\kappa^{-s} \theta) = \mathcal{T}_{f,s}(\theta \omega^s)$. Let $\mathcal{A}_{f,s} := \mathcal{T}_{f,s} \otimes E/\mathcal{O}_E = A_{f,k-s-2}$. We will write

$$A(s) := \mathcal{A}_{f,s}(\theta \omega^s).$$

Note that we slightly differ from the notation of [HL19, Proof of Prop. 3.3, line -1 in page 1274] where it is denoted as A , since they fix s .

We define the Γ_0 -Euler characteristic of $\mathrm{Sel}_i(A(s)/K_\infty)$ as

$$\chi(\Gamma_0, \mathrm{Sel}_i(A(s)/K_\infty)) = \frac{\#H^0(\Gamma_0, \mathrm{Sel}_i(A(s)/K_\infty))}{\#H^1(\Gamma_0, \mathrm{Sel}_i(A(s)/K_\infty))},$$

whenever the quantities are finite.

When $\mathrm{Sel}_i(A(s)/K_\infty)$ is Λ -cotorsion, let $f_{i,s} \in \Lambda$ be a characteristic power series that generates its Pontryagin dual.

Remark 3.1. The torsion assumption (tor) guarantees that $\mathrm{Sel}_i(A(s)/K_\infty)$ is Λ -cotorsion for all s . This follows from [HL19, Remark 2.6], and noting that twisting a finitely generated Λ -cotorsion module by the cyclotomic character results in a finitely generated Λ -cotorsion module.

Theorem 3.2. *Assume (tor), (inv), and (irred). Choose $i \in \{1, 2\}$. Then for all but finitely many s , $\chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty))$ is well-defined. Moreover, $\text{Sel}_i(A(s)/\mathbb{Q})$ is finite, $f_{i,s}(0) \neq 0$, and*

$$(3.1) \quad \# \left(\frac{\mathcal{O}_E}{f_{i,s}(0)\mathcal{O}_E} \right) = \chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty)) = \#\text{Sel}_i(A(s)/\mathbb{Q}) \cdot \prod_{\ell \in \Sigma, \ell \neq p} c_\ell(A(s)).$$

Further, suppose that $s \in \{0, \dots, k-2\}$. If $i = 1$, assume also that $a_p(f) \neq \varepsilon(p)p^s + p^{k-s-2}$. Then $\text{Sel}_i(A(s)/\mathbb{Q})$ and $\text{Sel}_{\text{BK}}(A(s)/\mathbb{Q})$ are finite, $f_{i,s}(0) \neq 0$, and

$$(3.2) \quad \# \left(\frac{\mathcal{O}_E}{f_{i,s}(0)\mathcal{O}_E} \right) = \chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty)) = \#\text{Sel}_{\text{BK}}(A(s)/\mathbb{Q}) \cdot \prod_{\ell \in \Sigma, \ell \neq p} c_\ell(A(s)).$$

Here $c_\ell(A(s)) := [H_{\text{ur}}^1(\mathbb{Q}_\ell, A(s)) : H_f^1(\mathbb{Q}_\ell, A(s))]$ is the p -part of the Tamagawa number of $A(s)$ at ℓ .

We record a proposition and two lemmas before proving the theorem.

Proposition 3.3 (Hatley-Lei). *Under the hypotheses (tor), (inv), and (irred), we have for each $i = 1$ or 2 that for all but finitely many integers s :*

- (1) $(\text{Sel}_i(A(s)/K_\infty))^{\Gamma_0}$ is finite,
- (2) The kernel and cokernel of the restriction map

$$\text{Sel}_i(A(s)/\mathbb{Q}) \rightarrow (\text{Sel}_i(A(s)/K_\infty))^{\Gamma_0}$$

are both finite.

Lemma 3.4. *Let $\ell \in \Sigma$ and v be any one of the finitely many places of K_∞ over ℓ .*

- (1) *In the commutative diagram below, h is an isomorphism and ρ is surjective.*
- (2) *In the case $\ell \neq p$, $\#\ker g_{/i,\ell} = c_\ell(A(s))$.*

$$(3.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Sel}_i(A(s)/\mathbb{Q}) & \longrightarrow & H^1(\mathbb{Q}_\Sigma/\mathbb{Q}, A(s)) & \xrightarrow{\rho} & \prod_{\ell \in \Sigma} H_{/i}^1(\mathbb{Q}_\ell, A(s)) \\ & & \downarrow a & & \downarrow h & & \downarrow g = \prod g_{/i,\ell} \\ 0 & \longrightarrow & \text{Sel}_i(A(s)/K_\infty)^{\Gamma_0} & \longrightarrow & H^1(\mathbb{Q}_\Sigma/K_\infty, A(s))^{\Gamma_0} & \longrightarrow & \prod_{v|\ell, \ell \in \Sigma} H_{/i}^1(K_{\infty,v}, A(s))^{\Gamma_0} \end{array}$$

Lemma 3.5 (Mini-Control Lemma). *The vertical maps in the commutative diagram below are isomorphisms:*

(3.4)

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_i^1(\mathbb{Q}_p, A(s)) & \longrightarrow & H^1(\mathbb{Q}_p, A(s)) & \xrightarrow{c} & H_{/i}^1(\mathbb{Q}_p, A(s)) \\
& & \downarrow g_{i,p} & & \downarrow g_p & & \downarrow g_{/i,p} \\
0 & \longrightarrow & H_i^1(K_{\infty,p}, A(s))^{\Gamma_0} & \longrightarrow & H^1(K_{\infty,p}, A(s))^{\Gamma_0} & \longrightarrow & H_{/i}^1(K_{\infty,p}, A(s))^{\Gamma_0}
\end{array}$$

Proof of Theorem 3.2. The assumption **(tor)** gives us that the Euler characteristic $\chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty))$ is well-defined: Indeed, proposition 3.3(1) shows that $\text{Sel}_i(A(s)/K_\infty)^{\Gamma_0}$ is finite. Hence by lemma 2.1, we know that $\text{Sel}_i(A(s)/K_\infty)_{\Gamma_0}$ is finite. Since $\Gamma_0 \cong \mathbb{Z}_p$, we can identify $H^1(\Gamma_0, \text{Sel}_i(A(s)/K_\infty))$ with $\text{Sel}_i(A(s)/K_\infty)_{\Gamma_0}$ and so the Euler characteristic equals

$$\chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty)) = \frac{\#\text{Sel}_i(A(s)/K_\infty)^{\Gamma_0}}{\#\text{Sel}_i(A(s)/K_\infty)_{\Gamma_0}}.$$

For the main part of the proof, choose s so that $\text{Sel}_i(A(s)/K_\infty)^{\Gamma_0}$ is finite. This only excludes finitely many s by proposition 3.3. Applying the snake lemma to diagram (3.3), we obtain from lemma 3.4(1) that $\text{coker } a$ is finite, $\text{ker } a$ is trivial, and $\text{Sel}_i(A(s)/\mathbb{Q})$ is finite. From lemma 3.4(2) and lemma 3.5, we then have

$$\#\text{Sel}_i(A(s)/K_\infty)^{\Gamma_0} = \#\text{Sel}_i(A(s)/\mathbb{Q}) \cdot \prod_{\ell \in \Sigma, \ell \neq p} c_\ell(A(s)).$$

But the left-hand side is the Euler characteristic, since from [HL19, Proof of Theorem 3.1, very end of Section 3], $\text{Sel}_i(A(s)/K_\infty)_{\Gamma_0} = 0$, i.e. we really have

$$(3.5) \quad \chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty)) = \#\text{Sel}_i(A(s)/\mathbb{Q}) \cdot \prod_{\ell \in \Sigma, \ell \neq p} c_\ell(A(s)),$$

which is the second equality in equation (3.1) of the theorem.

Greenberg's Lemma 2.1 for $M = \text{Sel}_i(A(s)/K_\infty)$ implies $f_{i,s}(0) \neq 0$ and

$$\# \left(\frac{\mathcal{O}_E}{f_{i,s}(0)\mathcal{O}_E} \right) = \chi(\Gamma_0, \text{Sel}_i(A(s)/K_\infty)),$$

giving us the first equality. (Note that this is the supersingular analogue of [LV21, Prop. 4.3]).

To obtain the equalities in equation (3.2), recall from [HL19, Prop. 2.14 and Rem. 2.15] that for $i = 2$ and $s \in \{0, \dots, k-2\}$,

$$\text{Sel}_2(A(s)/\mathbb{Q}) = \text{Sel}_{\text{BK}}(A(s)/\mathbb{Q}),$$

and that if $i = 1$ and if we further assume that $a_p(f) \neq \varepsilon(p)p^s + p^{k-s-2}$, then

$$\mathrm{Sel}_1(A(s)/\mathbb{Q}) \cong \mathrm{Sel}_{\mathrm{BK}}(A(s)/\mathbb{Q}).$$

□

Remark 3.6. In the last part of the proof of Theorem 3.2, we needed to restrict s in the range $\{0, \dots, k-2\}$ because this is needed in the proof of [HL19, Prop. 2.12]. In their proof they first use a result from [LZ14] where they need s to be nonnegative. Later in their proof they use a result from [LLZ17] where they need s to be in the range $\{0, \dots, k-2\}$ so that the twisted representation has appropriate Hodge-Tate weights².

3.1. Proofs of Proposition 3.3 , lemma 3.4, and lemma 3.5.

Proof of Proposition 3.3. In [HL19, Prop. 3.3], the authors prove the statements in the proposition hold for some integer s . But a closer look at their proof reveals that the same is true for all but finitely many s .

Here is a guide to their notation: In the proof of [HL19, Prop. 3.3], the authors set $\eta = \theta^{-1}\omega^{2-k}$ (independent of s), and write $\mathrm{Sel}_i(A(s)/K_\infty)$ as $\mathrm{Sel}_i(A_f/\mathbb{Q}_\infty)^\eta \otimes \kappa^{k-s-2}$, working with the following isomorphism of Λ -modules.

$$\mathrm{Sel}_i(A(s)/K_\infty) = \mathrm{Sel}_i(\mathcal{A}_{f,s}(\theta\omega^s)/K_\infty) \cong \mathrm{Sel}_i(A_f(\theta\omega^{k-2})/K_\infty) \otimes \kappa^{k-s-2}.$$

□

Proof of Lemma 3.4. $\ker(h)$ and $\mathrm{coker}(h)$ can be regarded as inside the inflation-restriction exact sequence, and are zero because of hypothesis **(inv)**. (Cf. also [HM99, Lemma 3.3].)

The surjectivity of ρ follows from [HL19, Prop. 3.3] and its proof.

The Greenberg local condition at v (see [LV21, Sec. 3.4.2]) coincides with the Bloch–Kato local condition defined in section 2.3. Hence, by [LV21, Lemma 5.3], we obtain

$$\#\ker g_{/i,\ell} = c_\ell(A(s)),$$

which is indeed finite by [LV21, Lemma 3.1]. (It is also interesting to note that if $v \notin \Sigma$ the Tamagawa factors are trivial (see [LV21, Lemma 3.3])). □

Proof of Lemma 3.5. The kernel of the map g_p is $H^1(K_{\infty,p}/\mathbb{Q}_p, A(s)^{G_{K_{\infty,p}}})$ and its cokernel maps into $H^2(K_{\infty,p}/\mathbb{Q}_p, A(s)^{G_{K_{\infty,p}}})$ which are both trivial because of our hypothesis **(inv)**. Hence the map g_p is an isomorphism.

²We thank Jeffrey Hatley for pointing this out to us.

Next, we analyze the kernel and the cokernel of the map $g_{/i,p}$.

We have the following isomorphism by duality

$$H_i^1(K_{\infty,p}, A(s)) = (\mathrm{Im}(\mathrm{Col}_{f,i})^\theta \otimes \kappa^{-s})^\vee.$$

Since $\mathrm{Im}(\mathrm{Col}_{f,i})^\theta \subset \Lambda$, it implies that

$$(\mathrm{Im}(\mathrm{Col}_{f,i})^\theta \otimes \kappa^{-s})^{\Gamma_0} = 0.$$

Hence we have

$$(3.6) \quad H_i^1(K_{\infty,p}, A(s))_{\Gamma_0} = 0.$$

Consider the short exact sequence

$$0 \rightarrow H_i^1(K_{\infty,p}, A(s)) \rightarrow H^1(K_{\infty,p}, A(s)) \rightarrow H_{/i}^1(K_{\infty,p}, A(s)) \rightarrow 0.$$

Combining (3.6) with the isomorphism

$$H_i^1(K_{\infty,p}, A(s))_{\Gamma_0} \cong H^1(\Gamma_0, H_i^1(K_{\infty,p}, A(s))),$$

(see e.g. [NSW00, Prop. 1.7.7]), we obtain

$$H_{/i}^1(K_{\infty,p}, A(s))^{\Gamma_0} \cong \frac{H^1(K_{\infty,p}, A(s))^{\Gamma_0}}{H_i^1(K_{\infty,p}, A(s))^{\Gamma_0}},$$

as in p. 1276 of [HL19].

But it follows from [HL19, p. 1276, line 10] that

$$\frac{H^1(K_{\infty,p}, A(s))^{\Gamma_0}}{H_i^1(K_{\infty,p}, A(s))^{\Gamma_0}} \cong H_{/i}^1(\mathbb{Q}_p, A(s))$$

using the Hochschild-Serre spectral sequence and the facts that

$$\begin{aligned} H^1(K_{\infty,p}, A(s))^{\mathrm{Gal}(K_{\infty,p}/\mathbb{Q}_p)} &= H^1(\mathbb{Q}_p, A(s)), \\ H_i^1(K_{\infty,p}, A(s))^{\mathrm{Gal}(K_{\infty,p}/\mathbb{Q}_p)} &= H_i^1(\mathbb{Q}_p, A(s)), \end{aligned}$$

cf. [HL19, Remark 2.5]. Finally, we obtain from this that

$$H_{/i}^1(K_{\infty,p}, A(s))^{\Gamma_0} \cong H_{/i}^1(\mathbb{Q}_p, A(s)),$$

which shows that the map $g_{/i,p}$ is also an isomorphism.

The map c is surjective by definition, and hence by the snake lemma we obtain that the map $g_{i,p}$ is also an isomorphism. (This generalizes an analogous property that has been proved by Ahmed and Lim in the case of elliptic curves (see [AL20, Lemma 2.6])). \square

REFERENCES

- [AL20] Suman Ahmed and Meng Fai Lim. On the Euler characteristics of signed Selmer groups. *Bulletin of the Australian Mathematical Society*, 101(2):238–246, 2020.
- [CS99] John Coates and Ramdorai Sujatha. Euler-Poincaré characteristics of abelian varieties. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, 329(4):309–313, 1999.
- [CS06] J. Coates and R. Sujatha. *Cyclotomic fields and zeta values*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2006.
- [CS10] John Coates and Ramdorai Sujatha. *Galois cohomology of elliptic curves*. Published by Narosa Publishing House, New Delhi; for the Tata Institute of Fundamental Research, Mumbai, second edition, 2010.
- [Del71] Pierre Deligne. Formes modulaires et représentations ℓ -adiques. In *Séminaire Bourbaki vol. 1968/69 Exposés 347-363*, pages 139–172. Springer, 1971.
- [DFG04] Fred Diamond, Matthias Flach, and Li Guo. The Tamagawa number conjecture of adjoint motives of modular forms. *Ann. Sci. École Norm. Sup. (4)*, 37(5):663–727, 2004.
- [Gre89] Ralph Greenberg. Iwasawa theory for p -adic representations. In *Algebraic number theory*, volume 17 of *Adv. Stud. Pure Math.*, pages 97–137. Academic Press, Boston, MA, 1989.
- [Gre99] Ralph Greenberg. Iwasawa theory for elliptic curves. In *Arithmetic theory of elliptic curves (Cetraro, 1997)*, volume 1716 of *Lecture Notes in Math.*, pages 51–144. Springer, Berlin, 1999.
- [HL19] Jeffrey Hatley and Antonio Lei. Arithmetic properties of signed Selmer groups at non-ordinary primes. *Annales de l'Institut Fourier*, 69(3):1259–1294, 2019.
- [HM99] Yoshitaka Hachimori and Kazuo Matsuno. An analogue of Kida's formula for the Selmer groups of elliptic curves. *J. Algebraic Geom.*, 8(3):581–601, 1999.
- [Kat04] Kazuya Kato. p -adic Hodge theory and values of zeta functions of modular forms. *Astérisque*, (295):ix, 117–290, 2004. Cohomologies p -adiques et applications arithmétiques. III.
- [Kim13] Byoung Du Kim. The plus/minus Selmer groups for supersingular primes. *Journal of the Australian Mathematical Society*, 95(2):189–200, 2013.
- [Kob03] Shin-ichi Kobayashi. Iwasawa theory for elliptic curves at supersingular primes. *Invent. Math.*, 152(1):1–36, 2003.
- [Lei11] Antonio Lei. Iwasawa theory for modular forms at supersingular primes. *Compos. Math.*, 147(3):803–838, 2011.
- [LLZ10] Antonio Lei, David Loeffler, and Sarah Livia Zerbes. Wach modules and Iwasawa theory for modular forms. *Asian J. Math.*, 14(4):475–528, 2010.
- [LLZ11] Antonio Lei, David Loeffler, and Sarah Livia Zerbes. Coleman maps and the p -adic regulator. *Algebra Number Theory*, 5(8):1095–1131, 2011.
- [LLZ17] Antonio Lei, David Loeffler, and Sarah Livia Zerbes. On the asymptotic growth of Bloch-Kato-Shafarevich-Tate groups of modular forms over cyclotomic extensions. *Canadian Journal of Mathematics*, 69(4):826–850, 2017.
- [LV19] Matteo Longo and Stefano Vigni. Plus/minus Heegner points and Iwasawa theory of elliptic curves at supersingular primes. *Bollettino dell'Unione Matematica Italiana*, 12(3):315–347, 2019.
- [LV21] Matteo Longo and Stefano Vigni. On Bloch-Kato Selmer groups and Iwasawa theory of p -adic Galois representations. *New York Journal of Mathematics*, 27:437–467, 2021.
- [LZ14] David Loeffler and Sarah Livia Zerbes. Iwasawa theory and p -adic L -functions over \mathbb{Z}_p^2 -extensions. *International Journal of Number Theory*, 10(08):2045–2095, 2014.
- [NSW00] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg. *Cohomology of number fields*, volume 323 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2000.

- [Pon20] Gautier Ponsinet. On the structure of signed Selmer groups. *Mathematische Zeitschrift*, 294(3):1635–1658, 2020.
- [Rub14] Karl Rubin. Euler Systems.(am-147), volume 147. In *Euler Systems.(AM-147)*, Volume 147. Princeton University Press, 2014.
- [Spr12] Florian E. Ito Sprung. Iwasawa theory for elliptic curves at supersingular primes: a pair of main conjectures. *J. Number Theory*, 132(7):1483–1506, 2012.
- [Spr16] Florian E. Ito Sprung. The Iwasawa main conjecture for odd supersingular primes. preprint, arxiv: 1610.10017, 2016.

(Ray) TCG CENTRES FOR RESEARCH AND EDUCATION IN SCIENCE AND TECHNOLOGY, 1ST FLOOR, TOWER 1, BENGAL ECO INTELLIGENT PARK (TECHNA BUILDING), BLOCK EM, PLOT NO 3, SECTOR V, SALT LAKE, KOLKATA 700091

Email address: jishnu.ray@tcgcrest.org; jishnuray1992@gmail.com

(Sprung) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY TEMPE, AZ 85287-1804, USA

Email address: florian.sprung@asu.edu