

# Fibered Universal Algebra for First-Order Logics

Colin Bloomfield, Yoshihiro Maruyama

August 9, 2022

## Abstract

We extend Lawvere-Pitts prop-categories (aka. hyperdoctrines) to develop a general framework for providing “algebraic” semantics for non-classical first-order logics. This framework includes a natural notion of substitution, which allows first-order logics to be considered as structural closure operators just as propositional logics are in abstract algebraic logic. We then establish an extension of the homomorphism theorem from universal algebra for generalized prop-categories and characterize two natural closure operators on the prop-categorical semantics. The first closes a class of structures (which are interpreted as morphisms of prop-categories) under the satisfaction of their common first-order theory and the second closes a class of prop-categories under their associated first-order consequence. It turns out these closure operators have characterizations that closely mirror Birkhoff’s characterization of the closure of a class of algebras under the satisfaction of their common equational theory and Blok and Jónsson’s characterization of closure under equational consequence, respectively. These “algebraic” characterizations of the first-order closure operators are unique to the prop-categorical semantics and do not have analogs, for example, in the Tarskian semantics for classical first-order logic. The prop-categories we consider are much more general than traditional intuitionistic prop-categories or triposes (i.e., topos representing indexed partially ordered sets). Nonetheless, to our knowledge, our results are still new even when restricted to these special classes of prop-categories.

## 1 Introduction

There is a vast and ever-growing expanse of nonclassical logics and with it, a need to develop a unified theoretical framework to classify, prove metalogical results, and develop semantics for *classes* of logics. For propositional logics, the field of abstract algebraic logic (AAL) provides a unified theoretical framework rooted in universal algebra. However, for first-order logics, there is not a clear consensus on what the unifying algebraic perspective should be [6, p. 104]. Two main challenges mentioned in the literature are (1) a lack of *general* algebraic approaches to providing semantics for first-order logics and (2) a lack of a notion of formula substitution which preserves logical consequence [6, p. 106].

To address (1), we use Lawvere’s theory of hyperdoctrines [13] to develop a general approach to providing “algebraic” semantics for nonclassical first-order logics. The key insight to this approach is to consider terms and formulas *in context*, that is with a list of variable/sort pairs that at least include those occurring in the formula. The use of contexts is ubiquitous in classical first-order model theory where it allows one to unambiguously interpret terms as term functions and formulas as definable sets. However, it is scarcely formally presented<sup>1</sup>, which is unfortunate because contexts allow one to give an elegant point-free/valuation-free definition of satisfaction.<sup>2</sup>

Since we model truth and not proof, we use a special class of hyperdoctrines for our semantics called prop-categories by Pitts [18]. A prop-category is a pair  $(\mathcal{C}, P)$ , where  $\mathcal{C}$  is a category with designated finite products and  $P: \mathcal{C} \rightarrow \mathbf{Pos}$  is a contravariant functor to  $\mathbf{Pos}$ , the category of partially ordered sets and monotone maps. The prop-categories in [18] are assumed to satisfy additional adjoint conditions so that they interpret the quantifiers and connectives of intuitionistic predicate logic. To model a broad variety of logics, we allow arbitrarily many quantifier and operation symbols in a first-order language  $\mathcal{L}$  and we replace the adjoint conditions with weaker algebraic ones. Despite this generalization, the prop-categorical semantics of a logic can still be encoded in a 2-category of prop-categories we denote  $\mathbf{FA}_{\mathcal{L}}$  (or just  $\mathbf{FA}$ ), where roughly theories are objects, structures are morphisms, and structure-preserving maps are 2-cells.

Moreover, the morphisms in  $\mathbf{FA}$  determine a natural action of formula substitution which also addresses issue (2). A footnote in a standard reference on AAL [6, p. 106] says:

In first-order languages we replace an individual (free) variable by a term; we do not replace an atomic formula, inside a more complex one, by and an arbitrary formula, which is what is done in a sentential language.

However, if one unpacks the action of morphisms on theories it does exactly that: atomic formulas are uniformly replaced by arbitrary formulas. Despite being an elementary property of classical first-order logic, Church’s Introduction to Mathematical Logic [5, p. 191] is the only textbook account we could find. Already, for first-order logic without function symbols, the account in [5] which does not use contexts is complicated and so this notion of formula substitution is even easier to overlook without contexts when considering first-order logics with function symbols.

Much like the study of algebraic semantics for first-order logic, currently, the study of prop-categorical semantics for logics (in contrast with type theories)

---

<sup>1</sup>The usual justification is that the formal presentation is trivial enough to be omitted, and the extension of the syntax is too burdensome.

<sup>2</sup>In Philosophy and Model Theory [4, p. 10], Tarski’s approach via valuations is presented and the philosophical issues of assigning objects to variables is discussed. In order to mend this, an alternative semantics due to Robinson is considered, which is also criticized for requiring one to flood the signature with additional constant symbols. To resolve these philosophical issues the authors adopt an approach which is a compromise between the two. However, Lawvere’s semantics (which is not discussed) avoids these issues.

appears to be focused on studying particular logics and not on establishing a general framework in the spirit of AAL. An early exception is [21] (also see [20]) which provides complete prop-categorical semantics for substructural predicate logics, and uses this semantics to prove their disjunctive and existence properties. In [15], complete hyperdoctrine semantics for substructural predicate logics is also given and used to present a unified account of logical translations, including Girard’s exponential translation in linear logic and Kolmogorov’s double negation translation in intuitionistic logic. In contrast to these existing works on general prop-categorical semantics we model nonclassical quantifiers and focus on (1) establishing minimal conditions on prop-categories and first-order logics so that structures are morphisms in a 2-category of prop-categories and (2) developing a toolkit of general algebraic/categorical results on the resulting prop-categorical semantics.

Towards (1), we define a weak logic  $\mathcal{L}^m$  such that whenever a logic  $\mathcal{L}$  is stronger than  $\mathcal{L}^m$ , each  $\mathcal{L}$ -theory  $T$  defines a classifying prop-category  $(\mathcal{C}_T, P_T)$  in  $\mathbf{FA}$ , which contains a generic  $T$ -model  $G$  in  $(\mathcal{C}_T, P_T)$ . Such a logic  $\mathcal{L}$  is called *adequate* and  $\mathcal{L} \supseteq \mathcal{L}^m$  is also necessary for  $\mathcal{L}$  to be adequate. Moreover,  $\mathcal{L}^m$  has a complete semantics with respect to the class of all of its classifying prop-categories (Theorem 2), as do many natural extensions of  $\mathcal{L}^m$  by the addition of various sequent rules (Theorem 3). We then develop an algebraic view of the prop-categorical semantics of a logic  $\mathcal{L}$  in analogy with the algebraic approach to providing semantics for propositional logics. In this view, a full sub-2-category  $\mathbf{FA}_{\mathcal{L}}$  of  $\mathbf{FA}$  replaces the usual quasivariety of algebras forming the algebraic semantics of a propositional logic and we define the kernel of a morphism (Definition 5.1) to give an algebraic/morphic view of logical entailment. These results are directly applied to formulate and prove the later “fibered” universal algebraic results.

After singling out the class of first-order logics which have nice categorical semantics, in the second part of the paper, we develop a “fibered universal algebra” for  $\mathbf{FA}$ . The internal logic of a prop-category  $(\mathcal{C}, P)$  in  $\mathbf{FA}$  is defined and shown to be an  $\mathcal{L}^m$ -theory  $T$ , such that  $(\mathcal{C}_T, P_T) \equiv (\mathcal{C}, P)$  in  $\mathbf{FA}$ . Subprop-categories and morphic images of prop-categories are defined and  $\mathbf{FA}$  is shown to have arbitrary products. The logical properties of products and subprop-categories are considered which mirror the logical properties of submatrices and products of matrices respectively in the matrix semantics of AAL [6, p. 183].

We then prove a “fibered” analogue of the homomorphism theorem from universal algebra in two parts (Theorem 6). The first part, says that each morphism  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  in  $\mathbf{FA}_{\mathcal{L}}$  can be factored through a “quotient” prop-category which is also in  $\mathbf{FA}_{\mathcal{L}}$ . The second part characterizes when, given morphisms  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  and  $K: (\mathcal{C}, P) \rightarrow (\mathcal{E}, R)$ , there exists a unique morphism  $H: (\mathcal{E}, R) \rightarrow (\mathcal{D}, Q)$  such that  $H \circ K = F$ . Part 2, says that under reasonable “surjectivity” assumptions on  $K$ , there is a unique completion  $H$  iff  $\ker K \leq \ker F$ . Our results on morphic images and the internal logic allow an intuitively clear “algebraic” proofs of these results. Moreover, the fibered homomorphism theorem, determines an orthogonal factorization system for  $\mathbf{FA}_{\mathcal{L}}$ .

In the final section, we use the fibered homomorphism theorem to character-

ize two natural closure operators. The first operator characterizes the closure of a class of structures under the satisfaction of their common first-order theory. Taking submodels, homomorphic images and products does not preserve the satisfaction of theories in the Tarskian semantics for classical first-order logic. However, for logics  $\mathcal{L}$  with prop-categorical semantics in  $\mathbf{FA}_{\mathcal{L}}$ , the closure operator is characterized by  $\mathbb{H}\mathbb{S}\mathbb{P}$  where  $\mathbb{H}$ ,  $\mathbb{S}$  and  $\mathbb{P}$  close a collection of structures under natural notions of homomorphic image, submodel and product of structures in the prop-categorical semantics (Theorem 7).

The second operator closes a collection  $\mathcal{X} \subseteq \text{Ob}(\mathbf{FA})$  under logical consequence: that is, the closure of  $\mathcal{X}$  is the largest  $\mathcal{Y} \supseteq \mathcal{X}$  such that  $\models_{\mathcal{Y}} = \models_{\mathcal{X}}$ , i.e. they define the same logic. Since  $\mathbf{FA}_{\mathcal{L}}$  is stable under products and subprop-categories,  $\mathcal{Y} \supseteq \mathbb{S}\mathbb{P}(\mathcal{X})$ , where here  $\mathbb{S}$  is closure under subprop-categories and  $\mathbb{P}$  is closure under products of prop-categories. It turns out that we need another operation which we call  $\mathbb{U}$ , and Theorem 9 shows that  $\mathcal{Y} = \mathbb{U}\mathbb{S}\mathbb{P}(\mathcal{X})$ . This result is a natural extension of a corresponding result by Blok and Jónsson in [3], which shows that for a collection of algebras  $\mathcal{V}$ , closure under the equational consequence is given by  $\mathbb{U}_{\lambda}\mathbb{S}\mathbb{P}(\mathcal{V})$ , where  $\lambda$  is the cardinality of the set of variables over which equations are defined, and  $B \in \mathbb{U}_{\lambda}(\mathcal{V})$  if every  $\lambda$ -generated subalgebra  $A \leq B$  is in  $\mathcal{V}$ . (See Theorem 8 for the directly analogous “fibered” result which considers a fixed first-order signature and uses an operator  $\mathbb{U}_{Sg}$ .)

## 2 First-Order Logics

This section begins by presenting the first-order syntax using *contexts*. Contexts allow us to unambiguously interpret well-formed logical expressions in prop-categories which are introduced in the next section. After presenting typed equational logic the section ends by defining a theories, first-order logics and derivability.

Throughout the paper, we fix a **first-order language**  $\mathcal{L}$ , which consists of a collection  $\mathcal{L}_q$  of quantifier symbols and a collection  $\mathcal{L}_{\omega}$  of propositional connectives each with a designated arity in  $\omega$ . Let  $\mathcal{L}_n$  denote the propositional connectives of arity  $n \in \omega$  in  $\mathcal{L}_{\omega}$ . We consider multi-sorted first-order logic, and so a **signature**  $Sg$  is specified by a collection of sort symbols  $\sigma, \tau, \gamma, \dots$ , typed function symbols  $f: \sigma_1, \dots, \sigma_n \rightarrow \tau$ , (we write  $f: \tau$  for  $n = 0$ ), and typed relation symbols  $R \subseteq \sigma_1, \dots, \sigma_n$ <sup>3</sup>. (we write  $R \subseteq \emptyset$  for  $n = 0$ )

To provide prop-categorical semantics, well-formed terms and formulas must be “in context”. A **context**  $\Gamma = [x_1 : \sigma_1, \dots, x_n : \sigma_n]$  is a list of variables,  $x_1, \dots, x_n$  where each variable  $x_i$  is assigned a type (sort symbol)  $\sigma_i$ . We let  $M : \tau [\Gamma]$  be the assertion that the term  $M$  is of type  $\tau$  and  $\Gamma$  is a valid context for  $M$ . The (well-formed) terms-in-context are defined inductively via the following typing rules:

$$\frac{}{x : \sigma [\Gamma, x : \sigma, \Gamma']} \quad \frac{M_1 : \sigma_1 [\Gamma] \quad \dots \quad M_n : \sigma_n [\Gamma]}{f(M_1, \dots, M_n) : \tau [\Gamma]},$$

<sup>3</sup> $f: \sigma_1, \dots, \sigma_n \rightarrow \tau$  and  $R \subseteq \sigma_1, \dots, \sigma_n$  are merely typing assignments and do not assert that  $f$  is a morphism or  $R$  is a subset of some set.

for each variable  $x$  and function symbol  $f: \sigma_1, \dots, \sigma_n \rightarrow \tau$  in  $Sg$ . These are the only typing rules for terms, and from these rules alone, one may prove that the typing rule

$$\frac{M : \sigma [\Delta] \quad N : \tau [\Gamma, x : \sigma, \Gamma']}{N[M/x] : \tau [\Gamma, \Gamma']} \quad (1)$$

is admissible, where  $N[M/x]$  denotes the operation of substituting each occurrence of  $x$  in  $N$  with  $M$  and the variable sort pairs in  $\Delta$  are contained in the set of those in  $\Gamma, \Gamma'$ . We denote this by  $\text{VS}(\Delta) \subseteq \text{VS}(\Gamma, \Gamma')$ .

An **equation-in-context**  $M_1 = M_2 : \tau [\Gamma]$  is well-formed if  $M_i : \tau [\Gamma]$  holds for  $i \in \{1, 2\}$ . In the sequel, all the logics we consider are built over the usual (typed) equational derivation system. The rules of this system are as follows:

### Equational Logic.

$$\frac{}{M = M : \sigma [\Gamma]} \quad \frac{M = M' : \sigma [\Gamma]}{M' = M : \sigma [\Gamma]} \quad \frac{M = M' : \sigma [\Gamma] \quad M' = M'' : \sigma [\Gamma]}{M = M'' : \sigma [\Gamma]}$$

$$\frac{M = M' : \sigma [\Delta] \quad N = N' : \tau [\Gamma, x : \sigma, \Gamma']}{N[M/x] = N'[M'/x] : \tau [\Gamma, \Gamma']} \text{VS}(\Delta) \subseteq \text{VS}(\Gamma, \Gamma').$$

For a formula  $\phi$ , we let  $\phi : \text{prop} [\Gamma]$  be the assertion that  $\phi [\Gamma]$  is a well-formed formula-in-context. The **atomic** well-formed formulas-in-context are constructed using the rules

$$\frac{M_1 : \sigma_1 [\Gamma] \quad \dots \quad M_n : \sigma_n [\Gamma]}{R(M_1, \dots, M_n) : \text{prop} [\Gamma]} \quad \frac{M_1 : \sigma [\Gamma] \quad M_2 : \sigma [\Gamma]}{M_1 =_{\sigma} M_2 : \text{prop} [\Gamma]},$$

for each  $n$ -ary relation symbol  $R \subseteq \sigma_1, \dots, \sigma_n$ . Note that  $M_1 =_{\sigma} M_2 [\Gamma]$  is considered a formula-in-context whereas  $M_1 = M_2 : \sigma [\Gamma]$  is not. Thus we have two different notions of equality and for many logics,  $=_{\sigma}$  is coarser than  $=$ .

The well-formed formulas-in-context are recursively generated from the atomic ones via the rules

$$\frac{\phi_1 : \text{prop} [\Gamma] \quad \dots \quad \phi_n : \text{prop} [\Gamma]}{\diamond(\phi_1, \dots, \phi_n) : \text{prop} [\Gamma]} \quad \frac{\phi : \text{prop} [\Gamma, x : \sigma]}{\Omega_{x:\sigma}(\phi) : \text{prop} [\Gamma]},$$

for each quantifier symbol  $\Omega \in \mathcal{L}_q$ , all  $n \in \omega$  and each  $n$ -ary propositional connective  $\diamond \in \mathcal{L}_n$ . We consider formulas equal up to  $\alpha$ -equivalence, i.e. up to renaming of bound variables. This allows us to define the action of simultaneous term substitution on ( $\alpha$ -equivalence classes of) formulas that avoids variable capture which we denote by  $\phi[M_1/x_1, \dots, M_n/x_n]$ .

The well-formed **sequents-in-context** are of the form  $\phi_1, \dots, \phi_n \vdash \phi_{n+1} [\Gamma]$ , where  $n \in \omega$  and for each  $i \leq n+1$ ,  $\phi_i : \text{prop} [\Gamma]$ . We call the sequents-in-context and equations-in-context **assertions** and let  $A_{Sg}$  denote the collection of all  $Sg$ -assertions. A **theory**  $T$  is an ordered pair  $T = (\text{Sg}(T), \text{A}(T))$ , where  $\text{Sg}(T)$  is a signature and  $\text{A}(T)$  is a collection of  $\text{Sg}(T)$ -assertions. For each signature  $Sg$ , we let  $\text{Th}_{Sg}$  denote the complete lattice of  $Sg$ -theories ordered by  $T_1 \leq T_2$  if and only if  $\text{A}(T_1) \subseteq \text{A}(T_2)$ .

Now that we have defined the well-formed expressions, a **(first-order) logic**<sup>4</sup> is a closure operator on the lattice of all  $Sg$ -theories for each signature  $Sg$ . In the sequel, we will also assume all logics we consider satisfy the rules of equational logic in Figure 2. Given  $T, T' \in \text{Th}_{Sg}$ , we let  $T_{\mathcal{L}}$  denote the  $\mathcal{L}$ -closure of  $T$  and  $T \vdash_{\mathcal{L}} T'$ , denote the assertion that  $T'$  is **derivable** from  $T$ , i.e. that  $T' \leq T_{\mathcal{L}}$ .

### 3 Prop-Categorical Semantics

In this section, we define the class of prop-categories that provide semantics for first-order logics and provide examples. Structures in prop-categories and the notion of a structure satisfying a theory are then defined and it is shown how a class of prop-categories defines a first-order logic.

A **prop-category**  $(\mathcal{C}, P)$ , is a category  $\mathcal{C}$  with designated finite products, and a contravariant functor  $P: \mathcal{C} \rightarrow \mathbf{Pos}$ , where  $\mathbf{Pos}$  is the category of all partially ordered sets and monotone maps. The objects of  $\mathcal{C}$  interpret the sorts, the morphisms interpret terms-in-context and the posets  $P(c)$  for each  $c \in \text{Ob}(\mathcal{C})$  interpret the formulas-in-context. We define  $\text{Ob}(\mathbf{FA})$  to be the collection of all prop-categories which additionally satisfy:

1. For each  $c \in \text{Ob}(\mathcal{C})$ ,  $P(c)$  is an  $\mathcal{L}_\omega$ -algebra<sup>5</sup> and for all  $f \in \text{Mor}(\mathcal{C})$ ,  $P(f)$  is an  $\mathcal{L}_\omega$ -algebra homomorphism.
2. For each  $c \in \text{Ob}(\mathcal{C})$ , there is a designated element  $Eq_c \in P(c \times c)$ .
3. For each  $c \in \text{Ob}(\mathcal{C})$ , and each  $\Omega \in \mathcal{L}_q$ , there is a natural transformation  $\Omega_{(\cdot),c}: UP(\_ \times c) \Rightarrow UP$ , where  $U: \mathbf{Pos} \rightarrow \mathbf{Set}$  is the forgetful functor.
4. For each  $c \in \text{Ob}(\mathcal{C})$ ,  $P(c)$  has a designated binary operation  $\otimes^{P(c)}$ , and nullary operation  $e_c \in P(c)$  so that  $(P(c), \otimes^{P(c)}, e_c)$  is a monoid.
5. For all  $\Omega \in \mathcal{L}_q$ , and all  $b, c, d \in \text{Ob}(\mathcal{C})$ ,

$$\Omega_{b,1} \circ P(\pi_1^{b,1}) = id_{P(b)} \quad \text{and} \quad \Omega_{b,c \times d} \circ P(a_{b,c,d}) = \Omega_{b,c} \circ \Omega_{b \times c, d}, \quad (2)$$

where  $a_{b,c,d}: b \times (c \times d) \rightarrow (b \times c) \times d$  is the change-in-product isomorphism.

6.  $Eq_1 = e_{1 \times 1}$ , and for all  $c_1, c_2 \in \text{Ob}(\mathcal{C})$ ,  $c = c_1 \times c_2$ ,

$$Eq_c = P(\langle \pi_1^{c_1, c_2} \pi_1^{c, c}, \pi_1^{c_1, c_2} \pi_2^{c, c} \rangle) Eq_{c_1} \otimes P(\langle \pi_2^{c_1, c_2} \pi_1^{c, c}, \pi_2^{c_1, c_2} \pi_2^{c, c} \rangle) Eq_{c_2}. \quad (3)$$

<sup>4</sup>In Section 5 we define an action by substitution which can be used to define a (first-order) logic as *structural* closure operator.

<sup>5</sup>By  $\mathcal{L}_\omega$ -algebra we mean any algebra in the signature  $\mathcal{L}_\omega$ . We do not assume these algebras satisfy any particular collection of equations, nor do we assume their operations are monotone.

Conditions 1-3 are necessary for structures in  $(\mathcal{C}, P)$  to interpret equations and formulas in context. Condition 4 and the requirement that the codomain of  $P$  is **Pos** are needed to interpret sequents<sup>6</sup>. Conditions 5 and 6 are properties satisfied by the classifying prop-categories associated to certain theories and are not strictly necessary to define the prop-categorical semantics. However, they must be assumed to interpret structures as morphisms out of classifying prop-categories, a key ingredient in the proofs of our main results.

*Remark 1.* There are adjoint conditions for the quantifiers  $\{\forall, \exists\}$  (See Example 1) originally due to Lawvere [13] which asserts that for all objects  $a, b \in \text{Ob}(\mathcal{C})$ ,  $\forall_{a,b}, \exists_{a,b}$  are the left and right adjoints of  $P(\pi_1^{a,b})$  respectively. One can show that Condition 5 follows from these adjoint conditions. In [14], Lawvere also provides an adjoint interpretation of the equality predicate. For an account in the context of intuitionistic logic, see [9, p. 190]. We use conditions 5 and 6 instead of the corresponding adjoint conditions, because they are simple, comparatively weak and in a form that will be directly used to prove essential properties of the fibered semantics<sup>7</sup>. Moreover, in the spirit of AAL, we seek to provide general semantics for as broad of a variety of logics as possible and to use the property of *having this semantics* – here, a semantics with classifying prop-categories – as a means to classify nonclassical first-order logics. However, for many important examples  $\mathcal{L}_q = \{\forall, \exists\}$  and the quantifiers and equality do satisfy these adjoint conditions. Thus, one may safely replace 5 and 6 with these adjoint conditions provided corresponding “adjoint” derivation rules are added to the minimal logic  $\mathcal{L}^m$  in Section 4 to preserve the completeness results therein.

**Example 1.** Let  $L$  be an algebra of signature  $\mathcal{L}_\omega$  which is also a complete lattice and let  $\mathcal{L}_q = \{\forall, \exists\}$ . Then the contravariant functor  $\mathbf{Set}(\_, L): \mathbf{Set} \rightarrow \mathbf{Pos}$  determines a prop-category in  $\text{Ob}(\mathbf{FA})$ . The operations on  $L$  extend pointwise to logical operations on  $\mathbf{Set}(A, L)$  for all sets  $A$ . For each set  $X$ , let  $e_X = \top$ ,  $\otimes := \wedge$  and

$$Eq_X(x_1, x_2) = \begin{cases} \top & x_1 = x_2, \\ \perp & x_1 \neq x_2. \end{cases}$$

For sets  $X, Y$ , the quantifiers  $\exists_{X,Y}, \forall_{X,Y}$  are defined as the left and right adjoints of  $\mathbf{Set}(\pi_1^{X,Y}, L)$  respectively. That is,

$$\exists_{X,Y}(R)(x) = \bigvee_{y \in Y} R(x, y), \quad \text{and} \quad \forall_{X,Y}(R)(x) = \bigwedge_{y \in Y} R(x, y).$$

In particular,  $\mathbf{Set}(\_, 2)$  is the prop-category whose semantics corresponds to the usual Tarskian semantics for classical first-order logic<sup>8</sup>.

<sup>6</sup>In AAL, a (logical) matrix  $\langle A, F \rangle$  consists of an algebra  $A$  in the propositional language and a subset  $F \subseteq A$  of “truth values” [6, p. 183]. One can develop a “fibered” matrix semantics by dropping Condition 4 on prop-categories and defining a fibered matrix as  $\langle (\mathcal{C}, P), \mathcal{F} \rangle$  where  $(\mathcal{C}, P)$  is a prop-category and  $\mathcal{F} \subseteq \sqcup_{c \in \text{Ob}(\mathcal{C})} P(c)$ .

<sup>7</sup>Proving 6 from the adjoint conditions requires some work and is proved in [9, pg. 198] for a slightly different setup.

<sup>8</sup>Except here we haven’t excluded structures which interpret sorts as the emptyset.

Another important example in this class is  $\mathbf{Set}(\_, [0, 1])$ , where for each set  $A$ , the poset  $\mathbf{Set}(A, [0, 1])$  is known as the set of all **fuzzy sets on  $A$** . There is a vast literature on fuzzy set theory (for an introduction see [7]) which generalizes set theory by considering partial or probabilistic membership to sets.

An important class of logical operations on fuzzy sets are  $T$ -norms [12, pg. 4] and dually  $S$ -norms. A  $T$ -norm is an operation  $\odot: [0, 1]^2 \rightarrow [0, 1]$  which makes  $([0, 1], \odot, 1)$  a partially ordered monoid. Canonical examples of  $T$ -norms include minimum, product and the Łukasiewicz  $T$ -norms which is defined for  $x, y \in [0, 1]$  by  $x \odot y := \max\{0, x + y - 1\}$ . In [16], to each  $T$ -norm  $\odot$ , a quantifier  $\Omega^\odot$ , is defined on sets  $A, B$  and  $R: A \times B \rightarrow [0, 1]$ , by

$$\Omega_{A,B}^\odot(R) := \inf_{U \in \text{fin}(B)} \left\{ \bigcirc_{u \in U} R(\_, u) \right\},$$

where  $\text{fin}(B)$  is the set of all finite subsets of  $B$ . In particular, note that if  $\odot(a, b) = \min\{a, b\}$  then  $\Omega^\odot = \forall$ . One may verify that for an arbitrary  $T$ -norm,  $\odot$ ,  $\Omega^\odot$  satisfies Conditions 3 and 5. Thus  $\mathbf{Set}(\_, [0, 1])$  may be extended with additional quantifiers  $\Omega^\odot$  for each  $T$ -norm  $\odot$ . Moreover,  $\Omega^\odot$  may also be extended with  $S$ -quantifiers corresponding to  $S$ -norms as defined in [16]. If  $\odot$  is the usual product, then for  $R: 1 \times 2 \rightarrow [0, 1]$ ,  $R(0, 0) = R(0, 1) = 1/2$ ,  $\forall_{1,B}(R)(0) = \exists_{1,B}(R)(0) = 1/2$ , whereas  $\Omega_{1,B}^\odot(R)(0) = 1/4$ . Moreover, observe that in this example  $\odot$  may be used for  $\otimes$  instead of  $\min$ .

The following is a toy example which suggests integral operators are an fruitful place to look for nonclassical quantifiers and metrics for different notions of equality:

**Example 2.** Let  $\mathcal{K}$  be the collection of compact subsets of  $\mathbb{R}^n$  for some  $n < \omega$ , each equipped with the Euclidean metric along with  $1 = \{*\}$ , where  $1$ , has the discrete metric. Let  $\mathcal{C}$  be the category whose objects are finite products of sets in  $\mathcal{K}$  equipped with the associated product metric and continuous functions between them. For  $A \in \text{Ob}(\mathcal{C})$ , let  $P(A)$  be the collection of continuous real-valued functions from  $A$  to  $\mathbb{R}$ , which we denote  $C(A, \mathbb{R})$ , and for  $f: A \rightarrow B$ , let  $P(f): C(B, \mathbb{R}) \rightarrow C(A, \mathbb{R})$  be precomposition by  $f$ . Then, post-composition by any continuous map  $h: \mathbb{R}^n \rightarrow \mathbb{R}$  can interpret each  $n$ -ary connective  $\diamond \in \mathcal{L}_n$ .

For  $A \in \text{Ob}(\mathcal{C})$ , we let  $e_A$  be the constant zero function and  $Eq_A: A \times A \rightarrow \mathbb{R}$  be defined by  $Eq_A(a_1, a_2) := -|a_1 - a_2|$ , where  $|\cdot|$  is the Manhattan norm. For  $f, g \in C(A, \mathbb{R})$ , we define  $f \otimes g := f + g$ . Then  $(A, \otimes^{P(A)}, e_A)$  is a monoid. Moreover, Condition 6 is satisfied.

For all objects  $A, B$  we have a map  $\int_{A,B}: P(A \times B) \rightarrow P(A)$ , which sends each continuous function  $p: A \times B \rightarrow \mathbb{R}$  to  $\int_B p(x, y) dy$ . One may verify that  $\int_{(\cdot), \mathcal{C}}$  is natural for each object  $C$  and since  $1$  was given the discrete metric,  $\int_{A,1} \circ P(\pi_1^{A,1}) = id_{P(A)}$ . from Fubini's Theorem,  $\int_{(\cdot), (\cdot)}$  also satisfies the second part of Condition 5. Thus  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$ . Note, in general,  $\int_{A,B}$  is not left or right adjoint to  $P(\pi_1^{A,B})$ . To see this, consider  $A = B = [0, 1]$ ,  $r(x, y): A \times B \rightarrow [0, \infty)$ ,  $r(x, y) = y$ . Then  $\int_{A,B}(r) = 1/2$  whereas  $\exists_{A,B}(r) = 0$  and  $\forall_{A,B}(r) = 1$ .



Let  $Sg$  be a signature and  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$ . An  $Sg$ -**structure**  $S$  in  $(\mathcal{C}, P)$  is an assignment of an object  $\llbracket \sigma \rrbracket$  to each sort symbol  $\sigma$ , a morphism  $\llbracket f \rrbracket: \llbracket \sigma_1 \rrbracket \times \cdots \times \llbracket \sigma_n \rrbracket \rightarrow \llbracket \tau \rrbracket$  to each function symbol  $f: \sigma_1, \dots, \sigma_n \rightarrow \tau$  and an element  $\llbracket R \rrbracket \in P(\llbracket \sigma_1 \rrbracket \times \cdots \times \llbracket \sigma_n \rrbracket)$  to each relation symbol  $R \subseteq \sigma_1, \dots, \sigma_n$  in  $Sg$ . (To disambiguate between structures, we sometimes write  $S[\cdot]$  instead of  $\llbracket \cdot \rrbracket$ .) For a context  $\Gamma = x_1 : \sigma_1, \dots, x_n : \sigma_n$ , we define  $\llbracket \Gamma \rrbracket := \llbracket \sigma_1 \rrbracket \times \cdots \times \llbracket \sigma_n \rrbracket$ , in particular if  $n = 0$ ,  $\llbracket \Gamma \rrbracket := 1$ . Each term-in-context  $M : \tau \llbracket \Gamma \rrbracket$  may be given a unique interpretation  $\llbracket M : \tau \llbracket \Gamma \rrbracket \rrbracket$  defined recursively as follows:

$$\begin{aligned} \llbracket x_i : \sigma_i \llbracket x_1 : \sigma_1, \dots, x_n : \sigma_n \rrbracket \rrbracket &:= \pi_i : \llbracket \sigma_1 \rrbracket \times \cdots \times \llbracket \sigma_n \rrbracket \rightarrow \llbracket \sigma_i \rrbracket \\ \llbracket c : \tau \llbracket \Gamma \rrbracket \rrbracket &:= \llbracket c \rrbracket \circ !_{\llbracket \Gamma \rrbracket} \\ \llbracket f(M_1, \dots, M_n) : \tau \llbracket \Gamma \rrbracket \rrbracket &:= \llbracket f \rrbracket \circ \langle \llbracket M_1 \llbracket \Gamma \rrbracket \rrbracket, \dots, \llbracket M_n \llbracket \Gamma \rrbracket \rrbracket \rangle, \end{aligned}$$

where  $\pi_i$  is the  $i$ -th projection map and  $!_{\llbracket \Gamma \rrbracket}$  denotes the unique morphism from  $\llbracket \Gamma \rrbracket$  to 1. Note we often abbreviate  $\llbracket M : \tau \llbracket \Gamma \rrbracket \rrbracket$  as  $\llbracket M \llbracket \Gamma \rrbracket \rrbracket$  as the type  $\tau$  can be inferred from the term  $M$  and context  $\Gamma$ .

$S$  interprets formulas-in-context recursively as follows:

$$\begin{aligned} \llbracket R(M_1, \dots, M_n) \llbracket \Gamma \rrbracket \rrbracket &:= P(\langle \llbracket M_1 \llbracket \Gamma \rrbracket \rrbracket, \dots, \llbracket M_n \llbracket \Gamma \rrbracket \rrbracket \rangle)(\llbracket R \rrbracket), \\ \llbracket M_1 =_{\tau} M_2 \llbracket \Gamma \rrbracket \rrbracket &:= P(\langle \llbracket M_1 \llbracket \Gamma \rrbracket \rrbracket, \llbracket M_2 \llbracket \Gamma \rrbracket \rrbracket \rangle)(\llbracket Eq_{\llbracket \tau \rrbracket} \rrbracket), \\ \llbracket \diamond(\phi_1, \dots, \phi_n) \llbracket \Gamma \rrbracket \rrbracket &:= \diamond^{P(\llbracket \Gamma \rrbracket)}(\llbracket \phi_1 \llbracket \Gamma \rrbracket \rrbracket, \dots, \llbracket \phi_n \llbracket \Gamma \rrbracket \rrbracket), \\ \llbracket \Omega_{x:\sigma}(\phi) \llbracket \Gamma \rrbracket \rrbracket &:= \Omega_{\llbracket \Gamma \rrbracket, \llbracket \sigma \rrbracket} \circ P(a_{\llbracket \Gamma, x:\sigma \rrbracket}^S)(\llbracket \phi \llbracket \Gamma, x:\sigma \rrbracket \rrbracket), \end{aligned}$$

where  $a_{\llbracket \Gamma, x:\sigma \rrbracket}^S : \llbracket \Gamma \rrbracket \times \llbracket x:\sigma \rrbracket \rightarrow \llbracket \Gamma, x:\sigma \rrbracket$  is the change-in-product isomorphism<sup>9</sup>.

Then,  $S$  **satisfies an equation-in-context**  $M_1 = M_2 : \tau \llbracket \Gamma \rrbracket$ , if

$$\llbracket M_1 : \tau \llbracket \Gamma \rrbracket \rrbracket = \llbracket M_2 : \tau \llbracket \Gamma \rrbracket \rrbracket$$

and  $S$  **satisfies a sequent-in-context**  $\phi_1, \dots, \phi_n \vdash \phi_{n+1} \llbracket \Gamma \rrbracket$ , if for  $n \geq 0$ ,

$$\llbracket \phi_1 \llbracket \Gamma \rrbracket \rrbracket \otimes \cdots \otimes \llbracket \phi_n \llbracket \Gamma \rrbracket \rrbracket \leq \llbracket \phi_{n+1} \llbracket \Gamma \rrbracket \rrbracket.$$

In particular, for  $n = 0$ , if  $e_{\llbracket \Gamma \rrbracket} \leq \llbracket \phi_{n+1} \llbracket \Gamma \rrbracket \rrbracket$ . We define the **theory of**  $S$ , denoted  $\text{Th}(S)$ , to be the  $Sg$ -theory where  $\text{A}(\text{Th}(S))$  is the collection of all assertions satisfied by  $S$ . We say  $S$  **satisfies**  $T$  or  $S$  **is a**  $T$ -**model** if  $T \leq \text{Th}(S)$ .

**Example 3.** In [17], Mostowski introduced a general class of quantifiers and studied classical first-order logic extended with these quantifiers. Instead of  $\mathbf{Set}(\_, 2)$ , we consider the isomorphic powerset functor  $\mathcal{P} : \mathbf{Set} \rightarrow \mathbf{Pos}$ , which sends a set  $A$  to its powerset  $\mathcal{P}(A)$  and a function  $f : A \rightarrow B$  to its preimage operator. A **Mostowski quantifier**<sup>10</sup>  $\Omega$ , is specified for each set  $A$ , by a subset  $\Omega(A) \subseteq \mathcal{P}(A)$ . Then for  $A, B$  sets, and  $R \subseteq A \times B$ , we define

$$\Omega_{A,B}(R) := \{a \in A : \{b \in B : (a, b) \in R\} \in \Omega(B)\}.$$

<sup>9</sup>This handles the case where  $\Gamma = \emptyset$  and  $\llbracket \Gamma, x:\sigma \rrbracket = \llbracket x:\sigma \rrbracket \neq \llbracket \Gamma \rrbracket \times \llbracket x:\sigma \rrbracket = 1 \times \llbracket x:\sigma \rrbracket$ .

<sup>10</sup>In Mostowski's original formulation is more restrictive and requires quantifiers to be cardinality invariant. Our presentation here is essentially that of [22].

Note that quantifiers  $\forall$  and  $\exists$  belong to the class, defined for a set  $A$ , by  $\forall(A) := \{A\}$  and  $\exists(A) := \mathcal{P}(A) \setminus \{\emptyset\}$  respectively. A nonstandard Mostowski quantifier of early interest [10] is the “there exists uncountably many” quantifier  $Q_1$ , where for each set  $A$ ,  $Q_1(A)$  is the collection of all uncountable subsets of  $A$ .

One may verify that  $(\mathbf{Set}, \mathcal{P})$  augmented with a Mostowski quantifier  $\Omega$  satisfies Condition (3) and so structures in  $(\mathbf{Set}, \mathcal{P})$  can interpret the language of classical logic augmented with  $\Omega$ . However, in general, many such natural quantifiers, fail to satisfy Conditions (5). The first part of Equation 2 requires that  $Q_1(1) = \{1\}$ , and the second part of Equation 2 requires for all sets  $A, B$  and  $R \subseteq A \times B$ , that

$$R(x, y) \in Q(A \times B) \iff \{a \in A : R(a, y) \in Q(B)\} \in Q(A). \quad (4)$$

One sees that this condition also fails for  $Q_1$ , by considering  $A = B = \mathbb{R}$  and  $R = \mathbb{R} \times \{0\}$ .

A subcollection  $\mathcal{X} \subseteq \text{Ob}(\mathbf{FA})$ , defines a logic  $\models_{\mathcal{X}}$ , where given  $Sg$ -theories  $T$  and  $T'$ ,  $T \models_{\mathcal{X}} T'$  if every  $T$ -model  $S \in (\mathcal{C}, P) \in \mathcal{X}$  is also a  $T'$ -model. Given a logic  $\mathcal{L}$ , we let  $\text{Ob}(\mathbf{FA}_{\mathcal{L}})$  be the collection of all prop-categories in  $\text{Ob}(\mathbf{FA})$  such that  $\mathcal{L}$  is sound with respect to  $\models_{\text{Ob}(\mathbf{FA}_{\mathcal{L}})}$ , that is, whenever  $T \vdash_{\mathcal{L}} T'$ , then  $T \models_{\text{Ob}(\mathbf{FA}_{\mathcal{L}})} T'$ .

## 4 Classifying Prop-Categories and General Completeness Theorems

In the subsequent sections, we suppose  $\mathcal{L}$  has designated operation symbols  $e \in \mathcal{L}_0$  and  $\otimes \in \mathcal{L}_2$ . Given a sufficiently rich theory  $T$ , we will construct the *classifying prop-category*  $(\mathcal{C}_T, P_T) \in \text{Ob}(\mathbf{FA})$ , which is the first-order analogue of the Lindenbaum-Tarski (LT) algebras of propositional logic. As with LT-algebras, each classifying prop-category  $(\mathcal{C}_T, P_T)$  contains a *generic*  $T$ -model  $G \in (\mathcal{C}_T, P_T)$ , i.e.  $\text{Th}(G) = T$ , which interprets each term and formula in context as its associated equivalence class obtained by “quotienting out” by the theory  $T$ .

We call a logic  $\mathcal{L}$  over  $\mathcal{L}$  **adequate** if for each  $\mathcal{L}$ -theory  $T$ ,  $(\mathcal{C}_T, P_T)$  can be constructed and  $(\mathcal{C}_T, P_T) \in \text{Ob}(\mathbf{FA})$ . Define  $\mathcal{L}^m$  to be the first-order logic whose rules are listed in Figure 1.<sup>11</sup>

**Theorem 1.** *If  $\mathcal{L} \supseteq \mathcal{L}^m$ , then  $\mathcal{L}$  is adequate.*

To construct  $\mathcal{C}_T$ , from an  $\mathcal{L}$ -theory  $T$ , we consider contexts, formulas and terms in context equal up to renaming of their free variables. Morphisms are

<sup>11</sup>In AAL, propositional logics which satisfy  $\diamond$ -Cong for each connective are called self-extensional, which is the weakest class of logics in the Fregean hierarchy. [23] [6, pg. 419] Kleene’s strong 3-valued logic  $K3$  [11, p. 332] is a simple example of a non self-extensional propositional logic. However, A first-order version of  $K3$  can still be modeled as the fragment of an adequate logic by restricting to sequents with empty antecedent.

$$\begin{array}{c}
\frac{}{\phi \vdash \phi [\Gamma]} \text{Ax} \quad \frac{\phi \vdash \psi [\Gamma] \quad \psi \vdash \theta [\Gamma]}{\phi \vdash \theta [\Gamma]} \text{Cut} \\
\\
\frac{\Phi \vdash \psi [\Gamma]}{\Phi \vdash \psi [\Gamma, x : \sigma]} \text{Cwk} \quad \frac{M = M' : \sigma [\Delta] \quad \Phi \vdash \psi [\Gamma, x : \sigma, \Gamma']}{\Phi[M/x] \vdash \psi[M'/x] [\Gamma, \Gamma']} \text{Sub} \\
\\
\frac{\phi \dashv\vdash \psi [\Gamma, x : \sigma]}{\Omega_{x:\sigma}(\phi) \vdash \Omega_{x:\sigma}(\psi) [\Gamma]} \Omega\text{-Con} \quad \frac{\phi_1 \dashv\vdash \phi'_1 [\Gamma] \quad \dots \quad \phi_n \dashv\vdash \phi'_n [\Gamma]}{\diamond(\phi_1, \dots, \phi_n) \vdash \diamond(\phi'_1, \dots, \phi'_n) [\Gamma]} \diamond\text{-Cong} \\
\\
\frac{\Phi, \alpha, \beta, \Psi \vdash \theta [\Gamma]}{\Phi, \alpha \otimes \beta, \Psi \vdash \theta [\Gamma]} \otimes\text{-Ref} \quad \frac{\Phi, e, \Psi \vdash \phi [\Gamma]}{\Phi, \Psi \vdash \phi [\Gamma]} e\text{-Ref}
\end{array}$$

Figure 1: Rules for  $\mathcal{L}^m$ . (Where  $\text{VS}(\Delta) \subseteq \text{VS}(\Gamma, \Gamma')$  and  $\forall \Omega \in \mathcal{L}_q, \forall n \in \omega$  and  $\forall \diamond \in \mathcal{L}_n$ ).

sequences of equivalence classes of terms in a common context:

$$\gamma: \Gamma' \rightarrow \Gamma = [N_1 : \sigma_1 [\Gamma']/\sim, \dots, N_n : \sigma_n [\Gamma']/\sim],$$

where,  $\Gamma = x_1 : \sigma_1, \dots, x_n : \sigma_n$  and

$$N : \tau [\Gamma] \sim M : \tau [\Gamma] \iff N = M : \tau [\Gamma] \in T.$$

Composition of morphisms is by component-wise substitution and one may show using the rules of equational logic that  $\mathcal{C}_T$  is a category with strictly associative finite products [18, p. 30].

For each context  $\Gamma$ , we define  $\phi [\Gamma] \sim \psi [\Gamma]$  if  $\phi \vdash \psi [\Gamma], \psi \vdash \phi [\Gamma] \in T$ . From Ax and Cut,  $\sim$  defines an equivalence relation on formulas-in-context  $\Gamma$  and we let  $P_T(\Gamma)$  be the partial order of all such classes where  $\phi [\Gamma]/\sim \leq \psi [\Gamma]/\sim$  iff  $\phi \vdash \psi [\Gamma] \in T$ . For each morphism  $\gamma: \Gamma' \rightarrow \Gamma$ , we define  $P_T(\gamma): P_T(\Gamma) \rightarrow P_T(\Gamma')$  by simultaneous substitution:

$$P_T([N_1 : \sigma_1 [\Gamma']/\sim, \dots, N_n : \sigma_n [\Gamma']/\sim])(\phi [\Gamma]/\sim) := \phi[N_1/x_1, \dots, N_n/x_n] [\Gamma']/\sim.$$

For each  $n \in \omega$  and  $\diamond \in \mathcal{L}_n$ , we define:

$$\diamond^\Gamma(\phi_1 [\Gamma]/\sim, \dots, \phi_n [\Gamma]/\sim) := \diamond(\phi_1, \dots, \phi_n) [\Gamma]/\sim.$$

For each context  $\Gamma$ , define  $e_\Gamma := e [\Gamma]/\sim$ . For each quantifier symbol  $\Omega \in \mathcal{L}_q$ , and  $\phi [\Gamma, \Gamma']$ , where  $\Gamma' = y_1 : \tau_1, \dots, y_m : \tau_m$ , we interpret  $\Omega_{\Gamma, \Gamma'}$  as

$$\Omega_{\Gamma, \Gamma'}(\phi [\Gamma, \Gamma']/\sim) := \Omega_{y_1:\tau_1} \dots \Omega_{y_m:\tau_m}(\phi) [\Gamma]/\sim.$$

For each  $\Gamma = x_1 : \sigma_1, \dots, x_n : \sigma_n$ , let  $\Gamma' = x'_1 : \sigma_1, \dots, x'_n, \sigma_n$ . We define

$$Eq_\Gamma := x_1 =_{\sigma_1} x'_1 \otimes \dots \otimes x_n =_{\sigma_n} x'_n [\Gamma, \Gamma'] / \sim.$$

In particular, when  $\Gamma = []$ ,  $Eq_{[]} = e[] / \sim = e_{[] \times []}$ . Using the rules in Figure 1 one shows that these operations are well-defined and  $(\mathcal{C}_T, P_T) \in \text{Ob}(\mathbf{FA})$ .

The generic  $T$ -model  $G \in (\mathcal{C}_T, P_T)$ , is defined on sorts by  $G[[\sigma]] := x : \sigma$ , on function symbols  $f : \sigma_1, \dots, \sigma_n \rightarrow \tau$ , by  $G[[f]] := [f(\bar{x}) : \tau [\Gamma] / \sim]$  and on relation symbols  $R \subseteq \sigma_1, \dots, \sigma_n$  by  $G[[R]] := R(\bar{x}) [\Gamma] / \sim$ . By induction one shows that  $G[[M : \tau [\Gamma]]] = [M : \tau [\Gamma] / \sim]$  and  $G[[\phi [\Gamma]]] = \phi [\Gamma] / \sim$ . From  $\otimes$ -Ref and  $e$ -Ref it follows that  $\text{Th}(G) = T$ .

**Theorem 2.** *The logic  $\vDash_{\mathcal{K}}$  provides a complete semantics for  $\vdash_{\mathcal{L}^m}$ , where  $\mathcal{K}$  is any subclass of  $\text{Ob}(\mathbf{FA})$  containing  $\{(\mathcal{C}_T, P_T) : T \text{ is an } \mathcal{L}^m\text{-theory}\}$ .*

*Proof.* To show that  $\vdash_{\mathcal{L}^m} \subseteq \vDash_{\text{Ob}(\mathbf{FA})}$ , one verifies that each  $S \in (\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$  satisfies every interpretation of the rules defining  $\mathcal{L}^m$ . Now suppose  $T \vDash_{\mathcal{K}} T'$ . Then the generic  $\langle T \rangle_{\mathcal{L}^m}$ -model satisfies  $T'$ , and so  $T \vdash_{\mathcal{L}^m} T'$ .  $\square$

**Corollary 1.** *If  $\mathcal{L}$  is adequate, then  $\vdash_{\mathcal{L}} \supseteq \vdash_{\mathcal{L}^m}$ .*

Since otherwise, the generic model of some  $\mathcal{L}$ -theory will fail to satisfy an instance of a rule of  $\mathcal{L}^m$ . As a consequence, we let  $(\mathcal{C}_{Sg}, P_{Sg}) := (\mathcal{C}_{Sg_{\mathcal{L}^m}}, P_{Sg_{\mathcal{L}^m}})$ . In  $(\mathcal{C}_{Sg}, P_{Sg})$ , morphisms  $\gamma : \Gamma \rightarrow \Gamma'$  are just lists of terms and the elements of  $P_{Sg}(\Gamma)$  are just formulas, both up to  $\alpha$ -equivalence and change in the name of free variables, which are equivalences considered in mathematical practice. Thus,  $(\mathcal{C}_{Sg}, P_{Sg})$  is a suitable “fibered” analogue of the formula algebra from *AAL*.

Suppose  $\mathcal{L}$  is adequate and let  $\mathcal{K} = \{(\mathcal{C}_T, P_T) : T \text{ is an } \mathcal{L}\text{-theory}\}$ . Then using generic models, one proves  $\vDash_{\mathcal{K}} \subseteq \vdash_{\mathcal{L}}$ . All that remains to prove completeness is to show soundness, i.e.  $\vdash_{\mathcal{L}} \subseteq \vDash_{\mathcal{K}}$ . If  $\mathcal{L}$  is defined by a sequent calculus, we only need to show each structure in a classifying prop-category satisfies the defining rules of  $\mathcal{L}$ . We have the following general result:

**Theorem 3.** *Suppose  $\mathcal{L}_q = \{\forall, \exists\}$  and  $\mathcal{L}$  is an extension of  $\mathcal{L}^m$  possibly by  $-$ Adj,  $\exists$ -Adj,  $\forall$ -Adj shown in Figure 2 and any number of structural and propositional connective rules<sup>12</sup>. Then  $\vdash_{\mathcal{L}} = \vDash_{\mathcal{K}}$ , where  $\mathcal{K} = \{(\mathcal{C}_T, P_T) : T \text{ is an } \mathcal{L}\text{-theory}\}$ .*

*Proof.* Showing that  $\vDash_{\mathcal{K}}$  satisfies the structural and propositional connective rules of  $\mathcal{L}$  is straightforward.

Suppose  $\mathcal{L}$  satisfies  $\exists$ -Adj and let

$$\phi_1, \dots, \phi_n, \psi \vdash \theta [\Gamma, x : \sigma],$$

---

<sup>12</sup>Propositional connective rules, are sequent rules whose metaformulas do not include equality or any quantifiers such as modus ponens. Structural rules additionally do not include propositional connectives such as cut.

be an interpretation of the top line of  $\exists$ -Adj. Let  $S \in (\mathcal{C}_T, P_T) \in \mathcal{K}$  and for each  $i \in \{1, \dots, n\}$ , let  $S[\phi_i[\Gamma]] = \phi_i^S[\Gamma_S]/\sim$ ,  $S[\theta[\Gamma]] = \theta^S[\Gamma_S] \sim$  and  $S[\psi[\Gamma, x : \sigma]] = \psi^S[\Gamma_S, \Gamma_\sigma]/\sim$ , where  $S[\sigma] = \Gamma_\sigma = x_1 : \sigma_1, \dots, x_n : \sigma_n$ . Then,

$$\begin{aligned}
& \bigotimes_{i=1}^n S[\phi_i[\Gamma, x : \sigma]] \otimes S[\psi[\Gamma, x : \sigma]] \leq S[\theta[\Gamma, x : \sigma]] \\
\iff & \bigotimes_{i=1}^n G[\phi_i^S[\Gamma_S, \Gamma_\sigma]] \otimes G[\psi^S[\Gamma_S, \Gamma_\sigma]] \leq G[\theta^S[\Gamma_S, \Gamma_\sigma]] \\
\iff & T \vdash_{\mathcal{L}} \phi_1^S, \dots, \phi_n^S, \psi^S \vdash \theta^S[\Gamma_S, \Gamma_\sigma] \\
\iff & T \vdash_{\mathcal{L}} \phi_1^S, \dots, \phi_n^S, \exists_{x_1:\sigma_1} \dots \exists_{x_n:\sigma_n} (\psi^S) \vdash \theta^S[\Gamma_S] \\
\iff & \bigotimes_{i=1}^n S[\phi_i[\Gamma]] \otimes S[\exists_{x:\sigma}(\psi)[\Gamma]] \vdash S[\theta[\Gamma]].
\end{aligned}$$

It follows that  $\models_{\mathcal{K}}$  satisfies  $\exists$ -Adj. The proof that  $\models_{\mathcal{K}}$  satisfies  $\forall$ -Adj is similar.

Now suppose  $\mathcal{L}$  satisfies  $=$ -Adj and let  $S$  be a structure in  $(\mathcal{C}_T, P_T)$ . Consider an interpretation of  $=$ -Adj:

$$\frac{\phi_1, \dots, \phi_n \vdash \psi[x/x'][\Gamma, x : \sigma]}{\phi_1, \dots, \phi_n, x =_{\sigma} x' \vdash \psi[\Gamma, x : \sigma, x' : \sigma]}.$$

Let  $S[\phi_i[\Gamma, x : \sigma]] = \phi_i^S[\Gamma_S, \Gamma_\sigma]/\sim$ , where  $\Gamma_S \in S[\Gamma]$ ,  $\Gamma_\sigma \in S[\sigma]$  and  $S[\psi[\Gamma, x : \sigma, x', \sigma']] = \psi^S[\Gamma_S, \Gamma_\sigma, \Gamma'_\sigma]/\sim$ , where  $\Gamma'_\sigma \in S[\sigma']$ . From Sub and Cwk, one may derive the rule

$$\frac{\Phi \vdash \psi[\Gamma, x : \sigma, y : \tau, \Gamma']}{\Phi \vdash \psi[\Gamma, y : \tau, x : \sigma, \Gamma']} \quad (5)$$

Let  $\Gamma_\sigma = x_1 : \sigma_1, \dots, x_n : \sigma_n$  and  $\Gamma'_\sigma = x'_1 : \sigma_1, \dots, x'_n : \sigma_n$ . Let  $\bar{x} = x_1, \dots, x_n$  and  $\bar{x}' = x'_1, \dots, x'_n$  be the lists of variables in  $\Gamma_\sigma$  and  $\Gamma'_\sigma$  respectively. Then  $S[\psi[x/x'][\Gamma, x : \sigma]] = \psi^S[\bar{x}/\bar{x}'][\Gamma_S, \Gamma_\sigma]/\sim$  and

$$\begin{aligned}
& S[\phi_1[\Gamma, x : \sigma]] \otimes \dots \otimes S[\phi_n[\Gamma, x : \sigma]] \leq S[\psi[x/x'][\Gamma, x : \sigma]] \\
\iff & G[\phi_1^S[\Gamma_S, \Gamma_\sigma]] \otimes \dots \otimes G[\phi_n^S[\Gamma_S, \Gamma_\sigma]] \leq G[\psi^S[\bar{x}/\bar{x}'][\Gamma_S, \Gamma_\sigma]] \\
\iff & T \vdash_{\mathcal{L}} \phi_1^S, \dots, \phi_n^S \vdash \psi^S[\bar{x}/\bar{x}'][\Gamma_S, \Gamma_\sigma] \\
\iff & T \vdash_{\mathcal{L}} \phi_1^S, \dots, \phi_n^S, x_1 =_{\sigma_1} x'_1, \dots, x_n =_{\sigma_n} x'_n \vdash \psi^S[\Gamma_S, \Gamma_\sigma, \Gamma'_\sigma] \\
& \hspace{15em} (=Adj \text{ and } 5) \\
\iff & T \vdash_{\mathcal{L}} \phi_1^S, \dots, \phi_n^S, \bigotimes_{i=1}^n x_i =_{\sigma_i} x'_i \vdash \psi^S[\Gamma_S, \Gamma_\sigma, \Gamma'_\sigma] \quad (\otimes\text{-Ref or } e\text{-Ref}) \\
\iff & S[\phi_1[\Gamma, x : \sigma, x' : \sigma]] \otimes \dots \otimes S[\phi_n[\Gamma, x : \sigma, x' : \sigma]] \otimes S[x =_{\sigma} x'[\Gamma, x : \sigma, x' : \sigma]] \\
& \leq S[\psi[\Gamma, x : \sigma, x' : \sigma]].
\end{aligned}$$

Therefore,  $S$  satisfies all interpretations of  $=$ -Adj.  $\square$

$$\frac{\phi_1, \dots, \phi_n \vdash \psi[x/x'] [\Gamma, x : \sigma]}{\phi_1, \dots, \phi_n, x =_\sigma x' \vdash \psi [\Gamma, x : \sigma, x' : \sigma]} =\text{-Adj}$$

$$\frac{\Phi \vdash \psi [\Gamma, x : \sigma]}{\Phi, \vdash \forall_{x:\sigma}(\psi) [\Gamma]} \forall\text{-Adj} \quad \frac{\Phi, \psi \vdash \theta [\Gamma, x : \sigma]}{\Phi, \exists_{x:\sigma}(\psi) \vdash \theta [\Gamma]} \exists\text{-Adj}$$

Figure 2: Adjoint rules for  $\forall$ ,  $\exists$  and  $=$ .

## 5 The 2-Categorical View of the Prop-Categorical Semantics

We may extend  $\text{Ob}(\mathbf{FA})$  to a 2-category by defining morphisms  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  by the following data: (1) a product preserving functor  $F^o: \mathcal{C} \rightarrow \mathcal{D}$  and (2) a natural transformation  $F^p: P \Rightarrow Q \circ F^o$ ,

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{P} & \mathbf{Pos} \\ & \searrow^{F^o} & \uparrow^Q \\ & & \mathcal{D} \end{array} \quad \begin{array}{c} F^p \\ \Downarrow \\ \end{array}$$

such that the following hold:

1. For each  $c \in \text{Ob}(\mathcal{C})$ ,  $F_c^p: P(c) \rightarrow Q \circ F^o(c)$  is an  $\mathcal{L}_\omega$ -algebra homomorphism.
2. For all  $\Omega \in \mathcal{L}_q$  and all  $b, c \in \text{Ob}(\mathcal{C})$ ,

$$F_b^p \circ \Omega_{b,c} = \Omega_{F^o b, F^o c} \circ Q(a_{F,b,c}^{-1}) \circ F_{b \times c}^p.$$

3. For all  $c \in \text{Ob}(\mathcal{C})$ ,

$$F_{c \times c}^p(Eq_c) = Q(a_{F,c,c})(Eq_{F^o c}),$$

where  $a_{F,b,c}: F^o(b \times c) \rightarrow F^o b \times F^o c$  is the change-in-product isomorphism. Given  $K: (\mathcal{D}, Q) \rightarrow (\mathcal{E}, R)$ , we define  $K \circ F$  by  $(K \circ F)^o := K^o \circ F^o$  and  $(K \circ F)^p := K_{F^o}^p \cdot F^p$ . For parallel morphisms  $F, H: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$ , we define a 2-cell  $\eta: F \Rightarrow H$  to be a natural transformation  $\eta: F^o \Rightarrow H^o$  such that  $F^p = Q\eta \cdot H^p$ . Compositions of 2-cells is just as it is in  $\mathbf{Cat}$ . If  $\eta$  is a natural isomorphism, we call  $\eta$  a 2-isomorphism. For a logic  $\mathcal{L}$ , we now let  $\mathbf{FA}_{\mathcal{L}}$  be the corresponding full sub-2-category of  $\mathbf{FA}$ .

**Theorem 4.** *Let  $(\mathcal{C}, P), (\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA})$  and  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  be a morphism. Then for each theory  $T$ , each  $T$ -model  $S$  in  $(\mathcal{C}, P)$  gives rise to a  $T$ -model  $F(S)$  in  $(\mathcal{D}, Q)$ .*

*Proof.* First we define the structure  $F(S)$ . For each sort symbol  $\sigma$ ,  $F(S)[\sigma] := F^o(S[\sigma])$ , for each function symbol  $f: \sigma_1, \dots, \sigma_n \rightarrow \tau$ ,  $F(S)[f] := F^o(S[f]) \circ a_\Gamma^{-1}$ , and for each relation symbol  $R \subseteq \sigma_1, \dots, \sigma_n$ ,  $F(S)[R] := Q(a_\Gamma^{-1}) \circ F_{S[\Gamma]}^p(S[R])$ , where  $a_\Gamma: F^o(S[\Gamma]) \rightarrow F(S)[\Gamma]$  is the change-of-product morphism and  $\Gamma = x_1: \sigma_1, \dots, x_n: \sigma_n$ . A product preserving functor preserves the satisfaction of equations-in-context since by induction on the complexity of an arbitrary term-in-context  $M: \tau[\Gamma]$ ,

$$F(S)[M: \tau[\Gamma]] = F^o(S[M: \tau[\Gamma]]) \circ a_\Gamma^{-1}.$$

If for each formula-in-context  $\phi[\Gamma]$ ,

$$F(S)[\phi[\Gamma]] = Q(a_\Gamma^{-1})(F_{S[\Gamma]}^p(S[\phi[\Gamma]])), \quad (6)$$

then if  $S$  satisfies a sequent-in-context  $\phi_1, \dots, \phi_n \vdash \psi[\Gamma]$ , since  $F_{S[\Gamma]}^p$  and  $Q(a_\Gamma^{-1})$  are monotone  $\mathcal{L}$ -algebra homomorphisms,

$$\begin{aligned} & \bigotimes_{i=1}^n S[\phi_i[\Gamma]] \leq S[\psi[\Gamma]] \\ \implies & \bigotimes_{i=1}^n Q(a_\Gamma^{-1}) \circ F_{S[\Gamma]}^p(S[\phi_i[\Gamma]]) \leq Q(a_\Gamma^{-1}) \circ F_{S[\Gamma]}^p(S[\psi[\Gamma]]) \\ \implies & \bigotimes_{i=1}^n F(S)[\phi_i[\Gamma]] \leq F(S)[\psi[\Gamma]], \end{aligned}$$

where if  $n = 0$ , then  $\bigotimes_{i=1}^n = e_{S[\Gamma]}$ . And so  $F(S)$  satisfies  $\phi_1, \dots, \phi_n \vdash \psi[\Gamma]$ . It follows that  $F(S)$  is a  $T$ -model and so we only need to prove that Equation 6 holds.

Consider a formula-in-context  $\phi[\Gamma]$ . Suppose  $\phi[\Gamma]$  is atomic. Case 1:  $\phi[\Gamma]$  is of the form  $R(M_1, \dots, M_n)[\Gamma]$ , where  $R \subseteq \tau_1, \dots, \tau_n$ , and each  $M_i: \tau_i[\Gamma]$  is well-formed. If  $\gamma$  is the list  $\gamma = [M_1[\Gamma], \dots, M_n[\Gamma]]$ , we define

$$S[\gamma] := \langle S[M_1[\Gamma]], \dots, S[M_n[\Gamma]] \rangle.$$

Note that

$$F(S)[\gamma] = a_{\Gamma'} \circ F^o(S[\gamma]) \circ a_\Gamma^{-1}$$

Then,

$$\begin{aligned} & F(S)[R(M_1, \dots, M_n)[\Gamma]] \\ &= Q(F(S)[\gamma])(F(S)[R]) \\ &= Q(F(S)[\gamma]) \circ Q(a_\Gamma^{-1})(F_{S[\Gamma']}^p(S[R])) \\ &= Q(a_\Gamma^{-1}) \circ Q(F^o(S[\gamma])) \circ F_{S[\Gamma']}^p(S[R]) \\ &= Q(a_\Gamma^{-1}) \circ F_{S[\Gamma]}^p \circ P(S[\gamma])(S[R]) \\ &= Q(a_\Gamma^{-1}) \circ F_{S[\Gamma]}^p(S[R(M_1, \dots, M_n)[\Gamma]]). \end{aligned}$$

Otherwise,  $\phi[\Gamma]$  is of the form  $M_1 =_\tau M_2[\Gamma]$ . Let  $\Gamma' = [x_1 : \tau, x_2 : \tau]$  and  $\gamma: \Gamma \rightarrow \Gamma'$  be the context morphism represented by  $[M_1[\Gamma], M_2[\Gamma]]$ . Then,

$$\begin{aligned}
& F(S)[[M_1 =_\tau M_2[\Gamma]]] \\
&= Q(F(S)[[\gamma]])(Eq_{F(S)[[\tau]]}) \\
&= Q(a_{\Gamma'}^{-1}) \circ Q(F^o(S[[\gamma]])) \circ Q(a_{\Gamma'}) \circ Eq_{F(S)[[\tau]]} \\
&= Q(a_{\Gamma'}^{-1}) \circ Q(F^o(S[[\gamma]])) \circ F_{S[[\tau]] \times S[[\tau]]}^p(Eq_{S[[\tau]]}) \\
&= Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p \circ P(S[[\gamma]])(Eq_{S[[\tau]]}) \\
&= Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[M_1 =_\tau M_2[\Gamma]]]).
\end{aligned}$$

Now suppose  $\phi[\Gamma]$  is of the form  $\diamond(\phi_1, \dots, \phi_n)[\Gamma]$ , where  $\diamond$  is some  $n$ -ary operation in  $\mathcal{L}_\omega$  and for each  $i \in \{1, \dots, n\}$ ,  $F(S)[[\phi_i[\Gamma]]] = Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[\phi_i[\Gamma]]])$ . Then,

$$\begin{aligned}
& F(S)[[\diamond(\phi_1, \dots, \phi_n)[\Gamma]]] \\
&= \diamond^{Q(F(S)[[\Gamma]])}(F(S)[[\phi_1[\Gamma]]], \dots, F(S)[[\phi_n[\Gamma]]]) \\
&= \diamond^{Q(F(S)[[\Gamma]])}(Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[\phi_1[\Gamma]]]), \dots, Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[\phi_n[\Gamma]]])) \\
&= Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(\diamond^{P(S[[\Gamma]])}(S[[\phi_1[\Gamma]]], \dots, S[[\phi_n[\Gamma]]])) \\
&= Q(a_{\Gamma'}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[\diamond(\phi_1, \dots, \phi_n)[\Gamma]]]).
\end{aligned}$$

Suppose  $\phi[\Gamma]$  is of the form  $\Omega_{x:\sigma}(\psi)[\Gamma]$  for  $\Omega \in \mathcal{L}_q$ . Let  $I = S[[\Gamma]]$  and  $X = S[[\sigma]]$ . For  $c = c_1 \times \dots \times c_n$ , we let  $a_c: F^o(c) \rightarrow F^o c_1 \times \dots \times F^o c_n$ , and  $a_{c_1, c_2}: F^o(c_1 \times c_2) \rightarrow F^o c_1 \times F^o c_2$  be the change in product isomorphisms. Then

$$\begin{aligned}
& F(S)[[\Omega_{x:\sigma}(\psi)[\Gamma]]] \\
&= \Omega_{F(S)[[\Gamma], F(S)[[\sigma]]} \circ Q(a_{\Gamma, x:\sigma}^{F(S)})(F(S)[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_I^{-1}) \circ \Omega_{F^o(I), X} \circ Q(a_I \times id_{F(X)}) \circ Q(a_{\Gamma, x:\sigma}^{F(S)})(F(S)[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_I^{-1}) \circ \Omega_{F^o(I), X} \circ Q(a_I \times id_{F(X)}) \circ Q(a_{\Gamma, x:\sigma}^{F(S)}) \circ Q(a_{S[[\Gamma, x:\sigma]]}^{-1}) \circ F_{S[[\Gamma, x:\sigma]]}^p(S[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_I^{-1}) \circ \Omega_{F^o(I), X} \circ Q(a_{I, X}^{-1}) \circ Q(F^o(a_{\Gamma, x:\sigma}^S)) \circ F_{S[[\Gamma, x:\sigma]]}^p(S[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_I^{-1}) \circ \Omega_{F^o(I), X} \circ Q(a_{I, X}^{-1}) \circ F_{I \times X}^p \circ P(a_{\Gamma, x:\sigma}^S)(S[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_I^{-1}) \circ F_I^p \circ \Omega_{I, X} \circ P(a_{\Gamma, x:\sigma}^S)(S[[\psi[\Gamma, x : \sigma]]]) \\
&= Q(a_{S[[\Gamma]]}^{-1}) \circ F_{S[[\Gamma]]}^p(S[[\Omega_{x:\sigma}(\psi)[\Gamma]]]).
\end{aligned}$$

□

The next result says we can identify  $T$ -models  $S \in (\mathcal{C}, P)$  with morphisms  $\overline{S}: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$  which we use to develop an “algebraic” view of entailment.

**Theorem 5.** *Let  $\mathcal{L}$  be adequate and  $T$  an  $\mathcal{L}$ -theory. For each  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$  and  $T$ -model  $S$  in  $(\mathcal{C}, P)$  there is a morphism  $\overline{S}: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$  in  $\mathbf{FA}$ ,*



unique up to a 2-isomorphism such that  $\overline{S}(G) = S$ , where  $G$  is the generic  $T$ -model in  $(\mathcal{C}_T, P_T)$ .

*Proof.* We define  $\overline{S}^o$  on objects by  $\overline{S}^o(\Gamma) := S[[\Gamma]]$ , and on morphisms by

$$\overline{S}^o([M_1[\Gamma]/\sim, \dots, M_n[\Gamma]/\sim]) := \langle S[[M_1[\Gamma]]], \dots, S[[M_n[\Gamma]]] \rangle.$$

Since  $S$  is a  $T$ -model,  $\overline{S}^o$  is well defined on morphisms and it is straightforward to show that  $\overline{S}^o: \mathcal{C}_T \rightarrow \mathcal{C}$  is a product preserving functor such that  $\overline{S}(G) = S$ , where  $G$  is the generic  $T$ -algebra in  $\mathcal{C}_T$ . We can extend  $\overline{S}^o$ , by defining  $\overline{S}_\Gamma^p(\phi[\Gamma]/\sim) := S[[\phi[\Gamma]]]$ , and  $\overline{S}_\Gamma^p$  well-defined and monotone because  $S$  is a  $T$ -model. By definition,  $\overline{S}(G)$  agrees with  $\overline{S}^o$  on relation symbols. It is also straightforward to verify that  $\overline{S}^p: P_T \rightarrow P \circ \overline{S}^o$  is a natural transformation, and that for each context  $\Gamma$ ,  $\overline{S}_\Gamma^p$  is an  $\mathcal{L}$ -homomorphism.

We now verify Condition 2. For  $m > 0$ , let  $\Gamma = x_1 : \sigma_1, \dots, x_n : \sigma_n$ ,  $\Gamma' = x_{n+1} : \sigma_{n+1}, \dots, x_{n+m} : \sigma_{n+m}$ , and for each  $i \in \{1, \dots, n+m\}$ , let  $\pi_i: S[[\Gamma, \Gamma']] \rightarrow S[[\sigma_i]]$  be the  $i$ th projection map. Let  $S[[\Gamma]] = I$ ,  $S[[\Gamma']] = I'$  and for  $i \in \{1, \dots, n+m\}$ , let  $S[[\sigma_i]] = X_i$ . Let  $a: S[[\Gamma, \Gamma']] \rightarrow S[[\Gamma]] \times S[[\Gamma']]$  be the change-of-product isomorphism. Then for each  $\Omega \in \mathcal{L}_q$ , if  $m > 0$ ,

$$\begin{aligned} & \overline{S}_\Gamma^p \circ \Omega_{\Gamma, \Gamma'}(\phi[\Gamma, \Gamma']/\sim) \\ &= \overline{S}_\Gamma^p(\Omega_{x_{n+1}:\sigma_{n+1}} \cdots \Omega_{x_{n+m}:\sigma_{n+m}}(\phi)[\Gamma]/\sim) \\ &= \Omega_{I, X_{n+1}} \circ \dots \circ \Omega_{I \times X_{n+1} \cdots X_{n+m-1}, X_{n+m}}(S[[\phi[\Gamma, \Gamma']]]) \\ &= \Omega_{S[[\Gamma]], S[[\Gamma']]} \circ a^{-1*}(S[[\phi[\Gamma, \Gamma']]]) \quad (\text{by 5}) \\ &= \Omega_{\overline{S}^o(\Gamma), \overline{S}^o(\Gamma')} \circ a^{-1*} \circ \overline{S}_{\Gamma \times \Gamma'}^p(\phi[\Gamma, \Gamma']/\sim). \end{aligned}$$

Otherwise,  $\Gamma' = []$  and

$$\begin{aligned} \overline{S}_\Gamma^p \circ \Omega_{\Gamma, []}(\phi[\Gamma]/\sim) &= \overline{S}_\Gamma^p(\phi[\Gamma]/\sim) = S[[\phi[\Gamma]]] \\ &= \Omega_{S[[\Gamma]], S[[]]} \circ P(\pi_1^{S[[\Gamma]], S[[]]})(S[[\phi[\Gamma]]]) \quad (\text{by 5}) \\ &= \Omega_{\overline{S}^o(\Gamma), \overline{S}^o([])} \circ P(a_{\overline{S}, \Gamma, []}^{-1}) \circ \overline{S}_{\Gamma \times []}^p(\phi[\Gamma]/\sim). \end{aligned}$$

Let  $\Gamma = x_1 : \sigma_1, \dots, x_n : \sigma_n, \Gamma' = x_{n+1} : \sigma_1, \dots, x_{2n} : \sigma_n$ . If  $n > 0$ ,

$$\begin{aligned}
& \overline{S}_{\Gamma \times \Gamma'}^p(Eq_\Gamma) \\
&= \bigotimes_{i=1}^n S[[x_i =_{\sigma_i} x_{n+i}[\Gamma, \Gamma']]] \\
&= \bigotimes_{i=1}^n P(\langle \pi_i^{S[\Gamma, \Gamma']}, \pi_{i+n}^{S[\Gamma, \Gamma']} \rangle) Eq_{S[[\sigma_i]]} \\
&= \bigotimes_{i=1}^n P(\langle \pi_i^{S[\Gamma]} \pi_1^{S[\Gamma], S[\Gamma']}, \pi_i^{S[\Gamma']} \pi_2^{S[\Gamma], S[\Gamma']} \rangle \circ a) Eq_{S[[\sigma_i]]} \\
&= P(a) \circ \bigotimes_{i=1}^n P(\langle \pi_i^{S[\Gamma]} \pi_1^{S[\Gamma], S[\Gamma']}, \pi_i^{S[\Gamma']} \pi_2^{S[\Gamma], S[\Gamma']} \rangle) Eq_{S[[\sigma_i]]} \\
&= P(a) \circ Eq_{\overline{S}^o(\Gamma)}. \tag{by 6}
\end{aligned}$$

And if  $n = 0$ ,

$$\begin{aligned}
\overline{S}_{[]}^p(Eq_{[]}) &= \overline{S}_{[]}^p(e_{[]} / \sim) = S[[e_{[]}]] = e_1 && \text{( by 6)} \\
&= P(a_{\overline{S}, [], []})(e_{1 \times 1}) = P(a_{\overline{S}, [], []})(Eq_{\overline{S}^o[]}).
\end{aligned}$$

Now all that remains to show is that if  $F: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$  is another morphism such that  $F(G) = \overline{S}(G) = S$  then they are 2-isomorphic.

Let  $a_{F, \Gamma}: F^o(\Gamma) \rightarrow \overline{F(G)}^o(\Gamma) = \overline{S}^o(\Gamma)$  be the change-of-product isomorphism. Then  $a_F: F^o \Rightarrow \overline{S}^o$  is a natural isomorphism, and  $P a_F \cdot \overline{S}^p = P a_F \cdot \overline{F(G)}^p = F^p$ , from Equation 6. Thus  $a_F$  is a 2-isomorphism from  $F$  to  $\overline{S}$ .  $\square$

*Remark 2.* In the proof of Theorem 5, we see that to interpret structures as morphisms, all we need is for the prop-category  $(\mathcal{C}, P)$  to satisfy the identities satisfied by the classifying prop-category  $(\mathcal{C}_T, P_T)$ . In turn these identities are determined the theory  $T$  and more generally, the syntax. In order to include Mostowski Quantifiers which failed to satisfy Condition 5, one can extend the syntax with the following quantifier rule

$$\frac{\phi : \text{prop } [\Gamma, \Gamma']}{\Omega_{\Gamma'}(\phi) : \text{prop } [\Gamma]},$$

and define the interpretation of  $\Omega_{\Gamma'}(\phi) [\Gamma]$  by a structure as

$$[[\Omega_{\Gamma'}(\phi) [\Gamma]]] := \Omega_{[[\Gamma], [\Gamma']]} \circ P(a)^{-1}([[ \phi [\Gamma, \Gamma'] ]]),$$

where  $a$  is the change-in-product isomorphism. Thus Condition 5 may be dropped and similar considerations hold for the equality predicate and Condition 6.

**Definition 5.1.** Let  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  be a morphism in **FA**. We define the **kernel of  $F$** , denoted  $\ker F$  to consist of the following data:

1. A relation on  $\text{Ob}(\mathcal{C})$ , such that  $c_1 \sim c_2$  iff  $F^o c_1 = F^o c_2$ .
2. A relation on  $\text{Mor}(\mathcal{C})$  such that  $f_1 \sim f_2$  iff  $F^o f_1 = F^o f_2$ .
3. A relation on  $\sqcup_{c \in \text{Ob}(\mathcal{C})} P(c)$  such that  $r_1 \prec r_2$  iff  $F_{c_1}^p(r_1) \leq F_{c_2}^p(r_2)$ , where  $r_i \in P(c_i)$ .

If  $K: (\mathcal{C}, P) \rightarrow (\mathcal{E}, R)$  is another morphism, we say  $\ker K \leq \ker F$ , if and only if the relations of  $\ker K$  are contained in the corresponding relations of  $\ker F$ .

Let  $S$  be an  $Sg$ -structure in  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$  and let  $a \in A_{Sg}$ . If  $a = \phi_1, \dots, \phi_n \vdash \phi [\Gamma]$ , define

$$S[[a]] := \left( \bigotimes_{i=1}^n S[[\phi_i[\Gamma]]], S[[\phi[\Gamma]]] \right),$$

and if  $a$  is an equation  $M_1 = M_2 : \tau [\Gamma]$ , define

$$S[[a]] := (S[[M_1 : \tau [\Gamma]]], S[[M_2 : \tau [\Gamma]]]).$$

For each  $Sg$ -theory  $T$ , we define  $S[[T]] := \{S[[a]] : a \in A(T)\}$  and for  $F: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$  where  $T$  is an  $\mathcal{L}^m$ -theory, let  $\text{Th}(F)$  be the  $Sg(T)$ -theory such that

$$A(\text{Th}(F)) := \{a \in A_{Sg} : G[[a]] \in \ker F\}.$$

It is straightforward to verify that  $\text{Th}(F) = \text{Th}(F(G))$ . Thus, for each structure  $S \in (\mathcal{C}, P)$ ,  $\text{Th}(S) = \text{Th}(\overline{S}(G)) = \text{Th}(\overline{S})$ . Moreover, whenever  $F$  is 2-isomorphic to a parallel 2-cell  $K$ , then  $\text{Th}(F) = \text{Th}(K)$ .

Let  $T$ , and  $T'$  be  $Sg$ -theories and  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$ . From our prior observations the following are equivalent:

1.  $T \vDash_{(\mathcal{C}, P)} T'$ .
2.  $\forall F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P)$  such that  $T \leq \text{Th}(F)$ , then  $T' \leq \text{Th}(F)$ .
3.  $\forall F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P)$  such that  $G[[T]] \subseteq \ker F$ , then  $G[[T']] \subseteq \ker F$ .

Thus in the sequel, we take  $Sg$ -structures in  $(\mathcal{C}, P)$  to be morphisms  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P)$ . In the remainder of the section, we show how the prop-categorical semantics provide a natural notion of structural action on theories. This material is not necessary for the development of subsequent sections.

Let  $H: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_{Sg'}, P_{Sg'})$  be a morphism and let  $T_1, T_2$  be  $Sg$ -theories. We can define an action of  $H$  on  $Sg$  equations and sequents in context which can be extended to  $T_1$  by taking the union: that is  $H \cdot T_1$  is the  $Sg'$  theory whose assertions are  $\cup_{a \in A(T_1)} H \cdot a$ . For equations-in-context  $M^1 = M^2 : \tau [\Gamma]$ ,

$$H \cdot (M^1 = M^2 : \tau [\Gamma]) := \{M_i^1 = M_i^2 : \tau_i [\Gamma_H]\}_{i=1}^n,$$

where  $H^o(M^i : \tau [\Gamma]) = [M_1^i : \tau_1 [\Gamma_H], \dots, M_n^i : \tau_n [\Gamma_H]]$  for  $i \in \{1, 2\}$ . And for sequents-in-context  $\phi_1, \dots, \phi_n \vdash \phi_{n+1} [\Gamma]$ ,

$$H \cdot (\phi_1, \dots, \phi_n \vdash \phi_{n+1} [\Gamma]) = \phi_1^H, \dots, \phi_n^H \vdash \phi_{n+1}^H [\Gamma_H],$$

where  $H_T^p(\phi_i[\Gamma]) = \phi_i^H[\Gamma_H]$ , for all  $i \in \{1, \dots, n+1\}$ . Since products in  $(\mathcal{C}_{Sg}, P_{Sg})$  are unique up to permutation of the list of their variable sort pairs, if  $K: (\mathcal{C}_{Sg'}, P_{Sg'}) \rightarrow (\mathcal{C}_{Sg''}, P_{Sg''})$ , then  $(K \circ F) \cdot T_1 = K \cdot (F \cdot T_1)$  and clearly,  $id_{(\mathcal{C}_{Sg}, P_{Sg})} \cdot T_1 = T_1$ . Moreover, suppose  $\mathcal{V} \subseteq \text{Ob}(\mathbf{FA}_{\mathcal{L}})$  and that  $T_1 \vDash_{\mathcal{V}} T_2$ . Let  $(\mathcal{C}, P) \in \mathcal{V}$  and  $F: (\mathcal{C}_{Sg'}, P_{Sg'}) \rightarrow (\mathcal{C}, P)$  such that  $G[[H \cdot T_1]] \subseteq \ker F$ . Then  $G[[T_1]] \subseteq \ker F \circ H$  and since  $T_1 \vDash_{\mathcal{V}} T_2$ ,  $G[[T_2]] \subseteq \ker F \circ H$ . It follows that  $G[[H \cdot T_2]] \subseteq \ker F$  and so  $H \cdot T_1 \vDash_{\mathcal{V}} H \cdot T_2$ . Therefore, all logics defined semantically by subcollections of  $\text{Ob}(\mathbf{FA}_{\mathcal{L}})$  are structural with respect to the actions by morphisms.

This allows one to define a (first-order) logic as a class of closure operators  $\vdash_{\mathcal{L}}^{Sg}$ , one for each signature that is structural with respect to some subcategory  $\mathcal{G}$  of  $\mathbf{FA}_{\mathcal{L}}$  whose objects are the prop-categories  $(\mathcal{C}_{Sg}, P_{Sg})$ . Particularly interesting candidates are when  $\mathcal{G}$  is the full subcategory and when  $\mathcal{G}$  is the subcategory of all morphisms  $H: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_{Sg'}, P_{Sg'})$  which map single sort contexts to single sort contexts.

## 6 Basic Constructions in FA

Let  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$  and consider the signature  $Sg$  whose sorts are  $\text{Ob}(\mathcal{C})$ , whose function symbols are  $f: c_1, \dots, c_n \rightarrow c$ , for each  $f: c_1 \times \dots \times c_n \rightarrow c \in \text{Mor}(\mathcal{C})$  and whose relation symbols are  $R \subseteq c_1, \dots, c_n$  for each  $R \in P(c_1 \times \dots \times c_n)$ . Note that  $f: c_1 \times \dots \times c_n \rightarrow c \in \text{Ob}(\mathcal{C})$  is included both as an  $n$ -ary operation symbol  $f: c_1, \dots, c_n \rightarrow c$  and as a unary operation symbol  $f: c_1 \times \dots \times c_n \rightarrow c$  in  $Sg$  and similarly for relation symbols. There is a canonical  $Sg$ -structure  $S$  in  $(\mathcal{C}, P)$  called the **internal structure of  $(\mathcal{C}, P)$** , where  $[[c]] = c$ ,  $[[f]] = f$  and  $[[R]] = R$  for each sort, function symbol and relation symbol respectively. We define  $\text{Th}(\mathcal{C}, P) := \text{Th}(S)$ .

**Proposition 1.** *Let  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA})$  and let  $S$  be the internal structure of  $(\mathcal{C}, P)$ , and  $T$  the theory of  $S$ . Then  $\bar{S}: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$  determines a 2-equivalence.*

*Proof.* Define  $\iota: (\mathcal{C}, P) \rightarrow (\mathcal{C}_T, P_T)$  by  $\iota^o(c) := x: c$  for  $c \in \text{Ob}(\mathcal{C})$ ,  $\iota^o(f: c_1 \rightarrow c_2) := f(x): c_2 [x: c_1]/\sim$  for  $f \in \text{Mor}(\mathcal{C})$  and  $\iota^p(R) := R(x) [x: c]/\sim$  for each  $c \in \text{Ob}(\mathcal{C})$  and each  $R \in P(c)$ . It is straightforward to verify that  $\iota^o$  is a finite product preserving functor. Let  $f: c_1 \rightarrow c_2 \in \text{Mor}(\mathcal{C})$ . Then for  $R \in P(c_2)$ ,

$$\begin{aligned} \iota_{c_1}^p \circ P(f)(R) &= P(f)(R)(x)[x: c_1]/\sim \\ &= R(f(x))[x: c_1]/\sim \\ &= P_T(f(x): c_2[x: c_1]/\sim)(R(x)[x: c_2]/\sim) \\ &= P_T(\iota^o(f)) \circ \iota_{c_2}^p(R). \end{aligned}$$

Thus  $\iota^p$  is a natural transformation. It is also straightforward to verify, for each  $c \in \text{Ob}(\mathcal{C})$  that  $\iota_c^p$  is an monotone  $\mathcal{L}$ -algebra homomorphism. Let  $\Omega \in \mathcal{L}_q$ ,

$c_1, c_2 \in \text{Ob}(\mathcal{C})$  and  $R \in P(c_1 \times c_2)$ . Then

$$\begin{aligned} & \Omega_{\iota^{\circ} c_1, \iota^{\circ} c_2} \circ P_T(a^{-1}) \circ \iota_{c_1 \times c_2}^p(R) \\ &= \Omega_{x_2 : c_2}(R(\langle \pi_1, \pi_2 \rangle(x_1, x_2)))[x_1 : c_1] / \sim \\ &= \Omega_{c_1, c_2}(R)(x)[x : c_1] / \sim \\ &= \iota_{c_1}^p \circ \Omega_{c_1, c_2}(R). \end{aligned}$$

Let  $c \in \text{Ob}(\mathcal{C})$ . Then

$$\begin{aligned} & P_T(a^{-1}) \circ \iota_{c \times c}^p(Eq_c) \\ &= Eq_c(\langle \pi_1, \pi_2 \rangle(x_1, x_2))[x_1 : c, x_2 : c] / \sim \\ &= Eq_c(x_1, x_2)[x_1 : c, x_2 : c] / \sim \\ &= x_1 =_c x_2 [x_1 : c, x_2 : c] / \sim \\ &= Eq_{x:c}. \end{aligned}$$

It follows that  $\iota$  is a morphism.

Now  $\bar{S}^o \circ \iota^o = id_{\mathcal{C}}$ , and for each  $c \in \text{Ob}(\mathcal{C})$  and  $R \in P(c)$ ,

$$(\bar{S}_{\iota^o}^p \cdot \iota^p)_c(R) = \bar{S}_{\iota^o c}^p \circ \iota_c^p(R) = \bar{S}_{\iota^o c}^p(R(x)[x : c] / \sim) = R.$$

It follows that  $\bar{S} \circ \iota = id_{(\mathcal{C}, P)}$ .

In the other direction,

$$\iota^o \circ \bar{S}^o([x_1 : c_1, \dots, x_n : c_n]) = [x : c_1 \times \dots \times c_n],$$

$$\begin{aligned} & \iota^o \circ \bar{S}^o([M_1 : c_1[\Gamma] / \sim, \dots, M_m : c_m[\Gamma] / \sim]) \\ &= \langle S[[M_1 : c_1[\Gamma]]], \dots, S[[M_m : c_m[\Gamma]]] \rangle(x) : c_1 \times \dots \times c_m [x : S[[\Gamma]]] / \sim \end{aligned}$$

and

$$(\iota_{\bar{S}^o}^p \cdot \bar{S}^p)_{\Gamma}(\phi[\Gamma] / \sim) = S[[\phi[\Gamma]]](x)[x : S[[\Gamma]]] / \sim.$$

For each context  $\Gamma = x_1 : c_1, \dots, x_n : c_n$ , define  $\eta : \iota^o \cdot \bar{S}^o \Rightarrow id_{\mathcal{C}_T}$  by

$$\eta_{\Gamma} = [\pi_1(x) : c_1[x : S[[\Gamma]]] / \sim, \dots, \pi_n(x) : c_n[x : S[[\Gamma]]] / \sim].$$

One may verify that  $\eta$  is a natural isomorphism and that  $P_T \eta \cdot id_{(\mathcal{C}_T, P_T)}^p = \iota_{\bar{S}^o}^p \cdot \bar{S}^p$ .

It follows that  $\eta : \iota \circ \bar{S} \Rightarrow id_{(\mathcal{C}_T, P_T)}$  is a 2-isomorphism and so  $(\mathcal{C}_T, P_T)$  and  $(\mathcal{C}, P)$  are equivalent.  $\square$

Thus  $(\mathcal{C}_T, P_T) \equiv (\mathcal{C}, P)$  and so  $(\mathcal{C}_T, P_T)$  is a syntactic representation of  $(\mathcal{C}, P)$  similar to the representation of an algebra as the free algebra over its elements quotiented out by its equational theory. Moreover, it can be shown that if  $(\mathcal{C}, P), (\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA})$  are equivalent in  $\mathbf{FA}$ , then  $\models_{(\mathcal{C}, P)} = \models_{(\mathcal{D}, Q)}$  and so  $\models_{(\mathcal{C}, P)} = \models_{(\mathcal{C}_T, P_T)}$ .

Let  $(\mathcal{C}, P), (\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA})$ . We say  $(\mathcal{D}, Q)$  is a **subprop-category** of  $(\mathcal{C}, P)$ , if there exists a morphism  $\iota : (\mathcal{D}, Q) \rightarrow (\mathcal{C}, P)$ , such that  $\iota^o$  is faithful and for each  $c \in \text{Ob}(\mathcal{C})$ ,  $\iota_c^p$  is an order embedding. We call the morphism  $\iota$  a **subprop-morphism**.

**Proposition 2.** *If  $(\mathcal{D}, Q)$  is a subprop-category of  $(\mathcal{C}, P)$ , then  $\vDash_{(\mathcal{C}, P)} \subseteq \vDash_{(\mathcal{D}, Q)}$ .*

*Proof.* Suppose  $T \vDash_{(\mathcal{C}, P)} T'$  and consider  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{D}, Q)$ , such that  $G[[T]] \subseteq \ker F$ . Then  $G[[T]] \subseteq \ker(\iota \circ F)$ , and since  $T \vDash_{(\mathcal{C}, P)} T'$ ,  $G[[T']] \subseteq \ker(\iota \circ F)$ . Since  $\iota$  is a subprop-morphism,  $G[[T']] \subseteq \ker(F)$  and so  $T \vDash_{(\mathcal{D}, Q)} T'$ .  $\square$

Let  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  be a morphism in **FA**. In general,  $(F(\mathcal{C}), F(P))$ , is not a subprop-category of  $(\mathcal{D}, Q)$  where  $F(\mathcal{C})$  is the image of  $\mathcal{C}$  under  $F^\circ$ , and for all  $g: d_1 \rightarrow d_2 \in \text{Mor}(F(\mathcal{C}))$ ,  $F(P)(g) := Q(g)|_{F_{c_2}^p P(c_2)}: F_{c_2}^p P(c_2) \rightarrow F_{c_1}^p P(c_1)$ , where  $F^\circ c_i = d_i$  for  $i \in \{1, 2\}$ . When  $(F(\mathcal{C}), F(P))$  does define a subprop-category of  $(\mathcal{D}, Q)$  we call it the **image of  $F$** .

**Lemma 1.** *Let  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  be a morphism in **FA** such that  $F^\circ$  is injective on objects. Then  $(F(\mathcal{C}), F(P))$  is a subprop-category of  $(\mathcal{D}, Q)$  and there exists a unique morphism  $H: (\mathcal{C}, P) \rightarrow (F(\mathcal{C}), F(P))$  such that  $\iota \circ H = F$ , where,  $\iota: (F(\mathcal{C}), F(P)) \hookrightarrow (\mathcal{D}, Q)$  is the inclusion morphism. We call  $H$ , the **corestriction of  $F$  to  $(F(\mathcal{C}), F(P))$** . Furthermore,  $H$  is strictly finite product preserving, full, surjective on objects, and for each  $c \in \text{Ob}(\mathcal{C})$ ,  $H_c^p$  is surjective.*

*Proof.* Since  $F^\circ$  is injective on objects,  $F(\mathcal{C})$  is a sub-category of  $\mathcal{D}$ . Let  $d_1, d_2 \in \text{Ob}(F(\mathcal{C}))$  and  $F^\circ(c_i) = d_i$ . Then the designated product diagram of  $d_1$  and  $d_2$  in  $F(\mathcal{C})$  which we denote  $(d_1 \hat{\times} d_2, \hat{\pi}_1, \hat{\pi}_2)$  (to distinguish it from the designated diagram of  $d_1 \times d_2$  in  $\mathcal{D}$ ) is  $(F^\circ(c_1 \times c_2), F^\circ(\pi_1), F^\circ(\pi_2))$  where  $(c_1 \times c_2, \pi_1, \pi_2)$  is the designated product diagram of  $c_1 \times c_2$  in  $\mathcal{C}$ . Given  $g_1: d \rightarrow d_1$  and  $g_2: d \rightarrow d_2$  we let  $\langle g_1, g_2 \rangle := F^\circ(\langle f_1, f_2 \rangle)$ , which is the unique morphism such that  $\hat{\pi}_i \circ \langle g_1, g_2 \rangle = g_i$  for  $i \in \{1, 2\}$ . We also define the designated terminal object in  $F(\mathcal{C})$  to be  $1_{F(\mathcal{C})} := F^\circ(1_{\mathcal{C}})$ . Define  $F(P)(d_1) := F^p(P(c_1)) \leq Q(d_1)$  and for  $g \in F(\mathcal{C})(d_1, d_2)$ , if  $F^\circ(f) = g$ , define  $F(P)(g) := Q(g)|_{F(P)(d_2)} = Q(F^\circ f)|_{F_{c_2}^p P(c_2)}$ . Since  $F^p: P \Rightarrow Q \circ F^\circ$ , we have  $F(P)(g)(F(P)(d_2)) \subseteq F(P)(d_1)$ . For  $\Omega \in \mathcal{L}_q$  we denote the interpretation of  $\Omega$  in  $(F(\mathcal{C}), F(P))$  by  $\Omega^{F(P)}$ , which we define for  $r \in F(P)(d_1 \hat{\times} d_2)$  as

$$\Omega_{F^\circ c_1, F^\circ c_2}^{F(P)}(r) := \Omega_{F^\circ c_1, F^\circ c_2} \circ Q(a_{F, c_1, c_2}^{-1})(r).$$

Let  $r' \in P(c_1 \times c_2)$  such that  $F_{c_1 \times c_2}^p(r') = r$ . Then,

$$\Omega_{d_1, d_2}^{F(P)}(r) = \Omega_{d_1, d_2} \circ Q(a^{-1}) \circ F_{c_1 \times c_2}^p(r) = F_{c_1}^p(\Omega_{c_1, c_2}(r')) \in F(P)(d_1).$$

Equality  $Eq^{F(P)}$  is defined by

$$Eq_{F^\circ c}^{F(P)} := Q(a_{F, c, c})(Eq_{F^\circ c}) = F_{c \times c}^p(Eq_c) \in F(P)(F^\circ c \hat{\times} F^\circ c).$$

First we show that  $(F(\mathcal{C}), F(P)) \in \mathbf{FA}$ . Conditions 1 and 2 are immediate. Condition 3 follows from the fact that for each  $F^\circ c \in \text{Ob}(F(\mathcal{C}))$ ,

$$\Omega_{(\cdot), d}^{F(P)} = \Omega_{(\cdot), F^\circ c} \cdot Q(a_{F, (\cdot), c}^{-1})|_{F(P)((\cdot) \hat{\times} F^\circ c)}.$$

Condition 4 is satisfied as we take  $\otimes^{F(P)(d_1)} := \otimes^{Q(d_1)}|_{F(P)(d_1)}$  and  $e_{d_1}^{F(P)} := F_{c_1}^p(e_{c_1}) = e_{d_1}$ .

Next we consider Condition 5. Let  $d = F^o c \in F(\mathcal{C})$  and  $\Omega \in \mathcal{L}_q$ . Then

$$\begin{aligned} \Omega_{d, F^o(1)}^{F(P)} \circ F(P)(\hat{\pi}_1^{d, F^o(1)}) &= \Omega_{d, F^o(1)} \circ Q(a_{F, c, 1}^{-1}) \circ Q(\hat{\pi}_1^{d, F^o(1)})|_{F(P)(d)} \\ \Omega_{d, F^o(1)} \circ Q(\hat{\pi}_1^{d, F^o(1)})|_{F(P)(d)} &= id_{F(P)(d)}. \end{aligned}$$

Let  $d_1, d_2, d_3 \in F(\mathcal{C})$ , and  $F^o(c_i) = d_i$ . Let  $r \in F(P)((d_1 \hat{\times} d_2) \hat{\times} d_3)$  and  $r' \in P((c_1 \times c_2) \times c_3)$  such that  $F_{(c_1 \times c_2) \times c_3}^p(r') = r$ . Then

$$\begin{aligned} &= \Omega_{d_1, d_2 \hat{\times} d_3}^{F(P)} \circ Q(\hat{a}_{d_1, d_2, d_3})(r) \\ &= \Omega_{d_1, d_2 \hat{\times} d_3} \circ Q(a_{F, c_1, c_2 \times c_3}^{-1}) \circ Q(F^o(a_{c_1, c_2, c_3})) \circ F_{(c_1 \times c_2) \times c_3}^p(r') \\ &= \Omega_{F^o c_1, F^o(c_2 \times c_3)} \circ Q(a_{F, c_1, c_2 \times c_3}^{-1}) \circ F_{c_1 \times (c_2 \times c_3)}^p \circ P(a_{c_1, c_2, c_3})(r') \\ &= F^p \circ \Omega_{c_1, c_2 \times c_3} \circ P(a_{c_1, c_2, c_3})(r') \\ &= F_{c_1}^p \circ \Omega_{c_1, c_2} \circ \Omega_{c_1 \times c_2, c_3}(r') \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a_{F, c_1, c_2}^{-1}) \circ F_{c_1 \times c_2}^p \circ \Omega_{c_1 \times c_2, c_3}(r') \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a_{F, c_1, c_2}^{-1}) \circ \Omega_{F^o(c_1 \times c_2), F^o c_3} \circ Q(a_{F, c_1 \times c_2, c_3}^{-1}) \circ F_{(c_1 \times c_2) \times c_3}^p(r') \\ &= \Omega_{d_1, d_2}^{F(P)} \circ \Omega_{(d_1 \hat{\times} d_2), d_3}^{F(P)}(r). \end{aligned}$$

Thus Condition 5 is satisfied.

We now consider Condition 6.

$$Eq_{F^o(1)}^{F(p)} = F_{1 \times 1}^p(Eq_1) = F_{1 \times 1}^p(e_{1 \times 1}) = e_{F^o 1 \hat{\times} F^o 1}.$$

Let  $d_1, d_2 \in \text{Ob}(F(\mathcal{C}))$  and  $F^o c_i = d_i$ . Let  $c = c_1 \times c_2$  and  $d = d_1 \hat{\times} d_2$ . Then,

$$\begin{aligned} Eq_d^{F(P)} &= F_{c \times c}^p(Eq_c) \\ &= F_{c \times c}^p(\bigotimes_{i=1}^2 P(\langle \pi_i^c \pi_1^{c, c}, \pi_i^c \pi_2^{c, c} \rangle) Eq_{c_i}) \\ &= \bigotimes_{i=1}^2 F_{c \times c}^p \circ P(\langle \pi_i^c \pi_1^{c, c}, \pi_i^c \pi_2^{c, c} \rangle) Eq_{c_i} \\ &= \bigotimes_{i=1}^2 Q(F^o \langle \pi_i^c \pi_1^{c, c}, \pi_i^c \pi_2^{c, c} \rangle) \circ F_{c_i \times c_i}^p Eq_{c_i} \\ &= \bigotimes_{i=1}^2 F(P)(\langle \hat{\pi}_i^d \hat{\pi}_1^{d, d}, \hat{\pi}_i^d \hat{\pi}_2^{d, d} \rangle) Eq_{d_i}^{F(P)}. \end{aligned}$$

It follows that  $(F(\mathcal{C}), F(P)) \in \mathbf{FA}$ .

Let  $\iota^o: F(\mathcal{C}) \hookrightarrow \mathcal{D}$  be the inclusion functor and for each  $d \in \text{Ob}(F(\mathcal{C}))$ , let  $\iota_d^p: F(P)(d) \hookrightarrow Q(d)$  be the inclusion homomorphism. It is immediate that  $\iota^p: F(P) \Rightarrow Q \circ \iota^o$  is natural. Let  $\Omega \in \mathcal{L}_q$ ,  $d_1, d_2 \in \text{Ob}(F(\mathcal{C}))$  and  $F^o(c_i) = d_i$ .

For  $r \in F(P)(d_1 \hat{\times} d_2)$ ,

$$\begin{aligned} \iota_{d_1}^p \circ \Omega_{d_1, d_2}^{F(P)}(r) &= \Omega_{d_1, d_2} \circ Q(a_{F, c_1, c_2}^{-1})(r) \\ &= \Omega_{\iota_{d_1}^o, \iota_{d_2}^o} \circ Q(a_{F, c_1, c_2}^{-1}) \circ \iota_{d_1 \hat{\times} d_2}^p(r). \end{aligned}$$

And,

$$\iota_{d \hat{\times} d}^p(Eq^{F(P)}) = Q(a_{F, c, c}) \circ Eq_{\iota^o d}.$$

It follows that  $\iota = (\iota^o, \iota^p) \in \mathbf{FA}((F(\mathcal{C}), F(P)), (\mathcal{D}, Q))$ .

Define  $H = (H^o, H^p)$  by  $H^o: \mathcal{C} \rightarrow F(\mathcal{C})$ , where  $H^o(c) := F^o(c)$ ,  $H^o(f) := F^o(f)$  and  $H_c^p$  is the corestriction of  $F_c^p$  to  $F(P)(F^o c)$ . The naturality of  $H^p: P \Rightarrow F(P) \circ H^o$  follows from the naturality of  $F^p: P \Rightarrow Q \circ F^o$ . By definition  $H^o$  strictly preserves finite products and  $\iota \circ H = F$ . Since  $\iota^o$  is injective on objects and faithful, and  $\iota_d^p$  is injective, for each  $d \in \text{Ob}(F(\mathcal{C}))$ ,  $H$  is uniquely defined so that  $H \circ \iota = F$ . Let  $c_1, c_2 \in \text{Ob}(\mathcal{C})$  and  $r \in P(c_1 \times c_2)$ . That  $H$  commutes with the quantifiers and equality follows from the fact that  $F$  does. Thus,  $H \in \mathbf{FA}((\mathcal{C}, P), (F(\mathcal{C}), F(P)))$ .  $\square$

**Proposition 3.**  $\mathbf{FA}$  has all products.

*Proof.* The terminal object  $\mathbf{1}_{\mathbf{FA}}$  is  $(*, \mathbf{1}_*)$  where  $*$  is the terminal category and  $\mathbf{1}_*$  is its unique endofunctor. It is straightforward to verify that  $(*, \mathbf{1}_*) \in \mathbf{FA}$  and that for each prop-category  $(\mathcal{C}, P) \in \mathbf{FA}$ , there is a unique morphism  $!(\mathcal{C}, P): (\mathcal{C}, P) \rightarrow (*, \mathbf{1}_*)$ , where  $!(\mathcal{C}, P) = (!_{\mathcal{C}}, !_P)$ .

Let  $\{(\mathcal{C}_i, P_i)\}_{i \in I} \subseteq \text{Ob}(\mathbf{FA})$  be non-empty. Let  $\prod_I \mathcal{C}_i$  denote the product of the categories  $\{\mathcal{C}_i\}_{i \in I}$ , that is, the category whose objects  $a$  specify for each  $i \in I$ , an object  $a_i \in \text{Ob}(\mathcal{C}_i)$ , and whose morphisms  $f: a \rightarrow b$  specify for each  $i \in I$  a morphism  $f_i \in \mathcal{C}_i(a_i, b_i)$ . Composition  $g \circ f$  of morphisms  $f: a \rightarrow b$  and  $g: b \rightarrow c$  is defined for each  $i \in I$  as  $(g \circ f)_i := g_i \circ f_i: a_i \rightarrow c_i$ . Since each  $\mathcal{C}_i$  has finite products, we define  $\mathbf{1}$  so that  $\mathbf{1}_i = \mathbf{1}_{\mathcal{C}_i}$  and for  $a, b \in \text{Ob}(\prod_I \mathcal{C}_i)$ , we define  $a \times b$  so that  $(a \times b)_i := a_i \times b_i$ . For  $k \in \{1, 2\}$ , we define the projection maps  $\pi_k^{a, b}$  by  $\pi_{k, i}^{a, b} = \pi_k^{a_i, b_i}$ . One verifies that this determines a product diagram for  $a \times b$ , where for every pair of maps  $f: c \rightarrow a$  and  $g: c \rightarrow b$ ,  $\langle f, g \rangle_i := \langle f_i, g_i \rangle$ .

Now we define  $\prod_I P_i: \prod_I \mathcal{C}_i \rightarrow \mathbf{Pos}$ , on objects  $a \in \text{Ob}(\prod_I \mathcal{C}_i)$  by  $(\prod_I P_i)(a) := \prod_I P_i(a_i)$  and on morphisms  $f \in \prod_I \mathcal{C}_i(a, b)$  by  $(\prod_I P_i)(f) := \prod_I P_i(f_i)$ . Then for  $f: a \rightarrow b$  and  $g: b \rightarrow c$ ,  $(\prod_I P_i)(g \circ f) = \prod_I P_i(g_i \circ f_i) := \prod_I P_i(f_i) P_i(g_i) := \prod_I P_i(f_i) \prod_I P_i(g_i) = (\prod_I P_i)(f) \circ (\prod_I P_i)(g)$ . One also verifies that  $\prod_I P_i(id_a) = id_{\prod_I P_i(a)}$ , so that  $\prod_I P_i$  is a contravariant functor and thus  $(\prod_I \mathcal{C}_i, \prod_I P_i)$  is a prop-category.

For each  $a \in \text{Ob}(\prod_I \mathcal{C}_i)$ , we define  $Eq_a$  and  $e_a$  so that  $Eq_{a, i} := Eq_{a_i}$  and  $e_{a, i} := e_{a_i}$ . Let  $b \in \text{Ob}(\prod_I \mathcal{C}_i)$  and  $c = a \times b$ . Then  $Eq_{\mathbf{1}} = e_{\mathbf{1} \times \mathbf{1}}$  since for each  $i \in I$ ,

$$Eq_{\mathbf{1}, i} = Eq_{\mathbf{1}_{\mathcal{C}_i}} = e_{\mathbf{1}_{\mathcal{C}_i} \times \mathbf{1}_{\mathcal{C}_i}} = e_{\mathbf{1} \times \mathbf{1}, i}.$$

Also,

$$Eq_c = \left( \prod_I P_i \right) \langle \pi_1^{a, b} \pi_1^{c, c}, \pi_1^{a, b} \pi_1^{c, c} \rangle Eq_a \otimes \left( \prod_I P_i \right) \langle \pi_2^{a, b} \pi_1^{c, c}, \pi_2^{a, b} \pi_1^{c, c} \rangle Eq_b,$$



since for each  $j \in I$ ,

$$\begin{aligned}
& ((\prod_I P_i) \langle \pi_1^{a,b} \pi_1^{c,c}, \pi_1^{a,b} \pi_1^{c,c} \rangle Eq_a \otimes (\prod_I P_i) \langle \pi_2^{a,b} \pi_1^{c,c}, \pi_2^{a,b} \pi_1^{c,c} \rangle Eq_b)_j \\
&= P_j \langle \pi_1^{a_j, b_j} \pi_1^{c_j, c_j}, \pi_1^{a_j, b_j} \pi_1^{c_j, c_j} \rangle Eq_{a_j} \otimes P_j \langle \pi_2^{a_j, b_j} \pi_1^{c_j, c_j}, \pi_2^{a_j, b_j} \pi_1^{c_j, c_j} \rangle Eq_{b_j} \\
&= Eq_{c,j}
\end{aligned}$$

For  $\Omega \in \mathcal{L}_q$ ,  $\Omega_{a,b}: \prod_I P_i(a \times b) \rightarrow \prod_I P_i(a)$  is defined so that  $\Omega_{a,b,i} := \Omega_{a_i, b_i}$ . As with the conditions for equality, the remaining conditions for the quantifiers follow from the fact that they hold coordinate-wise. Therefore,  $(\prod_I \mathcal{C}_i, \prod_I P_i) \in \text{Ob}(\mathbf{FA})$ .

For  $j \in I$ , the projection morphism  $\pi_j: (\prod_I \mathcal{C}_i, \prod_I P_i) \rightarrow (\mathcal{C}_j, P_j)$  consists of the projection functor  $\pi_j^o: \prod_I \mathcal{C}_i \rightarrow \mathcal{C}_j$  and the natural transformation  $\pi_j^p: \prod_I P_i \Rightarrow P_j \circ \pi_j^o$  defined for each object  $a \in \text{Ob}(\prod_I \mathcal{C}_i)$ , by  $\pi_{j,a}^p: \prod_I P_i(a) \rightarrow P_j(a_j)$ , i.e.  $\pi_{j,a}^p$  is the (monotone) projection homomorphism. For all  $f: a \rightarrow b \in \text{Mor}(\prod_I \mathcal{C}_i)$ ,

$$\begin{array}{ccc}
b & & \prod_I P_i(b_i) \xrightarrow{\pi_{j,b}^p} P_j(b_j) \\
\downarrow f^{op} & & \downarrow \prod_I P_i(f_i) \quad \quad \quad \downarrow P_j(f_j) \\
a & & \prod_I P_i(a_i) \xrightarrow{\pi_{j,a}^p} P_j(a_j)
\end{array}$$

commutes and so  $\pi_j^p: \prod_I P_i \rightarrow P_j \circ \pi_j^o$  is a natural transformation. For all  $a \in \text{Ob}(\prod_I \mathcal{C}_i)$ ,

$$\pi_{j,a \times a}^p(Eq_a) = Eq_{a,j} = Eq_{a_j} = Eq_{\pi_j^o(a)}.$$

For  $\Omega \in \mathcal{L}_q$ ,  $a, b \in \text{Ob}(\prod_I \mathcal{C}_i)$  and  $A \in (\prod_I P_i)(a \times b)$ ,

$$\pi_{j,a}^p \circ \Omega_{a,b}(A) = \Omega_{a_j, b_j}(A_j) = \Omega_{\pi_j^o(a), \pi_j^o(b)} \circ \pi_{j,a \times b}^p(A).$$

It follows that  $\pi_j: (\prod_I \mathcal{C}_i, \prod_I P_i) \rightarrow (\mathcal{C}_j, P_j)$  is a prop-category morphism.

Let  $(\mathcal{D}, Q) \in \mathbf{FA}$  and for each  $j \in I$ , let  $F_j: (\mathcal{D}, Q) \rightarrow (\mathcal{C}_j, P_j)$  be a morphism in  $\mathbf{FA}$ . Define  $\langle F_i \rangle_I: (\mathcal{D}, Q) \rightarrow (\prod_I \mathcal{C}_i, \prod_I P_i)$ , so that  $\langle F_i \rangle_I^o := \langle F_i^o \rangle_I$  and define  $\langle F_i \rangle_I^p: Q \Rightarrow \prod_I P_i \circ \langle F_i \rangle_I^o$  for  $d \in \text{Ob}(\mathcal{D})$ , by  $\langle F_i \rangle_{I,d}^p := \langle F_{i,d}^p \rangle_I$ . Then for all  $g: d_1 \rightarrow d_2 \in \text{Mor}(\mathcal{D})$ ,

$$\begin{array}{ccc}
d_2 & & Q(d_2) \xrightarrow{\langle F_{i,d_2}^p \rangle_I} \prod_I (P_i \circ F_i^o(d_2)) \\
\downarrow g^{op} & & \downarrow Q(g) \quad \quad \quad \downarrow \prod_I (P_i \circ F_i^o(g)) \\
d_1 & & P_{\mathcal{D}}(d_1) \xrightarrow{\langle F_{i,d_1}^p \rangle_I} \prod_I (P_i \circ F_i^o(d_1))
\end{array}$$

commutes since

$$\begin{aligned} \prod_I (P_i \circ F_i^o(g)) \circ \langle F_{i,d_2}^p \rangle_I &= \langle P_i \circ F_i^o(g) \circ F_{i,d_2}^p \rangle_I \\ &= \langle F_{i,d_1}^p \circ Q(g) \rangle_I \\ &= \langle F_{i,d_1}^p \rangle_I \circ Q(g). \end{aligned}$$

It follows that  $\langle F_i \rangle_I^p: Q \Rightarrow \prod_I P_i \circ \langle F_i \rangle_I^o$  is a natural transformation.

Let  $d \in \text{Ob}(\mathcal{D})$  and  $a \in \text{Mor}(\prod_I C_i)$  such that for each  $j \in I$ ,  $a_j: F_j^o(d \times d) \rightarrow F_j^o d \times F_j^o d$  is the change-of-product isomorphism. Then for each  $j \in I$ ,

$$\langle F_{i,d \times d}^p \rangle_I (Eqd)_j = F_{j,d \times d}^p Eqd = P_j(a_j) Eq_{F_j^o d},$$

and so

$$\langle F_i \rangle_{I,d \times d}^p Eqd = \prod_I P_i(a) Eq_{\langle F_i \rangle_I^o d}$$

Let  $d_1, d_2 \in \text{Ob}(\mathcal{D})$  and  $a \in \text{Mor}(\prod_I C_i)$  such that for each  $j \in I$ ,  $a_j: F_j^o(d_1 \times d_2) \rightarrow F_j^o d_1 \times F_j^o d_2$  is the change-of-product isomorphism. Then, for  $\Omega \in \mathcal{L}_q$ ,

$$\begin{aligned} \langle F_i \rangle_{I,d_1}^p \circ \Omega_{d_1,d_2} &= \langle F_{i,d_1}^p \circ \Omega_{d_1,d_2} \rangle_I \\ &= \langle \Omega_{F_i^o d_1, F_i^o d_2} \circ P(a_i^{-1}) \circ F_{i,d_1 \times d_2}^p \rangle_I \\ &= \Omega_{\langle F_i \rangle_I^o d_1, \langle F_i \rangle_I^o d_2} \circ \prod_I P_i(a^{-1}) \circ \langle F_i \rangle_{I,d_1 \times d_2}^p. \end{aligned}$$

Therefore,  $\langle F_i \rangle_I$  is a morphism. Also, for each  $j \in I$  and  $d \in \text{Ob}(\mathcal{D})$ ,

$$(\pi_{j, \langle F_i \rangle_I^o}^p \cdot \langle F_i \rangle_I^p)_d = \pi_{j, \langle F_i \rangle_I^o d}^p \circ \langle F_{i,d}^p \rangle_I = F_{j,d}^p,$$

and

$$\pi_j^o \circ \langle F_i \rangle_I^o = F_j^o.$$

Thus for each  $j \in I$ ,  $\pi_j \circ \langle F_i \rangle_I = F_j$ . If  $\phi: (\mathcal{D}, Q) \rightarrow (\prod_I C_i, \prod_I P_i)$  is another morphism such that for each  $j \in I$ ,  $\pi_j \circ \phi = F_j$ , then  $\phi^o = F_i^o$  and for each  $d \in \text{Ob}(\mathcal{D})$ ,

$$(\pi_{j, \phi^o}^p \cdot \phi^p)_d = \pi_{j, \langle F_i^o \rangle_I d}^p \circ \phi_d^p = F_{j,d}^p.$$

It follows that  $\phi_d^p = \langle F_i \rangle_{I,d}^p$ , and so  $\phi = \langle F_i \rangle_I$ .  $\square$

**Proposition 4.** *Let  $\mathcal{L}$  be a first-order language and  $\{(C_i, P_i)\}_{i \in I} \subseteq \text{Ob}(\mathbf{FA})$ . Then*

$$\models_{(\prod_I C_i, \prod_I P_i)} \supseteq \models_{\{(C_i, P_i)\}_{i \in I}}.$$

*Proof.* If  $I = \emptyset$ , then  $(\prod_I C_i, \prod_I P_i) = 1_{\mathbf{FA}} = (*, 1_*)$  which satisfies every sequent-in-context and equation-in-context. Thus,  $\models_{1_{\mathbf{FA}}} = \models_{\emptyset}$ . Otherwise, suppose  $T \models_{\{(C_i, P_i)\}_{i \in I}} T'$  and let  $S \in (\prod_I C_i, \prod_I P_i)$  be a  $T$ -model. Then for each  $i \in I$ ,  $\pi_i(S)$  is a  $T$ -model in  $(C_i, P_i)$ . By assumption  $\pi_i(S)$  satisfies  $T'$  for all  $i \in I$  and it follows that  $S$  satisfies  $T'$ .  $\square$

As a consequence, for each logic  $\mathcal{L}$ ,  $\mathbf{FA}_{\mathcal{L}}$  has all products.

## 7 Fibered Homomorphism Theorem

In this section we present a fibered homomorphism theorem which closely mirrors the corresponding result from universal algebra. These results give an orthogonal factorization system for  $\mathbf{FA}_{\mathcal{L}}$ . For a general reference on factorization systems see [19].

Fix a logic  $\mathcal{L}$  and let  $\mathcal{E}$  be the class of morphisms  $\epsilon$  in  $\mathbf{FA}_{\mathcal{L}}$  such that  $\epsilon^o$  is bijective on objects, full and for each  $c \in \text{Ob}(\mathcal{C})$ ,  $\epsilon_c^p$  is surjective. Let  $\mathcal{M}$  be the collection of subprop-morphisms in  $\mathbf{FA}_{\mathcal{L}}$ .

**Theorem 6** (Fibered Homomorphism Theorem). *Let  $F: (\mathcal{C}, P) \rightarrow (\mathcal{D}, Q)$  be a morphism in  $\mathbf{FA}_{\mathcal{L}}$ .*

1.  $F$  factors as  $\psi \circ \epsilon$ , for some  $\epsilon \in \mathcal{E}$  and  $\psi \in \mathcal{M}$ .
2. If  $K: (\mathcal{C}, P) \rightarrow (\mathcal{E}, R) \in \text{Mor}(\mathbf{FA}_{\mathcal{L}})$  where  $K$  is a full, surjective on objects and for each  $e \in \text{Ob}(\mathcal{E})$ ,  $K_e^p$  is surjective, then there exists a unique morphism  $H: (\mathcal{E}, R) \rightarrow (\mathcal{D}, Q)$  such that  $HK = F$  if and only if  $\ker K \leq \ker F$ .

*Proof of Part 1 of Theorem 6.* Let  $T$  be the theory of  $(\mathcal{C}, P)$  and  $S$  the internal structure of  $(\mathcal{C}, P)$ . Let  $T'$  be the theory of  $F(S)$  (which is an  $\mathcal{L}^m$ -theory) and  $\overline{G}$  the generic  $T'$ -model in  $(\mathcal{C}_T, P_T)$ . Let  $\iota: (\mathcal{C}, P) \rightarrow (\mathcal{C}_T, P_T)$  be the morphism which with  $\overline{S}$  witnesses the equivalence of  $(\mathcal{C}, P)$  and  $(\mathcal{C}_T, P_T)$ . We have the following diagram:

$$\begin{array}{ccc} (\mathcal{C}, P) & \xrightarrow{F} & (\mathcal{D}, Q) \\ \iota \downarrow & & \nearrow \overline{F(S)} \\ (\mathcal{C}_T, P_T) & \xrightarrow{\overline{G}} & (\mathcal{C}_{T'}, P_{T'}) \end{array}$$

It is straightforward to verify that the above diagram commutes. Since  $\overline{G} \circ \iota$  is injective on objects, from Proposition 1, the following diagram commutes

$$\begin{array}{ccc} (\mathcal{C}, P) & \xrightarrow{F} & (\mathcal{D}, Q) \\ \searrow \overline{G} \circ \iota & & \nearrow \overline{F(S)} \\ & (\mathcal{C}_{T'}, P_{T'}) & \\ \downarrow H & \uparrow \lambda & \nearrow \overline{F(S)} \circ \lambda \\ & (\overline{G} \circ \iota(\mathcal{C}), \overline{G} \circ \iota(P)) & \end{array}$$

where  $\lambda$  is the inclusion morphism and  $H$  the corestriction. Let  $\epsilon = H$ ,  $\psi = \overline{F(S)} \circ \lambda$  and  $(\overline{G} \circ \iota(\mathcal{C}), \overline{G} \circ \iota(P)) = (\mathcal{E}, R)$ . By construction,  $\epsilon$  and  $\psi$  meet the conditions of the theorem. Since  $\iota$  is a subprop-morphism, from Proposition 2,  $(\mathcal{E}, R) \in \mathbf{FA}_{\mathcal{L}}$ .  $\square$

*Proof of part 2 of Theorem 6.* ( $\Leftarrow$ ) Let  $e_1 \in \text{Ob}(\mathcal{E})$ . Since  $K^o$  is surjective on objects, there exists  $c_1 \in \text{Ob}(\mathcal{C})$ , such that  $K^o(c_1) = e_1$ . Define  $H^o e_1 := F^o c_1$ .

Let  $e_2 \in \text{Ob}(\mathcal{E})$ , and  $h_1: e_1 \rightarrow e_2$ . Then there exists  $c_2 \in \text{Ob}(\mathcal{C})$  such that  $K^o c_2 = e_2$  and since  $K^o$  is full, there exists  $f_1: c_1 \rightarrow c_2$  such that  $K^o f_1 = h_1$ . Define  $H^o h_1 := F^o f_1$ . Since  $\ker K \leq \ker F$ ,  $H^o$  is well defined. If  $h_2: e_2 \rightarrow e_3$ , let  $f_2: c_2 \rightarrow c_3$  such that  $K^o f_2 = h_2$ . Then

$$H^o(h_2 h_1) = H^o(K^o f_2 K^o f_1) = H^o(K^o(f_2 f_1)) = F^o(f_2 f_1) = H^o h_2 H^o h_1.$$

Also,  $H^o(id_{e_1}) = H^o(K^o id_{c_1}) = F^o id_{c_1} = id_{F^o c_1} = id_{H^o e_1}$  and so  $H^o$  is a functor.

Let  $1_{\mathcal{E}}$  be the terminal element in  $\mathcal{E}$ . Then there exists  $c \in \text{Ob}(\mathcal{C})$  such that  $K^o c = 1_{\mathcal{E}}$ . Since  $K^o$  preserves finite products  $K^o 1_{\mathcal{C}}$  is a terminal element in  $\mathcal{E}$  and so  $K^o 1_{\mathcal{C}} \cong 1_{\mathcal{E}}$ . Since  $H^o$  is a functor, it preserves isomorphisms and so  $H^o 1_{\mathcal{E}} \cong H^o K^o 1_{\mathcal{C}} = F^o 1_{\mathcal{C}}$ , and so  $H^o 1_{\mathcal{E}}$  is terminal in  $\mathcal{D}$  since  $F^o$  preserves finite products. Let  $e_1, e_2 \in \text{Ob}(\mathcal{E})$  and  $c_1, c_2, c \in \text{Ob}(\mathcal{C})$ , such that  $K^o c_i = e_i$  and  $K^o c = e_1 \times e_2$ . For  $i \in \{1, 2\}$ , let  $f_i: c \rightarrow c_i$  such that  $K^o f_i = \pi_i^{e_1, e_2}$  and let  $\alpha = \langle K^o \pi_1^{c_1, c_2}, K^o \pi_2^{c_1, c_2} \rangle$ . Since  $K^o$  preserves finite products,  $\alpha$  is an isomorphism. For  $i \in \{1, 2\}$ ,  $K^o \pi_i^{c_1, c_2} K^o \langle f_1, f_2 \rangle = K^o f_i = \pi_i^{e_1, e_2}$  and so  $K^o \langle f_1, f_2 \rangle = \alpha^{-1}$ . Then  $H^o(\alpha^{-1}) = F^o \langle f_1, f_2 \rangle$  is an isomorphism and for  $i \in \{1, 2\}$ ,  $F^o \pi_i^{c_1, c_2} \circ F^o \langle f_1, f_2 \rangle = F^o f_i$ . Thus since  $F^o$  preserves finite products,  $F^o \langle f_1, f_2 \rangle$  is a change-of-product isomorphism from  $(F^o c, F^o f_1, F^o f_2)$  to  $(F^o(c_1 \times c_2), F^o \pi_1^{c_1, c_2}, F^o \pi_2^{c_1, c_2})$ . It follows that  $(H^o(e_1 \times e_2), H^o \pi_1^{e_1, e_2}, H^o \pi_2^{e_1, e_2}) = (F^o c, F^o f_1, F^o f_2)$  is a product diagram for  $H^o e_1, H^o e_2$  and so  $H^o$  preserves finite products.

Let  $e \in \text{Ob}(\mathcal{E})$ , and  $r \in R(e)$ . Then there exists  $c_1 \in \text{Ob}(\mathcal{C})$  and  $r_1 \in P(c_1)$  such that  $K_{c_1}^p(r_1) = r$ . Define  $H_e^p(r) := F_{c_1}^p(r_1)$ . If  $r_2 \in P(c_2)$ , such that  $K_{c_2}^p(r_2) = r$ , then  $(r_1, r_2), (r_2, r_1) \in \text{Oker } K \subseteq \text{Oker } F$  and so  $F_{c_1}^p(r_1) = F_{c_2}^p(r_2)$ . It follows that  $H_e^p$  is well-defined and by definition,  $H_e^p \circ K_{c_1}^p = F_{c_1}^p$ . Let  $r_1, r_2 \in R(e)$  such that  $r_1 \leq r_2$ . Then there exists  $c \in \text{Ob}(\mathcal{C})$  and  $r'_1, r'_2 \in P(c)$  such that  $K_c^p(r'_i) = r_i$  for  $i \in \{1, 2\}$ . Then  $(r'_1, r'_2) \in \text{Oker } K \subseteq \text{Oker } F$ , and so  $H_e^p(r_1) = F_c^p(r'_1) \leq F_c^p(r'_2) = H_e^p(r_2)$ . Thus  $H_e^p$  is monotone.

Let  $h: e_1 \rightarrow e_2$  and  $f: c_1 \rightarrow c_2$  such that  $K^o f = h$ . Then

$$\begin{aligned} Q(H^o h) \circ H_{e_2}^p \circ K_{c_2}^p &= Q(F^o f) \circ F_{c_2}^p = F_{c_1}^p \circ P(f) \\ &= H_{e_1}^p \circ K_{c_1}^p \circ P(f) = H_{e_1}^p \circ R(h) \circ K_{c_2}^p, \end{aligned}$$

and since  $K_{c_2}^p$  is surjective,  $Q(H^o h) \circ H_{e_2}^p = H_{e_1}^p \circ R(h)$ . Thus,  $HP: R \Rightarrow QH^o$  is a natural transformation.

Let  $e \in \text{Ob}(\mathcal{E})$ . Then there exists  $c, c' \in \text{Ob}(\mathcal{C})$ , such that  $K^o c' = e \times e$  and  $K^o c = e$ . Then for  $i \in \{1, 2\}$ , there exists  $f_i: c' \rightarrow c$  such that  $K^o f_i = \pi_i^{e, e}$ . Then for  $i \in \{1, 2\}$ ,

$$K^o \pi_i^{c, c'} K^o \langle f_1, f_2 \rangle = K^o f_i = \pi_i^{e, e}.$$

It follow that  $K^o \langle f_1, f_2 \rangle = \alpha^{-1}$ , where  $\alpha = \langle K^o \pi_1^{c, c'}, K^o \pi_2^{c, c'} \rangle$ . Define:

$$a := \langle F^o \pi_1^{c, c'}, F^o \pi_2^{c, c'} \rangle \quad b := \langle H^o \pi_1^{e, e}, H^o \pi_2^{e, e} \rangle.$$

Then

$$a F^o \langle f_1, f_2 \rangle = \langle F^o f_1, F^o f_2 \rangle = \langle H^o \pi_1^{e, e}, H^o \pi_2^{e, e} \rangle = b,$$

and so  $F^o\langle f_1, f_2 \rangle = a^{-1}b$ . Also note that

$$F_{c \times c}^p(Eq_c) = Q(a)(Eq_{F^o c}) \quad \text{and} \quad K_{c \times c}^p(Eq_c) = R(\alpha)(Eq_{K^o c}).$$

Then

$$\begin{aligned} Q(b)(Eq_{H^o e}) &= Q(b)(Eq_{F^o c}) = Q(b)(Q(a^{-1})F_{c \times c}^p Eq_c) \\ &= Q(a^{-1}b)F_{c \times c}^p(Eq_c) = Q(F^o\langle f_1, f_2 \rangle)F_{c \times c}^p(Eq_c) \\ &= F_{c' \times c'}^p P(\langle f_1, f_2 \rangle)(Eq_c) = H_{e \times e}^p K_{c' \times c'}^p P(\langle f_1, f_2 \rangle)(Eq_c) \\ &= H_{e \times e}^p R(K^o\langle f_1, f_2 \rangle)K_{c \times c}^p(Eq_c) = H_{e \times e}^p (R(\alpha^{-1})K_{c \times c}^p(Eq_c)) \\ &= H_{e \times e}^p (Eq_{K^o c}) = H_{e \times e}^p (Eq_e). \end{aligned}$$

Let  $e_1, e_2 \in \text{Ob}(\mathcal{E})$ , and  $c_1, c_2, c \in \text{Ob}(\mathcal{C})$  such that  $K^o c_i = e_i$  and  $K^o c = e_1 \times e_2$ . Let  $f_i: c \rightarrow c_i$  such that  $K^o f_i = \pi_i^{e_1, e_2}$  and define

$$a := \langle F^o \pi_1^{c_1, c_2}, F^o \pi_2^{c_1, c_2} \rangle \quad b := \langle H^o \pi_1^{e_1, e_2}, H^o \pi_2^{e_1, e_2} \rangle.$$

Also, let  $\alpha := \langle K^o \pi_1^{c_1, c_2}, K^o \pi_2^{c_1, c_2} \rangle$ . Then for  $\Omega \in \mathcal{L}_q$

$$\begin{aligned} &= \Omega_{H^o e_1, H^o e_2} \circ Q(b^{-1}) \circ H_{e_1 \times e_2}^p \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a^{-1}) \circ Q(b^{-1}a) \circ H_{e_1 \times e_2}^p \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a^{-1}) \circ Q(H^o \alpha) \circ H_{e_1 \times e_2}^p \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a^{-1}) \circ H_{K^o(c_1 \times c_2)}^p \circ R(\alpha) \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a^{-1}) \circ H_{K^o(c_1 \times c_2)}^p \circ K_{c_1 \times c_2}^p \\ &= \Omega_{F^o c_1, F^o c_2} \circ Q(a^{-1}) \circ F_{c_1 \times c_2}^p \\ &= F_{c_1}^p \circ \Omega_{c_1, c_2} \\ &= H_{e_1}^p \circ K_{c_1}^p \circ \Omega_{c_1, c_2} \\ &= H_{e_1}^p \circ \Omega_{K^o c_1, K^o c_2} \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p \\ &= H_{e_1}^p \circ \Omega_{e_1, e_2} \circ R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p. \end{aligned}$$

Since  $R(\alpha^{-1}) \circ K_{c_1 \times c_2}^p$  is surjective it follows that

$$\Omega_{H^o e_1, H^o e_2} \circ Q(b^{-1}) \circ H_{e_1 \times e_2}^p = H_{e_1}^p \circ \Omega_{e_1, e_2}.$$

Thus  $H$  is a morphism in **FA** and  $H \circ K = F$ .

( $\implies$ ). Since  $\mathcal{K}$  is full, surjective on objects and for each  $e \in \text{Ob}(\mathcal{E})$ ,  $K_e^p$  is surjective, there is only one possible definition for  $H$ . One may verify that if  $\ker K \not\subseteq \ker F$ , then either  $H^o$  is ill-defined or for some  $e \in \text{Ob}(\mathcal{E})$ ,  $H_e^p$  is ill-defined or not monotone.  $\square$

*Remark 3.* Like part 1 of Theorem 6, the internal logic can also be used to prove part 2. In what follows we sketch the proof of the ( $\longleftarrow$ ) direction. Let

$S$  be the internal structure in  $(\mathcal{E}, R)$  and  $T$  its theory. We construct a  $T$ -model  $A \in (\mathcal{D}, Q)$ , so that

$$\begin{array}{ccc}
 (\mathcal{C}, P) & \xrightarrow{F} & (\mathcal{D}, Q) \\
 & \searrow K & \swarrow \bar{A} \\
 & & (\mathcal{E}, R) \xrightarrow{\bar{A} \circ \iota} (\mathcal{C}_T, P_T) \\
 & & \uparrow \iota
 \end{array}$$

commutes.

For each  $e \in \text{Ob}(\mathcal{E})$ , let  $c_e \in \text{Ob}(\mathcal{C})$  such that  $K^o c_e = e$ . For each  $h \in \text{Mor}(\mathcal{E})$ , let  $f_h \in \text{Mor}(\mathcal{C})$ , such that  $K^o f_h = h$  and for each  $e \in \text{Ob}(\mathcal{E})$  and  $r \in R(e)$ , let  $q_r \in P(c)$  for some  $c \in \text{Ob}(\mathcal{C})$  such that  $K_c^p(q_r) = r$ . For each each sort  $e$ ,  $A[[e]] := F^o c_e$ , for each unary function symbol  $h: e_1 \times \dots \times e_n \rightarrow e$ ,  $A[[h]] := F^o f_h$  and each unary relation symbol  $r \in R(e_1 \times \dots \times e_n)$ ,  $A[[r]] := F_{c_e}^p q_r$ . For each  $n$ -ary function symbol  $h: e_1, \dots, e_n \rightarrow e$ , we define  $A[[h]] := f_h \circ a_{\bar{e}}^{-1}$  where  $a_{\bar{e}}: F^o(c_{\bar{e}}) \rightarrow A[[e_1]] \times \dots \times A[[e_n]]$  is the change-in-product isomorphism. Similarly, for each  $n$ -ary relation symbol  $r \subseteq e_1, \dots, e_n$ , we define  $A[[r]] := Q(a_{\bar{e}})^{-1} \circ F_{c_e}^p(q_r)$ .

By induction, one may prove that for each term  $M: e[\Gamma]$ , where  $\Gamma = x_1: e_1, \dots, x_n: e_n$ , that  $A[[M: e[\Gamma]]] = F^o f_{S[[M: e[\Gamma]]]} \circ a_{\bar{e}}^{-1}$ , and for each formula-in-context  $\phi[\Gamma]$ , that  $A[[\phi[\Gamma]]] = Q(a_{\bar{e}}^{-1}) \circ F_{c_e}^p(q_{S[[\phi[\Gamma]]]})$ . It follows that  $A$  satisfies each equation-in-context in  $T$ , and if  $\phi_1, \dots, \phi_n \vdash \phi_{n+1}[\Gamma]$  is in  $T$ , then

$$\begin{aligned}
 & S[[\phi_1 \otimes \dots \otimes \phi_n[\Gamma]]] \leq S[[\phi_{n+1}[\Gamma]]] \\
 \implies & K_c^p(q_{S[[\phi_1 \otimes \dots \otimes \phi_n[\Gamma]]]}) \leq K_c^p(q_{S[[\phi_{n+1}[\Gamma]]]}) \\
 \implies & F_c^p(q_{S[[\phi_1 \otimes \dots \otimes \phi_n[\Gamma]]]}) \leq F_c^p(q_{S[[\phi_{n+1}[\Gamma]]]}) \\
 \implies & F_c^p(q_{S[[\phi_1[\Gamma]]]}) \otimes \dots \otimes F_c^p(q_{S[[\phi_n[\Gamma]]]}) \leq F_c^p(q_{S[[\phi_{n+1}[\Gamma]]]}) \\
 \implies & A[[\phi_1[\Gamma]]] \otimes \dots \otimes A[[\phi_n[\Gamma]]] \leq A[[\phi_{n+1}[\Gamma]]].
 \end{aligned}$$

It follows that  $A$  is a  $T$ -model. By construction, we have  $K \circ \bar{A} \circ \iota = F$ .

**Proposition 5.** *The classes  $\mathcal{E}$  and  $\mathcal{M}$  form a factorization system in  $\mathbf{FA}_{\mathcal{L}}$ .*

*Proof.* Note that both  $\mathcal{M}$  and  $\mathcal{E}$  contain all the isomorphisms in  $\mathbf{FA}_{\mathcal{L}}$  and are closed under composition. From Part 1 of Theorem 6, each  $F \in \text{Mor}(\mathbf{FA}_{\mathcal{L}})$  can be factored as  $m \circ e$  with  $m \in \mathcal{M}$  and  $e \in \mathcal{E}$ . Now consider the following commuting solid diagram of morphisms in  $\mathbf{FA}_{\mathcal{L}}$ :

$$\begin{array}{ccc}
 \cdot & \xrightarrow{u} & \cdot \\
 e \downarrow & & \downarrow e' \\
 \cdot & \xrightarrow{\exists! w} & \cdot \\
 m \downarrow & & \downarrow m \\
 \cdot & \xrightarrow{v} & \cdot
 \end{array}$$

We want to show there is a unique morphism  $w$  making the small squares commute. Note that  $\text{Th}(e) \subseteq \text{Th}(vme) = \text{Th}(m'e'u) = \text{Th}(e'u)$ , and since  $e$

is bijective on objects,  $\ker(e) \leq \ker(e'u)$ . From part 2 of Theorem 6, there exists a unique morphism  $w$  such that  $we = e'u$ . Moreover it is straightforward to verify that the morphisms in  $\mathcal{E}$  are epimorphisms in  $\mathbf{FA}_{\mathcal{L}}$  and so  $e$  is an epimorphism. From this fact and the fact that the large rectangle and small top square commute, the small bottom square must commute as well.  $\square$

## 8 Algebraic Characterizations of Logical Closure Operators

Let  $Sg$  be a single-sorted algebraic signature,  $\mathbf{Alg}_{Sg}$  the category of set-valued  $Sg$ -algebras and  $Eq_{Sg}$  the class of  $Sg$ -equations. Then there is an adjunction

$$\begin{array}{ccc} & \text{Alg}(\cdot) & \\ & \curvearrowright & \\ \mathcal{P}(Eq_{Sg}) & \perp & \mathcal{P}(\mathbf{Alg}_{Sg})^{op} \\ & \curvearrowleft & \\ & \text{Eq}(\cdot) & \end{array} \quad (7)$$

Where  $\mathcal{P}$  is the operation of taking the powerset and both  $\mathcal{P}(Eq_{Sg})$  and  $\mathcal{P}(\mathbf{Alg}_{Sg})$  are ordered by inclusion. This adjunction determines a closure operator  $Eq \circ \text{Alg}$  on  $\mathcal{P}(Eq_{Sg})$ , and a closure operator  $\text{Alg} \circ Eq$  on  $\mathcal{P}(\mathbf{Alg}_{Sg})$ . From Birkhoff's Completeness Theorem for equational logic,  $Eq \circ \text{Alg}(\Theta)$  is the closure of  $\Theta$  under the derivation rules of equational logic. Birkhoff's HSP Theorem asserts  $\text{Alg} \circ Eq(\cdot)$  is  $\mathbb{H}\mathbb{S}\mathbb{P}(\cdot)$  where  $\mathbb{H}$ ,  $\mathbb{S}$ ,  $\mathbb{P}$  are the operations which close a class of algebras under the operations of taking homomorphic images, subalgebras and products respectively [2].

In the categorical semantics for equational logic,  $\mathbf{Alg}_{Sg} \equiv \mathbf{FP}(\mathcal{C}_{Sg}, \mathbf{Set})$  where  $\mathbf{FP}$  is the 2-category of categories with finite products and product preserving functors. Birkhoff's HSP Theorem interpreted in the categorical semantics says  $\mathcal{Y} \subseteq \text{Ob}(\mathbf{FP}(\mathcal{C}_{Sg}, \mathbf{Set}))$  is an equational class iff  $\mathbb{H}\mathbb{S}\mathbb{P}(\mathcal{Y}) = \mathcal{Y}$ , where

1.  $\mathbb{P}(\mathcal{Y})$  is the smallest class containing  $\mathcal{Y}$  stable under products.
2.  $\mathbb{S}(\mathcal{Y})$  is the smallest class containing  $\mathcal{Y}$  such that whenever  $F \in \mathcal{Y}$  and  $\eta: K \Rightarrow F$  is a monomorphism, then  $K \in \mathbb{S}(\mathcal{Y})$ .
3.  $\mathbb{H}(\mathcal{Y})$  is the smallest class containing  $\mathcal{Y}$  such that whenever  $F \in \mathcal{Y}$ , and  $\eta: F \Rightarrow K$  is a regular epimorphism, then  $K \in \mathbb{H}(\mathcal{Y})$ .

For an analogous characterization for a multi-sorted algebraic signature see [1]. Since we are considering first-order logics, we are interested in a similar result where  $\mathcal{P}(Eq_{Sg})$  is replaced by the complete lattice of all  $Sg$ -theories  $\text{Th}_{Sg}$  and algebras by first-order structures. If  $Sg$  is single sorted and  $\mathcal{L}$  is a language for classical first-order logic, then the Tarskian  $Sg$ -structures of classical first-order logic can be identified with morphisms  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathbf{Set}, \mathcal{P})$ , where  $\mathcal{P}$  is the preimage functor. It is well-known that closing a class of Tarskian

structures under their common theory, is equivalent to closing the class under ultraproducts, isomorphic copies and ultraroots [8, p. 454]. In what follows, we develop an analogous result for an arbitrary first-order logic  $\mathcal{L}$  and its general prop-categorical semantics.

Let  $\text{Mod}_{Sg}^{\mathcal{L}}$  be the collection of all  $Sg$ -structures in  $\mathbf{FA}_{\mathcal{L}}$ . Then we are interested in characterizing  $\text{Mod}_{(\cdot)} \circ \text{Th}$ , where

$$\begin{array}{ccc} & \xrightarrow{\text{Mod}_{(\cdot)}} & \\ \text{Th}_{Sg} & \perp \mathcal{P}(\text{Mod}_{Sg}^{\mathcal{L}})^{op} & \\ & \xleftarrow{\text{Th}_{(\cdot)}} & \end{array}$$

We take  $\text{Mod}_{Sg}^{\mathcal{L}}$  to be the collection of all morphisms  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P)$  such that  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA}_{\mathcal{L}})$ . Then for each  $\mathcal{Y} \subseteq \text{Mod}_{Sg}^{\mathcal{L}}$ ,  $\text{Th}(\mathcal{Y})$  is the  $Sg$ -theory whose assertions are  $\{a \in A_{Sg} : \forall F \in \mathcal{Y}, G[[a]] \in \ker F\}$  and for each  $T \in \text{Th}_{Sg}$ ,  $\text{Mod}_T = \{F \in \text{Mod}_{Sg}^{\mathcal{L}} : G[[T]] \subseteq \ker F\}$ . If  $\mathbf{FA}_{\mathcal{L}}$  forms a complete semantics for  $\mathcal{L}$ ,  $\text{Th}(\text{Mod}_T)$  is the  $Sg$ -theory whose assertions are  $\{a \in A_{Sg} : T \vdash_{\mathcal{L}} a\}$ . For  $\mathcal{Y} \subseteq \text{Mod}_{Sg}^{\mathcal{L}}$ , we wish to characterize  $\text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ . Let  $\{F_i: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_i, P_i)\}_{i \in I} \subseteq \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ . Then each  $(\mathcal{C}_i, P_i) \in \text{Ob}(\mathbf{FA}_{\mathcal{L}})$ , and from Proposition 4,  $(\prod \mathcal{C}_i, \prod P_i) \in \text{Ob}(\mathbf{FA}_{\mathcal{L}})$ , and so  $\langle F_i \rangle_{i \in I} \in \text{Mod}_{Sg}^{\mathcal{L}}$ . Moreover,  $\langle F_i \rangle_{i \in I}$  satisfies each assertion satisfied by all  $F_i$  and so  $\langle F_i \rangle_{i \in I} \in \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ . We define the **(external) product** of  $\{F_i\}_{i \in I}$  to be  $\langle F_i \rangle_{i \in I}$  and let  $\mathbb{P}(\mathcal{Y})$  denote the closure of  $\mathcal{Y}$  under external products. And so,  $\text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$  is stable under taking external products.

Let  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P) \in \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ . Then for each  $H \in \mathbf{FA}_{\mathcal{L}}((\mathcal{C}, P), (\mathcal{D}, Q))$ ,  $H \circ F \in \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ . We say  $H \circ F$  is an **(external) homomorphic image** of  $F$  and let  $\mathbb{H}(\mathcal{Y})$  denote the closure of  $\mathcal{Y}$  under external homomorphic images. If  $H: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{E}, R)$  and  $\iota: (\mathcal{E}, R) \rightarrow (\mathcal{C}, P)$  are morphisms such that  $F = \iota \circ H$  and  $\iota$  is a subprop-morphism, then we call  $H$  an **(external) submodel** of  $F$  and let  $\mathbb{S}(\mathcal{Y})$  denote the closure of  $\mathcal{Y}$  under taking external submodels. Since  $G[[\text{Th}(\mathcal{Y})]] \subseteq \ker F$ ,  $\iota \circ H = F$  and  $\iota$  is a subprop-morphism,  $G[[\text{Th}(\mathcal{Y})]] \subseteq \ker H$  and so  $H \in \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ .

*Remark 4.* If we consider  $T$ -algebras as product preserving functors  $\{F_i: \mathcal{C}_T \rightarrow \mathcal{C}\}_{i \in I}$ , where  $\mathcal{C}$  has arbitrary products then,

$$\begin{array}{ccc} \mathcal{C}_T & \xrightarrow{F_{\times}} & \mathcal{C} \\ & \searrow \langle F_i \rangle_{i \in I} & \nearrow \times \\ & & \prod_{i \in I} \mathcal{C} \end{array}$$

$a \Downarrow$

commutes up to a change-in-product natural isomorphism  $a$ , where  $\times$  is the right adjoint to the diagonal functor  $\text{Diag}: \mathcal{C} \rightarrow \prod_{i \in I} \mathcal{C}$  and  $F_{\times}$  is the usual “internal” product of the algebras  $\{F_i\}_{i \in I}$ . Since  $\times$  is faithful,  $\langle F_i \rangle_{i \in I}$  and  $F_{\times}$  satisfy the same equations. Then the external product  $\langle F_i \rangle_{i \in I}$  satisfies the same equational theory as the internal product  $F_{\times}$ . For first-order models, the



internal product need not correspond to the external. The issue is that for  $c \in \text{Ob}(\prod_{i \in I} \mathcal{C})$ , in general,  $\prod_{i \in I} P(c_i) \not\cong P(\times(c))$ .

**Theorem 7.** *Let  $\mathcal{L}$  be a logic and  $Sg$  a signature. For each  $\mathcal{Y} \subseteq \text{Mod}_{Sg}^{\mathcal{L}}$ ,  $\text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y}) = \mathbb{HSP}(\mathcal{Y})$ .*

*Proof.* If  $\mathcal{Y}$  is the class of all  $Sg$ - $\mathcal{L}$  structures of a given theory  $T$ , then we showed  $\mathcal{Y}$  is closed under taking (external) products, submodels and homomorphic images and so  $\mathbb{HSP}(\mathcal{Y}) \subseteq \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ .

In the other direction, let  $T = \text{Th}(\mathcal{Y})$ ,  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}, P)$  be a  $T$ -model, where  $(\mathcal{C}, P) \in \text{Ob}(\mathbf{FA}_{\mathcal{L}})$  and let  $\{T_i\}_{i \in I}$  be the collection of all  $Sg$ - $\mathcal{L}$  theories such that there exists  $F_i: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_i, P_i) \in \mathcal{Y}$  such that  $\text{Th}(F_i) = T_i$ . Then  $T = \bigcap_{i \in I} T_i$ , and define  $\epsilon_T := \overline{G}: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_T, P_T)$ , where  $G \in (\mathcal{C}_T, P_T)$  is the generic  $T$ -model. Then  $\ker \epsilon_T \leq \ker \langle F_i \rangle_I$ , and so from Theorem 6, there exists  $\iota: (\mathcal{C}_T, P_T) \rightarrow (\prod_I \mathcal{C}_i, \prod_I P_i)$  such that

$$\begin{array}{ccc} (\mathcal{C}_{Sg}, P_{Sg}) & \xrightarrow{\langle F_i \rangle_I} & (\prod_I \mathcal{C}_i, \prod_I P_i) \\ & \searrow \epsilon_T & \nearrow \iota \\ & (\mathcal{C}_T, P_T) & \end{array}$$

commutes. Moreover, since  $\text{Th}(\langle F_i \rangle_I) = T$ ,  $\iota$  is a subprop-morphism. From Proposition 2,  $(\mathcal{C}_T, P_T) \in \text{Ob}(\mathbf{FA}_{\mathcal{L}})$  and so  $\epsilon_T \in \mathbb{SP}(\mathcal{Y}) = \mathcal{Y}$ . Since  $F$  is a  $T$ -model,  $\ker \epsilon_T \leq \ker F$ . From Theorem 6, there exists a morphism  $\iota: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{C}, P)$ , such that

$$\begin{array}{ccc} (\mathcal{C}_{Sg}, P_{Sg}) & \xrightarrow{F} & (\mathcal{C}, P) \\ & \searrow \epsilon & \nearrow \iota \\ & (\mathcal{C}_T, P_T) & \end{array}$$

commutes and so  $F \in \mathbb{HSP}(\mathcal{Y})$ . Therefore,  $\mathbb{HSP}(\mathcal{Y}) \supseteq \text{Mod}_{(\cdot)} \circ \text{Th}(\mathcal{Y})$ .  $\square$

Let  $\text{Log}$  be the partial order of all logics. We now consider the following adjunction:

$$\begin{array}{ccc} & \text{Ob}(\mathbf{FA}_{(\cdot)}) & \\ & \xrightarrow{\quad} & \\ \text{Log} & \perp \mathcal{P}(\text{Ob}(\mathbf{FA}))^{op} & (8) \\ & \xleftarrow{\quad} & \\ & \text{F}_{(\cdot)} & \end{array}$$

In order to give a characterization of the closure operator  $\text{Ob}(\mathbf{FA}_{\text{F}_{(\cdot)}})$ , we first restrict the logics in  $\text{Log}$  to some fixed signature  $Sg$ . Let  $\text{Log}^{Sg}$  be the collection of all logics restricted to  $Sg$ -assertions and we denote the corresponding

restrictions of  $\text{Ob}(\mathbf{FA}_{(\cdot)})$  and  $\vDash_{(\cdot)}$ ,  $\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg})$  and  $\vDash_{(\cdot)}^{Sg}$  respectively.

$$\begin{array}{ccc} & \text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) & \\ & \curvearrowright & \\ \text{Log}^{Sg} & \perp & \mathcal{P}(\text{Ob}(\mathbf{FA}))^{op} \\ & \curvearrowleft & \\ & \vDash_{(\cdot)}^{Sg} & \end{array} \quad (9)$$

In the context of the (untyped) equational logic Adjunction 9 corresponds to the following: Let  $\mathcal{L}_\omega$  be an algebraic signature and  $V$  a set of variables of cardinality  $\lambda$ . Let  $Eq_V$  be the collection of  $\mathcal{L}_\omega$ -equations over  $V$  which we identify with  $Fm_V^2$ , the square of the formula-algebra. A collection of  $\mathcal{L}_\omega$ -algebras  $\mathcal{A}$  determines a structural closure operator  $\vDash_{\mathcal{A}}$ , defined by  $\Theta \vDash_{\mathcal{A}} \epsilon = \delta$  if and only if for all homomorphisms  $h: Fm_V \rightarrow A$ , where  $A \in \mathcal{A}$ , if  $\Theta \subseteq \ker h$ , then  $\epsilon = \delta \in \ker h$ . Observe that  $\vDash_{\mathbb{SP}(\mathcal{A})} = \vDash_{\mathcal{A}}$ , but  $\mathbb{SP}(\mathcal{A})$  may not be the largest collection of algebras defining the same consequence. In [3] it is shown that the largest such class of  $\mathcal{L}_\omega$ -algebras is  $\mathbb{U}_\lambda \mathbb{SP}(\mathcal{A})$ , where  $B \in \mathbb{U}_\lambda(\mathcal{A})$  if every  $\lambda$ -generated subalgebra of  $B$  is in  $\mathcal{A}$ .

Taking the operation  $\mathbb{U}_\lambda$  as inspiration, where  $\lambda$  up to renaming specifies the propositional signature, given a signature  $Sg$ , and  $\mathcal{X}$  a collection of prop-categories, we define  $\mathbb{U}_{Sg}(\mathcal{X})$  so that  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}(\mathcal{X})$ , if each classifying subprop-category  $(\mathcal{C}_T, P_T)$  of  $(\mathcal{D}, Q)$  is in  $\mathcal{X}$ , where  $\text{Sg}(T) = Sg$ . For  $\mathcal{X} \subseteq \text{Ob}(\mathbf{FA})$ , we take  $\mathbb{P}(\mathcal{X})$  to be the closure of  $\mathcal{X}$  under taking products and  $\mathbb{S}(\mathcal{X})$  to be the closure of  $\mathcal{X}$  under taking subprop-categories.

**Theorem 8.** *For each  $\mathcal{X} \subseteq \text{Ob}(\mathbf{FA})$ ,  $\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X}) = \mathbb{U}_{Sg} \mathbb{SP}(\mathcal{X})$ .*

*Proof.* From Proposition 2 and Proposition 4,  $\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X})$  is stable under  $\mathbb{S}$  and  $\mathbb{P}$ . Let  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}(\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X}))$  and suppose  $T \vDash_{\mathcal{X}} a$ . Let  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{D}, Q)$  be a  $T$ -model in  $(\mathcal{D}, Q)$  and  $T' = \text{Th}(F)$ . From Theorem 6 there exists  $\iota: (\mathcal{C}_{T'}, P_{T'}) \rightarrow (\mathcal{D}, Q)$  which makes the diagram

$$\begin{array}{ccc} (\mathcal{C}_{Sg}, P_{Sg}) & \xrightarrow{F} & (\mathcal{D}, Q) \\ & \searrow \epsilon_{T'} & \nearrow \iota \\ & (\mathcal{C}_{T'}, P_{T'}) & \end{array}$$

commute. Moreover, since  $\text{Th}(F) = \text{Th}(\epsilon_{T'})$ ,  $\iota$  is a subprop-morphism. Then  $(\mathcal{C}_{T'}, P_{T'}) \in \text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X})$ , because  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}(\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X}))$ . Since  $T \vDash_{\mathcal{X}} a$  and  $G[[T]] \subseteq \ker \epsilon_{T'}$ ,  $G[[a]] \in \ker \epsilon_{T'}$ . Thus  $G[[a]] \in \ker F$  and so  $T \vDash_{(\mathcal{D}, Q)} a$ . It follows that  $(\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X})$  and so  $\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X})$  is stable under  $\mathbb{U}_{Sg}$ . Therefore,  $\text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X}) \supseteq \mathbb{U}_{Sg} \mathbb{SP}(\mathcal{X})$ .

Now let  $(\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA}_{(\cdot)}^{Sg}) \circ \vDash_{(\cdot)}^{Sg}(\mathcal{X})$  and consider a subprop-morphism  $\iota: (\mathcal{C}_T, P_T) \rightarrow (\mathcal{D}, Q)$  such that  $\text{Sg}(T) = Sg$ . If  $a \in A_{Sg} \setminus A(T)$ , then the generic

$T$ -model  $G$  in  $(\mathcal{C}_T, P_T)$  does not satisfy  $a$  and so  $T \not\models_{(\mathcal{C}_T, P_T)} a$ . Since  $\models_{(\mathcal{C}_T, P_T)} \supseteq \models_{(\mathcal{D}, Q)} \supseteq \models_{\mathcal{X}}$  for each  $a \notin T$ , there exists a  $T$ -model  $F_a: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{C}_a, P_a)$  such that  $(\mathcal{C}_a, P_a) \in \mathcal{X}$  and  $G[[a]] \notin \ker F_a$ . Then  $\ker(\epsilon_T) \leq \ker(\langle F_a \rangle_{a \notin T})$  and so there exists a morphism  $\lambda: (\mathcal{C}_T, P_T) \rightarrow \prod_{a \notin T} (\mathcal{C}_a, P_a)$  such that

$$\begin{array}{ccc} (\mathcal{C}_{Sg}, P_{Sg}) & \xrightarrow{\langle F_a \rangle_{a \notin T}} & \prod_{a \notin T} (\mathcal{C}_a, P_a) \\ & \searrow \epsilon_T & \nearrow \lambda \\ & & (\mathcal{C}_T, P_T) \end{array}$$

commutes. Moreover,  $\text{Th}(\epsilon_T) = \text{Th}(\langle F_a \rangle_{a \notin T})$  and so  $\lambda$  is a subprop-morphism. Thus  $(\mathcal{C}_T, P_T) \in \mathbb{S}\mathbb{P}(\mathcal{X})$  and so  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}\mathbb{S}\mathbb{P}(\mathcal{X})$ .  $\square$

Adjunction 9 is relevant when one wants to consider a logic over a fixed signature. However, in categorical logic, where internal logic is used, it makes sense to have a logic be independent of a particular signature. Thus we also provide a characterization of the closure  $\text{Ob}(\mathbf{FA}_{\models_{(\cdot)}})$  in Adjunction 8. For  $\mathcal{X} \subseteq \text{Ob}(\mathbf{FA})$ , we define  $\mathbb{U}(\mathcal{X})$  by  $(\mathcal{D}, Q) \in \mathbb{U}(\mathcal{X})$ , if each classifying subprop-category  $(\mathcal{C}_T, P_T)$  of  $(\mathcal{D}, Q)$  is in  $\mathcal{X}$ . That is,  $(\mathcal{D}, Q) \in \mathbb{U}(\mathcal{X})$  if and only if for each signature  $Sg$ ,  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}(\mathcal{X})$ .

**Theorem 9.**  $\text{Ob}(\mathbf{FA}_{\models_{(\cdot)}}) = \mathbb{U}\mathbb{S}\mathbb{P}(\cdot)$ .

*Proof.* Clearly  $\text{Ob}(\mathbf{FA}_{\models_{\mathcal{X}}})$  is stable under  $\mathbb{S}$  and  $\mathbb{P}$ . Suppose  $(\mathcal{D}, Q) \in \mathbb{U}(\text{Ob}(\mathbf{FA}_{\models_{\mathcal{X}}}))$ . Suppose  $T \models_{\mathcal{X}} T'$  and let  $Sg = \text{Sg}(T)$ . Consider  $F: (\mathcal{C}_{Sg}, P_{Sg}) \rightarrow (\mathcal{D}, Q)$  such that  $T \subseteq \text{Th}(F)$ . Then  $F$  factors as

$$\begin{array}{ccc} (\mathcal{C}_{Sg}, P_{Sg}) & \xrightarrow{F} & (\mathcal{D}, Q) \\ & \searrow \epsilon_{T''} & \nearrow \iota \\ & & (\mathcal{C}_{T''}, P_{T''}) \end{array}$$

where  $T'' = \text{Th}(F) \supseteq T$  and  $\iota$  is a subprop-morphism. Then  $(\mathcal{C}_{T''}, P_{T''}) \in \text{Ob}(\mathbf{FA}_{\models_{\mathcal{X}}})$ , and so  $T' \subseteq T'' = \text{Th}(F)$  and so  $T \models_{(\mathcal{D}, Q)} T'$ . Thus  $\text{Ob}(\mathbf{FA}_{\models_{(\cdot)}})$  is stable under  $\mathbb{U}$ .

Now suppose  $(\mathcal{D}, Q) \in \text{Ob}(\mathbf{FA}_{\models_{\mathcal{X}}})$ . Then  $\models_{(\mathcal{D}, Q)} \supseteq \models_{\mathcal{X}}$ . Following the proof of Theorem 8, one shows for each signature  $Sg$ ,  $(\mathcal{D}, Q) \in \mathbb{U}_{Sg}\mathbb{S}\mathbb{P}(\mathcal{X})$  and so  $(\mathcal{D}, Q) \in \mathbb{U}\mathbb{S}\mathbb{P}(\mathcal{X})$ .  $\square$

We would like to thank Adam Přenosil for his many insights and helpful suggestions on this paper.

## References

- [1] J. Adámek, J. Rosický, and E. M. Vitale. *Algebraic theories*. Vol. 184. Cambridge Tracts in Mathematics. A categorical introduction to general algebra, With a foreword by F. W. Lawvere. Cambridge University Press, Cambridge, 2011, pp. xviii+249. ISBN: 978-0-521-11922-1.

- [2] Garrett Birkhoff. “On the Structure of Abstract Algebras”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 31.4 (1935), pp. 433–454. DOI: 10.1017/S0305004100013463.
- [3] W. J. Blok and Bjarni Jónsson. “Equivalence of consequence operations”. In: *Studia Logica* 83.1-3 (2006). With a preface by Jónsson, pp. 91–110. ISSN: 0039-3215. DOI: 10.1007/s11225-006-8299-z. URL: <https://doi-org.proxy.library.vanderbilt.edu/10.1007/s11225-006-8299-z>.
- [4] Tim Button and Sean Walsh. *Philosophy and model theory*. With a historical appendix by Wilfrid Hodges. Oxford University Press, Oxford, 2018, pp. xvi+517. ISBN: 978-0-19-879040-2; 978-0-19-879039-6. DOI: 10.1093/oso/9780198790396.001.0001.
- [5] Alonzo Church. *Introduction to mathematical logic. Vol. I*. Princeton University Press, Princeton, N. J., 1956, pp. x+376.
- [6] Josep Maria Font. *Abstract algebraic logic*. Vol. 60. Studies in Logic (London). An introductory textbook, Mathematical Logic and Foundations. College Publications, London, 2016, pp. xxix+521. ISBN: 978-1-84890-207-7.
- [7] Petr Hájek. *Metamathematics of fuzzy logic*. Vol. 4. Trends in Logic—Studia Logica Library. Kluwer Academic Publishers, Dordrecht, 1998, pp. viii+297. ISBN: 0-7923-5238-6. DOI: 10.1007/978-94-011-5300-3. URL: <https://doi-org.proxy.library.vanderbilt.edu/10.1007/978-94-011-5300-3>.
- [8] Wilfrid Hodges. *Model theory*. Vol. 42. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 1993, pp. xiv+772. ISBN: 0-521-30442-3. DOI: 10.1017/CB09780511551574. URL: <https://doi.org/10.1017/CB09780511551574>.
- [9] Bart Jacobs. *Categorical logic and type theory*. Vol. 141. Studies in Logic and the Foundations of Mathematics. North-Holland Publishing Co., Amsterdam, 1999, pp. xviii+760. ISBN: 0-444-50170-3.
- [10] M. Kaufmann. “The quantifier “there exist uncountably many” and some of its relatives”. In: *Model-theoretic logics*. Perspect. Math. Logic. Springer, New York, 1985, pp. 123–176.
- [11] Stephen Cole Kleene. *Introduction to metamathematics*. D. Van Nostrand Co., Inc., New York, N. Y., 1952, pp. x+550.
- [12] Erich Peter Klement, Radko Mesiar, and Endre Pap. *Triangular norms*. Vol. 8. Trends in Logic—Studia Logica Library. Kluwer Academic Publishers, Dordrecht, 2000, pp. xx+385. ISBN: 0-7923-6416-3. DOI: 10.1007/978-94-015-9540-7. URL: <https://doi-org.proxy.library.vanderbilt.edu/10.1007/978-94-015-9540-7>.
- [13] F. William Lawvere. “Adjointness in foundations”. In: *Repr. Theory Appl. Categ.* 16 (2006). Reprinted from *Dialectica* 23 (1969), pp. 1–16.
- [14] F. William Lawvere. “Equality in hyperdoctrines and comprehension schema as an adjoint functor”. In: *Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968)*. Amer. Math. Soc., Providence, R.I., 1970, pp. 1–14.

- [15] Yoshihiro Maruyama. “Fibred algebraic semantics for a variety of non-classical first-order logics and topological logical translation”. In: *J. Symb. Log.* 86.3 (2021), pp. 1189–1213. ISSN: 0022-4812. DOI: 10.1017/jsl.2021.41. URL: <https://doi-org.proxy.library.vanderbilt.edu/10.1017/jsl.2021.41>.
- [16] Radko Mesiar and Helmut Thiele. “On  $T$ -quantifiers and  $S$ -quantifiers”. In: *Discovering the world with fuzzy logic*. Vol. 57. Stud. Fuzziness Soft Comput. Physica, Heidelberg, 2000, pp. 310–326.
- [17] A. Mostowski. “On a generalization of quantifiers”. In: *Fund. Math.* 44 (1957), pp. 12–36. ISSN: 0016-2736. DOI: 10.4064/fm-44-1-12-36. URL: <https://doi-org.proxy.library.vanderbilt.edu/10.4064/fm-44-1-12-36>.
- [18] Andrew M. Pitts. “Categorical logic”. In: *Handbook of logic in computer science, Vol. 5*. Vol. 5. Handb. Log. Comput. Sci. Oxford Univ. Press, New York, 2000, pp. 39–128.
- [19] Emily Riehl. “Factorization Systems”. In: Algebraic Topology and Category Theory Proseminar in Fall 2008 at the University of Chicago.
- [20] Hiroyuki Shirasu. “Duality in superintuitionistic and modal predicate logics”. In: *Advances in modal logic, Vol. 1 (Berlin, 1996)*. Vol. 87. CSLI Lecture Notes. CSLI Publ., Stanford, CA, 1998, pp. 223–236.
- [21] Hiroyuki Shirasu. “Glueing of algebras for substructural logics”. In: 927. Problems concerning nonclassical logics and their Kripke semantics (Japanese) (Kyoto, 1995). 1995, pp. 127–139.
- [22] Dag Westerståhl. “Generalized Quantifiers”. In: *The Stanford Encyclopedia of Philosophy*. Ed. by Edward N. Zalta. Winter 2019. Metaphysics Research Lab, Stanford University, 2019.
- [23] Ryszard Wójcicki. “Referential matrix semantics for propositional calculi”. In: *Logic, methodology and philosophy of science, VI (Hannover, 1979)*. Vol. 104. Studies in Logic and the Foundations of Mathematics. North-Holland, Amsterdam-New York, 1982, pp. 325–334.