

An Equivalence Principle for the Spectrum of Random Inner-Product Kernel Matrices

Yue M. Lu and Horng-Tzer Yau

Harvard University

May 16, 2022

Abstract

We consider random matrices whose entries are obtained by applying a (nonlinear) kernel function to the pairwise inner products between n independent data vectors drawn uniformly from the unit sphere in \mathbb{R}^d . Our study of this model is motivated by problems in machine learning, statistics, and signal processing, where such inner-product kernel random matrices and their spectral properties play important roles. Under mild conditions on the kernel function, we establish the weak-limit of the empirical spectral distribution of these matrices when $d, n \rightarrow \infty$ such that $n/d^\ell \rightarrow \kappa \in (0, \infty)$, for some fixed $\ell \in \mathbb{N}$ and $\kappa \in \mathbb{R}$. This generalizes an earlier result of Cheng and Singer [9], who studied the same model in the linear scaling regime (with $\ell = 1$ and $n/d \rightarrow \kappa$). The main insight of our work is a general *equivalence principle*: the spectrum of the random kernel matrix is asymptotically equivalent to that of a simpler matrix model, constructed as the linear combination of a (shifted) Wishart matrix and an independent matrix drawn from the Gaussian orthogonal ensemble. The aspect ratio of the Wishart matrix and the coefficients of the linear combination are determined by ℓ and by the expansion of the kernel function in the orthogonal Hermite polynomial basis. Consequently, the limiting spectrum of the random kernel matrix can be characterized as the free additive convolution between a Marchenko-Pastur law and a semicircle law.

Contents

1	Introduction	2
1.1	Notation	3
2	An Asymptotic Equivalence Principle	4
2.1	Polynomial Kernel Functions	4
2.2	A Heuristic Explanation of the Equivalence Principle	5
2.3	General Nonlinear Kernels	7
2.4	Numerical Experiments	9
3	Related Work	10
4	Proof of the Main Result	11
4.1	Outline of the Proof	11
4.2	Schur Complement: Reparameterization	12
4.3	Schur Complement: the Leading Term	15
4.4	Proof of Theorem 1	19
5	Summary and Discussions	20
A	Gegenbauer Polynomials and Spherical Harmonics	21

B	Proof of Proposition 1	22
C	Review of the Marchenko-Pastur Law	26
D	Moment Bounds and Concentration Inequalities	27
E	The Operator Norms of $\{A_k\}$	32
E.1	The Low-Order Components	32
E.2	Moment Calculations for the High-Order Components	33
E.3	Bounds on the Operator Norms	37
F	Perturbations of Resolvents and Stieltjes Transforms	38
G	Stability Analysis of the Self-Consistent Equation	40

1 Introduction

Let $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^d$ be a set of independent data vectors drawn uniformly from the unit sphere $\mathcal{S}^{(d-1)}$. Consider a random matrix $A \in \mathbb{R}^{n \times n}$ with entries

$$A_{ij} := \begin{cases} \frac{1}{\sqrt{n}} f_d(\sqrt{d} \mathbf{x}_i^\top \mathbf{x}_j) & \text{if } i \neq j \\ 0 & \text{if } i = j \end{cases}, \quad (1.1)$$

where $f_d : \mathbb{R} \mapsto \mathbb{R}$ is a (nonlinear) “kernel” function. Its empirical spectral distribution (ESD) is

$$\varrho_A(x) = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}(x)$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A , and δ_λ denotes the counting measure. In this work, we establish the weak-limit of the ESD ϱ_A when $d, n \rightarrow \infty$ such that $n/d^\ell \rightarrow \kappa \in (0, \infty)$, for some fixed $\kappa \in \mathbb{R}$ and $\ell \in \mathbb{N}$.

Our study of this model is motivated by recent problems in machine learning, statistics, and signal processing, where random matrices in the form of (1.1) and their spectral properties play important roles. Examples include kernel methods (such as kernel-PCA [33] and kernel-SVM [32]), covariance thresholding procedures [5, 12], nonlinear dimension reduction [3], and probabilistic matrix factorization [29]. Moreover, the closely-related non-Hermitian version of (1.1), where $A_{ij} = \frac{1}{n} f_d(\sqrt{d} \mathbf{x}_i \mathbf{y}_j)$ for two sets of vectors $\{\mathbf{x}_i\}_{i \leq n}$ and $\{\mathbf{y}_j\}_{j \leq p}$, appears in the random feature model [31, 26, 20, 27, 21] that has received a lot of recent attention as an interesting theoretical model for large neural networks.

In this work, we consider a general high-dimensional polynomial scaling regime, where the number of samples n can grow to infinity proportional to d^ℓ , for some positive integer ℓ . The linear regime, with $\ell = 1$, was previously studied by Cheng and Singer [9], who showed that the ESD of A converges to a deterministic limit distribution whose Stieltjes transform can be characterized as the solution to a cubic equation. (See Section 3 for further discussions on related work in the literature.) An interesting observation was made by Fan and Montanari [18], who pointed out that the limiting ESD of A (as determined by the aforementioned cubic equation) is equal to that of

$$B = \frac{\mu\sqrt{d}}{\sqrt{n}}(X^\top X - I) + \gamma H, \quad (1.2)$$

where $X = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{d \times n}$ is a matrix whose columns are the data vectors, H is an independent matrix drawn from the Gaussian orthogonal ensemble (GOE), and μ, γ are two constants that depend on f_d . Comparing the models in (1.1) and (1.2), one may interpret the effect of the *nonlinear* kernel function f_d as a *noisy linear* transformation applied to the shifted Gram matrix $X^\top X - I$.

The main result of this work is to show that the above characterization obtained under the linear scaling regime is a special case of a general *equivalence principle* that holds when $n \asymp d^\ell$ for any $\ell \in \mathbb{N}$ and under

mild conditions on the kernel f_d . Specifically, we show in Theorem 2 that the ESD of A is asymptotically equal to that of

$$B = \frac{\mu_\ell}{\sqrt{nN_\ell}}(W^\top W - N_\ell I) + \gamma_\ell H$$

where $W \in \mathbb{R}^{N_\ell \times n}$ is an i.i.d. standard Gaussian matrix with aspect ratio $N_\ell/n \rightarrow 1/(\kappa\ell!)$, and H is a GOE matrix independent of W . The constants μ_ℓ and γ_ℓ both depend on ℓ , and they can be computed by expanding f_d in the orthogonal Hermite polynomial basis. As an immediate consequence of this equivalence principle, the limiting ESD of A can be simply characterized as a free additive convolution between a (shifted) Marchenko-Pastur (MP) law and a semicircle law.

The rest of the paper is organized as follows. We start Section 2 by considering the special case when the kernel function f_d is a polynomial. We formalize the equivalence principle for these polynomial kernels in Theorem 1, with a simple heuristic explanation provided in Section 2.2. The case when f_d is a general nonlinear function is addressed in Section 2.3, where the corresponding asymptotic characterizations are stated in Theorem 2. We present in Section 2.4 several numerical experiments to illustrate our theory. Section 3 contains further discussions on related work in the literature. We devote Section 4 to the proof of Theorem 1, and collect several auxiliary results in the appendix. Finally, we conclude the paper in Section 5 with some discussions on possible extensions of our results as well as open problems.

1.1 Notation

Before starting our technical discussions, we first collect some notations that are used throughout this paper.

Common sets: \mathbb{N} denotes the set of positive integers, and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For each $n \in \mathbb{N}$, $[n] := \{1, 2, \dots, n\}$, and $(n)_k := n(n-1)\dots(n-k+1)$ is the falling factorial. For $i, j \in \mathbb{N}$, $\mathbb{1}_{ij}$ denotes the Kronecker delta function, *i.e.*, $\mathbb{1}_{ij} = 1$ if $i = j$ and $\mathbb{1}_{ij} = 0$ otherwise. We always use z to denote a complex number in the upper-half plane \mathbb{C}_+ , and we consistently write

$$\eta := \Im(z) > 0.$$

The dimension of the data vectors is denoted by d . The unit sphere in \mathbb{R}^d is written as \mathcal{S}^{d-1} .

Probability distributions: $\text{Unif}(\mathcal{S}^{d-1})$ denotes the uniform probability measure on \mathcal{S}^{d-1} . For two independent vectors $\mathbf{x}, \mathbf{y} \sim \text{Unif}(\mathcal{S}^{d-1})$, define

$$\mathbf{s}_d^1 := \text{Law}(\sqrt{d}\mathbf{x}^\top \mathbf{y}). \quad (1.3)$$

By the rotational symmetry of $\text{Unif}(\mathcal{S}^{d-1})$, we also have $\sqrt{d}\mathbf{x}^\top \mathbf{e}_1 \sim \mathbf{s}_d^1$, where $\mathbf{e}_1 = (1, 0, \dots, 0)$. Given a random variable X , its L^p norm is denoted by $\|X\|_{L^p} := (\mathbb{E}|X|^p)^{1/p}$. We say a symmetric random matrix is drawn from the GOE, if its law is equal to that of $(F + F^\top)/\sqrt{2n}$, where F is an $n \times n$ matrix with i.i.d. standard normal entries.

Stochastic order notation: In our proof, we will use a notion of high-probability bounds known as *stochastic domination*. This notion was first introduced in [17, 16], and it provides a convenient way to keep track of low-probability exceptional events where some bounds do not hold. Let

$$X = (X^{(d)}(u) : d \in \mathbb{N}, u \in U^{(d)}), \quad Y = (Y^{(d)}(u) : d \in \mathbb{N}, u \in U^{(d)})$$

be two families of nonnegative random variables, where $U^{(d)}$ is a possibly d -dependent parameter set. We say that X is *stochastically dominated* by Y , uniformly in u , if for all (small) $\varepsilon > 0$ and (large) $D > 0$ we have

$$\sup_{u \in U^{(d)}} \mathbb{P}[X^{(d)}(u) > d^\varepsilon Y^{(d)}(u)] \leq d^{-D}$$

for large enough $d \geq d_0(\varepsilon, D)$. If X is stochastically dominated by Y , uniformly in u , we use the notation $X \prec Y$. Moreover, if for some complex family X we have $|X| \prec Y$, we also write $X = \mathcal{O}_\prec(Y)$.

Vectors and matrices: For a vector $\mathbf{v} \in \mathbb{R}^n$, its ℓ_2 norm is denoted by $\|\mathbf{v}\|$. For a matrix $A \in \mathbb{R}^{n \times n}$, $\|A\|_{\text{op}}$ and $\|A\|_{\text{F}}$ denote the operator (spectral) norm and the Frobenius norm of A , respectively. On the other hand, $\|A\|_\infty := \max_{i,j \in [n]} |A(i,j)|$ denotes the entry-wise ℓ_∞ norm. The trace of A is written as $\text{tr}(A)$. Finally, $\mathbf{1}$ denotes $(1, 1, \dots, 1)$, and I is an identity matrix. Their dimensions can be inferred from the context.

2 An Asymptotic Equivalence Principle

2.1 Polynomial Kernel Functions

We start by considering the special case when the kernel function f_d in (1.1) is a degree- L polynomial, for some $L \in \mathbb{N}$. For reasons that will be made clear later (see Section 2.2), we expand f_d in the following form:

$$f_d(x) = \sum_{0 \leq k \leq L} \mu_k q_k^{(d)}(x) \quad (2.1)$$

where $\{\mu_k\}_k$ is a set of expansion coefficients, and $q_k^{(d)}(x)$ denotes the k th Gegenbauer polynomial [11, 14]. The Gegenbauer polynomials, also known as the ultraspherical polynomials in the literature [22], form a set of orthogonal polynomial basis with respect to the probability measure \mathfrak{s}_d^1 in (1.3). Specifically, we have $\deg q_k^{(d)}(x) = k$ and

$$\mathbb{E}_{\xi \sim \mathfrak{s}_d^1} [q_i^{(d)}(\xi) q_j^{(d)}(\xi)] = \mathbb{1}_{ij}, \quad (2.2)$$

where $\mathbb{1}_{ij}$ is the Kronecker delta.

The coefficients of these polynomials can be determined by performing the Gram-Schmidt procedure on the monomial basis $\{0, 1, x, x^2, \dots\}$ and by using the explicit formula for the moments of ξ [see (D.3)]. The first few polynomials in the sequence are

$$q_0^{(d)}(x) = 1, \quad q_1^{(d)}(x) = x, \quad q_2^{(d)}(z) = \frac{1}{\sqrt{2}} \sqrt{\frac{d+2}{d-1}} (x^2 - 1), \quad (2.3)$$

and

$$q_3^{(d)}(x) = \frac{1}{\sqrt{6}} \sqrt{\frac{d+4}{d-1}} \left(\frac{d+2}{d} x^3 - 3x \right). \quad (2.4)$$

Notice that the coefficients of the Gegenbauer polynomials depend on the dimension d . To lighten the notation, we will suppress this dependence by writing throughout the paper

$$q_k(x) := q_k^{(d)}(x) \quad \text{and} \quad \tilde{q}_k(x) := q_k^{(d-1)}(x). \quad (2.5)$$

In Appendix A, we collect a list of properties of Gegenbauer polynomials that will be used in our analysis.

For each $k \in \mathbb{N}_0$, define a matrix $A_k \in \mathbb{R}^{n \times n}$ such that

$$(A_k)_{ij} := \begin{cases} \frac{1}{\sqrt{n}} q_k(\sqrt{d} \mathbf{x}_i^\top \mathbf{x}_j) & \text{if } i \neq j \\ 0 & \text{if } i = j \end{cases}, \quad (2.6)$$

where $\{\mathbf{x}_i\}_{i \in [n]}$ is a collection of independent vectors drawn from $\text{Unif}(\mathcal{S}^{d-1})$. In light of (2.1), we can write the inner-product kernel matrix in (1.1) as

$$A = \sum_{k=0}^L \mu_k A_k = \left(\sum_{k=0}^{\ell-1} + \sum_{k=\ell}^L \right) \mu_k A_k, \quad (2.7)$$

where $\ell \in \mathbb{N}$ and $\ell \leq L$. The main result of this paper is to prove an asymptotic *equivalence principle* for the empirical spectral distribution (ESD) of A : when $n = \kappa d^\ell + o(d^\ell)$ for some $\kappa > 0$, the empirical spectral distribution (ESD) of A is asymptotically equivalent to that of a matrix $B \in \mathbb{R}^{n \times n}$, defined as

$$B := \sum_{k=\ell}^L \mu_k B_k. \quad (2.8)$$

The first component in the sum in (2.8) is a (shifted) Wishart matrix, constructed as

$$B_\ell = \frac{1}{\sqrt{n N_\ell}} W^\top W - \sqrt{\frac{N_\ell}{n}} I, \quad (2.9)$$

where $W \in \mathbb{R}^{N_\ell \times n}$ is an i.i.d. Gaussian matrix with $N_\ell := d^\ell/\ell! + o(d^\ell)$; for each $k > \ell$, B_k is an $n \times n$ GOE matrix; moreover, $B_\ell, B_{\ell+1}, \dots, B_L$ are jointly *independent*.

For $z \in C_+$, let $s_A(z)$ and $s_B(z)$ denote the Stieltjes transforms of the ESDs of A and B , respectively. The following theorem formalizes the asymptotic equivalence between the matrix models A and B .

Theorem 1. Fix $\kappa, \tau > 0$, two positive integers $\ell \leq L$, and a set of coefficients $\{\mu_k\}_{0 \leq k \leq L}$. Suppose that $n/d^\ell = \kappa + \mathcal{O}(d^{-1/2})$, and $z = E + i\eta$ with $|E| \leq \tau^{-1}$ and $\tau \leq \eta \leq \tau^{-1}$. Then, almost surely as $d \rightarrow \infty$, we have $s_A(z) \rightarrow m(z)$ and $s_B(z) \rightarrow m(z)$, where $m(z)$ is the unique solution in C_+ to the equation

$$m(z) \left(z + \frac{\gamma_a m(z)}{1 + \gamma_b m(z)} + \gamma_c m(z) \right) + 1 = 0. \quad (2.10)$$

Here, $\gamma_a := \mu_\ell^2$, $\gamma_b := \mu_\ell \sqrt{\ell! \kappa}$, and $\gamma_c := \sum_{\ell+1 \leq k \leq L} \mu_k^2$. Moreover, for every $\varepsilon > 0$ and $D > 0$, we have

$$\mathbb{P} \left(|s_A(z) - m(z)| \geq d^{-(1/2-\varepsilon)} \right) \leq d^{-D} \quad (2.11)$$

for all sufficiently large d .

Remark 1. As B is the linear combination of a (shifted) Wishart matrix and independent GOE matrices, its limiting eigenvalue density is given by an additive free convolution of a Marchenko-Pastur law with a semicircle law. Explicit formulas for the limiting density function

$$\varrho_{\text{fc}}(x) := \lim_{\eta \downarrow 0} \frac{1}{\pi} \Im m(x + i\eta) \quad (2.12)$$

can be found in [9, Appendix A]. As a consequence of Theorem 1, the spectral distribution of A converges weakly almost surely to the same limit $\rho_{\text{fc}}(E)$ as $d \rightarrow \infty$.

Remark 2. Note that those components of A associated with indices $k < \ell$ do not play any role in the equivalent model given by B . This is because those “low-order” components $\{A_k\}_{k < \ell}$ are essentially low-rank matrices whose ranks are of order $\mathcal{O}(d^{\ell-1})$. Thus they do not contribute to the limiting spectrum density of the $n \times n$ matrix with $n \asymp d^\ell$. See Lemma 2 for details.

2.2 A Heuristic Explanation of the Equivalence Principle

The equivalence conjecture stated above has a simple heuristic explanation. To see that, we first recall the following important property of the Gegenbauer polynomials: For any $\mathbf{x}_i, \mathbf{x}_j \in \mathcal{S}^{d-1}$,

$$q_k(\sqrt{d} \mathbf{x}_i^\top \mathbf{x}_j) = \frac{1}{\sqrt{N_k}} \sum_{a \in [N_k]} Y_{k,a}(\mathbf{x}_i) Y_{k,a}(\mathbf{x}_j), \quad (2.13)$$

where $\{Y_{k,a}(\mathbf{x}_i)\}_{a \in [N_k]}$ is a set of orthonormal degree- k spherical harmonics associated with \mathbf{x}_i , and

$$N_k = d^k/k!(1 + \mathcal{O}(d^{-1})) \quad (2.14)$$

is the cardinality of the set. We refer the reader to Appendix A for more details, including an exact expression for N_k . The above identity is useful because it allows us to “linearize” the term $q_k(\sqrt{d} \mathbf{x}^\top \mathbf{y})$ as an inner product of two N_k -dimensional vectors made of the spherical harmonics.

Using (2.13) and an additional identity that $q_k(\sqrt{d}) = \sqrt{N_k}$, we can now rewrite the matrices $\{A_k\}$ in a factorized form:

$$A_k = \frac{1}{\sqrt{n N_k}} S_k^\top S_k - \sqrt{\frac{N_k}{n}} I, \quad (2.15)$$

where S_k is a $N_k \times n$ matrix whose entries are the spherical harmonics, *i.e.*

$$(S_k)_{ai} = Y_{k,a}(\mathbf{x}_i), \quad \text{for } a \in [N_k], i \in [n] \quad (2.16)$$

and $\{\mathbf{x}_i\}_{i \in [n]}$ are the data vectors in the definition in (2.6). By construction, the i th column of S_k only depends on \mathbf{x}_i . It follows from the independence of $\{\mathbf{x}_i\}_i$ that the columns of S_k are fully independent. The entries within each column are indeed dependent, but one can check from the orthonormality of the spherical harmonics (see (A.4) in Appendix A) that these entries are uncorrelated random variables with unit variances. In other words, the columns of S_k are independent and isotropic random vectors in \mathbb{R}^{N_k} .

Heuristically, if we replace the entries of S_k by i.i.d. standard Gaussians, we can then expect that the limiting ESD of A_k to be characterized by the Marchenko-Pastur (MP) law with an aspect ratio parameter

$$\phi_k := N_k/n. \quad (2.17)$$

For $k = \ell$, this is indeed the intuition behind the equivalent model B_ℓ constructed in (2.9). (See Appendix C for a review of relevant properties of the MP law.) Note that, since we set the diagonal entries of A_k to zero, there is an additional shift (by $\sqrt{\phi_k}$) of the eigenvalues in (2.9).

According to the above heuristic arguments, the limiting ESD of A_k for every k should be characterized by the MP law. However, in our equivalent model given by B , the higher order components, *i.e.*, B_k with $k > \ell$, are associated with the semicircle law. This discrepancy can be explained by examining the density function associated with the (shifted) MP law. By setting $\phi = \phi_k$ and $t = 1/\sqrt{\phi_k}$ in (C.4), we get

$$\varrho_k(x) dx := \frac{\sqrt{[(2+x-1/\sqrt{\phi_k})(2-x-1/\sqrt{\phi_k})]_+}}{2\pi(1+x/\sqrt{\phi_k})} dx + (1-\phi_k)_+ \delta(x+\sqrt{\phi_k}) dx. \quad (2.18)$$

This density function is completely determined by the aspect ratio parameter ϕ_k defined in (2.17). Recall the expression for N_k in (2.14) and that $n = \kappa d^\ell + o(d^\ell)$. For $k = \ell$, we have $\phi_k \rightarrow (\ell!\kappa)^{-1}$ as $d \rightarrow \infty$. In this case, (2.18) corresponds to an MP law with an aspect ratio of size $\mathcal{O}(1)$. However, for $k > \ell$, we have $\phi_k \asymp d^{k-\ell}$ and thus $1/\sqrt{\phi_k} = \mathcal{O}(1/\sqrt{d})$. In this case, MP law in (2.18) degenerates to the standard semicircle law

$$\varrho_{sc} := \frac{1}{2\pi} \sqrt{(4-x^2)_+} dx. \quad (2.19)$$

Theorem 1 rigorously establishes the above heuristic arguments. In addition, it shows that, despite the apparent statistical dependence among $\{A_k\}_{k \geq \ell}$, they can be treated as a collection of asymptotically independent matrices. Consequently, the self-consistent equation in (2.10) corresponds to the additive free convolution of an MP law with a semicircle law. As a heuristic way to obtain (2.10), we can use (2.15) and (2.16) to write

$$A_{ij} = \frac{1}{n} \sum_{0 \leq k \leq L} t_k \sum_{a \in [N_k]} Y_{k,a}(\mathbf{x}_i) Y_{k,a}(\mathbf{y}_j) - \left(\frac{1}{n} \sum_{0 \leq k \leq L} t_k N_k \right) I, \quad t_k := \frac{\mu_k \sqrt{n}}{\sqrt{N_k}}. \quad (2.20)$$

Observe that this has the form of the matrix given in (C.1) (in Appendix C). Again, by assuming that the family $\{Y_{k,a}(\mathbf{x})\}$ are not just uncorrelated but are indeed independent random variables, A satisfies the MP law in (C.2), *i.e.*,

$$\begin{aligned} \frac{1}{m} &= -z - \frac{1}{n} \left[\sum_{k < \ell} + \sum_{k = \ell} + \sum_{k > \ell} \right] \frac{N_k t_k^2 m}{1 + t_k m} \\ &= -z - \left[\sum_{k < \ell} + \sum_{k = \ell} + \sum_{k > \ell} \right] \frac{\mu_k^2 m}{1 + \mu_k \sqrt{n/N_k} m} \\ &= -z - \frac{\mu_\ell^2 m}{1 + \mu_\ell \sqrt{\ell! \kappa} m} + \left(\sum_{k > \ell} \mu_k^2 \right) m + \mathcal{O}(d^{-1/2}), \end{aligned} \quad (2.21)$$

where in the last step we have used the fact that $n/N_\ell \rightarrow \kappa \ell!$, $n/N_k \rightarrow \infty$ for $k < \ell$ and $n/N_k \rightarrow 0$ for $k > \ell$. Observe that, after dropping the $\mathcal{O}(d^{-1/2})$ error term, (2.21) is exactly the self-consistent equation in (2.10).

2.3 General Nonlinear Kernels

In this section, we extend the equivalence principle presented in Theorem 1 to the case when the kernel f_d in (1.1) is a general function beyond polynomials. Our analysis uses the following comparison inequality.

Lemma 1. *Let A be the matrix in (1.1), and \hat{A} be another matrix constructed in the same way, but with f_d replaced by \hat{f}_d . For $z \in C_+$ with $\eta = \Im m[z]$, let $s(z)$ and $\hat{s}(z)$ denote the Stieltjes transforms of A and \hat{A} , respectively. Then we have*

$$|s(z) - \hat{s}(z)| \leq \frac{1}{\eta^2} (\mathbb{E}_{\xi \sim s_d^1} [f_d(\xi) - \hat{f}_d(\xi)]^2)^{1/2} + \mathcal{O}_{\prec} \left(\frac{1}{\eta \sqrt{n}} \right).$$

Proof. By the triangular inequality,

$$\begin{aligned} |s(z) - \hat{s}(z)| &\leq |\mathbb{E}s(z) - \mathbb{E}\hat{s}(z)| + |s(z) - \mathbb{E}s(z)| + |\hat{s}(z) - \mathbb{E}\hat{s}(z)| \\ &\leq \mathbb{E}|s(z) - \hat{s}(z)| + \mathcal{O}_{\prec}((\eta \sqrt{n})^{-1}), \end{aligned}$$

where the second step uses the standard concentration inequality collected in Lemma 13. To complete the proof, we apply the perturbation estimate (F.7) in Lemma 18 and Hölder's inequality to write

$$\mathbb{E}|s(z) - \hat{s}(z)| \leq \frac{1}{\sqrt{n}\eta^2} \mathbb{E}\|A - \hat{A}\|_{\mathbb{F}} \leq \frac{1}{\sqrt{n}\eta^2} (\mathbb{E}\|A - \hat{A}\|_{\mathbb{F}}^2)^{1/2} \leq \frac{1}{\eta^2} (\mathbb{E}[f_d(\xi) - \hat{f}_d(\xi)]^2)^{1/2}.$$

In the last step, we have used the fact that, for any $i, j \in [n]$ with $i \neq j$, the probability distribution of $\sqrt{n}(A_{ij} - \hat{A}_{ij})$ equals to that of $f_d(\xi) - \hat{f}_d(\xi)$, with $\xi \sim s_d^1$. \square

In what follows, we establish an equivalence principle for the matrix in (1.1) under the following assumption on the function f_d .

Assumption 1. *Let $\xi^{(d)} \sim s_d^1$. The function f_d in (1.1) satisfies the following conditions:*

(a) *For each $k \in \mathbb{N}_0$, we have*

$$\mathbb{E}[f_d(\xi^{(d)}) q_k^{(d)}(\xi^{(d)})] \xrightarrow{d \rightarrow \infty} \mu_k,$$

for some finite numbers $\{\mu_k\}$.

(b) $\sigma^2 := \sum_{k=0}^{\infty} \mu_k^2 \leq \infty$, and $\mathbb{E}[f_d(\xi^{(d)})]^2 \rightarrow \sigma^2$ as $d \rightarrow \infty$.

In the above expressions, we have temporarily brought back the superscripts in $\xi^{(d)}$ and $q_k^{(d)}$ to emphasize the fact that both depend on the dimension d . As in the rest of the paper, we will omit the superscripts and make the d -dependence implicit in the following discussions.

Remark 3. *It is possible to further relax the conditions in Assumption 1 and consider a larger class of functions that can be approximated by a finite number of Gegenbauer polynomials uniformly over d . See [9, Sec. 3.2] for details. On the other hand, when $f_d = f$ is a function that does not depend on d , there are simple sufficient conditions one can use. Specifically, by [9, Lemma C.2, Lemma C.5] (see also Lemma 8 in Appendix A), a function f satisfies Assumption 1 if (a) f is locally bounded, i.e., $\sup_{|x| \leq t} |f(x)| < \infty$ for every $t > 0$; and (b) $\mathbb{E}[f^2(z) \mathbb{1}(|z| \geq t)] \rightarrow 0$ as $t \rightarrow \infty$, where $z \sim \mathcal{N}(0, 1)$. Moreover, in this case the limits in Assumption 1 can be explicitly computed as*

$$\mu_k = \mathbb{E}[f(z) h_k(z)] \quad \text{and} \quad \sigma^2 = \sum_{k=0}^{\infty} \mu_k^2 = \mathbb{E}f^2(z), \tag{2.22}$$

where $h_k(x)$ is the k th (normalized) Hermite polynomial (see Appendix A).

Theorem 2. *Let A be the matrix in (1.1) with the function f_d satisfying the conditions in Assumption 1. $s_A(z)$ denotes its Stieltjes transform. Fix $\kappa, \tau > 0$, and suppose that $n/d^\ell = \kappa + \mathcal{O}(d^{-1/2})$, and $z = E + i\eta$*

with $|E| \leq \tau^{-1}$ and $\tau \leq \eta \leq \tau^{-1}$. Then, $s_A(z) \rightarrow m(z)$ almost surely as $d \rightarrow \infty$, where $m(z)$ is the unique solution in C_+ to the equation

$$m(z) \left(z + \frac{\gamma_a m(z)}{1 + \gamma_b m(z)} + \hat{\gamma}_c m(z) \right) + 1 = 0. \quad (2.23)$$

Here, $\gamma_a = \mu_\ell^2$, $\gamma_b = \mu_\ell \sqrt{\ell! \kappa}$, $\hat{\gamma}_c = \sigma^2 - \sum_{0 \leq k \leq \ell} \mu_k^2$, and $\{\mu_k\}, \sigma^2$ are the constants defined in Assumption 1.

Remark 4. Recall from Section 2.2 that the self-consistent equation (2.23) characterizes the free additive convolution of a (shifted) MP law and the semicircle law. It follows that the statement of Theorem 2 can still be interpreted in terms of an equivalence principle: the limiting ESD of A is equivalent to that of

$$B = \mu_\ell B_\ell + (\sigma^2 - \sum_{0 \leq k \leq \ell} \mu_k^2)^{1/2} H$$

where B_ℓ is the matrix defined in (2.9) and H is an independent GOE matrix.

Proof. By Assumption 1(b), for any fixed (small) $c > 0$, there exists some integer $L \geq \ell + 1$ such that

$$\sigma^2 - (\eta^4 c^2 / 24) \leq \sum_{0 \leq k \leq L-1} \mu_k^2 \leq \sigma^2. \quad (2.24)$$

For $\xi \sim s_d^1$, we write $\langle f, g \rangle := \mathbb{E}[f(\xi)g(\xi)]$ and $\|f\| := [\mathbb{E}f^2(\xi)]^{1/2}$. Now define two functions

$$P_{f_d, L}(x) = f_d(x) - \sum_{0 \leq k \leq L} \langle f_d, q_k \rangle q_k(x)$$

and

$$\hat{f}_d(x) = \sum_{0 \leq k \leq L-1} \mu_k q_k(x) + \hat{\mu}_L q_L(x),$$

where $\hat{\mu}_L := (\sigma^2 - \sum_{0 \leq k \leq L-1} \mu_k^2)^{1/2}$. We then have

$$\begin{aligned} \|f_d - \hat{f}_d\|^2 &= \left\| \sum_{0 \leq k \leq L-1} (\langle f_d, q_k \rangle - \mu_k) q_k(x) + (\langle f_d, q_L \rangle - \hat{\mu}_L) q_L(x) + P_{f_d, L}(x) \right\|^2 \\ &= \sum_{0 \leq k \leq L-1} (\langle f_d, q_k \rangle - \mu_k)^2 + (\langle f_d, q_L \rangle - \hat{\mu}_L)^2 + (\|f_d\|^2 - \sum_{0 \leq k \leq L} \langle f_d, q_k \rangle^2) \\ &\leq (\mu_L - \hat{\mu}_L)^2 + (\sigma^2 - \sum_{0 \leq k \leq L} \mu_k^2) + \eta^4 c^2 / 24, \quad \text{for all sufficiently large } d. \end{aligned} \quad (2.25)$$

In the second step, we have used the property that the Gegenbauer polynomials are orthonormal [see (2.2)] In the last step, we have used Assumption 1(a), which guarantees that the inequality (2.25) holds for all $d \geq d(L, c)$, where $d(L, c)$ is some integer that might depend on L and c . By (2.24), $\hat{\mu}_L^2 \leq \eta^4 c^2 / 24$ and $\mu_L^2 \leq \eta^4 c^2 / 24$. We can then further bound the right-hand side of (2.25) as

$$\|f_d - \hat{f}_d\|^2 \leq 2\mu_L^2 + 2\hat{\mu}_L^2 + (\sigma^2 - \sum_{0 \leq k \leq L} \mu_k^2) + \eta^4 c^2 / 24 \leq \eta^4 c^2 / 4, \quad (2.26)$$

for all sufficiently large d .

Let \hat{A} be a matrix constructed according to (1.1) but with the function f_d replaced by \hat{f}_d . Let $s_{\hat{A}}$ be its Stieltjes transform. By construction, \hat{f}_d is a linear combination of $L + 1$ Gegenbauer polynomials. Thus, we can apply Theorem 1 to get

$$|s_{\hat{A}}(z) - m(z)| \prec \frac{1}{\sqrt{d}}, \quad (2.27)$$

where $m(z) \in \mathbb{C}_+$ is the solution to (2.23). By using the triangular inequality in the first step, and by applying Lemma 1 and (2.27) in the second step, we have

$$\begin{aligned} |s_A(z) - m(z)| &\leq |s_A(z) - s_{\hat{A}}(z)| + |s_{\hat{A}}(z) - m(z)| \\ &\leq \frac{1}{\eta^2} [\mathbb{E}(f_d(\xi) - \hat{f}_d(\xi)^2)]^{1/2} + \mathcal{O}_\prec\left(\frac{1}{\eta\sqrt{n}}\right) + \mathcal{O}_\prec\left(\frac{1}{\sqrt{d}}\right) \\ &\leq c/2 + \mathcal{O}_\prec\left(\frac{1}{\sqrt{d}}\right), \end{aligned}$$

where the last inequality follows from (2.26). Thus, for any $D > 0$,

$$\mathbb{P}(|s_A(z) - m(z)| > c) < d^{-D}$$

for all sufficiently large d . Applying the Borel-Cantelli lemma (for any fixed $D > 1$), we can then conclude that $s_A(z)$ converges to $m(z)$ almost surely. \square




Figure 1: Demonstration of the equivalence principle in the quadratic scaling regime, where $n \asymp \kappa d^2$. In this experiment, $d = 300$ and $\kappa = 0.15$. The ESDs of A_2 , A_3 , and A_4 are plotted against their corresponding limiting spectral densities given by Theorem 1. Observe that the ESD of A_2 is characterized by the MP law in (C.4) (with shape parameters $\phi = 1/(2\kappa)$ and $t = 1/\sqrt{\phi}$), whereas the ESDs of A_3 and A_4 both follow the standard semicircle law given in (2.19).




Figure 2: Linear combinations of the Gegenbauer polynomial matrices. (a) The ESD of the matrix defined in (2.28). In this experiment, $d = 100$ and $n = 1.8 \times 10^4$, and thus $n/d^2 = 1.8$. (b) The ESD of the matrix defined in (2.29). Here, we set $d = 200$, $n = 2 \times 10^4$. The truncation threshold is set to $M = 3$ in (2.30).

2.4 Numerical Experiments

In this subsection we present some numerical experiments to demonstrate the equivalence principle stated in Theorems 1 and 2. The first example considers the ESD of the individual polynomial component matrix A_k defined in (2.6). In the quadratic scaling regime, where $n \asymp \kappa d^2$ for some fixed κ , A_2 is asymptotically equivalent to B_2 in (2.9). Thus, the ESD of A_2 converges to the MP law, with a density function given in (C.4). Both A_3 and A_4 are asymptotically equivalent to a GOE matrix, and thus their ESDs converge to the standard semicircle law given in (2.19). We observe from Figure 1 that the ESDs of A_2 , A_3 and A_4 indeed closely match their corresponding limiting spectral densities.

In the second example, we consider linear combinations of the polynomial matrices $\{A_k\}$. The equivalence principle formalized in Theorem 1 states that the ESD of any fixed linear combination of A_k can be obtained by a free additive convolution between the MP law and the semicircle law. In Figure 2a, we plot the ESD of

$$-A_2 + \frac{1}{\sqrt{20}}A_3 + \frac{1}{\sqrt{20}}A_4 \quad (2.28)$$

against the limiting spectral density. The theoretical curve (red solid line in the figure) is computed by solving the self-consistent equation (2.10) and then evaluating (2.12) numerically. In Figure 2b, we show the ESD of another matrix in the form of

$$A_2 + \frac{1}{4}A_3 + \frac{1}{4}A_4 + \frac{1}{4}A_5^M + \frac{1}{4}A_6^M. \quad (2.29)$$

Here, A_5^M and A_6^M are the truncated version of A_5 and A_6 , respectively. Specifically, for $M > 0$, we have

$$(A_5^M)_{ij} = (A_5)_{ij} \cdot \mathbf{1}(|(A_5)_{ij}| \leq M/\sqrt{n}) \quad i, j \in [n], \quad (2.30)$$

and A_6^M is defined similarly. Note that we apply this truncation to remove a small number of outliers in the entries of A_5 and A_6 that have very large magnitudes. The presence of these outliers will exacerbate




Figure 3: The ESD of the inner-product kernel matrix (1.1) where f_d is the soft-thresholding function in (2.31) with $\tau = 1$. In the experiment, we choose $\tau = 1$, $\kappa = 0.15$, and set the dimension to $d = 1.2 \times 10^4$, $d = 300$ and $d = 50$ in the linear, quadratic, and cubic scaling regimes, respectively.

the finite-size effect, making the ESD deviate from the limiting spectral density when the dimension d is not too large. Due to the truncation step, the matrix in (2.29) is no longer a finite linear combination of $\{A_k\}$. Thus, we use Theorem 2 and the formulas in (2.22) to calculate the theoretical curve.

In the last example, we consider the ESD of the matrix in (1.1), where the nonlinear function is a soft-thresholding operator, *i.e.*,

$$f_d(x) = \text{sign}(x)(|x| - \tau)_+ \quad (2.31)$$

for some threshold $\tau > 0$. Figure 3 compares the ESDs of this matrix against the limiting spectral densities in the linear ($n \asymp \kappa d$), quadratic ($n \asymp \kappa d^2$), and cubic ($n \asymp \kappa d^3$) scaling regimes, respectively. Since $\mathbb{E}_{\xi \sim \mathcal{S}_d^1}[f_d(\xi)q_1(\xi)] > 0$, the matrix A has a nonzero “projection” in A_1 . As we show in Lemma 14 in Appendix E, A_1 is essentially a rank- d matrix with d large eigenvalues of size $\mathcal{O}(\sqrt{n/d})$. It follows that, in the quadratic and cubic scaling regimes, the spectrum of A contains d outlier eigenvalues that are separated from the bulk.

3 Related Work

In this section, we discuss several related lines of work in the literature.

Random kernel matrices: As one of the earlier studies on the spectrum of random kernel matrices, Koltchinskii and Giné [25] considered a setting where the data vectors $\{\mathbf{x}_i\}$ are sampled from a fixed manifold in \mathbb{R}^d . They showed that, when $n \rightarrow \infty$ with d fixed, the spectrum of the kernel matrix converges to that of an integral operator on the manifold. The high-dimensional setting, where $n, d \rightarrow \infty$ with $n/d \rightarrow \kappa \in (0, \infty)$, was first studied by El Karoui [15]. That work considered a scaling where the matrix elements A_{ij} are $f(\mathbf{x}_i^\top \mathbf{x}_j)$. When the data vectors $\{\mathbf{x}_i\}$ are sampled independently from $\text{Unif}(\mathcal{S}^{d-1})$, the typical size of $\mathbf{x}_i^\top \mathbf{x}_j$ is $\mathcal{O}(1/\sqrt{d})$ for $i \neq j$. Consequently, the kernel matrix only depends on the properties of the kernel function $f(x)$ in a local neighborhood near $x = 0$. Indeed, El Karoui [15] showed that, under mild local smoothness assumption on $f(x)$, the kernel matrix is asymptotically equivalent to a simpler matrix obtained via a first-order Taylor expansion. Our work, as well as the work of [9], considers a different scaling: the presence of the \sqrt{d} factor in (1.1) means that the kernel function f_d is applied to values of size $\mathcal{O}(1)$ rather than $\mathcal{O}(1/\sqrt{d})$. Consequently, the spectrum of the matrix in (1.1) depends on the global properties of the kernel function.

The linear asymptotic regime of the model in (1.1) was first studied by Cheng and Singer [9], who established the weak limit of the ESD of the kernel matrix. The idea of “decorrelation” by expanding the nonlinear kernel function via orthogonal polynomial basis was first introduced in that work, and it plays important roles in our paper. The results of [9] were obtained for data vectors sampled from the isotropic Gaussian and spherical distributions. Do and Vu [13] generalized these results to other distributions (such as the Bernoulli case). The spectral norm of the kernel matrix was studied by Fan and Montanari [18], who also made the interesting observation that the limiting distribution obtained in [9] is the free additive convolution of an MP law and a semicircle law.

Asymptotics with polynomial scaling: Studies of high-dimensional statistical problems often consider the linear asymptotic regime. In random matrix theory, the more general polynomial asymptotic regime was studied by Bloemendal et al. [7], who proved the local MP law for sample covariance matrices $X^\top X$, where

X is an $N \times n$ with $\log N \asymp \log n$. In that work, the matrix X is assumed to have independent entries. Although the kernel matrix in this work can also be written in the form of a (generalized) sample covariance matrix (see (2.15) and (2.20)), the matrix entries in our problem are the spherical harmonics, which are uncorrelated but dependent random variables. Thus, the technical approach of [7] cannot be directly applied here. In the context of kernel methods, exact asymptotics in the polynomial scaling regime were studied in Oppor and Urbanczik [30] and Bordelon et al. [8] via nonrigorous statistical physics methods. The heuristics used by these authors (the so-called Gaussian equivalence conjecture) were similar to the scheme sketched in Section 2.2, namely, treating the uncorrelated spherical harmonics as if they were independent standard normal random variables.

Ghorbani et al. [19] studied the high-dimensional limit of kernel regression (as well as several other related models) under the scaling where $d^{\ell+\delta} \leq n \leq d^{\ell+1-\delta}$ for some (small) $\delta > 0$. In other words, the number of samples is assumed to fall between two consecutive integer powers of d . This is different from the setting of this work, where we assume $n/d^\ell \rightarrow \kappa \in (0, \infty)$. Finally, upon the completion of this paper, we became aware of a very recent preprint Misiakiewicz [28] that studies the limit of kernel ridge regression in the above-mentioned exact polynomial asymptotic regime. As a main result of that study, it shows that the limiting spectrum of the kernel matrix is the MP law when the kernel function f_d is the ℓ -th Gegenbauer polynomial. This corresponds to a special case in our equivalence principle, with matrix A in (2.7) consisting only of the component $\mu_\ell A_\ell$. The presence of the high-order components $\{A_k\}_{k>\ell}$ in our model leads to more general limiting distributions (the free convolution between the MP law and the semicircle law) as well as additional technical challenges in the proof.

4 Proof of the Main Result

This section is devoted to the proof of Theorem 1. We start by presenting a high-level outline of the proof in Section 4.1. The technical details are given in Sections 4.2 and 4.3, and in the appendix.

4.1 Outline of the Proof

Recall that $n = \kappa d^\ell + o(d^\ell)$ for some fixed $\kappa > 0$ and $\ell \in \mathbb{N}$. Accordingly, we split the matrix A in (2.7) into two parts: the low-order term $A_{\text{LO}} := \sum_{0 \leq k < \ell} \mu_k A_k$, and the high-order term $A_{\text{HI}} := \sum_{\ell \leq k \leq L} \mu_k A_k$. We start by observing that the contribution of the low-order term to the Stieltjes transform is negligible.

Lemma 2. *Let $s(z)$ and $\hat{s}(z)$ denote the Stieltjes transforms of A and A_{HI} , respectively. Under the same settings of Theorem 1, we have*

$$|s(z) - \hat{s}(z)| = \mathcal{O}\left(\frac{1}{\eta^2 \sqrt{d}}\right).$$

Proof. See Appendix F. □

In light of Lemma 2, we will assume without loss of generality that $A_{\text{LO}} = 0$ and $A = A_{\text{HI}}$ in our following discussions. For each $i \in [n]$, we use $A^{[i]} \in \mathbb{R}^{(n-1) \times (n-1)}$ to denote the minor matrix obtained by removing the i th column and row of A , *i.e.*,

$$A_{ab}^{[i]} = A_{ab}, \quad a, b \neq i.$$

Note that, rather than mapping the indices of $A^{[i]}$ to $[n-1]$, we use $a, b \in [n] \setminus \{i\}$ to index the entries of $A^{[i]}$. Let $G(z) = (A - zI)^{-1}$ and $G^{[i]}(z) = (A^{[i]} - zI)^{-1}$ be the resolvents of A and $A^{[i]}$, respectively.

We start our proof by applying Schur's complement formula, which gives us

$$\frac{1}{G_{ii}(z)} = -z - \sum_{a, b \neq i} A_{ia} G_{ab}^{[i]}(z) A_{ib}. \quad (4.1)$$

Recall that the Stieltjes transform of A can be obtained as $s(z) = \frac{1}{n} \sum_i G_{ii}(z)$. The main technical step in our proof is to show that

$$Q_i(z) := \sum_{a, b \neq i} A_{ia} G_{ab}^{[i]}(z) A_{ib} - \left[\frac{\gamma_a s(z)}{1 + \gamma_b s(z)} + \gamma_c s(z) \right] = \mathcal{O}_{\prec}(d^{-1/2}), \quad (4.2)$$

where $\gamma_a, \gamma_b, \gamma_c$ are the three constants defined in Theorem 1. We will prove this estimate in Section 4.3 (see Proposition 2).

Using (4.2), we can now split the right-hand side of (4.1) into a leading term and an error term. Multiplying both sides of (4.1) by $G_{ii}(z)$ and averaging over i , we get

$$s(z) \left(z + \frac{\gamma_a s(z)}{1 + \gamma_b s(z)} + \gamma_c s(z) \right) + 1 = \underbrace{-\frac{1}{n} \sum_i Q_i(z) G_{ii}(z)}_{\text{(error term)}}. \quad (4.3)$$

By (F.2) in the first step, and by applying (4.2) and the union bound in the second step,

$$|(\text{error term})| \leq \frac{1}{\eta} \max_i |Q_i(z)| = \frac{1}{\eta} \mathcal{O}_{\prec}(d^{-1/2}). \quad (4.4)$$

Observe that the left-hand side of (4.3) has exactly the same functional form as the left-hand side of (2.10). The error estimate in (4.4) then implies that the Stieltjes transform $s(z)$ approximately satisfies the nonlinear equation in (2.10). By analyzing the uniqueness and stability of the solution $m(z)$ to this nonlinear equation (see Proposition 8 in Appendix G), we can then conclude that $s(z) \approx m(z)$.

4.2 Schur Complement: Reparameterization

We devote this and the next subsections to establishing the estimate in (4.2). Observe that, by construction, different coordinates of A are statistically exchangeable. Thus, we only need to show (4.2) for a single index $i \in [n]$. We choose to do so for $i = n$.

Recall from (2.7) and (2.6) that

$$A_{na} = \frac{1}{\sqrt{n}} \sum_{\ell \leq k \leq L} \mu_k q_k(\sqrt{d} \mathbf{x}_n^\top \mathbf{x}_a), \quad (4.5)$$

and the entries of the minor $A^{[n]}$ are of the form

$$A_{ab}^{[n]} = \frac{1}{\sqrt{n}} \sum_{\ell \leq k \leq L} \mu_k q_k(\sqrt{d} \mathbf{x}_a^\top \mathbf{x}_b) \mathbb{1}(a \neq b), \quad \text{for } a, b \in [n-1]. \quad (4.6)$$

A reminder to the reader: due to Lemma 2, we assume that A contains no lower order component, *i.e.*, $\mu_k = 0$ for $0 \leq k < \ell$.

The random variables $\{A_{na}\}$ and $\{A_{ab}^{[n]}\}$ are weakly correlated. To make this correlation explicit, we use the following reparameterization. For every $a \in [n-1]$, let

$$\xi_a := \sqrt{d} \mathbf{x}_n^\top \mathbf{x}_a \quad \text{and} \quad \tilde{\mathbf{x}}_a := (1 - \xi_a^2/d)^{-1/2} R_n^\top \mathbf{x}_a,$$

where $R_n \in \mathbb{R}^{d \times (d-1)}$ is a matrix whose columns are unit-norm vectors orthogonal to \mathbf{x}_n . We can now rewrite (4.5) as

$$A_{na} = \frac{1}{\sqrt{n}} \sum_{\ell \leq k \leq L} \mu_k q_k(\xi_a). \quad (4.7)$$

By definition, $(R_n R_n^\top + \mathbf{x}_n \mathbf{x}_n^\top) = I$, and hence $\mathbf{x}_a^\top \mathbf{x}_b = \mathbf{x}_a^\top (R_n R_n^\top + \mathbf{x}_n \mathbf{x}_n^\top) \mathbf{x}_b$ for $a, b \in [n-1]$. We can then verify that $\sqrt{d} \mathbf{x}_a^\top \mathbf{x}_b = r(\xi_a) r(\xi_b) \sqrt{d-1} \tilde{\mathbf{x}}_a^\top \tilde{\mathbf{x}}_b + \xi_a \xi_b / \sqrt{d}$, where

$$r(\xi) := (1 - 1/d)^{-1/4} (1 - \xi^2/d)^{1/2}. \quad (4.8)$$

This allows us to rewrite (4.6) as

$$A_{ab}^{[n]} = \frac{1}{\sqrt{n}} \sum_{\ell \leq k \leq L} \mu_k q_k \left(r(\xi_a) r(\xi_b) \sqrt{d-1} \tilde{\mathbf{x}}_a^\top \tilde{\mathbf{x}}_b + \xi_a \xi_b / \sqrt{d} \right) \mathbb{1}(a \neq b). \quad (4.9)$$

The advantage of the new parameterizations in (4.7) and (4.9) is that the families $\{\xi_a\}$ and $\{\tilde{\mathbf{x}}_a\}$ are independent, as we show below.

Lemma 3. $\{\xi_a\}_{a < n}$ is an i.i.d. family with $\xi_a \sim \mathfrak{S}_d^1$, as defined in (1.3). $\{\tilde{\mathbf{x}}_a\}_{a < n}$ is an i.i.d. family of random vectors with $\tilde{\mathbf{x}}_a \sim \text{Unif}(\mathcal{S}^{d-2})$. Moreover, $\{\xi_a\}$ and $\{\tilde{\mathbf{x}}_a\}$ are mutually independent.

Proof. Fix \mathbf{x}_n , and let $E = [\mathbf{x}_n R_n]$. By construction, E is an orthogonal matrix. Define

$$\mathbf{v}_a = E^\top \mathbf{x}_a, \quad 1 \leq a \leq n-1.$$

As $\mathbf{x}_a \sim_{\text{i.i.d.}} \text{Unif}(\mathcal{S}^{d-1})$ and E is independent of $\{\mathbf{x}_a\}_{a < n}$, we can conclude from the orthogonal invariant property of the spherical distribution that $\mathbf{v}_a \sim_{\text{i.i.d.}} \text{Unif}(\mathcal{S}^{d-1})$. The statement of the lemma then follows from the observation that

$$\xi_a = \sqrt{d} \mathbf{e}_1^\top \mathbf{v}_a \quad \text{and} \quad \tilde{\mathbf{x}}_a = \frac{\mathbf{v}_{a, \setminus 1}}{\|\mathbf{v}_{a, \setminus 1}\|},$$

where $\mathbf{v}_{a, \setminus 1}$ is the $(d-1)$ -dimensional vector obtained from \mathbf{v}_a after removing its first element. Then $\xi_a \sim \mathfrak{S}_d^1$. Meanwhile, $\tilde{\mathbf{x}}_a$ is uniformly distributed over \mathcal{S}^{d-2} , and is independent of ξ_a . Note that the above derivations are done for fixed \mathbf{x}_n (i.e. when conditioned on \mathbf{x}_n). However, since the distributions of $\{\xi_a\}$ and $\{\tilde{\mathbf{x}}_a\}$ are invariant to the choice of \mathbf{x}_n , we can conclude that $\{\xi_a\}$ and $\{\tilde{\mathbf{x}}_a\}$ are actually independent of \mathbf{x}_n . \square

It is easy to check that $\max_a |\xi_a| = \mathcal{O}_{\prec}(1)$ [see (D.12)], and thus $\xi_a \xi_b / \sqrt{d} = \mathcal{O}_{\prec}(\frac{1}{\sqrt{d}})$. It is then natural to consider a Taylor expansion of the right-hand side of (4.9) to try to bring the term $\xi_a \xi_b / \sqrt{d}$ out of the polynomial $q_k(\cdot)$. In fact, a more careful analysis will give us the following expansion formula, which plays an important role in our proof. First, we need to introduce a set of functions: for each $t \in \mathbb{N}_0$, let

$$\mathcal{E}_t = \left\{ \sum_{0 \leq \alpha, \beta \leq t} c_{\alpha\beta}(d) \frac{q_\alpha(\xi_1) q_\beta(\xi_2)}{d^{\max\{\alpha, \beta\}/2 + 1}} : c_{\alpha\beta}(d) = \mathcal{O}(1) \text{ for } 0 \leq \alpha, \beta \leq t \right\}. \quad (4.10)$$

In what follows, we will use $\mathcal{E}_t(\xi_1, \xi_2)$ to denote *any* function in \mathcal{E}_t . The exact form of $\mathcal{E}_t(\xi_1, \xi_2)$ can change from one expression to another.

Proposition 1. For each $k = 0, 1, 2, \dots, L$, we have

$$q_k\left(r(\xi_1)r(\xi_2)x + \xi_1\xi_2/\sqrt{d}\right) = \sum_{t=0}^k \tilde{q}_{k-t}(x) r^{k-t}(\xi_1) r^{k-t}(\xi_2) \left[\sqrt{(k)_t} \frac{q_t(\xi_1) q_t(\xi_2)}{d^{t/2}} + \mathcal{E}_t(\xi_1, \xi_2) \right], \quad (4.11)$$

where $(k)_t = k!/(k-t)!$ denotes the falling factorial, and $\tilde{q}_k(x)$ is the k th Gegenbauer polynomial in dimension $d-1$ as defined in (2.5).

Proof. See Appendix B. \square

For each $k \in \mathbb{N}_0$, define an $(n-1) \times (n-1)$ matrix \tilde{A}_k [corresponding to $t=0$ in (4.11)], whose entries are given by

$$(\tilde{A}_k)_{ab} = \frac{1}{\sqrt{n}} \tilde{q}_k(\sqrt{d-1} \tilde{\mathbf{x}}_a^\top \tilde{\mathbf{x}}_b) \mathbf{1}(a \neq b), \quad \text{for } a, b \in [n-1]. \quad (4.12)$$

We write their linear combination as

$$\tilde{A} := \sum_{\ell \leq k \leq L} \mu_k \tilde{A}_k.$$

For the first term on the right side of (4.11) with $t=k$, we define the vector

$$\mathbf{v}_k(\xi) := (q_k(\xi_1), q_k(\xi_2), \dots, q_k(\xi_{n-1}))^\top. \quad (4.13)$$

Using the expansion in (4.11), we can now write the matrix in (4.9) as

$$A^{[n]} = \tilde{A} + \frac{\mu_\ell}{\sqrt{nd^\ell/\ell!}} \mathbf{v}_\ell(\xi) \mathbf{v}_\ell(\xi)^\top + \sum_{1 \leq i \leq 5} \Delta_i, \quad (4.14)$$

where $\Delta_1, \dots, \Delta_5$ are matrices defined as follows: for $a, b \in [n-1]$,

$$(\Delta_1)_{ab} = \frac{-\mu_\ell q_\ell^2(\xi_a)}{\sqrt{nd^\ell/\ell!}} \delta_{ab}, \quad (4.15a)$$

$$(\Delta_2)_{ab} = \sum_{\ell \leq k \leq L} \mu_k \sum_{0 \leq t \leq k} \sqrt{\binom{k}{t}} \cdot (\tilde{A}_{k-t})_{ab} (r^{k-t}(\xi_a) r^{k-t}(\xi_b) - 1) \frac{q_t(\xi_a) q_t(\xi_b)}{d^{t/2}}, \quad (4.15b)$$

$$(\Delta_3)_{ab} = \mu_\ell \sum_{1 \leq t \leq \ell-1} \sqrt{\binom{\ell}{t}} \cdot (\tilde{A}_{\ell-t})_{ab} \frac{q_t(\xi_a) q_t(\xi_b)}{d^{t/2}}, \quad (4.15c)$$

$$(\Delta_4)_{ab} = \sum_{\ell \leq k \leq L} \mu_k \sum_{1 \leq t \leq k} \sqrt{\binom{k}{t}} \cdot (\tilde{A}_{k-t})_{ab} \frac{q_t(\xi_a) q_t(\xi_b)}{d^{t/2}}, \quad (4.15d)$$

$$(\Delta_5)_{ab} = \sum_{\ell \leq k \leq L} \mu_k \sum_{0 \leq t \leq k} (\tilde{A}_{k-t})_{ab} \cdot r^{k-t}(\xi_a) r^{k-t}(\xi_b) \mathcal{E}_t(\xi_a, \xi_b). \quad (4.15e)$$

Here Δ_2 is the error for replacing $r^{k-t}(\xi_a) r^{k-t}(\xi_b)$ by 1; the error terms for $k = \ell$ with $t \geq 1$ are collected in Δ_3 ; the error terms for $\ell < k \leq L$ are collected in Δ_4 ; and all the terms involving \mathcal{E} are collected in Δ_5 .

Later, we will show that the matrices $\{\Delta_i\}$ can all be treated as small noise terms. Moreover, by the construction of \tilde{A}_k in (4.12) and by Lemma 3, both \tilde{A} and its resolvent $\tilde{G}(z) := (\tilde{A} - zI)^{-1}$ are independent of $\{\xi_a\}$. These observations lead us to consider the following expansion.

Lemma 4. *Let $G^{[n]}(z)$ and $\tilde{G}(z)$ be the resolvents of $A^{[n]}$ and \tilde{A} , respectively. We have*

$$\begin{aligned} \sum_{a, b \neq n} A_{na} G_{ab}^{[n]}(z) A_{nb} &= \sum_{\ell \leq \alpha, \beta \leq L} \mu_\alpha \mu_\beta \chi_{\alpha\beta}(z) - \theta(z) \sum_{\ell \leq \beta \leq L} \mu_\beta \chi_{\ell\beta}(z) \\ &\quad - \frac{1}{\sqrt{n}} \left(\sum_{\ell \leq \alpha \leq L} \mu_\alpha \mathbf{v}_\alpha^\top(\xi) - \theta(z) \mathbf{v}_\ell^\top(\xi) \right) \tilde{G}(z) \left(\sum_{1 \leq c \leq 5} \Delta_c \right) G^{[n]}(z) \left(\frac{1}{\sqrt{n}} \sum_{\ell \leq \beta \leq L} \mu_\beta \mathbf{v}_\beta(\xi) \right), \end{aligned} \quad (4.16)$$

where, for $\alpha, \beta \in \mathbb{N}$, $\mathbf{v}_\alpha(\xi)$ is the vector defined in (4.13),

$$\chi_{\alpha\beta}(z) := \frac{1}{n} \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \mathbf{v}_\beta(\xi), \quad (4.17)$$

and

$$\theta(z) := \frac{\mu_\ell \sum_{\ell \leq \alpha \leq L} \mu_\alpha \chi_{\alpha\ell}}{\sqrt{d^\ell/(\ell!n)} + \mu_\ell \chi_{\ell\ell}}. \quad (4.18)$$

Proof. We start by recalling the simple resolvent identity in (F.6). Applying this identity on the two matrices $A^{[n]}$ and $\tilde{A} + \frac{\mu_\ell \mathbf{v}_\ell(\xi) \mathbf{v}_\ell(\xi)^\top}{\sqrt{nd^\ell/\ell!}}$, and by using (4.14), we get

$$\begin{aligned} G^{[n]}(z) &= \left(\tilde{A} + \frac{\mu_\ell \mathbf{v}_\ell(\xi) \mathbf{v}_\ell(\xi)^\top}{\sqrt{nd^\ell/\ell!}} - zI \right)^{-1} \left(I - \sum_{1 \leq i \leq 5} \Delta_i G^{[n]}(z) \right) \\ &= \left(\tilde{G}(z) - \frac{\mu_\ell \tilde{G}(z) \mathbf{v}_\ell(\xi) \mathbf{v}_\ell^\top(\xi) \tilde{G}(z)}{\sqrt{nd^\ell/\ell!} + \mu_\ell \mathbf{v}_\ell^\top(\xi) \tilde{G}(z) \mathbf{v}_\ell(\xi)} \right) \left(I - \sum_{1 \leq i \leq 5} \Delta_i G^{[n]}(z) \right), \end{aligned} \quad (4.19)$$

where the second equality follows from a special case of the Woodbury matrix identity, i.e.,

$$(A + \tau q q^*)^{-1} = A^{-1} - \frac{\tau A^{-1} q \cdot q^* A^{-1}}{1 + \tau q^* A^{-1} q}.$$

By (4.7) and (4.13),

$$A_{na} = \frac{1}{\sqrt{n}} \sum_{\ell \leq k \leq L} \mu_k [\mathbf{v}_k(\xi)]_a.$$

Substituting this into (4.19) then leads to the desired result. \square

4.3 Schur Complement: the Leading Term

Next, we show that the right-hand side of (4.16) can indeed be split into a leading term, the form of which is given in (4.2), and a small (random) error term.

Proposition 2. *Let $A = \sum_{\ell \leq k \leq L} \mu_k A_k$. Under the same conditions of Theorem 1, we have*

$$\sum_{a,b \neq i} A_{ia} G_{ab}^{[i]}(z) A_{ib} = \left[\frac{\gamma_a s(z)}{1 + \gamma_b s(z)} + \gamma_c s(z) \right] + \mathcal{O}_{\prec}(d^{-1/2}), \quad (4.20)$$

where $s(z)$ is the Stieltjes transform of the spectrum of A .

To prepare for the proof of Proposition 2, we first establish several intermediate results.

Lemma 5. *For $1 \leq \alpha, \beta \leq L$, we have*

$$\chi_{\alpha\beta}(z) = s(z) \delta_{\alpha\beta} + \mathcal{O}_{\prec} \left(\frac{1}{\eta^2 \sqrt{d}} \right). \quad (4.21)$$

Proof. Let $\tilde{s}(z) := \frac{1}{n} \text{tr} \tilde{G}$ and $s^{[n]}(z) := \frac{1}{n} \text{tr} G^{[n]}$ be the Stieltjes transforms of \tilde{G} and $G^{[n]}$, respectively. We will establish (4.21) in three steps, by showing (a) $\chi_{\alpha\beta}(z) \approx \tilde{s}(z) \delta_{\alpha\beta}$; (b) $\tilde{s}(z) \approx s^{[n]}(z)$; and (c) $s^{[n]}(z) \approx s(z)$.

For step (a), we first recall the Ward identity (F.3), which gives us

$$\frac{1}{n} \sum_{1 \leq a, b < n} |\tilde{G}_{ab}|^2 = \frac{\Im(\tilde{s}(z))}{\eta} \leq \frac{1}{\eta^2}.$$

The second step uses (F.4). By the definitions in (4.13) and (4.17), $\chi_{\alpha\beta}(z) = \frac{1}{\sqrt{n}} \sum_{a,b} [\frac{1}{\sqrt{n}} \tilde{G}_{ab}(z)] q_{\alpha}(\xi_a) q_{\beta}(\xi_b)$. By applying (D.24b) in Proposition 6 with $\Psi_c = 1/\eta$, we have

$$|\chi_{\alpha\beta}(z) - \tilde{s}(z) \delta_{\alpha\beta}| \prec \frac{1}{\eta \sqrt{n}}. \quad (4.22)$$

For step (b), we use the expansion formula in (4.14). By (F.8), (F.7), and the triangular inequality,

$$|s^{[n]}(z) - \tilde{s}(z)| \leq \frac{C}{n\eta} + \frac{\sum_{i=1}^5 \|\Delta_i\|_{\text{F}}}{\sqrt{n}\eta^2} \leq \frac{C}{n\eta} + \frac{\sqrt{n}}{\eta^2} \sum_{i=1}^5 \|\Delta_i\|_{\infty}, \quad (4.23)$$

where C is an absolute constant. Next, we bound the entry-wise ℓ_{∞} norm of $\{\Delta_i\}$, whose constructions are given in (4.15a) – (4.15e). Our derivations will frequently use the following property of stochastic domination, which can be verified by a simple union bound: If $X_1 \prec Y_1$ and $X_2 \prec Y_2$, then

$$X_1 X_2 \prec Y_1 Y_2 \quad \text{and} \quad X_1 + X_2 \prec Y_1 + Y_2. \quad (4.24)$$

By (D.12), and the property stated above, we have $\max_a |q_k(\xi_a)|^2 \prec 1$. It follows that $\|\Delta_1\|_{\infty} \prec \frac{1}{\sqrt{nd}^t}$. Similarly, by using (D.13), we can easily verify that

$$\|\Delta_3\|_{\infty} \prec \frac{1}{\sqrt{nd}} \quad \text{and} \quad \|\Delta_4\|_{\infty} \prec \frac{1}{\sqrt{nd}}.$$

To bound Δ_5 , we first check from the definition in (4.10) that $\max_{a,b \in [n-1]} |\mathcal{E}_t(\xi_a, \xi_b)| \prec 1/d$. Moreover, by its construction in (4.8), the function $r(\xi) \leq 1 + \mathcal{O}(1/d)$ for all ξ . Combining these estimates then gives us $\|\Delta_5\|_{\infty} \prec \frac{1}{d\sqrt{n}}$.

The matrix Δ_2 in (4.15b) requires some additional care, as it contains terms with index $t = 0$. First write

$$r^{k-t}(\xi_a) r^{k-t}(\xi_b) - 1 = (r^{k-t}(\xi_a) - 1)(r^{k-t}(\xi_b) - 1) + (r^{k-t}(\xi_a) - 1) + (r^{k-t}(\xi_b) - 1). \quad (4.25)$$

Applying the high-probability bound (D.14) and the property in (4.24), we get

$$\max_{a,b} \left| r^{k-t}(\xi_a) r^{k-t}(\xi_b) - 1 \right| \prec 1/d,$$

which then implies that $\|\Delta_2\|_\infty \prec \frac{1}{d\sqrt{n}}$. Substituting these bounds on $\|\Delta_c\|_\infty$ for $1 \leq c \leq 5$ into (4.23) then leads to

$$\left| s^{[n]}(z) - \tilde{s}(z) \right| \prec \frac{1}{\eta^2 \sqrt{d}}. \quad (4.26)$$

Now we move to step (c), where we compare $s^{[n]}(z)$ with $s(z)$. By the interlacing properties of the eigenvalues of $A^{[n]}$ and A , we have the following standard result (see [16, Lemma 7.5] for a proof):

$$\left| s^{[n]}(z) - s(z) \right| \leq \frac{C}{n\eta}, \quad (4.27)$$

where C is an absolute constant. Finally, the statement of the lemma can be obtained by applying the triangular inequality and by using the estimates in (4.22), (4.26), and (4.27). \square

Next, we show that the terms involving $\{\Delta_c\}_{1 \leq c \leq 5}$ on the right-hand side of (4.16) are small. This is done in the following two lemmas.

Lemma 6. *Let α, β, k be integers such that $0 \leq \alpha, \beta, k \leq L$ and $\alpha \neq \beta$. We have*

$$\left\| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_\beta(\xi) \tilde{A}_k \right\| \prec \frac{\sqrt{n}}{\eta}, \quad (4.28)$$

where $\Lambda_\beta(\xi)$ denotes a diagonal matrix defined as

$$\Lambda_\beta(\xi) := \text{diag} \{ q_\beta(\xi_1), q_\beta(\xi_2), \dots, q_\beta(\xi_{n-1}) \}. \quad (4.29)$$

Proof. Write $u := \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_\beta(\xi) \tilde{A}_k$. This is a vector with $n-1$ entries. In our proof, we will show that

$$\max_{a \in [n-1]} |u_a| \prec 1/\eta, \quad (4.30)$$

of which (4.28) is then an immediate consequence. For each $a \in [n-1]$, define a matrix

$$C_a := \tilde{G}(z) \text{diag} \{ (\tilde{A}_k)_{1a}, (\tilde{A}_k)_{2a}, \dots, (\tilde{A}_k)_{n-1,a} \}.$$

One can verify that

$$u_a = \mathbf{v}_\alpha^\top(\xi) C_a \mathbf{v}_\beta(\xi).$$

We start by proving (4.30) for the special case when $\alpha > \beta = 0$. Recall that $\mathbf{v}_0(\xi) = \mathbf{1}$. We then have $\|C_a \mathbf{v}_0(\xi)\| \leq \|\tilde{G}\|_{\text{op}} \|\tilde{A}_k\|_\infty \sqrt{n} \prec 1/\eta$, where the second step uses (F.1) and (D.13). Since C_a is independent of $\mathbf{v}_\alpha(\xi)$, we can apply (D.24a) in Proposition 6 with $\Psi_b = 1/\eta$ to get

$$u_a = \mathbf{v}_\alpha^\top(\xi) C_a \mathbf{v}_0(\xi) \prec 1/\eta. \quad (4.31)$$

By the same arguments, $\mathbf{v}_0^\top(\xi) C_a \mathbf{v}_\beta(\xi) \prec 1/\eta$ for $\beta > 0$. In what follows, we assume $1 \leq \alpha, \beta \leq L$ with $\alpha \neq \beta$. Using the Ward identity (F.3) in the second step, and applying (F.4), (D.13) in the third step, we get

$$\|C_a\|_{\text{F}} \leq \|\tilde{G}\|_{\text{F}} \|\tilde{A}_k\|_\infty \leq \frac{\sqrt{n \cdot \Im \tilde{s}(z)}}{\sqrt{\eta}} \|\tilde{A}_k\|_\infty \prec \frac{1}{\eta}.$$

By appealing to (D.24b) in Proposition 6 with $\Psi_c = 1/\eta$, we have

$$u_a = \mathbf{v}_\alpha^\top(\xi) C_a \mathbf{v}_j(\xi) \prec 1/\eta. \quad (4.32)$$

One can check that the high-probability bounds in (4.31) and (4.32) are both uniform in $a \in [n-1]$. (To see this, note that we always bound $\max_i |(\tilde{A}_k)_{ia}|$ via $\|\tilde{A}_k\|_\infty$ in the above derivations.) Applying the union bound over a then gives us (4.30). \square

Lemma 7. For $\alpha, \beta \in \mathbb{N}$ with $\ell \leq \alpha, \beta \leq L$, and for $1 \leq c \leq 5$, we have

$$\frac{1}{n} \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Delta_c G^{[n]}(z) \mathbf{v}_\beta(\xi) = \mathcal{O}_{\prec} \left(\frac{1}{\eta^2 \sqrt{d}} \right). \quad (4.33)$$

Proof. By using (F.1) in the first step and (D.12) in the second step, we have $\|G^{[n]} \mathbf{v}_\beta(\xi)\| \leq \frac{1}{\eta} \|\mathbf{v}_\beta(\xi)\| \prec \frac{\sqrt{n}}{\eta}$. The same arguments will also give us $\|\mathbf{v}_\alpha^\top(\xi) \tilde{G}(z)\| \prec \sqrt{n}/\eta$. It follows that

$$\left| \frac{1}{n} \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Delta_c G^{[n]}(z) \mathbf{v}_\beta(\xi) \right| \prec \frac{1}{\eta^2} \|\Delta_c\|_{\text{op}}. \quad (4.34)$$

Thus, one way to bound the left-hand side of (4.33) is to obtain estimates on the operator norm of Δ_c . We start from Δ_1 in (4.15a). Since it is a diagonal matrix,

$$\|\Delta_1\|_{\text{op}} \leq \frac{1}{\sqrt{nd}^\ell} \max_a q_\ell^2(\xi_a) \prec \frac{1}{\sqrt{nd}^\ell}, \quad (4.35)$$

where the second step is due to (D.12). For Δ_2 in (4.15b), write $u_k := \max_a |r^k(\xi_a) - 1|$. Using the expansion in (4.25) and the triangular inequality, we have

$$\|\Delta_2\|_{\text{op}} \leq C \sum_{\ell \leq k \leq L} \sum_{0 \leq t \leq k} \frac{1}{d^{t/2}} \|\tilde{A}_{k-t}\|_{\text{op}} (u_{k-t}^2 + 2u_{k-t}) \max_a |q_t(\xi_a)|^2$$

for some constant C that depends on $\{\mu_k\}$. Note that \tilde{A}_k has exactly the same construction as A_k . The only difference is to change the dimension parameter from d to $d-1$. Thus, we can apply Proposition 7 to get $\|\tilde{A}_{k-t}\|_{\text{op}} \leq d^{\max\{(\ell-k+t)/2, 0\}}$. Combining this estimate with those in (D.14) and (D.12) leads to

$$\|\Delta_2\|_{\text{op}} \prec C \sum_{\ell \leq k \leq L} \sum_{0 \leq t \leq k} d^{\max\{(\ell-k+t)/2, 0\}} d^{-t/2} d^{-1} \prec d^{-1}, \quad (4.36)$$

where the second step is due to $\max\{(\ell-k+t)/2, 0\} \leq t/2$ for $k \geq \ell$. Similarly, for Δ_4 in (4.15d), we have

$$\|\Delta_4\|_{\text{op}} \prec C \sum_{\ell < k \leq L} \sum_{1 \leq t \leq k} d^{\max\{(\ell-k+t)/2, 0\}} d^{-t/2} \prec d^{-1/2}. \quad (4.37)$$

The second step follows from the fact that, for $k > \ell$ and $t \geq 1$, $\max\{(\ell-k+t)/2, 0\} \leq \max\{\frac{t-1}{2}, 0\} = \frac{t-1}{2}$. By substituting (4.35), (4.36), and (4.37) into (4.34), we verify the statement in (4.33) for the cases of $c = 1, 2$ and 4.

The cases of Δ_3 and Δ_5 require a different approach. We start with Δ_3 . Recall the diagonal matrix notation introduced in (4.29). Write

$$\mathbf{u}_\beta := G^{[n]}(z) \mathbf{v}_\beta(\xi). \quad (4.38)$$

By (D.12) and (F.1), we have

$$\|\mathbf{u}_\beta\| \leq \|G^{[n]}(z)\|_{\text{op}} \cdot \|\mathbf{v}_\beta(\xi)\| \prec \frac{\sqrt{n}}{\eta}. \quad (4.39)$$

and

$$\|\Lambda_t(\xi) \mathbf{u}_\beta\| \leq \left(\max_a |q_t(\xi_a)| \right) \cdot \|\mathbf{u}_\beta\| \prec \frac{\sqrt{n}}{\eta}. \quad (4.40)$$

From the construction of Δ_3 in (4.15c), one can check that

$$\begin{aligned} \left| \frac{1}{n} \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Delta_3 G^{[n]}(z) \mathbf{v}_\beta(\xi) \right| &\leq C \sum_{1 \leq t \leq \ell-1} \frac{1}{d^{t/2} n} \left| \left(\mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_t(\xi) \tilde{A}_{\ell-t} \right) \Lambda_t(\xi) \mathbf{u}_\beta \right| \\ &\leq C \sum_{1 \leq t \leq \ell-1} \frac{1}{d^{t/2} n} \left\| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_t(\xi) \tilde{A}_{\ell-t} \right\| \cdot \|\Lambda_t(\xi) \mathbf{u}_\beta\| \\ &\prec \frac{1}{\eta^2 \sqrt{d}}. \end{aligned}$$

The second step uses the Cauchy-Schwarz inequality; in the last step, we have applied Lemma 6 (note that $\alpha \geq \ell > t$), and the estimate in (4.39).

We complete the proof by showing (4.33) for Δ_5 . By construction, Δ_5 is the linear combination of a finite number of matrices $\{\Delta_5^{(k,t)}\}$, indexed by k, t . The entries of $\Delta_5^{(k,t)}$ are

$$\begin{aligned}
(\Delta_5^{(k,t)})_{ab} &= (\tilde{A}_{k-t})_{ab} \cdot r^{k-t}(\xi_a)r^{k-t}(\xi_b)\mathcal{E}_t(\xi_a, \xi_b) \\
&\stackrel{(a)}{=} (\tilde{A}_{k-t})_{ab}(\mathcal{E}_{t+2\ell}(\xi_a, \xi_b) + f_2(\xi_a, \xi_b)) \\
&\stackrel{(b)}{=} (\tilde{A}_{k-t})_{ab} \left(\underbrace{\sum_{\substack{0 \leq i, j \leq t+2\ell}} c_{ij}(d) d^{-\max\{i,j\}/2-1} q_i(\xi_a) q_j(\xi_b)}_{(*)} + f_2(\xi_a, \xi_b) \right) \\
&\stackrel{(c)}{=} (\tilde{A}_{k-t})_{ab} \sum_{\substack{0 \leq i \leq \ell-1 \\ 0 \leq j \leq t+2\ell}} (*) + \underbrace{(\tilde{A}_{k-t})_{ab} \left(\sum_{\substack{\ell \leq i \leq t+2\ell \\ 0 \leq j \leq t+2\ell}} (*) \right)}_{=: R_{ab}} + f_2(\xi_a, \xi_b). \tag{4.41}
\end{aligned}$$

In step (a), we apply Proposition 5 to split $r^{k-t}(\xi_a)r^{k-t}(\xi_b)\mathcal{E}_t(\xi_a, \xi_b)$ into the sum of two functions, with $f_1(\xi_a, \xi_b) \in \mathcal{E}_{t+2\ell}$. Step (b) uses the definition in (4.10); In step (c), we split the double sum over $0 \leq i, j \leq t+2\ell$ into two parts, with the index i ranging from 0 to $\ell-1$ in the first partial sum. Observe that each term in the partial sum in R_{ab} includes a factor $d^{-\max\{i,j\}/2-1} \leq d^{-\ell/2-1} = \mathcal{O}(n^{-1/2}d^{-1})$, because $i \geq \ell$. By (D.13), (D.12), and (D.18), and by the union bound, we can verify that

$$\max_{a,b} |R_{ab}| \prec \frac{1}{dn} \quad \text{and thus} \quad \|R\|_{\text{op}} \leq \|R\|_{\text{F}} \prec \frac{1}{d}. \tag{4.42}$$

Now consider the left-hand side of (4.33), with Δ_c replaced by $\Delta_5^{(k,t)}$. By using (4.41) and the notation introduced in (4.29) and (4.38), we have

$$\begin{aligned}
\frac{1}{n} \left| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Delta_5^{(k,t)} G^{[n]}(z) \mathbf{v}_\beta(\xi) \right| &\leq \frac{1}{nd} \sum_{\substack{0 \leq i \leq \ell-1 \\ 0 \leq j \leq t+2\ell}} \left| c_{ij}(d) \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_i(\xi) \tilde{A}_{k-t} \Lambda_j(\xi) \mathbf{u}_\beta \right| + \frac{1}{n} \left| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) R \mathbf{u}_\beta \right| \\
&\leq \frac{1}{nd} \sum_{\substack{0 \leq i \leq \ell-1 \\ 0 \leq j \leq t+2\ell}} |c_{ij}(d)| \cdot \left\| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \Lambda_i(\xi) \tilde{A}_{k-t} \right\| \cdot \left\| \Lambda_j(\xi) \mathbf{u}_\beta \right\| + \frac{1}{n} \left\| \mathbf{v}_\alpha^\top(\xi) \tilde{G}(z) \right\| \|R\|_{\text{op}} \|\mathbf{u}_\beta\| \tag{4.43}
\end{aligned}$$

$$\prec \frac{1}{\eta^2 d}. \tag{4.44}$$

In the last step, we bound the right-hand side of (4.43) as follows: for the terms in the sum over i, j , we have used the fact that $c_{ij}(d) = \mathcal{O}(1)$ and have applied (4.40) and (4.28) in Lemma 6. (Note that $\beta \geq \ell > i$ and thus (4.28) is applicable.) For the last term on the right-hand side of (4.43), we use (4.42), (4.39), and the estimate that $\|\mathbf{v}_\alpha^\top(\xi) \tilde{G}(z)\| \prec \sqrt{n}/\eta$.

The large deviation bound in (4.44) holds for all k, t over the range $\ell \leq k \leq L$ and $0 \leq t \leq k$. Applying the triangular inequality then yields the estimate in (4.33) for Δ_5 . \square

We may now complete the proof of Proposition 2.

Proof of Proposition 2. Observe that the different coordinates of A are statistically exchangeable. Thus, it is sufficient to show (4.20) for any particular choice of the index i . We choose to do so for $i = n$. With the decomposition given in (4.16), our task boils down to verifying that the right-hand side of (4.16) concentrates around the leading term given in (4.20). To that end, we first show that the term $\theta(z)$ in (4.18) is close to

$$\theta^*(z) := \frac{\mu_\ell^2 s(z)}{\sqrt{d^\ell/(\ell n)} + \mu_\ell s(z)}.$$

Indeed, we have

$$\left| \theta(z) - \theta^*(z) \right| \leq \frac{\sum_{\ell < \alpha \leq L} |\mu_\ell \mu_\alpha \chi_{\alpha\ell}|}{\left| \sqrt{d^\ell/(\ell n)} + \mu_\ell \chi_{\ell\ell} \right|} + \frac{\mu_\ell^2 \sqrt{d^\ell/(\ell n)} |\chi_{\ell\ell} - s(z)|}{\left| \sqrt{d^\ell/(\ell n)} + \mu_\ell \chi_{\ell\ell}(z) \right| \cdot \left| \sqrt{d^\ell/(\ell n)} + \mu_\ell s(z) \right|}. \tag{4.45}$$

The numerators on the right-hand side are small due to the estimates provided in Lemma 5. The denominators can be bounded as follows. To lighten the notation, write $C_{d,\ell} := \sqrt{d^\ell/(\ell!n)}$. Recall the definition of $\chi_{\ell\ell}(z)$ in (4.17). By (F.1),

$$|\mu_\ell \chi_{\ell\ell}(z)| \leq \frac{|\mu_\ell| \cdot \|\mathbf{v}_\ell(\xi)\|^2}{n\eta}.$$

Consider two cases: (a) If $|\mu_\ell| \cdot \|\mathbf{v}_\ell(\xi)\|^2/(n\eta) \leq C_{d,\ell}/2$, we have $|\mu_\ell \chi_{\ell\ell}(z)| \leq C_{d,\ell}/2$ and thus

$$\frac{1}{|C_{d,\ell} + \mu_\ell \chi_{\ell\ell}|} \leq \frac{2}{C_{d,\ell}}. \quad (4.46)$$

(b) Suppose that $|\mu_\ell| \cdot \|\mathbf{v}_\ell(\xi)\|^2/(n\eta) > C_{d,\ell}/2$. Let $\{\lambda_j\}_{j \in [n-1]}$ denote the eigenvalues of \tilde{A} . One can verify from (4.17) that

$$|\mu_\ell| \Im \chi_{\ell\ell}(z) \geq \frac{|\mu_\ell| \cdot \|\mathbf{v}_\ell\|^2}{n\eta} \min_{j \in [n-1]} \frac{\eta^2}{(\lambda_j - E)^2 + \eta^2} \geq \left(\frac{C_{d,\ell}}{2}\right) \frac{\eta^2}{2\|\tilde{A}\|_{\text{op}}^2 + 2E^2 + \eta^2},$$

where $E = \Re \mathfrak{c}(z)$. This then gives us

$$\frac{1}{|C_{d,\ell} + \mu_\ell \chi_{\ell\ell}|} \leq \frac{2}{C_{d,\ell}} \left(1 + \frac{2(\|\tilde{A}\|_{\text{op}}^2 + E^2)}{\eta^2}\right). \quad (4.47)$$

Since the right-hand side of (4.47) dominates that of (4.46), the bound in (4.47) in fact is valid for both cases. Note that $|\mu_\ell s(z)| \leq |\mu_\ell|/\eta$. By considering two cases depending on whether $|\mu_\ell|/\eta \geq C_{d,\ell}/2$ and by following analogous arguments as above, we can also verify that

$$\frac{1}{|C_{d,\ell} + \mu_\ell s(z)|} \leq \frac{2}{C_{d,\ell}} \left(1 + \frac{2(\|A\|_{\text{op}}^2 + E^2)}{\eta^2}\right). \quad (4.48)$$

We already have controls on the operator norm of A . Indeed, the estimates in (E.27) give us

$$\|A\|_{\text{op}} \leq \sum_{k=\ell}^L |\mu_k| \cdot \|A_k\|_{\text{op}} \prec 1.$$

Similarly, $\|\tilde{A}\|_{\text{op}} \prec 1$, as \tilde{A} is just the $(d-1)$ -dimensional version of A . Substituting (4.47) and (4.48) into (4.45), and using the large deviation estimates in (4.21), we get

$$|\theta(z) - \theta^*(z)| \prec \frac{1}{\sqrt{d}}. \quad (4.49)$$

Finally, by applying (4.49), (4.21), and (4.33) to the right-hand side of (4.16), and by using the assumption that $n/d^\ell = \kappa + \mathcal{O}(d^{-1/2})$, we reach the statement (4.20) of the proposition. \square

4.4 Proof of Theorem 1

The asymptotic limit of the ESD of the B matrix can be obtained by standard methods in random matrix theory. For example, since B_ℓ is a shifted Wishart matrix, its limiting ESD follows the Marchenko-Pastur (MP) law with a ratio parameter equal to $(\ell!\kappa)^{-1}$. Let $s_\ell(z)$ denote the Stieltjes transform of the limiting spectral density of the linearly-scaled version $\mu_\ell B_\ell$. By using properties of the MP law (see Appendix C), and after taking into account the linear transformation (namely, the scaling by μ_ℓ and shifting by $(\ell!\kappa)^{-1/2}$) of the eigenvalues, we can characterize $s_\ell(z)$ as the unique solution to

$$\frac{1}{s_\ell(z)} = -z - \frac{\gamma_a s_\ell(z)}{1 + \gamma_b s_\ell(z)}, \quad (4.50)$$

where γ_a, γ_b are the constants defined in Theorem 1.

By construction, the law of B is equal to that of $\mu_\ell B_\ell + \sqrt{\gamma_c} H$, where $H \in \mathbb{R}^{n \times n}$ is a GOE matrix independent of B_ℓ . Thus, the Stieltjes transform $s_B(z)$ of its ESD converges almost surely to a limiting function $m(z)$. Moreover, $m(z)$ can be obtained via a free additive convolution between the MP law and the semicircle law, which gives us $m(z) = s_\ell(z + \gamma_c m(z))$. It then follows from (4.50) that

$$\begin{aligned} \frac{1}{m(z)} &= \frac{1}{s_\ell(z + \gamma_c m(z))} = -z - \gamma_c m(z) + \frac{\gamma_a s_\ell(z + \gamma_c m(z))}{1 + \gamma_b s_\ell(z + \gamma_c m(z))} \\ &= -z + \frac{\gamma_a m(z)}{1 + \gamma_b m(z)} - \gamma_c m(z), \end{aligned}$$

which is exactly (2.10).

The rest of the proof is devoted to establishing the estimate in (2.11) for the matrix A . To lighten the notation, we drop the subscript in $s_A(z)$ in the following discussions. We start by recalling Lemma 2, which shows that removing the low-order components from A will only incur an error of size $\mathcal{O}(1/\sqrt{d})$ in the Stieltjes transform. As this error is of the same size as the right-hand side of (2.11), it is sufficient for us to establish (2.11) for the special case when A contains only the high-order components, *i.e.*, $A = \sum_{k=\ell}^L \mu_k A_k$.

By (4.3), (4.4), (4.2), and the high-probability estimate given in Proposition 2, we have

$$s(z) \left(z + \frac{\gamma_a s(z)}{1 + \gamma_b s(z)} + \gamma_c s(z) \right) + 1 = \omega,$$

where the error term

$$|\omega| \prec \frac{1}{\eta \sqrt{d}}. \quad (4.51)$$

Thus, $s(z)$ is an approximate solution to (2.10). To show $s(z) \approx m(z)$, we will apply the stability bound given in Proposition 8. To that end, we first establish a high-probability upper bound for $1/\Im[s(z)]$. Let $\{\lambda_i\}_{i \in [n]}$ denotes the eigenvalues of A , and $z = E + i\eta \in C_+$. We have

$$\Im[s(z)] = \frac{1}{n} \sum_i \frac{\eta}{(\lambda_i - E)^2 + \eta^2} \geq \frac{\eta/2}{\|A\|_{\text{op}}^2 + E^2 + \eta^2}. \quad (4.52)$$

Applying the triangular inequality and Proposition 7, we have $\|A\|_{\text{op}} \leq \sum_{k=\ell}^L |\mu_k| \cdot \|A_k\|_{\text{op}} \prec 1$. Moreover, by the assumption made in Theorem 1, $|E| \leq \tau^{-1}$ and $\tau \leq \eta \leq \tau^{-1}$ for some fixed $\tau > 0$. Substituting these bounds into (4.52) then gives us

$$\frac{1}{\Im[s(z)]} \leq \frac{\|A\|_{\text{op}}^2 + E^2 + \eta^2}{\eta/2} \prec 1. \quad (4.53)$$

For any $\varepsilon > 0$ and $D > 0$, we have

$$\begin{aligned} \mathbb{P}(|s(z) - m(z)| \geq d^{\varepsilon-1/2}) &\leq \mathbb{P}(|\omega/s(z)| > \eta/2) + \mathbb{P}(|s(z) - m(z)| \geq d^{\varepsilon-1/2}, |\omega/s(z)| \leq \eta/2) \\ &\stackrel{(a)}{\leq} \mathbb{P}(|\omega| (\Im[s(z)])^{-1} \geq \eta/2) + \mathbb{P}(|\omega| \geq (\eta^2/4) d^{\varepsilon-1/2}) \\ &\stackrel{(b)}{\leq} d^{-(D+1)} + d^{-(D+1)}. \end{aligned}$$

In step (a), we have applied Proposition 8 to obtain the second term on the right-hand side of the inequality. Step (b) follows from the high-probability estimates given in (4.51) and (4.53). As ε and D can be chosen arbitrarily, we have verified (2.11). The almost sure convergence of $s(z)$ to $m(z)$ then follows from (2.11) and the Borel-Cantelli lemma.

5 Summary and Discussions

Our main results, Theorem 1 and Theorem 2, establish the weak-limit of the spectrum of random inner-product kernel matrices in the polynomial scaling regime, when $d, n \rightarrow \infty$ such that $n/d^\ell \rightarrow \kappa \in (0, \infty)$, for

some fixed $\kappa \in \mathbb{R}$ and $\ell \in \mathbb{N}$. The main insight of our work is an asymptotic equivalence principle, which states that the limiting spectral distribution of the kernel matrix is equivalent to that of a linear combination of a (shifted) Wishart matrix and an independent GOE matrix. Consequently, the limiting distribution is characterized by a free additive convolution between an MP law and a semicircle law.

Our results were stated only when the distributions of the data vectors are spherical. It is not difficult to generalize this result to isotropic Gaussian distributions. In fact, Theorem 1 is expected to hold (with certain modifications in error bounds) for essentially any distributions with reasonably fast decay at infinity. The error bounds on Theorem 1 are clearly not optimal. Furthermore, our current proof approach restricts the imaginary part of the spectral parameter $z = E + i\eta$ to $\eta \geq d^{-c}$ for some small constant c . This is far from the regime $\eta \sim n^{-1+\epsilon}$ needed to get to the locations of individual eigenvalues. However, extending Theorem 1 to the entire range $n^{-1+\epsilon} \leq \eta \leq 1$ may not be a simple matter since there seem to be multiscale structures in the eigenvalues due to that the matrix A in (2.7) is a sum of component matrices $\{A_k\}$ over different scales. Therefore, characterizing the precise behavior of individual eigenvalues, including the extremal eigenvalues, is an interesting open problem. Related questions such as eigenvector statistics and universality of eigenvalues will also depend on the resolution of this question. We remark that the possibility of multiscale structures is a feature of the nonlinear model, which does not appear in standard mean field models such as Wigner matrices or sample covariance matrices [16]. We will address some of these questions in the forthcoming papers.

A Gegenbauer Polynomials and Spherical Harmonics

In this appendix, we collect a few useful properties of the (normalized) Gegenbauer polynomials $\{q_k(x)\}_{k \geq 0}$ defined in (2.2). All of these results are standard, and their proofs can be found in *e.g.*, [11, 14].

Three-term recurrence relation: For each $k \in \mathbb{N}$, we have

$$xq_k(x) = a_k q_{k+1}(x) + a_{k-1} q_{k-1}(x), \quad (\text{A.1})$$

where

$$a_k = \sqrt{\frac{(k+1)(d+k-2)d}{(d+2k)(d+2k-2)}} = \sqrt{k+1} (1 + \mathcal{O}(d^{-1})). \quad (\text{A.2})$$

We will repeatedly apply this recurrence relation in our proof of Proposition 1.

Spherical harmonics and the addition theorem: We recall that a spherical harmonic of degree k is a harmonic and homogeneous polynomial of degree k in d variables that are restricted to \mathcal{S}^{d-1} . The set of all degree k harmonics forms a linear subspace, whose dimension is denoted by N_k . We have $N_0 = 1$, $N_1 = d$, and in general, for $k \in \mathbb{N}$,

$$N_k := \frac{2k+d-2}{k} \binom{k+d-3}{k-1} = \frac{d^k}{k!} (1 + \mathcal{O}(1/d)). \quad (\text{A.3})$$

Using the Gram-Schmidt procedure, we can construct an orthonormal set of spherical harmonics. Let $Y_{k,i}(\mathbf{x})$, for $i \in [N_k]$, denote the i th degree- k spherical harmonic in this orthonormal set. We then have

$$\mathbb{E}_{\mathbf{x} \sim \text{Unif}(\mathcal{S}^{d-1})} Y_{k,a}(\mathbf{x}) Y_{k',b}(\mathbf{x}) = \mathbf{1}_{kk'} \mathbf{1}_{ab}, \quad (\text{A.4})$$

for any $k, k' \in \mathbb{N}_0$ and any $a \in [N_k], b \in [N_{k'}]$.

The Gegenbauer polynomials and spherical harmonics are deeply connected. In particular, we have the following identity, which can be viewed as a high-dimensional generalization of the classical addition theorem [10]: for any $\mathbf{x}, \mathbf{y} \in \mathcal{S}^{d-1}$,

$$q_k(\sqrt{d} \mathbf{x}^\top \mathbf{y}) = \frac{1}{\sqrt{N_k}} \sum_{a \in [N_k]} Y_{k,a}(\mathbf{x}) Y_{k,a}(\mathbf{y}). \quad (\text{A.5})$$

In addition, for the special case of $\mathbf{x} = \mathbf{y}$, we have

$$q_k(\sqrt{d}) = \frac{1}{\sqrt{N_k}} \sum_{a \in [N_k]} Y_{k,a}^2(\mathbf{x}) = \sqrt{N_k}. \quad (\text{A.6})$$

The identity in (A.5) is useful because it allows us to “linearize” the term $q_k(\sqrt{d}\mathbf{x}^\top\mathbf{y})$ as an inner product of two vectors made of the spherical harmonics. Moreover, (A.4) implies that, if $\mathbf{x} \sim \text{Unif}(S^{d-1})$, the vector of spherical harmonics $(Y_{k,1}(\mathbf{x}), \dots, Y_{k,N_k}(\mathbf{x}))$ is an isotropic random vector.

Using (A.5) and (A.6), we can rewrite the matrix A_k defined in (2.6) as

$$A_k = \frac{1}{\sqrt{nN_k}} S_k^\top S_k - \sqrt{\frac{N_k}{n}} I, \quad (\text{A.7})$$

where S_k is a $N_k \times n$ matrix whose entries are the spherical harmonics, *i.e.*

$$(S_k)_{ai} = Y_{k,a}(\mathbf{x}_i), \quad (\text{A.8})$$

and $\{\mathbf{x}_i\}_{i \in [n]}$ are the data vectors in the definition in (2.6).

Relationship with the Hermite polynomials: The weight function of the Gegenbauer polynomials is the probability distribution \mathfrak{s}_d^1 given in (1.3). Since $\mathfrak{s}_d^1 \rightarrow \mathcal{N}(0,1)$ when $d \rightarrow \infty$, the Gegenbauer polynomials asymptotically become the Hermite polynomials. In this work, we consider normalized versions of the Hermite polynomials, denoted by $\{h_k(x)\}_k$. They are constructed as follows: $\deg h_k(x) = k$ and

$$\mathbb{E}h_i(z)h_j(z) = \mathbf{1}_{ij},$$

where $z \sim \mathcal{N}(0,1)$ and $\mathbf{1}_{ij}$ is the Kronecker delta. The first four (normalized) Hermite polynomials are

$$h_0(x) = 1, \quad h_1(x) = x, \quad h_2(x) = \frac{1}{\sqrt{2}}(x^2 - 1), \quad h_3(x) = \frac{1}{\sqrt{6}}(x^3 - 3x). \quad (\text{A.9})$$

Comparing (A.9) with (2.3) and (2.4), we can see that, for $k = 0, 1, 2, 3$, $h_k(x)$ is indeed the asymptotic limit of $q_k(x)$. For a general $k \in \mathbb{N}_0$, one can show (see [9, Lemma 4.1]) that

$$q_k(x) = h_k(x) + \sum_{i=0}^k c_{i,k}(d)x^i,$$

where $\max_i |c_{i,k}(d)| = \mathcal{O}_k(1/d)$.

In our discussions of the equivalence principle for general nonlinear functions f_d in Section 2.3, we need to use the following result, which is a consequence of [9, Lemma C.2, Lemma C.5]. We state the result in a lemma for easy reference.

Lemma 8. *Let $q_k(x)$ and $h_k(x)$ denote the k th Gegenbauer and the k th Hermite polynomials, respectively. Suppose that a function $f : \mathbb{R} \mapsto \mathbb{R}$ is locally bounded, *i.e.*, $\sup_{|x| \leq t} |f(x)| < \infty$ for all $t > 0$ and satisfies*

$$\mathbb{E}[f^2(z)\mathbf{1}(|z| \geq t)] \xrightarrow{t \rightarrow \infty} 0,$$

where $z \sim \mathcal{N}(0,1)$. For $\xi \sim \mathfrak{s}_d^1$, we have

$$\begin{aligned} \mathbb{E}[f^2(\xi)] &\xrightarrow{t \rightarrow \infty} \sigma^2 = \mathbb{E}[f^2(z)] < \infty, \\ \mathbb{E}[f(\xi)q_k(\xi)] &\xrightarrow{t \rightarrow \infty} \mu_k = \mathbb{E}[f(z)h_k(z)], \end{aligned}$$

for all $k \in \mathbb{N}_0$, and

$$\sum_{k=0}^{\infty} \mu_k^2 = \sigma^2.$$

B Proof of Proposition 1

We first state several simple properties of the set \mathcal{E}_m introduced in (4.10).

Lemma 9. For $m \in \mathbb{N}_0$, we have

$$\mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+1}, \quad (\text{B.1a})$$

$$\frac{\xi_1 \xi_2}{\sqrt{d}} \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+1}, \quad (\text{B.1b})$$

$$\frac{\xi_1^2}{d} \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}, \quad \frac{\xi_2^2}{d} \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2} \quad (\text{B.1c})$$

and

$$r^2(\xi_1) r^2(\xi_2) \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}, \quad (\text{B.1d})$$

where $r(\xi)$ is the function defined in (4.8).

Proof. The property (B.1a) follows immediately from the definition of \mathcal{E}_m . To show (B.1b), (B.1c), and (B.1d), we can assume without loss of generality that

$$\mathcal{E}_m(\xi_1, \xi_2) = \frac{q_a(\xi_1) q_b(\xi_2)}{d^{\max\{a,b\}/2+1}} \quad (\text{B.2})$$

for some a, b such that $0 \leq a, b \leq m$. The more general case, where $\mathcal{E}_m(\xi_1, \xi_2)$ is a linear combination of terms like (B.2), can be handled by using the linearity of \mathcal{E}_{m+1} and \mathcal{E}_{m+2} .

The recurrent relation (A.1) implies that $\xi_1 q_a(\xi_1) = \sum_{i \in [0, a+1]} \mu_i q_i(\xi_1)$ and $\xi_2 q_b(\xi_2) = \sum_{j \in [0, b+1]} \nu_j q_j(\xi_2)$, where $\mu_i = \mathcal{O}(1), \nu_j = \mathcal{O}(1)$ are some fixed expansion coefficients. It follows that

$$\frac{\xi_1 \xi_2}{\sqrt{d}} \mathcal{E}_m(\xi_1, \xi_2) = \frac{\sum_{i=0}^{a+1} \sum_{j=0}^{b+1} \mu_i \nu_j q_i(\xi_1) q_j(\xi_2)}{d^{\max\{a+1, b+1\}/2+1}} = \frac{\sum_{i=0}^{a+1} \sum_{j=0}^{b+1} \frac{\mu_i \nu_j}{d^{\pi_{ij}}} q_i(\xi_1) q_j(\xi_2)}{d^{\max\{i,j\}/2+1}},$$

where $\pi_{ij} = \max\{a+1, b+1\}/2 - \max\{i, j\}/2 \geq 0$. Thus, we $\frac{\xi_1 \xi_2}{\sqrt{d}} \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+1}$.

The properties stated in (B.1c) can be proved similarly. Consider the function in (B.2). We have $\xi_1^2 q_a(\xi_1) = \sum_{i=0}^{a+2} \mu_i q_i(\xi_1)$ for some expansion coefficients $\mu_i = \mathcal{O}(1)$. It follows that

$$\left(\xi_1^2/d\right) \mathcal{E}_m(\xi_1, \xi_2) = \frac{\sum_{i=0}^{a+2} \frac{\mu_i}{d^{\pi_i}} q_i(\xi_1) q_b(\xi_2)}{d^{\max\{i,b\}/2+1}},$$

where $\pi_i = \max\{a+2, b+2\}/2 - \max\{i, b\}/2 \geq 0$. Since $\mu_i/d^{\pi_i} = \mathcal{O}(1)$, we have $(\xi_1^2/d) \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}$. That $(\xi_2^2/d) \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}$ follows from analogous arguments.

Recall the definition of $r(\xi)$ in (4.8). We have

$$\begin{aligned} r^2(\xi_1) r^2(\xi_2) \mathcal{E}_m(\xi_1, \xi_2) &= (1 - 1/d)^{-1} (1 - \xi_1^2/d) (1 - \xi_2^2/d) \mathcal{E}_m(\xi_1, \xi_2) \\ &= (1 + \mathcal{O}(1/d)) \left(1 + (\xi_1 \xi_2 / \sqrt{d})^2 - \xi_1^2/d - \xi_2^2/d\right) \mathcal{E}_m(\xi_1, \xi_2). \end{aligned} \quad (\text{B.3})$$

Note that $\mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}$ by (B.1a). Applying the property in (B.1b) twice, we have $(\xi_1 \xi_2 / \sqrt{d})^2 \mathcal{E}_m(\xi_1, \xi_2) \in \mathcal{E}_{m+2}$. By (B.1c), the last two terms of (B.3) also belong to \mathcal{E}_{m+2} . By the linearity of \mathcal{E}_{m+2} , we can conclude that the left-hand side of (B.1d) indeed belongs to \mathcal{E}_{m+2} . \square

Lemma 10. For each $k \in \mathbb{N}$, we have

$$\frac{\xi_1 \xi_2 q_k(\xi_1) q_k(\xi_2)}{d^{(k+1)/2}} = (k+1) \frac{q_{k+1}(\xi_1) q_{k+1}(\xi_2)}{d^{(k+1)/2}} + \sqrt{k(k+1)} \frac{q_{k-1}(\xi_1) q_{k+1}(\xi_2) + q_{k+1}(\xi_1) q_{k-1}(\xi_2)}{d^{(k+1)/2}} + \mathcal{E}_{k+1}(\xi_1, \xi_2), \quad (\text{B.4})$$

and

$$\frac{r^2(\xi_1) r^2(\xi_2) q_{k-1}(\xi_1) q_{k-1}(\xi_2)}{d^{(k-1)/2}} = \frac{q_{k-1}(\xi_1) q_{k-1}(\xi_2)}{d^{(k-1)/2}} - \sqrt{k(k+1)} \frac{q_{k-1}(\xi_1) q_{k+1}(\xi_2) + q_{k+1}(\xi_1) q_{k-1}(\xi_2)}{d^{(k+1)/2}} + \mathcal{E}_{k+1}(\xi_1, \xi_2). \quad (\text{B.5})$$

Proof. To lighten the notation, we write, for any $a, b \in \mathbb{N}_0$,

$$q_{a,b} = q_a(\xi_1)q_b(\xi_2) \quad \text{and} \quad r_{1,2} = r(\xi_1)r(\xi_2). \quad (\text{B.6})$$

To show (B.4), we use the recurrent relation in (A.1), which gives us $\xi q_k(\xi) = a_k q_{k+1}(\xi) + a_{k-1} q_{k-1}(\xi)$. It follows that

$$\frac{\xi_1 \xi_2 q_{k,k}}{d^{(k+1)/2}} = a_k^2 \frac{q_{k+1,k+1}}{d^{(k+1)/2}} + a_k a_{k-1} \frac{q_{k-1,k+1} + q_{k+1,k-1}}{d^{(k+1)/2}} + a_{k-1}^2 \frac{q_{k-1,k-1}}{d^{(k-1)/2+1}}.$$

Note that the last term on the right-hand side of above expression is in \mathcal{E}_{k-1} and thus in \mathcal{E}_{k+1} (by (B.1a)). From (A.2), we have $a_k^2 = (k+1)(1 + \mathcal{O}(1/d))$ and $a_k a_{k-1} = \sqrt{k(k+1)}(1 + \mathcal{O}(1/d))$. Thus, $a_k^2 \frac{q_{k+1,k+1}}{d^{(k+1)/2}} = (k+1) \frac{q_{k+1,k+1}}{d^{(k+1)/2}} + \mathcal{E}_{k+1}(\xi_1, \xi_2)$, and similarly, $a_k a_{k-1} \frac{q_{k-1,k+1} + q_{k+1,k-1}}{d^{(k+1)/2}} = \sqrt{k(k+1)} \frac{q_{k-1,k+1} + q_{k+1,k-1}}{d^{(k+1)/2}} + \mathcal{E}_{k+1}(\xi_1, \xi_2)$.

Next, we show (B.5). Recall from the definition in (4.8) that $\sqrt{1 + d^{-1}r^2}(\xi) = 1 - \xi^2/d$. Thus,

$$\begin{aligned} \sqrt{1 + d^{-1}r^2}(\xi)q_{k-1}(\xi) &= q_{k-1}(\xi) - [\xi^2 q_{k-1}(\xi)]/d \\ &= q_{k-1}(\xi) - [a_{k-1}a_k q_{k+1}(\xi) + (a_{k-1}^2 + a_{k-2}^2)q_{k-1}(\xi) + a_{k-2}a_{k-3}q_{k-2}(\xi)]/d \end{aligned} \quad (\text{B.7})$$

where to reach (B.7), we have applied the recurrent formula (A.1) twice. Here, we have assumed that $k \geq 3$. In fact, the formula is valid for all $k \in \mathbb{N}$, if we define $a_{-1} = a_{-2} = 0$ and $q_{-1}(\xi) = q_{-2}(\xi) = 0$. Using (B.7), we can write

$$\frac{r_{1,2}^2 q_{k-1,k-1}}{d^{(k-1)/2}} = [d/(d+1)] \frac{q_{k-1,k-1}}{d^{(k-1)/2}} - [d/(d+1)] a_{k-1} a_k \frac{q_{k-1,k+1} + q_{k+1,k-1}}{d^{(k+1)/2}} + \mathcal{E}_{k+1}(\xi_1, \xi_2),$$

which then leads to (B.5), as $d/(d+1) = 1 + \mathcal{O}(1/d)$ and $[d/(d+1)] a_{k-1} a_k = \sqrt{k(k+1)} + \mathcal{O}(1/d)$. \square

Next, we prove Proposition 1 by induction on the polynomial degree k . Throughout the proof, we use the shorthand notation introduced in (B.6). Recall that $q_0(x) = \tilde{q}_0(x) = 1$ and $q_1(x) = \tilde{q}_1(x) = x$. It is then straightforward to verify the formula (4.11) for $k = 0, 1$. Specifically,

$$q_0(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) = \tilde{q}_0(x)$$

and

$$q_1(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) = \tilde{q}_1(x)r_{1,2} + \tilde{q}_0(x) \frac{q_{1,1}}{\sqrt{d}}.$$

Now we carry out the induction. Assume that (4.11) holds for k and $k-1$, with some $k \geq 1$. To prove it for $k+1$, we apply the recurrence relation in (A.1), which gives us

$$\begin{aligned} a_k q_{k+1}(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) &= (r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) q_k(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) - a_{k-1} q_{k-1}(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) \\ &= (r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}}) \sum_{m=0}^k \tilde{q}_{k-m}(x) r_{1,2}^{k-m} \left[\sqrt{(k)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right] \\ &\quad - a_{k-1} \sum_{m=0}^{k-1} \tilde{q}_{k-1-m}(x) r_{1,2}^{k-1-m} \left[\sqrt{(k-1)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right], \end{aligned} \quad (\text{B.8})$$

where in reaching the last step we have used (4.11) to expand $q_k(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}})$ and $q_{k-1}(r_{1,2}x + \frac{\xi_1 \xi_2}{\sqrt{d}})$.

On the right-hand side of (B.8) there are factors related to x in the form of $x \tilde{q}_{k-m}(x)$. They are polynomials of x , and can thus be rewritten as a linear combination of the orthogonal polynomials. To that end, we first recall from (2.5) that $\{\tilde{q}_k(x)\}$ denote the orthogonal polynomials defined for dimension $d-1$. Similar to (A.1), they also satisfy a recurrence relation

$$x \tilde{q}_k(x) = \tilde{a}_k \tilde{q}_{k+1}(x) + \tilde{a}_{k-1} \tilde{q}_{k-1}(x),$$

where the coefficients are

$$\tilde{a}_k = \sqrt{\frac{(k+1)(d+k-3)(d-1)}{(d+2k-1)(d+2k-3)}} = \sqrt{k+1} (1 + \mathcal{O}(1/d)). \quad (\text{B.9})$$

Applying this formula, and by setting $\tilde{a}_{-1} = 0$ and $\tilde{q}_{-1}(x) = 0$, we can write

$$x\tilde{q}_{k-m}(x) = \tilde{a}_{k-m}\tilde{q}_{k+1-m}(x) + \tilde{a}_{k-1-m}\tilde{q}_{k-1-m}(x) \quad \text{for } 0 \leq m \leq k. \quad (\text{B.10})$$

Replacing all the factors of $x\tilde{q}_{k-m}(x)$ in (B.8) by the right-hand side of (B.10), we can rewrite (B.8) as a linear combination of the orthogonal polynomials $\{\tilde{q}_{k+1-m}(x)\}_{m=0}^{k+1}$, *i.e.*,

$$\begin{aligned} q_{k+1}(r_{1,2}x + \frac{\xi_1\xi_2}{\sqrt{d}}) &= \sum_{m=0}^k \tilde{q}_{k+1-m}(x)r_{1,2}^{k+1-m} \frac{\tilde{a}_{k-m}}{a_k} \left[\sqrt{(k)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right] \\ &\quad + \sum_{m=0}^{k-1} \tilde{q}_{k-1-m}(x)r_{1,2}^{k+1-m} \frac{\tilde{a}_{k-1-m}}{a_k} \left[\sqrt{(k)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right] \\ &\quad + \sum_{m=0}^k \tilde{q}_{k-m}(x)r_{1,2}^{k-m} \frac{\xi_1\xi_2}{a_k\sqrt{d}} \left[\sqrt{(k)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right] \\ &\quad - \sum_{m=0}^{k-1} \tilde{q}_{k-1-m}(x)r_{1,2}^{k-1-m} \frac{a_{k-1}}{a_k} \left[\sqrt{(k-1)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right] \\ &= \sum_{m=0}^{k+1} \tilde{q}_{k+1-m}(x)C_m(\xi_1, \xi_2), \end{aligned} \quad (\text{B.11})$$

where $\{C_m(\xi_1, \xi_2)\}_{m=0}^{k+1}$ are some functions that only depend on ξ_1, ξ_2 but not on x . Next, we identify the exact expressions for $C_m(\xi_1, \xi_2)$.

By examining (B.11), we can see that

$$C_0(\xi_1, \xi_2) = r_{1,2}^{k+1} \frac{\tilde{a}_k}{a_k} [1 + \mathcal{E}_0(\xi_1, \xi_2)] = r_{1,2}^{k+1} [1 + \mathcal{E}_0(\xi_1, \xi_2)], \quad (\text{B.12})$$

where the last equality follows from (A.2) and (B.9).

$$\begin{aligned} C_1(\xi_1, \xi_2) &= r_{1,2}^k \frac{\tilde{a}_{k-1}}{a_k} \left[\sqrt{k} \frac{q_{1,1}}{\sqrt{d}} + \mathcal{E}_1(\xi_1, \xi_2) \right] + r_{1,2}^k \frac{\xi_1\xi_2}{a_k\sqrt{d}} \left[1 + \mathcal{E}_0(\xi_1, \xi_2) \right] \\ &= r_{1,2}^k \frac{\sqrt{k}}{\sqrt{k+1}} \left[\sqrt{k} \frac{q_{1,1}}{\sqrt{d}} + \mathcal{E}_1(\xi_1, \xi_2) \right] + r_{1,2}^k \frac{1}{\sqrt{k+1}} \left[\frac{q_{1,1}}{\sqrt{d}} + \mathcal{E}_1(\xi_1, \xi_2) \right] \quad \text{by (A.2), (B.9) and (B.1b)} \\ &= r_{1,2}^k \left[\sqrt{k+1} \frac{q_{1,1}}{\sqrt{d}} + \mathcal{E}_1(\xi_1, \xi_2) \right]. \end{aligned} \quad (\text{B.13})$$

For each m in the range $2 \leq m \leq k+1$, we have

$$\begin{aligned} C_m(\xi_1, \xi_2) &= r_{1,2}^{k+1-m} \underbrace{\left[\frac{\tilde{a}_{k-m}}{a_k} \sqrt{(k)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right]}_{\mathcal{F}_1} \\ &\quad + r_{1,2}^{k+1-m} \underbrace{\left[\frac{\tilde{a}_{k+1-m}}{a_k} \sqrt{(k)_{m-2}} (r_{1,2}^2 \frac{q_{m-2,m-2}}{d^{(m-2)/2}}) + r_{1,2}^2 \mathcal{E}_{m-2}(\xi_1, \xi_2) \right]}_{\mathcal{F}_2} \\ &\quad + r_{1,2}^{k+1-m} \underbrace{\left[\frac{1}{a_k} \sqrt{(k)_{m-1}} \frac{\xi_1\xi_2 q_{m-1,m-1}}{d^{m/2}} + \frac{\xi_1\xi_2}{\sqrt{d}} \mathcal{E}_{m-1}(\xi_1, \xi_2) \right]}_{\mathcal{F}_3} \\ &\quad - r_{1,2}^{k+1-m} \underbrace{\left[\frac{a_{k-1}}{a_k} \sqrt{(k-1)_{m-2}} \frac{q_{m-2,m-2}}{d^{(m-2)/2}} + \mathcal{E}_{m-2}(\xi_1, \xi_2) \right]}_{\mathcal{F}_4}. \end{aligned} \quad (\text{B.14})$$

Using the formulas for a_k and \tilde{a}_k in (A.2) and (B.9), we have

$$\mathcal{F}_1 = \sqrt{\frac{(k+1-m)(k)_m}{k+1}} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) = \frac{k+1-m}{k+1} \sqrt{(k+1)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2). \quad (\text{B.15})$$

Applying (B.5) and (B.1d) gives us

$$\mathcal{F}_2 = \sqrt{\frac{(k)_m}{k+1}} \left(\frac{q_{m-2,m-2}}{d^{(m-2)/2}} - \sqrt{m(m-1)} \frac{q_{m-2,m} + q_{m,m-2}}{d^{m/2}} \right) + \mathcal{E}_m(\xi_1, \xi_2). \quad (\text{B.16})$$

By (B.4) and (B.1b),

$$\mathcal{F}_3 = \sqrt{\frac{(k)_m}{k+1}} \left(m \frac{q_{m,m}}{d^{m/2}} + \sqrt{m(m-1)} \frac{q_{m-2,m} + q_{m,m-2}}{d^{m/2}} \right) + \mathcal{E}_m(\xi_1, \xi_2). \quad (\text{B.17})$$

Similarly, one can verify that

$$\mathcal{F}_4 = \sqrt{\frac{(k)_m}{k+1}} \frac{q_{m-2,m-2}}{d^{(m-2)/2}} + \mathcal{E}_m(\xi_1, \xi_2). \quad (\text{B.18})$$

Substituting (B.15), (B.16), (B.17), (B.18) into (B.14), we have

$$C_m(\xi_1, \xi_2) = r_{1,2}^{k+1-m} \left(\sqrt{(k+1)_m} \frac{q_{m,m}}{d^{m/2}} + \mathcal{E}_m(\xi_1, \xi_2) \right). \quad (\text{B.19})$$

Note that the expressions in (B.12), (B.13) and (B.19) exactly match the right-hand side of (4.11) for $k+1$. Thus, by induction, the formula (4.11) holds for all k .

C Review of the Marchenko-Pastur Law

In this appendix, we review some basic properties of the Marchenko-Pastur (MP) law that will be used in our work. Let X be an $M \times n$ matrix whose entries $X_{\mu i}$ are independent complex-valued random variables satisfying

$$\mathbb{E}X_{\mu i} = 0 \quad \mathbb{E}|X_{\mu i}|^2 = \frac{1}{n}.$$

We assume that M and n satisfy the bounds

$$n^{1/C} \leq M \leq n^C$$

for some positive constant C , and define the aspect ratio

$$\phi := \phi_n = \frac{M}{n},$$

which may depend on n . Now consider an $n \times n$ matrix

$$R = X^\top \Sigma X - \frac{\text{tr} \Sigma}{n} I, \quad (\text{C.1})$$

where Σ is a diagonal $M \times M$ matrix with matrix elements σ_i on the diagonal. Note that the subtraction by $\frac{\text{tr} \Sigma}{n} I$ in (C.1) makes sure that the diagonal elements of R are approximately equal to 0. We do this to match the construction in (1.1).

Assuming that the law of the diagonal entries of Σ is given by a probability law $\pi_M = M^{-1} \sum \delta_{\sigma_i} \rightarrow \pi$. The Marchenko-Pastur law asserts that the limiting Stieltjes transform of the eigenvalues of R satisfies the self-consistent equation

$$\frac{1}{m} = -z - \phi \int \frac{x^2 m}{1 + xm} \pi(dx). \quad (\text{C.2})$$

See, e.g., eq. (2.10) in Knowles and Yin [24]. (Notice that our equation is slightly different due to a constant shift of the eigenvalues). In the special case of $\Sigma = tI$ and $R = tX^\top X - t\phi I$, the above formula reduces to

$$\frac{1}{m} = -z - \phi \frac{t^2 m}{1 + tm}. \quad (\text{C.3})$$

The limiting eigenvalue density associated with (C.3) is given by

$$\varrho_{\phi,t}(x) dx := \frac{\sqrt{[(2\sqrt{\phi} + x/t - 1)(2\sqrt{\phi} - x/t + 1)]_+}}{2\pi \text{sign}(t)[x + t\phi]} dx + (1 - \phi)_+ \delta(x + t\phi). \quad (\text{C.4})$$

As a side note, we remark that a different convention was used in Bloemendal et al. [6], where Y is an $M \times n$ matrix whose entries $Y_{i\mu}$ are independent random variables satisfying $\mathbb{E}Y_{i\mu} = 0$, $\mathbb{E}Y_{i\mu}^2 = \frac{1}{\sqrt{nM}}$. One can check easily that $\varrho_{tH+r}(x) dx = \varrho_H\left(\frac{x-r}{t}\right) \frac{dx}{|t|}$ where ϱ_H denotes the density of a symmetric matrix H . Hence $R = t[\sqrt{\phi}Y^\top Y - \phi I]$ and we have $\varrho_R(x) dx = \varrho_{Y^\top Y}\left(\frac{x}{t\sqrt{\phi}} + \sqrt{\phi}\right) / (|t|\sqrt{\phi}) dx$. Thus our formula is identical to the one that appeared in [6, eq. (3.3)].

D Moment Bounds and Concentration Inequalities

In this appendix, we derive several moment and concentration inequalities that will be used in our proof.

Lemma 11. *Let $\xi \sim s_d^1$, and $q_k(x)$ be the k th Gegenbauer polynomial for $k \in \mathbb{N}_0$. For each $p \in \mathbb{N}$, we have*

$$\|\xi\|_{L^p} = (\mathbb{E}|\xi|^p)^{1/p} \leq 2\sqrt{p} \quad (\text{D.1})$$

and

$$\|q_k(\xi)\|_{L^p} \leq C(k, p) \quad (\text{D.2})$$

where $C(k, p)$ is some constant that only depends on k and p .

Remark 5. *Both the probability measure s_d^1 and the Gegenbauer polynomials $q_k(x)$ depend on the dimension d . However, the upper bounds in (D.1) and (D.2) hold uniformly for all d .*

Proof. The probability distribution of ξ is given by

$$w_d(\xi) = \begin{cases} \frac{\Gamma(d/2)}{\sqrt{d\pi} \Gamma((d-1)/2)} (1 - \xi^2/d)^{\frac{d-3}{2}} & \text{if } |\xi| \leq \sqrt{d} \\ 0 & \text{otherwise,} \end{cases}$$

where $\Gamma(\cdot)$ is the gamma function. Thus, we have

$$\begin{aligned} \mathbb{E}|\xi|^p &= \frac{2\Gamma(d/2)}{\sqrt{d\pi} \Gamma((d-1)/2)} \int_0^{\sqrt{d}} \xi^p (1 - \xi^2/d)^{\frac{d-3}{2}} d\xi \\ &= \frac{d^{p/2} \Gamma(d/2) \Gamma((p+1)/2)}{\Gamma((d+p)/2) \Gamma(1/2)} \\ &\leq (4p)^{p/2}, \end{aligned} \quad (\text{D.3})$$

where the second line uses the relationship between the beta function and the gamma function [1], and the last line is obtained by applying Stirling's formula for the gamma function [23].

Next, we show (D.2). Two special cases are easy: for $k = 0$, we have $\|q_0(\xi)\|_{L^p} = 1$; for $p = 1$, $\|q_k(\xi)\|_{L^1} \leq \|q_k(\xi)\|_{L^2} = 1$. Thus, in what follows we assume $k \geq 1$ and $p \geq 2$. Denote by c_k the leading coefficient of the polynomial $q_k(\xi)$. Since $q_k(\xi) - c_k \xi^k$ is a polynomial of degree $k-1$, it can be written as linear combination of the lower order Gegenbauer polynomials, i.e.,

$$q_k(\xi) = c_k \xi^k + \sum_{i=0}^{k-1} b_{k,i} q_i(\xi),$$

where $\{b_{k,i}\}_i$ are the expansion coefficients. By using the orthogonality of the Gegenbauer polynomials (see (2.2)), we have, for $0 \leq i \leq k-1$,

$$0 = \mathbb{E}[q_k(\xi)q_i(\xi)] = c_k \mathbb{E}[\xi^k q_i(\xi)] + b_{k,i},$$

and thus

$$q_k(\xi) = c_k \xi^k - c_k \sum_{i=0}^{k-1} \mathbb{E}[\xi^k q_i(\xi)] q_i(\xi).$$

From the triangular inequality,

$$\|q_k(\xi)\|_{L^p} \leq c_k \|\xi^k\|_{L^p} + c_k \sum_{i=0}^{k-1} \left| \mathbb{E}[\xi^k q_i(\xi)] \right| \cdot \|q_i(\xi)\|_{L^p}. \quad (\text{D.4})$$

Applying the Cauchy-Schwarz inequality, we have

$$\left| \mathbb{E}[\xi^k q_i(\xi)] \right| \leq [\mathbb{E} \xi^{2k} \mathbb{E} q_i^2(\xi)]^{1/2} \leq (8k)^{k/2},$$

where the last inequality is due to (D.1) and the fact that $\mathbb{E} q_i^2(\xi) = 1$. In addition, $\|\xi^k\|_{L^p} \leq (4pk)^{k/2}$ by (D.1). Substituting these two bounds into (D.4) then gives us, for $p \geq 2$,

$$\|q_k(\xi)\|_{L^p} \leq c_k (4pk)^{k/2} \left(1 + \sum_{i=0}^{k-1} \|q_i(\xi)\|_{L^p} \right). \quad (\text{D.5})$$

We now provide a bound for c_k , the leading coefficient of $q_k(\xi)$. Recall from (2.3) that $c_1 = 1$ and $c_2 = (1/\sqrt{2})\sqrt{(d+2)/(d-1)}$. A general formula for c_k can be obtained by examining the three-term recurrence relation in (A.1), which gives us $c_{k+1} = c_k/a_k$ for every $k \in \mathbb{N}$. This implies that $c_k = (a_0 a_1 a_2 \dots a_{k-1})^{-1}$, and from the explicit formula for a_k in (A.2),

$$c_k = \prod_{i=0}^{k-1} \left(\frac{1 + (2/d)i}{1+i} \right)^{1/2} \left(1 + \frac{i}{i+d-2} \right)^{1/2} \leq 2^{k/2}$$

for all $d \geq 2$. Using this estimate in (D.5) leads to

$$\|q_k(\xi)\|_{L^p} \leq (8pk)^{k/2} \left(1 + \sum_{i=0}^{k-1} \|q_i(\xi)\|_{L^p} \right), \quad \text{for all } k \in \mathbb{N}.$$

Starting from $\|q_0(\xi)\|_{L^p} = 1$, we can apply the above bound recursively to verify (D.2). \square

Next, we derive several useful moment and high probability bounds for linear and quadratic functions of independent random variables. We first recall the following estimates, whose proof can be found in [16, Lemma 7.8, Lemma 7.9].

Lemma 12. *Let $(X_i)_{i \in [n]}$ and $(Y_j)_{j \in [m]}$ be independent families of random variables, and $(b_i)_{i \in [n]}$ and $(c_{ij})_{i \in [n], j \in [m]}$ be deterministic complex-valued coefficients; here $n, m \in \mathbb{N}$. Suppose that all entries of (X_i) and (Y_j) are independent and satisfy*

$$\mathbb{E}X = \mathbb{E}Y = 0, \quad \mathbb{E}|X|^2 = \mathbb{E}|Y|^2 = 1, \quad \|X\|_{L^p} \leq \kappa_p, \quad \|Y\|_{L^p} \leq \gamma_p,$$

for all $p \in \mathbb{N}$ and some constants κ_p, γ_p . Then, we have

$$\left\| \sum_i b_i X_i \right\|_{L^p} \leq (Cp)^{1/2} \kappa_p \left(\sum_i |b_i|^2 \right)^{1/2} \quad (\text{D.6a})$$

and

$$\left\| \sum_{i,j} c_{ij} X_i Y_j \right\|_{L^p} \leq Cp \kappa_p \gamma_p \left(\sum_{i,j} |c_{ij}|^2 \right)^{1/2}, \quad (\text{D.6b})$$

where C is an absolute constant.

Proposition 3. Let $(c_{ij})_{i,j \in [n]}$ be a deterministic matrix with complex-valued coefficients, and $(\xi_i)_{i \in [n]}$ be an independent family of random variables with $\xi_i \sim \mathfrak{s}_d^1$. For every $a, b \in \mathbb{N}$, we have

$$\left\| \sum_{i,j} c_{ij} q_a(\xi_i) q_b(\xi_j) - (\sum_i c_{ii}) \delta_{ab} \right\|_{L^p} \leq C(a, b, p) (\sum_{i,j} |c_{ij}|^2)^{1/2}, \quad (\text{D.7})$$

where $C(a, b, p)$ is a quantity that depends on a, b , and p , but not on n .

Proof. By the triangular inequality, we have

$$\left\| \sum_{i,j} c_{ij} q_a(\xi_i) q_b(\xi_j) - (\sum_i c_{ii}) \delta_{ab} \right\|_{L^p} \leq \left\| \sum_{i \neq j} c_{ij} q_a(\xi_i) q_b(\xi_j) \right\|_{L^p} + \left\| \sum_i c_{ii} (q_a(\xi_i) q_b(\xi_i) - \delta_{ab}) \right\|_{L^p}. \quad (\text{D.8})$$

To bound the first term on the right-hand side, we use the standard decoupling technique. Observe that, for every $i \neq j$, the following identity holds:

$$1 = \frac{1}{Z_n} \sum_{I \subset [n]} \mathbf{1}(i \in I) \mathbf{1}(j \in I^c),$$

where the sum ranges over all subsets of $[n]$, and $Z_n = 2^{n-2}$. Applying this identity and the triangular inequality allows us to write

$$\left\| \sum_{i \neq j} c_{ij} q_a(\xi_i) q_b(\xi_j) \right\|_{L^p} \leq \frac{1}{Z_n} \sum_{I \subset [n]} \left\| \sum_{i \in I, j \in I^c} c_{ij} q_a(\xi_i) q_b(\xi_j) \right\|_{L^p}. \quad (\text{D.9})$$

Note that the families of random variables $\{q_a(\xi_i)\}_{i \in I}$ and $\{q_b(\xi_j)\}_{j \in I^c}$ have mean-zero independent components with unit variances; they are also mutually independent. We can then apply (D.6b) to bound each term in the sum in (D.9). Moreover, the sum involves a total of $2^n - 2$ terms. Thus,

$$\left\| \sum_{i \neq j} c_{ij} q_a(\xi_i) q_b(\xi_j) \right\|_{L^p} \leq 4Cp \|q_a(\xi)\|_{L^p} \|q_b(\xi)\|_{L^p} \left(\sum_{i \neq j} |c_{ij}|^2 \right)^{1/2}. \quad (\text{D.10})$$

Now we bound the second term on the right-hand side of (D.8). Write $X_i = q_a(\xi_i) q_b(\xi_i) - \delta_{ab}$ and $\sigma = \|X_i\|_{L^2}$. Then $(X_i/\sigma)_i$ is a family of independent random variables that satisfy the condition of Lemma 12. Using (D.6a) gives us

$$\begin{aligned} \left\| \sum_i c_{ii} (q_a(\xi_i) q_b(\xi_i) - \delta_{ab}) \right\|_{L^p} &= \sigma \left\| \sum_i c_{ii} (X_i/\sigma) \right\|_{L^p} \\ &\leq (CP)^{1/2} \|X_i\|_{L^p} (\sum_i |c_{ii}|^2)^{1/2}. \end{aligned}$$

By the triangular inequality (in the first line) and the Cauchy-Schwarz inequality (in the third line),

$$\begin{aligned} \|X_i\|_{L^p} &\leq \|q_a(\xi_i) q_b(\xi_i)\|_{L^p} + 1 \\ &= (\mathbb{E} |q_a(\xi_i)|^p |q_b(\xi_i)|^p)^{1/p} + 1 \\ &\leq \|q_a(\xi_i)\|_{L^{2p}} \|q_b(\xi_i)\|_{L^{2p}} + 1. \end{aligned} \quad (\text{D.11})$$

Substituting (D.10) and (D.11) into (D.8), and by applying the moment bound in (D.2), we complete the proof of the statement in (D.7). \square

The moment estimates obtained above can be easily turned into high probability bounds, as follows.

Proposition 4. Suppose $n = \mathcal{O}(d^\ell)$ for some fixed $\ell \in \mathbb{N}$. Let $(\xi_i)_{i \in [n]}$ be an independent family of random variables with $\xi_i \sim \mathfrak{s}_d^1$, and let A_k be the matrix defined in (2.6). For each fixed $k \in \mathbb{N}_0$, we have the bounds

$$\max_{i \in [n]} |q_k(\xi_i)| \prec 1, \quad (\text{D.12})$$

$$\|A_k\|_\infty \prec \frac{1}{\sqrt{n}}. \quad (\text{D.13})$$

$$\max_{i \in [n]} |r^k(\xi_i) - 1| \prec \frac{1}{d}, \quad (\text{D.14})$$

where $r(\xi)$ is the function defined in (4.8).

Proof. For any $\epsilon > 0$ and $p \in \mathbb{N}$,

$$\begin{aligned} \mathbb{P}(\max_i |q_a(\xi_i)| \geq d^\epsilon) &\leq n \mathbb{P}(|q_a(\xi)| \geq d^\epsilon) \\ &\leq C d^\ell \mathbb{P}(|q_a(\xi)|^p \geq d^{\epsilon p}) \\ &\leq C d^{\ell - \epsilon p} C(a, p), \end{aligned}$$

where the last step is by the Markov inequality and by the moment bound in (D.2). For any $D > 0$, we can always find $p = p(\epsilon, D)$ such that $C d^{\ell - \epsilon p} C(a, p) < d^{-D}$ for all sufficiently large d . Since ϵ and D were arbitrary, we have verified (D.12). To show (D.13), observe that the law of $(A_k)_{ij}$, for any $i \neq j$, is equal to that of $q_k(\xi)/\sqrt{n}$ with $\xi \sim \mathfrak{s}_d^1$. By (D.12), we have $|(A_k)_{ij}| \prec 1/\sqrt{n}$. Applying the union bound over $\mathcal{O}(d^{2\ell})$ entries of A_k then yields (D.13).

Now we prove (D.14). The case of $k = 0$ is trivial, so we assume $k \geq 1$. From the definition in (4.8), $r(\xi) = (1 - \xi^2/d)^{1/2} + \mathcal{O}(1/d)$, and thus

$$\begin{aligned} |r(\xi) - 1| &\leq \left| 1 - (1 - \xi^2/d)^{1/2} \right| + \mathcal{O}(1/d) \\ &\leq \xi^2/d + \mathcal{O}(1/d). \end{aligned}$$

Combining this deterministic bound with the high-probability bound in (D.12), we have that, for $\xi_i \sim \mathfrak{s}_d^1$,

$$r(\xi_i) - 1 = \mathcal{O}_{\prec}(1/d). \quad (\text{D.15})$$

For each $k \in \mathbb{N}$,

$$\left| r^k(\xi_i) - 1 \right| \leq \sum_{t=1}^k \binom{k}{t} |r(\xi_i) - 1|^t.$$

Applying (D.15) and the union bound then allows us to conclude (D.14). \square

Proposition 5. *Suppose that $n = \mathcal{O}(d^\ell)$ for some fixed $\ell \in \mathbb{N}$. Let $(\xi_i)_{i \in [n]}$ be an independent family of random variables with $\xi_i \sim \mathfrak{s}_d^1$. For every $k, t \in \mathbb{N}_0$, and every function $\mathcal{E}_t(\xi_1, \xi_2)$ from the function class defined in (4.10), we can write*

$$r^k(\xi_1) r^k(\xi_2) \mathcal{E}_t(\xi_1, \xi_2) = f_1(\xi_1, \xi_2) + f_2(\xi_1, \xi_2), \quad (\text{D.16})$$

where the two functions f_1, f_2 are such that

$$f_1(\xi_1, \xi_2) \in \mathcal{E}_{t+2\ell} \quad (\text{D.17})$$

and

$$\max_{a, b \in [n]} |f_2(\xi_a, \xi_b)| \prec \frac{1}{d\sqrt{n}}. \quad (\text{D.18})$$

Proof. The special case when $k = 0$ is trivial, as we can simply choose $f_1(\xi_1, \xi_2) = \mathcal{E}_t(\xi_1, \xi_2)$ and $f_2(\xi_1, \xi_2) = 0$. In what follows we assume $k \geq 1$. For any $\nu \in \mathbb{N}$, consider the ν -th order Taylor expansion of the function $(1-x)^{k/2}$ around $x = 0$. This allows us to write

$$(1-x)^{k/2} = P_\nu(x) + R_\nu(x), \quad (\text{D.19})$$

where

$$P_\nu(x) = \sum_{i=0}^{\nu-1} \frac{(-1)^i (k/2)_i}{i!} x^i, \quad (\text{D.20})$$

and $R_\nu(x)$ is the remainder term. (In (D.20), $(k/2)_i = (k/2)(k/2-1)\dots(k/2-i+1)$ denotes the falling factorial.) For any $0 \leq x \leq 1$, we can verify from the Lagrange form of the the remainder that

$$|R_\nu(x)| \leq (k/2)_\nu \max \left\{ (1-x)^{(k/2)_{\nu+1}}, 1 \right\} x^\nu. \quad (\text{D.21})$$

Now recall the definition of $r(\xi)$ in (4.8). By using (D.19), we can indeed decompose $r^k(\xi_1)r^k(\xi_2)\mathcal{E}_t(\xi_1, \xi_2)$ into the form of (D.16), where

$$f_1(\xi_1, \xi_2) = (1 - 1/d)^{-k/2} P_\nu(\xi_1^2/d) P_\nu(\xi_2^2/d) \mathcal{E}_t(\xi_1, \xi_2)$$

and

$$f_2(\xi_1, \xi_2) = (1 - 1/d)^{-k/2} [R_\nu(\xi_1^2/d) P_\nu(\xi_2^2/d) + P_\nu(\xi_1^2/d) R_\nu(\xi_2^2/d) + R_\nu(\xi_1^2/d) R_\nu(\xi_2^2/d)] \mathcal{E}_t(\xi_1, \xi_2).$$

Note that $(1 - 1/d)^{-k/2} = 1 + \mathcal{O}(1/d)$. For an independent family of random variables $\{\xi_a\}_{a \in [n]}$ with $\xi \sim \mathfrak{s}_d^1$, we can use (D.12) to verify that $\max_a |P_\nu(\xi_a^2/d)| \prec 1$. Similarly, by the definition in (4.10) and (D.12), we have $\max_{a,b} |\mathcal{E}_t(\xi_a, \xi_b)| \prec 1/d$. It follows from these estimates that

$$\max_{a,b} |f_2(\xi_a, \xi_b)| \prec \frac{1}{d} (1 + \max_a |R_\nu(\xi_a^2/d)|) \max_a |R_\nu(\xi_a^2/d)|. \quad (\text{D.22})$$

Next, we establish a high-probability upper bound for $\max_a |R_\nu(\xi_a^2/d)|$. For any $a \in [n]$, $\epsilon \in (0, 1)$, and $D > 0$, and for sufficiently large d , we have

$$\begin{aligned} \mathbb{P}(|R_\nu(\xi_a^2/d)| \geq d^{\epsilon-\nu}) &\leq \mathbb{P}((1 - \xi_a^2/d)^{(k/2)_{\nu+1}} \xi_a^{2\nu} \geq d^{\epsilon/2}) + \mathbb{P}(\xi_a^{2\nu} \geq d^{\epsilon/2}) \\ &\leq \mathbb{P}((1 - \xi_a^2/d)^{(k/2)_{\nu+1}} \xi_a^{2\nu} \geq d^{\epsilon/2}, \xi_a^2 < d/2) + \mathbb{P}(\xi_a^2 \geq d/2) + \mathbb{P}(\xi_a^{2\nu} \geq d^{\epsilon/2}) \\ &\leq \mathbb{P}(\xi_a^{2\nu} \geq \max\{2^{(k/2)_{\nu+1}}, 1\} d^{\epsilon/2}) + \mathbb{P}(\xi_a^2 \geq d/2) + \mathbb{P}(\xi_a^{2\nu} \geq d^{\epsilon/2}) \\ &\leq d^{-D}. \end{aligned}$$

In the first step, we use the deterministic bound in (D.21) and the fact that $(k/2)_\nu < d^{\epsilon/2}$ for all sufficiently large d . The last step follows from the property that $\xi_a = \mathcal{O}_\prec(1)$. Since the above inequality is uniform in $a \in [n]$, we have $\max_a |R_\nu(\xi_a^2/d)| \prec d^{-\nu}$. Substituting this bound into (D.22) and choosing $\nu = \lceil \ell/2 \rceil$, we then reach the statement in (D.18).

To show (D.17), we use the definition of the Taylor polynomial in (D.20), which gives us

$$f_1(\xi_1, \xi_2) = (1 - 1/d)^{-k/2} \sum_{0 \leq i, j \leq \nu-1} c_i c_j (\xi_1^2/d)^i (\xi_2^2/d)^j \mathcal{E}_t(\xi_1, \xi_2),$$

where $\{c_i\}$ are some fixed coefficients. The statement in (D.17) then follows from a repeated application of the property in (B.1c). \square

Proposition 6. *Let $(b_i)_{i \in [n]}$, $(c_{ij})_{i, j \in [n]}$ and $(\xi_i)_{i \in [n]}$ be three independent families of random variables. Moreover, (ξ_i) is i.i.d., with $\xi_i \sim \mathfrak{s}_d^1$. Suppose that*

$$\left(\sum_i |b_i|^2\right)^{1/2} \prec \Psi_b \quad \text{and} \quad \left(\sum_{i, j} |c_{ij}|^2\right)^{1/2} \prec \Psi_c \quad (\text{D.23})$$

for some random variables Ψ_b and Ψ_c . Then, for every fixed $L \in \mathbb{N}$, and any $a, b \in [L]$, we have

$$\left|\sum_i b_i q_a(\xi_i)\right| \prec \Psi_b, \quad (\text{D.24a})$$

and

$$\left|\sum_{i, j} c_{ij} q_a(\xi_i) q_b(\xi_j) - \left(\sum_i c_{ii}\right) \delta_{ab}\right| \prec \Psi_c. \quad (\text{D.24b})$$

Proof. For any $\epsilon > 0$, $D > 0$,

$$\begin{aligned} &\mathbb{P}\left(\left|\sum_{i, j} c_{ij} q_a(\xi_i) q_b(\xi_j) - \left(\sum_i c_{ii}\right) \delta_{ab}\right| \geq d^\epsilon \Psi\right) \\ &\leq \mathbb{P}\left(\left|\sum_{i, j} c_{ij} q_a(\xi_i) q_b(\xi_j) - \left(\sum_i c_{ii}\right) \delta_{ab}\right| \geq d^\epsilon \Psi \text{ and } \left(\sum_{i, j} |c_{ij}|^2\right)^{1/2} \leq d^{\epsilon/2} \Psi\right) \\ &\quad + \mathbb{P}\left(\left(\sum_{i, j} |c_{ij}|^2\right)^{1/2} > d^{\epsilon/2} \Psi\right) \\ &\leq \mathbb{P}\left(\left|\sum_{i, j} c_{ij} q_a(\xi_i) q_b(\xi_j) - \left(\sum_i c_{ii}\right) \delta_{ab}\right| \geq d^{\epsilon/2} \left(\sum_{i, j} |c_{ij}|^2\right)^{1/2}\right) + d^{-(D+1)} \\ &\leq \left(\frac{C(a, b, p)}{d^{\epsilon/2}}\right)^p + d^{-(D+1)}. \end{aligned}$$

In the second step, we have used the definition of the stochastic dominance inequality (D.23) with parameters $\epsilon/2$ and $D + 1$. The last step follows from the Markov inequality and the moment bound (D.7). For any $D > 0$, there is a large enough p such that $\epsilon p/2 \geq D + 1$. Thus, the right-hand side can be bounded by d^{-D} for all sufficiently large d . This then established the bound in (D.24b).

The proof of (D.24a) follows exactly the same arguments. The only difference is that, instead of using the moment bound in (D.7), we appeal to the bound in (D.6a) and the moment estimate of $q_a(\xi)$ given in (D.2) when applying the Markov inequality. We omit the details. \square

Finally, we recall a concentration result for the Stieltjes transforms of matrices in the form of (1.1).

Lemma 13. *Let A be the matrix in (1.1), and $s(z)$ its Stieltjes transform. We have*

$$s(z) = \mathbb{E}s(z) + \mathcal{O}_{\prec}\left(\frac{1}{\eta\sqrt{n}}\right). \quad (\text{D.25})$$

Proof. The estimate in (D.25) can be established by standard arguments in random matrix theory. One approach is to view $s(z)$, for fixed z , as a function of the data vectors $\{\mathbf{x}_i\}_{i \in [n]}$. It is easy to verify that this function has bounded variations. This then allows one to apply McDiarmid's inequality to reach (D.25).

Another approach, as used in [9], is to apply the Burkholder's inequality [2, Lemma 2.12] to directly bound the moments of the difference between $s(z)$ and $\mathbb{E}s(z)$. By [9, Lemma 2.4], for any $p \in \mathbb{N}$,

$$\mathbb{E}|s(z) - \mathbb{E}s(z)|^{2p} \leq \frac{C_p}{\eta^{2p}n^p},$$

where C_p is a constant that depends on p . For any $\epsilon > 0$, applying Markov's inequality gives us

$$\mathbb{P}\left(|s(z) - \mathbb{E}s(z)| \geq \frac{d^\epsilon}{\eta\sqrt{n}}\right) = \mathbb{P}\left(|s(z) - \mathbb{E}s(z)|^{2p} \geq \frac{d^{2\epsilon p}}{\eta^{2p}n^p}\right) \leq C_p d^{-2\epsilon p}.$$

For any $D > 0$, we can always find a p such that the right-hand side of the above inequality is less than d^{-D} for all sufficiently large d . This then completes the proof. \square

E The Operator Norms of $\{A_k\}$

In this appendix, we present some high probability bounds for the operator norms of the matrices $\{A_k\}$ defined in (2.6).

E.1 The Low-Order Components

We start with the low-order components, *i.e.*, those A_k with $0 \leq k < \ell$.

Lemma 14. *Suppose that $n = \alpha d^\ell + o(d^\ell)$ for some $\ell \in \mathbb{N}$ and $\alpha > 0$. For each $k \in \mathbb{N}_0$ with $k < \ell$, let $\lambda_1(A_k) \geq \lambda_2(A_k) \geq \dots \geq \lambda_n(A_k)$ denote the eigenvalues of A_k in non-increasing order. For all sufficiently large d , we have*

$$\max_{i \in [N_k]} \left| \lambda_i(A_k) - \sqrt{n/N_k} \right| = \mathcal{O}_{\prec}(1) \quad (\text{E.1})$$

and

$$\lambda_i(A_k) = -\sqrt{N_k/n} \quad \text{for all } i > N_k. \quad (\text{E.2})$$

Remark 6. *Recall from (A.3) that $N_k = d^k/k! + \mathcal{O}(d^{k-1})$. Thus, the condition $k < \ell$ ensures that $N_k \ll n$ for all sufficiently large d . The lemma states that the spectrum of A_k can be divided into two clusters: the top N_k eigenvalues are in an interval of width $\mathcal{O}_{\prec}(1)$ centered at $\sqrt{n/N_k} = \sqrt{\alpha k! d^{\ell-k}(1 + o(1))}$; the remaining $n - N_k$ eigenvalues are all equal to $\sqrt{N_k/n} = o(1)$.*

Proof. Consider the factorized representation of A_k in (A.7). Observe that the positive-semidefinite matrix $Y_k^\top Y_k$ is rank-deficient. In fact, $\text{rank}(Y_k^\top Y_k) = N_k \ll n$, which then immediately gives us (E.2).

Next, we study the top N_k eigenvalues of A_k . In what follows, we use $\lambda_i(\mathbf{M})$ to represent the i th largest eigenvalue of any symmetric matrix \mathbf{M} . By (A.7), and for each $i \in [N_k]$, we have

$$\lambda_i(A_k) + \sqrt{N_k/n} = \frac{1}{\sqrt{nN_k}} \lambda_i(Y_k^\top Y_k) = \frac{1}{\sqrt{nN_k}} \lambda_i(Y_k Y_k^\top).$$

It follows that

$$\left| \lambda_i(A_k) - \sqrt{n/N_k} \right| \leq \frac{1}{\sqrt{nN_k}} \left| \lambda_i(Y_k Y_k^\top - nI) \right| + \sqrt{N_k/n}$$

and thus

$$\max_{i \in [N_k]} \left| \lambda_i(A_k) - \sqrt{n/N_k} \right| \leq \frac{1}{\sqrt{nN_k}} \left\| Y_k Y_k^\top - nI \right\|_{\text{op}} + \mathcal{O}(d^{-(\ell-k)}). \quad (\text{E.3})$$

The case of $k = 0$ is trivial: Recall that $N_0 = 1$ and $Y_{0,1}(\mathbf{x}) = 1$. Thus, $Y_0 = (1, 1, \dots, 1)$ is a constant row vector of length n , and $Y_0 Y_0^\top - n = 0$. For $1 \leq k < \ell$, we bound the operator norm of $Y_k Y_k^\top - nI$ by using the matrix Bernstein inequality. To that end, we first define a sequence of matrices $\{\mathbf{Z}_i = \mathbf{u}_i \mathbf{u}_i^\top - I\}_{i \in [n]}$ where

$$\mathbf{u}_i = (Y_{k,1}(\mathbf{x}_i), Y_{k,2}(\mathbf{x}_i), \dots, Y_{k,N_k}(\mathbf{x}_i))$$

is an N_k -dimensional random vector. One can verify from the definition of S_k in (A.8) that

$$S_k S_k^\top - nI = \sum_{i \in [n]} \mathbf{Z}_i. \quad (\text{E.4})$$

Since the data vectors $\mathbf{x}_i \sim_{\text{iid}} \text{Unif}(\mathcal{S}^{d-1})$, the matrices $\{\mathbf{Z}_i\}$ are independent. By (A.4), \mathbf{u}_i is an isotropic random vector and thus $\mathbb{E} \mathbf{Z}_i = \mathbf{0}$. Moreover, (A.6) implies that $\|\mathbf{u}_i\| = \sqrt{N_k}$. It follows that

$$K = \|\mathbf{Z}_i\|_{\text{op}} = N_k - 1$$

and

$$\sigma^2 = \left\| \sum_{i \in [n]} \mathbb{E} \mathbf{Z}_i^2 \right\|_{\text{op}} = \left\| \sum_{i \in [n]} \mathbb{E}[(N_k - 2)\mathbf{u}_i \mathbf{u}_i^\top + I] \right\|_{\text{op}} = n(N_k - 1).$$

For every $t > 0$, the standard matrix Bernstein inequality [34] gives us

$$\begin{aligned} \mathbb{P}\left(\frac{1}{\sqrt{nN_k}} \left\| \sum_{i \in [n]} \mathbf{Z}_i \right\|_{\text{op}} \geq t\right) &\leq 2N_k \exp\left(-\frac{nN_k t^2/2}{\sigma^2 + K\sqrt{nN_k}(t/3)}\right) \\ &\leq 2N_k \exp\left(-2 \min(t^2, \sqrt{n/N_k}t)\right). \end{aligned}$$

For every $0 < \epsilon < 1/2$, let $t = d^\epsilon$. Since $N_k = \mathcal{O}(d^k)$ with $k < \ell$, we have $\sqrt{N_k/n}(t/3) < 1$ for all sufficiently large d . It follows that, for every $D > 0$,

$$\mathbb{P}\left(\frac{1}{\sqrt{nN_k}} \left\| \sum_{i \in [n]} \mathbf{Z}_i \right\|_{\text{op}} \geq t\right) \leq d^{-D}$$

for sufficiently large d , and thus $\frac{1}{\sqrt{nN_k}} \left\| \sum_{i \in [n]} \mathbf{Z}_i \right\|_{\text{op}} = \mathcal{O}_{\prec}(1)$. The statement (E.1) of the lemma then follows from (E.4) and (E.3). \square

E.2 Moment Calculations for the High-Order Components

Next, we bound the operator norms of the high-order matrices, *i.e.*, those A_k with $k \geq \ell$. The approach taken in our proof of Lemma 14, which is based on the matrix Bernstein inequality, can be modified to show that $\|A_\ell\|_{\text{op}} = \mathcal{O}_{\prec}(1)$, but it does not seem to yield useful results for cases when $k > \ell$. In what follows, we

control the operator norm of A_k by using the standard moment method. Observe that, for each $\epsilon > 0$ and $p \in \mathbb{N}$, we have

$$\begin{aligned} \mathbb{P}(\|A_k\|_{\text{op}} \geq d^\epsilon) &= \mathbb{P}(\|A_k\|_{\text{op}}^{2p} \geq d^{2\epsilon p}) \\ &\leq d^{-2\epsilon p} \mathbb{E}\|A_k\|_{\text{op}}^{2p} \\ &\leq d^{-2\epsilon p} \mathbb{E} \text{tr}(A_k^{2p}). \end{aligned} \tag{E.5}$$

Thus, to control the probability on the left-hand side, it suffices to bound $\mathbb{E} \text{tr}(A_k^{2p})$ for a sufficiently large integer p . Bounds for the moments have been previously studied in the literature. Cheng and Singer [9] provides a bound for the 4th moment $\mathbb{E} \text{tr}(A_k^4)$. For general $p \in \mathbb{N}$, Ghorbani et al. [19, Lemma 3] showed that

$$\mathbb{E} \text{tr}(A_k^{2p}) \leq (Cp)^{3pn} (1 + d^{p(k-\ell) + \ell - 2k}) \tag{E.6}$$

for some positive constant C . Substituting (E.6) into (E.5), we can see that, when $k > \ell$, this moment bound is not sufficient to make (E.5) smaller than d^{-D} for any $D > 0$. In this work, we adapt the moment calculations in [19] and obtain a new bound for $\mathbb{E} \text{tr}(A_k^{2p})$ that is tight up to its leading order (see Lemma 17).

As in [19], our moment calculations hinge on the following property of Gegenbauer polynomials.

Lemma 15. *Let $\mathbf{z} \sim \text{Unif}(\mathcal{S}^{d-1})$. For each $k \in \mathbb{N}_0$ and any deterministic vectors \mathbf{x}, \mathbf{y} in \mathcal{S}^{d-1} , we have*

$$\mathbb{E}_{\mathbf{z}} q_k(\sqrt{d}\mathbf{x}^\top \mathbf{z}) q_k(\sqrt{d}\mathbf{z}^\top \mathbf{y}) = \frac{1}{\sqrt{N_k}} q_k(\sqrt{d}\mathbf{x}^\top \mathbf{y}). \tag{E.7}$$

Proof. Using (A.5), we can write

$$\begin{aligned} \mathbb{E}_{\mathbf{z}} q_k(\sqrt{d}\mathbf{x}^\top \mathbf{z}) q_k(\sqrt{d}\mathbf{z}^\top \mathbf{y}) &= \frac{1}{N_k} \sum_{a,b \in [N_k]} Y_{k,a}(\mathbf{x}) Y_{k,b}(\mathbf{y}) [\mathbb{E} Y_{k,a}(\mathbf{z}) Y_{k,b}(\mathbf{z})] \\ &= \frac{1}{N_k} \sum_{a,b \in [N_k]} Y_{k,a}(\mathbf{x}) Y_{k,b}(\mathbf{y}) \delta_{ab} \quad [\text{by (A.4)}] \\ &= \frac{1}{N_k} \sum_{a \in [N_k]} Y_{k,a}(\mathbf{x}) Y_{k,a}(\mathbf{y}). \end{aligned}$$

Applying (A.5) one more time then gives us (E.7). □

Lemma 16. *Let $\mathbf{x}_1, \dots, \mathbf{x}_n \sim_{\text{iid}} \text{Unif}(\mathcal{S}^{d-1})$. For $k \in \mathbb{N}_0$ and $i, j \in [n]$, write*

$$Q_{ij} := q_k(\sqrt{d}\mathbf{x}_i^\top \mathbf{x}_j). \tag{E.8}$$

Then, for every length $2p$ sequence of indices $\mathbf{i} = [i_1, i_2, \dots, i_{2p}] \in [n]^{2p}$, we have

$$|\mathbb{E} Q_{i_1 i_2} Q_{i_2 i_3} \dots Q_{i_{2p-1} i_{2p}} Q_{i_{2p} i_1}| \leq [\mathbb{E}_{\xi \sim \mathfrak{S}_d^1} q_k^{2p}(\xi)] N_k^{p-\nu(\mathbf{i})+1}, \tag{E.9}$$

where $\nu(\mathbf{i}) := \#\{i_1, i_2, \dots, i_{2p}\}$ is the number of distinct indices in \mathbf{i} , and \mathfrak{S}_d^1 is the probability measure defined in (1.3).

Remark 7. *This lemma is essentially a restatement of [19, Lemma 3], after one adjusts for the different definitions and scalings used in that work and this paper. For the sake of readability and completeness, we present the results in a form that is more convenient for our subsequent arguments, and provide a slightly simplified proof below.*

Proof. We can view the product $Q_{i_1 i_2} Q_{i_2 i_3} \dots Q_{i_{2p-1} i_{2p}} Q_{i_{2p} i_1}$ graphically, as a length $2p$ cycle on the vertex set $[n]$. First, consider two special cases: (I) $\nu(\mathbf{i}) = 1$, i.e., all the indices are identical. Recall from (A.6) that

$$Q_{ii} = \sqrt{N_k} \quad \text{for all } i. \tag{E.10}$$

Thus

$$\mathbb{E}Q_{i_1 i_2} Q_{i_2 i_3} \cdots Q_{i_{2p}, i_1} = Q_{i_1 i_1}^{2p} = N_k^p. \quad (\text{E.11})$$

(II) $\nu(\mathbf{i}) = 2p$, *i.e.*, all the indices are distinct. Mapping the indices to the canonical set $[2p]$, we have

$$\begin{aligned} \mathbb{E}Q_{12}Q_{23}Q_{34} \cdots Q_{2p,1} &= \mathbb{E}_{\mathbf{x}_1, \mathbf{x}_3, \dots, \mathbf{x}_{2p}} \left([\mathbb{E}_{\mathbf{x}_2} Q_{12}Q_{23}] Q_{34} \cdots Q_{2p,1} \right) \\ &= \frac{1}{\sqrt{N_k}} \mathbb{E}_{\mathbf{x}_1, \mathbf{x}_3, \dots, \mathbf{x}_{2p}} (Q_{13}Q_{34} \cdots Q_{2p,1}), \end{aligned} \quad (\text{E.12})$$

where the last step follows from (E.7). The above reduction procedure can be applied repeatedly: at each step, it reduces the length of the cycle by 1 and produces an extra factor of $1/\sqrt{N_k}$. Doing this for $2p - 1$ times then gives us

$$\mathbb{E}Q_{12}Q_{23} \cdots Q_{2p,1} = N_k^{-(2p-1)/2} \mathbb{E}Q_{11} = N_k^{1-p}, \quad (\text{E.13})$$

where the last equality is due to (E.10). Note that, since the Gegenbauer polynomials are normalized (recall (2.2)), we have

$$\mathbb{E}q_k^{2p}(\xi) \geq (\mathbb{E}q_k^2(\xi))^p = 1. \quad (\text{E.14})$$

Thus, with (E.11) and (E.13), we have verified the bound in (E.9) for the cases of $\nu(\mathbf{i}) = 1$ and $\nu(\mathbf{i}) = 2p$, respectively.

In fact, the procedures leading to (E.11) and (E.13) are special cases of a general and systematic reduction process. Suppose we are given a length t cycle on indices $\mathbf{i} = (i_1, i_2, \dots, i_t)$. To simplify the notation, define

$$M(\mathbf{i}) = \mathbb{E}Q_{i_1 i_2} Q_{i_2 i_3} \cdots Q_{i_{t-1} i_t} Q_{i_t i_1}.$$

We now introduce two reduction mappings, each of which converts a length t cycle to a length $(t - 1)$ cycle.

Type-A reduction: Given an index sequence $\mathbf{i} = (i_1, i_2, \dots, i_t)$ for $t \geq 2$. Suppose there exists at least one $j \in [t]$ such that $i_j \neq i_a$ for all $a \in [t] \setminus \{j\}$. In other words, the index i_j appears exactly once in the cycle. (It is possible that there are more than one such “singleton” indices in the sequence, in which case we will choose any one of them.) We remove i_j from the original sequence \mathbf{i} , and call the resulting length $t - 1$ sequence $R_A(\mathbf{i})$. More precisely,

$$R_A(\mathbf{i}) = (i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_t)$$

where the indices $j - 1$ and $j + 1$ are interpreted modulo $[t]$. Using the same conditioning technique that leads to (E.12), we can easily verify that

$$M(\mathbf{i}) = \frac{1}{\sqrt{N_k}} M(R_A(\mathbf{i})). \quad (\text{E.15})$$

Thus, a type-A reduction step $R_A(\mathbf{i})$ reduces the cycle length by 1 and contributes a factor of $1/\sqrt{N_k}$.

Type-B reduction: Given an index sequence $\mathbf{i} = (i_1, i_2, \dots, i_t)$ for $t \geq 2$. Suppose there exists at least one $j \in [t]$ such that $i_{j+1} = i_j$, where $j + 1$ is to be interpreted modulo $[t]$. (If more than one such indices exist, we choose any one of them.) We define

$$R_B(\mathbf{i}) = (i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_t)$$

as a length $t - 1$ sequence obtained by removing i_j from \mathbf{i} . By (E.10), we must have

$$M(\mathbf{i}) = \sqrt{N_k} M(R_B(\mathbf{i})). \quad (\text{E.16})$$

Thus, a type-B reduction step $R_B(\mathbf{i})$ reduces the cycle length by 1 and contributes a factor of $\sqrt{N_k}$.

Given an index sequence $\mathbf{i} = (i_1, i_2, \dots, i_t)$, we can simplify it by iteratively using the Type-A and Type-B reductions, until it cannot be further simplified. Clearly, this process stops after a finite number of steps. Let $\bar{\mathbf{i}} = (\bar{i}_1, \bar{i}_2, \dots, \bar{i}_{t(\bar{\mathbf{i}})})$ denote the final outcome of the iterative reduction process.

Example 1. *As an illustration, let us consider two concrete index sequences and demonstrate how they can be simplified through the above process.*

(a) $\mathbf{i} = (1, 1, 2, 2, 2, 3)$: We can start by using a Type-A step to remove the “singleton” 3. Then, the “redundant” copies of 1 and 2 can be removed by applying the Type-B step three times. This then gives us a shorter sequence (1, 2), in which both 1 and 2 are singletons. Removing 1 by a Type-A step gives us the final output $\bar{\mathbf{i}} = (2)$.

(b) $\mathbf{i} = (1, 2, 1, 2, 2, 3)$: Removing the “singleton” 3 (via Type-A reduction) and one redundant copy of 2 (via Type-B reduction), we get $\bar{\mathbf{i}} = (1, 2, 1, 2)$, which cannot be further simplified.

By (E.15) and (E.16), we must have

$$M(\mathbf{i}) = N_k^{(n_B - n_A)/2} M(\bar{\mathbf{i}}), \quad (\text{E.17})$$

where n_A (resp. n_B) is the total number of Type-A (resp. Type-B) steps used in the reduction process that leads to the final outcome $\bar{\mathbf{i}}$. Denote by $t(\mathbf{i})$ and $\nu(\mathbf{i})$ the length and number of unique indices in \mathbf{i} , respectively. We also define $t(\bar{\mathbf{i}})$ and $\nu(\bar{\mathbf{i}})$ in the same way for the simplified sequence $\bar{\mathbf{i}}$. By construction, we must have

$$n_A + n_B = t(\mathbf{i}) - t(\bar{\mathbf{i}}).$$

One can also check that

$$n_A = \nu(\mathbf{i}) - \nu(\bar{\mathbf{i}}).$$

Substituting these two identities into (E.17) then gives us

$$M(\mathbf{i}) = N_k^{(t(\mathbf{i}) - 2\nu(\mathbf{i}) + 2\nu(\bar{\mathbf{i}}) - t(\bar{\mathbf{i}}))/2} M(\bar{\mathbf{i}}), \quad (\text{E.18})$$

Again, by examining the constructions of the Type-A and Type-B reduction steps, we can conclude that there can only be two possibilities for the final output $\bar{\mathbf{i}}$:

Case 1: $t(\bar{\mathbf{i}}) = \nu(\bar{\mathbf{i}}) = 1$, i.e., $\bar{\mathbf{i}} = (\bar{i}_1)$ is a length 1 cycle. Recall from (E.10) that $M(\bar{\mathbf{i}}) = \sqrt{N_k}$. It then follows from (E.18) that

$$M(\mathbf{i}) = N_k^{(t(\mathbf{i}) - 2\nu(\mathbf{i}) + 2)/2}.$$

By applying this identity to a sequence of length $t(\mathbf{i}) = 2p$ and recalling (E.14), we reach the statement in (E.9).

Case 2: The only other possibility is for $\bar{\mathbf{i}}$ to contain at least two unique indices, i.e., $\nu(\bar{\mathbf{i}}) \geq 2$. Moreover, every unique index must appear at least twice in the cycle, as otherwise the sequence $\bar{\mathbf{i}}$ can be further simplified by a Type-A step. These two conditions imply $t(\bar{\mathbf{i}}) \geq 2\nu(\bar{\mathbf{i}})$, in which case (E.17) gives us

$$|M(\mathbf{i})| \leq N_k^{(t(\mathbf{i}) - 2\nu(\mathbf{i}))/2} |M(\bar{\mathbf{i}})|. \quad (\text{E.19})$$

Moreover, consecutive indices in $\bar{\mathbf{i}} = (\bar{i}_1, \bar{i}_2, \dots, \bar{i}_{t(\bar{\mathbf{i}})})$ cannot have repetitions, i.e., $\bar{i}_j \neq \bar{i}_{j+1}$ for all $j \in [t(\bar{\mathbf{i}})]$ with $j+1$ to be interpreted as modulo $t(\bar{\mathbf{i}})$. (If this were not true, then $\bar{\mathbf{i}}$ would not be the final output as it could be further simplified by a Type-B step.) This condition allows us to apply Hölder’s inequality to get

$$|M(\bar{\mathbf{i}})| = \left| \mathbb{E} Q_{\bar{i}_1 \bar{i}_2} Q_{\bar{i}_2 \bar{i}_3} \dots Q_{\bar{i}_{t(\bar{\mathbf{i}})} \bar{i}_1} \right| \leq \mathbb{E}_{\xi \sim s_d^1} |q_k(\xi)|^{t(\bar{\mathbf{i}})}. \quad (\text{E.20})$$

Now consider a sequence \mathbf{i} with length $t(\mathbf{i}) = 2p$. Since $t(\bar{\mathbf{i}}) \leq 2p$, we can use Hölder’s inequality to further bound the right-hand side as

$$\mathbb{E}_{\xi \sim s_d^1} |q_k(\xi)|^{t(\bar{\mathbf{i}})} \leq [\mathbb{E}_{\xi \sim s_d^1} q_k^{2p}(\xi)]^{t(\bar{\mathbf{i}})/(2p)} \leq \mathbb{E}_{\xi \sim s_d^1} q_k^{2p}(\xi), \quad (\text{E.21})$$

where the last step follows from (E.14), i.e., $\mathbb{E}_{\xi \sim s_d^1} q_k^{2p} \geq 1$. By substituting (E.20) and (E.21) into (E.19), we can verify the bound in (E.9). \square

Lemma 17. For every $k \in \mathbb{N}_0$ and $p \in \mathbb{N}$, we have

$$\mathbb{E} \text{tr}(A_k^{2p}) \leq n(2p)^{2p-2} [p/n + \sum_{i=0}^{p-1} (n/N_k)^i] \cdot \mathbb{E}_{\xi \sim s_d^1} q_k^{2p}(\xi). \quad (\text{E.22})$$

Proof. Using the shorthand notation in (E.8), we can write $A_k(i, j) = \frac{1}{\sqrt{n}} Q_{ij} \cdot (1 - \delta_{ij})$ for $i, j \in [n]$. Let

$$\mathcal{C} := \left\{ \mathbf{i} = (i_1, i_2, \dots, i_{2p}) \in [n]^{2p} : i_j \neq i_{j+1} \text{ for all } j \in [2p-1] \text{ and } i_1 \neq i_{2p} \right\}$$

denote the set of all length $2p$ cycles in which no two consecutive indices are equal. We then have

$$\begin{aligned} \mathbb{E} \operatorname{tr}(A_k^{2p}) &= \frac{1}{n^p} \sum_{\mathbf{i} \in \mathcal{C}} \mathbb{E} Q_{i_1 i_2} Q_{i_2 i_3} \cdots Q_{i_{2p-1} i_{2p}} Q_{i_{2p} i_1} \\ &= \frac{1}{n^p} \left(\sum_{\mathbf{i} \in \mathcal{C}_1} M(\mathbf{i}) + \sum_{\mathbf{i} \in \mathcal{C}_2} M(\mathbf{i}) + \dots + \sum_{\mathbf{i} \in \mathcal{C}_{2p}} M(\mathbf{i}) \right), \end{aligned} \quad (\text{E.23})$$

where, for each $a \in [2p]$,

$$\mathcal{C}_a := \mathcal{C} \cap \left\{ \mathbf{i} \in [n]^{2p} : \nu(\mathbf{i}) = a \right\}$$

and $M(\mathbf{i}) = \mathbb{E} Q_{i_1 i_2} Q_{i_2 i_3} \cdots Q_{i_{2p-1} i_{2p}} Q_{i_{2p} i_1}$. The terms $M(\mathbf{i})$ in (E.23) can be bounded in two different ways. First, recall that, for any $i_j \neq i_{j+1}$, we have $Q_{i_j i_{j+1}} \sim q_k(\xi)$ with $\xi \sim \mathfrak{s}_d^1$. Thus, for every $\mathbf{i} \in \mathcal{C}$, we can apply Hölder's inequality to get

$$|M(\mathbf{i})| \leq \mathbb{E} q_k^{2p}(\xi). \quad (\text{E.24})$$

On the other hand, for each $a \in [2p]$, Lemma 16 gives us

$$|M(\mathbf{i})| \leq \mathbb{E} q_k^{2p}(\xi) N_k^{p-a+1} \quad \text{for all } \mathbf{i} \in \mathcal{C}_a. \quad (\text{E.25})$$

We use (E.24) to bound the $M(\mathbf{i})$ terms in $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_{p+1}$ and use (E.25) for the $M(\mathbf{i})$ terms in $\mathcal{C}_{p+2}, \dots, \mathcal{C}_{2p}$. It then follows from (E.23) that

$$\left| \mathbb{E} \operatorname{tr}(A_k^{2p}) \right| \leq \frac{1}{n^p} \left(\sum_{a=1}^{p+1} |\mathcal{C}_a| + \sum_{a=p+2}^{2p} |\mathcal{C}_a| N_k^{p-a+1} \right) \mathbb{E} q_k^{2p}(\xi), \quad (\text{E.26})$$

where $|\mathcal{C}_a|$ denotes the cardinality of \mathcal{C}_a . For each $a \in [2p]$, we have

$$|\mathcal{C}_a| \leq a^{2p-2} n(n-1) \cdots (n-a+1) \leq (2p)^{2p-2} n^a.$$

This is a crude bound, but it is sufficient for our purpose. Substituting this bound into (E.26) give us

$$\left| \mathbb{E} \operatorname{tr}(A_k^{2p}) \right| \leq (2p)^{2p-2} \left(\sum_{a=1}^{p+1} n^{a-p} + n \sum_{a=p+2}^{2p} (n/N_k)^{a-p-1} \right) \mathbb{E} q_k^{2p}(\xi),$$

which then implies (E.22). \square

E.3 Bounds on the Operator Norms

Proposition 7. *Under the conditions of Theorem 1, we have, for every $k \in \mathbb{N}_0$,*

$$\|A_k\|_{\text{op}} \prec d^{\max\{\ell-k, 0\}/2}. \quad (\text{E.27})$$

Proof. The statement of the proposition means that $\|A_k\|_{\text{op}} \prec 1$ for all the high-order components with $k \geq \ell$. In contrast, for the low-order components, $\|A_k\|_{\text{op}} \prec d^{(\ell-k)/2}$ when $0 \leq k < \ell$.

To show (E.27) for the case of $k \geq \ell$, suppose that we are given an arbitrary $\epsilon > 0$. Substituting (E.22) into (E.5) yields

$$\mathbb{P}(\|A_k\|_{\text{op}} \geq d^\epsilon) \leq d^{-2\epsilon p} n(2p)^{2p-2} \left[p/n + \sum_{i=0}^{p-1} (n/N_k)^i \right] \cdot \mathbb{E}_{\xi \sim \mathfrak{s}_d^1} q_k^{2p}(\xi)$$

Recall that $n = \alpha d^\ell + o(d^\ell)$ and $k \geq \ell$. In addition, we have $N_k = \frac{d^k}{k!}(1 + \mathcal{O}(1/d))$. Thus, for sufficiently large d , we have, by Markov's inequality,

$$\mathbb{P}(\|A_k\|_{\text{op}} \geq d^\epsilon) \leq C(\alpha, k, p)d^{\ell-2\epsilon p},$$

where $C(\alpha, k, p)$ is some constant that depends on α, k, p but not on d . For any $D > 0$, choose a p such that $\ell - 2\epsilon p < -D - 1$. It follows that

$$\mathbb{P}(\|A_k\|_{\text{op}} \geq d^\epsilon) \leq d^{-D}$$

for all sufficiently large d .

Next, we consider the low-order components, with $0 \leq k < \ell$. Lemma 14 implies that

$$\|A\|_{\text{op}} \leq \sqrt{n/N_k} + \mathcal{O}_{\prec}(1) \prec d^{(\ell-k)/2},$$

where the second step uses the fact that $n = \alpha n^\ell + o(n^\ell)$ and $N_k = d^k/k!(1 + \mathcal{O}(1/d))$. \square

F Perturbations of Resolvents and Stieltjes Transforms

In this appendix, we recall some standard properties of the resolvent and the stieltjes transform of Hermitian matrices. Let H be an $n \times n$ Hermitian matrix. Its resolvent is denoted by $G(z) := (H - zI)^{-1}$, where $z \in \mathbb{C}$ is the spectral parameter with $\eta = \Im m[z] > 0$. From the trivial inequality

$$\left| \frac{1}{\lambda - z} \right| \leq \frac{1}{\eta} \quad \text{for all } \lambda \in \mathbb{R},$$

one can easily verify the following bounds on the operator and the entry-wise ℓ_∞ norms of $G(z)$:

$$\|G(z)\|_{\text{op}} \leq \frac{1}{\eta} \tag{F.1}$$

$$\|G(z)\|_{\infty} \leq \frac{1}{\eta}. \tag{F.2}$$

In our proof, we will also need the *Ward identity* [16, Lemma 8.3]:

$$\sum_{ij} |G_{ij}(z)|^2 = \frac{\Im m(\text{tr } G(z))}{\eta}. \tag{F.3}$$

Let $s(z)$ be the Stieltjes transform of the empirical spectral distribution of H . Since $s(z) = \frac{1}{n} \text{tr } G(z)$, (F.2) immediately implies the pointwise bound

$$|s(z)| \leq \frac{1}{\eta}. \tag{F.4}$$

One can also easily check that

$$|s(z_1) - s(z_2)| \leq \frac{|z_1 - z_2|}{\Im m(z_1)\Im m(z_2)}. \tag{F.5}$$

Both the resolvent and the Stieltjes transform are stable with respect to matrix perturbations. In our proof of Theorem 1, we will use the following standard perturbation estimates.

Lemma 18. *Let H_1, H_2 be $n \times n$ Hermitian matrices, and $G_1(z), G_2(z)$ their resolvents. Then,*

$$G_1(z) = G_2(z) - G_2(z)(H_1 - H_2)G_1(z). \tag{F.6}$$

Moreover, for the Stieltjes transforms of H_1 and H_2 , we have

$$|s_1(z) - s_2(z)| \leq \frac{\|H_1 - H_2\|_{\text{F}}}{\sqrt{n} \eta^2} \tag{F.7}$$

and

$$|s_1(z) - s_2(z)| \leq \frac{C \text{rank}(H_1 - H_2)}{n \cdot \eta}, \tag{F.8}$$

where C is an absolute constant.

Proof. The formula in (F.6) can be easily verified by using the identities $I = (H_1 - zI)G_1 = (H_2 - zI + H_1 - H_2)G_1$ and $G_2(H_2 - zI) = I$. Applying (F.6) gives us

$$\begin{aligned} |s_1(z) - s_2(z)| &= \frac{1}{n} |\operatorname{tr}[G_2(z)(H_1 - H_2)G_1(z)]| \\ &= \frac{1}{n} |\operatorname{tr}[G_1(z)G_2(z)(H_1 - H_2)]| \\ &\leq \frac{1}{n} \|G_1(z)G_2(z)\|_{\mathbb{F}} \|H_1 - H_2\|_{\mathbb{F}}, \end{aligned} \quad (\text{F.9})$$

where the last step uses the Cauchy-Schwarz inequality. By using (F.1) in the last step, we have

$$\|G_1(z)G_2(z)\|_{\mathbb{F}} \leq \sqrt{n} \|G_1(z)G_2(z)\|_{\text{op}} \leq \sqrt{n} \|G_1(z)\|_{\text{op}} \|G_2(z)\|_{\text{op}} \leq \frac{\sqrt{n}}{\eta^2}.$$

Substituting this bound into (F.9) then leads to (F.7).

Next, we show (F.8). First consider the special case when $\operatorname{rank}(H_1 - H_2) = 1$. As the eigenvalues are invariant under unitary transforms, we can assume without loss of generality that

$$H_1 - H_2 = a e_1 e_1^{\top}, \quad (\text{F.10})$$

for some $a \neq 0$. Let $H_1^{[1]}$ (resp. $H_2^{[1]}$) denote the $(n-1) \times (n-1)$ minor matrix obtained by removing the first column and row of H_1 (resp. H_2). By (F.10), $H_1^{[1]} = H_2^{[1]}$. Let $s^{[1]}(z)$ denote the Stieltjes transform of the eigenvalues of $H_1^{[1]}$. We have

$$|s_1(z) - s_2(z)| \leq |s_1(z) - s^{[1]}(z)| + |s^{[1]}(z) - s_2(z)| \leq \frac{C}{n \cdot \eta},$$

where the second step uses [16, Lemma 7.5], and C is an absolute constant. For the case when $\operatorname{rank}(H_1 - H_2) > 1$, we can always write $s_1(z) - s_2(z)$ as a telescoping sum of rank-one perturbations. The inequality in (F.8) can then be obtained by applying the triangular inequality and by using the result for the rank-one case. \square

Proof of Lemma 2. By using the factorized representation of A_k in (A.7), we can write the low-order term as

$$A_{\text{LO}} = \sum_{0 \leq k < \ell} \frac{\mu_k}{\sqrt{n N_k}} S_k^{\top} S_k - \sigma_d I,$$

where S_k is an $N_k \times n$ matrix, with $N_k = \mathcal{O}(d^k)$, and $\sigma_d := \sum_{0 \leq k < \ell} \mu_k \sqrt{N_k/n}$. Since $n \asymp d^\ell$, we have $\sigma_d = \mathcal{O}(d^{-1/2})$ and that

$$\sigma_d \operatorname{rank}(A_{\text{LO}} + \sigma_d I) \leq \sum_{0 \leq k < \ell} N_k = \mathcal{O}(d^{\ell-1}). \quad (\text{F.11})$$

Write $\check{A} := A + \sigma_d I = (A_{\text{LO}} + \sigma_d I) + A_{\text{HI}}$, whose Stieltjes transform is denoted by $\check{s}(z)$. By the triangular inequality,

$$\begin{aligned} |s(z) - \hat{s}(z)| &\leq |s(z) - \check{s}(z)| + |\check{s}(z) - \hat{s}(z)| \\ &\leq \frac{\sigma_d}{\eta^2} + \frac{C \operatorname{rank}(\check{A} - A_{\text{HI}})}{n \cdot \eta}, \end{aligned} \quad (\text{F.12})$$

where the last step follows from the observation that $s(z) = \check{s}(z + c)$ and (F.5), and from the perturbation inequality in (F.8). Substituting the estimates $\sigma_d = \mathcal{O}(d^{-1/2})$ and (F.11) into (F.12), we complete the proof. \square

G Stability Analysis of the Self-Consistent Equation

In this appendix, we present a stability analysis of the self-consistent equation in (2.10) satisfied by the limiting Stieltjes transform. We start by rewriting (2.10) as

$$\frac{1}{m} = -z - \frac{\gamma_a m}{1 + \gamma_b m} - \gamma_c m. \quad (\text{G.1})$$

Now suppose $s \in C_+$ is an approximate solution to (G.1) such that

$$\frac{1}{s} = -z - \frac{\gamma_a s}{1 + \gamma_b s} - \gamma_c s + \omega, \quad (\text{G.2})$$

with some error term ω . We show in the following proposition that $s \approx m$ as long as the error ω is small.

Proposition 8. *For any $z \in C_+$, there exists a unique solution $m \in C_+$ to (G.1). Moreover, let $s \in C_+$ be an approximate solution to (2.10) in the sense of (G.2), with the error term satisfying*

$$|\omega| \leq \frac{\eta}{2}, \quad \eta = \Im m[z]. \quad (\text{G.3})$$

We have

$$|s - m| \leq \frac{4|\omega|}{\eta^2}. \quad (\text{G.4})$$

Proof. The existence and uniqueness of the solution $m \in C_+$ is a well-known result. One way to establish this is to observe that (G.1) is just the free additive convolution of the Marchenko-Pastur law and the semicircle law. See, e.g., [4] for more details. An alternative approach, as used in [9, Lemma A.1], is to rewrite (G.1) as a third-degree polynomial in m and to analyze the roots of that polynomial.

Next, we establish the stability bound in (G.4). Recall the family of MP density functions given in (C.4), with two shape parameters ϕ and t . In what follows, we set $\phi = \gamma_a/\gamma_b^2$ and $t = \mu_\ell/\sqrt{\phi}$, and write $\varrho_{\text{MP}} := \varrho_{\phi,t}$. The Stieltjes transform of ϱ_{MP} is specified as the unique solution to the following (G.6). Substituting our particular choices of ϕ and t into (G.6), and by the uniqueness of the solution, we obtain the following characterization: for any $z, \Gamma \in C_+$,

$$\frac{1}{\Gamma} = -z - \frac{\gamma_a \Gamma}{1 + \gamma_b \Gamma} \quad \Leftrightarrow \quad \Gamma = \int \frac{\varrho_{\text{MP}}(dx)}{x - z}. \quad (\text{G.5})$$

Using (G.5) and (G.1), we can now write the exact solution m as

$$m = \int \frac{\varrho_{\text{MP}}(dx)}{x - z - \gamma_c m}. \quad (\text{G.6})$$

By the assumption in (G.3), $\Im m[z + \gamma_c s - \omega] \geq \eta/2$. Thus, it follows from (G.5) and (G.2) that

$$s = \int \frac{\varrho_{\text{MP}}(dx)}{x - z - \gamma_c s + \omega}. \quad (\text{G.7})$$

Computing the imaginary part of the equation for m , we get

$$\Im m[m] = \int \frac{\eta + \gamma_c \Im m[m]}{|x - z - \gamma_c m|^2} \varrho_{\text{MP}}(dx), \quad (\text{G.8})$$

which also gives us the trivial bound $\Im m[m] \leq 1/\eta$. Similarly,

$$\Im m[s] = \int \frac{\eta + \gamma_c \Im m[s] - \Im m[\omega]}{|x - z - \gamma_c s + \omega|^2} \varrho_{\text{MP}}(dx) \quad (\text{G.9})$$

$$\geq \int \frac{\eta/2 + \gamma_c \Im m[s]}{|x - z - \gamma_c s + \omega|^2} \varrho_{\text{MP}}(dx), \quad (\text{G.10})$$

where the second step uses (G.3). Note that (G.9) and (G.3) also imply that $\Im m[s] \leq 2/\eta$. Taking the difference of (G.6) and (G.7) gives us

$$(s - m) \left[1 - \int \frac{\gamma_c \varrho_{\text{MP}}(dx)}{(x - z - \gamma_c m)(x - z - \gamma_c s + \omega)} \right] = \int \frac{-\omega \varrho_{\text{MP}}(dx)}{(x - z - \gamma_c m)(x - z - \gamma_c s + \omega)}. \quad (\text{G.11})$$

By the Cauchy-Schwarz inequality in the first step, and by (G.8), (G.10) in the second step,

$$\begin{aligned} \left| \int \frac{\varrho_{\text{MP}}(dx)}{(x - z - \gamma_c m)(x - z - \gamma_c s + \omega)} \right| &\leq \left[\int \frac{\varrho_{\text{MP}}(dx)}{|x - z - \gamma_c m|^2} \int \frac{\varrho_{\text{MP}}(dx)}{|x - z - \gamma_c s + \omega|^2} \right]^{1/2} \\ &\leq \left[\frac{\Im m[m]}{\eta + \gamma_c \Im m[m]} \cdot \frac{\Im m[s]}{\eta/2 + \gamma_c \Im m[s]} \right]^{1/2} \\ &\leq \frac{1}{\eta^2/4 + \gamma_c}, \end{aligned} \quad (\text{G.12})$$

where the last step uses the bound that $\max \{ \Im m[m], \Im m[s] \} \leq 2/\eta$. This allows us to get

$$\left| 1 - \int \frac{\gamma_c \varrho_{\text{MP}}(dx)}{(x - z - m)(x - z - s + \omega)} \right| \geq 1 - \frac{\gamma_c}{\eta^2/4 + \gamma_c} = \frac{\eta^2}{\eta^2 + 4\gamma_c}. \quad (\text{G.13})$$

Substituting (G.12) and (G.13) into (G.11), we then reach the statement (G.4) of the proposition. \square

References

- [1] E. Artin. *The Gamma Function*. Dover Books on Mathematics. Dover Publications, Inc, Mineola, New York, dover edition edition, 2015. ISBN 978-0-486-78978-1.
- [2] Z. Bai and J. W. Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*. Springer Series in Statistics. Springer New York, New York, NY, 2010.
- [3] M. Belkin and P. Niyogi. Laplacian Eigenmaps for Dimensionality Reduction and Data Representation. *Neural Computation*, 15(6):1373–1396, June 2003.
- [4] P. Biane. On the free convolution with a semi-circular distribution. *Indiana University Mathematics Journal*, 46(3):0–0, 1997. ISSN 0022-2518.
- [5] P. J. Bickel and E. Levina. Covariance regularization by thresholding. *The Annals of Statistics*, 36(6):2577–2604, 2008.
- [6] A. Bloemendal, A. Knowles, H.-T. Yau, and J. Yin. On the principal components of sample covariance matrices. *arXiv:1404.0788 [math-ph, stat]*, Apr. 2014.
- [7] A. Bloemendal, L. Erdos, A. Knowles, H.-T. Yau, and J. Yin. Isotropic Local Laws for Sample Covariance and Generalized Wigner Matrices. *arXiv:1308.5729 [math-ph]*, July 2015.
- [8] B. Bordenon, A. Canatar, and C. Pehlevan. Spectrum Dependent Learning Curves in Kernel Regression and Wide Neural Networks. *arXiv:2002.02561 [cs, stat]*, Feb. 2021.
- [9] X. Cheng and A. Singer. The spectrum of random inner-product kernel matrices. *Random Matrices: Theory and Applications*, 02(04), Oct. 2013.
- [10] J. Coster and H. B. Hart. A simple proof of the addition theorem for spherical harmonics. *American Journal of Physics*, 59(4):371–373, Apr. 1991.
- [11] F. Dai and Y. Xu. *Approximation Theory and Harmonic Analysis on Spheres and Balls*. Springer Monographs in Mathematics. Springer New York, New York, NY, 2013.

- [12] Y. Deshpande and A. Montanari. Sparse pca via covariance thresholding. In *Advances in Neural Information Processing Systems*, pages 334–342, 2014.
- [13] Y. Do and V. Vu. The spectrum of random kernel matrices: Universality results for rough and varying kernels. *arXiv:1206.3763 [math]*, May 2013.
- [14] C. Efthimiou and C. Frye. *Spherical Harmonics in p Dimensions*. World Scientific, New Jersey, 2014. ISBN 978-981-4596-69-5.
- [15] N. El Karoui. The spectrum of kernel random matrices. *The Annals of Statistics*, 38(1):1–50, Feb. 2010. ISSN 0090-5364. doi: 10.1214/08-AOS648.
- [16] L. Erdős and H.-T. Yau. *Dynamical Approach to Random Matrix Theory*. The AMS and the Courant Institute of Mathematical Sciences at New York University, 2017.
- [17] L. Erdos, A. Knowles, H.-T. Yau, and J. Yin. The Local Semicircle Law for a General Class of Random Matrices. *arXiv:1212.0164 [math-ph]*, May 2013.
- [18] Z. Fan and A. Montanari. The Spectral Norm of Random Inner-Product Kernel Matrices. *arXiv:1507.05343 [math, stat]*, Feb. 2017.
- [19] B. Ghorbani, S. Mei, T. Misiakiewicz, and A. Montanari. Linearized two-layers neural networks in high dimension. *arXiv:1904.12191 [cs, math, stat]*, Feb. 2020.
- [20] T. Hastie, A. Montanari, S. Rosset, and R. J. Tibshirani. Surprises in High-Dimensional Ridgeless Least Squares Interpolation. *arXiv:1903.08560 [cs, math, stat]*, Dec. 2020.
- [21] H. Hu and Y. M. Lu. Universality Laws for High-Dimensional Learning with Random Features. *arXiv:2009.07669 [cs, math]*, Mar. 2021.
- [22] M. E. H. Ismail. *Classical and Quantum Orthogonal Polynomials in One Variable*. Cambridge University Press, first edition, Nov. 2005.
- [23] G. J. O. Jameson. A simple proof of Stirling’s formula for the gamma function. *The Mathematical Gazette*, 99(544):68–74, Mar. 2015.
- [24] A. Knowles and J. Yin. Anisotropic local laws for random matrices. *arXiv:1410.3516 [math-ph]*, Aug. 2016.
- [25] V. Koltchinskii and E. Giné. Random Matrix Approximation of Spectra of Integral Operators. *Bernoulli*, 6(1):113, Feb. 2000. ISSN 13507265. doi: 10.2307/3318636.
- [26] C. Louart, Z. Liao, and R. Couillet. A Random Matrix Approach to Neural Networks. *arXiv:1702.05419 [cs, math]*, June 2017.
- [27] S. Mei and A. Montanari. The generalization error of random features regression: Precise asymptotics and double descent curve. *arXiv:1908.05355 [math, stat]*, Oct. 2019.
- [28] T. Misiakiewicz. Spectrum of inner-product kernel matrices in the polynomial regime and multiple descent phenomenon in kernel ridge regression. *arXiv:2204.10425 [math, stat]*, Apr. 2022.
- [29] A. Mnih and R. R. Salakhutdinov. Probabilistic Matrix Factorization. In J. Platt, D. Koller, Y. Singer, and S. Roweis, editors, *Advances in Neural Information Processing Systems*, volume 29, 2007.
- [30] M. Opper and R. Urbanczik. Support vector machines learning noisy polynomial rules. *Physica A*, page 9, 2001.
- [31] A. Rahimi and B. Recht. Random Features for Large-Scale Kernel Machines. In *Proceedings of the 20th International Conference on Neural Information Processing Systems*, page 8, 2007.

- [32] B. Schölkopf and A. J. Smola. *Learning with Kernels: Support Vector Machines, Regularization, Optimization, and Beyond*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, Mass, 2002.
- [33] B. Schölkopf, A. Smola, and K.-R. Müller. Nonlinear Component Analysis as a Kernel Eigenvalue Problem. *Neural Computation*, 10(5):1299–1319, July 1998. ISSN 0899-7667, 1530-888X.
- [34] J. A. Tropp. An Introduction to Matrix Concentration Inequalities. *arXiv:1501.01571 [cs, math, stat]*, Jan. 2015.