

REVERSING MONOID ACTIONS AND DOMINATION IN GRAPHS

MEHMET AKIF ERDAL

ABSTRACT. Given a graph $G = (V, E)$, a set of vertices $D \subseteq V$ is called a dominating set if every vertex in $V \setminus D$ is adjacent to a vertex in D , and a subset $B \subseteq V$ is called a nonblocking set if $V - B$ is a dominating set. In this paper, we introduce a graph dynamical systems detecting vertex sets that are simultaneously dominating and nonblocking sets via reversing the action of the system. Moreover, by using actions of multiple such graph dynamical systems we define actions of free monoid on two letters for which elements in the reverse action corresponds to more special dominating sets.

1. INTRODUCTION

The notion of graph dynamical system was first introduced in [2]. Later many applications on modeling and simulation of complex systems, such as biological and social networks, have been studied, see e.g. [6, 3, 12]. For a given graph $G = (V, E)$, a graph dynamical system on G consists of a set S of states and an update function $F : S^V \rightarrow S^V$ on the set of functions from V to S , where F is defined via a local update functions $f_v : S^{N[v]} \rightarrow S$. Here $N[v]$ denotes the closed neighborhood of $v \in V$. The map F induces an action of the monoid \mathbb{N} of natural numbers as $n \cdot s = F^{\circ n}(s)$, where $F^{\circ n}$ denotes the n -fold composition of F by itself. Besides, if we are given a collection $\{F_i\}_{i \in K}$ of graph dynamical systems on the same graph with the same state set, then we can induce an action of the free monoid on K .

In [5] a functorial construction Inv was introduced for actions of monoids on finite sets. For a finite set A on which a monoid M acts, $\text{Inv}(A)$ is given by

$$\text{Inv}(A) = \{\sigma : M \rightarrow A \mid m \cdot \sigma(mn) = \sigma(m), \forall n, m \in M\}.$$

The set $\text{Inv}(A)$ admits a right M action defined by $(\sigma \cdot m)(m_0) = \sigma(mm_0)$ for every $m, m_0 \in M$. There is an evaluation function $\epsilon : \text{Inv}(A) \rightarrow A$ given by $\epsilon(\sigma) = \sigma(1)$, where $1 \in M$ is the identity. This map is injective and its image $\text{Im}(\epsilon)$ corresponds to the maximal invariant subset of A on which M acts by bijections.

For a given connected finite simple graph $G = (V, E)$, we present an application of graph dynamical systems, together with reverse action as given in [5], on locating special dominating sets. Relations between dominating sets of some graphs and graph dynamical systems previously established in [1, Thm. 3.5, Cor. 4.7], for which the number of fixed points in a certain parallel sequential dynamical system and the number of a special kind of dominating sets of a bipartite graph coincide.

In this paper, for an arbitrary finite simple connected graph $G = (V, E)$, we construct a graph dynamical system $F : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ by $F(s) = sP_s + P_s + s + 1$ where $P_s(v) = \prod_{x \in N(v)} (1 + s(x))$ for $v \in V$. This system has no fixed point

Key words and phrases. Semigroup action, Reverse action, Graph Dynamical System, Dominating set.

other than the constant function c_0 at 0 and its periodic points are those non-zero elements in $\text{Im}(\epsilon)$. We prove the following theorem.

Theorem 1.1. *Let $G = (V, E)$ be a graph and $F : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ be the graph dynamical system given by $F(s) = sP_s + P_s + s + 1$ for $s \in \mathbb{Z}_2^V$. Let $\epsilon : \text{Inv}(A) \rightarrow A$ be the evaluation map given by $\epsilon(s) = s(0)$. For a surjection $s \in \mathbb{Z}_2^V$, we have $s \in \text{Im}(\epsilon)$ if and only if both $s^{-1}(0)$ and $s^{-1}(1)$ are dominating sets of G . Moreover, in this case $F \circ F(s) = s$.*

The theorem establishes a one to one correspondence between the non-zero elements in $\text{Im}(\epsilon)$ and the subsets of V that are simultaneously dominating and nonblocking sets. It is known that complement of a minimal dominating set is also a dominating set, see [11]. Hence, in particular, all minimal dominating sets and maximal independent sets can be obtained as $s^{-1}(1)$ or $s^{-1}(0)$ for some s in $\text{Im}(\epsilon)$.

In Section 3.1, we extend the action of the system in Theorem 3.4 to actions of the free monoid on two letters, and establish one to one correspondences between more special dominating sets of G as fixed points of these actions, see Corollary 3.8 and Proposition 3.9.

2. A FEW PRELIMINARIES

Throughout this paper $G = (V, E)$ denotes an undirected finite simple connected graph with vertex set V and edge set E .

2.1. Dominating and nonblocking sets. Dominating sets of graphs and related problems are well studied in the literature. For general references about classical and recent advances we refer to [9] and [8]. A dominating set for G is a subset A of V for which every vertex that is not in A is adjacent to a vertex in A . A nonblocking set (also known as nonblocker) for G is a subset B of V such that every vertex in B is adjacent to vertices outside of the subset. By definition B is a nonblocking set if and only if $V \setminus B$ is a dominating set, see [4]. When a graph is given there are special subsets of the set of vertices that are simultaneously dominating and nonblocking sets. For example any minimal dominating set; that is, a dominating set that does not properly contains any other dominating set, is also a nonblocking set. In fact, due to [11, 13.1.14], if $D \subset V$ is a minimal dominating set of D then $V \setminus D$ is also a dominating set.

2.2. Graph dynamical systems. For a vertex $v \in V$ the open neighborhood of v is given by

$$N(v) = \{x \in V \mid \{x, v\} \in E\}$$

and the closed neighborhood of v is $N[v] = N(v) \cup \{v\}$. A graph dynamical system consists of a graph $G = (V, E)$, a finite state set S and an update map $F : S^V \rightarrow S^V$ where F is defined via a local update function $f_v : S^{N[v]} \rightarrow S$. If F satisfies $F(s)(v) = f_v(s|_{N[v]})$ for each v in V , then such a system is referred as generalized cellular automata (see [13]). In this paper, we are interested in a special case where $S = \mathbb{Z}_2$ the field with 2 elements $\{0, 1\}$.

2.3. Reversing monoid actions. A central construction we use in this paper is the reverse action of a monoid action. Let M be a monoid acting on a finite set A . We define the reverse action on A as follows

$$\text{Inv}(A) = \{\sigma : M \rightarrow A \mid \forall n, m \in M, m \cdot \sigma(nm) = \sigma(n)\}.$$

The set $\text{Inv}(A)$ admits a right M action given by $(\sigma \cdot m_0)(m) = \sigma(m_0 \cdot m)$. Note that

$$m \cdot (\sigma \cdot m_0)(nm) = m_0 \cdot \sigma(m_0nm) = \sigma(m_0n) = (\sigma \cdot m_0)(n);$$

i.e., $\sigma \cdot m_0 \in \text{Inv}(A)$ and the action is well defined. This is a special case of the construction Inv_r^l in [5] for left actions. There is an associated evaluation function $\epsilon : \text{Inv}(A) \rightarrow A$ given by $\epsilon(\sigma) = \sigma(1)$. This function satisfies

$$m \cdot \epsilon(\sigma \cdot m) = m \cdot ((\sigma \cdot m)(1)) = m \cdot \sigma(m) = \sigma(1) = \epsilon(\sigma).$$

Let $\mu_m : A \rightarrow A$ is the function given by $\mu_m(a) = m \cdot a$. We say M acts on A by bijections if for every $m \in M$, μ_m is a bijection.

The following proposition and the subsequent lemma follow from Proposition 5 of Section 4.2 and Lemma 3 of Section 4.4 in [5], respectively. However, as we do not consider two sided actions here, we prove both statements for completeness and notational convenience.

Proposition 2.1. *For a finite M -set A , M acts on A by bijections if and only if $\epsilon : \text{Inv}(A) \rightarrow A$ is a bijection.*

Proof. Assume that μ_m is a bijection for every $m \in M$. For each $a \in A$, define $\tau(a) : M \rightarrow A$ by $\tau(a)(m) = \mu_m^{-1}(a)$. Then

$$m \cdot \tau(a)(nm) = \mu_m(\mu_{nm})^{-1}(a) = \mu_m \circ \mu_m^{-1} \circ \mu_n^{-1}(a) = \mu_n^{-1}(a) = \tau(a)(n);$$

i.e., $\tau(a) \in \text{Inv}(A)$ and τ defines a function from A to $\text{Inv}(A)$. Besides, for every $a \in A$, $\epsilon \circ \tau(a) = \tau(a)(1) = a$ and for every $\sigma \in \text{Inv}(A)$,

$$(\tau \circ \epsilon)(\sigma)(m) = \tau(\sigma(1))(m) = \mu_m^{-1}(\sigma(1)) = \mu_m^{-1}(m \cdot \sigma(m)) = \sigma(m).$$

Thus, ϵ is a bijection with inverse τ . Conversely, assume ϵ is a bijection with inverse τ . For each $m \in M$, define $\nu_m : A \rightarrow A$ by $\nu_m(a) = \tau(a)(m)$ for $a \in A$. Then,

$$\mu_m \circ \nu_m(a) = m \cdot \tau(a)(m) = \tau(a)(1) = \epsilon(\tau(a)) = a;$$

i.e., μ_m admits a section. But since A is finite, this means μ_m is a bijection. \square

Lemma 2.2. *Let A be a finite M -set. Then $\epsilon : \text{Inv}(A) \rightarrow A$ is injective with image*

$$\text{Im}(\epsilon) = \{a \in A \mid \forall m \in M, \mu_m|_{Ma} : Ma \rightarrow Ma \text{ is bijection}\},$$

where $Ma = \{m \cdot a \mid m \in M\}$.

Proof. Let $\sigma, \tau \in \text{Inv}(A)$ with $\sigma(1) = \tau(1)$ and let $m \in M$. Then, as A is finite, there exist positive natural numbers s and t with $\sigma(m^s) = \sigma(1)$ and $\tau(m^t) = \tau(1)$. Assume that t and s are smallest such natural numbers and without loss of generality assume $t \leq s$. Then $s = t$, as otherwise we have

$$\sigma(1) = \tau(1) = m^t \cdot \tau(m^t) = m^t \cdot \sigma(m^s) = m^t \cdot \sigma(m^{s-t}m^t) = \sigma(m^{s-t})$$

which gives $\sigma(m^{s-t}) = \sigma(1)$, contradicting to the minimality of s . Hence for every $m \in M$ there exists a positive integer t with $\tau(m^t) = \sigma(m^t)$. This implies

$$m^{t-1} \cdot \tau(m^t) = m^{t-1} \cdot \sigma(m^t)$$

and so $\tau(m) = \sigma(m)$. Therefore, we have $\tau = \sigma$; i.e., ϵ is injective. Now, if $a \in \text{Im}(\epsilon)$; that is, $a = \sigma(1)$ for some $\sigma \in \text{Inv}(A)$, then $m \cdot ((\sigma \cdot m)(1)) = \sigma(1)$. Thus, $\mu_m|_{\text{Im}(\epsilon)}$ is surjective; and as A is finite, we have $\mu_m|_{\text{Im}(\epsilon)}$ is bijective. In particular, $\mu_m|_{M\sigma(1)}$ is bijective. Conversely, if $a \in A$ such that $\mu_m|_{Ma} : Ma \rightarrow Ma$ is bijective for every $m \in M$, we can define $\tau(a) : M \rightarrow A$ by $\tau(a)(m) = \mu_m|_{Ma}^{-1}(a)$.

Then, as in the proof of proposition above, $\tau(a) \in \text{Inv}(A)$ and $\tau(a)(1) = a$; i.e., $a \in \text{Im}(\epsilon)$. \square

The lemma above, combined with the previous proposition, implies that $\text{Inv}(A)$ is finite and every element of M acts on each orbit by bijections. Besides, every such orbit is in the image of ϵ . Thus, as A is finite, the action on $\text{Inv}(A)$ is reversible; that is, every element of M acts by bijections on $\text{Inv}(A)$. Therefore, $\text{Im}(\epsilon)$ is the maximal invariant subset of A on which M acts by bijections; i.e., $\mu_m|_{\text{Im}(\epsilon)} : \text{Im}(\epsilon) \rightarrow \text{Im}(\epsilon)$ is a bijection for every $m \in M$. This, in particular, implies $F(\text{Im}(\epsilon)) = \text{Im}(\epsilon)$.

We use the construction Inv , first in the case when $M = \mathbb{N}$, the monoid of natural numbers, and then $M = \mathcal{F}(a, b)$, the free monoid on two letters a and b . The latter will be denoted by Inv_M for convenience. Note that in the case when $M = \mathbb{N}$, the set $\text{Im}(\epsilon)$ consists of all fixed points and periodic points of the action.

3. MAIN RESULTS

For any set X denote by \mathbb{Z}_2^X the set of functions from X to \mathbb{Z}_2 . The addition and multiplication of elements in \mathbb{Z}_2^X is defined point-wise. For each $s : V \rightarrow \mathbb{Z}_2$ let $P_s : V \rightarrow \mathbb{Z}_2$ be the function given by

$$P_s(v) = \prod_{x \in N(v)} (1 + s(x)).$$

We define $F : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ as

$$F(s) = sP_s + P_s + s + 1$$

for $s \in \mathbb{Z}_2^V$. The function F induces a generalized cellular automata with local functions $f_v : \mathbb{Z}_2^{N[v]} \rightarrow \mathbb{Z}_2$ given by

$$f_v(s) = 1 + s(v) + \prod_{x \in N(v)} (1 - s(x)) + (s(v) \prod_{x \in N(v)} (1 - s(x))).$$

We can consider this system as an \mathbb{N} -action on \mathbb{Z}_2^V , which is defined by $n \cdot s = F^{\circ n}(s)$, where $F^{\circ n}$ denotes the n -fold composition of F and $F^{\circ 0}$ is the identity map.

Lemma 3.1. *For any $s \in \mathbb{Z}_2^V$,*

$$F(s)^{-1}(1) = \{v \in s^{-1}(0) \mid \exists u \in s^{-1}(1), \{u, v\} \in E\}.$$

Proof. Note that $F(s)(v) = 1$ if and only if $s(v) = 0 = \prod_{x \in N(v)} (1 + s(x))$. Assume that $v \in F(s)^{-1}(1)$. Then $s(v) = 0$ and $\prod_{x \in N(v)} (1 + s(x)) = 0$. But this means there is $u \in N(v)$ with $s(u) = 1$. Therefore, $u \in s^{-1}(1)$ and $\{u, v\} \in E$. Conversely, if $s(v) = 0$ and $\exists u \in s^{-1}(1)$ such that $\{u, v\} \in E$, then $u \in N(v)$ and $1 + s(u) = 0$, which implies $\prod_{x \in N(v)} (1 + s(x)) = 0$. \square

Let $\text{Surj}(\mathbb{Z}_2^V) = \mathbb{Z}_2^V \setminus \{c_0, c_1\}$ where c_i is the constant function at $i \in \{0, 1\}$; i.e., the subset of \mathbb{Z}_2^V of all surjections from V to \mathbb{Z}_2 . A subset A of an \mathbb{N} -set is called forward invariant if for every $n \in \mathbb{N}$, $n \cdot A \subseteq A$. We have $F(c_0) = c_0$ and $F(c_1) = c_0$, so that $\{c_0, c_1\}$ is forward invariant under the \mathbb{N} -action. We also have the following proposition.

Proposition 3.2. *$\text{Surj}(\mathbb{Z}_2^V)$ is forward invariant under the \mathbb{N} -action.*

Proof. Let $s \in \text{Surj}(\mathbb{Z}_2^V)$ such that $F(s) \in \{c_0, c_1\}$. Recall that $F(s)(v) = 1$ if and only if $s(v) = 0 = \prod_{x \in N(v)} (1 + s(x))$. Since s is not constant, we have $F(s) \neq c_1$. Assume $F(s) = c_0$. Then, by lemma 3.1, $\emptyset = F(s)^{-1}(1) = \{v \in s^{-1}(0) \mid \exists u \in s^{-1}(1), \{u, v\} \in E\}$. But as G is connected, there exist two adjacent vertices $v, w \in V$ such that $s(v) = 1$ and $s(w) = 0$, so a contradiction. Thus, $F(s)$ does not belong to $\{c_0, c_1\}$; or equivalently, F maps surjections in \mathbb{Z}_2^V to surjections. Therefore, $F(\text{Surj}(\mathbb{Z}_2^V)) \subseteq \text{Surj}(\mathbb{Z}_2^V)$. \square

Proposition 3.3. *The only fixed point of F is c_0 . In particular, $\text{Surj}(\mathbb{Z}_2^V)$ does not have any fixed points under F .*

Proof. Assume for some $s \in \mathbb{Z}_2^V$ we have $F(s) = sP_s + s + P_s + 1 = s$. If $s(v) = 1$ for some $v \in V$, then

$$s(v)P_s(v) + P_s(v) + 1 = 2P_s(v) + 1 = 0,$$

which is a contradiction. Thus $s(v) = 0$ for every $v \in V$. \square

Now we state and prove our main result.

Theorem 3.4. *Let $G = (V, E)$ be a graph and $F : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ be the graph dynamical system given by $F(s) = sP_s + P_s + s + 1$ where $P_s(v) = \prod_{x \in N(v)} (1 + s(x))$ for $v \in V$. Let $\epsilon : \text{Inv}(A) \rightarrow A$ be the evaluation map given by $\epsilon(s) = s(0)$. For a surjection $s \in \mathbb{Z}_2^V$, we have $s \in \text{Im}(\epsilon)$ if and only if both $s^{-1}(0)$ and $s^{-1}(1)$ are dominating sets of G . Moreover, in this case $F \circ F(s) = s$.*

Proof. Suppose that both $s^{-1}(0)$ and $s^{-1}(1)$ are dominating sets. Then for every $v \in s^{-1}(0)$ there exists $u \in s^{-1}(1)$ such that $\{u, v\} \in E$. By lemma 3.1, $F(s)^{-1}(1) = s^{-1}(0)$. Similarly, $F(s)^{-1}(0) = s^{-1}(1)$. This means $F(s) = 1 + s$ and $F(1 + s) = s$ (where $(1 + s)(x) = 1 + s(x)$ for $x \in V$). Thus, $\mathbb{N}_s = \{s, 1 + s\}$ and $F|_{\mathbb{N}_s}$ is a bijection.

Conversely, assume that at least one of $s^{-1}(0)$ and $s^{-1}(1)$ is not a dominating set. First assume $s^{-1}(1)$ is not a dominating set. Then

$$F(s)^{-1}(1) = \{v \in s^{-1}(0) \mid \exists u \in s^{-1}(1), \{u, v\} \in E\} \neq s^{-1}(0).$$

Then there exists a function $\tilde{s} : V \rightarrow \mathbb{Z}_2$ satisfying


$$\tilde{s}(x) = \begin{cases} 1 & \text{if } s(x) = 1 \text{ or } F(s)(x) = 0 \\ 0 & \text{if } F(s)(x) = 1 \end{cases}.$$

Since $F(s)^{-1}(1) \subsetneq s^{-1}(0)$, there exist $v \in V$ with $s(v) = 0 = F(s)(v)$; i.e., $\tilde{s} \neq s$. By definition of \tilde{s} , we have $F(\tilde{s})(v) = 1$ if and only if $F(s)(x) = 1$; i.e., $F(\tilde{s}) = F(s)$. But this implies $s \notin \text{Im}(\epsilon)$ (otherwise the restriction of μ_1 on $\text{Im}(\epsilon)$ can not be invertible). Now, assume that $s^{-1}(1)$ is a dominating set but $s^{-1}(0)$ is not. Then $F(s)^{-1}(1) = s^{-1}(0)$, which means $F(s) = 1 + s \notin \text{Im}(\epsilon)$. But this means $F(s) \notin \text{Im}(\epsilon)$ which gives $s \notin \text{Im}(\epsilon)$. \square

Observe that the theorem implies if $s : V \rightarrow \mathbb{Z}_2$ is a surjective function, then $F^{\text{on}}(s)$ is a surjective element in $\text{Im}(\epsilon)$ for sufficiently large n ; i.e., any element in $\text{Surj}(\mathbb{Z}_2^V)$ converges to a surjective function $V \rightarrow \mathbb{Z}_2$ for which pre-images of 0 and 1 are both dominating sets.

Corollary 3.5. *The set of surjective functions in $\text{Im}(\epsilon)$ admits an involution and its orbit space is in one to one correspondance with the set of all domatic 2-partitions.*

As an example, consider the imperfect graph $G_3 = (V, E)$ with vertex set $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and edge set $E = \{v_1v_2, v_1v_4, v_2v_3, v_2v_5, v_3v_4, v_4v_5, v_3v_6\}$; see the figure below




Given an integer k with $0 \leq k \leq 63$, one uniquely writes

$$k = 32i_1 + 16i_2 + 8i_3 + 4i_4 + 2i_5 + i_6$$

with $i_j \in \{0, 1\}$ for $1 \leq j \leq 6$ (that is, k written in the base 2). Let $s_k : V \rightarrow \mathbb{Z}_2$ denote the function defined as $s_k(v_j) = i_j$, when k is written as above. Then

$$\mathbb{Z}_2^V = \{s_k \mid k \in \mathbb{Z}, 0 \leq k \leq 63\}.$$

The system F associated to this graph is as follows:



The shaded nodes denote the elements in $\text{Im}(\epsilon)$. The set of all surjective elements in $\text{Im}(\epsilon)$ is

$$\{s_7, s_{12}, s_{14}, s_{19}, s_{21}, s_{24}, s_{26}, s_{28}, s_{35}, s_{37}, s_{39}, s_{42}, s_{44}, s_{49}, s_{51}, s_{56}\}$$

and the only fixed point is s_0 , the constant map at 0. Note that all non-zero elements in $\text{Im}(\epsilon)$ are in one to one correspondence with dominating sets that are also nonblockers.

3.1. Graph induced monoid actions and further consequences. We can generalize the notion of graph dynamical system to action of monoids as follows. Let M be a finitely generated monoid, $G = (V, E)$ be a graph and S be a set of states. We say an action of M on S^V graph induced if for every generator $m \in M$, the map $\mu_m : S^V \rightarrow S^V$ given by $\mu_m(s) = m \cdot s$ is a graph dynamical system with dependency graph $G = (V, E)$. If further μ_m is a generalized cellular automata, then we say the M -action is cellular.

Conversely, by a given collection of graph dynamical systems on the same graph, we can define an action of a free monoid. For any given set K , let $\mathcal{F}(K)$ denote the free monoid generated by elements of K . Let $\{F_t : S^V \rightarrow S^V \mid t \in K\}$ a collection of graph dynamical systems each of which having dependency graph $G = (V, E)$. For $t \in K$, one defines the action of $t \in \mathcal{F}(K)$ (considered as a word with length one) as $t \cdot s = F_t(s)$ for every $s \in S$. Therefore, given a word $w = t_{i_1} t_{i_2} \dots t_{i_n} \in \mathcal{F}(K)$, we have $w \cdot s = F_{i_1} \circ F_{i_2} \dots F_{i_n}(s)$.

Here we construct several graph induced action for the free monoid on two letters and establish a relation between their fixed points and more special dominating sets; such as, independent or perfect dominating sets, via reversing their actions. For the rest of the paper, we let $G = (V, E)$ be a connected finite simple graph and $F : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ be as in Theorem 3.4.

3.1.1. On independent domination. A subset $A \subseteq V$ is called an independent set of $G = (V, E)$ if no two vertices in A is adjacent. An independent set that is also dominating is called an independent dominating set. These subsets are also called maximal independent sets as such a set can not be properly contained in a larger independent set. Let $M = \mathcal{F}(a, b)$ be the free monoid with two generators a, b . Consider the following action of M on \mathbb{Z}_2^V defined on generators by $a \cdot s = F^{\circ 2}(s)$ and $b \cdot s = sP_s$; so that, for every $v \in V$

$$(b \cdot s)(v) = s(v) \prod_{x \in N(v)} (1 + s(x)).$$

It is known that any independent dominating set must be minimal, so that its complement is also a dominating set. Thus, the following corollary follows from Theorem 3.4.

Corollary 3.6. *If $s^{-1}(1)$ is an independent dominating set of G , then $a \cdot s = s$.*

We have the following lemma:

Lemma 3.7. *For $s \in \mathbb{Z}_2^V$, we have $b \cdot s = s$ if and only if $s^{-1}(1)$ is an independent set of G .*

Proof. Assume $s^{-1}(1)$ is an independent set; so that, for $v \in V$ if $s(v) = 1$, then $s(x) = 0$ whenever $x \in N(v)$. Thus, if $s(v) = 1$, then $(b \cdot s)(v) = s(v) \prod_{x \in N(v)} (1 + s(x)) = 1$. Note that if $s(v) = 0$, then $(b \cdot s)(v) = 0$. Thus, $b \cdot s = s$. Conversely, assume $s^{-1}(1)$ is not an independent set. Then there exists $v, w \in V$ such that $\{v, w\} \in E$ and $s(v) = s(w) = 1$. This implies $\prod_{x \in N(v)} (1 + s(x)) = 0$ as $w \in N(v)$; and thus, $(b \cdot s)(v) = 0$. Therefore $b \cdot s \neq s$ since $s(v) = 1$. \square

Since we previously used the notation $\text{Inv}(\mathbb{Z}_2^V)$ for the reverse action for the \mathbb{N} -set on \mathbb{Z}_2^V , in this section we use the notation $\text{Inv}_M(\mathbb{Z}_2^V) = \{\sigma : M \rightarrow \mathbb{Z}_2^V \mid \forall n, m \in M, m \cdot \sigma(nm) = \sigma(n)\}$, for the reverse action construction for the M -set \mathbb{Z}_2^V . Similarly, we denote by $\epsilon_M : \text{Inv}_M(\mathbb{Z}_2^V) \rightarrow \mathbb{Z}_2^V$ the evaluating function in this case; i.e. $\epsilon_M(\sigma) = \sigma(e)$ where e is the identity element (the empty word) in M . By Lemma 2.2, ϵ_M is injective. We have the following corollary.

Corollary 3.8. *For $s \in \mathbb{Z}_2^V$, $s \in \text{Im}(\epsilon_M)$ if and only if $s^{-1}(1)$ is a maximal independent set of G .*

Proof. Assume $s^{-1}(1)$ is a maximal independent set of G . Then by Lemma 3.7 $b \cdot s = s$. Since any maximal independent set is also a dominating set, by Corollary 3.6, $a \cdot s = s$. This means s is a fixed point of the action, and by Lemma 2.2, $s \in \text{Im}(\epsilon_M)$.

Conversely, assume $s \in \text{Im}(\epsilon_M)$. Then by Lemma 2.2, $\mu_m|_{Ms} : Ms \rightarrow Ms$ is a bijection for each $m \in M$, or equivalently μ_a and μ_b are bijections (where $\mu_m : \mathbb{Z}_2^V \rightarrow \mathbb{Z}_2^V$ is the function given by $\mu_m(s) = m \cdot s$). By Theorem 3.4 and as $\langle a \rangle s \subseteq Ms$, $\mu_a|_{Ms}$ is a bijection implies both $s^{-1}(0)$ and $s^{-1}(1)$ are dominating sets of G . In this case, $\mu_a(s) = a \cdot s = F^{o2}(s) = s$. Besides, as Ms is finite, if $\mu_b|_{Ms}$ is a bijection then there exists $k \in \mathbb{N}^+$ with $b^k \cdot s = s$. Note that if $s(v) = 0$, then $(b \cdot s)(v) = 0$ and so $(b^k \cdot s)(v) = 0$. Hence, if $b^k \cdot s = s$, then $k = 1$; i.e., $b \cdot s = s$. By Lemma 3.7 this implies that $s^{-1}(1)$ is an independent set. Therefore, we need $s^{-1}(1)$ to be simultaneously a dominating set and independent set. But this means $s^{-1}(1)$ is a independent dominating set, or equivalently, a maximal independent set. \square

3.1.2. Dominating sets with dominated vertex conditions. There are several different types of dominating sets where dominated vertices satisfy certain conditions. A list of such special dominating sets can be found in [7, Sec. 4]. We can use the system F in Theorem 3.4 along with another system to obtain correspondences between such dominating sets that are also nonblockers. For each vertex $v \in V$ let $C(v)$ be a condition on v . We say a dominating set D is a C -dominating set if $C(v)$ holds for every $v \in V \setminus D$. For a given graph $G = (V, E)$ one defines an action of $M = \mathcal{F}(a, b)$ on \mathbb{Z}_2^V as follows. For $s \in \mathbb{Z}_2^V$ define $a \cdot s = F^{o2}(s)$ and $b \cdot s = 1 + T_s + sT_s$ where

$$T_s(v) = \begin{cases} 1 & \text{if } C(v) \text{ holds} \\ 0 & \text{if } C(v) \text{ does not hold} \end{cases}$$

Then, we have the following proposition.

Proposition 3.9. *The graph $G = (V, E)$ has a C -dominating and nonblocking set if and only if $\text{Im}(\epsilon_M) \neq \emptyset$. Besides, if $s \in \text{Im}(\epsilon_M)$ exists, then s is a fixed point of this action and $s^{-1}(1)$ is a C -dominating set. Conversely, if D is a C -dominating set of G that is also nonblocker, then there exists $s \in \text{Im}(\epsilon_M)$ with $s^{-1}(1) = D$.*

Proof. Let $D \subset V$ be a C -dominating and nonblocking set of G . Define $s : V \rightarrow \mathbb{Z}_2$ by

$$s(v) = \begin{cases} 1 & \text{if } v \in D \\ 0 & \text{if } v \notin D \end{cases}$$

Now, as $s^{-1}(1)$ is a C -dominating set of G , $s(v) = 0$ implies v satisfies $C(v)$ and so $T_s(v) = 1$, which implies $(b \cdot s)(v) = 1 + T_s(v) + s(v)T_s(v) = 0$. Conversely, if

$(b \cdot s)(v) = 1 + T_s(v) + s(v)T_s(v) = 0$ then $(1 + s(v))T_s(v) = 1$ and so $s(v) = 0$. Therefore, in the case when $s^{-1}(1)$ is a C -dominating set, $s(v) = 0$ if and only if $(b \cdot s)(v) = 0$; i.e., $b \cdot s = s$. By assumption D is also a nonblocking set, so $a \cdot s = s$. This means s is a fixed point of the M -action as described above; and thus, $s \in \text{Im}(\epsilon_M)$.

Assume now $s \in \text{Im}(\epsilon_M)$. Then by Theorem 3.4 we have $a \cdot s = s$. This means $s^{-1}(1)$ is a dominating and nonblocking set. Since by Lemma 2.2 $\mu_b|_{Ms} : Ms \rightarrow Ms$ is a bijection, there exists a positive integer k with $b^k \cdot s = s$. Note that if $s(v) = 1$, then $(b \cdot s)(v) = 1$. Thus, $b^k \cdot s = s$ gives $k = 1$ and s is a fixed point of the action. Now assume that $s^{-1}(1)$ is not a C -dominating set. Then there exists a vertex v with $s(v) = T_s(v) = 0$. This means $(b \cdot s)(v) = 1 + T_s(v) + s(v)T_s(v) = 1$, i.e. $b \cdot s \neq s$. But this is a contradiction, so $s^{-1}(1)$ must be a C -dominating set. \square

The proposition can be applied to several types of special dominating sets mentioned in [7, Sec. 4] (e.g., perfect and even/odd dominating sets) by choosing T_s appropriately. For example, if T_s is given by

$$T_s(v) = \begin{cases} 1 & \text{if } |N(v) \cap s^{-1}(1)| = 1 \\ 0 & \text{if } |N(v) \cap s^{-1}(1)| \neq 1 \end{cases},$$

then one gets all perfect dominating sets. Note that any perfect dominating set has to be minimal, see [10], so must also be a nonblocker.

Alternatively, we can define T_s as

$$T_s(v) = \begin{cases} 1 & \text{if } |N(v) \cap s^{-1}(1)| = 1 \pmod{2} \\ 0 & \text{if } |N(v) \cap s^{-1}(1)| = 0 \pmod{2} \end{cases}.$$

In this case, $\text{Im}(\epsilon_M)$ consists of elements s such that $s^{-1}(1)$ is an odd dominating set that is also nonblocker. Note that by Sutner's Theorem [13] every finite graph has an odd dominating set.

Observe that a similar proposition can be obtained for dominating sets whose vertices satisfy a given condition.

REFERENCES

- [1] Juan A. Aledo, Ali Barzanouni, Ghazaleh Malekbala, Leila Sharifan, and Jose C. Valverde. Fixed points in generalized parallel and sequential dynamical systems induced by a minterm or maxterm Boolean functions. *J. Comput. Appl. Math.*, 408:13, 2022. Id/No 114070.
- [2] C. L. Barrett, H. S. Mortveit, and C. M. Reidys. Elements of a theory of simulation. II: Sequential dynamical systems. *Appl. Math. Comput.*, 107(2-3):121–136, 2000.
- [3] Chris Barrett, Harry B. III Hunt, Madhav V. Marathe, S. S. Ravi, Daniel J. Rosenkrantz, and Richard E. Stearns. Modeling and analyzing social network dynamics using stochastic discrete graphical dynamical systems. *Theor. Comput. Sci.*, 412(30):3932–3946, 2011.
- [4] Frank Dehne, Michael Fellows, Henning Fernau, Elena Prieto, and Frances Rosamond. nonblocker : Parameterized algorithmics for minimum dominating set. In *SOFSEM 2006: Theory and practice of computer science. 32nd conference on current trends in theory and practice of computer science, Měříň, Czech Republic, January 21–27, 2006. Proceedings*, pages 237–245. Berlin: Springer, 2006.
- [5] Mehmet Akif Erdal and Özgün Ünlü. Semigroup actions on sets and the Burnside ring. *Appl. Categ. Struct.*, 26(1):7–28, 2018.
- [6] Stephen Eubank, Hasan Guclu, VS Anil Kumar, Madhav V Marathe, Aravind Srinivasan, Zoltan Toroczkai, and Nan Wang. Modelling disease outbreaks in realistic urban social networks. *Nature*, 429(6988):180–184, 2004.
- [7] Teresa W. Haynes, Stephen T. Hedetniemi, and Michael A. Henning. Models of domination in graphs. In *Topics in domination in graphs*, pages 13–30. Cham: Springer, 2020.

- [8] Teresa W. Haynes, Stephen T. Hedetniemi, and Michael A. Henning, editors. *Topics in domination in graphs*, volume 64 of *Dev. Math.* Cham: Springer, 2020.
- [9] Teresa W. Haynes, Stephen T. Hedetniemi, and Peter J. Slater. *Fundamentals of domination in graphs*, volume 208 of *Pure Appl. Math., Marcel Dekker*. New York, NY: Marcel Dekker, Inc., 1998.
- [10] Marilyn Livingston and Quentin F. Stout. Perfect dominating sets. *Congr. Numerantium*, 79:187–203, 1990.
- [11] Øystein Ore. *Theory of graphs*, volume 38 of *Colloq. Publ., Am. Math. Soc.* American Mathematical Society (AMS), Providence, RI, 1962.
- [12] Assieh Saadatpour, István Albert, and Réka Albert. Attractor analysis of asynchronous Boolean models of signal transduction networks. *J. Theor. Biol.*, 266(4):641–656, 2010.
- [13] K. Sutner. Linear cellular automata and the garden-of-eden. *Math. Intell.*, 11(2):49–53, 1989.

YEDITEPE UNIVERSITY, FACULTY OF ARTS AND SCIENCES, DEPARTMENT OF MATHEMATICS,
34755, İSTANBUL, TURKEY

Email address: mehmet.erdal@yeditepe.edu.tr