

VARIATIONS OF PROPERTY (A) CONSTANTS AND LEBESGUE-TYPE INEQUALITIES FOR THE WEAK THRESHOLDING GREEDY ALGORITHMS

HÙNG VIỆT CHU

ABSTRACT. Albiac and Wojtaszczyk introduced property (A) to characterize 1-greedy bases. Later, Dilworth et al. generalized the concept to C -property (A), where the case $C = 1$ gives property (A). They (among other results) characterized greedy bases by unconditionality and C -property (A). In this paper, we extend the definition of the so-called A-property constant to (A, τ) -property constants and use the extension to obtain new estimates for various Lebesgue parameters. Furthermore, we study the relation among (A, τ) -property constants and other well-known constants when τ varies.

CONTENTS

1. Introduction	1
2. Preliminary results	6
3. Bounds for Lebesgue parameters	8
4. Lebesgue constants, Property (A, τ), and partial symmetry	12
5. Property (A, τ), uniform property (A), and quasi-greedy bases	15
6. From classical greedy-type bases to weak greedy-type bases	22
7. Future research	26
Appendix A. Repeated proofs	26
References	27

1. INTRODUCTION

1.1. **Lebesgue-type parameters.** Let $(\mathbb{X}, \|\cdot\|)$ be an infinite-dimensional Banach space over the field $\mathbb{F} = \mathbb{R}$ or \mathbb{C} with dual $(\mathbb{X}^*, \|\cdot\|_*)$. We say that $\mathcal{B} = (e_n)_{n=1}^\infty$ is a *semi-normalized Markushevich* basis (or M-basis or simply a basis, for short) if the following hold

- (1) There exists a unique collection of biorthogonal functionals $(e_n^*)_{n=1}^\infty$ such that $e_i^*(e_j) = \delta_{i,j}$.
- (2) $0 < \inf_n \{\|e_n\|, \|e_n^*\|_*\} \leq \sup_n \{\|e_n\|, \|e_n^*\|_*\} < \infty$.
- (3) $\mathbb{X} = \overline{\text{span}\{e_n : n \in \mathbb{N}\}}$.
- (4) $\mathbb{X}^* = \overline{\text{span}\{e_n^* : n \in \mathbb{N}\}}^{w^*}$.

2020 *Mathematics Subject Classification.* 41A65; 46B15.

Key words and phrases. Greedy, almost greedy, partially greedy, Property (A).

The author is thankful to Timur Oikhberg for helpful feedback on earlier drafts of this paper.

With respect to an M-basis \mathcal{B} , every $x \in \mathbb{X}$ can be represented by the formal sum (possibly divergent) $\sum_{n=1}^{\infty} e_n^*(x)e_n$ with $\lim_{n \rightarrow \infty} e_n^*(x) = 0$. If, in addition,

- (5) the partial sum operator $(S_m)_{m=1}^{\infty}$ defined as $S_m(x) = \sum_{n=1}^m e_n^*(x)e_n$ is uniformly bounded, i.e, $\sup_m \|S_m\| \leq C$ for some $C > 0$,

we say that \mathcal{B} is a *Schauder basis*. In this case, we let $\mathbf{K}_b := \sup_m \|S_m\|$, which is called the *basis constant*. We set $\text{supp}(x) := \{n : e_n^*(x) \neq 0\}$, $\mathbb{X}_c = \{x \in \mathbb{X} : |\text{supp}(x)| < \infty\}$, and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

1.1.1. *Thresholding Greedy Algorithm (TGA)*. We wish to approximate each vector $x \in \mathbb{X}$ with a finite linear combination from the basis \mathcal{B} . Among all, the greedy algorithm is probably the most natural, which we now describe. Consider $x \sim \sum_{n=1}^{\infty} e_n^*(x)e_n$. A finite set $\Lambda \subset \mathbb{N}$ is called a *greedy set* of x if $\min_{n \in \Lambda} |e_n^*(x)| \geq \max_{n \notin \Lambda} |e_n^*(x)|$. For $m \in \mathbb{N}_0$, let

$$\mathcal{G}(x) := \{\Lambda : \Lambda \text{ is a greedy set of } x\} \text{ and } \mathcal{G}(x, m) := \{\Lambda \in \mathcal{G}(x) : |\Lambda| = m\}.$$

A *greedy operator* of order m is defined as

$$G_m(x) := \sum_{n \in \Lambda_x} e_n^*(x)e_n, \text{ for some } \Lambda_x \in \mathcal{G}(x, m).$$

Let \mathcal{G}_m denote the set of all greedy operators of order m , and $\mathcal{G} = \cup_{m \geq 1} \mathcal{G}_m$. We have $G_0(x) = 0$ and $\mathcal{G}(x, 0) = \{\emptyset\}$. We capture the error term from the greedy algorithm by

$$\gamma_m(x) := \sup_{G_m \in \mathcal{G}_m} \|x - G_m(x)\|,$$

and quantify the efficiency of the greedy algorithm by comparing $\gamma_m(x)$ to the smallest possible error of an arbitrary m -term approximation

$$\sigma_m(x) := \inf \left\{ \left\| x - \sum_{n \in A} a_n e_n \right\| : A \subset \mathbb{N}, |A| \leq m, (a_n) \subset \mathbb{F} \right\}.$$

We can also compare $\gamma_m(x)$ to the smallest projection error

$$\tilde{\sigma}_m(x) := \inf \{ \|x - P_A(x)\| : A \subset \mathbb{N}, |A| = m \},$$

where $P_A(x) = \sum_{n \in A} e_n^*(x)e_n$. We have $\gamma_0(x) = \sigma_0(x) = \tilde{\sigma}_0(x) = \|x\|$. In particular, for each $m \in \mathbb{N}_0$, let \mathbf{L}_m (and $\tilde{\mathbf{L}}_m$, respectively) be the smallest constant such that for all $x \in \mathbb{X}$,

$$\gamma_m(x) \leq \mathbf{L}_m \sigma_m(x) \text{ (and } \gamma_m(x) \leq \tilde{\mathbf{L}}_m \tilde{\sigma}_m(x), \text{ respectively.)}$$

There have been many estimates and ongoing improvements for these Lebesgue parameters \mathbf{L}_m and $\tilde{\mathbf{L}}_m$ under various settings. For example, see [1, 3, 8, 10, 11, 14, 18].

1.1.2. *Weak Thresholding Greedy Algorithm (WTGA)*. Throughout this paper, let τ be a number in $(0, 1]$. Temlyakov [17] considered the WTGA, which allows more flexibility in forming greedy sums. Given $x \in \mathbb{X}$, a finite set $\Lambda \subset \mathbb{N}$ is τ -*weak greedy* with respect to x if $\min_{n \in \Lambda} |e_n^*(x)| \geq \tau \max_{n \notin \Lambda} |e_n^*(x)|$. For $m \in \mathbb{N}_0$, let

$$\mathcal{G}(x, \tau) := \{\Lambda : \Lambda \text{ is a } \tau\text{-weak greedy set of } x\}, \text{ and}$$

$$\mathcal{G}(x, m, \tau) := \{\Lambda \in \mathcal{G}(x, \tau) : |\Lambda| = m\}.$$

A τ -greedy operator of order m is defined as

$$G_m^\tau(x) := \sum_{n \in \Lambda_x} e_n^*(x) e_n, \text{ for some } \Lambda_x \in \mathcal{G}(x, m, \tau).$$

Let \mathcal{G}_m^τ denote the set of all τ -greedy operators of order m , and $\mathcal{G}^\tau = \cup_{m \geq 1} \mathcal{G}_m^\tau$. We capture the error term from the weak greedy algorithm by

$$\gamma_{m,\tau}(x) := \sup_{G_m^\tau \in \mathcal{G}_m^\tau} \|x - G_m^\tau(x)\|.$$

Similar to \mathbf{L}_m and $\tilde{\mathbf{L}}_m$, for $m \in \mathbb{N}_0$, we define $\mathbf{L}_{m,\tau}$ to be the smallest constant such that

$$\gamma_{m,\tau}(x) \leq \mathbf{L}_{m,\tau} \sigma_m(x), \forall x \in \mathbb{X},$$

and $\tilde{\mathbf{L}}_{m,\tau}$ to be the smallest constant such that

$$\gamma_{m,\tau}(x) \leq \tilde{\mathbf{L}}_{m,\tau} \tilde{\sigma}_m(x), \forall x \in \mathbb{X}.$$

We call $\mathbf{L}_{m,\tau}$ the *weak greedy Lebesgue parameter* and call $\tilde{\mathbf{L}}_{m,\tau}$ the *weak almost greedy Lebesgue parameter*.

1.1.3. *Chebyshev weak thresholding greedy algorithms (CWTGA)*. We also study the Lebesgue parameter for the CWTGA, which was introduced by Dilworth et al. [13] to improve the rate of convergence of the TGA. A *Chebyshev τ -greedy operator* of order m ($m \in \mathbb{N}_0$), denoted by $CG_m^\tau : \mathbb{X} \rightarrow \mathbb{X}$, satisfies the following: for every $x \in \mathbb{X}$, there exists $\Lambda \in \mathcal{G}(x, m, \tau)$ such that

- (1) $\text{supp}(CG_m^\tau(x)) \subset \Lambda$ and
- (2) we have

$$\|x - CG_m^\tau(x)\| = \min \left\{ \left\| x - \sum_{n \in \Lambda} a_n e_n \right\| : (a_n) \subset \mathbb{F} \right\}.$$

For $m \in \mathbb{N}_0$, let $\mathbf{L}_{m,\tau}^{ch}$ denote the smallest constant such that

$$\|x - CG_m^\tau(x)\| \leq \mathbf{L}_{m,\tau}^{ch} \sigma_m(x), \forall x \in \mathbb{X}, \forall CG_m^\tau \in \mathcal{CG}_m^\tau,$$

where \mathcal{CG}_m^τ is the set of all Chebyshev τ -greedy operators of order m and $\mathcal{CG}^\tau = \cup_{m \geq 1} \mathcal{CG}_m^\tau$. We call $\mathbf{L}_{m,\tau}^{ch}$ the *weak greedy Chebyshev Lebesgue parameter*.

Tight estimates for both parameters $\mathbf{L}_{m,\tau}$ and $\mathbf{L}_{m,\tau}^{ch}$ were established in [7, 9].

1.1.4. *Partially greedy parameter*. The *residual Lebesgue parameter* was introduced in [9] to compare the performance of greedy algorithms $(G_m^\tau(x))_{m=1}^\infty$ to that of the partial sums $(S_m(x))_{m=1}^\infty$. Specifically, for $m \in \mathbb{N}_0$, let $\mathbf{L}_{m,\tau}^{re}$ be the smallest constant such that

$$\gamma_{m,\tau}(x) \leq \mathbf{L}_{m,\tau}^{re} \|x - S_m(x)\|, \forall x \in \mathbb{X}.$$

In [5], the authors established bounds for the so-called *strong residual Lebesgue parameter*. For each $m \in \mathbb{N}_0$, they defined $\widehat{\mathbf{L}}_{m,\tau}^{re}$ as the smallest constant verifying

$$\gamma_{m,\tau}(x) \leq \widehat{\mathbf{L}}_{m,\tau}^{re} \widehat{\sigma}_m(x), \forall x \in \mathbb{X}, \quad (1.1)$$

where $\widehat{\sigma}_m(x) = \inf_{0 \leq n \leq m} \|x - S_n(x)\|$. Clearly, $\mathbf{L}_{m,\tau}^{re} \leq \widehat{\mathbf{L}}_{m,\tau}^{re}$; if our basis is Schauder, then $(\mathbf{K}_b + 1)\mathbf{L}_{m,\tau}^{re} \geq \widehat{\mathbf{L}}_{m,\tau}^{re}$, and so $\mathbf{L}_{m,\tau}^{re} \approx \widehat{\mathbf{L}}_{m,\tau}^{re}$. For estimates of $\mathbf{L}_{m,\tau}^{re}$ and $\widehat{\mathbf{L}}_{m,\tau}^{re}$, see [5, 9].

1.2. Notation and relevant constants. For two functions $f(a_1, a_2, \dots)$ and $g(a_1, a_2, \dots)$, we write $f \lesssim g$ to indicate that there exists an absolute constant $C > 0$ (independent of a_1, a_2, \dots) such that $f \leq Cg$. Similarly, $f \gtrsim g$ means that $Cf \geq g$ for some constant C . For two sets $A, B \subset \mathbb{N}$, we write $A < B$ for $\max A < \min B$ and write $A < m$ ($m \in \mathbb{N}$) for $\max A < m$. We shall use the following notation

$$1_A := \sum_{n \in A} e_n \text{ and } 1_{\varepsilon A} := \sum_{n \in A} \varepsilon_n e_n,$$

where $\varepsilon = (\varepsilon_n)_{n=1}^\infty$ and $|\varepsilon_n| = 1$. For $x \in \mathbb{X}$, $\|x\|_\infty := \sup_n |e_n^*(x)|$, and we write $A \sqcup B \sqcup x$ to indicate that A, B , and $\text{supp}(x)$ are pairwise disjoint. We now present a list of relevant constants, most of which have appeared frequently in the literature, except that we may generalize them to accommodate the concept of τ -weak greedy sets:

- (A, τ) -property constants: for $m \in \mathbb{N}_0$,

$$\nu_{m, \tau} = \sup_{\substack{(\varepsilon), (\delta) \\ \|x\|_\infty \leq 1/\tau}} \left\{ \frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} : |A| = |B| \leq m, A \sqcup B \sqcup x \right\}. \quad (1.2)$$

In the above definition of $\nu_{m, \tau}$, we can replace the condition “ $|A| = |B| \leq m$ ” by “ $|B| \leq |A| \leq m$ ” due to norm convexity.

- Left (A, τ) -property constants: for $m \in \mathbb{N}_0$,

$$\nu_{m, \tau, \ell} = \sup_{\substack{(\varepsilon), (\delta) \\ \|x\|_\infty \leq 1/\tau}} \left\{ \frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} : |A| = |B| \leq m, B < A, A \sqcup B \sqcup x \right\}, \text{ and} \quad (1.3)$$

$$\nu'_{m, \tau, \ell} = \sup_{\substack{(\varepsilon), (\delta) \\ \|x\|_\infty \leq 1/\tau}} \left\{ \frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} : |B| \leq |A| \leq m, B < \text{supp}(x) \sqcup A, B \leq m \right\}. \quad (1.4)$$

Note that while $\nu'_{m, \tau, \ell}$ puts more restrictions on the positions of sets A and B than $\nu_{m, \tau, \ell}$ does, $\nu'_{m, \tau, \ell}$ allows A and B to be of different cardinalities, but $\nu_{m, \tau, \ell}$ does not.

- Unconditional constants: for $m \in \mathbb{N}_0$,

$$k_m = \sup_{|A| \leq m} \|P_A\| \text{ and } k_m^c = \sup_{|A| \leq m} \|I - P_A\|.$$

- Quasi-greedy constants: for $m \in \mathbb{N}_0$,

$$g_{m, \tau} = \sup \{ \|G_n^\tau\| : G_n^\tau \in \cup_{k \leq m} \mathcal{G}_k^\tau \}, \text{ and} \\ g_{m, \tau}^c = \sup \{ \|I - G_n^\tau\| : G_n^\tau \in \cup_{k \leq m} \mathcal{G}_k^\tau \}.$$

We have $g_{m, \tau}^c - 1 \leq g_{m, \tau} \leq g_{m, \tau}^c + 1$.

- Super-democracy constants: for $m \in \mathbb{N}$,

$$\mu_m = \sup \left\{ \frac{\|1_{\delta B}\|}{\|1_{\varepsilon A}\|} : |A| = |B| \leq m, (\varepsilon), (\delta) \right\}. \quad (1.5)$$

- Super-conservancy constants: for $m \in \mathbb{N}$,

$$\psi_m = \sup_{(\varepsilon), (\delta)} \left\{ \frac{\|1_{\delta B}\|}{\|1_{\varepsilon A}\|} : |A| = |B| \leq m, B \subset A \right\}. \quad (1.6)$$

1.3. **Main results.** Three main goals of this paper are to

- use (left) (A, τ) -property constants to generalize tight estimates for $\mathbf{L}_{m, \tau}$, $\tilde{\mathbf{L}}_{m, \tau}$, $\mathbf{L}_{m, \tau}^{ch}$, $\mathbf{L}_{m, \tau}^{re}$, and $\widehat{\mathbf{L}}_{m, \tau}^{re}$.
- show how our estimates help generalize various existing results in the literature.
- study the relation among (A, τ) -property constants and other well-known constants when τ varies.

Theorem 1.1. *Let \mathcal{B} be an M -basis in \mathbb{X} . Then for $m \in \mathbb{N}$,*

$$\max \left\{ k_m^c, \tilde{\mathbf{L}}_{m, \tau}, \frac{\nu_{m, \tau}}{\tau} \right\} \leq \mathbf{L}_{m, \tau} \leq \frac{k_{2m-1}^c \nu_{m, \tau}}{\tau}. \quad (1.7)$$

Theorem 1.2. *Let \mathcal{B} be an M -basis in \mathbb{X} . Then for $m \in \mathbb{N}$,*

$$\max \left\{ g_{m, \tau}^c, \frac{\nu_{m, \tau}}{\tau} \right\} \leq \tilde{\mathbf{L}}_{m, \tau} \leq \frac{g_{m-1, \tau}^c \nu_{m, \tau}}{\tau}. \quad (1.8)$$

Proposition 1.3. *For $m \in \mathbb{N}_0$,*

$$\frac{\nu'_{m, \tau, \ell}}{\tau} \leq \max_{0 \leq k \leq m} \widehat{\mathbf{L}}_{k, \tau}^{re}. \quad (1.9)$$

Theorem 1.4. *Let \mathcal{B} be an M -basis in \mathbb{X} . Then*

$$\mathbf{L}_{m, \tau}^{re} \leq \frac{g_{m-1, \tau}^c \nu_{m, \tau, \ell}}{\tau}, \forall m \in \mathbb{N}, \quad (1.10)$$

$$g_{m, \tau}^c \leq \widehat{\mathbf{L}}_{m, \tau}^{re} \leq \frac{g_{m-1, \tau}^c \nu'_{m, \tau, \ell}}{\tau}, \forall m \in \mathbb{N}. \quad (1.11)$$

As a result,

$$\widehat{\mathbf{L}}_{1, \tau}^{re} = \frac{\nu'_{1, \tau, \ell}}{\tau}. \quad (1.12)$$

If \mathcal{B} is Schauder, then

$$\mathbf{L}_{m, \tau}^{re} \geq \frac{g_{m, \tau}^c}{\mathbf{K}_b + 1}, \forall m \in \mathbb{N}_0. \quad (1.13)$$

Theorem 1.5. *Let \mathcal{B} be a Schauder basis with basis constant \mathbf{K}_b . Then for $m \geq \mathbb{N}_0$,*

$$g_{m, \tau}^c \leq \mathbf{K}_b \mathbf{L}_{m, \tau}^{ch} (1 + \mathbf{L}_{m, \tau}^{ch} + \mathbf{L}_{m, \tau}^{ch} \mathbf{K}_b). \quad (1.14)$$

Consequently,

$$\mathbf{L}_{m, \tau}^{ch} \geq \frac{1}{\mathbf{K}_b} \left(\frac{g_{m, \tau}^c}{3} \right)^{1/2}. \quad (1.15)$$

Remark 1.6. When $\tau = 1$, Theorems 1.1 and 1.2 give [8, Inequalities (1.3) and (1.5)]. Similarly, Proposition 1.3 implies [5, Proposition 1.13]. Finally, (1.11) and (1.12) give [5, Theorem 1.14].

We describe the outline of this paper:

- Section 2 presents important results that will be used in due course;

- Section 3 establishes bounds for Lebesgue parameters;
- Section 4 studies the relation among Lebesgue constants, (left) Property (A, τ), and partial symmetry for largest coefficients;
- Section 5 studies the relation among Property (A, τ), uniform property (A), and quasi-greedy bases;
- Section 6 estimates Lebesgue constants when we go from the classical greedy setting to the weak greedy setting;
- Section 7 contains several questions for further investigation.

2. PRELIMINARY RESULTS

We prove an analog of [3, Proposition 2.1] for the unconditionality constant k_m^c .

Proposition 2.1. *Fix $m \in \mathbb{N}$. Let $J \subset \mathbb{N}$ with $|J| \leq m$. Assume $(a_n)_{n \in J}, (b_n)_{n \in J}$ are scalars so that $|a_n| \leq |b_n|$ for all $n \in J$, and $\text{sgn}(a_n) = \text{sgn}(b_n)$ whenever $a_n b_n \neq 0$. Then*

$$\left\| x + \sum_{n \in J} a_n e_n \right\| \leq k_m^c \left\| x + \sum_{n \in J} b_n e_n \right\|,$$

for all $x \in \mathbb{X}$, where $\text{supp}(x) \cap J = \emptyset$. If, in addition, there exists $j \in J$ such that $a_j = b_j$, then

$$\left\| x + \sum_{n \in J} a_n e_n \right\| \leq k_{m-1}^c \left\| x + \sum_{n \in J} b_n e_n \right\|,$$

Proof. For each $n \in J$, we have

$$\frac{a_n}{b_n} = \int_0^{\frac{a_n}{b_n}} 1 dt;$$

hence,

$$x + \sum_{n \in J} a_n e_n = x + \sum_{n \in J} \left(\int_0^1 b_n \chi_{(0, \frac{a_n}{b_n})}(t) dt \right) e_n = \int_0^1 \left(x + \sum_{n \in J} b_n \chi_{(0, \frac{a_n}{b_n})}(t) e_n \right) dt.$$

For each $t \in (0, 1)$, we have

$$\left\| x + \sum_{n \in J} b_n \chi_{(0, \frac{a_n}{b_n})}(t) e_n \right\| \leq k_m^c \left\| x + \sum_{n \in J} b_n e_n \right\|.$$

Therefore,

$$\left\| x + \sum_{n \in J} a_n e_n \right\| \leq \int_0^1 \left\| x + \sum_{n \in J} b_n \chi_{(0, \frac{a_n}{b_n})}(t) e_n \right\| dt \leq k_m^c \left\| x + \sum_{n \in J} b_n e_n \right\|,$$

where in the last inequality, we can replace k_m^c by k_{m-1}^c if $a_j = b_j$ for some $j \in J$. This completes our proof. \square

Lemma 2.2. For $m \in \mathbb{N}_0$, let

$$\Omega_{m,\tau} := \sup_{\substack{(\varepsilon) \\ \|x\|_\infty \leq t/\tau}} \left\{ \frac{\|x\|}{\|x - P_B(x) + t1_{\varepsilon A}\|} : |A| = |B| \leq m, (\text{supp}(x) \cup B) \cap A = \emptyset \right\}. \quad (2.1)$$

We have

$$\Omega_{m,\tau} = \frac{\nu_{m,\tau}}{\tau}, \forall m \in \mathbb{N}_0.$$

Proof. Let $A, B, x, (\varepsilon), t$ be chosen as in (2.1). By norm convexity, we have

$$\begin{aligned} \|x\| &= \left\| x - P_B(x) + \sum_{n \in B} e_n^*(x) e_n \right\| \leq \sup_{(\delta)} \left\| x - P_B(x) + \frac{t}{\tau} 1_{\delta B} \right\| \\ &= \frac{1}{\tau} \sup_{(\delta)} \|\tau(x - P_B(x)) + t1_{\delta B}\| \\ &\leq \frac{\nu_{m,\tau}}{\tau} \|x - P_B(x) + t1_{\varepsilon A}\|. \end{aligned}$$

Hence, $\Omega_{m,\tau} \leq \nu_{m,\tau}/\tau$.

For the reverse inequality, let $A, B, x, (\varepsilon), (\delta)$ be chosen as in (1.2). Let $y = \tau x + 1_{\delta B}$. Then $\|y\|_\infty \leq 1$. By (2.1),

$$\|\tau x + 1_{\delta B}\| = \|y\| \leq \Omega_{m,\tau} \|y - P_B(y) + \tau 1_{\varepsilon A}\| = \Omega_{m,\tau} \tau \|x + 1_{\varepsilon A}\|.$$

Hence, $\Omega_{m,\tau} \geq \nu_{m,\tau}/\tau$. \square

Similarly, we have two following lemmas, whose proofs are moved to the Appendix.

Lemma 2.3. For $m \in \mathbb{N}_0$, let

$$\Omega_{m,\tau,\ell} := \sup_{\substack{(\varepsilon) \\ \|x\|_\infty \leq t/\tau}} \left\{ \frac{\|x\|}{\|x - P_B(x) + t1_{\varepsilon A}\|} : |A| = |B| \leq m, B < A, \text{supp}(x) \cap A = \emptyset \right\}. \quad (2.2)$$

We have

$$\Omega_{m,\tau,\ell} = \frac{\nu_{m,\tau,\ell}}{\tau}, \forall m \in \mathbb{N}_0.$$

Lemma 2.4. For $m \in \mathbb{N}_0$, let

$$\Omega'_{m,\tau,\ell} := \sup_{\substack{(\varepsilon) \\ \|x\|_\infty \leq t/\tau \\ B \leq m}} \left\{ \frac{\|x\|}{\|x - P_B(x) + t1_{\varepsilon A}\|} : |B| \leq |A| \leq m, B < \text{supp}(x - P_B(x)) \sqcup A \right\}. \quad (2.3)$$

We have

$$\Omega'_{m,\tau,\ell} = \frac{\nu'_{m,\tau,\ell}}{\tau}, \forall m \in \mathbb{N}_0.$$

The next proposition shares the same spirit with Proposition 2.1 for the constant $g_{m,\tau}^c$.

Proposition 2.5. *Let $m \in \mathbb{N}$ and $\alpha \geq 0$. Then for any $x \in \mathbb{X}$ with $\|x\|_\infty \leq \alpha/\tau$ and any (ε) , we have*

$$\|x + \alpha 1_{\varepsilon A}\| \leq g_{m,\tau}^c \left\| x + \sum_{n \in A} \varepsilon_n a_n e_n \right\|,$$

for all real scalars (a_n) with $a_n \geq \alpha$ and any $A \subset \mathbb{N}$ with $|A| \leq m$ and $A \cap \text{supp}(x) = \emptyset$. If, in addition, there exists $j \in A$ such that $a_j = \alpha$, then

$$\|x + \alpha 1_{\varepsilon A}\| \leq g_{m-1,\tau}^c \left\| x + \sum_{n \in A} \varepsilon_n a_n e_n \right\|.$$

Proof. Let $z = \sum_{n \in A} \varepsilon_n a_n e_n$, $y = x + z$, and $\Lambda_{\alpha,s} = \{n \in A : |e_n^*(z)| > \alpha/s\}$ for each $s \in (0, 1]$. Clearly, $\Lambda_{\alpha,s}$ is a τ -weak greedy set of y and $|\Lambda_{\alpha,s}| \leq |A| \leq m$. We have

$$\begin{aligned} \|x + \alpha 1_{\varepsilon A}\| &= \left\| x + \int_0^1 \left(\sum_{n \in A} \chi_{[0, \frac{\alpha}{|e_n^*(z)|}]}(s) e_n^*(z) e_n \right) ds \right\| \\ &= \left\| \int_0^1 (x + (I - P_{\Lambda_{\alpha,s}})z) ds \right\| \\ &\leq \int_0^1 \|x + (I - P_{\Lambda_{\alpha,s}})z\| ds \\ &= \int_0^1 \|(I - P_{\Lambda_{\alpha,s}})y\| ds \leq g_{m,\tau}^c \|y\|. \end{aligned}$$

The proof is completed.

The second assertion is obvious because if there exists $j \in A$ such that $a_j = \alpha$, then $|\Lambda_{\alpha,s}| \leq m - 1$. \square

3. BOUNDS FOR LEBESGUE PARAMETERS

Proof of Theorem 1.1. First, we establish the upper bound for $\mathbf{L}_{m,\tau}$. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$ and $A \in \mathcal{G}(x, m, \tau)$. Let $z = \sum_{n \in B} b_n e_n$ with $B \subset \mathbb{N}$, $|B| = m$, and $(b_n)_{n \in B} \subset \mathbb{F}$. Set $\alpha := \min_{n \in A} |e_n^*(x)|$. By definition, $\|x - P_A(x)\|_\infty \leq \alpha/\tau$. By Lemma 2.2, we have

$$\|x - P_A(x)\| \leq \frac{\nu_{m,\tau}}{\tau} \left\| x - P_A(x) - P_{B \setminus A}(x) + \alpha \sum_{n \in A \setminus B} \text{sgn}(e_n^*(x)) e_n \right\|.$$

Case 1: if $|A \cup B| \leq 2m - 1$, then by the first assertion of Proposition 2.1, we have

$$\|x - P_A(x)\| \leq \frac{\nu_{m,\tau} k_{2m-1}^c}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in A \setminus B} e_n^*(x) e_n + \sum_{n \in B} (e_n^*(x) - b_n) e_n \right\|. \quad (3.1)$$

Case 2: if $|A \cup B| = 2m$, then A is disjoint from B . By the second assertion of Proposition 2.1, we also have (3.1)

Taking the infimum over all B and $(b_n)_{n \in B}$ to obtain

$$\|x - P_A(x)\| \leq \frac{\nu_{m,\tau} k_{2m}^c}{\tau} \sigma_m(x).$$

As A is arbitrary, $\gamma_{m,\tau}(x) \leq \frac{\nu_{m,\tau} k_{2m-1}^c}{\tau} \sigma_m(x)$ and so, $\mathbf{L}_{m,\tau} \leq \nu_{m,\tau} k_{2m-1}^c / \tau$.

We now bound $\mathbf{L}_{m,\tau}$ from below. Observe that $\mathbf{L}_{m,\tau} \geq \mathbf{L}_m$ and $\mathbf{L}_m \geq k_m^c$ by [8, Proposition 1.1]; hence, $\mathbf{L}_{m,\tau} \geq k_m^c$. By definition, $\mathbf{L}_{m,\tau} \geq \tilde{\mathbf{L}}_{m,\tau}$. As we shall show in the proof of Theorem 1.2, $\tilde{\mathbf{L}}_{m,\tau} \geq \nu_{m,\tau} / \tau$. Hence,

$$L_{m,\tau} \geq \max \left\{ k_m^c, \tilde{\mathbf{L}}_{m,\tau}, \frac{\nu_{m,\tau}}{\tau} \right\}.$$

□

Proof of Theorem 1.2. First, we establish the upper bound for $\tilde{\mathbf{L}}_{m,\tau}$. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$ and $A \in \mathcal{G}(x, m, \tau)$. Let $B \subset \mathbb{N}$ with $|B| = m$. Set $\alpha := \min_{n \in A} |e_n^*(x)|$. By definition, $\|x - P_A(x)\|_\infty \leq \alpha / \tau$. By Lemma 2.2, we have

$$\begin{aligned} \|x - P_A(x)\| &\leq \frac{\nu_{m,\tau}}{\tau} \left\| x - P_A(x) - P_{B \setminus A}(x) + \alpha \sum_{n \in A \setminus B} \operatorname{sgn}(e_n^*(x)) e_n \right\| \\ &= \frac{\nu_{m,\tau}}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in A \setminus B} \frac{\alpha}{|e_n^*(x)|} e_n^*(x) e_n \right\|. \end{aligned}$$

Case 1: if $|A \setminus B| \leq m - 1$, then the first assertion of Proposition 2.5 gives

$$\|x - P_A(x)\| \leq \frac{\nu_{m,\tau} g_{m-1,\tau}^c}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in A \setminus B} e_n^*(x) e_n \right\| = \frac{\nu_{m,\tau} g_{m-1,\tau}^c}{\tau} \|x - P_B(x)\|. \quad (3.2)$$

Case 2: if $|A \setminus B| = m$, then $A \setminus B = A$, and the second assertion of Proposition 2.5 again gives (3.2).

Taking the infimum over all B , we obtain

$$\|x - P_A(x)\| \leq \frac{\nu_{m,\tau} g_{m-1,\tau}^c}{\tau} \tilde{\sigma}_m(x).$$

Since A is arbitrary, we have proved $\tilde{\mathbf{L}}_{m,\tau} \leq \nu_{m,\tau} g_{m-1,\tau}^c / \tau$.

Next, we bound $\tilde{\mathbf{L}}_{m,\tau}$ from below. Choose $A, B, x, (\varepsilon), (\delta)$ as in (1.2) with $x \in \mathbb{X}_c$. Set

$$z := 1_{\varepsilon A} + x + \frac{1}{\tau} 1_{\delta B} + 1_C,$$

where $C > A \cup B \cup \operatorname{supp}(x)$ and $|C| = m - |A|$. Since $A \cup C \in \mathcal{G}(x, m, \tau)$,

$$\begin{aligned} \left\| x + \frac{1}{\tau} 1_{\delta B} \right\| &= \|z - P_{A \cup C}(z)\| \leq \tilde{\mathbf{L}}_{m,\tau} \tilde{\sigma}_m(z) \\ &\leq \tilde{\mathbf{L}}_{m,\tau} \left\| z - \frac{1}{\tau} 1_{\delta B} - 1_C \right\| = \tilde{\mathbf{L}}_{m,\tau} \|1_{\varepsilon A} + x\|. \end{aligned}$$

Therefore,

$$\frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} \leq \tau \tilde{\mathbf{L}}_{m,\tau}.$$

We conclude that $\nu_{m,\tau} / \tau \leq \tilde{\mathbf{L}}_{m,\tau}$. Finally, we show that $\tilde{\mathbf{L}}_{m,\tau} \geq g_m^c$. Let $x \in \mathbb{X}_c$ and $\Lambda \in \mathcal{G}(x, k, \tau)$ with $k \leq m$. It suffices to prove $\|x - P_\Lambda(x)\| \leq \tilde{\mathbf{L}}_{m,\tau} \|x\|$. Let

$\alpha = \min_{n \in \Lambda} |e_n^*(x)|$. Choose $C > \text{supp}(x)$ with $|C| = m - k$. Set $y := x + 1_C$. Clearly, $\Lambda \cup C \in \mathcal{G}(y, m, \tau)$ and so,

$$\begin{aligned} \|x - P_\Lambda(x)\| &= \|y - P_{\Lambda \cup C}(y)\| \leq \tilde{\mathbf{L}}_{m,\tau} \tilde{\sigma}_m(y) \\ &\leq \tilde{\mathbf{L}}_{m,\tau} \|y - P_C(y)\| = \tilde{\mathbf{L}}_{m,\tau} \|x\|, \end{aligned}$$

as desired. \square

Proof of Proposition 1.3. We follow the argument in the proof of [5, Proposition 1.13]. Let $A, B, x, (\varepsilon), (\delta)$ be chosen as in (1.4). Let $m_1 := \max B$. By the conditions put on sets A and B , there exists a set D (possibly empty) such that $\max D \leq m_1, D \cap B = \emptyset$, and $m_1 \leq m_2 := |D \cup A| \leq m$. Set

$$y := \frac{1}{\tau} 1_{\delta B} + 1_D + x + 1_{\varepsilon A}.$$

Clearly, $D \cup A \in \mathcal{G}(y, m_2, \tau)$ and $S_{m_1}(y) = 1_D + \frac{1}{\tau} 1_{\delta B}$. We, therefore, have

$$\left\| x + \frac{1}{\tau} 1_{\delta B} \right\| = \|y - P_{D \cup A}(y)\| \leq \hat{\mathbf{L}}_{m_2,\tau}^{re} \|y - P_{m_1}(y)\| \leq \max_{0 \leq k \leq m} \hat{\mathbf{L}}_{k,\tau} \|x + 1_{\varepsilon A}\|.$$

We conclude that $\nu'_{m,\tau,\ell}/\tau \leq \max_{0 \leq k \leq m} \hat{\mathbf{L}}_{k,\tau}$. \square

Proof of Theorem 1.4. We prove each inequality below.

- (1) Proof of (1.10): We first prove the right inequality. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$, $A \in \mathcal{G}(x, m, \tau)$. Set $B = \{1, \dots, m\} \setminus A$, $F = A \setminus \{1, \dots, m\}$, and $\alpha := \min_{n \in A} |e_n^*(x)|$. By definition, $\|x - P_A(x)\|_\infty \leq \alpha/\tau$. Observe that $|B| = |F|$ and $B < F$. By Lemma 2.3 and Proposition 2.5, we have

$$\begin{aligned} &\|x - P_A(x)\| \\ &\leq \frac{\nu_{m,\tau,\ell}}{\tau} \left\| x - P_A(x) - P_B(x) + \alpha \sum_{n \in F} \text{sgn}(e_n^*(x)) e_n \right\| \\ &= \frac{\nu_{m,\tau,\ell}}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in F} \frac{\alpha}{|e_n^*(x)|} e_n^*(x) e_n \right\| \\ &\leq \frac{\nu_{m,\tau,\ell} g_{m-1,\tau}^c}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in F} e_n^*(x) e_n \right\| = \frac{\nu_{m,\tau,\ell} g_{m-1,\tau}^c}{\tau} \|x - S_m(x)\|, \end{aligned}$$

where we obtain the second inequality by case analysis as in the proof of Theorems 1.1 and 1.2. Therefore, $\mathbf{L}_{m,\tau}^{re} \leq g_{m-1,\tau}^c \nu_{m,\tau,\ell}/\tau$.

- (2) Proof of (1.11): The left inequality is immediate from (1.1) by noticing that $\hat{\sigma}_m(x) \leq \|x\|$. We prove the right inequality. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$, $A \in \mathcal{G}(x, m, \tau)$. Fix $k \leq m$. Set $B = \{1, \dots, k\} \setminus A$, $F = A \setminus \{1, \dots, k\}$, and $\alpha := \min_{n \in A} |e_n^*(x)|$. By definition, $\|x - P_A(x)\|_\infty \leq \alpha/\tau$. Observe that $|B| \leq |F| \leq m$ and $B < \text{supp}(x - P_A(x) - P_B(x)) \sqcup F$. By Lemma 2.4 and

Proposition 2.5, we have

$$\begin{aligned}
& \|x - P_A(x)\| \\
& \leq \frac{\nu'_{m,\tau,\ell}}{\tau} \left\| x - P_A(x) - P_B(x) + \alpha \sum_{n \in F} \operatorname{sgn}(e_n^*(x)) e_n \right\| \\
& = \frac{\nu'_{m,\tau,\ell}}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in F} \frac{\alpha}{|e_n^*(x)|} e_n^*(x) e_n \right\| \\
& \leq \frac{\nu'_{m,\tau,\ell} g_{m-1,\tau}^c}{\tau} \left\| P_{A \cup B}^c(x) + \sum_{n \in F} e_n^*(x) e_n \right\| = \frac{\nu'_{m,\tau,\ell} g_{m-1,\tau}^c}{\tau} \|x - S_k(x)\|,
\end{aligned}$$

where we obtain the second inequality by case analysis as in the proof of Theorems 1.1 and 1.2. Therefore, $\widehat{\mathbf{L}}_{m,\tau}^{re} \leq g_{m-1,\tau}^c \nu'_{m,\tau,\ell} / \tau$.

(3) Proof of (1.12): This inequality follows directly from (1.9) and (1.11).

(4) Proof of (1.13): This inequality is due to $\widehat{\mathbf{L}}_{m,\tau}^{re} \leq (\mathbf{K}_b + 1) \mathbf{L}_{m,\tau}^{re}$ and (1.11). \square

Proof of Theorem 1.5. Fix $x \in \mathbb{X}_c$, $m \in \mathbb{N}_0$, and $A \in \mathcal{G}(x, m, \tau)$. Choose $D \subset \mathbb{N}$ with $D > \operatorname{supp}(x)$, $|D| = m$ and set $y := x - P_A(x) + \alpha 1_D$ and $z := x + \alpha 1_D$, where $\alpha := \min_{n \in A} |e_n^*(x)|$. Then $D \in \mathcal{G}(y, m, \tau)$ and $A \in \mathcal{G}(z, m, \tau)$. Choose $CG_m^\tau \in \mathcal{CG}_m^\tau$ such that $\operatorname{supp}(CG_m^\tau(y)) \subset D$ and $\operatorname{supp}(CG_m^\tau(z)) \subset A$. We have

$$\|y - CG_m^\tau(y)\| = \left\| x - P_A(x) + \sum_{n \in D} a_n e_n \right\|,$$

for some scalars $(a_n) \subset \mathbb{F}$. Hence,

$$\begin{aligned}
\|x - P_A(x)\| & \leq \mathbf{K}_b \|y - CG_m^\tau(y)\| \leq \mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} \sigma_m(y) \\
& \leq \mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} \|x + \alpha 1_D\| \leq \mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} (\|x\| + \|\alpha 1_D\|). \tag{3.3}
\end{aligned}$$

We now bound $\|\alpha 1_D\|$:

$$\|z - CG_m^\tau(z)\| = \left\| \sum_{n \in A} b_n e_n + \sum_{n \notin A} e_n^*(x) e_n + \alpha 1_D \right\|,$$

for some scalars $(b_n) \subset \mathbb{F}$. Therefore,

$$\|\alpha 1_D\| \leq (\mathbf{K}_b + 1) \|z - CG_m^\tau(z)\| \leq (\mathbf{K}_b + 1) \mathbf{L}_{m,\tau}^{ch} \sigma_m(z) \leq (\mathbf{K}_b + 1) \mathbf{L}_{m,\tau}^{ch} \|x\|. \tag{3.4}$$

From (3.3) and (3.4), we conclude that

$$\|x - P_A(x)\| \leq \mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} (1 + \mathbf{L}_{m,\tau}^{ch} + \mathbf{L}_{m,\tau}^{ch} \mathbf{K}_b) \|x\|. \tag{3.5}$$

Hence, $g_{m,\tau}^c \leq \mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} (1 + \mathbf{L}_{m,\tau}^{ch} + \mathbf{L}_{m,\tau}^{ch} \mathbf{K}_b)$, which is (1.14). To obtain (1.15), we use the trivial estimate,

$$\mathbf{K}_b \mathbf{L}_{m,\tau}^{ch} (1 + \mathbf{L}_{m,\tau}^{ch} + \mathbf{L}_{m,\tau}^{ch} \mathbf{K}_b) \leq 3(\mathbf{K}_b \mathbf{L}_{m,\tau}^{ch})^2.$$

This completes our proof. \square

4. LEBESGUE CONSTANTS, PROPERTY (A, τ) , AND PARTIAL SYMMETRY

In this section, using our bounds for Lebesgue parameters, we generalize several existing theorems in the literature. We write “ \sup_m ” to indicate “ $\sup_{m \geq 0}$ ”.

4.1. Estimates on the τ -greedy constant (Generalization of [3, Theorem 3.4] and [10, Theorem 2]).

Definition 4.1. A basis \mathcal{B} is said to be (C, τ) -greedy if $C := C(\tau) \geq 1$ verifies

$$\sup_m \mathbf{L}_{m,\tau} \leq C < \infty. \quad (4.1)$$

The least constant satisfying (4.1) is denoted by $\mathbf{C}_{g,\tau}$. In particular,

$$\mathbf{C}_{g,\tau} = \sup_m \mathbf{L}_{m,\tau} < \infty.$$

Definition 4.2. A basis \mathcal{B} is said to be K -suppression unconditional if $K \geq 1$ verifies

$$\sup_m k_m \leq K < \infty. \quad (4.2)$$

The least constant satisfying (4.2) is denoted by \mathbf{K}_s ; that is,

$$\mathbf{K}_s = \sup_m k_m = \sup_m k_m^c < \infty.$$

Definition 4.3. A basis \mathcal{B} is said to have C -property (A, τ) if $C := C(\tau) \geq 1$ verifies

$$\sup_m \nu_{m,\tau} \leq C < \infty. \quad (4.3)$$

The least constant satisfying (4.3) is denoted by $\mathbf{C}_{b,\tau}$; specifically,

$$\mathbf{C}_{b,\tau} = \sup_m \nu_{m,\tau} < \infty.$$

Theorem 4.4. Let \mathcal{B} be an M -basis. The following hold.

- (1) If \mathcal{B} is $(\mathbf{C}_{g,\tau}, \tau)$ -greedy, then \mathcal{B} is $\mathbf{C}_{g,\tau}$ -suppression unconditional and has $\tau \mathbf{C}_{g,\tau}$ -property (A, τ) .
- (2) Conversely, if \mathcal{B} is \mathbf{K}_s -suppression unconditional and has $\mathbf{C}_{b,\tau}$ -property (A, τ) , then \mathcal{B} is $(\frac{\mathbf{K}_s \mathbf{C}_{b,\tau}}{\tau}, \tau)$ -greedy.

Remark 4.5. Observe that setting $\mathbf{C}_{g,\tau} = \tau = \mathbf{C}_{b,\tau} = \mathbf{K}_s = 1$ in Theorem 4.4 gives [3, Theorem 3.4]. Instead, if we set $\tau = 1$, we have [10, Theorem 2].

Proof of Theorem 4.4. (1) By (1.7), if \mathcal{B} is $(\mathbf{C}_{g,\tau}, \tau)$ -greedy,

$$\begin{aligned} \mathbf{C}_{g,\tau} &= \sup_m \mathbf{L}_{m,\tau} \geq \sup_m \frac{\nu_{m,\tau}}{\tau} = \frac{\mathbf{C}_{b,\tau}}{\tau}, \text{ and} \\ \mathbf{C}_{g,\tau} &= \sup_m \mathbf{L}_{m,\tau} \geq \sup_m k_m^c = \mathbf{K}_s. \end{aligned}$$

Therefore, \mathcal{B} is $\mathbf{C}_{g,\tau}$ -suppression unconditional and has $\tau \mathbf{C}_{g,\tau}$ -property (A, τ) .

(2) By (1.7), if \mathcal{B} is \mathbf{K}_s -suppression unconditional and has $\mathbf{C}_{b,\tau}$ -property (A, τ) ,

$$\sup_{\geq 1} \mathbf{L}_{m,\tau} \leq \sup_{m \geq 1} \frac{k_{2m-1}^c \nu_{m,\tau}}{\tau} \leq \frac{1}{\tau} \sup_{m \geq 1} k_{2m-1}^c \sup_{m \geq 1} \nu_{m,\tau} \leq \frac{\mathbf{K}_s \mathbf{C}_{b,\tau}}{\tau}.$$

Hence, \mathcal{B} is $(\frac{\mathbf{K}_s \mathbf{C}_{b,\tau}}{\tau}, \tau)$ -greedy. □

4.2. Estimates on the τ -almost greedy constant (Generalization of [1, Theorem 3.3]).

Definition 4.6. A basis is said to be (C, τ) -almost greedy if $C := C(\tau) \geq 1$ verifies

$$\sup_m \tilde{\mathbf{L}}_{m,\tau} \leq C < \infty. \quad (4.4)$$

The least constant C satisfying (4.4) is denoted by $\mathbf{C}_{a,\tau}$, which is equal to $\sup_m \tilde{\mathbf{L}}_{m,\tau}$.

Definition 4.7. A basis is said to be (C, τ) -quasi-greedy if $C := C(\tau) \geq 1$ verifies

$$\sup_m g_{m,\tau}^c \leq C < \infty. \quad (4.5)$$

The least constant C satisfying (4.5) is denoted by $\mathbf{C}_{\ell,\tau}$, which is equal to $\sup_m g_{m,\tau}^c$. Note that the term ‘‘suppression quasi-greedy’’ is also used when $\tau = 1$ (see Definition 5.1).

Theorem 4.8. *Let \mathcal{B} be an M -basis. The following hold.*

- (1) *If \mathcal{B} is $(\mathbf{C}_{a,\tau}, \tau)$ -almost greedy, then \mathcal{B} is $(\mathbf{C}_{a,\tau}, \tau)$ -quasi-greedy and has $\tau \mathbf{C}_{\alpha,\tau}$ -property (A, τ) .*
- (2) *Conversely, if \mathcal{B} is $(\mathbf{C}_{\ell,\tau}, \tau)$ -quasi-greedy and has $\mathbf{C}_{b,\tau}$ -property (A, τ) , then \mathcal{B} is $(\frac{\mathbf{C}_{\ell,\tau} \mathbf{C}_{b,\tau}}{\tau}, \tau)$ -almost greedy.*

Remark 4.9. Setting $\tau = 1$ in Theorem 4.8, we obtain [1, Theorem 3.3].

Proof of Theorem 4.8. (1) By (1.8), if \mathcal{B} is $(\mathbf{C}_{a,\tau}, \tau)$ -almost greedy,

$$\begin{aligned} \mathbf{C}_{a,\tau} &= \sup_m \tilde{\mathbf{L}}_{m,\tau} \geq \sup_m g_{m,\tau}^c = \mathbf{C}_{\ell,\tau}, \text{ and} \\ \mathbf{C}_{a,\tau} &= \sup_m \tilde{\mathbf{L}}_{m,\tau} \geq \sup_m \frac{\nu_{m,\tau}}{\tau} = \frac{\mathbf{C}_{b,\tau}}{\tau}. \end{aligned}$$

Hence, \mathcal{B} is $(\mathbf{C}_{a,\tau}, \tau)$ -quasi-greedy and has $\tau \mathbf{C}_{\alpha,\tau}$ -property (A, τ) .

(2) By (1.8), if \mathcal{B} is $(\mathbf{C}_{\ell,\tau}, \tau)$ -quasi-greedy and has $\mathbf{C}_{b,\tau}$ -property (A, τ) ,

$$\sup_{m \geq 1} \tilde{\mathbf{L}}_{m,\tau} \leq \sup_{m \geq 1} \frac{g_{m-1,\tau}^c \nu_{m,\tau}}{\tau} \leq \frac{1}{\tau} \sup_{m \geq 1} g_{m-1,\tau}^c \sup_{m \geq 1} \nu_{m,\tau} \leq \frac{\mathbf{C}_{\ell,\tau} \mathbf{C}_{b,\tau}}{\tau}.$$

Therefore, \mathcal{B} is $(\frac{\mathbf{C}_{\ell,\tau} \mathbf{C}_{b,\tau}}{\tau}, \tau)$ -almost greedy. \square

4.3. Estimates on the τ -partially greedy constant (Generalization of [15, Theorem 3.11]).

Definition 4.10. A basis \mathcal{B} is (C, τ) -partially greedy if $C := C(\tau) \geq 1$ verifies

$$\sup_m \mathbf{L}_{m,\tau}^{re} \leq C < \infty.$$

In this case, we let $\mathbf{C}_{p,\tau} := \sup_m \mathbf{L}_{m,\tau}^{re}$.

A basis \mathcal{B} is (C, τ) -strong partially greedy if $C := C(\tau) \geq 1$ verifies

$$\sup_m \hat{\mathbf{L}}_{m,\tau}^{re} \leq C < \infty.$$

We let $\mathbf{C}_{sp,\tau} := \sup_m \hat{\mathbf{L}}_{m,\tau}^{re}$.

The classical definition of partially greedy bases was due to Dilworth et al. [12], where they characterized such a basis as being quasi-greedy and conservative (defined later.)

Definition 4.11. A basis \mathcal{B} is said to have C -left property (A, τ) if $C := C(\tau) \geq 1$ verifies

$$\sup_m \nu_{m,\tau,\ell} \leq C < \infty.$$

In this case, we let $\mathbf{C}_{L,\tau} := \sup_m \nu_{m,\tau,\ell}$.

Definition 4.12. A basis \mathcal{B} is C -super-conservative if

$$\sup_{m \geq 1} \psi_m \leq C < \infty.$$

We let $\mathbf{C}_{sc} := \sup_{m \geq 1} \psi_m$.

Definition 4.13. A basis \mathcal{B} is said to be (C, τ) -partially symmetric for largest coefficients if $C := C(\tau) \geq 1$ verifies

$$\sup_m \nu'_{m,\tau,\ell} \leq C < \infty.$$

In this case, we let $\mathbf{C}_{pl,\tau} := \sup_m \nu'_{m,\tau,\ell}$.

In [15], Khurana characterized partially greedy Schauder bases by quasi-greediness and left-property (A), while in [4], Berná characterized strong partially greedy M-bases by partial symmetry for largest coefficients and quasi-greediness. We now generalize these results to the setting of τ -weak greedy sets.

Theorem 4.14. *Let \mathcal{B} be an M-basis.*

- (1) *If \mathcal{B} is $(\mathbf{C}_{sp,\tau}, \tau)$ -strong partially greedy, then \mathcal{B} is $(\mathbf{C}_{sp,\tau}, \tau)$ -quasi-greedy and is $(\tau \mathbf{C}_{sp,\tau}, \tau)$ -partially symmetric for largest coefficients.*
- (2) *Conversely, if \mathcal{B} is $(\mathbf{C}_{\ell,\tau}, \tau)$ -quasi-greedy and is $(\mathbf{C}_{pl,\tau}, \tau)$ -partially symmetric for largest coefficients, then \mathcal{B} is $(\frac{\mathbf{C}_{\ell,\tau} \mathbf{C}_{pl,\tau}}{\tau}, \tau)$ -strong partially greedy.*

Proof. (1) By (1.9) and (1.11), if \mathcal{B} is $(\mathbf{C}_{sp,\tau}, \tau)$ -strong partially greedy,

$$\begin{aligned} \sup_m g_{m,\tau}^c &\leq \sup_m \widehat{\mathbf{L}}_{m,\tau}^{re} = \mathbf{C}_{sp,\tau}, \\ \sup_m \frac{\nu'_{m,\tau,\ell}}{\tau} &\leq \sup_m \widehat{\mathbf{L}}_{m,\tau}^{re} = \mathbf{C}_{sp,\tau}. \end{aligned}$$

So, \mathcal{B} is $(\mathbf{C}_{sp,\tau}, \tau)$ -quasi-greedy and $(\tau \mathbf{C}_{sp,\tau}, \tau)$ -partially symmetric for largest coefficients.

(2) This is immediate from the upper bound of $\widehat{\mathbf{L}}_{m,\tau}^{re}$ in (1.11). \square

Theorem 4.15. *Let \mathcal{B} be an M-basis.*

- (1) *If \mathcal{B} is Schauder and $(\mathbf{C}_{p,\tau}, \tau)$ -partially greedy, then \mathcal{B} is $((\mathbf{K}_b + 1)\mathbf{C}_{p,\tau}, \tau)$ -quasi-greedy and has $\mathbf{C}_{L,\tau}$ -left property (A, τ) with*

$$\mathbf{C}_{L,\tau} \leq \tau \mathbf{C}_{p,\tau} (2 + \mathbf{K}_b + \mathbf{C}_{p,\tau} + \mathbf{C}_{p,\tau} \mathbf{K}_b). \quad (4.6)$$

- (2) *Conversely, if \mathcal{B} is $(\mathbf{C}_{\ell,\tau}, \tau)$ -quasi-greedy and has $(\mathbf{C}_{L,\tau}, \tau)$ -left property (A, τ) , then \mathcal{B} is $(\frac{\mathbf{C}_{\ell,\tau} \mathbf{C}_{L,\tau}}{\tau}, \tau)$ -partially greedy.*

Before proving Theorem 4.15, we need the following lemma.

Lemma 4.16. *Let \mathcal{B} be $(C_{\ell,\tau}, \tau)$ -quasi-greedy. Then \mathcal{B} is super-conservative if and only if it has left property (A, τ) . In particular,*

$$C_{sc} \leq C_{L,\tau} \leq \tau C_{\ell,\tau} + C_{sc} + C_{\ell,\tau} C_{sc}.$$

Proof. Since $\nu_{m,\tau,\ell} \geq \psi_m$ for all $m \geq 1$, if \mathcal{B} has left property (A), then it is super-conservative, and the left inequality is obvious. Assume that \mathcal{B} is C_{sc} -super-conservative. Let $x, A, B, (\varepsilon), (\delta)$ be chosen as in (1.3). We have

$$\begin{aligned} \|\tau x + 1_{\delta B}\| &\leq \tau \|x\| + \|1_{\delta B}\| \leq \tau C_{\ell,\tau} \|x + 1_{\varepsilon A}\| + C_{sc} \|1_{\varepsilon A}\| \\ &\leq (\tau C_{\ell,\tau} + C_{sc}(C_{\ell,\tau} + 1)) \|x + 1_{\varepsilon A}\|. \end{aligned}$$

Hence, $C_{L,\tau} \leq \tau C_{\ell,\tau} + C_{sc} + C_{\ell,\tau} C_{sc}$. \square

Proof of Theorem 4.15. (1) Assume that \mathcal{B} is $(C_{p,\tau}, \tau)$ -partially greedy. From (1.13), we know that \mathcal{B} is $(C_{p,\tau}(\mathbf{K}_b + 1), \tau)$ -quasi-greedy.

Fix $A, B, (\varepsilon), (\delta)$ as in (1.6). Let $y := 1_{\delta B} + 1_D + \tau 1_{\varepsilon A}$, where $D = \{1, \dots, \max B\} \setminus B$, then $D \cup A \in G(y, \max B, \tau)$. We have

$$\|1_{\delta B}\| = \|y - P_{D \cup A}(y)\| \leq C_{p,\tau} \|y - S_{\max B}(y)\| = C_{p,\tau} \tau \|1_{\varepsilon A}\|.$$

Hence, \mathcal{B} is $(C_{p,\tau} \tau)$ -super-conservative. By Lemma 4.16 and due to $(C_{p,\tau}(\mathbf{K}_b + 1), \tau)$ -quasi-greediness, we obtain (4.6).

(2) This is immediate from the upper bound of $L_{m,\tau}^{re}$ in (1.10). \square

5. PROPERTY (A, τ), UNIFORM PROPERTY (A), AND QUASI-GREEDY BASES

Let us retrieve some classical definitions when $\tau = 1$ in the definitions given in Section 4:

Definition 5.1.

- *Property (A):* If \mathcal{B} has $C_{b,1}$ -property (A, 1), we say \mathcal{B} has C_b -property (A) (here $C_b := C_{b,1}$) or property (A) if we do not want to specify C_b .
- *Greedy:* If \mathcal{B} is $(C_{g,1}, 1)$ -greedy, we say \mathcal{B} is C_g -greedy (here $C_g := C_{g,1}$.) We may also say \mathcal{B} is greedy without specifying C_g .
- *Almost greedy:* If \mathcal{B} is $(C_{a,1}, 1)$ -almost greedy, we say \mathcal{B} is C_a -almost greedy (here $C_a := C_{a,1}$) or is almost greedy without specifying the constant.
- *Quasi-greedy*¹: If \mathcal{B} is $(C_{\ell,1}, 1)$ -quasi-greedy, we say \mathcal{B} is C_ℓ -suppression quasi-greedy (here $C_\ell := C_{\ell,1}$) (or simply, is quasi-greedy). In this case, we may also say \mathcal{B} is quasi-greedy without specifying the constant. While $C_\ell = \sup_m g_{m,1}^c$, we let $C_w := \sup_m g_{m,1}$ and say \mathcal{B} is C_w -quasi-greedy. Clearly, $C_\ell - 1 \leq C_w \leq C_\ell + 1$.
- *Super-democratic:* \mathcal{B} is said to be C -super-democratic if $\sup_m \mu_m \leq C < \infty$. In this case, we let $C_{sd} := \sup_{m \geq 1} \mu_m$.

We now study the relation among property (A, τ) for different values of τ .

¹Wojtaszczyk [19] claimed that $\sup_m \|G_m\| < \infty$ is equivalent to the convergence of greedy algorithms $(G_m(x))_{m=1}^\infty$ to x for each $x \in \mathbb{X}$. Such an equivalence holds if and only if our M-basis is strong (see [7, Remark 6.2].)

5.1. Truncation operator. For each $\alpha > 0$, we define the truncation function T_α as follows: for $b \in \mathbb{F}$,

$$T_\alpha(b) = \begin{cases} \operatorname{sgn}(b)\alpha, & \text{if } |b| > \alpha, \\ b, & \text{if } |b| \leq \alpha. \end{cases}$$

We define the truncation operator $T_\alpha : \mathbb{X} \rightarrow \mathbb{X}$ as

$$T_\alpha(x) = \sum_{n=1}^{\infty} T_\alpha(e_n^*(x))e_n = \alpha 1_{\varepsilon\Gamma_\alpha(x)} + P_{\Gamma_\alpha^c(x)}(x),$$

where $\Gamma_\alpha(x) = \{n : |e_n^*(x)| > \alpha\}$ and $\varepsilon_n = \operatorname{sgn}(e_n^*(x))$ for all $n \in \Gamma_\alpha(x)$.

Theorem 5.2. [8, Lemma 2.5] *Let \mathcal{B} be \mathbf{C}_ℓ -suppression quasi-greedy. Then for any $\alpha > 0$, $\|T_\alpha\| \leq \mathbf{C}_\ell$.*

Proof. Simply observe that this lemma is a special of Proposition 2.5 when $\tau = 1$. \square

5.2. Relation among property (A, τ) for different τ .

Proposition 5.3. *Let \mathcal{B} be an M -basis. The following hold.*

- (1) *If \mathcal{B} is \mathbf{K}_s -suppression unconditional and has \mathbf{C}_b -property (A), then \mathcal{B} has $\mathbf{C}_b\mathbf{K}_s$ -property (A, τ) for all $\tau \in (0, 1]$.*
- (2) *Let $0 < \tau_2 < \tau_1 \leq 1$. If \mathcal{B} has \mathbf{C}_{b,τ_2} -property (A, τ_2), then \mathcal{B} has $\mathbf{C}_{b,\tau_2}\tau_1/\tau_2$ -property (A, τ_1).*

Proof. We prove (1). Assume that \mathcal{B} is \mathbf{K}_s -suppression unconditional and has \mathbf{C}_b -property (A). Choose $A, B, x, (\varepsilon), (\delta)$ as in (1.2). We have

$$\|x + 1_{\varepsilon A}\| \geq \frac{1}{\mathbf{K}_s} \|\tau x + 1_{\varepsilon A}\| \geq \frac{1}{\mathbf{C}_b\mathbf{K}_s} \|\tau x + 1_{\delta B}\|.$$

The first inequality is due to Proposition 2.1, while the second inequality is due to \mathbf{C}_b -property (A).

Next, we prove (2). Assume that \mathcal{B} has \mathbf{C}_{b,τ_2} -property (A, τ_2). Let us choose $A, B, x, (\varepsilon), (\delta)$ as in (1.2). Here $\|x\|_\infty \leq 1/\tau_1$. By \mathbf{C}_{b,τ_2} -property (A, τ_2), we have

$$\|x + 1_{\varepsilon A}\| \geq \frac{1}{\mathbf{C}_{b,\tau_2}} \|\tau_2 x + 1_{\theta B}\|, \forall \operatorname{signs}(\theta).$$

Hence,

$$\|x + 1_{\varepsilon A}\| \geq \frac{\tau_2}{\tau_1\mathbf{C}_{b,\tau_2}} \sup_{(\theta)} \left\| \tau_1 x + \frac{\tau_1}{\tau_2} 1_{\theta B} \right\| \geq \frac{\tau_2}{\tau_1\mathbf{C}_{b,\tau_2}} \|\tau_1 x + 1_{\delta B}\|,$$

where the last inequality is due to norm convexity. \square

In going from property (A) to property (A, τ) in the above proposition, we need unconditionality to “flatten out” the vector x so that $\|x\|_\infty \leq 1$, which makes it possible to apply property (A). Like unconditionality, quasi-greediness can obtain the same result thanks to the boundedness of the truncation operator T_1 (see Theorem 5.2.) We can therefore relax unconditionality in Proposition 5.3 as follows:

Proposition 5.4. *Let $\tau \in (0, 1]$. If an M -basis \mathcal{B} is \mathbf{C}_ℓ -suppression quasi-greedy and has $\mathbf{C}_{b,\tau}$ -property (A, τ), then \mathcal{B} has $2\mathbf{C}_{b,\tau}^2\mathbf{C}_\ell$ -property (A, γ) for all $\gamma \in (0, 1]$.*

Proof. Assume that our basis \mathcal{B} has $\mathbf{C}_{b,\tau}$ -property (A, τ) and is \mathbf{C}_ℓ -suppression quasi-greedy. Fix $\gamma \in (0, 1]$. Let $x, A, B, (\varepsilon), (\delta)$ be chosen as in (1.2), where $x \in \mathbb{X}_c$ and $\|x\|_\infty \leq 1/\gamma$. Let $E = \{n : |e_n^*(x)| \geq 1\}$ and T_1 be the 1-truncation operator. We shall show that

$$\|x + 1_{\varepsilon A}\| \geq \frac{1}{2\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\gamma x + 1_{\delta B}\|. \quad (5.1)$$

Case 1: $\gamma \neq \tau^2$. Write

$$\|x + 1_{\varepsilon A}\| = \|P_E x + 1_{\varepsilon A} + P_E^c x\| \text{ and } \|\gamma x + 1_{\delta B}\| = \|\gamma P_E x + \gamma P_E^c x + 1_{\delta B}\|. \quad (5.2)$$

Since $P_E x + 1_{\varepsilon A}$ is a greedy sum of $x + 1_{\varepsilon A}$, we have

$$\|x + 1_{\varepsilon A}\| \geq \frac{1}{\mathbf{C}_\ell} \|P_E^c x\| = \frac{1}{|\gamma - \tau^2| \mathbf{C}_\ell} \|(\gamma - \tau^2) P_E^c x\| \geq \frac{1}{\mathbf{C}_\ell} \|(\gamma - \tau^2) P_E^c x\|. \quad (5.3)$$

Choose a sign (ψ) such that $\psi_n = \text{sgn}(e_n^*(x))$ for all $n \in \text{supp}(x)$. Choose $D \subset \mathbb{N}$ with $\max(A \cup B \cup \text{supp}(x)) < \min D$ and $|D| = |A| + |E|$. By Theorem 5.2 and $\mathbf{C}_{b,\tau}$ -property (A, τ), we obtain

$$\begin{aligned} \|x + 1_{\varepsilon A}\| &\geq \frac{1}{\mathbf{C}_\ell} \|1_{\psi E} + 1_{\varepsilon A} + P_E^c x\| \geq \frac{1}{\mathbf{C}_\ell \mathbf{C}_{b,\tau}} \|\tau P_E^c x + 1_D\| \\ &\geq \frac{1}{\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \sup_{(\theta)} \|\tau^2 P_E^c x + 1_{\theta E} + 1_{\delta B}\| \\ &\geq \frac{1}{\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\tau^2 P_E^c x + \gamma P_E x + 1_{\delta B}\|. \end{aligned} \quad (5.4)$$

where the last inequality is due to norm convexity. It follows from (5.2), (5.3), and (5.4) that

$$\begin{aligned} \|x + 1_{\varepsilon A}\| &\geq \frac{1}{2\mathbf{C}_\ell} \|(\gamma - \tau^2) P_E^c x\| + \frac{1}{2\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\tau^2 P_E^c x + 1_{\delta B} + \gamma P_E x\| \\ &\geq \frac{1}{2\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\gamma P_E^c x + \gamma P_E x + 1_{\delta B}\| \\ &= \frac{1}{2\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\gamma x + 1_{\delta B}\|, \end{aligned}$$

which is (5.1), as desired.

Case 2: $\gamma \neq \tau^2$. Simply use (5.4) to get $\|x + 1_{\varepsilon A}\| \geq \frac{1}{\mathbf{C}_\ell \mathbf{C}_{b,\tau}^2} \|\gamma x + 1_{\delta B}\|$. \square

Corollary 5.5. *If an M -basis \mathcal{B} has 1-property (A), then for all $\tau \in (0, 1)$, \mathcal{B} has 2-property (A, τ).*

Proof. Let $\tau \in (0, 1)$. Assume \mathcal{B} has 1-property (A). By [1, Proposition 2.5], \mathcal{B} is 1-suppression quasi-greedy. Plugging in $\mathbf{C}_\ell = \tau_1 = \mathbf{C}_{b,\tau_1} = 1$ into Proposition 5.4, we know that \mathcal{B} has 2-property (A, τ). \square

Proposition 5.3 item (2) suggests that we may need to increase the property (A, τ) constant when τ increases from τ_2 to τ_1 . On the other hand, an interesting feature of Proposition 5.3 item (1), Proposition 5.4, and Corollary 5.5 is that there exists a constant C (independent of τ) such that \mathcal{B} has C -property (A, τ) for all $\tau \in (0, 1]$. Such a basis is said to have *uniform property (A)*.

The following corollary is immediate from Proposition 5.4.

Corollary 5.6. *Fix $\tau \in (0, 1]$. If an M -basis \mathcal{B} has property (A, τ) and is quasi-greedy, then \mathcal{B} has uniform property (A).*

A natural question is whether the converse of Corollary 5.6 holds; that is, whether uniform property (A) implies quasi-greediness. We know property (A) alone does not imply quasi-greediness (see [8, Proposition 5.7].) Below we construct a basis that has uniform property (A) but is not quasi-greedy. Our construction is somewhat simpler than [8, Proposition 5.7] since we do not attempt to optimize various parameters.

Theorem 5.7. *There exists a Banach space \mathbb{X} with a monotone ($\mathbf{K}_b = 1$) Schauder basis that is non-quasi-greedy and has uniform property (A).*

The next subsection establishes Theorem 5.7.

5.3. Non-quasi-greedy basis with uniform property (A).

Proposition 5.8. *Let \mathbb{X} be the completion of c_{00} with respect to the norm $\|\cdot\| = \max\{\|\cdot\|_1, \|\cdot\|_{\ell_2}\}$, where $\|\cdot\|_1$ is a semi-norm on c_{00} . Assume $(\mathbb{X}, \|\cdot\|)$ has a basis \mathcal{B} . If there exists $\lambda \geq 1$ such that for all signs (δ) and nonempty, finite sets $A \subset \mathbb{N}$, we have $\|1_{\delta A}\|_1 \leq \lambda|A|^{1/2}$, then \mathcal{B} has (3λ) -property (A, τ) for all $\tau \in (0, 1]$ (with respect to $\|\cdot\|$.)*

Proof. Let $x \in c_{00}$ and $A, B \subset \mathbb{N}$ be finite with $|A| = |B| = n$ and $A \sqcup B \sqcup x$. Pick signs $(\varepsilon), (\delta)$. We want to show that $\|\tau x + 1_{\delta B}\| \leq 3\lambda\|x + 1_{\varepsilon A}\|$. Assume, for a contradiction, that $\|\tau x + 1_{\delta B}\| > 3\lambda\|x + 1_{\varepsilon A}\|$. Since $\|\tau x + 1_{\delta B}\|_{\ell_2} \leq \|x + 1_{\varepsilon A}\|_{\ell_2}$, it must be that $\|\tau x + 1_{\delta B}\| = \|\tau x + 1_{\delta B}\|_1$. We have

$$\begin{aligned} \|x + 1_{\varepsilon A}\|_1 &\geq \|x\|_1 - \|1_{\varepsilon A}\|_1 \geq \|x\|_1 - \lambda\sqrt{n} \\ &\geq \|\tau x\|_1 + \|1_{\delta B}\|_1 - 2\lambda\sqrt{n} \geq \|\tau x + 1_{\delta B}\| - 2\lambda\sqrt{n}. \end{aligned}$$

Therefore,

$$\|\tau x + 1_{\delta B}\| = \|\tau x + 1_{\delta B}\|_1 > 3\lambda\|x + 1_{\varepsilon A}\|_1 \geq 3(\|\tau x + 1_{\delta B}\| - 2\lambda\sqrt{n}),$$

which gives

$$\|\tau x + 1_{\delta B}\| < 3\lambda\sqrt{n}. \quad (5.5)$$

On the other hand, by assumption,

$$\|\tau x + 1_{\delta B}\| > 3\lambda\|x + 1_{\varepsilon A}\|_{\ell_2} \geq 3\lambda\sqrt{n},$$

contradicting (5.5). \square

5.3.1. Construct the weight sequence. For $n \geq 1$, let $t_n = \frac{1}{\sqrt{n}}$, $L_n = e^{(\log n)^2}$ and $a_n = \frac{1}{\sqrt{n} \log(n+1)}$. Define an increasing sequence $(N_n)_{n=0}^\infty$ recursively. Set $N_0 = 0$. Choose $N_1 > 10$ to be the smallest such that

$$b_1 := a_1 t_1 \left(\sum_{n=1}^{N_1} t_n \right)^{-1} < \frac{a_1}{L_1}.$$

Once N_j and b_j are defined for $j \geq 1$, choose N_{j+1} to be the smallest such that

$$b_{j+1} := a_{j+1} t_{j+1} \left(\sum_{n=N_j+1}^{N_j+N_{j+1}} t_n \right)^{-1} < \min \left\{ \frac{a_{j+1}}{L_{j+1}}, b_j \right\} \text{ and } \frac{N_{j+1}}{N_j} > 10. \quad (5.6)$$

For $j \geq 1$, denote the finite sequence $(t_n)_{n=N_{j-1}+1}^{N_j-1+N_j} = (t_{N_{j-1}+1}, \dots, t_{N_{j-1}+N_j})$ by B_j .

We now define weight $\omega := (w_n)_{n=1}^\infty$ on \mathbb{N} as follows:

$$(w_n) = (t_1, B_1, t_2, B_2, t_3, B_3, t_4, \dots).$$

In words, the weight (w_n) is chosen such that the first one is t_1 ; the next N_1 weights are taken from B_1 in the same order; the next weight is t_2 , then the next N_2 weights are taken from B_2 in the same order, and so on.

5.3.2. Construct the desired space and verify its properties. Let \mathbb{X} be the completion of c_{00} under the following norm

$$\left\| \sum_n x_n e_n \right\| = \max \left\{ \sup_N \left| \sum_{n=N}^\infty w_n x_n \right|, \left(\sum_{n=1}^\infty |x_n|^2 \right)^{1/2} \right\}.$$

For ease of notation, set

$$\left\| \sum_n x_n e_n \right\|_1 = \sup_N \left| \sum_{n=N}^\infty w_n x_n \right| \text{ and } \left\| \sum_n x_n e_n \right\|_2 = \left(\sum_{n=1}^\infty |x_n|^2 \right)^{1/2}.$$

Clearly, the standard unit vector basis \mathcal{B} is Schauder and normalized in $(\mathbb{X}, \|\cdot\|)$.

Proposition 5.9. *The basis \mathcal{B} has uniform property (A).*

Proof. It suffices to verify that the condition in Proposition 5.8 is satisfied. Choose $A \subset \mathbb{N}$ with $|A| = m$ and sign (δ) . By construction, each value of t_n appears exactly twice in ω . Hence,

$$\|1_{\delta A}\|_1 < 2 \sum_{n=1}^m t_n = 2 + 2 \sum_{n=2}^m \frac{1}{\sqrt{n}} \leq 2 + 2 \int_1^m \frac{dx}{\sqrt{x}} = 2\sqrt{m},$$

as desired. \square

Proposition 5.10. *The basis \mathcal{B} is not quasi-greedy.*

Lemma 5.11. *The following vector is in \mathbb{X} :*

$$x = (a_1, \underbrace{-b_1, \dots, -b_1}_{N_1 \text{ times}}, a_2, \underbrace{-b_2, \dots, -b_2}_{N_2 \text{ times}}, \dots, a_j, \underbrace{-b_j, \dots, -b_j}_{N_j \text{ times}}, \dots).$$

Proof. Let S_N be the partial sum of order N and $T_N = I - S_N$ be the N th tail operator. Fix $\varepsilon > 0$. Call $(a_j, \underbrace{-b_j, \dots, -b_j}_{N_j \text{ times}})$ the j th block. Pick k such that $a_k t_k = \frac{1}{k \log(k+1)} < \varepsilon$.

Let N_0 be sufficiently large such that $\min \text{supp}(T_{N_0}(x))$ does not lie before the k th block. By construction of (w_n) , (a_n) , and (b_n) , if a block fully appears in a tail, then it contributes nothing to the norm $\|\cdot\|_1$. Hence, for all $N \geq N_0$, $\|T_N x\|_1 \leq a_k t_k < \varepsilon$, and we have $S_N(x) \rightarrow x$ in $\|\cdot\|_1$.

Next, we show that $\|x\|_2 < \infty$. Observe that

$$\sum_{n=1}^{\infty} (a_n)^2 = \sum_{n=1}^{\infty} \frac{1}{n(\log(n+1))^2} < 4.$$

We also have

$$\begin{aligned} \sum_{n=1}^{\infty} N_n (b_n)^2 &= \sum_{n=1}^{\infty} N_n a_n^2 t_n^2 \left(\sum_{j=N_{n-1}+1}^{N_{n-1}+N_n} \frac{1}{\sqrt{j}} \right)^{-2} \leq \sum_{n=1}^{\infty} N_n a_n^2 t_n^2 \left(\int_{N_{n-1}+1}^{N_{n-1}+N_n+1} \frac{dx}{\sqrt{x}} \right)^{-2} \\ &\leq \sum_{n=1}^{\infty} \frac{1}{n^2 \log^2(n+1) (\sqrt{N_{n-1}/N_n + 1} + 1/N_n - \sqrt{N_{n-1}/N_n + 1/N_n})^2} \\ &\leq 3 \sum_{n=1}^{\infty} \frac{1}{n^2 \log^2(n+1)}, \end{aligned}$$

where the last inequality is due to $1/N_n + N_{n-1}/N_n < 0.2$. We have shown that $\|x\|_2 < \infty$ and so, $x \in \mathbb{X}$. \square

Proof of Proposition 5.10. Consider the vector x as in Lemma 5.11:

$$x = (a_1, \underbrace{-b_1, \dots, -b_1}_{N_1 \text{ times}}, a_2, \underbrace{-b_2, \dots, -b_2}_{N_2 \text{ times}}, \dots, a_j, \underbrace{-b_j, \dots, -b_j}_{N_j \text{ times}}, \dots).$$

Let $k \in \mathbb{N}$. Pick $\varepsilon \in (b_{k+1}, b_k)$ and consider $T_\varepsilon(x)$, where T_ε is the thresholding operator. In particular, for each $x = \sum_n x_n e_n \in \mathbb{X}$,

$$T_\varepsilon(x) := \sum_{n: |x_n| > \varepsilon} x_n e_n.$$

Since (b_n) is strictly decreasing, $T_\varepsilon(x)$ does not contain any coefficient b_n for $n \geq k+1$; that is, from the $(k+1)$ th block onward, only certain values of a_n at the start of the n th block appear in $T_\varepsilon(x)$. Specifically, let us establish a range of n such that a_n still appears in the n th block. By construction,

$$\varepsilon < b_k < \frac{a_k}{L_k} = \frac{1}{(\log(k+1))\sqrt{k}e^{(\log(k))^2}}.$$

We want to find n such that

$$a_n > \frac{1}{(\log k)\sqrt{k}e^{(\log(k))^2}} > \varepsilon.$$

Equivalently,

$$\frac{1}{\sqrt{n} \log(n+1)} > \frac{1}{(\log k)\sqrt{k}e^{(\log(k))^2}}.$$

It is easy to check that for sufficiently large k , we have $n \leq \frac{1}{2}e^{(\log(k))^2/2}$ work. Therefore,

$$\begin{aligned} \|T_\varepsilon(x)\|_1 &\geq \sum_{n=k+1}^{\frac{1}{2}e^{(\log(k))^2/2}} t_n a_n = \sum_{n=k+1}^{\frac{1}{2}e^{(\log(k))^2/2}} \frac{1}{n \log(n+1)} \geq \frac{1}{2} \int_{k+1}^{\frac{1}{2}e^{(\log(k))^2/2}} \frac{dx}{x \log x} \\ &= \frac{1}{2} \left(\log \log \left(\frac{1}{2}e^{(\log(k))^2/2} \right) - \log \log(k+1) \right) \rightarrow \infty \end{aligned}$$

as $k \rightarrow \infty$. Therefore, \mathcal{B} is not quasi-greedy. \square

5.4. Yet another characterization of property (A). Lemma 2.2 gives a characterization of property (A, τ) for M -bases. We now give another characterization of property (A, τ), which in turn gives a seemingly new characterization of property (A).

Lemma 5.12. *If a basis \mathcal{B} has $\mathbf{C}_{b,\tau}$ -property (A, τ), then \mathcal{B} is $\min\{2\mathbf{C}_{b,\tau}/\tau, \mathbf{C}_{b,\tau}^2\}$ -super-democratic.*

Proof. By Proposition 2.1 item (2), \mathcal{B} has $\mathbf{C}_{b,\tau}/\tau$ -property (A). Let $A, B, (\varepsilon), (\delta)$ be chosen as in (1.5). Without loss of generality, we can assume that $\varepsilon \equiv 1$. By [9, Lemma 6.4], we know that $1_{\delta B} \in 2S$, where $S = \{\sum_{D \subset B} a_D 1_D : (a_D) \subset \mathbb{R}, \sum_{D \subset B} |a_D| \leq 1\}$. We shall show that

$$\|1_D\| \leq \frac{\mathbf{C}_{b,\tau}}{\tau} \|1_A\|, \forall D \subset B.$$

We have

$$\|1_D\| = \|1_{D \setminus A} + 1_{D \cap A}\| \leq \frac{\mathbf{C}_{b,\tau}}{\tau} \|1_{A \setminus D} + 1_{D \cap A}\| = \frac{\mathbf{C}_{b,\tau}}{\tau} \|1_A\|.$$

The inequality is due to $\mathbf{C}_{b,\tau}/\tau$ -property (A) and the fact that $|D \setminus A| \leq |A \setminus D|$. Therefore, \mathcal{B} is $2\mathbf{C}_{b,\tau}/\tau$ -super-democratic.

To see that \mathcal{B} is $\mathbf{C}_{b,\tau}^2$ -super-democratic, pick any set $E \subset \mathbb{N}$ with $|E| = |A| = |B|$ and $E > A \cup B$. By $\mathbf{C}_{b,\tau}$ -property (A, τ), we know that $\|1_{\delta B}\| \leq \mathbf{C}_{b,\tau} \|1_E\|$ and $\|1_E\| \leq \mathbf{C}_{b,\tau} \|1_{\varepsilon A}\|$. Hence,

$$\|1_{\delta B}\| \leq \mathbf{C}_{b,\tau}^2 \|1_{\varepsilon A}\|.$$

\square

Theorem 5.13. *For an M -basis \mathcal{B} , the following are equivalent:*

- (1) \mathcal{B} has property (A, τ).
- (2) There exists a function $f : \mathbb{N} \rightarrow \mathbb{R}_+$ and constants $c_1, c_2 > 0$ such that

$$\|1_{\varepsilon \Lambda}\| \leq c_1 f(|\Lambda|) \text{ and } \|1_{\varepsilon \Lambda} + x\| \geq c_2 f(|\Lambda|), \quad (5.7)$$

for any $x \in \mathbb{X}$ with $\|x\|_\infty \leq 1/\tau$, any nonempty, finite set $\Lambda \subset \mathbb{N}$ with $\text{supp}(x) \cap \Lambda = \emptyset$, and sign (ε) . Furthermore, f is unique in the following sense: if g is another function verifying (5.7), then $0 < \inf g(n)/f(n) \leq \sup g(n)/f(n) < \infty$.

Proof. (1) \Rightarrow (2): Assume that \mathcal{B} has $\mathbf{C}_{b,\tau}$ -property (A, τ). Define $f : \mathbb{N} \rightarrow \mathbb{R}_+$ as follows:

$$f(n) := \sup_{(\varepsilon), \Lambda \subset \mathbb{N}, |\Lambda|=n} \|1_{\varepsilon \Lambda}\|.$$

We show that f verifies (5.7). Fix a nonempty, finite set $A \subset \mathbb{N}$, $\text{sign}(\varepsilon)$, and $x \in \mathbb{X}_c$ with $\|x\|_\infty \leq 1/\tau$ and $\text{supp}(x) \cap A = \emptyset$. By definition, $\|1_{\varepsilon A}\| \leq f(|A|)$. We have

$$\|1_{\varepsilon A}\| \leq \frac{1}{2}(\|1_{\varepsilon A} + x\| + \|1_{\varepsilon A} - x\|). \quad (5.8)$$

Choose $B \supset A \cup \text{supp}(x)$ with $|B| = |A|$. Then

$$\|1_{\varepsilon A} - x\| \leq \mathbf{C}_{b,\tau} \|1_B + x\| \leq \mathbf{C}_{b,\tau}^2 \|1_{\varepsilon A} + x\|. \quad (5.9)$$

By (5.8), (5.9), and Lemma 5.12, we obtain

$$\|1_{\varepsilon A} + x\| \geq \frac{2}{1 + \mathbf{C}_{b,\tau}^2} \|1_{\varepsilon A}\| \geq \frac{2}{\mathbf{C}_{b,\tau}^2(1 + \mathbf{C}_{b,\tau}^2)} f(|A|).$$

Setting $c_2 = \frac{2}{\mathbf{C}_{b,\tau}^2(1 + \mathbf{C}_{b,\tau}^2)}$, we are done.

(2) \Rightarrow (1): Let $x, A, B, (\varepsilon), (\delta)$ be chosen as in (1.2).

Case 1: $\|x\| \geq 2c_1 f(|A|)$. Then $\|1_{\varepsilon A}\|, \|1_{\delta B}\| \leq \|x\|/2$. We have

$$\frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} \leq \frac{\tau \|x\| + \|1_{\delta B}\|}{\|x\| - \|1_{\varepsilon A}\|} \leq \frac{(\tau + 1/2)\|x\|}{\|x\|/2} = 2\tau + 1.$$

Case 2: $\|x\| < 2c_1 f(|A|)$. We have

$$\frac{\|\tau x + 1_{\delta B}\|}{\|x + 1_{\varepsilon A}\|} \leq \frac{\tau \|x\| + \|1_{\delta B}\|}{c_2 f(|A|)} < \frac{2c_1 \tau f(|B|) + c_1 f(|B|)}{c_2 f(|A|)} = \frac{(2\tau + 1)c_1}{c_2}.$$

Therefore, \mathcal{B} has $(2\tau + 1)c_1/c_2$ -property (A, τ) .

Finally, we prove that if g is another function verifying (5.7), then

$$0 < \inf \frac{f(n)}{g(n)} \leq \sup \frac{f(n)}{g(n)} < \infty.$$

By symmetry, it suffices to prove that $\inf f(n)/g(n) > 0$. Suppose not. Let $c_3 > 0$ be such that

$$\|1_{\varepsilon A} + x\| \geq c_3 g(|A|),$$

for any $x \in \mathbb{X}$ with $\|x\|_\infty \leq 1/\tau$, any finite set $A \subset \mathbb{N}$ with $\text{supp}(x) \cap A = \emptyset$, and $\text{sign}(\varepsilon)$. Let N be chosen such that $f(N)/g(N) < c_3$. Choose $B \subset \mathbb{N}$ with $|B| = N$. By construction of f , we have $\|1_B\| \leq f(N)$. On the other hand, $\|1_B\| \geq c_3 g(N)$, and so, $f(N) \geq c_3 g(N)$, which contradicts our choice of N . \square

6. FROM CLASSICAL GREEDY-TYPE BASES TO WEAK GREEDY-TYPE BASES

It is obvious that a (C, τ) -(quasi, almost) greedy basis is (quasi, almost) greedy (see Definition 5.1.) Thanks to the work of Konyagin and Temlyakov [16], we know that the converse is true as well; that is, for any fixed $\tau \in (0, 1]$, a C_1 -(quasi, almost) greedy basis is (C_2, τ) -(quasi, almost) greedy for C_2 dependent on τ and C_1 . For example, if a basis is \mathbf{C}_w -quasi-greedy, then $\mathbf{C}_{\ell,\tau}$ can be bounded above by a quantity involving \mathbf{C}_w and τ . Corresponding results hold for $\mathbf{C}_{a,\tau}$ and $\mathbf{C}_{g,\tau}$. In this section, we shall compute and improve these upper bounds.

6.1. From quasi-greedy to $(C_{\ell, \tau}, \tau)$ -quasi-greedy.

Theorem 6.1 (Konyagin and Temlyakov [16]). *Let \mathcal{B} be a C_w -quasi-greedy M -basis. Then for $\tau \in (0, 1]$, we have*

$$\gamma_{m, \tau}(x) \leq C(C_w, \tau) \|x\|, \forall x \in \mathbb{X}, \forall m \in \mathbb{N}. \quad (6.1)$$

That is, \mathcal{B} is $(C(C_w, \tau), \tau)$ -quasi-greedy.

The constant $C(C_w, \tau)$ in Theorem 6.1 was shown to be of order $O(C_w^5/\tau)$. We reduce the order of $C(C_w, \tau)$ to $O(C_w^4/\tau)$. The improvement comes from the following estimate. For its proof, see [6, Lemma 6.1].

Lemma 6.2. *Let \mathcal{B} be a C_w -quasi-greedy basis. Fix $\tau \in (0, 1]$ and $x \in \mathbb{X}$. Let A_1 and A_2 be finite sets of positive integers such that $A_1 \subset A_2$ and for all $n \in A_2$, we have $\tau \leq |e_n^*(x)| \leq 1$. Then*

$$\|P_{A_1}x\| \leq \frac{8C_w^3}{\tau} \|P_{A_2}x\|.$$

Proof of Theorem 6.1. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$, and $A \in \mathcal{G}(x, m, \tau)$. Set $\alpha := \max_{n \notin A} |e_n^*(x)|$. The idea is to sandwich our τ -weak greedy set A between two greedy sets. If $\alpha = 0$, then $x - P_A(x) = 0$ and there is nothing to prove. Assume $\alpha > 0$. Define two sets

$$A_1 = \{n : |e_n^*(x)| > \alpha\} \text{ and } A_2 = \{n : |e_n^*(x)| \geq \tau\alpha\}.$$

Clearly, $A \subset A_2$, and since $A_1 \cap A^c = \emptyset$, we know that $A_1 \subset A$. We write

$$P_A(x) = P_{A_1}(x) + P_A(P_{A_2 \setminus A_1}(x)). \quad (6.2)$$

Since A_1 is a greedy set of x , by C_w -quasi-greediness, we obtain

$$\|P_{A_1}(x)\| \leq C_w \|x\|. \quad (6.3)$$

For each $n \in A_2 \setminus A_1$, we have $\tau \leq \frac{|e_n^*(x)|}{\alpha} \leq 1$. Due to Lemma 6.2, we obtain

$$\begin{aligned} \|P_A(P_{A_2 \setminus A_1}(x))\| &\leq \frac{8C_w^3}{\tau} \|P_{A_2 \setminus A_1}(x)\| = \frac{8C_w^3}{\tau} \|P_{A_2}(x) - P_{A_1}(x)\| \\ &\leq \frac{8C_w^3}{\tau} (\|P_{A_2}(x)\| + \|P_{A_1}(x)\|) \leq \frac{16C_w^4}{\tau} \|x\|. \end{aligned} \quad (6.4)$$

Together, (6.2), (6.3), and (6.4) give

$$\|P_A(x)\| \leq \left(C_w + \frac{16C_w^4}{\tau} \right) \|x\|.$$

Therefore, $\|x - P_A(x)\| \leq \left(1 + C_w + \frac{16C_w^4}{\tau} \right) \|x\|$. \square

6.2. From almost greedy to $(C_{a, \tau}, \tau)$ -almost greedy. The improvement in the upper bound in Theorem 6.1 leads to an improvement for the theorem below, which shows that an almost greedy basis is $(C_{a, \tau}, \tau)$ -almost greedy for any fixed $\tau \in (0, 1]$. First, we state a useful estimate.

Lemma 6.3. *Suppose that \mathcal{B} is C_w -quasi-greedy. Let $x \in \mathbb{X}$. For finite sets $A, B \subset \mathbb{N}$ with $A \subset B$, $(a_n)_{n \in A} \subset \mathbb{F}$, and any sign (ε) , we have that*

(1)

$$\left\| \sum_{n \in A} a_n e_n \right\| \leq 2\mathbf{C}_w \max_{n \in A} |a_n| \|1_{\varepsilon A}\|. \quad (6.5)$$

(2) if A is a greedy set of x , then

$$\min_{n \in A} |e_n^*(x)| \left\| \sum_{n \in A} \delta_n e_n \right\| \leq 2 \min\{\mathbf{C}_\ell, \mathbf{C}_w\} \|x\|, \quad (6.6)$$

where $\delta_n = \text{sgn}(e_n^*(x))$.*Proof.* For (6.5), see [2, Corollary 10.2.11]; for (6.6), see [8, Lemma 2.3]. \square

Theorem 6.4 (Konyagin and Temlyakov [16]). *Let \mathcal{B} be a \mathbf{C}_a -almost greedy M -basis. Then for $\tau \in (0, 1]$, we have*

$$\gamma_{m,\tau}(x) \leq C(\mathbf{C}_a, \tau) \tilde{\sigma}_m(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}. \quad (6.7)$$

Following the proof by Konyagin and Temlyakov, we see that $\mathbf{C}_{a,\tau} \lesssim \mathbf{C}_a^9/\tau^2$. As a byproduct of Theorem 6.1 and Lemma 6.3, we shall show that $\mathbf{C}_{a,\tau} \lesssim \mathbf{C}_a^7/\tau^2$.

Proof. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$, and $\varepsilon > 0$. Let $A \subset \mathbb{N}$ be such that $|A| = m$ and $\|x - P_A x\| \leq \tilde{\sigma}_m(x) + \varepsilon$. Pick $B \in \mathcal{G}(x, m, \tau)$. By the triangle inequality, we have

$$\begin{aligned} \|x - P_B(x)\| &\leq \|x - P_A(x)\| + \|P_A(x) - P_B(x)\| \\ &= \|x - P_A(x)\| + \|P_{A \setminus B}(x) - P_{B \setminus A}(x)\| \\ &\leq \|x - P_A(x)\| + \|P_{A \setminus B}(x)\| + \|P_{B \setminus A}(x)\|. \end{aligned} \quad (6.8)$$

Note that $|A \setminus B| = |B \setminus A|$. If $|B \setminus A| = 0$, then $\|x - P_B(x)\| = \|x - P_A(x)\| \leq \tilde{\sigma}_m(x) + \varepsilon$. Assume that $|B \setminus A| \neq 0$. We bound $\|P_{B \setminus A}(x)\|$. Set $y := x - P_A(x)$. Then $B \setminus A \in \mathcal{G}(y, |B \setminus A|, \tau)$. Since a \mathbf{C}_a -almost greedy basis is \mathbf{C}_a -suppression quasi-greedy (by Theorem 4.8), we can use Theorem 6.1 to obtain

$$\|P_{B \setminus A}(x)\| = \|P_{B \setminus A}(y)\| \leq C_1(\mathbf{C}_a, \tau) \|y\| = C_1(\mathbf{C}_a, \tau) \|x - P_A(x)\|, \quad (6.9)$$

where $C_1(\mathbf{C}_a, \tau) \lesssim \mathbf{C}_a^4/\tau$.

Next, we bound $\|P_{A \setminus B}(x)\|$. By Theorem 4.8, our basis has \mathbf{C}_a -property (A). Observe that

$$\tau \max_{n \in A \setminus B} |e_n^*(x)| \leq \tau \max_{n \notin B} |e_n^*(x)| \leq \min_{n \in B} |e_n^*(x)| \leq \min_{n \in B \setminus A} |e_n^*(x)|.$$

Let $\varepsilon_n = \text{sgn}(e_n^*(x))$ for all $n \in \text{supp}(x)$. We have

$$\begin{aligned}
 \|P_{A \setminus B}(x)\| &\leq 2(\mathbf{C}_a + 1) \max_{n \in A \setminus B} |e_n^*(x)| \left\| \sum_{n \in A \setminus B} \varepsilon_n e_n \right\| \text{ due to (6.5)} \\
 &\leq \frac{2(\mathbf{C}_a + 1)\mathbf{C}_a}{\tau} \min_{n \in B \setminus A} |e_n^*(x)| \left\| \sum_{n \in B \setminus A} \varepsilon_n e_n \right\| \text{ due to } \mathbf{C}_a\text{-property (A)} \\
 &\leq \frac{4(\mathbf{C}_a + 1)\mathbf{C}_a^2}{\tau} \|P_{B \setminus A}(x)\| \text{ due to (6.6)} \\
 &\leq \frac{4(\mathbf{C}_a + 1)\mathbf{C}_a^2}{\tau} C_1(\mathbf{C}_a, \tau) \|x - P_A(x)\| \lesssim \frac{\mathbf{C}_a^7}{\tau^2} \|x - P_A(x)\|. \quad (6.10)
 \end{aligned}$$

From (6.8), (6.9), and (6.10), we obtain

$$\|x - P_B(x)\| \lesssim \frac{\mathbf{C}_a^7}{\tau^2} (\tilde{\sigma}_m(x) + \varepsilon).$$

Letting $\varepsilon \rightarrow 0$, we are done. \square

6.3. From greedy to $(\mathbf{C}_{g,\tau}, \tau)$ -greedy bases. Temlyakov [17] proved that when $\mathbb{X} = L_p$, $1 < p < \infty$, the normalized Haar basis is $(\mathbf{C}_{g,\tau}, \tau)$ -greedy for all $\tau \in (0, 1]$. Later, Konyagin and Temlyakov [16] noted that the same argument works for any greedy basis. In particular, we have

Theorem 6.5 (Konyagin and Temlyakov). *Let \mathcal{B} be a \mathbf{C}_g -greedy M -basis. Then for any fixed $\tau \in (0, 1]$, we have*

$$\gamma_{m,\tau}(x) \leq \left(\frac{\mathbf{C}_g^4}{\tau} + \mathbf{C}_g \right) \sigma_m(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}. \quad (6.11)$$

Proof. Let $x \in \mathbb{X}$, $m \in \mathbb{N}$, and $\varepsilon > 0$. Let $A \subset \mathbb{N}$ and $(a_n)_{n \in A}$ be such that $|A| = m$ and $\|x - \sum_{n \in A} a_n e_n\| \leq \sigma_m(x) + \varepsilon$. Pick $B \in \mathcal{G}(x, m, \tau)$. By the triangle inequality, we have

$$\|x - P_B(x)\| = \|P_{A \cup B}^c(x) + P_{A \setminus B}(x)\| \leq \|P_{A \cup B}^c(x)\| + \|P_{A \setminus B}(x)\|. \quad (6.12)$$

We bound $\|P_{A \setminus B}(x)\|$. By Theorem 4.4, our basis is \mathbf{C}_g -suppression unconditional and satisfies \mathbf{C}_g -property (A). Observe that

$$\tau \max_{n \in A \setminus B} |e_n^*(x)| \leq \tau \max_{n \notin B} |e_n^*(x)| \leq \min_{n \in B} |e_n^*(x)| \leq \min_{n \in B \setminus A} |e_n^*(x)|.$$

Let $\varepsilon_n = \text{sgn}(e_n^*(x))$ for all $n \in \text{supp}(x)$. We have

$$\begin{aligned}
 \|P_{A \setminus B}(x)\| &\leq \mathbf{C}_g \max_{n \in A \setminus B} |e_n^*(x)| \|1_{\varepsilon A \setminus B}\| \text{ due to Proposition 2.1} \\
 &\leq \frac{\mathbf{C}_g^2}{\tau} \min_{n \in B \setminus A} |e_n^*(x)| \|1_{\varepsilon B \setminus A}\| \text{ by } \mathbf{C}_g\text{-property (A)} \\
 &\leq \frac{\mathbf{C}_g^3}{\tau} \|P_{B \setminus A}(x)\| \text{ due to Proposition 2.1.} \quad (6.13)
 \end{aligned}$$

Next, we bound $\|P_{B \setminus A}(x)\|$ and $\|P_{A \cup B}^c(x)\|$ as follows:

$$\|P_{B \setminus A}(x)\| = \left\| P_{B \setminus A} \left(x - \sum_{n \in A} a_n e_n \right) \right\| \leq \mathbf{C}_g \left\| x - \sum_{n \in A} a_n e_n \right\|, \text{ and} \quad (6.14)$$

$$\|P_{A \cup B}^c(x)\| = \left\| P_{B \cup A}^c \left(x - \sum_{n \in A} a_n e_n \right) \right\| \leq \mathbf{C}_g \left\| x - \sum_{n \in A} a_n e_n \right\|. \quad (6.15)$$

From (6.12) to (6.15), we obtain

$$\|x - P_B(x)\| \leq \left(\frac{\mathbf{C}_g^4}{\tau} + \mathbf{C}_g \right) (\sigma_m(x) + \varepsilon).$$

Letting $\varepsilon \rightarrow 0$, we are done. \square

7. FUTURE RESEARCH

We list several problems for future research:

Problem 7.1. Can we improve the power $1/2$ in (1.15)?

Problem 7.2. Is there a Schauder basis \mathcal{B} such that $\mathbf{C}_{b,\tau} < \infty$ for all $\tau \in (0, 1]$ (see Definition 4.3) and $\sup_\tau \mathbf{C}_{b,\tau} = \infty$? By Corollary 5.6, \mathcal{B} must be non-quasi-greedy.

Problem 7.3. Given $0 < \tau_2 < \tau_1 \leq 1$, for an M-basis, does property (A, τ_1) imply property (A, τ_2) without any additional assumption?

Problem 7.4. Is the estimate in Proposition 5.3 optimal? Specifically, for $0 < \tau_2 < \tau_1 \leq 1$, can we construct a basis \mathcal{B} with \mathbf{C}_{b,τ_2} -property (A, τ_2) and \mathbf{C}_{b,τ_1} -property (A, τ_1) so that $\mathbf{C}_{b,\tau_1} = \mathbf{C}_{b,\tau_2} \tau_1 / \tau_2$?

APPENDIX A. REPEATED PROOFS

Proof of Lemma 2.3. Let $A, B, x, (\varepsilon), t$ be chosen as in (2.2). By norm convexity, we have

$$\begin{aligned} \|x\| &= \left\| x - P_B(x) + \sum_{n \in B} e_n^*(x) e_n \right\| \leq \sup_{(\delta)} \left\| x - P_B(x) + \frac{t}{\tau} 1_{\delta B} \right\| \\ &= \frac{1}{\tau} \sup_{(\delta)} \|\tau(x - P_B(x)) + t 1_{\delta B}\| \\ &\leq \frac{\nu_{m,\tau,\ell}}{\tau} \|x - P_B(x) + t 1_{\varepsilon A}\|. \end{aligned}$$

Hence, $\Omega_{m,\tau,\ell} \leq \nu_{m,\tau,\ell} / \tau$.

For the reverse inequality, let $A, B, x, (\varepsilon), (\delta)$ be chosen as in (1.2). Let $y = \tau x + 1_{\delta B}$. Then $\|y\|_\infty \leq 1$. By (2.1),

$$\|\tau x + 1_{\delta B}\| = \|y\| \leq \Omega_{m,\tau,\ell} \|y - P_B(y) + \tau 1_{\varepsilon A}\| = \Omega_{m,\tau,\ell} \tau \|x + 1_{\varepsilon A}\|.$$

Hence, $\Omega_{m,\tau,\ell} \geq \nu_{m,\tau,\ell} / \tau$. \square

Proof of Lemma 2.4. Let $A, B, x, (\varepsilon), t$ be chosen as in (2.3). By norm convexity, we have

$$\begin{aligned} \|x\| &= \left\| x - P_B(x) + \sum_{n \in B} e_n^*(x) e_n \right\| \leq \sup_{(\delta)} \left\| x - P_B(x) + \frac{t}{\tau} 1_{\delta B} \right\| \\ &= \frac{1}{\tau} \sup_{(\delta)} \|\tau(x - P_B(x)) + t 1_{\delta B}\| \\ &\leq \frac{\nu'_{m,\tau,\ell}}{\tau} \sup_{(\delta)} \|x - P_B(x) + t 1_{\varepsilon A}\|. \end{aligned}$$

Hence, $\Omega'_{m,\tau,\ell} \leq \nu'_{m,\tau,\ell}/\tau$.

For the reverse inequality, let $A, B, x, (\varepsilon), (\delta)$ be chosen as in (1.4). Let $y = \tau x + 1_{\delta B}$. Then $\|y\|_\infty \leq 1$. We have

$$\|\tau x + 1_{\delta B}\| = \|y\| \leq \Omega'_{m,\tau,\ell} \|y - P_B(y) + \tau 1_{\varepsilon A}\| = \tau \Omega'_{m,\tau,\ell} \|x + 1_{\varepsilon A}\|.$$

Hence, $\Omega'_{m,\tau,\ell} \geq \nu'_{m,\tau,\ell}/\tau$. \square

REFERENCES

- [1] F. Albiac and J. L. Ansorena, Characterization of 1-almost greedy bases, *Rev. Mat. Complut.* **30** (2017), 13–24.
- [2] F. Albiac and N. Kalton, *Topics in Banach Space Theory*, second edition, ISBN 978-3-319-31555-3 (2016).
- [3] F. Albiac and P. Wojtaszczyk, Characterization of 1-greedy bases, *J. Approx. Theory* **138** (2006), 65–86.
- [4] P. M. Berná, A note on partially-greedy bases in quasi-Banach spaces, *Studia Math.* **259** (2021), 225–239.
- [5] M. Berasategui, P. M. Berná, and S. Lassalle, Strong partially greedy bases and Lebesgue-type inequalities, *Constr. Approx.* **54** (2021), 507–528.
- [6] P. M. Berná and H. V. Chu, On some characterizations of greedy-type bases, preprint. Available at <https://arxiv.org/abs/2201.12029>.
- [7] P. M. Berná, Ó. Blasco, G. Garrigós, E. Hernández, and T. Oikhberg, Lebesgue inequalities for Chebyshev thresholding greedy algorithms, *Rev. Mat. Complut.* **33** (2020), 695–722.
- [8] P. M. Berná, O. Blasco, and G. Garrigós, Lebesgue inequalities for greedy algorithm in general bases, *Rev. Mat. Complut.* **30** (2017), 369–392.
- [9] S. J. Dilworth, D. Kutzarova, and T. Oikhberg, Lebesgue constants for the weak greedy algorithm, *Rev. Mat. Complut.* **28** (2015), 393–409.
- [10] S. J. Dilworth, D. Kutzarova, E. Odell, Th. Schlumprecht, and A. Zsák, Renorming spaces with greedy bases, *J. Approx. Theory* **188** (2014), 39–56.
- [11] S. J. Dilworth, M. Soto-Bajo, and V. N. Temlyakov, Quasi-greedy bases and Lebesgue-type inequalities, *Studia Math.* **211** (2012), 41–69.
- [12] S. J. Dilworth, N. J. Kalton, D. Kutzarova, and V. N. Temlyakov, The thresholding greedy algorithm, greedy bases, and duality, *Constr. Approx.* **19** (2003), 575–597.
- [13] S. J. Dilworth, N. J. Kalton, and D. Kutzarova, On the existence of almost greedy bases in Banach spaces, *Studia Math.* **159** (2003), 67–101.
- [14] G. Garrigós, E. Hernández, and T. Oikhberg, Lebesgue-type inequalities for quasi-greedy bases, *Constr. Approx.* **38** (2013), 447–470.
- [15] D. Khurana, Weight-partially greedy bases and weight-property (A), *Ann. Funct. Anal.* **11** (2020), 101–117.
- [16] S. V. Konyagin and V. N. Temlyakov, Greedy approximation with regard to bases and general minimal systems, *Serdica Math. J.* **28** (2002), 305–328.

- [17] V. N. Temlyakov, The best m -term approximation and greedy algorithms, *Advances in Comp. Math.* **8** (1998), 249–265.
- [18] V. N. Temlyakov, M. Yang, and P. Ye, Lebesgue-type inequalities for greedy approximation with respect to quasi-greedy bases, *East J. Approx.* **17** (2011), 127–138.
- [19] P. Wojtaszczyk, Greedy algorithms for general biorthogonal systems. *J. Approx. Theory* **107** (2000), 293–314.

Email address: hungchu2@illinois.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA,
IL 61820, USA