

ON SHADOWING AND STOCHASTIC STABILITY

HECTOR SUNI PUMA AND CHRISTIAN S. RODRIGUES

ABSTRACT. In this paper, we study stochastic stability of a dynamical system with shadowing property, which evolves under small random perturbation. We prove that time averages along the pseudo-trajectory converge with respect to stationary measure for the randomly perturbed dynamics. In particular, we prove that stationary measures converge to physical measures when the noise level goes to zero if dynamical system has a shadowing property.

1. INTRODUCTION

The study of statistical properties of Dynamical Systems is currently present in almost all fields of science, from fundamental Mathematics to applied modelling. In its essence, one wants to classify long term behaviour of processes undergoing time transformation and understand how stable they are. This task is particularly intricate in the case of randomly perturbed dynamics, where several appearing phenomena have no counter part in the deterministic case.

On the one hand, when the dynamical system evolves under small random perturbations, a way to characterise robust behaviour is by means of the so called *stochastic stability*. We say that a system is stochastically stable if the stationary measure for the randomly perturbed dynamics is close, in some sense, to the invariant physical or Sinai-Ruelle-Bowen (SRB) measure, which in turn encodes important dynamical information. A more precise definition will be timely presented.

Since the 1980's, several results have been obtained under different hypotheses leading to proving stochastic stability of some classes of systems. To mention a few, Kifer has first proved stochastic stability of diffeomorphisms having (uniformly) hyperbolic attractors and expanding maps [Kif86, Kif88]. Under slightly different technical assumptions L.-S. Young then showed that the stability of (uniformly) hyperbolic attractors of C^2 - diffeomorphisms follows from the persistence of hyperbolic structures [You86]. Viana has shown how to use spectral properties of transfer operators to infer stochastic stability of several classes of maps [Via97]. Later Araújo

Date: May 17, 2022.

2020 *Mathematics Subject Classification.* 37C50(primary), 37C05 (secondary), 37H10, 37C40, 49K45.

Key words and phrases. Markov chain, random dynamics, shadowing, stochastic stability.

has proved the existence and finiteness of physical measures for randomly perturbed dynamics considering some special type of perturbations [Ara00, Ara01]. Then Alves and Araújo, based on some construction already present in [Ara00], gave sufficient and necessary, conditions for stochastic stability of some non-uniformly expanding maps [AlA03]. More recently, Benedicks and Viana have proved the stochastic stability of Hénon like maps [BeV06], and Alves, Araújo, and Vásquez the stochastic stability of a type of non-uniformly hyperbolic diffeomorphisms with dominating splitting [AAV07].

On the other hand, another common way to characterise robustness of dynamics is through the idea of *structural stability*. A dynamical system is said to be structurally stable if it is equivalent to some other dynamical system in its vicinity, choosing an appropriate topology. Among different consequences of structural stability, one may investigate the *shadowing property*. Vaguely speaking, the shadowing property says that every pseudo-orbit, i.e. every sequence of points whose elements are close enough to an orbit of a dynamical system, can be described by a real orbit itself. Several authors have studied different types of shadowing under different technical assumptions, mainly, because of its close connection to structural stability [Bow75, Kat80, Pil99, PiT10]. See [Pil99] for a extensive historical account.

These two apparently distinct approaches to stability lead us to an important question: **how structural properties of dynamical systems are related to their statistical ones?**

The concept of shadowing is clearly related to stochastic stability because the orbits evolving under random perturbations are pseudo-orbits. In this context, proving stochastic stability equals to proving that most random orbits mimics the behaviour of real orbits (the unperturbed ones). The connection of shadowing with stochastic stability, however, has never been satisfactorily addressed. This problem has also been raised by Bonatti, Díaz and Viana in [BDV05, Problem D10]. To the best of our knowledge, the only attempt to study this connection was presented in the manuscript [Tod14], where the author imposes several hypotheses and a much more restrictive context than ours.

In this paper, we study stochastic stability of a dynamical system with shadowing property, which evolves under small random perturbation. We define time averages with respect to continuous observables along the trajectory which converges to their averages with respect to an invariant measure. Using this framework we study time averages along the pseudo-trajectory that converge with respect to stationary measure for the randomly perturbed dynamics. Finally, we proof that stationary measures converge to physical measures when the noise level goes to zero if dynamical system has a shadowing property.

2. ERGODICITY AND PHYSICAL MEASURES

To get started, let M be a compact smooth manifold of finite dimension with or without border. We endow M with the distance d induced by a Riemannian metric and a normalised Riemannian volume form m , which we fix once and for all. When it is not stated otherwise, absolute continuity will be taken with respect to the probability m .

We set $\mathbf{x} = (x_j)_{j \in \mathbb{Z}_0^+}$ to be a sequence of points in M from the non-negative integers. We define the following probability measure

$$S_n(\mathbf{x}) = \frac{1}{n+1} \sum_{j=0}^n \delta_{x_j} \quad (1)$$

for $x_j \in M$, where for every $y \in M$, the measure δ_y is the Dirac delta probability measure supported on y .

We shall consider the dynamics generated by the iteration of a continuous function $f : M \rightarrow M$, where we define $z_j = f^j(z_0) = \underbrace{f \circ \cdots \circ f}_{j \text{ times}}(z_0)$ for any

initial condition $z_0 \in M$. Hereafter we set $f^0(z) = z$, for every $z \in M$. We also define the probability measure

$$S_n(f) = S_n(\{f^j(z)\}_{j=0}^\infty) = \frac{1}{n+1} \sum_{j=0}^n \delta_{f^j(z)}. \quad (2)$$

As usual, we say that a probability measure μ is invariant by f when $\mu(f^{-1}(A)) = \mu(A)$, for every Borel set $A \subset M$. Recall that, since M endowed with the distance induced by the Riemannian metric is a compact metric space and f is continuous, by the Krylov-Bogolyubov Theorem, there exists at least one f -invariant probability measure on M . So, if μ is such an f -invariant probability measure and $\varphi \in L^1(\mu)$ then, by the Birkhoff Ergodic Theorem, the limit

$$\tilde{\varphi} = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n \varphi(f^j(x))$$

exists for μ -almost every point $x \in M$. Note that, on the one hand, for any $\varphi \in L^1(M)$, from (2) we have

$$\begin{aligned} \int \varphi dS_n(f) &= \int \varphi d \left(\frac{1}{n+1} \sum_{j=0}^n \delta_{f^j(x)} \right) \\ &= \frac{1}{n+1} \sum_{j=0}^n \int \varphi d\delta_{f^j(x)} \\ &= \frac{1}{n+1} \sum_{j=0}^n \varphi(f^j(x)). \end{aligned}$$

Thus, there exists the limit

$$E_x(\varphi) \equiv \lim_{n \rightarrow \infty} \int \varphi dS_n(f)$$

for almost every $x \in M$. On the other hand, we also have that $\varphi \mapsto E(\varphi) = E_x(\varphi)$, for each $x \in B \subset M$, where B is some set of positive measure, defines a non-negative linear functional on the space $L^1(M)$. Therefore, by the Riesz-Markov-Kakutani Representation Theorem, there exists a unique Radon measure ν on M , such that,

$$\int \varphi d\nu = E(\varphi) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n \varphi(f^j(x)). \quad (3)$$

Hence, the measure μ is ergodic if for μ -almost every point $x \in M$, it holds that

$$S_n(x) \rightarrow \mu, \text{ as } n \rightarrow \infty$$

in the *weak**-topology on the space of probability measures on M . These considerations lead us to the idea of physical measures.

Definition 2.1 (Physical measure). *An invariant probability measure μ on M is called physical measure if equation (3) holds true for a set $\mathcal{B}(\mu) \subset M$, of positive Lebesgue measure, and for all continuous function $\varphi : M \rightarrow \mathbb{R}$.*

Therefore, μ being ergodic equals to (3) holding true for almost every point in M , so that μ has full measure. Thus, in this case, μ is physical for f .

Note that we have not assumed *ad hoc* hypotheses for the existence of a unique ergodic physical measure, as it appears to be the case in [Tod14].

3. SHADOWING PROPERTY

Using the notation of Section 2, we shall consider the dynamics generated by the iteration of a continuous function $f : M \rightarrow M$. For a trajectory

$z_j = f^j(z_0)$ we can consider sequences of points which are close enough to this trajectory, what is formalised in the following definition.

Definition 3.1 (Pseudo-trajectory). *Given $\delta > 0$, we say that a sequence of points $\mathbf{x} = (x_j)_{j \in \mathbb{Z}_0^+}$ in M is a δ -pseudo-trajectory of f if*

$$d(f(x_j), x_{j+1}) \leq \delta, \text{ for all } j \geq 0. \quad (4)$$

We can now define the shadowing property we will be considering in this paper.

Definition 3.2 (Shadowing). *We say that a dynamical system generated by f has the shadowing property if, for any $\varepsilon > 0$, we can find a $\delta > 0$, such that, for each δ -pseudo-trajectory $\mathbf{x} = (x_j)_{j \in \mathbb{Z}_0^+}$ of f , there exists a point $z \in M$, such that,*

$$d(f^j(z), x_j) < \varepsilon, \text{ for all } j \geq 0.$$

4. RANDOM PERTURBATIONS

Given any small enough $\varepsilon_0 > 0$, as before, we consider the dynamics generated by the iteration of a continuous function $f : M \rightarrow M$. A Markov chain is defined by a family $\{P_\varepsilon(\cdot|x) : x \in M, 0 < \varepsilon < \varepsilon_0\}$ of Borel probability measures, such that, for each $x \in M$, we have $\text{supp}(P_\varepsilon(\cdot|x)) \subset B_\varepsilon(f(x))$, where $B_\varepsilon(f(x))$ is the open ball of radius ε centred in $f(x)$. Henceforth, the orbit of f subject to small perturbation will be called random orbits. It is given by the sequence of random variables, $(x_j)_j$ for all $j = 0, 1, 2, \dots$, where each x_{j+1} has probability distribution $P_\varepsilon(\cdot|x_j)$. This family of Markov chains will be called small random perturbation of f . Under quite general assumptions, it is shown to exist a family of random maps representing the Markov chain [JKR15, JMPRI9].

In addition, in this paper we assume that every realisation, $(x_j)_{j \in \mathbb{Z}_0^+}$, of the family of Markov chain for any $x_0 \in M$, is a $\delta(\varepsilon)$ -pseudo-trajectory of f , for any $\delta(\varepsilon) > 0$. That is

$$d(f(x_j), x_{j+1}) \leq \delta(\varepsilon) \text{ for all } j \geq 0.$$

Remark 4.1. *The quantifier $\delta(\varepsilon)$ should be considered as a continuous function controlling the type of shadowing, $\delta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that, $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. It is the same appearing in (4), but we introduce an explicit dependence on the size of the random perturbation. Therefore, the approach taken here allows for different versions of shadowing property by choosing different functions $\delta(\varepsilon)$. This includes, for example, the Lipschitz shadowing [PiT10] as well as some other types of shadowing properties.*

Similarly to the importance of invariant measure for an application, the stationary measures for the Markov chain encodes important ergodic properties of the randomly perturbed dynamics.

Definition 4.2. A Borel probability measure μ_ε on M is a stationary measure for the Markov chain if

$$\mu_\varepsilon(G) = \int P_\varepsilon(G|x)d\mu_\varepsilon$$

for every Borel set $G \subset M$.

We can define a shift map $F : M \times M^\mathbb{N} \rightarrow M \times M^\mathbb{N}$, $(x_0, \mathbf{x}) \mapsto (x_1, \sigma(\mathbf{x}))$, where σ is the left shift on the space of random orbit, such that, $\sigma(\mathbf{x}) = (x_1, x_2, \dots)$ for any $\mathbf{x} = (x_0, x_1, x_2, \dots)$ and x_0 . The stationary measure defined above is equivalent to the skew-product measure $\mu_\varepsilon \times P_\varepsilon^\mathbb{N}$, given by

$$d(\mu_\varepsilon \times P_\varepsilon^\mathbb{N})(x_0, x_1, \dots, x_j, \dots) = \mu_\varepsilon(x_0)P_\varepsilon(dx_1|x_0)\dots P_\varepsilon(dx_j|x_{j-1})\dots,$$

to be invariant under the shift map. Since M is a compact metric space, by Tychonoff Theorem, the space $M \times M^\mathbb{N}$ is also compact. Moreover, F is a continuous map, then by the Krylov-Bogolyubov Theorem, there exists at least one F -invariant probability measure on $M \times M^\mathbb{N}$. So, if $\mu_\varepsilon \times P_\varepsilon^\mathbb{N}$ is such an F -invariant probability measure and ψ is an integrable function, then, by the Birkhoff Ergodic Theorem,

$$\tilde{\psi}(\mathbf{x}) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n \psi(x_j) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n (\psi \circ \pi_0)(F^j(\mathbf{x}))$$

exists for $\mu_\varepsilon \times P_\varepsilon^\mathbb{N}$ -a.e $(x, \mathbf{x}) \in M \times M^\mathbb{N}$, where $\pi_0 : M \times M^\mathbb{N} \rightarrow M$ is the projection on the first factor. Thus, there exists the limit

$$E_x(\psi) := \lim_{n \rightarrow \infty} \int \psi dS_n(\mathbf{x})$$

for μ_ε -almost every $x \in M$. Moreover, we have that $\psi \mapsto E(\psi)$, for any $x \in B \subset M$, where B is some set of positive measure, defines a non-negative linear functional on the space $L^1(M)$. Thus, by the Riesz-Markov Kakutani Representation Theorem, there exists a unique Radon measure ν_ε on M , such that,

$$\int \psi d\nu_\varepsilon = E(\psi) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n \psi(x_j)$$

Hence, the measure μ_ε is ergodic if μ_ε -almost every point $x \in M$, it holds that

$$S_n(\mathbf{x}) \rightarrow \mu_\varepsilon, \text{ as } n \rightarrow \infty \quad (5)$$

in the *weak**-topology sense in the space of probability measures on M .

Based on these, we can define the stability against random perturbation as following.

Definition 4.3. The system (f, μ) is stochastically stable under the small perturbation scheme $\{P_\varepsilon(\cdot|x) : x \in M, 0 < \varepsilon < \varepsilon_0\}$ if μ_ε converges, in the *weak** sense, to the physical measure μ , when the noise level ε goes to zero.

We shall consider the space of continuous functions from M to \mathbb{R} , denoted by $C(M)$, which we endow with the *sup*-norm. Furthermore, let $\mathcal{B} \subset C(M)$ be any set of bounded continuous functions on M . These functions will be called *observables* from \mathcal{B} . Then, we can prove the following easy lemmata for later use.

Lemma 4.4. *The measure μ is ergodic if for μ -a.e point $x \in M$, it holds that $S_n(f) \rightarrow \mu$, as $n \rightarrow \infty$. Then, for any $\varphi \in \mathcal{B}$, and for any $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$, such that,*

$$\left| \int \varphi dS_n(f) - \int \varphi d\mu \right| \leq \varepsilon, \text{ for all } n \geq n_0.$$

Proof. Suppose that $S_n(x) \rightarrow \mu$ in M and take $\varphi \in \mathcal{B}$. Given $\varepsilon > 0$, consider the neighbourhood

$$N_{\varepsilon, \varphi}(\mu) = \left\{ \nu : \left| \int \varphi d\nu - \int \varphi d\mu \right| < \varepsilon \right\}.$$

By hypothesis, there exists $n_0 \in \mathbb{N}$, such that, for all $n \geq n_0$, we have $S_n(f) \in N_{\varepsilon, \varphi}(\mu)$. That is, $\left| \int \varphi dS_n(f) - \int \varphi d\mu \right| < \varepsilon$, for all $n \geq n_0$. \square

Lemma 4.5. *The measure μ_ε is ergodic, if for μ_ε -a.e point $x \in M$, it holds that $S_n(\mathbf{x}) \rightarrow \mu_\varepsilon$, as $n \rightarrow \infty$. Then, for any $\varphi \in \mathcal{B}$, and for any $\delta(\varepsilon) > 0$, there exists $n_0 \in \mathbb{N}$, such that,*

$$\left| \int \varphi dS_n(\mathbf{x}) - \int \varphi d\mu_\varepsilon \right| \leq \delta(\varepsilon), \forall n \geq n_0$$

Proof. Suppose that $S_n(\mathbf{x}) \rightarrow \mu_\varepsilon$ in M and take $\varphi \in \mathcal{B}$. Given $\delta(\varepsilon) > 0$, consider the neighborhood $N_{\varepsilon, \varphi}(\mu_\varepsilon) = \left\{ \nu_\varepsilon : \left| \int \varphi d\nu_\varepsilon - \int \varphi d\mu_\varepsilon \right| < \delta(\varepsilon) \right\}$. By hypothesis, there exists $n_0 \in \mathbb{N}$, such that, for all $n \geq n_0$ we have $S_n(\mathbf{x}) \in N_{\varepsilon, \varphi}(\mu_\varepsilon)$, that is $\left| \int \varphi dS_n(\mathbf{x}) - \int \varphi d\mu_\varepsilon \right| < \delta(\varepsilon)$, for all $n \geq n_0$. \square

5. SHADOWING AND STOCHASTIC STABILITY

We can now state and prove the main result of this manuscript.

Theorem 5.1. *If the dynamics generated by the map $f : M \rightarrow M$ has the shadowing property and $\text{supp}(\mu_\varepsilon) \subset B_\varepsilon(f(x))$, then it is stochastically stable under the perturbation scheme $\{P_\varepsilon(\cdot|x) : x \in B_\varepsilon(f), \varepsilon > 0\}$ in the weak*-topology sense.*

Proof. Given any small $\varepsilon_0 > 0$, for $0 < \varepsilon < \varepsilon_0$, let $B_{\delta, \varepsilon}$ be a set of realizations of the Markov chain. By hypothesis, f has the shadowing property, which means that, for any $0 < \varepsilon < \varepsilon_0$, we can find a $\delta(\varepsilon) > 0$, such that, for any $\delta(\varepsilon)$ -pseudo-trajectory $\mathbf{x} \in B_{\delta, \varepsilon}$ of f , there exists a point $x \in M$, such that,

$$d(f^j(x), x_j) < \varepsilon, \text{ for all } j \geq 0.$$

For any fixed $\varepsilon > 0$, $\varphi \in \mathcal{B}$, and for any $N \in \mathbb{N}$, there exists a constant $C(\varphi)$, such that, holds

$$\left| \int \varphi dS_n(f) - \int \varphi dS_n(\mathbf{x}) \right| \leq C(\varphi)\varepsilon, \text{ for all } n > N. \quad (6)$$

Indeed,

$$\begin{aligned} & \left| \int \varphi dS_n(f) - \int \varphi dS_n(\mathbf{x}) \right| \\ &= \left| \int \varphi d \left(\frac{1}{n+1} \sum_{j=0}^n \delta_{f^j(x)} \right) - \int \varphi d \left(\frac{1}{n+1} \sum_{j=0}^n \delta_{x_j} \right) \right| \\ &\leq \frac{1}{n+1} \sum_{j=0}^n \left| \int \varphi d\delta_{f^j(x)} - \int \varphi d\delta_{x_j} \right| \\ &= \frac{1}{n+1} \sum_{j=0}^n |\varphi(f^j(x)) - \varphi(x_j)| \\ &\leq \frac{1}{n+1} \sum_{j=0}^n \sup_{x \in M} |\varphi(f^j(x)) - \varphi(x_j)|. \end{aligned} \quad (7)$$

Since $\varphi \in \mathcal{B}$, there exists a constant $C(\varphi) > 0$, such that,

$$\sup_{x \in M} |\varphi(x)| \leq C(\varphi).$$

Then, we can write the equation (7) as,

$$\sup_{x \in M} |\varphi(f^j(x)) - \varphi(x_j)| \leq \sup_{x \in M} |\varphi(x)| d(f^j(x), x_j) \leq C(\varphi)\varepsilon.$$

Thus, we have

$$\left| \int \varphi dS_n(x) - \int \varphi dS_n(\mathbf{x}) \right| \leq C(\varphi)\varepsilon \text{ for all } n \geq N.$$

Now, we show stochastic stability. In effect, note that $|\int \varphi d\mu - \int \varphi d\mu_\varepsilon|$ one can rewrite in terms of $\int \varphi dS_n(f)$ and $\int \varphi dS_n(\mathbf{x})$ and, then applying triangle inequality, we have the following

$$\begin{aligned} \left| \int \varphi d\mu - \int \varphi d\mu_\varepsilon \right| &\leq \left| \int \varphi d\mu - \int \varphi dS_n(f) \right| + \left| \int \varphi dS_n(\mathbf{x}) - \int \varphi d\mu_\varepsilon \right| \\ &\quad + \left| \int \varphi dS_n(f) - \int \varphi dS_n(\mathbf{x}) \right| \\ &\leq (C(\varphi) + 1)\varepsilon + \delta(\varepsilon). \end{aligned}$$

Observe that the first and second terms follow from Lemmata 4.4 and 4.5. The last term is a consequence from (6). Moreover, since $B_{\delta,\varepsilon}$ be a set realization of Markov chain and the measure $S_n(\mathbf{x})$, defined as (1), then from (5) and ergodicity, we have that in the limit we have $\mu_\varepsilon(B_{\delta,\varepsilon}) \rightarrow 1$, as $N \rightarrow \infty$. This concludes the proof. \square

It is possible to formulate a similar result to Theorem 5.1 in terms of random maps, given that there is an equivalent formulation of the perturbation scheme through iteration of random functions via the representation of Markov chains [JKR15, JMPR19].

ACKNOWLEDGEMENTS

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. C. S. R. would like to acknowledge support from the Max Planck Society, Germany, through the award of a Max Planck Partner Group for Geometry and Probability in Dynamical Systems. C. S. R. has also been supported by the Brazilian agency: grant #2016/00332-1, grant #2018/13481-0, grant #2020/04426-6, São Paulo Research Foundation (FAPESP). The opinions, hypotheses and conclusions or recommendations expressed in this work are the responsibility of the authors and do not necessarily reflect the views of FAPESP.

REFERENCES

- [AAV07] J. F. Alves, V. Araújo, and C. H. Vásquez, *Stochastic stability of non-uniformly hyperbolic diffeomorphisms*, Stoch. Dyn. **7** (2007), 299-333.
- [AIA03] J. F. Alves and V. Araújo, *Random perturbations of nonuniformly expanding maps*, Astérisque **286**, (2003), 25-62.
- [Ara00] V. Araújo, *Attractors and time averages for random maps*, Ann. de l'Inst. H. Poincaré - Anal. Non linéaire **17** (2000), 307-369.
- [Ara01] V. Araújo, *Infinitely Many Stochastically Stable Attractors*, Nonlinearity **14** (2001), 583-596.
- [BeV06] M. Benedicks and M. Viana, *Random perturbations and statistical properties of Hénon-like maps.*, Non Linéaire **23** (2006), no. 5, 713-752.
- [BDV05] C. Bonatti, L. J. Díaz, and M. Viana, *Dynamics beyond uniform hyperbolicity: a global geometric and probabilistic perspective*, Encyclopedia of Mathematical Sciences, vol 102, Springer, Berlin, 2005.
- [Bow75] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lect. Notes in Math. Vol 470, Springer-Verlag (1975).
- [JKR15] J. Jost, M. Kell, and C. S. Rodrigues, *Representation of Markov chains by random maps: existence and regularity conditions*, Calculus of Variations and Partial Differential Equations **54** (2015), 2637-2655.
- [JMPR19] J. Jost, R. Rostislav, J. W. Portegies, and C. S. Rodrigues, *On the regular representation of measures*, Communications in Analysis and Geometry **27** (2019) 1799-1823.
- [Kat80] A. Katok, *Lyapunov exponents, entropy and periodic points of diffeomorphisms*, Publ. Math. IHES, **51** (1980), 137-173.
- [Kif86] Yu. Kifer, *Ergodic theory of random transformations*, Progress in Probability and Statistics, vol 10, Birkhäuser, Boston, 1986.
- [Kif86b] Yu. Kifer, *General random perturbations of hyperbolic and expanding transformations*, Jour. D'Anal. Mathématique **47** (1986), 112-150.
- [Kif88] Yu. Kifer, *Random perturbations of dynamical systems*, Progress in Probability and Statistics, vol 16, Birkhäuser, Boston, 1988.

- [Tod14] Dmitry Todorov, Stochastic shadowing and stochastic stability, arXiv:1411.7604v1 [math.DS], (2014).
- [Pil99] Yu. S. Pilyugin, *Shadowing in Dynamical Systems*, Lect. Notes in Math. Vol 1706, Springer-Verlag (1999).
- [PiT10] Yu. S. Pilyugin and S. Tikhomirov, *Lipschitz shadowing implies structural stability*, Nonlinearity **23** (2010), 2509-2515.
- [Via97] M. Viana, *Stochastic dynamics of deterministic systems*, 21st - Braz. Math. Colloq., IMPA - Rio de Janeiro (1997).
- [You86] L.-S. Young, *Stochastic stability of hyperbolic attractors*, Erg. Th. & Dyn. Sys. **6** (1986), 311-319.

HECTOR SUNI PUMA, INSTITUTE OF MATHEMATICS, DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSIDADE ESTADUAL DE CAMPINAS, 13.083-859 CAMPINAS - SP, BRAZIL
Email address: `hector.suni@upr.edu`

CHRISTIAN S. RODRIGUES, INSTITUTE OF MATHEMATICS, DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSIDADE ESTADUAL DE CAMPINAS, 13.083-859 CAMPINAS - SP, BRAZIL
Email address: `rodrigues@ime.unicamp.br`