

GORENSTEIN ENDOMORPHISM RINGS ON CURVE SINGULARITIES

PHILIPP KORELL

ABSTRACT. We characterize the Gorensteinness of endomorphism rings of a fractional ideal on a curve singularity by stability of the ideal and a condition on its value semigroup ideal. Moreover, the Gorenstein algebroid curves with only Gorenstein integral extensions are classified.

1. INTRODUCTION

By the Cayley–Hamilton theorem, endomorphism rings of modules are integral extensions of the base ring. Consequently, endomorphism rings appear in the contexts of (algorithmic) normalization or non-commutative resolutions [1]. Grauert and Remmert showed that a reduced Noetherian ring is normal if and only if it is the endomorphism ring of a so-called *test ideal* [2]. This criterion for normality is the foundation of de Jong’s algorithm for normalization [3, 4]: Computing successively the endomorphism rings of test ideals yields a sequence of integral extensions, eventually stabilizing at the normalization. For one-dimensional rings, Leuschke related the sequence of endomorphism rings obtained by this procedure to non-commutative crepant resolutions [5].

In dimension one, normalization is the same as desingularization. The unique test ideal of a local algebroid curve is the maximal ideal. In this setting, Böhm, Decker, and Schulze studied the intermediate rings produced in the algorithmic normalization of algebroid curves according to de Jong [6]. As a main tool for computing the endomorphism ring of the maximal ideal they considered the *semigroup of values* of an algebroid curve.

In this article, we consider endomorphism rings of fractional ideals of a certain class of one-dimensional Cohen–Macaulay rings which we call *admissible* (see Definition 2.15). These admissible rings generalize algebroid curves while still allowing for the construction of a semigroup of values.

For an admissible ring R , there exist only finitely many (discrete) valuation rings of the total ring of fractions Q_R which contain R . Considering simultaneously the corresponding discrete valuations $\nu = (\nu_1, \dots, \nu_s): Q_R \rightarrow (\mathbb{Z} \cup \infty)^s$, we define the semigroup of values as the set of values of non-zero divisors of R : $\Gamma_R = \nu(R^{\text{reg}}) \subset \mathbb{N}^s$. Similarly, to any fractional ideal \mathfrak{J} of R we may associate a *value semigroup ideal* $\Gamma_{\mathfrak{J}}$.

It is a particularly useful feature of value semigroups that a certain symmetry condition determines whether an admissible ring is Gorenstein. This symmetry was first discovered by Kunz for irreducible curves [7] and later extended by Delgado to reducible curves [8]. From Kunz’ symmetry condition, Jäger derived a characterization of canonical ideal in terms of their value semigroup ideals [9]. Following Delgado’s notion, D’Anna extended this characterization to reducible admissible rings [10].

Barucci, D’Anna, and Fröberg took a closer look at the semiring structure of value semigroups [11]. They introduced the class of *good semigroups* which are submonoids of \mathbb{N}^s satisfying certain properties of value semigroups first identified by Delgado [8]. Notably, they also gave an example of a good semigroup which is not the semigroup of values of an admissible ring.

Nonetheless, good semigroups together with the so-called *good semigroup ideals* can be regarded as “combinatorial versions” of admissible rings and their fractional ideals. In this spirit, in [12] it was shown that every good semigroup possesses a *canonical semigroup ideal* which induces a purely combinatorial version of the Cohen–Macaulay duality on fractional ideals. In fact, this canonical semigroup ideal is characterized by its dualizing property. This is in particular not obvious as the set of good semigroup ideals is in general

2020 *Mathematics Subject Classification*. Primary 13H10; Secondary 20M12, 14H20.

Key words and phrases. curve singularity, value semigroup, endomorphism ring, Gorenstein.

not closed under taking differences (which is the corresponding operation to ideal quotients). Moreover, this “combinatorial duality” is compatible with the algebraic one under taking values.

We apply these combinatorial methods to determine the Gorensteinness of the endomorphism ring $\text{End}(\mathfrak{J})$ of a fractional ideal \mathfrak{J} of an admissible ring R . The case of \mathfrak{J} being the maximal ideal \mathfrak{m}_R of a local admissible ring R is solved already: Barucci and Fröberg stated that $\text{End}(\mathfrak{m}_R)$ is Gorenstein if and only if R is almost Gorenstein of maximal embedding dimension [13]. A full proof of this equivalence was later provided by Goto, Matsuoka, and Phuong [14].

As shown by Lipman, R has maximal embedding dimension if and only if \mathfrak{m}_R is stable [15]. In fact, we show that for any fractional ideal \mathfrak{J} , the Gorensteinness of $\text{End}(\mathfrak{J})$ implies stability of \mathfrak{J} . Incorporating value semigroups provides the condition to turn this implication into an equivalence:

Theorem (Theorem 5.1). *The endomorphism ring $\text{End}(\mathfrak{J})$ of a fractional ideal \mathfrak{J} of an admissible ring is Gorenstein if and only if \mathfrak{J} is stable and the difference (see Definition 3.21) $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ of the good semigroup ideal $\Gamma_{\mathfrak{J}}$ is symmetric.*

For the second main result of this paper we turn to the case of $\mathfrak{J} = \mathfrak{m}_R$ being the maximal ideal of a local admissible ring R . Then we classify the simultaneous Gorensteinness of R and $\text{End}(\mathfrak{m}_R)$:

Theorem (Theorem 7.1). *Let R be a local admissible ring with maximal ideal \mathfrak{m}_R . The following are equivalent.*

- (a) *Every integral extension of R in its total ring of fractions Q_R is Gorenstein.*
- (b) *R and $\mathfrak{m}_R : \mathfrak{m}_R$ are Gorenstein.*
- (c) *R is Gorenstein and has maximal embedding dimension.*
- (d) *Γ_R and $\Gamma_R \setminus \{0\} - \Gamma_R \setminus \{0\}$ are symmetric.*

Moreover, if R contains a field, and if $k = R/\mathfrak{m}_R$ is algebraically closed, then the above statements are equivalent to $\widehat{R} \cong k[[x, y]]/\langle x^{n+1} - y^2 \rangle$ for some $n \in \mathbb{N}$.

Sections 2 – 4 provide the necessary foundations on admissible rings, value semigroups, reps. good semigroups, and the symmetry condition. Section 5 is concerned with stating and proving Theorem 5.1. In Section 6, we establish a combinatorial version of Theorem 7.1 which we then use to prove Theorem 7.1 in Section 7.

Notations. By a *ring* R we mean a commutative unitary ring. The set of maximal ideals of R is denoted by $\text{Max}(R)$, and the set of minimal prime ideals by $\text{Min}(R)$. We write R^{reg} for the set of non-zero divisors (i.e. the *regular* elements) of R . The *total ring of fractions* of R is the localization $Q_R = (R^{\text{reg}})^{-1}R$. For a subset \mathfrak{J} of Q_R we use the notation $\mathfrak{J}^{\text{reg}} = \mathfrak{J} \cap Q_R^{\text{reg}}$.

The integral closure of R in Q_R is denoted by \overline{R} , also called the *normalization* of R if R is reduced. Finally, if R is semilocal, then \widehat{R} is the *completion* of R at the *Jacobson radical* $\mathfrak{J}_R = \bigcap_{\mathfrak{m} \in \text{Max}(R)} \mathfrak{m}$.

2. PRELIMINARIES

A *complex algebroid curve* is a complete reduced Noetherian local \mathbb{C} -algebra R of dimension one. Therefore, its normalization \overline{R} is finite, and by *splitting of normalization* it is the product of finitely many discrete valuation rings:

$$\begin{array}{ccc}
 R & \xrightarrow{\quad\quad\quad} & \overline{R} \\
 \searrow & & \downarrow \cong \\
 & & \prod_{\mathfrak{p} \in \text{Min}(R)} R/\mathfrak{p} \xrightarrow{\quad\quad\quad} \prod_{\mathfrak{p} \in \text{Min}(R)} \overline{R}/\mathfrak{p} \\
 \searrow \phi & & \downarrow \cong \\
 & & \prod_{\mathfrak{p} \in \text{Min}(R)} \mathbb{C}[[t_{\mathfrak{p}}]].
 \end{array}$$

This yields a *parametrization* ϕ of the curve. Since the valuation of each $\mathbb{C}[[t_{\mathfrak{p}}]]$ is the order $\text{ord}_{t_{\mathfrak{p}}}$, it furthermore allows for the definition of the classical *semigroup of values*

$$\Gamma_R = \text{ord} \circ \phi(R^{\text{reg}}) \subset \mathbb{N}^{\text{Min}(R)}.$$

When Delgado characterized Gorenstein algebroid curves by symmetry of their value semigroups [8], he identified the following important properties:

- (S0) There is a $\gamma \in \Gamma_R$ such that $\gamma + \mathbb{N}^{\text{Min}(R)} \subset \Gamma_R$.
- (S1) If $\alpha, \beta \in \Gamma_R$, then $\inf(\alpha, \beta) = (\min\{\alpha_{\mathfrak{p}}, \beta_{\mathfrak{p}}\})_{\mathfrak{p} \in \text{Min}(R)} \in \Gamma_R$.
- (S2) For any $\alpha, \beta \in \Gamma_R$ with $\alpha_{\mathfrak{p}} = \beta_{\mathfrak{p}}$ for some $\mathfrak{p} \in \text{Min}(R)$, there is an $\varepsilon \in \Gamma_R$ such that $\varepsilon_{\mathfrak{p}} > \alpha_{\mathfrak{p}} = \beta_{\mathfrak{p}}$ and $\varepsilon_{\mathfrak{q}} \geq \min\{\alpha_{\mathfrak{q}}, \beta_{\mathfrak{q}}\}$ for all $\mathfrak{q} \in \text{Min}(R)$, where equality is obtained whenever $\alpha_{\mathfrak{q}} \neq \beta_{\mathfrak{q}}$.

The reason behind (S0) is that \overline{R} is finite. For (S1) consider the sum of two elements $x, y \in R$ with values α and β such that in no component of $\phi(x) + \phi(y)$ a term of least order is canceled. Complementary, for (S2) we choose x and y such that there is cancellation in component \mathfrak{p} .

Barucci, D’Anna, and Fröberg considered conditions (S0), (S1), and (S2) as defining properties of the so-called *good semigroups* [11] which will be treated in Section 3. Before, the goal of this section is to enlarge the class of available rings which admit a semigroup of values satisfying the above properties. This will lead to the notion of *admissible rings* (see Definition 2.15).

We begin, however, with recalling some basic properties of fractional ideals, i.e. “ideals with denominators”, for later reference.

2.1. (Regular) Fractional Ideals.

Definition 2.1. Let R be a ring with total ring of fractions Q_R .

- (1) An R -submodule \mathfrak{I} of Q_R is called a *fractional ideal* of R if there is an $x \in R^{\text{reg}}$ such that $x\mathfrak{I} \subset R$.
- (2) A fractional ideal \mathfrak{I} of R is called *regular* if $\mathfrak{I}^{\text{reg}} \neq \emptyset$.
- (3) We denote the set of regular fractional ideals of R by \mathcal{R}_R .

Remark 2.2. Let R be a Noetherian ring. Then an R -submodule \mathfrak{I} of Q_R is a fractional ideal of R if and only if it is finitely generated.

Definition 2.3. Let R be a ring, and let \mathfrak{I} and \mathfrak{J} be R -submodules of Q_R .

- (1) The *product* of \mathfrak{I} and \mathfrak{J} is $\mathfrak{I}\mathfrak{J} = \left\{ \sum_{(x,y) \in \Lambda} xy \mid \Lambda \subset \mathfrak{I} \times \mathfrak{J} \text{ finite} \right\}$.
- (2) The *quotient* of \mathfrak{I} and \mathfrak{J} in Q_R is $\mathfrak{I} : \mathfrak{J} = \{x \in Q_R \mid x\mathfrak{J} \subset \mathfrak{I}\}$.

Lemma 2.4. Let R be a ring, and let $\mathfrak{I}, \mathfrak{J} \in \mathcal{R}_R$. Then $\mathfrak{I}\mathfrak{J}, \mathfrak{I} : \mathfrak{J} \in \mathcal{R}_R$.

Proof. See [16, Proposition 2.7]. □

Remark 2.5. Let R be a ring, let $x \in Q_R^{\text{reg}}$, and let $\mathfrak{I}, \mathfrak{J}', \mathfrak{J}, \mathfrak{J}'$, and \mathfrak{J} be R -submodules of Q_R . Then

- (1) $(x\mathfrak{I}) : \mathfrak{J} = x(\mathfrak{I} : \mathfrak{J}) = \mathfrak{I} : (x^{-1}\mathfrak{J})$,
- (2) $\mathfrak{I} : \mathfrak{J}' \subset \mathfrak{I} : \mathfrak{J} \subset \mathfrak{I}' : \mathfrak{J}$ if $\mathfrak{I} \subset \mathfrak{I}'$ and $\mathfrak{J} \subset \mathfrak{J}'$, and
- (3) $\mathfrak{I} : R = \mathfrak{I}$.

See [16, Lemma 2.3].

Lemma 2.6. Let R be a ring, and let \mathfrak{I} and \mathfrak{J} be regular R -submodules of Q_R . Then there is a natural R -module isomorphism

$$\begin{aligned} \text{Hom}_R(\mathfrak{I}, \mathfrak{J}) &\rightarrow \mathfrak{J} : \mathfrak{I}, \\ \phi &\mapsto \frac{\phi(x)}{x}, \end{aligned}$$

which is independent of the choice of a regular element $x \in \mathfrak{I}^{\text{reg}}$. In particular, any $\phi \in \text{Hom}_R(\mathfrak{I}, \mathfrak{J})$ is multiplication by an element of $\mathfrak{J} : \mathfrak{I}$, and it can be extended uniquely to an endomorphism of Q_R .

Proof. See [17, Lemma 2.1] and [18, Lemma 3.1]. □

Proposition 2.7. Let R be a ring, and let $\mathfrak{I} \in \mathcal{R}_R$. Then $\mathfrak{I} : \mathfrak{I}$ is an integral extension of R in Q_R .

Proof. This follows from Lemmas 2.4 and 2.6 and the Cayley–Hamilton Theorem (see [19, Theorem 4.3]). \square

Definition 2.8. Let R be a ring, and let $\mathfrak{J} \in \mathcal{R}_R$. The *conductor* of \mathfrak{J} is $\mathfrak{C}_R = \mathfrak{J} : \overline{R}$.

2.2. Discrete Valuation Rings. In this section, let Q be a ring with $Q^{\text{reg}} = Q^*$ having a *large Jacobson radical*, i.e. any prime ideal of Q containing the Jacobson radical is maximal (see [20, Proposition 19] for equivalent characterizations).

Definition 2.9.

- (1) A *valuation ring* of Q is a proper subring V of Q such that the set $Q \setminus V$ is multiplicatively closed.
- (2) Let V be a valuation ring of Q . Then for any subring R of V with $Q_R = Q$ we call V a *valuation ring over R* .
- (3) If R is a subring of Q with $Q_R = Q$, the set of valuation rings of Q over R is denoted by \mathcal{V}_R .

Remark 2.10. A valuation ring has by [21, Chapter I, Theorem 2.2] a unique regular maximal ideal \mathfrak{m}_V containing all regular non-units of V .

Definition 2.11. A valuation ring V of Q with regular maximal ideal \mathfrak{m}_V is called a *discrete valuation ring* if \mathfrak{m}_V is finitely generated.

Definition 2.12. Let A be a ring. A *discrete valuation* of A is a surjective map $\nu: A \rightarrow \mathbb{Z}_\infty = \mathbb{Z} \cup \{\infty\}$ satisfying $\nu(xy) = \nu(x) + \nu(y)$ and $\nu(x+y) \geq \min\{\nu(x), \nu(y)\}$ for every $x, y \in \mathbb{Z}_\infty$. Here we set $x + \infty = \infty + x = \infty$ and $\infty \geq x$ for all $x \in \mathbb{Z}_\infty$.

Proposition 2.13. *Let Q be a ring with $Q^{\text{reg}} = Q^*$, and let V be a discrete valuation ring of Q . Then there is a discrete valuation ν_V of Q such that the following hold.*

- (1) $V = \{x \in Q \mid \nu_V(x) \geq 0\}$, $\mathfrak{m}_V = \{x \in Q \mid \nu_V(x) > 0\}$, and $V^* = \{x \in Q^{\text{reg}} \mid \nu_V(x) = 0\}$.
- (2) The regular maximal ideal \mathfrak{m}_V is generated by any $t \in Q^{\text{reg}}$ with $\nu_V(t) = 1$. Such an element is called a *uniformizing parameter* V .
- (3) Let t be a uniformizing parameter for V . Then for any $x \in Q^{\text{reg}}$ there is a unique $a \in V^*$ such that $x = at^{\nu_V(x)}$.

Proof. See [16, Propositions 2.22, 2.23, and D.13 and Corollary 2.25]. \square

2.3. Admissible Rings. The theory of discrete valuation rings provides a general framework for the definition of value semigroups. So let R be a one-dimensional equidimensional semilocal Cohen–Macaulay ring with total ring of fractions Q_R . Then the set \mathcal{V}_R is non-empty and finite, and it contains only discrete valuation rings. Moreover, $\overline{R} = \bigcap_{V \in \mathcal{V}_R} V$ (see [21, Chapter II, Theorem 2.11]). We write ν_V for the discrete valuation $Q_R \rightarrow \mathbb{Z}_\infty$ corresponding to $V \in \mathcal{V}_R$ (see Proposition 2.13). Moreover, we denote

$$\nu_R = (\nu_V)_{V \in \mathcal{V}_R}: Q_R \rightarrow (\mathbb{Z}_\infty)^{\mathcal{V}_R}.$$

Definition 2.14. Let R be a one-dimensional equidimensional semilocal Cohen–Macaulay ring.

- (1) The *semigroup of values* (or *value semigroup*) of R is

$$\Gamma_R = \nu_R(R^{\text{reg}}) \subset \mathbb{N}^{\mathcal{V}_R}.$$

- (2) For a regular fractional ideal \mathfrak{J} of R we call

$$\Gamma_{\mathfrak{J}} = \nu_R(\mathfrak{J}^{\text{reg}}) \subset \mathbb{Z}^{\mathcal{V}_R}$$

the *value semigroup ideal* of \mathfrak{J} .

However, if we require value semigroups to satisfy conditions (S0) – (S2), we need to further restrict the available rings. For example, (S0) means that the normalization is finite. In order to obtain sums with or without cancellation of values we need sufficiently large and compatible residue fields. As explained in [12, Section 3] this leads to the following definition.

Definition 2.15. A one-dimensional equidimensional semilocal Cohen–Macaulay ring R is called *admissible* if the following hold.

- (1) R is *analytically reduced*, i.e. \widehat{R} is reduced.

- (2) R is *residually rational*, i.e. $R/\mathfrak{m} = \overline{R}/\mathfrak{n}$ for any $\mathfrak{m} \in \text{Max}(R)$ and for every $\mathfrak{n} \in \text{Max}(\overline{R})$ with $\mathfrak{n} \cap R = \mathfrak{m}$.
- (3) R has *large residue fields*, i.e. $|R/\mathfrak{m}| \geq |\mathcal{V}_{R/\mathfrak{m}}|$ for all $\mathfrak{m} \in \text{Max}(R)$.

Remark 2.16. Let R be a one-dimensional equidimensional semilocal reduced Cohen–Macaulay ring. The map

$$\begin{aligned} \mathcal{V}_R &\rightarrow \text{Min}(\widehat{R}), \\ V &\mapsto \{x \in R \mid \nu_V(x) = \infty\}, \end{aligned}$$

is a bijection, see [16, Remark 3.39].

Integral extensions of local ring are not necessarily local. For example, if R is reduced, then $|\text{Max}(\overline{R})| = |\text{Min}(R)|$. Therefore, we allow algebroid curves to be semilocal. Since they are complete, an algebroid curve is the product of finitely many local algebroid curves.

Definition 2.17. Let k be a field. An *algebroid curve* over k is a complete equidimensional reduced Noetherian semilocal k -algebra R of dimension one such that all residue fields of R are isomorphic to k (under the canonical surjections $R \rightarrow R/\mathfrak{m}$ for $\mathfrak{m} \in \text{Max}(R)$). We call an algebroid curve *admissible* if it is an admissible ring.

Proposition 2.18. *Let k be a field, and let R be an algebroid curve over k . If $|k| \geq |\text{Min}(R)|$, then R is admissible.*

Proof. See [16, Proposition 3.41]. □

Proposition 2.19. *Let R be a ring, and let A be an integral extension of R in Q_R .*

- (1) *If R is admissible, then A is admissible with $\mathcal{V}_R = \mathcal{V}_A$ and $\nu_R = \nu_A$. In particular, the value semigroup ideal of A as a regular fractional ideal of R coincides with its semigroup of values as an admissible ring.*
- (2) *If R is an (admissible) algebroid curve over a field k , then A is an (admissible) algebroid curve over k .*

Proof. See [16, Theorem 3.45]. □

Proposition 2.20. *Let k be a field, and let R be an algebroid curve over k . For every $\mathfrak{p} \in \text{Min}(R)$ and any uniformizing parameter $t_{\mathfrak{p}}$ for the discrete valuation ring $\overline{R}/\mathfrak{p}$ there is an isomorphism*

$$\begin{aligned} \phi_{\mathfrak{p}} : Q_R/\mathfrak{p}Q_R = Q_{R/\mathfrak{p}} &\rightarrow k[[T_{\mathfrak{p}}]][T_{\mathfrak{p}}^{-1}], \\ t_{\mathfrak{p}} &\mapsto T_{\mathfrak{p}}, \end{aligned}$$

with $\phi_{\mathfrak{p}}(\overline{R}/\mathfrak{p}) = k[[t_{\mathfrak{p}}]]$. This induces an isomorphism

$$\phi : Q_R = \prod_{\mathfrak{p} \in \text{Min}(R)} Q_R/\mathfrak{p}Q_R \rightarrow \prod_{\mathfrak{p} \in \text{Min}(R)} k[[T_{\mathfrak{p}}]][T_{\mathfrak{p}}^{-1}]$$

satisfying

$$\nu_R = \text{ord}_T \circ \phi.$$

Proof. See [16, Theorems 3.44 and A.74]. □

2.4. Blowups and Stable Ideals. Let again R be a one-dimensional semilocal Cohen–Macaulay ring. *Blowing up* a regular fractional ideal \mathfrak{J} of R , we obtain a sequence of integral extensions of R (see Proposition 2.7)

$$(1) \quad R \subset \mathfrak{J} : \mathfrak{J} \subset \mathfrak{J}^2 : \mathfrak{J}^2 \subset \dots$$

Note that by definition there is a regular element $x \in Q_R^{\text{reg}}$ such that $x\mathfrak{J}$ is a regular ideal of R . So Remark 2.5.(1) yields for any $n \in \mathbb{N}$

$$(x\mathfrak{J})^n : (x\mathfrak{J})^n = (x^n\mathfrak{J}^n) : (x^n\mathfrak{J}^n) = x^n(x^n)^{-1}(\mathfrak{J}^n : \mathfrak{J}^n) = \mathfrak{J}^n : \mathfrak{J}^n.$$

Thus, we may assume that \mathfrak{J} is a regular ideal of R .

We assign a name to the union of the integral extensions in (1).

Definition 2.21. Let R be a one-dimensional semilocal Cohen–Macaulay ring, and let \mathfrak{J} be a regular fractional ideal of R . The *blowup* of \mathfrak{J} is the ring $R^{\mathfrak{J}} = \bigcup_{n \in \mathbb{N}} (\mathfrak{J}^n : \mathfrak{J}^n)$.

In fact, the sequence (1) stabilizes after finitely many steps.

Proposition 2.22. Let R be a one-dimensional semilocal Cohen–Macaulay ring, and let \mathfrak{J} be a regular fractional ideal of R .

- (1) $R^{\mathfrak{J}}$ is a finitely generated R -module, and $R^{\mathfrak{J}} = \mathfrak{J}^n : \mathfrak{J}^n$ for all sufficiently large n .
- (2) There is a regular element $x \in R^{\mathfrak{J}}$ such that $\mathfrak{J}R^{\mathfrak{J}} = xR^{\mathfrak{J}}$.

Proof. See [15, Proposition 1.1]. □

It is of particular interest when the sequence (1) stabilizes at the first power of \mathfrak{J} .

Definition 2.23. Let R be a one-dimensional semilocal Cohen–Macaulay ring. A regular fractional ideal \mathfrak{J} of R is called *stable* if $R^{\mathfrak{J}} = \mathfrak{J} : \mathfrak{J}$.

Proposition 2.24. Let R be a one-dimensional semilocal Cohen–Macaulay ring. A regular fractional ideal \mathfrak{J} of R is stable if and only if there is an $x \in \mathfrak{J}$ such that $x(\mathfrak{J} : \mathfrak{J}) = \mathfrak{J}$.

Proof. If \mathfrak{J} is stable, then by Proposition 2.22.(2) there is an $x \in \mathfrak{J}$ such that $x(\mathfrak{J} : \mathfrak{J}) = \mathfrak{J}(\mathfrak{J} : \mathfrak{J}) = \mathfrak{J}$.

Conversely, suppose that there is an $x \in \mathfrak{J}$ such that $x(\mathfrak{J} : \mathfrak{J}) = \mathfrak{J}$. Then $\mathfrak{J}^2 = x(\mathfrak{J} : \mathfrak{J})\mathfrak{J} = x\mathfrak{J}$. Hence, \mathfrak{J} is stable by [15, Lemma 1.11(i)]. □

3. GOOD SEMIGROUPS

Based on conditions (S0) – (S1), Barucci, D’Anna, and Fröberg introduced the notion of *good semigroups* [11]. Before defining good semigroups, we describe these conditions more generally.

Definition 3.1. Let I be a finite set, and consider on \mathbb{Z}^I the natural partial order. We denote $N = \{\alpha \in \mathbb{Z}^I \mid \alpha \geq \mathbf{0}\}$. For a subset E of \mathbb{Z}^I we consider the following properties (see [8, Section 1] and [10, Section 2]).

- (E0) There exists an $\alpha \in \mathbb{Z}^I$ such that $\alpha + N \subset E$.
- (E1) If $\alpha, \beta \in E$, then $\inf(\alpha, \beta) = (\min\{\alpha_i, \beta_i\})_{i \in I} \in E$.
- (E2) For any $\alpha, \beta \in E$ with $\alpha_j = \beta_j$ for some $j \in I$, there exists an $\varepsilon \in E$ such that $\varepsilon_j > \alpha_j = \beta_j$ and $\varepsilon_i \geq \min\{\alpha_i, \beta_i\}$ for all $i \in I$, where equality is obtained whenever $\alpha_i \neq \beta_i$.

We are now equipped to introduce good semigroups.

Definition 3.2. Let S be a partially ordered cancellative commutative monoid with unique minimal element 0. Then S is called a *good semigroup* if it satisfies the following.

- (1) The group of differences D_S is generated by a finite set I such that the natural surjection $\mathbb{Z}^I \rightarrow D_S$ is an order preserving isomorphism. Note that I is then unique and contains only positive elements (see [12, Lemma 3.1.7]). In the following, we identify S with its image as a submonoid of \mathbb{Z}^I , and we refer to I without further explanation.
- (2) Properties (E1), (E1), and (E2) hold for S (as a subset of $D_S = \mathbb{Z}^I$).

These semigroups have a “tropical” semiring structure. In order to view this structure parallel to the structure of admissible rings, we restrict our considerations to certain semigroup ideals corresponding to the value sets of fractional ideals.

Definition 3.3. Let S be a good semigroup, embedded into its group of differences $D_S = \mathbb{Z}^I$.

- (1) We set $\overline{S} = \{\alpha \in \mathbb{Z}^I \mid \alpha \geq \mathbf{0}\} \cong \mathbb{N}^I$.
- (2) A good semigroup S is said to be a *numerical semigroup* if $|I| = 1$.
- (3) A *semigroup ideal* of a good semigroup S is a non-empty subset E of D_S such that $E + S \subset E$ and $\alpha + E \in S$ for some $\alpha \in S$.
- (4) A *good semigroup ideal* of a good semigroup S is a semigroup ideal E of S satisfying properties (E1) and (E2).
- (5) For a good semigroup S we denote by \mathcal{G}_S the set of all good semigroup ideals of S .

In fact, the definitions of good semigroups and good semigroup ideals describe value sets of admissible rings and their fractional ideals.

Proposition 3.4. *Let R be an admissible ring. Then Γ_R is a good semigroup, and $\Gamma_{\mathfrak{J}} \in \mathcal{G}_{\Gamma_R}$ for every $\mathfrak{J} \in \mathcal{R}_R$.*

Proof. See [12, Remark 4.1.2.(d)]. □

However, value semigroups of admissible rings are not characterized as good semigroups. An example of a good semigroup which is not the value semigroups of an admissible ring can be found in [11, Example 2.16].

Remark 3.5.

- (1) Let S be a good semigroup in \mathbb{Z}^I . Then its group of differences is $D_S = \mathbb{Z}^I$.
- (2) If S is a good semigroup, any semigroup ideal E of S satisfies property (E0) since S does and $E + S \subset E$.
- (3) If S is a numerical semigroup, then $E \in \mathcal{G}_S$ for every semigroup ideal E of S .
- (4) If S and S' are good semigroups with $S \subset S' \subset \overline{S}$, then $D_{S'} = D_S$, and hence $\overline{S'} = \overline{S}$. It follows that $\mathcal{G}_{S'} \subset \mathcal{G}_S$, and, in particular, $S' \in \mathcal{G}_S$.

See [12, Remark 4.1.2].

Definition 3.6. Let S be a good semigroup, and let $J \subset I$. We denote by $\text{pr}_J: \mathbb{Z}^I \rightarrow \mathbb{Z}^J$ the projection map. For a subset E of \mathbb{Z}^I we call $\text{pr}_J(E)$ the *projection* of E , and we denote it by E_J . We write $E_i = E_{\{i\}}$ for $i \in I$.

Proposition 3.7. *Let S be a good semigroup, and let $J \subset I$. Then the projection S_J is a good semigroup with $D_{S_J} = \mathbb{Z}^J$, and $E_J \in \mathcal{G}_{S_J}$ for every $E \in \mathcal{G}_S$.*

Proof. See [16, Proposition 4.59]. Also see [11, Proposition 2.2]. □

Definition 3.8.

- (1) Let R be an admissible ring, and let \mathfrak{J} be an R -submodule of Q_R . For $\alpha \in D_{\Gamma_R}$ we set

$$\mathfrak{J}^\alpha = \{x \in \mathfrak{J} \mid \nu(x) \geq \alpha\} = \mathfrak{J} \cap (Q_R)^\alpha.$$

- (2) Let S be a good semigroup, and let $E \subset D_S$. For $\alpha \in D_S$ we set

$$E^\alpha = \{\beta \in E \mid \beta \geq \alpha\} = E \cap (D_S)^\alpha.$$

Remark 3.9.

- (1) Let R be an admissible ring, and let $\mathfrak{J} \in \mathcal{R}_R$. For any $\alpha \in D_{\Gamma_R}$ the set \mathfrak{J}^α is a regular fractional ideal of R with $\Gamma_{\mathfrak{J}^\alpha} = (\Gamma_{\mathfrak{J}})^\alpha$.
- (2) Let S be a good semigroup, and let $E \in \mathcal{G}_S$. For any $\alpha \in D_S$ the set E^α is a good semigroup ideal of S .

3.1. Localization. Considering only local algebroid curves, Delgado found that $\mathbf{0}$ is the only element of the value semigroup with a zero component. Accordingly, a good semigroup S has been called *local* if $\mathbf{0}$ is the only element of S with a zero component [11, 12, 16].

In a local good semigroup S , the set $M_S = S \setminus \{\mathbf{0}\}$ is a good semigroup ideal, and it is called the *maximal ideal* of S . In fact, an admissible ring R is local if and only if its semigroup of values Γ_R is local, and if \mathfrak{m} is the maximal ideal of R , then $\Gamma_{\mathfrak{m}} = M_{\Gamma_R}$ (see [11, page 6]).

Any good semigroup decomposes uniquely into a product of finitely many local good semigroups. Moreover, this decomposition is compatible with localizations of admissible rings: If R is an admissible ring, then $\Gamma_R = \prod_{\mathfrak{m} \in \text{Max}(R)} \Gamma_{R_{\mathfrak{m}}}$, see [12, Theorems 3.2.2 and 4.1.6].

In this section, we establish an intrinsic notion of maximal ideals of good semigroups. This extends the above definition in the local case, and Corollary 3.15 yields again the above characterization of locality.

We furthermore remark that a similar intrinsic approach can be applied for minimal prime ideals of good semigroups. Then for an admissible ring R we can deduce a bijection $\text{Min } R \rightarrow \text{Min } \Gamma_R$, $\mathfrak{p} \mapsto \Gamma_{\mathfrak{p}}$, such that $\Gamma_{R/\mathfrak{p}} = \Gamma_R/\Gamma_{\mathfrak{p}}$ (using a “natural” definition of the *quotient semigroup* corresponding to the ring version), cf. [10, Proposition 1.2] and [16, Propositions 4.67 and 4.69].

Definition 3.10. Let S be a good semigroup.

- (1) A good semigroup ideal M of S with $M \subsetneq S$ is called *maximal* if for every good semigroup ideal E of S with $M \subset E \subsetneq S$ we have $E = M$.
- (2) We denote the set of maximal ideals of S by $\text{Max}(S)$.
- (3) S is called *local* if it has a unique maximal ideal M_S .

Remark 3.11. A numerical semigroup is a local good semigroup.

Lemma 3.12. Let S be a good semigroup. Then $\text{Max}(S) = \{S^{e_i} \mid i \in I\}$.

Proof. Let E be a good semigroup ideal of S with $E \subsetneq S$. Since E satisfies property (E1), there is an $i \in I$ such that $\alpha_i > 0$ for all $\alpha \in E$ (otherwise $\mathbf{0} \in E$). This implies $E \subset S^{e_i}$. Therefore, we have $\text{Max}(S) = \{S^{e_i} \mid i \in I\}$ (see Remark 3.9.(2)). \square

In particular, any good semigroup is “semilocal”.

Proposition 3.13. Let S be a good semigroup. For every $M \in \text{Max}(S)$ we denote $I_M = \{i \in I \mid \mathbf{0} \in M_i\}$.

- (1) $M = S^{e_{I \setminus I_M}}$ for every $M \in \text{Max}(S)$.
- (2) The family $(I \setminus I_M)_{M \in \text{Max}(S)}$ yields a partition of I .

Proof. On I we define the following equivalence relation: two elements i and j of I are equivalent if and only if $S^{e_i} = S^{e_j}$. Then $S^{e_i} = S^{e_{\bar{i}}}$ for every $i \in I$, where we denote by \bar{i} the equivalence class of i . Moreover, denoting by \bar{I} the set of equivalence classes there is by Lemma 3.12 a bijection

$$\begin{aligned} \text{Max}(S) &\rightarrow \bar{I}, \\ M &\mapsto I \setminus I_M, \\ S^{e_j} &\leftarrow J. \end{aligned} \quad \square$$

Remark 3.14. Let S be a good semigroup, and let $M \in \text{Max}(S)$. For every $i \in I_M$ pick an $\alpha^{(i)} \in M_S$ with $(\alpha^{(i)})_i = 0$, and set $\alpha = \inf(\alpha^{(i)} \mid i \in I_M)$. Then $\alpha_i > 0$ for all $i \in I \setminus I_M$ (see Proposition 3.13.(1)) and $\alpha_{I_M} = 0$.

The following immediate consequence of Proposition 3.13 relates our definition of maximal ideals and local good semigroups to the one used for example in [11, 12, 16].

Corollary 3.15. Let S be a good semigroup. Then S is local if and only if $\mathbf{0}$ is the only element in S with a zero component. Moreover, if S is local, then its maximal ideal is $M_S = S \setminus \{\mathbf{0}\} = S^1$. \square

Remark 3.16. Note that every numerical semigroup is local.

Proposition 3.17. Let R be an admissible ring. Then there is a bijection

$$\begin{aligned} \text{Max}(R) &\rightarrow \text{Max } S, \\ \mathfrak{m} &\mapsto \Gamma_{\mathfrak{m}}. \end{aligned}$$

(note that maximal ideals of R must be regular as R is equidimensional, see [16, Proposition B.27]). In particular, R is local if and only if Γ_R is local.

Proof. Let \mathfrak{i} be a proper ideal of R . Then there is an $V \in \mathcal{V}_R$ such that $\nu_V(x) > 0$ for all $x \in \mathfrak{i}$ since $\{x \in R \mid \nu(x) = \mathbf{0}\} = R^*$ (see [12, Equation (3.1)]). Therefore, $\text{Max}(R) = \{R^{e_V} \mid V \in \mathcal{V}_R\}$ (see Remark 3.9.(1)). Then claim follows from Lemma 3.12 since $\Gamma_{R^\alpha} = (\Gamma_R)^\alpha$ for any $\alpha \in D_{\Gamma_R}$ by Remark 3.9.(1) \square

With a notion of maximal ideals of good semigroups, we could use the semiring structure to define localizations in the “usual” way. It turns out, however, that it suffices to consider projections.

Definition 3.18. Let S be a good semigroup, and let $E \in \mathcal{G}_S$. For a maximal ideal M of S we define the *localization* of E with respect to M as $E_M = E_{I \setminus I_M}$, where $E_{I \setminus I_M}$ is the projection of Definition 3.6, and I_M is defined as in Proposition 3.13.

We can now formulate [12, Theorem 4.1.6] in terms of localizations.




FIGURE 1. The value semigroup (ideals) in Example 3.24.

Theorem 3.19. *Let S be a good semigroup.*

- (1) *For every maximal ideal M of S the good semigroup S_M (see Proposition 3.7) is local with maximal ideal M_M .*
- (2) $S = \prod_{M \in \text{Max}(S)} S_M$.
- (3) *Every good semigroup ideal E of S decomposes as $E = \prod_{M \in \text{Max}(S)} E_M$.*

Proof. See [11, (Proof of) Theorem 2.5 and Proposition 2.12] and Proposition 3.13.(2) and Remark 3.14. \square

Remark 3.20. Note that the decomposition of a good semigroup into its localizations as in Theorem 3.19.(2) is unique, see [11, Remark 2.6].

3.2. Differences. Products and quotients are important operations for fractional ideals (see Definition 2.3). The obvious combinatorial counterparts appear to be sums and differences.

Definition 3.21. Let S be a good semigroup, and let $E, F \subset_S$. The *difference* of E and F is

$$E - F = \{\alpha \in D_S \mid \alpha + F \subset E\}.$$

However, good semigroup ideals are not closed under these operations.

Remark 3.22. Let S be a good semigroup, and let E and F be two semigroup ideals of S . Then $E + F$ and $E - F$ are semigroup ideals of S (see [16, Lemma 4.18]). However, $E, F \in \mathcal{G}_S$ does in general neither imply $E + F \in \mathcal{G}_S$ (see Example 3.24 below) nor $E - F \in \mathcal{G}_S$ (see Example 3.25 below).

In particular, sums and differences are not compatible with there algebraic counterparts.

Remark 3.23. Let R be an admissible ring, and let $\mathfrak{I}, \mathfrak{J} \in \mathcal{R}_R$.

- (1) If $\mathfrak{I} \subset \mathfrak{J}$, then $\Gamma_{\mathfrak{I}} \subset \Gamma_{\mathfrak{J}}$.
- (2) $\Gamma_{\mathfrak{I}} + \Gamma_{\mathfrak{J}} \subset \Gamma_{\mathfrak{I} + \mathfrak{J}}$. In general, however, this inclusion is not an equality, see Example 3.24 below.
- (3) $\Gamma_{\mathfrak{I} : \mathfrak{J}} \subset \Gamma_{\mathfrak{I}} - \Gamma_{\mathfrak{J}}$. In general, however, this inclusion is not an equality, see Example 3.25 below.

See [12, Remark 3.1.10].

Example 3.24. Consider the admissible ring (see Proposition 2.18)

$$R = \mathbb{C} [[(-t_1^4, t_2), (-t_1^3, 0), (0, t_2), (t_1^5, 0)]] \subset \mathbb{C} [[t_1]] \times \mathbb{C} [[t_2]] = \overline{R},$$

and the R -submodules of Q_R

$$\begin{aligned} \mathfrak{I} &= \langle (t_1^3, t_2), (t_1^2, 0) \rangle_R, \\ \mathfrak{J} &= \langle (t_1^3, t_2), (t_1^4, 0), (t_1^5, 0) \rangle_R. \end{aligned}$$




FIGURE 2. The value semigroup (ideals) in Example 3.25, see [11, Example 3.3].

Then $\mathfrak{J}, \mathfrak{J} \in \mathcal{R}_R$ (see Remark 2.2). Moreover, Figure 1 shows that $\Gamma_{\mathfrak{J}} + \Gamma_{\mathfrak{J}}$ does not satisfy property (E2). Thus, $\Gamma_{\mathfrak{J}} + \Gamma_{\mathfrak{J}} \subsetneq \Gamma_{\mathfrak{J}\mathfrak{J}}$ by Remark 3.23.(2) and Proposition 3.4.

Example 3.25. Barucci, D’Anna and Fröberg showed in [11, Example 3.3] that for the local admissible ring (see Figure 2 and Propositions 2.18 and 3.15)

$$R = \mathbb{C}[[x_1, \dots, x_{11}]]$$

with $x_1 = (t_1^7, t_2^6)$, $x_2 = (t_1^6, t_2^7)$, $x_3 = (t_1^9, t_2^{11})$, $x_4 = (t_1^{10}, t_2^{10})$, $x_5 = (t_1^{11}, t_2^9)$, $x_6 = (t_1^{11}, t_2^{10})$, $x_7 = (t_1^{12}, t_2^{12})$, $x_8 = (t_1^{13}, -t_2^{13})$, $x_9 = (t_1^{20}, t_2^{12})$, $x_{10} = (t_1^{16}, t_2^{20})$, $x_{11} = (t_1^{12}, t_2^{20})$ with maximal ideal \mathfrak{m}_R the difference $\Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R}$ does not satisfy property (E2), see Figure 2. Thus, $\Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R} \subsetneq \Gamma_{\mathfrak{m}_R \cdot \mathfrak{m}_R}$ by Remark 3.23.(3) and Proposition 3.4.

Remark 3.26. Let S be a good semigroup, and let $\alpha \in D_S$.

- (1) Let E, F , and G be semigroup ideals of S . Then

$$(E - F) - G = E - (F + G) = (E - G) - F.$$

- (2) For any two semigroup ideals E and F of S we have

$$(\alpha + E) - F = \alpha + (E - F) = E - (-\alpha + F).$$

- (3) Let E, E', F , and F' be semigroup ideals of S . If $E \subset E'$ and $F \subset F'$, then

$$E - F' \subset E - F \subset E' - F.$$

- (4) For any $E \in \mathcal{G}_S$ we have $E - S = E$.

- (5) The map $\mathcal{G}_S \rightarrow \mathcal{G}_S, E \mapsto \alpha + E$ is a bijection.

Let S be a good semigroup, and let $E \in \mathcal{G}_S$. Since E satisfies property (E1), and since it is bounded from below, there is a unique minimal element of E (see [16, Lemma 4.12]). Moreover, the set $E - \overline{S}$ is a good semigroup ideal of S (see [12, Lemma 4.1.4]). This allows for the following definition.

Definition 3.27. Let S be a good semigroup, and let $E \in \mathcal{G}_S$.

- (1) We denote the *minimal element* of E by μ_E .
- (2) The *conductor ideal* of E is $C_E = E - \overline{S}$.
- (3) The *conductor* of E is $\gamma_E = \mu_{C_E} = \inf \{ \alpha \in D_S \mid \alpha + \overline{S} \subset E \}$.
- (4) We write $\tau_E = \gamma_E - \mathbf{1}$.

Remark 3.28. Let S be a good semigroup, and let $E \in \mathcal{G}_S$. Then

- (1) $E \subset \mu_E + \overline{S}$, and
- (2) $C_E = \gamma_E + \overline{S} \subset E$ (see [16, Remark 4.27]).

Lemma 3.29. *Let S be a good semigroup, and let E and F be good semigroup ideals of S . Then*

$$\gamma_{E-F} = \gamma_E - \mu_F.$$

Proof. See [12, Lemma 4.1.11] □

Lemma 3.30. *Let S be a good semigroup, and let $E \in \mathcal{G}_S$. For any $\alpha \in D_S$, we have $\alpha \in E$ if and only if $\inf(\alpha, \gamma_E) \in E$.*

Proof. See [12, Lemma 4.1.9]. □

Proposition 3.31. *Let R be an admissible ring, and let $\mathfrak{J}, \mathfrak{J}' \in \mathcal{R}_R$ with $\mathfrak{J} \subset \mathfrak{J}'$. Then $\mathfrak{J} = \mathfrak{J}'$ if and only if $\Gamma_{\mathfrak{J}} = \Gamma_{\mathfrak{J}'}$.*

Proof. See [12, Corollary 4.2.8]. □

Lemma 3.32. *Let R and R' be admissible rings such that $\mathfrak{C}_{R'} \subset R \subset R' \subset Q_R$. Then*

- (1) $R' \in \mathcal{R}_R$, and
- (2) if $\Gamma_R = \Gamma_{R'}$, then $R = R'$.

Proof.

- (1) Let $x \in (\mathfrak{C}'_R)^{\text{reg}} \subset R^{\text{reg}}$. Then $xR' \subset \mathfrak{C}_R \subset R$. Since $\emptyset \neq R^{\text{reg}} \subset (R')^{\text{reg}}$, this yields $R' \subset \mathcal{R}_R$.
- (2) Since $R' \in \mathcal{R}_R$ by (1), and since $R \in \mathcal{R}_R$, this follows from Proposition 3.31. □

Proposition 3.33. *Let R be an admissible ring, and let $\mathfrak{J} \in \mathcal{R}_R$. Then $\mathfrak{C}_{\mathfrak{J}} = (Q_R)^{\gamma_{\mathfrak{J}}}$.*

Proof. See [16, Proposition 4.56]. □

3.3. Stable Ideals. Stable semigroup ideals were introduced by Barucci, D'Anna, and Fröberg [11]. The definition reflects the characterization of stable fractional ideals in Proposition 2.24.

Definition 3.34. Let S be a good semigroup. A good semigroup ideal E of S is called *stable* if there is an $\alpha \in E$ such that $\alpha + (E - E) = E$. Note that then $\alpha = \mu_E$.

Stability of a fractional ideal \mathfrak{J} implies stability of the value semigroup ideal $\Gamma_{\mathfrak{J}}$. While the converse is in general not true, we have the following characterization.

Proposition 3.35. *Let R be an admissible ring, and let $\mathfrak{J} \in \mathcal{R}_R$. Then \mathfrak{J} is stable if and only if $\Gamma_{\mathfrak{J}; \mathfrak{J}} = \Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ and $\Gamma_{\mathfrak{J}}$ is stable.*

Proof. See [11, Proposition 3.14]. □

4. CANONICAL IDEALS AND GORENSTEIN PROPERTY

As seen in the previous section, good semigroup ideals do not behave well under differences in general. However, extending results by Jäger [9] and D'Anna [10], *canonical semigroup ideals* were introduced in [12].

Definition 4.1.

- (1) Let R be a one-dimensional equidimensional Cohen–Macaulay ring. A regular fractional ideal \mathfrak{K} is called a *canonical ideal* of R if $\mathfrak{K} : (\mathfrak{K} : \mathfrak{J}) = \mathfrak{J}$ for every $\mathfrak{J} \in \mathcal{R}_R$.
- (2) Let S be a good semigroup. A good semigroup ideal S is called a *canonical (semigroup) ideal* of S if $K \subset E$ implies $K = E$ for all $E \in \mathcal{G}_S$ with $\gamma_E = \gamma_K$.

It turns out that canonical semigroup ideals induce a duality on good semigroup ideals (see Theorem 4.5) in analogy to that on fractional ideals. So if S is a good semigroup, then a canonical semigroup ideal K is characterized by the property $K - (K - E) = E$ for every good semigroup ideal E of S . Moreover, the *dual* $K - E$ of E is in fact a good semigroup ideal.

Canonical fractional ideals and canonical semigroup ideals are compatible in the following sense: Canonical fractional ideals are characterized by their value semigroup ideals, and dualizing commutes with taking values.

Theorem 4.2. *Let R be an admissible ring.*

- (1) *A regular fractional ideal \mathfrak{K} of R is canonical if and only if $\Gamma_{\mathfrak{K}}$ is a canonical ideal of Γ_R .*
- (2) *If \mathfrak{K} is a canonical ideal of R , then $\Gamma_{\mathfrak{K};\mathfrak{J}} = \Gamma_{\mathfrak{K}} - \Gamma_{\mathfrak{J}}$ for every regular fractional ideal \mathfrak{J} of \mathfrak{K} .*

Proof. See [12, Theorems 5.3.2 and 5.3.4]. □

Before stating the main properties of canonical semigroup ideals in Theorem 4.5 we need a few technical definitions in order to describe canonical semigroup ideals explicitly.

Definition 4.3. Let I be a finite set, and let $\alpha \in \mathbb{Z}^I$.

- (1) For any subset J of I we set

$$\Delta_J(\alpha) = \{\beta \in \mathbb{Z}^s \mid \alpha_j = \beta_j \text{ for all } j \in J \text{ and } \alpha_i < \beta_i \text{ for all } i \in I \setminus J\}.$$

- (2) We set

$$\Delta(\alpha) = \bigcup_{i \in I} \Delta_i(\alpha),$$

where

$$\Delta_i(\alpha) = \Delta_{\{i\}}(\alpha)$$

for any $i \in I$.

- (3) For a subset E of \mathbb{Z}^s and any subset J of I we denote

$$\Delta_J^E(\alpha) = \Delta_J(\alpha) \cap E.$$

Similarly, we write $\Delta_i^E(\alpha)$ for given $i \in I$ as well as $\Delta^E(\alpha)$ when considering the union.

- (4) For any subset J of I we set

$$\overline{\Delta}_J(\alpha) = \{\beta \in \mathbb{Z}^s \mid \alpha_j = \beta_j \text{ for all } j \in J \text{ and } \alpha_i \leq \beta_i \text{ for all } i \in I \setminus J\},$$

and as in (1), (2), and (3) we define accordingly $\overline{\Delta}_i(\alpha)$, $\overline{\Delta}(\alpha)$, $\overline{\Delta}_J^E(\alpha)$, $\overline{\Delta}_i^E(\alpha)$, and $\overline{\Delta}^E(\alpha)$ for any $i \in I$ and for any subset E of \mathbb{Z}^I .

Turning Delgado's symmetry condition for value semigroups of Gorenstein algebroid curves [8] into the description of a semigroup ideal, D'Anna characterized "normalized" canonical ideals of admissible rings by having the following value semigroup ideal [10].

Definition 4.4. Let S be a good semigroup. The set

$$K_S^0 = \{\alpha \in D_S \mid \Delta^S(\tau_S - \alpha) = \emptyset\}$$

is called the (*normalized*) *canonical (semigroup) ideal* of S .

As for fractional ideals, canonical semigroup ideals induce a duality on good semigroup ideals, and they are characterized by this property.

Theorem 4.5. *Every good semigroup has a canonical ideal. Moreover, for any $K \in \mathcal{G}_S$ the following are equivalent.*

- (a) *K is a canonical ideal of S .*
- (b) *There is an $\alpha \in D_S$ such that $\alpha + K = K_S^0$.*
- (c) *For all $E \in \mathcal{G}_S$ we have $K - (K - E) = E$.*

If K is a canonical ideal of S , then the following hold.

- (1) *If $S \subset K \subset \overline{S}$, then $K = K_S^0$.*
- (2) *$K - E \in \mathcal{G}_S$ for every $E \in \mathcal{G}_S$.*
- (3) *If S' is a good semigroup with $S \subset S' \subset \overline{S}$, then $K' = K - S$ is a canonical ideal of S' .*

Proof. See [12, Theorem 5.2.6]. □

We are particularly interested in the case when an admissible ring, respectively a good semigroup, is its own canonical ideal.

Definition 4.6.

- (1) An admissible ring R is called *Gorenstein* if R is a canonical ideal of itself (see [12, Remark 5.1.4]).

(2) A good semigroup S is called *symmetric* if S is a canonical ideal of itself, i.e.

$$S = \{\alpha \in D_S \mid \Delta^S(\tau_S - \alpha) = \emptyset\},$$

see Remark 4.12.

Proposition 4.7. *An admissible ring R is Gorenstein if and only if Γ_R is symmetric.*

Proof. See [12, Proposition 5.3.6]. □

In the following we collect some results on canonical ideals for later reference.

Remark 4.8. Let R be a one-dimensional equidimensional Cohen–Macaulay ring, and let \mathfrak{K} be a canonical ideal of R . Then $\mathfrak{K} : \mathfrak{K} = \mathfrak{K} : (\mathfrak{K} : R) = R$ (see Remark 2.5.(3)).

Proposition 4.9. *Let R be a one-dimensional equidimensional semilocal Cohen–Macaulay ring with canonical ideal \mathfrak{K} . A regular fractional ideal \mathfrak{K}' of R is a canonical ideal of R if and only if there is an $x \in Q_R^{\text{reg}}$ such that $\mathfrak{K}' = x\mathfrak{K}$.*

Proof. See [17, Satz 2.8] and [12, Proposition 5.1.5]. □

Lemma 4.10. *Let R and A be one-dimensional equidimensional semilocal Cohen–Macaulay rings such that $A \in \mathcal{R}_R$. If \mathfrak{K} is a canonical ideal of R , then $\mathfrak{K} : A$ is a canonical ideal of A .*

Proof. See [12, Lemma 5.1.8]. □

Proposition 4.11. *Let S be a good semigroup, and let K be a canonical ideal of S . Then $E \subset K$ for every $E \in \mathcal{G}_S$ with $\gamma_E = \gamma_K$.*

Proof. See [12, Proof of Proposition 5.2.10]. □

Remark 4.12. The normalized canonical ideal K_S^0 of a good semigroup S is the unique canonical ideal K of S satisfying $S \subset K \subset \bar{S}$.

Lemma 4.13. *Let S be a good semigroup, and let K be a canonical semigroup ideal of S .*

- (1) $\Delta^K(\tau_K) = \emptyset$.
- (2) If S is local, then $\bar{\Delta}(\tau_K) \setminus \Delta(\tau_K) \subset K$.

Proof. By Theorem 4.5.(b) we may assume that $K = K_S^0$, and hence $\tau_K = \tau_S$.

- (1) Let $i \in I$, and let $\alpha \in \Delta_i^{K_S^0}(\tau_{K_S^0})$. Then

$$\begin{aligned} \alpha_i &= (\tau_S)_i, \\ \alpha_j &\geq (\tau_S)_j \text{ for all } j \in I \setminus \{i\}. \end{aligned}$$

Hence,

$$\begin{aligned} (\tau_S - \alpha)_i &= 0, \\ (\tau_S - \alpha)_j &\leq 0 \text{ for all } j \in I \setminus \{i\}. \end{aligned}$$

This implies $0 \in \Delta_i^S(\tau_S - \alpha)$, and therefore $\Delta^{K_S^0}(\tau_{K_S^0}) = \emptyset$ (see Definition 4.4). Also see [12, Lemma 5.2.8].

- (2) Note that $\bar{\Delta}(\tau_K) = \Delta(\tau_K)$ if $|I| = 1$. So we consider the case $|I| \geq 2$, and suppose that S is local. Let $i \in I$, and let

$$\alpha \in \bar{\Delta}_i^{K_S^0}(\tau_{K_S^0}) \setminus \Delta_i^{K_S^0}(\tau_{K_S^0}).$$

Then there is a $j \in I \setminus \{i\}$ such that

$$\begin{aligned} \alpha_i &= (\tau_S)_i, \\ \alpha_j &= (\tau_S)_j, \\ \alpha_k &\geq (\tau_S)_k \text{ for all } k \in I \setminus \{i, j\} \end{aligned}$$




FIGURE 3. The local algebroid curve $R = \mathbb{C}[[t^4, t^6 + t^7, t^{11}]]$ of Example 5.2 has semigroup of values $\Gamma_R = \langle 4, 6, 11, 13 \rangle$ with maximal ideal $\Gamma_{\mathfrak{m}_R} = M_{\Gamma_R} = \Gamma_R \setminus \{0\}$. Then $\Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R} = \langle 2, 7 \rangle$ and $\Gamma_{\mathfrak{m}_R:\mathfrak{m}_R} = \langle 4, 6, 7, 9 \rangle \subsetneq \Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R}$.

(see Definition 4.3). Hence,

$$\begin{aligned} (\tau_S - \alpha)_i &= 0, \\ (\tau_S - \alpha)_j &= 0, \\ (\tau_S - \alpha)_k &\leq 0 \text{ for all } k \in I \setminus \{i, j\}. \end{aligned}$$

Then $\Delta_i^S(\tau_S - \alpha) = \emptyset = \Delta_j^S(\tau_S - \alpha)$ by Corollary 3.15, and $\Delta_k^S(\tau_S - \alpha) = \emptyset$ for all $k \in I \setminus \{i, j\}$ since $S \geq \mathbf{0}$. Thus, $\Delta^S(\tau_S - \alpha) = \emptyset$, and therefore $\alpha \in K_S^0$. \square

Lemma 4.14. *Let S be a good semigroup, and let $E \in \mathcal{G}_S$. Then*

$$K_S^0 - E = \{\alpha \in D_S \mid \Delta^E(\tau_S - \alpha) = \emptyset\}.$$

Proof. See [10, Computation 3.3]. \square

5. GORENSTEIN ENDOMORPHISM RINGS

In this section, we prove the following characterization of the Gorensteinness of the endomorphism ring $\text{End}(\mathfrak{J})$, where \mathfrak{J} is a regular fractional ideal of an admissible ring.

Theorem 5.1. *Let R be an admissible ring, and let \mathfrak{J} be a fractional ideal of R . If $\mathfrak{J} : \mathfrak{J}$ is Gorenstein, then \mathfrak{J} is stable. More precisely, the following are equivalent.*

- (a) $\mathfrak{J} : \mathfrak{J}$ is Gorenstein
- (b) $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} = \Gamma_{\mathfrak{J}:\mathfrak{J}}$ and $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ is a symmetric semigroup.
- (c) \mathfrak{J} is stable and $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ is a symmetric semigroup.

The implication (a) \implies (c) relates Theorem 5.1 to the theory of almost Gorenstein rings. As shown in [14], for a local one-dimensional Cohen–Macaulay ring R , the endomorphism ring $\mathfrak{m} : \mathfrak{m}$ of the maximal ideal \mathfrak{m} is Gorenstein if and only if R is almost Gorenstein of maximal embedding dimension. However, R having maximal embedding dimension is equivalent to \mathfrak{m} being stable, see [15, Corollary 1.10].

From Section 3 we know that the difference $E - E$ is not necessarily good even if E is the value semigroup ideal of a fractional ideal. In particular, differences are in general not compatible with quotients. In Section 4, however, we learned that differences behave well in the context of dualizing, i.e. if canonical ideals are involved. Nonetheless, the following example shows that in Theorem 5.1.(b), only requiring symmetry of $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ would not be sufficient.

Example 5.2. Consider the local algebroid curve $R = \mathbb{C}[[t^4, t^6 + t^7, t^{11}]]$ with maximal ideal $\mathfrak{m}_R = \langle t^4, t^6 + t^7, t^{11} \rangle$. Then $\Gamma_R = \langle 4, 6, 11, 13 \rangle$ and $\Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R} = \langle 2, 7 \rangle$ are good semigroups (cf. Remark 3.5.(3)). Assume that $2 \in \Gamma_{\mathfrak{m}_R:\mathfrak{m}_R}$. Then there is by Proposition 2.13.(3) a unique $a \in (\mathbb{C}[[t]])^*$ such that $at^2 \in \mathfrak{m}_R : \mathfrak{m}_R$. Therefore, there is a $b \in \mathbb{C} \setminus \{0\}$ such that $at^2t^4 = b(t^6 + t^7)$. Hence, $a = b(1 + t)$. However, $b(t^2 + t^3)(t^6 + t^7) = b(t^8 + 2t^9 + t^{10})$ is not contained in \mathfrak{m}_R . This implies $2 \notin \Gamma_{\mathfrak{m}_R:\mathfrak{m}_R}$, and hence $\Gamma_{\mathfrak{m}_R:\mathfrak{m}_R} \subsetneq \Gamma_{\mathfrak{m}_R} - \Gamma_{\mathfrak{m}_R}$. See Figure 3.

In the following, we collect some preliminary results for the proof of Theorem 5.1.

Lemma 5.3. *Let S be a good semigroup, and let E be a good semigroup ideal of S with $\gamma_E = \gamma_S$. If $E - E$ is a symmetric good semigroup, then $E - E = K_S^0 - E$.*

Proof. Since $\gamma_E = \gamma_S = \gamma_{K_S^0}$, we have $E \subset K_S^0$ by Proposition 4.11, and hence $E - E \subset K_S^0 - E$ by Remark 3.26.(3).

Let now $\alpha \in D_S \setminus (E - E)$. Since $E - E$ is a symmetric good semigroup, we have

$$E - E = K_{E-E}^0 = \{\beta \in D_S \mid \Delta^{E-E}(\tau_{E-E} - \beta) = \emptyset\}$$

by Theorem 4.5.(1). Hence, there is a $\delta \in \Delta^{E-E}(\tau_{E-E} - \alpha)$. This implies

$$\delta + \mu_E \in \Delta(\tau_{E-E} + \mu_E - \alpha) \cap ((E - E) + E) \subset \Delta(\tau_E - \alpha) \cap E = \Delta^E(\tau_S - \alpha)$$

by Lemma 3.29. Therefore, $\alpha \notin K_S^0 - E$ by Lemma 4.14. This yields $K_S^0 - E \subset E - E$. Hence, $E - E = K_S^0 - E$. \square

The combinatorial version of implication (a) \implies (c) also holds true for value semigroup ideals.

Proposition 5.4. *Let S be a good semigroup, and let $E \in \mathcal{G}_S$. If $E - E$ is a symmetric good semigroup, then E is stable.*

Proof. First suppose that $\gamma_E = \gamma_S$. By Lemma 3.29 and Theorem 4.5.(1) and (3) we have

$$\begin{aligned} K_{E-E}^0 &= (\gamma_{E-E} - \gamma_S) + (K_S^0 - (E - E)) \\ &= (\gamma_E - \mu_E - \gamma_S) + (K_S^0 - (E - E)) \\ &= -\mu_E + (K_S^0 - (E - E)). \end{aligned}$$

Therefore, Theorem 4.5 and Lemma 5.3 yield

$$E = K_S^0 - (K_S^0 - E) = K_S^0 - (E - E) = \mu_E + K_{E-E}^0 = \mu_E + (E - E).$$

Thus, E is stable.

Let now γ_E be arbitrary. Then $\gamma_{\gamma_S - \gamma_E + E} = \gamma_S$, and $(\gamma_S - \gamma_E + E) - (\gamma_S - \gamma_E + E) = E - E$ is symmetric by Remark 3.26.(2). Thus, $(\gamma_S - \gamma_E + E)$ is stable. This implies

$$\begin{aligned} E - E &= (\gamma_S - \gamma_E + E) - (\gamma_S - \gamma_E + E) = -\mu_{\gamma_S - \gamma_E + E} + (\gamma_S - \gamma_E + E) \\ &= -(\gamma_S - \gamma_E + \mu_E) + \gamma_S - \gamma_E + E = -\mu_E + E. \end{aligned}$$

Therefore, E is stable. \square

Lemma 5.5. *Let R be an admissible ring, and let $\mathfrak{J} \in \mathcal{R}_R$. Then $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} \subset K_{\Gamma_{\mathfrak{J}, \mathfrak{J}}}^0$.*

Proof. By Remark 3.26.(2) we may assume that $\gamma_{\Gamma_{\mathfrak{J}}} = \gamma_{\Gamma_R}$. Suppose that there is an $\alpha \in (\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}) \setminus K_{\Gamma_{\mathfrak{J}, \mathfrak{J}}}^0$. Then there is a

$$\beta \in \Delta^{\Gamma_{\mathfrak{J}, \mathfrak{J}}}(\tau_{\Gamma_{\mathfrak{J}, \mathfrak{J}}} - \alpha)$$

(see Definition 4.4). So Lemma 3.29 yields

$$\beta + \mu_{\Gamma_{\mathfrak{J}}} \in \Delta(\tau_{\Gamma_{\mathfrak{J}, \mathfrak{J}}} + \mu_{\Gamma_{\mathfrak{J}}} - \alpha) \cap ((\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}) + \Gamma_{\mathfrak{J}}) \subset \Delta^{\Gamma_{\mathfrak{J}}}(\tau_{\Gamma_{\mathfrak{J}}} - \alpha).$$

This implies by Lemma 4.14 $\alpha \notin K_{\Gamma_R}^0 - \Gamma_{\mathfrak{J}}$, contradicting $\alpha \in \Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} \subset K_{\Gamma_R}^0 - \Gamma_{\mathfrak{J}}$ (see Remark 3.26.(3), Definition 4.1.(2), and Proposition 4.11). \square

Lemma 5.6. *Let R be an admissible ring, and let $\mathfrak{J} \in \mathcal{R}_R$. If $\mathfrak{J} : \mathfrak{J}$ is Gorenstein, then $\Gamma_{\mathfrak{J} : \mathfrak{J}} = \Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$.*

Proof. If $\mathfrak{J} : \mathfrak{J}$ is Gorenstein, then Remark 3.23.(3), Theorem 4.5.(1), Proposition 4.7, and Lemma 5.5 yield

$$\Gamma_{\mathfrak{J} : \mathfrak{J}} \subset \Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} \subset K_{\Gamma_{\mathfrak{J}, \mathfrak{J}}}^0 = \Gamma_{\mathfrak{J} : \mathfrak{J}}. \quad \square$$

Collecting the above results, we obtain the proof of Theorem 5.1.

Proof of Theorem 5.1.

(a) \implies (b): If $\mathfrak{J} : \mathfrak{J}$ is Gorenstein, then $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} = \Gamma_{\mathfrak{J} : \mathfrak{J}}$ by Lemma 5.6. Hence, $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ is symmetric by Proposition 4.7.

- (b) \implies (c): If $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ is symmetric, then $\Gamma_{\mathfrak{J}}$ is stable by Proposition 5.4. So if $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} = \Gamma_{\mathfrak{J}:\mathfrak{J}}$, then Proposition 3.35 implies that \mathfrak{J} is stable.
- (c) \implies (a): If \mathfrak{J} is stable, then $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}} = \Gamma_{\mathfrak{J}:\mathfrak{J}}$ by Proposition 3.35. So if $\Gamma_{\mathfrak{J}} - \Gamma_{\mathfrak{J}}$ is symmetric, then $\Gamma_{\mathfrak{J}:\mathfrak{J}}$ is symmetric, and hence $\mathfrak{J} : \mathfrak{J}$ is Gorenstein by Proposition 4.7. \square

6. SYMMETRIC SEMIGROUPS

The aim for the rest of the paper is to characterize in Theorem 7.1 the Gorenstein local admissible rings for which the endomorphism ring of the maximal ideal is Gorenstein again. To achieve this, we first prove the combinatorial version in Theorem 6.4.

6.1. Differences in Symmetric Semigroups. By Proposition 4.7, an admissible ring is Gorenstein if and only if its value semigroup is symmetric. We therefore begin by studying differences in symmetric semigroups more closely.

Lemma 6.1. *Let S be a good semigroup, and let $\alpha \in D_S$ with $\alpha \leq \gamma_S$. Then $S^\alpha - S^\alpha = S - S^\alpha$.*

Proof. Definition 3.27.(3) and Remark 3.28.(2) yield $\mu_{S-S^\alpha} \geq \gamma_S - \gamma_{S^\alpha} = \gamma_S - \gamma_S = 0$. This implies with Remark 3.28.(1)

$$(S - S^\alpha) + S^\alpha \subset (D_S)^\alpha \cap S = S^\alpha,$$

and hence $S - S^\alpha \subset S^\alpha - S^\alpha$.

The claim follows since $S^\alpha - S^\alpha \subset S - S^\alpha$ by Remark 3.26.(3). \square

Lemma 6.2. *Let S be a good symmetric semigroup, and let $\alpha \in D_S$ with $\alpha \leq \gamma_S$. Then $S^\alpha - S^\alpha$ is a good semigroup with $S \subset S^\alpha - S^\alpha \subset \overline{S}$. In particular, if $S = \overline{S}$, then $S^\alpha - S^\alpha = S$.*

Proof. A straightforward argument shows that $S^\alpha - S^\alpha$ is a partially ordered cancellative commutative monoid with $D_{S^\alpha - S^\alpha} = D_S$ and $S \subset S^\alpha - S^\alpha \subset \overline{S^\alpha - S^\alpha} = \overline{S}$ (see [16, Proposition 4.25]). Moreover, $S^\alpha - S^\alpha$ satisfies property (E0) by Remarks 3.5.(2), 3.22, and 3.9.(2).

Since S is symmetric, i.e. S is a canonical ideal of itself, Remark 3.9.(2), Theorem 4.5.(2), and Lemma 6.1 yield

$$S^\alpha - S^\alpha = S - S^\alpha \in \mathcal{G}_S.$$

Thus, $S^\alpha - S^\alpha$ satisfies properties (E1) and (E2). \square

The following Lemma was proved for numerical semigroups in [22, Satz 1.9.(e)].

Lemma 6.3. *Let S be a local symmetric semigroup with maximal ideal M_S . If $S \neq \overline{S}$, then $M_S - M_S = S \cup \Delta(\tau_S)$.*

Proof. Lemmas 3.29 and 6.2 imply $S \cup \Delta^S(\tau_S) \subset M_S - M_S$ since $\mu_{M_S} \geq \mathbf{1}$ by Corollary 3.15.

Now suppose that there is an $\alpha \in (M_S - M_S) \setminus (S \cup \Delta(\tau_S))$. Since S is symmetric, there is a $\beta \in \Delta^S(\tau_S - \alpha)$. By assumption and Lemma 4.13.(2) we have $\alpha \notin \overline{\Delta^S}(\tau_S)$. Therefore, $\tau_S - \alpha > \mathbf{0}$, and hence $\beta \in \Delta^{M_S}(\tau_S - \alpha)$. This implies

$$\alpha + \beta \in (\alpha + \Delta(\tau_S - \alpha)) \cap ((M_S - M_S) + M_S) \subset \Delta^{M_S}(\tau_S) \subset \Delta^S(\tau_S) = \emptyset,$$

where the last equality follows from Lemma 4.13.(1) since S is symmetric. \square

6.2. Local Symmetric Semigroups. Let S be a local good semigroup with maximal ideal M_S . We can now fully classify when S and $M_S - M_S$ are symmetric.

Theorem 6.4. *Let S be a local good semigroup with maximal ideal M_S . The following are equivalent.*

- (a) Every good semigroup S' with $S \subset S' \subset \overline{S}$ is symmetric.
- (b) S and $M_S - M_S$ are symmetric good semigroups.
- (c) There is an $n \in 2\mathbb{N}$ such that

$$S = \langle 2, n+1 \rangle,$$

or there is an $n \in 2\mathbb{N} + 1$ such that

$$S = \langle \mathbf{1} \rangle \cup \left(\left(\binom{n+1}{2} \right)_{i \in I} + \mathbb{N}^I \right),$$

where $|I| = 2$.

Note that the equivalence (a) \iff (c) of Theorem 6.4 was shown in [22, Lemma 1.15] for numerical semigroups.

Lemma 6.5. *Let S be a numerical symmetric semigroup. Then $\gamma_S = 2|\overline{S} \setminus S|$.*

Proof. See [22, Satz 1.9.(c)]. □

Proposition 6.6. *Let S be a numerical symmetric semigroup. If $M_S - M_S$ is symmetric (see Remark 3.16), then $S = \langle 2, n+1 \rangle$ for some $n \in 2\mathbb{N}$.*

Proof. Suppose that $S \neq \overline{S}$ (see Lemma 6.2), and that $M_S - M_S$ is symmetric. Since $M_S - M_S = S \cup \{\tau_S\}$ by Lemma 6.3, Lemma 6.5 yields

$$\gamma_{M_S - M_S} = 2|\overline{S} \setminus (M_S - M_S)| = 2(|\overline{S} \setminus S| - 1) = \gamma_S - 2.$$

Thus, we have $2 = \gamma_S - \gamma_{M_S - M_S} = \mu_{M_S} \in M_S \subset S$ by Lemma 3.29, and by Corollary 3.15 the only smaller element in S is 0. This implies $S = \langle 2, \gamma_S + 1 \rangle$, where $\gamma_S = 2|\overline{S} \setminus S|$ is even. □

Lemma 6.7. *Let S be a local symmetric semigroup. If $M_S - M_S$ is symmetric, then $\alpha - \mu_{M_S} \in S$ for every $\alpha \in M_S \setminus C_S$.*

Proof. By Proposition 5.4 and Lemma 6.3 we have

$$\alpha \in M_S = \mu_{M_S} + (M_S - M_S) = \mu_{M_S} + (S \cup \Delta(\tau_S)) = (\mu_{M_S} + S) \cup \Delta(\tau_S + \mu_{M_S}).$$

Assume that $\alpha \in \Delta(\tau_S + \mu_{M_S})$. Then $\alpha \geq \gamma_S$ since $\mu_{M_S} \geq 1$ by Corollary 3.15. But this is a contradiction to the choice of α . Hence, $\alpha - \mu_{M_S} \in S$. □

Lemma 6.8. *Let S be a local symmetric semigroup. If $M_S - M_S$ is symmetric, then for every $\alpha \in M_S \setminus C_S$ there is an $n \in \mathbb{N}$ such that $\alpha = n\mu_{M_S}$. In particular, $S = \langle \mu_{M_S} \rangle \cup C_S$.*

Proof. Let $\alpha \in M_S \setminus C_S$. Since $S = \{0\} \cup M_S$ by Corollary 3.15, and since $C_S = \gamma_S + \overline{S}$ by Remark 3.28.(2), repeatedly applying Lemma 6.7 yields $\alpha - m\mu_{M_S} \in S$ for all $m \in \mathbb{N}$ satisfying $(m-1)\mu_{M_S} < \alpha$. Since α is finite, there is $n = \max\{m \in \mathbb{N} \mid (m-1)\mu_{M_S} < \alpha\}$. Then $\alpha - n\mu_{M_S} \in S$ implies $\alpha - n\mu_{M_S} \geq 0$. Hence, $n\mu_{M_S} \not\prec \alpha$ yields $\alpha = n\mu_{M_S}$. The particular claim follows since $S = \{0\} \cup M_S$ by Corollary 3.15. □

Lemma 6.9. *Let S be a local symmetric semigroup. If $M_S - M_S$ is symmetric, then $|I| \leq 2$.*

Proof. If $|I| \geq 3$, then $\overline{\Delta}(\tau_S) \setminus \Delta(\tau_S) \subset S$ contradicts Lemma 6.8. □

Proposition 6.10. *Let S be a local symmetric semigroup with $|I| = 2$. If $M_S - M_S$ is symmetric, then $\mu_{M_S} = 1$. Moreover, there is an $n \in 2\mathbb{N} + 1$ such that*

$$S = \langle 1 \rangle \cup \left(\left(\frac{n+1}{2} \right)_{i \in I} + \mathbb{N}^I \right).$$

Proof. We may suppose that $I = \{1, 2\}$. Since $\mu_{M_S} \geq 1$ by Corollary 3.15, Lemma 3.29 yields $\gamma_{M_S - M_S} = \gamma_{M_S} - \mu_{M_S} \leq \gamma_S - 1 = \tau_S$ (note that $S \neq \overline{S}$ by Corollary 3.15). Assume that $\gamma_{M_S - M_S} < \tau_S$. Then $\overline{\Delta}_i(\tau_S - \mathbf{e}_i) \subset M_S - M_S$ for an $i \in I$. However, $\overline{\Delta}_i(\tau_S - \mathbf{e}_i) \not\subset S$ by Lemma 6.8, and $\overline{\Delta}_i(\tau_S - \mathbf{e}_i) \cap \Delta(\tau_S) = \emptyset$ by Definition 4.3. So with Lemma 6.3 we obtain the contradiction $\overline{\Delta}_i(\tau_S - \mathbf{e}_i) \not\subset S \cup \Delta(\tau_S) = M_S - M_S$. Therefore, $\gamma_{M_S - M_S} = \tau_S$, and Lemma 3.29 yields $\mu_{M_S} = \gamma_S - \gamma_{M_S - M_S} = \gamma_S - \tau_S = 1$. Thus, we have $S = \langle 1 \rangle \cup C_S$ by Lemma 6.8.

Without loss of generality we may assume that $(\gamma_S)_1 \leq (\gamma_S)_2$. Let $\alpha \in \mathbb{N}$ with $\alpha \geq (\gamma_S)_1$. Then Lemma 3.30 implies $(\alpha, \alpha) + \mathbf{Ne}_1 \subset S$. Let now $m \in \mathbb{N}$. Then $(\alpha + m, \alpha + m) \in S$. Moreover, $\alpha \geq (\gamma_S)_1$ and $\max\{\alpha + m, (\gamma_S)_2\} \geq (\gamma_S)_2$ imply $(\alpha, \max\{\alpha + m, (\gamma_S)_2\}) \in S$. Since S satisfies property (E1), this yields

$$(\alpha, \alpha + m) = \inf((\alpha + m, \alpha + m), (\alpha, \max\{\alpha + m, (\gamma_S)_2\})) \in S,$$

and hence $(\alpha, \alpha) + \mathbf{Ne}_2 \subset S$. Since also $(\alpha, \alpha) + \mathbf{Ne}_1 \subset S$ by Lemma 3.30, we obtain $((\gamma_S)_1, (\gamma_S)_1) + \overline{S} \subset S$, and hence $\gamma_S = ((\gamma_S)_1, (\gamma_S)_1)$.




FIGURE 4. The symmetric numerical semigroup $S = \langle 2, 6 + 1 \rangle$. Every good semigroup S' with $S \subset S' \subset \overline{S}$ is of the same form, e.g. $S' = \langle 2, 4 + 1 \rangle$.




FIGURE 5. Every good semigroup containing $S = \langle (1, 1) \rangle \cup \left(\left(\frac{9+1}{2}, \frac{9+1}{2} \right) + \mathbb{N}^2 \right)$ and any point of $\overline{\Delta}((3, 3)) \setminus \{(3, 3)\}$ must also contain the good semigroup $S' = \langle (1, 1) \rangle \cup \left(\left(\frac{5+1}{2}, \frac{5+1}{2} \right) + \mathbb{N}^2 \right)$ in order to satisfy properties (E1) and (E2).

Therefore, setting $n = 2(\gamma_S)_1 - 1 \in 2\mathbb{N} + 1$ we obtain

$$S = \langle \mathbf{1} \rangle \cup C_S = \langle \mathbf{1} \rangle \cup \left(\left(\frac{n+1}{2}, \frac{n+1}{2} \right) + \mathbb{N}^I \right)$$

(see Remark 3.28.(2)). □

Proof of Theorem 6.4.

- (a) \implies (b): Suppose that any good semigroup S' with $S \subset S' \subset \overline{S}$ is symmetric. Then, in particular, S is symmetric. Therefore, $M_S - M_S = S - M_S$ (see Corollary 3.15 and Lemma 6.1) is a good semigroup by Lemma 6.2. Since $S \subset M_S - M_S \subset \overline{S}$ by Lemma 6.2, it is symmetric by assumption.
- (b) \implies (c): Suppose that S and $M_S - M_S$ are symmetric semigroups. Then $|I| \leq 2$ by Lemma 6.9. The statement follows from Propositions 6.6 and 6.10.
- (c) \implies (a): If $S = \langle 2, n + 1 \rangle$ for some $n \in 2\mathbb{N}$, the proof is in [22, Lemma 1.15].

So suppose that $S = \langle \mathbf{1} \rangle \cup \left(\left(\frac{n+1}{2} \right)_{i \in I} + \mathbb{N}^I \right)$ for some $n \in 2\mathbb{N} + 1$. Let S' be a good semigroup with $S \subset S' \subset \overline{S}$. Since S' has to satisfy properties (E1) and (E2), it is easy to see that there is an $m \in 2\mathbb{N} + 1$ such that $S' = \langle \mathbf{1} \rangle \cup \left(\left(\frac{m+1}{2} \right)_{i \in I} + \overline{S} \right)$ (see Figure 5). □

7. GORENSTEIN ALGEBROID CURVES

We conclude the paper with the following characterization of Gorenstein admissible rings which only have Gorenstein integral extensions in their ring of fractions. The equivalence of statements (a), (e), and (e) in Theorem 7.1 was proved for irreducible admissible rings already in [22, Proposition 2.22].

Theorem 7.1. *Let R be a local admissible ring with maximal ideal \mathfrak{m}_R . The following are equivalent.*

- (a) *Every integral extension of R in its total ring of fractions Q_R is Gorenstein.*
- (b) *R and $\mathfrak{m}_R : \mathfrak{m}_R$ are Gorenstein.*
- (c) *R is Gorenstein and has maximal embedding dimension.*

- (d) Γ_R and $M_{\Gamma_R} - M_{\Gamma_R}$ are symmetric.
(e) There is an $n \in 2\mathbb{N}$ such that

$$\Gamma_R = \langle 2, n+1 \rangle,$$

or there is an $n \in 2\mathbb{N} + 1$ such that

$$\Gamma_R = \langle \mathbf{1} \rangle \cup \left(\left(\frac{n+1}{2} \right)_{V \in \mathcal{V}_R} + \mathbb{N}^{\mathcal{V}_R} \right),$$

where $|\mathcal{V}_R| = 2$.

If R contains a field, and if $k = R/\mathfrak{m}_R$ is algebraically closed, then the above statements are equivalent to the following.

- (f) There is a surjective k -algebra homomorphism

$$\phi: k[[x, y]] \rightarrow \widehat{R}$$

with

$$\ker \phi = \langle x^{n+1} - y^2 \rangle$$

for some $n \in \mathbb{N}$ (note that this is the same n as in (e)).

Proof.

- (a) \implies (b): This follows trivially as R and $\mathfrak{m}_R : \mathfrak{m}_R$ are integral extensions of R .
(b) \implies (d): If R is Gorenstein, then Γ_R is symmetric by Proposition 4.7. Hence, Remark 2.5.(2), Proposition 2.7, Theorem 4.2.(2), Proposition 3.17, and Lemma 6.1 yield

$$\Gamma_{\mathfrak{m}_R : \mathfrak{m}_R} = \Gamma_{R : \mathfrak{m}_R} = \Gamma_R - \Gamma_{\mathfrak{m}_R} = \Gamma_R - M_{\Gamma_R} = M_{\Gamma_R} - M_{\Gamma_R}.$$

Therefore, Γ_R and $M_{\Gamma_R} - M_{\Gamma_R}$ are symmetric by Propositions 2.19.(1) and 4.7.

- (d) \implies (e): This follows from Theorem 6.4.
(e) \implies (a): Suppose that

$$\Gamma_R = \langle 2, n+1 \rangle,$$

for some $n \in 2\mathbb{N}$ or

$$\Gamma_R = \langle \mathbf{1} \rangle \cup \left(\left(\frac{n+1}{2} \right)_{V \in \mathcal{V}_R} + \mathbb{N}^{\mathcal{V}_R} \right)$$

for some $n \in 2\mathbb{N} + 1$. Let A be an integral extension of R in Q_R . Then A is an admissible ring by Proposition 2.19.(1), and $A \in \mathcal{R}_R$ with $\Gamma_R \subset \Gamma_A \subset \overline{\Gamma_R}$ (see Remark 3.23.(1)). Thus, Γ_A is symmetric by Theorem 6.4 (also see Proposition 2.19.(1)), and hence A is Gorenstein by Proposition 4.7.

- (b) \implies (c): If $\mathfrak{m}_R : \mathfrak{m}_R$ is Gorenstein, then R has maximal embedding dimension by [14, Theorem 5.1].
(c) \implies (b): If R is Gorenstein, then it is almost Gorenstein, see [14, p. 364]. Since R has maximal embedding dimension, also $\mathfrak{m}_R : \mathfrak{m}_R$ is Gorenstein by [14, Theorem 5.1].

Now suppose that R contains a field, and that $k = R/\mathfrak{m}_R$ is algebraically closed. By [23, Corollary of Theorem 56] \widehat{R} is local with residue field k . Since k is algebraically closed, we have $\text{char } k = 0$. In particular, this means $|k| \geq |\text{Min}(\widehat{R})|$. By [24, Corollary 2.1.8] we have $\dim \widehat{R} = \dim R1$. In particular, \widehat{R} is equidimensional. Moreover, \widehat{R} is Noetherian by [25, Theorem 10.26], and it is reduced since R is admissible. Finally, \widehat{R} is a k -algebra by the Cohen Structure Theorem (see [19, Theorem 7.7]). Thus, \widehat{R} is an algebroid curve over R . Since $\Gamma_R = \Gamma_{\widehat{R}}$ by [12, Theorem 3.3.5], we may assume that R is an algebroid curve over k .

- (e) \implies (f): First suppose that there is an $n \in 2\mathbb{N}$ such that $\Gamma_R = \langle 2, n+1 \rangle$. We identify $R \subset Q_R = k[[t]][t^{-1}]$ (see Proposition 2.20). Since $2 \in \Gamma_R$, there is $x \in R$ with $\nu_R(x) = 2$. Then by Proposition 2.13.(2) there is an $a \in (k[[t]])^*$ such that $x = at^2$. Since k is algebraically closed, there is a $b \in (k[[t]])^*$ such that $b^2 = a$, and hence $x = (bt)^2$. By Propositions 2.13.(1) and (2) and 2.20 bt is a uniformizing parameter for $k[[t]]$. Therefore, we may by Proposition 2.20 assume that $x = t^2$.

Since $\mathfrak{C}_R = t^{\nu_R} k[[t]]$ by Proposition 3.33, we have $t^{\nu_R+1} \in R$. Then $R' = k[[t^2, t^{\nu_R+1}]]$ is an algebroid curve over k with $\overline{R'} = \overline{R} = k[[t]]$. Moreover, we have $\Gamma_R = \Gamma_{R'}$, and hence

$$\mathfrak{C}_R = \mathfrak{C}_{R'} \subset R' \subset R \subset Q_{R'}$$

by Proposition 3.33. Therefore, $R = k[[t^2, t^{\gamma_{\Gamma_R} + 1}]]$ by Lemma 3.32.(2). Then the k -algebra homomorphism

$$\begin{aligned}\phi: k[[x, y]] &\rightarrow R, \\ x &\mapsto t^2, \\ y &\mapsto t^{n+1},\end{aligned}$$

has kernel $\langle x^{n+1} - y^2 \rangle$ (see [26, Algorithm 3.6 and Corollary 3.8] or [16, Proposition 5.54]).

Now suppose that there is an $n \in 2\mathbb{N} + 1$ such that

$$\Gamma_R = \langle (1, 1) \rangle \cup \left(\left(\frac{n+1}{2}, \frac{n+1}{2} \right) + \mathbb{N}^{\nu_R} \right)$$

We identify $R \subset Q_R = k[[t_1]][t_1^{-1}] \times k[[t_2]][t_2^{-1}]$ (see Proposition 2.20). Since $(1, 1) \in \Gamma_R$, we may by Propositions 2.13 and 2.20 choose the parameters t_1 and t_2 such that $(t_1, t_2) \in R$.

Set $\gamma = \frac{n+1}{2}$. Then $\gamma_{\Gamma_R} = (\gamma, \gamma)$. So by Proposition 3.33 we have $(t_1^\gamma, -t_2^\gamma) \in \mathfrak{C}_R \subset R$. Then $R'' = k[[t_1, t_2], (t_1^\gamma, -t_2^\gamma)]]$ is an algebroid curve over k with $\overline{R''} = \overline{R} = k[[t_1]] \times k[[t_2]]$. Moreover, we have $\Gamma_R = \Gamma_{R''}$, and hence

$$\mathfrak{C}_R = \mathfrak{C}_{R''} \subset R'' \subset R \subset Q_{R''}$$

by Proposition 3.33. Therefore, $R = k[[t_1, t_2], (t_1^\gamma, -t_2^\gamma)]]$ by Lemma 3.32.(2). Then the k -algebra homomorphism

$$\begin{aligned}\psi: k[[x, y]] &\rightarrow R, \\ x &\mapsto (t_1, t_2), \\ y &\mapsto \left(t_1^{\frac{n+1}{2}}, -t_2^{\frac{n+1}{2}} \right),\end{aligned}$$

has kernel $\langle x^{n+1} - y^2 \rangle$ (see [26, Algorithm 3.6 and Corollary 3.8] or [16, Proposition 5.54]).

(f) \implies (e): This follows by computing the semigroup of values of $k[[x, y]]/\langle x^{n+1} - y^2 \rangle$ (see [27]). \square

REFERENCES

- [1] Graham J. Leuschke. Non-commutative crepant resolutions: scenes from categorical geometry. In *Progress in commutative algebra 1*, pages 293–361. de Gruyter, Berlin, 2012.
- [2] H. Grauert and R. Remmert. *Analytische Stellenalgebren*. Springer-Verlag, Berlin-New York, 1971. Unter Mitarbeit von O. Riemenschneider, Die Grundlehren der mathematischen Wissenschaften, Band 176.
- [3] Theo de Jong. An algorithm for computing the integral closure. *J. Symbolic Comput.*, 26(3):273–277, 1998.
- [4] Wolfram Decker, Theo de Jong, Gert-Martin Greuel, and Gerhard Pfister. The normalization: a new algorithm, implementation and comparisons. In *Computational methods for representations of groups and algebras (Essen, 1997)*, volume 173 of *Progr. Math.*, pages 177–185. Birkhäuser, Basel, 1999.
- [5] Graham J. Leuschke. Endomorphism rings of finite global dimension. *Canad. J. Math.*, 59(2):332–342, 2007.
- [6] Janko Böhm, Wolfram Decker, and Mathias Schulze. Local analysis of Grauert-Remmert-type normalization algorithms. *Internat. J. Algebra Comput.*, 24(1):69–94, 2014.
- [7] Ernst Kunz. The value-semigroup of a one-dimensional Gorenstein ring. *Proc. Amer. Math. Soc.*, 25:748–751, 1970.
- [8] Félix Delgado de la Mata. Gorenstein curves and symmetry of the semigroup of values. *Manuscripta Math.*, 61(3):285–296, 1988.
- [9] Joachim Jäger. Längenberechnung und kanonische Ideale in eindimensionalen Ringen. *Arch. Math. (Basel)*, 29(5):504–512, 1977.
- [10] Marco D’Anna. The canonical module of a one-dimensional reduced local ring. *Comm. Algebra*, 25(9):2939–2965, 1997.
- [11] V. Barucci, M. D’Anna, and R. Fröberg. Analytically unramified one-dimensional semilocal rings and their value semigroups. *J. Pure Appl. Algebra*, 147(3):215–254, 2000.
- [12] Philipp Korell, Mathias Schulze, and Laura Tozzo. Duality on value semigroups. *J. Commut. Algebra*, 11(1):81–129, 2019.
- [13] Valentina Barucci and Ralf Fröberg. One-dimensional almost Gorenstein rings. *J. Algebra*, 188(2):418–442, 1997.
- [14] Shiro Goto, Naoyuki Matsuoka, and Tran Thi Phuong. Almost Gorenstein rings. *J. Algebra*, 379:355–381, 2013.
- [15] Joseph Lipman. Stable ideals and Arf rings. *Amer. J. Math.*, 93:649–685, 1971.
- [16] Philipp Korell. *Combinatorics of Valuations on Curve Singularities*. Dissertation, Technische Universität Kaiserslautern, 2018.
- [17] Jürgen Herzog and Ernst Kunz, editors. *Der kanonische Modul eines Cohen-Macaulay-Rings*. Lecture Notes in Mathematics, Vol. 238. Springer-Verlag, Berlin-New York, 1971. Seminar über die lokale Kohomologietheorie von Grothendieck, Universität Regensburg, Wintersemester 1970/1971.

- [18] Gert-Martin Greuel, Santiago Laplagne, and Frank Seelisch. Normalization of rings. *J. Symbolic Comput.*, 45(9):887–901, 2010.
- [19] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [20] Malcolm Griffin. Valuations and Prüfer rings. *Canad. J. Math.*, 26:412–429, 1974.
- [21] K. Kiyek and J. L. Vicente. *Resolution of curve and surface singularities*, volume 4 of *Algebras and Applications*. Kluwer Academic Publishers, Dordrecht, 2004. In characteristic zero.
- [22] J. Herzog and E. Kunz. Die Wertehalbgruppe eines lokalen Rings der Dimension 1. *S.-B. Heidelberger Akad. Wiss. Math. Natur. Kl.*, pages 27–67, 1971.
- [23] Hideyuki Matsumura. *Commutative algebra*, volume 56 of *Mathematics Lecture Note Series*. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., second edition, 1980.
- [24] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [25] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [26] G.-M. Greuel, C. Lossen, and E. Shustin. *Introduction to singularities and deformations*. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [27] A. Campillo, F. Delgado, and S. M. Gusein-Zade. On generators of the semigroup of a plane curve singularity. *J. London Math. Soc. (2)*, 60(2):420–430, 1999.

PHILIPP KORELL, OVGU MAGDEBURG, GERMANY
 Email address: philipp.korell@ovgu.de