

BOHR'S PHENOMENON IN COMPLEX BANACH SPACE

VIBHUTI ARORA AND SHANKEY KUMAR

ABSTRACT. In this article, we investigate sharp Bohr's phenomenon to four classes of holomorphic functions mappings the bounded balanced domain G of a complex Banach space X into the right half-plane, the slit domain, the punctured unit disk, and the exterior of the closed unit disk.

1. Introduction

Let X be a complex Banach space. For domain $G \subset X$, let $H(G, D)$ be the set of holomorphic mappings from G into D . Every $f \in H(G, D)$ can be expanded, in a neighbourhood of any given $x_0 \in G$, into the series

$$(1.1) \quad f(x) = \sum_{m=0}^{\infty} \frac{1}{m!} D^m f(x_0) \underbrace{\left((x - x_0), (x - x_0), \dots, (x - x_0) \right)}_{m\text{-times}}$$

where $D^m f(x_0)$, $m \in \mathbb{N}$, denote the m -th Fréchet derivative of f at x_0 , which is a bounded symmetric m -linear mapping from $\prod_{i=1}^m X$ to \mathbb{C} . For convenience, in this article, further we write $D^m f(x_0)((x - x_0)^m)$ instead of

$$D^m f(x_0) \underbrace{\left((x - x_0), (x - x_0), \dots, (x - x_0) \right)}_{m\text{-times}}.$$

For more information on this topic, we refer [18] to the reader.

A domain $G \subset X$ is said to be *balanced*, if $zG \subset G$ for all $z \in \overline{\mathbb{D}}$, where $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. Given a balanced domain G , we denote higher dimensional Bohr radius by $K_X^G(D)$ the largest non-negative number r with the property that if the power series

$$\sum_{m=1}^{\infty} \left| \frac{1}{m!} D^m f(0)(x^m) \right| \leq d(f(0), \partial D)$$

holds for all $x \in rG$ and for all holomorphic functions $f \in H(G, D)$ with an expansion (1.1) about origin. Here $d(f(0), \partial D)$ denotes the Euclidean distance between $f(0)$ and the boundary ∂D of the domain D . In 1914, Bohr gave a remarkable result [14] (due to the work of Riesz, Schur, Wiener) states that $K_{\mathbb{D}}^{\mathbb{C}}(\mathbb{D}) = 1/3$. An application of this result in the Banach algebras given by Dixon in [16]. After this, Bohr's result have been very popular to many mathematician. To see the progress on this topic, we refer survey article [5] and for recent results related to the Bohr phenomenon with complex variables, reader can refer to [6, 11, 20, 22, 25, 26].

In recent years, a lot of attention has been paid to the multidimensional generalizations of Bohr's theorem which was first given by Boas and Khavinson [13] in 1997. The result $K_{\mathbb{C}^n}^{\mathbb{C}}(\mathbb{D}) \geq 1/3$ was proved by Aizenberg [7] for any balanced domain G in \mathbb{C}^n , in the same paper he gave the optimal result $K_{\mathbb{C}^n}^{\mathbb{C}}(\mathbb{D}) = 1/3$ by assuming G is convex. In 2009, Hamada et al. [19, Corollary 3.2] discussed a more general result $K_X^G(\mathbb{D}) = 1/3$ for any bounded

2010 *Mathematics Subject Classification.* 30A10, 30C45, 30C80.

Key words and phrases. Bohr phenomena, Holomorphic mappings, Subordination.

balanced domain G subset of complex Banach space X . Also, by taking a restriction on $f \in H(G, \mathbb{D})$ such that $f(0) = 0$ and $G \subset \mathbb{C}^n$, Liu and Ponnusamy [23] improved the quantity $K_{\mathbb{C}^n}^G(\mathbb{D}) = 1/\sqrt{2}$ and obtain a sharp radius $K_{\mathbb{C}^n}^G(\mathbb{D}) = 1/\sqrt{2}$ if G is convex. Moreover, Bhowmik and Das [12] calculated the following quantities

$$\inf\{K_X^G(\Omega) : \Omega \subset \mathbb{C} \text{ is simply connected}\} = 3 - 2\sqrt{2}$$

and

$$\inf\{K_X^G(\Omega) : \Omega \subset \mathbb{C} \text{ is convex}\} = \frac{1}{3}.$$

In higher dimension, there are many significant and attractive results (see for instance [7, 8]).

Before proceeding further we set some notations. Let $\mathbb{H} := \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ be the right half-plane, $T := \{z \in \mathbb{C} : |\arg z| < \pi\}$ be the slit region, $\mathbb{D}_0 := \{z \in \mathbb{C} : 0 < |z| < 1\}$ be the punctured unit disk, and $\overline{\mathbb{D}}^c := \{z \in \mathbb{C} : |z| > 1\}$ be the exterior of the closed unit disk $\overline{\mathbb{D}}$. The spherical distance between two complex numbers of extended complex plan is given by

$$\lambda(z, w) = \begin{cases} \frac{|z-w|}{\sqrt{1+|z|^2}\sqrt{1+|w|^2}}, & \text{if } z, w \in \mathbb{C}, \\ \frac{1}{\sqrt{1+|z|^2}}, & \text{if } w = \infty. \end{cases}$$

In [2], Abu-Muhanna and Ali observed that if $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D}, \overline{\mathbb{D}}^c)$ then

$$(1.2) \quad \lambda\left(\sum_{n=0}^{\infty} |a_n z^n|, |a_0|\right) \leq \lambda(f(0), \partial\overline{\mathbb{D}}^c)$$

for all $|z| \leq 1/3$, here the bound $1/3$ is sharp. In 2013, Abu-Muhanna and Ali in [3] obtained $K_{\mathbb{C}}^{\mathbb{D}}(\mathbb{H}) = 1/3$ and $K_{\mathbb{C}}^{\mathbb{D}}(T) = (\sqrt{2}-1)/(\sqrt{2}+1)$. Further, Abu-Muhanna et al. [4] determined $K_{\mathbb{C}}^{\mathbb{D}}(\mathbb{D}_0) = 1/3$.

The goal of this paper is establish sharp bound of $K_X^G(\mathbb{H})$, $K_X^G(\mathbb{D}_0)$, and $K_X^G(T)$, where G is a balanced domain in a Banach space X . Moreover, we obtain a radius under which the inequality (1.2) holds for functions of the class $H(G, \overline{\mathbb{D}}^c)$.

2. Preliminaries and Main Results

In this section, we will use the concept of subordination to prove our results. Therefore, it is worth to recall here the definition of subordination. Suppose f and g are analytic functions in \mathbb{D} . We say that g is *subordinate* to f , or $g \prec f$, if there is an analytic function $w : \mathbb{D} \rightarrow \mathbb{D}$ with $w(0) = 0$ such that $g = f \circ w$. To know more about subordination, reader can refer to [17, 24]. The Bohr phenomenon for the class of subordinations studied in [1, 9, 10, 21].

2.1. Half-plane and Convex domains. Let \mathcal{P} denote the class of all analytic functions p with $p(0) = 1$ having positive real part in \mathbb{D} , with the form

$$(2.1) \quad p(z) = 1 + c_1 z + c_2 z^2 + \cdots.$$

A member of \mathcal{P} is called a *Carathéodory function*. In order to establish our first main result, we need the following fundamental result for Carathéodory functions.

Lemma 2.1. *For every $p \in \mathcal{P}$ of the form (2.1), we have $|c_n| \leq 2$ for $n \geq 1$.*

We can now state and prove our first result on multidimensional Bohr's phenomenon for functions in the class $H(G, \mathbb{H})$.

Theorem 2.2. *Let $f \in H(G, \mathbb{H})$ be of the form (1.1) in a neighbourhood of the origin. Then*

$$\sum_{m=1}^{\infty} \left| \frac{1}{m!} D^m f(0)(x^m) \right| \leq d(f(0), \partial\mathbb{H})$$

for $x \in (1/3)G$. Moreover, the constant $1/3$ is best possible.

Proof. Note that $|f(0)| > 0$, for $f \in H(G, \mathbb{H})$. Assume any fixed $y \in G$ and let

$$F(z) := f(zy), \quad z \in \mathbb{D}.$$

Then $F : \mathbb{D} \rightarrow \mathbb{H}$ is holomorphic and

$$(2.2) \quad F(z) = f(0) + \sum_{m=1}^{\infty} \frac{1}{m!} D^m f(0)(y^m) z^m.$$

Consider a function

$$g(z) = \frac{F(z) - i\text{Im}(f(0))}{\text{Re}(f(0))} = 1 + \frac{1}{\text{Re}(f(0))} \sum_{m=1}^{\infty} \frac{1}{m!} D^m f(0)(y^m) z^m.$$

It easily follow that $g \in \mathcal{P}$. We now apply Lemma 2.1 on the Carathéodory function g and obtain

$$\left| \frac{D^m f(0)(y^m)}{m!} \right| \leq 2\text{Re}(f(0))$$

for $m \geq 1$. Thus we have

$$\sum_{m=1}^{\infty} \left| \frac{D^m f(0)(x^m)}{m!} \right| = \sum_{m=1}^{\infty} \left| \frac{D^m f(0)(y^m)}{m!} \right| \left(\frac{1}{3} \right)^m \leq \text{Re}(f(0)) = d(f(0), \partial\mathbb{H})$$

for $x \in (1/3)G$.

Now let us prove that $1/3$ is optimal number using the technique given in [19]. For any $1/3 < r_0 < 1$, there exist a $0 < c < 1$ and $v \in \partial G$ such that $cr_0 > 1/3$ and $c \sup_{x \in \partial G} \|x\| < \|v\|$. Next, we consider a function f_0 on G by

$$f_0(x) = L\left(\frac{c\psi_v(x)}{\|v\|}\right)$$

where $L(z) = (1+z)/(1-z)$, $z \in \mathbb{D}$, ψ_v is a bounded linear functional on X with $\psi_v(v) = \|v\|$ and $\|\psi_v\| = 1$. It is easy to check that $c\psi_v(x)/\|v\| \in \mathbb{D}$ and $f_0 \in H(G, \mathbb{H})$. Choose $x = r_0v$, we get

$$f_0(r_0v) = \frac{1 + cr_0}{1 - cr_0} = 1 + 2 \sum_{n=1}^{\infty} cr_0^n = f_0(0) + \sum_{m=1}^{\infty} \left| \frac{D^m f_0(0)(x^m)}{m!} \right| > 2 = 1 + d(f_0(0), \partial\mathbb{H}).$$

The proof of theorem is complete. □

It is now appropriate to remark that when $X = \mathbb{C}$ and $G = \mathbb{D}$, Theorem 2.2 coincides with [3, Theorem 2.1]. Furthermore, the next result, for holomorphic functions defined on a bounded balanced domain which takes values in a convex domain, can be derived from Theorem 2.2:

Corollary 2.3. *Let $f \in H(G, C)$, where C is a convex domain, be of the form (1.1) in a neighbourhood of the origin. Then*

$$\sum_{m=1}^{\infty} \left| \frac{D^m f(0)(x^m)}{m!} \right| \leq d(f(0), \partial C)$$

for $x \in (1/3)G$. This result is sharp.

Proof. Let $u \in \partial C$ be nearest to $f(0)$. Further let T_u be the tangent line at u , and H_u the half-plane containing C . Then $f \in H(G, H_u)$. Now choose t real so that $(H_u - u)e^{it} = \mathbb{H}$, the right half-plane and let $K = (C - u)e^{it} \subset \mathbb{H}$. Hence it yields that $g(z) = (f(z) - u)e^{it} \in H(G, K) \subset H(G, \mathbb{H})$ and $g(0) = (f(0) - u)e^{it} = |f(0) - u| = d(f(0), \partial C)$. Then we can apply Theorem 2.2 on g and it follows the desired result as well as the sharpness. \square

2.2. Punctured unit disk. Our next objective is to give multidimensional analogue of [4, Theorem 2.1] for the class $H(G, \mathbb{D}_0)$.

Theorem 2.4. *Let $f \in H(G, \mathbb{D}_0)$ be of the form (1.1) in a neighbourhood of the origin. Then*

$$(2.3) \quad \sum_{m=1}^{\infty} \left| \frac{1}{m!} D^m f(0)(x^m) \right| \leq d(f(0), \partial \mathbb{D}_0) = 1$$

for $x \in (1/3)G$. This result is sharp.

Proof. Because of the inclusion $H(G, \mathbb{D}_0) \subset H(G, \mathbb{D})$, the required inequality (2.3) for $x \in (1/3)G$ directly follows from [19, Corollary 3.2]. To prove sharpness part, we apply the same technique as in the proof of Theorem 2.2. We consider for any $r_0 \in (1/3, 1)$ there exist $0 < c < 1$ and $v \in \partial G$ such that $cr_0 > 1/3$ and $c \sup_{x \in \partial G} \|x\| < \|v\|$. Next, we define

$$H_t(z) = \exp\left(-t \frac{1+z}{1-z}\right) = \frac{1}{e^t} + \frac{1}{e^t} \sum_{n=1}^{\infty} \left[\sum_{m=1}^n \frac{(-2t)^m}{m!} \binom{n-1}{m-1} \right] z^n, \quad t > 0$$

which is in $H(\mathbb{D}, \mathbb{D}_0)$. Now consider a function $f_1 \in H(G, \mathbb{D}_0)$ defined by

$$f_1(x) = H_t\left(\frac{c\psi_v(x)}{\|v\|}\right),$$

where ψ_v is a bounded linear functional on X with $\psi_v(v) = \|v\|$ and $\|\psi_v\| = 1$. Thus, for $x = r_0 v$, we get

$$f_1(r_0 v) = H_t(cr_0) = \sum_{m=1}^{\infty} \left| \frac{D^m f_1(0)(x^m)}{m!} \right| > 1$$

by using the argument given in [4, Theorem 2.1]. This concludes the proof. \square

2.3. Slit mappings. The following well-known result of de Branges [15] on subordination is useful to prove our next result.

Lemma 2.5. *If $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$ are two analytic functions in \mathbb{D} such that $f(0) = g(0)$, g is univalent (one-to-one) and $f \prec g$, then $|a_n| \leq n|b_1|$.*

We now state and prove our next result which provides the sharp Bohr inequality for functions belonging to the class $H(G, T)$.

Theorem 2.6. *Let $f \in H(G, T)$ of the form (1.1) in a neighbourhood of the origin with $f(0) > 0$. Then*

$$(2.4) \quad \sum_{m=1}^{\infty} \left| \frac{D^m f(0)(x^m)}{m!} \right| \leq d(f(0), \partial T) = f(0)$$

for $x \in (3 - 2\sqrt{2})G$. This result is sharp.

Proof. Suppose that $f \in H(G, T)$. Now, we construct F as in the proof of Theorem 2.2 which is in $H(\mathbb{D}, T)$ having series expansion (2.2). Since $F \in H(\mathbb{D}, T)$, we may write the given condition as

$$F(z) \prec g(z)$$

where

$$g(z) = f(0) \left(\frac{1+z}{1-z} \right)^2 = f(0) + 4f(0) \sum_{n=1}^{\infty} n z^n.$$

Here g is univalent mapping in \mathbb{D} such that $F(\mathbb{D}) \subset g(\mathbb{D})$ and $F(0) = g(0)$. According to Lemma 2.5, we have

$$\left| \frac{D^m f(0)(y^m)}{m!} \right| \leq 4f(0)m, \quad m \geq 1.$$

This gives, for $z \in \mathbb{D}$ and $y \in G$, that

$$\sum_{m=1}^{\infty} \left| \frac{D^m f(0)(y^m)}{m!} \right| |z|^m \leq 4f(0) \sum_{m=1}^{\infty} m |z|^m = 4f(0) \frac{|z|}{(1-|z|)^2}$$

which is less than or equal to $f(0)$ for all $|z| \leq 3 - 2\sqrt{2} \approx 0.17157$. Hence the inequality (2.4) is hold in $(3 - 2\sqrt{2})G$.

To prove that the result is sharp, we adopt the approach of Theorem 2.2 and consider the following situation: for any $r_0 \in (3 - 2\sqrt{2}, 1)$, there exist a $c \in (0, 1)$ such that $cr_0 > 3 - 2\sqrt{2}$. Also there exist a $v \in \partial G$ such that

$$c \sup\{\|x\| : x \in \partial G\} < \|v\|.$$

Next, we consider a function f_2 on G defined by

$$f_2(x) = U \left(\frac{c\psi_v(x)}{\|v\|} \right),$$

where

$$U(z) = \left(\frac{1+z}{1-z} \right)^2, \quad z \in \mathbb{D}$$

and, ψ_v is a bounded linear functional on X with $\psi_v(v) = \|v\|$ and $\|\psi_v\| = 1$. It is easy to check that $c\psi_v(x)/\|v\| \in \mathbb{D}$ and $f_2 \in H(G, T)$. Then, for $x = r_0 v$, we have

$$f_2(x) = f_2(0) + \sum_{m=1}^{\infty} \left| \frac{D^m f_2(0)(y^m)}{m!} \right| |z|^m = \left(\frac{1+cr_0}{1-cr_0} \right)^2 = 1 + 4 \sum_{n=1}^{\infty} n (cr_0)^n > 2.$$

This shows the sharpness. □

2.4. Exterior of the closed unit disk. The following lemma by Abu-Muhanna and Ali [2] will be required to establish our next result.

Lemma 2.7. *Let $f \in H(\mathbb{D}, \overline{\mathbb{D}}^c)$. Then $f \prec W$, where*

$$W(z) = \exp\left(\frac{1 + \phi(z)}{1 - \phi(z)}\right),$$

with

$$\phi(z) = \frac{z + b}{1 + \bar{z}b} \text{ and } b = \frac{\log f(0) - 1}{\log f(0) + 1}.$$

The following theorem extends a result, given by Abu-Muhanna and Ali [2, Theorem 2.1], in higher dimension.

Theorem 2.8. *Let $f \in H(G, \overline{\mathbb{D}}^c)$ of the form (1.1) in a neighbourhood of the origin. Then*

$$(2.5) \quad \lambda\left(\sum_{m=1}^{\infty} \left|\frac{D^m f(0)(x^m)}{m!}\right|, |f(0)|\right) \leq \lambda(f(0), \partial\overline{\mathbb{D}}^c)$$

for $x \in (1/3)G$. This result is sharp.

Proof. Let $f \in H(G, \overline{\mathbb{D}}^c)$. Then we can construct a function F as in the proof of Theorem 2.2 which will be in $H(\mathbb{D}, \overline{\mathbb{D}}^c)$ and have the series expansion (2.2). Now, by using Lemma 2.7, we have $F \prec W$. It is easy to observe that, the $W(z)$ defined in Lemma 2.7 can be written as

$$W(z) = F(0) \exp\left(\frac{\log |F(0)|^2 z}{1 - z}\right).$$

Hence, from [2, Equation 2.7], we obtain

$$\sum_{m=0}^{\infty} \left|\frac{1}{m!} D^m f(0)(x^m)\right| |z|^m \leq |F(0)|^{\frac{1+|z|}{1-|z|}} = |f(0)|^{\frac{1+|z|}{1-|z|}}.$$

It leads to the fact that, for $|z| \leq 1/3$,

$$\sum_{m=0}^{\infty} \left|\frac{1}{m!} D^m f(0)(x^m)\right| |z|^m \leq |f(0)|^2.$$

Then a simple computation shows that

$$\lambda\left(\sum_{m=0}^{\infty} \left|\frac{1}{m!} D^m f(0)(x^m)\right|, |f(0)|\right) \leq \lambda(|f(0)|, |f(0)|^2) \leq \lambda(|f(0)|, 1)$$

holds for $x \in (1/3)G$.

Finally, we prove that inequality (2.5) is not holds true for $x \in r_0 G$, where $r_0 \in (1/3, 1)$. As we know that there exist a $c \in (0, 1)$ and $v \in \partial G$ such that $cr_0 > 1/3$ and

$$c \sup\{\|x\| : x \in \partial G\} < \|v\|.$$

Now, we consider a function f_3 on G defined by

$$f_3(x) = W\left(\frac{c\psi_v(x)}{\|v\|}\right),$$

where ψ_v is a bounded linear functional on X with $\psi_v(v) = \|v\|$ and $\|\psi_v\| = 1$. Thus

$$f_3(r_0 v) = W(cr_0) = |f(0)| \exp\left(\frac{cr_0}{1 - cr_0} \log |f(0)|^2\right) = |f(0)|^{(1+cr_0)/(1-cr_0)}.$$

Also, a simple computation gives that

$$\frac{\lambda(|f(0)|, |f(0)|^{(1+cr_0)/(1-cr_0)})}{\lambda(|f(0)|, 1)} = \frac{\sqrt{2}|f(0)|(|f(0)|^{(2cr_0)/(1-cr_0)} - 1)}{(|f(0)| - 1)\sqrt{1 + |f(0)|^{2(1+cr_0)/(1-cr_0)}}} \rightarrow \frac{2cr_0}{1 - cr_0}$$

as $|f(0)| \rightarrow 1$. Since $2x/(1-x) > 1$, for $x > 1/3$, it leads to the fact that

$$\lambda\left(\sum_{m=0}^{\infty} \left| \frac{1}{m!} D^m f_0(0)(x^m) \right| |z|^m, |f(0)|\right) = \lambda(|f(0)|, |f(0)|^2) > \lambda(|f(0)|, 1)$$

as $|f(0)| \rightarrow 1$. This concludes the theorem. \square

Acknowledgement. The authors would like to thank Professor Swadesh Kumar Sahoo for fruitful discussion on this topic. The first author thanks IIT Bhubaneswar for providing Institute Post Doctoral Fellowship. The work of the second author is supported by the Institute Post Doctoral Fellowship of NISER Bhubaneswar, India.

Conflict of Interests. The authors declare that there is no conflict of interests regarding the publication of this paper.

REFERENCES

- [1] Y. Abu-Muhanna, Bohr's phenomenon in subordination and bounded harmonic classes, *Complex Var. Elliptic Equ.* **55**(11), 1071–1078 (2010).
- [2] Y. Abu-Muhanna and R. M. Ali, Bohr's phenomenon for analytic functions into the exterior of a compact convex body, *J. Math. Anal. Appl.* **379** (2011), 512–517.
- [3] Y. Abu-Muhanna and R. M. Ali, Bohr's phenomenon for analytic functions and the hyperbolic metric, *Math. Nachr.* **286**(11-12) (2013), 1059–1065.
- [4] Y. Abu-Muhanna, R. M. Ali, and Z. C. Ng, Bohr radius for the punctured disk, *Math. Nachr.* **290** (2017), 2434–2443.
- [5] Y. Abu-Muhanna, R. M. Ali, and S. Ponnusamy, On the Bohr inequality. In "Progress in Approximation Theory and Applicable Complex Analysis" (Edited by N. K. Govil et al.), Springer Optimization and Its Applications, **117** (2016), 265–295.
- [6] Y. Abu-Muhanna, R. M. Ali, and S. K. Lee, The Bohr operator on analytic functions and sections, *J. Math. Anal. Appl.* **496**(2), (2021), 124837.
- [7] L. Aizenberg, Multidimensional analogues of Bohr's theorem on power series, *Proc. Amer. Math. Soc.* **128** (2000), no. 4, 1147–1155.
- [8] L. Aizenberg, Generalization of results about the Bohr radius for power series, *Stud. Math.* **180** (2007), 161–168.
- [9] V. Allu and V. Arora, Bohr-Rogosinski type inequalities for concave univalent functions, Preprint. [arXiv:2204.14085](https://arxiv.org/abs/2204.14085)
- [10] B. Bhowmik and N. Das, Bohr phenomenon for subordinating families of certain univalent functions, *J. Math. Anal. Appl.* **462**(2) (2018) 1087–1098.
- [11] B. Bhowmik and N. Das, Bohr phenomenon for locally univalent functions and logarithmic power series, *Comput. Methods Funct. Theory* **19**(4), 729–745 (2019).
- [12] B. Bhowmik and N. Das, Bohr radius and its asymptotic value for holomorphic functions in higher dimensions, *C. R. Math. Acad. Sci. Paris*, **359** (2021), 911–918.
- [13] H. P. Boas and D. Khavinson, Bohr's power series theorem in several variables, *Proc. Amer. Math. Soc.* **125** (1997), 2975–2979.
- [14] H. Bohr, A theorem concerning power series, *Proc. London Math. Soc.* **13**(2) (1914), 1–5.
- [15] L. de Branges, A proof of the Bieberbach conjecture, *Acta Math.* **154** (1985), 137–152.
- [16] P. G. Dixon, Banach algebras satisfying the non-unital von Neumann inequality, *Bull. Lond. Math. Soc.*, **27** (1995), pp. 359–362
- [17] P. L. Duren, *Univalent Functions* Springer-Verlag, New York, (1983).
- [18] I. Graham and G. Kohr, *Geometric function theory in one and higher dimensions*, Pure and Applied Mathematics, Marcel Dekker, 255, Marcel Dekker, 2003.

- [19] H. Hamada, T. Honda, and G. Kohr, Bohr's theorem for holomorphic mappings with the values in homogeneous balls, *Isr. J. Math.* **173** (2009), 177–187.
- [20] I. R. Kayumov and S. Ponnusamy, On a powered Bohr inequality, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **44** (2019), 301–310.
- [21] S. Kumar and S. K. Sahoo, A generalization of the Bohr-Rogosinski sum, Preprint. [arXiv:2106.06502](https://arxiv.org/abs/2106.11158)
<https://arxiv.org/abs/2106.11158>
- [22] G. Liu, Z. Liu, and S. Ponnusamy, Refined Bohr inequality for bounded analytic functions, *Bull. Sci. Math.* **173** (2021), Paper No. 103054, 20 pp.
- [23] M. S. Liu and S. Ponnusamy, Multidimensional analogues of refined Bohr's inequality, *Proc. Amer. Math. Soc.* **149** (2021), 2133–2146.
- [24] S. S. Miller and P. T. Mocanu, *Differential Subordinations-Theory and Applications*, Marcel Dekker, Inc., New York, 2000.
- [25] M. S. Liu, S. Ponnusamy, and J. Wang, Bohr's phenomenon for the classes of Quasi-subordination and K -quasiregular harmonic mappings, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.* **114** Article 115 (2020).
- [26] Z. H. Liu and S. Ponnusamy, Bohr radius for subordination and K -quasiconformal harmonic mappings, *Bull. Malays. Math. Sci. Soc.* **42** (2019), 2151–2168.

VIBHUTI ARORA, SCHOOL OF BASIC SCIENCE, INDIAN INSTITUTE OF TECHNOLOGY BHUBANESWAR, ODISHA, INDIA.

Email address: vibhuti.arora1991@gmail.com

SHANKEY KUMAR, SCHOOL OF MATHEMATICAL SCIENCES, NATIONAL INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, BHUBANESWAR, AN OCC OF HOMI BHABHA NATIONAL INSTITUTE, JATNI 752050, INDIA

Email address: shankeygarg93@gmail.com