

The Cavendish *Computors*: The women working in scientific computing for Radio Astronomy

Verity Allan

Abstract A discussion of the history of scientific computing for Radio Astronomy in the Cavendish Laboratory of the University of Cambridge in the decades after the Second World War. This covers the development of the aperture synthesis technique for Radio Astronomy and how that required using the new computing technology developed by the University's Mathematical Laboratory: the EDSAC, EDSAC 2 and TITAN computers. It looks at the scientific advances made by the Radio Astronomy group, particularly the assembling of evidence which contradicted the Steady State hypothesis. It also examines the software advances that allowed bigger telescopes to be built: the Fast Fourier Transform (FFT) and the degriding algorithm. Throughout, the contribution of women is uncovered, from the diagrams they drew for scientific publications, through programming and operating computers, to writing scientific papers.

1 Introduction

The years following the Second World War were a time of massive scientific and technological innovation in the University of Cambridge. The University's Mathematical Laboratory was set up prior to the War to investigate numerical methods, and after the war, Maurice Wiles became its director. Having visited the United States and seen the ENIAC computer, he started building EDSAC, putting Cambridge at the forefront of the computer revolution [2]. In the same post-War years, the Cavendish Radio Astronomy group assembled, with Martin Ryle at its head. The Cavendish Laboratory of the University of Cambridge had been doing fundamental physics research for decades, and staff had won Nobel Prizes.[6] The Radio Astronomy group would add to this collection, as Ryle and Hewish won the Nobel

Verity Allan
Battcock Centre for Experimental Astrophysics, Cavendish Laboratory, University of Cambridge
e-mail: v1a22@mrao.cam.ac.uk

Prize for Physics in 1974 for their innovative telescope design and the discovery of pulsars — work that relied on the computers of the Mathematical Laboratory [17].

This reliance on computers is exemplified by Ryle's request for a bigger computer:

One day, Ryle came to me to say that he was planning the erection of a much larger telescope and to ask whether the Mathematical Laboratory could undertake to provide the computing support required. [34], p. 193

I will explain why Radio Astronomy needed a bigger computer, and will elucidate the role of other members of the Radio Astronomy group and the Mathematical Laboratory beyond the relatively well-known names of Ryle, Hewish, and Wilkes.

It has become more widely known in recent years that a lot of early computing relied on women — thanks to work by Janet Abbate and Mar Hicks and Margo Lee Shetterley [1] [13] [27]. Women were clearly involved in the early years of computing at Cambridge, as can be seen from the archives of the Department of Computer Science and Technology, which is a successor to the Mathematical Laboratory. A photograph of staff from 1949 shows 20 staff, of whom 8 were women [9]. In this paper, we will uncover some of the contributions of women to science and scientific programming in Radio Astronomy in the years from 1950 to 1963.

The University computers that the Radio Astronomy group used during this period were EDSAC, the first electronic digital stored program computer in the world, EDSAC 2, which led to many innovations in microprogramming, and TITAN, which led to advances in computer graphics and computer aided design (CAD) [2]. TITAN was the computer requested by Ryle, which was needed for the One-Mile telescope.

2 The Women In Radio Astronomy Computing

The Radio Astronomy group became reliant on computing, and by 1965, over 40% of all papers produced by the group had required the use of a computer to process data [14]. And women were extensively involved in this endeavour. Around 50% of people acknowledged in papers for their contribution to the programming and operation of computers were women [14]. These women were employed in a variety of roles.

Some were hired as Computers in the 1950s and 1960s,¹ initially to do calculations, and then later to assist with programming. *Computer* or *Computer* as a job description was common in the 1940s, first to describe someone who was hired to do calculations by hand or with a mechanical or electro-mechanical computer, and then to describe a worker who used a computer in some capacity [13]. The Cavendish records also show that people (often women) were hired as *scanners* for High Energy Physics: their job was to analyse photographic plates from cloud chamber experiments. Women were also hired as Research Assistants: Dr Elizabeth Waldram (whom we'll meet again later) was hired as such [2], p. 63. Judy Bailey,

¹ Source: Cavendish Laboratory Archive Personnel Records.

who later became Deputy Director of the University Computing Service (another successor to the Mathematical Laboratory) worked for Radio Astronomy as a Technical Officer [31].

Women were also hired to prepare diagrams for scientific papers, drawing graphs and contour diagrams, as it was not yet possible to prepare these with computers. At the time of writing, it's not possible to look at University records to get a sense of the number of people in these roles, nor of how many of them were women. They were known as *the girls in the attic* [8]

Some of the women involved in scientific computing were PhD students and researchers themselves. As we'll see, Ann Neville, one of Ryle's PhD students, did a considerable amount of programming. She had to program for EDSAC 2, and to save space, many constants had to be combined into one constant, just to shave a few bytes off the size of the program.²

3 The first forays into scientific computing

While Charles Wynn-Williams was a pioneer of electro-mechanical computing for the Cavendish [35], the Radio Astronomy group didn't leap straight to digital computing for all their papers. We can trace the use of computers and the contributions of programmers primarily through the acknowledgements section of papers. They published their first paper using EDSAC in 1953, and use through the 1950s was fairly sporadic [14].

The first women we encounter in the acknowledgements probably were the computers. In 1951, a Mrs A. C. Hollis-Hallett is credited as having *analysed a large proportion of the records* [28]. It's not possible to verify this, as no records survive. Similarly, a Mrs Moore is credited in a paper by Scheuer and Ryle from 1953, *for help in the computation of results* [24]. It's likely that Mrs Moore was the computer, not EDSAC, as EDSAC was not listed in the acknowledgements (which was the common practice at the time).

However, the development of a new technique in Radio Astronomy was going to change the group's relationship with computers: the technique of aperture synthesis.

The basic principles were sketched out by Ryle in a 1957 paper [20], and implemented by John Blythe. Aperture synthesis, as the name suggests, allows for the synthesis of a larger aperture. By using two or more smaller telescopes, you can achieve the resolution (but not the sensitivity) of a telescope with diameter equivalent to the longest spacing between antennas.³ However, when you measure a signal with a particular pair of antennas, you are only observing part of the signal – you're observing a Fourier component of the signal [7], Chapter 8. To reconstruct the complete signal, you need to measure the other Fourier components, which can

² Donald Wilson, Personal Communication, 22 April 2021

³ The distance between a given pair of antennas is known as a *baseline*: the length of the longest baseline dictates the resolution achievable by the telescope.

be achieved by moving the antennas around to fill in the aperture. The Fourier components then need to be combined and transformed into a useful measurement of sky brightness.

Thus the Fourier Transform must be calculated for aperture synthesis telescopes, and this is not really feasible to do by hand. So the Radio Astronomy group was forced to turn to computers. Blythe's papers from 1957, where he implemented this technique, used EDSAC for the calculations [4] [5]. One of these papers used EDSAC to perform 15 hours of computation, performing over 2 million operations, to output data in a form suitable for making contour maps [4]. In both papers, Blythe credits a Mrs J. Sandos *who performed most of the analysis*. No-one in the Radio Astronomy group can recall Mrs Sandos (who may have been a member of the Mathematical Laboratory), but she undoubtedly helped with the complex data preparation process: error-free punched tapes were required as input, and all the data had to be arranged in a particular way in order to compensate for EDSAC's lack of storage space.

By 1960, we see EDSAC 2 appearing in paper acknowledgements. However, the workflow still retains a number of manual steps, as one paper notes:

It would be possible to record such information directly in digital form, but it has been found relatively easy to transfer the present records to punched tape in a record reader and manual perforator. [10]

The acknowledgements are silent about who did this, but it's quite likely to be either PhD students or *Computors*.

4 Earth Rotation Aperture Synthesis and the Steady State Hypothesis

We now move forward to 1961, and an extension of the aperture synthesis technique: you can use the rotation of the Earth to help fill in the aperture of your telescope, thus you are performing Earth Rotation Aperture Synthesis. This technique was prototyped using the Cambridge 178MHz telescope and EDSAC 2 — the new, more powerful, computer in the Mathematical Laboratory. This telescope and others were used to collect data that seriously undermined the Steady State hypothesis

In 1961, Scott, Ryle and Hewish performed earth-rotation aperture synthesis for the first time [26]. This paper has a number of interesting features. Firstly, we have this interesting paragraph:

The processes of filtering and convolution are basically equivalent and in this case where the output of the receiver was already recorded in digital form it was convenient to carry out a direct convolution with a time function whose Fourier transform provides the required polar diagram.

This is an early use of a digital filter rather than a piece of electronics; the work is moving from hardware to software. Note also that the *digital form* is paper tape!

To record the data, a *sample and hold* electronic circuit is implemented electro-mechanically, using photo-electric cells, to output two columns of 5-hole tape for each data point (giving ten bits of precision).⁴ Each tape starts at the same sidereal time: at the same time with respect to the fixed stars in the sky, which is most useful to astronomers, so non-local astronomical bodies can be located easily. The sidereal time arrangement meant that the data can be efficiently processed — while EDSAC 2 had more storage space than EDSAC, programmers still could not afford to be profligate. During the data input phase, the data were convolved with a function which allowed for error correction, and which also allowed the extraction of the sine and cosine components which were required for the main computation. At this point, the data were sorted into a more convenient order before being written onto magnetic tape. This magnetic tape was used as the input data for the main calculation, in which the data were put through the Fourier Transform. The final outputs were effectively a 2-dimensional map of the sky, sampled at discrete intervals. The probability distribution of the measured intensity was also calculated and output. The final values for the sky map needed to be interpolated, but this was done manually, possibly by the *girls in the attic* or junior members of the group.

As well as being the first practical demonstration of aperture synthesis using the earth to fill in the aperture, this paper also was important for providing evidence against the Steady State hypothesis. The authors note:

there is no evidence for a reduction in the density of radio sources... this observation implies an isotropy which extends to at least 10,000 sources steradian. [26]

This isotropy in extra-galactic faint sources conflicted with predictions from the Steady State model, which predicted that there should be many fewer faint sources, and that there should be no difference between galactic and extra-galactic sources. And once again we find a woman tucked away in the acknowledgments: Mrs R. Feinstein. Ruth Feinstein was employed by the Mathematical Laboratory [32], and to have been an assistant to Dr Lucy Slater, who became Head of Computing in the University's Department of Applied Economics [3]. She assisted her husband, Charles Feinstein, with

calculating and typing

for his PhD thesis[18]. Unfortunately, she left the Mathematical Laboratory shortly after marriage [32], and she seems to play no further part in computing for Radio Astronomy. She was, along with Dr David Wheeler,

responsible both for the running and design of the programme.⁵

⁴ Thanks to B.J. Harris for spotting that they were building a sample and hold circuit.

⁵ Note that at this time, the UK still used *programme* to refer to computer programs.

4.1 *The Steady State Hypothesis*

A summary of the steady state hypothesis is that the universe is expanding, but has no beginning and no end, and that its appearance does not change over time. Whereas the Big Bang theory states that the universe expanded from an initial state of high density and temperature, and thus its appearance changes over time.

Ryle's group was building up to attack the Steady State hypothesis. There are two papers from Harriet Tumner and Patricia Leslie, both PhD researchers in the group [15] [29]. Neither paper mentions using EDSAC 2, though both are using data from the Cambridge surveys that were processed using that computer. These papers look at the brightness, density, size, and clustering of extra-galactic radio sources — where the Steady State and Big Bang hypotheses generated different predictions.

Then there's a 1961 paper from Paul Scott and Martin Ryle [25]. This looks at the relationship between the number of radio sources, and their flux density (essentially their brightness). The Steady State hypothesis assumes that there's a particular relationship between the brightness of the source, and how many sources of that brightness there are.

The acknowledgements in this paper read:

We should like to thank Dr Hewish and Miss Patricia Leslie for useful discoveries, and Miss Ann Neville and Mr DR Marks for assistance in the reduction of observations.

Ann Neville was one of Ryle's PhD students, and she'll appear as an author on later papers.

Next there's a paper by Hewish. He's modelling the source number–flux density relationship, and uses EDSAC 2 to do it. This is one of the earliest uses of the Monte Carlo method in Radio Astronomy: successors to this method are being used today as a key modelling technique [11]. Hewish doesn't credit anyone else, so he most likely did a lot of the programming himself. This contrasts with Ryle, who didn't program at all, even though Ryle was well aware of how essential computers were for aperture synthesis.⁶

We turn to a Ryle and Clarke paper of 1961 in which they show that extra-galactic radio sources have no variation in density as you look at different parts of the sky — the only variation comes when you look at the galaxy (which is nearby in astronomical terms). [22] The observed source number-flux density relationship doesn't match the steady state hypothesis. They conclude:

These observations do, however, appear to provide conclusive evidence against the steady state model.

And any last hopes that the Steady State theorists had of salvaging their model was blown away conclusively in 1964 when Penzias and Wilson detected the Cosmic Microwave Background [19].

⁶ Dr Elizabeth Waldram, personal communication 4 May 2021

5 A radio survey of the North polar region

We now turn to analyse in some detail a paper from 1962, in which major advances were made in both computing and Radio Astronomy [23].

There are two major computational innovations in this paper. The first is Wheeler's Fast Fourier Transform (FFT) algorithm, tersely described in a single paragraph, plus an equation. From this, we can glean that the data from the sky needs to be placed onto a Cartesian grid (a grid like a chess board, or a Go board) with (x, y) co-ordinates. This means converting from spherical co-ordinates (similar to longitude and latitude), which are useful for working out where something is on the sky, into these Cartesian co-ordinates. While the mathematician Gauss developed the theory of FFT, this practical implementation pre-dates the one by Cooley and Tukey in 1965, which is usually credited as the first practical implementation [12].

The other programming innovation was the map degriding algorithm, which enables conversion back from Cartesian co-ordinates into spherical co-ordinates, so the radio sources are correctly located on the sky, which also ended up producing some of the world's first computer graphics. The output from the computer was plotted on a CRT monitor, where it lasted for a few seconds, just long enough to be photographed [2]. This work was done by Dr Elizabeth Waldram, then a Research Assistant in the group. This degriding algorithm (and its successors) were used for all subsequent interferometers built by the group.⁷

With the FFT and the degriding algorithm, they could build a map of the northern radio sky. It's very likely that the final construction of the map from the photographic plates was made by the *girls in the attic*. This kind of mapping was a large part of their job. They weren't working directly with computers themselves, but they were a vital part of the scientific workflow, often alongside secretarial staff who would type up manuscripts. We can see a rare acknowledgment for the women doing such work in a 1962 paper [30]:

We are grateful to Miss C. Dunn and Mrs M. Warren for helping with the analysis and the preparation of the contour maps.

The contour maps were drawn from EDSAC 2 output [30].

With this paper, we have the foundations for modern radio astronomy computing. The FFT saves a lot of computational effort compared to the Direct Fourier Transform which works with spherical co-ordinates. However, the use of the FFT also necessitates the use of gridding and degriding algorithms, which we need to first transform our data into the right shape for the FFT, and then put it back into spherical co-ordinates afterwards. FFT and gridding algorithms remain an important part of Radio Astronomy research [36] [33] [16].

It's clear from Wheeler's inclusion in the acknowledgments rather than as a co-author that the exclusion of women programmers from authorship wasn't purely a result of sexism. Sexism may still have played a part, as a majority of those acknowledged for programming in Radio Astronomy papers were women [14]. Mar

⁷ Dr Elizabeth Waldram, personal communication 04 May 2021.

Hicks has shown how work done by women, especially in the computing field, was devalued, which may also have been a contributing factor [13]. Wheeler was acknowledged in Ryle's Nobel Prize lecture for his Fast Fourier Transform algorithm [21]; none of the women get mentioned.

6 Discussion

It's clear from the preceding sections that women were deeply involved in the production of scientific papers in the 1950s and 60s, and that they were also involved in doing much of the programming and a good deal more of the science than is usually recognised. The programming also required good mathematical skills and an understanding of Radio Astronomy, so that programmers could work with sidereal time and Right Ascension and Declination (a spherical co-ordinate system used by astronomers) and their conversion into Cartesian co-ordinates for use in the FFT — something that mathematics undergraduates often struggle with. Perhaps the *girls in the attic* didn't require such advanced understanding, but they were expected to work to the highest standards of accuracy.

It is extraordinarily difficult to get a picture of many of the computational methods used. Programs were not routinely recorded at the time, though there are some contemporaneous program booklets stored in the archive of the Department of Computer Science and Technology.⁸ However, none of these survive for Radio Astronomy. The information presented in papers is minimal - usually just enough to allow for some mathematical recreation, but not enough to understand the algorithms. This may have been the result of the culture of the research group at the time; Ryle may have maintained more of a war-time culture of secrecy than we find desirable as scientists today.⁹

It is also worth noting that, while women may have left the Mathematical Laboratory on marriage, this wasn't true for members of the Radio Astronomy group. Indeed, programmers were in such high demand that they could work part-time and flexibly, many years ahead of this becoming widely available. Dr Waldram recalls working 10-15 hours a week in the early 1960s, to accommodate her childcare needs.¹⁰

7 Conclusion

The acknowledgements to papers provide one of the few ways of finding out about the work done in early scientific computing, as they often tell us that work was done

⁸ See for example, shelfmark V75-58, which lists programs used by the Department on Geology on TITAN.

⁹ Donald Wilson, personal communication 22 April 2021.

¹⁰ Elizabeth Waldram, personal communication 4 May 2021.

using a computer, and who did that work. We can get tantalising glimpses of what that work was, and the people behind it. This could usefully be supplemented by more archival work within the University, once this is possible again.

Acknowledgements My thanks go to Dr Elizabeth Waldram, one of the Cavendish’s computing pioneers, who has given her time very generously, to Dr Donald Wilson, who provided many valuable insights into this period, to Dr David Green, who helped me with my \LaTeX , and to Professor Malcolm Longair, who has supported this project.

References

1. Abbate, J.: *Recoding gender : women’s changing participation in computing* / Janet Abbate. History of computing. MIT Press, Cambridge, Mass. (2012)
2. Ahmed, H.: *Cambridge Computing: The First 75 Years*. Third Millennium Publishing, London, UK (2013)
3. Andrews, G.E.: The discovery of ramanujan’s lost notebook: In honour of the 125th birthday of ramanujan (2013, cited 2021 Apr 7). URL www.personal.psu.edu/lxv1/alg/antheory/preprint/andrews/295.pdf
4. Blythe, J.H.: A new type of pencil beam aerial for radio astronomy. *MNRAS***117**, 644 (1957). DOI 10.1093/mnras/117.6.644
5. Blythe, J.H.: Results of a survey of galactic radiation at 38 Mc/s. *MNRAS***117**, 652 (1957). DOI 10.1093/mnras/117.6.652
6. Nobel Prizes (2021, Cited 30 Sep 2021). URL <https://www.phy.cam.ac.uk/history/nobel>
7. Christiansen, W.N., Hogbom, J.A.: *Radiotelescopes*. Cambridge University Press (1985)
8. Clifford, F. E.: The formation of the iron peak elements (1964)
9. Computer Laboratory, University of Cambridge: Archive photos (1999). URL https://www.cl.cam.ac.uk/relics/archive_photos.html
10. Costain, C.H., Smith, F.G.: The radio telescope for 7.9 meteres wavelength at the Mullard Observatory. *MNRAS***121**, 405 (1960). DOI 10.1093/mnras/121.4.405
11. Feroz, F., Hobson, M.P., Cameron, E., Pettitt, A.N.: Importance nested sampling and the multi-nest algorithm. *The Open Journal of Astrophysics* **2**(1) (2019). DOI 10.21105/astro.1306.2144. URL <http://dx.doi.org/10.21105/astro.1306.2144>
12. Heideman, M.T., Johnson, D.H., Burrus, C.S.: Gauss and the History of the Fast Fourier Transform. *IEEE ASSP Magazine* pp. 15–21 (1984). URL https://www.cis.rit.edu/class/simg716/Gauss_History_FFT.pdf
13. Hicks, M.: *Programmed inequality : how Britain discarded women technologists and lost its edge in computing* / Marie Hicks. History of computing. The MIT Press (2018)
14. Leedham, C.S., Allan, V.L.: *Scientific computing in the cavendish laboratory*. In prep. (2021)
15. Leslie, P.R.R.: An investigation of the clustering of radio stars. *MNRAS***122**, 371 (1961). DOI 10.1093/mnras/122.5.371
16. Nishimura, A., Matsumoto, T., Yonetsu, T., Nakao, Y., Fujita, S., Maezawa, H., Onishi, T., Ogawa, H.: Observational demonstration of a low-cost fast Fourier transform spectrometer with a delay-line-based ramp-compare ADC implemented on FPGA. arXiv e-prints arXiv:2104.00252 (2021)
17. Nobel Media AB: The nobel prize in physics 1974 (1974 Apr 7 [cited 2021 Apr 7]). URL <https://www.nobelprize.org/prizes/physics/1974/summary/>
18. Offer, A.: Charles feinstein (1932–2004), and british historical national accounts. Discussion Papers in Economic and Social History (2008 Jun, [cited 2021 Apr 7]). URL www.nuff.ox.ac.uk/economics/history/Paper70/70offer.pdf

19. Penzias, A.A., Wilson, R.W.: A Measurement of Excess Antenna Temperature at 4080 Mc/s. *ApJ***142**, 419–421 (1965). DOI 10.1086/148307
20. Ryle, M.: The Mullard Radio Astronomy Observatory, Cambridge. *Nature***180**(4577), 110–112 (1957). DOI 10.1038/180110a0
21. Ryle, M.: Radio telescopes of large resolving power. In: S. Lundqvist (ed.) *Nobel Lectures, Physics 1971-1980*. World Scientific Publishing Co., Singapore (1992). URL https://www.nobelprize.org/nobel_prizes/physics/laureates/1974/ryle-lecture.pdf
22. Ryle, M., Clarke, R.W.: An examination of the steady-state model in the light of some recent observations of radio sources. *MNRAS***122**, 349 (1961). DOI 10.1093/mnras/122.4.349
23. Ryle, M., Neville, A.C.: A radio survey of the North Polar region with a 4.5 minute of arc pencil-beam system. *MNRAS***125**, 39 (1962). DOI 10.1093/mnras/125.1.39
24. Scheuer, P.A.G., Ryle, M.: An investigation of the H II regions by a radio method. *MNRAS***113**, 3 (1953). DOI 10.1093/mnras/113.1.3
25. Scott, P.F., Ryle, M.: The number-flux density relation for radio sources away from the galactic plane. *MNRAS***122**, 389 (1961). DOI 10.1093/mnras/122.5.389
26. Scott, P.F., Ryle, M., Hewish, A.: First results of radio star observations using the method of aperture synthesis. *MNRAS***122**, 95–111 (1961). DOI 10.1093/mnras/122.1.95
27. Shetterly, M.L.: *Hidden figures : the untold story of the African-American women who helped win the space race / Margot Lee Shetterly*. HarperCollins Publishers (2016)
28. Smith, F.G.: An Attempt to Measure the Annual Parallax or Proper Motion of Four Radio Stars. *Nature***168**(4283), 962–963 (1951). DOI 10.1038/168962a0
29. Tunmer, H.E.C.: On the dimensions, spatial density and power of the extra-galactic radio sources. *MNRAS***120**, 360 (1960). DOI 10.1093/mnras/120.4.360
30. Turtle, A.J., Baldwin, J.E.: A survey of galactic radiation at 178 Mc/s. *MNRAS***124**, 459 (1962). DOI 10.1093/mnras/124.6.459
31. University of Cambridge: Judith Ann Bailey 20 May 1934 - 6 May 2008 (2008 [cited 2021 Apr 7]). URL <https://www.cam.ac.uk/news/judith-ann-bailey-20-may-1934-6-may-2008>
32. University of Cambridge Computer Laboratory: Edsac 1 and after - a compilation of personal reminiscences (1999, [cited 2021 Apr 7]). URL <https://www.cl.cam.ac.uk/events/EDSAC99/reminiscences/>
33. Veenboer, B., Petschow, M., van der Tol, S., Offringa, A.R.: *IDG: Image Domain Gridding* (2019)
34. Wilkes, M.V.: *Memoirs of a Computer Pioneer*. The MIT Press series in the History of Computing. MIT Press, Cambridge (MA); London (1985)
35. Wynn-Williams, C.E.: The use of thyratrons for high speed automatic counting physical phenomena. *Proceedings of the Royal Society London A* pp. 295–310 (1931). URL <https://doi.org/10.1098/rspa.1931.0102>
36. Ye, H., Gull, S.F., Tan, S.M., Nikolic, B.: Optimal gridding and degrading in radio interferometry imaging. *MNRAS***491**(1), 1146–1159 (2020). DOI 10.1093/mnras/stz2970