

NON-INVERTIBLE PLANAR SELF-AFFINE SETS

ANTTI KÄENMÄKI AND PETTERI NISSINEN

ABSTRACT. We compare the dimension of a non-invertible self-affine set to the dimension of the respective invertible self-affine set. In particular, for generic planar self-affine sets, we show that the dimensions coincide when they are large and differ when they are small. Our study relies on thermodynamical formalism where, for dominated and irreducible matrices, we completely characterize the behavior of the pressures.

1. INTRODUCTION

Let J be a finite set and $(A_i + v_i)_{i \in J}$ a tuple of contractive affine self-maps on \mathbb{R}^2 , where we have written $A + v$ to denote the affine map $x \mapsto Ax + v$ defined on \mathbb{R}^2 for all matrices $A \in M_2(\mathbb{R})$ and translation vectors $v \in \mathbb{R}^2$. If the affine maps $A_i + v_i$ do not have a common fixed point, then we call such a tuple an *affine iterated function system*. We also write $f_i = A_i + v_i$ for all $i \in J$ and note that the associated tuple of matrices $(A_i)_{i \in J}$ is an element of $M_2(\mathbb{R})^J$.

A classical result of Hutchinson [14] shows that for each affine iterated function system $(f_i)_{i \in J}$ there exists a unique non-empty compact set $X' \subset \mathbb{R}^2$, called the *self-affine set*, such that

$$X' = \bigcup_{i \in J} f_i(X'). \quad (1.1)$$

In this article, if $I = \{i \in J : A_i \text{ is invertible}\}$ is non-empty, then the self-affine set $X \subset X'$ associated to $(f_i)_{i \in I}$ is called *invertible*, and if $J \setminus I$ is non-empty, then the self-affine set X' associated to $(f_i)_{i \in J}$ is called *non-invertible*.

In our main result, Theorem 1.1 below, the part (1) shows that generically under a separation condition the dimensions of X' and X agree when they are at least 1. Furthermore, if the dimension of X is strictly less than 1, then the part (2) demonstrates that generically the dimensions of X' and X are distinct. Regarding the part (3), let us first recall that Marstrand's projection theorem [19] gives $\dim_{\mathbb{H}}(\text{proj}_V(X')) = \min\{1, \dim_{\mathbb{H}}(X')\}$ for Lebesgue almost all $V \in \mathbb{RP}^1$. Although the equality holds for generic V , it is often difficult to say whether a particular V satisfies it. The purpose of the part (3) is to verify that the orthogonal complement of the kernel of one of the rank one matrices is such a direction.

The precise definitions of the assumptions used in the theorem will be given in coming sections.

Date: May 17, 2022.

2000 Mathematics Subject Classification. Primary 28A80; Secondary 37C45, 37D35.

Key words and phrases. Self-affine set, Hausdorff dimension.

Theorem 1.1. *Suppose that X' and X are the planar self-affine sets associated to affine iterated function systems $(A_i + v_i)_{i \in J}$ and $(A_i + v_i)_{i \in I}$ such that $A_i \in GL_2(\mathbb{R})$ for all $i \in I \subset J$, respectively.*

- (1) *If $(A_i)_{i \in I}$ is strictly affine and strongly irreducible such that $\dim_{\text{aff}}((A_i)_{i \in I}) \geq 1$ and X satisfies the strong open set condition, then*

$$\begin{aligned} \overline{\dim}_M(X') &= \dim_H(X), \\ \dim_H(\text{proj}_V(X')) &= 1 \end{aligned}$$

for all $V \in \mathbb{RP}^1$.

- (2) *If $(A_i)_{i \in J}$ is dominated or irreducible such that $\max_{i \in J} \|A_i\| < \frac{1}{2}$, contains a rank one matrix, and $\dim_{\text{aff}}((A_i)_{i \in I}) < 1$, then*

$$\dim_H(X'_v) > \overline{\dim}_M(X_v)$$

for $\mathcal{L}^{2\#J}$ -almost all translation vectors $\mathbf{v} = (v_i)_{i \in J} \in (\mathbb{R}^2)^{\#J}$.

- (3) *If $(A_i)_{i \in J}$ contains a rank one matrix, $(A_i)_{i \in I}$ is strictly affine and strongly irreducible such that $\dim_{\text{aff}}((A_i)_{i \in I}) < 1$, and X satisfies the strong open set condition, then there exists a rank one matrix A in \mathbf{A} such that*

$$\dim_H(X') = \dim_H(\text{proj}_{\ker(A)^\perp}(X')) \leq 1.$$

The remaining of the paper is organized as follows. In Section 2, we compare the behavior of the pressures and study the continuity. In particular, for dominated and irreducible matrices, we completely characterize the continuity of the pressure in the non-invertible case. In Section 3, we uncover how the study of non-invertible self-affine sets is connected to the theory of sub-self-affine and inhomogeneous self-affine sets, and prove the main result.

2. PRODUCTS OF MATRICES

2.1. Rank one matrices. We denote the collection of all 2×2 matrices with real entries by $M_2(\mathbb{R})$, the general linear group of degree 2 over \mathbb{R} by $GL_2(\mathbb{R}) \subset M_2(\mathbb{R})$, and the orthogonal group in dimension 2 over \mathbb{R} by $O_2(\mathbb{R}) \subset GL_2(\mathbb{R})$. A matrix $A \in GL_2(\mathbb{R})$ is called *proximal* if it has two real eigenvalues with different absolute values. If $A \in M_2(\mathbb{R})$, then the *singular values* of A are defined to be the non-negative square roots of the eigenvalues of the positive-semidefinite matrix $A^\top A$ and are denoted by $\alpha_1(A)$ and $\alpha_2(A)$ in non-increasing order. Recall that the rank of A is the number of non-zero singular values of A . The identities $\alpha_1(A) = \|A\|$ and $\alpha_1(A)\alpha_2(A) = |\det(A)|$ for all $A \in M_2(\mathbb{R})$ are standard, as is the identity $\alpha_2(A) = \|A^{-1}\|^{-1}$ in the case where A is invertible. For each $A \in M_2(\mathbb{R})$ and $s \geq 0$ we define the *singular value function* by setting

$$\varphi^s(A) = \begin{cases} \alpha_1(A)^s, & \text{if } 0 \leq s \leq 1, \\ \alpha_1(A)\alpha_2(A)^{s-1}, & \text{if } 1 < s \leq 2, \\ |\det(A)|^{s/2}, & \text{if } 2 < s < \infty, \end{cases}$$

where we interpret $0^0 = 1$. The value $\varphi^s(A)$ represents a measurement of the s -dimensional volume of the image of the Euclidean unit ball under A . Since $\alpha_1(A)\alpha_2(A)^{s-1} = \alpha_1(A)^{2-s}|\det(A)|^{s-1}$ for all $1 < s \leq 2$, the inequality $\varphi^s(AB) \leq \varphi^s(A)\varphi^s(B)$ is valid for all $s \geq 0$. In other words, the singular value function is sub-multiplicative.

Note that if $A \in M_2(\mathbb{R})$ has rank one, then $\varphi^s(A) = 0$ for all $s > 1$. Recalling that A has rank zero if and only if A is the zero matrix, we see that $\varphi^s(A) = 0$ for all $s > 0$. Let us next recall that rank one matrices are projections. Let \mathbb{RP}^1 be the real projective line, that is, the set of all lines through the origin in \mathbb{R}^2 . If $V, W \in \mathbb{RP}^1$, then the *projection* $\text{proj}_V^W: \mathbb{R}^2 \rightarrow V$ is the linear map such that $\text{proj}_V^W|_V = \text{Id}|_V$ and $\ker(\text{proj}_V^W) = W$. Furthermore, the *orthogonal projection* $\text{proj}_V^{V^\perp}$ onto the subspace V is denoted by proj_V . The following lemma is well-known. But, as the proof is short, we provide the reader with full details.

Lemma 2.1. *A matrix $A \in M_2(\mathbb{R})$ has rank one if and only if there exist $v, w \in \mathbb{R}^2 \setminus \{(0, 0)\}$ such that $A = vw^\top$. In this case,*

$$A = \begin{cases} \langle v, w \rangle \text{proj}_{\text{im}(A)}^{\ker(A)}, & \text{if } A \text{ is not nilpotent,} \\ |v||w|R \text{proj}_{\ker(A)^\perp}, & \text{if } A \text{ is nilpotent,} \end{cases}$$

where $R \in O_2(\mathbb{R})$ is a rotation by an angle $\pi/2$. In particular, $A(X)$ is bi-Lipschitz equivalent to $\text{proj}_{\ker(A)^\perp}(X)$ for all $X \subset \mathbb{R}^2$.

Proof. Let us first prove the characterization of rank one matrices. If $A = vw^\top$ for some $v, w \in \mathbb{R}^2 \setminus \{(0, 0)\}$, then $Ax = vw^\top x = \langle w, x \rangle v$ for all $x \in \mathbb{R}^2$. Therefore, A maps every x to a scalar multiple of v , $\text{rank}(A) = 1$, and $\text{im}(A) = \text{span}(v)$. If $x \in \text{span}(w)^\perp$, then $Ax = vw^\top x = \langle w, x \rangle v = 0$ and $\ker(A) = \text{span}(w)^\perp$. Conversely, if $\text{rank}(A) = 1$, then there is $v \in \mathbb{R}^2 \setminus \{(0, 0)\}$ such that Ax is a scalar multiple of v for all $x \in \mathbb{R}^2$. In particular, this is true when $x = (1, 0)$ and $x = (0, 1)$. That is, there are $w_1, w_2 \in \mathbb{R} \setminus \{0\}$ such that $A(1, 0) = w_1 v$ and $A(0, 1) = w_2 v$. In other words,

$$A = \begin{pmatrix} w_1 v_1 & w_2 v_1 \\ w_1 v_2 & w_2 v_2 \end{pmatrix} = vw^\top,$$

where $w = (w_1, w_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}$.

Let us then show that a rank one matrix A is a projection. If A is not nilpotent, then $\text{span}(v) = \text{im}(A) \neq \ker(A) = \text{span}(w)^\perp$. Since $Ax = vw^\top x = \langle x, w \rangle v$ and it is easy to see that

$$\text{proj}_{\text{im}(A)}^{\ker(A)}(x) = \frac{\langle x, w \rangle}{\langle v, w \rangle} v = \frac{1}{\langle v, w \rangle} Ax$$

for all $x \in \mathbb{R}^2$, we have shown the first case. If A is nilpotent, then $\text{span}(v) = \text{im}(A) = \ker(A) = \text{span}(w)^\perp$. Since $Rw/|w| = v/|v|$, where $R \in O_2(\mathbb{R})$ is a rotation by an angle $\pi/2$, we have

$$\text{proj}_{\ker(A)^\perp}(x) = \text{proj}_{\text{span}(w)}(x) = \frac{\langle x, w \rangle}{|w|^2} w = \frac{\langle x, w \rangle}{|v||w|} R^{-1}v$$

and hence,

$$R \text{proj}_{\ker(A)^\perp}(x) = \frac{\langle x, w \rangle}{|v||w|} v = \frac{1}{|v||w|} Ax$$

as claimed.

Since the last claim follows immediately from the the fact that a rank one matrix is a projection, we have finished the proof. \square

2.2. Pressure. Let J be a finite set and $A = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ be a tuple of matrices. We say that A is *irreducible* if there does not exist $V \in \mathbb{RP}^1$ such that $A_i V \subset V$ for all $i \in J$; otherwise A is *reducible*. Note that the irreducibility is equivalent to the property that the matrices in A do not have a common eigenvector. Therefore, A is reducible if and only if the matrices in A can simultaneously be presented (in some coordinate system) as upper triangular matrices. The tuple A is *strongly irreducible* if there does not exist a finite set $\mathcal{V} \subset \mathbb{RP}^1$ such that $A_i \mathcal{V} = \mathcal{V}$ for all $i \in J$.

We call a proper subset $\mathcal{C} \subset \mathbb{RP}^1$ a *multicone* if it is a finite union of closed non-trivial projective intervals. We say that A is *dominated* if each matrix A_i is non-zero and there exists a multicone $\mathcal{C} \subset \mathbb{RP}^1$ such that $A_i \mathcal{C} \subset \mathcal{C}^\circ$ for all $i \in J$, where \mathcal{C}° is the interior of \mathcal{C} . Conversely, if a multicone $\mathcal{C} \subset \mathbb{RP}^1$ satisfies such a condition, then we say that \mathcal{C} is a *strongly invariant multicone* for A . For example, the first quadrant is strongly invariant for any tuple of positive matrices. Note that a dominated tuple is not necessarily irreducible and vice versa. If $A \in GL_2(\mathbb{R})^J$ is dominated and irreducible, then, by [3, Lemma 2.10], A is strongly irreducible.

We let J^* denote the set of all finite words $\{\emptyset\} \cup \bigcup_{n \in \mathbb{N}} J^n$, where \emptyset satisfies $\emptyset \mathbf{i} = \mathbf{i} \emptyset = \mathbf{i}$ for all $\mathbf{i} \in J^*$. For notational convenience, we set $J^0 = \{\emptyset\}$. The set $J^\mathbb{N}$ is the collection of all infinite words. We define the *left shift* $\sigma: J^\mathbb{N} \rightarrow J^\mathbb{N}$ by setting $\sigma \mathbf{i} = i_2 i_3 \dots$ for all $\mathbf{i} = i_1 i_2 \dots \in J^\mathbb{N}$. The concatenation of two words $\mathbf{i} \in J^*$ and $\mathbf{j} \in J^* \cup J^\mathbb{N}$ is denoted by $\mathbf{ij} \in J^* \cup J^\mathbb{N}$ and the length of $\mathbf{i} \in J^* \cup J^\mathbb{N}$ is denoted by $|\mathbf{i}|$. If $\mathbf{j} \in J^* \cup J^\mathbb{N}$ and $1 \leq n < |\mathbf{j}|$, then we define $\mathbf{j}|_n$ to be the unique word $\mathbf{i} \in J^n$ for which $\mathbf{ik} = \mathbf{j}$ for some $\mathbf{k} \in J^* \cup J^\mathbb{N}$. Furthermore, if $\mathbf{i} \in J^n$ for some $n \in \mathbb{N}$, then we set $[\mathbf{i}] = \{\mathbf{j} \in J^\mathbb{N} : \mathbf{j}|_n = \mathbf{i}\}$. The set $[\mathbf{i}]$ is called a *cylinder set*. We write $A_{\mathbf{i}} = A_{i_1} \dots A_{i_n}$ for all $\mathbf{i} = i_1 \dots i_n \in J^n$ and $n \in \mathbb{N}$. We say that $A \in GL_2(\mathbb{R})^J$ is *strictly affine* if there is $\mathbf{i} \in J^*$ such that $A_{\mathbf{i}}$ is proximal. Recall that $A \in GL_2(\mathbb{R})$ is *proximal* if it has two real eigenvalues with different absolute values. By [2, Corollary 2.4], a dominated tuple in $GL_2(\mathbb{R})^J$ is strictly affine.

If $\Gamma \subset J^\mathbb{N}$ is a non-empty compact set such that $\sigma(\Gamma) \subset \Gamma$, then we define $\Gamma_n = \{\mathbf{i}|_n : \mathbf{i} \in \Gamma\}$ and $\Gamma_* = \bigcup_{n \in \mathbb{N}} \Gamma_n$. We keep denoting $(J^\mathbb{N})_n$ and $(J^\mathbb{N})_*$ by I^n and I^* , respectively, for all $I \subset J$ and $n \in \mathbb{N}$. Given a tuple $A = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ of matrices, we define for each such $\Gamma \subset J^\mathbb{N}$ and $s \geq 0$ the *pressure* by setting

$$P(\Gamma, A, s) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\mathbf{i} \in \Gamma_n} \varphi^s(A_{\mathbf{i}}) = \inf_{n \in \mathbb{N}} \frac{1}{n} \log \sum_{\mathbf{i} \in \Gamma_n} \varphi^s(A_{\mathbf{i}}) \in [-\infty, \infty).$$

The assumption $\sigma(\Gamma) \subset \Gamma$ guarantees that if $\mathbf{i} \in J^m$ and $\mathbf{j} \in J^n$ such that $\mathbf{ij} \in \Gamma_{m+n}$, then $\mathbf{i} \in \Gamma_m$ and $\mathbf{j} \in \Gamma_n$. Therefore, as the singular value function is sub-multiplicative, the sequence $(\log \sum_{\mathbf{i} \in \Gamma_n} \varphi^s(A_{\mathbf{i}}))_{n \in \mathbb{N}}$ is sub-additive and hence, the limit above exists or is $-\infty$ by Fekete's lemma.

Let A be a tuple of strictly contractive matrices and $\Gamma \subset J^\mathbb{N}$ be a non-empty compact set such that $\sigma(\Gamma) \subset \Gamma$. Since $\varphi^s(A_i) \leq \varphi^t(A_i) \max_{k \in J} \|A_k\|^{(s-t)}$ for all $i \in J$, we see that $P(\Gamma, A, s) \leq P(\Gamma, A, t) + (s-t) \log \max_{k \in J} \|A_k\|$ for all $s > t \geq 0$. Since A consists only of strictly contractive matrices, we have $\max_{k \in J} \|A_k\| < 1$ and hence, the pressure $P(\Gamma, A, s)$ is strictly decreasing as a function of s whenever it is finite. Notice also that $P(\Gamma, A, 0) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \#\Gamma_n \geq 0$ and $\lim_{s \rightarrow \infty} P(\Gamma, A, s) = -\infty$. In this case, we

define the *affinity dimension* by setting

$$\dim_{\text{aff}}(\Gamma, \mathbf{A}) = \inf\{s \geq 0 : P(\Gamma, \mathbf{A}, s) \leq 0\}.$$

Notice that if the pressure $s \mapsto P(\Gamma, \mathbf{A}, s)$ is continuous at $s_0 = \dim_{\text{aff}}(\Gamma, \mathbf{A})$, then $P(\Gamma, \mathbf{A}, s_0) = 0$.

We are interested in the properties of the pressure

$$P(\mathbf{A}, s) = P(J^{\mathbb{N}}, \mathbf{A}, s)$$

as a function of s and the affinity dimension $\dim_{\text{aff}}(\mathbf{A}) = \dim_{\text{aff}}(J^{\mathbb{N}}, \mathbf{A})$. To that end, let us introduce some further notation. Let $I = \{i \in J : A_i \text{ is invertible}\}$. In this case, we trivially have that

$$I^{\mathbb{N}} = \{\mathbf{i} \in J^{\mathbb{N}} : A_{\mathbf{i}|_n} \text{ is invertible for all } n \in \mathbb{N}\}$$

is a compact subset of $J^{\mathbb{N}}$ and satisfies $\sigma(I^{\mathbb{N}}) = I^{\mathbb{N}}$. Therefore, the pressure $P(I^{\mathbb{N}}, \mathbf{A}, s)$ is well-defined for all $s \geq 0$. We also define

$$\Sigma = \{\mathbf{i} \in J^{\mathbb{N}} : A_{\mathbf{i}|_n} \text{ is non-zero for all } n \in \mathbb{N}\}.$$

It is easy to see that Σ is a compact subset of $J^{\mathbb{N}}$ and satisfies $\sigma(\Sigma) \subset \Sigma$. Indeed, if $\mathbf{j} \in \sigma(\Sigma)$, then there is $\mathbf{i} \in \Sigma$ such that $\mathbf{j} = \sigma\mathbf{i}$ and $A_{\mathbf{i}|_n} \neq 0$ for all $n \in \mathbb{N}$. As clearly $A_{\sigma\mathbf{i}|_n} \neq 0$ for all $n \in \mathbb{N}$, we see that $\mathbf{j} = \sigma\mathbf{i} \in \Sigma$ as claimed. Hence, also the pressure $P(\Sigma, \mathbf{A}, s)$ is well-defined for all $s \geq 0$. Observe that the inclusion $\sigma(\Sigma) \subset \Sigma$ can be strict: if $J = \{0, 1\}$ and

$$A_0 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad A_1 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix},$$

then $\Sigma = \{0111 \dots, 111 \dots\}$ and $\sigma(\Sigma) = \{111 \dots\}$.

Lemma 2.2. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$, then*

$$P(\mathbf{A}, s) = \begin{cases} \log \#J, & \text{if } s = 0, \\ P(\Sigma, \mathbf{A}, s), & \text{if } 0 < s \leq 1, \\ P(I^{\mathbb{N}}, \mathbf{A}, s), & \text{if } 1 < s < \infty. \end{cases}$$

Furthermore, the function $s \mapsto P(\mathbf{A}, s)$ is uniformly continuous on $(0, \infty) \setminus \{1\}$ whenever it is finite.

Proof. Recall first that $\varphi^s(A) = \alpha_1(A)^s = \|A\|^s$ for all $0 \leq s \leq 1$. Therefore, as we interpreted $0^0 = 1$, we have

$$P(\mathbf{A}, 0) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\mathbf{i} \in J^n} \|A_{\mathbf{i}}\|^0 = \log \#J.$$

Since $\alpha_1(A) > 0$ if and only if $A \in M_2(\mathbb{R})$ is non-zero, we see that for each $0 < s \leq 1$ the singular value satisfies $\varphi^s(A_{\mathbf{i}}) = \|A_{\mathbf{i}}\|^s > 0$ if and only if $\mathbf{i} \in \Sigma_*$. Therefore, $P(\mathbf{A}, s) = P(\Sigma, \mathbf{A}, s)$ for all $0 < s \leq 1$ and, by [18, proof of Lemma 2.1], the function $s \mapsto P(\mathbf{A}, s)$ is uniformly continuous on $(0, 1)$ whenever it is finite. Furthermore, since $\alpha_2(A) > 0$ if and only if $A \in GL_2(\mathbb{R})$, we have $\varphi^s(A_{\mathbf{i}}) > 0$ if and only if $\mathbf{i} \in I^*$ whenever $1 < s < \infty$. This shows that $P(\mathbf{A}, s) = P(I^{\mathbb{N}}, \mathbf{A}, s)$ for all $1 < s < \infty$ and, by [18, Lemma 2.1], the function $s \mapsto P(\mathbf{A}, s)$ is uniformly continuous on $(1, \infty)$ whenever it is finite. \square

The following lemma characterizes the continuity of the function $s \mapsto P(\mathbf{A}, s)$ at 0.

Lemma 2.3. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$, then the function $s \mapsto P(\mathbf{A}, s)$ is right-continuous at 0 if and only if the semigroup $\{A_i : i \in J^*\}$ does not contain rank zero matrices.*

Proof. If the semigroup $\{A_i : i \in J^*\}$ does not contain rank zero matrices, then $\Sigma = J^{\mathbb{N}}$ and the right-continuity at 0 is guaranteed by [18, proof of Lemma 2.1].

If A_i has rank zero for some $i \in J^n$ and $n \in \mathbb{N}$, then clearly $\#\Sigma_n < \#J^n = (\#J)^n$. Fix $0 < s \leq 1$ and notice that Lemma 2.2 implies

$$P(\mathbf{A}, s) \leq \frac{1}{n} \log \sum_{i \in \Sigma_n} \|A_i\|^s$$

and

$$\lim_{s \downarrow 0} P(\mathbf{A}, s) \leq \frac{1}{n} \log \#\Sigma_n < \frac{1}{n} \log \#J^n = P(\mathbf{A}, 0),$$

where the limit exists by the uniform continuity on $(0, 1)$. In particular, the function $s \mapsto P(\mathbf{A}, s)$ is not right-continuous at 0. \square

The possible discontinuity at 1 has already been observed by Feng and Shmerkin [12, Remark 1.1]. In their example, the pressure is not finite when $s > 1$, but it is easy to see that this is not a necessity. If $J = \{0, 1\}$ and

$$A_0 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

then, by Lemma 2.2, for $\mathbf{A} = (A_0, A_1) \in M_2(\mathbb{R})^J$ we have $P(\mathbf{A}, 1) = \log 2$ and $P(\mathbf{A}, s) = 0$ for all $s > 1$. The continuity of the function $s \mapsto P(\mathbf{A}, s)$ at 1 will be characterized for dominated and irreducible tuples in Lemma 2.10.

Let us next determine when the pressure is finite. For that, we need the following definition. Given a tuple $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ of matrices, we define the *joint spectral radius* by setting

$$\varrho(\mathbf{A}) = \lim_{n \rightarrow \infty} \max_{i \in J^n} \|A_i\|^{1/n}.$$

As the operator norm is sub-multiplicative, the sequence $(\log \max_{i \in J^n} \|A_i\|)_{n \in \mathbb{N}}$ is sub-additive and hence, the limit above exists by Fekete's lemma.

Lemma 2.4. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ is dominated or irreducible, then $\varrho(\mathbf{A}) > 0$.*

Proof. Let us first assume that \mathbf{A} is dominated and $\mathcal{C} \subset \mathbb{RP}^1$ is a strongly invariant multicone for \mathbf{A} . Since there exists a multicone $\mathcal{C}_0 \subset \mathbb{RP}^1$ such that $\bigcup_{i \in J^n} A_i \mathcal{C} \subset \mathcal{C}_0 \subset \mathcal{C}^o$ for all $n \in \mathbb{N}$, we find, by applying [6, Lemma 2.2], a constant $\kappa > 0$ such that

$$\|A_i|V\| \geq \kappa \|A_i\| \tag{2.1}$$

for all $V \in \mathcal{C}_0$ and $i \in J^*$. It follows that if $V \in \mathcal{C}_0$, then $A_j V \in \mathcal{C}_0$ and $\|A_i A_j\| \geq \|A_i A_j|V\| = \|A_i|A_j V\| \|A_j|V\| \geq \kappa^2 \|A_i\| \|A_j\|$ for all $i, j \in J^*$. Therefore,

$$\varrho(\mathbf{A}) \geq \liminf_{n \rightarrow \infty} \max_{i_1 \dots i_n \in J^n} \kappa^{2(n-1)/n} \|A_{i_1}\|^{1/n} \dots \|A_{i_n}\|^{1/n} \geq \kappa^2 \min_{j \in J} \|A_j\| > 0$$

as claimed.

Although the proof in the irreducible case can be found in [15, Lemma 2.2], we present the full details for the convenience of the reader. Denote the unit circle by S^1 and suppose that for each $k \in \mathbb{N}$ there is $x_k \in S^1$ such that for every $i \in J$ we have $|A_i x_k| < \frac{1}{k}$. By the compactness of S^1 , there is $x \in S^1$ such that $|A_i x| = 0$ for all $i \in J$. Choosing $V = \text{span}(x) \in \mathbb{RP}^1$, we see that $A_i V = \{(0, 0)\} \subset V$ for all $i \in J$ and \mathbf{A} is reducible.

It follows that there is $\delta > 0$ such that for every $x \in S^1$ there exists $i \in J$ for which $|A_i x| \geq \delta$. Let us next apply this inductively. Fix $x_0 \in S^1$ and choose $i_1 \in J$ such that $|A_{i_1} x_0| \geq \delta$. Write $x_1 = A_{i_1} x_0$ and choose $i_2 \in J$ such that $|A_{i_2} \frac{x_1}{|x_1|}| \geq \delta$ whence $|A_{i_2} A_{i_1} x_0| = |A_{i_2} x_1| \geq \delta |x_1| = \delta |A_{i_1} x_0| \geq \delta^2$. Continuing in this manner, we find for each $n \in \mathbb{N}$ a word $\mathbf{i}_n \in J^n$ such that $\|A_{\mathbf{i}_n}\| \geq |A_{\mathbf{i}_n} x_0| \geq \delta^n$. Hence,

$$\varrho(\mathbf{A}) \geq \liminf_{n \rightarrow \infty} \|A_{\mathbf{i}_n}\|^{1/n} \geq \delta > 0$$

as wished. \square

The following two lemmas characterize the finiteness of the pressure.

Lemma 2.5. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ satisfies $\max_{i \in J} \|A_i\| < 1$, then the following four conditions are equivalent:*

- (1) $P(\mathbf{A}, s) > -\infty$ for all $0 \leq s \leq 1$,
- (2) $\lim_{s \downarrow 0} P(\mathbf{A}, s) > -\infty$,
- (3) there does not exist $n \in \mathbb{N}$ such that $A_{\mathbf{i}} = 0$ for all $\mathbf{i} \in J^n$,
- (4) there exists $\mathbf{j} \in J^{\mathbb{N}}$ such that $A_{\mathbf{j}|_n} \neq 0$ for all $n \in \mathbb{N}$.

Furthermore, all of these conditions hold if \mathbf{A} is dominated or irreducible.

Proof. Notice that the limit in (2) exists by Lemma 2.2 and the implications (1) \Rightarrow (2) and (4) \Rightarrow (3) are trivial. Let us first show the implication (2) \Rightarrow (3). If (3) does not hold, then there exists $n_0 \in \mathbb{N}$ such that $A_{\mathbf{i}} = 0$ for all $\mathbf{i} \in J^{n_0}$. Since now $\|A_{\mathbf{i}}\| = 0$ for all $\mathbf{i} \in J^n$ and $n \geq n_0$, we see that $P(\mathbf{A}, s) = -\infty$ for all $s > 0$ and (2) cannot hold.

Let us then show the implication (3) \Rightarrow (4). If (4) does not hold, then for every $\mathbf{j} \in J^{\mathbb{N}}$ there is $n(\mathbf{j}) \in \mathbb{N}$ such that $A_{\mathbf{j}|_{n(\mathbf{j})}} = 0$. By compactness of $J^{\mathbb{N}}$, there exist $M \in \mathbb{N}$ and $\mathbf{j}_1, \dots, \mathbf{j}_M \in J^{\mathbb{N}}$ such that $\{[\mathbf{j}_i|_{n(\mathbf{j}_i)}]\}_{i=1}^M$ still covers $J^{\mathbb{N}}$. Choosing $n = \max_{i \in \{1, \dots, M\}} n(\mathbf{j}_i)$, we see that for every $\mathbf{i} \in J^n$ there is $i \in \{1, \dots, M\}$ such that $A_{\mathbf{i}} = A_{\mathbf{j}_i|_{n(\mathbf{j}_i)}} A_{\sigma^{n(\mathbf{j}_i)} \mathbf{i}} = 0$ and (3) cannot hold.

Since \mathbf{A} is a tuple of strictly contractive matrices, the function $s \mapsto P(\mathbf{A}, s)$ is strictly decreasing whenever it is finite. Therefore, we have $P(\mathbf{A}, s) \geq P(\mathbf{A}, 1) \geq \log \varrho(\mathbf{A})$ for all $0 \leq s \leq 1$ and hence, $\varrho(\mathbf{A}) > 0$ implies the condition (1). Therefore, to conclude the proof, it suffices to show that the condition (3) implies $\varrho(\mathbf{A}) > 0$ and also verify $\varrho(\mathbf{A}) > 0$ when \mathbf{A} is dominated or irreducible. While the latter is immediately assured by Lemma 2.4, we also see that to prove the former, we may assume that \mathbf{A} is reducible. This means that, after possibly a change of basis, the matrices A_i in \mathbf{A} are of the form

$$A_i = \begin{pmatrix} a_i & b_i \\ 0 & c_i \end{pmatrix}$$

for all $i \in J$. Since $A_i(1, 0) = a_i(1, 0)$ and $A_i(\frac{b_i}{c_i - a_i}, 1) = c_i(\frac{b_i}{c_i - a_i}, 1)$ when $a_i \neq c_i$, we see that $\max\{|a_i|, |c_i|\} \leq \|A_i\|$ for all $i \in J$. As the product of upper triangular matrices is upper triangular with diagonal entries obtained as products of the corresponding

diagonal entries, we also have $\max\{|a_{i_1} \cdots a_{i_n}|, |c_{i_1} \cdots c_{i_n}|\} \leq \|A_{\mathbf{i}}\|$ for all $\mathbf{i} = i_1 \cdots i_n \in J^n$ and $n \in \mathbb{N}$. Therefore, if $\varrho(\mathbf{A}) = 0$, then

$$\max_{\mathbf{i} \in J} |a_{\mathbf{i}}| = \lim_{n \rightarrow \infty} \max_{i_1 \cdots i_n \in J^n} |a_{i_1} \cdots a_{i_n}|^{1/n} \leq \varrho(\mathbf{A}) = 0$$

and, similarly, $\max_{i \in J} |c_i| = 0$. In other words, the diagonal entries in all of the matrices $A_{\mathbf{i}}$ are zero. Thus, $A_{\mathbf{i}} = 0$ for all $\mathbf{i} \in J^2$ and (3) cannot hold. \square

Lemma 2.6. *If $\mathbf{A} = (A_{\mathbf{i}})_{\mathbf{i} \in J} \in M_2(\mathbb{R})^J$, then the following five conditions are equivalent:*

- (1) $P(I^{\mathbb{N}}, \mathbf{A}, s) > -\infty$ for all $s \geq 0$,
- (2) $P(\mathbf{A}, s) > -\infty$ for all $s \geq 0$,
- (3) $\lim_{s \downarrow 1} P(\mathbf{A}, s) > -\infty$,
- (4) there does not exist $n \in \mathbb{N}$ such that $A_{\mathbf{i}}$ has rank at most one for all $\mathbf{i} \in J^n$,
- (5) there exists $j \in J$ such that $A_j \in GL_2(\mathbb{R})$.

Proof. Notice that the limit in (3) exists by Lemma 2.2 and the implications (1) \Rightarrow (2) and (2) \Rightarrow (3) are trivial. Let us first show the implication (3) \Rightarrow (4). If (4) does not hold, then there exists $n_0 \in \mathbb{N}$ such that $A_{\mathbf{i}}$ has rank at most one for all $\mathbf{i} \in J^{n_0}$. It follows that for every $\mathbf{i} \in J^n$ and $n \geq n_0$ the rank of $A_{\mathbf{i}}$ is at most one as it is bounded above by the rank of $A_{\mathbf{i}|_{n_0}}$. Therefore, as $\varphi^s(A_{\mathbf{i}}) = 0$ for all $\mathbf{i} \in J^n$, $n \geq n_0$, and $s > 1$, we have $P(\mathbf{A}, s) = -\infty$ for all $s > 1$ and (3) cannot hold.

Let us then show the implication (4) \Rightarrow (5). If (5) does not hold, then A_j has rank at most one for all $j \in J$. It follows that for every $\mathbf{i} \in J^n$ and $n \in \mathbb{N}$ the rank of $A_{\mathbf{i}}$ is at most one and (4) cannot hold.

Finally, let us show the implication (5) \Rightarrow (1). The condition (5) implies that $A_{\mathbf{j}|_n} \in GL_2(\mathbb{R})$ for all $n \in \mathbb{N}$ where $\mathbf{j} = jj \cdots \in J^{\mathbb{N}}$. Since $\varphi^s(A_{\mathbf{j}|_n}) \geq \alpha_2(A_{\mathbf{j}|_n}) \geq \alpha_2(A_j)^n > 0$ for all $n \in \mathbb{N}$ and $s \geq 0$, we see that $P(I^{\mathbb{N}}, \mathbf{A}, s) \geq \log \alpha_2(A_j) > -\infty$ for all $s \geq 0$ as wished. \square

2.3. Equilibrium states. Let $\mathcal{M}_{\sigma}(J^{\mathbb{N}})$ be the collection of all σ -invariant Borel probability measures on $J^{\mathbb{N}}$. If $0 < s \leq 1$, then we say that a measure $\mu_K \in \mathcal{M}_{\sigma}(J^{\mathbb{N}})$ is *s-Gibbs-type* if there exists a constant $C \geq 1$ such that

$$C^{-1} e^{-nP(\mathbf{A}, s)} \|A_{\mathbf{i}}\|^s \leq \mu_K([\mathbf{i}]) \leq C e^{-nP(\mathbf{A}, s)} \|A_{\mathbf{i}}\|^s$$

for all $\mathbf{i} \in J^n$ and $n \in \mathbb{N}$.

Lemma 2.7. *If $\mathbf{A} = (A_{\mathbf{i}})_{\mathbf{i} \in J} \in M_2(\mathbb{R})^J$ satisfies $\max_{i \in J} \|A_i\| < 1$ and is dominated or irreducible, then for every $0 < s \leq 1$ there exist a unique ergodic s-Gibbs-type measure $\mu_K \in \mathcal{M}_{\sigma}(J^{\mathbb{N}})$.*

Proof. Recall first that, by Lemma 2.5, the pressure $P(\mathbf{A}, s)$ is finite for all $0 < s \leq 1$. If \mathbf{A} is irreducible, then the existence of the claimed measure $\mu_K \in \mathcal{M}_{\sigma}(J^{\mathbb{N}})$ follows immediately from [11, Proposition 1.2]. We may thus assume that \mathbf{A} is dominated. Fix $0 < s \leq 1$ and notice that, by (2.1), there exist $\kappa > 0$ and a multicone $\mathcal{C}_0 \subset \mathbb{R}\mathbb{P}^1$ such that $\|A_{\mathbf{i}}|V\| \geq \kappa \|A_{\mathbf{i}}\|$ for all $V \in \mathcal{C}_0$ and $\mathbf{i} \in J^*$. Fixing $V \in \mathcal{C}_0$ and defining a Hölder continuous function $g: J^{\mathbb{N}} \rightarrow \mathbb{R}$ by setting

$$g(\mathbf{i}) = \log \|A_{\mathbf{i}|_1} |A_{\sigma \mathbf{i}} V\|^s,$$

we see that

$$\log \|A_{\mathbf{i}|_n}\|^s + \log \kappa^s \leq \sum_{k=0}^{n-1} g(\sigma^k \mathbf{i}) \leq \log \|A_{\mathbf{i}|_n}\|^s$$

for all $\mathbf{i} \in J^{\mathbb{N}}$ and $n \in \mathbb{N}$. By [8, Theorems 1.7 and 1.16], there exist an ergodic measure $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ and a constant $C \geq 1$ such that

$$\kappa^s C^{-1} e^{-nP(\mathbf{A}, s)} \|A_{\mathbf{i}}\|^s \leq \mu_K([\mathbf{i}]) \leq C e^{-nP(\mathbf{A}, s)} \|A_{\mathbf{i}}\|^s$$

for all $\mathbf{i} \in J^n$ and $n \in \mathbb{N}$; see also [3, Lemma 2.12]. The uniqueness of μ_K is now evident as two different ergodic measures are mutually singular. \square

If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ is dominated, then it follows from (2.1) that $\|A_{\mathbf{i}}\| \geq \kappa^{2(n-1)} \|A_{i_1}\| \cdots \|A_{i_n}\| \geq \kappa^{2(n-1)} \min_{i \in J} \|A_i\|^n > 0$ for all $\mathbf{i} = i_1 \cdots i_n \in J^n$ and $n \in \mathbb{N}$. Hence the semigroup $\{A_{\mathbf{i}} : \mathbf{i} \in J^*\}$ does not contain rank zero matrices and, by Lemma 2.3, the function $s \mapsto P(\mathbf{A}, s)$ is right-continuous at 0. Furthermore, as in this case $\Sigma = J^{\mathbb{N}}$, the s -Gibbs-type measure $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ is fully supported on $J^{\mathbb{N}}$. If \mathbf{A} is irreducible, then μ_K is supported only on Σ .

Given $\mu \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ and $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$, we define for each $s \geq 0$ the *energy* by setting

$$\Lambda(\mu, \mathbf{A}, s) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mathbf{i} \in J^n} \mu([\mathbf{i}]) \log \varphi^s(A_{\mathbf{i}}) = \inf_{n \in \mathbb{N}} \frac{1}{n} \sum_{\mathbf{i} \in J^n} \mu([\mathbf{i}]) \log \varphi^s(A_{\mathbf{i}}).$$

The limit above exists or is $-\infty$ again by Fekete's lemma. Recall that the *entropy* of μ is

$$h(\mu) = - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mathbf{i} \in J^n} \mu([\mathbf{i}]) \log \mu([\mathbf{i}]).$$

It is well-known that

$$P(\mathbf{A}, s) \geq h(\mu) + \Lambda(\mu, \mathbf{A}, s) \tag{2.2}$$

for all $\mu \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ and $s \geq 0$; for example, see [18, §3]. A measure $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ is an *s-equilibrium state* if it satisfies

$$P(\mathbf{A}, s) = h(\mu_K) + \Lambda(\mu_K, \mathbf{A}, s) > -\infty. \tag{2.3}$$

The following lemma shows the uniqueness of the equilibrium state in dominated and irreducible cases.

Lemma 2.8. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ satisfies $\max_{i \in J} \|A_i\| < 1$ and is dominated or irreducible, then for every $0 < s \leq 1$ the ergodic s -Gibbs-type measure $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ is the unique s -equilibrium state.*

Proof. Fix $0 < s \leq 1$ and let $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ be the ergodic s -Gibbs-type measure. Since, by Lemmas 2.7 and 2.5,

$$\begin{aligned} h(\mu_K) + \Lambda(\mu_K, \mathbf{A}, s) &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mathbf{i} \in \Sigma_n} \mu_K([\mathbf{i}]) \log \frac{\|A_{\mathbf{i}}\|^s}{\mu_K([\mathbf{i}])} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mathbf{i} \in \Sigma_n} \mu_K([\mathbf{i}]) \log e^{nP(\mathbf{A}, s)} = P(\mathbf{A}, s) > -\infty, \end{aligned}$$

we see that μ_K is an s -equilibrium state. As μ_K is ergodic, the uniqueness follows from [18, Theorem 3.6]. \square

If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ contains an invertible matrix, then $I \neq \emptyset$ and, by Lemma 2.6, $P(I^{\mathbb{N}}, \mathbf{A}, s) > -\infty$ for all $s \geq 0$. In this case, regardless of domination and irreducibility, it follows from [16, Theorem 4.1] that for every $s > 0$ there exists an ergodic measure $\nu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ supported on $I^{\mathbb{N}}$ such that

$$P(I^{\mathbb{N}}, \mathbf{A}, s) = h(\nu_K) + \Lambda(\nu_K, \mathbf{A}, s). \quad (2.4)$$

Note that such a measure is not necessarily unique; see [7, 11, 17, 18].

Lemma 2.9. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ satisfies $\max_{i \in J} \|A_i\| < 1$, contains a rank one matrix, and is dominated or irreducible, then*

$$P(I^{\mathbb{N}}, \mathbf{A}, s) < P(\mathbf{A}, s)$$

for all $0 \leq s \leq 1$.

Proof. Since \mathbf{A} is dominated or irreducible, Lemma 2.5 shows that $P(\mathbf{A}, s) > -\infty$ for all $0 \leq s \leq 1$. Notice that, by Lemma 2.2, $P(I^{\mathbb{N}}, \mathbf{A}, 0) = \log \#I < \log \#J = P(\mathbf{A}, 0)$ and we may fix $0 < s \leq 1$. Therefore, by Lemma 2.8, there exists unique $\mu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ such that

$$P(\mathbf{A}, s) = h(\mu_K) + \Lambda(\mu_K, \mathbf{A}, s) > -\infty. \quad (2.5)$$

Furthermore, by Lemma 2.7, μ_K satisfies

$$\mu_K([i]) \geq C^{-1} e^{-nP(\mathbf{A}, s)} \|A_i\|^s > 0$$

for all $i \in \Sigma_n$ and $n \in \mathbb{N}$, where $C \geq 1$ is a constant. In particular, if A_k is a rank one matrix in \mathbf{A} , then $\mu_K([k]) > 0$.

If \mathbf{A} does not contain invertible matrices, then trivially $P(I^{\mathbb{N}}, \mathbf{A}, s) = -\infty$ for all $s > 0$ and there is nothing to prove. We may thus assume that \mathbf{A} contains an invertible matrix. Therefore, by (2.4), there exists a measure $\nu_K \in \mathcal{M}_\sigma(J^{\mathbb{N}})$ supported on $I^{\mathbb{N}}$ such that

$$P(I^{\mathbb{N}}, \mathbf{A}, s) = h(\nu_K) + \Lambda(\nu_K, \mathbf{A}, s). \quad (2.6)$$

Since A_k is not invertible and ν_K is supported on $I^{\mathbb{N}}$, we have $\nu_K([k]) = 0$. As μ_K is the unique measure in $\mathcal{M}_\sigma(J^{\mathbb{N}})$ satisfying (2.5) and $\mu_K([k]) > \nu_K([k])$, we see that ν_K does not satisfy (2.5) and therefore, by (2.6),

$$P(\mathbf{A}, s) > h(\nu_K) + \Lambda(\nu_K, \mathbf{A}, s) = P(I^{\mathbb{N}}, \mathbf{A}, s)$$

as claimed. \square

The following lemma characterizes the continuity of the function $s \mapsto P(\mathbf{A}, s)$ at 1.

Lemma 2.10. *If $\mathbf{A} = (A_i)_{i \in J} \in M_2(\mathbb{R})^J$ satisfies $\max_{i \in J} \|A_i\| < 1$ and is dominated or irreducible, then the function $s \mapsto P(\mathbf{A}, s)$ is continuous at 1 if and only if \mathbf{A} does not contain rank one matrices.*

Proof. If \mathbf{A} does not contain rank one matrices, then it contains only invertible or rank zero matrices. By Lemma 2.2, rank zero matrices do not have any effect on the value of the pressure $P(\mathbf{A}, s)$ when $s > 0$. Therefore, rank zero matrices have no impact on the

continuity at 1 and we may assume that $A \in GL_2(\mathbb{R})^J$. But in this case, the continuity follows from [18, Lemma 2.1].

Let us then assume that A contains a rank one matrix. If A does not contain invertible matrices, then, as the function $s \mapsto P(A, s)$ is strictly decreasing, Lemma 2.6 implies that $P(A, s) = -\infty$ for all $s > 1$. Furthermore, since A is dominated or irreducible, Lemma 2.5 shows that $P(A, s) > -\infty$ for all $0 \leq s \leq 1$ and the function $s \mapsto P(A, s)$ is discontinuous at 1. We may thus assume that A contains an invertible matrix. By Lemma 2.6, we thus have $P(I^{\mathbb{N}}, A, s) > -\infty$ for all $s \geq 0$. Recall that, by [18, Lemma 2.1], the function $s \mapsto P(I^{\mathbb{N}}, A, s)$ is continuous at 1. Therefore, by Lemma 2.2, showing

$$P(I^{\mathbb{N}}, A, 1) < P(A, 1)$$

proves the function $s \mapsto P(A, s)$ discontinuous at 1. But, as A contains a rank one matrix, this follows immediately from Lemma 2.9. \square

3. DIMENSION OF NON-INVERTIBLE SELF-AFFINE SETS

Recall that J is a finite set and the affine iterated function system is a tuple $(f_i)_{i \in J}$ a tuple of contractive affine self-maps on \mathbb{R}^2 not having a common fixed point. We write $f_i = A_i + v_i$ for all $i \in J$, where $A_i \in M_2(\mathbb{R})$ and $v_i \in \mathbb{R}^2$, and $f_{\mathbf{i}} = f_{i_1} \circ \cdots \circ f_{i_n}$ for all $\mathbf{i} = i_1 \cdots i_n \in J^n$ and $n \in \mathbb{N}$. We let $f_{\emptyset} = \text{Id}$ to be the identity map. Note that the associated tuple of matrices $(A_i)_{i \in J}$ is an element of $M_2(\mathbb{R})^J$ and satisfies $\max_{i \in J} \|A_i\| < 1$.

If $I = \{i \in J : A_i \text{ is invertible}\}$ is non-empty, then the invertible self-affine set X is associated to $(f_i)_{i \in I}$, and if $J \setminus I$ is non-empty, then the non-invertible self-affine set X' is associated to $(f_i)_{i \in J}$. Recall the defining property (1.1) of a self-affine set. We use the convention that whenever we speak about a self-affine set, then it is automatically accompanied with a tuple of affine maps which defines it. This makes it possible to write that e.g. “a non-invertible self-affine set is dominated” which obviously then means that “the associated tuple $A = (A_i)_{i \in J}$ of matrices in $M_2(\mathbb{R})^J$ is dominated”.

The study of non-invertible self-affine sets is connected to the theory of sub-self-affine sets. If the *canonical projection* $\pi: J^{\mathbb{N}} \rightarrow \mathbb{R}^2$ is defined such that

$$\pi(\mathbf{i}) = \lim_{n \rightarrow \infty} f_{\mathbf{i}|_n}(0) = \lim_{n \rightarrow \infty} \sum_{k=1}^n A_{\mathbf{i}|_{k-1}} v_{i_k}$$

for all $\mathbf{i} = i_1 i_2 \cdots \in J^{\mathbb{N}}$, then we write $X'' = \pi(\Sigma)$, where $\Sigma = \{\mathbf{i} \in J^{\mathbb{N}} : A_{\mathbf{i}|_n} \text{ is non-zero for all } n \in \mathbb{N}\}$. Observe that $X = \pi(I^{\mathbb{N}}) \subset X'' \subset \pi(J^{\mathbb{N}}) = X'$ and, as $\sigma(\Sigma) \subset \Sigma$, the set X'' is *sub-self-affine*, i.e.

$$X'' \subset \bigcup_{i \in J} f_i(X''); \quad (3.1)$$

see [18]. The study is also connected to inhomogeneous self-affine sets. If $C \subset \mathbb{R}^2$ is compact, then there exists a unique non-empty compact set $X_C \subset \mathbb{R}^2$ such that

$$X_C = \bigcup_{i \in I} f_i(X_C) \cup C.$$

The set X_C is called the *inhomogeneous self-affine set* with condensation C . Such sets were introduced by Barnsley and Demko [5] and studied by Barnsley [4]. Note that X_\emptyset is the invertible self-affine set X .

Lemma 3.1. *If X' and X are non-invertible and invertible planar self-affine sets, respectively, and X'' is the associated sub-self-affine set defined in (3.1), then $X' \setminus X''$ is countable and*

$$X' = X_C = X \cup \bigcup_{\mathbf{i} \in I^*} f_{\mathbf{i}}(C),$$

where $C = \bigcup_{i \in J \setminus I} f_i(X')$.

Proof. Let us first show that $X' \setminus X''$ is countable. Writing $v_{\mathbf{i}} = \sum_{k=1}^n A_{\mathbf{i}|_{k-1}} v_{i_k}$, we see that $f_{\mathbf{i}} = A_{\mathbf{i}} + v_{\mathbf{i}}$ for all $\mathbf{i} = i_1 \cdots i_n \in J^n$ and $n \in \mathbb{N}$. Let $\mathbf{i} \in J^{\mathbb{N}} \setminus \Sigma$ and choose $n_0(\mathbf{i}) = \min\{n \in \mathbb{N} : A_{\mathbf{i}|_n} \text{ is zero}\}$. Since $v_{\mathbf{i}|_{n+1}} = \sum_{k=1}^{n+1} A_{\mathbf{i}|_{k-1}} v_{i_k} = A_{\mathbf{i}|_n} v_{i_{n+1}} + \sum_{k=1}^n A_{\mathbf{i}|_{k-1}} v_{i_k} = v_{\mathbf{i}|_n}$ for all $n \geq n_0(\mathbf{i})$, a simple induction shows that

$$f_{\mathbf{i}|_n}(X') = \{v_{\mathbf{i}|_{n_0(\mathbf{i})}}\}$$

for all $n \geq n_0(\mathbf{i})$. As $J^{\mathbb{N}} \setminus \Sigma$ is clearly separable, there exist countably many infinite words $\mathbf{i}_1, \mathbf{i}_2, \dots \in J^{\mathbb{N}} \setminus \Sigma$ such that $J^{\mathbb{N}} \setminus \Sigma \subset \bigcup_{k \in \mathbb{N}} [\mathbf{i}_k|_{n_0(\mathbf{i}_k)}]$. It follows that

$$X' \setminus X'' \subset \{v_{\mathbf{i}_k|_{n_0(\mathbf{i}_k)}} : k \in \mathbb{N}\}$$

is countable.

Let us then prove the claimed equalities. Note that [21, Lemma 3.9] verifies

$$X_C = X \cup \bigcup_{\mathbf{i} \in I^*} f_{\mathbf{i}}(C). \quad (3.2)$$

To prove the remaining equality, let us first show that $X' \subset X_C$. To that end, fix $x \in X'$. By (3.2), we have $X \subset X_C$ and we may assume that $x \in X' \setminus X$. But this implies that there exist $\mathbf{i} \in I^*$ and $i \in J \setminus I$ such that $x \in f_{\mathbf{i}i}(X')$. Since, again by (3.2),

$$f_{\mathbf{i}i}(X') \subset f_{\mathbf{i}}(C) \subset \bigcup_{\mathbf{i} \in I^*} f_{\mathbf{i}}(C) \subset X_C$$

we have shown that $X' \subset X_C$. The inclusion $X_C \subset X'$ follows immediately from (3.2) since we trivially have $X \subset X'$ and $f_{\mathbf{i}}(C) \subset X'$ for all $\mathbf{i} \in I^*$. Thus $X' = X_C$ as claimed. \square

We are interested in the dimension of the non-invertible self-affine set. Relying on (1.1), the non-invertible self-affine set X' can naturally be covered by the sets $f_{\mathbf{i}}(B)$, where B is a ball containing X' . Note that such sets are ellipses or line segments, depending on whether the associated matrix is invertible or has rank one. Each set $f_{\mathbf{i}}(B)$ can be covered by one ball of radius $\alpha_1(A_{\mathbf{i}}) \text{diam}(B)$ or by $\alpha_1(A_{\mathbf{i}})/\alpha_2(A_{\mathbf{i}})$ many balls of radius $\alpha_2(A_{\mathbf{i}}) \text{diam}(B)$. This motivates us to study the limiting behavior of sums $\sum_{\mathbf{i} \in J^n} \varphi^s(A_{\mathbf{i}})$ and hence, the pressure $P(A, s)$.

Recall that the upper Minkowski dimension $\overline{\dim}_M$ is an upper bound for the Hausdorff dimension \dim_H for all compact sets; see [20, §5.3]. The following lemma shows that the affinity dimension is an upper bound for the upper Minkowski dimension for all non-invertible self-affine sets.

Lemma 3.2. *If X' is a planar self-affine set, then*

$$\dim_{\mathbb{M}}(X') \leq \dim_{\text{aff}}(\mathbf{A}).$$

Proof. The proof follows by a slight modification of [10, Theorem 5.4] which verifies the upper bound for invertible self-affine sets. \square

The Assouad dimension $\dim_{\mathbb{A}}$, the maximal dimension possible to obtain by looking at coverings, serves as an upper bound for the upper Minkowski dimension. It can happen that the Assouad dimension of a dominated self-affine set is strictly larger than the affinity dimension already in the invertible case; see [3, Theorem 1.4].

It is easy to construct examples of self-affine sets having dimension strictly less than the affinity dimension. For example, several self-affine carpets have this property. Nevertheless, the classical result of Falconer [10, Theorem 5.3] shows that, perhaps rather surprisingly, the Hausdorff dimension of a non-invertible self-affine set equals the affinity dimension for Lebesgue-almost every choice of translation vectors.

Theorem 3.3. *If $X'_{\mathbf{v}}$ is a planar self-affine set and \mathbf{A} satisfies $\max_{i \in J} \|A_i\| < \frac{1}{2}$, then*

$$\dim_{\mathbb{H}}(X'_{\mathbf{v}}) = \min\{2, \dim_{\text{aff}}(\mathbf{A})\}$$

for $\mathcal{L}^{2\#J}$ -almost all translation vectors $\mathbf{v} = (v_i)_{i \in J} \in (\mathbb{R}^2)^{\#J}$.

Originally, Falconer assumed that the matrix norms are bounded above by $\frac{1}{3}$. Solomyak [22] relaxed the bound to $\frac{1}{2}$ which, by the example of Edgar [9], is known to be the best possible.

Recently, it was found a deterministic class of invertible self-affine sets for which the Hausdorff dimension equals the affinity dimension. We say that X satisfies the *open set condition* if there exists an open set $U \subset \mathbb{R}^2$ such that $f_i(U) \cap f_j(U) = \emptyset$ and $f_i(U) \subset U$ for all $i, j \in I$ with $i \neq j$. If such a set U also intersects X , then we say that X satisfies the *strong open set condition*. The following breakthrough result for self-affine sets is proven by Bárány, Hochman, and Rapaport [1, Theorems 1.1 and 7.1]:

Theorem 3.4. *If X is an invertible strictly affine strongly irreducible planar self-affine set satisfying the strong open set condition, then*

$$\begin{aligned} \dim_{\mathbb{H}}(X) &= \min\{2, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\}, \\ \dim_{\mathbb{H}}(\text{proj}_V(X)) &= \min\{1, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\} \end{aligned}$$

for all $V \in \mathbb{RP}^1$.

We emphasize that Theorem 3.4 uses the assumption that the affine iterated function system consists only of invertible maps. It is currently not known whether the result holds also with non-invertible maps. We also remark that Hochman and Rapaport [13] have recently managed to relax the assumptions of the result. They showed that the strong open set condition can be replaced by exponential separation, a separation condition which allows overlapping.

The following three propositions collect our dimension results for non-invertible self-affine sets.

Proposition 3.5. *Suppose that X' and X are non-invertible and invertible planar self-affine sets, respectively. If*

$$\begin{aligned}\dim_{\mathbb{H}}(X) &= \min\{2, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\} \geq 1, \\ \dim_{\mathbb{H}}(\text{proj}_V(X)) &= \min\{1, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\} = 1\end{aligned}$$

for all $V \in \mathbb{RP}^1$, then $\overline{\dim}_{\mathbb{M}}(X') = \dim_{\mathbb{H}}(X)$ and $\dim_{\mathbb{H}}(\text{proj}_V(X')) = 1$ for all $V \in \mathbb{RP}^1$.

Proof. To simplify notation, write $s = \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})$. If $1 < s < \infty$, then Lemma 2.2 shows that $\dim_{\text{aff}}(\mathbf{A}) = s \geq 1$. If $s = 1$, then we get $P(\mathbf{A}, t) = P(I^{\mathbb{N}}, \mathbf{A}, t) < 0 = P(I^{\mathbb{N}}, \mathbf{A}, 1) \leq P(\mathbf{A}, 1)$ for all $1 < t < \infty$ and we again have $\dim_{\text{aff}}(\mathbf{A}) = s \geq 1$. Therefore, by Lemma 3.2, we have $\overline{\dim}_{\mathbb{M}}(X') \leq \min\{2, \dim_{\text{aff}}(\mathbf{A})\} = \min\{2, s\} = \dim_{\mathbb{H}}(X) \leq \dim_{\mathbb{H}}(X')$. To finish the proof, notice that $1 = \dim_{\mathbb{H}}(\text{proj}_V(X)) \leq \dim_{\mathbb{H}}(\text{proj}_V(X')) \leq 1$ for all $V \in \mathbb{RP}^1$. \square

Proposition 3.6. *Suppose that X' and X are non-invertible and invertible planar self-affine sets, respectively. If X' is dominated or irreducible, \mathbf{A} contains a rank one matrix, $\dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A}) < 1$, and*

$$\dim_{\mathbb{H}}(X') = \min\{2, \dim_{\text{aff}}(\mathbf{A})\},$$

then $\dim_{\mathbb{H}}(X') > \overline{\dim}_{\mathbb{M}}(X)$.

Proof. To simplify notation, write $s = \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})$. Since $s < 1$, Lemma 2.9 implies that $0 = P(I^{\mathbb{N}}, \mathbf{A}, s) < P(\mathbf{A}, s)$. Therefore, as Lemmas 2.5 and 2.2 guarantee the continuity of the pressure, we have $s < \dim_{\text{aff}}(\mathbf{A})$. Therefore, by Lemma 3.2, we have $\overline{\dim}_{\mathbb{M}}(X) \leq s < \min\{2, \dim_{\text{aff}}(\mathbf{A})\} = \dim_{\mathbb{H}}(X')$. \square

Proposition 3.7. *Suppose that X' and X are non-invertible and invertible planar self-affine sets, respectively. If \mathbf{A} contains a rank one matrix and*

$$\dim_{\mathbb{H}}(X) = \dim_{\mathbb{H}}(\text{proj}_V(X)) < 1$$

for all $V \in \mathbb{RP}^1$, then there exists a rank one matrix A in \mathbf{A} such that $\dim_{\mathbb{H}}(X') = \dim_{\mathbb{H}}(\text{proj}_{\ker(A)^\perp}(X')) \leq 1$.

Proof. To simplify notation, write $s = \dim_{\mathbb{H}}(X)$. By Lemma 3.1, the non-invertible self-affine set can be expressed as an inhomogeneous self-affine set,

$$X' = X_C = X \cup \bigcup_{i \in I^*} f_i(C),$$

where $C = \bigcup_{i \in J \setminus I} f_i(X')$. Therefore, by the countable stability of Hausdorff dimension,

$$\begin{aligned}\dim_{\mathbb{H}}(X') &= \max\{s, \sup_{i \in I^*} \dim_{\mathbb{H}}(f_i(C))\} \\ &= \max\{s, \dim_{\mathbb{H}}(C)\} = \max\{s, \max_{i \in J \setminus I} \dim_{\mathbb{H}}(A_i(X'))\}.\end{aligned}\tag{3.3}$$

Let A be a rank one matrix in \mathbf{A} such that $\dim_{\mathbb{H}}(A(X')) = \max_{i \in J \setminus I} \dim_{\mathbb{H}}(A_i(X'))$. Since, by the assumption and Lemma 2.1, $s = \dim_{\mathbb{H}}(\text{proj}_{\ker(A)^\perp}(X)) \leq \dim_{\mathbb{H}}(\text{proj}_{\ker(A)^\perp}(X')) = \dim_{\mathbb{H}}(A(X'))$, the claim follows from (3.3). \square

We are now ready to prove the main result. The proof basically just applies Theorems 3.3 and 3.4 in the above propositions.

Proof of Theorem 1.1. (1) Since, by Theorem 3.4, we have

$$\begin{aligned}\dim_{\mathbb{H}}(X) &= \min\{2, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\} \geq 1, \\ \dim_{\mathbb{H}}(\text{proj}_V(X)) &= \min\{1, \dim_{\text{aff}}(I^{\mathbb{N}}, \mathbf{A})\} = 1\end{aligned}$$

for all $V \in \mathbb{R}\mathbb{P}^1$, Proposition 3.5 implies $\overline{\dim}_{\mathbb{M}}(X') = \dim_{\mathbb{H}}(X)$ and $\dim_{\mathbb{H}}(\text{proj}_V(X')) = 1$ for all $V \in \mathbb{R}\mathbb{P}^1$.

(2) Since, by Theorem 3.3, we have

$$\dim_{\mathbb{H}}(X'_v) = \min\{2, \dim_{\text{aff}}(\mathbf{A})\}$$

for $\mathcal{L}^{2\#J}$ -almost all $v \in (\mathbb{R}^2)^{\#J}$, Proposition 3.6 implies $\dim_{\mathbb{H}}(X'_v) > \overline{\dim}_{\mathbb{M}}(X_v)$ for $\mathcal{L}^{2\#J}$ -almost all $v \in (\mathbb{R}^2)^{\#J}$.

(3) Since, by Theorem 3.4, we have

$$\dim_{\mathbb{H}}(X) = \dim_{\mathbb{H}}(\text{proj}_V(X)) < 1$$

for all $V \in \mathbb{R}\mathbb{P}^1$, Proposition 3.7 implies that there exists a rank matrix A in \mathbf{A} such that $\dim_{\mathbb{H}}(X') = \dim_{\mathbb{H}}(\text{proj}_{\ker(A)^\perp}(X')) \leq 1$. \square

REFERENCES

- [1] B. Bárány, M. Hochman, and A. Rapaport. Hausdorff dimension of planar self-affine sets and measures. *Invent. Math.*, 216(3):601–659, 2019.
- [2] B. Bárány, A. Käenmäki, and I. D. Morris. Domination, almost additivity, and thermodynamic formalism for planar matrix cocycles. *Israel J. Math.*, 239(1):173–214, 2020.
- [3] B. Bárány, A. Käenmäki, and H. Yu. Finer geometry of planar self-affine sets. Preprint, available at arXiv:2107.00983, 2021.
- [4] M. F. Barnsley. *Superfractals*. Cambridge University Press, Cambridge, 2006.
- [5] M. F. Barnsley and S. Demko. Iterated function systems and the global construction of fractals. *Proc. Roy. Soc. London Ser. A*, 399(1817):243–275, 1985.
- [6] J. Bochi and I. D. Morris. Continuity properties of the lower spectral radius. *Proc. Lond. Math. Soc.* (3), 110(2):477–509, 2015.
- [7] J. Bochi and I. D. Morris. Equilibrium states of generalised singular value potentials and applications to affine iterated function systems. *Geom. Funct. Anal.*, 28(4):995–1028, 2018.
- [8] R. Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, volume 470 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, revised edition, 2008. With a preface by David Ruelle, Edited by Jean-René Chazottes.
- [9] G. A. Edgar. Fractal dimension of self-affine sets: some examples. Number 28, pages 341–358. 1992. Measure theory (Oberwolfach, 1990).
- [10] K. J. Falconer. The Hausdorff dimension of self-affine fractals. *Math. Proc. Cambridge Philos. Soc.*, 103(2):339–350, 1988.
- [11] D.-J. Feng and A. Käenmäki. Equilibrium states of the pressure function for products of matrices. *Discrete Contin. Dyn. Syst.*, 30(3):699–708, 2011.
- [12] D.-J. Feng and P. Shmerkin. Non-conformal repellers and the continuity of pressure for matrix cocycles. *Geom. Funct. Anal.*, 24(4):1101–1128, 2014.
- [13] M. Hochman and A. Rapaport. Hausdorff dimension of planar self-affine sets and measures with overlaps. *J. Eur. Math. Soc. (JEMS)*, 24(7):2361–2441, 2022.
- [14] J. E. Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.*, 30(5):713–747, 1981.
- [15] R. Jungers. *The joint spectral radius*, volume 385 of *Lecture Notes in Control and Information Sciences*. Springer-Verlag, Berlin, 2009. Theory and applications.
- [16] A. Käenmäki. On natural invariant measures on generalised iterated function systems. *Ann. Acad. Sci. Fenn. Math.*, 29(2):419–458, 2004.

- [17] A. Käenmäki and I. D. Morris. Structure of equilibrium states on self-affine sets and strict monotonicity of affinity dimension. *Proc. Lond. Math. Soc. (3)*, 116(4):929–956, 2018.
- [18] A. Käenmäki and M. Vilppolainen. Dimension and measures on sub-self-affine sets. *Monatsh. Math.*, 161(3):271–293, 2010.
- [19] J. M. Marstrand. Some fundamental geometrical properties of plane sets of fractional dimensions. *Proc. London Math. Soc. (3)*, 4:257–302, 1954.
- [20] P. Mattila. *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*. Cambridge University Press, Cambridge, 1995.
- [21] N. Snigireva. Inhomogeneous self-similar sets and measures. PhD Dissertation, University of St Andrews, 2008.
- [22] B. Solomyak. Measure and dimension for some fractal families. *Math. Proc. Cambridge Philos. Soc.*, 124(3):531–546, 1998.

(Antti Käenmäki) RESEARCH UNIT OF MATHEMATICAL SCIENCES, P.O. BOX 8000, FI-90014 UNIVERSITY OF OULU, FINLAND

Email address: `antti.kaenmaki@oulu.fi`

(Petteri Nissinen) DEPARTMENT OF PHYSICS AND MATHEMATICS, UNIVERSITY OF EASTERN FINLAND, P.O. BOX 111, FI-80101 JOENSUU, FINLAND

Email address: `pettern@student.uef.fi`