

STRONG \mathbb{A}^1 -INVARIANCE OF \mathbb{A}^1 -CONNECTED COMPONENTS OF REDUCTIVE ALGEBRAIC GROUPS

CHETAN BALWE, AMIT HOGADI, AND ANAND SAWANT

ABSTRACT. We show that the sheaf of \mathbb{A}^1 -connected components of a reductive algebraic group over a perfect field is strongly \mathbb{A}^1 -invariant. As a consequence, torsors under such groups give rise to \mathbb{A}^1 -fiber sequences. We also show that sections of \mathbb{A}^1 -connected components of anisotropic, semisimple, simply connected algebraic groups over an arbitrary field agree with their R -equivalence classes, thereby removing the perfectness assumption in the previously known results about the characterization of isotropy in terms of affine homotopy invariance of Nisnevich locally trivial torsors.

CONTENTS

| | |
|--|----|
| 1. Introduction..... | 1 |
| 2. Birationality and Strong \mathbb{A}^1 -invariance..... | 3 |
| 3. \mathbb{A}^1 -connected components of reductive groups and their classifying spaces..... | 5 |
| 4. Semisimple, simply connected algebraic groups over a perfect field..... | 9 |
| 5. Strong \mathbb{A}^1 -invariance of the sheaf of \mathbb{A}^1 -connected components of a reductive group | 11 |
| References..... | 13 |

1. INTRODUCTION

Principal bundles (or in other words, torsors) under a group are among the most typical examples of fibrations in topology. The concept of a fibration is extremely useful, since a fibration gives rise to a long exact sequence consisting of homotopy groups of the total space, the base space and the fiber. Motivic homotopy theory developed by Morel and Voevodsky [26] provides a framework to import techniques from topology into algebraic geometry. Computations using motivic fiber sequences have been successfully used in the works of Asok and Fasel [3, 4] to address classical questions about splitting properties of vector bundles on smooth affine varieties using the \mathbb{A}^1 -homotopical interpretation of these questions. Unlike in classical topology, torsors under sheaves of groups do not automatically give rise to fiber sequences in \mathbb{A}^1 -homotopy theory, the reason behind which can be attributed to the peculiar behaviour of the sheaf of \mathbb{A}^1 -connected components. The set of connected components of a space in classical topology is a discrete space. The analogous statement in \mathbb{A}^1 -homotopy theory states that the sheaf of \mathbb{A}^1 -connected components of a space is \mathbb{A}^1 -invariant. This statement, conjectured to be true by Morel [25], is not known in full generality.

The authors acknowledge the support of India DST-DFG Project on Motivic Algebraic Topology DST/IBCD/GERMANY/DFG/2021/1, SERB Start-up Research Grant SRG/2020/000237 and the Department of Atomic Energy, Government of India, under project no. 12-R&D-TFR-5.01-0500.

Let us fix a base field k and denote by $\mathcal{H}(k)$ the unstable motivic homotopy category over k , the objects of which are simplicial Nisnevich sheaves of sets on the category Sm_k of smooth schemes over k . We refer the reader to Sections 2 and 3 for relevant definitions not mentioned in the introduction. Let G be a Nisnevich sheaf of groups on Sm_k and let $\mathcal{P} \rightarrow \mathcal{X}$ be a Nisnevich locally trivial G -torsor in $\mathcal{H}(k)$. This means that \mathcal{P} has a right G -action, the morphism $\mathcal{P} \rightarrow \mathcal{X}$ is equivariant for the trivial right G -action on \mathcal{X} and there is an isomorphism $G \times \mathcal{P} \xrightarrow{\sim} \mathcal{P} \times_{\mathcal{X}} \mathcal{P}$ of simplicial Nisnevich sheaves. By a result of Choudhury [12], it is known that $\pi_0^{\mathbb{A}^1}(G)$ is \mathbb{A}^1 -invariant. Morel showed in [25, Theorem 6.50] that if the sheaf $\pi_0^{\mathbb{A}^1}(G)$ of \mathbb{A}^1 -connected components of G is strongly \mathbb{A}^1 -invariant, then a G -torsor $\mathcal{P} \rightarrow \mathcal{X}$ yields an \mathbb{A}^1 -fiber sequence

$$\mathcal{P} \rightarrow \mathcal{X} \rightarrow BG.$$

We will be interested in the case when G is represented by an algebraic group over k . Strong \mathbb{A}^1 -invariance of $\pi_0^{\mathbb{A}^1}(G)$ is known when G is isotropic, semisimple and simply connected over a field due to the work of Asok, Hoyois and Wendt [5, 6]. The main result of this paper is to generalize this result to all reductive algebraic groups, at least when the base field is perfect.

Theorem 1.1 (Theorem 5.2 and Corollary 5.3). *If G is a reductive algebraic group over a perfect field k , then $\pi_0^{\mathbb{A}^1}(G)$ is strongly \mathbb{A}^1 -invariant. Consequently, for any Nisnevich locally trivial G -torsor $\mathcal{P} \rightarrow \mathcal{X}$ in $\mathcal{H}(k)$,*

$$\mathcal{P} \rightarrow \mathcal{X} \rightarrow BG$$

is an \mathbb{A}^1 -fiber sequence.

Remark 1.2. It is easy to see that the conclusion of Theorem 1.1 holds for any algebraic group over a perfect field. Indeed, strong \mathbb{A}^1 -invariance of $\pi_0^{\mathbb{A}^1}(G)$ for an arbitrary algebraic group G over a perfect field reduces to the case of affine linear algebraic group by Chevalley's structure theorem (see [31, Section 5.2]). One can reduce to the case of reductive groups by observing that the unipotent radical is \mathbb{A}^1 -contractible, being isomorphic to an affine space.

There are two main ingredients in the proof of Theorem 1.1. The first one states that if G as in Theorem 1.1 is moreover semisimple and simply connected, then $\pi_0^{\mathbb{A}^1}(G)$ is a birational sheaf (see Theorem 4.5). The perfectness assumption on k is needed here since our arguments require the study of base change of G to the coefficient field of a smooth henselian local ring, for which we require the coefficient field to contain a copy of k . Note that even in the case of isotropic, semisimple, simply connected and absolutely almost simple G , Theorem 4.5 provides a refinement of the result in [5, 6] about the strong \mathbb{A}^1 -invariance of $\pi_0^{\mathbb{A}^1}(G)$. The second key ingredient in the proof of Theorem 1.1 is the preservation of strong \mathbb{A}^1 -invariance under isogenies of simply connected groups (see Lemma 5.1). This step relies crucially on a tricky result of Choudhury and Hogadi [13, Theorem 1.3], which states that \mathbb{A}^1 -invariant quotients of strongly \mathbb{A}^1 -invariant sheaves are strongly \mathbb{A}^1 -invariant. Another key input in this step is the \mathbb{A}^1 -homotopical interpretation of classical results on torsors on the affine line in terms of \mathbb{A}^1 -connected components of classifying spaces.

We believe that it should be possible to prove Theorem 1.1 over an arbitrary field. The second main result of this paper (Theorem 3.2) concerns the comparison of R -equivalence classes and sections of $\pi_0^{\mathbb{A}^1}(G)$ over arbitrary fields, generalizing [8, Main Theorem]. This leads to the anisotropic analogue of the reinterpretation of [19, Théorème 5.8, Théorème 7.2] in [5, Corollary 4.3.6].

Theorem 1.3 (Theorem 3.2, Corollary 3.3 and Corollary 3.7). *Let k be an arbitrary field and let G be a semisimple, simply connected algebraic group over k . Then the natural map*

$$\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow G(k)/R$$

is a bijection. Moreover, the contraction $\pi_0^{\mathbb{A}^1}(G)_{-1}$ is the trivial point sheaf.

Theorem 1.3 vindicates the principle that for semisimple, simply connected groups, \mathbb{A}^1 -connected components provide the right framework as an anisotropic analogue of the Whitehead group of an isotropic group and can be used to put results about isotropic and anisotropic groups on the same footing.

Acknowledgements. We thank Matthias Wendt for helpful correspondence regarding the results of [16] and Swarnava Mukhopadhyay for pointing us to the literature on principal bundles on curves.

Notations and conventions. The letter k will always denote a field. We will denote by Sm_k the big Nisnevich site of finite type, separated schemes over k . Given a simplicial sheaf of sets \mathcal{X} on Sm_k and an affine scheme $\text{Spec } A$ over k , we will often write $\mathcal{X}(A)$ for $\mathcal{X}(\text{Spec } A)$ for the sake of brevity.

We will often make use of essentially smooth schemes, that is, schemes which are filtered inverse limits of diagrams of smooth schemes in which the transition maps are étale, affine morphisms. All presheaves on Sm_k will be extended to essentially smooth schemes by defining $\mathcal{F}(\varinjlim U_\alpha) = \varinjlim \mathcal{F}(U_\alpha)$. By a henselian local scheme over k , we mean an affine scheme of the form $U = \text{Spec}(R)$, where R is a henselian local k -algebra.

2. BIRATIONALITY AND STRONG \mathbb{A}^1 -INVARIANCE

In this section, we recall definitions relevant to birationality and strong \mathbb{A}^1 -invariance of sheaves on Sm_k and basic properties that will be used in the proofs of our main results.

Definition 2.1. Let \mathcal{F} be a presheaf of sets on Sm_k .

- We say that \mathcal{F} is \mathbb{A}^1 -invariant if for every $U \in Sm_k$, the map $\mathcal{F}(U) \rightarrow \mathcal{F}(U \times \mathbb{A}^1)$ induced by the projection $U \times \mathbb{A}^1 \rightarrow U$, is a bijection.
- We say that \mathcal{F} is *strongly* \mathbb{A}^1 -invariant if for every $U \in Sm_k$, the map $H_{\text{Nis}}^i(U, \mathcal{F}) \rightarrow H_{\text{Nis}}^i(U \times \mathbb{A}^1, \mathcal{F})$ induced by the projection $U \times \mathbb{A}^1 \rightarrow U$, is a bijection for $i = 0, 1$.

Definition 2.2. A presheaf \mathcal{F} of sets on Sm_k is said to be *birational* if

- (1) for every $X \in Sm_k$ with irreducible components X_1, \dots, X_n , the map

$$\mathcal{F}(X) \rightarrow \prod_{1 \leq i \leq n} \mathcal{F}(X_i)$$

is a bijection; and

- (2) for every $X \in Sm_k$ and a dense open subscheme U of X , the restriction map $\mathcal{F}(X) \rightarrow \mathcal{F}(U)$ is a bijection.

A sheaf of sets is said to be birational if its underlying presheaf is birational.

Remark 2.3. Let \mathcal{F} be a birational presheaf of (not necessarily abelian) groups on Sm_k . Then \mathcal{F} is already a Nisnevich sheaf by [7, Lemma 6.1.2]. Note that \mathcal{F} is flasque by definition. We claim that the Nisnevich cohomology $H_{\text{Nis}}^1(X, \mathcal{F})$ vanishes for all $X \in Sm_k$. Take an element of $H_{\text{Nis}}^1(X, \mathcal{F})$, which is represented by a Nisnevich locally trivial \mathcal{F} -torsor P on X . Let U be a maximal open subset of X on which this torsor is trivial. Suppose, if possible, that $U \neq X$. Let η be a generic point of $X \setminus U$. There exists an étale map $V \rightarrow X$ such that $U \amalg V$ is an elementary Nisnevich cover of an open subset U' of X containing U and η and such that the torsor P is trivial on V . Since $\mathcal{F}(V) \rightarrow \mathcal{F}(U \times_{U'} V)$ is surjective, a straightforward Čech cohomology calculation shows that P is trivial on U' , contradicting the choice of U . Hence, $H_{\text{Nis}}^1(X, \mathcal{F}) = 0$, for every $X \in Sm_k$. Thus, we conclude that if \mathcal{F} is a birational and \mathbb{A}^1 -invariant sheaf of groups, then it is strongly \mathbb{A}^1 -invariant.

Birationality of a Nisnevich sheaf of groups on Sm_k can be checked by verifying the condition given in Definition 2.2 on smooth henselian schemes.

Lemma 2.4. *Let \mathcal{F} be an \mathbb{A}^1 -invariant sheaf of groups on Sm_k . Suppose that for every smooth henselian local k -algebra R with quotient field K , the restriction map $\mathcal{F}(R) \rightarrow \mathcal{F}(K)$ is an isomorphism. Then \mathcal{F} is a birational sheaf.*

Proof. Let $X \in Sm_k$ and let U be a dense open subset of X . By [25, Lemma 3.45], we know that the restriction map $\mathcal{F}(X) \rightarrow \mathcal{F}(U)$ is injective. We now show that it is surjective.

Let $\alpha \in \mathcal{F}(U)$. For every point $x \in X$, we have a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(X) & \longrightarrow & \mathcal{F}(U) \\ \downarrow & & \downarrow \\ \mathcal{F}(\mathcal{O}_{X,x}^h) & \longrightarrow & \mathcal{F}(K_x^h) \end{array}$$

in which K_x^h denotes the quotient field of henselization $\mathcal{O}_{X,x}^h$ of the local ring $\mathcal{O}_{X,x}$. Since the bottom horizontal arrow is an isomorphism by hypothesis, we can lift $\alpha \in \mathcal{F}(U)$ to an element of $\mathcal{F}(\mathcal{O}_{X,x}^h)$, which by a standard limiting argument lifts to an element $\alpha_x \in \mathcal{F}(V_x)$ for some Nisnevich neighborhood V_x of x . Since X is noetherian, we can find a refinement $\tilde{X} \rightarrow X$ of the Nisnevich cover $\coprod_{x \in X} V_x \rightarrow X$ with $\tilde{X} \in Sm_k$ such that $\alpha \in \mathcal{F}(U)$ lifts to an element $\tilde{\alpha} \in \mathcal{F}(\tilde{X})$. We now show that $\tilde{\alpha}$ descends to an element of $\mathcal{F}(X)$ which restricts to α . Let \tilde{U} denote the inverse image of U under the map $\tilde{X} \rightarrow X$. We now have a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}(X) & \longrightarrow & \mathcal{F}(\tilde{X}) & \rightrightarrows & \mathcal{F}(\tilde{X} \times_X \tilde{X}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{F}(U) & \longrightarrow & \mathcal{F}(\tilde{U}) & \rightrightarrows & \mathcal{F}(\tilde{U} \times_U \tilde{U}) \end{array}$$

in which the rows are equalizer diagrams and all the vertical maps are injective. It remains to observe that the image of $\tilde{\alpha}$ under the two maps $\mathcal{F}(\tilde{X}) \rightrightarrows \mathcal{F}(\tilde{X} \times_X \tilde{X})$ is the same. But this follows from the injectivity of the rightmost vertical map and the fact that the restriction of $\tilde{\alpha}$ to \tilde{U} is the image of $\alpha \in \mathcal{F}(U)$. This shows that $\mathcal{F}(X) \rightarrow \mathcal{F}(U)$ is surjective, completing the proof. \square

Since the sheaf of \mathbb{A}^1 -connected components of any H -group is \mathbb{A}^1 -invariant by the results of [12], the restriction maps on such sheaves happen to be injective. This observation will prove useful in our analysis of \mathbb{A}^1 -connected components of a reductive algebraic group.

Lemma 2.5. *Let \mathcal{G} be a sheaf of groups on Sm_k . Then for any $X \in Sm_k$ and a dense open subset U of X , the restriction map $\pi_0^{\mathbb{A}^1}(\mathcal{G})(X) \rightarrow \pi_0^{\mathbb{A}^1}(\mathcal{G})(U)$ is injective. In particular, the induced map $\pi_0^{\mathbb{A}^1}(\mathcal{G})(X) \rightarrow \pi_0^{\mathbb{A}^1}(\mathcal{G})(k(X))$ is injective.*

Proof. This is a straightforward consequence of [12, Theorem 4.18] and [25, Lemma 3.45]. \square

3. \mathbb{A}^1 -CONNECTED COMPONENTS OF REDUCTIVE GROUPS AND THEIR CLASSIFYING SPACES

One of our main interests behind the study of the sheaf of \mathbb{A}^1 -connected components of a reductive group G over a field is its application to \mathbb{A}^1 -fiber sequences due to Morel [25, Theorem 6.50] (see also [2, Theorem 3.5]). We refer the reader to [25, Chapter 6] or [2, Section 2] for generalities on \mathbb{A}^1 -fiber sequences. In this section, we study the sheaf of \mathbb{A}^1 -connected components of reductive algebraic groups and their classifying spaces.

3.1. Reductive groups.

We refer the reader to [24] and [15, Appendix A] for basic facts from algebraic group theory. A semisimple, simply connected algebraic group G over a field k is called *isotropic* if every almost simple factor of G contains a non-central k -subgroup-scheme isomorphic to \mathbb{G}_m . A semisimple, simply connected algebraic group is *anisotropic* if it contains no subgroup isomorphic to \mathbb{G}_m .

Let G be a semisimple, simply connected and absolutely almost simple algebraic group over an arbitrary field k . The *Whitehead group* of G over k is defined to be

$$W(k, G) := G(k)/G(k)^+,$$

where $G(k)^+$ is the normal subgroup of $G(k)$ generated by k -rational points of subgroups of G isomorphic to \mathbb{G}_a . A related invariant of G is the group $G(k)/R$ of *R-equivalence* classes, defined to be the quotient of $G(k)$ by the normal subgroup consisting of elements $g \in G(k)$ for which there exists a rational map $h : \mathbb{P}^1 \dashrightarrow G$ defined at 0 and 1 such that $h(0) = e$ and $h(1) = g$, where e denotes the neutral element of $G(k)$. See [19] for a detailed discussion of Whitehead groups and *R-equivalence* classes. As a consequence of affine homotopy invariance of Nisnevich locally trivial G -torsors for isotropic groups G (see [5, Theorem 3.3.7] for the statement over infinite fields and [6, Theorem 2.4] for the statement over finite fields), Asok, Hoyois and Wendt have shown that the Morel-Voevodsky singular construction $\text{Sing}_*^{\mathbb{A}^1} G$ is \mathbb{A}^1 -local (see [5, Remark 3.3.8]). Thus, if G is in addition isotropic, the natural map $W(k, G) \rightarrow G(k)/R$ factors through $\pi_0^{\mathbb{A}^1}(G)(k)$ and induces bijections

$$(3.1) \quad W(k, G) \xrightarrow{\simeq} \pi_0^{\mathbb{A}^1}(G)(k) \xrightarrow{\simeq} G(k)/R,$$

by [19, Théorème 7.2].

In the case G is anisotropic, the group $G(k)^+$ is trivial. However, it was shown in [8] (see also [9, Theorem 3.6]) that over an infinite perfect field k , there is a bijection

$$\pi_0^{\mathbb{A}^1}(G)(k) \xrightarrow{\simeq} G(k)/R.$$

We first observe that the proof of this result given in [8] can be appropriately modified to make it work over an arbitrary base field. There are two key ingredients in the proof - the first is the existence of compactifications of algebraic groups over an arbitrary field due to Gabber (see [18, Théorème 5.2] and also [20, Theorem 4, Theorem 6]), which yields the following consequence.

Lemma 3.1. *Let G be an anisotropic semisimple group over an arbitrary field k . Then any rational map $h : \mathbb{P}_k^1 \dashrightarrow G$ is defined at all the k -rational points of \mathbb{P}_k^1 .*

Proof. Follow exactly the proof of [8, Lemma 3.6] with [11, 8.2] replaced by [18, Théorème 5.2] (or [20, Theorem 4, Theorem 6]). \square

The second key ingredient is the choice of a Nisnevich cover of the affine line over which an R -equivalence between two k -points can be completed to give an \mathbb{A}^1 -ghost homotopy connecting the two points. This is where the proof of Theorem 3.2 differs from the one given in [8].

Theorem 3.2. *Let k be an arbitrary field and let G be a semisimple, simply connected, absolutely almost simple and anisotropic group over k . Then the natural map*

$$\pi_0^{\mathbb{A}^1}(G)(F) \rightarrow G(F)/R$$

is a bijection, for every field extension F of k .

Proof. We closely follow the proof of [8, Theorem 4.2]. We are reduced to the case $F = k$ as in [8]. By [11, 8.2] (in characteristic 0) or [18, Théorème 5.2] (in arbitrary characteristic), there exists a compactification \overline{G} of G such that $G(k) = \overline{G}(k)$. If two elements p and q of $G(k)$ map to the same element in $\pi_0^{\mathbb{A}^1}(G)(k)$, then they map to the same element in $\pi_0^{\mathbb{A}^1}(\overline{G})(k)$ by functoriality. Since \overline{G} is proper over k , we see that p and q map to the same element in $\mathcal{S}(\overline{G})(k)$ by [7, Theorem 2.4.3]. Therefore, p and q are \mathbb{A}^1 -chain homotopic k -rational points of \overline{G} and hence, R -equivalent in $G(k) = \overline{G}(k)$. In other words, the natural surjective map $G(k) \rightarrow G(k)/R$ factors through $\pi_0^{\mathbb{A}^1}(G)(k)$.

We now show the injectivity of the map $\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow G(k)/R$. Let p and q be two elements of $G(k)$, which are R -equivalent through a rational map $h : \mathbb{P}_k^1 \dashrightarrow G$ defined on 0 and 1 such that $h(0) = p$ and $h(1) = q$. Choose a compactification \overline{G} of G such that $G(k) = \overline{G}(k)$ and uniquely extend the rational map h to a morphism $\overline{h} : \mathbb{P}_k^1 \rightarrow \overline{G}$. Now, h is not defined only at points of \mathbb{A}_k^1 having residue fields that are non-trivial finite extensions of k . We define $V := \overline{h}^{-1}(G) \cap \mathbb{A}_k^1$ which is a Zariski open subscheme of \mathbb{A}_k^1 . Let $\mathbb{A}_k^1 \setminus V = \{p_1, \dots, p_n\}$ and let the residue field at p_i be L_i . We define $h_V : V \rightarrow G$ by $h_V := \overline{h}|_V$. As observed in [8, Proof of Theorem 4.2], G_{L_i} is an isotropic group for each i .

For each i , let K_i denote the field of fractions of the henselization $\mathcal{O}_{\mathbb{A}_k^1, p_i}^h$ of the local ring of p_i at \mathbb{A}_k^1 . We set $W_i := \text{Spec} \left(\mathcal{O}_{\mathbb{A}_k^1, p_i}^h \right)$. Note that K_i is a separable extension of $k(t)$. Since G_{L_i} is isotropic, so is G_{K_i} by [14, Proposition A.2]. By [19, Lemme 4.5], we have

$$G(K_i) = G(K_i)^+ \cdot G(\mathcal{O}_{\mathbb{A}_k^1, p_i}^h),$$

so any morphism $\text{Spec}(K_i) \rightarrow G$ is \mathbb{A}^1 -chain homotopic to a morphism that factors as $\text{Spec}(K_i) \xrightarrow{\eta_i} W_i \rightarrow G$, where $\eta_i : \text{Spec}(K_i) \rightarrow W_i$ denotes the generic point of W_i . Now,

consider the pullback $h_V|_{W_i} : V \times_{\mathbb{A}_k^1} W_i \rightarrow G$ of h_V and denote its restriction to $\mathrm{Spec}(K_i)$ by h_i . By the above observation, $h_i : \mathrm{Spec}(K_i) \rightarrow G$ is \mathbb{A}^1 -chain homotopic to a morphism that factors as

$$\mathrm{Spec}(K_i) \xrightarrow{\eta_i} W_i \xrightarrow{\rho_i} G,$$

for some $\rho_i \in G(\mathcal{O}_{\mathbb{A}_k^1, p_i}^h)$. Define $W := \coprod_i W_i$ and $h_W := \coprod_i \rho_i : W \rightarrow G$.

We define $p_V : V \rightarrow \mathbb{A}_k^1$ to be the inclusion. For each i , let $p_i : W_i \rightarrow \mathbb{A}_k^1$ denote the natural map induced by $\mathrm{Spec}(\mathcal{O}_{\mathbb{A}_k^1, p_i}^h) \rightarrow \mathbb{A}_k^1$ and define $p_W := \coprod_i p_i : W \rightarrow \mathbb{A}_k^1$. By a standard limiting argument, we may replace W_i with a scheme that is finite étale over \mathbb{A}_k^1 . We may further shrink W_i suitably to ensure that

- W_i is an étale neighbourhood of p_i ;
- $\{p_V, p_W\}$ is an elementary Nisnevich cover of \mathbb{A}_k^1 ; and
- the \mathbb{A}^1 -chain homotopy between h_i and $\rho_i \circ \eta_i$ spreads over W_i .

We then modify W and h_W accordingly. We now apply [8, Lemma 2.4] to conclude that p and q map to the same element in $\pi_0^{\mathbb{A}^1}(G)(k)$. \square

Combining Theorem 3.2 with the results of Asok-Hoyois-Wendt and Gille, we get the following generalizations of [9, Theorem 3.6] and [9, Theorem 1], which we record below.

Corollary 3.3. *Let k be an arbitrary field and let G be a semisimple, simply connected algebraic group over k . Then the natural map $\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow G(k)/R$ is a bijection.*

Proof. As in [9, Proof of Theorem 3.6], we are reduced to showing that $\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow G(k)/R$ is a bijection, where G is semisimple, simply connected and absolutely almost simple over k . If G is isotropic, then the desired result follows from (3.1). If G is anisotropic, then we apply Theorem 3.2. \square

Corollary 3.4. *Let G be a reductive algebraic group over an arbitrary field k . Then the following conditions are equivalent:*

- (1) $\mathrm{Sing}_*^{\mathbb{A}^1} G$ is \mathbb{A}^1 -local;
- (2) G is isotropic;
- (3) the presheaf $H_{\mathrm{Nis}}^1(-, G)$ is \mathbb{A}^1 -invariant on smooth affine schemes over k .

Proof. This follows exactly as in [9, Proof of Theorem 1]. \square

3.2. Contractions of \mathbb{A}^1 -connected components of reductive groups.

In this subsection, we record the generalizations of [19, Théorème 5.8, Théorème 7.2] and [5, Corollary 4.3.6] to anisotropic and hence, to all semisimple, simply connected groups over any field.

Proposition 3.5. *Let G be a semisimple, simply connected algebraic group over a field k . Then the natural map*

$$\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow \pi_0^{\mathbb{A}^1}(G)(k(t))$$

is an isomorphism.

Proof. Clearly, it suffices to prove the statement in the case where G is absolutely almost simple. By Theorem 3.2, we see that the natural map

$$\pi_0^{\mathbb{A}^1}(G)(F) \rightarrow G(F)/R$$

is an isomorphism, for every field extension F of k . The proof in the case G is isotropic follows from [19, Théorème 5.8, Théorème 7.2]. Suppose that G is anisotropic. Note that the map $\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow \pi_0^{\mathbb{A}^1}(G)(k(t))$ is injective. For surjectivity, let $\alpha \in \pi_0^{\mathbb{A}^1}(G)(k(t)) = G(k(t))/R$. Now, α corresponds to a rational map $f_\alpha : \mathbb{A}^1 \dashrightarrow G$. Precomposing by the multiplication map $\mathbb{A}^1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^1$, we get a rational map

$$h : \mathbb{A}^1 \times \mathbb{A}^1 \dashrightarrow G; \quad h(s, t) := f_\alpha(st).$$

Let us denote by h_s the rational map $\mathbb{A}^1 \dashrightarrow G$ given by $t \mapsto h(s, t)$. Then h_0 and h_1 are R -equivalent as elements of $G(k(t))$ and h_0 lies in the image of the map $\pi_0^{\mathbb{A}^1}(G)(k) \rightarrow \pi_0^{\mathbb{A}^1}(G)(k(t))$. This completes the proof. \square

Remark 3.6. It is worthwhile to remark that the results [19, Théorème 5.8, Théorème 7.2] that play a crucial role in the proof of isotropic case of Proposition 3.5 also play a crucial role in the proof of the anisotropic case of Theorem 3.2.

Corollary 3.7. *If G is a semisimple, simply connected algebraic group over a field k , then $\pi_0^{\mathbb{A}^1}(G)_{-1} = 0$.*

Proof. We reduce to the case that G is absolutely almost simple as above. The result in the case where G is isotropic can be found in [5, Corollary 4.3.6]. The proof in the anisotropic case is immediate from Proposition 3.5, since we have

$$\pi_0^{\mathbb{A}^1}(G)_{-1}(U) = \ker \left(\pi_0^{\mathbb{A}^1}(G)(U \times \mathbb{G}_m) \xrightarrow{(id, 1)^*} \pi_0^{\mathbb{A}^1}(G)(U) \right),$$

for any smooth k -scheme U by definition. \square

3.3. Classifying spaces of reductive groups.

In this section, we briefly review the results of Elmanto, Kulkarni and Wendt [16] that will be used in Section 5. Let Sch_k denote the site of schemes of finite type over k with étale topology. Let $\mathbf{R}_{\text{ét}}$ denote the fibrant replacement functor for the Čech étale-local injective model structure. The inclusion functor $i : Sm_k \rightarrow Sch_k$ induces a restriction functor i^* from the category of simplicial étale sheaves on Sch_k to the category of simplicial Nisnevich sheaves on Sm_k .

Let G be a reductive algebraic group over a field k . We will denote by BG the pointed simplicial sheaf whose n -simplices are G^n with usual face and degeneracy maps. Let G also denote the underlying étale sheaf of groups on Sch_k . Then we define $B_{\text{ét}}G$ to be the simplicial Nisnevich sheaf defined by

$$B_{\text{ét}}G := i^* \circ \mathbf{R}_{\text{ét}}(BG).$$

The sheaf $\mathcal{H}_{\text{ét}}^1(G)$ is defined to be the Nisnevich sheafification of the presheaf $U \mapsto H_{\text{ét}}^1(U, G)$ on Sm_k . We will freely use the identification $\mathcal{H}_{\text{ét}}^1(G) = \pi_0^{\mathbb{A}^1}(B_{\text{ét}}G)$ obtained in [16, Theorem 1.5].

We now discuss a variant of a result due to Elmanto, Kulkarni and Wendt [16, Theorem 1.2, Proposition 3.7], which will play a key role in the proof of our main theorem. We note

below that under the assumption of simple connectedness, [16, Proposition 3.7] holds over an arbitrary field.

Theorem 3.8. *Let G be a semisimple, simply connected algebraic group over a field k . Then the sheaf $\mathcal{H}_{\text{ét}}^1(G)$ is \mathbb{A}^1 -invariant.*

Proof. The proof of [16, Proposition 3.7] works almost verbatim (with the use of [10] in place of [29]), so we only give an outline of the argument for the convenience of readers.

By the strong Grothendieck-Serre property over a field (see [17, Corollary 1] and [27, Corollary 1.2]) along with [16, Proposition 3.4], we are reduced to showing that the map $H_{\text{ét}}^1(F, G) \rightarrow H_{\text{ét}}^1(\mathbb{A}_F^1, G)$ induced by the projection map $\mathbb{A}_F^1 \rightarrow \text{Spec}(F)$ is bijective for any field extension F of k . Fix a separable closure F_{sep} of F . By [28, Theorem 1.1] (restated as [16, Theorem 3.6]), there is a bijection

$$H_{\text{ét}}^1(F, G) \rightarrow \ker \left(H_{\text{ét}}^1(\mathbb{A}_F^1, G) \rightarrow H_{\text{ét}}^1(\mathbb{A}_{F_{\text{sep}}}^1, G) \right).$$

It follows from [22, Satz 3.3] that generically trivial G -torsors on smooth affine curves are trivial. We now apply [10, 8.6] to conclude that $H_{\text{ét}}^1(\text{Spec}(F_{\text{sep}}(t)), G) = 0$, and consequently, every G -torsor on $\mathbb{A}_{F_{\text{sep}}}^1$ is trivial. \square

4. SEMISIMPLE, SIMPLY CONNECTED ALGEBRAIC GROUPS OVER A PERFECT FIELD

The aim of this section is to show that the sheaf of \mathbb{A}^1 -connected components of a semisimple, simply connected algebraic group over a perfect field is birational. The relationship between the sections of the sheaf of \mathbb{A}^1 -connected components over a discrete valuation ring and its quotient field is tantamount to the understanding of the birationality property. We treat the cases of isotropic and anisotropic groups separately below.

Lemma 4.1. *Let G be an isotropic, semisimple, simply connected, absolutely almost simple algebraic group over a field k . Let R be a discrete valuation ring with quotient field K and residue field k such that R is a k -algebra. Then the restriction map $\pi_0^{\mathbb{A}^1}(G)(R) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K)$ is an isomorphism.*

Proof. Let R^h denote the henselization of R . Let K^h denote the quotient field of R^h . By [19, Lemme 4.5],

$$G(K^h) = G(K^h)^+ \cdot G(R^h),$$

where $G(K^h)^+$ denotes the subgroup of $G(K^h)$ generated by unipotent elements. This implies that the natural map $\pi_0^{\mathbb{A}^1}(G)(R^h) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K^h)$ is an isomorphism. We now have a diagram

$$\begin{array}{ccccc} \pi_0^{\mathbb{A}^1}(G)(R) & \longrightarrow & \pi_0^{\mathbb{A}^1}(G)(R^h) & \xrightarrow{\cong} & \pi_0^{\mathbb{A}^1}(G)(R^h \otimes_R R^h) \\ \downarrow & & \downarrow \cong & & \downarrow \\ \pi_0^{\mathbb{A}^1}(G)(K) & \longrightarrow & \pi_0^{\mathbb{A}^1}(G)(K^h) & \xrightarrow{\cong} & \pi_0^{\mathbb{A}^1}(G)(K^h \otimes_K K^h) \end{array}$$

in which the rows are equalizer diagrams, all the vertical maps are injective and the middle vertical map is an isomorphism. Surjectivity of the leftmost vertical map now follows from an easy diagram chase. \square

Lemma 4.2. *Let G be a semisimple algebraic group over a field k . Let R be a regular local ring with quotient field K such that R is a k -algebra. If G_K is anisotropic, then the natural map $G(R) \rightarrow G(K)$ is an isomorphism. Moreover, if R is henselian, then the map $\pi_0^{\mathbb{A}^1}(G)(R) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K)$ is an isomorphism.*

Proof. First assume that R is a discrete valuation ring. Since G is anisotropic, we have $G(R) = G(K)$ by [14, Proposition A.3].

In the general case, write $S = \text{Spec}(R)$ and let $\alpha \in G(K)$. For every $x \in S^{(1)}$, there exists a unique $\alpha_x \in G(\mathcal{O}_{S,x})$ such that $\alpha_x|_K = \alpha$. Therefore, there exists an open subset U of S with $\text{codim}(S \setminus U) \geq 2$ such that α lies in the image of the natural map $G(U) \rightarrow G(K)$. This gives a rational map $\tilde{\alpha} : S \dashrightarrow G$, which is defined on all the codimension 1 points of S . But since G is affine, the rational map $\tilde{\alpha}$ must be a morphism. This can be seen by embedding G into an affine space and reducing to the statement that any rational map from a normal variety to \mathbb{A}^1 defined at codimension 1 points is a morphism. The latter statement easily follows from [23, Theorem 38], which says that any normal domain is the intersection of its localizations at height 1 prime ideals. Thus, we get an element $\tilde{\alpha} \in G(S)$ extending $\alpha \in G(K)$.

Assume now that R is moreover henselian. Injectivity of the map $\pi_0^{\mathbb{A}^1}(G)(R) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K)$ follows from Lemma 2.5. Surjectivity of the map $\pi_0^{\mathbb{A}^1}(G)(R) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K)$ follows from the surjectivity of the maps $G(R) \rightarrow \pi_0^{\mathbb{A}^1}(G)(R)$ and $G(K) \rightarrow \pi_0^{\mathbb{A}^1}(G)(K)$. \square

Remark 4.3. Statements related to those in Lemma 4.1 and Lemma 4.2 from the perspective of R -equivalence on group schemes have been studied in [21, Section 8]. An independent proof of Lemma 4.1 can be given using [21, Proposition 8.5] in view of (3.1).

Another ingredient required for the proof of birationality of \mathbb{A}^1 -connected components of a semisimple, simply connected group is the preservation of the birationality property under Weil restriction.

Lemma 4.4. *Let L/k be a finite separable extension of fields and let G be an algebraic group over L . If $\pi_0^{\mathbb{A}^1}(G)$ is a birational sheaf, then so is $\pi_0^{\mathbb{A}^1}(R_{L/k}(G))$, where $R_{L/k}$ denotes the Weil restriction functor.*

Proof. Let X be a smooth scheme over k and let U be a dense open subscheme of X . Without loss of generality we may assume that X is henselian and $U = \text{Spec } k(X)$. We then have $U_L = \text{Spec } K(X) \otimes_k L = \text{Spec } L(X_L)$. We need to show that the restriction map $\pi_0^{\mathbb{A}^1}(R_{L/k}(G))(X) \rightarrow \pi_0^{\mathbb{A}^1}(R_{L/k}(G))(U)$ is a bijection. It remains to show surjectivity of this map as this map is known to be injective (see [12, Corollary 4.17] or [25, Lemma 3.45]).

Let $\alpha \in \pi_0^{\mathbb{A}^1}(R_{L/k}(G))(U)$. Since $R_{L/k}(G) \rightarrow \pi_0^{\mathbb{A}^1}(R_{L/k}(G))$ is an epimorphism of Nisnevich sheaves, the element α is the image of an element $\tilde{\alpha} \in R_{L/k}(G)(U) = G(U_L)$. Since $\pi_0^{\mathbb{A}^1}(G)$ is a birational sheaf, there exists $\beta \in G(X_L)$ such that $\beta|_{U_L}$ and $\tilde{\alpha}$ map to the same element in $\pi_0^{\mathbb{A}^1}(G)(U_L)$. We will identify sections of $\pi_0^{\mathbb{A}^1}(G)$ over fields with R -equivalence classes, using [8, Main Theorem] and [9, Theorem 3.6]. Hence, $\beta|_{U_L}$ and $\tilde{\alpha}$ are R -equivalent. So there exists a rational map $h : \mathbb{P}_L^1 \dashrightarrow G$ defined at 0 and 1 such that $h(0) = \beta|_{U_L}$ and $h(1) = \tilde{\alpha}$. This corresponds to a rational map $h' : \mathbb{P}^1 \rightarrow R_{L/k}(G)$ connecting α to the restriction of the element in $R_{L/k}(G)(X)$ corresponding to $\beta \in G(X_L)$ by R -equivalence. Consequently, the restriction map $\pi_0^{\mathbb{A}^1}(R_{L/k}(G))(X) \rightarrow \pi_0^{\mathbb{A}^1}(R_{L/k}(G))(U)$ is surjective. \square

We are now set to prove the main theorem of this section.

Theorem 4.5. *If G is a semisimple, simply connected algebraic group over a perfect field k , then $\pi_0^{\mathbb{A}^1}(G)$ is a birational sheaf.*

Proof. Since G is the direct product of its almost simple factors, it suffices to prove the theorem in the case where G is semisimple, simply connected and almost simple. By [24, Chapter 24.a] (see also [30]), there exists a finite separable extension L/k and an absolutely almost simple group G' such that G is isomorphic to $R_{L/k}(G')$ as an algebraic group over k . In view of Lemma 4.4, it suffices to prove that $\pi_0^{\mathbb{A}^1}(G')$ is a birational sheaf.

In view of Lemma 2.4, it suffices to show that for every henselian local k -scheme $S = \text{Spec}(R)$, the restriction map $\pi_0^{\mathbb{A}^1}(G')(S) \rightarrow \pi_0^{\mathbb{A}^1}(G')(k(S))$ is a surjection (injectivity of this map follows from Lemma 2.5). Let ℓ denote the residue field of S . Since k is perfect, there exists an inclusion $\ell \hookrightarrow R$ as k -algebras. Hence, $\pi_0^{\mathbb{A}^1}(G')(S) = \pi_0^{\mathbb{A}^1}(G'_\ell)(S)$. Thus, without loss of generality, we may assume that $\ell = k$. If G' is anisotropic, then the desired statement is given by Lemma 4.2.

If G' is isotropic, then $\pi_0^{\mathbb{A}^1}(G')$ is strongly \mathbb{A}^1 -invariant by [5, Theorem 4.3.3]. In particular, $\pi_0^{\mathbb{A}^1}(G')$ is an unramified sheaf. Hence,

$$(4.1) \quad \pi_0^{\mathbb{A}^1}(G')(S) = \bigcap_{x \in S^{(1)}} \pi_0^{\mathbb{A}^1}(G')(\mathcal{O}_{S,x}) \subseteq \pi_0^{\mathbb{A}^1}(G')(k(S)).$$

Let $\alpha \in \pi_0^{\mathbb{A}^1}(G')(k(S))$. We will show that α “extends to all the codimension 1 points of S ”. Let $x \in S^{(1)}$. By Lemma 4.1 applied to $\text{Spec}(\mathcal{O}_{S,x})$, there exists $\alpha_x \in \pi_0^{\mathbb{A}^1}(G')(\mathcal{O}_{S,x})$ such that $\alpha_x|_{k(S)} = \alpha$. Since the restriction map $\pi_0^{\mathbb{A}^1}(G')(\mathcal{O}_{S,x}) \rightarrow \pi_0^{\mathbb{A}^1}(G')(k(S))$ is injective for every $x \in S^{(1)}$, it follows that $\alpha_x|_{k(S)} = \alpha_y|_{k(S)}$ for all $x, y \in S^{(1)}$. The surjectivity of $\pi_0^{\mathbb{A}^1}(G')(S) \rightarrow \pi_0^{\mathbb{A}^1}(G')(k(S))$ now follows from (4.1). \square

Corollary 4.6. *If G is a semisimple, simply connected algebraic group over a perfect field k , then $\pi_0^{\mathbb{A}^1}(G)$ is strongly \mathbb{A}^1 -invariant.*

Proof. This is a straightforward consequence of Theorem 4.5 and Remark 2.3, since $\pi_0^{\mathbb{A}^1}(G)$ is an \mathbb{A}^1 -invariant sheaf by [12, Theorem 4.18]. \square

5. STRONG \mathbb{A}^1 -INVARIANCE OF THE SHEAF OF \mathbb{A}^1 -CONNECTED COMPONENTS OF A REDUCTIVE GROUP

In this section, we build on the results of the previous section to show that the sheaf of \mathbb{A}^1 -connected components of any reductive algebraic group over a perfect field is strongly \mathbb{A}^1 -invariant. We begin by observing that strong \mathbb{A}^1 -invariance of \mathbb{A}^1 -connected components is preserved under isogenies.

Lemma 5.1. *Let k be a field. Let $\tilde{G} \rightarrow G$ be a central isogeny of semisimple algebraic groups, the kernel of which is a group μ of multiplicative type. If $\pi_0^{\mathbb{A}^1}(\tilde{G})$ is strongly \mathbb{A}^1 -invariant, then so is $\pi_0^{\mathbb{A}^1}(G)$.*

Proof. By [26, §4.3, Proposition 3.1], the space $B_{\text{ét}}\mu$ is \mathbb{A}^1 -local. We have a simplicial fiber sequence

$$\tilde{G} \rightarrow G \rightarrow B_{\text{ét}}\mu,$$

since $G \rightarrow B_{\text{ét}}\mu$ is a \tilde{G} -torsor. Since $\pi_0^{\mathbb{A}^1}(\tilde{G})$ is strongly \mathbb{A}^1 -invariant by hypothesis, by [25, Theorem 6.50], the sequence is an \mathbb{A}^1 -fiber sequence. This gives rise to an exact sequence of sheaves of sets

$$\cdots \rightarrow \pi_0^{\mathbb{A}^1}(\tilde{G}) \rightarrow \pi_0^{\mathbb{A}^1}(G) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu).$$

For every smooth henselian local k -scheme U , we have a commutative diagram

$$\begin{array}{ccccccc} G(U) & \longrightarrow & H_{\text{ét}}^1(U, \mu) & \longrightarrow & H_{\text{ét}}^1(U, \tilde{G}) & \longrightarrow & H_{\text{ét}}^1(U, G) \\ \downarrow & & \downarrow \simeq & & \downarrow \simeq & & \\ \pi_0^{\mathbb{A}^1}(G)(U) & \longrightarrow & \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu)(U) & \longrightarrow & \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\tilde{G})(U) & \longrightarrow & \end{array}$$

in which the top row is exact. Note that the cokernel of the map $\pi_0^{\mathbb{A}^1}(G)(U) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu)(U)$ is contained in $\pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu)(U) \simeq H_{\text{ét}}^1(U, \tilde{G})$. By Theorem 3.8, the sheaf $\pi_0^{\mathbb{A}^1}(B_{\text{ét}}\tilde{G}) = \mathcal{H}_{\text{ét}}^1(\tilde{G})$ is \mathbb{A}^1 -invariant. Thus, the cokernel of $\pi_0^{\mathbb{A}^1}(G) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu)$ is an \mathbb{A}^1 -invariant sheaf. Note that the sheaf $\pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu) = \mathcal{H}_{\text{ét}}^1(\mu)$ is strongly \mathbb{A}^1 -invariant by [1, Example 4.6]. Applying [13, Theorem 1.3], we see that $\text{Coker}(\pi_0^{\mathbb{A}^1}(G) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu))$ is a strongly \mathbb{A}^1 -invariant sheaf. Now [13, Theorem 1.5] implies that $\text{Im}(\pi_0^{\mathbb{A}^1}(G) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu))$ is a strongly \mathbb{A}^1 -invariant sheaf. Since $\pi_0^{\mathbb{A}^1}(\tilde{G})$ is strongly \mathbb{A}^1 -invariant, another application of [13, Theorem 1.3, Theorem 1.5] yields that $\text{Im}(\pi_0^{\mathbb{A}^1}(\tilde{G}) \rightarrow \pi_0^{\mathbb{A}^1}(G))$ is strongly \mathbb{A}^1 -invariant. We now conclude from the short exact sequence

$$1 \rightarrow \text{Im}(\pi_0^{\mathbb{A}^1}(\tilde{G}) \rightarrow \pi_0^{\mathbb{A}^1}(G)) \rightarrow \pi_0^{\mathbb{A}^1}(G) \rightarrow \text{Im}(\pi_0^{\mathbb{A}^1}(G) \rightarrow \pi_0^{\mathbb{A}^1}(B_{\text{ét}}\mu)) \rightarrow 1$$

that $\pi_0^{\mathbb{A}^1}(G)$ is strongly \mathbb{A}^1 -invariant. \square

We now use the standard structure theory of reductive groups along with Lemma 5.1 and the results of Section 4 to obtain our main theorem.

Theorem 5.2. *If G is a reductive algebraic group over a perfect field k , then $\pi_0^{\mathbb{A}^1}(G)$ is strongly \mathbb{A}^1 -invariant.*

Proof. Let G_{der} denote the derived group of G , which is a normal, semisimple subgroup scheme of G . Let G_{sc} denote the universal simply connected cover of G_{der} . By Corollary 4.6, the sheaf $\pi_0^{\mathbb{A}^1}(G_{\text{sc}})$ is strongly \mathbb{A}^1 -invariant. Lemma 5.1 then implies that $\pi_0^{\mathbb{A}^1}(G_{\text{der}})$ is strongly \mathbb{A}^1 -invariant.

There exists a central isogeny (see [32, Exposé XXII, Proposition 6.2.4]) $G_{\text{der}} \times T \rightarrow G$, where T is a torus, the radical of G . This is a faithfully flat, finitely presented morphism, whose kernel is a finite group of multiplicative type contained in the center of $G_{\text{der}} \times T$. It follows by another application of Lemma 5.1 that $\pi_0^{\mathbb{A}^1}(G)$ is strongly \mathbb{A}^1 -invariant, since $\pi_0^{\mathbb{A}^1}(G_{\text{der}} \times T) \simeq \pi_0^{\mathbb{A}^1}(G_{\text{der}}) \times T$ is strongly \mathbb{A}^1 -invariant. \square

A straightforward consequence of Theorem 5.2 and [25, Theorem 6.50] is the following application to \mathbb{A}^1 -fiber sequences.

Corollary 5.3. *Let G be a reductive algebraic group over a perfect field k . For any Nisnevich locally trivial G -torsor $\mathcal{P} \rightarrow \mathcal{X}$ in $\mathcal{H}(k)$,*

$$\mathcal{P} \rightarrow \mathcal{X} \rightarrow BG$$

is an \mathbb{A}^1 -fiber sequence.

REFERENCES

- [1] A. Asok: *Birational invariants and \mathbb{A}^1 -connectedness*, J. Reine Angew. Math. 681 (2013), 39–64. [12](#)
- [2] A. Asok: *Splitting vector bundles and \mathbb{A}^1 -fundamental groups of higher-dimensional varieties*, J. Topol. 6 (2013), no. 2, 311–348. [5](#)
- [3] A. Asok, J. Fasel: *A cohomological classification of vector bundles on smooth affine threefolds*, Duke Math. J. 163 (2014), 14, 2651–2601. [1](#)
- [4] A. Asok, J. Fasel: *Splitting vector bundles outside the stable range and \mathbb{A}^1 -homotopy sheaves of punctured affine spaces*, J. Amer. Math. Soc. 28 (2015), no. 4, 1031–1062. [1](#)
- [5] A. Asok, M. Hoyois, M. Wendt: *Affine representability results in \mathbb{A}^1 -homotopy theory, II: Principal bundles and homogeneous spaces*, Geom. Topol. 22 (2018), no. 2, 1181–1225. [2](#), [5](#), [7](#), [8](#), [11](#)
- [6] A. Asok, M. Hoyois, M. Wendt: *Affine representability results in \mathbb{A}^1 -homotopy theory, III: Finite fields and complements*, Algebr. Geom. 7 (2020), no. 5, 634–644. [2](#), [5](#)
- [7] A. Asok, F. Morel: *Smooth varieties up to \mathbb{A}^1 -homotopy and algebraic h-cobordisms*, Adv. Math. 227(2011), no. 5, 1990–2058. [4](#), [6](#)
- [8] C. Balwe and A. Sawant: *R-equivalence and \mathbb{A}^1 -connectedness in anisotropic groups*, Int. Math. Res. Not. IMRN 2015, No. 22, 11816–11827. [2](#), [5](#), [6](#), [7](#), [10](#)
- [9] C. Balwe, A. Sawant: *\mathbb{A}^1 -connectedness in reductive algebraic groups*, Trans. Amer. Math. Soc. 369 (2017), no. 8, 5999–6015. [5](#), [7](#), [10](#)
- [10] A. Borel, T. A. Springer: *Rationality properties of linear algebraic groups. II*, Tohoku Math. J. 20 (1968), 443–497. [9](#)
- [11] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. No. 27 (1965), 55–150. [6](#)
- [12] U. Choudhury: *Connectivity of motivic H-spaces*, Algebr. Geom. Topol. 14 (2014), no. 1, 37–55. [2](#), [5](#), [10](#), [11](#)
- [13] U. Choudhury, A. Hogadi: *The Hurewicz map in motivic homotopy theory*, Annals of K-theory, Vol. 7 (2022), No. 1, 179–190. [2](#), [12](#)
- [14] J.-L. Colliot-Thélène, D. Harbater, J. Hartmann, D. Krashen, R. Parimala, V. Suresh: *Local-global principles for constant reductive groups over semi-global fields*, Preprint: arXiv:2108.12349 [math.AG]. [6](#), [10](#)
- [15] B. Conrad, O. Gabber and G. Prasad: *Pseudo-reductive groups*, New Mathematical Monographs, 26, Cambridge University Press, Cambridge, 2015. [5](#)
- [16] E. Elmanto, G. Kulkarni, M. Wendt: *\mathbb{A}^1 -connected components of classifying spaces and purity for torsors*, Preprint: arXiv:2104.06273 [math.AG]. [3](#), [8](#), [9](#)
- [17] R. Fedorov, I. A. Panin: *A proof of the Grothendieck-Serre conjecture on principal bundles over regular local rings containing infinite fields*, Publ. Math. Inst. Hautes Études Sci. 122 (2015), 169–193. [9](#)
- [18] O. Gabber, P. Gille, L. Moret-Bailly: *Fibrés principaux sur les corps valués henséliens*, Algebr. Geom. 1 (2014), no. 5, 573–612. [6](#)
- [19] P. Gille: *Le problème de Kneser-Tits*, Séminaire Bourbaki. Vol. 2007/2008. Astérisque No. 326 (2009), Exp. No. 983, vii, 39–81 (2010). [2](#), [5](#), [6](#), [7](#), [8](#), [9](#)
- [20] P. Gille: *Gabber’s compactifications of algebraic groups and homogeneous spaces*, Oberwolfach reports, 31/2013. [6](#)
- [21] P. Gille, A. Stavrova: *R-equivalence on reductive group schemes*, Preprint, arXiv:2107.01950v3 [math.AG]. [10](#)
- [22] G. Harder: *Halbeinfache Gruppenschemata über Dedekindringen*, Invent. Math. 4 (1967), 165–191. [9](#)
- [23] H. Matsumura: *Commutative algebra*, Second edition. Mathematics Lecture Note Series, 56. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980. [10](#)
- [24] J.S. Milne: *Algebraic groups. The theory of group schemes of finite type over a field*, Cambridge Studies in Advanced Mathematics, 170, Cambridge University Press, 2017. [5](#), [11](#)

- [25] F. Morel: \mathbb{A}^1 -algebraic topology over a field, Lecture Notes in Mathematics, Vol. 2052, Springer, Heidelberg, 2012. [1](#), [2](#), [4](#), [5](#), [10](#), [12](#)
- [26] F. Morel, V. Voevodsky: \mathbb{A}^1 -homotopy theory of schemes, Inst. Hautes Études Sci. Publ. Math. 90 (1999), 45–143. [1](#), [12](#)
- [27] I.A. Panin: *Proof of the Grothendieck-Serre conjecture on principal bundles over regular local rings containing a field*, Izv. Ross. Akad. Nauk Ser. Mat. 84 (2020), no. 4, 169–186. [9](#)
- [28] M.S. Raghunathan, A. Ramanathan: *Principal bundles on the affine line*, Proc. Indian Acad. Sci. (Math. Sci.) 93, No. 2 & 3 (1984), 137–145. [9](#)
- [29] R. Steinberg: *Regular elements of semisimple algebraic groups*, Inst. Hautes Études Sci. Publ. Math. No. 25 (1965), 49–80. [9](#)
- [30] J. Tits: *Classification of algebraic semisimple groups*, Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965), Amer. Math. Soc. (1966), 33–62. [11](#)
- [31] M. Wendt: \mathbb{A}^1 -homotopy of Chevalley groups, J. K-Theory 5 (2) (2010), 245–287. [2](#)
- [32] *Schémas en groupes. III: Structure des schémas en groupes réductifs*, Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 153, Springer-Verlag, 1970. [12](#)

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH
MOHALI, KNOWLEDGE CITY, SECTOR-81, MOHALI 140306, INDIA.

Email address: cbalwe@iisermohali.ac.in

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH
PUNE, DR. HOMI BHABHA ROAD, PASHAN, PUNE 411008, INDIA.

Email address: amit@iiserpune.ac.in

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, CO-
LABA, MUMBAI 400005, INDIA.

Email address: asawant@math.tifr.res.in