

Filtrations on the globalization of twisted D-modules over Dedekind schemes

Takuma Hayashi*

Abstract

In [12], we established the theory of twisted D-modules over general base schemes. In this short note, we construct a K -invariant positive exhaustive filtration on the globalization of the twisted D-module on a smooth quasi-compact K -scheme over a Dedekind scheme S obtained by the direct image of a torsion-free and \mathcal{O}_Y -coherent K -equivariant twisted D-module along a K -equivariant closed immersion from a smooth proper integral K -scheme Y with K a smooth S -affine group scheme, whose p th associated graded \mathcal{O}_S -module is finite locally free for every integer p .

1 Introduction

In the 2010s, arithmetic structures of Harish-Chandra modules attract attention to some authors for applications to the study of special values of automorphic L -functions. In fact, the cohomology of adélic locally symmetric spaces is expected to know some special L -values. The origin of this idea goes back to [19]. One may wish that arithmetic structures of the decomposition of the cuspidal cohomology into cohomology of irreducible cuspidal automorphic representations, together with more arguments, for example, the study of the Eisenstein cohomology will lead to rationality or integrality results of special L -values. For this reason, one can think of the study of arithmetic structures of Harish-Chandra modules as an Archimedean part of the theory of automorphic representations in this context.

To get rational models of discrete series representations, M. Harris proposed in [6] and [7] to use the localization. F. Januszewski introduced the cohomological induction functor over fields of characteristic zero to get rational models of cohomologically induced modules in [14] Section 3 and Section 7. Then the author constructed the cohomological induction functor over general commutative rings in [9] and [10] to get models over Dedekind domains.

In [14], Januszewski also proposed to study the descent problem of the fields of definition of Harish-Chandra modules as a new aspect of representation theory

*Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, 1-5 Yamadaoka, Suita, Osaka 565-0871, Japan, hayashi-t@ist.osaka-u.ac.jp

of real reductive Lie groups. As one can find in [15], the descent theorems (potentially) imply stronger rationality of special L -values. Januszewski gave examples of smaller fields of definition by using the Galois descent ([14] Section 7.4).

In [12], Januszewski and the author improved Harris' idea to construct smaller arithmetic models of cohomologically induced modules. In fact, $A_q(\lambda)$ -modules are realized as the spaces of global sections of the twisted D-module theoretic direct image of equivariant line bundles on the corresponding closed K -orbits to partial flag varieties (see [13] Theorem 4.3 and [16] Theorem 5.1, Corollary 5.5). Therefore we should get smaller models of $A_q(\lambda)$ -modules from smaller models of their geometric materials via the theory of twisted D-modules. To follow this idea, they developed the theory of twisted D-modules over general base schemes in [12] Section 1–4. They also discussed the descent problem of base schemes of certain closed orbits on partial flag schemes in [12] Section 5. The author studied a cohomological classification of data of Galois descent for the rings of definition of equivariant line bundles on partial flag schemes in [11]. As a consequence, we achieved the construction of arithmetic models of $A_q(\lambda)$ -modules and the descent of their rings of definition simultaneously. However, it was nontrivial yet that the models over Dedekind domains of dimension one are essentially new. In fact, rational models are already integral models due to the isomorphism $\mathbb{Q} \otimes_{\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q}$. Here \mathbb{Z} and \mathbb{Q} are the ring of integers and the field of rational numbers respectively. In this paper, we distinguish the models of $A_q(\lambda)$ -modules over Dedekind domains of dimension one we constructed in [12] Section 6 from the rational models by proving the following result:

Theorem 1.1 (Corollary 3.2, Proposition 4.4, Proposition 4.5, Theorem 5.1). *Let K be a smooth S -affine group scheme over a Dedekind scheme S (i.e., a Noetherian integral regular scheme of dimension ≤ 1 , see [3] Section (7.13)), $i : Y \hookrightarrow X$ be a K -equivariant closed immersion of smooth K -schemes over S , and \mathcal{A} be a K -equivariant tdo on X . Write $x : X \rightarrow S$ for the structure morphism. Suppose that the following conditions are satisfied:*

- (i) X is quasi-compact over S ;
- (ii) Y is proper over S ;
- (iii) Y is integral.

Let \mathcal{V} be a K -equivariant right $i^\mathcal{A}$ -module which is coherent and torsion-free as an \mathcal{O}_Y -module in the sense of [4] Proposition (7.4.1). Then there exists a natural exhaustive K -invariant filtration $G_\bullet i_+ \mathcal{V}$ on $i_+ \mathcal{V}$ such that the p th associated graded sheaf $\mathrm{gr}_G^p x_* i_+ \mathcal{V} := x_* G_p i_+ \mathcal{V} / x_* G_{p-1} i_+ \mathcal{V}$ to the induced K -invariant filtration $x_* G_\bullet i_+ \mathcal{V}$ on $x_* i_+ \mathcal{V}$ is a finite locally free \mathcal{O}_S -module for every integer p .*

Our strategy for the proof is as follows: For each integer p , let $G_p \mathcal{A}_{Y \rightarrow X}$ be the left $i^* \mathcal{A}$ -module generated by $i^* F_p \mathcal{A}$, where $F_p \mathcal{A}$ is the p th filter of the given

filtration on \mathcal{A} . Define filtrations

$$\begin{aligned} G_\bullet(\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X}), \\ G_\bullet i_+ \mathcal{V}, \\ G_\bullet x_* i_+ \mathcal{V} \end{aligned}$$

on $\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X}$, $i_+ \mathcal{V}$, $x_* i_+ \mathcal{V}$ by

$$\begin{aligned} G_\bullet(\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X}) &= \mathcal{V} \otimes_{i_*\mathcal{A}} G_\bullet \mathcal{A}_{Y \rightarrow X} \\ G_\bullet i_+ \mathcal{V} &= i_* G_\bullet(\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X}) \\ G_\bullet x_* i_+ \mathcal{V} &= x_* G_\bullet i_+ \mathcal{V} \end{aligned}$$

respectively. One can prove that $G_\bullet i_+ \mathcal{V}$ is K -invariant by putting the K -equivariant structure on $G_\bullet i_+ \mathcal{V}$ in a similar way to [12] Theorem 3.11.2. The filtration $G_\bullet i_+ \mathcal{V}$ on $i_+ \mathcal{V}$ is exhaustive since i is quasi-compact. Since X is quasi-compact and quasi-separated over S , $G_\bullet x_* i_+ \mathcal{V}$ is an exhaustive quasi-coherent K -invariant filtration on the representation $x_* i_+ \mathcal{V}$ of K (cf. [12] Theorem 3.11.2, Theorem 3.12.1). The key idea for the algebraic properties of $G_\bullet x_* i_+ \mathcal{V}$ is to prove an isomorphism

$$\mathrm{gr}_G^p(\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X}) \cong \mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y}^p i^* \Theta_X / \Theta_{Y/S}$$

for each integer p as [18] Lemma 3.3 and the proof of [2] 7.7 Proposition. One can prove by comparing $\mathrm{gr}_G^p x_* i_+ \mathcal{V}$ and $y_* \mathrm{gr}_G^p(\mathcal{V} \otimes_{i_*\mathcal{A}} \mathcal{A}_{Y \rightarrow X})$ that $\mathrm{gr}_G^p x_* i_+ \mathcal{V}$ is a torsion-free and coherent \mathcal{O}_S -module, where y is the structure morphism $Y \rightarrow S$. The assertion now follows since S is Dedekind.

One can also prove the following result in the almost same line:

Variante 1.2 (Corollary 3.2, Theorem 5.1). *Let $i : Y \hookrightarrow X$ be a closed immersion of smooth schemes over a Noetherian integral domain k which is Dedekind in the sense of [3] Definition/Proposition B.87, and \mathcal{A} be a tdo on X . Suppose that Y is proper over k and integral. Let \mathcal{V} be a right $i_*\mathcal{A}$ -module which is coherent and torsion-free as an \mathcal{O}_Y -module. Then there exists a natural exhaustive k -linear filtration $G_\bullet \Gamma(X, i_+ \mathcal{V})$ on $\Gamma(X, i_+ \mathcal{V})$ such that the p th associated graded k -module $G_p \Gamma(X, i_+ \mathcal{V}) / G_{p-1} \Gamma(X, i_+ \mathcal{V})$ is finitely generated and projective for every integer p .*

We only see the global sections this time. Hence we need not to appeal to the quasi-coherence of $G_\bullet x_* i_+ \mathcal{V}$. This is why we do not assume x to be quasi-compact. The exhaustivity follows since y is quasi-compact. Besides these points, the assertion is proved in the same line as Theorem 1.1.

A standard argument on the Mittag-Leffler condition (Lemma 5.2) implies that our models are essentially new also in the following sense:

Corollary 1.3. *Let $i : Y \hookrightarrow X$ be a closed immersion of smooth schemes over a Dedekind domain k , and \mathcal{A} be a tdo on X . Suppose that Y is proper over k and integral. Let \mathcal{V} be a right $i_*\mathcal{A}$ -module which is coherent and torsion-free as an \mathcal{O}_Y -module. Then $\Gamma(X, i_+ \mathcal{V})$ is projective as a k -module.*

Finally, remark that one can prove all the results in this paper for left $i\mathcal{A}$ -modules \mathcal{U} in the same lines by replacing $\mathcal{A}_{Y \rightarrow X}$ with

$$\mathcal{A}_{X \leftarrow Y} = \omega_{Y/S} \otimes_{\mathcal{O}_Y} (\omega_{X/S}^\vee \otimes_{\mathcal{O}_X} \mathcal{A}^{\text{op}} \otimes_{\mathcal{O}_X} \omega_{X/S})_{Y \rightarrow X} \otimes_{i^{-1}\mathcal{O}_X} i^{-1}\omega_{X/S}^\vee$$

in [12] Section 3.9.1.

Acknowledgements

The author thanks Yoshiki Oshima for helpful comments.

This work was supported by JSPS KAKENHI Grant Number JP21J00023.

2 Notation

All the filtrations in this paper are increasing and positive. That is, any filtration $F_\bullet \mathcal{M}$ on an abelian group or a sheaf of abelian groups satisfies $F_p \mathcal{M} \subset F_{p+1} \mathcal{M}$ for every integer p and $F_{-1} \mathcal{M} = 0$. For an object \mathcal{M} , equipped with a filtration $F_\bullet \mathcal{M}$, we denote the associated graded object by

$$\text{gr}_F \mathcal{M} = \bigoplus_p \text{gr}_F^p \mathcal{M}, \quad \text{gr}_F^p \mathcal{M} = F_p \mathcal{M} / F_{p-1} \mathcal{M},$$

where p runs through all integers.

Let (X, \mathcal{O}_X) be a ringed space. We refer to the unbounded derived category of \mathcal{O}_X -modules as $D(\mathcal{O}_X)$. For an integer d , let $D^{\leq d}(\mathcal{O}_X) \subset D(\mathcal{O}_X)$ denote the full subcategory consisting of complexes \mathcal{F}^\bullet satisfying $H^n(\mathcal{F}^\bullet) = 0$ for $n > d$. Let \mathcal{F} be an \mathcal{O}_X -module. For an open subset $U \subset X$, let $\Gamma(U, \mathcal{F})$ denote the abelian group of U -sections of \mathcal{F} . For an integer p , we refer to the p th symmetric power of \mathcal{F} as $\text{Sym}_{\mathcal{O}_X}^p \mathcal{F}$. Set $\text{Sym}_{\mathcal{O}_X} \mathcal{F} = \bigoplus_p \text{Sym}_{\mathcal{O}_X}^p \mathcal{F}$ as the total symmetric power of \mathcal{F} .

For a scheme X , we write \mathcal{O}_X for its structure sheaf.

Let S be a scheme, and X be a smooth S -scheme. We denote the structure morphism $X \rightarrow S$ by the corresponding small letter x . Let $\Theta_{X/S}$ be the \mathcal{O}_X -module of S -vector fields on X . Write $\omega_{X/S}$ and $\omega_{X/S}^\vee$ for the canonical bundle on X and its dual.

In the rest of this paper, let S be a scheme, and $i: Y \hookrightarrow X$ be an immersion of smooth schemes over S unless specified otherwise. Let i_* and i^{-1} be the sheaf theoretic direct image and pullback respectively. For a left \mathcal{O}_X -module \mathcal{F} , set $i^* \mathcal{F} = \mathcal{O}_Y \otimes_{i^{-1}\mathcal{O}_X} i^{-1} \mathcal{F}$. Set $\mathcal{N}_{X/Y} = i^* \Theta_{X/S} / \Theta_{Y/S}$ as the normal sheaf.

For the basic notations on tdos and twisted D-modules over schemes, we follow [12]. Let \mathcal{A} be a tdo on X ([12] Definition 1.1.4). Write $F_\bullet \mathcal{A}$ for the given filtration on \mathcal{A} . Recall that we defined the pullback $i\mathcal{A}$ in [12] Section 1.3.5. Let $\mathcal{A}_{Y \rightarrow X}$ denote the $(i\mathcal{A}, i^{-1}\mathcal{A})$ -bimodule in [12] Lemma 3.6.5. We will fix a right $i\mathcal{A}$ -module \mathcal{V} . Write $i_+ \mathcal{V} = i_*(\mathcal{V} \otimes_{i\mathcal{A}} \mathcal{A}_{Y \rightarrow X})$ in this paper (even if i is not a closed immersion).

3 Filter G on $\mathcal{A}_{Y \rightarrow X}$

The aim of this section is to study the filtration on $\mathcal{A}_{Y \rightarrow X}$ explained in Section 1.

Regard \mathcal{A} as an \mathcal{O}_X -bimodule for the multiplication from the left and right sides. Then for each integer p , $F_p \mathcal{A}$ is a sub- \mathcal{O}_X -bimodule of \mathcal{A} by definition. Since $F_\bullet \mathcal{A}$ is an exhaustive filtration on \mathcal{A} , we obtain an initial (colimit) cocone

$$\begin{array}{ccccccc} \cdots & \hookrightarrow & F_{p-1} \mathcal{A} & \hookrightarrow & F_p \mathcal{A} & \hookrightarrow & F_{p+1} \mathcal{A} & \hookrightarrow & \cdots \\ & & & & \downarrow & & & & \\ & & & & \mathcal{A} & & & & \end{array}$$

Apply i^* to get an initial cocone of $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -bimodules

$$\begin{array}{ccccccc} \cdots & \hookrightarrow & i^* F_{p-1} \mathcal{A} & \hookrightarrow & i^* F_p \mathcal{A} & \hookrightarrow & i^* F_{p+1} \mathcal{A} & \hookrightarrow & \cdots \\ & & & & \downarrow & & & & \\ & & & & \mathcal{A}_{Y \rightarrow X} & & & & \end{array}$$

The horizontal arrows are monic since $\mathrm{gr}_F^p \mathcal{A}$ are finite locally free as left \mathcal{O}_X -modules for all integers p . Therefore $F_\bullet \mathcal{A}_{Y \rightarrow X} := i^* F_\bullet \mathcal{A}$ exhibits an exhaustive filtration on $\mathcal{A}_{Y \rightarrow X}$ as an $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -bimodule. Since i^* is right exact, we also have a canonical $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -bilinear isomorphism $i^* \mathrm{gr}_F^p \mathcal{A} \cong \mathrm{gr}_F^p \mathcal{A}_{Y \rightarrow X}$ for each integer p . Write ϕ for the total isomorphism $i^* \mathrm{gr}_F \mathcal{A} \cong \mathrm{gr}_F \mathcal{A}_{Y \rightarrow X}$.

Let $G_p \mathcal{A}_{Y \rightarrow X}$ be the left $i^* \mathcal{A}$ -submodule of $\mathcal{A}_{Y \rightarrow X}$ generated by $F_p \mathcal{A}_{Y \rightarrow X}$. Since $F_p \mathcal{A}$ is a right \mathcal{O}_X -submodule of \mathcal{A} , $G_p \mathcal{A}_{Y \rightarrow X}$ is a right $i^{-1} \mathcal{O}_X$ -submodule of $\mathcal{A}_{Y \rightarrow X}$.

Theorem 3.1. *There is a canonical graded $(i^* \mathcal{A}, i^{-1} \mathcal{O}_X)$ -bilinear isomorphism*

$$i^* \mathcal{A} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y} \cong \mathrm{gr}_G \mathcal{A}_{Y \rightarrow X},$$

where $\mathrm{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y}$ is regarded as a right $i^{-1} \mathcal{O}_X$ -module for the multiplication through the structure homomorphism $i^{-1} \mathcal{O}_X \rightarrow \mathcal{O}_Y$.

Proof. Recall that we are given an \mathcal{O}_X -algebra isomorphism

$$\sigma : \mathrm{Sym}_{\mathcal{O}_X} \Theta_{X/S} \cong \mathrm{gr}_F \mathcal{A}$$

by definition of tdos . Compose $i^* \sigma$ with the canonical isomorphism

$$\mathrm{Sym}_{\mathcal{O}_Y} i^* \Theta_{X/S} \cong i^* \mathrm{Sym}_{\mathcal{O}_X} \Theta_{X/S}$$

to get

$$\mathrm{Sym}_{\mathcal{O}_Y} i^* \Theta_{X/S} \cong \mathrm{Sym}_{\mathcal{O}_Y} i^* \Theta_{X/S} \stackrel{i^* \sigma}{\cong} i^* \mathrm{gr}_F \mathcal{A},$$

which we will denote by ψ . We regard ψ as an $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -bilinear isomorphism. Here each vertex in the above sequence is a right $i^{-1} \mathcal{O}_X$ -module for the structure homomorphism $i^{-1} \mathcal{O}_X \rightarrow \mathcal{O}_Y$.

Since $F_p \mathcal{A}_{Y \rightarrow X} \subset G_p \mathcal{A}_{Y \rightarrow X}$ for every p , we have a canonical $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -bilinear homomorphism $\pi : \text{gr}_F \mathcal{A}_{Y \rightarrow X} \rightarrow \text{gr}_G \mathcal{A}_{Y \rightarrow X}$. Let τ denote the composite map

$$\text{Sym}_{\mathcal{O}_Y} i^* \Theta_{X/S} \xrightarrow{\psi} i^* \text{gr}_F \mathcal{A} \xrightarrow{\phi} \text{gr}_F \mathcal{A}_{Y \rightarrow X} \xrightarrow{\pi} \text{gr}_G^p \mathcal{A}_{Y \rightarrow X}.$$

We wish to prove that τ descends to an $(\mathcal{O}_Y, i^{-1} \mathcal{O}_X)$ -module homomorphism $\text{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y} \rightarrow \text{gr}_G^p \mathcal{A}_{Y \rightarrow X}$. This problem is local in X and Y since the desired homomorphism $\text{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y} \rightarrow \text{gr}_G^p \mathcal{A}_{Y \rightarrow X}$ exists at most uniquely. Therefore we may take (x_i) , (∂_i) , and $r \leq n$ in [12] Theorem A.2.20. Then one can identify $\mathcal{N}_{X/Y}$ with $\bigoplus_{r+1 \leq j \leq n} \mathcal{O}_Y \partial_j \subset i^* \Theta_{X/S}$ as an \mathcal{O}_Y -module. The proof is completed by showing that the composite map

$$\mathcal{O}_Y [\partial_{r+1}, \partial_{r+2}, \dots, \partial_n] \subset \text{Sym}_{\mathcal{O}_Y} i^* \Theta_{X/S} \xrightarrow{\tau} \text{gr}_G^p \mathcal{A}_{Y \rightarrow X}$$

exhibits the desired homomorphism. For this, it will suffice to show

$$\sigma(a \partial_1^{l_1} \partial_2^{l_2} \dots \partial_n^{l_n}) = 0$$

for a local section $a \in \mathcal{O}_Y$ and nonnegative integers l_1, l_2, \dots, l_n with $\sum_{j=1}^r l_j > 0$. Choose a lift $e_j \in F_1 \mathcal{A}$ of ∂_j for each $1 \leq j \leq n$. Then $\sigma(a \partial_1^{l_1} \partial_2^{l_2} \dots \partial_n^{l_n})$ lifts to $a \otimes e_1^{l_1} e_2^{l_2} \dots e_n^{l_n} \in G_{\sum_{j=r+1}^n l_j} \mathcal{A}_{Y \rightarrow X} \subset G_{\sum_{j=1}^n l_j - 1} \mathcal{A}_{Y \rightarrow X}$. This shows $\sigma(a \partial_1^{l_1} \partial_2^{l_2} \dots \partial_n^{l_n}) = 0$ in $\text{gr}_G^{\sum_{j=1}^n l_j} \mathcal{A}_{Y \rightarrow X}$.

Finally, we show the scalar extension $\tilde{\tau} : i \mathcal{A} \otimes_{\mathcal{O}_Y} \text{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y} \rightarrow \text{gr}_G \mathcal{A}_{Y \rightarrow X}$ is an isomorphism. We may again choose (x_i) , (∂_i) , and $r \leq n$ in [12] Theorem A.2.20 and the lift (e_j) as above. Then

$$1 \otimes \partial_{r+1}^{l_{r+1}} \partial_{r+2}^{l_{r+2}} \dots \partial_n^{l_n}$$

with $l_{r+1}, l_{r+2}, \dots, l_n \geq 0$ form a free basis of $i \mathcal{A} \otimes_{\mathcal{O}_Y} \text{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y}$ as a left $i \mathcal{A}$ -module. The homomorphism $\tilde{\tau}$ sends them to $1 \otimes e_{r+1}^{l_{r+1}} e_{r+2}^{l_{r+2}} \dots e_n^{l_n}$, which form a free basis of $\text{gr}_G \mathcal{A}_{Y \rightarrow X}$ by the proof of [12] Lemma 3.9.5. This completes the proof. \square

Set

$$\begin{aligned} G_\bullet(\mathcal{V} \otimes_{i \mathcal{A}} \mathcal{A}_{Y \rightarrow X}) &:= \mathcal{V} \otimes_{i \mathcal{A}} G_\bullet \mathcal{A}_{Y \rightarrow X} \\ G_\bullet i_+ \mathcal{V} &:= i_* (\mathcal{V} \otimes_{i \mathcal{A}} G_\bullet \mathcal{A}_{Y \rightarrow X}) \\ G_\bullet x_* i_+ \mathcal{V} &:= x_* G_\bullet i_+ \mathcal{V} = y_* G_\bullet (\mathcal{V} \otimes_{i \mathcal{A}} \mathcal{A}_{Y \rightarrow X}). \end{aligned}$$

Corollary 3.2. (1) The $i^{-1} \mathcal{O}_X$ -modules $G_\bullet(\mathcal{V} \otimes_{i \mathcal{A}} \mathcal{A}_{Y \rightarrow X})$ form an exhaustive filtration on $\mathcal{V} \otimes_{i \mathcal{A}} \mathcal{A}_{Y \rightarrow X}$ with

$$\text{gr}_G(\mathcal{V} \otimes_{i \mathcal{A}} \mathcal{A}_{Y \rightarrow X}) \cong \mathcal{V} \otimes_{\mathcal{O}_Y} \text{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y}.$$

(2) The \mathcal{O}_X -modules $G_\bullet i_+ \mathcal{V}$ form a filtration on $i_+ \mathcal{V}$. Moreover, it is exhaustive if i is quasi-compact.

(3) The \mathcal{O}_S -modules $G_\bullet x_* i_+ \mathcal{V}$ form a filtration on $x_* i_+ \mathcal{V}$. Moreover, it is exhaustive if y is quasi-compact.

Proof. Part (1) is an easy consequence of Theorem 3.1 (cf. the argument at the beginning of this section). The first part of (2) (resp. (3)) follows since i_* (resp. y_*) respects monomorphisms. The latter part of (2) and (3) follows from the general fact that the direct image functor of sheaves along a quasi-compact morphism of schemes respects direct limits of sheaves with monic transition maps. This completes the proof. \square

Remark 3.3. If you wish to work with a left $i\mathcal{A}$ -module \mathcal{U} , you tentatively define $i_+\mathcal{U} = i_*(\mathcal{A}_{X \leftarrow Y} \otimes_{i\mathcal{A}} \mathcal{U})$. One can identify it with

$$i_*((\mathcal{U} \otimes_{\mathcal{O}_Y} \omega_{Y/S}) \otimes_{i\mathcal{B}} \mathcal{B}_{Y \rightarrow X}) \otimes_{\mathcal{O}_X} \omega_{X/S}^\vee$$

without any finiteness conditions by the projection formula for ringed spaces, where $\mathcal{B} := \omega_{X/S}^\vee \otimes_{\mathcal{O}_X} \mathcal{A}^{\text{op}} \otimes_{\mathcal{O}_X} \omega_{X/S}$. The sheaf \mathcal{A}^{op} stands for the opposite tdo ([12] Example 1.1.6).

4 Equivariant structure

In this section, we note equivariant structures on the filtrations. Let K be a smooth S -affine group scheme over S , $i: Y \rightarrow X$ be a K -equivariant immersion of smooth schemes over S , and \mathcal{A} be a K -equivariant tdo on X . Moreover, assume:

Condition 4.1. (i) i is affine;

(ii) there exist an affine open covering $X = \cup_\lambda U_\lambda$ and integers d_λ such that $\mathbb{R}(i|_{i^{-1}(U_\lambda)})_*$ send $D^{\leq 0}((i|_{i^{-1}(U_\lambda)})^{-1}\mathcal{O}_{U_\lambda})$ to $D^{\leq d_\lambda}(\mathcal{O}_{U_\lambda})$ for all indices λ .

Remark 4.2. There exists an integer d such that $\mathbb{R}i_*$ sends $D^{\leq 0}(i^{-1}\mathcal{O}_X)$ to $D^{\leq d}(\mathcal{O}_X)$ if one of the following conditions is satisfied:

(a) i is a closed immersion;

(b) Y is Noetherian of finite Krull dimension.

In fact, a similar argument to [17] Proposition (3.9.2) works if (b) holds. For the vanishing of $H^{m-n}(f^{-1}U, H^n(F))$ there, take the dimension of Y for d , and use [8] Chapter III, Theorem 2.7. Note that for the definition of the cohomology $H^\bullet(Z, \mathcal{F})$ of an \mathcal{R} -module \mathcal{F} on a topological space Z with \mathcal{R} a sheaf of commutative rings on Z , one can take the derived functors on the categories of sheaves of abelian groups and of \mathcal{R} -modules. The resulting abelian groups coincide since both of them can be computed by an \mathcal{R} -linear flasque resolution.

We note that in general, for a flat morphism $f: Z \rightarrow W$ of smooth schemes over a scheme S and a tdo \mathcal{B} on W , the pullback of right \mathcal{B} -modules to right $f\mathcal{B}$ -modules is defined by

$$f^*(- \otimes_{\mathcal{O}_W} \omega_{W/S}^\vee) \otimes_{\mathcal{O}_Z} \omega_{Z/S}.$$

We are now able to define the notion of K -equivariant right \mathcal{A} -modules in a similar way to [12] Definition 3.11.1.

Let \mathcal{V} be a K -equivariant right $i^*\mathcal{A}$ -module. One can prove the following result in the same line as [12] Claim 3.10.8 (1):

Proposition 4.3. *Suppose that we are given a Cartesian diagram of smooth S -schemes*

$$\begin{array}{ccc} \tilde{Y} & \xrightarrow{\tilde{p}} & Y \\ \tilde{i} \downarrow & & \downarrow i \\ \tilde{X} & \xrightarrow{p} & X. \end{array}$$

Assume that the following conditions are satisfied:

- (i) p is isomorphic in the category of morphisms of S -schemes to a projection morphism in the category of smooth S -schemes.
- (ii) i and \tilde{i} satisfy Condition 4.1.

Let p be an integer, and \mathcal{V} be a quasi-coherent right $i^*\mathcal{A}$ -module. Then there is a natural isomorphism

$$p^*G_p i_+ \mathcal{V} \cong G_p \tilde{i}_+ \tilde{p}^* \mathcal{V}.$$

We then obtain a K -equivariant structure on $G_\bullet i_+ \mathcal{V}$ by formal applications of Proposition 4.3:

Proposition 4.4. *For each integer p , $G_p i_+ \mathcal{V}$ is naturally equipped with the structure of a K -invariant quasi-coherent submodule of $i_+ \mathcal{V}$.*

Note that the quasi-coherence follows by a similar argument to [12] Lemma 3.9.6 and Theorem 3.8.4.

The following result is a consequence of [1] (6.7) Theorem and (5.6) Theorem:

Proposition 4.5. *Suppose that x is quasi-compact and quasi-separated. Then for each integer p , $x_* G_p i_+ \mathcal{V} \subset x_* i_+ \mathcal{V}$ is naturally equipped with the structure of a subrepresentation of K (cf. see the beginning of [12] Section 3.12 for the definition of representations of K).*

Remark 4.6. The structure morphism x is quasi-separated if X is locally Noetherian ([3] Corollary 10.24). A typical case in our paper is when S is locally Noetherian since x is locally of finite type ([3] Proposition 10.9).

5 Algebraic properties of $G_\bullet x_* i_+ \mathcal{V}$

In this section, we verify the algebraic properties of the filtered module $G_\bullet x_* i_+ \mathcal{V}$ stated in Section 1.

Let S be a Dedekind scheme, $i : Y \hookrightarrow X$ be a closed immersion of smooth schemes over S , and \mathcal{A} be a tdo on X . Let \mathcal{V} be a right $i^*\mathcal{A}$ -module. Moreover, suppose that the following conditions are satisfied:

- (i) Y is integral.
- (ii) y is proper.
- (iii) \mathcal{V} is torsion-free and coherent as an \mathcal{O}_Y -module.

Theorem 5.1. (1) The \mathcal{O}_S -module $\mathrm{gr}_G^p x_* i_+ \mathcal{V}$ is torsion-free for every integer p .

- (2) For each integer p and a nonempty affine open subscheme $U \subset S$ with coordinate ring k , $\Gamma(U, \mathrm{gr}_G^p x_* i_+ \mathcal{V})$ is finitely generated and projective as a k -module.
- (3) For each integer p , $\mathrm{gr}_G^p x_* i_+ \mathcal{V}$ is finite locally free as an \mathcal{O}_S -module if x is quasi-compact.

Proof. We first study $y_* \mathrm{gr}_G^p(\mathcal{V} \otimes_{i_* \mathcal{A}} \mathcal{A}_{Y \rightarrow X})$. Since \mathcal{V} is a torsion-free (resp. coherent) \mathcal{O}_Y -module, and $\mathcal{N}_{X/Y}$ is a finite locally free \mathcal{O}_Y -module,

$$\mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y}^p \mathcal{N}_{X/Y}$$

is torsion-free (resp. coherent as an \mathcal{O}_Y -module). Hence $y_* \mathrm{gr}_G^p(\mathcal{V} \otimes_{i_* \mathcal{A}} \mathcal{A}_{Y \rightarrow X})$ is torsion-free (resp. coherent) by Corollary 3.2 (1) and [4] Proposition (7.4.5) (resp. and [5] Théorème (3.2.1)).

Part (1) is now obtained by regarding $\mathrm{gr}_G^p x_* i_+ \mathcal{V}$ as an \mathcal{O}_S -submodule of $y_* \mathrm{gr}_G^p(\mathcal{V} \otimes_{i_* \mathcal{A}} \mathcal{A}_{Y \rightarrow X})$. For (2), let U be an arbitrary nonempty affine open subscheme of S with coordinate ring k . Then one can show in a similar way to (1) that $\Gamma(U, \mathrm{gr}_G^p x_* i_+ \mathcal{V})$ is a finitely generated and torsion-free k -module. Since k is a Dedekind domain, $\Gamma(U, \mathrm{gr}_G^p x_* i_+ \mathcal{V})$ is a finitely generated and projective k -module ([3] Proposition B.89 (4)). This shows (2).

Finally, we prove (3). Recall that the filtration $G_\bullet x_* i_+ \mathcal{V}$ consists of quasi-coherent \mathcal{O}_S -modules by Proposition 4.5 (put $K = S$). The assertion now follows from (2). \square

We deduce Corollary 1.3 by applying the following general fact:

Lemma 5.2. Let k be a commutative ring. Let V be a k -module, equipped with an exhaustive filtration $G_\bullet V$. If the associated graded k -modules $\mathrm{gr}_G^p V$ are projective for all p , so is V .

Indeed, $V = \Gamma(X, i_+ \mathcal{V})$ and $G_\bullet V = \Gamma(X, G_\bullet i_+ \mathcal{V})$ satisfy the conditions of Lemma 5.2 under the hypothesis of Corollary 1.3 by Theorem 5.1 and Corollary 3.2 (3).

A Compute $\mathrm{gr}_G i_+ \mathcal{V}$

Recall that F. V. Bien proved an isomorphism $\mathrm{gr}_G i_+ \mathcal{V} \cong i_*(\mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y})$ in [2] if i is a closed immersion. Indeed, this isomorphism holds over general bases if i is a closed immersion since i_* is then exact.

In this section, assume Condition 4.1 and \mathcal{V} to be quasi-coherent. We aim to prove the isomorphism $\mathrm{gr}_G i_+ \mathcal{V} \cong i_*(\mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y})$ under this setting.

Lemma A.1. *Let \mathcal{W} be an $(i\mathcal{A}, i^{-1}\mathcal{O}_X)$ -bimodule. Suppose that \mathcal{W} is flat and quasi-coherent as a left $i\mathcal{A}$ -module. Then $R^p i_*(\mathcal{V} \otimes_{i\mathcal{A}} \mathcal{W})$ vanishes unless $p = 0$.*

Proof. It follows by definition that the p th derived functor vanishes if $p < 0$. We see the assertion for $p > 0$. We may assume the following conditions by the hypothesis (ii) in Condition 4.1:

- (i) X is affine;
- (ii) there exists an integer d such that $\mathbb{R}i_*$ sends $D^{\leq 0}(i^{-1}\mathcal{O}_X)$ to $D^{\leq d}(\mathcal{O}_X)$.

One can see by taking a free resolution $\mathcal{F}^\bullet \rightarrow \mathcal{V} \rightarrow 0$ of right $i\mathcal{A}$ -modules that the proof is completed by showing $R^p i_*(\mathcal{F}^\bullet \otimes_{i\mathcal{A}} \mathcal{W}) = 0$ for $p > 0$ and complexes \mathcal{F}^\bullet of free right $i\mathcal{A}$ -modules concentrated in nonpositive degrees. We may and do assume $\mathcal{F}^\bullet = \mathcal{F}$ is concentrated in degree zero by applying stupid truncations. In this case, the assertion follows from [5] Théorème (1.3.1) since the sheaf $\mathcal{F} \otimes_{i\mathcal{A}} \mathcal{W}$ of abelian groups on Y admits the structure of a quasi-coherent \mathcal{O}_Y -module (see [12] Proposition 2.2.2 and the end of Remark 4.2). \square

Theorem A.2. *There is a canonical isomorphism*

$$\mathrm{gr}_G i_+ \mathcal{V} \cong i_*(\mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y} \mathcal{N}_{X/Y})$$

of \mathcal{O}_X -modules.

Proof. For each integer p , consider the short exact sequence

$$0 \rightarrow \mathcal{V} \otimes_{i\mathcal{A}} G_{p-1} \mathcal{A}_{Y \rightarrow X} \rightarrow \mathcal{V} \otimes_{i\mathcal{A}} G_p \mathcal{A}_{Y \rightarrow X} \rightarrow \mathcal{V} \otimes_{i\mathcal{A}} \mathrm{gr}_G^p \mathcal{A}_{Y \rightarrow X} \rightarrow 0.$$

It gives rise to an exact sequence

$$\begin{aligned} 0 \rightarrow i_*(\mathcal{V} \otimes_{i\mathcal{A}} G_{p-1} \mathcal{A}_{Y \rightarrow X}) \rightarrow i_*(\mathcal{V} \otimes_{i\mathcal{A}} G_p \mathcal{A}_{Y \rightarrow X}) \rightarrow i_*(\mathcal{V} \otimes_{i\mathcal{A}} \mathrm{gr}_G^p \mathcal{A}_{Y \rightarrow X}) \\ \rightarrow R^1 i_*(\mathcal{V} \otimes_{i\mathcal{A}} G_{p-1} \mathcal{A}_{Y \rightarrow X}). \end{aligned}$$

One can see from the proof of Theorem 3.1 that $G_{p-1} \mathcal{A}_{Y \rightarrow X}$ is locally free as a left $i\mathcal{A}$ -module. In particular, $G_{p-1} \mathcal{A}_{Y \rightarrow X}$ is a flat and quasi-coherent left $i\mathcal{A}$ -module. Hence we can apply Lemma A.1 to $\mathcal{W} = G_{p-1} \mathcal{A}_{Y \rightarrow X}$ to show that the first derived functor $R^1 i_*(\mathcal{V} \otimes_{i\mathcal{A}} G_{p-1} \mathcal{A}_{Y \rightarrow X})$ vanishes. The assertion now follows as

$$\mathrm{gr}_G^p i_+ \mathcal{V} \cong i_*(\mathcal{V} \otimes_{i\mathcal{A}} \mathrm{gr}_G^p \mathcal{A}_{Y \rightarrow X}) \cong i_*(\mathcal{V} \otimes_{\mathcal{O}_Y} \mathrm{Sym}_{\mathcal{O}_Y}^p \mathcal{N}_{X/Y})$$

(use Theorem 3.1 for the last isomorphism). \square

References

- [1] A. Altman, B., R. T. Hoobler, and S. L. Kleiman. A note on the base change map for cohomology. *Compositio Math.*, 27:25–38, 1973.

- [2] F. V. Bien. *\mathcal{D} -modules and spherical representations*, volume 39 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1990.
- [3] U. Görtz and T. Wedhorn. *Algebraic geometry I. Schemes*. Springer Studium Mathematik—Master. Springer Spektrum, Wiesbaden, second edition, [2020] ©2020. With examples and exercises.
- [4] A. Grothendieck. Éléments de géométrie algébrique. I. Le langage des schémas. *Inst. Hautes Études Sci. Publ. Math.*, (4):228, 1960.
- [5] A. Grothendieck. Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I. *Inst. Hautes Études Sci. Publ. Math.*, (11):167, 1961.
- [6] M. Harris. Beilinson-Bernstein localization over \mathbb{Q} and periods of automorphic forms. *Int. Math. Res. Not. IMRN*, (9):2000–2053, 2013.
- [7] M. Harris. Beilinson-Bernstein localization over \mathbb{Q} and periods of automorphic forms: erratum. *Int. Math. Res. Not. IMRN*, (3):957–960, 2020.
- [8] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [9] T. Hayashi. Flat base change formulas for (\mathfrak{g}, K) -modules over Noetherian rings. *J. Algebra*, 514:40–75, 2018.
- [10] T. Hayashi. Dg analogues of the Zuckerman functors and the dual Zuckerman functors I. *J. Algebra*, 540:274–305, 2019.
- [11] T. Hayashi. Half-integrality of line bundles on partial flag schemes of classical Lie groups, 2021. arXiv:2104.13304.
- [12] T. Hayashi and F. Januszewski. Families of twisted \mathcal{D} -modules and arithmetic models of Harish-Chandra modules, 2018. arXiv:1808.10709.
- [13] H. Hecht, D. Miličić, W. Schmid, and J. A. Wolf. Localization and standard modules for real semisimple Lie groups. I. The duality theorem. *Invent. Math.*, 90(2):297–332, 1987.
- [14] F. Januszewski. Rational structures on automorphic representations. *Math. Ann.*, 370(3-4):1805–1881, 2018.
- [15] F. Januszewski. On period relations for automorphic L -functions I. *Trans. Amer. Math. Soc.*, 371(9):6547–6580, 2019.
- [16] S. N. Kitchen. Cohomology of standard modules on partial flag varieties. *Represent. Theory*, 16:317–344, 2012.
- [17] J. Lipman. Notes on derived functors and Grothendieck duality. In *Foundations of Grothendieck duality for diagrams of schemes*, volume 1960 of *Lecture Notes in Math.*, pages 1–259. Springer, Berlin, 2009.

- [18] Y. Oshima. On the restriction of Zuckerman's derived functor modules $A_q(\lambda)$ to reductive subgroups. *Amer. J. Math.*, 137(4):1099–1138, 2015.
- [19] G. Shimura. Sur les intégrales attachées aux formes automorphes. *J. Math. Soc. Japan*, 11:291–311, 1959.