

**PROBABILITY OF GENERATION BY RANDOM
PERMUTATIONS OF GIVEN CYCLE TYPE**

SEAN EBERHARD AND DANIELE GARZONI

ABSTRACT. Suppose π and π' are two random elements of S_n with constrained cycle types such that π has $xn^{1/2}$ fixed points and $yn/2$ two-cycles, and likewise π' has $x'n^{1/2}$ fixed points and $y'n/2$ two-cycles. We show that the events that $G = \langle \pi, \pi' \rangle$ is transitive and $G \geq A_n$ both have probability approximately

$$(1 - yy')^{1/2} \exp\left(-\frac{xx' + \frac{1}{2}x^2y' + \frac{1}{2}x'^2y}{1 - yy'}\right),$$

provided (x, x') is not close to $(0, \infty)$ or $(\infty, 0)$. This formula is derived from some preliminary results in a recent paper of the authors. As an application, we show that two uniformly random elements of uniformly random conjugacy classes of S_n generate the group with probability about 51%.

1. INTRODUCTION

Consider a sequence of pairs of conjugacy classes $\mathcal{C}_n, \mathcal{C}'_n$ of S_n (one pair for each positive integer n). Let c_i and c'_i be the number of i -cycles of an element of \mathcal{C}_n and \mathcal{C}'_n , respectively. We are interested in the group G generated by uniformly random $\pi \in \mathcal{C}_n$ and $\pi' \in \mathcal{C}'_n$.

Theorem 1.1. *Assume*

$$\begin{aligned} c_1/n^{1/2} &\rightarrow x, & 2c_2/n &\rightarrow y, \\ c'_1/n^{1/2} &\rightarrow x', & 2c'_2/n &\rightarrow y', \end{aligned}$$

where $0 \leq x, x' \leq \infty$ and $0 \leq y, y' \leq 1$. If $yy' < 1$, assume $(x, x') \notin \{(0, \infty), (\infty, 0)\}$. Then

$$\mathbf{P}(G \geq A_n) \rightarrow (1 - yy')^{1/2} \exp\left(-\frac{xx' + \frac{1}{2}x^2y' + \frac{1}{2}x'^2y}{1 - yy'}\right). \quad (1)$$

The right-hand side is interpreted (continuously) as zero if $y = y' = 1$ or $x = \infty$ or $x' = \infty$. The same estimate holds for $\mathbf{P}(G \text{ transitive})$.

This theorem essentially generalizes [1, Theorem 1.1], which describes the boundary cases of (1). We will deduce Theorem 1.1 from some of the preliminary results of [1].

Remark 1.2. In the indeterminate case $(x, x') \in \{(0, \infty), (\infty, 0)\}$, $\mathbf{P}(G \geq A_n)$ can be close to 0 or to 1. Some examples are given in [1, Section 2.5].

2020 *Mathematics Subject Classification.* 20B30, 20P05.

Key words and phrases. symmetric group, random generation.

2. PROOF

By [1, Theorem 1.1(2)] we may assume $x+x' < \infty$ and $yy' < 1$. By [1, Lemma 4.5 and Lemma 4.9], $\mathbf{P}(G \text{ transitive})$ and $\mathbf{P}(G \geq A_n)$ differ by $e^{-\Omega(n)}$, so we may focus on $\mathbf{P}(G \text{ transitive})$. As in [1, Section 3] let N_k be the number of orbits of G of size k and let

$$N = \sum_{k=1}^{\lfloor n/2 \rfloor} N_k.$$

Then G is transitive if and only if $N = 0$. By [1, Lemma 3.3 and Lemma 3.4], $\mathbf{E}N_k \leq \epsilon_k$ for some constant ϵ_k such that $\sum_k \epsilon_k < \infty$, which allows us to apply the dominated convergence theorem. Now fix k . By [1, (3)],

$$\mathbf{E}N_k = \sum \binom{n}{k}^{-1} \binom{c_1}{d_1} \cdots \binom{c_k}{d_k} \binom{c'_1}{d'_1} \cdots \binom{c'_k}{d'_k} p(d_1, \dots, d_k; d'_1, \dots, d'_k),$$

where the sum goes over all solutions to

$$\begin{aligned} k &= d_1 1 + d_2 2 + \cdots + d_k k \\ k &= d'_1 1 + d'_2 2 + \cdots + d'_k k \end{aligned}$$

such that $0 \leq d_i \leq c_i$ and $0 \leq d'_i \leq c'_i$ for each i , and where $p(d_1, \dots, d_k; d'_1, \dots, d'_k)$ is the probability that random permutations $\tau, \tau' \in S_k$ with cycle types $1^{d_1} \cdots k^{d_k}$ and $1^{d'_1} \cdots k^{d'_k}$ generate a transitive subgroup. Write

$$\mathbf{E}N_k = \Sigma_1 + \Sigma_2,$$

where Σ_1 is the sum of all terms with $d_i = d'_i = 0$ for each $i \geq 3$ and Σ_2 is the sum of all other terms. If $d_i + d'_i > 0$ for any $i \geq 3$ then

$$\binom{c_1}{d_1} \cdots \binom{c_k}{d_k} \binom{c'_1}{d'_1} \cdots \binom{c'_k}{d'_k} \leq O_k(n^{d_1/2+d_2+\cdots+d_k} n^{d'_1/2+d'_2+\cdots+d'_k}) \leq O_k(n^{k-1/2}).$$

Hence

$$\Sigma_2 \leq O_k(n^{-1/2}).$$

Now consider Σ_1 . We have $p(d_1, d_2; d'_1, d'_2) = 0$ unless $d_1 + d'_1 \leq 2$, so if $k = 2m + 1$ is odd we have

$$\Sigma_1 = \binom{n}{k}^{-1} \binom{c_1}{1} \binom{c_2}{m} \binom{c'_1}{1} \binom{c'_2}{m} p(1, m; 1, m),$$

while if $k = 2m$ is even then Σ_1 is the sum of three terms:

$$\begin{aligned} \Sigma_1 &= \binom{n}{k}^{-1} \binom{c_1}{2} \binom{c_2}{m-1} \binom{c'_2}{m} p(2, m-1; 0, m) \\ &\quad + \binom{n}{k}^{-1} \binom{c_2}{m} \binom{c'_1}{2} \binom{c'_2}{m-1} p(0, m; 2, m-1) \\ &\quad + \binom{n}{k}^{-1} \binom{c_2}{m} \binom{c'_2}{m} p(0, m; 0, m). \end{aligned}$$

The three calculations

$$\begin{aligned} p(1, m; 1, m) &= \frac{(2m+1)!}{((2m+1)!/2^m m!)^2}, \\ p(2, m-1; 0, m) &= \frac{(2m)!/2}{((2m)!/2^{m-1} 2!(m-1)!) \times ((2m)!/2^m m!)}, \\ p(0, m; 0, m) &= \frac{(2m-1)!}{((2m)!/2^m m!)^2} \end{aligned}$$

can be left to the reader (cf. [1, Lemma 2.1]). Hence, as $n \rightarrow \infty$ with k fixed,

$$\Sigma_1 \rightarrow \begin{cases} xy^m x' y'^m & : k = 2m + 1 > 0, \\ (x^2/y + x'^2/y') y^m y'^m / 2 + y^m y'^m / (2m) & : k = 2m > 0. \end{cases}$$

It follows by the dominated convergence theorem that

$$\begin{aligned} \mathbf{E}N &= \sum_{k=1}^{\lfloor n/2 \rfloor} \mathbf{E}N_k \\ &\rightarrow \sum_{k=2m+1 > 0} xx'(yy')^m + \sum_{k=2m > 0} \left((x^2/y + x'^2/y')(yy')^m / 2 + (yy')^m / (2m) \right) \\ &= (xx' + x^2 y' / 2 + x'^2 y / 2) / (1 - yy') - \frac{1}{2} \log(1 - yy'). \end{aligned}$$

Hence, by [1, Theorem 3.10],

$$\mathbf{P}(N = 0) = e^{-\mathbf{E}N} + o(1) \rightarrow (1 - yy')^{1/2} \exp\left(-\frac{xx' + \frac{1}{2}x^2 y' + \frac{1}{2}x'^2 y}{1 - yy'}\right).$$

This completes the proof.

3. APPLICATION

Let \mathcal{C}_n and \mathcal{C}'_n be uniformly random conjugacy classes of S_n , and let $\pi \in \mathcal{C}_n$ and $\pi' \in \mathcal{C}'_n$ be uniformly random. Let $G = \langle \pi, \pi' \rangle$. What is the probability that $G \geq A_n$? The Hardy–Ramanujan asymptotic for the partition function states that

$$p(n) \sim \frac{a}{n} \exp(2bn^{1/2}),$$

where $a = 1/(4\sqrt{3})$ and $b = \pi/\sqrt{6}$ (see [2, Proposition VIII.6, p. 578]). Since the cycle type of π is that of a uniformly random partition of n , it follows that

$$\begin{aligned} \mathbf{P}(c_1 \geq xn^{1/2}, c_2 \geq yn^{1/2}) &\sim \frac{n}{n - (x+2y)n^{1/2}} \exp(2b(n - (x+2y)n^{1/2})^{1/2} - 2bn^{1/2}) \\ &\rightarrow \exp(-b(x+2y)). \end{aligned}$$

In particular note that $c_2 = o(n)$ with high probability. Hence Theorem 1.1 implies that $\mathbf{P}(G \geq A_n)$ and $\mathbf{P}(G \text{ transitive})$ are both asymptotically

$$b^2 \int_0^\infty \int_0^\infty \exp(-xx' - b(x+x')) dx dx' = b^2 e^{b^2} \int_{b^2}^\infty t^{-1} e^{-t} dt \approx 0.6889.$$

A random conjugacy class consists of even permutations with probability tending to 1/2. Hence $\mathbf{P}(G = A_n) \approx 0.1722$ and $\mathbf{P}(G = S_n) \approx 0.5167$.

REFERENCES

- [1] Sean Eberhard and Daniele Garzoni, *Random generation with cycle type restrictions*, *Algebr. Comb.* **4** (2021), no. 1, 1–25, <https://doi.org/10.5802/alco.149>. [1](#), [1.2](#), [2](#)
- [2] Philippe Flajolet and Robert Sedgewick, *Analytic combinatorics*, Cambridge University Press, Cambridge, 2009, <https://doi.org/10.1017/CB09780511801655>. [3](#)

CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB (U.K.)
Email address: `eberhard@maths.cam.ac.uk`

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 69978 (ISRAEL)
Email address: `danieleg@mail.tau.ac.il`