

GENERALIZED MULTIPLICATIVE HILBERT OPERATORS

JIANJUN JIN

ABSTRACT. In this note, by introducing some parameters, we define and study some generalized multiplicative Hilbert operators. We provide a sufficient and necessary condition for the boundedness of these operators in terms of the parameters. Also, for certain special cases, we determine the exact values of the norms of the generalized multiplicative Hilbert operators.

1. INTRODUCTION AND MAIN RESULTS

Let $p > 1$. We use p' to denote the conjugate of p , i.e., $\frac{1}{p} + \frac{1}{p'} = 1$. In a recent work [3], Brevig et al. studied the following operator

$$\mathbf{H}_p(a)(m) := \frac{1}{m^{\frac{1}{p}}} \sum_{n=2}^{\infty} \frac{a_n}{n^{\frac{1}{p'}} (\log mn)}, \quad a = \{a_n\}_{n=2}^{\infty} \in l^p, \quad m \geq 2,$$

which is induced by the *multiplicative Hilbert matrix*

$$\mathbf{M}_p := \left(\frac{1}{n^{\frac{1}{p'}} m^{\frac{1}{p}} (\log mn)} \right)_{m, n \geq 2}.$$

Here, l^p is the space of sequences of complex numbers, i.e.,

$$l^p := \left\{ a = \{a_n\}_{n=2}^{\infty} : \|a\|_p := \left(\sum_{n=2}^{\infty} |a_n|^p \right)^{\frac{1}{p}} < \infty \right\}.$$

The multiplicative Hilbert matrix \mathbf{M}_p is related to the study of the theory of Hardy spaces of Dirichlet series and Riemann zeta function. See [2], [3], [4], [6] for some recent results of this topic. We will call \mathbf{H}_p *multiplicative Hilbert operator*.

It was proved in [3] that

Theorem 1.1. *Let $p > 1$. Then \mathbf{H}_p is bounded on l^p and the norm of \mathbf{H}_p is $\pi \csc\left(\frac{\pi}{p}\right)$.*

In this note, we will study the *generalized multiplicative Hilbert operator* $\mathbf{H}_{p, \alpha, \beta, \gamma}^{\mu, \nu}$, which is defined as, for $a = \{a_n\}_{n=2}^{\infty} \in l^p$,

$$\mathbf{H}_{p, \alpha, \beta, \gamma}^{\mu, \nu}(a)(m) := \frac{(\log m)^{\frac{1}{p}[(\alpha-1)+\alpha\mu]}}{m^{\frac{1}{p}}} \sum_{n=2}^{\infty} \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{n^{\frac{1}{p'}} [(\log m)^\alpha + (\log n)^\beta]^\gamma} a_n, \quad m \geq 2,$$

where $\gamma > 0$, $0 < \alpha, \beta \leq 1$, $-1 < \mu, \nu < p - 1$.

2010 *Mathematics Subject Classification.* 26D15; 47A30.

Key words and phrases. Generalized multiplicative Hilbert operators; boundedness of operator; norm of operator.

The author was supported by National Natural Science Foundation of China(Grant Nos. 11501157).

When $\alpha = \beta = \gamma = 1$, $\mu = \nu = 0$, $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ reduces to the multiplicative Hilbert operator \mathbf{H}_p . $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ corresponds to the following *generalized multiplicative Hilbert matrix*

$$\mathbf{M}_{p,\alpha,\beta,\gamma}^{\mu,\nu} := \left(\frac{(\log m)^{\frac{1}{p}[(\alpha-1)+\alpha\mu]} \cdot (\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{m^{\frac{1}{p}} \cdot [(\log m)^\alpha + (\log n)^\beta]^\gamma \cdot n^{\frac{1}{p'}}} \right)_{m,n \geq 2}.$$

As our first main result of this paper, we will give a sufficient and necessary condition for the boundedness of $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ in terms of the parameters γ, μ, ν . We will prove that

Theorem 1.2. *Let $p > 1$, $\gamma > 0$, $0 < \alpha, \beta \leq 1$, $-1 < \mu, \nu < p-1$, and $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ be as above. Then $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ is bounded on l^p if and only if $p(\gamma-1) - (\mu-\nu) \geq 0$.*

Next, we consider the case when $p(\gamma-1) - (\mu-\nu) = 0$, i.e., $\gamma = 1 + \frac{\mu-\nu}{p}$. In this case, we denote by $\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}$ the operator instead of $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$. That is, for $a = \{a_n\}_{n=2}^\infty \in l^p$,

$$\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}(a)(m) := \frac{(\log m)^{\frac{1}{p}[(\alpha-1)+\alpha\mu]}}{m^{\frac{1}{p}}} \sum_{n=2}^\infty \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{n^{\frac{1}{p'} [(\log m)^\alpha + (\log n)^\beta]^{1+\frac{\mu-\nu}{p}}}} a_n, \quad m \geq 2.$$

We denote by $\|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}\|$ the norm of $\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}$. We will show the following result, which is an extension of Theorem 1.1.

Theorem 1.3. *Let $p > 1$, $0 < \alpha, \beta \leq 1$, $-1 < \mu, \nu < p-1$, and $\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}$ be as above. Then $\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}$ is bounded on l^p and*

$$(1.1) \quad \|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}\| = \frac{1}{\alpha^{\frac{1}{p}} \beta^{\frac{1}{p'}}} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right).$$

Here $B(\cdot, \cdot)$ is the well-known Beta function, which is defined as

$$B(u, v) = \int_0^\infty \frac{t^{u-1}}{(1+t)^{u+v}} dt, \quad u > 0, v > 0.$$

It is known that

$$B(u, v) = \int_0^1 t^{u-1} (1-t)^{v-1} dt = \frac{\Gamma(u)\Gamma(v)}{\Gamma(u+v)},$$

where $\Gamma(x)$ is the Gamma function, defined as

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt, \quad x > 0.$$

See [1] for more detailed introductions to these special functions.

The paper is organized as follows. We will first prove Theorem 1.3 in the next section. The proof of Theorem 1.2 will be give in Section 3. Some remarks will be finally present in the Section 4.

2. PROOF OF THEOREM 1.3

In the proof of Theorem 1.3, we need the following

Lemma 2.1. *Let $p > 1$, $0 < \alpha, \beta \leq 1$, $-1 < \mu, \nu < p - 1$. We define*

$$E(m) := \sum_{n=2}^{\infty} \frac{(\log n)^{\beta-1}}{n[(\log m)^{\alpha} + (\log n)^{\beta}]^{1+\frac{\mu-\nu}{p}}} \cdot \frac{(\log m)^{\frac{\alpha(1+\mu)}{p}}}{(\log n)^{\frac{\beta(1+\nu)}{p}}}, \quad m \geq 2;$$

$$F(n) := \sum_{m=2}^{\infty} \frac{(\log m)^{\alpha-1}}{m[(\log m)^{\alpha} + (\log n)^{\beta}]^{1+\frac{\mu-\nu}{p}}} \cdot \frac{(\log n)^{\frac{\beta(p-1-\nu)}{p}}}{(\log m)^{\frac{\alpha(p-1-\mu)}{p}}}, \quad n \geq 2.$$

Then we have

$$(2.1) \quad E(m) \leq \frac{1}{\beta} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right), \quad m \geq 2,$$

$$(2.2) \quad F(n) \leq \frac{1}{\alpha} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right), \quad n \geq 2.$$

Proof. In view of the assumptions $0 < \alpha, \beta \leq 1$, $-1 < \mu, \nu < p - 1$, we see that, for $m \geq 2$,

$$E(m) \leq \int_1^{\infty} \frac{(\log x)^{\beta-1}}{n[(\log m)^{\alpha} + (\log x)^{\beta}]^{1+\frac{\mu-\nu}{p}}} \cdot \frac{(\log m)^{\frac{\alpha(1+\mu)}{p}}}{(\log x)^{\frac{\beta(1+\nu)}{p}}} dx.$$

Consequently, by a change of variables $s = (\log x)^{\beta}$, we have

$$\begin{aligned} E(m) &\leq \frac{1}{\beta} \int_0^{\infty} \frac{1}{[(\log m)^{\alpha} + s]^{1+\frac{\mu-\nu}{p}}} \cdot \frac{(\log m)^{\frac{\alpha(1+\mu)}{p}}}{s^{\frac{1+\nu}{p}}} ds \\ &= \frac{1}{\beta} \int_0^{\infty} \frac{t^{-\frac{1+\nu}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt = \frac{1}{\beta} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right). \end{aligned}$$

By the similar way, we can obtain (2.2). The lemma is proved. \square

We start to prove Theorem 1.3. For $a = \{a_n\}_{n=2}^{\infty} \in l^p$, $m \geq 2$, we write

$$\begin{aligned} &\frac{(\log m)^{\frac{1}{p}[(\alpha-1)+\alpha\mu]}}{m^{\frac{1}{p}}} \left| \sum_{n=2}^{\infty} \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{n^{\frac{1}{p'}} [(\log m)^{\alpha} + (\log n)^{\beta}]^{1+\frac{\mu-\nu}{p}}} a_n \right| \\ &= \sum_{n=2}^{\infty} \left\{ [K(m, n)]^{\frac{1}{p}} P_1(m, n) \cdot [K(m, n)]^{\frac{1}{p'}} P_2(m, n) \right\} := I(m), \end{aligned}$$

where

$$\begin{aligned} K(m, n) &= \frac{1}{[(\log m)^{\alpha} + (\log n)^{\beta}]^{1+\frac{\mu-\nu}{p}}}; \\ P_1(m, n) &= \frac{(\log n)^{\frac{\beta(1+\nu)}{pp'} - \frac{\beta\nu}{p}}}{(\log m)^{\frac{\alpha(p-1-\mu)}{p^2}}} \cdot \frac{(\log m)^{\frac{1}{p}(\alpha-1)}}{m^{\frac{1}{p}}} \cdot |a_n|; \\ P_2(m, n) &= \frac{(\log m)^{\frac{\alpha(p-1-\mu)}{p^2} + \frac{\alpha\mu}{p}}}{(\log n)^{\frac{\beta(1+\nu)}{pp'}}} \cdot \frac{(\log n)^{\frac{1}{p'}(\beta-1)}}{n^{\frac{1}{p'}}}. \end{aligned}$$

Applying the Hölder's inequality on $I(m)$, we get from (2.1) that

$$\begin{aligned} I(m) &\leq \left[\sum_{n=2}^{\infty} K(m, n)[P_1(m, n)]^p \right]^{\frac{1}{p}} \left[\sum_{n=2}^{\infty} K(m, n)[P_2(m, n)]^{p'} \right]^{\frac{1}{p'}} \\ &= [E(m)]^{\frac{1}{p'}} \left[\sum_{n=2}^{\infty} K(m, n)[P_1(m, n)]^p \right]^{\frac{1}{p}}. \end{aligned}$$

It follows from (2.2) that

$$\begin{aligned} \|\tilde{\mathbf{H}}_{p, \alpha, \beta}^{\mu, \nu} a\|_p &= \left[\sum_{m=2}^{\infty} I^p(m) \right]^{\frac{1}{p}} \\ &\leq \frac{1}{\beta^{\frac{1}{p'}}} \left[B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right) \right]^{\frac{1}{p'}} \left[\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} K(m, n)[P_1(m, n)]^p \right]^{\frac{1}{p}} \\ &= \frac{1}{\beta^{\frac{1}{p'}}} \left[B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right) \right]^{\frac{1}{p'}} \left[\sum_{n=2}^{\infty} F(n)|a_n|^p \right]^{\frac{1}{p}} \\ &\leq \frac{1}{\alpha^{\frac{1}{p}} \beta^{\frac{1}{p'}}} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right) \|a\|_p. \end{aligned}$$

This means that $\tilde{\mathbf{H}}_{p, \alpha, \beta}^{\mu, \nu}$ is bounded on l^p and

$$(2.3) \quad \|\tilde{\mathbf{H}}_{p, \alpha, \beta}^{\mu, \nu}\| \leq \frac{1}{\alpha^{\frac{1}{p}} \beta^{\frac{1}{p'}}} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right).$$

For $\varepsilon > 0$, we take $\tilde{a} = \{\tilde{a}_n\}_{n=2}^{\infty}$ with $\tilde{a}_n = \varepsilon^{\frac{1}{p}} n^{-\frac{1}{p}} (\log n)^{-\frac{1+\beta\varepsilon}{p}}$. On one hand, we have

$$\|\tilde{a}\|_p^p = \varepsilon \sum_{n=2}^{\infty} n^{-1} (\log n)^{-1-\beta\varepsilon} \geq \varepsilon \int_2^{\infty} x^{-1} (\log x)^{-1-\beta\varepsilon} dx = \frac{1}{\beta(\log 2)^{\beta\varepsilon}}.$$

On the other hand, we have

$$\|\tilde{a}\|_p^p = \frac{\varepsilon}{2(\log 2)^{1+\beta\varepsilon}} + \sum_{n=3}^{\infty} n^{-1} (\log n)^{-1-\beta\varepsilon} \leq \frac{\varepsilon}{2(\log 2)^{1+\beta\varepsilon}} + \frac{1}{\beta(\log 2)^{\beta\varepsilon}}.$$

Thus, we have

$$(2.4) \quad \|\tilde{a}\|_p^p = \frac{1}{\beta}(1 + o(1)), \quad \varepsilon \rightarrow 0^+.$$

We write

$$(2.5) \quad \|\tilde{\mathbf{H}}_{p, \alpha, \beta}^{\mu, \nu} \tilde{a}\|_p^p = \varepsilon \sum_{m=2}^{\infty} \frac{(\log m)^{(\alpha-1)+\alpha\mu}}{m} \cdot [J(m)]^p.$$

Here

$$J(m) := \sum_{n=2}^{\infty} \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]} \cdot (\log n)^{-\frac{1+\beta\varepsilon}{p}}}{n[(\log m)^\alpha + (\log n)^\beta]^{1+\frac{\mu-\nu}{p}}}.$$

In view of the assumptions $0 < \beta \leq 1, -1 < \nu < p-1$, we have $1 + \beta\nu \geq 0$. Hence we get that

$$\frac{1}{p'}[(\beta-1) - (p'-1)\beta\nu] - \frac{1+\beta\varepsilon}{p} = \frac{1}{p'}(\beta-1) - \frac{1+\beta\nu+\beta\varepsilon}{p} < 0.$$

Consequently,

$$\begin{aligned}
(2.6) \quad J(m) &\geq \int_2^\infty \frac{(\log x)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]} \cdot (\log x)^{-\frac{1+\beta\varepsilon}{p}}}{n[(\log m)^\alpha + (\log x)^\beta]^{1+\frac{\mu-\nu}{p}}} dx \\
&= \frac{1}{\beta} \int_{\log 2}^\infty \frac{s^{-\frac{1+\nu+\varepsilon}{p}}}{[(\log m)^\alpha + s]^{1+\frac{\mu-\nu}{p}}} ds \\
&= \frac{1}{\beta} (\log m)^{-\frac{\alpha}{p}(1+\mu+\varepsilon)} \int_{\frac{\log 2}{(\log m)^\alpha}}^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt
\end{aligned}$$

Also, for $\varepsilon < p - 1 - \nu$, we have

$$\begin{aligned}
(2.7) \quad \int_{\frac{\log 2}{(\log m)^\alpha}}^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} &= \int_0^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt - \int_0^{\frac{\log 2}{(\log m)^\alpha}} \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt \\
&= B\left(\frac{1+\mu}{p} + \frac{\varepsilon}{p}, \frac{p-1-\nu}{p} - \frac{\varepsilon}{p}\right) - \int_0^{\frac{\log 2}{(\log m)^\alpha}} \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt. \\
&:= L(\varepsilon) - Q(m).
\end{aligned}$$

Combining (2.5), (2.6) and (2.7), we get that

$$(2.8) \quad \|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu} \tilde{a}\|_p^p \geq \frac{\varepsilon}{\beta p} \sum_{m=2}^\infty \frac{(\log m)^{-1-\alpha\varepsilon}}{m} \cdot [L(\varepsilon) - Q(m)]^p$$

By using the Bernoulli's inequality(see [5]), we obtain that

$$(2.9) \quad [L(\varepsilon) - Q(m)]^p \geq [L(\varepsilon)]^p \left[1 - \frac{p}{L(\varepsilon)} \int_0^{\frac{\log 2}{(\log m)^\alpha}} \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt\right]^p.$$

We also note that

$$(2.10) \quad \varepsilon \sum_{m=2}^\infty \frac{(\log m)^{-1-\alpha\varepsilon}}{m} = \frac{1}{\alpha} (1 + o(1)), \quad \varepsilon \rightarrow 0^+,$$

and, for $\varepsilon < p - 1 - \nu$,

$$\begin{aligned}
(2.11) \quad \sum_{m=2}^\infty \frac{(\log m)^{-1-\alpha\varepsilon}}{m} \int_0^{\frac{\log 2}{(\log m)^\alpha}} \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^{1+\frac{\mu-\nu}{p}}} dt \\
\leq \sum_{m=2}^\infty \frac{(\log m)^{-1-\alpha\varepsilon}}{m} \int_0^{\frac{\log 2}{(\log m)^\alpha}} t^{-\frac{1+\nu+\varepsilon}{p}} dt \\
= \frac{p(\log 2)^{\frac{p-1-\nu-\varepsilon}{p}}}{p-1-\nu-\varepsilon} \sum_{m=2}^\infty \frac{(\log m)^{-1-\alpha(\frac{\varepsilon}{p} + \frac{p-1-\nu}{p})}}{m} = O(1), \quad \varepsilon \rightarrow 0^+.
\end{aligned}$$

It follows from (2.8)-(2.11) that

$$\|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu} \tilde{a}\|_p^p \geq \frac{1}{\alpha\beta p} (1 + o(1)) \cdot [L(\varepsilon)]^p \cdot [1 - \varepsilon O(1)].$$

Hence, from (2.4), we get that

$$\|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu} \tilde{a}\| \geq \frac{\|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu} \tilde{a}\|_p}{\|\tilde{a}\|_p} \geq \frac{\frac{1}{\alpha\beta p} (1 + o(1)) \cdot [L(\varepsilon)] \cdot [1 - \varepsilon O(1)]^{\frac{1}{p}}}{\frac{1}{\beta^{\frac{1}{p}}} (1 + o(1))}.$$

Take $\varepsilon \rightarrow 0^+$, we see that

$$(2.12) \quad \|\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}\| \geq \frac{1}{\alpha^{\frac{1}{p}} \beta^{\frac{1}{p'}}} B\left(\frac{1+\mu}{p}, \frac{p-1-\nu}{p}\right).$$

Combining (2.3) and (2.12), we see that (1.1) is true and the proof of Theorem 1.3 is finished.

3. PROOF OF THEOREM 1.2

We first prove the "if" part. If $p(\gamma-1) + (\mu-\nu) \geq 0$, that is $\gamma \geq 1 + \frac{\mu-\nu}{p}$, then, for $a = \{a_n\}_{n=2}^\infty \in l^p$, $m \geq 2$, it is easy to see that

$$\left| \sum_{n=2}^\infty \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{n^{\frac{1}{p'}[(\log m)^\alpha + (\log n)^\beta]^\gamma}} a_n \right| \leq \left| \sum_{n=2}^\infty \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]}}{n^{\frac{1}{p'}[(\log m)^\alpha + (\log n)^\beta]^{1+\frac{\mu-\nu}{p}}}} a_n \right|.$$

Consequently, in view of the boundedness of $\tilde{\mathbf{H}}_{p,\alpha,\beta}^{\mu,\nu}$, we conclude that $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ is bounded on l^p when $p(\gamma-2) + (\mu-\nu) \geq 0$.

Next, we prove the "only if" part. We will show that, if $p(\gamma-1) + (\mu-\nu) < 0$, then $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ can not be bounded on l^p .

Actually, let $\varepsilon > 0$, we take, as is done in the proof of Theorem 1.3, $\tilde{a} = \{\tilde{a}_n\}_{n=2}^\infty$ with $\tilde{a}_n = \varepsilon^{\frac{1}{p}} n^{-\frac{1}{p}} (\log n)^{-\frac{1+\beta\varepsilon}{p}}$. We have

$$\|\tilde{a}\|_p^p = \frac{1}{\beta} (1 + o(1)), \quad \varepsilon \rightarrow 0^+.$$

Hence we have

$$\begin{aligned} \|\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu} \tilde{a}\|_p^p &= \sum_{m=2}^\infty \frac{(\log m)^{(\alpha-1)+\alpha\mu}}{m^{\frac{1}{p}}} \left[\sum_{n=2}^\infty \frac{(\log n)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]} \cdot (\log n)^{-\frac{1+\beta\varepsilon}{p}}}{n[(\log m)^\alpha + (\log n)^\beta]^\gamma} \right]^p \\ &:= \sum_{m=2}^\infty \frac{(\log m)^{(\alpha-1)+\alpha\mu}}{m} \cdot [R(m)]^p. \end{aligned}$$

On the other hand, we have, for $m \geq 2$,

$$\begin{aligned} R(m) &\geq \int_2^\infty \frac{(\log x)^{\frac{1}{p'}[(\beta-1)-(p'-1)\beta\nu]} \cdot (\log x)^{-\frac{1+\beta\varepsilon}{p}}}{n[(\log m)^\alpha + (\log x)^\beta]^\gamma} dx \\ &= \frac{1}{\beta} \int_{\log 2}^\infty \frac{s^{-\frac{1+\nu+\varepsilon}{p}}}{[(\log m)^\alpha + s]^\gamma} ds \\ &= \frac{1}{\beta} (\log m)^{-\frac{\alpha}{p}[p(\gamma-1)+(1+\nu+\varepsilon)]} \int_{\frac{\log 2}{(\log m)^\alpha}}^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^\gamma} dt \\ &\geq \frac{1}{\beta} (\log m)^{-\frac{\alpha}{p}[p(\gamma-1)+(1+\nu+\varepsilon)]} \int_{(\log 2)^{1-\alpha}}^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^\gamma} dt. \end{aligned}$$

Consequently,

$$\|\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu} \tilde{a}\|_p^p \geq \frac{1}{\beta^p} \left[\int_{(\log 2)^{1-\alpha}}^\infty \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^\gamma} dt \right]^p \left[\sum_{m=2}^\infty \frac{(\log m)^{\alpha[p(\gamma-1)+(\mu-\nu)-\varepsilon]-1}}{m} \right].$$

We suppose $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu} : l^p \rightarrow l^p$ is bounded. Then there exists a constant $C > 0$ such that

$$(3.1) \quad \begin{aligned} C &\geq \frac{\|\tilde{\mathbf{H}}_{p,\alpha,\beta,\gamma}^{\mu,\nu} \tilde{a}\|_p^p}{\|\tilde{a}\|_p^p} \\ &\geq \frac{1}{\beta^{p-1}} (1 + o(1)) \left[\int_{(\log 2)^{1-\alpha}}^{\infty} \frac{t^{-\frac{1+\nu+\varepsilon}{p}}}{(1+t)^\gamma} dt \right]^p \\ &\quad \times \left[\sum_{m=2}^{\infty} \frac{(\log m)^{\alpha[p(1-\gamma)+(\mu-\nu)-\varepsilon]-1}}{m^{\frac{1}{p}}} \right]. \end{aligned}$$

But, when $\varepsilon < p(1-\gamma) + (\mu-\nu)$, we see from $p(1-\gamma) + (\mu-\nu) - \varepsilon := \theta > 0$ that

$$\sum_{m=2}^{\infty} \frac{(\log m)^{\alpha[p(1-\gamma)+(\mu-\nu)-\varepsilon]-1}}{m} = \sum_{m=2}^{\infty} \frac{(\log m)^{\theta\alpha-1}}{m} = +\infty.$$

Hence we get that (3.1) is a contradiction. This proves that $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ can not be bounded on l^p , if $p(\gamma-1) + (\mu-\nu) < 0$. The theorem is proved.

4. REMARKS

Remark 4.1. We first remark that the assumptions $-1 < \mu, \nu < p-1$ in Theorem 1.2 and 1.3 are both necessary.

We consider the case when $\alpha = \beta = \gamma = 1, \mu = \nu := \delta$. That is to say, we will consider the operator

$$\mathbf{H}_{p,1,1,1}^{\delta,\delta}(a)(m) = \frac{(\log m)^{\frac{\delta}{p}}}{m^{\frac{1}{p}}} \sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\delta}{p}}}{n^{\frac{1}{p}} \log(mn)} a_n, \quad a = \{a_n\}_{n=2}^{\infty} \in l^p, \quad m \geq 2.$$

We next write $\widehat{\mathbf{H}}_p^\delta$ to denote $\mathbf{H}_{p,1,1,1}^{\delta,\delta}$. We will show that

Claim 4.2. $\widehat{\mathbf{H}}_p^\delta$ is not bounded on l^p , if $\delta \leq -1$, or $\delta \geq p-1$.

Proof. For $\varepsilon > 0$, we take $\widehat{a} = \{\widehat{a}_n\}_{n=2}^{\infty}$ with $\widehat{a}_n = \varepsilon^{\frac{1}{p}} n^{-\frac{1}{p}} (\log n)^{-\frac{1+\varepsilon}{p}}$. We see that

$$\|\widehat{a}\|_p^p = \frac{\varepsilon}{2(\log 2)^{1+\varepsilon}} + \varepsilon \sum_{n=3}^{\infty} n^{-1} (\log n)^{-1-\varepsilon} \leq \varepsilon + 1.$$

Then we have

$$\|\widehat{\mathbf{H}}_p^\delta \widehat{a}\|_p^p = \sum_{m=2}^{\infty} \frac{(\log m)^\delta}{m} \left[\sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\delta+1+\varepsilon}{p}}}{n \log(mn)} \right]^p.$$

(1) If $\delta < -1$, when $\varepsilon < -(\delta+1)$, we see from $\delta+1+\varepsilon < 0$ that, for a fixed $m \geq 2$, it holds that

$$\sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\delta+1+\varepsilon}{p}}}{n \log(mn)} \geq \sum_{n=m}^{\infty} \frac{(\log n)^{-\frac{\delta+1+\varepsilon}{p}}}{n \log(mn)} \geq \frac{1}{2} \sum_{n=m}^{\infty} \frac{(\log n)^{-1-\frac{\delta+1+\varepsilon}{p}}}{n} = +\infty.$$

This means that $\widehat{\mathbf{H}}_p^\delta$ is not bounded on l^p in this case.

(2) If $\delta = -1$ or $\delta \geq p-1$, we have

$$\|\widehat{\mathbf{H}}_p^\delta \widehat{a}\|_p^p = \sum_{m=2}^{\infty} \frac{(\log m)^\delta}{m} \left[\sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\delta+1+\varepsilon}{p}}}{n \log(mn)} \right]^p$$

$$\begin{aligned}
&\geq \sum_{m=2}^{\infty} \frac{(\log m)^\delta}{m} \left[\int_2^{\infty} \frac{(\log x)^{-\frac{\delta+1+\varepsilon}{p}}}{x \log(xm)} dx \right]^p \\
&= \sum_{m=2}^{\infty} \frac{1}{m(\log m)^{1+\varepsilon}} \left[\int_{\frac{\log 2}{\log m}}^{\infty} \frac{t^{-\frac{\delta+1+\varepsilon}{p}}}{1+t} dt \right]^p \\
&\geq \sum_{m=2}^{\infty} \frac{1}{m(\log m)^{1+\varepsilon}} \left[\int_1^{\infty} \frac{t^{-\frac{\delta+1+\varepsilon}{p}}}{1+t} dt \right]^p.
\end{aligned}$$

We note that

$$\sum_{m=2}^{\infty} \frac{1}{m(\log m)^{1+\varepsilon}} = \frac{1}{\varepsilon}(1 + o(1)), \varepsilon \rightarrow 0^+.$$

Therefore, we conclude that

$$\|\widehat{\mathbf{H}}_p^\delta \widehat{a}\|_p^p \geq \frac{1}{\varepsilon}(1 + o(1)) \left[\int_1^{\infty} \frac{t^{-\frac{\varepsilon}{p}}}{1+t} dt \right]^p.$$

Take $\varepsilon \rightarrow 0^+$, we get that $\|\widehat{\mathbf{H}}_p^\delta \widehat{a}\|_p^p \rightarrow +\infty$. This implies that $\widehat{\mathbf{H}}_p^\delta$ is not bounded on l^p when $\delta = -1$ or $\delta \geq p-1$. The claim is proved. \square

Remark 4.3. When $\alpha = \beta = 1$. From Theorem 1.2, we know that

$$\mathbf{H}_{p,1,1,\gamma}^{\mu,\nu}(a)(m) = \frac{(\log m)^{\frac{\mu}{p}}}{m^{\frac{1}{p}}} \sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\nu}{p}}}{n^{\frac{1}{p}}} \cdot \frac{a_n}{(\log mn)^\gamma}, \quad a = \{a_n\}_{n=2}^{\infty} \in l^p, \quad m \geq 2,$$

is not bounded on l^p .

On one hand, We note that, for $\gamma > 0$,

$$\int_{[0,1)} t^{(\log mn)-1} (1-t)^{\gamma-1} dt = \frac{\Gamma(\log mn)\Gamma(\gamma)}{\Gamma(\gamma + \log mn)}, \quad m, n \geq 2.$$

On the other hand, by using the fact that

$$(4.1) \quad \Gamma(x) = \sqrt{2\pi} x^{x-\frac{1}{2}} e^{-x} [1 + r(x)], \quad |r(x)| \leq e^{\frac{1}{12x}} - 1, \quad x > 0,$$

we see that there are two constants $C_1 > 0, C_2 > 0$, independent of γ, m, n , such that

$$\frac{C_1}{(\log mn)^\gamma} \leq \frac{\Gamma(\log mn)\Gamma(\gamma)}{\Gamma(\gamma + \log mn)} \leq \frac{C_2}{(\log mn)^\gamma}.$$

Thus,

$$\frac{C_1}{(\log mn)^\gamma} \leq \int_{[0,1)} t^{(\log mn)-1} (1-t)^{\gamma-1} dt \leq \frac{C_2}{(\log mn)^\gamma}.$$

Hence, in order for $\mathbf{H}_{p,1,1,\gamma}^{\mu,\nu} : l^p \rightarrow l^p$ to be bounded when $\gamma < 1 + \frac{\mu-\nu}{p}$, let λ be a positive measure in $[0, 1)$, we will consider the operator $\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}$, which is defined as

$$\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}(a)(m) := \frac{(\log m)^{\frac{\mu}{p}}}{m^{\frac{1}{p}}} \sum_{n=2}^{\infty} \frac{(\log n)^{-\frac{\nu}{p}}}{n^{\frac{1}{p}}} \lambda[mn] a_n, \quad a = \{a_n\}_{n=2}^{\infty} \in l^p, \quad m \geq 2,$$

where

$$\lambda[n] = \int_{[0,1)} t^{(\log n)-1} (1-t)^{\gamma-1} d\lambda(t), \quad n \geq 2.$$

We find that

Proposition 4.4. *Let $p > 1, \gamma > 0, -1 < \mu, \nu < p - 1$. If $(1 - t)^{\gamma-1}d\lambda(t)$ is a $[1 + \frac{\mu-\nu}{p}]$ -Carleson measure, then $\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}$ is bounded on l^p .*

Here, for $s > 0$, a positive Borel measure λ on $[0, 1]$, we say λ is an s -Carleson measure if there is a constant $C_3 > 0$ such that

$$\lambda([t, 1]) \leq C_3(1 - t)^s$$

holds for all $t \in [0, 1]$.

Proof. Set $d\tau(t) = (1 - t)^{\gamma-1}d\lambda(t)$. We get from integration by parts that, for $n \geq 8$,

$$\begin{aligned} \lambda[n] &= \int_0^1 t^{(\log n)-1} d\tau(t) = \tau([0, 1]) - [(\log n) - 1] \int_0^1 t^{(\log n)-2} \tau([0, t]) dt \\ &= [(\log n) - 1] \int_0^1 t^{(\log n)-2} \tau([t, 1]) dt. \end{aligned}$$

Since $d\tau(t) = (1 - t)^{\gamma-1}d\lambda(t)$ is a $[1 + \frac{\mu-\nu}{p}]$ -Carleson measure, then we see that there is a constant $C_4 > 0$ such that

$$\tau([t, 1]) \leq C_4(1 - t)^{1 + \frac{\mu-\nu}{p}}$$

holds for all $t \in [0, 1]$. It follows that, for $n \geq 8$,

$$\begin{aligned} \lambda[n] &\leq C_4 [(\log n) - 1] \int_0^1 t^{(\log n)-2} (1 - t)^{1 + \frac{\mu-\nu}{p}} dt \\ &= C_4 \frac{[(\log n) - 1] \Gamma((\log n) - 1) \Gamma(2 + \frac{\mu-\nu}{p})}{\Gamma((\log n) + 1 + \frac{\mu-\nu}{p})}. \end{aligned}$$

By using (4.1) again, we see that there is a constant $C_5 > 0$ such that

$$\lambda[n] \leq \frac{C_5}{(\log n)^{1 + \frac{\mu-\nu}{p}}}.$$

Consequently, we conclude that, for all $m, n \geq 2$, there is a constant $C_6 > 0$ such that

$$\lambda[mn] \leq \frac{C_6}{(\log mn)^{1 + \frac{\mu-\nu}{p}}}.$$

Then, by checking the proof of Theorem 1.3, we see that $\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}$ is bounded on l^p . The proposition is proved. \square

However, we do not know whether the condition $(1 - t)^{\gamma-1}d\lambda(t)$ is a $[1 + \frac{\mu-\nu}{p}]$ -Carleson measure is necessary for the boundedness of $\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}$ on l^p . It is natural to consider the following

Problem 4.5. *Characterize measures λ such that $\widehat{\mathbf{H}}_{p,\gamma,\lambda}^{\mu,\nu}$ is bounded on l^p .*

Remark 4.6. In Theorem 1.3, we give a sharp estimate for the norm of $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ when $\gamma = 1 + \frac{\mu-\nu}{p}$. It is interesting to study

Problem 4.7. *Determine the exact value of the norm of $\mathbf{H}_{p,\alpha,\beta,\gamma}^{\mu,\nu}$ when $\gamma > 1 + \frac{\mu-\nu}{p}$.*

REFERENCES

- [1] Andrews G., Askey R., Roy R., *Special functions*, Cambridge University Press, Cambridge, 1999.
- [2] Bayart F., Brevig O., *Composition operators and embedding theorems for some function spaces of Dirichlet series*, Math. Z., 293 (2019), no. 3-4, pp. 989-1014.
- [3] Brevig O., Perfekt K., Seip, K., Siskakis A., Vukotić D., *The multiplicative Hilbert matrix*, Adv. Math., 302 (2016), pp. 410-432.
- [4] Miheisi N., Pushnitski A., *A Helson matrix with explicit eigenvalue asymptotics*, . J. Funct. Anal. , 275 (2018), no. 4, pp. 967-987.
- [5] Mitrinović, D., *Analytic inequalities*, Springer-Verlag, New York-Berlin, 1970.
- [6] Perfekt K., Pushnitski A., *On the spectrum of the multiplicative Hilbert matrix*, Ark. Mat. , 56 (2018), no. 1, pp. 163-183.

SCHOOL OF MATHEMATICS SCIENCES, HEFEI UNIVERSITY OF TECHNOLOGY, XUANCHENG CAMPUS, XUANCHENG 242000, P.R.CHINA

Email address: jin@hfut.edu.cn, jinjjhb@163.com