

p -Adic quantum calculus and ideals of compact operators

Evgeny I. Zelenov

May 10, 2022

Abstract

The paper proposes a construction of a quantum differentiation operator defined on the spaces of complex-valued functions of p -adic argument and taking values in the algebra of bounded operators on a Hilbert space. The properties of this operator are investigated. In particular, it is proved that the differentiation operator maps p -adic Besov spaces into Schatten-von Neumann ideals in the algebra of compact operators.

1 Introduction

First of all, we will give a description of what will be called the p -adic quantum calculus. Consider the algebra $A = L^\infty(\mathbb{Z}_p, \mathbb{C})$ of almost everywhere bounded measurable on \mathbb{Z}_p functions taking value in the field of complex numbers. Let's define an irreducible representation of this algebra by multiplication operators in the Hilbert space $L^2 = L^2(\mathbb{Z}_p)$, $A \ni f \mapsto M_f \in \mathcal{B}(L^2)$,

$$M_f: L^2 \ni h \mapsto fh \in L^2.$$

On the space L^2 , the Hilbert operator S is defined (we will define it later), which is a self-adjoint operator and $S^2 = \text{Id}$. For each function $f \in A$, we define a quantum differentiation operator

$$df = [M_f, S].$$

The correspondence $f \mapsto df$ constructed in this way between functions on \mathbb{Z}_p and the operators df on L^2 we will call p -adic quantum calculus.

The group \mathbb{Z}_p of p -adic integers is the inverse limit of finite cyclic groups $\mathbb{Z}/p^n\mathbb{Z}$, $n \in \mathbb{Z}_+$:

$$\longleftarrow \mathbb{Z}/p^n\mathbb{Z} \longleftarrow \mathbb{Z}/p^{n+1}\mathbb{Z} \longleftarrow$$

In this sense, the proposed construction is a combinatorial analogue of the quantum calculus suggested by A. Connes[1].

The purpose of this article is to study this correspondence. In particular, the following statements will be proved.

- df is a finite rank operator if and only if the function f is locally constant;
- df is a compact operator if and only if the function f belongs to the space $VMO(\mathbb{Z}_p)$;
- df belongs to the Schatten-von Neumann ideal \mathfrak{S}^q if and only if f belongs to the p -adic Besov space $B_{q,q}^{1/q}$.

Note that the last two statements are completely analogous to the corresponding statements for algebras of complex-valued functions on the circle $\mathbb{T} \subset \mathbb{C}$.

The Connes quantum calculus (another name is the Connes quantization) has many interesting applications (see, for example, [2, 3]).

There is an important feature in the p -adic case considered in this paper. Namely, in all cases interesting for the application, there is no nontrivial "classical" differentiation (by differentiation on the algebra A we mean a linear map $A \rightarrow A$ satisfying the Leibniz rule). The reason for this is that there are no nontrivial differentiations on the algebra of locally constant functions, and this algebra is a dense subalgebra of the algebra of observables in most interesting cases ([4, 5]) (for example, locally constant functions form a dense subset in the algebra of continuous on \mathbb{Z}_p functions). At the same time, "quantum" differentiation, that is, differentiation with values in the algebra of bounded linear operators on a Hilbert space, is not trivial.

One of the possible applications of this paper is the construction of differential calculus on spaces of functions defined on totally disconnected spaces ([6, 7, 8]).

2 p -Adic numbers, functions, spaces

In this section, the necessary notations are introduced and the functional spaces in which we will work are defined. Everywhere further we consider the case of $p \neq 2$.

Any nonzero p -adic number can be represented in the following form (canonical decomposition):

$$\mathbb{Q}_p \ni x = \sum_{k=-n}^{+\infty} x_k p^k, \quad n \in \mathbb{Z}_+, \quad x_k \in \{0, 1, \dots, p-1\}.$$

The finite part of the canonical decomposition, consisting of the sum of terms with negative powers of p , is the fractional part of x and is denoted as $\{x\}_p$, respectively, the remaining part of the decomposition (with non-negative powers of p) is the integer part of x . Note that the fractional part is always a rational number, p -adic numbers with a trivial fractional part are p -adic integers \mathbb{Z}_p .

$$\underbrace{p^{-n}x_{-n} + p^{-n+1}x_{-n+1} + \dots + p^{-1}x_{-1}}_{\{x\}_p} + \underbrace{x_0 + px_1 + \dots + p^kx_k + \dots}_{[x]_p}$$

Let's define the order of a nonzero p -adic number x by the following rule: $\text{ord}(x) = \min\{k: x_k \neq 0\}$. Order has the following property: $\text{ord}(x+y) \geq \min\{\text{ord}(x), \text{ord}(y)\}$. p -adic norm is determined by the formula: $|x|_p = p^{-\text{ord}(x)}$. This norm is non-Archimedean, that is, a strong triangle inequality is valid: $|x+y|_p \leq \max\{|x|_p, |y|_p\}$. By χ we denote the additive character of the field \mathbb{Q}_p of the following form: $\chi(x) = \exp(2\pi i\{x\}_p)$.

For a nonzero p -adic number $x \in \mathbb{Q}_p^*$, the symbol $x_{\text{ord}(x)} \in \{0, 1, \dots, p-1\}$ denotes the coefficient at the first nonzero term of the canonical decomposition of x . Let's define the sign $\text{sgn}(x)$ of a nonzero p -adic number as the Legendre symbol of $x_{\text{ord}(x)}$: $\text{sgn}(x) = \left(\frac{x_{\text{ord}(x)}}{p}\right)$. The function $\text{sgn}(x)$ takes values in the set $\{-1, 1\}$ and is the multiplicative character of the field \mathbb{Q}_p : $\mathbb{Q}_p^* \rightarrow \{-1, 1\}$, $\text{sgn}(xy) = \text{sgn}(x)\text{sgn}(y)$. A disk centered at zero of radius p^{-n} is denoted by B_n : $B_n = \{x \in \mathbb{Z}_p : \text{ord}(x) \geq n \equiv |x|_p \leq p^{-n}\}$. In the following, we will consider functions defined on the ring \mathbb{Z}_p of p -adic integers (as a set, \mathbb{Z}_p is a unit disk, $\mathbb{Z}_p = B_0$) and taking values in the field \mathbb{C} of complex numbers.

A function f is locally constant of order n if it is invariant with respect to shifts: $f(x + u) = f(x)$, $x \in \mathbb{Z}_p$, $u \in B_n$. The set of such functions form a finite-dimensional vector space LC_n over complex numbers of dimension p^n . The basis vectors of this space are indicator functions of disks of radius p^{-n} in \mathbb{Z}_p . Since the topological space \mathbb{Z}_p is totally disconnected, locally constant functions are continuous. The following embeddings are valid: $m \leq n$, $LC_m \subseteq LC_n$. The set of all locally constant functions will be denoted by LC . This space has a natural structure of the direct limit of finite-dimensional subspaces: $LC = \varinjlim LC_n$. The Haar measure on \mathbb{Z}_p will be denoted by μ . Lebesgue spaces of measurable functions with respect to this measure are defined in a standard way: $L^q \equiv \{f: \mathbb{Z}_p \rightarrow \mathbb{C}, \|f\|_q = \left(\int_{\mathbb{Z}_p} |f|^q d\mu\right)^{1/q} < \infty\}$, $\|f\|_\infty = \text{ess sup}_{\mathbb{Z}_p} |f|$.

There is a chain of embeddings:

$$LC \subset L^\infty \subset \dots \subset L^q \subset \dots \subset L^1,$$

LC is dense in L^q for all $1 \leq q \leq \infty$.

3 Hilbert Transform

In this section, we define and consider the basic properties of the p -adic Hilbert transform (or operator). A large number of papers have been devoted to the study of the properties of this type of operators, here are just some, see [9, 10, 11].

The Hilbert transform S of the function f is defined by the following formula (the integral is understood in the sense of principal value):

$$((Sf)(x) = \frac{1}{\Gamma} \text{p.v.} \int_{\mathbb{Z}_p} \frac{\sin(x-y)}{|x-y|_p} f(y) dy, \Gamma = \begin{cases} \sqrt{p}, & p \equiv 1 \pmod{4} \\ i\sqrt{p}, & p \equiv 3 \pmod{4} \end{cases}$$

We use the notation S for the Hilbert operator, since in the future it plays the role of a symmetry operator. The Hilbert operator is bounded from L^q to L^q for all $1 < q < \infty$. For $f \in L^1$, the estimate is valid:

$$\mu \{x \in \mathbb{Z}_p : |(Sf)(x)| > \lambda\} \leq C \frac{\|f\|_1}{\lambda}.$$

4 Fourier transform and Hilbert transform

The following section presents some well-known facts from harmonic analysis over the field of p -adic numbers [10].

Any character of the group \mathbb{Z}_p of p -adic integers has the form $\chi_\alpha(x) = \chi(\alpha x)$, $x \in \mathbb{Z}_p$, $\alpha \in \mathbb{Q}_p$. Thus, the Pontryagin dual of the group \mathbb{Z}_p is $\mathbb{Q}_p/\mathbb{Z}_p$:

$$\hat{\mathbb{Z}}_p = \{\chi_\alpha(x) = \chi(\alpha x), \alpha \in \mathbb{Q}_p/\mathbb{Z}_p\}.$$

In the last formula in the expression $\chi(\alpha x)$ an arbitrary representative from the corresponding adjacency class is taken as α . Since the function $\chi(x)$ is identically equal to 1 on \mathbb{Z}_p , then the formula is correct. The group $\hat{\mathbb{Z}}_p$ is also known as the Prufer group $\mathbb{Z}(p^\infty)$, it is a quasi-cyclic group and can be represented as a subgroup in \mathbb{T} consisting of p^n -th roots of unity $n \in \mathbb{N}$.

The set of functions $\{\chi_\alpha(x), \alpha \in \hat{\mathbb{Z}}_p\}$ forms an orthonormal basis in L^2 . The Fourier transform of the function $\phi \in L^2$ is defined in the standard way:

$$\phi \in L^2, F[\phi](\alpha) = \langle \phi, \chi_\alpha \rangle = \hat{\phi}_\alpha = \int_{\mathbb{Z}_p} \phi(x) \chi(-\alpha x) dx.$$

Note that the function χ_α is a locally constant of order n , where n is determined by the norm α : $\chi_\alpha \in LC_n$, $|\alpha|_p = p^n$. The norm $|\cdot|_p$ can be correctly defined on $\mathbb{Q}_p/\mathbb{Z}_p$, assuming it is equal to zero on a zero adjacency class and equal to the norm of any representative from a non-zero adjacency class (due to the non-archimedean norm, such a definition does not depend on the choice of a representative). Moreover, the family of functions $\{\chi_\alpha(x), \alpha \in \hat{\mathbb{Z}}_p, |\alpha|_p \leq p^n\}$ forms an orthonormal basis in the space LC_n . It also follows that the function ϕ is locally constant if and only if it is represented as a finite Fourier series: $\phi \in LC \iff \hat{\phi}_\alpha = 0$ for almost all $\alpha \in \hat{\mathbb{Z}}_p$.

In this sense, locally constant functions are analogs of trigonometric polynomials. However, there is an important difference - the ratio of two locally constant functions (provided that the function in the denominator does not take zero values) is again a locally constant function.

Hilbert transform has the following properties..

- The Hilbert transform maps constant functions into a function identically equal to zero: $(SI_{\mathbb{Z}_p})(x) = 0$.

- The characters χ_α , $\alpha \in \hat{\mathbb{Z}}_p$ are eigenfunctions of the Hilbert operator with eigenvalues $\text{sgn}(\alpha)$: $(S\chi_\alpha)(x) = \text{sgn}(\alpha)\chi_\alpha(x)$, $\alpha \in \hat{\mathbb{Z}}_p$, $x \in \mathbb{Z}_p$. It should be noted here that the function sgn can be correctly defined on the group $\hat{\mathbb{Z}}_p$ as follows: $\text{sgn}(\alpha)$ is set equal to zero on the unit element in $\hat{\mathbb{Z}}_p$ and equal to the value on an arbitrary adjacency class representative otherwise. Obviously, this value does not depend on the choice of a representative.
- As can be easily seen from the previous property of the Hilbert operator, this operator is self-adjoint, and its square is equal to the identity operator. That is, the Hilbert operator is a symmetry operator: $S^* = S$, $S^2 = 1$.
- The space L_0^2 (the space of functions from L^2 orthogonal to constants) can be decomposed into a direct orthogonal sum of subspaces W^+ and W^- : $L^2 = W^+ \oplus W^-$, the space W^+ is linear span of the basis vectors χ_α , for which the condition $\text{sgn}(\alpha) = 1$ is met, the space W^- is linear span of the basis vectors χ_α , for which the condition $\text{sgn}(\alpha) = -1$ is met. By P^+ and P^- we denote orthogonal projectors into subspaces W^+ and W^- , respectively. Then the representation is valid for the Hilbert operator $S = P^+ - P^-$.

5 Differentiation operator

In this section, we will define the quantum differentiation operator in the p -adic case (hereinafter - the differentiation operator), and give its simplest properties.

Definition 1 *Let $f \in L^\infty$ and M_f be the multiplication operator in L^2 , $(M_f\phi)(x) = f(x)\phi(x)$. The operator in L^2 of the following form*

$$df = [M_f, S]$$

will be called the value of the differentiation operator of the function f .

The square brackets in the last formula denote the commutator. Thus, the differentiation operator is defined on the algebra L^∞ of functions on \mathbb{Z}_p and takes a value in the algebra $\mathcal{B} = \mathcal{B}(L^2)$ of bounded linear operators on the space L^2 . It is obvious that the differentiation operator satisfies the Leibniz

rule: $d(fg) = (df)g + f(dg)$. We will also call the value of the operator d on the function f the derivative of the function f .

It immediately follows from the definition that the derivative of a constant function is a trivial operator on \mathcal{B} (maps any function to the identically zero function). Let's give a less trivial example.

Example 1 *Let $f(x) = \chi_a(x)$. Let us find the derivative of this function. Since df is an operator in L^2 , we calculate the value of this operator on the basis vector χ_α . The answer is as follows:*

$$(d\chi_a)\chi_\alpha = (\operatorname{sgn}(\alpha) - \operatorname{sgn}(\alpha + a))\chi_{\alpha+a}.$$

The following equations are valid:

$$\begin{aligned} (d\chi_a)\chi_\alpha &= [M_{\chi_a}, S]\chi_\alpha = \chi_a S\chi_\alpha - S(\chi_a\chi_\alpha) = \\ &= \chi_a \operatorname{sgn}(\alpha)\chi_\alpha - \operatorname{sgn}(\alpha + a)\chi_{\alpha+a} = (\operatorname{sgn}(\alpha) - \operatorname{sgn}(\alpha + a))\chi_{\alpha+a}. \end{aligned}$$

The following theorem is valid.

Theorem 1 *Let $f \in L^\infty$. The derivative df is an operator of finite rank if and only if the function f is locally constant.*

Let $f \in LC$. Then, as noted above, the function f is represented as a finite linear combination of characters χ_a , that is, $f(x) = \sum_{a \in \Omega} \hat{f}_a \chi_a(x)$, while a belongs to a finite subset of Ω in $\hat{\mathbb{Z}}_p$. Therefore, the inequality $\operatorname{rank}(df) \leq \sum_{a \in \Omega} \operatorname{rank}(d\chi_a)$ is valid. Consider the operator $T = d\chi_a(d\chi_a)^*$. The rank of the operator T coincides with the rank of the operator $d\chi_a$. As follows from the Example 1, the operator T acts on the basis vectors χ_α according to the following formula:

$$T\chi_\alpha = (\operatorname{sgn}(\alpha + a) - \operatorname{sgn}(\alpha))^2 \chi_\alpha. \quad (1)$$

Thus, the characters χ_α are the eigenvalues of the operator T , the corresponding eigenvalues are

$$\lambda_\alpha = (\operatorname{sgn}(\alpha + a) - \operatorname{sgn}(\alpha))^2.$$

Let $a \neq 0$. Then it is easy to see that $\lambda_0 = \lambda_{-a} = 1$. In addition, as follows from the definition of the function sgn , $\operatorname{sgn}(\alpha + a) = \operatorname{sgn}(\alpha)$ for all α such that the inequality $|\alpha|_p > |a|_p$ is satisfied. Thus, the image of the operator

T is a subspace of the linear span of the basis vectors χ_α , for which the condition $|\alpha|_p \leq |a|_p$ is satisfied. The dimension of this subspace does not exceed $|a|_p$. Hence, $\text{rank}(d\chi_a) \leq |a|_p$. Hence the sufficiency of the condition of the theorem is proved.

It is possible to obtain an exact formula for the rank of the operator $d\chi_a$. Namely:

$$\text{rank}(d\chi_a) = \frac{|a|_p + 3}{2}. \quad (2)$$

Indeed, in a finite field of order p^n , the number of nonzero elements that are squares coincides with the number of elements that are not squares and is equal to $1/2(p^n - 1)$. Therefore, for $\alpha \neq 0$ and $\alpha \neq -a$, we have $1/2(|a|_p - 1)$ of nonzero eigenvalues of λ_α , and we need to add two more, namely, λ_0 and λ_{-a} . If the function $f \in L^\infty$ is not locally constant, then its Fourier series expansion contains nonzero terms $\hat{f}_a \chi_a$ with an index a of arbitrarily large norm. Taking into account the formula (2), the operator df cannot have a finite rank in this case.

Corollary 1 *If the function f is continuous, $f \in C(\mathbb{Z}_p)$, then its derivative df is a compact operator.*

Indeed, the mapping $f \mapsto M_f$ is continuous in a uniform topology on $C(\mathbb{Z}_p)$ and norm topologies on $\mathcal{B}(L^2)$. Mapping $A \mapsto [A, S]$, $A \in \mathcal{B}(L^2)$ is a differentiation on the algebra of bounded operators $\mathcal{B}(L^2)$ and is therefore continuous in the topology of the norm ([12]). Thus, the differentiation operator $f \mapsto df$ is continuous in a uniform topology on $C(\mathbb{Z}_p)$ and norm topologies on $\mathcal{B}(L^2)$. Since locally constant functions are dense in $C(\mathbb{Z}_p)$, the image of the algebra of continuous functions lies in the closure of the space of operators of finite rank in the topology of the norm in $\mathcal{B}(L^2)$, that is, in the algebra $\mathcal{K}(L^2)$ of compact operators on L^2 . Corollary 1 gives a sufficient condition for the compactness of the derivative, however, this condition is not necessary.

6 Hilbert-Schmidt operators

Let's define the p -adic Sobolev space $H^{1/2}$. This space is an analogue of the Sobolev space of semi-differentiable functions. To do this, on the space LC of locally constant functions, we define the $\|\cdot\|_{1/2}$ semi-norm as follows. Let $f \in LC$ be represented as its (finite) Fourier series: $f(x) = \sum_{a \in \hat{\mathbb{Z}}_p} \hat{f}_a \chi_a(x)$.

Then, by definition,

$$\|f\|_{1/2} = \left(\sum_{a \in \hat{\mathbb{Z}}_p} |a|_p |\hat{f}_a|^2 \right)^{1/2}. \quad (3)$$

The kernel of this semi-norm is a set of constant functions. The space $H^{1/2}$ is defined as the closure of the space LC with respect to the seminorm (3).

Lemma 1 *Let $f(x) = \sum_{a \in \hat{\mathbb{Z}}_p} \hat{f}(a) \chi_a(x)$. The following formula is valid:*

$$\text{Tr} (df)^* (df) = 2 \sum_{a \in \hat{\mathbb{Z}}_p} |a|_p |\hat{f}_a|^2.$$

Let's use the formula (1) to calculate the diagonal matrix element of the operator $(df)^* (df)$:

$$\begin{aligned} \langle (df)^* (df) \chi_\alpha, \chi_\alpha \rangle &= \\ &= \left\langle \sum_{a, b \in \hat{\mathbb{Z}}_p} \hat{f}_a \bar{\hat{f}}_b (\text{sgn}(\alpha) - \text{sgn}(a + \alpha)) (\text{sgn}(\alpha) - \text{sgn}(b + \alpha)) \chi_{\alpha+a-b}, \chi_\alpha \right\rangle = \\ &= \sum_{a \in \hat{\mathbb{Z}}_p} |\hat{f}_a|^2 (\text{sgn}(\alpha) - \text{sgn}(a + \alpha))^2. \quad (4) \end{aligned}$$

As noted above, $\lambda_\alpha = (\text{sgn}(\alpha) - \text{sgn}(a + \alpha))^2$ takes exactly two values equal to one (for $\alpha = 0$ and $\alpha = -a$) and the value 4 exactly for $1/2(|a|_p - 1)$ of different values of α , for the rest α the eigenvalues of λ_α are equal to zero. Therefore, taking into account the relations (4), the equalities are valid:

$$\begin{aligned} \text{Tr} (df)^* (df) &= \sum_{\alpha \in \hat{\mathbb{Z}}_p} \sum_{a \in \hat{\mathbb{Z}}_p} |\hat{f}_a|^2 (\text{sgn}(\alpha) - \text{sgn}(a + \alpha))^2 = \\ &= \sum_{a \in \hat{\mathbb{Z}}_p} |\hat{f}_a|^2 \left(2 + 4 \frac{|a|_p - 1}{2} \right) = 2 \sum_{a \in \hat{\mathbb{Z}}_p} |a|_p |\hat{f}_a|^2. \end{aligned}$$

Thus, we have proved the following theorem.

Theorem 2 *The derivative df is a Hilbert-Schmidt operator, $df \in \mathfrak{S}^2$, if and only if the function f belongs to the space $H^{1/2}$.*

7 Spaces $BMO(\mathbb{Z}_p)$ and $VMO(\mathbb{Z}_p)$

Initially, we defined the differentiation operator for the functions $f \in L^\infty$. In fact, this operator can naturally be extended to a wider space. We will give definitions of the corresponding spaces. Let the function f be integrable on \mathbb{Z}_p , $f \in L^1$ and B denote an arbitrary disk in \mathbb{Z}_p . Using f_B , we denote the average value of the function f on the disk B :

$$f_B = \frac{1}{\mu(B)} \int_B f(x) dx.$$

Let's calculate the average deviation of the function f from its average value on the disk B , and consider the maximum of such average deviations for all disks of radius not exceeding p^{-n} , $n \in \mathbb{Z}_+$:

$$M_n = \max_{\{B: \mu(B) \leq p^{-n}\}} \frac{1}{\mu(B)} \int_B |f(x) - f_B| dx.$$

We will say that the function f belongs to the space $BMO = BMO(\mathbb{Z}_p)$ (bound mean oscillation), if the sequence M_n is bounded, $\|f\|_{BMO} = \sup_n M_n$ sets a semi-norm on the BMO space, the kernel of this semi-norm consists of constant functions. A typical example of a function from the BMO space is the function $\log |x|_p$. A subspace of the BMO space consisting of functions for which the condition is met $\lim_{n \rightarrow \infty} M_n = 0$ denote $VMO = VMO(\mathbb{Z}_p)$ (vanishing mean oscillation). The BMO space contains the space L^∞ as its proper subspace and is the natural space of the definition of the differentiation operator.

Theorem 3 *The derivative df of the function f is a bounded operator on L^2 , $df \in \mathcal{B}(L^2)$, if and only if f belongs to the space BMO . The derivative df is a compact operator, $df \in \mathcal{K}(L^2)$, if and only if the function f belongs to the VMO space.*

These statements are completely analogous to the corresponding statement for quantum differentiation in the complex case and are proved in a similar way. Therefore, we will present only a sketch of the proof. It is known ([10, 13]) that the Hilbert operator is a continuous operator from BMO to BMO and a continuous operator from L^∞ to BMO. In addition, the following characterization of the BMO space is valid in terms of the action of the Hilbert operator on the space L^∞ . Namely, any function $f \in BMO$ is

represented as $f = g + Sh$, where the functions g and h belong to the space L^∞ , while the inequalities $\|f\|_{BMO} \geq C\|g\|_\infty$, $\|f\|_{BMO} \geq C\|h\|_\infty$ are valid for some constant C . This result was proved for functions defined on \mathbb{R}^N ([14]), for functions on \mathbb{Z}_p the proof is similar. Using this representation for a function from the BMO space, to continue the differentiation operator from L^∞ to the BMO space, it is enough to do this on SL^∞ , and this follows from the continuity of the Hilbert operator from L^∞ to BMO . Similarly, the second part of the theorem can be proved. Instead of the result of Fefferman and Stein ([14]), we will use the following result of Sarason ([15]), more precisely, the p -adic analogue of this result. A function $f \in BMO$ belongs to the VMO space if and only if it can be represented as $f = g + Sh$, where the functions g and h are continuous on \mathbb{Z}_p . Taking into account the last statement, the continuity of the Hilbert operator and Corollary 1 of the Theorem 1, we obtain the second statement of the theorem.

8 Besov spaces $B_{q,r}^s$

A considerable amount of work has been devoted to the Besov spaces of functions on the field \mathbb{Q}_p of p -adic numbers (and more generally to the spaces of functions on profinite groups and martingales), for example [16, 17, 18].

Let q, r, s be real numbers satisfying the inequalities $s > 0$, $1 \leq q, r \leq \infty$. The subspace of functions from L^1 for which the seminorm $\|\cdot\|_{B_{q,r}^s}$ defined below is finite, we will call the Besov space $B_{q,r}^s$:

$$B_{q,r}^s = \left\{ f \in L^1 : \|f\|_{B_{q,r}^s} = \left(\int_{\mathbb{Z}_p} \left(\frac{\|f(x-y) - f(x)\|_q}{|y|_p^s} \right)^r \frac{dy}{|y|_p} \right)^{1/r} < \infty \right\}.$$

An equivalent definition of the Besov space is useful ([16]). We introduce the notation $\Delta_n(x)$ for the normalized indicator function of the disk B_n : $\Delta_n(x) = \frac{1}{\mu(B_n)} \mathbb{I}_{B_n}(x)$. The mapping $L^1 \ni f \mapsto f * \Delta_n \in LC_n$ defines the projection of the space L^1 into the space LC_n ($*$ denotes convolution).

Proposition 1 *Let the conditions be met: $f \in L^1$, $s > 0$, $1 \leq q, r \leq \infty$.*

The seminorm $\|f\|_{B_{q,r}^s}$ is equivalent to the following seminorm:

$$\|f\|_{B_{q,r}^s} \asymp \left(\sum_{n=0}^{\infty} (p^{ns} \|f - f * \Delta_n\|_q)^r \right)^{1/r}.$$

Below, we will need the case $r = q$, $s = 1/q$. An obvious consequence follows from the last statement.

Corollary 2 *The function f belongs to the Besov space $B_{q,q}^{1/q}$, $f \in B_{q,q}^{1/q}$, if and only if the condition is satisfied $p^{n/q} \|f - f * \Delta_n\|_q \in \ell^q$.*

The connection of Besov and BMO spaces on Vilenkin groups was studied in [16]. In particular, a stronger statement follows from the results of this work. Namely:

Proposition 2 *The function f belongs to the Besov space $B_{q,q}^{1/q}$, $f \in B_{q,q}^{1/q}$, if and only if the following condition is satisfied $p^{n/q} \|f - f * \Delta_n\|_{BMO} \in \ell^q$.*

The following theorem is valid.

Theorem 4 *The derivative df belongs to the Schatten-von Neumann ideal \mathfrak{S}^q , $df \in \mathfrak{S}^q$, if and only if the function f belongs to the Besov space $B_{q,q}^{1/q}$, $f \in B_{q,q}^{1/q}$.*

Recall the definition of the ideal \mathfrak{S}^q . Let $K \in \mathcal{K}(L^2)$ is a compact operator. The singular number $s_n(K)$ of this operator is defined as the distance in the operator norm to the space of operators of rank no higher than n :

$$s_n(K) = \inf_R \{ \|K - R\| : \text{rank } R \leq n \}.$$

By definition, $K \in \mathfrak{S}^q$ if the condition $\{s_n(K)\} \in \ell^q$ is satisfied. The last condition is equivalent to the following condition: $\{p^{n/q} s_{p^n}(K)\} \in \ell^q$. Since $f * \Delta_n \in LC_n$, the derivative $d(f * \Delta_n)$ is an operator of rank no higher than p^n (Theorem 1). Therefore, the inequality $s_{p^n}(df) \leq \|d(f - f * \Delta_n)\|$ is valid. In fact, a two-way estimate of $s_{p^n}(df) \asymp \|d(f - f * \Delta_n)\|$ is valid, since the function $f * \Delta_n$ gives the best approximation in the topology of the VMO space of the function f by a locally constant function from LC_n , and the differentiation operator is a continuous mapping from VMO to the algebra of compact operators. Taking into account Theorem 3, the chain of relations is valid:

$$s_{p^n}(df) \asymp \|d(f - f * \Delta_n)\| \asymp \|f - f * \Delta_n\|_{BMO}.$$

Taking into account the Proposition 2, the statement of the theorem follows from the last chain of inequalities.

References

- [1] A. Connes, Noncommutative geometry, San Diego, CA: Academic Press, 1994.
- [2] A. G. Sergeev, “Quantization of the Sobolev Space of Half-Differentiable Functions, II”, Russ. J. Math. Phys., **26**:3 (2019), 401–405.
- [3] A. G. Sergeev, “Quantum Calculus and Quasiconformal Mappings”, Math. Notes, **100**:1 (2016), 123–131.
- [4] V. S. Vladimirov, I. V. Volovich, “p-adic quantum mechanics”, Comm. Math. Phys., **123**:4 (1989), 659–676.
- [5] V. S. Vladimirov, I. V. Volovich, E. I. Zelenov, p-Adic analysis and mathematical physics, Series on Soviet and East European Mathematics, 1, World Scientific Publishing Co. Inc., River Edge, NJ, 1994.
- [6] J. V. Bellissard, M. Marcolli, K. Reihani, “Dynamical systems on spectral metric spaces”, 2010, arXiv:1008.4617 [math.OA].
- [7] J. V. Bellissard, J. C. Pearson, Noncommutative Riemannian Geometry and Diffusion on Ultrametric Cantor Sets, J. Noncommut. Geo., **3** (2009), 447-480.
- [8] S. Klimek, M. McBride, S. Rathnayake, A p -adic spectral triple, 2014, arXiv:1403.7263v2 [math.OA].
- [9] K. Phillips, “Hilbert transforms for the p-adic and p-series”, Pacific J. Math., **23** (1967), 329–347.
- [10] M. H. Taibleson, Fourier Analysis on Local Fields, Princeton Univ. Press, 1975.
- [11] J.-A. Chao, M. H. Taibleson, “A Hilbert transform for non-homogeneous martingales”, Colloquium Mathematicum, **LVIII** (1989), 111–123.
- [12] S. Sakai, C^* -algebras and W^* -algebras, Springer-Verlag, Berlin, Heidelberg, New York, 1971.
- [13] J. E. Daly, D. S. Kurtz, “BMO and singular integrals over local fields”, J. Austral. Math. Soc. (Series A), **54** (1993), 321-333.

- [14] C. Fefferman, E. M. Stein, “ H^p spaces of several variables”, Acta Math., **129** (1972), 393-400.
- [15] D. Sarason, “Function of vanishing mean oscillation”, Transaction of the AMS, **207** (1975), 391-405.
- [16] C. W. Onneweer, S. Weiyi, “Homogeneous Besov spaces on locally compact Vilenkin group”, Studia Mathematica, **XCIII** (1989), 17-39.
- [17] J. Franke, “Besov spaces on local fields”, Approximation and function spaces, Banach center publications, v. 22, Polish scientific publishers, Warsaw, 1989.
- [18] C. Gu, M. Taibleson. “Besov spaces on non-homogeneous martingales”, In: Picardello, M. A. (eds) Harmonic Analysis and Discrete Potential Theory. Springer, Boston, MA ,1992.