

MULTIFUNCTORIAL K -THEORY IS AN EQUIVALENCE OF HOMOTOPY THEORIES

NILES JOHNSON AND DONALD YAU

ABSTRACT. We show that each of the three K -theory multifunctors from small permutative categories to \mathcal{G}_* -categories, \mathcal{G}_* -simplicial sets, and connective spectra, is an equivalence of homotopy theories. For each of these K -theory multifunctors, we describe an explicit homotopy inverse functor. As a separate application of our general results about pointed diagram categories, we observe that the right-induced homotopy theory of Bohmann-Osorno \mathcal{E}_* -categories is equivalent to the homotopy theory of pointed simplicial categories.

CONTENTS

1. Introduction	1
2. Γ -Categories and \mathcal{G}_* -Categories	3
3. Right-Induced Equivalences of Homotopy Theories	7
4. Applications to K -Theory Multifunctors	12
5. Homotopy Inverses of K -Theory Multifunctors	15
References	17

1. INTRODUCTION

The Segal K -theory functor [Seg74],

$$K^{\text{Se}} : \text{PermCat}^{\text{su}} \longrightarrow \text{Sp}_{\geq 0}$$

provides a construction of connective spectra from small permutative categories. Its domain, $\text{PermCat}^{\text{su}}$, is the category of small permutative categories and strictly unital symmetric monoidal functors. Its codomain, $\text{Sp}_{\geq 0}$, is the category of connective symmetric spectra.

In (1.1), below, K^{Se} factors as a composite of three functors. First, the functor we call Segal J -theory, J^{Se} , constructs Γ -categories from small permutative categories. Then, the levelwise nerve, N_* , takes Γ -categories to Γ -simplicial sets. Finally, there is a construction of symmetric spectra from Γ -simplicial sets, denoted $K^{\mathcal{F}}$.

The composite, K^{Se} , is known to be an equivalence of stable homotopy theories. Indeed, it follows from work of Segal [Seg74], Bousfield-Friedlander [BF78], Thomason [Tho95], and Mandell [Man10] that each of the intermediate constructions is so.

Date: 23 May 2022.

2020 Mathematics Subject Classification. 18M65, 55P42, 55P48, 18F25.

Key words and phrases. multifunctor, K -theory, connective spectra.

Thus, each arrow in (1.1), *except for* $K^{\mathcal{G}}$, is an equivalence of stable homotopy theories. In the triangle at left,

$$\Pi^* : \wedge^* \mathcal{J}^{\text{se}} \longrightarrow \mathcal{J}^{\text{EM}}$$

is a natural transformation that induces componentwise stable equivalences; see Theorem 4.8. In the middle square,

$$N_* \wedge^* = \wedge^* N_*, \quad \text{and} \quad i^* N_* = N_* i^*$$

by associativity of functor composition. The two adjunctions $L \dashv i^*$ follow from Proposition 3.11. The triangle at right involving \wedge^* commutes [JY ∞ , 9.3.16]:

$$K^{\mathcal{G}} \wedge^* = K^{\mathcal{F}}.$$

At the end of Section 4 we briefly describe a separate application to the Bohmann-Osorno K -theory construction from [BO20]. This application is independent from the rest of the work here, but follows from the same general results in Section 3. Corollary 4.11 shows that the right-induced homotopy theory of Bohmann-Osorno \mathcal{E}_* -categories is equivalent to the homotopy theory of pointed simplicial categories.

In Section 5 we describe explicit homotopy inverses of the K -theory multifunctors \mathcal{J}^{EM} , $N_* \mathcal{J}^{\text{EM}}$, and K^{EM} . The following is a summary table; its details are explained in Sections 4 and 5.

K -theory multifunctors	Homotopy inverses	Note
$\mathcal{J}^{\text{EM}} : \text{PermCat}^{\text{su}} \longrightarrow \mathcal{G}_* \text{-Cat}$ (4.6)	$\mathcal{P}i^* : \mathcal{G}_* \text{-Cat} \longrightarrow \text{PermCat}^{\text{su}}$ (5.5)	i^* is not a multifunctor
$N_* \mathcal{J}^{\text{EM}} : \text{PermCat}^{\text{su}} \longrightarrow \mathcal{G}_* \text{-sSet}$ (4.6, 4.7)	$\mathcal{P}S_* i^* : \mathcal{G}_* \text{-sSet} \longrightarrow \text{PermCat}^{\text{su}}$ (5.6)	S_* is not a multifunctor
$K^{\text{EM}} : \text{PermCat}^{\text{su}} \longrightarrow \text{Sp}_{\geq 0}$ (4.9)	$\mathcal{P}S_* \mathbb{A} : \text{Sp}_{\geq 0} \longrightarrow \text{PermCat}^{\text{su}}$ (5.7)	\mathbb{A} is not a multifunctor

Remark 1.2 (Open Questions About Alternatives). There are a number of alternative ways that one might define a class of stable equivalences in $\mathcal{G}_* \text{-sSet}$ and $\mathcal{G}_* \text{-Cat}$. Among them are the following:

- Say that a morphism f of \mathcal{G}_* -simplicial sets is a $K^{\mathcal{F}}$ -stable equivalence if $K^{\mathcal{F}}(f)$ is a stable equivalence.
- There is a functor $\# : \mathcal{G}_* \text{-sSet} \longrightarrow \Gamma \text{-sSet}$ that is left adjoint to the functor \wedge^* induced by the smash product of pointed finite sets. Say that f is a $\#$ -stable equivalence if $\#(f)$ is a stable equivalence of Γ -simplicial sets.

It is an open question whether these notions of stable equivalence result in homotopy theories that are equivalent to each other, or whether either is equivalent to that of the i^* -stable equivalences defined in Section 4. Our results hold for the i^* -stable equivalences, created by i^* . \diamond

2. Γ -CATEGORIES AND \mathcal{G}_* -CATEGORIES

In this section we recall the basic definitions of Γ -categories and \mathcal{G}_* -categories from [Seg74] and [EM06], respectively. A unified treatment is given in [JY ∞ , Chapters 8 and 9].

Γ -Diagrams. We begin by recalling the indexing category \mathcal{F} for Γ -diagrams.

Definition 2.1. A *zero object* in a category \mathcal{C} is an object $*$ that is both initial and terminal. Given a zero object $*$ in \mathcal{C} , a *zero morphism* in \mathcal{C} is a morphism that factors through $*$. A *nonzero morphism* is a morphism that does not factor through $*$. For objects a and b in \mathcal{C} , we write $\mathcal{C}^b(a, b)$ for the set of nonzero morphisms $a \rightarrow b$. \diamond

Definition 2.2. Let \mathcal{F} denote the category whose objects are pointed finite sets $\underline{n} = \{0, \dots, n\}$ for natural numbers $n \geq 0$ and whose morphisms are maps of sets that preserve the basepoint element 0. The pointed finite set $\underline{0}$ is an initial and terminal object for \mathcal{F} .

The lexicographic order of smash products gives an isomorphism

$$\underline{m} \wedge \underline{n} \cong \underline{mn}$$

making $(\mathcal{F}, \wedge, \underline{1})$ a small permutative category. The symmetry isomorphism is induced by that of the smash product of pointed finite sets. See, e.g., [JY ∞ , 8.1.6] for further explanation of this structure. \diamond

Definition 2.3. Let \mathbf{Cat} denote the category of small categories, and let $\mathbf{1}$ denote a chosen terminal category, with only a single object $*$ and only a single morphism 1_* . \diamond

Definition 2.4. A Γ -category is a basepoint-preserving functor

$$(\mathcal{F}, \underline{0}) \rightarrow (\mathbf{Cat}, \mathbf{1}).$$

Morphisms of Γ -categories are natural transformations. Note that the basepoint component of a Γ -category morphism is necessarily the identity, because $\mathbf{1}$ is terminal in \mathbf{Cat} . The category of Γ -categories and morphisms is denoted $\Gamma\text{-Cat}$.

Likewise, a Γ -simplicial set is a basepoint-preserving functor

$$(\mathcal{F}, \underline{0}) \rightarrow (\mathbf{sSet}, *)$$

and a morphism of such is given by a natural transformation. The category of Γ -simplicial sets is denoted $\Gamma\text{-sSet}$. \diamond

\mathcal{G}_* -Diagrams. Next we recall the indexing category \mathcal{G} for \mathcal{G}_* -diagrams.

Definition 2.5. Let Inj denote the category whose objects are *unpointed finite sets*

$$\bar{p} = \begin{cases} \{1, \dots, p\} & \text{if } p > 0, \\ \emptyset & \text{if } p = 0, \end{cases}$$

for each natural number $p \geq 0$, and whose morphisms are injections

$$f : \bar{q} \hookrightarrow \bar{p}.$$

For each such injection f , define a functor

$$f_* : \mathcal{F}^q \rightarrow \mathcal{F}^p$$

called the *reindexing injection* as follows. Suppose given q -tuples of pointed finite sets or pointed functions

$$\langle \underline{n} \rangle = (\underline{n}_1, \dots, \underline{n}_q) \quad \text{or} \quad \langle \psi \rangle = (\psi_1, \dots, \psi_q) \in \mathcal{F}^q,$$

respectively. Define $f_* \langle \underline{n} \rangle$ to be the p -tuple whose j th entry is $\underline{n}_{f^{-1}(j)}$ and $f_* \langle \psi \rangle$ to be the p -tuple whose j th entry is $\psi_{f^{-1}(j)}$, where

$$\underline{n}_\emptyset = \underline{1} \quad \text{and} \quad \psi_\emptyset = 1_\underline{1}. \quad \diamond$$

Definition 2.6. For each unpointed finite set \bar{q} with $q > 0$, define

$$\mathcal{F}^{(q)} = \mathcal{F}^{\wedge q},$$

the q -fold smash power of pointed categories, where \mathcal{F} has basepoint $\underline{0}$. We denote the objects and morphisms of $\mathcal{F}^{(q)}$ as q -tuples

$$\langle \underline{n} \rangle = (\underline{n}_1, \dots, \underline{n}_q) \quad \text{and} \quad \langle \psi \rangle = (\psi_1, \dots, \psi_q),$$

respectively, where $\langle \underline{n} \rangle$ is identified with the basepoint of $\mathcal{F}^{(q)}$ if any $\underline{n}_i = \underline{0}$. A tuple $\langle \psi \rangle$ is a zero morphism if any ψ_i factors through $\underline{0}$ in \mathcal{F} .

For $q = 0$, define

$$\text{Ob } \mathcal{F}^{(0)} = \{*, \langle \rangle\}$$

where $*$ is the basepoint object and $\langle \rangle$ is the empty tuple. Define the morphisms of $\mathcal{F}^{(0)}$ such that $*$ is both initial and terminal and the only nonzero morphism is the identity morphism of $\langle \rangle$. \diamond

Definition 2.7. We define a small pointed category \mathcal{G} as follows.

Objects: The set of objects is the wedge of pointed sets

$$\text{Ob } \mathcal{G} = \bigvee_{q \geq 0} \text{Ob } (\mathcal{F}^{(q)}).$$

Morphisms: The basepoint $*$ is both initial and terminal in \mathcal{G} . The set of morphisms from a q -tuple $\langle \underline{n} \rangle$ to a p -tuple $\langle \underline{m} \rangle$ is

$$(2.8) \quad \begin{aligned} \mathcal{G}(\langle \underline{n} \rangle, \langle \underline{m} \rangle) &= \bigvee_{f \in \text{Inj}(\bar{q}, \bar{p})} \left(\mathcal{F}^{(p)}(f_* \langle \underline{n} \rangle, \langle \underline{m} \rangle) \right) \\ &= \bigvee_{f \in \text{Inj}(\bar{q}, \bar{p})} \left(\bigwedge_{j=1}^p \mathcal{F}(\underline{n}_{f^{-1}(j)}, \underline{m}_j) \right). \end{aligned}$$

In (2.8) for $p > 0$ we denote a morphism by a pair $(f, \langle \psi \rangle)$, where

$$f : \bar{q} \hookrightarrow \bar{p} \quad \text{in} \quad \text{Inj}$$

and

$$\langle \psi \rangle : f_* \langle \underline{n} \rangle \longrightarrow \langle \underline{m} \rangle$$

is a morphism in $\mathcal{F}^{(p)}$. A morphism $(f, \langle \psi \rangle)$ is identified with the zero morphism in $\mathcal{G}(\langle \underline{n} \rangle, \langle \underline{m} \rangle)$ if there exists a component morphism

$$\psi_j : \underline{n}_{f^{-1}(j)} \longrightarrow \underline{m}_j$$

that is a zero morphism in \mathcal{F} , factoring through $\underline{0}$.

Identities: The identity on a q -tuple $\langle \underline{n} \rangle$ is given by the pair $(1_{\bar{q}}, 1_{\langle \underline{n} \rangle})$.

Composition: The composite of morphisms

$$\langle \underline{n} \rangle \xrightarrow{(f, \langle \psi \rangle)} \langle \underline{m} \rangle \xrightarrow{(g, \langle \phi \rangle)} \langle \underline{\ell} \rangle$$

is given by the pair $(gf, \langle \phi \rangle \circ g_* \langle \psi \rangle)$.

This finishes the definition of \mathcal{G} . The composition in \mathcal{G} is associative and unital since $(gf)_* = g_* \circ f_*$ for composable injections f and g . Moreover, we note the following.

- We call q the *length* of a q -tuple $\langle \underline{n} \rangle$.
- Each morphism set $\mathcal{G}(\langle \underline{n} \rangle, \langle \underline{m} \rangle)$ is a pointed set with basepoint the zero morphism.

- For readability, we sometimes use a semicolon and write $\mathcal{G}(\langle \underline{n} \rangle; \langle \underline{m} \rangle)$ for the set of morphisms in \mathcal{G} from $\langle \underline{n} \rangle$ to $\langle \underline{m} \rangle$. \diamond

We define \mathcal{G}_* -categories and \mathcal{G}_* -simplicial sets as pointed diagrams, similar to the corresponding versions for \mathcal{F} in Definition 2.4.

Definition 2.9. The category of \mathcal{G}_* -categories is denoted $\mathcal{G}_*\text{-Cat}$ and consists of basepoint-preserving functors

$$(\mathcal{G}, *) \longrightarrow (\text{Cat}, \mathbf{1})$$

together with natural transformations between them. Likewise, the category of \mathcal{G}_* -simplicial sets is denoted $\mathcal{G}_*\text{-sSet}$ and consists of basepoint-preserving functors

$$(\mathcal{G}, *) \longrightarrow (\text{sSet}, *)$$

together with natural transformations between them. \diamond

Relating \mathcal{F} and \mathcal{G} . We will use the following functors to compare Γ -diagrams and \mathcal{G}_* -diagrams.

Definition 2.10 ([JY ∞ , 9.1.15]). The smash product of pointed finite sets defines a strict symmetric monoidal basepoint-preserving functor

$$\wedge : (\mathcal{G}, \oplus, \langle \rangle, *) \longrightarrow (\mathcal{F}, \wedge, \underline{\mathbf{1}}, \underline{\mathbf{0}})$$

given on objects by the following assignments:

$$\begin{aligned} \wedge \langle \underline{n} \rangle &= \underline{n_1 \cdots n_q} \quad \text{for } q > 0, \\ \wedge \langle \rangle &= \underline{\mathbf{1}}, \quad \text{and} \\ \wedge * &= \underline{\mathbf{0}}. \end{aligned}$$

The value of \wedge on a morphism $(f, \langle \psi \rangle)$ is given by the smash product $\wedge_j \psi_j$ together with lexicographic ordering and a permutation of factors induced by f . See [JY ∞ , 9.1.15] for further details. There are functors

$$\wedge^* : \Gamma\text{-Cat} \longrightarrow \mathcal{G}_*\text{-Cat} \quad \text{and} \quad \wedge^* : \Gamma\text{-sSet} \longrightarrow \mathcal{G}_*\text{-sSet}$$

induced by precomposition and whiskering with \wedge . \diamond

Definition 2.11. The *length-one inclusion*

$$(2.12) \quad i : \mathcal{F} \longrightarrow \mathcal{G}$$

sends each pointed finite set \underline{n} to the length-one tuple (\underline{n}) and each morphism ψ in \mathcal{F} to the pair $(1_{\overline{\mathbf{1}}}, \langle \psi \rangle)$. There are functors

$$(2.13) \quad i^* : \mathcal{G}_*\text{-Cat} \longrightarrow \Gamma\text{-Cat} \quad \text{and} \quad i^* : \mathcal{G}_*\text{-sSet} \longrightarrow \Gamma\text{-sSet}$$

induced by precomposition and whiskering with i . \diamond

The functor i is fully faithful; this will be used in Corollary 4.2 below. Moreover, there is an equality

$$\wedge \circ i = 1_{\mathcal{F}} : \mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{F}.$$

This will be used in the proof of (4.4) and (4.5) below.

We also note that, unlike \wedge , the functor $i : \mathcal{F} \longrightarrow \mathcal{G}$ is neither monoidal nor oplax monoidal. In more detail, there are, in general, no morphisms

$$i(\underline{m}) \oplus i(\underline{n}) = (\underline{m}, \underline{n}) \longrightarrow (\underline{mn}) = i(\underline{m} \wedge \underline{n})$$

in \mathcal{G} , so i does not admit any monoidal constraint. There are also no morphisms

$$i(\underline{\mathbf{1}}) = (\underline{\mathbf{1}}) \longrightarrow \langle \rangle$$

in \mathcal{G} , so i does not admit any oplax unit constraint.

3. RIGHT-INDUCED EQUIVALENCES OF HOMOTOPY THEORIES

In this section we describe the general framework that we will apply in Corollary 4.2 to show that the functors

$$i^* : \mathcal{G}_* \text{-Cat} \longrightarrow \Gamma\text{-Cat} \quad \text{and} \quad i^* : \mathcal{G}_* \text{-sSet} \longrightarrow \Gamma\text{-sSet}$$

of (2.13) induce equivalences of homotopy theories.

Equivalences of Homotopy Theories. We briefly recall the following background concepts for homotopy theories and equivalences thereof. Our work below will use only a special case from [GJO17, 2.9], stated in Proposition 3.4 below. We refer the reader to [JY22] for details of the following. See [DK80, Hir03, Toë05, BK12] for further development of the theory.

Definition 3.1.

- (1) A *relative category* is a pair (C, \mathcal{W}) consisting of a category C and a subcategory \mathcal{W} containing all of the objects of C . A *relative functor* from (C, \mathcal{W}) to (C', \mathcal{W}') is a functor from C to C' that sends morphisms of \mathcal{W} to those of \mathcal{W}' .
- (2) We say that a subcategory $\mathcal{W} \subset C$ satisfies the *2-out-of-3* property if, for each composable pair of morphisms f and g in C , whenever any two of f , g , and gf are in \mathcal{W} , then so is the third.
- (3) We say that a functor $F : C \longrightarrow C'$ *creates morphisms in \mathcal{W}* if $\mathcal{W} = F^{-1}(\mathcal{W}')$.
- (4) For each $n \geq 0$, the *relative simplex category* $(C, \mathcal{W})^{\Delta[n]}$ has objects given by functors $\Delta[n] \longrightarrow C$. The morphisms are natural transformations whose components are morphisms in \mathcal{W} .
- (5) Each relative category (C, \mathcal{W}) has a bisimplicial *classification diagram* $N^{\text{cl}}(C, \mathcal{W})$ formed by taking levelwise nerve of the relative simplex categories $(C, \mathcal{W})^{\Delta[n]}$ for $n \geq 0$.
- (6) A *complete Segal Space* [Rez01, Section 6] is a bisimplicial set X that is fibrant in the Reedy model structure and satisfies further conditions related to the Segal maps and the zeroth simplicial set $X(0)$. The complete Segal Spaces are the fibrant objects in a localization of the Reedy model structure, called the *complete Segal space model structure* [Rez01, 7.2].
- (7) A relative functor

$$F : (C, \mathcal{W}) \longrightarrow (C', \mathcal{W}')$$

is an *equivalence of homotopy theories* if the induced morphism $R N^{\text{cl}} F$ is a weak equivalence in the complete Segal space model structure, where R denotes fibrant replacement. \diamond

Relative functors are closed under composition, but they do not satisfy the 2-out-of-3 property in general. However, the following observation follows from the 2-out-of-3 property of equivalences of homotopy theories.

Lemma 3.2. *Suppose (C, \mathcal{W}) , (C', \mathcal{W}') , and (C'', \mathcal{W}'') are relative categories, and suppose*

$$F : C \longrightarrow C' \quad \text{and} \quad F' : C' \longrightarrow C''$$

are functors such that the following two statements hold.

- (1) The functor $F' : (\mathcal{C}', \mathcal{W}') \longrightarrow (\mathcal{C}'', \mathcal{W}'')$ creates morphisms in \mathcal{W}' and is an equivalence of homotopy theories.
- (2) The composite $F'F : (\mathcal{C}, \mathcal{W}) \longrightarrow (\mathcal{C}'', \mathcal{W}'')$ is a relative functor and is an equivalence of homotopy theories.

Then $F : (\mathcal{C}, \mathcal{W}) \longrightarrow (\mathcal{C}', \mathcal{W}')$ is a relative functor and is an equivalence of homotopy theories.

Next we give two results that will be used below to show that certain functors are equivalences of homotopy theories. Both follow from the observation that a natural transformation between functors induces a simplicial homotopy on nerves.

Lemma 3.3. *Suppose*

$$F, F' : (\mathcal{C}, \mathcal{W}) \longrightarrow (\mathcal{C}', \mathcal{W}')$$

are relative functors and suppose $p : F \longrightarrow F'$ is a natural transformation with each component in \mathcal{W}' . Then F is an equivalence of homotopy theories if and only if F' is an equivalence of homotopy theories.

Proof. Because the components of p are in \mathcal{W}' , it induces, for each $n \geq 0$, a natural transformation

$$p^{\Delta[n]} : F^{\Delta[n]} \longrightarrow (F')^{\Delta[n]} : (\mathcal{C}, \mathcal{W})^{\Delta[n]} \longrightarrow (\mathcal{C}', \mathcal{W}')^{\Delta[n]}$$

between the induced functors on relative simplex categories. Therefore, taking nerves, $N(p^{\Delta[n]})$ is a simplicial homotopy for each $n \geq 0$. Thus,

$$N^{\text{cl}}(p) : N^{\text{cl}}(F) \longrightarrow N^{\text{cl}}(F') : N^{\text{cl}}(\mathcal{C}, \mathcal{W}) \longrightarrow N^{\text{cl}}(\mathcal{C}', \mathcal{W}')$$

is a levelwise simplicial homotopy between morphisms of bisimplicial sets. The result then follows from functoriality of fibrant replacements and 2-out-of-3 for weak equivalences in Rezk's complete Segal space model structure. \square

The previous result will be useful on its own, but is also used in the proof of the following, from [GJO17, 2.8, 2.9]. Its proof is similar to that of [JY22, 2.12].

Proposition 3.4 ([GJO17, 2.9]). *Suppose given relative functors*

$$F : (\mathcal{C}, \mathcal{W}) \rightleftarrows (\mathcal{C}', \mathcal{W}') : G$$

such that each of the composites GF and FG is connected to the respective identity functor by a zigzag of natural transformations whose components are in \mathcal{W} and \mathcal{W}' , respectively. Then F and G are equivalences of homotopy theories.

Adjunction Between Pointed Diagram Categories. Next we describe a general context where a right adjoint of diagram categories induces an equivalence of homotopy theories. We will apply this in Section 4 to the length-one inclusion (2.12)

$$i : \mathcal{F} \longrightarrow \mathcal{G}$$

with the codomain categories $(\text{Cat}, \mathbf{1})$ and $(\text{sSet}, *)$.

Definition 3.5. Suppose $(A, *)$ is a cocomplete category with a chosen terminal object $*$. A *pointed object* in A is a pair (x, ι^x) consisting of an object x in A and a morphism $\iota^x : * \longrightarrow x$ in A . A *pointed morphism* $f : (x, \iota^x) \longrightarrow (y, \iota^y)$ between pointed objects is a morphism $f : x \longrightarrow y$ such that $f\iota^x = \iota^y$. For pointed objects (x, ι^x) and (y, ι^y) , the *wedge* $x \vee y$ is the pointed object given by the pushout of

$x \leftarrow * \rightarrow y$ in A . A wedge indexed by an empty indexing set is defined to be the chosen terminal object $*$. \diamond

Convention 3.6. Suppose \mathcal{C} and \mathcal{D} are small categories, each with a chosen initial and terminal basepoint object denoted $*$. Suppose

$$i : \mathcal{C} \longrightarrow \mathcal{D}$$

is a basepoint-preserving functor, so $i^* = *$. Suppose A is a complete and co-complete category with chosen terminal object $*$. Let \mathcal{C}_*-A and \mathcal{D}_*-A denote the categories of basepoint-preserving functors

$$(\mathcal{C}, *) \longrightarrow (A, *) \quad \text{and} \quad (\mathcal{D}, *) \longrightarrow (A, *),$$

respectively, with morphisms given by natural transformations. Then there is a functor

$$i^* : \mathcal{D}_*-A \longrightarrow \mathcal{C}_*-A$$

given by precomposition and whiskering with i . \diamond

We assume the context of Convention 3.6 for the remainder of this section.

Definition 3.7 (Functor L). Define a functor

$$L : \mathcal{C}_*-A \longrightarrow \mathcal{D}_*-A$$

by the following coend construction in A . For $X \in \mathcal{C}_*-A$ and $t \in \mathcal{D}$,

$$(LX)t = \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n), t)} Xn,$$

where, recalling Definition 2.1, $\mathcal{D}^b(i(n), t)$ denotes the set of nonzero morphisms in \mathcal{D} from $i(n)$ to t . If $\mathcal{D}^b(i(n), t)$ is empty (for example, if $n = *$ or if $t = *$), then the displayed wedge above is the chosen terminal object $*$. This ensures that $(LX)^* = *$ in A . The definition of LX on morphisms $t \rightarrow s$ in \mathcal{D} is given by functoriality of the indexing sets $\mathcal{D}^b(i(n), t)$ and of the coend. \diamond

Definition 3.8 (Unit η). Define a unit natural transformation whose component at $X \in \mathcal{C}_*-A$,

$$X \xrightarrow{\eta_X} i^* LX,$$

is given by the following composite, for each $p \in \mathcal{C}$, where the first unlabeled arrow includes to the component at $i(p)$ and the second is the coend structure morphism for $n = p$.

$$\begin{array}{ccc} Xp & \xrightarrow{\eta_{X,p}} & (i^* LX)p = \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n), i(p))} Xn \\ & \searrow & \nearrow \\ & \bigvee_{\mathcal{D}^b(i(p), i(p))} & Xp \end{array}$$

Naturality of η_X with respect to morphisms $h : p \rightarrow p'$ in \mathcal{C} follows from the coend condition that the following square commutes, where the two arrows at left are induced by $i(h)$ and Xh , and the two arrows at right are the structure

morphisms for $n = p$ and $n = p'$.

$$\begin{array}{ccc}
 & \bigvee_{\mathcal{D}^b(i(p),i(p'))} Xp & \\
 & \nearrow & \searrow \\
 \bigvee_{\mathcal{D}^b(i(p'),i(p'))} Xp & & (i^* LX)p' = \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),i(p'))} Xn \\
 & \searrow & \nearrow \\
 & \bigvee_{\mathcal{D}^b(i(p'),i(p'))} Xp' &
 \end{array}$$

Naturality of η with respect to morphisms $f : X \rightarrow X'$ in \mathcal{C}_* -A follows from naturality of f with respect to morphisms in \mathcal{C} . \diamond

Definition 3.9 (Counit ε). Define a counit natural transformation whose component at $Y \in \mathcal{D}_*$ -A,

$$Li^*Y \xrightarrow{\varepsilon_Y} Y$$

is given as follows. In the diagram below, $\theta : i(n) \rightarrow t$ is a nonzero morphism in \mathcal{D} , for $n \in \mathcal{C}$ and $t \in \mathcal{D}$. The top vertical arrow is given by the identity on the summand at θ . Then the unlabeled dashed arrow is the universal morphism out of the wedge, induced by $Y\theta$ for each $\theta \in \mathcal{D}^b(i(n), t)$.

$$(3.10) \quad \begin{array}{ccc}
 Yi(n) & & \\
 \downarrow & \searrow^{Y\theta} & \\
 \bigvee_{\mathcal{D}^b(i(n),t)} Yi(n) & & \\
 \downarrow & \dashrightarrow^{\varepsilon_{Y,t}} & \\
 (Li^*Y)t = \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),t)} Yi(n) & \dashrightarrow^{\varepsilon_{Y,t}} & Yt
 \end{array}$$

The bottom vertical arrow in the above diagram is the structure morphism for the coend, and $\varepsilon_{Y,t}$ is the induced universal morphism. Naturality of ε_Y with respect to morphisms $t \rightarrow t'$ in \mathcal{D} follows from naturality of the universal morphisms above. Naturality of ε with respect to morphisms $g : Y \rightarrow Y'$ in \mathcal{D}_* -A follows from naturality of g with respect to morphisms in \mathcal{D} and uniqueness of the induced morphism

$$\int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),t)} Yi(n) \rightarrow Y't$$

given by either $\varepsilon_{Y',t}$ and g_t or by $\bigvee_{\theta} g_{i(n)}$ and $\varepsilon_{Y,t}$. \diamond

Proposition 3.11. *In the context of Convention 3.6, there is an adjunction of categories*

$$L : \mathcal{C}_* \text{-A} \rightleftarrows \mathcal{D}_* \text{-A} : i^*.$$

Proof. The unit and counit are those of Definitions 3.8 and 3.9, respectively. The two triangle identities are verified componentwise as follows.

First, for $X \in \mathcal{C}_* \text{-A}$ and $t \in \mathcal{D}$, consider the composite
(3.12)

$$(LX)t = \int^{p \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(p),t)} Xp \xrightarrow{L(\eta_{X,p})t} \int^{p \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(p),t)} \left[\int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),i(p))} Xn \right] \xrightarrow{\varepsilon_{LX,t}} \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),t)} Xn = (LX)t.$$

The above composite is the unique universal morphism out of the coend $(LX)t$ induced by the following composite, where we use the notation change $m = p$ on the left and $m = n$ on the right:

$$\bigvee_{\mathcal{D}^b(i(m),t)} Xm \longrightarrow \bigvee_{\mathcal{D}^b(i(m),t)} \bigvee_{\mathcal{D}^b(i(m),i(m))} Xm \longrightarrow \bigvee_{\mathcal{D}^b(i(m),t)} Xm.$$

In the above composite, the first morphism is induced by $\bigvee_{\mathcal{D}^b(i(m),t)} 1_{i(m)}$ on indexing sets, and the second by composition on indexing sets. Thus the above composite is the identity, and therefore so is (3.12).

Second, for $Y \in \mathcal{D}_* \text{-A}$ and $p \in \mathcal{C}$, consider the composite

$$(3.13) \quad \begin{array}{ccc} (i^*Y)p & \xrightarrow{\eta_{i^*Y,p}} & \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),i(p))} Yi(n) \\ & \searrow & \downarrow (i^* \varepsilon_Y)_p = \varepsilon_{i^*Y,i(p)} \\ & \bigvee_{\mathcal{D}^b(i(p),i(p))} Yi(p) & \\ & \nearrow & \\ & & Yi(p) = (i^*Y)p. \end{array}$$

The above composite is the identity because the first morphism is induced by including to the component at $1_{i(p)}$ and the composite to $Yi(p)$ is that defining $\varepsilon_{i^*Y,i(p)}$ in (3.10) with $\theta = 1_{i(p)}$. \square

Lemma 3.14. *Suppose, in the context of Convention 3.6, that i is fully faithful. Then the unit η of the adjunction $L \dashv i^*$ is a natural isomorphism.*

Proof. For X in $\mathcal{C}_* \text{-A}$ and $p \in \mathcal{C}$, the Yoneda Density Theorem [JY ∞ , III.3.7.8] gives an isomorphism

$$\int^{n \in \mathcal{C}} \bigvee_{\mathcal{C}^b(n,p)} Xn \xrightarrow{\cong} Xp$$

that is natural with respect to morphisms $p \rightarrow p'$ in \mathcal{C} . The assumption that i is fully faithful implies there is an isomorphism

$$\mathcal{D}^b(i(n),i(p)) \cong \mathcal{C}^b(n,p) \quad \text{for } n, p \in \mathcal{C}.$$

The above isomorphism, applied to indexing sets, gives the middle isomorphism below:

$$(3.15) \quad Xp \xrightarrow{\eta_{X,p}} \int^{n \in \mathcal{C}} \bigvee_{\mathcal{D}^b(i(n),i(p))} Xn \cong \int^{n \in \mathcal{C}} \bigvee_{\mathcal{C}^b(n,p)} Xn \xrightarrow{\cong} Xp.$$

The above composite is the identity on Xp because $\eta_{X,p}$ includes Xp at the component indexed by $1_{i(p)}$ and the Yoneda Density isomorphism is induced by applying X to the morphisms of $\mathcal{C}^b(n, p)$. Therefore $\eta_{X,p}$ is an isomorphism with inverse given by the other morphisms of (3.15). \square

Equivalences of Homotopy Theories Between Pointed Diagram Categories.

Definition 3.16. Suppose, in the context of Convention 3.6, that $\mathcal{S} \subset \mathcal{C}_*-A$ is a subcategory containing all objects of \mathcal{C}_*-A , so $(\mathcal{C}_*-A, \mathcal{S})$ is a relative category. The *right-induced subcategory* of \mathcal{D}_*-A is the subcategory

$$\mathcal{S}^i = (i^*)^{-1}(\mathcal{S}) \subset \mathcal{D}_*-A$$

consisting of morphisms f such that $i^*(f) \in \mathcal{S}$. Thus, $(\mathcal{D}_*-A, \mathcal{S}^i)$ is a relative category and

$$i^* : (\mathcal{D}_*-A, \mathcal{S}^i) \longrightarrow (\mathcal{C}_*-A, \mathcal{S})$$

is a relative functor with the morphisms of \mathcal{S}^i created by i^* . If \mathcal{S} contains all isomorphisms, then so does \mathcal{S}^i . If \mathcal{S} satisfies the 2-out-of-3 property, then so does \mathcal{S}^i . \diamond

Theorem 3.17. *In the context of Convention 3.6, suppose $\mathcal{S} \subset \mathcal{C}_*-A$ is a subcategory containing all objects and suppose, moreover, that the following three conditions hold.*

- *The functor $i : \mathcal{C} \longrightarrow \mathcal{D}$ is fully faithful.*
- *The subcategory \mathcal{S} contains all isomorphisms.*
- *The subcategory \mathcal{S} has the 2-out-of-3 property.*

Let $\mathcal{S}^i = (i^)^{-1}(\mathcal{S})$ be the right-induced subcategory of \mathcal{D}_*-A . Then the adjunction $L \dashv i^*$ from Definition 3.7 induces an equivalence of homotopy theories*

$$L : (\mathcal{C}_*-A, \mathcal{S}) \xleftarrow{\sim} (\mathcal{D}_*-A, \mathcal{S}^i) : i^*.$$

Proof. Since i is fully faithful, each component of the unit η is an isomorphism by Lemma 3.14. Therefore each η_X is a morphism of \mathcal{S} . This implies that L is a relative functor, by naturality of η and the 2-out-of-3 property for \mathcal{S} . For each component ε_Y , the right triangle identity

$$(i^* \varepsilon_Y) \circ \eta_{i^*Y} = 1_{i^*Y}$$

from (3.13), together with the 2-out-of-3 property for \mathcal{S} , implies that $i^* \varepsilon_Y$ is a morphism of \mathcal{S} and hence ε_Y is a morphism of $\mathcal{S}^i = (i^*)^{-1}(\mathcal{S})$. Since the components of η , respectively ε , are morphisms of \mathcal{S} , respectively \mathcal{S}^i , the result then follows from Proposition 3.4. \square

4. APPLICATIONS TO K -THEORY MULTIFUNCTORS

This section contains our main applications of Theorem 3.17, with $i : \mathcal{F} \longrightarrow \mathcal{G}$ the inclusion of length-one tuples from Definition 2.11. At the end of the section, we give a separate application to Bohmann-Osorno \mathcal{E}_* -categories.

Definition 4.1. Let $\mathcal{S} \subset \text{Sp}_{\geq 0}$ denote the subcategory of *stable equivalences* of connective spectra. Then, in each of the following, let \mathcal{S} denote the subcategory of morphisms created by the indicated functor:

$$(\text{PermCat}^{\text{su}}, \mathcal{S}) \xrightarrow{\text{J}^{\text{se}}} (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow{N_*} (\Gamma\text{-sSet}, \mathcal{S}) \xrightarrow{\text{K}^{\mathcal{F}}} (\text{Sp}_{\geq 0}, \mathcal{S}).$$

In each case, we call morphisms in \mathcal{S} *stable equivalences*. Let $\mathcal{S}^i \subset \mathcal{G}_* \text{-Cat}$ and $\mathcal{S}^i \subset \mathcal{G}_* \text{-sSet}$ denote the right-induced subcategories, created by

$$i^* : \mathcal{G}_* \text{-Cat} \longrightarrow \Gamma \text{-Cat} \quad \text{and} \quad i^* : \mathcal{G}_* \text{-sSet} \longrightarrow \Gamma \text{-sSet}.$$

The morphisms in \mathcal{S}^i are called *i^* -stable equivalences*. \diamond

Since $i : \mathcal{F} \longrightarrow \mathcal{G}$ is fully faithful and the stable equivalences of connective spectra include all isomorphisms and satisfy 2-out-of-3, we have the following application of Theorem 3.17.

Corollary 4.2. *The adjunctions $L \dashv i^*$ are equivalences of homotopy theories*

$$L : (\Gamma \text{-Cat}, \mathcal{S}) \xleftarrow{\sim} (\mathcal{G}_* \text{-Cat}, \mathcal{S}^i) : i^* \quad \text{and}$$

$$L : (\Gamma \text{-sSet}, \mathcal{S}) \xleftarrow{\sim} (\mathcal{G}_* \text{-sSet}, \mathcal{S}^i) : i^*.$$

Now we use Corollary 4.2 to prove our main results about equivalences of homotopy theories in (1.1).

Theorem 4.3. *The following are equivalences of homotopy theories:*

$$(4.4) \quad \wedge^* : (\Gamma \text{-Cat}, \mathcal{S}) \xrightarrow{\sim} (\mathcal{G}_* \text{-Cat}, \mathcal{S}^i),$$

$$(4.5) \quad \wedge^* : (\Gamma \text{-sSet}, \mathcal{S}) \xrightarrow{\sim} (\mathcal{G}_* \text{-sSet}, \mathcal{S}^i),$$

$$(4.6) \quad J^{\text{EM}} : (\text{PermCat}^{\text{su}}, \mathcal{S}) \xrightarrow{\sim} (\mathcal{G}_* \text{-Cat}, \mathcal{S}^i), \quad \text{and}$$

$$(4.7) \quad N_* : (\mathcal{G}_* \text{-Cat}, \mathcal{S}^i) \xrightarrow{\sim} (\mathcal{G}_* \text{-sSet}, \mathcal{S}^i).$$

Proof. Consider the following.

$$\begin{array}{ccc} & & (\Gamma \text{-Cat}, \mathcal{S}) \\ & \nearrow^{J^{\text{Se}}} & \uparrow \\ (\text{PermCat}^{\text{su}}, \mathcal{S}) & \xrightarrow{\sim} & \uparrow i^* \\ & \searrow_{J^{\text{EM}}} & \downarrow \wedge^* \\ & & (\mathcal{G}_* \text{-Cat}, \mathcal{S}^i) \end{array}$$

The functors J^{Se} and i^* create stable, respectively i^* -stable, equivalences, and therefore are relative functors. Work of Thomason [Tho95] shows that J^{Se} is an equivalence of homotopy theories. An independent proof by Mandell [Man10] constructs an inverse directly. Corollary 4.2 shows that i^* is an equivalence of homotopy theories.

For (4.4), note that we have

$$\wedge \circ i = 1 : \mathcal{F} \longrightarrow \mathcal{F}$$

and hence $i^* \wedge^* = 1$, the identity on $\Gamma \text{-Cat}$. This implies that \wedge^* is a relative functor and an equivalence of homotopy theories by Lemma 3.2 with $F' = i^*$ and $F = \wedge^*$. The proof for (4.5) is the same by replacing Cat with sSet.

For (4.6), a similar reduction for smash products over singletons shows that $J^{\text{Se}} = i^* J^{\text{EM}}$ by unpacking the definitions [JY ∞ , 10.3.1, 10.3.27]. This implies that J^{EM} is a relative functor and an equivalence of homotopy theories by Lemma 3.2 with $F' = i^*$ and $F = J^{\text{EM}}$.

For (4.7), consider the following, which commutes by associativity of functor composition.

$$\begin{array}{ccc} (\Gamma\text{-Cat}, \mathcal{S}) & \xrightarrow[\sim]{N_*} & (\Gamma\text{-sSet}, \mathcal{S}) \\ i^* \uparrow \sim & & \sim \uparrow i^* \\ (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) & \xrightarrow{N_*} & (\mathcal{G}_*\text{-sSet}, \mathcal{S}^i) \end{array}$$

The two instances of i^* create i^* -stable equivalences and are equivalences of homotopy theories by Corollary 4.2. The top functor N_* creates the stable equivalences of Γ -categories by definition. It induces an equivalence of homotopy theories by work of Mandell [Man10]. Therefore, by commutativity of the diagram and Lemma 3.2 again, the lower instance of N_* is a relative functor and an equivalence of homotopy theories. \square

Next we recall the following result due to Elmendorf-Mandell [EM06, JY ∞]. We will use this to prove that K^{EM} is an equivalence of homotopy theories.

Theorem 4.8 ([JY ∞ , 10.6.10]). *The natural transformation Π^* in (1.1) induces a componentwise stable equivalence from Segal K -theory, K^{Se} , to Elmendorf-Mandell K -theory, K^{EM} . That is,*

$$K^{\mathcal{G}} N_* \Pi^* : K^{\mathcal{F}} N_* J^{\text{Se}} C = K^{\mathcal{G}} N_* \wedge^* J^{\text{Se}} C \longrightarrow K^{\mathcal{G}} N_* J^{\text{EM}} C$$

is a stable equivalence of connective spectra for each small permutative category C .

The functor K^{EM} is the composite

$$K^{\text{EM}} = K^{\mathcal{G}} N_* J^{\text{EM}}.$$

While

$$J^{\text{EM}} : (\text{PermCat}^{\text{su}}, \mathcal{S}) \longrightarrow (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) \text{ and } N_* : (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) \longrightarrow (\mathcal{G}_*\text{-sSet}, \mathcal{S}^i)$$

are relative functors, $K^{\mathcal{G}}$ may not be. Thus the first step in proving the following result is to use naturality of Π^* to show that the composite K^{EM} is a relative functor.

Theorem 4.9. *The Elmendorf-Mandell K -theory functor is an equivalence of homotopy theories*

$$K^{\text{EM}} : (\text{PermCat}^{\text{su}}, \mathcal{S}) \xrightarrow{\sim} (\text{Sp}_{\geq 0}, \mathcal{S}).$$

Proof. For a strictly unital symmetric monoidal functor

$$F : C \longrightarrow D \text{ in } \text{PermCat}^{\text{su}}$$

we have the following naturality square.

$$\begin{array}{ccc} K^{\text{Se}}(C) & \xrightarrow{K^{\mathcal{G}} N_*(\Pi_C^*)} & K^{\text{EM}}(C) \\ K^{\text{Se}}(F) \downarrow & & \downarrow K^{\text{EM}}(F) \\ K^{\text{Se}}(D) & \xrightarrow{K^{\mathcal{G}} N_*(\Pi_D^*)} & K^{\text{EM}}(D) \end{array}$$

By Theorem 4.8, the components of

$$p = K^{\mathcal{G}} N_* \Pi^* : K^{\text{Se}} \longrightarrow K^{\text{EM}}$$

are stable equivalences of connective spectra. If F is a stable equivalence in $\text{PermCat}^{\text{su}}$, then $K^{\text{Se}}(F)$, and therefore also $K^{\text{EM}}(F)$, are stable equivalences.

Homotopy Inverse of J^{EM} . By Remark 4.10

$$(5.2) \quad \Pi^* : \wedge^* J^{\text{Se}} \longrightarrow J^{\text{EM}} : (\text{PermCat}^{\text{su}}, \mathcal{S}) \longrightarrow (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i)$$

induces a natural isomorphism in the stable homotopy categories, obtained by inverting the respective classes of stable equivalences. By Corollary 4.2 and Theorem 4.3, both relative functors

$$(5.3) \quad (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow[\sim]{\wedge^*} (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) \xrightarrow[\sim]{i^*} (\Gamma\text{-Cat}, \mathcal{S})$$

are equivalences of homotopy theories, and the analogous statement holds with sSet in place of Cat . The equality $i^* \wedge^* = 1$ implies that the functors \wedge^* and i^* are *homotopy inverses* of each other. This means that, passing to the stable homotopy categories, they induce mutually inverse equivalences of categories.

A homotopy inverse of Segal J -theory, J^{Se} , is given by Mandell's inverse K -theory functor [Man10, Theorem 4.5]

$$(5.4) \quad \mathcal{P} : (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow{\sim} (\text{PermCat}^{\text{su}}, \mathcal{S}).$$

This is a non-symmetric Cat -enriched multifunctor by [JY ∞ b, Theorem 1.3]. Combining (5.2) through (5.4), a homotopy inverse of Elmendorf-Mandell J -theory, J^{EM} , is given by the following composite.

$$(5.5) \quad (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) \xrightarrow[\sim]{i^*} (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow[\sim]{\mathcal{P}} (\text{PermCat}^{\text{su}}, \mathcal{S})$$

We note that—while J^{EM} and \mathcal{P} are Cat -enriched multifunctors (non-symmetric in the case of \mathcal{P})—the functor i^* does *not* appear to be a multifunctor, even in the non-symmetric sense. In particular, i^* does not admit any reasonable monoidal constraint that is compatible with the monoidal structures on $\Gamma\text{-Cat}$ and $\mathcal{G}_*\text{-Cat}$.

Homotopy Inverse of $N_* J^{\text{EM}}$. A homotopy inverse of

$$N_* : (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow{\sim} (\Gamma\text{-sSet}, \mathcal{S})$$

is given by Mandell's functor S_* in [Man10, Proposition 2.7], denoted \mathcal{S} there. This is *not* a multifunctor, as discussed in the introduction of [JY ∞ b]. By (5.2) and the equality

$$N_* \wedge^* J^{\text{Se}} = \wedge^* N_* J^{\text{Se}},$$

a homotopy inverse of the composite

$$(\text{PermCat}^{\text{su}}, \mathcal{S}) \xrightarrow[\sim]{J^{\text{EM}}} (\mathcal{G}_*\text{-Cat}, \mathcal{S}^i) \xrightarrow[\sim]{N_*} (\mathcal{G}_*\text{-sSet}, \mathcal{S}^i)$$

is given by the following composite.

$$(5.6) \quad (\mathcal{G}_*\text{-sSet}, \mathcal{S}^i) \xrightarrow[\sim]{i^*} (\Gamma\text{-sSet}, \mathcal{S}) \xrightarrow[\sim]{S_*} (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow[\sim]{\mathcal{P}} (\text{PermCat}^{\text{su}}, \mathcal{S})$$

Homotopy Inverse of K^{EM} . A homotopy inverse of

$$K^{\mathcal{F}} : (\Gamma\text{-sSet}, \mathcal{S}) \xrightarrow{\sim} (\text{Sp}_{\geq 0}, \mathcal{S})$$

is given by Segal's functor \mathbb{A} in [Seg74, Definition 3.1]. The adjoint pair $K^{\mathcal{F}} \dashv \mathbb{A}$ is a Quillen equivalence by [BF78, Theorem 5.8]. By (5.2) and the equalities

$$K^{\mathcal{G}} N_* \wedge^* J^{\text{Se}} = K^{\mathcal{G}} \wedge^* N_* J^{\text{Se}} = K^{\mathcal{F}} N_* J^{\text{Se}},$$

a homotopy inverse of Elmendorf-Mandell K -theory, $K^{\text{EM}} = K^{\mathcal{G}}N_*J^{\text{EM}}$, is given by the following composite along the top of (5.1).

$$(5.7) \quad (\text{Sp}_{\geq 0}, \mathcal{S}) \xrightarrow[\sim]{\mathbb{A}} (\Gamma\text{-sSet}, \mathcal{S}) \xrightarrow[\sim]{S_*} (\Gamma\text{-Cat}, \mathcal{S}) \xrightarrow[\sim]{\mathcal{P}} (\text{PermCat}^{\text{su}}, \mathcal{S})$$

We note that—while K^{EM} and \mathcal{P} are enriched multifunctors (non-symmetric in the case of \mathcal{P})—the functor \mathbb{A} does *not* appear to be a multifunctor, even in the non-symmetric sense. Similar to i^* , the functor \mathbb{A} does not admit any reasonable monoidal constraint that is compatible with the monoidal structures on $\Gamma\text{-sSet}$ and $\text{Sp}_{\geq 0}$.

REFERENCES

- [BK12] C. Barwick and D. M. Kan, *A characterization of simplicial localization functors and a discussion of DK equivalences*, *Indag. Math. (N.S.)* **23** (2012), no. 1-2, 69–79. doi:[10.1016/j.indag.2011.10.001](https://doi.org/10.1016/j.indag.2011.10.001) (cit. on p. 7).
- [BO20] A. M. Bohmann and A. M. Osorno, *A multiplicative comparison of Segal and Waldhausen K-theory*, *Math. Z.* **295** (2020), no. 3-4, 1205–1243. doi:[10.1007/s00209-019-02394-7](https://doi.org/10.1007/s00209-019-02394-7) (cit. on pp. 3, 15).
- [BF78] A. K. Bousfield and E. M. Friedlander, *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, *Lecture Notes in Math.*, vol. 658, Springer, Berlin, 1978, pp. 80–130. (cit. on pp. 1, 15, 16).
- [DK80] W. G. Dwyer and D. M. Kan, *Calculating simplicial localizations*, *J. Pure Appl. Algebra* **18** (1980), no. 1, 17–35. doi:[10.1016/0022-4049\(80\)90113-9](https://doi.org/10.1016/0022-4049(80)90113-9) (cit. on p. 7).
- [EM06] A. D. Elmendorf and M. A. Mandell, *Rings, modules, and algebras in infinite loop space theory*, *Adv. Math.* **205** (2006), no. 1, 163–228. doi:[10.1016/j.aim.2005.07.007](https://doi.org/10.1016/j.aim.2005.07.007) (cit. on pp. 2, 3, 14).
- [EM09] ———, *Permutative categories, multicategories and algebraic K-theory*, *Algebr. Geom. Topol.* **9** (2009), no. 4, 2391–2441. doi:[10.2140/agt.2009.9.2391](https://doi.org/10.2140/agt.2009.9.2391) (cit. on p. 2).
- [GJO17] N. Gurski, N. Johnson, and A. M. Osorno, *K-theory for 2-categories*, *Adv. Math.* **322** (2017), 378–472. doi:[10.1016/j.aim.2017.10.011](https://doi.org/10.1016/j.aim.2017.10.011) (cit. on pp. 7, 8).
- [Hir03] P. S. Hirschhorn, *Model categories and their localizations*, *Mathematical Surveys and Monographs*, vol. 99, American Mathematical Society, Providence, RI, 2003. (cit. on p. 7).
- [JY ∞] N. Johnson and D. Yau, *Bimonoidal Categories, E_n -Monoidal Categories, and Algebraic K-Theory. Volume III: From Categories to Structured Ring Spectra*, available at <https://nilesjohnson.net>. (cit. on pp. 2, 3, 4, 6, 11, 13, 14, 15).
- [JY ∞ b] ———, *Multifunctorial inverse K-theory*. arXiv:[2109.01430](https://arxiv.org/abs/2109.01430) (cit. on p. 16).
- [JY21] ———, *2-Dimensional Categories*, Oxford University Press, New York, 2021. doi:[10.1093/oso/9780198871378.001.0001](https://doi.org/10.1093/oso/9780198871378.001.0001) (cit. on p. 2).
- [JY22] ———, *Multicategories model all connective spectra*, To appear in *Homology, Homotopy and Applications* (2022). arXiv:[2111.08653](https://arxiv.org/abs/2111.08653) (cit. on pp. 7, 8).
- [Man10] M. A. Mandell, *An inverse K-theory functor*, *Doc. Math.* **15** (2010), 765–791. (cit. on pp. 1, 13, 14, 16).
- [Rez01] C. Rezk, *A model for the homotopy theory of homotopy theory*, *Trans. Amer. Math. Soc.* **353** (2001), no. 3, 973–1007 (electronic). doi:[10.1090/S0002-9947-00-02653-2](https://doi.org/10.1090/S0002-9947-00-02653-2) (cit. on p. 7).
- [Seg74] G. Segal, *Categories and cohomology theories*, *Topology* **13** (1974), 293–312. doi:[10.1016/0040-9383\(74\)90022-6](https://doi.org/10.1016/0040-9383(74)90022-6) (cit. on pp. 1, 3, 15, 16).
- [Tho95] R. W. Thomason, *Symmetric monoidal categories model all connective spectra*, *Theory Appl. Categ.* **1** (1995), No. 5, 78–118. (cit. on pp. 1, 13).
- [Toë05] B. Toën, *Vers une axiomatisation de la théorie des catégories supérieures*, *K-Theory* **34** (2005), no. 3, 233–263. doi:[10.1007/s10977-005-4556-6](https://doi.org/10.1007/s10977-005-4556-6) (cit. on p. 7).

Email address: johnson.5320@osu.edu, yau.22@osu.edu

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY AT NEWARK, 1179 UNIVERSITY DRIVE, NEWARK, OH 43055, USA