

ADMISSIBLE MODULES AND NORMALITY OF CLASSICAL NILPOTENT ORBITS II

DAN BARBASCH AND KAYUE DANIEL WONG

ABSTRACT. As a sequel to [BW1], we study the character formula of the Brylinski model $\mathcal{B}(\overline{\mathcal{O}})$ for classical nilpotent varieties $\overline{\mathcal{O}}$. As a consequence, one can compute the multiplicities of all K -types of the ring of regular functions $R(\overline{\mathcal{O}})$ for all classical nilpotent varieties.

1. INTRODUCTION

Let $G = Sp(2n, \mathbb{C})$ or $SO(2n + \delta, \mathbb{C})$ ($\delta \in \{0, 1\}$) be a complex classical Lie group. In [Br], Brylinski defined a (\mathfrak{g}, K) -module $\mathcal{B}(\overline{\mathcal{O}})$ for all classical nilpotent orbits, whose $G \cong K_c$ -spectrum (here the subscript c denotes complexification) satisfies

$$\mathcal{B}(\overline{\mathcal{O}})|_{K_c} \cong R(\overline{\mathcal{O}})|_G,$$

where $R(\overline{\mathcal{O}})$ is the ring of regular functions of $\overline{\mathcal{O}}$. In [BW1], it is proved that $\mathcal{B}(\overline{\mathcal{O}})$ is the cyclic submodule of a ‘deformed’ unipotent representation $\mathcal{B}(\mathcal{O})$ attached to \mathcal{O} satisfying $\mathcal{B}(\mathcal{O})|_{K_c} \cong R(\mathcal{O})$ (see (5) below). Moreover, the diminutive K_c -type multiplicities of $\mathcal{B}(\overline{\mathcal{O}})$ are obtained in [BW1], resulting in an alternative proof on the classification of (non-)normal classical nilpotent varieties in [KP].

In this paper, we obtain the character formula of $\mathcal{B}(\overline{\mathcal{O}})$. As a result, the multiplicities of all $G \cong K_c$ -types of $R(\overline{\mathcal{O}})$ can be effectively computed. An important tool of our study of $\mathcal{B}(\overline{\mathcal{O}})$ is a reduction theorem, which reduces the problem of studying to $\mathcal{B}(\overline{\mathcal{O}})$ to that of $\mathcal{B}(\overline{\mathcal{O}_{gen}})$, where \mathcal{O}_{gen} is the *generic part* of \mathcal{O} (Theorem 2.2).

1.1. Notations. We introduce the notations and terminologies needed for this paper. Many of them are introduced in [BW1]:

(a) Irreducible representations for complex Lie groups. Let G be a complex Lie group viewed as a real Lie group with Cartan involution θ . Consider $H = T \cdot A$ be the Cartan decomposition of the Cartan subgroup H of G . Then the *Langlands parameter* of any irreducible module (\mathfrak{g}_c, K_c) -modules is a pair $(\lambda_L; \lambda_R)$ such that $\mu := \lambda_L - \lambda_R$ is the parameter of a character of T , and $\nu := \lambda_L + \lambda_R$ the A -character.

The *principal series representation* associated to $(\lambda_L; \lambda_R)$ is the (\mathfrak{g}_c, K_c) -module

$$X(\lambda_L, \lambda_R) = \text{Ind}_B^G(e^\mu \otimes e^\nu \otimes 1)_{K\text{-finite}},$$

Date: May 18, 2022.

where the symbol Ind refers to Harish-Chandra induction, and let $\overline{X}(\lambda_L; \lambda_R)$ be the unique irreducible subquotient of $X(\lambda_L; \lambda_R)$ containing the K_c -type with extremal weight μ . This is called the *Langlands subquotient*. The infinitesimal character of $X(\lambda_L; \lambda_R)$ and $\overline{X}(\lambda_L; \lambda_R)$, when \mathfrak{g}_c is identified with $\mathfrak{g} \times \mathfrak{g}$, is $(\lambda_L; \lambda_R)$.

The case of the orthogonal group is dealt with via Clifford theory. We use Weyl's parametrization of the finite dimensional representations of the orthogonal groups (see page 6 of [AB]). The highest weight of a K_c -type will be denoted

$$(1) \quad \mu = (a_1 \geq \cdots \geq a_n | \pm 1)$$

whenever $a_n = 0$, and the Langlands quotients will acquire a \pm whenever the corresponding lowest K_c -types are in different irreducible quotients.

If we need to specify the group, the standard module and Langlands quotient will acquire a subscript, e.g. $X_G(\lambda_L; \lambda_R)$ or $\overline{X}_G(\lambda_L; \lambda_R)$ (respectively $X_G(\lambda_L; \lambda_R | \pm 1)$ or $\overline{X}_G(\lambda_L; \lambda_R | \pm 1)$ for orthogonal groups).

(b) Shorthand for parabolically induced modules. Let G' be the Lie group with the same type as G of lower rank, and Ψ be a representation of G' . We write

$$I^G \left(\begin{pmatrix} \lambda_L \\ \lambda_R \end{pmatrix} \boxtimes \cdots \boxtimes \begin{pmatrix} \lambda'_L \\ \lambda'_R \end{pmatrix} \boxtimes \Psi \right) := \text{Ind}_{GL \times \cdots \times GL \times G'}^G (\overline{X}_{GL}(\lambda_L; \lambda_R) \boxtimes \cdots \boxtimes \overline{X}_{GL}(\lambda'_L; \lambda'_R) \boxtimes \Psi).$$

Similarly, we write

$$I^G \left(\begin{pmatrix} \lambda_L \\ \lambda_R \end{pmatrix} \boxtimes \cdots \boxtimes \begin{pmatrix} \lambda'_L \\ \lambda'_R \end{pmatrix} \right) := \text{Ind}_{GL \times \cdots \times GL}^G (\overline{X}_{GL}(\lambda_L; \lambda_R) \boxtimes \cdots \boxtimes \overline{X}_{GL}(\lambda'_L; \lambda'_R)).$$

(c) Diminutive K_c -types. Let G be a classical complex Lie group, and V_μ be the irreducible, finite-dimensional K_c -type with highest weight μ (or Weyl's parametrization (1) for orthogonal groups). The **diminutive K_c -types** of G are:

$$\begin{cases} V_{(1^k, 0^{n-2k}, -1^k)} & (0 \leq k \leq \lfloor \frac{n}{2} \rfloor) & \text{for } G = GL(n, \mathbb{C}) \\ V_{(1^k, 0^{n-k} | (-1)^k)} & (0 \leq k \leq n) & \text{for } G = O(2n+1, \mathbb{C}) \\ V_{(1^{2k}, 0^{n-2k})} & (0 \leq k \leq \lfloor \frac{n}{2} \rfloor) & \text{for } G = Sp(2n, \mathbb{C}) \\ V_{(1^{2k}, 0^{n-2k} | \pm 1)} & (0 \leq k \leq \lfloor \frac{n}{2} \rfloor) & \text{for } G = O(2n, \mathbb{C}) \end{cases}.$$

In particular, the diminutive K_c -types are equal to $\wedge^{2\ell} \mathbb{C}^{2n+\delta}$ ($\delta \in \{0, 1\}$) for orthogonal groups, and $\wedge^{2\ell} \mathbb{C}^{2n} / \wedge^{2\ell-1} \mathbb{C}^{2n+\delta}$ for symplectic groups.

Let X, Y be two admissible (\mathfrak{g}_c, K_c) -modules. We write

$$X \approx Y$$

if X and Y have the same composition factors with diminutive lowest K_c -types (with multiplicities). Note that this is a weaker notion than \xrightarrow{dm} defined in [BW1].

(d) Shorthand for Langlands parameters. Let $a, A \in \frac{1}{2}\mathbb{Z}$ be such that $A - a \in \mathbb{Z}$. A *string* is an ascending sequence of numbers $(a \dots A) := (a, a+1, \dots, A-1, A)$ (if $A - a < 0$ then the string is empty).

For $x, y \in \mathbb{Z}_{\geq -1}$ of the same parity, define $\lambda[x, y]^+$ by the spherical Langlands parameter

$$(2) \quad \lambda[x, y]^+ := \begin{pmatrix} -\frac{y}{2} + 1 \dots \frac{x}{2} \\ -\frac{y}{2} + 1 \dots \frac{x}{2} \end{pmatrix}$$

If $x \geq y$, we denote by $\lambda[x, y]^-$ the **non-spherical** parameter

$$\lambda[x, y]^- := \begin{pmatrix} -\frac{y}{2} + 1 \dots \frac{y}{2}, & \frac{y}{2} + 1 \dots \frac{x}{2} \\ -\frac{y}{2} \dots \frac{y}{2} - 1, & \frac{y}{2} + 1 \dots \frac{x}{2} \end{pmatrix}.$$

Note that

$$(3) \quad \begin{aligned} \overline{X}_{GL}(\lambda[x, y]^+) &= |\det|^{x-y}, \\ \overline{X}_{GL}(\lambda[x, y]^-) &= \text{Ind}_{GL(y) \times GL(\frac{x-y}{2})}^{GL(\frac{x+y}{2})} (\det \boxtimes |\det|^y) \end{aligned}$$

(e) Nilpotent orbits. A classical nilpotent orbit \mathcal{O} is denoted by the column sizes of its corresponding partition

$$(4) \quad \mathcal{O} = \begin{cases} (c_0 \geq c_1 \geq \dots \geq c_{2p} \geq c_{2p+1}) & \text{if } G = Sp(2n, \mathbb{C}), \\ (c_1 \geq c_2 \geq \dots \geq c_{2p} \geq c_{2p+1}) & \text{if } G = O(2n + \delta, \mathbb{C}) \ (\delta \in \{0, 1\}) \end{cases},$$

such that $c_{2i-1} + c_{2i}$ is an even integer for all i . We say \mathcal{O} is even or odd if c_i are all even or odd integers respectively.

Let $\tau(\mathcal{O}) := \{i \mid c_{2i-1} = c_{2i}\}$. Define \mathcal{O}' by

$$\mathcal{O}' := \begin{cases} (c'_0 \geq c'_1 \geq \dots \geq c'_{2q} \geq c'_{2q+1}) & \text{if } G' = Sp(2n', \mathbb{C}), \\ (c'_1 \geq c'_2 \geq \dots \geq c'_{2q} \geq c'_{2q+1}) & \text{if } G' = O(2n' + \delta, \mathbb{C}) \end{cases}$$

where \mathcal{O}' is obtained from \mathcal{O} by removing all c_{2i}, c_{2i+1} for $i \in \tau(\mathcal{O})$ in \mathcal{O} . Then \mathcal{O}' is a nilpotent orbit in \mathfrak{g}' of the same type as \mathfrak{g} but of smaller rank.

For example, let $\mathcal{O} = (9, 9, 9, 9, 8, 6, 6, 6, 5, 5, 4, 2, 2, 2, 2, 0)$ be a symplectic nilpotent orbit, then $\tau(\mathcal{O}) = \{1, 3, 6, 7\}$ and $\mathcal{O}' = (9, 9, 8, 6, 5, 5, 4, 0)$.

Using the above notations,

$$(5) \quad \mathcal{B}(\mathcal{O}) := \text{Ind}_{\prod_{i \in \tau(\mathcal{O})} GL(c_{2i}) \times G'}^G (|\det| \boxtimes \dots \boxtimes |\det| \boxtimes \mathcal{U}(\mathcal{O}')),$$

where $\mathcal{U}(\mathcal{O}')$ is the *spherical unipotent representation* attached to a classical nilpotent orbit \mathcal{O}' given by

$$\mathcal{U}(\mathcal{O}') := \begin{cases} \overline{X}_{G'}(\lambda[c'_0, c'_1]^+, \dots, \lambda[c'_{2q}, c'_{2q+1}]^+) & \text{if } G' = Sp(2n', \mathbb{C}) \\ \overline{X}_{G'}(\lambda[-\epsilon, c'_1]^+, \lambda[c'_2, c'_3]^+, \dots, \lambda[c'_{2q}, c'_{2q+1}]^+ | + 1) & \text{if } G' = O(2n' + \delta, \mathbb{C}) \end{cases}$$

We now state the main results of [BW1]:

Theorem 1.1. *Let \mathcal{O} be as given in (4). The Brylinski model $\mathcal{B}(\overline{\mathcal{O}})$ in [Br] is the cyclic submodule of $\mathcal{B}(\mathcal{O})$ defined in (5).*

To compute the diminutive K_c -type multiplicities of $\mathcal{B}(\overline{\mathcal{O}})$, one has

Theorem 1.2. *Let \mathcal{O} be as given in (4), and*

$$(6) \quad \Gamma(\mathcal{O}) := \begin{cases} I^G (\lambda[c_0, c_1]^+ \boxtimes \cdots \boxtimes \lambda[c_{2p}, c_{2p+1}]^+) & \text{if } G = Sp(2n, \mathbb{C}), \\ I^G (\lambda[c_2, c_3]^+ \boxtimes \cdots \boxtimes \lambda[c_{2p}, c_{2p+1}]^+ \boxtimes T(c_1|+1)) & \text{if } G = O(2n + \delta, \mathbb{C}) \end{cases}$$

where $T(x|+1)$ and $T(x|-1)$ are the trivial and sign representation of $O(x, \mathbb{C})$ respectively. Then

$$\Gamma(\mathcal{O}) \approx \mathcal{B}(\overline{\mathcal{O}}).$$

As proved in the Appendix B of [BW1], we also record the following, which will be studied in full detail in Section 3 below:

Theorem 1.3. *The lowest K_c -types of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ and $\mathcal{B}(\mathcal{O})$ are diminutive.*

2. REDUCTION TO GENERIC ORBITS

In this section, we define the notion of generic orbits, and prove that our study of $\mathcal{B}(\overline{\mathcal{O}})$ can be reduced to that of generic orbits.

Definition 2.1. *Let \mathcal{O} be a classical nilpotent orbit given in (4). We say \mathcal{O} is **non-generic** if there exists $c_i = c_{i+1} = c_{i+2}$ for some i . Otherwise, it is called **generic**. In other words, \mathcal{O} is generic if and only if the number of repeated columns of \mathcal{O} is at most two.*

Let $N_{\mathcal{O}}(\ell) := \#\{i \mid c_i = \ell\}$ be the number of columns of \mathcal{O} with size equal to ℓ . The *generic part* \mathcal{O}_{gen} of \mathcal{O} is defined by the following: For each $\ell \in \mathbb{N}$,

$$N_{\mathcal{O}_{gen}}(\ell) = \begin{cases} 1 & \text{if } N_{\mathcal{O}}(\ell) > 2 \text{ is odd;} \\ 2 & \text{if } N_{\mathcal{O}}(\ell) > 2 \text{ is even;} \\ N_{\mathcal{O}}(\ell) & \text{otherwise} \end{cases}$$

That is, \mathcal{O}_{gen} is obtained from \mathcal{O} by removing as few column pairs $c_{2i-1} = c_{2i}$ from \mathcal{O} as possible such that \mathcal{O}_{gen} is generic.

For example, for the symplectic orbit $\mathcal{O} = (9, \underline{9}, 9, 9, 8, \underline{6}, 6, 6, 5, 5, 4, \underline{2}, 2, 2, 2, 0)$ above,

$$\mathcal{O} = \underline{(9, 9)} \cup \underline{(6, 6)} \cup \underline{(2, 2)} \cup (9, 9, 8, 6, 5, 5, 4, 2, 2, 0),$$

i.e. $\mathcal{O}_{gen} = (9, 9, 8, 6, 5, 5, 4, 2, 2, 0)$. Note that $\mathcal{O}'_{gen} = \mathcal{O}'$, both equal to $(9, 9, 8, 6, 5, 5, 4, 0)$.

Theorem 2.2. *Let \mathcal{O} be classical nilpotent given in (4) with generic part \mathcal{O}_{gen} . Suppose $\mathcal{O} = (g_1, g_1) \cup \cdots \cup (g_k, g_k) \cup \mathcal{O}_{gen}$. Then*

$$\mathcal{B}(\overline{\mathcal{O}}) \cong I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\overline{\mathcal{O}_{gen}})).$$

Proof. By Theorem 1.1, $\mathcal{B}(\overline{\mathcal{O}_{gen}}) \subseteq \mathcal{B}(\mathcal{O}_{gen})$ as a cyclic submodule. Using induction in stages, we have

$$\begin{aligned} & I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\overline{\mathcal{O}_{gen}})) \\ & \subseteq I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\mathcal{O}_{gen})) \cong \mathcal{B}(\mathcal{O}) \end{aligned}$$

The last \cong holds since $\mathcal{O}'_{gen} = \mathcal{O}'$, and all $\lambda[g_i, g_i]^+$ comes from $\lambda[c_{2i}, c_{2i-1}]^+$ for some $i \in \tau(\mathcal{O})$. Therefore, we have the inclusion

$$(7) \quad \mathcal{B}(\overline{\mathcal{O}}) \subseteq I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\overline{\mathcal{O}_{gen}})) \quad (\subseteq \mathcal{B}(\mathcal{O})).$$

since $\mathcal{B}(\overline{\mathcal{O}})$ is a cyclic submodule of $\mathcal{B}(\mathcal{O})$.

To see the above inclusion is an equality, note that $\Gamma(\mathcal{O}_{gen}) \approx \mathcal{B}(\overline{\mathcal{O}_{gen}})$ by Theorem 1.2 and hence

$$\Gamma(\mathcal{O}) = I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \Gamma(\mathcal{O}_{gen})) \approx I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\overline{\mathcal{O}_{gen}}))$$

On the other hand, we have $\Gamma(\mathcal{O}) \approx \mathcal{B}(\overline{\mathcal{O}})$ by Theorem 1.2 again. Consequently,

$$\mathcal{B}(\overline{\mathcal{O}}) \approx I^G (\lambda[g_1, g_1]^+ \boxtimes \cdots \boxtimes \lambda[g_k, g_k]^+ \boxtimes \mathcal{B}(\overline{\mathcal{O}_{gen}}))$$

have the same diminutive K_c -type multiplicities.

As a result, the inclusion (7) is an isomorphism on the level of diminutive K_c -types. Now the result follows from Theorem 1.3, since the lowest K_c -types of all composition factors of both modules of (7) are diminutive. \square

From now on, we assume our orbit $\mathcal{O} = \mathcal{O}_{gen}$ is generic, which has at most two repeated columns.

3. COMPOSITION FACTORS OF $\mathcal{B}(\overline{\mathcal{O}})$

In Appendix B of [BW1], a list of candidates of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ (and $\mathcal{B}(\mathcal{O})$) is given. We recall them as follows:

Definition 3.1. *Let $(b_0 \geq b_1 = b_2 \geq b_3)$ be four positive integers of the same parity. A **fundamental degeneration** is defined by:*

- $(b_0 > b_1 = b_2 > b_3) \rightarrow (b_0, b_1 + 2, b_2 - 2, b_3)$.
- $(b_0 = b_1 = b_2 > b_3) \rightarrow (b_0 + 1, b_1 + 1, b_2 - 2, b_3)$.
- $(b_0 > b_1 = b_2 = b_3) \rightarrow (b_0, b_1 + 2, b_2 - 1, b_3 - 1)$.
- $(b_0 = b_1 = b_2 = b_3) \rightarrow (b_0 + 1, b_1 + 1, b_2 - 1, b_3 - 1)$.

In the first two cases, we omit the columns $b_2 - 2, b_3$ if both terms are equal to zero. Note that when $b_0 > b_1$, the size of b_0 remains unchanged after degeneration. Similarly, if $b_2 > b_3$, the size of b_3 is the same after degeneration.

Definition 3.2. *Let \mathcal{O} be a classical orbit given in (4) (take $c_0 = \infty$ for orthogonal orbits). We construct a collection of orbits as follows:*

(1) *For each $\dots c_{2i} \geq c_{2i+1} = c_{2i+2} = \dots = c_{2j-1} = c_{2j} \geq c_{2j+1} \dots$ appearing in \mathcal{O} , perform fundamental degeneration on the columns $c_{2i} \geq c_{2i+1} = c_{2j} \geq c_{2j+1}$ and get a new orbit:*

$$(c_0 \geq \dots \geq c'_{2i} \geq c'_{2i+1} \geq c_{2i+2} = \dots = c_{2j-1} \geq c'_{2j} \geq c'_{2j+1} \geq \dots \geq c_{2p+1}).$$

(2) *For each new orbit obtained in Step (1), repeat Step (1) on them until there are no more $c_{2j+1} = c_{2j+2}$'s.*

Denote the collection of all such orbits by $\text{Norm}(\mathcal{O})$.

Note that from the definition, $\mathcal{P} \in \text{Norm}(\mathcal{O})$ may not be generic. This can also be seen in the example below:

Example 3.3. Let $\mathcal{O} = (8, 6, 6, 4, 4, 2, 2, 0)$ be a symplectic orbit. Then the $\mathcal{P} \in \text{Norm}(\mathcal{O})$ are given by

$$(8) \quad \begin{array}{ccc} & (86644220) & \\ & (88444220) & (86662220) & (866444) \\ & (88552220) & (884444) & (866633) \\ & & (885533) & \end{array} .$$

We now list the candidates of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$. For simplicity, we only study the case when \mathcal{O} is **even, generic**. In such a case, each $\mathcal{P} \in \text{Norm}(\mathcal{O})$ is a special orbit in the sense of Lusztig.

The following Lemma is essential in the determination of the candidates of composition factors:

Definition 3.4. Let \mathcal{O} be an **even, generic** orbit (so that $\delta = 0$ for orthogonal groups). For each

$$\mathcal{P} := \begin{cases} (d_0 \geq d_1 \geq \cdots \geq d_{2s} \geq d_{2s+1}) & \text{if } G = Sp(2n, \mathbb{C}), \\ (d_1 \geq d_2 \geq \cdots \geq d_{2s} \geq d_{2s+1}) & \text{if } G = O(2n, \mathbb{C}) \end{cases}$$

in $\text{Norm}(\mathcal{O})$, let $\tau_0(\mathcal{P}) := \{i \mid d_{2i-1} = d_{2i} \text{ is even}\} = \{i_1, \dots, i_r\}$ and

$$\mathcal{P}^* := \begin{cases} (d_0^* \geq d_1^* \geq \cdots \geq d_{2t}^* \geq d_{2t+1}^*) & \text{if } G^* = Sp(2n^*, \mathbb{C}), \\ (d_1^* \geq d_2^* \geq \cdots \geq d_{2t}^* \geq d_{2t+1}^*) & \text{if } G^* = O(2n^*, \mathbb{C}) \end{cases}$$

is obtained from \mathcal{P} by removing the columns $d_{2i-1} = d_{2i}$ for $i \in \tau_0(\mathcal{P})$. The **parameter group** of \mathcal{P} is given by

$$\mathcal{A}(\mathcal{P}) \cong (\mathbb{Z}/2\mathbb{Z})^{a_0} \times \cdots \times (\mathbb{Z}/2\mathbb{Z})^{a_t}, \quad a_i := \begin{cases} 1 & \text{if } d_{2i+1}^* \text{ is even and positive,} \\ 0 & \text{if } d_{2i+1}^* \text{ is odd or } = 0 \end{cases} .$$

Consequently, all irreducible representations $\epsilon \in \mathcal{A}(\mathcal{P})^\vee$ of the parameter group is

$$\epsilon = (\epsilon_1, \dots, \epsilon_{t+1}), \quad \epsilon_i \in \begin{cases} \{+, -\} & \text{if } a_i = 1, \\ \{+\} & \text{if } a_i = 0. \end{cases}$$

When \mathcal{P} is a symplectic orbit, $\mathcal{A}(\mathcal{P}) \cong \overline{\mathcal{A}}(\mathcal{P})$ is isomorphic to the Lusztig's quotient group of \mathcal{P} (as well as \mathcal{P}^*). On the other hand, if \mathcal{P} is an orthogonal orbit, then $\mathcal{A}(\mathcal{P}) \cong \overline{\mathcal{A}}(\mathcal{P}) \times \mathbb{Z}/2\mathbb{Z}$. The extra $\mathbb{Z}/2\mathbb{Z}$ is due to the fact that $O(2n, \mathbb{C})$ has two connected components.

Proposition 3.5. Retain the notations of Definition 3.4. For each $\mathcal{P} \in \text{Norm}(\mathcal{O})$ and $\epsilon \in \mathcal{A}(\mathcal{P})^\vee$, consider the induced modules

$$(9) \quad \mathcal{M}(\mathcal{P}, \epsilon) := I^G (\lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \cdots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \mathcal{U}(\mathcal{P}^*; \epsilon))$$

where $\mathcal{U}(\mathcal{P}^*; \epsilon)$ are the special unipotent representations attached to the special orbit \mathcal{P}^* given by

$$\mathcal{U}(\mathcal{P}^*, \epsilon) := \begin{cases} \overline{X}(\lambda[d_0^*, d_1^*]^{\epsilon_0}, \dots, \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon_t}) & \text{if } G^* = Sp(2n^*, \mathbb{C}), \\ \overline{X}(\lambda[d_2^*, d_3^*]^{\epsilon_1}, \dots, \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon_t}, \lambda[0, d_1^*]^+ | \epsilon_0) & \text{if } G^* = O(2n^*, \mathbb{C}) \end{cases}$$

where $\lambda[x, y]^\pm$ are as defined in (2) if x and y are even, and

$$\lambda[x, y]^+ := \left(-\frac{x-1}{2} \dots \frac{x-1}{2} ; -\frac{x-1}{2} \dots \frac{x-1}{2} \right)$$

if $x = y$ is odd.

Then each $\mathcal{M}(\mathcal{P}, \epsilon)$ is an irreducible module with diminutive lowest K_c -type, and

$$\{\mathcal{M}(\mathcal{P}, \epsilon) \mid \epsilon \in \mathcal{A}(\mathcal{P})^\vee\}$$

are the candidates of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$.

Example 3.6. Let $\mathcal{O} = (8, 6, 6, 4, 4, 2, 2, 0)$ as in the previous example. For each $\mathcal{P} \in \text{Norm}(\mathcal{O})$, the columns $\mathbf{d}_{2i-1} = \mathbf{d}_{2i}$ for $i \in \tau_0(\mathcal{P})$ are in **bold**, so that the unbolded columns gives \mathcal{P}^* . Then the 17 induced modules $\mathcal{M}(\mathcal{P}, \epsilon)$ in Equation (9) are:

$$\begin{aligned} & \mathcal{M}(\mathbf{86644220}; +) = \\ & I^G(\lambda[\mathbf{66}]^- \boxtimes \lambda[\mathbf{44}]^- \boxtimes \lambda[\mathbf{22}]^- \boxtimes \mathcal{U}(80; +)) \end{aligned}$$

$$\mathcal{M}(88444220; \pm+) = I^G(\lambda[\mathbf{44}]^- \boxtimes \lambda[\mathbf{22}]^- \boxtimes \mathcal{U}(8840; \pm+))$$

$$\mathcal{M}(\mathbf{86662220}; \pm+) = I^G(\lambda[\mathbf{66}]^- \boxtimes \lambda[\mathbf{22}]^- \boxtimes \mathcal{U}(8620; \pm+))$$

$$\mathcal{M}(\mathbf{866444}; \pm) = I^G(\lambda[\mathbf{66}]^- \boxtimes \lambda[\mathbf{44}]^- \boxtimes \mathcal{U}(84; \pm))$$

$$\mathcal{M}(88552220; \pm++) = I^G(\lambda[\mathbf{22}]^- \boxtimes \mathcal{U}(885520; \pm++))$$

$$\mathcal{M}(884444; \pm\pm) = I^G(\lambda[\mathbf{44}]^- \boxtimes \mathcal{U}(8844; \pm\pm))$$

$$\mathcal{M}(\mathbf{866633}; \pm+) = I^G(\lambda[\mathbf{66}]^- \boxtimes \mathcal{U}(8633; \pm+))$$

$$\begin{aligned} & \mathcal{M}(885533; \pm++) = \\ & \mathcal{U}(885533; \pm++) \end{aligned}$$

Remark 3.7. The $\mathcal{M}(\mathcal{P}, \epsilon)$ in the proposition above satisfies the following:

For $G = Sp(2n, \mathbb{C})$,

(10)

$$\mathcal{M}(\mathcal{P}, \epsilon) \approx I^G(\lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \dots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \lambda[d_0^*, d_1^*]^{\epsilon_0} \boxtimes \dots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon_t}).$$

For $G = O(2n, \mathbb{C})$,

$$(11) \quad \mathcal{M}(\mathcal{P}, \epsilon) \approx I^G \left(\begin{array}{c} \lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \dots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \\ \lambda[d_2^*, d_3^*]^{\epsilon_1} \boxtimes \dots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon_t} \boxtimes \mathbf{T}(d_1^* | \epsilon_0) \end{array} \right).$$

This can be seen by the calculations of the diminutive K_c -types of any unipotent representations of classical groups in [Wo3], or Equation (16) of [BW1]. For example,

$$\begin{aligned}
\mathcal{M}(88552220; - + +) &= I^G(\lambda[2, 2]^- \boxtimes \mathcal{U}(885520; - + +)) \\
&\approx I^G(\lambda[2, 2]^- \boxtimes \lambda[8, 8]^- \boxtimes \lambda[5, 5]^+ \boxtimes \lambda[2, 0]^+). \\
&= \text{Ind}_{GL(2) \times GL(8) \times GL(5) \times GL(1)}^{Sp(32)}(\det \boxtimes \det \boxtimes \text{triv} \boxtimes |\det|).
\end{aligned}$$

where the last = is given by (3) (in fact the \approx here happens also to be =, but we do not need this fact).

4. DISTINGUISHED MODULES

In the previous section, we have defined the set $Norm(\mathcal{O})$ and listed the parameters attached to each $\mathcal{P} \in Norm(\mathcal{O})$ for all even, generic \mathcal{O} . They are the candidates of the parameters of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$. We now define a distinguished parameter for each $\mathcal{P} \in Norm(\mathcal{O})$. It will be proved in Theorem 4.3 that they are precisely the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$.

Definition 4.1. *Let \mathcal{O} be an even, generic nilpotent orbit. For each $\mathcal{P} \in Norm(\mathcal{O})$ with $\tau_0(\mathcal{P})$ and \mathcal{P}^* defined as in Definition 3.4, we say*

$$\epsilon(\mathcal{P}) := (\epsilon(\mathcal{P})_0, \dots, \epsilon(\mathcal{P})_t) \in \mathcal{A}(\mathcal{P})^\vee$$

is **distinguished** in \mathcal{P} if the following holds for all $i = 0, \dots, t$:

$$\text{If } d_{2i+1}^* \text{ even and positive, then } \epsilon(\mathcal{P})_i = (-1)^{\#\{j \in \tau_0(\mathcal{P}) \mid d_{2i}^* \geq d_{2j} \geq d_{2i+1}^*\}}$$

(take $d_0^* = \infty$ for orthogonal orbits). The irreducible module $\mathcal{M}(\mathcal{P}, \epsilon(\mathcal{P}))$ is **distinguished** in $\mathcal{P} \in Norm(\mathcal{O})$.

One can check directly from the definition that each $\mathcal{P} \in Norm(\mathcal{O})$ contains exactly one distinguished module.

Example 4.2. *Let $\mathcal{O} = (8 \geq 6 \geq 6 \geq 4 \geq 4 \geq 2 \geq 2 \geq 0)$ as above. Then the distinguished modules are:*

$$\begin{aligned}
&\mathcal{M}(86644220; +) \\
&\mathcal{M}(88444220; ++), \quad \mathcal{M}(86662220; -+), \quad \mathcal{M}(866444; +) \\
&\mathcal{M}(88552220; +++) \quad \mathcal{M}(884444; +-), \quad \mathcal{M}(866633; --) \\
&\mathcal{M}(885533; +++)
\end{aligned}$$

For example, take $\mathcal{P} = (d_0, d_1, d_2, \mathbf{d}_3, \mathbf{d}_4, d_5) = (8, 8, 4, \mathbf{4}, \mathbf{4}, 4) \in Norm(\mathcal{O})$ with

- $\tau_0(\mathcal{P}) = \{2\}$ with $d_3 = d_4 = 4$.
- $\mathcal{P}^* = (d_0^*, d_1^*, d_2^*, d_3^*) = (8, 8, 4, 4)$ with $\mathcal{A}(\mathcal{P})^\vee = \{++, +-, -+, --\}$.

Now there are no d_{2j} , $j \in \tau_0(\mathcal{P})$ lying between d_0^* and $d_1^* = 8$. Hence the sign is +. On the other hand, $d_4 = 4$ lies in between d_2^* and $d_3^* = 4$. Hence the sign is -. So the module $\mathcal{M}(884444; +-)$ is distinguished.

Theorem 4.3. *Let \mathcal{O} be an even, generic nilpotent orbit. Then the sum of all distinguished modules satisfies:*

$$\mathcal{B}(\overline{\mathcal{O}}) \approx \Gamma(\mathcal{O}) \approx \bigoplus_{\mathcal{P} \in \text{Norm}(\mathcal{O})} \mathcal{M}(\mathcal{P}, \epsilon(\mathcal{P})).$$

Since the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are all of the form $\mathcal{M}(\mathcal{P}, \epsilon)$ in Proposition 3.5. The above theorem implies:

Corollary 4.4. *Retain the above notations. Then the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are precisely*

$$\{\mathcal{M}(\mathcal{P}, \epsilon(\mathcal{P})) \mid \mathcal{P} \in \text{Norm}(\mathcal{O})\},$$

all appearing with multiplicity one.

To prove Theorem 4.3, we need the following three lemmas, all of which are (slight generalizations of) some special cases of Theorem 4.3:

Lemma 4.5 (“3–column Lemma”). *Let G be an orthogonal group, and $(a > b = c \geq d)$ be even, non-negative integers with $a = \infty$. Suppose $(a' > b' > c' \geq d')$ is obtained from $(a > b = c \geq d)$ by a fundamental degeneration (Definition 3.1). Then*

$$I^G(\lambda[c, d]^+ \boxtimes \mathbb{T}(b \pm 1)) \approx I^G(\lambda[c, b]^- \boxtimes \mathbb{T}(d \mp 1)) \oplus I^G(\lambda[c', d']^+ \boxtimes \mathbb{T}(b' \pm 1)).$$

Proof. Consider the orthogonal orbit $\mathcal{O} = (b, c, d)$. Then by Theorem 1.2,

$$\Gamma(\mathcal{O}) = I^G(\lambda[c, d]^+ \boxtimes \mathbb{T}(b + 1)) \approx \mathcal{B}(\overline{\mathcal{O}})$$

Let $\mathcal{P} := (b', c', d')$ so that $\text{Norm}(\mathcal{O}) = \{\mathcal{O}, \mathcal{P}\}$, then the candidates of composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are

$$(12) \quad \begin{array}{ll} \mathcal{M}(\mathcal{O}, \theta) & (\approx I^G(\lambda[c, b]^- \boxtimes \mathbb{T}(d|\theta))) \quad \text{where } \theta \in \mathcal{A}(\mathcal{O})^\vee; \\ \mathcal{M}(\mathcal{P}, \chi) & (\approx I^G(\lambda[c', d']^{\chi_1} \boxtimes \mathbb{T}(b'|\chi_2))) \quad \text{where } \chi \in \mathcal{A}(\mathcal{P})^\vee \end{array}$$

(the \approx above comes from Remark 3.7). Then the result follows by comparing the diminutive K_c -type multiplicities of $\Gamma(\mathcal{O}) \approx \mathcal{B}(\overline{\mathcal{O}})$ and the modules in (12). The proof is similar for $I^G(\lambda[c, d]^+ \boxtimes \mathbb{T}(b - 1))$. \square

Example 4.6. *Let $\mathcal{O} = (6, 6, 2)$. Then $\mathcal{P} = (8, 4, 2)$ and we would like to study the composition factors of*

$$I^G(\lambda[6, 2]^+ \boxtimes \mathbb{T}(6 \pm 1)).$$

The modules in the brackets of (12) are

$$(13) \quad I^G(\lambda[6, 6]^- \boxtimes \mathbb{T}(2 \pm 1)) \quad \text{and} \quad I^G(\lambda[4, 2]^\pm \boxtimes \mathbb{T}(8 \pm 1)),$$

By Frobenius reciprocity, one can calculate the diminutive K_c -type multiplicities of these modules:

	$\wedge^0 \mathbb{C}^{14}$	$\wedge^2 \mathbb{C}^{14}$	$\wedge^4 \mathbb{C}^{14}$	$\wedge^6 \mathbb{C}^{14}$	$\wedge^8 \mathbb{C}^{14}$	$\wedge^{10} \mathbb{C}^{14}$	$\wedge^{12} \mathbb{C}^{14}$	$\wedge^{14} \mathbb{C}^{14}$
$I^G(\lambda[6, 2]^+ \boxtimes T(6 +1))$	1	1	1	1	1	0	0	0
$I^G(\lambda[6, 2]^+ \boxtimes T(6 -1))$	0	0	0	1	1	1	1	1
$I^G(\lambda[6, 6]^- \boxtimes T(2 +1))$	0	0	0	1	0	0	0	0
$I^G(\lambda[6, 6]^- \boxtimes T(2 -1))$	0	0	0	0	1	0	0	0
$I^G(\lambda[4, 2]^+ \boxtimes T(8 +1))$	1	1	1	1	0	0	0	0
$I^G(\lambda[4, 2]^+ \boxtimes T(8 -1))$	0	0	0	0	1	1	1	1
$I^G(\lambda[4, 2]^- \boxtimes T(8 +1))$	0	1	1	0	0	0	0	0
$I^G(\lambda[4, 2]^- \boxtimes T(8 -1))$	0	0	0	0	0	1	1	0

It is easy to see that

$$I^G(\lambda[6, 2]^+ \boxtimes T(6|+1)) \approx I^G(\lambda[6, 6]^- \boxtimes T(2|-1)) \oplus I^G(\lambda[4, 2]^+ \boxtimes T(8|+1))$$

and

$$I^G(\lambda[6, 2]^+ \boxtimes T(6|-1)) \approx I^G(\lambda[6, 6]^- \boxtimes T(2|+1)) \oplus I^G(\lambda[4, 2]^+ \boxtimes T(8|-1)).$$

Lemma 4.7 (“4-column Lemma”). *Let G be a symplectic group, and $a \geq b = c \geq d$ be even, non-negative integers. Suppose $a' \geq b' \geq c' \geq d'$ is obtained from $a \geq b = c \geq d$ by a fundamental degeneration (Definition 3.1). Then*

$$I^G(\lambda[a, b]^+ \boxtimes \lambda[c, d]^+) \approx I^G(\lambda[c, b]^- \boxtimes \lambda[a, d]^{\sigma_1}) \oplus I^G(\lambda[a', b']^+ \boxtimes [c', d']^+),$$

where $\sigma_1 := -$ if $d > 0$, or $+$ if $d = 0$.

Moreover, if $a > b$, then the composition factors of

$$I^G(\lambda[a, b]^- \boxtimes \lambda[c, d]^+) \approx I^G(\lambda[c, b]^- \boxtimes \lambda[a, d]^{\sigma_2}) \oplus I^G(\lambda[a', b']^- \boxtimes [c', d']^+),$$

where $\sigma_2 := +$ if $d > 0$, or $+$ if $d = 0$.

Proof. Consider $\mathcal{O} = (a, b, c, d)$. Then

$$\Gamma(\mathcal{O}) = I^G(\lambda[a, b]^+ \boxtimes \lambda[c, d]^+) \approx \mathcal{B}(\overline{\mathcal{O}}).$$

As in the previous Lemma, $\text{Norm}(\mathcal{O}) = \{\mathcal{O}, \mathcal{P} := (a', b', c', d')\}$ and the candidates of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are

$$(14) \quad \begin{aligned} \mathcal{M}(\mathcal{O}, \theta) &\approx I^G(\lambda[c, b]^- \boxtimes \lambda[a, d]^\theta), & \text{where } \theta \in \overline{\mathcal{A}}(\mathcal{O})^\vee; \\ \mathcal{M}(\mathcal{P}, \chi) &\approx I^G(\lambda[a', b']^{\chi_1} \boxtimes \lambda[c', d']^{\chi_2}) & \text{where } \chi \in \overline{\mathcal{A}}(\mathcal{P})^\vee. \end{aligned}$$

Then the result follows by comparing the diminutive K_c -type multiplicities of $\Gamma(\mathcal{O}) \approx \mathcal{B}(\overline{\mathcal{O}})$ and the modules in (14). \square

Example 4.8. *Let $\mathcal{O} = (6, 2, 2, 2)$ be non-generic orbit in $\mathfrak{sp}(12, \mathbb{C})$. Then $\mathcal{P} = (6, 4, 1, 1)$ and we study the composition factors of*

$$I^G(\lambda[6, 2]^\pm \boxtimes \lambda[2, 2]^+)$$

The modules in (14) are

$$(15) \quad I^G(\lambda[2, 2]^- \boxtimes \lambda[6, 2]^\pm) \quad \text{and} \quad I^G(\lambda[6, 4]^\pm \boxtimes \lambda[1, 1]^+),$$

As before, one can calculate the diminutive K_c -type multiplicities of these modules:

	$\wedge^0 \mathbb{C}^{12}$	$\wedge^2 \mathbb{C}^{12} / \wedge^0 \mathbb{C}^{12}$	$\wedge^4 \mathbb{C}^{12} / \wedge^2 \mathbb{C}^{12}$	$\wedge^6 \mathbb{C}^{12} / \wedge^4 \mathbb{C}^{12}$
$I^G(\lambda[6, 2]^+ \boxtimes \lambda[2, 2]^+)$	1	1	1	0
$I^G(\lambda[6, 2]^- \boxtimes \lambda[2, 2]^+)$	0	1	1	1
$I^G(\lambda[2, 2]^- \boxtimes \lambda[6, 2]^+)$	0	1	0	0
$I^G(\lambda[2, 2]^- \boxtimes \lambda[6, 2]^-)$	0	0	1	0
$I^G(\lambda[6, 4]^+ \boxtimes [1, 1]^+)$	1	1	0	0
$I^G(\lambda[6, 4]^- \boxtimes [1, 1]^+)$	0	0	1	1

Once again, one can see easily that

$$I^G(\lambda[6, 2]^+ \boxtimes \lambda[2, 2]^+) \approx I^G(\lambda[2, 2]^- \boxtimes \lambda[6, 2]^-) \oplus I^G(\lambda[6, 4]^+ \boxtimes [1, 1]^+)$$

and

$$I^G(\lambda[6, 2]^- \boxtimes \lambda[2, 2]^+) \approx I^G(\lambda[2, 2]^- \boxtimes \lambda[6, 2]^+) \oplus I^G(\lambda[6, 4]^- \boxtimes [1, 1]^+).$$

For the purpose of the induction arguments below, we record the following Lemma, whose proof is identical to that of Lemma 4.7:

Lemma 4.9 (“5-column Lemma”). *Let G be an orthogonal group, and $z > a \geq b = c \geq d$ be even, non-negative integers. Suppose $a' \geq b' \geq c' \geq d'$ is obtained from $a \geq b = c \geq d$ by a fundamental degeneration (Definition 3.1). Then*

$$\begin{aligned} & I^G(\lambda[a, b]^+ \boxtimes \lambda[c, d]^+ \boxtimes T(z|+1)) \\ & \approx I^G(\lambda[c, b]^- \boxtimes \lambda[a, d]^{\sigma_1} \boxtimes T(z|+1)) \oplus I^G(\lambda[a', b']^+ \boxtimes [c', d']^+ \boxtimes T(z|+1)), \end{aligned}$$

where $\sigma_1 := -$ if $d > 0$, or $+$ if $d = 0$.

Moreover, if $a > b$, then the composition factors of

$$\begin{aligned} & I^G(\lambda[a, b]^- \boxtimes \lambda[c, d]^+ \boxtimes T(z|+1)) \\ & \approx I^G(\lambda[c, b]^- \boxtimes \lambda[a, d]^{\sigma_2} \boxtimes T(z|+1)) \oplus I^G(\lambda[a', b']^- \boxtimes [c', d']^+ \boxtimes T(z|+1)), \end{aligned}$$

where $\sigma_2 := +$ if $d > 0$, or $-$ if $d = 0$.

Proof of Theorem 4.3. We now prove Theorem 4.3 by induction on the length of

$$\mathcal{O} = \begin{cases} (c_0, c_1, \dots, c_{2p}, c_{2p+1}) & \text{if } G = Sp(2n, \mathbb{C}), \\ (c_1, c_2, \dots, c_{2p}, c_{2p+1}) & \text{if } G = O(2n, \mathbb{C}) \end{cases}.$$

4.1. Initial Step. Let $p = 1$ and $\mathcal{O}_1 = (c_1, c_2, c_3)$ be an orthogonal orbit. Then

$$Norm(\mathcal{O}_1) = \begin{cases} \{\mathcal{O}_1\} & \text{if } c_1 \neq c_2; \\ \{\mathcal{O}_1, \mathcal{P}_1\} & \text{if } c_1 = c_2 \end{cases}$$

where $\mathcal{P}_1 := (c'_1, c'_2, c'_3)$ is obtained from \mathcal{O}_1 by a fundamental degeneration (with $c_0 = \infty$). The distinguished modules are

$$\begin{cases} \{\mathcal{M}(\mathcal{O}_1, \epsilon(\mathcal{O}_1))\} & \text{if } c_1 \neq c_2; \\ \{\mathcal{M}(\mathcal{O}_1, \epsilon(\mathcal{O}_1)), \mathcal{M}(\mathcal{P}_1, \epsilon(\mathcal{P}_1))\} & \text{if } c_1 = c_2. \end{cases}$$

By Remark 3.7, the sum of the distinguished modules are \approx to

$$\begin{cases} I^G(\lambda[c_2, c_3]^+ \boxtimes \mathbb{T}(c_1|+1)) & \text{if } c_1 \neq c_2; \\ I^G(\lambda[c_2, c_1]^- \boxtimes \mathbb{T}(c_3|-1)) \oplus I^G(\lambda[c'_2, c'_3]^+ \boxtimes \mathbb{T}(c'_1|+1)) & \text{if } c_1 = c_2. \end{cases}$$

In both cases, the sum is $I^G(\lambda[c_2, c_3]^+ \boxtimes \mathbb{T}(c_1|+1)) = \Gamma(\mathcal{O}_1)$ by the 3-column lemma (Lemma 4.5).

Let $\mathcal{O}_1 = (c_0, c_1, c_2, c_3)$ be a symplectic orbit. Then

$$\text{Norm}(\mathcal{O}_1) = \begin{cases} \{\mathcal{O}_1\} & \text{if } c_1 \neq c_2; \\ \{\mathcal{O}_1, \mathcal{P}_1\} & \text{if } c_1 = c_2 \end{cases}$$

where $\mathcal{P}_1 = (c'_0, c'_1, c'_2, c'_3)$ is obtained from \mathcal{O}_1 by a fundamental degeneration. As before, the sum of the distinguished modules are \approx to

$$\begin{cases} I^G(\lambda[c_0, c_1]^+ |\lambda[c_2, c_3]^+ & \text{if } c_1 \neq c_2; \\ I^G(\lambda[c_2, c_1]^- |\lambda[c_0, c_3]^-) \oplus I^G(\lambda[c'_0, c'_1]^+ |\lambda[c'_2, c'_3]^+) & \text{if } c_1 = c_2. \end{cases}$$

In both cases, the sum is $I^G(\lambda[c_0, c_1]^+ \boxtimes \lambda[c_2, c_3]^+) = \Gamma(\mathcal{O}_1)$ by the 4-column lemma (Lemma 4.7).

4.2. Induction Step. Now suppose the proposition holds for $p = k-1$. We give a complete proof of the case of $p = k$ for orthogonal orbits, while the arguments for symplectic orbits is similar.

Let $\mathcal{O}_k = (c_1, c_2, \dots, c_{2k}, c_{2k+1})$ be an orthogonal orbit with

$$(16) \quad \Gamma(\mathcal{O}_k) = I^G(\lambda[c_2, c_3]^+ \boxtimes \dots \boxtimes \lambda[c_{2k}, c_{2k+1}]^+ \boxtimes \mathbb{T}(c_1|+)) \approx \bigoplus_{\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)} \mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k)).$$

It is useful to keep in mind that the following holds:

- (i) $\mathcal{P}_k = (d_1, d_2, \dots, d_{2s}, d_{2s+1})$;
- (ii) $\tau_0(\mathcal{P}_k) = \{i_1, \dots, i_r\}$;
- (iii) $\mathcal{P}_k^* = (d_1^*, d_2^*, \dots, d_{2t}^*, d_{2t+1}^*)$;
- (iv) $c_{2k+1} = d_{2s+1} = d_{2t+1}^*$;
- (v) $\mathcal{A}(\mathcal{P}_k) = (\mathbb{Z}/2\mathbb{Z})^{a_0} \times \dots \times (\mathbb{Z}/2\mathbb{Z})^{a_t}$.

By Remark 3.7, $\mathcal{M}(\mathcal{P}_k; \epsilon(\mathcal{P}_k))$ satisfies

$$\mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k)) \approx I^G \left(\begin{array}{c} \lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \dots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \\ \lambda[d_2^*, d_3^*]^{\epsilon(\mathcal{P}_k)_1} \boxtimes \dots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \mathbb{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \end{array} \right)$$

Now consider $\mathcal{O}_{k+1} = (c_1, c_2, \dots, c_{2k}, c_{2k+1}, c_{2k+2}, c_{2k+3})$. By (16),

$$\Gamma(\mathcal{O}_{k+1}) = I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \Gamma(\mathcal{O}_k)) \approx \bigoplus_{\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)} I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k)))$$

So we wish to study the sum on the right. There are two cases:

(i) $c_{2k-1} \neq c_{2k}$: For each $\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)$, let

$$\mathcal{P}_{k+1} := (d_1, d_2, \dots, d_{2s}, d_{2s+1}, c_{2k+2}, c_{2k+3}).$$

Then it is easy to see from the definition of $\text{Norm}(\mathcal{O})$ that

$$\text{Norm}(\mathcal{O}_{k+1}) = \bigcup_{\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)} \{\mathcal{P}_{k+1}\}.$$

We now look at \mathcal{P}_{k+1} more closely: $\tau_0(\mathcal{P}_{k+1}) = \tau_0(\mathcal{P}_k)$ since $d_{2s+1} = c_{2k+1} \neq c_{2k+2}$. Therefore,

$$(17) \quad \mathcal{P}_{k+1}^* = (d_1^*, d_2^*, \dots, d_{2t}^*, d_{2t+1}^*, c_{2k+2}, c_{2k+3}), \quad \epsilon(\mathcal{P}_{k+1}) = (\epsilon(\mathcal{P}_k)_0, \dots, \epsilon(\mathcal{P}_k)_t, +)$$

Now

$$\begin{aligned} & I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k))) \\ & \approx I^G \left(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \dots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \right. \\ & \quad \left. \lambda[d_2^*, d_3^*]^{\epsilon(\mathcal{P}_k)_1} \boxtimes \dots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \text{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \right) \\ & \approx I^G \left(\lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \dots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \right. \\ & \quad \left. \lambda[d_2^*, d_3^*]^{\epsilon(\mathcal{P}_k)_1} \boxtimes \dots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \text{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \right) \\ & \approx \mathcal{M}(\mathcal{P}_{k+1}, \epsilon(\mathcal{P}_{k+1})). \end{aligned}$$

Here the last \approx comes from Remark 3.7 and (17).

(ii) $c_{2k-1} = c_{2k}$: For each $\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)$, let

$$\mathcal{P}_{k+1}^u := (d_1, d_2, \dots, d_{2s}, d_{2s+1}, c_{2k+2}, c_{2k+3}),$$

with $d_{2s+1} = c_{2k+1} = c_{2k+2}$. So we also let

$$\mathcal{P}_{k+1}^d := (d_1, d_2, \dots, d'_{2s}, d'_{2s+1}, c'_{2k+2}, c'_{2k+3}).$$

by performing a fundamental degeneration on $d_{2s} \geq d_{2s+1} = c_{2k+2} \geq c_{2k+3}$. Then

$$\text{Norm}(\mathcal{O}_{k+1}) = \bigcup_{\mathcal{P}_k \in \text{Norm}(\mathcal{O}_k)} \{\mathcal{P}_{k+1}^u, \mathcal{P}_{k+1}^d\}.$$

As before, we look at \mathcal{P}_{k+1}^u and \mathcal{P}_{k+1}^d more closely: Firstly, $\tau_0(\mathcal{P}_{k+1}^u) = \tau_0(\mathcal{P}_k) \cup \{k+1\}$ since $d_{2s+1} = c_{2k+1} = c_{2k+2}$. Therefore,

$$(18) \quad (\mathcal{P}_k^u)^* = (d_1^*, d_2^*, \dots, d_{2t}^*, c_{2k+3}), \quad \epsilon(\mathcal{P}_{k+1}^u) = (\epsilon(\mathcal{P}_k)_0, \dots, -\epsilon(\mathcal{P}_k)_t).$$

On the other hand, $\tau_0(\mathcal{P}_{k+1}^d) = \tau_0(\mathcal{P}_k)$ since $d'_{2s+1} \neq c'_{2k+2}$, and hence

$$(19) \quad (\mathcal{P}_k^d)^* = (d_1^*, d_2^*, \dots, d_{2t}^*, d'_{2s+1}, c'_{2k+2}, c'_{2k+3}), \quad \epsilon(\mathcal{P}_{k+1}^d) = (\epsilon(\mathcal{P}_k)_0, \dots, \epsilon(\mathcal{P}_k)_t, +).$$

Consider

$$\begin{aligned}
(20) \quad & I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k))) \\
& \approx I^G \left(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \cdots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \right. \\
& \quad \left. \lambda[d_2^*, d_3^*]^{\epsilon(\mathcal{P}_k)_1} \boxtimes \cdots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \mathrm{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \right) \\
& \approx I^G \left(\lambda[d_{2i_1}, d_{2i_1-1}]^- \boxtimes \cdots \boxtimes \lambda[d_{2i_r}, d_{2i_r-1}]^- \boxtimes \right. \\
& \quad \left. \lambda[d_2^*, d_3^*]^{\epsilon(\mathcal{P}_k)_1} \boxtimes \cdots \boxtimes \lambda[d_{2t}^*, d_{2t+1}^*]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \mathrm{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \right)
\end{aligned}$$

Look at the last part of the last module in (20). If $t = 0$, then $d_1^* = d_{2t+1}^* = c_{2k+1} = c_{2k+2}$. By applying the 3-column lemma (Lemma 4.5) with $(d_1^* = c_{2k+2} \geq c_{2k+3})$, the last module of (20) is \approx to

$$\begin{aligned}
& I^G(\cdots \boxtimes \lambda[c_{2k+2}, d_1^*]^- \boxtimes \mathrm{T}(c_{2k+3} | -\epsilon(\mathcal{P}_k)_1)) \oplus I^G(\cdots \boxtimes \lambda[c'_{2k+2}, c'_{2k+3}]^+ \boxtimes \mathrm{T}(d'_1 | \epsilon(\mathcal{P}_k)_1)) \\
& \approx \mathcal{M}(\mathcal{P}_{k+1}^u; \epsilon(\mathcal{P}_{k+1}^u)) \oplus \mathcal{M}(\mathcal{P}_{k+1}^d; \epsilon(\mathcal{P}_{k+1}^d))
\end{aligned}$$

where the last \approx follows from (18) and (19).

On the other hand, if $t > 0$, then $d_{2t+1}^* = c_{2k+1} = c_{2k+2}$. By applying the 5-column lemma (Lemma 4.9) with $(d_1^* > d_{2t}^* \geq d_{2t+1}^* = c_{2k+2} \geq c_{2k+3})$, the last module is \approx to

$$\begin{aligned}
& I^G \left(\cdots \boxtimes \lambda[c_{2k+2}, d_{2t+1}^*]^- \boxtimes \lambda[d_{2t}^*, c_{2k+3}]^{-\epsilon(\mathcal{P}_k)_t} \boxtimes \mathrm{T}(d_1^* | \epsilon(\mathcal{P}_k)_0) \right) \oplus \\
& I^G \left(\cdots \boxtimes \lambda[d'_{2t}, d'_{2t+1}]^{\epsilon(\mathcal{P}_k)_t} \boxtimes \lambda[c'_{2k+2}, c'_{2k+3}]^+ \boxtimes \mathrm{T}(d'_1 | \epsilon(\mathcal{P}_k)_0) \right)
\end{aligned}$$

which is also \approx to $\mathcal{M}(\mathcal{P}_{k+1}^u, \epsilon(\mathcal{P}_{k+1}^u)) \oplus \mathcal{M}(\mathcal{P}_{k+1}^d, \epsilon(\mathcal{P}_{k+1}^d))$.

By summing up all $\mathcal{P}_k \in \mathrm{Norm}(\mathcal{O}_k)$ in all the cases above, one always have

$$\begin{aligned}
& \bigoplus_{\mathcal{P}_{k+1} \in \mathrm{Norm}(\mathcal{O}_{k+1})} \mathcal{M}(\mathcal{P}_{k+1}, \epsilon(\mathcal{P}_{k+1})) \\
& \approx \bigoplus_{\mathcal{P}_k \in \mathrm{Norm}(\mathcal{O}_k)} I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \mathcal{M}(\mathcal{P}_k, \epsilon(\mathcal{P}_k))) \\
& \approx I^G(\lambda[c_{2k+2}, c_{2k+3}]^+ \boxtimes \lambda[c_2, c_3]^+ \boxtimes \cdots \boxtimes \lambda[c_{2k}, c_{2k+1}]^+ \boxtimes \mathrm{T}(c_1 | +)) \\
& \approx \Gamma(\mathcal{O}_{k+1})
\end{aligned}$$

for all even, generic orthogonal orbits. The proof for symplectic orbits is similar (and easier) using the 4-column lemma (Lemma 4.7). Therefore, Theorem 4.3 follows. \square

As in [BW1], the same arguments apply to odd orbits, and for general orbits involving both odd and even columns. We skip the proofs there.

Finally, we discuss how one can obtain the character formula of $\mathcal{B}(\overline{\mathcal{O}})$ for all classical nilpotent orbits. By Theorem 2.2, the character formula of $\mathcal{B}(\overline{\mathcal{O}})$ can be obtained from that of $\mathcal{B}(\overline{\mathcal{O}}_{gen})$. Using (9) and the character formulas of unipotent representations given in [BV2], one can get the character formulas for the distinguished modules $\mathcal{M}(\mathcal{P}; \epsilon(\mathcal{P}))$

for all $\mathcal{P} \in \text{Norm}(\mathcal{O}_{gen})$. Consequently, the character formula of $\mathcal{B}(\overline{\mathcal{O}_{gen}})$ can be obtained by summing up the formulas $\mathcal{B}(\overline{\mathcal{O}_{gen}}) \cong \bigoplus_{\mathcal{P} \in \text{Norm}(\mathcal{O}_{gen})} \mathcal{M}(\mathcal{P}; \epsilon(\mathcal{P}))$. As a consequence, one

can calculate the multiplicities of **all** K_c -types in $\mathcal{B}(\overline{\mathcal{O}}) \cong R(\overline{\mathcal{O}})$. Explicit calculations are carried out in [Wol].

We end this paper by an example of the generic orbit $\mathcal{O} = (4, 2, 2, 0)$ in $G = Sp(8, \mathbb{C})$ (with non-normal closure):

Example 4.10. *Let $\mathcal{O} = (4, 2, 2, 0)$ be a generic orbit in $G = Sp(8, \mathbb{C})$. The infinitesimal character for $\mathcal{B}(\overline{\mathcal{O}})$ is $\lambda_{\mathcal{O}} = (2, 1, 1, 0)$. The character formula of $\mathcal{B}(\overline{\mathcal{O}})$ is given by:*

$$\frac{1}{2} \left[\sum_{w_1 \in C_2 \times D_1 \times C_1} \text{sgn}(w_1) X((21, 0, 1); w_1(21, 0, 1)) + \sum_{w_2 \in D_3 \times C_1} \text{sgn}(w_2) X((210, 1); w_2(210, 1)) \right]$$

Upon restricting to K , the formula can be rewritten as:

$$\begin{aligned} \mathcal{B}(\overline{\mathcal{O}}) \cong & \text{Ind}_T^K(\mathbb{C}_{0000}) - 2\text{Ind}_T^K(\mathbb{C}_{1100}) - \text{Ind}_T^K(\mathbb{C}_{2000}) + 4\text{Ind}_T^K(\mathbb{C}_{2110}) \\ & - 2\text{Ind}_T^K(\mathbb{C}_{2211}) - \text{Ind}_T^K(\mathbb{C}_{2200}) + \text{Ind}_T^K(\mathbb{C}_{2222}) + 2\text{Ind}_T^K(\mathbb{C}_{3210}) \\ & - 2\text{Ind}_T^K(\mathbb{C}_{3221}) - \text{Ind}_T^K(\mathbb{C}_{4110}) + \text{Ind}_T^K(\mathbb{C}_{4211}) + \text{Ind}_T^K(\mathbb{C}_{4200}) - \text{Ind}_T^K(\mathbb{C}_{4220}). \end{aligned}$$

On the other hand, the character formula of $\mathcal{B}(\mathcal{O})$ in (5) can be written as:

$$\sum_{w \in C_2 \times A_1} \text{sgn}(w) X((21, 10); w(21, 10))$$

whose restriction to K is given by

$$\begin{aligned} \mathcal{B}(\mathcal{O}) \cong & \text{Ind}_T^K(\mathbb{C}_{0000}) - 2\text{Ind}_T^K(\mathbb{C}_{1100}) + \text{Ind}_T^K(\mathbb{C}_{1111}) - \text{Ind}_T^K(\mathbb{C}_{2000}) \\ & + \text{Ind}_T^K(\mathbb{C}_{2110}) + 2\text{Ind}_T^K(\mathbb{C}_{3100}) - 2\text{Ind}_T^K(\mathbb{C}_{3111}) - \text{Ind}_T^K(\mathbb{C}_{3300}) \\ & + \text{Ind}_T^K(\mathbb{C}_{3311}) - \text{Ind}_T^K(\mathbb{C}_{4000}) + \text{Ind}_T^K(\mathbb{C}_{4110}) + \text{Ind}_T^K(\mathbb{C}_{4200}) - \text{Ind}_T^K(\mathbb{C}_{4211}). \end{aligned}$$

The difference between the formulas occurs at \mathbb{C}_{1111} . This implies the diminutive K_c -type $V_{(1111)}$ has different multiplicities in $R(\overline{\mathcal{O}})$ and in $R(\mathcal{O})$. More precisely, $\text{Norm}(\mathcal{O}) = \{\mathcal{O}, (4, 4)\}$, and the difference between $R(\overline{\mathcal{O}})$ and $R(\mathcal{O})$ is exactly equal to

$$\mathcal{M}(44, -)|_K = \text{Ind}_{GL(4)}^{Sp(8)}(\det)|_K \cong \sum_{w \in A_3} \text{sgn}(w) \text{Ind}_T^K((210 - 1) - w(10 - 1 - 2)).$$

REFERENCES

- [AB] J. Adams, D. Barbasch *The Reductive Dual Pairs Correspondence for Complex Groups*, J. of Func. An. vol 132, 1995
- [B1] D. Barbasch *The unitary dual for complex classical Lie groups*, Invent. Math., vol 96, 1989, pp 103-176
- [B2] ——— *Unipotent representations for real reductive groups*, Proceedings of ICM 1990, Springer Verlag, Tokyo, 1991, pp. 769-777
- [B3] ——— *Orbital integrals of nilpotent orbits*, volume in honor of Harish-Chandra, Proc. Symp. Pure Math. 68 (2000) pp. 97-110.

- [B4] ——— *The unitary spherical spectrum for split classical groups*, Journal of Inst. of Math. Jussieu, 2010 9(2) pp 265-356
- [B5] ——— *Unipotent Representations and the Dual Pairs Correspondence*, Progress in Mathematics, vol. 323, 2017
- [BSS] D. Barbasch, S. Sahi, B. Speh, *Degenerate series representations for $GL(2n, \mathbb{R})$ and Fourier Analysis* Symposia Mathematica, vol XXXI, 45-69, Rome 1988, Academic Press, London, 1990
- [BW1] D. Barbasch, K.D. Wong, *Admissible modules and normality of classical nilpotent orbits I*, arXiv:1801.06909
- [BV] N. Berline, M. Vergne *Fourier transforms of orbits of the coadjoint representation*, Representation theory of reductive groups, Birkhäuser-Boston, Progress in Mathematics vol 40, 1983, pp. 53-69
- [BV1] D. Barbasch, D. Vogan *The local structure of characters*, J. Funct. Analysis, vol. 34 no. 1, 1980, pp. 27-55
- [BV2] D. Barbasch, D. Vogan *Unipotent representations of complex semisimple Lie groups*, Ann. of Math., 1985, vol 121, pp. 41-110
- [BV3] D. Barbasch, D. Vogan *Weyl group representations and nilpotent orbits*, Representation theory of reductive groups, Birkhäuser-Boston, Progress in mathematics vol 40, 1983, pp. 21-35
- [Br] R. Brylinski *Line bundles on the cotangent bundle of the flag variety*, Invent. Math., vol 113 Fasc. 1, 1993, pp. 1-20
- [BZ] I.N. Bernstein, A.V. Zelevensky *Induced representations of reductive p -adic groups. I*, Ann. Sci. École Norm. Sup. (4) 10 (1977), no. 4, 441–472.
- [CM] D. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Norstrand Reinhold Mathematics Series, 1993
- [G] W. Graham, *Functions on the universal cover of the principal nilpotent orbit*, Inv. Math. **108** (1992), 15-27.
- [HC] Harish-Chandra *Fourier transform on a semisimple Lie algebra II*, Amer. J. of Math., vol 79, 1957, pp. 193-257
- [HC1] Harish-Chandra *Harmonic analysis on semisimple Lie groups*, Bulletin AMS. vol 76, 1970, pp. 529-551
- [Kn] A. Knapp *Representation Theory of Semisimple Groups*, Princeton University Press, Princeton NJ, 1986
- [KnV] A. Knapp, D. Vogan *Cohomological induction and unitary representations*, Princeton University Press, Princeton NJ, 1995
- [KP] H-P. Kraft, C. Procesi *On the geometry of conjugacy classes in classical groups*, Comm. Math. Helv., 57, 1982, pp. 539-601
- [J] J. C. Jantzen *Nilpotent orbits in representation theory*, in volume Lie Theory, Lie algebras and Representations by J.-P Anker and B. Orsted, Progress in Mathematics, Birkhäuser, 2004
- [L] G. Lusztig, *Characters of Reductive Groups over a Finite Field*, Ann. Math. Studies, vol. 107, Princeton University Press, 1984
- [LS] G. Lusztig, N. Spaltenstein *Induced unipotent classes*, J. London Math. Soc., vol. 19, 1979, pp. 41-52
- [McG] M. McGovern *Rings of regular functions on nilpotent orbits and their covers*, Invent. Math., 97, 1989, pp. 209-217
- [Pz] T. Przebinda *The dual correspondence of infinitesimal characters*, Colloq. Math. 70 (1996), pp. 93–102
- [R] R. Rao *Orbital integrals in reductive groups*, Annals of Math., vol. 96 no. 3, 1972, pp. 505-510
- [Ro] W. Rossmann *Nilpotent orbital integrals in a real semisimple Lie algebra and representations of Weyl groups*, Operator algebras, unitary representations, enveloping algebras and invariant theory, Progress in Mathematics 92, Birkhäuser Boston, 1990, pp. 263-287
- [ScVi1] W. Schmid, K. Vilonen *Two geometric character formulas for reductive Lie groups*, J. Amer. math. Soc., vol. 11 no. 4, 1998, pp. 799-867
- [ScVi2] W. Schmid, K. Vilonen *Characteristic cycles and wave front cycles of representations of reductive groups*, Annals of Math. 151 (2000), pp. 1071 - 1118.

- [SV] B. Speh, D. Vogan *Reducibility of generalized principal series representations*, Acta Math. 145 (1980), pp. 227-299
- [V1] D. Vogan *The Algebraic Structure of the Representations of Semisimple Lie Groups I*, Annals of Mathematics, Second Series, Vol. 109, No. 1 (Jan., 1979), pp. 1-60
- [V2] D. Vogan *The orbit method and primitive ideals for semisimple Lie algebras*, Canad. Math. Soc. Conf. Proc. Vol. 5 (1986), pp. 281-316
- [V3] D. Vogan *The unitary dual of $GL(n)$ over an archimedean field*, Invent. Math. 83 (1986), pp. 449-505
- [V4] D. Vogan *Associated varieties and unipotent representations*, Harmonic Analysis on reductive groups, Progress in Mathematics vol. 101, Birkhäuser, Boston-Basel-Berlin, 1991, pp. 315-388
- [V5] D. Vogan *The method of coadjoint orbits for real reductive groups*, Representation Theory of Lie Groups, IAS/Park City Mathematics Series 8 (1999), pp. 179-238
- [V6] D. Vogan *Representation of Real Reductive Lie Groups*, Birkhauser, 1981
- [W] N. Wallach *Invariant differential operators on a reductive Lie algebra and Weyl group representations*, J. Amer. Math. Soc., vol. 6 no. 4, 1993, pp. 779-816
- [Wo1] K. D. Wong, *Dixmier Algebras on Complex Classical Nilpotent Orbits and their Representation Theories*, Ph.D. Thesis, Cornell University, 2013
- [Wo2] K.D. Wong *Regular functions on symplectic spherical nilpotent orbits and their quantizations*, Represent. Theory 19 (2015), pp. 333-346
- [Wo3] K.D. Wong *Quantization of special symplectic nilpotent orbits and normality of their closures*, J. Algebra 462 (2016), pp. 37-53

(Barbasch) DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853, U.S.A.
Email address: barbasch@math.cornell.edu

(Wong) SCHOOL OF SCIENCE AND ENGINEERING, THE CHINESE UNIVERSITY OF HONG KONG, SHENZHEN, GUANGDONG 518172, P. R. CHINA
Email address: kayue.wong@gmail.com