

The overfull conjecture on graphs of odd order and large minimum degree

Songling Shan
 Illinois State University, Normal, IL 61790
 sshan12@ilstu.edu

June 28, 2022

Abstract. Let G be a simple graph with maximum degree $\Delta(G)$. A subgraph H of G is overfull if $|E(H)| > \Delta(G) \lfloor \frac{1}{2}|V(H)| \rfloor$. Chetwynd and Hilton in 1986 conjectured that a graph G with $\Delta(G) > \frac{1}{3}|V(G)|$ has chromatic index $\Delta(G)$ if and only if G contains no overfull subgraph. Let $0 < \varepsilon < 1$ and G be a large graph on n vertices with minimum degree at least $\frac{1}{2}(1 + \varepsilon)n$. It was shown that the conjecture holds for G if n is even. In this paper, the same result is proved if n is odd. As far as we know, this is the first result on the conjecture for graphs of odd order and with a minimum degree constraint.

Keywords. Chromatic index; 1-factorization; Overfull Conjecture; Overfull graph.

1 Introduction

In this paper, the terminology graph is used to represent a simple graph. A multigraph may contain multiple edges but contains no loop. Let G be a multigraph. We use $V(G)$ and $E(G)$ to denote the vertex set and the edge set of G , respectively, and let $e(G) = |E(G)|$. For $v \in V(G)$, $N_G(v)$ is the set of neighbors of v in G , and $d_G(v)$, the degree of v in G , is the number of edges of G that are incident with v . We let $d_G^s(v) = |N_G(v)|$, and call it the *simple degree* of v in G . When G is simple, we have $d_G(v) = d_G^s(v)$. Let $V_1, V_2 \subseteq V(G)$ be two disjoint vertex sets. Then $E_G(V_1, V_2)$ is the set of edges in G with one end in V_1 and the other end in V_2 , and $e_G(V_1, V_2) := |E_G(V_1, V_2)|$. We write $E_G(v, V_2)$ and $e_G(v, V_2)$ if $V_1 = \{v\}$ is a singleton. We also use $G[V_1, V_2]$ to denote the bipartite subgraph of G with vertex set $V_1 \cup V_2$ and edge set $E_G(V_1, V_2)$. For $S \subseteq V(G)$ and $v \in V(G)$, $N_G(v, S) := N_G(v) \cap S$ and $d_S(v) := e_G(v, S \setminus \{v\})$, the subgraph of G induced by S is $G[S]$, and $G - S := G[V(G) \setminus S]$. If $F \subseteq E(G)$, then $G - F$ is obtained from G by deleting all the edges of F . Let $\mu(G) = \max\{e_G(u, v) : u, v \in V(G)\}$ be the multiplicity of G , and for $x \in V(G)$, let $\mu(x) = \max\{e_G(x, v) : v \in V(G)\}$ be the multiplicity of x .

For two integers p, q , let $[p, q] = \{i \in \mathbb{Z} : p \leq i \leq q\}$. For a nonnegative integer k , an *edge k -coloring* of a multigraph G is a mapping φ from $E(G)$ to the set of integers $[1, k]$, called *colors*, such that no two adjacent edges receive the same color with respect to φ . The *chromatic index* of G , denoted $\chi'(G)$, is the smallest integer k so that G has an edge k -coloring. We denote by $\mathcal{C}^k(G)$ the set of all edge k -colorings of G .

In 1960's, Gupta [10] and, independently, Vizing [25] showed that for all graphs G , $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$. This leads to a natural classification of graphs. Following Fiorini and Wilson [8], a graph G is of *class 1* if $\chi'(G) = \Delta(G)$ and of *class 2* if $\chi'(G) = \Delta(G) + 1$. Holyer [12] showed that it is NP-complete to determine whether an arbitrary graph is of class 1. However, there are sufficient conditions for a graph to be class 2. For example, any graph G with $|E(G)| > \Delta(G) \lfloor |V(G)|/2 \rfloor$ is class 2, since each color class of G is a matching and each matching in G has size at most $\lfloor |V(G)|/2 \rfloor$, we have $\chi'(G) \geq |E(G)| / \lfloor |V(G)|/2 \rfloor > \Delta(G)$ but $\chi'(G) \leq \Delta(G) + 1$ by the result of Gupta and Vizing. Such graphs G are called *overfull* (we can define a multigraph to be overfull in the same way). An overfull subgraph H of G with $\Delta(H) = \Delta(G)$ is called a $\Delta(G)$ -*overfull subgraph* of G . A number of long-standing conjectures listed in *Twenty Pretty Edge Coloring Conjectures* in [24] lie in deciding when a graph is overfull. Chetwynd and Hilton [3, 4], in 1986, proposed the following conjecture.

Conjecture 1.1 (Overfull Conjecture). *Let G be a graph with $\Delta(G) > \frac{1}{3}|V(G)|$. Then $\chi'(G) = \Delta(G)$ if and only if G contains no $\Delta(G)$ -overfull subgraph.*

The 3-critical graph P^* , obtained from the Petersen graph by deleting one vertex, has $\chi'(P^*) = 4$, satisfies $\Delta(P^*) = \frac{1}{3}|V(P^*)|$ but contains no 3-overfull subgraph. Thus the degree condition $\Delta(G) > \frac{1}{3}|V(G)|$ in the conjecture above is best possible. Applying Edmonds' matching polytope theorem, Seymour [22] showed that whether a graph G contains an overfull subgraph of maximum degree $\Delta(G)$ can be determined in polynomial time. Thus if the Overfull Conjecture is true, then the NP-complete problem of determining the chromatic index becomes polynomial-time solvable for graphs G with $\Delta(G) > \frac{1}{3}|V(G)|$. There have been some fairly strong results supporting the Overfull Conjecture in the case when G is regular. It is easy to verify that when G is regular with even order, G has no $\Delta(G)$ -overfull subgraphs if its vertex degrees are at least $\frac{1}{2}|V(G)|$. Thus the well-known 1-Factorization Conjecture stated below is a special case of the Overfull Conjecture.

Conjecture 1.2 (1-Factorization Conjecture). *Let G be a regular graph of order $2n$ with degree at least n if n is odd, or at least $n-1$ if n is even. Then G is 1-factorable; equivalently, $\chi'(G) = \Delta(G)$.*

Hilton and Chetwynd [2] verified the 1-Factorization Conjecture if the vertex degree is at least $0.823|V(G)|$. Perković and Reed [18] showed in 1997 that the 1-Factorization Conjecture is true for large regular graphs with vertex degree at least $|V(G)|(2-\varepsilon)$ for any $0 < \varepsilon < 1$. In 2016, Csaba, Kühn, Lo, Osthus and Treglown [6] verified the conjecture for sufficiently large $|V(G)|$. Much less is known about the truth of the Overfull Conjecture if we

no longer require that G is regular. It was confirmed for graphs with $\Delta(G) \geq |V(G)| - 3$ by Chetwynd and Hilton in 1989 [4]. Plantholt [19] in 2004 verified the conjecture for graphs of even order and minimum degree at least $0.8819|V(G)|$. More recently, Plantholt [20] showed the conjecture for sufficiently large even order graphs with minimum degree at least $\frac{2}{3}|V(G)|$. Plantholt [23] and the author extended these results and gave an asymptotic result for general graphs that is similar to the Perković-Reed result for regular graphs, by showing the following result: For all $0 < \varepsilon < 1$, there exists n_0 such that the following statement holds: if G is a graph on $2n \geq n_0$ vertices with $\delta(G) \geq (1 + \varepsilon)n$, then $\chi'(G) = \Delta(G)$ if and only if G contains no $\Delta(G)$ -overfull subgraph. Furthermore, there is a polynomial time algorithm that finds an optimal coloring. All the results mentioned above on graphs with minimum degree constraints are for graphs of even order. In this paper, we obtain the first result (as far as we know) on the Overfull Conjecture with large minimum degree constraints for graphs of odd order.

Theorem 1.3. *For all $0 < \varepsilon < 1$, there exists n_0 such that the following statement holds: if G is a graph on $2n - 1 \geq n_0$ vertices with $\delta(G) \geq (1 + \varepsilon)n$, then $\chi'(G) = \Delta(G)$ if and only if G is not overfull. Furthermore, there is a polynomial time algorithm that finds an optimal coloring.*

The remainder of this paper is organized as follows. In the next section, we introduce some notation and preliminary results. In Section 3, we prove Theorem 1.3 by applying Theorem 3.1 from Section 3. Theorem 3.1 is then proved in the last section.

2 Notation and preliminaries

2.1 Results on degree sequences, paths, matchings, and overfull graphs

We need the following classic result of Hakimi [11] on multigraphic degree sequence.

Theorem 2.1. *Let $0 \leq d_n \leq \dots \leq d_1$ be integers. Then there exists a multigraph G on vertices x_1, \dots, x_n such that $d_G(x_i) = d_i$ for all i if and only if $\sum_{i=1}^n d_i$ is even and $\sum_{i>1} d_i \geq d_1$.*

Though it is not explicitly stated in [11], the inductive proof yields a polynomial time algorithm which finds an appropriate multigraph if it exists.

Theorem 2.2 ([7]). *If G is a graph on $n \geq 3$ vertices with $\delta(G) \geq \frac{n}{2}$, then G is hamiltonian.*

Following the proof of Dirac [7], a hamiltonian cycle can be constructed in polynomial time in n if $\delta(G) \geq \frac{n}{2}$. In fact, there is a polynomial time algorithm that constructs the closure of a graph G and finds a hamiltonian cycle of G if its closure is a complete graph (see [1, Exercise 4.2.15, page 62]).

A path P connecting two vertices u and v is called a (u, v) -path, and we write uPv or vPu in order to specify the two endvertices of P . Let uPv and xQy be two disjoint paths. If vx is an edge, we write $uPvxQy$ as the concatenation of P and Q through the edge vx . If P is a path and $x, y \in V(P)$, then xPy is the subpath of P with endvertices x and y .

We will use the following notation: $0 < a \ll b \leq 1$. Precisely, if we say a claim is true provided that $0 < a \ll b \leq 1$, then this means that there exists a non-decreasing function $f : (0, 1] \rightarrow (0, 1]$ such that the statement holds for all $0 < a, b \leq 1$ satisfying $a \leq f(b)$.

Lemma 2.3 ([23]). *Let $0 < \frac{1}{n_0} \ll \varepsilon < 1$, and G be graph on $n \geq n_0$ vertices such that $\delta(G) \geq \frac{1}{2}(1 + \varepsilon)n$. Moreover, let $M = \{a_1b_1, \dots, a_tb_t\}$ be a matching in the complete graph on $V(G)$ of size at most $\frac{\varepsilon n}{8}$. Then there exist vertex-disjoint paths P_1, \dots, P_t in G such that $\bigcup V(P_i) = V(G)$ and P_i joins a_i to b_i , and these paths can be found in polynomial time.*

A multigraph G is a *star-multigraph* if G has a vertex x that is incident with all multiple edges of G . In other words, $G - x$ is a simple graph. The vertex x is called the *multi-center* of G .

Lemma 2.4. *Let $0 < \frac{1}{n_0} \ll \eta \ll \varepsilon < 1$, and G be star-multigraph graph on $n \geq n_0$ vertices such that $\delta(G) \geq \frac{1}{2}(1 + \varepsilon)n$ and $\mu(x) < \eta n$, where x is the multi-center of G . Moreover, let $M = \{a_1b_1, \dots, a_tb_t\}$ be a matching in the complete graph on $V(G)$ of size at most $\frac{\varepsilon n}{13}$. If $|N_G(x) \setminus \{a_1, b_1, \dots, a_t, b_t\}| \geq 2$, then there exist vertex-disjoint paths P_1, \dots, P_t in G such that $\bigcup V(P_i) = V(G)$ and P_i joins a_i to b_i , and these paths can be found in polynomial time.*

Proof. By relabeling the matching edges if necessary, assume that if $x \in \{a_1, b_1, \dots, a_t, b_t\}$, then $x = a_t$.

If $x \in \{a_1, b_1, \dots, a_t, b_t\}$ and so $x = a_t$, then let $a'_t \in N_G(x) \setminus \{a_1, b_1, \dots, a_t, b_t\}$, and $M' = (M \setminus \{a_tb_t\}) \cup \{a'_tb_t\}$. If $x \notin \{a_1, b_1, \dots, a_t, b_t\}$, then let $x_1, x_2 \in N_G(x) \setminus \{a_1, b_1, \dots, a_t, b_t\}$ be distinct, and let $M' = (M \setminus \{a_tb_t\}) \cup \{a_tx_1, x_2b_t\}$.

Note that $\delta(G - x) \geq \frac{1}{2}(1 + \varepsilon)n - \eta n \geq \frac{1}{2}(1 + \frac{2\varepsilon}{3})n$ and $|M'| \leq \frac{\varepsilon n}{13} + 1 < \frac{\varepsilon n}{12}$. Applying Lemma 2.3 to $G - x$ with matching M' , we find vertex-disjoint paths P'_1, \dots, P'_t in $G - x$ such that $\bigcup V(P'_i) = V(G - x)$. Furthermore, if $x = a_t$, P'_i joins a_i to b_i for $i \in [1, t - 1]$, and P'_t joins a'_t to b_t ; if $x \notin \{a_1, b_1, \dots, a_t, b_t\}$, P'_i joins a_i to b_i for $i \in [1, t - 1]$, P'_t joins a_t to x_1 , and P'_{t+1} joins x_2 to b_t . Letting $P_i = P'_i$ for $i \in [1, t - 1]$, and $P_t = a_t a'_t P'_t b_t$ if $x = a_t$ and $P_t = a_t P'_t x_1 x_2 P'_{t+1} b_t$ if $x \notin \{a_1, b_1, \dots, a_t, b_t\}$ gives the desired paths for G and M . By Lemma 2.3 and the simple adjustment on M above, it is clear that these paths can be found also in polynomial time. \square

Lemma 2.5. *Let G be a graph of order $2n$ and minimum degree at least 1. Assume that except at most one vertex all other vertices have degree at least $n + 1$ in G . Then G has a perfect matching, and the matching can be found in polynomial time.*

Proof. Let $u \in V(G)$ such that $d_G(u) = \delta(G)$. Since $\delta(G) \geq 1$, we let v be a neighbor of u , and let $G_1 = G - \{u, v\}$. Since $\delta(G - u) \geq n$, we have $\delta(G_1) \geq n - 1 = \frac{1}{2}(2n - 2)$.

Applying Theorem 2.2, G_1 has a perfect matching M . Thus $M \cup \{uv\}$ is a perfect matching of G . By the construction of G_1 and the comments immediately below Theorem 2.2, such a perfect matching of G can be found in polynomial time. \square

Lemma 2.6. *Let $G[X, Y]$ be bipartite star-multigraph with $|X| = |Y| = n$ and multi-center x . Suppose that $\delta(G) \geq 1$ and $\delta(G - x) \geq t$ for some $t \in [1, n - 1]$, and except at most t vertices all other vertices of G have simple degree at least $\frac{n}{2} + 1$ in G . Then G has a perfect matching. Furthermore, such a matching can be found in polynomial time.*

Proof. Suppose, without loss of generality, that $x \in X$. As $\delta(G) \geq 1$, we let $y \in Y$ be a neighbor of x . Let $X' = X \setminus \{x\}$ and $Y' = Y \setminus \{y\}$. It suffices to show that $G' := G[X', Y']$ satisfies Hall's Condition. If not, we let $S \subseteq X'$ with smallest cardinality such that $|S| > |N_{G'}(S)|$. Since $\delta(G') \geq 1$, it follows that $|S| \geq 1$. Then by the choice of S , we have $|S| = |N_{G'}(S)| + 1$ and $|N_{G'}(S)| < |Y'|$. As $|S| > |N_{G'}(S)|$, it follows that $|S| \geq \delta(G') + 1 \geq t + 1$. As G' has at most t vertices of degree less than $\frac{n}{2}$, S contains a vertex of degree at least $\frac{n}{2}$ in G' and so $|N_{G'}(S)| \geq \frac{n}{2}$. Thus $|S| > \frac{n}{2}$ and consequently $|X' \setminus S| < \frac{n}{2}$. Since $|N_{G'}(S)| < |Y'|$, there exists $z \in Y' \setminus N_{G'}(S)$ such that $N_{G'}(z) \subseteq X' \setminus S$. As $\delta(G') \geq t$, we have $|X' \setminus S| \geq t$. As $|Y' \setminus N_{G'}(S)| = |Y'| - |S| + 1 = |X'| - |S| + 1 \geq t + 1$ and G' has at most t vertices of degree less than $\frac{n}{2}$, $Y' \setminus N_{G'}(S)$ contains a vertex of degree at least $\frac{n}{2}$ in G' . As a result, $|X' \setminus S| \geq \frac{n}{2}$, contradicting the earlier assertion that $|X' \setminus S| < \frac{n}{2}$. Hence G' has a perfect matching.

There are polynomial time algorithms such as the Hopcroft-Karp algorithm [13] in finding a maximum matching in any bipartite graph, thus a perfect matching of G can be found in polynomial time. \square

Lemma 2.7. *Let $0 < \frac{1}{n_0} \ll \varepsilon < 1$ and G be a star-multigraph with multi-center x on $2n \geq n_0$ vertices. Suppose that $d_G(x) = \delta(G)$, $\delta(G - x) \geq (1 + \varepsilon)n$, and $G - x$ contains no $\Delta(G)$ -overfull subgraph. Then G contains no $\Delta(G)$ -overfull subgraph.*

Proof. Let $X \subseteq V(G)$ with $|X|$ odd. We show that $G[X]$ is not $\Delta(G)$ -overfull. Assume first that $3 \leq |X| \leq 2n - 3$. By symmetry between X and $V(G) \setminus X$, we further assume $|X| \leq n$. Then as $\delta(G - x) \geq (1 + \varepsilon)n$ and $G - x$ is simple, we have

$$e_G(X, V(G) \setminus X) \geq (|X| - 1)((1 + \varepsilon)n - |X| + 1) > 2n > \Delta(G).$$

Thus $2e(G[X]) < \Delta(G)|X| - \Delta(G)$ and so $G[X]$ is not $\Delta(G)$ -overfull. Thus, the only possible $\Delta(G)$ -overfull subgraph of G is obtained from G by deleting a single vertex. Note that if $G - u$ is $\Delta(G)$ -overfull for some $u \in V(G)$, then so is $G - x$ as x has minimum degree in G . However, $G - x$ is not $\Delta(G)$ -overfull by our assumption. Thus $G - u$ is not $\Delta(G)$ -overfull for any $u \in V(G)$. Now as $G[X]$ is not $\Delta(G)$ -overfull for any $X \subseteq V(G)$ with $|X|$ odd, it follows that G contains no $\Delta(G)$ -overfull subgraph. \square

2.2 Results on edge colorings

Let $\varphi \in \mathcal{C}^k(G)$ for some integer $k \geq 0$. For any $v \in V(G)$, the set of colors *present* at v is $\varphi(v) = \{\varphi(e) : e \in E(G) \text{ is incident to } v\}$, and the set of colors *missing* at v is $\bar{\varphi}(v) = [1, k] \setminus \varphi(v)$. For a subset X of $V(G)$ and a color $i \in [1, k]$, define $\bar{\varphi}_X^{-1}(i) = \{v \in X : i \in \bar{\varphi}(v)\}$. We simply write $\bar{\varphi}^{-1}(i)$ for $\bar{\varphi}_{V(G)}^{-1}(i)$.

For two distinct colors $\alpha, \beta \in [1, k]$, the components of the subgraph induced by edges with colors α or β are called (α, β) -chains. Each (α, β) -chain is either a path or an even cycle. If we interchange the colors α and β on an (α, β) -chain C of G , we get a new edge k -coloring of G , which is denoted by φ/C . This operation is called a *Kempe change*. For an (α, β) -chain P , if it is a path with an endvertex u , we also denote it by $P_u(\alpha, \beta, \varphi)$ to emphasize the endvertex u . Let $u, v \in V(G)$. If u and v are contained in the same (α, β) -chain of G with respect to φ , we say that u and v are (α, β) -*linked* with respect to φ .

In 1960's, Gupta [10] and, independently, Vizing [25] provided an upper bound on the chromatic index of multigraphs, and König [15] gave an exact value of the chromatic index for bipartite multigraphs.

Theorem 2.8 ([10, 25]). *Every multigraph G satisfies $\chi'(G) \leq \Delta(G) + \mu(G)$.*

Theorem 2.9 ([15]). *Every bipartite multigraph G satisfies $\chi'(G) = \Delta(G)$.*

An edge k -coloring of a multigraph G is said to be *equalized* if each color class contains either $\lfloor |E(G)|/k \rfloor$ or $\lceil |E(G)|/k \rceil$ edges. McDiarmid [16] observed the following result.

Theorem 2.10. *Let G be a multigraph with chromatic index $\chi'(G)$. Then for all $k \geq \chi'(G)$, there is an equalized edge-coloring of G with k colors.*

Let G be a multigraph, $k \geq 0$ be an integer and $\varphi \in \mathcal{C}^k(G)$. There is a polynomial time algorithm to modify φ into an equalized edge-coloring of G with k colors. To see this, suppose φ is not equalized and so we take two colors $i, j \in [1, k]$ such that $||\bar{\varphi}^{-1}(i)| - |\bar{\varphi}^{-1}(j)||$ is largest. Since φ is not equalized, $||\bar{\varphi}^{-1}(i)| - |\bar{\varphi}^{-1}(j)|| \geq 4$. Assume by symmetry that $|\bar{\varphi}^{-1}(i)| - |\bar{\varphi}^{-1}(j)| \geq 4$. Consider the submultigraph of G induced on the set of edges colored by i or j , then the submultigraph must have a component that is a path P starting at an edge colored by j and ending at an edge colored by i . By swapping the colors i and j along this path P , we decreased $|\bar{\varphi}^{-1}(i)| - |\bar{\varphi}^{-1}(j)|$ by 4. Repeating this process, we can obtain an equalized edge-coloring of G with k colors in $O(k^2|V(G)|^2)$ time.

Lemma 2.11 ([23]). *Let G be a star-multigraph on $2n$ vertices with multi-center x , and let A and B be a partition of $V(G)$ with $|A| = |B|$, where we assume $x \in A$. If $e(A) = e(B)$, $E_G(A, B) = E_G(x, B)$, and G has an edge coloring using k colors, then there exists an edge coloring φ using k colors such that for each $i, j \in [1, k]$, $|\bar{\varphi}_A^{-1}(i)| = |\bar{\varphi}_B^{-1}(i)|$ and $||\bar{\varphi}_A^{-1}(i)| - |\bar{\varphi}_A^{-1}(j)|| \leq 2$. Furthermore, such a coloring φ can be found in $O(k^2n^2)$ -time.*

Lemma 2.12. *Let G be a star-multigraph on $2n$ vertices with multi-center x , and let A and B be a partition of $V(G)$ with $|A| = |B|$, where we assume $x \in A$. If $E_G(A, B) = E_G(x, B)$, then any edge k -coloring of G can be modified into an edge k -coloring φ of G such that the restrictions of φ on both $G[A]$ and $G[B]$ are equitable. Furthermore, such a coloring φ can be found in $O(k^2n^2)$ -time.*

Proof. We choose an edge k -coloring φ of G such that

$$g_\varphi := \max_{C \in \{A, B\}} \max_{i, j} \left| |\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)| \right|$$

is smallest and subject to this, the number h_φ of color pairs (i, j) with $\left| |\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)| \right| = g_\varphi$ is smallest. If $g_\varphi \leq 2$, then we are done. Thus $g_\varphi \geq 3$. We assume, without loss of generality, that there exist $i, j \in [1, k]$ such that $|\overline{\varphi}_A^{-1}(i)| - |\overline{\varphi}_A^{-1}(j)| = g_\varphi \geq 3$. As $E_G(A, B) = E_G(x, B)$ and different (i, j) -chains are vertex-disjoint, at most one vertex from $\overline{\varphi}_A^{-1}(i)$ is (i, j) -lined with a vertex from B . As $|\overline{\varphi}_A^{-1}(i)| - |\overline{\varphi}_A^{-1}(j)| \geq 3$, there exist distinct $u, v \in \overline{\varphi}_A^{-1}(i)$ such that $P_u(i, j, \varphi) = P_v(i, j, \varphi)$. We now let $\psi = \varphi / P_u(i, j, \varphi)$. Then $g_\psi \leq g_\varphi$ and $h_\psi < h_\varphi$, showing a contradiction to the choice of φ . Thus we can find a coloring φ such that $\left| |\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)| \right| \leq 2$ for any $C \in \{A, B\}$ and all $i, j \in [1, k]$.

There are at most $\binom{k}{2}$ color pairs, each color pair takes $O(n)$ -time to reduce the difference of the two corresponding color classes by 2, and $O(n)$ steps to make the two corresponding color classes close in size. Thus a desired edge coloring φ can be found in $O(k^2n^2)$ -time. \square

Lemma 2.13 ([23]). *If G is a star-multigraph, then $\chi'(G) \leq \Delta(G) + 1$. Furthermore, an edge coloring of G using at most $\Delta(G) + 1$ colors can be found in polynomial time.*

A multigraph G is a *near star-multigraph* with multi-center x if, in $G - x$, there is at most one pair of vertices u and v such that $e_{G-x}(u, v) \geq 2$.

Lemma 2.14. *Let G be a near star-multigraph with multi-center x such that $\mu(G - x) = e_{G-x}(y, z)$ for some $y, z \in V(G - x)$. Then $\chi'(G) \leq \max\{\Delta(G) + e_G(y, z), \Delta(G) + 1\}$. Furthermore, an edge $\max\{\Delta(G) + e_G(y, z), \Delta(G) + 1\}$ -coloring of G can be found in polynomial time.*

Proof. We may assume $e_G(y, z) \geq 2$, as otherwise the statement follows from Lemma 2.13. Let G' be obtained from G by deleting $e_G(y, z) - 1$ edges joining y and z . Applying Lemma 2.13 on G' , we have $\chi'(G') \leq \Delta(G') + 1 \leq \Delta(G) + 1$. Now based on any edge $(\Delta(G') + 1)$ -coloring of G' , coloring each of the deleted edges using a distinct color gives an edge $(\Delta(G') + 1 + e_G(y, z) - 1)$ -coloring of G . As $\Delta(G') + 1 + e_G(y, z) - 1 \leq \Delta(G) + e_G(y, z)$, we have $\chi'(G) \leq \Delta(G) + e_G(y, z)$. By Lemma 2.13, there is a polynomial time algorithm to edge color G' using at most $\Delta(G) + 1$ colors. Thus there is a polynomial time algorithm to edge color G using at most $\max\{\Delta(G) + e_G(y, z), \Delta(G) + 1\}$ colors. \square

Given an edge coloring of G and a given color i , since vertices presenting i are saturated by the matching consisting of all edges colored by i , we have the Parity Lemma below, see [9, Lemma 2.1].

Lemma 2.15 (Parity Lemma). *Let G be a multigraph and $\varphi \in \mathcal{C}^k(G)$ for some integer $k \geq \Delta(G)$. Then $|\varphi^{-1}(i)| \equiv |V(G)| \pmod{2}$ for every color $i \in [1, k]$.*

3 Proof of Theorem 1.3

For $u \in V(G)$, the *deficiency* of u in G is $\text{def}_G(u) := \Delta(G) - d_G(u)$. For $U \subseteq V(G)$, $\text{def}_G(U) = \sum_{u \in U} \text{def}_G(u)$. We simply write $\text{def}_G(V(G))$ as $\text{def}(G)$. For an integer $i \geq 1$, define $V_i(G) = \{v \in V(G) : d_G(v) = i\}$, and we write V_i for $V_i(G)$ if G is clear. The proof of Theorem 1.3 is based on the following result, which will be proved in the last section.

Theorem 3.1. *For all $0 < \varepsilon < 1$, there exists n_0 and η with $0 < \frac{1}{n_0} \ll \eta \ll \varepsilon$ such that the following statement holds. Let G be a near star-multigraph on $2n \geq n_0$ vertices with multi-center x , and let $U = \{v \in V(G) : \Delta(G) - d_G(v) \geq \eta n\}$. Suppose that G satisfies one of the following five conditions:*

- (a) *G is a star-multigraph and is regular with $\delta(G) \geq \delta(G - x) \geq (1 + \frac{\varepsilon}{2})n$ and $\mu(x) < \eta n$.*
- (b) *G is regular with $\delta(G) \geq (1 + \varepsilon)n$, and there exist distinct $y, z \in V(G) \setminus \{x\}$ with the following properties:*
 - (i) *$d_G^s(x) \geq 2$ and there are at most $n^{\frac{1}{2}}$ neighbors w of x satisfying $e_G(x, w) \geq \eta n$;*
 - (ii) *y and z are the only two possible vertices in $G - x$ with $e_{G-x}(y, z) \geq 2$ but $\mu(G - x) < n^{\frac{1}{2}}$; and*
 - (iii) *$d_G^s(w) \geq (1 + \varepsilon)n$ for all $w \in V(G) \setminus \{x\}$.*
- (c) *G is a star-multigraph and is regular with $\delta(G) \geq (1 + \varepsilon)n$, and there exists $y \in V(G) \setminus \{x\}$ such that*
 - (i) *$2 \leq d^s(x) < n^{\frac{1}{2}}$;*
 - (ii) *$d_G^s(y) \geq 2\varepsilon n$; and*
 - (iii) *$d_G^s(w) \geq (1 + \varepsilon)n$ for all $w \in V(G) \setminus \{x, y\}$.*
- (d) *G is a star-multigraph and G has two distinct vertices y, z such that*
 - (i) *$\mu(x) < \eta n$;*
 - (ii) *$d_G(y) = d_G(z) = \delta(G) \geq (\frac{1}{2} + \frac{3\varepsilon}{2})n$ and $\min\{d_G^s(y), d_G^s(z)\} > \Delta(G) - \frac{1}{2}(1 - 0.9\varepsilon)n$;*
and
 - (iii) *$d_G(w) = \Delta(G) \geq (1 + \varepsilon)n$ and $d_G^s(w) \geq (1 + \varepsilon)n - \mu(x) \geq (1 + 0.9\varepsilon)n$ for all $w \in V(G) \setminus \{y, z\}$.*
- (e) *G is a star-multigraph, $|U| \geq \eta n$, $\delta(G) \geq \delta(G - x) \geq (1 + \varepsilon)n$, and $\mu(x) \leq \frac{2}{\eta}$.*

Then $\chi'(G) = \Delta(G)$. Furthermore, there is a polynomial time algorithm that finds an optimal coloring.

Theorem 1.3. *For all $0 < \varepsilon < 1$, there exists n_0 such that the following statement holds: if G is a graph on $2n - 1 \geq n_0$ vertices with $\delta(G) \geq (1 + \varepsilon)n$, then $\chi'(G) = \Delta(G)$ if and only if G is not overfull. Furthermore, there is a polynomial time algorithm that finds an optimal coloring.*

Proof. The general strategy for proving the first part of the statement is to add a vertex to G to produce a star-multigraph G' and then take off some matchings or linear forests from G' such that the resulting multigraph satisfies one of the conditions listed in Theorem 3.1. Choose integer n_0 and a real number η so that $0 < \frac{1}{n_0} \ll \eta \ll \varepsilon < 1$.

If $\chi'(G) = \Delta(G)$, then clearly G contains no $\Delta(G)$ -overfull subgraph. Thus we assume that G is not overfull and show $\chi'(G) = \Delta(G)$. Since n is odd and G is not overfull, we have $\text{def}(G) \geq \Delta(G)$. We call a vertex of degree less than $\Delta(G)$ but greater than $\delta(G)$ a *middle degree* vertex. Let

$$W = \{v \in V(G) : \Delta(G) - d_G(v) \geq \eta n\}.$$

We consider the following four cases regarding the size of W .

Case 1. $|W| \geq 2\eta n$.

We let $W' \subseteq W$ be a set of $\lfloor \eta n \rfloor$ vertices, and let x be a new vertex. We form a new graph G' as follows: add at most $\lfloor \frac{2}{\eta} \rfloor$ edges between x and each vertex of W' one by one until the degree of x in the resulting multigraph reaches to $\delta(G)$. Since $\lfloor \frac{2}{\eta} \rfloor \times \lfloor \eta n \rfloor = 2n > \delta(G)$ (we can choose η so that $\frac{1}{\eta}$ is an integer), the star-multigraph G' exists. As $\eta n > \lfloor \frac{2}{\eta} \rfloor$, we have $\Delta(G') = \Delta(G)$. By Lemma 2.7, G' contains no $\Delta(G')$ -overfull subgraph.

For each vertex $u \in V(G') \setminus \{x\}$, we have $d_{G'}^s(u) \geq \delta(G) \geq (1 + \varepsilon)n$. For each vertex $w \in W \setminus W'$, we have $\Delta(G') - d_{G'}(w) = \Delta(G) - d_G(w) \geq \eta n$. Furthermore, $|W \setminus W'| \geq \eta n$. Now with $W \setminus W'$ playing the role of U in Theorem 3.1(e) and applying Theorem 3.1, we see that $\chi'(G') = \Delta(G') = \Delta(G)$ and so $\chi'(G) = \Delta(G)$.

Case 2. $|W| = 0$.

The proof of this case is similar to the beginning part in the proof of Theorem 1.3 in [23], but we repeat it to be self-contained. Note that $|W| = 0$ implies that $\Delta(G) - \delta(G) < \eta n$. Let x be a new vertex and let G' be obtained from G by adding x to G and adding some edges between x and vertices $y \in V(G)$ with $d_G(y) < \Delta(G)$ with the following constraints:

- (1) $d_{G'}(x) = \delta(G') = \delta(G)$ and $\Delta(G') = \Delta(G)$;
- (2) subject to (1), at most one neighbor of x in G' is not a maximum degree vertex of G' .

Let $V(G) = \{x_1, \dots, x_{2n-1}\}$ and assume $d_G(x_1) \leq \dots \leq d_G(x_{2n-1})$. Note $d_G(x_{2n-1}) = \Delta(G)$. Let $s \in [1, 2n - 2]$ be the smallest integer such that $\sum_{i=1}^s \text{def}_G(x_i) \geq \delta(G)$; such an integer s exists since $\text{def}(G) \geq \Delta(G)$. For each $i \in [1, s - 1]$, we add exactly $\text{def}_G(x_i)$

edges between x and x_i , and we add $\delta(G) - \sum_{i=1}^{s-1} \text{def}_G(x_i)$ edges between x and x_s , and call G' the resulting multigraph. It is clear that G' satisfies constraints (1) and (2). Since $\Delta(G) - \delta(G) < \eta n$, we have

$$|N_{G'}(x)| \geq \frac{(1 + \varepsilon)n}{\eta n} > 3.$$

By Lemma 2.7, G' contains no $\Delta(G')$ -overfull subgraph.

Since x has the smallest degree in G' and $G = G' - x$ is not overfull, we get $\sum_{i \geq 1} \text{def}_{G'}(x_i) \geq \text{def}_{G'}(x)$. Since $|V(G')| = 2n$ is even, $\text{def}_{G'}(x) + \sum_{i \geq 1} \text{def}_{G'}(x_i)$ is even. Then by Theorem 2.1, there exists a multigraph H on $V(G')$ such that $d_H(x) = \text{def}_{G'}(x)$ and $d_H(x_i) = \text{def}_{G'}(x_i)$ for each $i \in [1, 2n - 1]$. The multigraph H will aid us in finding a spanning regular subgraph of G' .

Note that $\Delta(H) = \text{def}_{G'}(x) = \Delta(G) - \delta(G) < \eta n$ and H contains isolated vertices. Thus $\chi'(H) \leq \Delta(H) + \mu(H) \leq 2\Delta(H) < 2\eta n$ by Theorem 2.8. Hence we can greedily partition $E(H)$ into $k \leq 52 \frac{m}{\varepsilon}$ matchings M_1, \dots, M_k each of size at most $\frac{\varepsilon n}{26}$. Now we take out linear forests (union of vertex-disjoint paths) from G' by applying Lemma 2.4 with M_1, \dots, M_k . As at most one neighbor of x is not a maximum degree vertex of G' , it follows that x is adjacent in G' to at most one vertex of those vertices covered by M_i for each $i \in [1, k]$. More precisely, define spanning subgraphs G_0, \dots, G_k of G' and edge-disjoint linear forests F_1, \dots, F_k such that

- (1) $G_0 := G'$ and $G_i = G_{i-1} - E(F_i)$ for $i \in [1, k]$,
- (2) F_i is a spanning linear forest in G_{i-1} whose leaves are precisely the vertices in M_i .

Let $G_0 = G'$ and suppose that for some $i \in [1, k]$, we already defined G_0, \dots, G_{i-1} and F_1, \dots, F_{i-1} . As $\Delta(F_1 \cup \dots \cup F_{i-1}) \leq 2(i-1) \leq 104 \frac{m}{\varepsilon}$, it follows that $\delta(G_{i-1}) \geq (1 + \varepsilon)n - 104 \frac{m}{\varepsilon} \geq (1 + \frac{\varepsilon}{2})n$. Furthermore, by the construction of F_1, \dots, F_{i-1} , we know that maximum degree vertices of G' are still maximum degree vertices of G_{i-1} . Thus in G_{i-1} , still at most one neighbor of x is not a maximum degree vertex of G_{i-1} . As $\mu(x) < \eta n$ and $\delta(G_{i-1}) \geq (1 + \frac{\varepsilon}{2})n$, x has at least two neighbors in G_{i-1} that are not covered by M_i . Since M_i has size at most $\frac{\varepsilon n}{26}$, we can apply Lemma 2.4 to G_{i-1} and M_i and obtain a spanning linear forest F_i in G_{i-1} whose leaves are precisely the vertices in M_i . Set $G_i := G_{i-1} - E(F_i)$.

We claim that G_k is regular. Consider any vertex $u \in V(G_k)$. For every $i \in [1, k]$, $d_{F_i}(u) = 1$ if u is an endvertex of some edge of M_i and $d_{F_i}(u) = 2$ otherwise. Since M_1, \dots, M_k partition $E(H)$, we know that $\sum_{i=1}^k d_{F_i}(u) = 2k - d_H(u) = 2k - \text{def}_{G'}(u)$. Thus

$$d_{G_k}(u) = d_{G'}(u) - \sum_{i=1}^k d_{F_i}(u) = d_{G'}(u) - (2k - \text{def}_{G'}(u)) = \Delta(G') - 2k.$$

Note that $\mu(x) < \eta n$ and $\delta(G_k - x) \geq (1 + \varepsilon)n - 104 \frac{m}{\varepsilon} - \mu(x) > (1 + \frac{\varepsilon}{2})n$. Thus the multigraph G_k satisfies Condition (a) of Theorem 3.1. Now $\chi'(G_k) = \Delta(G_k)$ by Theorem 3.1. We color

the edges of F_i using 2 distinct colors from $[\Delta(G') - 2k + 1, \Delta(G')]$ for each $i \in [1, k]$. It is clear that any edge $\Delta(G_k)$ -coloring of G_k together with this coloring of $\bigcup_{i=1}^k F_i$ gives an edge coloring of G' using $\Delta(G_k) + 2k = \Delta(G')$ colors. Since G is a subgraph of G' and $\Delta(G) = \Delta(G')$, it follows that $\chi'(G) = \Delta(G)$.

Case 3. $n^{\frac{1}{2}} \leq |W| < 2\eta n$.

We add a new vertex x . For each vertex $v \in V(G) \setminus (V_\Delta \cup W)$, we add $\text{def}_G(v)$ edges between x and v until the degree of x reaches to $\delta(G)$. If the degree of x reaches to $\delta(G)$ already in this process, we denote the resulting multigraph by G' . If the degree of x is still less than $\delta(G)$ after we adding edges between x and all vertices of $V(G) \setminus (V_\Delta \cup W)$, then we add at most $2n^{\frac{1}{2}}$ edges between x and each vertex of W until the degree of x reaches to $\delta(G)$. Denote the resulting multigraph by G' . As $\text{def}(G) \geq \text{def}_G(W) > \Delta$ and $d_{G'}(x) = \delta(G)$, it follows that G' is not regular.

Note that $V_\delta(G') \subseteq W \cup \{x\}$ by the construction of G' and so $|V_\delta(G')| \leq 2\eta n + 1$. Thus for any $v \in V(G') \setminus V_\delta(G')$, $\delta(G' - v - V_\delta(G')) \geq n$ and so both $G' - V_\delta(G')$ and $G' - v - V_\delta(G')$ are hamiltonian by Theorem 2.2. Thus if $G' - V_\delta(G')$ and $G' - v - V_\delta(G')$ have even order, then they each has a perfect matching. Hence if $|V_\delta(G')|$ is even, we can decrease $\Delta(G') - \delta(G')$ but preserve $\delta(G')$ in deleting a perfect matching M of $G' - V_\delta(G')$. If $|V_\delta(G')|$ is odd but G' has a middle degree vertex v , we can decrease $\Delta(G') - \delta(G')$ but preserve $\delta(G')$ in deleting a perfect matching M of $G' - v - V_\delta(G')$. Denote by G_1 the reduced multigraph from G' by deleting M in either of these two cases. We show in the following that we can consider G_1 in the place of G' . As $\delta(G_1 - x) = \delta(G' - x) \geq (1 + \varepsilon)n$, $G_1[X]$ is not $\Delta(G_1)$ -overfull for any $X \subseteq V(G_1)$ with $3 \leq |X| \leq 2n - 3$. Thus to show that G_1 contains no $\Delta(G_1)$ -overfull subgraph, we show that $G_1 - u$ is not $\Delta(G_1)$ -overfull for any $u \in V(G_1)$. If $|V_\delta(G')| \geq 2$, then for any $u \in V(G_1)$ and $v \in V_\delta(G_1)$, we have

$$\begin{aligned} \sum_{w \in V(G_1 - u)} (\Delta(G_1) - d_{G_1 - u}(w)) &= d_{G_1}(u) + (\Delta(G_1) - d_{G_1}(v)) + \text{def}_{G_1}(V(G_1) \setminus \{u, v\}) \\ &\geq \Delta(G_1) + \text{def}_{G_1}(V(G_1) \setminus \{u, v\}) \geq \Delta(G_1). \end{aligned}$$

Thus $G_1 - u$ is not $\Delta(G_1)$ -overfull. If $|V_\delta(G')| = 1$, then $V_\delta(G') = \{x\}$. Note that $x \in V_\delta(G_1)$ and the middle degree vertex, say v , exists in G' . Then

$$\begin{aligned} \text{def}(G_1 - x) &= d_{G'}(x) + (\Delta(G_1) - d_{G'}(v)) + \text{def}_{G_1}(V(G_1) \setminus \{x, v\}) \\ &= d_{G'}(x) + (\Delta(G') - 1 - d_{G'}(v)) + \text{def}_{G'}(V(G') \setminus \{x, v\}) \\ &= \sum_{w \in V(G' - x)} (\Delta(G') - d_{G' - x}(w)) - 1. \end{aligned}$$

Since G' contains no $\Delta(G')$ -overfull subgraph, we have $\sum_{w \in V(G' - x)} (\Delta(G') - d_{G' - x}(w)) \geq \Delta(G')$. Thus $\text{def}(G_1 - x) \geq \Delta(G') - 1 = \Delta(G_1)$ and so G_1 contains no $\Delta(G_1)$ -overfull subgraph. Furthermore, $\chi'(G_1) = \Delta(G_1)$ implies that $\chi'(G') = \Delta(G')$. Thus when $|V_\delta(G')|$ is even or G' has a middle degree vertex, we can consider G_1 in the place of G' and show that G_1 is a class 1 graph.

Thus we assume that $|V_\delta(G')|$ is odd and G' has no middle degree vertex. This in particular, implies that $\delta(G')$ and $\Delta(G')$ have the same parity. As G' has no $\Delta(G')$ -overfull subgraph, $|V_\delta(G')| \geq 3$. Let $y, z \in V_\delta(G') \setminus \{x\}$ be distinct. We find a perfect matching M_{11} in $G - (V_\delta(G') \setminus \{y\})$ and a perfect matching M_{12} in $G - (V_\delta(G') \setminus \{z\})$. The matchings exist by Theorem 2.2. Let $G_1 = G' - M_{11} - M_{12}$. We repeat this same process and find a perfect matching M_{21} in $G_1 - (V_\delta(G') \setminus \{y\})$ and a perfect matching M_{22} in $G_1 - (V_\delta(G') \setminus \{z\})$. For $i \in [2, (\Delta(G') - \delta(G'))/2]$, we let $G_i = G_{i-1} - M_{i1} - M_{i2}$. Since $d_{G'}(y) \geq \delta(G') + e_{G'}(x, y) \geq (1 + \varepsilon)n + e_{G'}(x, y)$ and $d_{G'}(z) \geq \delta(G') + e_{G'}(x, z) \geq (1 + \varepsilon)n + e_{G'}(x, z)$, we have $d_{G_i}^s(y), d_{G_i}^s(z) \geq \delta(G') - i$. As $\Delta(G') - \delta(G') < 2n - (1 + \varepsilon)n = (1 - \varepsilon)n$, we see that $d_{G_i}^s(y) = d_{G_i}^s(z) = \delta(G') - i \geq (1 + \varepsilon)n - \frac{1}{2}(1 - \varepsilon)n \geq (\frac{1}{2} + \frac{3\varepsilon}{2})n$. For any vertex $w \in V(G_i) \setminus \{x, y, z\}$, $d_{G_i}^s(w) \geq \delta(G) - \mu(x) \geq (1 + 0.9\varepsilon)n$. As $|V_\delta(G')| + 1 < 2\eta n + 1 \ll 0.9\varepsilon n$, both $G_i - (V_\delta(G') \setminus \{y\})$ and $G_i - (V_\delta(G') \setminus \{z\})$ satisfy the condition of Lemma 2.5, and so each has a perfect matching. Thus for each $i \in [2, (\Delta(G') - \delta(G'))/2]$, we find matchings M_{i1} and M_{i2} respectively from $G_{i-1} - (V_\delta(G') \setminus \{y\})$ and $G_{i-1} - (V_\delta(G') \setminus \{z\})$. We claim that $G^* := G_{(\Delta(G') - \delta(G'))/2}$ satisfies Condition (d) of Theorem 3.1. We have

- (i) $\mu(x) < \eta n$, as there might be edges in G' between x and vertices of $V(G) \setminus (V_\Delta \cup W)$;
- (ii) $d_{G^*}(y) = d_{G^*}(z) = \delta(G^*) \geq (\frac{1}{2} + \frac{3\varepsilon}{2})n$;
- (iii) $\min\{d_{G^*}^s(y), d_{G^*}^s(z)\} \geq \Delta(G^*) - (\Delta(G') - \delta(G'))/2 - \mu(x) \geq \Delta(G^*) - \frac{1}{2}(1 - \varepsilon)n - \mu(x) > \Delta(G^*) - \frac{1}{2}(1 - 0.9\varepsilon)n$;
- (iv) $d_{G^*}(w) = \Delta(G^*) \geq (1 + \varepsilon)n$ and $d_{G^*}^s(w) \geq (1 + \varepsilon)n - \mu(x) \geq (1 + 0.9\varepsilon)n$ for all $w \in V(G^*) \setminus \{y, z\}$.

By Theorem 3.1, $\chi'(G^*) = \Delta(G^*) = \delta(G')$. Taking an edge $\delta(G')$ -coloring of G^* , coloring edges in M_{i1} with color $\delta(G') + 2i - 1$ and coloring edges in M_{i2} with color $\delta(G') + 2i$ for each $i \in [1, (\Delta(G') - \delta(G'))/2]$, we obtain an edge $\Delta(G')$ -coloring of G' and so an edge $\Delta(G)$ -coloring of G .

Case 4. $0 < |W| < n^{\frac{1}{2}}$.

We may assume that $|V_\delta| \geq 2$. Otherwise, suppose $|V_\delta| = 1$; we let M be a perfect matching of $G - V_\delta$ and let $G_1 = G - M$. As $\text{def}(G) \geq \Delta(G)$ by our assumption, we have $\text{def}(G_1) = \text{def}(G) - 1 \geq \Delta(G) - 1 = \Delta(G_1)$. We claim that G_1 contains no $\Delta(G_1)$ -overfull subgraph. As $\delta(G_1) \geq (1 + \varepsilon)n$, $G_1[X]$ is not $\Delta(G_1)$ -overfull for any $X \subseteq V(G_1)$ with $3 \leq |X| \leq 2n - 3$. Thus the only possible $\Delta(G_1)$ -overfull subgraph in G_1 is G_1 itself. However $\text{def}(G_1) \geq \Delta(G_1)$ implies that G_1 is not overfull. Since $\delta(G_1) = \delta(G)$, and $\chi'(G_1) = \Delta(G_1)$ implies that $\chi'(G) = \Delta(G)$, we can consider G_1 in the place of G .

Next we may assume that $\text{def}(G) \geq \Delta(G) + |W| + 1$. For otherwise, we let $y, z \in V_\delta$ be distinct and add $(\text{def}(G) - \Delta(G))/2$ edges between y and z to get G^* (note that $\text{def}(G) - \Delta(G)$ is even as $\text{def}(G) = (2n - 1)\Delta(G) - 2|E(G)|$), and now add a new vertex x and $\Delta(G)$ edge between x and the vertices of degree less $\Delta(G)$ in G^* to get a Δ -regular

graph G' . Note that G' may not be star-multiple, but all the possible multiple edges in $G' - x$ are between y and z and $e_{G'}(y, z) \leq \frac{1}{2}|W| + 1 < n^{\frac{1}{2}}$. As $\text{def}_{G^*}(w) < \Delta(G^*)$ for any $w \in V(G^*)$, we know that $d_{G'}^s(x) \geq 2$. By Theorem 3.1(b), we have $\chi'(G') = \Delta(G') = \Delta(G)$ and so $\chi'(G) = \Delta(G)$.

Thus $\text{def}(G) \geq \Delta(G) + |W| + 1$. Note that $V_\delta \subseteq W$ and so $|V_\delta| \leq |W| < n^{\frac{1}{2}}$. We may further assume that $|V_\delta|$ is even and G has no middle degree vertex. For otherwise, if $|V_\delta|$ is odd, then $G - V_\delta$ has a perfect matching M ; if $|V_\delta|$ is even and G has a middle degree vertex v , then $G - v - V_\delta$ has a perfect matching M . In both cases, let $G_1 = G - M$. As $\text{def}(G) \geq \Delta(G) + |W| + 1$, we have $\text{def}(G_1) \geq \Delta(G) - |W| - 1 \geq \Delta(G) > \Delta(G_1)$. Thus G_1 is not overfull. Furthermore, as $\delta(G_1) = \delta(G) \geq (1 + \varepsilon)n$, $G_1[X]$ is not $\Delta(G_1)$ -overfull for any $X \subseteq V(G_1)$ with $3 \leq |X| \leq 2n - 3$. Thus G_1 contains no $\Delta(G_1)$ -overfull subgraph. Furthermore $\chi'(G_1) = \Delta(G_1)$ implies that $\chi'(G) = \Delta(G)$. Thus we can consider G_1 in the place of G .

Thus we may assume that $|V_\delta| \geq 2$ is even, $\text{def}(G) \geq \Delta(G) + |W| + 1$, and G has no middle degree vertex. Let x be a new vertex and $y \in V_\delta$. We first add $\text{def}_G(y)$ edges between x and y and denote the resulting graph by G' . Note that $V_\delta(G') = \{x\}$. Denote by δ' the second minimum degree of G' . Then we add edges between x and vertices of $V_{\delta'}(G') = V_\delta \setminus \{y\}$ as follows:

Step 1 If $|V_{\delta'}(G')| + d_{G'}(x) \leq \delta' + 1$, then add an edge between x and each vertex from $V_{\delta'}(G')$ and update G' to be the resulting multigraph; otherwise, $|V_{\delta'}(G')| + d_{G'}(x) \geq \delta' + 2$, then we add an edge between x and each vertex of a selected set of $\delta' - d_{G'}(x)$ vertices from $V_{\delta'}(G')$ and update G' to be the resulting multigraph.

Step 2 If $|V_\delta(G')| \geq 2$, then stop; otherwise, return to Step 1.

Since $\text{def}(G) > \Delta(G)$ and $|V_\Delta| \geq 1$, the process above stops before or when the degree of x reaches to $\Delta(G)$. Thus $\Delta(G') = \Delta(G)$. Furthermore, we have $d_{G'}(x) = \delta(G') \geq \delta(G)$ and $d_{G'}^s(x) \geq 2$. Lastly by the process, we have that $x \in V_\delta(G')$, $|V_\delta(G')| \geq 2$, and $0 \leq d_{G'}(w) - d_{G'}(x) \leq 1$ for any $w \in N_{G'}(x) \setminus \{y\}$. Actually, as $\text{def}_G(y) + |W| \leq (1 - \varepsilon)n + n^{\frac{1}{2}} < \delta(G)$, the construction of G' implies $N_{G'}(x) = W$. As $G = G' - x$ is not overfull and $d_{G'}(x) = \delta(G)$, we know that $G' - u$ is not overfull for any $u \in V(G')$. Also, as $\delta(G' - x) \geq (1 + \varepsilon)n$, by calculation, $G'[X]$ is not $\Delta(G')$ -overfull for any $X \subseteq V(G')$ with $3 \leq |X| \leq 2n - 3$. Thus G' contains no $\Delta(G')$ -overfull subgraph. If G' is regular, then as $2 \leq d_{G'}^s(x) \leq |W|$ and $d_{G'}^s(w) \geq (1 + \varepsilon)n$ for any $w \in V(G') \setminus \{x\}$, we are done by Theorem 3.1(c).

Thus we assume that G' is not regular. Since G has no middle degree vertex, we have $|V(G') \setminus V_\Delta(G')| \leq |(V_\delta(G) \setminus \{y\}) \cup \{x\}| < n^{\frac{1}{2}}$. Since $\delta^s(G' - x) \geq (1 + \varepsilon)n$, it follows that for any $v \in V(G') \setminus V_\delta(G')$, in the graph $G' - v - V_\delta(G')$, every vertex has degree at least $n + 1$ except possibly for the vertex y , which has degree at least $(1 + \varepsilon)n - (2n - (1 + \varepsilon)n) - |W| - 1 > 1$. Thus both $G' - V_\delta(G')$ and $G' - v - V_\delta(G')$ have a perfect matching by Lemma 2.5. Thus if $G' - V_\delta(G')$ and $G' - v - V_\delta(G')$ have even order, then they each has a perfect matching.

Hence if $|V_\delta(G')|$ is even, we can decrease $\Delta(G') - \delta(G')$ but preserve $\delta(G')$ in deleting a perfect matching M of $G' - V_\delta(G')$. If $|V_\delta(G')|$ is odd but G' has a middle degree vertex v , we can decrease $\Delta(G') - \delta(G')$ but preserve $\delta(G')$ in deleting a perfect matching M of $G' - v - V_\delta(G')$. Denote by G_1 the reduced graph from G' by deleting M in either of these two cases. As $\delta(G_1) = \delta(G') \geq (1 + \varepsilon)n$, by the same argument as in Case 3, we know that G_1 contains no $\Delta(G_1)$ -overfull subgraph. Furthermore, $\chi'(G_1) = \Delta(G_1)$ implies that $\chi'(G') = \Delta(G')$. Thus we can consider G_1 in the place of G' and show that G_1 is a class 1 graph.

Thus G' has no middle degree vertex. This in particular, implies that $\delta(G')$ and $\Delta(G')$ have the same parity. Also, as $0 \leq d_{G'}(w) - d_{G'}(x) \leq 1$ for any $w \in N_{G'}(x) \setminus \{y\}$, it follows that $d_{G'}(w) = d_{G'}(x) = \delta(G')$ for any $w \in N_{G'}(x) \setminus \{y\}$. Since $W = V_\delta = N_{G'}(x)$, we know that $d_{G'}(w) = d_{G'}(x) = \delta(G')$ for any $w \in V_\delta \setminus \{y\}$. As $V(G) = V_\Delta \cup V_\delta$, and $d_{G'}(y) = \Delta(G')$, it follows that $V_\delta(G') = (V_\delta \setminus \{y\}) \cup \{x\}$ and so $|V_\delta(G')|$ is even and $|V_\Delta(G')|$ is even. As $\Delta(G') - \delta(G') \leq (1 - \varepsilon)n$ and $d_{G'}^s(y) \geq (1 + \varepsilon)n$, by Lemma 2.5, $G' - V_\delta(G')$ has a perfect matching M_1 . Let $G_1 = G' - V_\delta(G') - M_1$. By Lemma 2.5 again, we can find a perfect matching M_2 in G_1 . In general, we let $G_i = G_{i-1} - M_i$ for each $i \in [2, \Delta(G') - \delta(G')]$. We claim that G_i has a perfect matching M_i . Note that $d_{G_i}(y) \geq (1 + \varepsilon)n - (\Delta(G') - \delta(G')) - |V_\delta(G')| \geq (1 + \varepsilon)n - (1 - \varepsilon)n - n^{\frac{1}{2}} \geq \varepsilon n > 1$ and $d_{G_i}(z) \geq \delta(G') - |V_\delta(G')| \geq (1 + \frac{\varepsilon}{2})n$. Thus by Lemma 2.5, G_i has a perfect matching M_{i+1} . The graph $G^* := G_{\Delta(G') - \delta(G')}$ is $\delta(G')$ -regular satisfying the following conditions:

- (i) $2 \leq d_{G^*}^s(x) \leq |W| < n^{\frac{1}{2}}$;
- (ii) $d_{G^*}^s(y) \geq (1 + \varepsilon)n - (\Delta(G') - \delta(G')) \geq 2\varepsilon n$; and
- (iii) $d_{G^*}^s(w) \geq (1 + \varepsilon)n$ for all $w \in V(G^*) \setminus \{x, y\}$, because $w \notin N_{G'}(x)$ if $w \in V(G^*) \setminus (W \cup \{x\})$ and $d_{G^*}^s(w) = d_{G'}^s(w) \geq (1 + \varepsilon)n$ for any $w \in W \setminus \{y\}$.

Now by Theorem 3.1(c), $\chi'(G^*) = \Delta(G^*)$. An edge $\chi'(G^*)$ -coloring of G^* together with a coloring with the edges in M_i colored by i for each $i \in [\Delta(G^*) + 1, \Delta(G')]$ gives an edge $\Delta(G')$ -coloring of G' and so of G .

We lastly check that the procedure above yields a polynomial time algorithm. If G is overfull, then $\chi'(G) = \Delta(G) + 1$ and G can be edge colored using $\Delta(G) + 1$ colors in polynomial time [17]. Thus G is not overfull. In each of the Cases 1 to 4, it is polynomial time (all the results applied in the proof give appropriate running time statements) to reduce G into a multigraph that satisfies one of the Conditions (a) to (e) as listed in Theorem 3.1. Then we find an edge $\Delta(G)$ -coloring of G in polynomial time by Theorem 3.1. \square

4 Proof of Theorem 3.1

We will need the following result, which was proved using Chernoff bound.

Lemma 4.1 ([23], Lemma 3.2). *There exists a positive integer n_0 such that for all $n \geq n_0$ the following holds. Let G be a graph on $2n$ vertices, and $N = \{x_1, y_1, \dots, x_t, y_t\} \subseteq V(G)$, where $1 \leq t \leq n$ is an integer. Then $V(G)$ can be partitioned into two parts A and B satisfying the properties below:*

- (i) $|A| = |B|$;
- (ii) $|A \cap \{x_i, y_i\}| = 1$ for each $i \in [1, t]$;
- (iii) $|d_A(v) - d_B(v)| \leq n^{\frac{2}{3}} - 1$ for each $v \in V(G)$.

Furthermore, one such partition can be constructed in $O(2n^3 \log_2(2n^3))$ -time.

Theorem 1.4. *For all $0 < \varepsilon < 1$, there exists n_0 and η with $0 < \frac{1}{n_0} \ll \eta \ll \varepsilon$ such that the following statement holds. Let G be a near star-multigraph on $2n \geq n_0$ vertices with multi-center x , and let $U = \{v \in V(G) : \Delta(G) - d_G(v) \geq \eta n\}$. Suppose that G satisfies one of the following five conditions:*

- (a) *G is a star-multigraph and is regular with $\delta(G) \geq \delta(G - x) \geq (1 + \frac{\varepsilon}{2})n$ and $\mu(x) < \eta n$.*
- (b) *G is regular with $\delta(G) \geq (1 + \varepsilon)n$, and there exist distinct $y, z \in V(G) \setminus \{x\}$ with the following properties:*
 - (i) $d_G^s(x) \geq 2$ and there are at most $n^{\frac{1}{2}}$ neighbors w of x satisfying $e_G(x, w) \geq \eta n$;
 - (ii) y and z are the only two possible vertices in $G - x$ with $e_{G-x}(y, z) \geq 2$ but $\mu(G - x) < n^{\frac{1}{2}}$; and
 - (iii) $d_G^s(w) \geq (1 + \varepsilon)n$ for all $w \in V(G) \setminus \{x\}$.
- (c) *G is a star-multigraph and is regular with $\delta(G) \geq (1 + \varepsilon)n$, and there exists $y \in V(G) \setminus \{x\}$ such that*
 - (i) $2 \leq d^s(x) < n^{\frac{1}{2}}$;
 - (ii) $d_G^s(y) \geq 2\varepsilon n$; and
 - (iii) $d_G^s(w) \geq (1 + \varepsilon)n$ for all $w \in V(G) \setminus \{x, y\}$.
- (d) *G is a star-multigraph and G has two distinct vertices y, z such that*
 - (i) $\mu(x) < \eta n$;
 - (ii) $d_G(y) = d_G(z) = \delta(G) \geq (\frac{1}{2} + \frac{3\varepsilon}{2})n$ and $\min\{d_G^s(y), d_G^s(z)\} > \Delta(G) - \frac{1}{2}(1 - 0.9\varepsilon)n$; and
 - (iii) $d_G(w) = \Delta(G) \geq (1 + \varepsilon)n$ and $d_G^s(w) \geq (1 + \varepsilon)n - \mu(x) \geq (1 + 0.9\varepsilon)n$ for all $w \in V(G) \setminus \{y, z\}$.
- (e) *G is a star-multigraph, $|U| \geq \eta n$, $\delta(G) \geq \delta(G - x) \geq (1 + \varepsilon)n$, and $\mu(x) \leq \frac{2}{\eta}$.*

Then $\chi'(G) = \Delta(G)$. Furthermore, there is a polynomial time algorithm that finds an optimal coloring.

Proof. We choose n_0 and η so that $0 < \frac{1}{n_0} \ll \eta \ll \varepsilon$.

Let $x_1 = x$, and $y_1 \in V(G) \setminus \{x\}$. Define

$$N^b(x) = \begin{cases} \emptyset & \text{Conditions (a), (d), and (e);} \\ \{v \in V(G) \setminus \{y_1\} : e_G(x, v) \geq \eta n\} & \text{Condition (b);} \\ N_G(x) \setminus \{y_1\} & \text{Condition (c).} \end{cases}$$

We then define a set N according to the different conditions as follows:

Condition (a): let $N = \{x_1, y_1\}$.

Condition (b): let $N^b(x) = \{x_2, \dots, x_t\}$. By the assumption of Condition (b), we have $t - 1 < n^{\frac{1}{2}}$. We take distinct $y_2, \dots, y_t \in V(G) \setminus \{x_1, y_1, x_2, \dots, x_t\}$ and define $N = \{x_1, y_1, \dots, x_t, y_t\}$.

Condition (c): let $N_G(x) \setminus \{y_1\} = \{x_2, \dots, x_t\}$. By the assumption of Condition (c), we have $t - 1 < n^{\frac{1}{2}}$. We take distinct $y_2, \dots, y_t \in V(G) \setminus \{x_1, y_1, x_2, \dots, x_t\}$ and define $N = \{x_1, y_1, \dots, x_t, y_t\}$.

Condition (d): let $N = \{x_1, y_1, x_2, y_2\}$, where $x_2 = y$ and $y_2 = z$.

Condition (e): take $2\lfloor(2n - |V_\Delta|)/2\rfloor$ vertices from $V(G) \setminus V_\Delta$ and name them as $x_2, y_2, \dots, x_t, y_t$, where $t - 1 := \lfloor(2n - |V_\Delta|)/2\rfloor$ and we assume that the first $\lfloor|U|/2\rfloor$ pairs of vertices x_i, y_i are all from U . Let $N = \{x_1, y_1, \dots, x_t, y_t\}$.

We say that y_i and x_i from the set N are *partners* of each other.

Applying Lemma 4.1 on the underlying graph of G and N , we obtain a partition $\{A, B\}$ of $V(G)$ satisfying the following properties in the underlying graph of G : $|A| = |B|$; $|A \cap \{x_i, y_i\}| = 1$ for each $i \in [1, t]$; and $|d_A(v) - d_B(v)| \leq n^{\frac{2}{3}} - 1$ for each $v \in V(G)$.

For the pair (x_1, y_1) , where recall $x_1 = x$, by switching the two vertices from their current partitions and renaming the resulting partitions if necessary, we may assume that $x \in A$ and $d_B(x) \geq d_A(x)$. When G is in Conditions (b) or (c), if $N^b(x) \cap A \neq \emptyset$, we move vertices in $N^b(x) \cap A$ from A to B and move the partners of vertices in $N^b(x) \cap A$ from B to A . Still denote by A and B the resulting partition of $V(G)$. Then we still have $d_B(x) \geq d_A(x)$ as $e_G(x, x_i) \geq e_G(x, y_i)$ for any $x_i \in N^b(x)$. Considering now the partition $\{A, B\}$ in the multigraph G , it has the properties below, where ‘‘Pa’’ indicates the property when G is in Condition (a), and similar meaning for ‘‘Pb’’ to ‘‘Pe.’’

Pa

$$|d_A(v) - d_B(v)| \leq n^{\frac{2}{3}} + e_G(x, v) \quad \text{for each } v \in V(G) \setminus \{x\}.$$

Pb

$$\begin{aligned} N^b(x) &\subseteq B; \\ |d_A(v) - d_B(v)| &\leq n^{\frac{2}{3}} + 2n^{\frac{1}{2}} + n^{\frac{1}{2}} + e_G(x, v) \quad \text{for each } v \in V(G) \setminus \{x\}, \end{aligned}$$

where the term “ $n^{\frac{1}{2}}$ ” is to take account the the possible number of multiple edges joining x and z .

Pc

$$\begin{aligned} N_G(x) &\subseteq B; \\ |d_A(v) - d_B(v)| &\leq n^{\frac{2}{3}} + 2n^{\frac{1}{2}} + e_G(x, v) \quad \text{for each } v \in V(G) \setminus \{x\}. \end{aligned}$$

Pd

$$\begin{aligned} |\{y, z\} \cap A| &= |\{y, z\} \cap B| = 1; \\ |d_A(v) - d_B(v)| &\leq n^{\frac{2}{3}} + e_G(x, v) \quad \text{for each } v \in V(G) \setminus \{x\}. \end{aligned}$$

Pe

$$\begin{aligned} |U \cap A| &\geq \lfloor \frac{|U|}{2} \rfloor, \quad |U \cap B| \geq \lfloor \frac{|U|}{2} \rfloor; \\ |d_A(v) - d_B(v)| &\leq n^{\frac{2}{3}} + e_G(x, v) \quad \text{for each } v \in V(G) \setminus \{x\}. \end{aligned}$$

Let

$$G_A = G[A], \quad G_B = G[B], \quad \text{and} \quad H = G[A, B].$$

Define $G_{A,B}$ to be the union of $G[A]$, $G[B]$ together with $\lfloor (d_B(x) - d_A(x))/2 \rfloor$ edges incident with x from $E(H)$. When G is in Conditions (b) or (c), we take at most $\lceil e_G(x, u)/2 \rceil$ edges between x and each $u \in B$. In particular, when G is in Conditions (c), as $N_G(x) \subseteq B$, we take either $\lfloor e_G(x, u)/2 \rfloor$ or $\lceil e_G(x, u)/2 \rceil$ edges between x and each $u \in N_G(x)$. By this definition of $G_{A,B}$ and the Properties Pa to Pe above, we have

$$\begin{aligned} d_{G_{A,B}}(x) &= \lfloor d_G(x)/2 \rfloor, \\ d_{G_{A,B}}(v) &\geq \frac{1}{2} \left(d_G(v) - e_G(v, x) - n^{\frac{2}{3}} - 2n^{\frac{1}{2}} - n^{\frac{1}{2}} \right) + e_G(v, x) \quad \text{for } v \in A \setminus \{x\}, \quad (1) \\ d_{G_{A,B}}(v) &\geq \frac{1}{2} \left(d_G(v) - e_G(v, x) - n^{\frac{2}{3}} - 2n^{\frac{1}{2}} - n^{\frac{1}{2}} \right) \quad \text{for } v \in B. \end{aligned}$$

For vertex $v \in V(G) \setminus (N_b(x) \cup \{x\})$, as $e_G(x, v) < \eta n$ when G is in Conditions (a) to (d) and $e_G(x, v) \leq \frac{2}{\eta}$ when G is in Condition (e), we will use the following bound:

$$d_{G_{A,B}}(v) \geq \begin{cases} \frac{1}{2} (d_G(v) - 1.2\eta n) & \text{when } G \text{ is in Conditions (a) to (d);} \\ \frac{1}{2} (d_G(v) - 2n^{\frac{2}{3}}) & \text{when } G \text{ is in Condition (e).} \end{cases} \quad (2)$$

For $v \in N^b(x)$, we have

$$d_{G_{A,B}}(v) \geq \frac{1}{2} \left(d_G^s(v) - 2n^{\frac{2}{3}} \right). \quad (3)$$

Furthermore, when G is in Conditions (a) to (d), by the assumptions on G and the partition $\{A, B\}$, we have $\sum_{v \in A} d_G(v) = \sum_{v \in B} d_G(v)$ and so $e(G_A) = e(G_B)$.

To prove the theorem, we will construct an edge coloring of G using $\Delta(G)$ colors. We provide below an overview of the steps. At the start of the process, $E(G)$ is assumed to be uncolored, and throughout the process, the partial edge coloring of G is always denoted by φ , which is updated step by step.

Step 1 Define $S = \{v \in V(G) : \Delta(G) - d_G(v) \geq 7n^{\frac{2}{3}}\}$. Let $k = \Delta(G_{A,B}) + \lfloor n^{\frac{1}{2}} \rfloor$. Since G is a near star-multigraph with $\mu(G - x) < n^{\frac{1}{2}}$, by Theorem 2.14, we find an edge k -coloring φ of $G_{A,B}$. If there exist distinct $u, v \in S \cap A$ or distinct $u, v \in S \cap B$ such that $\overline{\varphi}(u) \cap \overline{\varphi}(v) \neq \emptyset$, we add an edge joining u and v and color the new edge by a color in $\overline{\varphi}(u) \cap \overline{\varphi}(v)$. The edge coloring φ is updated and we still call it φ . We iterate this process of adding and coloring edges and call the multigraphs resulting from G_A, G_B and $G_{A,B}$, respectively, as G_A^*, G_B^* , and $G_{A,B}^*$, and call G^* the union of $G_{A,B}^*$ and H . It can be verified that $\Delta(G^*) = \Delta(G)$. We will modify the current edge coloring, which is still named φ , such that the following properties are satisfied:

S1.1 Let $i, j \in [1, k]$ and $C \in \{A, B\}$. When G is in Conditions (a) to (d), we have

$$|\overline{\varphi}_A^{-1}(i)| = |\overline{\varphi}_B^{-1}(i)| \quad \text{and} \quad \left| |\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)| \right| \leq 2$$

When G is in Condition (e), we have

$$\left| |\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)| \right| \leq 2.$$

S1.2 Let $i \in [1, k]$. When G is in Conditions (a) to (d), we have

$$|\overline{\varphi}_A^{-1}(i)| < 3\eta n \quad \text{and} \quad |\overline{\varphi}_B^{-1}(i)| < 3\eta n.$$

When G is in Condition (e), we have

$$|\overline{\varphi}_A^{-1}(i)| < 7n^{\frac{2}{3}} \quad \text{and} \quad |\overline{\varphi}_B^{-1}(i)| < 7n^{\frac{2}{3}}.$$

Step 2 Modify the partial edge-coloring of G^* obtained in Step 1 by exchanging alternating paths. Upon the completion of Step 2, each of the k color class will be a 1-factor of G^* . In the process of Step 2, a few edges of H will be colored and a few edges of $G_A^* \cup G_B^*$ will be uncolored. Denote by R_A and R_B , respectively, the submultigraphs of G_A^* and G_B^* consisting of the uncolored edges. The two multigraphs R_A and R_B

will initially be empty, but one, two, three, or four edges will be added to at least one of them when each time we exchange an alternating path. The calculations involved when G is in Condition (e) is essentially different than the other cases, so we need define parameters correspondingly as follows: let

$$s = \begin{cases} 3\eta n^2 & \text{when } G \text{ is in Conditions (a) to (d),} \\ 7n^{\frac{5}{3}} & \text{when } G \text{ is in Condition (e),} \end{cases}$$

and

$$r = \begin{cases} \eta^{\frac{1}{2}}n & \text{when } G \text{ is in Conditions (a) to (d),} \\ n^{\frac{5}{6}} & \text{when } G \text{ is in Condition (e).} \end{cases}$$

The conditions below will be satisfied at the completion of this step:

S2.1 The number of uncolored edges in each of G_A^* and G_B^* is less than $4s$. When G is in Conditions (a) to (d), G_A^* and G_B^* have the same number of uncolored edges; and when G is in Condition (e), the number of uncolored edges in G_B^* is greater than or equal to the number of uncolored edges in G_A^* (this follows if we assume by symmetry that $e(G_A^*) \leq e(G_B^*)$).

S2.2 $\Delta(R_A)$ and $\Delta(R_B)$ are less than r .

S2.3 For $C \in \{A, B\}$, let $S_C = \{u \in S \cap C : d_{G_{A,B}^*}(u) \leq k - 2n^{\frac{2}{3}}\}$. We require every vertex $u \in V(G^*) \setminus (S_A \cup S_B \cup N^b(x) \cup \{x\})$ is incident in G^* with fewer than $|\bar{\varphi}(u)| + r$ colored edges of H ; and every vertex $u \in S_A \cup S_B \cup N^b(x) \cup \{x\}$ is incident in G^* with at most $|\bar{\varphi}(u)|$ colored edges of H .

Step 3 We will edge color each of R_A and R_B and color a few uncolored edges of H using another ℓ colors, where $\ell = 2\lceil r \rceil$. The goal is to ensure that each of these ℓ new color classes obtained at the completion of Step 3 presents at all vertices from $V(G^*) \setminus U$ while preserve the k 1-factors already obtained through Steps 1 and 2.

Step 4 At the start of Step 4, all of the uncolored edges of G^* belong to H . Denote by R the subgraph of G^* consisting of the uncolored edges. We will show that $\Delta(R) = \Delta(G^*) - k - \ell$. This submultigraph is bipartite, so we can color its edges using $\Delta(G^*) - k - \ell$ colors by Theorem 2.9.

When Step 4 is completed, we obtain an edge coloring of G^* using exactly $\Delta(G^*)$ colors. We now give the details of each step, and for notation that was already defined in the outline above, we will use them directly.

Step 1: Coloring $G_{A,B}$

Recall $S = \{v \in V(G) : \Delta(G) - d_G(v) \geq 7n^{\frac{2}{3}}\}$. Note that when G is in Conditions (a) to (c), $S = \emptyset$; when G is in Condition (d), $S \subseteq \{y, z\}$; and when G is in Condition (e), $U \subseteq S$. When G is in Condition (d), we have $|\{y, z\} \cap A| = |\{y, z\} \cap B| = 1$ by Property Pd. Thus, when G is in Conditions (a) to (d), we have $G_A^* = G_A$, $G_B^* = G_B$, $G_{A,B}^* = G_{A,B}$, and $G^* = G$.

Following the operations described in the outline of Step 1, for the current edge coloring φ of $G_{A,B}^*$, the following statement holds: $\overline{\varphi}(u) \cap \overline{\varphi}(v) = \emptyset$ for any two distinct $u, v \in S \cap A$ or any two distinct $u, v \in S \cap B$. Therefore, for any $C \in \{A, B\}$, we have

$$\sum_{u \in C \cap S} |\overline{\varphi}(u)| = \sum_{u \in C \cap S} (k - d_{G_{A,B}^*}(u)) \leq k. \quad (4)$$

Note that $k \leq \Delta(G_{A,B}) + n^{\frac{1}{2}}$, and by the partition $\{A, B\}$ of $V(G)$, we have $\Delta(G_{A,B}) \leq \max_{v \in V(G)} \{\frac{1}{2}(\Delta(G) - e_G(x, v) + n^{\frac{2}{3}} + 2n^{\frac{1}{2}} + n^{\frac{1}{2}}) + e^*\}$, where $e^* = e_G(x, v) < \eta n$ if $v \in A$ and $e^* = \lceil e_G(x, v)/2 \rceil$ if $v \in B$. Thus when G is in Conditions (a) to (d), we have

$$k \leq \frac{1}{2}(\Delta(G) + n^{\frac{2}{3}} + 2n^{\frac{1}{2}} + n^{\frac{1}{2}}) + \frac{1}{2}\eta n + n^{\frac{1}{2}} < \frac{1}{2}\Delta(G) + 0.6\eta n. \quad (5)$$

When G is in Condition (e), as $\mu(G) < \frac{2}{\eta}$, we have

$$k \leq \frac{1}{2}(\Delta(G) + n^{\frac{2}{3}} + 2n^{\frac{1}{2}} + n^{\frac{1}{2}}) + \frac{1}{\eta} + n^{\frac{1}{2}} < \frac{1}{2}\Delta(G) + n^{\frac{2}{3}}. \quad (6)$$

We will in the rest of the proof show that $\chi'(G^*) = \Delta(G^*)$, this is because G is a subgraph of G^* and $\Delta(G^*) = \Delta(G)$. To see $\Delta(G^*) = \Delta(G)$, note that the edge additions in Step 1 happen only when G is in Condition (e). Under Condition (e), for any $u \in S \cap A$, we have

$$d_{G^*}(u) \leq k + e_G(u, B) \leq \frac{1}{2}\Delta(G) + n^{\frac{2}{3}} + \frac{1}{2}(\Delta(G) - 7n^{\frac{2}{3}} + n^{\frac{2}{3}} + \frac{2}{\eta}) < \Delta(G).$$

Similarly, we have $d_{G^*}(u) \leq \Delta(G)$ for any $u \in S \cap B$. In particular, if $u \in U$, as $\Delta(G) - d_G(u) \geq \eta n$, we have

$$d_{G^*}(u) \leq \frac{1}{2}\Delta(G) + n^{\frac{2}{3}} + \frac{1}{2}(\Delta(G) - \eta n + n^{\frac{2}{3}} + \frac{2}{\eta}) < \Delta(G) - \frac{\eta n}{3}. \quad (7)$$

When G is in Conditions (a) to (d), since $e(G_A^*) = e(G_A) = e(G_B) = e(G_B^*)$, by Lemma 2.11, we modify φ into an edge k -coloring of $G_{A,B}^*$, and still call φ the resulting coloring, such that

$$|\overline{\varphi}_A^{-1}(i)| = |\overline{\varphi}_B^{-1}(i)| \quad \text{and} \quad ||\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)|| \leq 2 \quad \text{for } i, j \in [1, k] \text{ and } C \in \{A, B\}.$$

When G is in Condition (e), by Lemma 2.12, we modify φ so that

$$||\overline{\varphi}_C^{-1}(i)| - |\overline{\varphi}_C^{-1}(j)|| \leq 2 \quad \text{for } i, j \in [1, k] \text{ and } C \in \{A, B\}.$$

Note that under the modified coloring φ , it is possible that $\overline{\varphi}(u) \cap \overline{\varphi}(v) \neq \emptyset$ for some distinct $u, v \in S \cap A$ or distinct $u, v \in S \cap B$. However the inequalities in (4) still hold. For $C \in \{A, B\}$, define

$$S_C = \{u \in S \cap C : d_{G_{A,B}^*}(u) \leq k - 2n^{\frac{2}{3}}\}.$$

Next, we verify that every color $i \in [1, k]$ is missing at a small number of vertices. We first provide an upper bound on $|\overline{\varphi}(u)|$ for each $u \in V(G^*)$. By the definition of $G_{A,B}$, (??), (3), (5), and (6), as $|\overline{\varphi}(u)| = k - d_{G_{A,B}^*}(u)$, we have

$$|\overline{\varphi}(u)| \leq \begin{cases} \frac{1}{2}\Delta(G) + 0.6\eta n - \lfloor \Delta(G)/2 \rfloor < \eta n & u = x, \text{ Conditions (a) to (d);} \\ \frac{1}{2}\Delta(G) + n^{\frac{2}{3}} - \frac{1}{2}(1 + \varepsilon)n < \frac{1}{2}(1 - \varepsilon)n + n^{\frac{2}{3}} & u = x, \text{ Condition (e);} \\ \frac{1}{2}\Delta(G) + 0.6\eta n - \frac{1}{2}(\Delta(G) - 7n^{\frac{2}{3}} - 1.2\eta n) < 2\eta n & u \in V(G) \setminus (S \cup N^b(x) \cup \{x\}), \\ & \text{and Conditions (a) to (d);} \\ \frac{1}{2}\Delta(G) + n^{\frac{2}{3}} - \frac{1}{2}(\Delta(G) - 7n^{\frac{2}{3}} - 2n^{\frac{2}{3}}) < 6n^{\frac{2}{3}} & u \in V(G) \setminus S, \text{ Condition (e);} \\ \frac{1}{2}\Delta(G) + 0.6\eta n - \left(\frac{1}{2}(1 + \varepsilon)n - 2n^{\frac{2}{3}}\right) < \left(\frac{1}{2} - \frac{1}{3}\varepsilon\right)n & u \in N^b(x), \text{ and} \\ & \text{Conditions (b) and (c);} \\ \frac{1}{2}\Delta(G) + 0.6\eta n - \left(\frac{1}{2}(\Delta(G) - e_G(x, u) - 2n^{\frac{2}{3}}) + \lfloor e_G(x, u)/2 \rfloor\right) < \eta n & u = y, \text{ Condition (c);} \\ \frac{1}{2}\Delta(G) + 0.6\eta n - \frac{1}{2}(d_G^s(u) - 1.2\eta n) = \frac{1}{2}\Delta(G) + 1.2\eta n - \frac{1}{2}(\Delta(G) - \frac{1}{2}(1 - 0.9\varepsilon)n) < \left(\frac{1}{4} - \frac{1}{5}\varepsilon\right)n & u \in \{y, z\}, \text{ Condition (d);} \\ \frac{1}{2}\Delta(G) + n^{\frac{2}{3}} - \frac{1}{2}((1 + \varepsilon)n - 2n^{\frac{2}{3}}) < \left(\frac{1}{2} - \frac{1}{3}\varepsilon\right)n & u \in S_A \cup S_B, \text{ Condition (e);} \\ 2n^{\frac{2}{3}} & u \in S \setminus (S_A \cup S_B) \text{ and} \\ & \text{Condition (e).} \end{cases}$$

Since $|\overline{\varphi}(u)| < 2n$ for each $u \in N^b(x)$ and $|N^b(x)| < n^{\frac{1}{2}}$, the inequalities above together with the fact in (4) give the following inequalities for any $C \in \{A, B\}$:

$$\sum_{u \in C} |\overline{\varphi}(u)| \leq 2nn^{\frac{1}{2}} + 2\eta n^2 + k < 3\eta n^2 - 2n \quad G \text{ is in Conditions (a) to (d);} \quad (8)$$

$$\sum_{u \in C} |\overline{\varphi}(u)| \leq 2nn^{\frac{1}{2}} + 6n^{\frac{5}{3}} + k < 7n^{\frac{5}{3}} - 2n \quad G \text{ is in Condition (e).} \quad (9)$$

By SI.1, the Statement S1.2 stated in the outline of Step 1 holds.

Step 2: Extending existing color classes into 1-factors

We will extend each of the k color classes obtained in Step 1 into a 1-factor of G^* by exchanging alternating paths. Each alternating path starts and also ends at an uncolored edge of H and is alternating between colored edges of $G_A^* \cup G_B^*$ and uncolored edges of H . Thus during the procedure of Step 2, we will uncolor some of the edges of $G_A^* \cup G_B^*$, and will color some of the edges of H . Recall that R_A and R_B are the submultigraphs of G_A^* and G_B^* consisting of the uncolored edges, which are initially empty.

To ensure Condition S2.2 is satisfied, we say that an edge $e = uv \in E(G_A^* \cup G_B^*)$ is *good* if $e \notin E(R_A \cup R_B)$ and the degree of u and v in both R_A and R_B is less than r (actually, note that when $uv \in E(G_A^*)$, then the degree of u and v is zero in R_B and vice versa). Thus a good edge can be added to R_A or R_B without violating S2.2.

By S1.1, for each color $i \in [1, k]$, we pair up a vertex from $\overline{\varphi}_A^{-1}(i)$ and a vertex from $\overline{\varphi}_B^{-1}(i)$, and then pair up the remaining unpaired vertices from $\overline{\varphi}_A^{-1}(i)$ or $\overline{\varphi}_B^{-1}(i)$ as $|\overline{\varphi}_A^{-1}(i)| - |\overline{\varphi}_B^{-1}(i)|$ is even by the Parity Lemma. Each of those pairs is called a *missing-common-color pair* or *MCC-pair* in short with respect to the color i . In particular, when G is in Conditions (a) to (d), every vertex from $\overline{\varphi}_A^{-1}(i)$ is paired up with a vertex from $\overline{\varphi}_B^{-1}(i)$.

For every MCC-pair (a, b) with respect to some color $i \in [1, k]$, we will exchange an alternating path P from a to b with at most 13 edges, where, if exist, the first, third, fifth, seventh, ninth, eleventh, and thirteenth edges are uncolored and the second, fourth, sixth, eighth, tenth, and twelfth edges are good edges colored by i . After P is exchanged, a and b will be incident with edges colored by i , and at most four good edges will be added to each of R_A and R_B . With this information at hand, before demonstrating the existence of such paths, we show that Conditions S2.1, S2.2 and S2.3 can be guaranteed at the end of Step 2.

After the completion of Step 1, by (8) and (9), the total number of missing colors from vertices in A or from vertices in B is at most s , which is $3\eta n^2$ when G is in Conditions (a) to (d), and is $7n^{\frac{5}{3}}$ when G is in Condition (e). Thus there are at most s MCC-pairs. For each MCC-pair (a, b) with $a, b \in V(G^*)$, at most four edges will be added to each of R_A and R_B when we exchange an alternating path from a to b . Thus there will always be fewer than $4s$ edges in each of R_A and R_B . At the completion of Step 2, each of the k color classes is

a 1-factor of G^* so each of G_A^* and G_B^* have the same number of colored edges. When G is in Conditions (a) to (d), since $e(G_A^*) = e(G_A) = e(G_B) = e(G_B^*)$, we have $e(R_A) = e(R_B)$. When G is in Condition (e), suppose, without loss of generality, that $e(G_B^*) \geq e(G_A^*)$, then we get $e(R_B) \geq e(R_A)$. Thus Condition S2.1 will be satisfied at the end of Step 2. And as we only ever add good edges to R_A and R_B , Condition S2.2 will hold automatically. We now show that Condition S2.3 will also be satisfied. Recall that

$$S_A = \{u \in S \cap A : d_{G_{A,B}^*}(u) \leq k - 2n^{\frac{2}{3}}\} \quad \text{and} \quad S_B = \{u \in S \cap B : d_{G_{A,B}^*}(u) \leq k - 2n^{\frac{2}{3}}\}.$$

Since $\sum_{u \in C \cap S} \text{def}_{G_{A,B}^*}(u) \leq k \leq \frac{1}{2}\Delta(G) + 0.6\eta n < 2n$ for any $C \in \{A, B\}$, it follows that

$$|S_A| < n^{1/3} \quad \text{and} \quad |S_B| < n^{1/3}.$$

In the process of Step 2, the number of newly colored edges of H that are incident with a vertex $u \in V(G^*) \setminus (S_A \cup S_B \cup N^b(x) \cup \{x\})$ will equal the number of alternating paths containing u that have been exchanged. The number of such alternating paths of which u is the first vertex will equal the number of colors that missed at u at the end of Step 1. The number of alternating paths in which u is not the first vertex will equal the degree of u in $R_A \cup R_B$, and so will be less than r . Hence the number of colored edges of H that are incident with u will be less than $|\overline{\varphi}(u)| + r$. The number of newly colored edges of H that are incident with a vertex $u \in S_A \cup S_B \cup N^b(x) \cup \{x\}$ will equal the number of colors that missed at u at the end of Step 1. Thus Condition S2.3 will be satisfied.

As the vertex x may have a quite small simple degree, we deal with colors in $\overline{\varphi}(x)$ first. By the upper bound on $|\overline{\varphi}(x)|$ given in Step 1, we have $|\overline{\varphi}(x)| < \eta n$ when G is in Conditions (a) to (d), and $|\overline{\varphi}(x)| \leq \frac{1}{2}(1 - \varepsilon)n + n^{\frac{2}{3}}$ when G is in Condition (e). When G is in Condition (e), as $\mu(G) \leq \frac{2}{\eta}$ and $e_G(x, B) - e_{G_{A,B}^*}(x, B) \geq \frac{\eta}{2}$, we know that x is adjacent in G with at least $\frac{n/2}{2/\eta} \geq \frac{\eta}{4}n$ vertices from B that each joins x with an uncolored edge of H .

For each $i \in \overline{\varphi}(x)$, if there is a vertex $w \in N_G(x, B)$ with xw uncolored and $i \in \overline{\varphi}(w)$, we simply color xw by the color i . Thus we assume that for every $w \in N_G(x, B)$ with xw uncolored, it holds that $i \notin \overline{\varphi}(w)$. We choose $w \in N_G(x, B)$ with xw uncolored and with $d_{R_B}(w)$ smallest. Let $w^* \in N_G(w, B)$ such that ww^* is colored by i (the vertex w^* exists as we only added colored edges between x and $N_G(x, B)$ so far and $i \in \varphi(w)$). Then we color xw by i and uncolor ww^* . (We say w^* is *originated* from x with respect to the color i). This procedure guarantees all edges of R_B to be good, as when G is in Conditions (a) to (d), we have $|\overline{\varphi}(x)| < \eta n < \eta^{\frac{1}{2}}n$; and when G is in Condition (e), although $|\overline{\varphi}(x)| \leq \frac{1}{2}(1 - \varepsilon)n + n^{\frac{2}{3}}$, we have at least $\frac{1}{4}\eta n - |\overline{\varphi}(x)|/n^{\frac{5}{6}} > 1$ choices each time to pick a desired vertex $w \in N_G(x, B)$. After this procedure, all the k colors present at x . Thus x is not contained in any MCC-pair with respect to the current edge coloring.

Let $S_A^* = S_A \cup \{x\}$ and $S_B^* = S_B \cup N^b(x)$. We now show below the existence of alternating paths for the current MCC-pairs. For a given color $i \in [1, k]$, and vertices $a \in A$ and $b \in B$, let $N_B(a)$ be the set of vertices in B that are joined with a by an uncolored edge and are incident with a good edge colored i such that the good edge is not incident

with any vertex of S_B^* , and let $N_A(b)$ be the set of vertices in A that are joined with b by an uncolored edge and are incident with a good edge colored i such that the good edge is not incident with any vertex of S_A^* . In order to estimate the sizes of $N_A(b)$ and $N_B(b)$, we show that A and B contain only a few vertices that either miss the color i or are incident with a non-good edge colored i . By S2.1, there are at most $4s$ edges in R_B , so there are fewer than $\frac{8s}{r}$ vertices of degree at least r in R_B . Each non-good edge is incident with one or two vertices of R_B through the color i , so there are fewer than $\frac{16s}{r}$ vertices in B that are incident with a non-good edge colored i . Furthermore, there are at most $2|S_B^*| \leq 3n^{\frac{1}{3}}$ vertices in B that are either contained in S_B^* or adjacent to a vertex from S_B^* through an edge with color i . Finally, there are fewer than $\max\{3\eta n, 7n^{\frac{2}{3}}\} = 3\eta n$ vertices in B that are missed by the color i by S1.2. So the number of vertices in B that are not incident with a good edge colored i such that the good edge is not incident with any vertex from S_B^* is less than

$$\frac{16s}{r} + 3n^{\frac{1}{3}} + 3\eta n < 49\eta^{\frac{1}{2}}n.$$

Similarly, the number of vertices in A that are not incident with a good edge colored i such that the good edge is not incident with any vertex from S_A^* is less than $49\eta^{\frac{1}{2}}n$. Note that, by the partition $\{A, B\}$ of $V(G)$, we have

$$\begin{aligned} |N_G(v, B)| &\geq \frac{1}{2}(d_G^s(v) - n^{\frac{2}{3}} - 2n^{\frac{1}{2}} - n^{\frac{1}{2}}) \quad \text{for } v \in A \setminus \{x\}; \\ &> \frac{1}{2}(d_G^s(v) - 2n^{\frac{2}{3}}) \\ & \\ |N_G(v, A)| &\geq \frac{1}{2}(d_G^s(v) - n^{\frac{2}{3}} - 2n^{\frac{1}{2}} - n^{\frac{1}{2}}) \quad \text{for } v \in B. \\ &> \frac{1}{2}(d_G^s(v) - 2n^{\frac{2}{3}}) \end{aligned} \tag{10}$$

By S2.3 and the upper bound on $|\overline{\varphi}(u)|$ for any $u \in V(G^*)$ provided in Step 1, we have the following lower bounds on $|N_B(a)|$ and $|N_A(b)|$:

When $\{a, b\} \cap (S_A^* \cup S_B^*) = \emptyset$,

$$|N_A(b)|, |N_B(a)| \geq \frac{1}{2} \left((1 + \frac{1}{2}\varepsilon)n - 2n^{\frac{2}{3}} \right) - (2\eta n + \eta^{\frac{1}{2}}n) - 49\eta^{\frac{1}{2}}n > \left(\frac{1}{2} + \frac{1}{5}\varepsilon \right)n. \tag{11}$$

When G is in Conditions (b) or (c) and $\{a, b\} \cap N^b(x) \neq \emptyset$, we have

$$|N_A(b)|, |N_B(a)| \geq \frac{1}{2} \left((1 + \varepsilon)n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{2} - \frac{1}{3}\varepsilon \right)n - 49\eta^{\frac{1}{2}}n > \frac{1}{2}\varepsilon n. \tag{12}$$

When G is in Condition (c) and $b = y$, we have

$$|N_A(b)| \geq \frac{1}{2} \left(2\varepsilon n - 2n^{\frac{2}{3}} \right) - \eta n - 49\eta^{\frac{1}{2}}n > \frac{3}{4}\varepsilon n. \tag{13}$$

When G is in Condition (d) and $\{a, b\} \cap \{y, z\} \neq \emptyset$, we have

$$|N_A(b)|, |N_B(a)| \geq \frac{1}{2} \left(\left(\frac{1}{2} + \frac{3\varepsilon}{2} \right)n - \eta n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{4} - \frac{1}{5}\varepsilon \right)n - 49\eta^{\frac{1}{2}}n > \frac{3}{4}\varepsilon n. \tag{14}$$

When G is in Condition (e) and $\{a, b\} \cap (S_A^* \cup S_B^*) \neq \emptyset$, we have

$$|N_A(b)|, |N_B(a)| \geq \frac{1}{2} \left((1 + \varepsilon)n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{2} - \frac{1}{3}\varepsilon \right)n - 49\eta^{\frac{1}{2}}n > \frac{1}{2}\varepsilon n. \quad (15)$$

Let $M_B(a)$ be the set of vertices in B that are joined with a vertex in $N_B(a)$ by an edge of color i , and let $M_A(b)$ be the set of vertices in A that are joined with a vertex in $N_A(b)$ by an edge of color i . Note that $(S_A^* \cup S_B^*) \cap (M_A(b) \cup M_B(a)) = \emptyset$ by the choice of $N_A(b)$ and $N_B(a)$. Note also that $|M_B(a)| = |N_B(a)|$ but some vertices may be in both. Similarly $|M_A(b)| = |N_A(b)|$.

For a MCC-pair (a, b) , in order to have a unified discussion as in the case that $\{a, b\} \cap (S_A^* \cup S_B^*) = \emptyset$, if necessary, by exchanging an alternating path of length 2 from a to another vertex a^* , and exchanging an alternating path from b to another vertex b^* , we will replace the pair (a, b) by (a^*, b^*) such that $\{a^*, b^*\} \cap (S_A^* \cup S_B^*) = \emptyset$. Precisely, we will implement the following operations to vertices in $S_A \cup S_B^*$. For any vertex $a \in S_A$, and for each color $i \in \overline{\varphi}(a)$, we take an edge b_1b_2 with $b_1 \in N_B(a)$ and $b_2 \in M_B(a)$ such that b_1b_2 is colored by i , where the edge b_1b_2 exists by (11)-(15) and the fact that $|M_B(a)| = |N_B(a)|$. Then we exchange the path ab_1b_2 by coloring ab_1 with i and uncoloring the edge b_1b_2 (See Figure 1(a)). After this, the edge ab_1 of H is now colored by i , and the uncolored edge b_1b_2 is added to R_B . We then update the original MCC-pair that contains a with respect to the color i by replacing the vertex a with b_2 . We do this at the vertex a for every color $i \in \overline{\varphi}(a)$ and then repeat the same process for every vertex in S_A . Similarly, for any vertex $b \in S_B^*$, and for each color $i \in \overline{\varphi}(b)$, we take an edge a_1a_2 with $a_1 \in N_A(b)$ and $a_2 \in M_A(b)$ such that a_1a_2 is colored by i , where the edge a_1a_2 exists by (11)-(15) and the fact that $|M_B(b)| = |N_B(b)|$. Then we exchange the path ba_1a_2 by coloring ba_1 with i and uncoloring the edge a_1a_2 . The same, we update the original MCC-pair that contains b with respect to the color i by replacing the vertex b with a_2 .

After the procedure above, we have now three types MCC-pair (u, v) : $u, v \in A$, $u, v \in B$, and A contains exactly one of u and v and B contains the other. However, in either case, $\{u, v\} \cap (S_A^* \cup S_B^*) = \emptyset$. We will exchange alternating path for each of such pairs.

We deal with each of the colors from $[1, k]$ in turn. Let $i \in [1, k]$ be a color. We consider first an MCC-pair (a, a^*) with respect to i such that $a, a^* \in A$. By (11), we have $|M_B(a^*)| > (\frac{1}{2} + \frac{1}{5}\varepsilon)n$. We take an edge $b_1^*b_2^*$ colored by i with $b_1^* \in N_B(a^*)$ and $b_2^* \in M_B(a^*)$. Then again, by (11), we have $|M_B(a)|, |M_A(b_2^*)| > (\frac{1}{2} + \frac{1}{5}\varepsilon)n$. Therefore, as each vertex $c \in M_A(b_2^*)$ satisfies $|N_B(c)| > (\frac{1}{2} + \frac{1}{5}\varepsilon)n$, we have $|N_B(c) \cap M_B(a)| \geq \frac{2}{5}\varepsilon n$. We take $a_2a_2^*$ colored by i with $a_2^* \in N_A(b_2^*)$ and $a_2 \in M_A(b_2^*)$. Then we let $b_2 \in N_B(a_2) \cap M_B(a)$, and let b_1 be the vertex in $N_B(a)$ such that b_1b_2 is colored by i . Now we get the alternating path $P = ab_1b_2a_2a_2^*b_1^*a^*$ (See Figure 1(c)). We exchange P by coloring $ab_1, b_2a_2, a_2^*b_2^*$ and $b_1^*a^*$ with color i and uncoloring the edges $b_1b_2, b_1^*b_2^*$ and $a_2a_2^*$. After the exchange, the color i appears on edges incident with a and a^* , the edges b_1b_2 and $b_1^*b_2^*$ are added to R_B and the edge $a_2a_2^*$ is added to R_A . We added at most one edges to each of R_A and R_B when we updated the original MCC-pair corresponding to (a, a^*) , where a and a^* could be originated

from some vertices $b'_1, b'_2 \in S_B^*$. Also, exactly one of b'_1 and b'_2 could be originated from x with respect to the color i . Thus we added at most three edges to R_A and at most four edges to R_B when we modify φ to have the color i present at both of the vertices in the original MCC-pair corresponding to (a, a^*) . The maximum length of an alternating path combined from the three procedures (dealing missing colors at x , dealing missing colors at vertices from $S_A \cup S_B^*$, and dealing missing colors at vertices from $V(G^*) \setminus (S_A^* \cup S_B^*)$) together is at most $2 + 2 + 2 + 7 = 13$. By symmetry, we can deal with an MCC-pair (b, b^*) with respect to i such that $b, b^* \in B$ similarly as above.

Then we consider an MCC-pair (a, b) with respect to i such that $a \in A$ and $b \in B$. By (11), we have $|M_B(a)|, |M_A(b)| > (\frac{1}{2} + \frac{1}{5}\varepsilon)n$. We choose $a_1 a_2$ with color i such that $a_1 \in N_A(b)$ and $a_2 \in M_A(b)$. Now as $|M_B(a)|, |N_B(a_2)| > (\frac{1}{2} + \frac{1}{5}\varepsilon)n$ by (11), we know that $N_B(a_2) \cap M_B(a) \neq \emptyset$. We choose $b_2 \in N_B(a_2) \cap M_B(a)$ and let $b_1 \in N_B(a)$ such that $b_1 b_2$ is colored by i . Then $P = ab_1 b_2 a_2 a_1 b$ is an alternating path from a to b (See Figure 1(b)). We exchange P by coloring $ab_1, b_2 a_2$ and $a_1 b$ with color i and uncoloring the edges $a_1 a_2$ and $b_1 b_2$. After the exchange, the color i appears on edges incident with a and b , the edge $a_1 a_2$ is added to R_A and the edge $b_1 b_2$ is added to R_B . We added at most one edge to each of R_A and R_B when we updated the original MCC-pair corresponding to (a, b) . Thus we added at most one edge to R_A and at most two edges to R_B when we modify φ to have the color i present at both of the vertices in the original MCC-pair corresponding to (a, b) . By finding such paths for all MCC-pairs with respect to the color i , we can increase the number of edges colored i until the color class is a 1-factor of G^* . By doing this for all colors, we can make each of the k color classes a 1-factor of G^* .

Step 3: Coloring R_A and R_B and extending the new color classes

Each of the color classes for the colors from $[1, k]$ is now a 1-factor of G^* . We now consider the multigraphs R_A and R_B that consist of the uncolored edges of G_A^* and G_B^* . By Condition S2.1, R_A and R_B each has fewer than $4s$ edges, and $\Delta(R_A), \Delta(R_B) < r$. Note that R_A and R_B might contain parallel edges with endvertices in S or between y and z when G is in Condition (d). By Theorem 2.8 and Theorem 2.10, R_A and R_B each has an equalized edge-coloring with exactly $\ell := 2\lfloor r \rfloor$ colors $k + 1, \dots, k + \ell$.

If G is in Conditions (a) to (d), then we have $e(R_A) = e(R_B)$. Under these conditions, by renaming some color classes of R_A if necessary, we can assume that in the edge colorings of R_A and R_B , each color appears on the same number of edges in R_A as it does in R_B . When G is in Condition (e), by our assumption that G_B^* has more edges than G_A^* does, we have $e(R_A) \leq e(R_B)$. In this case, we can assume that in the edge colorings of R_A and R_B , the number of edges with a color $i \in [k + 1, k + \ell]$ in R_B is at least the number of edges with a color $i \in [k + 1, k + \ell]$ in R_A .

There are fewer than $4s$ edges in each of R_A and R_B , and $\ell > r$, so each of the color $i \in [k + 1, k + \ell]$ appears on fewer than $\frac{4s}{r}$ edges in each of R_A and R_B . We will now color some of the edges of H with the ℓ colors from $[k + 1, k + \ell]$ so that each of these color classes




Figure 1: The alternating path P . Dashed lines indicate uncoloured edges, and solid lines indicate edges with color i .

present at vertices from $V(G^*) \setminus U$. We perform the following procedure for each of the ℓ colors in turn.

Given a color i with $i \in [k+1, k+\ell]$, we let A_i and B_i be the sets of vertices in A and B respectively that are incident with edges colored i . Note that $|A_i| \leq |B_i| < 2 \times \frac{4s}{r} = \frac{8s}{r}$, as R_A and R_B each contains fewer than $\frac{4s}{r}$ edges colored i . Note that if G is in Conditions (a) to (d), we have $|A_i| = |B_i|$; and we might have $|B_i| > |A_i|$ when G is in Condition (e). When G is in Condition (e) and $|B_i| > |A_i|$, we let

$$A_i^* \subseteq (U \cap A) \setminus A_i$$

such that $|A_i^*| + |A_i| = |B_i|$, and just let $A_i^* = \emptyset$ otherwise. Note that such A_i^* exists as $|U \cap A| \geq \frac{1}{2}\eta n - 1$ and $|A_i|, |B_i| < \frac{8s}{r} = 56n^{\frac{5}{6}}$. Let H_i be the subgraph of H obtained by deleting the vertex sets $A_i \cup A_i^*$ and B_i and removing all colored edges. We will show next that H_i has a perfect matching and we will color the edges in the matching by the color i .

For each $u \in V(G^*)$, the number of colored edges of H incident with u is specified by

S2.3 and at most ℓ edges incident with u have been colored in Step 3. Also each vertex in G^* has fewer than $\frac{8s}{r} < 24\eta^{\frac{1}{2}}n$ edges that join it with a vertex in A_i or B_i . So by (10), each vertex $u \in V(H_i) \setminus (S_A^* \cup S_B^*)$ satisfies

$$d_{H_i}^s(u) > \frac{1}{2} \left(\left(1 + \frac{1}{2}\varepsilon\right)n - 2n^{\frac{2}{3}} \right) - (2\eta n + \eta^{\frac{1}{2}}n) - \ell - 24\eta^{\frac{1}{2}}n > \left(\frac{1}{2} + \frac{\varepsilon}{5}\right)n.$$

Note that $V(R_A \cup R_B) \cap (S_A^* \cup S_B^*) = \emptyset$. When G is in Conditions (b) or (c), each of the vertex $u \in N^b(x)$, where $u \neq y$ when G is in Condition (c), satisfies

$$d_{H_i}^s(u) > \frac{1}{2} \left((1 + \varepsilon)n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{2} - \frac{1}{3}\varepsilon\right)n - \ell - 24\eta^{\frac{1}{2}}n > \frac{1}{2}\varepsilon n.$$

When G is in Condition (c), the vertex y satisfies

$$d_{H_i}^s(y) > \frac{1}{2} \left(2\varepsilon n - 2n^{\frac{2}{3}} \right) - \eta n - \ell - 24\eta^{\frac{1}{2}}n > \frac{\varepsilon}{4}\varepsilon n.$$

When G is in Condition (d), each $u \in \{y, z\}$ satisfies

$$d_{H_i}^s(u) > \frac{1}{2} \left(\left(\frac{1}{2} + \frac{3\varepsilon}{2}\right)n - \eta n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{4} - \frac{1}{5}\varepsilon\right)n - \ell - 24\eta^{\frac{1}{2}}n > \frac{3}{4}\varepsilon n.$$

When G is in Condition (e), each $u \in S_A \cup S_B$ satisfies

$$d_{H_i}^s(u) > \frac{1}{2} \left((1 + \varepsilon)n - 2n^{\frac{2}{3}} \right) - \left(\frac{1}{2} - \frac{1}{3}\varepsilon\right)n - \ell - 24\eta^{\frac{1}{2}}n > \frac{1}{2}\varepsilon n.$$

For the vertex x , we have

$$d_{H_i}(x) \geq \delta(G) - k - \ell - 24\eta^{\frac{1}{2}}n \geq (1 + \varepsilon)n - \left(\frac{1}{2}\Delta(G) + 0.6\eta n\right) - \ell - 24\eta^{\frac{1}{2}}n > \frac{1}{2}\varepsilon n.$$

Thus $\delta(H_i) \geq 1$ and $\delta(H_i - x) \geq \frac{1}{2}\varepsilon n - 1$ in any case and H_i has at most $|S_A^* \cup S_B^*| \leq 3n^{1/3} < \frac{1}{2}\varepsilon n - 1$ vertices of simple degree less than $\frac{1}{2}n + 1$. So H_i has a 1-factor F by Lemma 2.6. If we color the edges of F with the color i , then every vertex in $V(G^*) \setminus A_i^*$ is incident with an edge of color i . We repeat this procedure for each of the colors from $[k + 1, k + \ell]$. After this has been done, each of these ℓ colors presents at all vertices from $V(G^*) \setminus U$. So at the conclusion of Step 3, all of the edges in G_A^* and G_B^* are colored, some of the edges of H are colored, each of the k color classes for colors from $[1, k]$ is a 1-factor of G^* , and each of the ℓ colors from $[k + 1, k + \ell]$ presents at all vertices from $V(G^*) \setminus U$.

Step 4: Coloring the graph R

Let R be the submultigraph of G^* consisting of the remaining uncolored edges. These edges all belong to H , so R is a submultigraph of H and hence is bipartite. We claim that

$\Delta(R) = \Delta(G^*) - k - \ell$. This is obvious when G is in Conditions (a) to (d). Thus we assume that G is in Condition (e). Note that every vertex from $V(G^*) \setminus U$ presents every color from $[1, k + \ell]$ and so those vertices have degree at most $\Delta(G^*) - k - \ell$ in R . For some vertices from U , they present all the colors from $[1, k]$. Thus by (7), those vertices have degree at most

$$\Delta(G^*) - \frac{1}{3}\eta n - k < \Delta(G^*) - k - \ell = \Delta(G^*) - k - 2\lfloor n^{\frac{5}{6}} \rfloor$$

in R . By Theorem 2.9 we can color the edges of R with $\Delta(R)$ colors from $[k + \ell + 1, \Delta(G^*)]$. Thus $\chi'(G^*) \leq k + \ell + (\Delta(G^*) - k - \ell) = \Delta(G^*)$ and so $\chi'(G^*) = \Delta(G^*)$, as desired.

Lastly, we check that there is a polynomial time algorithm to obtain an edge $\Delta(G)$ -coloring of G . By Lemma 4.1, we can obtain a desired partition $\{A, B\}$ of $V(G)$ in polynomial time. Also, it is polynomial time to edge color $G_{A,B}$ by an algorithm described in [17]. Modifying G_A and G_B into G_A^* and G_B^* and the corresponding edge colorings into the desired edge-colorings can be done in polynomial time too. In Step 2, the construction of the alternating paths and swaps of the colors on the paths can be done in $O(n^3)$ -time, as the total number of colors missing at vertices is $O(n^2)$ and it takes $O(n)$ -time to find an alternating path for a MCC-pair. In Step 3, there is polynomial time algorithm (see e.g. [14]) to edge color R_A and R_B using at most ℓ colors. Then by doing Kempe changes as mentioned in the comments immediately after Theorem 2.10, these edge colorings can be modified into equalized edge-colorings in polynomial time. The last step is to edge color the bipartite multigraph R using $\Delta(R)$ colors, which can be done in polynomial time in n , for example, using an algorithm from [5]. Thus, there is a polynomial time algorithm that gives an edge coloring of G using $\Delta(G)$ colors. \square

References

- [1] J. A. Bondy and U. S. R. Murty. Graph theory with applications. *American Elsevier Publishing Co., Inc., New York*, (Elsevier North-Holland), 1976.
- [2] A. G. Chetwynd and A. J. W. Hilton. 1-factorizing regular graphs of high degree—an improved bound. *Discrete Math.*, 75(1-3): 103–112, 1989.
- [3] A. G. Chetwynd and A. J. W. Hilton. Star multigraphs with three vertices of maximum degree. *Math. Proc. Cambridge Philos. Soc.*, 100(2):303–317, 1986.
- [4] A. G. Chetwynd and A. J. W. Hilton. The edge-chromatic class of graphs with maximum degree at least $|V| - 3$. In *Graph theory in memory of G. A. Dirac (Sandbjerg, 1985)*, volume 41 of *Ann. Discrete Math.*, pages 91–110. North-Holland, Amsterdam, 1989.
- [5] R. Cole and J. Hopcroft. On edge coloring bipartite graphs. *SIAM J. Comput.*, 11(3):540–546, 1982.

- [6] B. Csaba, D. Kühn, A. Lo, D. Osthus, and A. Treglown. Proof of the 1-factorization and Hamilton decomposition conjectures. *Mem. Amer. Math. Soc.*, 244(1154):v+164, 2016.
- [7] G. A. Dirac. Some theorems on abstract graphs. *Proc. London Math. Soc. (3)*, 2:69–81, 1952.
- [8] S. Fiorini and R. J. Wilson. Edge-colourings of graphs, Research notes in Maths. Pitman, London, 1977.
- [9] S. Grünewald and E. Steffen. Independent sets and 2-factors in edge-chromatic-critical graphs. *J. Graph Theory*, 45(2):113–118, 2004.
- [10] R. G. Gupta. *Studies in the Theory of Graphs*. PhD thesis, Tata Institute of Fundamental Research, Bombay, 1967.
- [11] S. L. Hakimi. On realizability of a set of integers as degrees of the vertices of a linear graph. I. *J. Soc. Indust. Appl. Math.*, 10:496–506, 1962.
- [12] I. Holyer. The NP-completeness of edge-coloring. *SIAM J. Comput.*, 10(4):718–720, 1981.
- [13] J. E. Hopcroft, and R. M. Karp. An $n^{5/2}$ algorithm for maximum matchings in bipartite graphs. *SIAM J. Comput.*, 2(4):225–231, 1973.
- [14] H. J. Karloff and D. B. Shmoys. Efficient parallel algorithms for edge coloring problems. *J. Algorithms*, 8(1):39–52, 1987.
- [15] D. König. Über Graphen und ihre Anwendung auf Determinantentheorie und Mengenlehre. *Math. Ann.*, 77(4):453–465, 1916.
- [16] C. J. H. McDiarmid. The solution of a timetabling problem. *J. Inst. Math. Appl.*, 9:23–34, 1972.
- [17] J. Misra and D. Gries. A constructive proof of Vizing’s theorem. *Inform. Process. Lett.*, 41(3):131–133, 1992.
- [18] L. Perković and B. Reed. Edge coloring regular graphs of high degree. *Discrete Math.*, 165/166: 567–578, 1997.
- [19] M. Plantholt. Overfull conjecture for graphs with high minimum degree. *J. Graph Theory*, 47(2):73–80, 2004.
- [20] M. Plantholt. The chromatic index of graphs with large even order n and minimum degree at least $2n/3$. *Discrete Math.*, 345(7): 112880, 2022.
- [21] M. Plantholt and S. Shan. Edge coloring graphs with large minimum degree. [arXiv:2105.05286](https://arxiv.org/abs/2105.05286), 2021.

- [22] P. D. Seymour. On multicolourings of cubic graphs, and conjectures of Fulkerson and Tutte. *Proc. London Math. Soc. (3)*, 38(3):423–460, 1979.
- [23] S. Shan. Chromatic index of dense quasirandom graphs. [arXiv:2104.06253](https://arxiv.org/abs/2104.06253), 2021.
- [24] M. Stiebitz, D. Scheide, B. Toft, and L. M. Favrholdt. Graph Edge Coloring: Vizing’s Theorem and Goldberg’s Conjecture. *Wiley Series in Discrete Mathematics and Optimization*. John Wiley & Sons, Inc., Hoboken, NJ, 2012.
- [25] V. G. Vizing. Critical graphs with given chromatic class. *Diskret. Analiz No.*, 5:9–17, 1965.