

On adjacency and Laplacian cospectral non-isomorphic signed graphs

Tahir Shamsher^a, S. Pirzada^b, Mushtaq A. Bhat^c,

^{a,b} *Department of Mathematics, University of Kashmir, Srinagar, Kashmir, India*

^c*Department of Mathematics, National Institute of Technology, Srinagar, India*

^atahir.maths.uok@gmail.com, ^bpirzadasd@kashmiruniversity.ac.in,

^cmushtaqab@nitsri.net

Abstract. Let $\Gamma = (G, \sigma)$ be a signed graph, where σ is the sign function on the edges of G . In this paper, we use the operation of partial transpose to obtain non-isomorphic Laplacian cospectral signed graphs. We will introduce two new operations on signed graphs. These operations will establish a relationship between the adjacency spectrum of one signed graph with the Laplacian spectrum of another signed graph. As an application, these new operations will be utilized to construct several pairs of cospectral non-isomorphic signed graphs. Finally, we construct integral signed graphs.

Keywords: Signed graph, partial transpose, cospectral signed graph, Laplacian cospectral signed graph, equienergetic signed graphs, integral signed graph.

AMS subject classification: 05C22, 05C50.

1 Introduction

Let $G = (V(G), E(G))$ be a simple connected graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. A signed graph is defined to be a pair $\Gamma = (G, \sigma)$, with $G = (V(G), E(G))$ as the underlying graph and $\sigma : E(G) \rightarrow \{-1, 1\}$ as the signing function. In this manuscript, bold lines denote positive edges, and dashed lines denote negative edges.

If all the edges of a signed graph have either a positive or negative sign, then it is said to be homogenous; otherwise, it is said to be heterogeneous. Signed graphs are a generalization of graphs, since they are homogeneous signed graphs with each edge positive. The sign of a signed graph is the product of the signs of its edges. A signed graph is said to be positive (or negative) if its sign is positive (or negative), that is, it has an even (or odd) number of negative edges. If all of the edges of a signed graph are positive (or negative), it is said to be all-positive (or all-negative) respectively. A signed graph is said to be balanced if none of its cycles is negative, otherwise unbalanced.

In a signed graph $\Gamma = (G, \sigma)$, the degree of a vertex v is the same as its degree in the underlying graph G (denoted by $d_v(G)$). Let Γ be a signed graph with vertex set $V(G)$ and let $X \subset V(G)$ be a nonempty set. Let Γ^X denote the signed graph obtained from Γ by reversing signs of edges between X and $V(G) - X$. Then, we say Γ^X is switching equivalent to Γ . Here, we note that the switching is an equivalence relation and preserves the eigenvalues of the adjacency and the Laplacian matrix including their multiplicities. A switching class is represented by a single signed graph.

The adjacency matrix of a signed graph Γ , whose vertices are v_1, v_2, \dots, v_n , is the $n \times n$ matrix $A(\Gamma) = (a_{ij})$, where

$$a_{ij} = \begin{cases} \sigma(v_i, v_j), & \text{if there is an edge from } v_i \text{ to } v_j, \\ 0, & \text{otherwise.} \end{cases}$$

For a graph G , the Laplacian matrix is $L(G) = D(G) - A(G)$ and signless Laplacian matrix is $Q(G) = D(G) + A(G)$, where $A(G)$ and $D(G)$ are respectively the adjacency matrix and the diagonal matrix of vertex degrees of G . The Laplacian matrix of Γ is $L(\Gamma) = L(G, \sigma) = D(G) - A(\Gamma)$. Note that $L(G, +) = L(G)$ and $L(G, -) = Q(G)$. The characteristic polynomial $|xI - A(\Gamma)|$ and eigenvalues of the adjacency matrix $A(\Gamma)$ of the signed graph Γ are denoted by $\phi_\Gamma(x)$ and $\lambda_1, \lambda_2, \dots, \lambda_n$ respectively. The characteristic polynomial $|xI - L(\Gamma)|$ and eigenvalues of the Laplacian matrix $L(\Gamma)$ of the signed graph Γ are denoted by $\psi_\Gamma(x)$ and $\mu_1, \mu_2, \dots, \mu_n$ respectively. For a graph G (resp. signed graph Γ), eigenvalues of its adjacency matrix and Laplacian matrix are called adjacency and Laplacian eigenvalues of G (resp. Γ). Clearly, $A(\Gamma)$ and $L(\Gamma)$ are real symmetric and so all their eigenvalues are real. Let the signed graph Γ of order n has distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$ and let their respective multiplicities be m_1, m_2, \dots, m_k . The adjacency spectrum of Γ is written as $Spec(\Gamma) = \{\lambda_1^{(m_1)}, \lambda_2^{(m_2)}, \dots, \lambda_k^{(m_k)}\}$. A signed graph is said to be an integral signed graph if its adjacency spectrum consists of integers only.

Given a graph G , its subdivision graph $S(G)$ is obtained from G by replacing each of its edge

by a path of length 2, or, equivalently, by inserting an additional vertex into each edge of G . If two signed graphs have the same adjacency spectrum (Laplacian spectrum), they are said to be cospectral (Laplacian cospectral); otherwise, they are noncospectral (Laplacian noncospectral). Any two isomorphic signed graphs are cospectral (Laplacian cospectral). A signed graph is said to be determined by its adjacency spectrum if cospectral signed graphs are isomorphic signed graphs. It is well-known that in general the adjacency spectrum does not determine the signed graph, and this problem has attracted to identify, if any, cospectral non-isomorphic signed graphs for a given class of signed graphs. For open problems in signed graphs we refer to [3].

The energy of a graph G is the sum of the absolute values of its adjacency eigenvalues. This concept was extended to signed graphs by Germina, Hameed and Zaslavsky [8]. The energy of a signed graph Γ with eigenvalues x_1, x_2, \dots, x_n is defined as $\mathcal{E}(\Gamma) = \sum_{j=1}^n |x_j|$. Two signed graphs of same order are said to be equienergetic if they have the same energy.

Harary [13] pioneered the use of signed graphs in connection with the study of social balance theory. Signed graphs have been intensively explored in a variety of fields such as group theory, topological graph theory and classical root system. The reader is referred to [18] for a complete bibliography on signed graphs.

The rest of the paper is organized as follows. In Section 2, we give some preliminary results which will be used in the sequel. In Section 3, we define the concept of partial transpose to signed graphs and use it to obtain non-isomorphic Laplacian cospectral signed graphs. In section 4, we introduce two new operations on signed graphs and which will be utilized to construct cospectral non-isomorphic signed graphs, noncospectral equienergetic signed graphs and integral signed graphs.

2 Preliminaries

In this section, we recall some previously established results which will be required in the subsequent sections.

Definition 2.1 [7] Let $P = (p_{ij}) \in M_{m \times n}(\mathbb{R})$ and $Q \in M_{p \times q}(\mathbb{R})$. The Kronecker product of P and Q , denoted by $P \otimes Q$, is defined as

$$P \otimes Q = \begin{pmatrix} p_{11}Q & p_{12}Q & \cdots & p_{1n}Q \\ p_{21}Q & p_{22}Q & \cdots & p_{2n}Q \\ \vdots & \vdots & \ddots & \vdots \\ p_{m1}Q & p_{m2}Q & \cdots & p_{mn}Q \end{pmatrix}.$$

Lemma 2.2 [7] Let $P, Q \in M_n(\mathbb{R})$ be two square matrices of order n . Let λ be an eigenvalue of matrix P with corresponding eigenvector x and μ be an eigenvalue of matrix Q with corresponding eigenvector y . Then $\lambda\mu$ is an eigenvalue of $P \otimes Q$ with corresponding eigenvector $x \otimes y$.

Lemma 2.3 [8] Let Γ_1 and Γ_2 be two signed graphs with respective eigenvalues x_1, x_2, \dots, x_{n_1} and y_1, y_2, \dots, y_{n_2} . Then

- (i) the eigenvalues of $\Gamma_1 \times \Gamma_2$ are $x_i + y_j$, for all $i = 1, 2, \dots, n_1$ and $j = 1, 2, \dots, n_2$,
- (ii) the eigenvalues of $\Gamma_1 \otimes \Gamma_2$ are $x_i y_j$, for all $i = 1, 2, \dots, n_1$ and $j = 1, 2, \dots, n_2$.

The Cartesian product (or sum) of two signed graphs $\Gamma_1 = (V(G_1), E(G_1), \sigma_1)$ and $\Gamma_2 = (V(G_2), E(G_2), \sigma_2)$, denoted by $\Gamma_1 \times \Gamma_2$, is the signed graph $(V(G_1) \times V(G_2), E, \sigma)$, where the edge set is that of the Cartesian product of underlying unsigned graphs and the sign function is defined by

$$\sigma((u_i, v_j), (u_k, v_l)) = \begin{cases} \sigma_1(u_i, u_k), & \text{if } j = l, \\ \sigma_2(v_j, v_l), & \text{if } i = k. \end{cases}$$

The Kronecker product (or conjunction) of two signed graphs $\Gamma_1 = (V(G_1), E(G_1), \sigma_1)$ and $\Gamma_2 = (V(G_2), E(G_2), \sigma_2)$, denoted by $\Gamma_1 \otimes \Gamma_2$, is the signed graph $(V(G_1) \otimes V(G_2), E, \sigma)$, where the edge set is that of the Kronecker product of underlying unsigned graphs and the sign function is defined by $\sigma((u_i, v_j), (u_k, v_l)) = \sigma_1(u_i, u_k)\sigma_2(v_j, v_l)$.

Lemma 2.4 [4] Let Γ be an unbalanced signed graph with at least one edge, whose spectrum is symmetric about the origin, having eigenvalues $\xi_1, \xi_2, \dots, \xi_n$. Then $\Gamma \times K_2$ and $\Gamma \otimes K_2$ are unbalanced, noncospectral and equienergetic if and only if $|\xi_j| \geq 1$, for all $j = 1, 2, \dots, n$.

Lemma 2.5 [12] Let $P(\bullet)$ be a given polynomial. If μ is an eigenvalue of $A \in M_n$, while y is an associated eigenvector, then $P(\mu)$ is an eigenvalue of the matrix $P(A)$ and y is an associated eigenvector with $P(\mu)$.

Lemma 2.6 [4] *Let Γ be a signed graph of order n . Then the following statements are equivalent.*

- (i) *The spectrum of Γ is symmetric about the origin,*
- (ii) *$\phi_\Gamma(x) = x^n + \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor} (-1)^k b_{2k} x^{n-2k}$, where b_{2k} are non negative integers for all $k = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor$,*
- (iii) *Γ and $-\Gamma$ are cospectral, where $-\Gamma$ is the signed graph obtained by negating sign of each edge of Γ .*

Lemma 2.7 [17] *For infinitely many n , there exists a family of 2^k pairwise nonisomorphic Laplacian integral, Laplacian cospectral graphs on n vertices, where $k > \frac{n}{(2\log_2(n))}$.*

3 Constructing Laplacian cospectral non-isomorphic signed graphs

Dutta [10] constructed large families of non-isomorphic signless Laplacian cospectral graphs using partial transpose on graphs. In this section, we define partial transpose for signed graphs. Let $\Gamma = (G, \sigma)$ be a signed graph on $2n$ vertices with vertex set $V(G) = V_1 \cup V_2$, such that $V_1 \cap V_2 = \emptyset$, and $V_1 = \{u_1, u_2, \dots, u_n\}$, $V_2 = \{v_1, v_2, \dots, v_n\}$. We denote $\langle V_1 \rangle_\Gamma$ and $\langle V_2 \rangle_\Gamma$ as the induced signed subgraphs of Γ generated by V_1 and V_2 , respectively. The spanning signed subgraph of Γ consisting of the signed edge set $\{(u_i, v_j) \in E(\Gamma) : u_i \in V_1, v_j \in V_2\}$ is denoted by $\langle V_1, V_2 \rangle_\Gamma$. Let $E(\widehat{\langle V_1, V_2 \rangle_\Gamma}) = \{(u_j, v_i), \text{ where } (u_i, v_j) \in E(\langle V_1, V_2 \rangle_\Gamma)\}$ be the set of edges, which suggests that given any signed edge $(u_i, v_j) \in E(\langle V_1, V_2 \rangle_\Gamma)$ there is a unique signed edge $(u_j, v_i) \in E(\widehat{\langle V_1, V_2 \rangle_\Gamma})$ with the same sign as the sign of edge (u_j, v_i) in $\langle V_1, V_2 \rangle_\Gamma$. It is easy to see that the set $E(\widehat{\langle V_1, V_2 \rangle_\Gamma})$ consists of all edges of the form (u_i, v_j) in Γ , but the existence of an edge (u_i, v_j) in Γ does not assure the existence of (u_j, v_i) in Γ .

The partial transpose of a signed graph Γ , denoted by Γ^τ , is defined as $\Gamma^\tau = \Gamma - E(\langle V_1, V_2 \rangle_\Gamma) + E(\widehat{\langle V_1, V_2 \rangle_\Gamma})$. Note that, subtracting $E(\langle V_1, V_2 \rangle_\Gamma)$ indicates to remove all the existing signed edges in Γ of the form $(u_i, v_j) \in E(\langle V_1, V_2 \rangle_\Gamma)$. Then we include the non-existing edges $(u_j, v_i) \in E(\widehat{\langle V_1, V_2 \rangle_\Gamma})$ to construct Γ^τ . If $i = j$, then the edge (u_i, v_i) will be removed and added again, that is the edge (u_i, v_i) is unaltered under partial transpose. Therefore, partial transpose of a signed graph Γ is an operation on the edge set which replaces the signed edge (u_i, v_j) with the sign $\sigma = \pm 1$, with the corresponding signed edge (u_j, v_i) with the same sign σ .

Consider the signed graphs Γ_1 and Γ_1^τ as shown in Figure 1. Here, we have $V_1 = \{u_1, u_2, u_3\}$, $V_2 = \{v_1, v_2, v_3\}$ and $E(\langle V_1, V_2 \rangle_{\Gamma_1}) = \{(u_1, v_1), (u_1, v_3)\}$. Thus, $E(\widehat{\langle V_1, V_2 \rangle_{\Gamma_1}}) = \{(u_1, v_1), (u_3, v_1)\}$.

Here, we replace the existing signed edge (u_1, v_3) with the non-existing signed edge (u_3, v_1) .

Remark 3.1 The partial transpose of a signed graph is labelling dependent. Therefore, isomorphic signed graphs may have non-isomorphic partial transposes, depending on the labellings. The partial transpose keeps $\langle V_1 \rangle$ and $\langle V_2 \rangle$ unaltered. The total number of vertices remains unchanged. If degree of a vertex w_i in the signed graph Γ is $d_\Gamma(w_i)$, then

$$\sum_{i=1}^n (d_\Gamma(u_i) + d_\Gamma(v_i)) = \sum_{i=1}^n (d_{\Gamma^\tau}(u_i) + d_{\Gamma^\tau}(v_i)).$$


A cycle $C_l^\sigma(v_1, v_2, \dots, v_l, v_1)$ in a signed graph $\Gamma = (G, \sigma)$ is a finite sequence of distinct vertices such that $(v_i, v_{i+1}) \in E(\Gamma)$ for all $i = 1, 2, \dots, l-1$ and $(v_l, v_1) \in E(\Gamma)$. We denote the negative edges in the signed cycle $C_l^\sigma(v_1, v_2, \dots, v_l, v_1)$ by putting the bar over the corresponding adjacent vertices. For example, the cycle $C_4^\sigma(v_1, v_2, v_3, v_4, v_1)$ on four vertices such that the only edge $(v_1, v_2) \in E(\Gamma)$ has negative sign will be denoted by $C_4^-(\overline{v_1}, \overline{v_2}, v_3, v_4, v_1)$. Similarly if only two consecutive edges $(v_1, v_2), (v_2, v_3) \in E(\Gamma)$ have negative signs, then the cycle $C_4^\sigma(v_1, v_2, v_3, v_4, v_1)$ will be denoted by $C_4^+(\overline{v_1}, \overline{v_2}, \overline{v_3}, v_4, v_1)$. In a signed graph Γ , a signed TU -subgraph Γ' is a signed subgraph whose components are trees or unbalanced unicyclic graphs, namely the unique cycle contains an odd number of negative edges. Thus, if H is a signed TU -subgraph, then $H = T_1 \cup T_2 \cup \dots \cup T_p \cup U_1 \cup U_2 \cup \dots \cup U_q$, where T_i 's are trees and U_i 's are unbalanced unicyclic graphs. The weight of the signed TU -subgraph H is defined as $w(H) = 4^q \prod_{i=1}^p |T_i|$, where $|T_i|$ is the number of vertices in the tree T_i . Note that we define $\prod_{i=1}^p |T_i| = 1$ when $p = 0$. The relation between the coefficients of the Laplacian characteristic polynomial with the TU -subgraphs of a signed graph can be seen in [[1], Theorem 3.9]. Let Γ be a signed graph with Laplacian characteristic polynomial $\psi(\Gamma, x) = x^n + a_1 x^{n-1} + \dots + a_{n-1} x + a_n$. Then its coefficients are given by

$$a_i = (-1)^i \sum_{H \in \mathcal{H}_i(\Gamma)} w(H) \quad (i = 1, 2, \dots, n), \quad (3.1)$$

where $\mathcal{H}_i(\Gamma)$ denotes the set of signed TU -subgraphs of Γ containing i edges. Two sets of signed TU -subgraphs $\mathcal{H}_i(\Gamma)$ and $\mathcal{H}_i(\Gamma')$ are comparable if

$$\sum_{H \in \mathcal{H}_i(\Gamma)} w(H) = \sum_{H \in \mathcal{H}_i(\Gamma')} w(H).$$

Now, Eq. (3.1) suggests that if Γ and Γ' are Laplacian cospectral, then the sets of their signed TU -subgraphs are comparable for all $i = 1, 2, \dots, m$, where m is the number of edges in the


 Figure 1: Signed graph Γ_1 and its partial transpose Γ_1^τ .

signed graph Γ . We say two signed graphs Γ_1 and Γ_2 are comparable if $\mathcal{H}_i(\Gamma_1)$ and $\mathcal{H}_i(\Gamma_2)$ are comparable for all i . As an example, two signed paths with equal number of vertices are comparable.


Example 3.1 Consider the signed graphs Γ_1 and Γ_1^τ as shown in Figure 1. We observe that Γ_1 contains two cycles $C_3^+(u_1, u_2, u_3, u_1)$ and $C_4^-(\overline{u_1, v_1}, v_2, v_3, u_1)$. The partial transpose Γ_1^τ of Γ_1 which is obtained by replacing the signed edge (u_1, v_3) with (u_3, v_1) preserves both these cycles $C_3^+(u_1, u_2, u_3, u_1)$ and $C_4^-(\overline{u_1, v_1}, u_3, u_2, u_1)$ in Γ_1^τ . As balanced cycle do not contribute to the coefficients of the Laplacian characteristic polynomial of a signed graph, therefore the signed TU -subgraphs generated by $C_4^-(\overline{u_1, v_1}, v_2, v_3, u_1)$ and $C_4^-(\overline{u_1, v_1}, u_3, u_2, u_1)$ in Γ_1 and Γ_1^τ respectively, are isomorphic, and have same contribution in $\psi_{\Gamma_1}(x)$ and $\psi_{\Gamma_1^\tau}(x)$.

The signed edges $K_{1,3} = \{(u_1, v_1), (u_1, v_3), (u_1, u_3)\}$ form a tree in Γ_1 . It is replaced by an unbalanced unicyclic TU -subgraph $C_3^-(\overline{u_1, v_1}, u_3, u_1)$ in Γ_1^τ . Clearly, the signed TU -subgraphs generated by $K_{1,3}$ and $C_3^-(\overline{u_1, v_1}, u_3, u_1)$ in Γ_1 and Γ_1^τ respectively have equal contribution in $\psi_{\Gamma_1}(x)$ and $\psi_{\Gamma_1^\tau}(x)$. Therefore, all the signed TU -subgraphs of Γ_1 and Γ_1^τ are comparable. Thus they have same Laplacian characteristic polynomial, which can be easily calculated by Eq. (3.1) and is given by

$$\psi_{\Gamma_1}(x) = \psi_{\Gamma_1^\tau}(x) = x^6 - 14x^5 + 73x^4 - 176x^3 + 196x^2 - 88x + 12.$$

If we add a signed edge $(w_i, w_j) : w_i, w_j \in V(G)$ in a signed graph $\Gamma = (G, \sigma)$, then the resultant signed graph will be denoted by $\Gamma' = \Gamma + \{(w_i, w_j)\}$. Similarly, $\Gamma' = \Gamma - \{(w_i, w_j)\}$ denotes the signed graph obtained by removing an edge $(w_i, w_j) : w_i, w_j \in V(G)$ in a signed graph $\Gamma = (G, \sigma)$. Whether the added/removed edge (w_i, w_j) is positive or negative, we denote a negative edge by $\overline{(w_i, w_j)}$, and a positive edge without a bar over the edge (w_i, w_j) .

Theorem 3.1 *Let the signed subgraphs $\langle V_1 \rangle_\Gamma$ and $\langle V_2 \rangle_\Gamma$ of the signed graph Γ be two paths on n vertices with each edge being positive. Let $\langle V_1, V_2 \rangle_\Gamma$ be an empty signed graph. For the new*

Figure 2: Signed graphs Γ_1 , Γ_1^τ , Γ_2 and Γ_3 .

constructed signed graph $\Gamma_1 = \Gamma + \left\{ (u_1, u_n), \overline{(u_1, v_1)}, (u_1, v_n) \right\}$,

(i) the signed graph Γ_1 is non-isomorphic and Laplacian cospectral to its partial transpose Γ_1^τ .

(ii) the signed graphs $\Gamma_2 = \Gamma_1 - \left\{ (u_{n-1}, u_n), \overline{(u_1, v_1)} \right\} + \left\{ \overline{(u_{n-1}, u_n)}, (u_1, v_1) \right\}$ and $\Gamma_3 = \Gamma_1^\tau - \left\{ (u_{n-1}, u_n) \right\} + \left\{ \overline{(u_{n-1}, u_n)} \right\}$ are non-isomorphic and Laplacian cospectral.


Proof. (i) The cycles generated by additional three edges and their incidence with existing edges in Γ_1 are $C_n^+(u_1, u_2, u_3, \dots, u_n, u_1)$ and $C_{n+1}^-(\overline{u_1, v_1}, v_2, \dots, v_n, u_1)$, respectively. The signed spanning subgraph $\langle V_1, V_2 \rangle_\Gamma$ contains only two signed edges which are $\overline{(u_1, v_1)}$ and (u_1, v_n) . Partial transpose replaces (u_1, v_n) with (u_n, v_1) . The cycles $C_n^+(u_1, u_2, u_3, \dots, u_n, u_1)$ and $C_{n+1}^-(\overline{u_1, v_1}, v_2, \dots, v_n, u_1)$ remain invariant (into an isomorphic cycles) under partial transpose on Γ_1 . As balanced cycles do not contribute to the coefficients of the Laplacian characteristic polynomial of a signed graph, therefore the signed TU -subgraphs generated by $C_{n+1}^-(\overline{u_1, v_1}, v_2, \dots, v_n, u_1)$ and $C_{n+1}^-(\overline{u_1, v_1}, u_n, \dots, u_2, u_1)$ in Γ_1 and Γ_1^τ , respectively, are isomorphic and have same contribution in $\psi_{\Gamma_1}(x)$ and $\psi_{\Gamma_1^\tau}(x)$. Now, the edges $\{(u_1, v_1), (u_1, v_n), (u_1, u_n)\}$ form a signed tree in Γ_1 . It is replaced by an unbalanced unicyclic TU -subgraph $C_3^-(\overline{u_1, v_1}, u_n, u_1)$ in Γ_1^τ and have same contribution in $\psi_{\Gamma_1}(x)$ and $\psi_{\Gamma_1^\tau}(x)$. Therefore, all the signed TU -subgraphs of Γ_1 and Γ_1^τ are comparable. Thus, by Eq. (3.1), they have same Laplacian characteristic polynomial, which proves the result in this case.

(ii) The proof of (ii) is similar to that of (i). ■

Example 3.2 Consider the signed graphs Γ_1 , Γ_1^τ , Γ_2 and Γ_3 as given in Figure 2. They are constructed by using Theorem 3.1. Their Laplacian characteristic polynomials are respectively given as below.

$$\psi_{\Gamma_1}(x) = \psi_{\Gamma_1^\tau}(x) = x^8 - 18x^7 + 131x^6 - 498x^5 + 1061x^4 - 1256x^3 + 764x^2 - 200x + 16,$$

$$\psi_{\Gamma_2}(x) = \psi_{\Gamma_3}(x) = x^8 - 18x^7 + 131x^6 - 498x^5 + 1065x^4 - 1288x^3 + 848x^2 - 280x + 36.$$


 Figure 3: Signed graphs Γ_1 , Γ_1^τ , Γ_2 , Γ_3 , Γ_4 and Γ_5 .


Clearly, the signed graphs Γ_1 and Γ_1^τ are non-isomorphic and Laplacian cospectral. Also, Γ_2 and Γ_3 are non-isomorphic and Laplacian cospectral signed graphs.

Theorem 3.2 *Let the signed subgraphs $\langle V_1 \rangle_\Gamma$ and $\langle V_2 \rangle_\Gamma$ of Γ be two cycles on n vertices with each edge being positive. Let $\langle V_1, V_2 \rangle_\Gamma$ be an empty signed graph. Given two non-adjacent vertices u_i and u_j with $i < j$, construct a new signed graph $\Gamma_1 = \Gamma + \left\{ (u_i, u_j), \overline{(u_i, v_i)}, (u_i, v_j) \right\}$. Then*

- (i) *the signed graph Γ_1 is non-isomorphic and Laplacian cospectral to its partial transpose Γ_1^τ ,*
- (ii) *the signed graphs $\Gamma_2 = \Gamma_1 - \left\{ (u_{n-1}, u_n), \overline{(u_i, v_i)}, (u_{j-1}, u_j) \right\} + \left\{ \overline{(u_{n-1}, u_n)}, (u_1, v_1), \overline{(u_{j-1}, u_j)} \right\}$ and $\Gamma_3 = \Gamma_1^\tau - \left\{ (u_{n-1}, u_n), (u_{j-1}, u_j) \right\} + \left\{ \overline{(u_{n-1}, u_n)}, \overline{(u_{j-1}, u_j)} \right\}$ are non-isomorphic and Laplacian cospectral,*
- (iii) *the signed graphs $\Gamma_4 = \Gamma_1 - \left\{ (v_{n-1}, v_n), \overline{(u_i, v_i)}, (u_{j-1}, u_j) \right\} + \left\{ \overline{(v_{n-1}, v_n)}, (u_1, v_1), \overline{(u_{j-1}, u_j)} \right\}$ and $\Gamma_5 = \Gamma_1^\tau - \left\{ (v_{n-1}, v_n), (u_{j-1}, u_j) \right\} + \left\{ \overline{(v_{n-1}, v_n)}, \overline{(u_{j-1}, u_j)} \right\}$ are non-isomorphic and Laplacian cospectral.*

Proof. The proof is similar to that of Theorem 3.1. ■

Example 3.3 Consider the signed graphs Γ_1 , Γ_1^τ , Γ_2 , Γ_3 , Γ_4 and Γ_5 as shown in Figure 3. Here $n = 4$, $i = 1$ and $j = 3$. The signed graph Γ_1 , which is generated by Theorem 3.2 is non-isomorphic and Laplacian cospectral to its partial transpose Γ_1^τ . We obtain the signed graph Γ_2 from Γ_1 by replacing the positive edges (u_2, u_3) and (u_3, u_4) with negative edges $\overline{(u_2, u_3)}$ and $\overline{(u_3, u_4)}$ and negative edge $\overline{(u_1, v_1)}$ with the positive edge (u_1, v_1) . Also, the signed graph Γ_3 is obtained from Γ_1^τ by replacing the positive edges (u_2, u_3) and (u_3, u_4) with negative edges $\overline{(u_2, u_3)}$ and $\overline{(u_3, u_4)}$. The signed graphs Γ_2 and Γ_3 are non-isomorphic and Laplacian cospectral. Similarly the non-isomorphic and Laplacian cospectral signed graphs Γ_4 and Γ_5 are obtained

Figure 4: Signed graphs Γ and Γ^τ .

from Γ_1 and Γ_1^τ , respectively, as in Theorem 3.2.

Remark 3.2 In Example 3.3, we have seen that Γ_1 and Γ_1^τ are Laplacian cospectral signed graphs. Also, we have mentioned that Γ_1^τ is the partial transpose of Γ_1 . But, not all signed graphs are Laplacian cospectral to their partial transpose, for instance, consider the signed graphs Γ and Γ^τ as given in Figure 4. It is easy to calculate that the Laplacian characteristic polynomials of Γ and Γ^τ are $\psi_\Gamma(x) = x^6 - 12x^5 + 52x^4 - 105x^3 + 104x^2 - 48x + 8$ and $\psi_{\Gamma^\tau}(x) = x^6 - 12x^5 + 51x^4 - 94x^3 + 72x^2 - 18x$.

Let G be a graph and $\Gamma = (G, \sigma)$ be a signed graph on G . Hou et al. [16] raised the following two problems.

Problem 1. Let G be a graph, $\Gamma_1 = (G, \sigma_1), \Gamma_2 = (G, \sigma_2)$ be two signed graphs on G , and $\det L(\Gamma_1) = \det L(\Gamma_2)$. Are $L(\Gamma_1)$ and $L(\Gamma_2)$ cospectral?

Problem 2. Do there exist pairs $\Gamma_1 = (G_1, \sigma_1), \Gamma_2 = (G_2, \sigma_2)$ of signed graphs that have either of the following properties (i) and (ii)?

- (i) Γ_1 and Γ_2 are not balanced but Laplacian cospectral such that G_1 and G_2 are nonisomorphic.
- (ii) Γ_1 and Γ_2 are not balanced but Laplacian cospectral such that G_1 and G_2 are not cospectral.

The statement of Problem 1 is not always true. To see this, let Γ_1 be a signed graph as shown in Figure 3. Let Γ' be the signed graph obtained from Γ_1 by replacing the negative edge $\overline{(u_1, v_1)}$ with positive edge (u_1, v_1) and positive edge (v_3, v_4) with negative edge $\overline{(v_3, v_4)}$. The Laplacian characteristic polynomials of Γ_1 and Γ' are respectively given by

$$\psi_{\Gamma_1}(x) = x^8 - 22x^7 + 197x^6 - 928x^5 + 2476x^4 - 3736x^3 + 2976x^2 - 1056x + 128,$$

$$\psi_{\Gamma'}(x) = x^8 - 22x^7 + 197x^6 - 928x^5 + 2476x^4 - 3748x^3 + 3048x^2 - 1152x + 128.$$

The underlying graphs of Γ_1 and Γ' are isomorphic and $\det L(\Gamma_1) = \det L(\Gamma')$. It is clear that the signed graphs Γ_1 and Γ' are not Laplacian cospectral and this answers Problem 1.

For Problem 2, consider the signed graph Γ_1 and its partial transpose Γ_1^τ as given in Figure 3. Clearly, the underlying graphs of Γ_1 and Γ_1^τ are non-isomorphic. The unbalanced signed graphs Γ_1 and Γ_1^τ are Laplacian cospectral. Also, it is easy to see that the underlying graph of Γ_1 and Γ_1^τ are not cospectral and this answers Problem 2.

4 Constructing cospectral(adjacency) non-isomorphic signed graphs, integral signed graphs and equienergetic(adjacency) signed graphs

The novel cospectral non-isomorphic signed graph constructions have implications for the complexity of the graph isomorphism problem. This necessitates the creation of methods for detecting and/or creating cospectral non-isomorphic graphs. Seidel switching, Godsil–McKay (GM) switching, and others are well-known approaches for constructing cospectral graphs. In 2019, Belardo et al. [2] used the Godsil-Mckay-type procedures developed for graphs to construct the pairs of cospectral switching non-isomorphic signed graphs. In this section, we will introduce two new operations in signed graphs. These operations establish the relationship of the adjacency spectrum of one signed graph with the Laplacian spectrum of another signed graph. Furthermore, these operations will be utilized to construct the pairs of cospectral non-isomorphic signed graphs and integral signed graphs.

The usual orientation of edges in digraphs differs slightly from the orientation of signed graphs. In fact in signed graphs, instead of one arrow, we can use two arrows assigned to edges. Bidirected graphs are the result of this. An orientated signed graph, more exactly, is an ordered pair $\Gamma_{\vartheta} = (\Gamma, \vartheta)$, where

$$\vartheta : V(G) \times E(G) \rightarrow \{0, 1, -1\} \quad (4.2)$$

satisfying the following three conditions.

- (a) $\vartheta(u, vw) = 0$ whenever $u \neq v, w; u, v, w \in V(G)$ and $vw \in E(G)$,
- (b) $\vartheta(v, vw) = 1$ (or -1) if an arrow at v is going into (rep. out of) v . For illustration, see Figure 5,
- (c) $\vartheta(v, vw) \vartheta(w, vw) = -\sigma(vw)$.

As a result, positive edges are oriented edges, whereas negative edges are unoriented (see Figure 5). Therefore, every bidirected graph is also a signed graph. The converse is likewise true, however, one arrow (at any end) can be taken at random, whereas the other arrow (in light of (c) above) cannot. For an oriented signed graph Γ_{ϑ} , its incidence matrix $B_{\vartheta} = (b_{ij})$ is a matrix, whose rows correspond to vertices and columns to edges of G , with $b_{ij} = \vartheta(v_i, e_j)$ (here $v_i \in V(G)$, $e_j \in E(G)$). Usually, when only Γ is given, then we use the arbitrary orientation. So each row of the incidence matrix corresponding to vertex v_i contains d_{v_i} non-zero entries, each




Figure 5: Bidirected edges in signed graphs.

equal to $+1$ or -1 . On the other hand, each column of the incidence matrix corresponding to edge e_j contains two non-zero entries, each equal to $+1$ or -1 . Therefore, even in the case that multiple edges exist, we easily obtain

$$B_{\vartheta} B_{\vartheta}^T = D(G) - A(\Gamma_{\vartheta}) = L(\Gamma_{\vartheta}), \quad (4.3)$$

where $D(G)$ is the diagonal matrix of vertex degrees of G . It is easy to observe that $L(\Gamma_{\vartheta})$ is positive-semidefinite.

The subdivision signed graph $A(S(\Gamma_{\vartheta}))$ is the signed graph whose underlying graph is $S(G)$ with vertex set $V(G) \cup E(G)$. It preserves the orientation ϑ and can be represented in the block form as follows.

$$A(S(\Gamma_{\vartheta})) = \begin{pmatrix} O_n & B_{\vartheta} \\ B_{\vartheta}^T & O_m \end{pmatrix},$$

where $O_r \in M_r(\mathbb{R})$. It is easy to see that the signature σ of the subdivision signed graph is defined by $\sigma(v_i e_j) = \vartheta_{ij}$. An example of subdivision graph of a signed graph is shown in Figure 6.

Remark 4.1 Any orientation (random) ϑ to the edges of Γ gives rise to the same matrices $A(\Gamma_{\vartheta}) = A(\Gamma)$ and $L(\Gamma_{\vartheta}) = L(\Gamma)$, while the matrix $A(S(\Gamma_{\vartheta}))$ does depend on ϑ . Let S be a ± 1 diagonal matrix such that $B'_{\vartheta} = B_{\vartheta} S$. It can be easily seen that $A(S(\Gamma_{\vartheta'})) = [I_n \dot{+} S] A(S(\Gamma_{\vartheta})) [I_n \dot{+} S]$, where $\dot{+}$ denotes the direct sum of two matrices. From now on, the index ϑ will be not specified anymore.

Lemma 4.1 [1] *If B is the incident matrix of a connected signed graph $\Gamma = (G, \sigma)$ having n vertices. Then*

$$\text{rank}(B) = \begin{cases} n - 1, & \text{if } \Gamma \text{ is balanced,} \\ n, & \text{if } \Gamma \text{ is unbalanced.} \end{cases}$$




Figure 6: A signed graph and the corresponding signed subdivision graph.

Operation 4.1 Let Γ be a signed graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. Corresponding to each signed edge e_i , $1 \leq i \leq m$, in Γ , introduce a set U_i^p of p (positive integer) isolated vertices and make every vertex in U_i^p adjacent to the vertices incident with e_i , $i = 1, 2, \dots, m$ (in the same way as in subdivision signed graph $S(\Gamma)$) and remove edges of Γ only. The resultant signed graph is denoted by $S_p(\Gamma)$. The number of vertices and edges of the signed graph $S_p(\Gamma)$ are $n + pm$ and $2pm$, respectively. If $p = 1$, then $S_p(\Gamma)$ coincides with the subdivision graph $S(\Gamma)$. Figure 7 illustrates the above operation.

Theorem 4.2 Let Γ be a signed graph with n vertices and m edges. Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n \geq 0$ be the Laplacian eigenvalues of the signed graph Γ . Then the adjacency spectrum of $S_p(\Gamma)$ is


$$\text{Spec}(S_p(\Gamma)) = \begin{cases} \{0^{(pm-n+2)}, \pm\sqrt{p\mu_1^{(1)}}, \pm\sqrt{p\mu_2^{(1)}}, \dots, \pm\sqrt{p\mu_{n-1}^{(1)}}\} & \text{if } \Gamma \text{ is balanced,} \\ \{0^{(pm-n)}, \pm\sqrt{p\mu_1^{(1)}}, \pm\sqrt{p\mu_2^{(1)}}, \dots, \pm\sqrt{p\mu_{n-1}^{(1)}}, \pm\sqrt{p\mu_n^{(1)}}\}, & \text{if } \Gamma \text{ is unbalanced.} \end{cases}$$

Proof. Using the suitable labelling of the vertices of $S_p(\Gamma)$, the adjacency matrix of $S_p(\Gamma)$ can be written as

$$A(S_p(\Gamma)) = \begin{pmatrix} O & B & B & \dots & B \\ B^T & O & O & \dots & O \\ B^T & O & O & \dots & O \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ B^T & O & O & \dots & O \end{pmatrix}.$$

Therefore, we have

$$A(S_p(\Gamma))^2 = \begin{pmatrix} O & B & B & \dots & B \\ B^T & O & O & \dots & O \\ B^T & O & O & \dots & O \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ B^T & O & O & \dots & O \end{pmatrix} \begin{pmatrix} O & B & B & \dots & B \\ B^T & O & O & \dots & O \\ B^T & O & O & \dots & O \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ B^T & O & O & \dots & O \end{pmatrix}$$

Figure 7: Signed graphs Γ , $S(\Gamma)$ and $S_2(\Gamma)$.

$$\begin{aligned}
&= \begin{pmatrix} pBB^T & O & O & \dots & O \\ O & B^T B & B^T B & \dots & B^T B \\ O & B^T B & B^T B & \dots & B^T B \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ O & B^T B & B^T B & \dots & B^T B \end{pmatrix} \\
&= \begin{pmatrix} pBB^T & O_{1 \times p} \\ O_{p \times 1} & J_{p \times p} \otimes B^T B \end{pmatrix},
\end{aligned}$$

where $J_{p \times p} \otimes B^T B$ denotes the Kronecker product of the matrices $J_{p \times p}$ and $B^T B$, and $J_{p \times p}$ is a square matrix whose all entries are equal to 1. Thus

$$\text{Spec}(A(S_p(\Gamma))^2) = \text{Spec}(pBB^T) \cup \text{Spec}(J_{p \times p} \otimes B^T B).$$

As $B^T B$ is a real symmetric matrix of order m , so all its eigenvalues are real. Let $x_1 \geq x_2 \geq \dots \geq x_m$ be the eigenvalues of the matrix $B^T B$. Note that $\text{rank}(BB^T) = \text{rank}(B^T B) = \text{rank}(B)$. Therefore, by Lemma 4.1, we have

$$\text{Spec}(B^T B) = \begin{cases} \{0^{(m-n+1)}, x_1, x_2, \dots, x_{n-1}\} & \text{if } \Gamma \text{ is balanced,} \\ \{0^{(m-n)}, x_1, x_2, \dots, x_{n-1}, x_n\} & \text{if } \Gamma \text{ is unbalanced} \end{cases}$$

and

$$\text{Spec}(BB^T) = \text{Spec}(L(\Gamma)) = \begin{cases} \{0, \mu_1, \mu_2, \dots, \mu_{n-1}\} & \text{if } \Gamma \text{ is balanced,} \\ \{\mu_1, \mu_2, \dots, \mu_{n-1}, \mu_n\} & \text{if } \Gamma \text{ is unbalanced,} \end{cases}$$

where $x_n \neq 0$ and $\mu_n \neq 0$. As $\text{Spec}(J_{p \times p})$ is $\{0^{p-1}, p\}$, then by Lemma 2.2, we have

$$\text{Spec}(J_{p \times p} \otimes B^T B) = \begin{cases} \{0^{(pm-n+1)}, px_1, px_2, \dots, px_{n-1}\} & \text{if } \Gamma \text{ is balanced,} \\ \{0^{(pm-n)}, px_1, px_2, \dots, px_{n-1}, px_n\} & \text{if } \Gamma \text{ is unbalanced.} \end{cases}$$

We know that the underlying graph of a subdivision signed graph is always bipartite. Therefore the underlying graph of $S_p(\Gamma)$ is always bipartite. Note that the eigenvalues of $B^T B$ are given by the eigenvalues of $B B^T$, together with 0 of multiplicity $m - n$. Therefore, by Lemmas 2.5 and 2.6, we have

$$\text{Spec}(S_p(\Gamma)) = \begin{cases} \{0^{(pm-n+2)}, \pm\sqrt{p\mu_1^{(1)}}, \pm\sqrt{p\mu_2^{(1)}}, \dots, \pm\sqrt{p\mu_{n-1}^{(1)}}\} & \text{if } \Gamma \text{ is balanced,} \\ \{0^{(pm-n)}, \pm\sqrt{p\mu_1^{(1)}}, \pm\sqrt{p\mu_2^{(1)}}, \dots, \pm\sqrt{p\mu_{n-1}^{(1)}}, \pm\sqrt{p\mu_n^{(1)}}\} & \text{if } \Gamma \text{ is unbalanced.} \end{cases}$$

■

The following result can also be seen in [[1], Theorem 2.2].

Corollary 4.3 *Let Γ be a signed graph with n vertices and m edges. Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n \geq 0$ be the Laplacian eigenvalues of the signed graph Γ . Then the adjacency spectrum of $S(\Gamma)$ is*

$$\text{Spec}(S(\Gamma)) = \begin{cases} \{0^{(m-n+2)}, \pm\sqrt{\mu_1^{(1)}}, \pm\sqrt{\mu_2^{(1)}}, \dots, \pm\sqrt{\mu_{n-1}^{(1)}}\} & \text{if } \Gamma \text{ is balanced,} \\ \{0^{(m-n)}, \pm\sqrt{\mu_1^{(1)}}, \pm\sqrt{\mu_2^{(1)}}, \dots, \pm\sqrt{\mu_{n-1}^{(1)}}, \pm\sqrt{\mu_n^{(1)}}\} & \text{if } \Gamma \text{ is unbalanced.} \end{cases}$$

Operation 4.2 Let Γ be a signed graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. Let $S(\Gamma)$ be a signed subdivision graph of a signed graph Γ with vertex set $V(G) \cup E(G)$. Corresponding to every vertex v_i , $1 \leq i \leq n$, in $S(\Gamma)$, introduce a set V_i^p of p (positive integer) isolated vertices and join each vertex of V_i^p to the neighbors of v_i with the same sign as the vertex v_i in $S(\Gamma)$. Then in the resulting signed graph, corresponding to each vertex e_i , ($i = 1, 2, \dots, m$) introduce a set of k isolated vertices U_j^k , $1 \leq j \leq m$, where $k = p$ or $k = p - 1$, join each vertex in U_j^k to the neighbors of e_i with the same sign as the vertex e_i in $S(\Gamma)$. The resultant signed graph is denoted by $S_p^k(\Gamma)$. The number of vertices and edges of the graph $S_p^k(\Gamma)$ are $(p+1)n + (k+1)m$ and $2(p+1)(k+1)m$, respectively. Figure 8 illustrates the above operation.

Theorem 4.4 *Let Γ be a signed graph with n vertices and m edges. Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n \geq 0$ be the Laplacian eigenvalues of the signed graph Γ . Then the adjacency spectrum of $S_p^k(\Gamma)$ is*

$$\text{Spec}(S_p^k(\Gamma)) = \{0^{((p-1)n+(k+1)m+2)}, \pm\sqrt{(p+1)(k+1)\mu_1^{(1)}}, \dots, \pm\sqrt{(p+1)(k+1)\mu_{n-1}^{(1)}}\}$$

if Γ is balanced, and

$$\text{Spec}(S_p^k(\Gamma)) = \{0^{((p-1)n+(k+1)m)}, \pm\sqrt{(p+1)(k+1)\mu_1^{(1)}}, \dots, \pm\sqrt{(p+1)(k+1)\mu_{n-1}^{(1)}}, \pm\sqrt{(p+1)(k+1)\mu_n^{(1)}}\}$$

if Γ is unbalanced.

Proof. By the suitable labelling of the vertices of $S_p^k(\Gamma)$, the adjacency matrix $S_p^k(\Gamma)$ can be written as

$$A(S_p^k(\Gamma)) = \begin{pmatrix} O & B & O & \dots & O & B \\ B^T & O & B^T & \dots & B^T & O \\ O & B & O & \dots & O & B \\ \vdots & \vdots & \ddots & \vdots & & \\ O & B & O & \dots & O & B \\ B^T & O & B^T & \dots & B^T & O \end{pmatrix},$$


when $k = p$. If $k = p - 1$, then we have

$$A(S_p^k(\Gamma)) = \begin{pmatrix} O & B & O & \dots & B & O \\ B^T & O & B^T & \dots & O & B^T \\ O & B & O & \dots & B & O \\ \vdots & \vdots & \ddots & \vdots & & \\ B^T & O & B^T & \dots & O & B^T \\ O & B & O & \dots & B & O \end{pmatrix}.$$

To prove the result, the following two cases arise.

Case 1. Let Γ be a balanced signed graph with n vertices and m edges. Let $X \in M_{n \times 1}(\mathbb{R})$ and $Y \in M_{m \times 1}(\mathbb{R})$ be two non-zero column vectors. Let $Z = \begin{pmatrix} X \\ Y \end{pmatrix} \in M_{(n+m) \times 1}(\mathbb{R})$ be the eigenvector corresponding to the non-zero eigenvalue λ_i , $1 \leq i \leq 2n - 2$, of $S(\Gamma)$. Then $A(S(\Gamma))Z = \lambda_i Z$ implies that $BY = \lambda_i X$ and $B^T X = \lambda_i Y$. To find the eigenvalues of $S_p^k(\Gamma)$, consider the following two subcases.

Subcase 1.1. If $k = p$, then let $U = \begin{pmatrix} X \\ Y \\ \vdots \\ X \\ Y \end{pmatrix} \in M_{((p+1)n+(k+1)m) \times 1}(\mathbb{R})$ be a non-zero column


 Figure 8: Signed graphs Γ , Γ_ϑ , $S(\Gamma)$, $S_1^0(\Gamma)$ and $S_1^1(\Gamma)$.

vector such that

$$\begin{aligned}
 A(S_p^k(\Gamma))U &= \begin{pmatrix} O & B & O & \dots & O & B \\ B^T & O & B^T & \dots & B^T & O \\ O & B & O & \dots & O & B \\ \vdots & \vdots & \ddots & \vdots & & \\ O & B & O & \dots & O & B \\ B^T & O & B^T & \dots & B^T & O \end{pmatrix} \begin{pmatrix} X \\ Y \\ \vdots \\ X \\ Y \end{pmatrix} = \begin{pmatrix} (p+1)\lambda_i X \\ (p+1)\lambda_i Y \\ \vdots \\ (p+1)\lambda_i X \\ (p+1)\lambda_i Y \end{pmatrix} \\
 &= (p+1)\lambda_i U.
 \end{aligned}$$

Therefore $(p+1)\lambda_i$ is an eigenvalue of $S_p^k(\Gamma)$ corresponding to an eigenvector U . Thus the result follows by Corollary 4.3.

Subcase 1.2. If $k = p - 1$, then let $U = \begin{pmatrix} \sqrt{k+1}X \\ \sqrt{p+1}Y \\ \vdots \\ \sqrt{k+1}X \\ \sqrt{p+1}Y \\ \sqrt{k+1}X \end{pmatrix} \in M_{((p+1)n+(k+1)m) \times 1}(\mathbb{R})$ be a non-zero

column vector such that

$$\begin{aligned}
A(S_p^k(\Gamma))U &= \begin{pmatrix} O & B & O & \dots & B & O \\ B^T & O & B^T & \dots & O & B^T \\ O & B & O & \dots & B & O \\ \vdots & \vdots & \ddots & \vdots & & \\ B^T & O & B^T & \dots & O & B^T \\ O & B & O & \dots & B & O \end{pmatrix} \begin{pmatrix} \sqrt{k+1}X \\ \sqrt{p+1}Y \\ \vdots \\ \sqrt{k+1}X \\ \sqrt{p+1}Y \\ \sqrt{k+1}X \end{pmatrix} \\
&= \begin{pmatrix} (k+1)\lambda_i\sqrt{p+1}X \\ (p+1)\lambda_i\sqrt{k+1}Y \\ \vdots \\ (k+1)\lambda_i\sqrt{p+1}X \\ (p+1)\lambda_i\sqrt{k+1}Y \\ (k+1)\lambda_i\sqrt{p+1}X \end{pmatrix} \\
&= \sqrt{(k+1)(p+1)}\lambda_i U.
\end{aligned}$$

Therefore $\sqrt{(k+1)(p+1)}\lambda_i$ is an eigenvalue of $S_p^k(\Gamma)$ corresponding to an eigenvector U . Therefore the result follows by Corollary 4.3.

Case 2. When Γ is an unbalanced signed graph with n vertices and m edges, the proof is similar to that of Case 1. ■

Various constructions for cospectral non-isomorphic regular graphs, cospectral non-isomorphic Laplacian graphs and cospectral non-isomorphic signless Laplacian graphs can be seen in [6, 9–11, 14, 15]. The following results shows that these constructions including the constructions obtained in the last section can be utilized to obtain infinite families of cospectral non-isomorphic signed graphs.

Corollary 4.5 *Let Γ_1 and Γ_2 be two non-isomorphic signed graphs which are Laplacian cospectral. Then*

- (i) *the signed graphs $S_p(\Gamma_1)$ and $S_p(\Gamma_2)$ are cospectral and non-isomorphic,*
- (ii) *the signed graphs $S_p^k(\Gamma_1)$ and $S_p^k(\Gamma_2)$ are cospectral and non-isomorphic.*

Proof. Let Γ_1 and Γ_2 be two non-isomorphic signed graphs. Then, clearly $S_p(\Gamma_1)$ and $S_p(\Gamma_2)$ are non-isomorphic signed graphs and $S_p^k(\Gamma_1)$ and $S_p^k(\Gamma_2)$ are non-isomorphic signed graphs. Hence the result follows by Theorems 4.2 and 4.4.




Figure 9: Cospectral signed graphs $S_2(\Gamma_1)$ and $S_2(\Gamma_2)$.

Example 4.1 Consider the two non-isomorphic signed graphs Γ_1 and Γ_2 , which are Laplacian cospectral, as shown in Figure 9. Their Laplacian spectrum is respectively given by $Spec_L(\Gamma_1) = \{0, 2, 3^{(2)}, 3+\sqrt{5}, 3-\sqrt{5}\}$ and $Spec_L(\Gamma_2) = \{0, 2, 3^{(2)}, 3+\sqrt{5}, 3-\sqrt{5}\}$. It is easy to see that $S_2(\Gamma_1)$ and $S_2(\Gamma_2)$ are non-isomorphic signed graphs which are cospectral as their adjacency spectrum are respectively given by $Spec(S_2(\Gamma_1)) = \{0^{(10)}, \pm 2, \pm\sqrt{6}^{(2)}, \pm(\sqrt{6+\sqrt{20}}), \pm(\sqrt{6-\sqrt{20}})\}$ and $Spec(S_2(\Gamma_2)) = \{0^{(10)}, \pm 2, \pm\sqrt{6}^{(2)}, \pm(\sqrt{6+\sqrt{20}}), \pm(\sqrt{6-\sqrt{20}})\}$.

Corollary 4.6 Let Γ_1 and Γ_2 be two cospectral non-isomorphic r -regular signed graphs. Then
 (i) the signed graphs $S_p(\Gamma_1)$ and $S_p(\Gamma_2)$ are cospectral and non-isomorphic,
 (ii) The signed graphs $S_p^k(\Gamma_1)$ and $S_p^k(\Gamma_2)$ are cospectral and non-isomorphic.

Proof. Let Γ_1 and Γ_2 be two non-isomorphic cospectral regular signed graphs. Then $L(\Gamma_1) = D(\Gamma_1) - A(\Gamma_1)$ and $L(\Gamma_2) = D(\Gamma_2) - A(\Gamma_2)$ are cospectral. Hence the result follows by Corollary 4.5. ■

Corollary 4.7 Let Γ be a signed graph whose all Laplacian eigenvalues are perfect squares. Then

- (i) the signed graph $S_p(\Gamma)$ is integral, if p is a perfect square,
- (ii) the signed graph $S_p^k(\Gamma)$ is integral, if $(p+1)(k+1)$ is a perfect square.

Example 4.2 Let K_n be a balanced complete signed graph on n vertices, where $n = t^2$, $t \geq 2$ is a positive integer. Then

- (i) the signed graph $S_p(K_n)$ is integral, if p is a perfect square,
(ii) the signed graph $S_p^k(K_n)$ is integral, if $(p+1)(k+1)$ is a perfect square.

The following result is the graceful implication of Lemma 2.7 and Corollaries 4.5 and 4.6.

Theorem 4.8 *For infinitely many n , there exists a family of 2^k pairwise cospectral nonisomorphic signed graphs on n vertices, where $k > \frac{n}{(2\log_2(n))}$.*

The following result directly follows from Theorems 4.2 and 4.4.

Theorem 4.9 *Let Γ be a signed graph with n vertices and m edges. Then*

- (i) $\mathcal{E}(S_p(\Gamma)) = \sqrt{p}\mathcal{E}(S(\Gamma))$,
(ii) $\mathcal{E}(S_p^k(\Gamma)) = \sqrt{(p+1)(k+1)}\mathcal{E}(S(\Gamma))$.

Theorem 4.10 *Let Γ be an unbalanced unicyclic signed graph with at least one edge and having Laplacian eigenvalues $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n > 0$. Then $S(\Gamma) \times K_2$ and $S(\Gamma) \otimes K_2$ are noncospectral and equienergetic if and only if $\mu_n \geq 1$.*

Proof. Let Γ be an unbalanced unicyclic signed graph. Then, by Theorem 4.2, we have

$$\text{Spec}(S(\Gamma)) = \{\pm\sqrt{\mu_1}^{(1)}, \pm\sqrt{\mu_2}^{(1)}, \dots, \pm\sqrt{\mu_{n-1}}^{(1)}, \pm\sqrt{\mu_n}^{(1)}\}.$$

First, assume that $\mu_n \geq 1$. This implies that $|\sqrt{\mu_j}| \geq 1$, for all $j = 1, 2, \dots, n$. Also,

$$E(S(\Gamma) \times K_2) = 2 \sum_{j=1}^n (|\sqrt{\mu_j} + 1| + |\sqrt{\mu_j} - 1|).$$

As $|\sqrt{\mu_j}| \geq 1$, for all $j = 1, 2, \dots, n$, we have

$$\begin{aligned} E(S(\Gamma) \times K_2) &= 2 \sum_{j=1}^n (|\sqrt{\mu_j}| + 1 + |\sqrt{\mu_j}| - 1) \\ &= 2E(S(\Gamma)) \\ &= E(S(\Gamma))E(K_2) = E(S(\Gamma) \otimes K_2). \end{aligned}$$

Note that $\sqrt{\mu_1} + 1 \in \text{Spec}(S(\Gamma) \times K_2)$ but $\sqrt{\mu_1} + 1 \notin \text{Spec}(S(\Gamma) \otimes K_2)$. Therefore $S(\Gamma) \times K_2$ and $S(\Gamma) \otimes K_2$ are noncospectral. The converse is similar to that of the converse in Lemma 2.4. ■

Example 4.3. Let $C_3^- = (C_3, -)$ be an unbalanced unicyclic signed graph on 3 vertices. Its Laplacian spectrum is given by $\text{Spec}_L(C_3^-) = \{4, 1, 1\}$. Therefore C_3^- meets the requirement of




Figure 10: Signed graphs Γ_1 , Γ_2 , Γ_3 and Γ_4 .

Theorem 4.10. Hence $S(C_3^-) \times K_2$ and $S(C_3^-) \otimes K_2$ are noncospectral and equienergetic.

The following corollary directly follows from Theorems 4.2, 4.4 and 4.9.

Corollary 4.11 *Let Γ_1 and Γ_2 be two signed graphs whose signed subdivision graphs are noncospectral and equienergetic. Then*

- (i) *the signed graphs $S_p(\Gamma_1)$ and $S_p(\Gamma_2)$ are noncospectral and equienergetic,*
- (ii) *the signed graphs $S_p^k(\Gamma_1)$ and $S_p^k(\Gamma_2)$ are noncospectral and equienergetic.*

Example 4.4. Consider the signed graphs Γ_1 , Γ_2 , Γ_3 and Γ_4 as shown in Figure 10. The adjacency spectrum of their signed subdivision graphs is respectively given by $Spec(S(\Gamma_1)) = \{\pm 2, \pm 1^{(2)}, 0\}$, $Spec(S(\Gamma_2)) = \{\pm 2, \pm 1^{(2)}\}$, $Spec(S(\Gamma_3)) = \{\pm 2^{(3)}, \pm \sqrt{2}^{(3)}, \pm \sqrt{6}, 0^{(6)}\}$ and $Spec(S(\Gamma_4)) = \{\pm 1^{(2)}, \pm 2^{(2)}, \pm \sqrt{2}, \pm 2\sqrt{2}, \pm \sqrt{6}, 0^{(7)}\}$. Clearly, the signed graphs $S(\Gamma_1)$ and $S(\Gamma_2)$ are noncospectral and equienergetic. Similarly, the signed graphs $S(\Gamma_3)$ and $S(\Gamma_4)$ are noncospectral and equienergetic. Thus, by Corollary 4.11, we have

- (i) *the signed graphs $S_p(\Gamma_1)$ and $S_p(\Gamma_2)$ are noncospectral and equienergetic,*
- (ii) *the signed graphs $S_p^k(\Gamma_1)$ and $S_p^k(\Gamma_2)$ are noncospectral and equienergetic,*
- (iii) *the signed graphs $S_p(\Gamma_3)$ and $S_p(\Gamma_4)$ are noncospectral and equienergetic,*
- (iv) *the signed graphs $S_p^k(\Gamma_3)$ and $S_p^k(\Gamma_4)$ are noncospectral and equienergetic.*

Acknowledgements. This research is supported by SERB-DST research project number CRG/2020/000109. The research of Tahir Shamsher is supported by SRF financial assistance by Council of Scientific and Industrial Research (CSIR), New Delhi, India.

Data availability Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- [1] F. Belardo and S. K. Simic, On the Laplacian coefficients of signed graphs, *Linear Algebra Appl.* **475** (2015) 94-113.
- [2] F. Belardo, M. Brunetti, M. Cavaleri and A. Donno, Constructing cospectral signed graphs, *Linear Multilinear Algebra* (2019) <https://doi.org/10.1080/03081087.2019.1694483>.
- [3] F. Belardo, S. M. Cioaba, J. Koolen and J. Wang, Open problems in the spectral theory of signed graphs, *The Art of Discrete and Applied Mathematics* 1 (2018) P2.10 <https://doi.org/10.26493/2590-9770.1286.d7b>.
- [4] M. A. Bhat and S. Pirzada, On equienergetic signed graphs, *Discrete. Appl. Math.* **189** (2015) 1-7.
- [5] M. A. Bhat and S. Pirzada, Unicyclic signed graphs with minimal energy, *Discrete. Appl. Math.* **226** (2017) 32-39.
- [6] Z. L. Blazsik, J. Cummings, and W. H. Haemers, Cospectral regular graphs with and without a perfect matching, *Discrete Math.* **338**(2015) 199-201.
- [7] D. Cvetkovic, S. Simic, and P. Rowlinson, *An Introduction to the Theory of Graph Spectra*, Cambridge University Press, 2009.
- [8] K. A. Germina, S. Hameed and T. Zaslavsky, On products and line graphs of signed graphs, their eigenvalues and energy, *Linear Algebra Appl.* **435** (2010) 2432-2450.
- [9] A. Dehghan and A. H. Banihashemi, Cospectral bipartite graphs with the same degree sequences but with different number of large cycles, *Graphs Combin.* **35** (2019) 1673-1693.
- [10] S. Dutta, Constructing non-isomorphic signless Laplacian cospectral graphs, *Discrete Math.* **343** (2020) 111783.
- [11] C. D. Godsil and B. D. McKay, Constructing cospectral graphs, *Aequationes Math.* **25** (1982) 257-268.
- [12] R. A. Horn and C. R. Johnson, *Matrix Analysis*, Cambridge University Press, 23-Feb-1990.
- [13] F. Harary, On the notion of balanced in a signed graph, *Michigan Math. J.* **2** (1953) 143-146.

- [14] M. Haythorpe and A. Newcombe, Constructing families of cospectral regular graphs, *Combin. Probab. Comput.* **29** (2020) 664-671.
- [15] W. H. Haemers and E. Spence, Enumeration of cospectral graphs, *European J. Combin.* **25** (2004)199-211.
- [16] Y. Hou, J. Li and Y. Pan, On the Laplacian eigenvalues of signed graphs. *Linear Multilinear Algebra* **51** (2003) 21-30.
- [17] R. Merris, Large Families of Laplacian isospectral graphs, *Linear Multilinear Algebra*, **43** (1997) 201 -205.
- [18] T. Zaslavsky, A mathematical bibliography of signed and gain graphs and allied areas, *Electron. J. Combin., Dyn. Surv.* (1999) DS8, <http://www3.combinatorics.org/Surveys/ds8.pdf>.