

PRIME IDEALS IN CATEGORIES OF REPRESENTATIONS OF QUIVERS OF TYPE A

SHUNSUKE TADA

ABSTRACT. We study prime tensor ideals in tensor abelian categories of quiver representations. Specifically, we classify the prime tensor ideals in the category of representations of zigzag quivers (with bounded path length) whose vertex set is the set of integers. We show that prime tensor ideals in these categories are in canonical bijection with ultrafilters on the set of integers. We also discuss connections to tensor triangulated geometry and potential applications to the multiparameter persistent modules.

1. INTRODUCTION

We will say a quiver (directed graph) is *of type A* if its vertex set is the set of integers, and there is a unique arrow between i and $i + 1$ for each i like below

$$\dots \longleftrightarrow (-3) \longleftrightarrow (-2) \longleftrightarrow (-1) \longleftrightarrow 0 \longleftrightarrow 1 \longleftrightarrow 2 \longleftrightarrow 3 \longleftrightarrow \dots,$$

where \longleftrightarrow means that we can chose the direction of each arrow. Such quivers' representations are used to study persistent homology which has topological information about the structure of data sets, [16]. The representations help visualize the topological information via the “barcode”, [13]. The theory of persistent homology is used in many areas. For example, see [10] and [5] for applications to biology, [8] for an application to chemistry, and [1] for an application to mathematics.

The category of representations of a quiver is a tensor (monoidal) category, that is, an abelian category with a monoidal structure. In this category, we can consider prime \otimes -ideals (see Definition 2.15) as defined by Tobias J. Peter in [14]. This concept is related to the tensor triangulated geometry developed by Paul Balmer [4].

In this thesis, we classify the prime \otimes -ideals of the category $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ of point-wise finite-dimensional representations (Definition 2.10) of any quiver $A_{\mathbb{Z}}$ of type A which has bounded path length. These prime \otimes -ideals are related to the prime ideals of the ring $\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$. We will obtain the relationship between the prime \otimes -ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and the prime ideals of $\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$ using the power set $\mathcal{P}(\mathbb{Z})$ of the set of integers \mathbb{Z} . We will obtain following theorem.

Theorem 1.1. *Let $A_{\mathbb{Z}}$ be a quiver of type A with bounded path length. That is, there exists $n \geq 1$ in \mathbb{N} such that any path in $A_{\mathbb{Z}}$ has length smaller than n . Then, there is a one-to-one correspondence between each of the four sets*

- (1) *prime tensor ideals of the category $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$,*
- (2) *prime ideals of the ring $\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$,*
- (3) *prime ideals of boolean algebra $\mathcal{P}(\mathbb{Z})$ and*

Key words and phrases. quiver, quiver representation, prime tensor ideal, tensor triangulated geometry.

(4) *ultrafilters of the boolean algebra $\mathcal{P}(\mathbb{Z})$.*

In this theorem, we need the condition

bounded path length

to have above relations. Without it, we could not have the classification of the prime \otimes -ideals of the category $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. For example, if the quiver has an infinite length path like

$$\cdots \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \cdots,$$

then we will have a inclusion map from the prime ideals of $\mathcal{P}(\mathbb{Z})$ to the prime \otimes -ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$, but the map is not surjective. We will prove this in Corollary 3.24.

Outline. This paper is organized as follows. In Section 2, we will see the basics of the quiver representations and lattice theory. In Section 3, we see the classification of prime \otimes -ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and some topological properties of the spectrum of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. In Section 4, we will discuss the relation to persistence homology and tensor triangulated geometry.

Acknowledgements. I would like to thank Shane Kelly for suggesting me the problem and discussing it.

2. PRELIMINARIES

2.1. Quiver and its representation. Here are some basics about quivers and their representations. See [2] for more details. Throughout, K will be a field.

Definition 2.1 (Quiver).

A *quiver* $Q = (Q_0, Q_1, s, t)$ is a quadruple consisting of two sets: Q_0 called the set of vertices and Q_1 called the set of arrows, and two maps $s, t : Q_1 \rightarrow Q_0$. For every arrow $\alpha \in Q_1$, $s(\alpha) \in Q_0$ and $t(\alpha) \in Q_0$ are called *source* and *target* of α , respectively. We usually denote the arrow α by $\alpha : s(\alpha) \rightarrow t(\alpha)$.

Definition 2.2 (Path).

Let $Q = (Q_0, Q_1, s, t)$ be a quiver with $a, b \in Q_0$. A path from source a to target b of length $l \geq 1$ is a sequence of arrows

$$(a \mid \alpha_1, \dots, \alpha_l \mid b)$$

such that $s(\alpha_1) = a$, $t(\alpha_l) = b$ and $t(\alpha_i) = s(\alpha_{i+1})$ for any $i \in \{1, \dots, l-1\}$. If there is no confusion, we write a path $(a \mid \alpha_1, \dots, \alpha_l \mid b)$ as

$$a \rightarrow b.$$

Definition 2.3 (subquiver / convex subquiver / connected quiver).

A *subquiver* of a quiver $Q = (Q_0, Q_1, s, t)$ is a quiver $Q' = (Q'_0, Q'_1, s', t')$ such that $Q'_0 \subset Q_0$, $Q'_1 \subset Q_1$ and the restrictions of the maps s and t to Q'_1 are respectively equal to s', t' .

If the arrows of the subquiver are

$$Q'_1 = \{ \alpha \in Q_1 : s(\alpha) \in Q'_0 \text{ and } t(\alpha) \in Q'_0 \},$$

then the subquiver $Q' = (Q'_0, Q'_1, s', t')$ is called a *full subquiver* of Q . A subquiver is uniquely determined by its sets of vertices.

The full subquiver Q' of a quiver Q is defined to be *convex* in Q if, for any path $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_t$ in Q with $x_0, x_t \in Q'$, we have $x_i \in Q'$ for all $i \in \{0, \dots, t\}$.

If the underlying graph of the quiver Q is connected, we call the quiver *connected*.

Example 2.4. Let $Q = (Q_0, Q_1, s, t)$ be a quiver of the form

$$\cdots \longleftrightarrow (-3) \longleftrightarrow (-2) \longleftrightarrow (-1) \longleftrightarrow 0 \longleftrightarrow 1 \longleftrightarrow 2 \longleftrightarrow 3 \longleftrightarrow \cdots$$

where \longleftrightarrow means that we can chose the direction of each arrows. For example,

$$\cdots \longleftarrow (-3) \longrightarrow (-2) \longrightarrow (-1) \longrightarrow 0 \longleftarrow 1 \longleftarrow 2 \longleftarrow 3 \longleftarrow \cdots$$

This quiver is a connected quiver and there is a path from source -3 to target 0 . Moreover, the subquiver

$$(-3) \longrightarrow (-2) \longrightarrow (-1) \quad 1 \longleftarrow 2 \longleftarrow 3$$

with vertices $\{\pm 1, \pm 2, \pm 3\}$ is not connected.

Definition 2.5 (Type A).

We will say a quiver is *of type A* if its vertex set is the set of integers, and there is a unique arrow between i and $i + 1$ for each i .

Definition 2.6 (Sink and source).

Let $Q = (Q_0, Q_1, s, t)$ be quiver. A vertex a in Q_0 is called a *sink* if there are no arrows whose source is a . A vertex a in Q_0 is called a *source* if there are no arrow whose target is a .

Definition 2.7 (Category of representations of quiver).

Let $Q = (Q_0, Q_1, s, t)$ be a quiver. The category $\text{Rep}(Q)$ of representations of Q is defined as follows. An object is pair consisting of a family of K -vector spaces indexed by Q_0 , and a family of K -homomorphisms indexed by Q_1 :

$$V := ((V_a)_{a \in Q_0}, (V_\alpha : V_{s(\alpha)} \rightarrow V_{t(\alpha)})_{\alpha \in Q_1}).$$

Let $V = ((V_a)_{a \in Q_0}, (V_\alpha)_{\alpha \in Q_1})$ and $W = ((W_a)_{a \in Q_0}, (W_\alpha)_{\alpha \in Q_1})$ be representations. A morphism in $\text{Rep}(Q)$ from V to W is a family of K -homomorphisms $(f_a : V_a \rightarrow W_a)_{a \in Q_0}$ satisfying

$$f_{t(\alpha)} \circ V_\alpha = W_\alpha \circ f_{s(\alpha)}$$

for every arrow $\alpha \in Q_1$.

Definition 2.8 (Sink and source).

Let $Q = (Q_0, Q_1, s, t)$ be quiver. A vertex a in Q_0 is called a *sink* if there are no arrows whose source is a . A vertex a in Q_0 is called a *source* if there are no arrow whose target is a .

Definition 2.9 (Interval representation).

Let $Q = (Q_0, Q_1, s, t)$ be a quiver, and let $I = (I_0, I_1, s', t')$ be a subquiver of Q . Define $K_I \in \text{Rep}(Q)$ by

$$(K_I)_a := \begin{cases} K & (a \in I_0) \\ 0 & (\text{otherwise}), \end{cases}$$

$$(K_I)_\alpha := \begin{cases} id_K & (\alpha \in I_1) \\ 0 & (\text{otherwise}). \end{cases}$$

If I is connected and convex (and therefore also full), then K_I is called an *interval representation*.

Definition 2.10 (Point-wise finite-dimensional representations).

Let Q be a quiver. A representation V of $\text{Rep}(Q)$ is said to be *point-wise finite-dimensional* if the dimension of vector space $\dim_K(V_a)$ is finite for every vertex a in Q .

We write $\text{Rep}^{\text{pfd}}(Q)$ for the full subcategory of $\text{Rep}(Q)$ whose objects are point-wise finite-dimensional representations.

Definition 2.11 (Direct sum in the category of representations).

Let $Q = (Q_0, Q_1, s, t)$ be a quiver and

$$V := ((V_a)_{a \in Q_0}, (V_\alpha : V_{s(\alpha)} \rightarrow V_{t(\alpha)})_{\alpha \in Q_1}) \text{ and}$$

$$W := ((W_a)_{a \in Q_0}, (W_\alpha : W_{s(\alpha)} \rightarrow W_{t(\alpha)})_{\alpha \in Q_1})$$

be representations. Then the *direct sum* of V and W in $\text{Rep}(Q)$ is defined by

$$V \oplus W := ((V_a \oplus W_a)_{a \in Q_0}, (V_\alpha \oplus W_\alpha : V_{s(\alpha)} \oplus W_{s(\alpha)} \rightarrow V_{t(\alpha)} \oplus W_{t(\alpha)})_{\alpha \in Q_1}).$$

To show that any representation in $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ is interval decomposable, we consider continuous quivers whose vertices are \mathbb{R} . Such quivers are defined in [9]. We will see representations of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ from the viewpoint of continuous quivers and we use the interval decomposition theorem for continuous quivers which has recently been proven in the paper [9] to show the proposition.

Proposition 2.12. Let $A_{\mathbb{Z}}$ be a quiver of type A . Then every point-wise finite-dimensional representation of $\text{Rep}(A_{\mathbb{Z}})$ is direct sum of interval representations.

Proof. The flow of our proof is as follows. To show that any representation V of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ is direct sum of interval representations, we will see V as the restriction of a representation of a continuous quiver. Then we apply the decomposition theorem for the representations of the continuous quiver and we will obtain the proof of the proposition. At first we will prepare subsets of \mathbb{Z} which is helpful to see the representation of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ as the representation of the continuous quiver.

Let S' be the set of sinks and sources of $A_{\mathbb{Z}}$. For example, if the quiver $A_{\mathbb{Z}}$ is one direction (we do not always draw all vertices)

$$\cdots \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \cdots,$$

then the set S' is empty set and if the quiver $A_{\mathbb{Z}}$ is

$$\cdots \longrightarrow \longrightarrow \longrightarrow \longrightarrow 0 \longleftarrow 1 \longrightarrow 2 \longleftarrow 3 \longrightarrow 4 \longleftarrow \cdots,$$

then the set S' is $\mathbb{Z}_{\geq 0}$. In this case, the $S' = \mathbb{Z}_{\geq 0}$ is unbounded above and bounded below.

Next, define the subset S of $\mathbb{Z} \cup \{-\infty, +\infty\}$ by

$$S := \begin{cases} S' & \text{(if } S' \text{ is not bounded)} \\ S' \cup \{+\infty\} & \text{(if } S' \text{ is bounded above and unbounded below)} \\ S' \cup \{-\infty\} & \text{(if } S' \text{ is bounded below and unbounded above)} \\ S' \cup \{+\infty, -\infty\} & \text{(if } S' \text{ is bounded)} \end{cases}$$

To show the proposition, we consider two different cases

- (1) $A_{\mathbb{Z}}$ does not have sources and sinks, i.e., $S' = \emptyset$,
- (2) $A_{\mathbb{Z}}$ contains sources or sinks, i.e., $S' \neq \emptyset$.

In the case of (1), the quiver $A_{\mathbb{Z}}$ can be one of the following quivers

$$\cdots \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \cdots ,$$

$$\cdots \longleftarrow \longleftarrow \longleftarrow \longleftarrow \longleftarrow \longleftarrow \cdots .$$

Now we can assume that $A_{\mathbb{Z}}$ is

$$\cdots \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \cdots$$

(we can assume the other quiver, but the proof is similar). Consider the quiver $\overline{A_{\mathbb{Z}}}$ whose vertices are \mathbb{R} such that arrows are defined by

$$a \longrightarrow b : \iff a \leq b$$

for $a, b \in \mathbb{R}$. For any representation V in $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$, we now show that the representation V is an interval decomposable representation using the quiver $\overline{A_{\mathbb{Z}}}$. Let \overline{V} be a representation of $\text{Rep}^{\text{pfd}}(\overline{A_{\mathbb{Z}}})$ such that for every $a \in \mathbb{R}$,

$$\overline{V}_a := V_{[a]}$$

where $[a]$ is the largest integer less than or equal to a (floor function of a) and arrows are defined to be

$$\overline{V}_{b \rightarrow c} := V_{[b] \rightarrow [c]},$$

where b and c is in \mathbb{R} . It is known that any point-wise finite-dimensional representation of $\overline{A_{\mathbb{Z}}}$ is direct sum of interval representations, [9, Theorem B]. Thus there is an isomorphism of $\text{Rep}^{\text{pfd}}(\overline{A_{\mathbb{Z}}})$

$$(f_a)_{a \in \mathbb{R}} : \overline{V} \rightarrow \bigoplus_I K_I$$

for intervals $I \subset \mathbb{R}$. Then we have an isomorphism of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$

$$(f_a)_{a \in \mathbb{Z}} : V \rightarrow \bigoplus_I K_{I \cap \mathbb{Z}}.$$

Hence, representations V in $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ are interval decomposable.

In the case of (2), assume that there is a source in $A_{\mathbb{Z}}$ and take a source arbitrarily and let the source be s_0 . If there is a sink larger than s_0 , let s_1 be the smallest sink larger than s_0 . If there is not such a sink, then let s_1 be $+\infty$. If $s_1 \neq +\infty$, define $s_2 \in S$ as the smallest source larger than s_1 . If there is not such a source, define s_2 be $+\infty$. We can inductively define s_3, s_4, \dots . If $s_n = +\infty$ for some $n \in \mathbb{Z}$, we define $s_{n \pm m} := +\infty$ for any $m \in \mathbb{Z}_{\geq 0}$. Similarly, we can define s_{-1} to the largest source smaller than s_0 , and inductively we can define s_{-2}, s_{-3}, \dots

Using the set S , we define a poset (\mathbb{R}, \preceq) called a *continuous quiver of type A* in [9]. Define intervals on \mathbb{R} by

$$\langle s_n, s_{n+1} \rangle := \begin{cases} [s_n, s_{n+1}] & (s_n, s_{n+1} \in S') \\ [s_n, s_{n+1}) & (s_n \in S', s_{n+1} = +\infty) \\ (s_n, s_{n+1}] & (s_n = -\infty, s_{n+1} \in S') \end{cases}$$

and define the order on each $\langle s_n, s_{n+1} \rangle$ by

$$a \preceq b := \begin{cases} a \leq b \text{ and } s_n \text{ is a source} \\ b \leq a \text{ and } s_n \text{ is a sink} \end{cases}$$

where s_n and s_{n+1} are in S . We obtain a poset (\mathbb{R}, \preceq) and it can be seen as a quiver if arrows are defined by

$$a \longrightarrow b : \iff a \preceq b.$$

We write $\overline{A_{\mathbb{Z}}}$ for the continuous quiver (\mathbb{R}, \preceq) . To put it roughly, the direction of arrows in $\overline{A_{\mathbb{Z}}}$ are same as the direction of arrows in $A_{\mathbb{Z}}$.

For example, if the quiver $A_{\mathbb{Z}}$ is

$$\cdots \longrightarrow \longrightarrow \longrightarrow \longrightarrow -1 \longleftarrow 0 \longrightarrow \longleftarrow \longrightarrow \longleftarrow \cdots,$$

then we have $0 \preceq -\frac{1}{2}$ and therefore there is an arrow from 0 to $-\frac{1}{2}$ in $\overline{A_{\mathbb{Z}}}$.

Finally, we show that any representation $V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ can be decomposed into interval representations. For any $a \in \mathbb{R}$, there exist $k \in \mathbb{Z}$ such that a is in $[k, k+1]$. Define a representation \overline{V} of $\overline{A_{\mathbb{Z}}}$ by

$$\overline{V}_a := \begin{cases} V_k & (\text{if } k \preceq k+1) \\ V_{k+1} & (\text{if } k+1 \preceq k) \end{cases}$$

and for any b and c in \mathbb{R} with $b \preceq c$, there exists an interval $\langle s_n, s_{n+1} \rangle$ such that b and c are in $\langle s_n, s_{n+1} \rangle$. Then we can define

$$\overline{V}_{b \rightarrow c} := \begin{cases} V_{\lfloor b \rfloor \rightarrow \lfloor c \rfloor} & (\text{if } s_n \text{ is a source}) \\ V_{\lceil b \rceil \rightarrow \lceil c \rceil} & (\text{if } s_n \text{ is a sink}) \end{cases}$$

where $\lfloor x \rfloor$ is the largest integer less than or equal to x (floor function of x) and $\lceil x \rceil$ is the smallest integer more than or equal to x (ceiling function of x). Here, we write

$$V_{l \rightarrow l'} := \begin{cases} V_{l'-1 \rightarrow l'} \circ \cdots \circ V_{l \rightarrow l+1} & (l < l') \\ id_{V_l} & (l = l') \\ V_{l+1 \rightarrow l'} \circ \cdots \circ V_{l \rightarrow l-1} & (l' < l) \end{cases}$$

if there is a path from l to l' in $A_{\mathbb{Z}}$.

It is known that any point-wise finite-dimensional representation is direct sum of interval representations, [9, Theorem B]. Thus there is an isomorphism of $\text{Rep}^{\text{pfd}}(\overline{A_{\mathbb{Z}}})$

$$(f_a)_{a \in \mathbb{R}} : \overline{V} \rightarrow \bigoplus_I K_I$$

for intervals $I \subset \mathbb{R}$. Then we have an isomorphism of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$

$$(f_a)_{a \in \mathbb{Z}} : V \rightarrow \bigoplus_I K_{I \cap \mathbb{Z}}.$$

Hence, for any representations in $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ are interval decomposable. \square

Definition 2.13 (Tensor category [11]).

A *tensor category* (or *monoidal category*) is $\nu = (\nu_0, \otimes, I, a, \ell, r)$ consists of a category ν_0 , a functor $\otimes : \nu_0 \times \nu_0 \rightarrow \nu_0$, an object I of ν_0 and natural isomorphisms $a_{XYZ} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$, $\ell_X : I \otimes X \rightarrow X$, $r_X : X \otimes I \rightarrow X$, subject to two coherence axioms expressing the commutativity of the following diagrams:

$$\begin{array}{ccc} ((W \otimes X) \otimes Y) \otimes Z & \xrightarrow{a} & (W \otimes X) \otimes (Y \otimes Z) & \xrightarrow{a} & W \otimes (X \otimes (Y \otimes Z)) \\ & \searrow^{a \otimes 1} & & & \nearrow_{1 \otimes a} \\ (W \otimes (X \otimes Y)) \otimes Z & \xrightarrow{a} & W \otimes ((X \otimes Y) \otimes Z) & & \end{array}$$

and

$$\begin{array}{ccc} (X \otimes I) \otimes Y & \xrightarrow{a} & X \otimes (I \otimes Y) \\ r \otimes I \searrow & & \swarrow r \otimes I \\ & X \otimes Y & \end{array}$$

Example 2.14.

The tuple $(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}), \otimes, K_{A_{\mathbb{Z}}}, a, \ell, r)$ is a tensor category. Let V, V', W and W' be objects of $\text{Rep}(A_{\mathbb{Z}})$. We define a functor

$$\otimes : \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) \times \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) \rightarrow \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$$

by

$$\otimes(V, W) := ((V_n \otimes_K W_n)_{n \in (A_{\mathbb{Z}})_0}, (V_\alpha \otimes_K W_\alpha)_{\alpha \in (A_{\mathbb{Z}})_1})$$

for objects and

$$\begin{aligned} \otimes((f, g) : (V, W) \rightarrow (V', W')) \\ := (f_n \otimes g_n)_{n \in (A_{\mathbb{Z}})_0} : ((V_n \otimes_K W_n)_{n \in (A_{\mathbb{Z}})_0} \rightarrow ((V'_n \otimes_K W'_n)_{n \in (A_{\mathbb{Z}})_0} \end{aligned}$$

for arrows. Then, we have a natural isomorphisms $a_{XYZ} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$, $\ell_X : K_{A_{\mathbb{Z}}} \otimes X \rightarrow X$ and $r_X : X \otimes K_{A_{\mathbb{Z}}} \rightarrow X$ for any X, Y and Z in $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. These natural isomorphisms are subject to two coherence axioms of tensor category.

Definition 2.15 (Prime tensor ideal [14]).

Let \mathcal{A} be an abelian category. A full subcategory \mathcal{M} of \mathcal{A} is called *coherent* if

- (1) it contains the zero object,
- (2) it is closed under extensions and
- (3) kernels and cokernels in \mathcal{A} of morphisms between objects of \mathcal{M} are in \mathcal{M} .

Suppose now that \mathcal{A} is a tensor abelian category. A (*coherent*) *tensor ideal* of \mathcal{A} is a coherent subcategory $\mathcal{M} \subset \mathcal{A}$ such that $\mathcal{M} \otimes \mathcal{A} \subset \mathcal{M}$. A proper ideal $\mathcal{M} \subsetneq \mathcal{A}$ is called a (*coherent*) *prime ideal* if $A \otimes B \in \mathcal{M}$ implies $A \in \mathcal{M}$ or $B \in \mathcal{M}$.

We write the set of all the prime tensor ideals by $\text{Spc}(\mathcal{A})$.

Example 2.16.

Let Q be a quiver, and consider the tensor category $(\text{Rep}^{\text{pfd}}(Q), \otimes, K_Q, a, \ell, r)$ and let \mathcal{M} be the full subcategory whose objects are $\{V \in Q : V_n = 0\}$ for some vertex $n \in Q_0$. We will show that the full subcategory \mathcal{M} is a prime \otimes -ideal.

First, we show that \mathcal{M} is closed under kernels and cokernels. Let V and W be representations in \mathcal{M} and $(f_n : V_n \rightarrow W_n)_{n \in Q}$ be a morphism from a representation V to W . Since V_n and W_n are isomorphic to zero, the kernel and the cokernel of $f_n : V_n \rightarrow W_n$ is isomorphic to zero. Thus, $\text{Ker}((f_n)_{n \in A_{\mathbb{Z}}})$ and $\text{Coker}((f_n)_{n \in A_{\mathbb{Z}}})$ is in \mathcal{M} . So the full subcategory \mathcal{M} is closed under kernels and cokernels.

Next, we show that \mathcal{M} is closed under extension. Take an exact sequence

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$$

where V' and V'' are objects of \mathcal{M} . Then, there is an exact sequence of K -homomorphisms

$$0 \rightarrow V'_n \rightarrow V_n \rightarrow V''_n \rightarrow 0.$$

Since the representations V' and V'' are in \mathcal{M} , we have $V'_n = 0 = V''_n$, so V_n is isomorphic to 0. Thus the representation V is in \mathcal{M} . So the full subcategory \mathcal{M} is closed under extensions.

Then, for any representations $V \in \text{Rep}^{\text{pfd}}(Q)$ and $W \in \mathcal{M}$, we have $(V \otimes W)_n = V_n \otimes W_n = 0$. Thus \mathcal{M} is a \otimes -ideal.

Finally, we show that \mathcal{M} is a prime \otimes -ideal. Let V and W be the representations of $\text{Rep}^{\text{pfd}}(Q)$. If $V \otimes W$ is in \mathcal{M} , then we have $0 = (V \otimes W)_n = V_n \otimes W_n$. Thus V_n or W_n are isomorphic to 0 and therefore V or W are in \mathcal{M} .

2.2. Boolean rings and Boolean algebras. Here are some basics about Boolean rings and Boolean algebras. See [3] for more details.

Definition 2.17 (Boolean ring).

A ring B with an identity is called a *Boolean ring* if $x^2 = x$ for all $x \in B$.

Example 2.18.

The ring $\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$ is a Boolean ring.

Proposition 2.19.

Any Boolean ring is a commutative ring, every prime ideal of a Boolean ring is a maximal ideal, and every residue field is isomorphic to $\mathbb{Z}/2$.

Proof. Let B be a Boolean ring. To show that the ring B is commutative, we take two elements x and y of B arbitrarily and show $xy = yx$ holds. Since the ring B is a Boolean ring, the equation $0 = (x+y)^2 - (x+y) = xy+yx = xy+(-yx)^2 = xy-yx$ holds. Thus we obtain $xy = yx$ and therefore the Boolean ring B is commutative.

Next, we will show that prime ideals of a Boolean ring are maximal ideals. Let P be a prime ideal of the Boolean ring B . It is enough to show that the quotient ring B/P is isomorphic to $\mathbb{Z}/2$. Take a nonzero $[x]$ in B/P where $[x]$ is an equivalence class of $x \in B$. Since $[x(x-1)] = [x^2-x] = [x-x] = [0]$ holds, we have $x(x-1) \in P$. Thus x or $(x-1)$ is in P because P is a prime ideal. If x is in P , then $[x] = [0]$ and it contradicts to that $[x]$ is not zero. Therefore, $(x-1)$ is in P and it implies $[x]-[1] = [x-1] = [0]$. We obtain $[x] = [1]$. Hence B/P is $\mathbb{Z}/2$, and P is a maximal ideal. \square

Lemma 2.20. *Let B be a Boolean ring. We can define the partially order on B by*

$$a \leq b : \iff a(b+1) = 0.$$

Thus the Boolean ring B has a structure of poset.

Proof. At first, we show the reflexivity. For any $a \in B$, we have $a(a+1) = a^2 + a = a + a = 0$. we can show that $a + a = 0$ by calculating $-a = (-a)^2 = a^2 = a$.

Next, we show the transitivity. If $a \leq b$ and $b \leq c$, then we have $a(b+1) = 0$ and $b(c+1) = 0$. Thus $a(c+1) = ac + a = ac - ab = ac - a(-bc) = ac(1+b) = ca(b+1) = 0$.

Finally, we will show the antisymmetry. If $a \leq b$ and $b \leq a$, then we have $a(b+1) = 0$ and $b(a+1) = 0$. Thus we obtain $a = -ab = b$.

Thus we defined the partially order on the Boolean ring B . \square

Definition 2.21 (Boolean algebras).

Let B be a poset. The *Boolean algebra* is defined to be a tuple $(B; \vee, \wedge, ', 0, 1)$ such that

- (i) $x \vee y := \sup\{x, y\}$ and $x \wedge y := \inf\{x, y\}$ exist for all $x, y \in B$,
- (ii) $(B; \vee, \wedge)$ is a distributive, i.e.,

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c),$$

- (iii) 0 is the minimal element, and 1 is the maximal element of B ,
- (iv) every element x admits a unique complement, i.e., an element satisfying $x \wedge x' = 0$ and $x \vee x' = 1$.

We introduce some properties about Boolean algebras which are helpful to show Lemma 2.25.

Lemma 2.22. *Let B be a Boolean algebra. Then*

- (i) $0' = 1$ and $1' = 0$,
- (ii) $a'' = a$ for all $a \in B$,
- (iii) de Morgan's laws hold: for all $a, b \in B$,

$$(a \vee b)' = a' \wedge b' \text{ and } (a \wedge b)' = a' \vee b',$$

- (iv) $a \wedge b' = 0$ if and only if $a \leq b$ for all $a, b \in B$.

Proof. See Lemma 4.15 of [3]. □

Corollary 2.23. *Let B be a Boolean algebra. Then $a \leq b$ if and only if $b' \leq a'$ for all $a, b \in B$.*

Proof. By Lemma 2.22 (iv), we have $a \leq b \Leftrightarrow a \wedge b' = 0 \Leftrightarrow b' \wedge a = 0 \Leftrightarrow b' \leq a'$. □

Definition 2.24 (Homomorphisms of Boolean algebras).

Given Boolean algebras B and C , a map $f : B \rightarrow C$ is a Boolean homomorphism if

- (i) f is join-preserving and meet-preserving, that is, for all $a, b \in B$,

$$f(a \vee b) = f(a) \vee f(b) \text{ and } f(a \wedge b) = f(a) \wedge f(b).$$

- (ii) f preserves 0, 1 and $(-)'$, that is, we have $f(0) = 0$, $f(1) = 1$, and $f(a') = (f(a))'$ for all $a \in B$.

A bijective Boolean homomorphism is called a Boolean isomorphism.

Lemma 2.25.

Let $(B; \vee, \wedge, ', 0, 1)$ be a Boolean algebra and B^{op} be the poset (with $a \leq_{B^{op}} b$ iff $b \leq_B a$). Then B^{op} is a Boolean algebra, and $(-)' : B \rightarrow B^{op}$ is an isomorphism of Boolean algebras.

Proof. At first, we will show that $B^{op} := (B^{op}; \vee^{op}, \wedge^{op}, ', 0', 1')$ is a Boolean algebra where \vee^{op} and \wedge^{op} are defined by

$$a \vee^{op} b := a \wedge b, \quad a \wedge^{op} b := a \vee b$$

for all $a, b \in B^{op}$.

Since the Boolean algebra B has $x \wedge y := \inf\{x, y\}$ and $x \vee y := \sup\{x, y\}$, the poset B^{op} has $x \wedge^{op} y = \sup\{x, y\}$ and $x \vee^{op} y = \inf\{x, y\}$ for all $a, b \in B^{op}$. Moreover $(B; \vee^{op}, \wedge^{op})$ is distributive. Indeed, we have the equation

$$\begin{aligned}
a \wedge^{op} (b \vee^{op} c) &= (a \wedge^{op} (b \vee^{op} c))'' \\
&= (a \vee (b \wedge c))'' \\
&= (a' \wedge (b \wedge c)')' \\
&= (a' \wedge (b' \vee c'))' \\
&= ((a' \wedge b') \vee (a' \wedge c'))' \\
&= ((a \vee b)' \vee (a \vee c)')' \\
&= (a \vee b)'' \wedge (a \vee c)'' \\
&= (a \vee b) \wedge (a \vee c) \\
&= (a \wedge^{op} b) \vee^{op} (a \wedge^{op} c).
\end{aligned}$$

The element $1'$ is maximal and $0'$ is the minimal element in the poset (B^{op}, \leq^{op}) , and for every $x \in B^{op}$ we have $x \wedge^{op} x' = x \vee x' = 1 = 0'$ and $x \vee^{op} x' = x \wedge x' = 0 = 1'$. Thus, the poset B^{op} is a Boolean algebra.

Next, we will show that the map $(-)' : B \rightarrow B^{op}$ is an isomorphism of the Boolean algebras. The map $(-)'$ preserves join and meet, that is, for all $a, b \in B$ we have,

$$(a \wedge b)' = (a' \wedge^{op} b') \text{ and } (a \vee b)' = (a' \vee^{op} b').$$

Moreover, the map $(-)'$ preserves $0, 1$ and $(-)'$ i.e., the two elements $0', 1'$ are the minimal and the maximal elements of B^{op} , and we have $(a')' = a = (a)''$. Since the map $(-)'$ is bijective, it is an isomorphism of Boolean algebras. \square

Example 2.26.

Let us write $\mathcal{P}(\mathbb{Z})$ for the power set of \mathbb{Z} . Then $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ is a Boolean algebra where c is the operation of taking the complement of an element.

Definition 2.27 (Ideal/prime ideal of Boolean algebra).

Let B be a Boolean algebra. A non-empty subset J of B is called an *ideal* if

- (i) $a, b \in J$ implies $a \vee b \in J$,
- (ii) $a \in B, b \in J$ and $a \leq b$ imply $a \in J$.

An ideal is *proper* if $J \subsetneq B$. A proper ideal J of B is said to be a *prime ideal* if $a, b \in B$ and $a \wedge b \in J$ imply $a \in J$ or $b \in J$. We denote the sets of prime ideals by $\text{Spc}(B)$.

Example 2.28.

Let $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ be the Boolean algebra associated to the power set $\mathcal{P}(\mathbb{Z})$ of the set of integers. Then $\mathcal{P}_a := \{I \in \mathcal{P}(\mathbb{Z}) : a \notin I\}$ is the prime ideal for every $a \in \mathbb{Z}$.

Remark 2.29 ([3] Theorem.10.12).

Prime ideals in a Boolean algebra are maximal ideals: We will see that there is a one-to-one inclusion preserving correspondence between the set of prime ideals of a Boolean algebra and the set of prime ideals of its associated Boolean ring in Proposition 2.35 (iv). Then by Proposition 2.19, it follows that prime ideals in Boolean algebras are maximal ideals.

Definition 2.30 (Filter/prime filter of a Boolean algebra).

Let B be a Boolean algebra. A non-empty subset \mathcal{F} of B is called a *filter* if the following are satisfied.

- (i) If $a, b \in \mathcal{F}$, then $a \wedge b \in \mathcal{F}$.
- (ii) If $a \in B, b \in \mathcal{F}$ and $b \leq a$, then $a \in \mathcal{F}$.

A filter is *proper* if $\mathcal{F} \subsetneq B$. A proper filter \mathcal{F} of B is said to be *prime* if $a, b \in B$ and $a \vee b \in \mathcal{F}$ imply $a \in \mathcal{F}$ or $b \in \mathcal{F}$. We denote the sets of prime filters by $\text{Ult}(B)$.

Remark 2.31.

Note, a filter (resp. prime filter) of B is exactly the same thing as a ideal (resp. prime ideal) of B^{op} , cf. Lemma 2.25.

Example 2.32.

Let $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ be the Boolean algebra associated to the power set of the set of integers. Then $\mathcal{F}_a := \{I \in \mathcal{P}(\mathbb{Z}) : a \in I\}$ is a prime filter for any $a \in \mathbb{Z}$.

Lemma 2.33. *Let B be a Boolean algebra. Then, there is a bijection between $\text{Spc}(B)$ and $\text{Ult}(B)$ given by*

$$\text{Spc}(B) \rightarrow \text{Ult}(B), \mathcal{Q} \mapsto \mathcal{Q}'$$

where $\mathcal{Q}' := \{a \in B : a' \in \mathcal{Q}\}$.

Proof. By Lemma 2.25, complement $(-)' : B \rightarrow B^{op}$ is an isomorphism of Boolean algebras, and by the definition, prime ideals of B are prime filters of B^{op} . So

$$\text{Spc}(B) = \text{Ult}(B^{op}) \xrightarrow{\sim} \text{Ult}(B).$$

□

Corollary 2.34.

Let $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ be the Boolean algebra associated to the power set of the set of integers. Then, there is a bijection between $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ and $\text{Ult}(\mathcal{P}(\mathbb{Z}))$ given by

$$\text{Spc}(\mathcal{P}(\mathbb{Z})) \rightarrow \text{Ult}(\mathcal{P}(\mathbb{Z})), \mathcal{Q} \mapsto \mathcal{Q}^c$$

where $\mathcal{Q}^c := \{I \in \mathcal{P}(\mathbb{Z}) : I^c \in \mathcal{Q}\}$.

Proof. It follows immediately from Lemma 2.33. □

Proposition 2.35.

- (i) Let B be a Boolean algebra and define $+$ and \cdot on B by

$$x + y := (x \wedge y') \vee (x' \wedge y), \quad x \cdot y := x \wedge y.$$

Then $(B; +, \cdot)$ is a Boolean ring.

- (ii) Let B be a Boolean ring and define \wedge and \vee on B by

$$x \vee y := x + y + xy, \quad x \wedge y := xy, \quad x' := 1 + x.$$

Then $(B; \wedge, \vee, ', 0, 1)$ is a Boolean algebra.

- (iii) The correspondence between Boolean algebras and Boolean rings established in (i) and (ii) is a bijective one.
- (iv) There is a one-to-one inclusion preserving correspondence between prime ideals of a Boolean algebra and the prime ideals of the associated Boolean ring.

Proof. For the proof of (i) \sim (iii) see [3, Ex.4.27].

(iv) Take an arbitrary prime ideal \mathcal{Q} of the Boolean algebra $(B; \vee, \wedge, ', 0, 1)$. We will show that the \mathcal{Q} is a prime ideal of the Boolean ring defined in Proposition 2.35(i).

For any $x, y \in \mathcal{Q}$, the sum $x + y := (x \wedge y') \vee (x' \wedge y)$ is in \mathcal{Q} because $x \wedge y' \in \mathcal{Q}$ and $x' \wedge y \in \mathcal{Q}$ for $x, y \in \mathcal{Q}$ (recall that \wedge is defined as \inf) and ideals of Boolean algebras are closed under join and \leq . For any $x \in B$ and for any $y \in \mathcal{Q}$, the product $x \cdot y := x \wedge y$ is in \mathcal{Q} . Thus \mathcal{Q} is an ideal of the Boolean ring. If $x \cdot y := x \wedge y \in \mathcal{Q}$, then x or y is in \mathcal{Q} because \mathcal{Q} is a prime ideal of the Boolean algebra. Therefore \mathcal{Q} is a prime ideal of the Boolean ring.

Let \mathcal{P} be a prime ideal of the Boolean ring associated to the Boolean algebra B . We will show that \mathcal{P} is a prime ideal of the Boolean algebra.

For any $a, b \in \mathcal{P}$, we have $a \vee b := a + b + ab \in \mathcal{P}$. Also, for any $a \in B$ and $b \in \mathcal{P}$, if $a \leq b$ then we have $a(b+1) = ab' = 0$. Thus $a \in \mathcal{P}$. So \mathcal{P} is an ideal of the Boolean algebra. Finally, we will show that the \mathcal{P} is a prime ideal as the Boolean algebra. If $a \wedge b \in \mathcal{P}$ for $a, b \in B$, we have $a \wedge b := ab \in \mathcal{P}$. Since \mathcal{P} is the prime ideal of the Boolean ring B , we have $a \in \mathcal{P}$ or $b \in \mathcal{P}$. Thus we have shown that \mathcal{P} is a prime ideal as the Boolean algebra. \square

By this proposition, we can see Boolean algebras and Boolean rings are equivalent.

Example 2.36.

Consider the Boolean algebra $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ corresponding to the power set of the set of integers. The Boolean ring corresponding to $\mathcal{P}(\mathbb{Z})$ is isomorphic to $\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$ via the map

$$\phi : \mathcal{P}(\mathbb{Z}) \rightarrow \prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}), \quad I \mapsto (a_i)_{i \in \mathbb{Z}}$$

where $a_i = 1 + 2\mathbb{Z}$ if $i \in I$ otherwise $a_i = 0 + 2\mathbb{Z}$. Note ϕ is a bijective ring homomorphism, i.e., a ring isomorphism. Here, for each $I, J \in \mathcal{P}(\mathbb{Z})$, the sum $I + J := (I \cup J) \setminus (I \cap J)$ and the product $I \cdot J := I \cap J$ on the left are those defined in Proposition 2.35(i). By Proposition 2.35(iv), The map ϕ gives a bijection between $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ and $\text{Spc}(\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}))$.

3. PRIME TENSOR IDEALS OF $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$

3.1. Classification of the prime tensor ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. Here, we will classify the prime \otimes -ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ when the quiver $A_{\mathbb{Z}}$ has bounded path length.

Denote the class of prime ideals of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and $(\mathcal{P}(\mathbb{Z}); \cap, \cup, ^c, \emptyset, \mathbb{Z})$ by $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ respectively. We will consider a subset Q'_0 of the vertices of a quiver $Q = (Q_0, Q_1, s, t)$ as a full subquiver of Q . A full subquiver is determined by the set of vertices (see Definition 2.3). For example, we will be able to see $\mathbb{Z} = (A_{\mathbb{Z}})_0$ as $A_{\mathbb{Z}}$.

Proposition 3.1. Assume that there is a natural number $n \geq 1$ such that the length of paths in the quiver $A_{\mathbb{Z}}$ is less than n . Then, for any representation $V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and a prime \otimes -ideal $\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$, we have $K_{\text{supp}(V)} \in \mathcal{M}$ if and only if $V \in \mathcal{M}$.

Proof. (\Rightarrow). Since \mathcal{M} is a prime \otimes -ideal, we have $V = V \otimes K_{\text{supp}(V)} \in \mathcal{M}$.
(\Leftarrow). Let the representation V be in \mathcal{M} . We have $0 = K_{\text{supp}(V)} \otimes K_{\text{supp}(V)^c} \in \mathcal{M}$. Since \mathcal{M} is a prime \otimes -ideal, either $K_{\text{supp}(V)}$ or $K_{\text{supp}(V)^c}$ are in \mathcal{M} . We show that $K_{\text{supp}(V)} \in \mathcal{M}$ by contradiction. More precisely, we will show that $K_{\text{supp}(V)^c} \in \mathcal{M}$ implies $K_{\mathbb{Z}} \in \mathcal{M}$, and this contradicts the assumption that \mathcal{M} is prime.

Since \mathcal{M} is a prime \otimes -ideal, \mathcal{M} is closed under extension. Thus, for an exact sequence,

$$0 \rightarrow V \rightarrow V \oplus K_{\text{supp}(V)^c} \rightarrow K_{\text{supp}(V)^c} \rightarrow 0$$

the representation $V \oplus K_{\text{supp}(V)^c}$ is in \mathcal{M} . Then, we have $K'_{\mathbb{Z}} \otimes (V \oplus K_{\text{supp}(V)^c}) \in \mathcal{M}$ where the representation $K'_{\mathbb{Z}}$ is defined by

$$(K'_{\mathbb{Z}})_a := K \text{ (} a \text{ is any integer)}$$

$$(K'_{\mathbb{Z}})_{\alpha} := 0 \text{ (} \alpha \text{ is any arrow)}$$

Then, the representation $K'_{\mathbb{Z}}$ is in \mathcal{M} because there exists a morphism

$$K'_{\mathbb{Z}} \otimes (V \oplus K_{\text{supp}(V)^c}) \rightarrow K'_{\mathbb{Z}} \otimes (V \oplus K_{\text{supp}(V)^c})$$

such that its kernel is $K'_{\mathbb{Z}}$. Here, note that support of $K'_{\mathbb{Z}} \otimes (V \oplus K_{\text{supp}(V)^c})$ is \mathbb{Z} .

Recall that we assume that all paths of the quiver $A_{\mathbb{Z}}$ have length smaller than or equal to n . This implies that the quiver has infinitely many sinks and sources. Take an arbitrary sink and let the sink be t_0 (t comes from ‘‘target’’). And define a source s_1 as the smallest source larger than t_0 . Define a sink t_2 as the smallest sink larger than s_1 . Inductively, we can define sinks t_{2i} and sources s_{2i+1} for all non negative integers i . Similarly, we can define sinks t_{2i} and sources s_{2i-1} for all non positive integers i .

Let $B \subset \mathbb{Z}$ be $\bigcup_{i \in \mathbb{Z}} [s_{4i-1}, s_{4i+1}]$, and $D \subset \mathbb{Z}$ be $\bigcup_{i \in \mathbb{Z}} [s_{4i+1}, s_{4i+3}]$ and C be $\bigcup_{i \in \mathbb{Z}} \{s_{4i-1}, s_{4i+1}\}$. Then we obtain

$$K_C = K_B \otimes K_D$$

If we write pictures of the sets B , D and C , the sets will be the integers marked by $\underbrace{\hspace{1.5cm}}$ where B is like

$$\cdots \leftarrow \underbrace{s_{-5} \rightarrow t_{-4} \leftarrow s_{-3} \rightarrow t_{-2} \leftarrow s_{-1} \rightarrow t_0 \leftarrow s_1 \rightarrow t_2 \leftarrow s_3 \rightarrow \cdots}_{\text{B}}$$

and D is like

$$\cdots \leftarrow \underbrace{s_{-5} \rightarrow t_{-4} \leftarrow s_{-3} \rightarrow t_{-2} \leftarrow s_{-1} \rightarrow t_0 \leftarrow s_1 \rightarrow t_2 \leftarrow s_3 \rightarrow \cdots}_{\text{D}}$$

and C is like

$$\cdots \leftarrow \underbrace{s_{-5}} \rightarrow t_{-4} \leftarrow \underbrace{s_{-3}} \rightarrow t_{-2} \leftarrow \underbrace{s_{-1}} \rightarrow t_0 \leftarrow \underbrace{s_1} \rightarrow t_2 \leftarrow \underbrace{s_3} \rightarrow \cdots$$

Since $K_B \otimes K_D = K_C = K_C \otimes K'_{\mathbb{Z}} \in \mathcal{M}$, we have K_B or K_D are in \mathcal{M} . We will assume that K_B is in \mathcal{M} . (we can assume $K_D \in \mathcal{M}$ instead of $K_B \in \mathcal{M}$. The proof is similar.)

For this B , we will construct subsets B' , B'' of \mathbb{Z} where $B' := B \cup \bigcup_{i \in \mathbb{Z}} [s_{4i+1}, t_{4i+2}-1]$ and $B'' := B' \cup \bigcup_{i \in \mathbb{Z}} [t_{4i-2}+1, s_{4i+2}-1]$ such that $K_{B'} \in \mathcal{M}$ and $K_{B''} \in \mathcal{M}$. And using this B'' , we will show that $K_{\mathbb{Z}} \in \mathcal{M}$.

(Construction of B')

For the B , we show $K_{B'} \in \mathcal{M}$ where $B' := B \cup \bigcup_{i \in \mathbb{Z}} [s_{4i+1}, t_{4i+2} - 1]$. Define a subset $B_1 := B \cup \{s_{4i+1} + 1 : i \in \mathbb{Z}, s_{4i+1} + 1 \text{ is not a sink}\}$. Then, we have an exact sequence

$$0 \rightarrow K_{B_1 \setminus B} \rightarrow K_{B_1} \rightarrow K_B \rightarrow 0$$

where $K_{B_1 \setminus B} = K_{B_1 \setminus B} \otimes K'_\mathbb{Z} \in \mathcal{M}$ and since \mathcal{M} is closed under extensions, the representation K_{B_1} is in \mathcal{M} . Inductively, we can define subsets of \mathbb{Z} by

$$B_{\ell+1} := B_\ell \cup \{s_{4i+1} + (\ell + 1) : i \in \mathbb{Z}, s_{4i+1} + 1, \dots, s_{4i+1} + \ell \text{ are not sinks}\}$$

for non negative integer ℓ . Then we can show that $K_\ell \in \mathcal{M}$ by mathematical induction. For $\ell = 1$, we have already shown that $K_{B_1} \in \mathcal{M}$. If we have $K_{B_\ell} \in \mathcal{M}$, then we can show $K_{B_{\ell+1}} \in \mathcal{M}$ because for $K_{B_\ell} \in \mathcal{M}$, there is an exact sequence

$$0 \rightarrow K_{B_{\ell+1} \setminus B_\ell} \rightarrow K_{B_{\ell+1}} \rightarrow K_{B_\ell} \rightarrow 0$$

We obtain $K_{B_{\ell+1}}$ in \mathcal{M} because \mathcal{M} is closed under extensions with K_{B_ℓ} and $K_{B_{\ell+1} \setminus B_\ell} \in \mathcal{M}$. Thus, by mathematical induction, we have $K_{B_\ell} \in \mathcal{M}$ for any $\ell \geq 1$. Note that there exists $m \leq n$ such that $B_m = B_{m+1} = \dots$ because the length of paths are smaller than or equal to n . Then, the set B_m contains $s_{4i+2} - 1$ for any $i \in \mathbb{Z}$ and we have $B' = B_m$ and $K_{B'} \in \mathcal{M}$.

(Construction of B'')

For the B' defined above, we show $K_{B''} \in \mathcal{M}$ where $B'' := B' \cup \bigcup_{i \in \mathbb{Z}} [s_{4i-2} + 1, s_{4i+2} - 1]$. Define a subset $B'_1 := B' \cup \{s_{4i-1} - 1 : i \in \mathbb{Z}, s_{4i-1} - 1 \text{ is not a sink}\}$. Then, we have an exact sequence

$$0 \rightarrow K_{B'_1 \setminus B'} \rightarrow K_{B'_1} \rightarrow K_{B'} \rightarrow 0$$

where $K_{B'_1 \setminus B'} = K_{B'_1 \setminus B'} \otimes K'_\mathbb{Z} \in \mathcal{M}$ and since \mathcal{M} is closed under extensions, $K_{B'_1}$ is in \mathcal{M} . Inductively, we can define subsets of \mathbb{Z} as

$$B'_{(\ell+1)} := B'_\ell \cup \{s_{4i-1} - (\ell + 1) : i \in \mathbb{Z}, s_{4i-1} - 1, \dots, s_{4i-1} - (\ell + 1) \text{ are not sinks}\}$$

for non negative integer ℓ . Then we can show that $K_{B'_\ell} \in \mathcal{M}$ for any non-negative integers by mathematical induction. For $\ell = 1$, we have already shown that $K_{B'_1} \in \mathcal{M}$. If we have $K_{B'_\ell} \in \mathcal{M}$, then we can show $K_{B'_{(\ell+1)}} \in \mathcal{M}$ because for $K_{B'_\ell} \in \mathcal{M}$, there is an exact sequence

$$0 \rightarrow K_{B'_{(\ell+1)} \setminus B'_\ell} \rightarrow K_{B'_{(\ell+1)}} \rightarrow K_{B'_\ell} \rightarrow 0$$

We obtain $K_{B'_{(\ell+1)}}$ in \mathcal{M} because \mathcal{M} is closed under extensions. Thus, by mathematical induction, we have $K_{B'_\ell} \in \mathcal{M}$ for any $\ell \geq 1$. Note that there exists $m \leq n$ such that $B'_m = B'_{(m+1)} = \dots$ because the length of paths are smaller than or equal to n . Then, the set B'_m contains $t_{4i-2} + 1$ for any $i \in \mathbb{Z}$ and we have $B'' = B'_m$ and $K_{B''} \in \mathcal{M}$.

(proof of $K_\mathbb{Z} \in \mathcal{M}$)

Define $E = \{t_{4i+2} : i \in \mathbb{Z}\}$ and note that $B'' = \mathbb{Z} \setminus \{t_{4i+2} : i \in \mathbb{Z}\} = \mathbb{Z} \setminus E$ and $K_E = K_E \otimes K'_\mathbb{Z} \in \mathcal{M}$. We have the exact sequence

$$0 \rightarrow K_E \rightarrow K_\mathbb{Z} \rightarrow K_{B''} \rightarrow 0.$$

Since \mathcal{M} is closed under extensions and $K_{B''}, K_E \in \mathcal{M}$, the representation $K_\mathbb{Z} \in \mathcal{M}$. Thus, $K_\mathbb{Z} \in \mathcal{M}$ and it leads $\mathcal{M} = \text{Rep}^{\text{pfd}}(A_\mathbb{Z})$. This is a contradiction. Hence, we have $K_{\text{supp}(V)} \in \mathcal{M}$. \square

Proposition 3.2.

There is a bijection between $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$.

Proof. Define two maps

$$\begin{aligned}\phi &: \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})) \rightarrow \text{Spc}(\mathcal{P}(\mathbb{Z})), \\ \mathcal{M} &\mapsto \{\text{supp}(V) \in \mathcal{P}(\mathbb{Z}) : V \in \mathcal{M}\}, \\ \psi &: \text{Spc}(\mathcal{P}(\mathbb{Z})) \rightarrow \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})), \\ \mathcal{Q} &\mapsto \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) \in \mathcal{Q}\}.\end{aligned}$$

First, we check $\phi(\mathcal{M}) \in \text{Spc}(\mathcal{P}(\mathbb{Z}))$ for $\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\psi(\mathcal{Q}) \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ for $\mathcal{Q} \in \text{Spc}(\mathcal{P}(\mathbb{Z}))$. After that, we will show ϕ and ψ are inverse maps of each other.

The following (i), (ii) and (iii) show that $\phi(\mathcal{M})$ is an ideal of $\mathcal{P}(\mathbb{Z})$ and (iv) shows that $\phi(\mathcal{M})$ is a prime ideal of $\mathcal{P}(\mathbb{Z})$.

(i) $\phi(\mathcal{M})$ is not empty because $\emptyset = \text{supp}(0) \in \phi(\mathcal{M})$ where 0 is the zero representation.

(ii) For any $\text{supp}(V)$ and $\text{supp}(V') \in \phi(\mathcal{M})$ (where V and $V' \in \mathcal{M}$), we will show $\text{supp}(V) \cup \text{supp}(V') \in \phi(\mathcal{M})$. Since \mathcal{M} is a prime \otimes -ideal of $\text{Spc}(\mathcal{P}(\mathbb{Z}))$, for the exact sequence

$$0 \rightarrow V \rightarrow V \oplus V' \rightarrow V' \rightarrow 0,$$

the direct sum $V \oplus V'$ of V and V' is in \mathcal{M} . Thus, we have $\text{supp}(V) \cup \text{supp}(V') = \text{supp}(V \oplus V') \in \phi(\mathcal{M})$.

(iii) For any $I \in \mathcal{P}(\mathbb{Z})$ and for any $\text{supp}(V) \in \phi(\mathcal{M})$, we will show that if $I \subset \text{supp}(V)$, then $I \in \phi(\mathcal{M})$. Assume that $I \subset \text{supp}(V)$. Then we have $\text{supp}(K_I) = \text{supp}(K_I \otimes V)$. Since $K_I \otimes V \in \mathcal{M}$, we obtain $K_I \in \mathcal{M}$ and $I \in \phi(\mathcal{M})$.

By (i), (ii) and (iii), $\phi(\mathcal{M})$ is an ideal of $\mathcal{P}(\mathbb{Z})$.

(iv) If $I \cap J \in \phi(\mathcal{M})$ for $I, J \in \mathcal{P}(\mathbb{Z})$, we will show that I or J are in $\phi(\mathcal{M})$. Since $I \cap J \in \phi(\mathcal{M})$, there exists $W \in \mathcal{M}$ such that $I \cap J = \text{supp}(W)$. Define $V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ by

$$V_a := \begin{cases} W_a & a \in \text{supp}(W) \\ K & a \in I \setminus \text{supp}(W) \\ 0 & \text{otherwise} \end{cases}$$

$$V_\alpha := \begin{cases} W_\alpha & s(\alpha), t(\alpha) \in \text{supp}(W) \\ 0 & \text{otherwise} \end{cases}$$

where a is a vertex and α is an arrow of $A_{\mathbb{Z}}$.

Then we have $V \otimes K_{\text{supp}(W)} = W \in \mathcal{M}$. Since \mathcal{M} is a prime \otimes -ideal, the representations V or $K_{\text{supp}(W)}$ are in \mathcal{M} . If $V \in \mathcal{M}$, then we have $I = \text{supp}(V) \in \phi(\mathcal{M})$. On the other hand, if $K_{\text{supp}(W)} \in \mathcal{Q}$, then $K_I \otimes K_J = K_{\text{supp}(W)} \in \mathcal{M}$. Therefore K_I or K_J are in \mathcal{M} because \mathcal{M} is a prime \otimes -ideal. This implies I or J are in $\phi(\mathcal{M})$.

Next, the following (i), (ii) and (iii) show that $\psi(\mathcal{Q})$ is a \otimes -ideal of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and (iv) shows that $\psi(\mathcal{Q})$ is a prime \otimes -ideal of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$.

(i) We will show that $\psi(\mathcal{Q})$ is closed under kernels and cokernels. For any $f : V \rightarrow W$ in $\psi(\mathcal{Q})$, the kernel and the cokernel of f satisfy $\text{supp}(\text{Ker}(f)) \subseteq \text{supp}(V)$ and $\text{supp}(\text{Coker}(f)) \subseteq \text{supp}(W)$ where $\text{supp}(V)$ and $\text{supp}(W)$ are in the prime ideal \mathcal{Q} of $\mathcal{P}(\mathbb{Z})$. Thus, we have $\text{supp}(\text{Ker}(f)) \in \mathcal{Q}$ and $\text{supp}(\text{Coker}(f)) \in \mathcal{Q}$. It means

that $\text{Ker}(f)$ and $\text{Coker}(f)$ are in $\psi(\mathcal{Q})$. Hence, $\psi(\mathcal{Q})$ is closed under kernels and cokernels.

(ii) We will show that $\psi(\mathcal{Q})$ is closed under extensions. For $V', V'' \in \psi(\mathcal{Q})$, take an exact sequence

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$$

where $V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. Then $\text{supp}(V) = \text{supp}(V') \cup \text{supp}(V'') \in \mathcal{Q}$ holds because $\text{supp}(V')$ and $\text{supp}(V'')$ are in the prime ideal \mathcal{Q} . we have $V \in \psi(\mathcal{Q})$ and therefore, $\psi(\mathcal{Q})$ is closed under extensions.

(iii) We will show that for any $M \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ and $V \in \psi(\mathcal{Q})$, the representation $M \otimes V$ is in $\psi(\mathcal{Q})$. Since $\text{supp}(M \otimes V) = \text{supp}(M) \cap \text{supp}(V) \in \mathcal{Q}$ holds, we have $M \otimes V \in \psi(\mathcal{Q})$.

By (i), (ii), and (iii), we have shown that $\psi(\mathcal{Q})$ is a \otimes -ideal of $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$. Now we show that it is prime.

(iv) If $V \otimes W \in \psi(\mathcal{Q})$ for V and $W \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$, then $\text{supp}(V) \cap \text{supp}(W) = \text{supp}(V \otimes W) \in \mathcal{Q}$. Since \mathcal{Q} is a prime ideal of $\mathcal{P}(\mathbb{Z})$, we have $\text{supp}(V) \in \mathcal{Q}$ or $\text{supp}(W) \in \mathcal{Q}$. Hence, V or $W \in \psi(\mathcal{Q})$ and therefore $\psi(\mathcal{Q})$ is a prime \otimes -ideal of $\mathcal{P}(\mathbb{Z})$.

Finally, we show ϕ and ψ are inverse maps of each other. For any $\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\mathcal{Q} \in \text{Spc}(\mathcal{P}(\mathbb{Z}))$, we have

$$\begin{aligned} \phi \circ \psi(\mathcal{Q}) &= \phi(\{V \in \text{Rep}^{\text{pfd}}(\mathbb{Z}) : \text{supp}(V) \in \mathcal{Q}\}) \\ &= \{\text{supp}(V) \in \mathcal{P}(\mathbb{Z}) : V \in \{V' \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V') \in \mathcal{Q}\}\} \\ &= \{\text{supp}(V) \in \mathcal{P}(\mathbb{Z}) : V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}), \text{supp}(V) \in \mathcal{Q}\} \\ &= \mathcal{Q}. \end{aligned}$$

The last equality holds because every $I \subseteq \mathbb{Z}$ is the support of some representation (e.g., K_I).

$$\begin{aligned} \psi \circ \phi(\mathcal{M}) &= \psi(\{\text{supp}(V) \in \mathcal{P}(\mathbb{Z}) : V \in \mathcal{M}\}) \\ &= \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) \in \{\text{supp}(V') \in \mathcal{P}(\mathbb{Z}) : V' \in \mathcal{M}\}\} \\ &= \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) = \text{supp}(V') \exists V' \in \mathcal{M}\} \\ &= \mathcal{M}. \quad (\because \text{Proposition 3.1}) \end{aligned}$$

Here, we use Proposition 3.1 to show the relation

$$\mathcal{M} \supseteq \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) = \text{supp}(V') \exists V' \in \mathcal{M}\}.$$

We will prove it. For any $V \in \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) = \text{supp}(V') \exists V' \in \mathcal{M}\}$, there exists $V' \in \mathcal{M}$ such that $\text{supp}(V) = \text{supp}(V')$. By Proposition 3.1, the interval representation $K_{\text{supp}(V')} \in \mathcal{M}$. Thus, we obtain $K_{\text{supp}(V)} = \text{supp}(V) \otimes K_{\text{supp}(V')} \in \mathcal{M}$. Therefore, the representation $V = V \otimes K_{\text{supp}(V)}$ is in \mathcal{M} . We have shown

$$\mathcal{M} = \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) = \text{supp}(V') \exists V' \in \mathcal{M}\}.$$

The two maps ϕ and ψ are inverse maps of each other. Therefore there is a bijection between $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$. \square

The next corollary is our main theorem.

Corollary 3.3. *There is a one-to-one correspondence between the following four sets, $\text{Ult}(\mathcal{P}(\mathbb{Z}))$, $\text{Spc}(\mathcal{P}(\mathbb{Z}))$, $\text{Spc}(\prod_{i \in \mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}))$ and $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$.*

Proof. There is a bijection between $\text{Ult}(\mathcal{P}(\mathbb{Z}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ by Lemma 2.34. There is a bijection between $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ and $\text{Spc}(\prod_{i \in \mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}))$ by Example 2.36. There is a bijection between $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$ by Proposition 3.2. \square

Remark 3.4. ([6, Corollary of 2.4])

The cardinality of $\text{Ult}(\mathcal{P}(\mathbb{Z}))$ is same as the cardinality of the power set of the power set of \mathbb{Z} . This is equal to the cardinality of the power set of \mathbb{R} . Thus, the cardinality of $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and the power set of \mathbb{R} is equal.

3.2. Topological properties of $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$. In this subsection, we will see that $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ is a compact Hausdorff space homeomorphic to the topological space $\text{Spc}(\mathcal{P}(\mathbb{Z})) \cong \text{Spc}(\prod_{i \in \mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}))$ when the quiver $A_{\mathbb{Z}}$ has bounded length of paths. That is, we show the following proposition.

Proposition 3.5.

Let $\mathcal{P}(\mathbb{Z})$ be the power set of \mathbb{Z} and consider it as a Boolean ring (isomorphic to $(\prod_{i \in \mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}))$ by Example 2.36). Then we have a homeomorphism between $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ and $\text{Spc}(\mathcal{P}(\mathbb{Z}))$, where both sets are equipped with their respective Zariski topologies (defined below).

We begin with the definition of the Zariski topology on $\text{Spc}(\mathcal{A})$ for a general tensor abelian category \mathcal{A} .

Definition 3.6 (Zariski topology on tensor categories [4], [14]).

Let \mathcal{A} be a tensor category. For every family of objects $\mathcal{S} \subset \mathcal{A}$ we denote by $Z(\mathcal{S})$ the following subset of $\text{Spc}(\mathcal{A})$:

$$Z(\mathcal{S}) := \{\mathcal{M} \in \text{Spc} \mathcal{A} : \mathcal{S} \cap \mathcal{M} = \emptyset\}.$$

The collection $\{Z(\mathcal{S}) : \mathcal{S} \subset \mathcal{A}\}$ defines closed subset of $\text{Spc}(\mathcal{A})$, called the *Zariski topology*. The subsets $Z(\{V\}) = \{\mathcal{M} \in \text{Spc}(\mathcal{A}) : V \notin \mathcal{M}\}$ for $V \in \mathcal{A}$ form a basis of closed subsets.

Proposition 3.7.

The collection of sets $Z(\mathcal{S})$ from Definition 3.6 are the closed subsets of a topology.

Proof. Let $(\mathcal{S}_j)_{j \in J}$ be a collection of families of objects of \mathcal{A} and let \mathcal{S}_1 and \mathcal{S}_2 be the families of objects of \mathcal{A} . Then we have $\bigcap_{j \in J} Z(\mathcal{S}_j) = Z(\bigcup_{j \in J} \mathcal{S}_j)$ and $Z(\mathcal{S}_1) \cup Z(\mathcal{S}_2) = Z(\{V_1 \oplus V_2 : V_1 \in \mathcal{S}_1, V_2 \in \mathcal{S}_2\})$. Moreover, we have $Z(\mathcal{A}) = \emptyset$ and $Z(\emptyset) = \mathcal{A}$. Thus the collection $\{Z(\mathcal{S}) : \mathcal{S} \subset \mathcal{A}\}$ are the closed subsets of a topology on $\text{Spc}(\mathcal{A})$. \square

Lemma 3.8.

Consider the tensor category $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$, where $A_{\mathbb{Z}}$ is a quiver of type A with bounded path length. Then, for any representation $V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$ the closed set $Z(\{V\}) = \{\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})) : V \notin \mathcal{M}\}$ is open.

Proof. For any prime \otimes -ideal \mathcal{M} in $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$, the representation $K_{\text{supp } V}$ is in \mathcal{M} if and only if $K_{\mathbb{Z} \setminus \text{supp } V}$ is not in \mathcal{M} . This is because if $K_{\text{supp } V} \in \mathcal{M}$ and $K_{\mathbb{Z} \setminus \text{supp } V} \in \mathcal{M}$, then we have $K_{\text{supp } V} \oplus K_{\mathbb{Z} \setminus \text{supp } V} \in \mathcal{M}$. It means $K_{\mathbb{Z}} =$

$K_{\text{supp}(K_{\text{supp } V} \oplus K_{\mathbb{Z} \setminus \text{supp } V})} \in \mathcal{M}$ by Lemma 3.1. This contradicts that \mathcal{M} is a prime \otimes -ideal. Thus $K_{\text{supp } V}$ or $K_{\mathbb{Z} \setminus \text{supp } V}$ is not in \mathcal{M} . Also, one of the representations $K_{\mathbb{Z} \setminus \text{supp } V}$ and $K_{\text{supp } V}$ is included. This is because $K_{\text{supp } V} \otimes K_{\mathbb{Z} \setminus \text{supp } V} = 0$ in \mathcal{M} and since \mathcal{M} is the prime \otimes -ideal, either $K_{\text{supp } V}$ or $K_{\mathbb{Z} \setminus \text{supp } V}$ is in \mathcal{M} . Thus we have shown that the representation $K_{\text{supp } V}$ is in \mathcal{M} if and only if $K_{\mathbb{Z} \setminus \text{supp } V}$ is not in \mathcal{M} .

Now by Lemma 3.1, we have the equation $Z(\{V\}) := \{\mathcal{M} : V \notin \mathcal{M}\} = \{\mathcal{M} : K_{\text{supp } V} \notin \mathcal{M}\} =: Z(\{K_{\text{supp } V}\})$. Thus we have

$$\begin{aligned} Z(\{V\})^c &= Z(\{K_{\text{supp } V}\})^c \\ &:= \{\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})) : K_{\text{supp } V} \notin \mathcal{M}\}^c \\ &= \{\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})) : K_{\text{supp } V} \in \mathcal{M}\} \\ &= \{\mathcal{M} \in \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})) : K_{\mathbb{Z} \setminus \text{supp } V} \notin \mathcal{M}\} \\ &\quad (\because K_{\text{supp}(V)} \in \mathcal{M} \text{ iff } K_{\mathbb{Z} \setminus \text{supp}(V)} \notin \mathcal{M}) \\ &=: Z(\{K_{\mathbb{Z} \setminus \text{supp } V}\}). \end{aligned}$$

Therefore $Z(\{V\})$ is open. \square

Corollary 3.9. *Consider the tensor category $\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})$, where $A_{\mathbb{Z}}$ is a quiver of type A with bounded path length. Then, the topological space $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ is a Hausdorff space.*

Proof. Let \mathcal{M}_1 and \mathcal{M}_2 be different prime \otimes -ideals. Then there exists a representation V such that V is in $\mathcal{M}_1 \setminus \mathcal{M}_2$. Then $\mathcal{M}_2 \in Z(\{V\})$ and $\mathcal{M}_1 \in Z(\{V\})^c$. By Lemma 3.8, the closed set $Z(\{V\})$ is open. Thus $\text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}))$ is a Hausdorff space. \square

Definition 3.10 (Zariski topology on a spectrum of a ring).

Let R be a commutative ring and I be an ideal. We denote by $V(I)$ the following subset of $\text{Spc}(R)$:

$$V(I) := \{\mathcal{P} \in \text{Spc}(R) : I \subseteq \mathcal{P}\},$$

the collection $\{V(I) : I \text{ is an ideal}\}$ are the closed subsets of a topology on $\text{Spc}(R)$, called the *Zariski topology*. Its has a basis of opens consisting of the $D_f := \{\mathcal{P} \in \text{Spc}(R) : f \notin \mathcal{P}\}$ where $f \in R$.

Lemma 3.11. *Let R be a commutative ring. The topological space $\text{Spc}(R)$ is compact.*

Proof. Let $\{V(I_j)^c\}_{j \in J}$ be an open covering of $\text{Spc}(R)$. Then we have $\emptyset = \bigcap_{j \in J} V(I_j) = V(\sum_{j \in J} I_j)$. Then, we have $(1) = \sum_{j \in J} I_j$ because if $\sum_{j \in J} I_j \subsetneq (1)$, then there exist a maximal ideal that contains $\sum_{j \in J} I_j$. It implies $\emptyset \neq V(\sum_{j \in J} I_j)$ which is a contradiction. Thus we have $(1) = \sum_{j \in J} I_j$. Then there exist $n \in \mathbb{Z}_{\geq 1}$ such that $1 = \sum_{i=1}^{i=n} f_{j_i}$ for $f_{j_i} \in I_{j_i}$. Hence we obtain $(1) = \sum_{i=1}^{i=n} I_{j_i}$, and we have $\text{Spc}(R)^c = \emptyset = V(\sum_{i=1}^{i=n} I_{j_i}) = \bigcap_{i=1}^{i=n} V(I_{j_i})$. We got $\text{Spc}(R) = \bigcup_{i=1}^{i=n} V(I_{j_i})^c$ and therefore the topological space $\text{Spc}(R)$ is compact. \square

We rewrite the statement of Proposition 3.5 again.

Proposition (3.5).

Let $\mathcal{P}(\mathbb{Z})$ be the power set of \mathbb{Z} and consider it as a Boolean ring (isomorphic

to $(\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}))$ by Example 2.36). Then we have a homeomorphism between $\mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}))$ and $\mathrm{Spc}(\mathcal{P}(\mathbb{Z}))$, where both sets are equipped with their respective Zariski topologies (defined below).

Proof. We will show that the bijective map

$$\begin{aligned} \psi : \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) &\rightarrow \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}})), \\ \mathcal{Q} &\mapsto \{V \in \mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}) : \mathrm{supp}(V) \in \mathcal{Q}\} \end{aligned}$$

defined in Proposition 3.2 is a homeomorphism. Since we know that $\mathrm{Spc}(\mathcal{P}(\mathbb{Z})) = \mathrm{Spc}(\prod_{i \in \mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}))$ is compact (see Lemma 3.11) and $\mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}))$ is a Hausdorff space (see Lemma 3.9), it is enough to show that the bijective map $\psi : \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) \rightarrow \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}))$ is continuous (a continuous bijective map from compact space to a Hausdorff space is homeomorphism, for example, [15, Proposition 13.26] for the proof).

Let $Z(\mathcal{S}) = \bigcap_{W \in \mathcal{S}} Z(\{W\})$ be a closed set of $\mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}))$. Then we obtain $\psi^{-1}(Z(\mathcal{S})) = \bigcap_{W \in \mathcal{S}} \psi^{-1}(Z(\{W\}))$. For each $\psi^{-1}(Z(\{W\}))$, we have the equation

$$\begin{aligned} \psi^{-1}(Z(\{W\})) &= \psi^{-1}(\{\mathcal{M} \in \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}})) : W \notin \mathcal{M}\}) \\ &= \psi^{-1}(\{\mathcal{M} \in \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}})) : K_{\mathrm{supp}(W)} \notin \mathcal{M}\}) \quad (\because \text{by Proposition 3.1}) \\ &= \{\psi^{-1}(\mathcal{M}) \in \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) : K_{\mathrm{supp}(W)} \notin \mathcal{M}\} \\ &= \{\psi^{-1}(\mathcal{M}) \in \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) : \mathrm{supp}(W) \notin \psi^{-1}(\mathcal{M})\} \\ &= \{\mathcal{Q} \in \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) : \mathrm{supp}(W) \notin \mathcal{Q}\} \quad (\text{Replace } \psi^{-1}(\mathcal{M}) \text{ with } \mathcal{Q}.) \\ &= \{\mathcal{Q} \in \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) : \mathbb{Z} \setminus \mathrm{supp}(W) \in \mathcal{Q}\} \quad (\because \mathcal{Q} \text{ is a prime ideal}) \\ &= \{\mathcal{Q} \in \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) : \mathbb{Z} \setminus \mathrm{supp}(W) \notin \mathcal{Q}\}^c \\ &= D_{\mathbb{Z} \setminus \mathrm{supp}(W)}^c. \end{aligned}$$

Since $D_{\mathbb{Z} \setminus \mathrm{supp}(W)}$ is open, the complement $D_{\mathbb{Z} \setminus \mathrm{supp}(W)}^c = \psi^{-1}(Z(\{W\}))$ is a closed set. Thus we can conclude $\psi^{-1}(Z(\mathcal{S})) = \bigcap_{W \in \mathcal{S}} \psi^{-1}(Z(\{W\}))$ is a closed set and therefore the map ψ is a homeomorphism. \square

3.3. About the representations of the quiver $\vec{A}_{\mathbb{Z}}$.

In the proof of Proposition 3.1, we need the condition

the length of paths are bounded.

Without it, we can not prove $K_{\mathrm{supp}(V)} \in \mathcal{M}$ if and only if $V \in \mathcal{M}$ for any $V \in \mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}})$ and $\mathcal{M} \in \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}))$. For example, there may be a representation such that its support is \mathbb{Z} , but the representation is contained in a prime \otimes -ideal. We will provide such a representation when $A_{\mathbb{Z}}$ is the quiver

$$\cdots \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \diamond \longrightarrow \cdots$$

which will be denote by $\vec{A}_{\mathbb{Z}}$. The proof seems to generalise, to any quiver of type A containing an unbounded path, but we restrict our attention to $\vec{A}_{\mathbb{Z}}$.

Using this, we show that the map

$$\begin{aligned} \psi : \mathrm{Spc}(\mathcal{P}(\mathbb{Z})) &\rightarrow \mathrm{Spc}(\mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}})), \\ \mathcal{Q} &\mapsto \{V \in \mathrm{Rep}^{\mathrm{pfd}}(A_{\mathbb{Z}}) : \mathrm{supp}(V) \in \mathcal{Q}\} \end{aligned}$$

which appeared in the proof of Proposition 3.2 is injective but not surjective.

We will prepare some lemmas and definitions to show the existence of the representation whose support is \mathbb{Z} .

Definition 3.12. (An ideal generated by a set)

Let Q be a quiver and S a subset of objects of $\mathrm{Rep}^{\mathrm{pfd}}(Q)$ which contains the representation 0. Set $S_{-1} := S$, and for $n \geq 0$ define full subcategories S_n, S'_n, S''_n of $\mathrm{Rep}^{\mathrm{pfd}}(Q)$ inductively as follows.

$$\begin{aligned} S_n &:= \left\{ V \in \mathrm{Rep}^{\mathrm{pfd}}(Q) : \begin{array}{l} \text{there exists an exact sequence} \\ 0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0 \\ \text{with } V', V'' \in S_{n-1} \end{array} \right\}, \\ S'_n &:= \{W \otimes V : W \in \mathrm{Rep}^{\mathrm{pfd}}(Q), V \in S_n\}, \\ S''_n &:= \bigcup_{V, W \in S'_n} \{\mathrm{Ker}(f), \mathrm{Coker}(f) : V \xrightarrow{f} W\}. \end{aligned}$$

We then define an ideal $\langle S \rangle$ of $\mathrm{Rep}^{\mathrm{pfd}}(Q)$ by

$$\langle S \rangle := \bigcup_{n \in \mathbb{Z}_{\geq 0}} S_n$$

and call it the \otimes -ideal generated by S . Here, we have

$$S \subseteq S_0 \subseteq S'_0 \subseteq S''_0 \subseteq S_1 \subseteq \cdots \subseteq S_n \subseteq S'_n \subseteq S''_n \subseteq S_{n+1} \subseteq \cdots \subseteq \langle S \rangle.$$

Proposition 3.13. The $\langle S \rangle := \bigcup_{n \in \mathbb{Z}_{\geq 0}} S_n$ defined above is a \otimes -ideal of $\mathrm{Rep}^{\mathrm{pfd}}(Q)$. In fact, it is the smallest \otimes -ideal containing the set S .

Proof. First we show that $\langle S \rangle$ is closed under extensions. Let V' and V'' be representations in $\langle S \rangle$. Then, there exists $n \in \mathbb{Z}_{\geq 0}$ such that V' and V'' are in S_n . Take an arbitrary exact sequence

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0,$$

and we have $V \in S_{n+1} \subseteq \langle S \rangle$. Thus $\langle S \rangle$ is closed under extensions.

Next, we show that $\langle S \rangle$ is closed under kernels and cokernels. Let $f : V \rightarrow W$ be a morphism in $\langle S \rangle$. Then, there exists $n \in \mathbb{Z}_{\geq 0}$ such that V and W are in S_n and its kernels and cokernels are in $S_{n+1} \subseteq \langle S \rangle$.

Finally, we show that $\langle S \rangle$ is closed under tensor products. Let V and W be in $\mathrm{Rep}^{\mathrm{pfd}}(Q)$ and $\langle S \rangle$ respectively. There exists $n \in \mathbb{Z}_{\geq 0}$ such that $W \in S_n$. We have $V \otimes W \in S_{n+1} \subseteq \langle S \rangle$. Therefore $\langle S \rangle$ is a \otimes -ideal of $\mathrm{Rep}^{\mathrm{pfd}}(Q)$.

Next, we will show that the $\langle S \rangle$ is the smallest \otimes -ideal containing S . Let \mathcal{I} be a \otimes -ideal containing S . By the construction of S_n and the definition of \otimes -tensor ideal, S_n is included by \mathcal{I} for every $n \in \mathbb{Z}_{\geq 0}$. Thus we have $\langle S \rangle = \bigcup_{n \in \mathbb{Z}_{\geq 0}} S_n \subseteq \mathcal{I}$ and therefore $\langle S \rangle$ is the smallest \otimes -tensor ideal containing S . \square

Lemma 3.14. *Let $[a, b]$ be an interval in \mathbb{Z} . If there exists a monomorphism*

$$\phi : K_{[a,b]} \rightarrow \bigoplus_{\lambda \in \Lambda} K_{I_\lambda},$$

then there exists $\lambda \in \Lambda$ such that $[a, b] \subseteq I_\lambda$ where $K_{[a,b]}$ and $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}$ are representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$.

Similarly, if there exists an epimorphism

$$\psi : \bigoplus_{\lambda \in \Lambda} K_{I_\lambda} \rightarrow K_{[a,b]},$$

then there exists $\lambda \in \Lambda$ such that $[a, b] \subseteq I_\lambda$ where $K_{[a,b]}$ and $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}$ are representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$.

Proof. We will show the monomorphism case by contradiction. The other case can be shown in the same way.

Assume that for any $\lambda \in \Lambda$, we have $[a, b] \not\subseteq I_\lambda$. Then we obtain non-commutative diagram

$$\begin{array}{ccc} (K_{[a,b]})_a & \xrightarrow{\phi_a} & (\bigoplus_{\lambda \in \Lambda} K_{I_\lambda})_a \\ id_K \downarrow & & \downarrow 0 \\ (K_{[a,b]})_b & \xrightarrow{\phi_b} & (\bigoplus_{\lambda \in \Lambda} K_{I_\lambda})_b, \end{array}$$

and it is the contradiction. Therefore there must be $\lambda \in \Lambda$ such that $[a, b] \subseteq I_\lambda$. \square

Lemma 3.15. *Let I be an interval in \mathbb{Z} . If there exists a monomorphism*

$$\phi : K_I \rightarrow \bigoplus_{\lambda \in \Lambda} K_{J_\lambda},$$

then there exists $\lambda \in \Lambda$ such that $I \subseteq J_\lambda$ where K_I and $\bigoplus_{\lambda \in \Lambda} K_{J_\lambda}$ are representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$.

Similarly, if there exists an epimorphism

$$\psi : \bigoplus_{\lambda \in \Lambda} K_{J_\lambda} \rightarrow K_I,$$

then there exists $\lambda \in \Lambda$ such that $I \subseteq J_\lambda$ where K_I and $\bigoplus_{\lambda \in \Lambda} K_{J_\lambda}$ are the representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$.

Proof. We will show the monomorphism case. The other case can be shown in the same way.

If I is bounded, then we have already shown in Lemma 3.14. So we will assume that I is not bounded.

In case of $I = \mathbb{Z}$, let I_n be the interval $[-n, n]$ for $n \in \mathbb{Z}_{\geq 0}$. By restricting the interval $I = \mathbb{Z}$ to I_n , we obtain the monomorphism

$$\phi : K_{I_n} \rightarrow \bigoplus_{\lambda \in \Lambda} K_{J_\lambda \cap I_n}.$$

By Lemma 3.14, there exists $\lambda \in \Lambda$ such that $I_n \subseteq J_\lambda \cap I_n$. Thus for any $n \in \mathbb{Z}_{\geq 0}$ we obtain $I_n \subseteq J_{\lambda_n}$ for some λ_n . Thus the set $C_n := \{J_\lambda : I_n \subseteq J_\lambda, \lambda \in \Lambda\} \subseteq \{J_\lambda : 0 \in J_\lambda, \lambda \in \Lambda\}$ is not empty and finite because the cardinality of

the set $\{J_\lambda : 0 \in J_\lambda, \lambda \in \Lambda\}$ is equal to or smaller than the finite dimension $\dim_K(\bigoplus_{\lambda \in \Lambda} K_{J_\lambda})_0$. Thus, for the descending sequence

$$C_0 \supseteq C_1 \supseteq C_2 \supseteq C_3 \supseteq \cdots,$$

there exists $m \in \mathbb{Z}_{\geq 0}$ such that:

$$C_m = C_{m+1} = \cdots.$$

We can conclude that there exists $\lambda \in \Lambda$ such that $I = \mathbb{Z} = J_\lambda \in \bigcap_{n \in \mathbb{Z}_{\geq 0}} C_n$.

If I is bounded above and unbounded below, then there exists $a \in \mathbb{Z}$ such that $I = (-\infty, a]$. Let I'_n be the interval $[a - n, a]$ for $n \in \mathbb{Z}_{\geq 0}$. We can show that $C'_n := \{J_\lambda : I'_n \subseteq J_\lambda, \lambda \in \Lambda\}$ is not empty because we have a monomorphism

$$\phi : K_{I'_n} \rightarrow \bigoplus_{\lambda \in \Lambda} K_{J_\lambda \cap I'_n},$$

such that there exists $\lambda \in \Lambda$ which satisfies $I'_n \subset J_\lambda \cap I'_n \subseteq J_\lambda$ by Lemma 3.14. Also, we can show $C'_n \subseteq \{J_\lambda : a \in J_\lambda, \lambda \in \Lambda\}$ is finite because $\{J_\lambda : a \in J_\lambda, \lambda \in \Lambda\}$ is equal to or smaller than the $\dim_K(\bigoplus_{\lambda \in \Lambda} K_{J_\lambda})_a$. We will obtain the descending sequence that stops for a $m \in \mathbb{Z}_{\geq 0}$

$$C'_0 \supseteq C'_1 \supseteq C'_2 \supseteq C'_3 \supseteq \cdots \supseteq C'_m = C'_{m+1} = \cdots.$$

We can conclude that there exists $\lambda \in \Lambda$ such that $I' = (-\infty, a] = J_\lambda \in \bigcap_{n \in \mathbb{Z}_{\geq 0}} C'_n$.

In case of I is bounded below and unbounded above, the proof is similar to the case that I is bounded above and unbounded below. \square

Lemma 3.16. *Let V and W be representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ whose interval decomposition are $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}$ and $\bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}}$. If there exists a monomorphism*

$$\phi : V \rightarrow W,$$

then for any interval I_λ , there exists $\lambda' \in \Lambda'$ such that $I_\lambda \subseteq J_{\lambda'}$.

Similarly if there exists an epimorphism

$$\psi : V \rightarrow W,$$

then for any interval I_λ , there exists $\lambda' \in \Lambda'$ such that $I_\lambda \subseteq J_{\lambda'}$.

Proof. We will show the monomorphism case. The other case can be shown in the same way.

Since we have the natural monomorphism

$$K_{I_\lambda} \rightarrow \bigoplus_{\lambda \in \Lambda} K_{I_\lambda} \cong V \xrightarrow{\phi} W \cong \bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}},$$

by Lemma 3.15, there exists $\lambda' \in \Lambda'$ such that $I_\lambda \subseteq J_{\lambda'}$. \square

Corollary 3.17. *Let V and W be representations of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ whose interval decomposition are $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}$ and $\bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}}$. If there exists a monomorphism (resp. epimorphism)*

$$\phi : V \rightarrow W,$$

then V (resp. W) is a direct summand of $V \otimes W$.

Proof. Now we have $V \otimes W \cong (\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}) \otimes (\bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}}) \cong \bigoplus_{\lambda \in \Lambda, \lambda' \in \Lambda'} K_{I_\lambda} \otimes K_{J_{\lambda'}} = \bigoplus_{\lambda \in \Lambda, \lambda' \in \Lambda'} K_{I_\lambda \cap J_{\lambda'}}$.

If the map ϕ is a monomorphism, for any $\lambda \in \Lambda$ there exists $\lambda' \in \Lambda'$ such that $I_\lambda \subseteq J_{\lambda'}$, by Lemma 3.16. For these λ and λ' , we have $K_{I_\lambda} \otimes K_{J_{\lambda'}} = K_{I_\lambda \cap J_{\lambda'}} = K_{I_\lambda}$. Therefore $V \otimes W \cong \bigoplus_{\lambda \in \Lambda, \lambda' \in \Lambda'} K_{I_\lambda \cap J_{\lambda'}}$ has the direct summand $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda} \cong V$.

In the similar way, we can show the case of ϕ is epi. \square

Lemma 3.18. *Let V' and V'' be representations in $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$. If these are decomposed into interval representations whose intervals are bounded, then for any exact sequence*

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0,$$

the representation V is decomposed into interval representations whose intervals are bounded. That is, if V' and V'' are of the form $\bigoplus_{j \in J} K_{I_j}$ for some collection $\{I_j\}_{j \in J}$ of bounded $I_i \subseteq \mathbb{Z}$, then V is also of this form.

Proof. We prove this lemma by a contradiction. Note that V is of the form $\bigoplus_{j \in J} K_{I_j}$ for some collection $\{I_j\}_{j \in J}$ of $I_i \subseteq \mathbb{Z}$ by Proposition 2.12. We will show that if some I_i is unbounded, then there is a contradiction. Assume that there is an unbounded interval I in $\vec{A}_{\mathbb{Z}}$ such that the interval representation K_I is a direct summand of V and choose an isomorphism $V \cong W \oplus K_I$. Then we can rewrite the exact sequence by

$$0 \rightarrow V' \xrightarrow{(\phi, \psi)} W \oplus K_I \xrightarrow{(f, g)} V'' \rightarrow 0$$

Now, assume that I is unbounded above. (We can also assume I is unbounded below, but the proof is similar.) Then, $\psi_a = 0$ for every $a \in \mathbb{Z}$. This is because if there exists $a \in \mathbb{Z}$ such that $\psi_a \neq 0$, we obtain non-commutative square

$$\begin{array}{ccc} V'_a & \xrightarrow{(\phi, \psi)_a} & (W \oplus K_I)_a \\ \downarrow & & \downarrow \\ V'_{a+m} & \xrightarrow{(\phi, \psi)_{a+m}} & (W \oplus K_I)_{a+m} \end{array}$$

where m is large enough for $V'_a \rightarrow V'_{a+m}$ to be zero. Thus $\psi_a = 0$ for all $a \in \mathbb{Z}$. Then, since $0 = \text{Im}(\psi) = \text{Ker}(g)$, we have that $g : K_I \rightarrow V''$ is injective. This implies V'' has a unbounded interval representation as a direct summand. However, the representation V'' is decomposed into interval representations whose intervals are bounded. It is a contradiction. Hence, we can conclude that V is decomposed into interval representations whose intervals are bounded. \square

Proposition 3.19. *Let \mathcal{I} be the full subcategory of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ whose representations are decomposed into bounded interval representations $\bigoplus_{j \in J} K_{I_j}$ for some collection $\{I_j\}_{j \in J}$ of bounded $I_i \subseteq \mathbb{Z}$. Then \mathcal{I} is a tensor ideal of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$.*

Proof. We have already shown that \mathcal{I} is closed under extensions in Lemma 3.18. Moreover \mathcal{I} is closed under kernels and cokernels by Lemma 3.16.

Next, we will show that the representation $V \otimes W$ is in \mathcal{I} for any $V \in \text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ and $W \in \mathcal{I}$. Let $V \in \text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ and $W \in \mathcal{I}$ be isomorphic to the representations $\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}$ and $\bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}}$ where $J_{\lambda'} \subseteq \mathbb{Z}$ is bounded for every $\lambda' \in \Lambda'$. Then we obtain $(\bigoplus_{\lambda \in \Lambda} K_{I_\lambda}) \otimes (\bigoplus_{\lambda' \in \Lambda'} K_{J_{\lambda'}}) \cong \bigoplus_{\lambda \in \Lambda, \lambda' \in \Lambda'} K_{I_\lambda \cap J_{\lambda'}} \in \mathcal{I}$.

We have shown that \mathcal{I} is a tensor ideal. \square

Definition 3.20. Define the representation $K'_\mathbb{Z} \in \text{Rep}^{\text{pfd}}(A_\mathbb{Z})$ by

$$(K'_\mathbb{Z})_a := K, \quad (K'_\mathbb{Z})_\alpha := 0,$$

for every $a \in \mathbb{Z}$ and every arrow α .

Lemma 3.21. *Let S be the set $\{0, K'_\mathbb{Z}\}$ of objects of $\text{Rep}^{\text{pfd}}(\vec{A}_\mathbb{Z})$ and let $\langle S \rangle$ be the \otimes -ideal generated by S . Then we have $K_\mathbb{Z} \notin \langle S \rangle$.*

Proof. We show $K_\mathbb{Z} \notin \langle S \rangle$ by a contradiction. Assume that $K_\mathbb{Z} \in \langle S \rangle$. Then there exists $n \in \mathbb{Z}_{\geq 0}$ such that $K_\mathbb{Z} \in S_n$ where S_n is defined in Definition 3.12. By Lemma 3.18, we can see that any representations of S_0 are decomposed into bounded interval representations. Since any representations of S'_0 and S''_0 are obtained by tensoring and taking kernels and cokernels, their representations are decomposed into bounded interval representations. Therefore any representation in S_1 are also decomposed into bounded interval representations. Inductively, every representation of S_n are decomposed into bounded interval representations. Since $K_\mathbb{Z} \in S_n$, the interval representation $K_\mathbb{Z}$ are decomposed into bounded interval representations. However, the representation $K_\mathbb{Z}$ is the indecomposable interval representation whose interval is infinite length. This is a contradiction. Hence we have $K_\mathbb{Z} \notin \langle S \rangle$. \square

Proposition 3.22. There exists a maximal \otimes -ideal in $\text{Rep}^{\text{pfd}}(\vec{A}_\mathbb{Z})$ that contains a representation whose support is \mathbb{Z} .

Proof. Let S be the set $\{0, K'_\mathbb{Z}\}$ and $\langle S \rangle$ be the ideal generated by S . Define a partially ordered set $(\Sigma, \subseteq) := \{ \text{proper ideals containing } \langle S \rangle \}$. Then Σ is not an empty set because $\langle S \rangle \in \Sigma$. Let $\sigma \subseteq \Sigma$ be a totally ordered subset. Then $\bigcup_{\mathcal{I} \in \sigma} \mathcal{I}$ is a proper ideal (it does not contain $K_\mathbb{Z}$) and upper bound of σ . Thus by Zorn's lemma, there exists a maximal ideal containing $K'_\mathbb{Z} \in \langle S \rangle$ whose supports are \mathbb{Z} . \square

Lemma 3.23. *Let \mathcal{M} be a maximal \otimes -ideal of $\mathcal{A} = \text{Rep}^{\text{pfd}}(\vec{A}_\mathbb{Z})$. Then, the maximal \otimes -ideal is a prime \otimes -ideal.*

Proof. Let \mathcal{M} be a maximal \otimes -ideal and suppose that $W \otimes W' \in \mathcal{M}$. We will show that if $W \notin \mathcal{M}$ then $W' \in \mathcal{M}$. Consider the set

$$\mathcal{I} := \{V \in \mathcal{A} : V \otimes W' \in \mathcal{M}\}.$$

Then $W \in \mathcal{I}$ and $\mathcal{M} \subseteq \mathcal{I}$ hold. Since we have assumed $W \notin \mathcal{M}$, the set \mathcal{I} is strictly larger than \mathcal{M} . If we show \mathcal{I} is a \otimes -ideal, then the tensor unit I is in $\mathcal{A} = \mathcal{I}$ by the maximality of \mathcal{M} , and therefore, we can show $W' = I \otimes W' \in \mathcal{M}$. So it is enough to show that \mathcal{I} is a \otimes -ideal to prove \mathcal{M} is a prime \otimes -ideal.

First, we will show that \mathcal{I} is closed under extensions. Let V' and V'' be in \mathcal{I} . For any short exact sequence

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0,$$

there exists an exact sequence

$$0 \rightarrow V' \otimes W' \rightarrow V \otimes W' \rightarrow V'' \otimes W' \rightarrow 0$$

such that $V' \otimes W'$ and $V'' \otimes W'$ are in the \otimes -ideal \mathcal{M} . Thus, the representation $V \otimes W'$ is in \mathcal{M} and therefore V is in \mathcal{I} .

Next, we will show that \mathcal{I} is closed under multiplication. Let V be in \mathcal{A} and V' be in \mathcal{I} . We have $(V \otimes V') \otimes W' = V \otimes (V' \otimes W') \in \mathcal{M}$ and therefore $(V \otimes V') \in \mathcal{I}$.

Then, we will show that \mathcal{I} is closed under kernels and cokernels. Let $f : V \rightarrow V'$ be in \mathcal{I} . Then, we have the monomorphism $\ker(f) \otimes id_{W'} : \text{Ker}(f) \otimes W' \hookrightarrow V \otimes W'$ such that $V \otimes W'$ is in \mathcal{M} . Thus $(\text{Ker}(f) \otimes W') \otimes (V \otimes W')$ is in \mathcal{M} . By Lemma 3.17, we have $\text{Ker}(f) \otimes W'$ as a direct summand of $(\text{Ker}(f) \otimes W') \otimes (V \otimes W')$ and since \mathcal{M} is a \otimes -ideal, the representation $\text{Ker}(f) \otimes W'$ is in \mathcal{M} . Hence $\text{Ker}(f)$ is in \mathcal{I} and therefore \mathcal{I} is closed under kernels. Similarly we can show that \mathcal{I} is closed under cokernels.

We have showed \mathcal{I} is a \otimes -ideal. \square

Next corollary is what we wanted.

Corollary 3.24. *There exists a prime \otimes -ideal of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ which contains a representation whose supports is \mathbb{Z} .*

Thus the map

$$\begin{aligned} \psi : \text{Spc}(\mathcal{P}(\mathbb{Z})) &\rightarrow \text{Spc}(\text{Rep}^{\text{pfd}}(A_{\mathbb{Z}})), \\ \mathcal{Q} &\mapsto \{V \in \text{Rep}^{\text{pfd}}(A_{\mathbb{Z}}) : \text{supp}(V) \in \mathcal{Q}\} \end{aligned}$$

constructed in Proposition 3.2 is not surjective.

Proof. By Proposition 3.22, there exists a maximal \otimes -ideal of $\text{Rep}^{\text{pfd}}(\vec{A}_{\mathbb{Z}})$ that contains a representation whose supports is \mathbb{Z} and the maximal \otimes -ideal is prime \otimes -ideal by Lemma 3.23. \square

4. FUTURE WORK

It would be nice if there is a relation between tensor triangulated geometry and multidimensional persistence modules (or topological data analysis) where multidimensional persistence modules are defined as follows (we can refer to Lesnick's paper [12]).

For $n \geq 1$, define a partial order on \mathbb{N}^n by taking $(a_1, \dots, a_n) \leq (b_1, \dots, b_n)$ if and only if $a_i \leq b_i$ for all i , and let \mathbb{N}^n denote the associated poset category. The objects of the functor category $(\mathbf{Vect})^{\mathbb{N}^n}$ are called n -dimensional persistence modules. Then, we have a categorical equivalence between $(\mathbf{Vect})^{\mathbb{N}^n}$ and $(n\text{-Mod})$ where $(n\text{-Mod})$ is defined as follows [12]. Let $P_n := K[x_1, \dots, x_n]$ be a polynomial ring in n variables. We define an n -graded module to be a P_n module M with a direct sum decomposition as a K -vector space $M \cong \bigoplus_{a \in \mathbb{N}^n} M_a$, such that

$$x^b(M_a) \subset M_{a+b}$$

for all $a, b \in \mathbb{N}^n$ where $x^b = x_1^{b_1} x_2^{b_2} \dots x_n^{b_n}$. An n -graded homomorphism $f : M \rightarrow N$ is a P_n homomorphism such that for all $a \in \mathbb{N}^n$ we have $f(M_a) \subset N_a$. Since there is a categorical equivalent between the functor category $(\mathbf{Vect})^{\mathbb{N}^n}$ and $(n\text{-Mod})$, we can consider n -modules instead of the persistence modules. In the paper [7], the (associated) prime ideals of the ring P_n and its graded modules are studied and they tell some information of the persistence modules. We hope there is a connection between tensor triangulated geometry and persistence modules through prime ideals and modules.

REFERENCES

- [1] Henry Adams, Manuchehr Aminian, Elin Farnell, Michael Kirby, Joshua Mirth, Rachel Neville, Chris Peterson, and Clayton Shonkwiler, *A fractal dimension for measures via persistent homology*, Abel Symposia (2020), 1–31.

- [2] Ibrahim Assem, Andrzej Skowronski, and Daniel Simson, *Elements of the representation theory of associative algebras: Techniques of representation theory*, London Mathematical Society Student Texts, vol. 1, Cambridge University Press, 2006.
- [3] B.A.Davey and H.A.Priestley, *Introduction to lattices and order second edition*, Cambridge University Press, Cambridge, 2002.
- [4] Paul Balmer, *The spectrum of prime ideals in tensor triangulated categories*, Journal für die reine und angewandte Mathematik Volume 2005 Issue 588. (2005), 149–168.
- [5] Joseph Minhow Chan, Gunnar Carlsson, and Raul Rabadan, *Topology of viral evolution*, Proceedings of the National Academy of Sciences **110 (46)** (Nov 2013), 18566–18571.
- [6] W. W. Comfort, *Ultrafilters: Some old and some new results*, Bulletin of The American Mathematical Society **83** (1977), no. 4, 417–455.
- [7] Heather A. Harrington, Nina Otter, Hal Schenck, and Ulrike Tillmann, *Stratifying multiparameter persistent homology*, 2019.
- [8] Yasuaki Hiraoka, Takenobu Nakamura, Akihiko Hirata, Emerson G Escolar, Kaname Matsue, and Yasumasa Nishiura, *Amorphous solids and persistent homology*, Proceedings of the National Academy of Sciences **113 (26)** (Jun 2016), 7035–7040.
- [9] Kiyoshi Igusa, Job D. Rock, and Gordana Todorov, *Continuous quivers of type a (i) foundations*, 2021.
- [10] Xia Kelin and Guo-Wei, *Multidimensional persistence in biomolecular data*, Journal of computational chemistry **36** (2015), 1502–20.
- [11] G. Kelly, *The basic concepts of enriched category theory*, Reprints in Theory and Categories[electronic only] **2005** (2005).
- [12] Michael Lesnick, *The theory of the interleaving distance on multidimensional persistence modules*, Foundations of Computational Mathematics **15** (2015), no. 3, 613–650.
- [13] Steve Y Oudot, *Persistence theory: From quiver representations to data analysis*, Mathematical Surveys and Monographs, no. 209, American Mathematical Society, 2015.
- [14] Tobias J Peter, *Prime ideals of mixed artin-tate motives*, Journal of K-Theory **11** (2013), 331–349.
- [15] Wulson A Sutherland, *Introduction to metric and topological spaces second edition*, Oxford University Press, Oxford, 2009.
- [16] Andrew Tausz and Gunnar Carlsson, *Applications of zigzag persistence to topological data analysis*, 2011.

Email address: 205d851d@stu.kobe-u.ac.jp