

NOTE ON THE SUM OF THE SMALLEST AND LARGEST EIGENVALUES OF A TRIANGLE-FREE GRAPH

PÉTER CSIKVÁRI

ABSTRACT. Let G be a triangle-free graph on n vertices with adjacency matrix eigenvalues $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$. In this paper we study the quantity

$$\mu_1(G) + \mu_n(G).$$

We prove that for any triangle-free graph G we have

$$\mu_1(G) + \mu_n(G) \leq (3 - 2\sqrt{2})n.$$

This was proved for regular graphs by Brandt, we show that the condition on regularity is not necessary. We also prove that among triangle-free strongly regular graphs the Higman-Sims graph achieves the maximum of

$$\frac{\mu_1(G) + \mu_n(G)}{n}.$$

1. INTRODUCTION

In this paper every graph is simple. Motivated by the papers [2] and [4] we study the following problem. Let \mathcal{G}_3 be the family of triangle-free graphs, and for a graph G on $v(G) = n$ vertices let $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$ be the eigenvalues of the adjacency matrix of G . The problem is to determine

$$c_3 = \sup_{G \in \mathcal{G}_3} \frac{\mu_1(G) + \mu_n(G)}{v(G)}.$$

Brandt [2] proved that for regular triangle-free graphs we have

$$\mu_1(G) + \mu_n(G) \leq (3 - 2\sqrt{2})n.$$

Very recently Balogh, Clemen, Lidický, Norin and Volec proved that for regular triangle-free graphs we have $\mu_1(G) + \mu_n(G) \leq \frac{15}{94}n < 0.1596n$ and they mention in their paper that a similar but larger computation also gives the result 0.15467 instead of 0.1596. In fact, they study the smallest eigenvalue $q_n(G)$ of the so-called signless laplacian matrix $\bar{L} = D + A$, where D is the diagonal matrix consisting of the degrees of the vertices and A is the adjacency matrix of the graph G . The quantity $q_n(G)$ coincides with $\mu_1(G) + \mu_n(G)$ if G is regular. Balogh, Clemen, Lidický, Norin and Volec mentions that in case of regular graphs they can further improve their result to prove $q_n(G) \leq 0.15442n$. Our first result is to prove that in Brandt's theorem one can drop the condition of regularity.

Theorem 1.1. *Let G be a triangle-free graph on n vertices, and let $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$ be the eigenvalues of its adjacency matrix. Then*

$$\mu_1(G) + \mu_n(G) \leq (3 - 2\sqrt{2})n.$$

2010 *Mathematics Subject Classification.* Primary: 05C35. Secondary:

Key words and phrases. eigenvalues, triangle-free graphs.

The author is supported by the Counting in Sparse Graphs Lendület Research Group.

The proof of Theorem 1.1 heavily relies on the following lemma which might be of independent interest.

Lemma 1.2. *Let G be a triangle-free graph on n vertices, and let $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$ be the eigenvalues of its adjacency matrix. Then*

$$\mu_1(G) \leq \frac{-n\mu_n(G)}{\mu_1(G) - \mu_n(G)}.$$

Brandt [2] also realized that for the so-called Higman-Sims graph H we have

$$\frac{\mu_1(H) + \mu_n(H)}{v(H)} = \frac{22 + (-8)}{100} = 0.14$$

which gives a rather good lower bound for c_3 . Higman-Sims graph is the unique strongly regular graph with parameters $(100, 22, 0, 6)$. Recall that a graph G is a strongly regular graph with parameters (n, k, a, b) if it has n vertices, k -regular, any two adjacent vertices have exactly a common neighbors, and any two non-adjacent vertices have exactly b common neighbors. In this paper we show that among the strongly regular graphs, it is indeed the Higman-Sims graph which maximizes the quantity

$$\frac{\mu_1(G) + \mu_n(G)}{v(G)}.$$

Note that only finitely many triangle-free strongly regular graphs are known currently, but we do not rely on this fact.

Theorem 1.3. *Let G be a triangle-free strongly regular graph on n vertices. Then*

$$\mu_1(G) + \mu_n(G) \leq 0.14n$$

with equality if and only if G is the Higman-Sims graph.

2. PROOF OF THEOREM 1.1

We begin with proving Lemma 1.2. Before we actually start it let us mention that for regular graphs this lemma is a simple consequence of the Hoffman-Delsarte bound for independent sets. Indeed, let $\alpha(G)$ denote the size of the largest independent set of a d -regular graph. Then by the Hoffman-Delsarte bound we have

$$\alpha(G) \leq \frac{-n\mu_n(G)}{d - \mu_n(G)}.$$

Since G is triangle-free, the neighbors of a vertex determine an independent set, whence $d \leq \alpha(G)$. Since $d = \mu_1(G)$ we get that

$$\mu_1(G) = d \leq \alpha(G) \leq \frac{-n\mu_n(G)}{d - \mu_n(G)} \leq \frac{-n\mu_n(G)}{\mu_1(G) - \mu_n(G)}.$$

Based on this inequality Brandt proved that

$$\mu_1(G) + \mu_n(G) \leq (3 - 2\sqrt{2})n.$$

So after proving Lemma 1.2 we practically copy the proof of Brandt.

Proof of Lemma 1.2. Let μ_s, \dots, μ_n be the set of non-positive eigenvalues. Then

$$0 = 6 \cdot \text{number of triangles} = \sum_{i=1}^n \mu_i^3 \geq \mu_1^3 + \sum_{i=s}^n \mu_i^3.$$

Hence

$$\sum_{i=s}^n (-\mu_i)^3 \geq \mu_1^3.$$

On the other hand, we have

$$\sum_{i=s}^n (-\mu_i)^3 \leq (-\mu_n) \sum_{i=s}^n (-\mu_i)^2 \leq (-\mu_n)(2e(G) - \mu_1^2) \leq (-\mu_n)(n\mu_1 - \mu_1^2).$$

Hence $\mu_1^3 \leq (-\mu_n)(n\mu_1 - \mu_1^2)$, thus $\mu_1^2 \leq (-\mu_n)(n - \mu_1)$, or in other words,

$$\mu_1 \leq \frac{-n\mu_n}{\mu_1 - \mu_n}.$$

□

Proof of Theorem 1.1. As we mentioned earlier this proof practically follows the argument of [2].

We only need to solve the constrained maximization problem:

$$\max \left\{ \frac{\mu_1 + \mu_n}{n} \mid \mu_1 \leq \frac{-n\mu_n}{\mu_1 - \mu_n} \right\}.$$

Let $a = \mu_1$, $b = -\mu_n$ then we have $a \leq \frac{nb}{a-b}$ which is equivalent to $\frac{a^2}{n-a} \leq b$. Hence

$$\frac{a-b}{n} \leq \frac{1}{n} \left(a - \frac{a^2}{n-a} \right) = \frac{an - 2a^2}{n(n-a)}.$$

So with the notation $\alpha = a/n$ we need to maximize $f(\alpha) := \frac{\alpha - 2\alpha^2}{1-\alpha}$. Its derivative is $\frac{1-4\alpha+2\alpha^2}{(1-\alpha)^2}$ which is 0 at $\alpha = 1 \pm 1/\sqrt{2}$. Note that $\mu_1 \leq \Delta \leq n-1$, where Δ is the largest degree, so $0 \leq \alpha \leq 1$. So we only need to consider $\alpha = 1 - 1/\sqrt{2}$ and the extreme points of the interval, $\alpha = 0$ and 1, to see that $f(\alpha)$ is indeed maximal at $1 - 1/\sqrt{2}$ and $f(\alpha) = 3 - 2\sqrt{2}$.

Hence $\mu_1 + \mu_n \leq (3 - 2\sqrt{2})n$.

□

3. PROOF OF THEOREM 1.3

In this section we prove Theorem 1.3.

Proof of Theorem 1.3. Suppose for contradiction that G is a strongly regular graph with eigenvalues $(k, \mu_2^{(m_2)}, \mu_n^{(m_n)})$ such that $\frac{k+\mu_n}{n} > 0.14$. Let $-\mu_n = r$ and $r/k = x$. Again we use that $k \leq \alpha(G) \leq \frac{-n\mu_n}{k-\mu_n}$. Hence $\frac{r}{k} \geq \frac{k+r}{n}$. Then

$$0.14 < \frac{k-r}{n} = \frac{k-r}{k+r} \cdot \frac{k+r}{n} \leq \frac{k-r}{k+r} \cdot \frac{r}{k} = \frac{x(1-x)}{1+x}.$$

From which we get that $x > 1/5$. Secondly, $m_n \geq \alpha(G) \geq k$ since we can assume that $\mu_2 > 0$. (Note that $\mu_2 > 0$ if G is not a blow-up of a complete graph.) Hence $kn = 2e(G) \geq m_n \mu_n^2 \geq kr^2$. So we have $n > r^2$. So we have two inequalities: $0.14 < \frac{k-r}{n} = \frac{k(1-x)}{n}$ and $n > r^2$. Then $n > r^2 = (kx)^2 > x^2 \frac{0.14^2}{(1-x)^2} n^2$. Thus $\frac{1}{0.14^2} \frac{(1-x)^2}{x^2} > n$. Since $x > 1/5$ we have $\frac{(1-x)^2}{x^2} < 16$. Hence $n < \frac{16}{0.14^2} \approx 816.33$. Now we can finish the proof since we know all possible strongly regular graph parameters up to 816, see Brouwer's website [3] and for the triangle-free strongly regular graph parameters the table on the next page. One can check that indeed the Higman-Sims graph achieves the maximum of $(\mu_1(G) + \mu_n(G))/n$. □

Remark 3.1. An interesting thing arises from the table on Andries Brouwer's website. If there were a strongly regular graph G with parameters $(28, 9, 0, 4)$ then for this graph G we would have

$$\frac{\mu_1(G) + \mu_n(G)}{v(G)} = \frac{9 + (-5)}{28} = \frac{1}{7} > 0.14.$$

It is known that there is no such strongly regular graph just as there is no strongly regular graph with parameters $(64, 21, 0, 10)$. For this graph we would have

$$\frac{\mu_1(G) + \mu_n(G)}{v(G)} = \frac{21 + (-11)}{64} = \frac{10}{64} > \frac{1}{7} > 0.14.$$

n	k	a	b	ϑ_1	ϑ_2	m_1	m_2	$\frac{k+\vartheta_2}{n}$	Appr.	Existence
5	2	0	1	$\frac{\sqrt{5}-1}{2}$	$\frac{-\sqrt{5}-1}{2}$	2	2	$\frac{3-\sqrt{5}}{10}$	0.076	Yes
10	3	0	1	1	-2	4	5	$\frac{1}{10}$	0.1	Yes
16	5	0	2	1	-3	10	5	$\frac{2}{16}$	0.125	Yes
28	9	0	4	1	-5	21	6	$\frac{4}{28}$	0.142	No
50	7	0	1	2	-3	28	21	$\frac{4}{50}$	0.08	Yes
56	10	0	2	2	-4	35	20	$\frac{6}{56}$	0.106	Yes
64	21	0	10	1	-11	56	7	$\frac{10}{64}$	0.156	No
77	16	0	4	2	-6	55	21	$\frac{10}{77}$	0.129	Yes
100	22	0	6	2	-8	77	22	$\frac{14}{100}$	0.14	Yes
162	21	0	3	3	-6	105	56	$\frac{15}{162}$	0.092	?
176	25	0	4	3	-7	120	55	$\frac{18}{176}$	0.102	?
210	33	0	6	3	-9	154	55	$\frac{24}{210}$	0.114	?
266	45	0	9	3	-12	209	56	$\frac{33}{266}$	0.124	?
324	57	0	12	3	-15	266	57	$\frac{42}{324}$	0.129	No
352	26	0	2	4	-6	208	143	$\frac{20}{352}$	0.056	?
352	36	0	4	4	-8	231	120	$\frac{28}{352}$	0.079	?
392	46	0	6	4	-10	276	115	$\frac{36}{392}$	0.091	?
552	76	0	12	4	-16	437	114	$\frac{60}{552}$	0.108	?
638	49	0	4	5	-9	406	231	$\frac{40}{638}$	0.062	?
650	55	0	5	5	-10	429	220	$\frac{50}{650}$	0.076	?
667	96	0	16	4	-20	551	115	$\frac{76}{667}$	0.113	?
704	37	0	2	5	-7	407	296	$\frac{30}{704}$	0.042	?
784	116	0	20	4	-24	667	116	$\frac{92}{784}$	0.117	?
800	85	0	10	5	-15	595	204	$\frac{70}{800}$	0.087	?

REFERENCES

- [1] József Balogh, Felix Christian Clemen, Bernard Lidický, Sergey Norin, Jan Volec: *The spectrum of triangle-free graphs*, ArXiv preprint 2204.00093
- [2] S. Brandt: *The local density of triangle-free graphs*, Discrete Math. **183** (1998), 17–25.
- [3] A. Brouwer: *Parameters of strongly regular graphs*, <https://www.win.tue.nl/~aeb/graphs/srg/srgtab.html>
- [4] L. de Lima, V. Nikiforov, C. Oliveira: *The clique number and the smallest Q -eigenvalue of graphs*, Discrete Mathematics **339**(6) (2016), 1744–1752.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS & EÖTVÖS LORÁND UNIVERSITY, DEPARTMENT
OF COMPUTER SCIENCE

Email address: `peter.csikvari@gmail.com`