

Correlations in typicality and an affirmative solution to the exact catalytic entropy conjecture

Henrik Wilming

Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany

It is well known that if a (finite-dimensional) density matrix ρ has smaller entropy than ρ' , then the tensor product of sufficiently many copies of ρ majorizes a quantum state arbitrarily close to the tensor product of correspondingly many copies of ρ' . In this short note I show that if additionally $\text{rank}(\rho) \leq \text{rank}(\rho')$, then n copies of ρ also majorize a state where all single-body marginals are *exactly identical* to ρ' but arbitrary correlations are allowed (for some sufficiently large n). An immediate application of this is an affirmative solution of the exact catalytic entropy conjecture introduced by Boes et al. [PRL 122, 210402 (2019)]: If $H(\rho) < H(\rho')$ and $\text{rank}(\rho) \leq \text{rank}(\rho')$ there exists a finite dimensional density matrix σ and a unitary U such that $U\rho \otimes \sigma U^\dagger$ has marginals ρ' and σ *exactly*. All the results transfer to the classical setting of probability distributions over finite alphabets with unitaries replaced by permutations.

Introduction and results

The exact catalytic entropy conjecture (CEC) is the following conjecture from Ref. [1] about the von Neumann entropy $H(\rho) := -\text{Tr}[\rho \log(\rho)]$ (log with base 2) illustrated in Fig. 1 (top):

Conjecture 1 (CEC). *Let ρ, ρ' be d -dimensional density matrices that are not unitarily equivalent. Then there exists a finite-dimensional density matrix σ and a unitary U such that*

$$\text{Tr}_1[U\rho \otimes \sigma U^\dagger] = \sigma, \text{Tr}_2[U\rho \otimes \sigma U^\dagger] = \rho' \quad (1)$$

if and only if $H(\rho) < H(\rho')$ and $\text{rank}(\rho) \leq \text{rank}(\rho')$.

An affirmative solution to the CEC provides a clean single-shot interpretation of von Neumann entropy without external randomness using the "catalyst" σ , see Refs. [1, 2] for discussions of the conjecture and their physical implications. It follows directly from the proof of the approximate version of the CEC in Ref. [2], that the CEC is true if the following Lemma holds true, illustrated in Fig. 1 (bottom):

Lemma 1. *Let ρ, ρ' be d -dimensional density matrices that are not unitarily equivalent. Then there exist $n \in \mathbb{N}$ and a density matrix ρ'_n on n copies of \mathbb{C}^d such that*

$$\rho^{\otimes n} \succeq \rho'_n, \quad \text{Tr}_{\{1, \dots, n\} \setminus i}[\rho'_n] = \rho' \quad \forall i = 1, \dots, n \quad (2)$$

if and only if $H(\rho) < H(\rho')$ and $\text{rank}(\rho) \leq \text{rank}(\rho')$.

Here, for any two density matrices ρ and σ , $\rho \succeq \sigma$ (" ρ majorizes σ ") means that there exists a collection of m unitary matrices U_i with associated probabilities p_i such that $\sigma = \sum_i p_i U_i \rho U_i^\dagger$. Equivalently, if λ_ρ denotes the vector of eigenvalues of ρ (including multiplicities) and λ_σ the corresponding vector of eigenvalues of σ , then there exists a collection of m permutation matrices Π_i and a probability distribution q_i such that $\lambda_\sigma = \sum_i q_i \Pi_i \lambda_\rho$.

In words the Lemma reads: if the entropy of ρ' is higher than that of ρ (and the rank not smaller), then for some finite n the state $\rho^{\otimes n}$ majorizes a state whose one-body marginals are all equal to ρ' . It is well known that for sufficiently large n the state $\rho^{\otimes n}$ majorizes an ϵ -approximation of $\rho'^{\otimes n}$. Lemma 1 instead shows that $\rho^{\otimes n}$ majorizes a state whose marginals coincide exactly with ρ' , but where arbitrary correlations are allowed.




FIG. 1. Top (CEC): A unitary operation U is applied to systems S and C in the state $\rho \otimes \sigma$. The resulting reduced state on S is exactly ρ' , while the reduced state on C is preserved exactly, but correlated to S (indicated by the dashed lines). Both σ and U may depend on ρ and ρ' . Bottom (Lemma 1): For some finite n , the state $\rho^{\otimes n}$ majorizes a state ρ'_n with all marginals equal to ρ' . The results of this paper show that if ρ and ρ' are not unitarily equivalent, then both situations occur if and only if $H(\rho) < H(\rho')$ and $\text{rank}(\rho) \leq \text{rank}(\rho')$. Both results transfer to the classical setting as well.

Our aim here is to prove Lemma 1 and thereby the exact catalytic entropy conjecture. In fact, the arguments below can also all be transferred to the classical setting, where density matrices are replaced by finite-dimensional probability vectors and unitary transformations by permutations of the entries of the vectors. Using the corresponding classical construction in Ref. [2] this implies that the classical version of the CEC also holds (contrarily to some prior beliefs):

Theorem 2 (classical CEC). *Let $\mathbf{p}, \mathbf{p}' \in \mathbb{R}^d$ be probability vectors that are not related by a permutation of their entries. Then there exists a finite-dimensional probability vector \mathbf{q} and a permutation matrix Π such that*

$$\text{Tr}_1[\Pi \mathbf{p} \otimes \mathbf{q}] = \mathbf{q}, \quad \text{Tr}_2[\Pi \mathbf{p} \otimes \mathbf{q}] = \mathbf{p}' \quad (3)$$

if and only if $H(\mathbf{p}) < H(\mathbf{p}')$ and $\text{rank}(\mathbf{p}) \leq \text{rank}(\mathbf{p}')$, with the Shannon entropy $H(\mathbf{p}) = -\sum_i p_i \log(p_i)$.

In the formulation of the theorem Tr_x denotes marginalization over tensor-factor x and $\text{rank}(\mathbf{x})$ denotes the number of non-zero elements of a vector \mathbf{x} .

In fact, contrary to the usual situation where results in quantum theory imply corresponding classical results, Theorem 2 implies the quantum version of the CEC but not vice-versa. However, as in the approximate case, the dimension required for the catalyst can be significantly smaller in the quantum case compared to the classical case, see Ref. [2].

Resource theories

In Refs. [3–6] catalysts that become correlated to the system of interest have been used to provide single-shot interpretations of the non-equilibrium free energy in quantum thermodynamics and the entanglement entropy in the framework of local operations and classical communication. These results were approximate, in the sense that the output state of a protocol only approximated the desired state up to arbitrary accuracy as the catalyst dimension diverged. Our result may suggest that a *strict* decrease in the free energy or entanglement entropy could imply the possibility of an exact transformation with a finite-dimensional catalyst also in those settings. However, Ref. [7] showed already that this is not the case, at least if one demands that the final correlations between catalyst and system can be made arbitrarily small. In the setting of this note, we cannot make this assumption since the mutual information between catalyst and final state of the system is always given by $H(\rho') - H(\rho)$. It would be interesting to know whether in the context of quantum thermodynamics a strict decrease in free energy allows for an exact conversion with finite-dimensional catalyst if one does not put a condition on the final correlations.

Open problems

Lemma 5 does not claim that the state ρ'_n is close to $\rho'^{\otimes n}$. It is an open problem whether one can both arrange that the marginals of ρ'_n are exactly equal to ρ' and ρ'_n is arbitrarily close to $\rho'^{\otimes n}$ in trace-distance while keeping the majorization condition $\rho'^{\otimes n} \succeq \rho'_n$. If true, this statement could likely have many further applications in physics and information theory.

A second open problem concerns the case $H(\rho) = H(\rho')$. It is known from lower bounds on the catalyst dimension that in this case a finite-dimensional catalyst in general cannot be used to implement the state-transition exactly [8, 9]. However, it is presently unknown whether a properly infinite dimensional catalyst could be used to implement the state-transition $\rho \rightarrow \rho'$ exactly in this case.

The proof

We formulate the proof in the quantum language, but, as mentioned above, everything transfers to the classical case by replacing density matrices with probability vectors.

Definition 3. For every finite-dimensional density matrix ρ and $n \in \mathbb{N}$, we define the following set:

$$S_n(\rho) := \left\{ \frac{1}{n} \sum_i \sigma_{\{i\}} \mid \rho^{\otimes n} \succeq \sigma, \sigma \geq 0, \text{Tr}[\sigma] = 1 \right\}, \quad (4)$$

where $\sigma_{\{i\}} = \text{Tr}_{\{1, \dots, n\} \setminus \{i\}}[\sigma]$.

Since randomly permuting subsystems is a random unitary process (in the sense of majorization), whenever $\rho' \in S_n(\rho)$ there exists a state ρ'_n with $\rho'^{\otimes n} \succeq \rho'_n$ as in Lemma 1. So we

want to show that whenever $H(\rho') > H(\rho)$ and $\text{rank}(\rho) \leq \text{rank}(\rho')$ then $\rho' \in S_n(\rho)$ for some $n \in \mathbb{N}$.

Lemma 4. For every ρ on \mathbb{C}^d , $S_n(\rho)$ is a compact convex set within the set of density matrices on \mathbb{C}^d and $\mathbb{1}/d \in S_n(\rho)$. Furthermore, $S_n(\rho)$ has full dimension within the set of density matrices.

Proof. The set of density matrices σ such that $\rho^{\otimes n} \succeq \sigma$ is closed and convex and hence compact. $S_n(\rho)$ is convex and compact, because it is its image under a linear (and hence continuous) map. It has full dimension since in particular it includes an open ball around $\mathbb{1}/d$ (and hence also $\mathbb{1}/d$). \square

We now introduce a form of typicality. Note that the statement is not in the usual form, as we claim a statement for all density matrices with certain entropies. In other words, we require an error bound that is uniform over *all* density matrices with a certain minimal entropy.

Lemma 5 (Typicality). For every $\delta > 0$ and $\epsilon > 0$ there exists $n_\epsilon \in \mathbb{N}$ such that for all states ρ' with $H(\rho') > H(\rho) + 2\delta$,

$$d(\rho', S_{n_\epsilon}(\rho)) \leq \epsilon, \quad (5)$$

where $d(\rho, \sigma) = \|\rho - \sigma\|_1 / 2$ denotes the trace-distance.

Proof. In general, a density matrix ρ majorizes an ϵ -approximation of ρ' if $H_{\max}^{\epsilon/2}(\rho) < H_{\min}^{\epsilon/2}(\rho')$, where H_{\max}^ϵ is the smooth max-entropy and H_{\min}^ϵ the smooth min-entropy, see [10, 11]. By Theorem 1 of Ref. [12] (in Ref. [12], the bounds are stated for conditional entropies of classical random variables, but in the unconditional case, they generalize immediately to the quantum case), it follows that for every $\delta \geq 0$ we have

$$H_{\max}^{\epsilon/2}(\rho^{\otimes n}) \leq n(H(\rho) + \delta), \quad (6)$$

$$H_{\min}^{\epsilon/2}(\rho'^{\otimes n}) \geq n(H(\rho') - \delta) \quad (7)$$

for $\epsilon = 2^{-\frac{n\delta^2}{2 \log^2(d+3)}}$ (which does not depend on ρ' or ρ) [13]. Thus, if we take

$$n_\epsilon = \left\lceil \frac{2 \log^2(d+3)}{\delta^2} \log\left(\frac{1}{\epsilon}\right) \right\rceil, \quad (8)$$

we find that $\rho^{\otimes n_\epsilon}$ majorizes an ϵ -approximation $\rho'_{\epsilon, n_\epsilon}$ of $\rho'^{\otimes n_\epsilon}$ as long as $H(\rho') > H(\rho) + 2\delta$. By the triangle inequality and monotonicity of the trace-distance under quantum channels, we find:

$$d\left(\frac{1}{n_\epsilon} \sum_{i=1}^{n_\epsilon} \text{Tr}_{\{1, \dots, n_\epsilon\} \setminus \{i\}}[\rho'_{\epsilon, n_\epsilon}], \rho'\right) \leq \epsilon. \quad (9)$$

Thus $d(\rho', S_{n_\epsilon}(\rho)) \leq \epsilon$. \square

The crucial additional ingredient to our proof apart from the typicality Lemma is the following simple observation about convex geometry.

Lemma 6. *Let $y \in \mathbb{R}^d$. Let $B_y(\delta)$ denote an open ball of radius δ around y and let C_ϵ be a parametrized family of closed and convex sets with the following property:*

$$\forall x \in B_y(\delta) \quad d(x, C_\epsilon) \leq \epsilon, \quad (10)$$

where d is the euclidean distance on \mathbb{R}^d . Then $y \in C_\epsilon$ for all $\epsilon < \delta/2$.

Proof. Let $\epsilon < \delta/2$ and suppose contrarily that $y \notin C_\epsilon$. Let c be the element in C_ϵ closest to y (which is unique because of convexity of C_ϵ). But since y is the center of the ball, there must then be an element $z \in B_y(\delta)$ with $\epsilon \geq d(z, C_\epsilon) \geq d(z, y) > \delta/2$, which is a contradiction. To find z , shoot a ray through y and c . The part of the ray inside of $B_y(\delta)$ has length at least δ , so in particular there must be a corresponding element z with $d(z, c) > \delta/2$. \square

Corollary 7. *Let C be a full-dimensional convex set in a normed, finite-dimensional, real vector space V and let $C_\epsilon \subseteq V$ be a parametrized family of closed and convex sets with the following property:*

$$\forall x \in C \quad d(x, C_\epsilon) < \epsilon, \quad (11)$$

where d is the distance induced by the norm. Then for any $y \in \text{int } C$ there exists an ϵ_0 such that $y \in C_\epsilon$ for all $\epsilon < \epsilon_0$.

Proof. Since y is an interior point, there is an open ball $B_y(\delta)$ around it and we can apply Lemma 6 (after embedding everything in some \mathbb{R}^d). \square

We can now prove Lemma 1: We start with the "if" part. By unitary freedom, we can choose ρ and ρ' so that $\text{supp}(\rho) \subseteq \text{supp}(\rho')$ and instead of the rank-condition we can then assume $\rho' > 0$, since we can simply shrink Hilbert-space to the support of ρ' . We will do so in the following. So let $H(\rho') - H(\rho) = 2\delta > 0$ and $\rho' > 0$. Then ρ' is an interior point of the convex set C of density matrices with entropy $\geq H(\rho) + \delta$ (which is full-dimensional within the convex set of density matrices). For any $\epsilon > 0$ set $C_\epsilon = S_{n_\epsilon}(\rho)$ with n_ϵ from Lemma 5. Then by the same Lemma, the conditions of

Corollary 7 are fulfilled if we set $y = \rho'$. Hence there exists some n such that $\rho' \in S_n(\rho)$.

For the "only if" part let us start with the condition on the von Neumann entropy: We use that H is sub-additive, $H(\rho_{12}) \leq H(\rho_1) + H(\rho_2)$ with equality if and only if $\rho_{12} = \rho_1 \otimes \rho_2$, and strictly Schur-concave, meaning that if $\rho \succeq \rho'$ but ρ and ρ' are not unitarily equivalent, then $H(\rho) < H(\rho')$. So suppose that $\rho^{\otimes n} \succeq \rho'^{\otimes n}$ and ρ and ρ' are not unitarily equivalent. Then we have

$$nH(\rho) = H(\rho^{\otimes n}) \leq H(\rho'^{\otimes n}) \leq nH(\rho'). \quad (12)$$

We now assume $H(\rho) = H(\rho')$ to arrive at a contradiction. Then (12) implies $H(\rho'^{\otimes n}) = nH(\rho')$, which in turn implies $\rho'^{\otimes n} = \rho^{\otimes n}$. But then $\rho^{\otimes n} \succeq \rho'^{\otimes n}$ and $H(\rho) = H(\rho')$, which by strict Schur-concavity implies that $\rho^{\otimes n}$ and $\rho'^{\otimes n}$ are unitarily equivalent. But then ρ and ρ' are unitarily equivalent, which yields the contradiction. Hence $H(\rho) < H(\rho')$. To conclude $\text{rank}(\rho) \leq \text{rank}(\rho')$, we use that $H_0(\rho) := \log(\text{rank}(\rho))$ is sub-additive [14] and additive over-tensor factors as well as non-decreasing under majorization. Hence

$$H_0(\rho^{\otimes n}) = nH_0(\rho) \leq H_0(\rho'^{\otimes n}) \leq nH_0(\rho'), \quad (13)$$

which finishes the proof.

Acknowledgements

I would like to thank Niklas Galke for rekindling my interest in this problem and pointing out a mistake in a previous version of the argument. I would also like to thank Paul Boes, Thomas Cope, Patryk Lipka-Bartosik, Nelly H.Y. Ng and Reinhard F. Werner for discussions. Support by the DFG through SFB 1227 (DQ-mat), Quantum Valley Lower Saxony, and funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germanys Excellence Strategy EXC-2123 QuantumFrontiers 390837967 is also acknowledged.

-
- [1] P. Boes, J. Eisert, R. Gallego, M. P. Müller, and H. Wilming, *Physical Review Letters* **122**, 210402 (2019).
[2] H. Wilming, *Physical Review Letters* **127**, 260402 (2021).
[3] M. P. Müller, *Physical Review X* **8**, 041051 (2018).
[4] N. Shiraishi and T. Sagawa, *Physical Review Letters* **126**, 150502 (2021).
[5] T. V. Kondra, C. Datta, and A. Streltsov, *Physical Review Letters* **127**, 150503 (2021).
[6] P. Lipka-Bartosik and P. Skrzypczyk, *Physical Review Letters* **127**, 080502 (2021).
[7] R. Rubboli and M. Tomamichel, (2021), <https://doi.org/10.48550/arXiv.2111.13356>, arXiv:2111.13356 [quant-ph].
[8] P. Boes, N. H. Ng, and H. Wilming, *PRX Quantum* **3**, 010325 (2022).
[9] V. Blakaj and M. M. Wolf, (2021), <https://doi.org/10.48550/arXiv.2111.10363>, arXiv:2111.10363 [math-ph].
[10] R. Renner, *Security of Quantum Key Distribution*, Ph.D. thesis, ETH Zurich (2005).
[11] M. Tomamichel, *Quantum Information Processing with Finite Resources* (Springer International Publishing, 2016).
[12] T. Holenstein and R. Renner, *IEEE Transactions on Information Theory* **57**, 1865 (2011).
[13] We define the smoothing of the entropies in terms of trace-distance instead of 1-norm, which accounts for a factor of 1/2.
[14] N. Linden, M. Mosonyi, and A. Winter, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **469**, 20120737 (2013).