

Gravitational Analysis of Rotating Charged Black Hole-Like Solution in Einstein-Gauss-Bonnet Gravity

Riasat Ali,^{1,*} Rimsha Babar,^{2,†} and Muhammad Asgher^{3,‡}

¹*Department of Mathematics, GC, University Faisalabad Layyah Campus, Layyah-31200, Pakistan*

²*Division of Science and Technology, University of Education, Township, Lahore-54590, Pakistan*

³*Department of Mathematics, The Islamia University of Bahawalpur, Bahawalpur-63100, Pakistan*

(Dated: May 19, 2022)

This work analyzes the Einstein-Gauss-Bonnet gravity of charged black hole solutions through Newman-Janis approach. We also compute the Hawking temperature of Einstein-Gauss-Bonnet gravity of charged rotating black hole. The solution depends upon rotation parameter a (occurs due to Newman-Janis algorithm), black hole mass, black hole charge, black hole horizon and arbitrary constant. Moreover, we discuss the graphical behavior of temperature via event horizon to analyze the stability of black hole under the effects of rotation parameter. We plot the graphs in the presence and absence of rotation parameter and charge of BH and observe that BH depicts more stable form in the absence of charge. Furthermore, we study the Hawking temperature under the effects of quantum gravity by using the semi-classical method as well as for different values of quantum gravity parameter, we analyze the graphical behaviour of corrected temperature with respect to the event horizon. In the plots, we also observe that the maximum temperature at non-zero horizon depicts the BH remnant. Finally, we compute the logarithmic corrected entropy for Einstein-Gauss-Bonnet gravity of rotating charged black hole and also study the logarithmic corrected entropy with the effects of the rotation parameter and correction parameter.

Keywords: Einstein-Gauss-Bonnet gravity theory of black hole; Newman-Janis algorithmic rule; Lagrangian equation; Hawking Temperature; Logarithmic corrected entropy.

I. INTRODUCTION

In order to study the gravity theories of higher-order curvature [1, 2], a few modifications have been introduced in general relativity (GR) theory, a well known theory is called Einstein Gauss–Bonnet (EGB) gravity theory. This theory is actually a rare case of Lovelock gravity and also a modification of GR. The EBG gravity Lagrangian is a special type of the Lovelock gravity Lagrangian [3]. One interesting feature of EGB is that it maintains the second-order equation of motion in self-assertive number of dimensions $D \geq 5$. This significant feature gives the premise to research on gravity in the other structure. Although, because of the metric definition in dimensions $D > 4$, the extent of exploration is restricted to space-time with higher dimensions. By re-scaling the coupling constant α , Glavan and Lin constructed 4-dimensions of EGB gravity [4]. They showed if we take the product of GB action with a factor $\frac{\alpha}{D-4}$ rather than α , then we will consider an unchanged contribution to Einstein's equations in dimension $D = 4$. After this presented method, several investigations about the EGB gravity and its thermodynamics have been done. Many investigations about the GB gravity as well as its properties have been made after presented this valuable method [5, 6]. The topological BH solutions of GB gravity have been derived using two kinds of nonlinear electrodynamics [7] and also discussed that the stability of heat capacity. Clifton and his colleagues [8] searched for the observational constraints on regular EGB gravity in 4 dimensions and find out the range of coupling constant. The following features about black holes (BHs) i.e., radiation, instability, shadows, quasinormal modes and grey-body factors in 4 dimensions EGB gravity have been studied in [9]-[13]. The most recent twenty years had a great deal of treat for theoretical cosmologists, coming from both cosmological scale information and furthermore from astrophysical scales occasions. Especially, the perception of the right now speeding up Universe coming from the standard candles SNe Ia [14], has completely changed our discernment of how the Universe evolves. The gravitational wave recognition coming from the neutron star consolidating GW170817 event [15], has totally influenced modified speculations of gravity, barring a few of these from being reasonable depictions of our Universe at astrophysical scales. Especially, the GW170817 occasion showed that the proliferating rate of the gravitational waves c_T is equivalent to unity to

*Electronic address: riasatyasin@gmail.com

†Electronic address: rimsha.babar10@gmail.com

‡Electronic address: m.asgher145@gmail.com

that of light, specifically $c_T^2 \simeq 1$, in standard units. This factor as we specified, avoided quickly numerous elective speculations of gravity. Although, there exist many altered gravity speculations that actually stay strong against the GW170817 event results. Oikonomou and his colleagues [16]-[20] have explored how the EBG theories can be rendered reasonable and viable with the GW170817 event. The gravitational wave speed for an EBG theory is equal to, $c_T^2 = 1 - \frac{Q_f}{2Q_t}$ with $Q_f = 8(\ddot{\xi} - H\dot{\xi})c_1$, while c_1 represents the dimensionless constant multiplication factor that work in the Lagrangian with GB term, and the function Q_t is a function of H where H stands for Hubble rate. So as indicated by our supposition, if $Q_f = 0$ the gravitational wave speed would be equivalent to one. This infers that the GB coupling must holds the differential condition $\ddot{\xi} - H\dot{\xi} = 0$, which suggests that $\dot{\xi} = e^N$, here N denotes the e-foldings number.

This technique empowered us to communicate all the slow-roll as well as the observational indices as elements of the e-foldings number, and in the long run concentrate on the phenomenology of the demonstrate.

Semiclassical techniques based on Hawking radiation as a tunneling impact were introduced in the course of the last two decade and have gained a great deal of interest. This approach has been presented by Hawking at the point when he proposed his interesting theoretical disclosure named "Hawking radiation" [21] and after this, it has been clarified by Parikh and Wilczek [22, 23] that how Hawking radiation occurs. The physical significance of this emission action shows that vacuum fluctuations accelerate particle (positive mass) anti-particle (negative mass) pairs towards the BH horizon. Hawking studied that positive mass particles have enough energy to radiate from the BH, while negative mass particles have no capability to emit from the BH. Parikh and Wilczek provided a complete path to this theoretical discovery in the form of WKB approximation. This method utilize geometric optic estimation which is another perspective of eikonal estimation in wave explanation [24]. Particles are set in front of a boundary, when particles tunnel from this boundary the BH mass reduces in the form of particles energy. In spite of the fact that with the Parikh-Wilczek approach, a more precise strategy of the tunneling was given, but there were as yet unsolved issues like information loss, temperature divergence and unitary. Numerous investigations have been done on the radiation and tunneling approach from the different BH horizons; Some of these most significant investigations can be seen in [25]-[55]. The tunneling radiation for diferrent BHs have been studied and also examined the tunneling radiation with the effects of the BH geometry and various parameter's. It is possible to discuss quantum-corrected thermodynamical features of BH by integrating generalised uncertainty principle (GUP) effects [56]. The GUP gives high-energy solution to BH thermodynamics, allowing quantum gravity theory to have a minimal length. By taking into account the GUP effects, it is very feasible to study the quantum modified thermodynamics of BHs. The GUP relation satisfies the following expression [50, 57]

$$\Delta x \Delta p \geq \frac{1}{2} \left(1 + \alpha (\Delta p)^2 \right).$$

here $\alpha = \frac{\alpha_0}{M_p^2}$, M_p shows the Plank's mass and $\alpha_0 < 10^{34}$ represents the dimensionless parameter.

The main purpose of this paper is to investigate the charged BH in EGB gravity for rotating case and then to discuss its thermodynamics i.e., Hawking temperature. Moreover, to check the stability condition of charged rotating BH in EGB gravity by graphical interpretation of Hawking temperature with horizon. In order to meet our aim, we attempt to construct the rotating charged BH solutions of EBG gravity with the help of Newman-Janis algorithm and then investigate their Hawking temperature and study the effects of rotation parameter with the help of plots. Furthermore, by using the quantum tunneling approach, we also study their quantum corrected Hawking temperature by using the semi-classical method with the help of WKB approximation. At last, we analyze the gravity effects by plotting the graphs of corrected temperature with horizon, which depicts the stability of BH in rotating case. This article is arranged as follows, Sec. **II** contains the spacetime information of D -dimensional BHs. Subsection **II. A**, investigate the solution of rotating charged BH in EBG gravity by using the Newman-Janis method and study the temperature for the associated BH. Sec. **III**, gives the graphical temperature behaviour w.r.t event horizon and analyze the effects of rotation parameter. Sec. **IV**, check the quantum corrected temperature for EGB gravity of charged rotating BH by using quantum tunneling method for boson particles in 4 dimension spaces. In Sec. **V**, we study the quantum gravity graphical effects on corrected Hawking temperature for boson particles. Section **VI** describes the logarithmic corrected entropy for EGB gravity of rotating charged BH. In Sec. **VII**, we summarize the result of our work.

II. EINSTEIN GAUSS BONNET GRAVITY THEORY OF BLACK HOLES

In this section, we examine the Einstein-Gauss-Bonnet D -dimension charged BHs gravity theory. To do so, we consider the EGB BHs gravity theory with Lagrangian equation,

$$\mathbb{L} = (R^2 + R_{abcd}R^{abcd}\hat{\beta} - 4R_{ab}R^{ab}) + R.$$

The solution for spherically symmetric charged BH with this gravity theory in D -dimension space-time can be given as [58]

$$ds^2 = -G(r)dt^2 + G^{-1}(r)dr^2 + r^2d\Omega_{d-2}^2, \quad (1)$$

where

$$G(r) = 1 + \left[1 - \sqrt{1 + \frac{16\beta M}{r^{d-1}} - \frac{8\beta q^2}{r^{2d-4}}} \right] \frac{r^2}{4\beta'}$$

where M , q and β are BH mass, BH charge and the BH coupling constant $\hat{\beta}$ as $\beta = (d-3)(d-4)\hat{\beta}/2$, respectively. We study the space-time by $d\Omega_{d-2}^2$ of the unit $(d-2)$ area of sphere A_{d-2} . The only nonzero component of the electromagnetic potential has the following form

$$A_t(r) = -\frac{q}{r^{d-3}}. \quad (2)$$

Here, M and q are associated to the BH mass \hat{M} and Arnowitt-Deser-Misner BH charge(\hat{q}) as

$$q^2 = \frac{2\hat{q}^2(d-3)}{d-2}, \quad M = \frac{8\pi}{A_{d-2}(d-2)}\hat{M}.$$

The horizon solution at $r = r_h$ can be obtained from the zeros of the function by setting $G(r_h) = 0$

$$r_h^{2(d-3)} + 2\beta r_h^{2(d-4)} - 2Mr_h^{d-3} + q^2 = 0. \quad (3)$$

The above equation gives the two real root ($r_h = r_{\pm}$), the positive sign denotes the largest root as well as negative sign describes the lowest root.

A. Hawking Temperature for Rotating Charged Black Hole

The spherically symmetric of charged BH solution in 4-dimensional space-time has the following form

$$ds^2 = -G(r)dt^2 + G^{-1}(r)dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\psi. \quad (4)$$

By the coordinates transformation (t, r, θ, ϕ) to (u, r, θ, ϕ) , we have

$$du = dt - \frac{dr}{G(r)}. \quad (5)$$

After applying the transformation to the metric Eq. (4), we get

$$ds^2 = -G(r)du^2 - 2dudr + r^2d\theta^2 + r^2\sin^2\theta d\phi^2. \quad (6)$$

The metric in the null (EF) frame can be expressed as

$$g^{\mu\nu} = -l^\nu n^\mu - l^\mu n^\nu + m^\nu \bar{m}^\mu + m^\mu \bar{m}^\nu. \quad (7)$$

The corresponding components are given as

$$\begin{aligned} l^\mu &= \delta_r^\mu, \quad n^\nu = \delta_u^\nu - \frac{1}{2}G\delta_r^\nu, \\ m^\mu &= \frac{1}{\sqrt{2}r}\delta_\theta^\mu + \frac{i}{\sqrt{2}r\sin\theta}\delta_\phi^\mu, \\ \bar{m}^\mu &= \frac{1}{\sqrt{2}r}\delta_\theta^\mu - \frac{i}{\sqrt{2}r\sin\theta}\delta_\phi^\mu. \end{aligned}$$

For all point in the BH metric the null vectors of the null tetrad satisfy the conditions of $l_\mu l^\mu = n_\mu n^\mu = m_\mu m^\mu = \bar{m}_\mu \bar{m}^\mu = 0$ and $l_\mu n^\mu = -m_\mu \bar{m}^\mu = 1$ in the place of (u, r) coordinate transformation are $u \rightarrow u - ia \cos \theta$, $r \rightarrow r + ia \cos \theta$, then we perform the transformation $G(r) \rightarrow \tilde{G}(r, a, \theta)$ and $\Sigma^2 = a^2 \cos^2 \theta + r^2$. The null vectors in the (u, r) space can be written as

$$\begin{aligned} l^\mu &= \delta_r^\mu, \quad n^\mu = \delta_u^\mu - \frac{1}{2} \tilde{G} \delta_r^\mu, \\ m^\mu &= \frac{1}{\sqrt{2\Sigma}} \left(\delta_\theta^\mu + ia \sin \theta (\delta_u^\mu - \delta_r^\mu) + \frac{i}{\sin \theta} \delta_\phi^\mu \right), \\ \bar{m}^\mu &= \frac{1}{\sqrt{2\Sigma}} \left(\delta_\theta^\mu - ia \sin \theta (\delta_u^\mu - \delta_r^\mu) - \frac{i}{\sin \theta} \delta_\phi^\mu \right), \end{aligned}$$

From the null tetrad definition of the metric tensor $g^{\mu\nu}$ are given by

$$\begin{aligned} g^{uu} &= \frac{a^2 \sin^2 \theta}{\Sigma^2}, \quad g^{ur} = g^{ru} = -1 - \frac{a^2 \sin^2 \theta}{\Sigma^2}, \quad g^{rr} = \tilde{G} + \frac{a^2 \sin^2 \theta}{\Sigma^2}, \quad g^{\theta\theta} = \frac{1}{\Sigma^2}, \\ g^{\phi\phi} &= \frac{1}{\Sigma^2 \sin^2 \theta}, \quad g^{u\phi} = g^{\phi u} = \frac{a}{\Sigma^2}, \quad g^{r\phi} = g^{\phi r} = -\frac{a}{\Sigma^2}. \end{aligned}$$

The new metric according to null tetrad can be obtained as

$$ds^2 = -\tilde{G}(r) du^2 - 2dudr - 2a \sin^2 \theta (1 - \tilde{G}) dud\phi + 2a \sin^2 \theta drd\phi + \Sigma^2 d\theta^2 + \sin^2 \theta (\Sigma^2 - a^2 \sin^2 \theta (\tilde{G} - 2)) d\phi^2. \quad (8)$$

Now, we consider the coordinate transformation from the Eddington Finkelstein (EF) to Boyer Lindquist (BL) coordinates as

$$du = dt + y(r)dr, \quad d\phi = d\phi + F(r)dr, \quad (9)$$

where

$$y(r) = \frac{a^2 + r^2}{r^2 \tilde{G} + a^2}, \quad F(r) = -\frac{a}{r^2 \tilde{G} + a^2}, \quad \tilde{G}(r, \theta) = \frac{a^2 \cos^2 \theta + r^2 G(r)}{\Sigma^2}. \quad (10)$$

Finally, we get the BH spacetime in the form

$$\begin{aligned} ds^2 &= - \left[1 + \frac{\frac{r^4}{4\beta} \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right)}{\Sigma^2} \right] dt^2 + \frac{\Sigma^2}{\Delta_r} dr^2 - 2a \sin^2 \theta \left[\frac{\frac{r^4}{4\beta} \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right)}{\Sigma^2} \right] dt d\phi \\ &+ \Sigma^2 d\theta^2 + a^2 \sin^2 \theta \left[\Sigma^2 - a^2 \sin^2 \theta \frac{\frac{r^4}{4\beta} \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right)}{\Sigma^2} \right] d\phi^2, \end{aligned} \quad (11)$$

here

$$\Delta_r = r^2 + a^2 + \frac{r^4}{4\beta} \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right).$$

Now, we discuss the thermodynamical property (i.e., Hawking temperature) for corresponding BH. In order to derive the Hawking temperature, we use the following formula for spherically symmetric BHs

$$T_H = \frac{\tilde{G}'(r_+)}{4\pi}. \quad (12)$$

The Hawking temperature for

$$T_H = \frac{r_+^5 (Y - r_+^2) - 4Mr_+^4 \beta + 2a^2 r_+ (4q^2 \beta + r_+^4 (Y - r_+^2) - 10Mr_+ \beta)}{8\pi Y \beta (r_+^2 + a^2)}, \quad (13)$$

where

$$Y = \sqrt{r_+^2 - 8q^2\beta + 16Mr_+\beta}.$$

The Hawking temperature depends upon the BH mass M , arbitrary constant β , BH charge q , spin parameter a and BH horizon radius r_+ .

III. STABILITY ANALYSIS VIA GRAPHICAL INTERPRETATION

This section investigate the stability of charged rotating BH with EBG gravity in the presence and absence of spin parameter a . We observe the horizon structure of BH by plotting the graphs of Δ_r with respect to horizon r_+ . as well as discuss the stability condition for BH under the effects of different parameters i.e., BH charge q and rotation parameter a in the range $0 \leq r_+ \leq 15$.

Fig. 1. shows the behavior of Δ_r with r_+ for fixed value of $M = 1$ (BH mass) and $\beta = 0.01$ (arbitrary parameter) in the presence and absence of spin parameter. In (i) the different values of spin parameter a for fixed values of BH charge $q = 1$ satisfy the relation of Δ_r with horizon equation by showing the positive behavior. In (ii) one can observe that in the absence of rotation parameter $a = 0$ and for different values of charge q , the behavior also satisfies the horizon relation.




Figure 1: Δ_r via horizon r_+ for $M = 1$, $\beta = 0.01$. Left $q = 1$, $a = 10$ (solid green), $a = 11$ (solid blue), $a = 12$ (solid red), $a = 13$ (solid cyan). Right $a = 0$, $q = 15$ (solid cyan), $q = 16$ (solid red), $q = 17$ (solid blue), $q = 18$ (solid green).




Figure 2: T_H via r_+ for $M = 10$. Left $\beta = 1$, $q = 0.5$, $a = 3$ (dashed), $a = 4$ (dot-dashed), $a = 5$ (small-dashed). Right $\beta = 0.1$, $a = 1$, $q = 5$ (dashed), $q = 5.5$ (dot-dashed), $q = 6$ (small-dashed).

Fig. 2. represents the temperature T_H behavior w.r.t horizon r_+ for fixed value of BH mass $M = 10$. In (i) the T_H shows the behavior for fixed values of BH charge $q = 0.5$, arbitrary parameter $\beta = 1$ and different values of rotation parameter a . One can observe that the temperature slowly goes on increasing with the increasing values of horizon but after a height it gradually goes on decreasing with the increasing horizon which satisfies the Hawking's phenomenon, so grantee's the physical and stable form of BH. One can also observe that with the increasing value of spin parameter the values of temperature increases.

The plot (ii) depicts the behavior of T_H for constants values of spin parameter $a = 1$, arbitrary parameter $\beta = 0.1$ and different values of charge q . We can observe at initial the BH is not stable but as time goes on BH attains its stable form, when it eventually drops down from a height and obtains an asymptotically flat state till $r_+ \rightarrow \infty$. The

decreasing temperature with increasing horizon also shows the stable condition of BH. One can observe with the increasing values of charge the values of temperature decreases.

Fig. 3. actually represents the behavior of temperature with event horizon for special case in the absence of spin parameter a or charge q . (i) shows the behavior of T_H for constant values of mass $M = 15$, arbitrary parameter $\beta = 0.01$ and different values of q (charge) in the absence of a (rotation parameter). One can analyze in the absence of rotation BH parameter also shows its stable form after passing some time. From Fig. 2 and 3, we can observe in the presence of rotation BH parameter the temperature increases as compare in the absence of a .

(ii) depicts the T_H behavior for changing values of rotation BH parameter a and fixed the parameter of $\beta = 1$, $M = 25$ in the absence of charge q . One can observe that in the absence of charge the BH is also in its stable form. We can also see that as compare to Fig. 2 (ii) the temperature is higher in Fig. 3 (ii) in the absence of charge.




Figure 3: T_H via r_+ . Left $M = 15, \beta = 0.01, a = 0, q = 1$ (dashed), $q = 2$ (dot-dashed), $q = 3$ (small-dashed). Right $M = 25, \beta = 1, q = 0, a = 5$ (dashed), $a = 6$ (dot-dashed), $a = 7$ (small-dashed).

IV. CORRECTED TUNNELING

The metric Eq. (11) can be re-expressed as

$$ds^2 = -\tilde{G}dt^2 + \tilde{H}dr^2 + \tilde{X}d\theta^2 + \tilde{Y}d\phi^2 + 2\tilde{Z}dtd\phi, \quad (14)$$

where

$$\tilde{G} = \left[1 + \frac{r^4 \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right)}{\Sigma^2} \right], \quad \tilde{Y} = a^2 \sin^2 \theta \left[\Sigma^2 - a^2 \sin^2 \theta \frac{r^4 \left(1 - \sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right)}{\Sigma^2} \right],$$

$$\tilde{Z} = a \sin^2 \theta \left[\frac{r^4 \left(\sqrt{1 - \frac{8\beta q^2}{r^4} + \frac{16\beta M}{r^3}} \right) - 1}{\Sigma^2} \right], \quad \tilde{H} = \frac{\Sigma^2}{\Delta_r}, \quad \tilde{X} = \Sigma^2.$$

The modified field equation is expressed by [50, 54]

$$\partial_\mu \left(\sqrt{-g} \phi^{\nu\mu} \right) + \sqrt{-g} \frac{m^2}{\hbar^2} \phi^\nu + \sqrt{-g} \frac{i}{\hbar} A_\mu \phi^{\nu\mu} + \sqrt{-g} \frac{i}{\hbar} e F^{\nu\mu} \phi_\mu + \hbar^2 \alpha \partial_0 \partial_0 \partial_0 \left(\sqrt{-g} g^{00} \phi^{0\nu} \right) - \alpha \hbar^2 \partial_i \partial_i \left(\sqrt{-g} g^{ii} \phi^{i\nu} \right) = 0, \quad (15)$$

here $\phi^{\nu\mu}$, m and g represent the anti-symmetric tensor, mass of particle and coefficient matrix of its determinant, since

$$\phi_{\nu\mu} = \left(1 - \alpha \hbar^2 \partial_\nu^2 \right) \partial_\nu \phi_\mu - \left(1 - \alpha \hbar^2 \partial_\mu^2 \right) \partial_\mu \phi_\nu + \left(1 - \alpha \hbar^2 \partial_\nu^2 \right) \frac{i}{\hbar} e A_\nu \phi_\mu - \left(1 - \alpha \hbar^2 \partial_\nu^2 \right) \frac{i}{\hbar} e A_\mu \phi_\nu, \quad \text{and } F_{\nu\mu} = \nabla_\nu A_\mu - \nabla_\mu A_\nu,$$

where α , e , A_μ and ∇_μ are the correction parameter, particle charge BH vector potential and covariant derivative, respectively. The anti-symmetric tensor with its non-zero components can be derived in the form

$$\begin{aligned}\phi^0 &= \frac{-\tilde{Y}\phi_0 + \tilde{Z}\phi_3}{\tilde{G}\tilde{Y} + \tilde{Z}^2}, \quad \phi^1 = \tilde{G}\phi_1, \quad \phi^2 = \frac{1}{\tilde{X}}\phi_2, \quad \phi^3 = \frac{\tilde{Z}\phi_0 + \tilde{G}\phi_3}{\tilde{G}\tilde{Y} + \tilde{Z}^2}, \quad \phi^{01} = \frac{\tilde{G}(-\tilde{D}\phi_{01} + \tilde{Z}\phi_{13})}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)}, \\ \phi^{02} &= \frac{-\tilde{Y}\phi_{02}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)}, \quad \phi^{03} = \frac{(-\tilde{G}\tilde{Y} + \tilde{G}^2)\phi_{03}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2}, \quad \phi^{12} = \frac{\tilde{G}}{\tilde{X}}\phi_{12}, \quad \phi^{13} = \frac{\tilde{G}}{\tilde{G}\tilde{Y} + \tilde{Z}^2}\phi_{13}, \quad \phi^{23} = \frac{\tilde{Z}\phi_{02} + \tilde{G}\phi_{23}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)}.\end{aligned}$$

The WKB approximation is

$$\phi_\nu = k_\nu \exp \left[\frac{i}{\hbar} I_0(t, r, \phi, \theta) + \Sigma \hbar^n I_n(t, r, \phi, \theta) \right]. \quad (16)$$

By putting all the components in Eq. (15), we get the following set of field equations in the form

$$\begin{aligned}& \frac{\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)(\partial_1 I_0)k_1 - (\partial_1 I_0)^2 k_0 - \alpha(\partial_1 I_0)^4 k_0 + eA_0(\partial_1 I_0)k_1 + \alpha(\partial_0 I_0)^3 (\partial_1 I_0)k_1 \right] \\ & + \alpha eA_0(\partial_0 I_0)^2 (\partial_1 I_0)k_1 \Big] - \frac{\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\alpha(\partial_1 I_0)^4 k_3 + (\partial_1 I_0)^2 k_3 - (\partial_1 I_0)(\partial_3 I_0)k_1 - (\partial_1 I_0)(\partial_3 I_0)^2 \alpha k_1 \right] \\ & + \frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)(\partial_2 I_0)k_2 + (\partial_0 I_0)^3 (\partial_2 I_0)\alpha k_2 - (\partial_2 I_0)^2 k_0 - (\partial_2 I_0)^4 \alpha k_0 + eA_0(\partial_2 I_0)k_2 \right. \\ & + eA_0\alpha(\partial_0 I_0)^2 (\partial_1 I_0)k_2 \Big] + \frac{\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[(\partial_0 I_0)(\partial_3 I_0)k_3 + (\partial_0 I_0)^3 (\partial_3 I_0)\alpha k_3 - (\partial_3 I_0)^2 k_0 - (\partial_3 I_0)^4 \alpha k_0 \right. \\ & + eA_0(\partial_3 I_0)k_3 + eA_0(\partial_0 I_0)^2 (\partial_3 I_0)k_3 \Big] - m^2 \left(\frac{\tilde{Y}k_0 - \tilde{Z}k_3}{\tilde{G}\tilde{Y} + \tilde{Z}^2} \right) = 0, \quad (17)\end{aligned}$$

$$\begin{aligned}& - \frac{\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)^2 k_1 + (\partial_0 I_0)^4 \alpha k_1 - (\partial_0 I_0)(\partial_1 I_0)k_0 - (\partial_0 I_0)(\partial_1 I_0)^3 \alpha k_0 + eA_0(\partial_0 I_0)k_1 \right. \\ & + eA_0(\partial_0 I_0)^3 \alpha k_1 \Big] + \frac{\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)(\partial_1 I_0)k_3 + (\partial_0 I_0)(\partial_1 I_0)^3 \alpha k_3 - (\partial_0 I_0)(\partial_3 I_0)k_1 \right. \\ & - (\partial_0 I_0)(\partial_3 I_0)^3 \alpha k_1 \Big] + \frac{\tilde{G}}{\tilde{X}} \left[(\partial_1 I_0)(\partial_2 I_0)k_2 + (\partial_1 I_0)(\partial_2 I_0)^3 \alpha k_2 - (\partial_2 I_0)^2 k_1 - (\partial_2 I_0)^4 \alpha k_1 \right] \\ & + \frac{\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_1 I_0)(\partial_3 I_0)k_3 + (\partial_1 I_0)(\partial_3 I_0)^3 \alpha k_3 - (\partial_3 I_0)^2 k_1 - (\partial_3 I_0)^4 \alpha k_1 \right] - \tilde{G}m^2 k_1 \\ & + \frac{eA_0\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)k_1 + (\partial_0 I_0)^3 \alpha k_1 - (\partial_1 I_0)k_0 - (\partial_1 I_0)^3 \alpha k_0 + eA_0k_1 + eA_0(\partial_0 I_0)^2 \alpha k_1 \right] \\ & + \frac{eA_0\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_1 I_0)k_3 + (\partial_1 I_0)^3 \alpha k_3 - (\partial_3 I_0)k_1 - (\partial_1 I_0)^3 \alpha k_1 \right] = 0, \quad (18)\end{aligned}$$

$$\frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)^2 k_2 + (\partial_0 I_0)^4 \alpha k_2 - (\partial_0 I_0)(\partial_2 I_0)k_0 - (\partial_0 I_0)(\partial_2 I_0)^3 \alpha k_0 + eA_0(\partial_0 I_0)k_2 \right]$$

$$\begin{aligned}
& + eA_0(\partial_0 I_0)^3 \alpha k_2 \Big] + \tilde{G}\tilde{X} \left[(\partial_1 I_0)^2 k_2 + (\partial_1 I_0)^4 \alpha k_2 - (\partial_1 I_0)(\partial_2 I_0)k_1 - (\partial_1 I_0)(\partial_2 I_0)^3 \alpha k_1 \right] - \frac{m^2 k_2}{\tilde{X}} \\
& - \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)(\partial_3 I_0)k_2 + (\partial_0 I_0)^3 (\partial_3 I_0)\alpha k_2 - (\partial_0 I_0)(\partial_3 I_0)k_0 - (\partial_0 I_0)^3 (\partial_3 I_0)\alpha k_0 + eA_0(\partial_3 I_0)k_2 \right. \\
& + eA_0(\partial_3 I_0)^3 \alpha k_2 \Big] + \frac{\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_2 I_0)(\partial_3 I_0)k_3 + (\partial_2 I_0)^3 (\partial_3 I_0)\alpha k_3 - (\partial_3 I_0)^2 k_2 - (\partial_3 I_0)^4 \alpha k_2 \right] \\
& + \frac{eA_0\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)k_2 + (\partial_0 I_0)^3 \alpha k_2 - (\partial_2 I_0)k_0 - (\partial_2 I_0)^3 \alpha k_0 + eA_0k_2 + eA_0\alpha(\partial_0 I_0)^2 k_2 \right] = 0, \tag{19} \\
& \frac{\tilde{G}\tilde{Y} - \tilde{G}^2}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[(\partial_0 I_0)^2 k_3 + (\partial_0 I_0)^4 \alpha k_3 - (\partial_0 I_0)(\partial_3 I_0)k_0 - (\partial_0 I_0)(\partial_3 I_0)^3 \alpha k_0 + eA_0(\partial_0 I_0)k_3 \right. \\
& + eA_0(\partial_0 I_0)^3 \alpha k_3 \Big] - \frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_1 I_0)^2 k_3 + (\partial_1 I_0)^4 \alpha k_3 - (\partial_1 I_0)(\partial_3 I_0)k_1 - (\partial_1 I_0)(\partial_3 I_0)^3 \alpha k_1 \right] \\
& - \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_0 I_0)(\partial_2 I_0)k_2 + (\partial_0 I_0)^3 (\partial_2 I_0)\alpha k_2 - (\partial_2 I_0)^2 k_0 + (\partial_2 I_0)^4 \alpha k_0 + eA_0(\partial_2 I_0)k_2 + eA_0\alpha \right. \\
& \left. (\partial_0 I_0)^2 (\partial_2 I_0)k_2 \right] - \frac{eA_0\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[(\partial_2 I_0)^2 k_3 + (\partial_2 I_0)^4 \alpha k_3 - (\partial_2 I_0)(\partial_3 I_0)k_2 - (\partial_0 I_0)(\partial_3 I_0)^3 \alpha k_2 \right] \\
& + \frac{eA_0(\tilde{G}\tilde{Y} - \tilde{G}^2)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[(\partial_0 I_0)k_3 + (\partial_0 I_0)^3 \alpha k_3 - (\partial_3 I_0)k_0 - (\partial_3 I_0)^3 \alpha k_0 + eA_0k_3 + eA_0(\partial_0 I_0)^2 \alpha k_3 \right] \\
& - \frac{m^2(\tilde{Z}k_0 - \tilde{G}k_3)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} = 0. \tag{20}
\end{aligned}$$

By utilizing the technique of separation of variables, we obtain

$$I_0 = - \left(E - \sum_{i=1}^2 j_i \Omega_i \right) t + W(r) + J\phi + \nu(\theta), \tag{21}$$

where E and J represent the particle energy, the particle angular momentum corresponding angle ϕ . After using the equation (21) into equations (17)-(20), we get a matrix of order 4×4 labeled as "U"

$$U(k_0, k_1, k_2, k_3)^T = 0.$$

The non-zero elements are expressed as follows

$$\begin{aligned}
U_{00} &= \frac{-\tilde{G}\tilde{D}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1^2 + \alpha W_1^4 \right] - \frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} [J^2 + \alpha J^4] - \frac{\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[\nu_1^2 + \alpha \nu_1^4 \right] - \frac{m^2 \tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)}, \\
U_{01} &= \frac{-\tilde{G}\tilde{D}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha \tilde{E}^3 + eA_0 + \alpha eA_0 \tilde{E}^2 \right] W_1 + \frac{\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} + [\nu_1 + \alpha \nu_1^3], \\
U_{02} &= \frac{-\tilde{D}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha \tilde{E}^3 - eA_0 - \alpha eA_0 \tilde{E}^2 \right] J, \quad K_{12} = \frac{\tilde{G}}{\tilde{X}} [W_1 + \alpha W_1^3] J, \\
H_{03} &= \frac{-\tilde{G}\tilde{E}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1^2 + \alpha W_1^4 \right] - \frac{\tilde{G}\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[\tilde{E} + \alpha \tilde{E}^3 - eA_0 - \alpha eA_0 \tilde{E}^2 \right] \nu_1 + \frac{m^2 \tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2},
\end{aligned}$$

$$\begin{aligned}
U_{10} &= \frac{-\tilde{G}\tilde{D}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}W_1 + \alpha\tilde{E}W_1^3 \right] - \tilde{G}m^2 - \frac{eA_0\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1 + \alpha W_1^3 \right], \\
U_{11} &= \frac{-\tilde{G}\tilde{D}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}^2 + \alpha\tilde{E}^4 - eA_0\tilde{E} - \alpha eA_0\tilde{E}W_1^2 \right] + \frac{\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1 + \alpha\nu_1^3 \right] \tilde{E} - \frac{\tilde{G}}{\tilde{X}} \left[J^2 + \alpha J^4 \right] \\
&\quad - \frac{\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1 + \alpha\nu_1^3 \right] - m^2\tilde{G} - \frac{eA_0\tilde{G}\tilde{Y}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha\tilde{E}^3 - eA_0 - \alpha eA_0\tilde{E}^2 \right] + \frac{eA_0\tilde{G}\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1 + \alpha\nu_1^3 \right], \\
U_{13} &= \frac{-\tilde{G}\tilde{E}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1 + \alpha W_1^3 \right] \tilde{E} + \frac{\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[W_1 + \alpha W_1^3 \right] \nu_1 + \frac{\tilde{G}\tilde{Z}eA_0}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1 + \alpha W_1^3 \right], \\
U_{20} &= \frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}J + \alpha\tilde{E}J^3 \right] + \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha\tilde{E}^3\nu_1^2 \right] - \frac{\tilde{Y}eA_0}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[J + \alpha J^3 \right], \\
U_{21} &= \frac{\tilde{G}}{\tilde{X}} \left[J + \alpha J^3 \right] W_1, \quad U_{23} = \frac{\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[J + \alpha J^3 \right] \nu_1, \\
U_{22} &= \frac{\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}^2 + \alpha\tilde{E}^4 - eA_0\tilde{E} - \alpha eA_0\tilde{E} \right] - \tilde{G}\tilde{X} + \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha\tilde{E}^3 - eA_0 - \alpha eA_0\tilde{E}^2 \right] \nu_1 \\
&\quad - \frac{m^2}{\tilde{X}} - \frac{\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1^2 + \alpha\nu_1^4 \right] - \frac{eA_0\tilde{Y}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha\tilde{E}^3 - eA_0 - \alpha eA_0\tilde{E}^2 \right], \\
U_{30} &= \frac{(\tilde{G}\tilde{Y} - \tilde{A}^2)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[\nu_1 + \alpha\nu_1^3 \right] \tilde{E} + \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[J^2 + \alpha J^4 \right] - \frac{m^2\tilde{Z}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} - \frac{eA_0(\tilde{G}\tilde{Y} - \tilde{A}^2)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)^2} \left[\nu_1 + \alpha\nu_1^3 \right], \\
U_{31} &= \frac{\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1 + \alpha\nu_1^3 \right] W_1, \quad U_{32} = \frac{\tilde{Z}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[J + \alpha J^3 \right] \tilde{E} + \frac{\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1 + \alpha\nu_1^3 \right] J, \\
U_{33} &= \frac{(\tilde{G}\tilde{Y} - \tilde{A}^2)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}^2 + \alpha\tilde{E}^4 - eA_0\tilde{E} - \alpha eA_0\tilde{E}^3 \right] - \frac{\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[W_1^2 + \alpha W_1^4 \right] - \frac{\tilde{G}}{\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[J^2 + \alpha J^4 \right] \\
&\quad - \frac{m^2\tilde{G}}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} - \frac{eA_0(\tilde{G}\tilde{Y} - \tilde{A}^2)}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} + \alpha\tilde{E}^3 - eA_0\tilde{E}^2 \right],
\end{aligned}$$

where $\tilde{E} = E - \sum_{i=1}^2 j_i \Omega_i$, $J = \partial_\phi I_0$, $W_1 = \partial_r I_0$ and $\nu_1 = \partial_\theta I_0$. We take the solution for non-trivial $|\mathbf{U}| = 0$ and get

$$\begin{aligned}
ImW^\pm &= \pm \int \sqrt{\frac{(\tilde{E} - j\Omega - eA_0)^2 + X_1 \left[1 + \alpha \frac{X_2}{X_1} \right]}{\frac{\tilde{G}(\tilde{G}\tilde{Y} + \tilde{Z}^2)}{\tilde{Y}}}} dr, \\
&= \pm i\pi \frac{(\tilde{E} - j\Omega - A_0 e) + \left[1 + \alpha \Xi \right]}{2\kappa(r_+)}, \tag{22}
\end{aligned}$$

where

$$\begin{aligned}
X_1 &= \frac{\tilde{Z}}{\tilde{G}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E} - eA_0 \right] \nu_1 + \frac{1}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1^2 \right] - \frac{1}{\tilde{G}} m^2, \\
X_2 &= \frac{\tilde{Y}}{\tilde{G}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}^4 - 2eA_0\tilde{E}^3 + (eA_0)^2\tilde{E}^2 \right] + \frac{\tilde{Z}}{\tilde{G}\tilde{X}(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\tilde{E}^3 - eA_0\tilde{E}^2 \right] \nu_1 - \frac{1}{(\tilde{G}\tilde{Y} + \tilde{Z}^2)} \left[\nu_1^4 \right] - W_1^4.
\end{aligned}$$

We observe the tunneling radiation without back-reaction and self-gravity interaction and also explore the corrected tunneling probability rate. The tunneling phenomenon is computed under the consideration of charge-energy conservation and quantum gravity effects. The modified tunneling probability depends on the geometry of BHs as well as quantum gravity. The modified tunneling probability for charged rotating BH with EBG gravity can be defined as

$$\Gamma = \frac{\Gamma_{emission}}{\Gamma_{absorption}} = \exp \left[-2\pi \frac{(\tilde{E} - j\Omega - A_0 e)}{\kappa(r_+)} \right] [1 + \alpha \Xi], \quad (23)$$

here

$$\kappa(r_+) = \frac{r_+^5 (Y - r_+^2) - 4Mr_+^4 \beta + 2a^2 r_+ (4q^2 \beta + r_+^4 (Y - r_+^2) - 10Mr_+ \beta)}{4\pi Y \beta (r_+^2 + a^2)}, \quad (24)$$

$$Y = \sqrt{r_+^2 - 8q^2 \beta + 16Mr_+ \beta}.$$

In the presence of gravity terms, we compute the T'_H of the 4-dimensional rotating charged BH in EGB gravity theory by considering the factor $\Gamma_B = \exp [(E - j\omega - eA_0)/T'_H]$ (Boltzmann factor) as

$$T'_H = \frac{r_+^5 (Y - r_+^2) - 4Mr_+^4 \beta + 2a^2 r_+ (4q^2 \beta + r_+^4 (Y - r_+^2) - 10Mr_+ \beta)}{8\pi Y \beta (r_+^2 + a^2)} [1 - \alpha \Xi]. \quad (25)$$

This solution shows that the T'_H depends on the BH geometry. Also, it is alike form that of the other scalar and fermion particles. The modified temperature depends on the correction parameter α , spin parameter a , BH charge q , arbitrary constant β , BH mass M and BH radius r_+ . Quantum corrections decelerate the increase in T'_H during the radiation phenomenon. These corrections make the radiation cease at some particular Hawking temperature, leaving the remnant mass. When this consideration holds, the temperature stops rising [59]

$$(M - dM)(1 + \alpha \varphi) \approx M, \quad (26)$$

where $\omega = dM$, $\alpha_0 (\frac{1}{M_{Planks}}) = \alpha$ and $M_{Planks} = \omega$. Here, α_0 and M_{Planks} are represent the dimensionless parameter and Planck mass and quantum gravity effects for $\alpha_0 < 10^5$ in [50, 60, 61].

V. GRAPHICAL T'_H ANALYSIS

This section describes the stability conditions of rotating charged BH with EBG gravity with the quantum gravity effects. We analyze the physical behavior and stability of the corresponding BH by plotting the graphs of corrected temperature T'_H via horizon r_+ for fixed value of arbitrary parameter $\Xi = 1$.




Figure 4: T'_H via r_+ for $\Xi = 1$ and $\beta = 0.5$. Left $M = 5 = a$, $q = 0.5$, $\alpha = 0.5$ (dashed), $\alpha = 0.6$ (dot-dashed), $\alpha = 0.7$ (small-dashed). Right $M = 1 = q$, $\alpha = 0.1$, $a = 0.1$ (dashed), $a = 0.2$ (dot-dashed), $a = 0.3$ (small-dashed).

Fig. 4 shows the corrected temperature T'_H behavior w.r.t horizon r_+ under the effects of quantum gravity.

(i) describes the temperature behavior for different the correction parameter α values and constant values of mass $M = 5$, arbitrary parameter and charge $\beta = 0.5 = q$ and spin parameter $a = 5$. One can observe the physical behavior (positive temperature) of BH under the effects of correction parameter. Moreover, the T'_H decreases as r_+ increases, which also satisfies the Hawking's Phenomenon.

(ii) depicts the T'_H behavior for constant values of mass $M = 1$, arbitrary parameter $\beta = 0.5$, charge $q = 1$, correction parameter $\alpha = 0.1$ and different values of rotation parameter a . At initial, the BH shows the non-physical behavior (negative temperature) but as time runs out and after a maximum height the temperature slowly decreases as r_+ increases as well as obtain an asymptotically flat condition to represents the complete stable form till $r_+ \rightarrow \infty$.

Fig. 5 represents the T'_H behavior in the presence of correction parameter α for fixed value of mass $M = 1$ and in the absence of charge and spin parameter ($q = 0 = a$), respectively.

(i) gives the temperature behavior in the absence of $q = 0$ (charge) for constant values of $\beta = 0.5, a = 10$ and different values of gravity parameter α . One can observe that the temperature gradually increases and then slowly decreases with the increasing r_+ that depicts the stable form of BH.

(ii) represents the T'_H behavior in the absence of $a = 0$ (spin parameter) for fixed $\beta = 0.1, q = 5$ and for varying α . We can examine that the T'_H behavior slowly decreases as horizon increases and gets an asymptotically flat condition that shows the stability of BH.

We can conclude from both plots that the BH is stable in the presence as well as absence of charge and spin parameter under the influence of quantum gravity parameter effects.




Figure 5: T'_H via r_+ for $\Xi = 1$ and $M = 1$. Left $\beta = 0.5, q = 0, a = 10, \alpha = 0.5$ (dashed), $\alpha = 0.6$ (dot-dashed), $\alpha = 0.7$ (small-dashed). Right $\beta = 0.1, q = 5, a = 0, \alpha = 5$ (dashed), $\alpha = 6$ (dot-dashed), $\alpha = 7$ (small-dashed).

VI. LOGARITHMIC CORRECTION OF ENTROPY FOR ROTATING CHARGED BH

In this section, the Bekenstein-Hawking entropy is considered to be corrected by particular correction terms in the quantum loop expansion. The logarithmic terms in BH entropy occur as a one-loop addition to the classical BH entropy. The BH entropy plays a significant role in the account of primordial BHs [66]. It imposes a minimum mass of the primordial BHs so the small ones have enough time to evaporate totally until the present form. The modified entropy with the logarithmic terms demonstrate, this conclusion is no more in place and the BHs evaporation decompresses as soon as mass decreases below the critical mass ($BH \text{ mass} \simeq \sqrt{s_0}(\text{planks mass})$) that extends to much longer BHs life time. In this form the small primordial BHs do not evaporate totally. Now, we analyze the corrections of entropy for EGB gravity of charged rotating BH. Banerjee and Majhi [62]-[64] have studied the corrections of Hawking temperature and entropy by taking into account the back-reaction effects via null geodesic method. We calculate the corrections of entropy for EGB gravity of charged rotating BH by considering the Bekenstein-Hawking entropy formula for first order corrections [65]. We derive the logarithmic corrections of entropy by using corrected temperature T'_H and standard entropy S_0 for EGB gravity of charged rotating BH. The corrections of entropy can be computed by the following formula

$$S = S_0 - \frac{1}{2} \ln \left| T_H^2, S_0 \right| + \dots \quad (27)$$

The standard entropy for EGB gravity of rotating charged BH can be calculated by the following formula

$$S_o = \frac{A_+}{4}, \quad (28)$$

here

$$\begin{aligned} A_+ &= \int_0^{2\pi} \int_0^\pi \sqrt{g_{\theta\theta}g_{\phi\phi}} d\theta d\phi, \\ &= \frac{\pi \left[2(r_+^2 + a^2)^2 - a^2q^2 + r_+Ma^2 \right]}{(r_+^2 + a^2)}. \end{aligned} \quad (29)$$

So the entropy term for standard can be derived as

$$S_o = \frac{\pi \left[2(r_+^2 + a^2)^2 - a^2q^2 + r_+Ma^2 \right]}{4(r_+^2 + a^2)}. \quad (30)$$

After substituting the values from Eq. (25) and (30) into Eq. (27), we compute the correction of entropy as

$$\begin{aligned} S &= \frac{\pi \left[2(r_+^2 + a^2)^2 - a^2q^2 + r_+Ma^2 \right]}{4(r_+^2 + a^2)} \\ &- \frac{1}{2} \ln \left| \frac{\left[(r_+^5(Y - r_+^2) - 4Mr_+^4\beta + 2a^2r_+(4q^2\beta + r_+^4(Y - r_+^2) - 10Mr_+\beta)) (1 - \alpha\Xi) \right]^2 \chi}{256\pi Y^2 \beta^2 (r_+^2 + a^2)^3} \right| + \dots, \end{aligned} \quad (31)$$

where

$$\chi = \left[2(r_+^2 + a^2)^2 - a^2q^2 + r_+Ma^2 \right].$$

The Eq. (31) represents the correction of entropy for EGB gravity of rotating charged BH.

VII. CONCLUSIONS

In this article, we have used the Newman-Janis algorithm to study the charged rotating BH with EBG gravity solution. We have derived the thermodynamics (Hawking temperature) at the outer horizon for the corresponding BH by using a general formula for symmetric space-time. The Hawking temperature depends upon rotation parameter a (occurs due to Newman-Janis algorithm), BH mass M , BH charge q , BH horizon r_+ and arbitrary constant β . Moreover, we analyzed the graphical behavior of temperature w.r.t horizon to analyze the stability of BH under the effects of spin parameter. We have also investigated the stability of BH in the presence and absence of charge and spin parameter $q = 0 = a$, respectively. We have also investigated the physical significance of our plots. We conclude the main results from the graphical analysis of Hawking temperature via horizon as follows:

- For different values of spin parameter a the T_H slowly goes on increasing with the increasing values of horizon but after a height it gradually goes on decreasing with the increasing horizon which satisfies the Hawking's phenomenon, so grantee's the physical and stable form of BH. The temperature T_H also increases with the increase in a .
- For varying values of charge q the BH at initial shows the non-physical form with negative temperature but as time goes on BH attains its stable form, when it eventually drops down from a height and obtains an asymptotically flat state till $r_+ \rightarrow \infty$. The decreasing temperature with increasing horizon also shows the stable condition of BH. The temperature decreases with the increasing values of charge.

- In the absence of spin parameter a , after a maximum height the T_H behavior slowly decreases as horizon increases and gets an asymptotically flat condition till $r_+ \rightarrow \infty$ that shows the stability of BH. From Fig. 2 and 3, it is clear that in the presence of rotation parameter the temperature increases as compare in the absence of a .
- In the absence of charge the temperature T_H gradually increases and then slowly decreases with the increasing r_+ that depicts the stable form of BH. But it can be observed from the plots that BH is in more stable form in the absence of charge q .

In all plots, the BH depicts its stable condition by satisfying the Hawking's idea (the size of BH reduces when more radiations emit), the temperature is high at small horizon. We observe maximum temperature at minimum value of horizon. Furthermore, we have calculated the Hawking temperature under the influence of quantum gravity by using the semi-classical method and also examined the graphical corrected temperature T'_H behavior w.r.t horizon r_+ to check the physical and stable conditions of BH under the effects of α (quantum gravity parameter). We also discussed the stability conditions of BH in the absence of charge and spin parameter with or without gravity effects. We have concluded that with or without quantum gravity the BH represents its stable form. In our analysis we have found that the quantum corrections decelerate the increase in temperature during the radiation process. This correction causes the radiation ceased at some specific temperature, leaving the remnant mass. The remnant will obtain at specific condition

$$M_{Res} \simeq \frac{M_p^2}{\beta_0 \omega} \gtrsim \frac{M_p}{\beta_0}. \quad (32)$$

The results from the graphical analysis of corrected Hawking temperatures with respect to the horizon for the given BH is summarized as follows:

- The T'_H decreases with the increasing horizon and BH reflects the stable state for varying values of rotation parameter a and correction parameter α . The corrected temperature T'_H also increases with the increase in a and α . The BH remnant can be obtained at nonzero horizon with maximum temperature in the domain $0 \leq r_+ \leq 15$.
- In the absence of charge the corrected temperature T'_H gradually increases and then slowly decreases with the increasing r_+ that depicts the stable form of BH.
- In the absence of spin parameter a the T'_H behavior slowly decreases as horizon increases and gets an asymptotically flat condition till $r_+ \rightarrow \infty$ that shows the stability of BH.

From our plots, we conclude that the tunneling emission rate increases (gives high temperature) with the increasing values of α . The corrected temperature of charged rotating BH with EBG gravity satisfies the both GUP and Hawking's conditions, that guarantee the physical and stable states of BH. According GUP condition the next order corrections must be small as compared to the standard term of Uncertainty relation. The positive value of temperature in these plots also satisfies the GUP relation, when GUP conditions does not satisfies temperature becomes negative (shows non-physical state of BH). Now by comparing the plots in the presence/absence of quantum gravity parameter, we can observe that the corrected temperature is lower that the original temperature. In conclusion, the Newman-Janis algorithm gravity has attracted more attention for different space-time. In this paper, we have evaluated the BH space-time in Newman-Janis algorithm and computed the Hawking temperature with the effects of the rotation parameter and correction parameter. Finally, we computed the logarithmic corrected entropy for EBG gravity of rotating charged BH and also checked the logarithmic corrected entropy with the effects of the rotation parameter and correction parameter.

In future, we will study on the other objects(i.e., wormholes and black rings) in the Newman-Janis algorithm.

-
- [1] D. G. Boulware and S. Deser, Phys. Rev. Lett. **55**(1985)2656.
[2] B. Zumino, Phys. Rev. Lett. **137**(1986)109.
[3] D. Lovelock, J. Math. Phys. **12**(1971)498.
[4] D. Glavan, Ch. Lin, Phys. Rev. Lett. **124**(2020)081301.
[5] G. G. L. Nashed, S. D. Odintsov and V. K. Oikonomou, Symmetry **14**(2022)545.
[6] C. H. Wu, Y. P. Hu, H. Xu, Eur. Phys. J. C **81**(2021)351.
[7] S. H. Hendi, S. Panahiyani and E. Mahmoudi, Eur. Phys. J. C **74**(2014)3079.

- [8] T. Clifton, et al, Phys. Rev. D **102**(2020)084005.
- [9] F. W. Shu, Phys. Lett. **B 811**(2020)135907 .
- [10] Ch.Y. Zhang, Ch. Li, M. Guo, Eur. Phys. J. **C 80**(2020)874.
- [11] R. A. Konoplya, A. F. Zinhailo, Phys. Lett. **B 810**(2020)135793.
- [12] R. A. Konoplya, A. F. Zinhailo, Eur. Phys. J. **C 80**(2020)1049.
- [13] S. G. Ghosh, S. D. Maharaj, Phys. Dark Univ. **30**(2020)100687.
- [14] A. G. Riess et al., Astron. J. **116**(1998)1009.
- [15] B. P. Abbott et al., Astrophys. J. **848**(2017)L12.
- [16] V. K. Oikonomou, Class. Quantum Grav. **38**(2021)195025.
- [17] S. D. Odintsov, V. K. Oikonomou, Phys. Lett. **B 805**(2020)135437.
- [18] S. D. Odintsov, V. K. Oikonomou, Phys. Lett. **B 797**(2019)134874.
- [19] V. K. Oikonomou, F. P. Fronimos, EPL **131**(2020)30001.
- [20] S.D. Odintsov, V.K. Oikonomou, F.P. Fronimos, Nucl. Phys. **B 958**(2020)115135.
- [21] S. W. Hawking, Commun. Math. Phys. **43**(1975)199.
- [22] M. K. Parikh and F. Wilczek, Phys. Rev. Lett. **85**(2000)5042.
- [23] M. K. Parikh, Int. J. Mod. Phys. **D 13**(2004)2351.
- [24] J. T. Firouzjaee and G. F. R. Ellis, Gen. Rel. Grav. **47**(2015)6.
- [25] W. Javed, G. Abbas, R. Ali, Eur. Phys. J. **C 77**(2017)296.
- [26] W. Javed, R. Babar, Adv. High Energy Phys. **2019**(2019)2759641.
- [27] W. Javed, R. Babar, Chinese Journal of Phys. **61**(2019)138.
- [28] W. Javed, R. Babar, Proceedings of the 15th Marcel Grossmann Meeting, <http://robot.icranet.org:8080/store/l380.pdf>.
- [29] W. Javed, R. Babar, Punjab University Journal of Mathematics **52**(2020)6.
- [30] W. Javed, R. Babar, A. Övgün, Mod. Phys. Lett. **A 34**(2019)1950057.
- [31] R. Babar, W. Javed, A. Övgün, Mod. Phys. Lett. **A 35**(2020)2050104.
- [32] M. Sharif, W. Javed, Eur. Phys. J. **C 72**(2012)1997.
- [33] M. Sharif, W. Javed, J. Korean Phys. Soc. **57**(2010)217.
- [34] A. Övgün, K. Jusufi, Eur. Phys. J. Plus. **132**(2017)298.
- [35] W. Javed, R. Ali, G. Abbas, Can. J. Phys. **97**(2018)176.
- [36] A. Övgün, W. Javed, R. Ali, Adv. High Energy Phys. **2018**(2018)11.
- [37] A. Övgün, Int. J. Theor. Phys. **55**(2016)2919.
- [38] A. Övgün, K. Jusufi, Eur. Phys. J. Plus **132**(2017)298.
- [39] K. Jusufi, A. Övgün, G. Apostolovska, Adv. High Energy Phys. **2017**(2017)8798657.
- [40] I. Sakalli, A. Övgün, K. Jusufi, Astrophys Space Sci. **361**(2016)330.
- [41] I. Sakalli, A. Övgün, General Relativity and Gravitation **48**(2016)1.
- [42] A. Övgün, I. Sakalli, Int. J. Theor. Phys. **57**(2018)322.
- [43] A. Övgün, I. Sakalli, J. Saavedra, C. Leiva, Mod. Phys. Lett. **A 35**(2020)2050163.
- [44] I. Sakalli, A. Övgün, J. Exp. Theor. Phys. **121**(2015)404.
- [45] R. Ali, K. Bamba, S. A. A. Shah, Symmetry. **631**(2019)11.
- [46] R. Ali, K. Bamba, M. Asgher, M. F. Malik, S. A. A. Shah, Symmetry. **12**(2020)1165.
- [47] R. Ali, K. Bamba, M. Asgher, S. A. A. Shah, Int. J. Mod. Phys. **D 30**(2021)2150002.
- [48] R. Ali, M. Asgher, M. F. Malik, Mod. Phys. Lett. **A 35**(2020)2050225.
- [49] W. Javed, R. Ali, R. Babar, A. Övgün, Eur. Phys. J. Plus **134**(2019)511.
- [50] W. Javed, R. Ali, R. Babar, A. Övgün, Chinese Phys. **C 44**(2020)015104.
- [51] K. Jusufi, A. Övgün, Astrophys Space Sci. **361**(2016)207.
- [52] R. Ali, R. Babar, M. Asgher, and S.A.A. Shah, Annals of Physics, **432**(2021)168572.
- [53] R. Ali, M. Asgher, New Astronomy **93**(2022)101759
- [54] R. Ali, R. Babar and P.K. Sahoo, Physics of the Dark Universe **35**(2022)100948.
- [55] R. Ali, et al. Int. J. Geom. Methods Mod. Phys. **19**(2022)2250017.
- [56] J. M. Bardeen, in Conference Proceedings of GR5 (Tbilisi, URSS, 1968), p. 174.
- [57] D. Chen, H. Wu, H. Yang, and S. Yang, JCAP, **03**(2014)036.
- [58] R. Ghosh, C. Fairros, S. Sarkar, Phys. Rev. D **100**(2019)124019.
- [59] D. Chen, H. Wu and H. Yang, Adv. High Energy Phys. **2013**(2013)432412.
- [60] D. Chen, H. Wu and H. Yang, JCAP, **1403**(2014)036.
- [61] S. Liberati, L. Maccione and T. P. Sotiriou, Phys. Rev. Lett. **109**(2012)151602.
- [62] R. Banerjee and B. R. Majhi, Phys. Lett. **B 662**(2018)62.
- [63] R. Banerjee and B. R. Majhi, J. Adv. High Energy Phys. **2008**(2008)095.
- [64] R. Banerjee, B. R. Majhi, and S. Samanta, Phys. Rev. D **77**(2008)124035.
- [65] P. Pradhan, Adv. High Energy Phys. **2017**(2017)2367387.
- [66] S. N. Solodukhin, Phys. Lett. **B 802**(2020)135225.