

Category of Quantizations and Inverse Problem

Akifumi Sako

Tokyo University of Science,
1-3 Kagurazaka, Shinjuku-ku, Tokyo, 162-8601, Japan

Abstract

We introduce a category composed of all quantizations of all Poisson algebras. By the category, we can treat in a unified way the various quantizations for all Poisson algebras and develop a new classical limit formulation. This formulation proposes a new method for the inverse problem, that is, the problem of finding the classical limit from a quantized space. Equivalence of quantizations is defined by using this category, and the conditions under which the two quantizations are equivalent are investigated. Two types of classical limits are defined as the limits in the context of category theory, and they are determined by giving a sequence of objects. Using these classical limits, we discuss the inverse problem of determining the classical limit from some noncommutative Lie algebra. From a Lie algebra, we construct a sequence of quantized spaces, from which we determine a Poisson algebra. We also present a method to obtain this sequence of quantizations from the principle of least action by using matrix regularization. Apart from the above category of quantizations of all Poisson algebras, a category of quantizations of a fixed single Poisson algebra is also introduced. In this category, the other classical limit is defined, and it is automatically determined for the category.

1 Introduction

In M-theory and string theory, it has been proposed for a long time that matrix models give their constructive formulation [5, 22]. Noncommutative manifolds are obtained as classical solutions of such matrix models. (See for example [17, 42] and references therein.) Assuming that the universe as a noncommutative manifold is realized, the space we inhabit is perceived as a smooth manifold, at least to an approximation. Therefore, it is important to obtain smooth manifolds as classical limits of noncommutative manifolds.

In many cases, noncommutative geometry is some kind of a noncommutative deformation of a smooth manifold. In other words, it is often made as a quantization of a smooth manifold. So, it is often called an inverse problem to construct a usual commutative manifold or commutative algebra as a classical limit of a noncommutative manifold.

Taking the classical limit of a noncommutative manifold is often fraught with difficulty. The images of the quantization maps have information of the classical geometry of manifolds, but the algebras generated from them lose information of the original geometry. For example, as a well-known phenomenon, matrix regularization generates the same matrix algebra whether a two-dimensional torus is transformed into a

fuzzy torus or a two-dimensional sphere into a fuzzy sphere. (See Appendix C and [18, 3, 21, 31, 32].) Put another way, when reading a classical geometry (Poisson algebra) from a matrix algebra, the Poisson algebra obtained depends on what classical limit is taken [12, 15, 16]. Various approaches have been taken to the inverse problem of how to extract geometric properties from matrix algebras [41, 11, 38, 23, 24, 4]. Therefore, a framework including any Poisson algebra and its quantizations is important in considering such issues to be investigated in a unified manner.

Another important lesson implied by the difficulty described above is that the commutative limit cannot be identified by information in the algebra obtained as the target of a quantization map. Rather, information about the modules or vector spaces that are images of quantization maps before their generating algebras is important to identify the classical limits. This tendency can be read from, for example, fuzzy spheres and fuzzy torus in Appendix C. Note that an image of a quantization map is not an algebra but a module in general since a quantization map is not always an algebra homomorphism but a module homomorphism (linear map). Furthermore, the classical limit depends on what classical limit is taken as already mentioned. Accordingly, it is suggested that we should prepare a sequence of modules to determine a Poisson algebra as a classical limit, not just one of them.

In light of the above, the first purpose of this study is to construct a category composed of the all quantizations of all Poisson algebras. This is actually performed in Section 3. We denote this category by QW (Quantum World), and its strict definition is given in Section 3. Quantization here refers to a noncommutative deformation in the sense of approximate canonical quantization of a Poisson algebra. The definition and properties of this quantization are given in Section 2. Quantizations in this paper are not necessarily identified with them in the sense of quantum mechanics, nor in the sense of spectral discretization, but in the sense of many noncommutative geometries. The morphisms that define this QW consist of Poisson morphisms, quantization maps, and algebra homomorphisms whose domain is restricted. Their compositions are constructed in such a way that they satisfy the axioms of the category without contradiction. Furthermore, in this context, it is also important to show that quantization can be properly classified. To this end, we construct a functor from QW to some comma category and show that the isomorphism of the object of the comma category gives the isomorphism pair of the Poisson algebra and the algebra generated by its quantized space. QW allows any quantization of any Poisson algebra to be treated as a single framework.

The second purpose is to define the classical limit within the framework of this category of the all quantizations QW in a non-traditional way. Of course, the question of what classical limit is best within the QW framework depends on what classical limit is needed. In this paper, we propose three types of classical limits and investigate their properties by considering concrete examples in Section 5. We define the classical limit using the limit in the category theory or a slight modified limit. The first two types are defined by using sequence of objects (modules) in QW and morphisms between them. One is a naive adaptation of the category theory limit to QW . When that category theory limit corresponds to a Poisson algebra, it is defined to be the classical limit. To define another classical limit, it is necessary to modify the definition of the limit of

category theory. We impose universality within the framework that the vertex of the cone is a Poisson algebra. The classical limit so defined is called the weak classical limit. These limits depends on how to choose the sequence of objects, and the existence of the limits are not guaranteed. As concrete examples, we investigate the classical limits for a sequence of deformation quantizations of a polynomial ring and for a sequence of fuzzy spheres. By considering the all quantizations as a single category, we can make a structure that a sequence of modules as a sequence of objects in the category determines a Poisson algebra as the classical limit from all Poisson algebras. To make the sequence of objects, we use matrix regularization as an example. Furthermore, developing this argument we study how to obtain the classical limit by using principle of least action in Section 6.

The third type of the classical limit is made by focusing a single Poisson algebra. When we choose a Poisson algebra, this classical limit is determined automatically. It is constructed in the framework of a category made from QW with a fixed Poisson algebra whose objects are quantization maps from the Poisson algebra. We prove that the classical limit coincides with the fixed Poisson algebra. We call this classical limit the strong classical limit.

We would like to make a few comments on recent previous studies that discuss quantization as noncommutative deformation of a Poisson algebra by using category theory approach. The construction of a quantization category was done by [19, 20]. The quantization category in [19, 20] was pioneering in the sense that it was a category-theoretic version of quantization, but it contained two problems. The first problem is that the category in [19, 20] is constructed as a whole of quantization with respect to a single fixed Poisson algebra. As already mentioned, this is a weakness for the inverse problem because it is important to treat the all quantizations for the all Poisson algebras, in a unified manner. The second problem is that the quantization maps treated in [19, 20] are maps from some Poisson algebra to Lie algebras. Target spaces of the maps do not necessarily reflect the reality of the image of the mapping. A Lie algebra that is a target space of some quantization map may have lost some of their original information of the image of quantization map, as already mentioned. So, to classify quantization maps, we need more complex structure in the formulation. On the other hand, the category constructed in this article conquers these two problems.

This paper is organized as follows. In Section 2, we study quantization maps. Using the properties of quantization maps, we construct a category of the all quantizations of all Poisson algebras QW in Section 3. In Section 4, a way to classify the quantization maps is established by using the context of QW . In Section 5, we discuss how to find the classical limit in QW . Three types of classical limits are discussed. Examples of these classical limits are studied. In Section 6, the inverse problem is discussed in the context of the classical limit defined in Section 5. Especially, a way to obtain the classical limit by the principle of least action with a sequence of actions is proposed. We make a summary of this article in Section 7. A symbol like Landau notation, used throughout this paper, is defined in Appendix A. Many new definitions appear in this paper, and a list of the symbols is available in Appendix B. We often use a fuzzy sphere or a fuzzy torus to make examples. So, we make a short introduction to them in Appendix C.

2 Quantization Maps

We define the quantization maps in this section. To formulate them, we use R -modules and R -algebras. R denote a commutative ring over \mathbb{C} , in this paper. Only finitely generated R -modules and R -algebras are considered. Existence of multiplicative identity does not claimed, and the condition for the identity in the axiom of the R -algebra homomorphism is removed.

2.1 Definition of Weak Quantization Maps

We define a category of Poisson algebras whose morphisms are restricted into surjective Poisson morphisms as follows.

Definition 2.1. Let $(A, \cdot_A, \{ , \}_A)$ and $(B, \cdot_B, \{ , \}_B)$ be Poisson algebras over \mathbb{C} . When a linear $\phi_{A,B} : A \rightarrow B$ satisfies

$$\phi_{A,B}(f \cdot_A g) = \phi_{A,B}(f) \cdot_B \phi_{A,B}(g) \quad (2.1)$$

$$\phi_{A,B}(\{f, g\}_A) = \{\phi_{A,B}(f), \phi_{A,B}(g)\}_B, \quad (2.2)$$

for any $f, g \in A$, $\phi_{A,B}$ is called a Poisson morphism. \mathcal{Poisss} is a category whose objects are Poisson algebras over \mathbb{C} and its morphisms are “surjective” Poisson morphisms.

The reason why the morphisms were restricted to surjective maps will become clear later. Often the subscripts of Poisson brackets, the symbol for the product \cdot_A , and so on are omitted in the following.

Definition 2.2 (Weak Quantization map wkQ). Let A be a Poisson algebra over \mathbb{C} , and let T_i be an R -module that is given by a subset of some R -algebra $(M, *_M)$. If an R -module homomorphism (linear map) $t_{A_i} \in Hom_{R\text{-Mod}}(A, T_i)$ equips a constant $\hbar(t_{A_i}) \in \mathbb{C}$ and satisfies

$$[t_{A_i}(f), t_{A_i}(g)]_M = \sqrt{-1}\hbar(t_{A_i})t_{A_i}(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_{A_i})) \quad (\epsilon > 0) \quad (2.3)$$

for arbitrary $f, g \in A$, where $[a, b]_M := a *_M b - b *_M a$, we call t_i a weak quantization map. For the case that $[t_{A_i}(f), t_{A_i}(g)]_M = 0$ and $t_{A_i}(\{f, g\}) = 0$ for $\forall f, g \in A$, we put $\hbar(t_{A_i}) = 0$. \tilde{O} is defined in the Appendix A. We denote the set of all weak quantization maps by wkQ :

$$wkQ := \bigsqcup_{\substack{A \in \mathcal{Poisss} \\ T_i \subset M \in R\text{-alg}}} \{t_{A_i} \in Hom_{R\text{-Mod}}(A, T_i) \mid t_{A_i} \text{ satisfies (2.3)}\}$$

Note that the target of t_{A_i} is not an algebra in general, but it is a subset of an algebra M , so the left-hand side of (2.3) is defined through the product in M . We call $\hbar(t_{A_i}) \in \mathbb{C}$ a noncommutativity parameter. As we will see in Proposition 2.14, the \hbar can take any value of \mathbb{C}^\times for fixed A and T_i if $\hbar(t_{A_i}) \neq 0$. In the following, $*_M$ is omitted as appropriate. For simplicity, $[a, b]_M$ is abbreviated to $[a, b]$ when there is no risk of confusion.

Definition 2.2 includes many kinds of quantizations. For example, matrix regularization [21, 3], fuzzy spaces [31], and Berezin-Toeplitz quantization [12, 15, 39] which

have original ideas of the matrix regularization, satisfy Definition 2.2. In fact, the fuzzy sphere appears frequently as an example in this paper. In addition, the strict deformation quantization introduced by Rieffel [36, 37, 26], the prequantization [8, 28, 45], and Poisson enveloping algebras [33, 34, 44, 46] are also in wkQ . These facts are derived immediately from these definitions. (In [19, 20], we can see organized discussions about these quantization maps. The conditions for wkQ is a part of the definition of pre- \mathcal{Q} in [19, 20].)

2.2 Properties for Weak Quantization Maps

For the later part, we summarize basic properties of R -algebra homomorphisms used in this paper, and we derive some propositions of the weak quantization maps. In order to make this article a self-contained article, proofs of basic matters are not omitted.

Proposition 2.3. *Let $(M, *M)$ and $(N, *N)$ be R -algebras, and let T be a generating set of M , i.e. $\langle T \rangle = M$. For any R -algebra homomorphisms h_{MN}, h'_{MN} from M to N ,*

$$h_{MN}|_T = h'_{MN}|_T \Leftrightarrow h_{MN} = h'_{MN}, \quad (2.4)$$

where $h_{NM}|_T$ and $h'_{NM}|_T$ are h_{MN} and h'_{MN} whose domains are restricted to T , respectively.

Proof. For a multi index $I_k = (i_1, \dots, i_k)$, we denote $a_{i_1} *M \dots *M a_{i_k}$ by a_{I_k} , where $a_{i_j} \in T$. Since T generates M , any $b \in M$ can be written as a polynomial

$$b = \sum_k \sum_{I_k} b_{I_k} a_{I_k} \quad (b_{I_k} \in R).$$

If $h_{MN}|_T = h'_{MN}|_T$, $h_{MN}(a_{i_j}) = h'_{MN}(a_{i_j})$ for each a_{i_j} , and

$$\begin{aligned} h_{NM}(b) &= \sum_k \sum_{I_k} b_{I_k} h_{NM}(a_{I_k}) \\ &= \sum_k \sum_{I_k} b_{I_k} h'_{NM}(a_{I_k}) = h'_{NM}(b). \end{aligned}$$

The proof for the opposite direction is trivial. □

Proposition 2.4. *Let M and $\langle T \rangle$ be R -algebras, where T is a subset of $\langle T \rangle$ that generates algebra $\langle T \rangle$. For any R -algebra homomorphism $h : \langle T \rangle \rightarrow M$,*

$$\langle h(T) \rangle = h(\langle T \rangle). \quad (2.5)$$

Proof. For any $m \in \langle h(T) \rangle$ can be written as a polynomial

$$m = \sum_k \sum_{I_k} m_{I_k} e_{I_k} \quad (m_{I_k} \in R),$$

where $e_{I_k} = e_{i_1} * \dots * e_{i_k}$ with $e_{i_j} \in h(T)$. There exists $e'_{i_j} \in T$ such that $e_{i_j} = h(e'_{i_j})$ for each e_{i_j} . Then,

$$\begin{aligned} m &= \sum_k \sum_{I_k} m_{I_k} h(e'_{i_1}) * \dots * h(e'_{i_k}) \\ &= h\left(\sum_k \sum_{I_k} m_{I_k} e'_{i_1} * \dots * e'_{i_k}\right). \end{aligned}$$

□

Proposition 2.5. *Let A be a Poisson algebra. Let $t_i : A \rightarrow M_i$ be a weak quantization map whose image $t_i(A)$ generates an R -algebra M_i , i.e. $\langle t_i(A) \rangle = M_i$. For an arbitrary R -algebra M_j and any R -algebra homomorphism $h_{ij} : M_i \rightarrow M_j$, an R -module homomorphism $t_j : A \rightarrow t_j(A) \subset M_j$ defined by*

$$t_j = h_{ij} \circ t_i, \quad (2.6)$$

satisfies

$$[t_j(f), t_j(g)]_{M_j} = \sqrt{-1}\hbar(t_i)t_j(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_i)) \quad (\forall f, g \in A). \quad (2.7)$$

Proof. For any $f, g \in A$, from (2.6) and the fact that h_{ij} is an R -algebra homomorphism,

$$\begin{aligned} & [t_j(f), t_j(g)]_{M_j} \\ &= (h_{ij}|_{t_i(A)} \circ t_i)(f) *_{M_j} (h_{ij}|_{t_i(A)} \circ t_i)(g) - (h_{ij}|_{t_i(A)} \circ t_i)(g) *_{M_j} (h_{ij}|_{t_i(A)} \circ t_i)(f) \\ &= h_{ij}(t_i(f) *_{M_i} t_i(g) - t_i(f) *_{M_i} t_i(g)) \\ &= h_{ji}([t_i(f), t_i(g)]_{M_i}) \end{aligned}$$

Since t_i is in wkQ , the above equation is written as

$$\begin{aligned} [t_j(f), t_j(g)]_{M_j} &= h_{ji}(\sqrt{-1}\hbar(t_i)t_i(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_i))) \\ &= \sqrt{-1}\hbar(t_i)t_j(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_i)). \end{aligned}$$

□

Note that (2.7) does not mean that t_j is a weak quantization map with $\hbar(t_j) = \hbar(t_i)$. For example, $\hbar(t_i) \neq 0$ case, if $t_j(A)$ generates commutative algebra, (2.7) means $\{f, g\}$ is in $\text{Ker } t_j$. In this case $\hbar(t_j) = 0$ by the definition of the weak quantization.

Corollary 2.6. *Let A be a nontrivial Poisson algebra, i.e. $\exists f, g \in A$ such that $\{f, g\} \neq 0$. Let $t_i, t_j, h_{ij}, M_i, M_j$ be them in Proposition 2.5. If $\langle t_i(A) \rangle$ is noncommutative and if $\langle t_j(A) \rangle = \langle h_{ij} \circ t_i(A) \rangle \subset M_j$ is commutative, then*

$$\{A, A\} := \{ \{f, g\} \in A \mid f, g \in A \} \subset \text{Ker } t_j, \quad (2.8)$$

where $\{A, A\}$ is non-empty. $t_i(\{A, A\})$ is not empty, too, and

$$t_i(\{A, A\}) \subset \text{Ker } h_{ij}. \quad (2.9)$$

Proof. When $\langle t_j(A) \rangle$ is a commutative algebra, for any $f, g \in A$,

$$0 = [t_j(f), t_j(g)] = i\hbar(t_i)t_j(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}). \quad (2.10)$$

Since $\langle t_i(A) \rangle$ is noncommutative, existence of $f, g \in A$ such that $[t_i(f), t_i(g)] \neq 0$ derives $\hbar(t_i) \neq 0$. Then we get $t_j(\{f, g\}) = h_{ij} \circ t_i(\{f, g\}) = 0$ for $\forall f, g \in A$. $\hbar(t_i) \neq 0$ also means that there exist $f, g \in A$ such that $t_i(\{f, g\}) \neq 0$, and $t_i(\{A, A\}) \subset \text{Ker } h_{ij}$ is also obtained. □

Proposition 2.7. *Suppose that a noncommutativity parameter of a weak quantization map $t_i : A \rightarrow t_i(A) \subset M_i$ is non-zero, i.e. $\hbar(t_i) \neq 0$, where M_i is some noncommutative R -algebra. For an R -algebra homomorphism $h_{ij} : M_i \rightarrow M_j$ and $t_j := h_{ij} \circ t_i$, $\langle t_j(A) \rangle$ is a commutative algebra if and only if*

$$\{[t_i(f), t_i(g)] \mid f, g \in A\} \subset \text{Ker } h_{ij}.$$

Proof. For any $f, g \in A$ with $[t_i(f), t_i(g)] \neq 0$,

$$h_{ij}([t_i(f), t_i(g)]) = [h_{ij} \circ t_i(f), h_{ij} \circ t_i(g)] = [t_j(f), t_j(g)].$$

Therefore, if $\langle t_j(A) \rangle$ is commutative, $\forall [t_i(f), t_i(g)] \in \text{Ker } h_{ij}$.

Conversly, suppose that $[t_i(f), t_i(g)]$ is in $\text{Ker } h_{ij}$ for $\forall f, g \in A$. $\forall m_l \in \langle t_j(A) \rangle$ is expressed as $m_l = \sum_k \sum_{I_k} m_{l_{I_k}} a_{I_k}$, where $a_{I_k} = a_{i_1} \cdots a_{i_k}$, $a_{i_p} \in \text{Im } h_{ij}$ and $m_{l_{I_k}} \in R$.

There exists $e_{i_p} \in t_i(A)$ such that $a_{i_p} = h_{ij}(e_{i_p})$. Any commutator $[m_l, m_p]$ is written as

$$[m_l, m_p] = \sum_{k,q} \sum_{I_k, J_q} m_{l_{I_k}} m_{p_{J_q}} [a_{I_k}, a_{J_q}],$$

and $[a_{I_k}, a_{J_q}] = \sum_{1 \leq s \leq k} \sum_{1 \leq n \leq q} (\cdots [a_{i_s}, a_{j_n}] \cdots)$. So,

$$\begin{aligned} [m_l, m_p] &= \sum_{k,q} \sum_{I_k, J_q} m_{l_{I_k}} m_{p_{J_q}} \sum_{1 \leq s \leq k} \sum_{1 \leq n \leq q} (\cdots [a_{i_s}, a_{j_n}] \cdots). \\ &= \sum_{k,q} \sum_{I_k, J_q} m_{l_{I_k}} m_{p_{J_q}} \sum_{1 \leq s \leq k} \sum_{1 \leq n \leq q} (\cdots h_{ij}([e_{i_s}, e_{j_n}]) \cdots). \end{aligned}$$

Since $[e_{i_s}, e_{j_n}] \in \text{Ker } h_{ij}$, we find $[m_l, m_p] = 0$. □

To avoid commutative algebras from quantum theories, we will restrict R -algebra homomorphisms to injective ones later.

We found that $t_j = h_{ij} \circ t_i$ in Proposition 2.5 is a weak quantization map. $\langle t_j(A) \rangle = \langle h_{ij} \circ t_i \rangle \subset M_j$, $\text{Im } h_{ij}$, and M_j are not always equal. So, “when $\langle h_{ij} \circ t_i(A) \rangle$ generates M_j ?” is a natural question. The following discussion derives a useful tool to judge this problem.

Proposition 2.8. *Let $A, B, C \in \text{ob}(R\text{-Mod})$ be R -modules satisfying $\langle A \rangle = M_i$, $\langle B \rangle = \langle C \rangle = M_j$, where M_i and M_j are R -algebras. If an R -algebra homomorphism $h_{ji} : M_j \rightarrow M_i$ satisfies $h_{ji}|_B(B) = A$, where $h_{ji}|_B$ is defined by restriction of its domain to B , then the image of C by h_{ji} generates M_i , i.e. $\langle h_{ji}|_C(C) \rangle = M_i$.*

Proof. From $\langle B \rangle = \langle C \rangle = M_j$, an arbitrary $b \in B$ is expressed by elements of C . By $h_{ji}|_B(B) = A$ and $\langle A \rangle = M_i$, $h_{ji}|_B(B)$ generates M_i . Then $\langle h_{ji}|_C(C) \rangle = M_i$ is given. □

We introduce a sequence of weak quantization maps $t_{\mu j}^{(l)} : A^\mu \rightarrow t_{\mu j}^{(l)}(A^\mu) \subset M_j$, ($l \in I_j, \mu = 1, 2, \dots$) that generate an R -algebra M_j

$$\begin{aligned} M_j &= \langle t_{\mu j}^{(l)}(A^\mu) \rangle = \dots = \langle t_{\mu j}^{(n)}(A^\mu) \rangle = \dots \\ &= \langle t_{\nu j}^{(l)}(A^\nu) \rangle = \dots = \langle t_{\nu j}^{(n)}(A^\nu) \rangle = \dots \\ &\vdots \end{aligned} \tag{2.11}$$


where $A^\mu, A^\nu, \dots \in \text{ob}(\mathcal{P}oiss)$ and I_j is an index set.

Proposition 2.9. *Let A^μ, A^τ be Poisson algebras. Let $t_{\mu j}^{(k)} : A^\mu \rightarrow t_{\mu j}^{(k)}(A^\mu)$ and $t_{\mu i}^{(l)} : A^\mu \rightarrow t_{\mu i}^{(l)}(A^\mu)$ be weak quantization maps such that $\langle t_{\mu j}^{(k)}(A^\mu) \rangle = M_j$, $\langle t_{\mu i}^{(l)}(A^\mu) \rangle = M_i$. If an R -algebra homomorphism $m_{ji} : M_j \rightarrow M_i$ satisfies commutativity $m_{ji} \circ t_{\mu j}^{(k)} = t_{\mu i}^{(l)}$, then $\forall t_{\tau j}^{(n)} \in \text{wkQ}$ such that $\langle t_{\tau j}^{(n)}(A^\tau) \rangle = M_j$, $t_i := m_{ji} \circ t_{\tau j}^{(n)} : A^\tau \rightarrow t_i(A^\tau) \subset M_i$ satisfies*

$$[t_i(f), t_i(g)]_{M_i} = \sqrt{-1}\hbar(t_{\tau j}^{(n)})t_i(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_{\tau j}^{(n)})) \quad (f, g \in A^\tau). \tag{2.12}$$

and

$$\langle t_i(A^\tau) \rangle = M_i. \tag{2.13}$$



Proof. This proposition follows from Proposition 2.5 and Proposition 2.8. \square


Remark. Note that the existence of Poisson maps between A^μ and A^τ are not requested in Proposition 2.9.

Example 2.10. *As an example, let us consider fuzzy spheres and fuzzy tori. These are obtained by matrix regularization for S^2 and T^2 , and their processes are summarized in Appendix C. \mathcal{A} and \mathcal{B} denote the Poisson algebras defined as a set of functions on S^2 and T^2 , respectively. $t_k : \mathcal{A} \rightarrow T_k := t_k(\mathcal{A}) \subset M_k = \text{Mat}_k(\mathbb{C})$ and $q_k : \mathcal{B} \rightarrow Y_k := q_k(\mathcal{B}) \subset M_k = \text{Mat}_k(\mathbb{C})$ are defined as weak quantization maps. The details are given in Appendix C. It is known that both $t_k(\mathcal{A})$ and $q_k(\mathcal{B})$ generate the same matrix algebra Mat_k .*

Let introduce a map $t_{k \oplus k} : \mathcal{A} \rightarrow T_k \oplus T_k \subset M_k \oplus M_k$ as

$$t_{k \oplus k}(f) = \left(\begin{array}{c|c} t_k(f) & 0 \\ \hline 0 & t_k(f) \end{array} \right) \quad (f \in \mathcal{A}). \quad (2.14)$$

This satisfies $[t_{k \oplus k}(f), t_{k \oplus k}(g)]_M = \sqrt{-1} \hbar(t_k) t_{k \oplus k}(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(t_k))$. In other words, $t_{k \oplus k}$ is a weak quantization map. We define $q_{k \oplus k} : \mathcal{B} \rightarrow Y_k \oplus Y_k$ as a weak quantization map, in the same way. Let $\pi : M_k \oplus M_k \rightarrow M_k$ be a projection map to one M_k . From Proposition 2.9, by using a projection π and commutativity $\pi \circ t_{k \oplus k} = t_k$, we obtain $q_k = \pi \circ q_{k \oplus k}$ as a weak quantization map and $\langle q_k(\mathcal{B}) \rangle = M_k$. This result reproduces known facts.



To prevent later confusion, a note is made here. From the fact that this projection π is an R -algebra homomorphism but not injective one, this π is removed from the set of R -algebra homomorphisms for QW we will consider in the following parts of this paper. As we will see in Section 3, only injective R -algebra homomorphisms are in $\text{Mor}(QW)$.

2.3 Quantization Maps

The parameter of noncommutativity \hbar is determined by the choice of weak quantization map i.e. $\hbar : wkQ \rightarrow \mathbb{C}$. The set of the weak quantization maps wkQ is the disjoint union as

$$wkQ = \bigsqcup_{x \in \mathbb{R}_{\geq 0}} Qh_x,$$

where

$$Qh_x := \{q \in wkQ \mid |\hbar(q)| = x \in \mathbb{R}_{\geq 0}\}.$$

We denote the set of all targets of Qh_x by

$$Qh_x(\mathcal{Poi}ss) := \{t(q) \mid q \in Qh_x\}, \quad (2.15)$$

where $t(q)$ is a target of q . Particularly, $Qh_0 := \{q \in wkQ \mid |\hbar(q)| = 0\}$ is a set of maps to commutative algebras including the identity Id_A for $A \in U_p(\mathcal{Poi}ss)$, where U_p is a forgetful functor from $\mathcal{Poi}ss$ to QW defined in Section 3.

Definition 2.11. The set of quantization maps Q is defined by

$$Q := wkQ \setminus Qh_0 = \bigsqcup_{x \neq 0} Qh_x, \quad (2.16)$$

and the set of all targets of the quantization maps is defined by

$$Q(\mathcal{P}oiss) := \{t(q) \mid q \in Q\}. \quad (2.17)$$

We call $q \in Q$ a quantization map.

Here, let us explain why we define the Poisson morphism in $\mathcal{P}oiss$ is surjective. At first, we see the following proposition.

Proposition 2.12. *Let A^μ and A^ν be Poisson algebras and let T_i be an R -module as a subset of an R -algebra M_i . For a weak quantization map $t_{\nu i} : A^\nu \rightarrow T_i$ and a Poisson morphism $\phi : A^\mu \rightarrow A^\nu$, $t_{\nu i} \circ \phi : A^\mu \rightarrow T_i$ is also a weak quantization map.*

Proof.

$$\begin{aligned} \sqrt{-1}\hbar(t_{\nu i})t_{\nu i} \circ \phi(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}) &= \sqrt{-1}\hbar(t_{\nu i})t_{\nu i}(\{\phi(f), \phi(g)\}) + \tilde{O}(\hbar^{1+\epsilon}) \\ &= [t_{\nu i}(\phi(f)), t_{\nu i}(\phi(g))] + \tilde{O}(\hbar^{1+\epsilon}). \end{aligned}$$

□

Note that Proposition 2.12 does not show that $t_{\nu i} \circ \phi$ is always in Q even if $t_{\nu i} \in Q$. For example, suppose that $C \subset T_i \subset M_i$ is a commutative subalgebra of M_i , and $\text{Im}\phi \subset t_{\nu i}^{-1}(C)$. Then the image of $t_{\nu i} \circ \phi$ is in C and $\hbar(t_{\nu i} \circ \phi) = 0$, i.e. $t_{\nu i} \circ \phi \notin Q$. On the other hand, if ϕ is surjective $t_{\nu i} \circ \phi(A^\mu) = t_{\nu i}(A^\nu)$. This fact and Proposition 2.12 derives the following.

Proposition 2.13. *Let A^μ and A^ν be Poisson algebras. Let $t_{\nu i} \in Q$ be a quantization map whose source is A^ν . If a Poisson morphism $\phi : A^\mu \rightarrow A^\nu$ is surjective, then $t_{\nu i} \circ \phi \in Q$.*

This property is important to make the category QW , so an ancillary condition that the morphisms are surjective is added to the conditions for $Mor(\mathcal{P}oiss)$.

Next, we make sure that we can treat \hbar as a continuous parameter in Q .

Proposition 2.14. *Let A be a Poisson algebra. Let T_i be an R -module. For $\forall q_i \in Q : A \rightarrow T_i$ and $\forall x \in \mathbb{C}^\times$,*

$$q_i^x := \frac{x}{\hbar(q_i)} q_i \quad (2.18)$$

is a quantization map from A to T_i , such that $\hbar(q_i^x) = x$, $q_i(A) = q_i^x(A)$.

Proof. From $[q_i(f), q_i(g)] = \sqrt{-1}\hbar(q_i)q_i(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon})$, we obtain

$$[q_i^x(f), q_i^x(g)] = ixq_i^x(\{f, g\}) + \frac{x^2}{\hbar^2(q_i)} \tilde{O}(\hbar^{1+\epsilon}).$$

Because $\frac{x^2}{\hbar^2(q_i)} \tilde{O}(\hbar^{1+\epsilon}) = \tilde{O}(x^{1+\epsilon})$, the proof is finished. □

This proposition teaches us that there are uncountable infinite number quantizations for each target T_i .

3 Category of Quantizations

In this section, we construct a category QW that describes whole space of the all quantizations of all Poisson algebras.

We denote the target and the source of a morphism $q \in Q$ by $t(q)$ and $s(q)$, respectively. Recall that $t(q)$ is a subspace of some R -algebra for each q .

At the first step, we introduce a category of target spaces $\{t(q) \mid q \in Q\}$.

Definition 3.1. QP is a subcategory of $R\text{-Mod}$ defined as follows.

- $ob(QP) := Q(\mathcal{P}oiss) = \{t(q) \mid q \in Q\}$.
- Suppose that two objects $t(q_i), t(q_j)$ are subsets of R -algebras M_i, M_j , respectively. The morphism $Mor(t(q_i), t(q_j))$ is the set of all $h_{ij}|_{t(q_i)} : t(q_i) \rightarrow t(q_j)$. Here, $h_{ij}|_{t(q_i)}$ is given as an injective R -algebra homomorphism $h_{ij} : M_i \rightarrow M_j$ whose source is restricted into $t(q_i)$, and $\text{Im}(h_{ij}|_{t(q_i)}) \subset t(q_j)$:

$$Mor(t(q_i), t(q_j)) := \{h_{ij}|_{t(q_i)} : t(q_i) \rightarrow t(q_j) \mid h_{ij} : M_i \rightarrow M_j \text{ is an injective } R\text{-algebra homomorphism}\}.$$

The composition of $m_{ij} = h_{ij}|_{t(q_i)} \in Mor(t(q_i), t(q_j))$ and $m_{jk} = h_{jk}|_{t(q_j)} \in Mor(t(q_j), t(q_k))$ is well-defined by

$$m_{jk} \circ m_{ij} = h_{jk}|_{t(q_j)} \circ h_{ij}|_{t(q_i)} = (h_{jk} \circ h_{ij})|_{t(q_i)}.$$

Next, we consider a category $\mathcal{P}oiss \sqcup QP$ defined by

- $ob(\mathcal{P}oiss \sqcup QP) = ob(\mathcal{P}oiss) \sqcup ob(QP)$
- $Mor(\mathcal{P}oiss \sqcup QP) = Mor(\mathcal{P}oiss) \sqcup Mor(QP)$.

Since $ob(\mathcal{P}oiss) \cap ob(QP) = \emptyset$ and $Mor(\mathcal{P}oiss) \cap Mor(QP) = \emptyset$, it is trivial that $\mathcal{P}oiss \sqcup QP$ is a category.

The final step to introduce a category QW that describe whole space of the all quantizations of all Poisson algebras is to define its morphism by

$$Mor(QW) := Mor(\mathcal{P}oiss) \sqcup Mor(QP) \sqcup Q. \quad (3.1)$$

Proposition 3.2. *If $\forall f, g \in Mor(QW)$ satisfy $s(f) = t(g)$, then $f \circ g \in Mor(QW)$.*

Proof. For the case $f, g \in Mor(\mathcal{P}oiss)$ or for the case $f, g \in Mor(QP)$, this statement is trivial. For $\forall f, g \in Q$, $f \circ g$ never exists, because $s(f) \in ob(\mathcal{P}oiss)$ and $t(g) \in ob(QP)$. So, we have to show this statement in the case i) $g \in Mor(\mathcal{P}oiss)$, $f \in Q$ and the case ii) $g \in Q$, $f \in Mor(QP)$. For the case i), Proposition 2.13 shows that $f \circ g \in Q$. For the case ii), Proposition 2.5, Corollary 2.6, and Proposition 2.7, show that $f \circ g \in Q$, since every morphism in QP is injective. Therefore, this proposition is proved. \square

Proposition 3.2 guarantees the following definition of a category of whole space of the quantizations makes sense.

Definition 3.3 (Quantization World QW). The category of quantization QW is a subcategory of $R\text{-Mod}$ whose object is defined by

$$ob(QW) := ob(\mathcal{P}oiss) \cup ob(QP) \quad (3.2)$$

and its morphism is defined by (3.1).

A schematic diagram of this category of quantization QW is shown below.

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & M_1 & \xrightarrow{h_{12}} & M_2 & \xrightarrow{h_{23}} & M_3 & \longrightarrow & \cdots \\
& & \downarrow & & \downarrow & & \downarrow & & \\
\cdots & \longrightarrow & \langle q_1(A) \rangle & \xrightarrow{h_{12}|_{\langle q_1(A) \rangle}} & \langle q_2(A) \rangle & \xrightarrow{h_{23}|_{\langle q_2(A) \rangle}} & \langle q_3(B) \rangle & \longrightarrow & \cdots \\
& & \downarrow & & \downarrow & & \downarrow & & \\
\cdots & \longrightarrow & q_1(A) & \xrightarrow{h_{12}|_{q_1(A)}} & q_2(A) & \xrightarrow{h_{23}|_{q_2(A)}} & q_3(B) & \longrightarrow & \cdots \\
& & \uparrow & \nearrow^{q_2} & \uparrow & \nearrow^{q_3} & & & \\
\cdots & \longrightarrow & A & \xrightarrow{\phi_{AB}} & B & \longrightarrow & \cdots & &
\end{array} \tag{3.3}$$

Here M_i is an R -algebra, and $\langle q_i(A) \rangle, \dots$ are subalgebras of M_i, \dots that generated by $q_i(A), \dots$, respectively. A, B, \dots are Poisson algebras, and q_i is a quantization map. ϕ_{AB} is a surjective Poisson morphism, and h_{ij} is an injective R -algebra homomorphism. This figure (3.3) is not strict but just for help to understand relations between objects in QW .

It would not be waste to emphasize the importance of injectivity of $Mor(t(q_j), t(q_i))$ here, even though this is an overlap with what was previously stated around Proposition 2.5, Corollary 2.6, and Proposition 2.7. Consider some $q_j : A \rightarrow T_j \in Q$ i.e. $\hbar(q_j) \neq 0$, where $\langle T_j \rangle = M_j$ is a noncommutative R -algebra. Suppose there exists an R -algebra homomorphism $h_{ji} : M_j \rightarrow M_i$ such that its image $\text{Im}h_{ji} \subset M_i$ is a commutative subalgebra. For $\forall f, g \in A$,

$$\begin{aligned}
0 &= [h_{ji} \circ q_j(f), h_{ji} \circ q_j(g)] \\
&= h_{ji}([q_j(f), q_j(g)]) \\
&= \hbar(q_j) h_{ji} \circ q_j(\{f, g\}) + \tilde{O}(\hbar^{1+\epsilon}(q_j)),
\end{aligned}$$

since $\text{Im}h_{ji}$ is commutative. From $\hbar(q_j) \neq 0$,

$$h_{ji} \circ q_j(\{f, g\}) = 0, \quad \tilde{O}(\hbar^{1+\epsilon}(q_j)) = 0.$$

Furthermore, we find that there exists a pair of f, g such that $q_j(\{f, g\}) \neq 0$ since $\hbar(q_j) \neq 0$. Then we obtain that

$$\text{Ker}h_{ji}|_{T_j} \supset \{q_j(\{f, g\}) \mid f, g \in A\} \neq \{0\}. \tag{3.4}$$

The presence of this kernel prevents the construction of a category QW using Q . To avoid this, injectivity is added to the definition for $Mor(t(q_j), t(q_i))$.

4 Classification of Quantizations

As the classical limits of matrix regularization and so on teach us, what characterizes quantum geometry is not only the algebra generated by the target space of the quantization map, but also the information about the map of quantization itself. Even if we confine our considerations to algebra, $\langle q_1(A) \rangle, \langle q_2(A) \rangle, \dots$ are more important than M_1, M_2, \dots in the diagram (3.3) to characterize the quantization maps. In this section, we introduce a framework that classifies quantizations.

4.1 Equivalence of Quantization Maps

Recall that Poisson category \mathcal{Poisss} is a category whose objects are Poisson algebras, and morphisms are surjective Poisson homomorphisms.

Let us introduce U_P as a forgetful functor from \mathcal{Poisss} to QW forgetting multiplication and Poisson structure;

$$U_P : \mathcal{Poisss} \rightarrow QW.$$

For simplicity, we abbreviate $U_P(A)$ and $U_P(\phi)$ as A and ϕ for $A \in ob(\mathcal{Poisss})$, $\phi \in Mor(\mathcal{Poisss})$, respectively. Next, let I_{QP} be a functor from QP to QW such that all objects and morphisms are identically embedded in QW as $I_{QP}(QP) = QP \subset QW$ i.e. $I_{QP}(T_i) = T_i \in ob(QW)$ for any object $T_i \in ob(QP)$ and $I_{QP}(h_{ij}) = h_{ij} \in Mor(QW)$ for any morphism $h_{ij} \in Mor(QP)$. As similar to the case of U_P , we abbreviate $I_{QP}(T_i)$ and $I_{QP}(h)$ as T_i and h for $T_i \in ob(QP)$, $h \in Mor(QP)$, respectively.

Using these functors a comma category is defined as follows.

Definition 4.1 (Comma category $(U_P \downarrow I_{QP})$). $(U_P \downarrow I_{QP})$ is a comma category whose object is a triple

$$\begin{aligned} & ob(U_P \downarrow I_{QP}) \\ & := \{(A, q_{A_i}, T_i) \mid A \in ob(\mathcal{Poisss}), T_i \in ob(QP), (q_{A_i} : A \rightarrow T_i) \in Mor(QW)\} \end{aligned} \quad (4.1)$$

and its morphism $(\phi_{AB}, t_{ij}) : (A, q_{A_i}, T_i) \rightarrow (B, q_{B_j}, T_j)$ is a pair of a Poisson map $(\phi_{AB} : A \rightarrow B) \in Mor(\mathcal{Poisss})$ and an R -module homomorphism $(t_{ij} : T_i \rightarrow T_j) \in Mor(QP)$ such that

$$t_{ij} \circ q_{A_i} = q_{B_j} \circ \phi_{AB}. \quad (4.2)$$

$$\begin{array}{ccc} U_P(A) = A & \xrightarrow{\phi_{AB}} & B = U_P(B) \\ q_{A_i} \downarrow & & \downarrow q_{B_j} \\ I_{QP}(T_i) = T_i & \xrightarrow{t_{ij}} & T_j = I_{QP}(T_j) \end{array}$$

Using this comma category, quantization maps are classified. For every $(q_{A_i} : A \rightarrow T_i) \in Q$ there is a unique map $(q'_{A_i} : A \rightarrow q_{A_i}(A)) \in Q$ defined by $q'_{A_i}(f) := q_{A_i}(f)$ for any $f \in A$. For simplicity, q'_{A_i} will simply be abbreviated as q_{A_i} in the following.

Definition 4.2. When $(q_{A_i} : A \rightarrow T_i) \in Q$ and $(q_{B_j} : B \rightarrow T_j) \in Q$ satisfy

$$(A, q_{A_i}, q_{A_i}(A)) \simeq (B, q_{B_j}, q_{B_j}(B))$$

in $(U_P \downarrow I_{QP})$, that is, there exists an isomorphism $(\phi_{AB}, t_{ij}) : (A, q_{A_i}, q_{A_i}(A)) \rightarrow (B, q_{B_j}, q_{B_j}(B))$, then we say that the quantization q_{A_i} and q_{B_j} are equivalent and we denote $q_{A_i} \simeq q_{B_j}$.

This is natural definition for equivalence of quantizations. However, the relation between algebras generating from $q_{A_i}(A)$ and $q_{B_j}(B)$ are not mentioned in this definition. Note that $q_{A_i}(A)$ and $q_{B_j}(B)$ are not algebras in general but just R -modules, subsets of some R -algebras. So, we investigate this relation, next.

Let us introduce the following functor.

Definition 4.3. A functor $F : (U_P \downarrow I_{QP}) \rightarrow \mathcal{P}oiss \times R\text{-alg}$ is defined by the map between these objects:

$$(A, q_{A_i}, T_i) \mapsto (A, \langle q_{A_i}(A) \rangle), \quad (4.3)$$

and the map between these morphisms:

$$(\phi_{AB}, t_{ij}) \mapsto (\phi_{AB}, h_{ij}|_{\langle q_{A_i}(A) \rangle}), \quad (4.4)$$

where h_{ij} is an R -algebra homomorphism from $\langle T_i \rangle \rightarrow \langle T_j \rangle$ such that $h_{ij}|_{T_i} = t_{ij}$.

Let us make sure that this F is correctly defined as a functor. We have to check three points : 1) $h_{ij}|_{\langle q_{A_i}(A) \rangle}$ is uniquely determined from t_{ij} , 2) $h_{ij}(\langle q_{A_i}(A) \rangle) \subset \langle q_{B_j}(B) \rangle$, and 3) compositions are defined in a consistent manner.

1) At first, we check that $h_{ij}|_{\langle q_{A_i}(A) \rangle}$ is uniquely determined from t_{ij} .

From $h_{ij}|_{T_i} = t_{ij}$ and $q_{A_i}(A) \subset T_i$, $h_{ij}|_{q_{A_i}(A)}$ uniquely exists. Therefore, we found $h_{ij}|_{\langle q_{A_i}(A) \rangle}$ is uniquely determined from Proposition 2.3.

2)Next, we would like to show $h_{ij}(\langle q_{A_i}(A) \rangle) \subset \langle q_{B_j}(B) \rangle$, because if the following proposition does not hold, then the composition of morphisms is not defined.

Proposition 4.4. *Let A and B be Poisson algebras with a surjective Poisson morphism $\phi_{AB} : A \rightarrow B$. Let $q_{A_i} : A \rightarrow T_i$ and $q_{B_j} : B \rightarrow T_j$ be quantization maps. Suppose that there exists $t_{ij} : T_i \rightarrow T_j$ defined by $t_{ij} = h_{ij}|_{T_i}$, where h_{ij} is some R -algebra homomorphism $h_{ij} : \langle T_i \rangle \rightarrow \langle T_j \rangle$. If they satisfy the commutativity (4.2), then,*

$$h_{ij}(\langle q_{A_i}(A) \rangle) = \langle q_{B_j}(B) \rangle.$$

Proof. Because ϕ_{AB} is surjective,

$$\langle q_{B_j}(B) \rangle = \langle q_{B_j} \circ \phi_{AB}(A) \rangle. \quad (4.5)$$

From the commutativity (4.2),

$$q_{B_j} \circ \phi_{AB} = t_{ij} \circ q_{A_i} = h_{ij}|_{q_{A_i}(A)} \circ q_{A_i}. \quad (4.6)$$

Proposition 2.4 shows that

$$h_{ij}(\langle q_{A_i}(A) \rangle) = \langle h_{ij}|_{q_{A_i}(A)}(q_{A_i}(A)) \rangle. \quad (4.7)$$

From (4.5),(4.6), and (4.7), we obtain the conclusion that we want. \square

By this proposition, $h_{ij}(\langle q_{A_i}(A) \rangle) \subset \langle q_{B_j}(B) \rangle$ is shown.

3) Finally, we make sure that F is defined as a functor. Consider the compositions of morphisms $(\phi_{AB}, t_{ij}) : (A, q_{A_i}, T_i) \rightarrow (B, q_{B_j}(A), T_j)$ and $(\phi_{BC}, t_{jk}) : (B, q_{B_j}(A), T_j) \rightarrow (C, q_{C_k}(A), T_k)$ in $(U_P \downarrow I_{QP})$. From the results of 1) and 2),

$$\begin{aligned} F(\phi_{BC}, t_{jk}) \circ F(\phi_{AB}, t_{ij}) &= (\phi_{BC}, h_{jk}|_{\langle q_{B_j}(B) \rangle}) \circ (\phi_{AB}, h_{ij}|_{\langle q_{A_i}(A) \rangle}) \\ &= (\phi_{BC} \circ \phi_{AB}, h_{jk} \circ h_{ij}|_{\langle q_{A_i}(A) \rangle}) = F((\phi_{BC}, t_{jk}) \circ (\phi_{AB}, t_{ij})). \end{aligned}$$

From the above, it is shown that F is a functor.

$F(U_P \downarrow I_{QP})$ is a subcategory of $\mathcal{Poisss} \times R\text{-alg}$. We call $F(U_P \downarrow I_{QP})$ $\mathcal{Poisss} - Q(\mathcal{Poisss})$ pair category. In the following, this is abbreviated as $P - QP := F(U_P \downarrow I_{QP})$. The object of $P - QP$ is a pair of a Poisson algebra and a generated algebra from an image of a quantization map, so we call the object of $P - QP$ quantization pair. Using $P - QP$, equivalence of quantization is expressed as follows.

Theorem 4.5. *Let $q_{Ai} : A \rightarrow T_i$ and $q_{Bj} : B \rightarrow T_j$ be quantization maps in Q . In $P - QP := F(U_P \downarrow I_{QP})$, if and only if there exists an isomorphism of $\mathcal{Poisss} \times R\text{-alg}$ (ϕ_{AB}, h_{ij}) from $(A, \langle q_{Ai}(A) \rangle)$ to $(B, \langle q_{Bj}(B) \rangle)$, q_{Ai} and q_{Bj} are equivalent quantization:*

$$(A, \langle q_{Ai}(A) \rangle) \simeq (B, \langle q_{Bj}(B) \rangle) \Leftrightarrow q_{Ai} \simeq q_{Bj}. \quad (4.8)$$

Proof. When $(\phi_{AB}, h_{ij}) : (A, \langle q_{Ai}(A) \rangle) \rightarrow (B, \langle q_{Bj}(B) \rangle)$ is an isomorphism in $\mathcal{Poisss} \times R\text{-alg}$, we put $t_{ij} = h_{ij}|_{q_{Ai}(A)}$ and $t_{ji} := h_{ij}^{-1}|_{q_{Bj}(B)}$. Then for $\forall a \in q_{Ai}(A) \subset \langle q_{Ai}(A) \rangle$

$$t_{ji} \circ t_{ij}(a) = h_{ij}^{-1} \circ h_{ij}(a) = a,$$

i.e. $t_{ji} \circ t_{ij} = id_{q_{Ai}(A)}$. $t_{ij} \circ t_{ji} = id_{q_{Bj}(B)}$ is obtained similarly. These show that $q_{Ai} \simeq q_{Bj}$.

Conversely, if $q_{Ai} \simeq q_{Bj}$, there exists an isomorphism $(\phi_{AB}, t_{ij}) : (A, q_{Ai}, q_{Ai}(A)) \rightarrow (B, q_{Bj}, q_{Bj}(B))$ and some R -algebra homomorphism $h_{ij} : \langle q_{Ai}(A) \rangle \rightarrow \langle q_{Bj}(B) \rangle$ such that $h_{ij}|_{q_{Ai}(A)} = t_{ij}$. Similarly, there exists $t_{ji} = t_{ij}^{-1} : q_{Bj}(B) \rightarrow q_{Ai}(A)$ given as some restriction of an R -algebra homomorphism h_{ji} . $\forall a \in \langle q_{Ai}(A) \rangle$ is expressed as $a = \sum_k \sum_{I_k} a_{I_k} e_{i_1} \cdots e_{i_k}$ by using $e_{i_l} \in q_{Ai}(A)$. And,

$$\begin{aligned} h_{ji} \circ h_{ij}(a) &= \sum_k \sum_{I_k} a_{I_k} h_{ji} \circ h_{ij}(e_{i_1}) \cdots h_{ji} \circ h_{ij}(e_{i_k}) \\ &= \sum_k \sum_{I_k} a_{I_k} t_{ji} \circ t_{ij}(e_{i_1}) \cdots t_{ji} \circ t_{ij}(e_{i_k}) \\ &= \sum_k \sum_{I_k} a_{I_k} e_{i_1} \cdots e_{i_k} = a. \end{aligned}$$

So, we find that $h_{ji}|_{q_{Bj}(B)} \circ h_{ij}|_{q_{Ai}(A)} = id_{q_{Ai}(A)}$. $h_{ij}|_{q_{Ai}(A)} \circ h_{ji}|_{q_{Bj}(B)} = id_{q_{Bj}(B)}$ is shown in the same way. Therefore, we obtain $(A, \langle q_{Ai}(A) \rangle) \simeq (B, \langle q_{Bj}(B) \rangle)$. \square

Remark. We can introduce a functor $\langle - \rangle : QP \rightarrow R\text{-alg}$, by $T_i \in ob(QP) \mapsto \langle T_i \rangle \in ob(R\text{-alg})$ and $t_{ij} \in Mor(T_i, T_j) \mapsto h_{ij} \in Mor(\langle T_i \rangle, \langle T_j \rangle)$ where $t_{ij} = h_{ij}|_{T_i}$. Proposition 2.3 guarantees the existence of this functor. We denote the image of this functor by $\langle QP \rangle$. There are natural projection functors $\pi_p : P - QP \rightarrow R\text{-alg}$ and $\pi_{(QP)} : P - QP \rightarrow \langle QP \rangle$.

4.2 Restriction to a Single Poisson Algebra A

Until now, we have considered whole space of all Poisson algebras and their all quantizations. However, it is also important to focus on quantizations on a single Poisson algebra. In this subsection, we will discuss about it.

Let $\mathbf{1}$ be the trivial category with the only one object $*$ and the only one morphism Id_* . For a Poisson algebra A , we define a functor $A : \mathbf{1} \rightarrow QW$ by $A(*) = A \in ob(QW)$ and $A(Id_*) = Id_A \in Mor(QW)$. Let us consider the comma category associated with A and I_{QP} .

Definition 4.6 ($(A \downarrow I_{QP})$). Let A be a fixed Poisson algebra. $(A \downarrow I_{QP})$ is a category whose objects and morphisms are given by

$$ob(A \downarrow I_{QP}) = \{(q_i, T_i) := (*, q_i, T_i) \mid (q_i : A \rightarrow T_i) \in Q, T_i \in ob(QP)\}, \quad (4.9)$$

$$Mor((q_i, T_i), (q_j, T_j)) = \{t_{ij} \in Mor_{QW}(T_i, T_j) \mid t_{ij} \circ q_i = q_j\}. \quad (4.10)$$

Roughly speaking, $(A \downarrow I_{QP})$ is a category of all quantizations of A . Note that $(A \downarrow I_{QP})$ is similar to a co-slice category but different. In the same way as $P - QP$, we define the following functor.

Definition 4.7 (F_A). The functor $F_A : (A \downarrow I_{QP}) \rightarrow R\text{-alg}$ is defined by

$$F_A((q_i, T_i)) = \langle q_i(A) \rangle$$

for $\forall (q_i, T_i) \in ob(A \downarrow I_{QP})$, and for $t_{ij} \in Mor_{(A \downarrow I_{QP})}((q_i, T_i), (q_j, T_j))$

$$F_A(t_{ij}) = h_{ij}|_{\langle q_i(A) \rangle},$$

where h_{ij} is an R -algebra homomorphism satisfying $h_{ij}|_{T_i} = t_{ij}$.

From Theorem 4.5 with $A = B$, we obtain the following.

Corollary 4.8. *Let q_i and q_j be quantization maps whose source is a Poisson algebra A . q_i and q_j are equivalent quantization of A in $F_A(A \downarrow I_{QP})$ if and only if object $\langle q_i(A) \rangle \in R\text{-alg}$ is isomorphic to $\langle q_j(A) \rangle \in R\text{-alg}$ in $R\text{-alg}$.*

From this corollary, we found that $F_A(A \downarrow I_{QP})$ classifies quantization of A .

Example 4.9. *Let $\mathcal{P}(\mathbb{C}^2)$ be the algebra of polynomials of a coordinate $(x, y) \in \mathbb{C}^2$. By introducing a Poisson bracket by*

$$\{f, g\} = \partial_x f \partial_y g - \partial_y f \partial_x g,$$

for any $f, g \in \mathcal{P}(\mathbb{C}^2)$, $(\mathcal{P}(\mathbb{C}^2), \cdot, \{, \})$ is regarded as a Poisson algebra.

We use the notation $x^1 := x$, $x^2 := y$, $\partial_1 = \frac{\partial}{\partial x}$, $\partial_2 = \frac{\partial}{\partial y}$. Let $H = (H_{ij})$ be a 2×2 complex matrix. We introduce a bi-differential operator \overleftrightarrow{H} as

$$f \overleftrightarrow{H} g = H_{ij} \partial_i f \partial_j g, \quad (f, g \in \mathcal{P}(\mathbb{C}^2)). \quad (4.11)$$

Here $H_{ij} \partial_i f \partial_j g$ is an abbreviation for $\sum_{i,j=1}^2 H_{ij} \partial_i f \partial_j g$. Moyal product $*_H$ is defined by

$$f *_H g = f(\exp \nu \overleftrightarrow{H})g = \sum_{l=0}^{\infty} \frac{\nu^l}{l!} H_{i_1 j_1} \cdots H_{i_l j_l} (\partial_{i_1} \cdots \partial_{i_l} f) \cdot (\partial_{j_1} \cdots \partial_{j_l} g)$$

where ν is a complex number. Then, by the Moyal product $*_H$, the strict deformation quantization $(\mathcal{P}(\mathbb{C}^2), *_H)$ is an R -algebra [35]. We define $q_H : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_H)$ by the inclusion map i.e. $q_H(\sum_{ij} a_{ij} x^i y^j) = \sum_{ij} a_{ij} x^i y^j$. It is clear that q_H is a quantization map.

Let us decompose $H = (H_{ij})$ into a symmetric matrix and a skew symmetric matrix as

$$H = J + K, \quad J = (J_{ij}), K = (K_{ij}), \text{ where } J_{ij} = -J_{ji}, \quad K_{ij} = K_{ji}.$$

Here we introduce a differential operator

$$T = \exp\left(-\frac{1}{2}\nu K_{ij}\partial_i\partial_j\right) = \sum_{l=0}^{\infty} \frac{1}{l!} \left(-\frac{1}{2}\nu K_{ij}\partial_i\partial_j\right)^l \quad (4.12)$$

that act a polynomial f as

$$T(f) = f + \sum_{l=1}^{\infty} \nu^l \frac{1}{l!} \left(-\frac{1}{2}K_{mj}\partial_m\partial_j\right)^l f. \quad (4.13)$$

To compare two algebra $(\mathcal{P}(\mathbb{C}^2), *_J)$ $(\mathcal{P}(\mathbb{C}^2), *_H)$, the following theorem is useful.

Theorem 4.10 (Tomihisa-Yoshioka [43]). $(\mathcal{P}(\mathbb{C}^2), *_H)$ and $(\mathcal{P}(\mathbb{C}^2), *_J)$ are R -algebra isomorphic. T defined by (4.12) satisfies

1. $T : (\mathcal{P}(\mathbb{C}^2), *_H) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_J)$ is linear and bijective.
2. $T(1) = 1$
3. For $f, g \in (\mathcal{P}(\mathbb{C}^2), *_H)$, $T(\tilde{f} *_H \tilde{g}) = T(\tilde{f}) *_J T(\tilde{g})$.

In other words, T is an R -algebra isomorphism.

We define $q_J : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_J)$ as an inclusion map i.e. $q_J(\sum_{ij} a_{ij} x^i y^j) = \sum_{ij} a_{ij} x^i y^j$, too. Then $\langle q_H(\mathcal{P}(\mathbb{C}^2)) \rangle = (\mathcal{P}(\mathbb{C}^2), *_H) \simeq \langle q_J(\mathcal{P}(\mathbb{C}^2)) \rangle = (\mathcal{P}(\mathbb{C}^2), *_J)$. From Corollary 4.8, q_H and q_J are equivalent quantizations.

5 Classical Limit

One of the most important problem is : how can we find the classical limit from the spaces given by the quantizations of Poisson algebras? To make the point of this problem easier to understand, let us consider matrix regularization of S^2 and T^2 . (In Appendix C, we summarize the matrix regularization of S^2 and T^2 .) As an well-known fact, both images of quantization maps from \mathcal{A} and \mathcal{B} generate the same matrix algebras. This example shows that the classical limit is not determined by observation of the algebra obtained from a single quantization. This shows that the Poisson algebra cannot be easily determined as the classical limit from the algebra obtained by quantization. But this example also teaches that the classical limit may be identified from the sequence of modules before generating algebras.

Since a new formulation of quantization using category theory was introduced in Section 3, we can define various classical limits in the framework. The purpose of this

section is to define some classical limits, study their properties by examining them with concrete examples, and look for useful definitions of classical limits for approaching inverse problems. For this purpose, we pay particular attention to the way the sequence of R -modules obtained by the quantization maps determines the Poisson algebra.

In the following $\lim_{\leftarrow \mathcal{I}} D$ means the categorical limit in this article. For example we can see its definition in Chapter V in [30] or Chapter 5 in [27].

5.1 Naive Classical Limit

For a start, the classical limit is defined as follows using a naive definition of the limit (in the sense of category theory).

Definition 5.1 (Naive classical limit of $D(\mathcal{I})$). Let \mathcal{I} be an index category, and let $D : \mathcal{I} \rightarrow QW$ be a diagram of shape \mathcal{I} . Let A be a Poisson algebra. Suppose that \mathbf{t}_A is a set of $t_i : A \rightarrow D(i) \in Q$ ($i \in \mathcal{I}$) such that t_i and t_j in \mathbf{t}_A satisfy $t_j = D(u_{ij}) \circ t_i$ for $\forall u_{ij} \in Mor_{\mathcal{I}}(i, j)$, that is (A, \mathbf{t}_A) is a cone over D . When \mathcal{I} and D have a limit (A, \mathbf{t}_A) :

$$\lim_{\leftarrow \mathcal{I}} D = (A \xrightarrow{t_i} D(i))_{i \in \mathcal{I}}, \quad (5.1)$$

and $\lim_{\leftarrow \mathcal{I}} D$ is called the naive classical limit of $D(\mathcal{I})$.

This definition may lead some readers to think that the naive classical limit is unrelated to the conventional classical limit. However, the following theorem shows that the naive classical limit is a generalization that includes the classical limit when only one type of quantization for one Poisson algebra is considered conventionally.

Theorem 5.2. *Let \mathcal{I} be an index category, and let $D : \mathcal{I} \rightarrow QW$ be a diagram of shape \mathcal{I} such that there is only one Poisson algebra A in $D(\mathcal{I})$. If (A, \mathbf{t}_A) is a cone over D , then (A, \mathbf{t}_A) is the naive classical limit.*

Proof. In the following, we denote $(A, \cdot, \{ \cdot, \cdot \})$ by A for simplicity, in contexts where there is no risk of confusion. The cone (A, \mathbf{t}) is expressed as:

$$\begin{array}{ccccccc} (A, \{, \}) & & & & & & \\ \downarrow Id_A & \searrow^{t_2 \dots} & & & & & \\ (A, \{, \}) & \xrightarrow{t_1} & (T_1, *1) & \xrightarrow{t_{12}} & (T_2, *2) & \longrightarrow & \dots \end{array}$$

Here each $t_i : A \rightarrow T_i$ is a quantization map, and $t_{ij} = D(u_{ij}) : T_i \rightarrow T_j$ is in $Mor_{QP}(T_i, T_j)$. $D(\mathcal{I})$ is expressed as

$$(A, \{, \}) \xrightarrow{t_1} (T_1, *1) \xrightarrow{t_{12}} (T_2, *2) \longrightarrow \dots$$

If there is another cone of (B, \mathbf{q}') , B is a Poisson algebra, since A is a Poisson algebra, and only Poisson algebras have possibility to be sources of morphisms in QW which

target is A . We denote this Poisson morphism by $\phi : B \rightarrow A$. $\mathbf{q}' = \{\phi, q'_{B1}, q'_{B2}, \dots\}$ is a set of morphisms that satisfy the following commutative diagram:

$$\begin{array}{ccccccc}
 & & B & & & & \\
 & & \downarrow \phi & \searrow q'_{B1} & \searrow q'_{B2} \cdots & & \\
 (A, \{, \}) & \xrightarrow{t_1} & (T_1, *1) & \xrightarrow{t_{12}} & (T_2, *2) & \rightleftharpoons & \cdots
 \end{array}$$

Suppose two $\phi, \phi' : B \rightarrow A$ satisfy the following commutative diagram:

$$\begin{array}{ccccccc}
 & & & \phi & & & \\
 & & & \downarrow & & & \\
 (A, \{, \}) & \xleftarrow{\phi} & B & \xrightarrow{q'_{B1}} & (A, *p) & \xrightarrow{t_{pr}} & (A, *r) \rightleftharpoons \cdots \\
 Id_A \downarrow & & \downarrow \phi & & \downarrow q'_{B1} & & \\
 (A, \{, \}) & \xrightarrow{q_p} & (A, *p) & \xleftarrow{t_{rp}} & (A, *r) & \rightleftharpoons & \cdots
 \end{array}$$

From the condition that $(A, \{, \})$ is an object in $D(\mathcal{I})$, Id_A is also a morphism in $D(\mathcal{I})$ because D is a functor. From the commutativity $\phi = Id_A \circ \phi' = \phi'$, we obtain the uniqueness of ϕ . \square

In the following subsections, we will actually consider two examples of quantization families corresponding to only one type of quantization for a single Poisson algebra and check that their naive limits are in fact what is expected.

5.2 Example of Naive Classical Limit ($\mathcal{P}(\mathbb{C}^2)$)

We consider $\mathcal{P}(\mathbb{C}^2)$, the set of all polynomials of x and y , and its quantization by deformation quantization, again. In the following, we denote $(\mathcal{P}(\mathbb{C}^2), \cdot, \{, \})$ by $\mathcal{P}(\mathbb{C}^2)$ for simplicity, in contexts where there is no risk of confusion. (See Example 4.9.)

Let introduce sequence of Moyal products with p ($p \in \mathbb{Q}$) by

$$f *_p g = f \exp \left(\overleftarrow{\partial}_x \frac{ip}{2} \overrightarrow{\partial}_y - \overleftarrow{\partial}_y \frac{ip}{2} \overrightarrow{\partial}_x \right) g,$$

where $f, g \in \mathcal{P}(\mathbb{C}^2)$. We define $q_p : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_p)$ by inclusion map i.e. $q_p(f) = f$, as similar to Example 4.9.

Let \mathcal{I}_{moyal} be an index category defined by $ob(\mathcal{I}_{moyal}) = \mathbb{Q}$ and

$$\begin{aligned}
 Mor(p, r) &:= \{(p \mapsto r =: +r - p)\} \text{ for } r \neq 0, p \in \mathbb{Q}, \\
 Mor(p, 0) &:= \emptyset \text{ (} p \neq 0 \text{)}, \quad Mor(0, 0) := \{Id_0 = +0 - 0\}.
 \end{aligned}$$

Note that $+p - p$ is an identity for p .

Let $D_{moyal} : \mathcal{I}_{moyal} \rightarrow QW$ be a diagram of shape \mathcal{I}_{moyal} defined by

$$\begin{aligned}
 D_{moyal}(p) &= (\mathcal{P}(\mathbb{C}^2), *_p) \text{ for } \forall p \neq 0 \in \mathbb{Q} \\
 D_{moyal}(0) &= (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \text{ for } 0 \in \mathbb{Q}
 \end{aligned}$$

and

$$\begin{aligned}
 D_{moyal}(+r - p) &= t_{p,r} \\
 D_{moyal}(+r - 0) &= q_r.
 \end{aligned}$$

Here $t_{p,r}$ is an intertwiner between $(\mathcal{P}(\mathbb{C}^2), *_{p})$ and $(\mathcal{P}(\mathbb{C}^2), *_{r})$. The intertwiner is given as an R -algebra isomorphism. The existence of such intertwiner is known [9, 10, 13, 14], and the existence of this $t_{p,r}$ is not essential to the following discussion. We choose here each isomorphism $t_{p,r}$ satisfying $t_{p,r} \circ q_p = q_r$.

The diagram $D_{moyal}(\mathcal{I}_{moyal})$ is given as:

$$(\mathcal{P}(\mathbb{C}^2), \{, \}) \xrightarrow{q_p} (\mathcal{P}(\mathbb{C}^2), *_{p}) \begin{array}{c} \xrightarrow{t_{pr}} \\ \xleftarrow{t_{rp}} \end{array} (\mathcal{P}(\mathbb{C}^2), *_{r}) \rightleftharpoons \dots$$

Then, $(\mathcal{P}(\mathbb{C}^2), \mathbf{q})$ is a cone:

$$\begin{array}{ccc} (\mathcal{P}(\mathbb{C}^2), \{, \}) & & \\ \downarrow Id_{\mathcal{P}(\mathbb{C}^2)} & \searrow q_p & \\ (\mathcal{P}(\mathbb{C}^2), \{, \}) & \xrightarrow{q_p} & (\mathcal{P}(\mathbb{C}^2), *_{p}) \begin{array}{c} \xrightarrow{t_{pr}} \\ \xleftarrow{t_{rp}} \end{array} (\mathcal{P}(\mathbb{C}^2), *_{r}) \rightleftharpoons \dots \end{array}$$

From Theorem 5.2, $((\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}), \mathbf{q})$ is the naive classical limit.

Corollary 5.3. *Let \mathbf{q} be a set of the above quantization $q_p : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_{p})$ ($p \in \mathbb{Q}$) and identity $Id_{\mathcal{P}(\mathbb{C}^2)}$ for Poisson algebra $(\mathcal{P}(\mathbb{C}^2), \cdot, \{, \})$ i.e. $\mathbf{q} = \{q_p \in Q \mid p \in \mathbb{Q}\} \cup \{Id_{\mathcal{P}(\mathbb{C}^2)}\}$. For the above \mathcal{I}_{moyal} , D_{moyal} , $((\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}), \mathbf{q})$ is the naive classical limit.*

5.3 Example of Naive Classical Limit (Fuzzy sphere)

In this subsection, let us consider the naive classical limit of fuzzy spheres. Basic knowledge for the fuzzy sphere and the notations used in this subsection are summarized in the Appendix C.

Let $P[x^a]$ be the algebra of polynomial generated by x^a . For an ideal I which is generated by the relation (C.1), we consider a Poisson algebra $\mathcal{A} = P[x^a]/I$ with Poisson bracket $\{x^a, x^b\} = \epsilon^{abc}x^c$. For arbitrary $f \in \mathcal{A}$ is given as

$$f = f_0 + f_a x^a + \frac{1}{2} f_{ab} x^a x^b + \dots$$

where $f_{a_1 \dots a_i} \in \mathbb{C}$ is completely symmetric and trace-free. In the case of $k \geq 2$, quantization maps $t_k : \mathcal{A} \rightarrow T_k := t_k(\mathcal{A}) \subset M_k = \text{Mat}_k$ are defined by

$$\begin{aligned} t_k(f) &:= f_0 \mathbf{1}_k + f_{a_1} t_k(x^{a_1}) + \dots + f_{a_1 \dots a_{k-1}} t_k(x^{a_1} \dots x^{a_{k-1}}) \\ t_k(x^a) &:= \hbar_k J_k^a = X_k^a, \end{aligned}$$

where J_a ($a = 1, 2, 3$) satisfy $[J_k^a, J_k^b] = i\epsilon^{abc} J_k^c$, that is J_a are generators of k -dimensional irreducible representation of $\mathfrak{su}(2)$. Its Casimir relation $J_k^a J_{ka} = \frac{1}{4}(k^2 - 1)\mathbf{1}_k$ and (C.1) derive the relation $\frac{4}{k^2 - 1} = \hbar_k^2$. The commutation relation of J_k^a or (C.4) shows that $t_k \in Q$ ($k \geq 2$).

Let \mathcal{I}_{fzS2} be an index category defined by $ob(\mathcal{I}_{fzS2}) = \mathbb{Z}_{>0}$ and

$$\begin{aligned} Mor(1, k) &:= \{(1 \mapsto k) =: \hat{k}\} \text{ for } k \in \mathbb{Z}_{>0}, \\ Mor(p, q) &:= \emptyset \text{ for integers } p, q > 1, p \neq q, \\ Mor(p, p) &:= Id_p. \end{aligned}$$

The absence of a morphism between p and q ($p, q > 1$ and $p \neq q$) in the index category is due to the fact that there is no morphism between T_p and T_q . Since $\ker t_p = \{f_{a_1 \dots a_p} x^{a_1} \dots x^{a_p} + f_{a_1 \dots a_{p+1}} x^{a_1} \dots x^{a_{p+1}} + \dots\}$, there is no injective morphism between T_p and T_q that satisfies the commutative diagram with t_p and t_q .

Let $D_{fzS2} : \mathcal{I}_{fzS2} \rightarrow QW$ be a diagram of shape \mathcal{I}_{fzS2} defined by

$$\begin{aligned} D_{fzS2}(1) &= \mathcal{A} \text{ for } 1, \\ D_{fzS2}(k) &= T_k \text{ for } \forall k \neq 1 \in \mathbb{Z}_{>0}, \end{aligned}$$

and

$$\begin{aligned} D_{fzS2}(\hat{1}) &= Id_{\mathcal{A}}, \\ D_{fzS2}(\hat{k}) &= t_k \text{ for } \forall k > 1, \\ D_{fzS2}(Id_k) &= Id_{T_k} \text{ for } \forall k > 1, \end{aligned}$$

$$\left(\begin{array}{c} 1 \\ \downarrow 2 \\ 2 \end{array} \begin{array}{c} \searrow 3 \\ \searrow 3 \\ 3 \end{array} \begin{array}{c} \searrow 4 \dots \\ \searrow 4 \dots \\ 4 \dots \end{array} \right) \xrightarrow{D_{fzS2}} \left(\begin{array}{c} \mathcal{A} \\ \downarrow t_2 \\ T_2 \end{array} \begin{array}{c} \searrow t_3 \\ \searrow t_3 \\ T_3 \end{array} \begin{array}{c} \searrow t_4 \dots \\ \searrow t_4 \dots \\ T_4 \dots \end{array} \right).$$

For the \mathcal{I}_{fzS2} and D_{fzS2} , the naive classical limit is determined.

Corollary 5.4. *Let \mathbf{t} be a set of the above quantizations $\mathbf{t} = \{t_k \in Q \mid k > 1\}$. For the above \mathcal{I}_{fzS2} , D_{fzS2} , $(\mathcal{A}, \mathbf{t})$ is the naive classical limit.*

This follows trivially from Theorem 5.2.

5.4 Weak Classical Limit

As we have seen, in the naive classical limit, the limit is trivially determined for some kinds of diagrams $D(\mathcal{I})$. From the viewpoint of discussing inverse problems, the range of the naive classical limit is too wide. In this subsection, we give a definition of the other classical limit with inverse problems in mind. We introduce it by restricting the diagram to QP and by using a limit that differs from the limit of category theory.

Let us introduce a quantization family and a kind of classical limit.

Definition 5.5 (Quantization Family of A , weak classical limit of $D(\mathcal{I})$). Let \mathcal{I} be an index category, and let $D : \mathcal{I} \rightarrow QP (= I_{QP}(QP)) \subset QW$ be a diagram of shape \mathcal{I} . Let A be a Poisson algebra. Suppose that \mathbf{t}_A is a set of $(t_i : A \rightarrow D(i)) \in Q$ ($i \in \mathcal{I}$) such that t_i and t_j in \mathbf{t}_A satisfy $t_j = D(u_{ij}) \circ t_i$ for $\forall u_{ij} \in Mor_{\mathcal{I}}(i, j)$, that is, (A, \mathbf{t}_A) is a cone over D . $D(\mathcal{I})$ is called a quantization family of A , and (A, \mathbf{t}_A) is called the weak classical limit of $D(\mathcal{I})$, when the cone (A, \mathbf{t}_A) satisfies the following condition:

- For any other cone (B, \mathbf{q}_B) over D , where $B \in ob(\mathcal{P}oiss) \subset ob(QW)$, there exists a unique map $(\phi_{BA} : B \rightarrow A) \in Mor_{QW}(B, A)$ such that $q_i : B \rightarrow D(i) \in \mathbf{q}_B$ satisfies

$$q_i = t_i \circ \phi_{BA}$$

for all $i \in \mathcal{I}$.

$$\begin{array}{ccc}
A & \xleftarrow{\phi_{BA}} & B & & : \mathcal{P}oiss(= U_p(\mathcal{P}oiss)) \subset QW \\
\downarrow & \searrow & \downarrow & \searrow & \dots \\
D(i) & \longrightarrow & D(j) & \longrightarrow & D(k) \dots & : QP(= I_{QP}(QP)) \subset QW
\end{array}$$

5.5 Examples of Candidates of Weak Classical limits

We consider $\mathcal{P}(\mathbb{C}^2)$, again. As in Subsection 5.2, the sequence of Moyal products with p ($p \in \mathbb{Q}^\times$) by $f *_p g = f \exp\left(\overleftarrow{\partial}_x \frac{ip}{2} \overrightarrow{\partial}_y - \overleftarrow{\partial}_y \frac{ip}{2} \overrightarrow{\partial}_x\right) g$, where $f, g \in \mathcal{P}(\mathbb{C}^2)$, and $q_p : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_p)$ is defined by $q_p(f) = f$.

\mathcal{I}'_{moyal} differs from the one in Subsection 5.2 in that 0 is excluded from the objects in \mathcal{I}_{moyal} . Let \mathcal{I}_{moyal} be an index category defined by $ob(\mathcal{I}'_{moyal}) = \mathbb{Q}^\times$ and

$$Mor(p, r) := \{(p \mapsto r =: +r - p)\} \text{ for } r, p \in \mathbb{Q}^\times.$$

Note that $+r - p$ is an identity for p .

Let $D'_{moyal} : \mathcal{I}'_{moyal} \rightarrow QP \subset QW$ be a diagram of shape \mathcal{I}'_{moyal} defined by $D'_{moyal}(p) = (\mathcal{P}(\mathbb{C}^2), *_p)$ for $\forall p \in \mathbb{Q}^\times$ and $D'_{moyal}(+r - p) = t_{p,r}$. Here $t_{p,r}$ is an intertwiner as the same one in Subsection 5.2.

The diagram $D'_{moyal}(\mathcal{I}'_{moyal})$ is given as:

$$(\mathcal{P}(\mathbb{C}^2), *_p) \begin{array}{c} \xleftarrow{t_{pr}} \\ \xrightarrow{t_{rp}} \end{array} (\mathcal{P}(\mathbb{C}^2), *_r) \rightleftharpoons \dots$$

Let \mathbf{q}' be a set of the above quantization $q_p : (\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}) \rightarrow (\mathcal{P}(\mathbb{C}^2), *_p)$ ($p \in \mathbb{Q}^\times$). Even with these change, $(\mathcal{P}(\mathbb{C}^2), \mathbf{q}')$ is still a cone:

$$\begin{array}{ccc}
(\mathcal{P}(\mathbb{C}^2), \{, \}) & & \\
q_p \downarrow & \searrow^{q_r \dots} & \\
(\mathcal{P}(\mathbb{C}^2), *_p) & \begin{array}{c} \xleftarrow{t_{pr}} \\ \xrightarrow{t_{rp}} \end{array} & (\mathcal{P}(\mathbb{C}^2), *_r) \rightleftharpoons \dots
\end{array}$$

At this case, the following proposition holds.

Proposition 5.6. *If there exist the weak classical limit for the above \mathcal{I}'_{moyal} , D'_{moyal} , the weak classical limit is isomorphic to $((\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}), \mathbf{q}')$.*

Proof. Let (L, \mathbf{q}_{lim}) be the weak quantization. Then there exists a unique Poisson morphism $\phi_{PL} : \mathcal{P}(\mathbb{C}^2) \rightarrow L$ such that $q_p = q_{Lp} \circ \phi_{PL}$ for any $(q_{Lp} : L \rightarrow (\mathcal{P}(\mathbb{C}^2), *_p)) \in \mathbf{q}_{lim}$ and any $q_p \in \mathbf{q}'$. Recall that this ϕ_{PL} is a surjective Poisson morphism. By definition of q_p , $f = q_p(f) = q_{Lp} \circ \phi_{PL}(f)$ for $\forall f \in \mathcal{P}(\mathbb{C}^2)$. Then ϕ_{PL} is injective. \square

In fact, if there is another cone, from which there is a morphism to $\mathcal{P}(\mathbb{C}^2)$ satisfying commutativity, then it is the unique one.

Proposition 5.7. *Let $((\mathcal{P}(\mathbb{C}^2), \cdot, \{, \}), \mathbf{q}')$ be a cone defined above. Let (B, \mathbf{t}) be the other cone where B is an object in QW . If there is $\phi : B \rightarrow \mathcal{P}(\mathbb{C}^2) \in Mor(QW)$*

satisfies the following commutative diagram:

$$\begin{array}{ccc}
\mathcal{P}(\mathbb{C}^2) & \xleftarrow{\phi} & B \\
q_p \downarrow & \searrow^{t_p} & \downarrow t_r \quad \dots \\
(\mathcal{P}(\mathbb{C}^2), *_{p}) & \xrightleftharpoons[t_{rp}]{t_{pr}} & (\mathcal{P}(\mathbb{C}^2), *_{r}) \rightleftharpoons \dots
\end{array}
,$$

then ϕ is unique.

Proof. Suppose two different $\phi, \phi' : B \rightarrow \mathcal{P}(\mathbb{C}^2)$ satisfy the following commutative diagram:

$$\begin{array}{ccc}
\mathcal{P}(\mathbb{C}^2) & \xleftarrow{\phi, \phi'} & B \\
q_p \downarrow & \searrow^{t_p} & \downarrow t_r \quad \dots \\
(\mathcal{P}(\mathbb{C}^2), *_{p}) & \xrightleftharpoons[t_{rp}]{t_{pr}} & (\mathcal{P}(\mathbb{C}^2), *_{r}) \rightleftharpoons \dots
\end{array}
.$$

There exists $b \in B$ such that $\phi(b) \neq \phi'(b)$. From the commutativity $t_p = q_p \circ \phi = q_p \circ \phi'$. By definition of each q_p , $\phi(b) = q_p \circ \phi(b) = q_p \circ \phi'(b) = \phi'(b)$. This is a contradiction. \square

Unfortunately, we do not know if the weak classical limit actually exists for this \mathcal{I}'_{moyal} , D'_{moyal} .

Next, as similar in Subsection 5.3, let us consider the example of fuzzy sphere, again. Let $(\mathcal{A} = P[x^a]/I, \{, \})$ be a Poisson algebra, where $\{x^a, x^b\} = \epsilon^{abc}x^c$ ($a, b, c = 1, 2, 3$). $f \in \mathcal{A}$ is given as $f = f_0 + f_a x^a + \frac{1}{2}f_{ab}x^a x^b + \dots$ where $f_{a_1 \dots a_i} \in \mathbb{C}$ is completely symmetric and trace-free. Quantization maps $t_k : \mathcal{A} \rightarrow T_k := t_k(\mathcal{A}) \subset M_k = \text{Mat}_k$ are defined by $t_k(f) := f_0 \mathbf{1}_k + f_{a_1} X_k^{a_1} + \dots + f_{a_1 \dots a_{k-1}} X_k^{a_1} \dots X_k^{a_{k-1}}$.

Let \mathcal{I}'_{fzS2} be an index category defined by $ob(\mathcal{I}'_{fzS2}) = \{2, 3, 4, \dots\}$ and

$$\begin{aligned}
Mor(p, q) &:= \emptyset \text{ for integers } p, q > 1, p \neq q, \\
Mor(p, p) &:= Id_p.
\end{aligned}$$

In other words, \mathcal{I}'_{fzS2} is a discrete category. Let $D'_{fzS2} : \mathcal{I}'_{fzS2} \rightarrow QP \subset QW$ be a diagram of shape \mathcal{I}'_{fzS2} defined by

$$D'_{fzS2}(k) = T_k \text{ for } \forall k \in \{2, 3, 4, \dots\},$$

and

$$D'_{fzS2}(Id_k) = Id_{T_k} \text{ for } \forall k \in \{2, 3, 4, \dots\}.$$

$$(2 \quad 3 \quad 4 \quad \dots) \xrightarrow{D'_{fzS2}} (T_2 \quad T_3 \quad T_4 \quad \dots).$$

Of course, $(\mathcal{A}, \mathbf{t})$ is a cone, where $\mathbf{t} = \{t_k \in Q \mid k > 1\}$.

For the \mathcal{I}'_{fzS2} and D'_{fzS2} , the following proposition is derived.

Proposition 5.8. *If there exists the weak classical limit for the \mathcal{T}'_{fzS2} and D'_{fzS2} , then the weak classical limit is isomorphic to $(\mathcal{A}, \mathfrak{t})$.*

Proof. Let (L, \mathbf{q}_{lim}) be the weak quantization. Then there exists a unique Poisson morphism $\phi_{AL} : \mathcal{A} \rightarrow L$ such that $t_k = q_{Lk} \circ \phi_{AL}$ for any $q_{Lk} : L \rightarrow T_k \in \mathbf{q}_{lim}$ and any $t_k \in \mathfrak{t}$. Recall that this ϕ_{AL} is a surjective Poisson morphism. Let us show that this ϕ_{AL} is injective by contradiction. Suppose that ϕ_{AL} is not injective. There are $f, g \in \mathcal{A}$ such that $f \neq g$ and $\phi_{AL}(f) = \phi_{AL}(g)$. For any k , $t_k(f) = t_k(g)$ since $t_k = q_{Lk} \circ \phi_{AL}$. For sufficiently large k , this is contradiction, and ϕ_{AL} is injective. \square

As a matter of fact, there is stronger uniqueness of the Poisson morphism in the case of fuzzy spheres.

Proposition 5.9. *Let \mathfrak{t} and \mathfrak{q} be sets of the quantizations $\mathfrak{t} = \{(t_k : \mathcal{A} \rightarrow T_k) \in Q \mid k > 1\}$ and $\mathfrak{q} = \{(q_k : B \rightarrow T_k) \in Q \mid k > 1\}$, where B is arbitrary object in QW . If there is $\phi : B \rightarrow \mathcal{A} \in Mor(QW)$ satisfies the following commutative diagram:*

$$\begin{array}{ccccc} \mathcal{A} & \xleftarrow{\phi} & B & & \\ \downarrow & \searrow & \downarrow & \searrow & \dots \\ T_2 & & T_3 & & T_4 \dots \end{array},$$

then ϕ is unique.

Proof. B is a Poisson algebra, since \mathcal{A} is a Poisson algebra, and only Poisson algebras have possibility to be sources of morphisms in QW which target is \mathcal{A} . Suppose that there exist two morphisms $\phi, \phi' : B \rightarrow \mathcal{A}$ such that $t_k \circ \phi = t_k \circ \phi' = q_k$.

$$\begin{array}{ccccc} \mathcal{A} & \xleftarrow{\phi, \phi'} & B & & \\ \downarrow & \searrow & \downarrow & \searrow & \dots \\ T_2 & & T_3 & & T_4 \dots \end{array}$$

If $\phi \neq \phi'$, there exist $b \in B$ such that $\phi(b) \neq \phi'(b)$ and $t_n(\phi(b)) = t_n(\phi'(b))$ for $\forall n \in \mathbb{Z}_{>0}$. We denote the coimage of t_N by $H_N := \{f_0 + f_{a_1}x^{a_1} + \dots + f_{a_1 \dots a_{N-1}}x^{a_1} \dots x^{a_{N-1}}\}$. For sufficiently large N , $\phi(b), \phi'(b) \in H_N$, and $t_N(\phi(b)) \neq t_N(\phi'(b))$. This is a contradiction, so the uniqueness of the map ϕ is shown. \square

If \mathcal{I} and D are chosen to satisfy Theorem 5.2, the naive classical limit is obviously determined, as shown in the examples above. On the other hand, to get the weak classical limits we have to choose appropriate pairs of an index category and a diagram. If we can not find an appropriate pair of them, the weak classical limit is not determined in general. Moreover, since any Poisson algebra is a candidate for this weak classical limit, proving the existence of the weak classical limit may be difficult in general.

5.6 Strong Classical Limit through $(A \downarrow I_{QP})$

Apart from the weak classical limit that we have examined in the previous subsections, it is also natural to assume that the classical limit should be determined from the all quantization spaces corresponding to a single Poisson algebra. In particular, it is

expected that the fixed Poisson algebra is a candidate for the classical limit. In this subsection, we investigate such classical limits.

To prepare for this, we first introduce a way to construct an index category from a category with total order.

Definition 5.10. Let \mathcal{C} be a category with total order denoted by $c_i \leq c_j$ ($c_i, c_j \in \text{ob}(\mathcal{C})$). J^\bullet is a functor like correspondent (but not functor) between \mathcal{C} and an index category $\mathcal{I} := J^\bullet(\mathcal{C})$ defined by the following conditions:

1. Between objects of the two categories,

$$\begin{aligned} J^\bullet : \text{ob}(\mathcal{C}) &\xrightarrow{\cong} \text{ob}(\mathcal{I}) = \text{ob}(J^\bullet(\mathcal{C})) \\ c &\longmapsto i_c := J^\bullet(c) \end{aligned}$$

is a one to one correspondence.

2. $\forall t_{ij} \in \text{Mor}_{\mathcal{C}}(c_i, c_j)$ if $c_i \leq c_j$ then there exists a $f_{ij} = J^\bullet(t_{ij}) \in \text{Mor}_{\mathcal{I}}(i_{c_i}, i_{c_j})$.
3. $\forall f_{kl} \in \text{Mor}_{\mathcal{I}}(i_{c_k}, i_{c_l})$, there exists a unique $t_{kl} \in \text{Mor}_{\mathcal{C}}(c_k, c_l)$ such that $c_k \leq c_l$ and $f_{kl} = J^\bullet(t_{kl})$.

Note that if $c_k > c_l$ then $\text{Mor}_{\mathcal{I}}(i_{c_k}, i_{c_l}) = \emptyset$ even when $\text{Mor}_{\mathcal{C}}(c_k, c_l) \neq \emptyset$. In short, J^\bullet acts to remove morphisms from bigger object to smaller object in \mathcal{C} . This is the reason that J^\bullet is not a functor.

Therefore, we have to show the following.

Proposition 5.11. $\mathcal{I} := J^\bullet(\mathcal{C})$ is a category.

Proof. The only condition that needs to be checked is consistency regarding $\text{Mor}_{\mathcal{I}}(i_{c_k}, i_{c_l}) = \emptyset$ when $\text{Mor}_{\mathcal{C}}(c_k, c_l) \neq \emptyset$ with $c_k > c_l$. For any two $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ their composition $g \circ f : X \rightarrow Z$ have to be a morphism of $\text{Mor}(X, Z)$. In our situation, if the morphism corresponding to $g \circ f$ is removed without removing the morphism corresponding to g or the morphism corresponding to f , then it is impossible for $\mathcal{I} := J^\bullet(\mathcal{C})$ to be a category. Let c_X, c_Y, c_Z be objects of \mathcal{C} and $i_X = J^\bullet(c_X), i_Y = J^\bullet(c_Y), i_Z = J^\bullet(c_Z)$ be objects of \mathcal{I} . For the case $c_X > c_Z$, $\text{Mor}_{\mathcal{I}}(i_X, i_Z)$ becomes \emptyset , so if $\text{Mor}_{\mathcal{I}}(i_X, i_Y) \neq \emptyset$ and $\text{Mor}_{\mathcal{I}}(i_Y, i_Z) \neq \emptyset$, then it is inconsistent for \mathcal{I} to be a category. However, such a situation cannot arise. The reasons are as follows. Suppose that $\text{Mor}_{\mathcal{I}}(i_X, i_Z)$ becomes \emptyset by the condition $c_X > c_Z$. For the case $c_Y \geq c_X$, $c_Y > c_Z$ also holds, then $\text{Mor}_{\mathcal{I}}(i_Y, i_Z) = \emptyset$. For the case $c_X > c_Y$, $\text{Mor}_{\mathcal{I}}(i_X, i_Y) = \emptyset$. Thus, it is shown that the consistency for compositions of morphisms is guaranteed. \square

We introduce a total order to $(A \downarrow I_{QP})$, as follows.

Definition 5.12 (total order of $(A \downarrow I_{QP})$). For $(q_i, T_i), (q_j, T_j) \in \text{ob}(A \downarrow I_{QP})$, we define $(q_i, T_i) \leq (q_j, T_j)$ by $|\hbar(q_i)|^2 \leq |\hbar(q_j)|^2$.

By this total order, an index category $\mathcal{I}_\hbar(A) := J^\bullet(A \downarrow I_{QP})$ is determined from $(A \downarrow I_{QP})$.

$$\left(\begin{array}{c} A \\ \downarrow q_1 \\ T_1 \end{array} \begin{array}{c} \searrow q_2 \\ \searrow q_3 \dots \\ \longrightarrow T_2 \end{array} \begin{array}{c} \searrow q_3 \dots \\ \longrightarrow T_3 \\ \longleftarrow T_2 \\ \longleftarrow \dots \end{array} \right) \rightarrow \left(i_1 \xrightarrow{f_{12}} i_2 \xrightarrow{f_{23}} i_3 \longrightarrow \dots \right) = \mathcal{I}_\hbar(A).$$

Here $J^\bullet(q_k, T_k) = i_k$ and $J^\bullet(t_{kl}) = f_{kl}$ when $|\hbar(q_k)|^2 \leq |\hbar(q_l)|^2$. Next, the diagram $D_{\hbar} : \mathcal{I}_{\hbar}(A) \rightarrow (A \downarrow I_{QP})$ is defined by $D_{\hbar}(i_k) = (q_k, T_k)$ and $D_{\hbar}(f_{kl}) = t_{kl}$.

$$\left(i_1 \xrightarrow{f_{12}} i_2 \xrightarrow{f_{23}} i_3 \longrightarrow \dots \right) \xrightarrow{D_{\hbar}} \left((q_1, T_1) \xrightarrow{t_{12}} (q_2, T_2) \xrightarrow{t_{23}} (q_3, T_3) \longrightarrow \dots \right).$$

Notice that it is possible to make a category by putting a natural initial object (Id_A, A) into this $D_{\hbar}(\mathcal{I}_{\hbar}(A))$.

Definition 5.13. A/QW_{\hbar} is a category defined as follows. Its objects are given as

$$ob(A/QW_{\hbar}) := ob(D_{\hbar}(\mathcal{I}_{\hbar}(A))) \cup \{(Id_A, A)\}.$$

Its morphisms are given as

$$\begin{aligned} Mor_{A/QW_{\hbar}}((q_i, T_i), (q_j, T_j)) &= Mor_{D_{\hbar}(\mathcal{I}_{\hbar}(A))}((q_i, T_i), (q_j, T_j)) \\ &\quad \text{for } \forall (q_i, T_i), (q_j, T_j) \in ob(D_{\hbar}(\mathcal{I}_{\hbar}(A))), \\ Mor_{A/QW_{\hbar}}((Id_A, A), (q_i, T_i)) &= \{q_i\} \quad \text{for } \forall (q_i, T_i) \in ob(D_{\hbar}(\mathcal{I}_{\hbar}(A))), \end{aligned}$$

and identity morphisms.

$$\begin{array}{ccccc} (Id_A, A) & & & & \\ \downarrow q_1 & \searrow q_2 & \searrow q_3 \dots & & \\ (q_1, T_1) & \xrightarrow{t_{12}} & (q_2, T_2) & \xrightarrow{t_{23}} & (q_3, T_3) \longrightarrow \dots \end{array}$$

Note that the index $\mathcal{I}_{\hbar}(A)$ and the diagram D_{\hbar} are still valid in A/QW_{\hbar} .

Proposition 5.14. A/QW_{\hbar} has a limit $((Id_A, A), \mathbf{q})$ for the index category $\mathcal{I}_{\hbar}(A)$ with the diagram D_{\hbar} . Here, $((Id_A, A), \mathbf{q})$ is a cone of $D_{\hbar}(\mathcal{I}_{\hbar}(A))$ with $\mathbf{q} = \{q_i \in Q \mid s(q_i) = A\}$.

Proof. (Id_A, A) is the initial object in A/QW_{\hbar} . By definition, $((Id_A, A), \mathbf{q})$ is a cone of $D_{\hbar}(\mathcal{I}_{\hbar}(A))$. Because every morphism $t_{ij} : (q_i, T_i) \rightarrow (q_j, T_j)$ satisfies commutativity $q_j = t_{ij} \circ q_i$ by the definition of $(A \downarrow I_{QP})$. In A/QW_{\hbar} , there is at most one morphism between any two objects. So, $((Id_A, A), \mathbf{q})$ is one of the candidates of the limit. Next, we prove that there are no other candidates for the limit other than $((Id_A, A), \mathbf{q})$, by contradiction. If the other candidate of the limit of the diagram D_{\hbar} exists, it is given by a cone with the form $((q_k, T_k), \mathbf{t}_{k-})$, where $q_k : A \rightarrow T_k$ is in Q and $\mathbf{t}_{k-} = \{t_{kl} : q_k \rightarrow q_l = t_{kl} \circ q_k \mid t_{kl} \in Mor_{QP}(T_k, T_l)\}$. From Proposition 2.14, there exist $q_x : A \rightarrow T_x \in Q$ such that $|q_x|^2 < |q_k|^2$. (q_x, T_x) is an object in $D_{\hbar}(\mathcal{I}_{\hbar}(A))$, and $(q_k, T_k) > (q_x, T_x)$. Then $Mor_{A/QW_{\hbar}}((q_k, T_k), (q_x, T_x)) = \emptyset$, and this contradicts $((q_k, T_k), \mathbf{t}_{k-})$ is a cone. Therefore, we find that $((Id_A, A), \mathbf{q})$ is the limit. \square

From this proposition, we found that for any Poisson algebra A in QW the limit for the index category $\mathcal{I}_{\hbar}(A)$ with the diagram D_{\hbar} is determined without additional information. We call this limit $((Id_A, A), \mathbf{q})$ ‘‘strong classical limit of A ’’.

This Proposition 5.14 shows that the strong classical limit satisfies at least the properties that the limit should have that we wanted in this subsection.

5.7 Example of Strong Classical Limit

Let us consider \mathcal{A} we studied in the fuzzy sphere in Appendix C. The index $\mathcal{I}_\hbar(\mathcal{A})$, the diagram $D_\hbar(\mathcal{I}_\hbar(\mathcal{A}))$ and \mathcal{A}/QW_\hbar are determined without extra information. We obtain the following fact.

Example 5.15. *Let \mathcal{A} be a Poisson algebra studied in Appendix C. The strong classical limit of \mathcal{A} is $((Id_{\mathcal{A}}, \mathcal{A}), \mathbf{q})$, where $\mathbf{q} = \{q_i \in Q \mid s(q_i) = \mathcal{A}\}$.*

From Proposition 5.14 there is no need to prove this Example 5.15. However, this claim can be shown just from the properties of some quantization maps of matrix regularization. This is so suggestive that we will also provide the proof below.

Proof. As we saw in the proof of Proposition 5.14, $((Id_{\mathcal{A}}, \mathcal{A}), \mathbf{q})$ is a cone for the diagram D_\hbar in \mathcal{A}/QW_\hbar . We show that there are no other cones.

Suppose that there is another cone. Then, the cone is given as some $((q_{cone}, T_{cone}), \mathbf{t}_{cone,-})$ where $q_{cone} \in Q$ with $s(q_{cone}) = \mathcal{A}$ and $\mathbf{t}_{cone,-} = \{t_{cone,k} : (q_{cone}, T_{cone}) \rightarrow (q_k, T_k) \mid q_k \in Q, s(q_k) = \mathcal{A}, q_k = t_{cone,k} \circ q_{cone}\}$. Consider $t_{cone,2} \in Mor((q_{cone}, T_{cone}), (t_2, T_2))$, where $t_2 \in Q$ is defined in Appendix C. Since $t_{cone,2}$ is an injection and T_2 is a 4-dimensional vector space, $\dim T_{cone} \leq 4$ is derived. For $t_3 : \mathcal{A} \rightarrow T_3$ defined in Appendix C, $\dim \text{Im } t_3 > 4$. On the other hand, $\dim \text{Im } t_{cone,3} \leq 4$ for $t_{cone,3} : T_{cone} \rightarrow T_3$. Therefore $t_3 \neq t_{cone,3} \circ q_{cone}$. This is contradiction to the assumption that $((q_{cone}, T_{cone}), \mathbf{t}_{cone,-})$ is a cone.

Thus, it is shown that $((Id_{\mathcal{A}}, \mathcal{A}), \mathbf{q})$ is the only cone and the limit for the diagram D_\hbar in \mathcal{A}/QW_\hbar . \square

6 One Attempt to Address the Inverse Problem

As mentioned at the beginning of this paper, the inverse problems of quantizations are important for physics related to noncommutative geometries. The inverse problem of quantizations is the problem of how to determine the classical limits (classical manifolds) based on the information of the spaces appearing in the quantizations. In general, classical limits are not uniquely determined from algebras as target spaces of quantization maps in a naive way, so we need to experiment with various subtle techniques. This problem is closely related to the question of how we can identify manifolds in the membrane theory, thus, it was often investigated in that context. For example, in [12, 15, 16] it is discussed how to construct the classical limit in geometric quantization. In [41, 11, 38], how membrane topology is distinguished in the context of matrix regularization. In [23, 24, 4], inverse problems of Berezin-Toeplitz quantizations are discussed.

Unlike in the past, the purpose of this section is to study this inverse problem as a problem of the classical limit discussed in the previous section. Until now, we have studied whole world of the quantizations of all Poisson algebras. In particular, the naive classical limit and the weak classical limits introduced in Subsection 5.1 and Subsection 5.4 are chosen from among all Poisson algebras in QW . To determine the naive classical limit or weak classical limit from the spaces appearing in the quantizations, we are going to examine how to construct a sequence of objects in QW and are going to discuss how to give the sequence by a process found in physics.

6.1 Matrix Regularization and Inverse Problem

The goal of this section is making an example of a method to obtain a Poisson algebra as a classical limit from a noncommutative associative algebra. Here we make this example using a fuzzy sphere as a role model.

At first, let \mathfrak{g} be a noncommutative algebra over \mathbb{C} regarded as a Lie algebra whose commutator is defined by its associative product i.e. $[a, b] := ab - ba$. Only semisimple Lie algebras are treated in this section. This \mathfrak{g} is the origin to get the sequence of objects in QW that may provide a classical limit. Such sequence of objects in QW is named the quantization family for the weak classical limit in Subsection 5.4. Let $e = \{e_1, e_2, \dots, e_d\}$ be a fixed base of \mathfrak{g} satisfying commutation relations $[e_i, e_j] = f_{ij}^k e_k$, where f_{ij}^k are structure constants of \mathfrak{g} . For this Lie algebra \mathfrak{g} we introduce a sequence of representations $\rho^\mu : \mathfrak{g} \rightarrow gl(V_\mu)$ and a sequence of numbers $\hbar(\mu) \neq 0$, ($\mu = 1, 2, 3, \dots$). Here V_μ is a finite dimensional vector space chosen as appropriate. We denote the corresponding basis of e by

$$e^{(\mu)} = \{\hbar(\mu)\rho^\mu(e_1), \hbar(\mu)\rho^\mu(e_2), \dots, \hbar(\mu)\rho^\mu(e_d)\} = \{e_1^{(\mu)}, e_2^{(\mu)}, \dots, e_d^{(\mu)}\}. \quad (6.1)$$

Then they satisfy

$$[e_i^{(\mu)}, e_j^{(\mu)}] = \hbar(\mu)f_{ij}^k e_k^{(\mu)}. \quad (6.2)$$

The Lie algebra $\rho^\mu(\mathfrak{g})$ or $\langle e^{(\mu)} \rangle$ are constructed by this basis. For each μ , if we add N_μ relations $f_i^\mu(e^{(\mu)}, \nu^{(\mu)}, \hbar(\mu))$, ($i = 1, 2, \dots, N_\mu$) by Casimir invariants, where $\nu^{(\mu)} = \{\nu_1^{(\mu)}, \nu_2^{(\mu)}, \dots\}$ is a set of parameters, and $N_\mu \leq \text{rank } \mathfrak{g}$, we can consider an algebra $\rho^\mu(\mathfrak{g})/\mathcal{I}_\mu$. Here \mathcal{I}_μ is an ideal defined by these relations

$$\mathcal{I}_\mu = \langle f_1^\mu(e^{(\mu)}, \nu^{(\mu)}, \hbar(\mu)), \dots, f_{N_\mu}^\mu(e^{(\mu)}, \nu^{(\mu)}, \hbar(\mu)) \rangle. \quad (6.3)$$

Next, let T_μ be the vector space that is $\langle e^{(\mu)} \rangle$ forgetting multiplication structure. We choose a basis of $T_\mu := \langle e^{(\mu)} \rangle$, E_1, E_2, \dots, E_D , as polynomials of $e^{(\mu)}$. The highest degree is denoted by n_μ , i.e. $n_\mu = \max\{\deg E_1, \dots, \deg E_D\}$. For later use, let us consider symmetrized polynomial

$$c_{i_1, \dots, i_k} e_{i_1}^{(\mu)} \cdots e_{i_k}^{(\mu)} = c_{i_1, \dots, i_k} e_{(i_1, \dots, i_k)}^{(\mu)} \quad (k = 0, 1, 2, \dots, n_\mu),$$

where the fixed coefficients $c_{i_1, \dots, i_k} \in \mathbb{C}$ are complete symmetric, and

$$e_{(i_1, \dots, i_k)}^{(\mu)} := \frac{1}{k!} \sum_{\sigma \in \text{Sym}(k)} e_{i_{\sigma(1)}}^{(\mu)} \cdots e_{i_{\sigma(k)}}^{(\mu)}.$$

Here $\text{Sym}(k)$ is a symmetric group. Obviously, each $e_{(i_1, \dots, i_k)}^{(\mu)}$ is given as a linear combination of E_1, E_2, \dots, E_D and its degree is smaller than or equal to n_μ . For $m > n_\mu$ $e_{(i_1, \dots, i_m)}^{(\mu)}$ is written by some linear combination of lower degree polynomials, then $\hbar(\mu)$ appear because of the definition (6.1). From the set of representation ρ^μ (and relations f_i^μ), we obtain sets of T_μ . We denote a set of all T_μ by $\{T_\mu\}$. To show that $\{T_\mu\}$ is a sequence of objects in QW , we have to find one Poisson algebra and to make quantization maps from the Poisson algebra to $\{T_\mu\}$.

The next step we introduce a way to obtain one Poisson algebra from a Lie algebra \mathfrak{g} as a candidate of a weak classical limit. There is a well-known way known as Kirillov-Kostant Poisson bracket, that is the way constructing a Poisson algebra. (See also [29, 40, 1].) We focus the following fact (See [21, 25]), here.

Theorem 6.1. *Let \mathfrak{g} be a d -dimensional Lie algebra. Let $e = \{e_1, e_2, \dots, e_d\}$ be a basis of \mathfrak{g} satisfying commutation relations $[e_i, e_j] = f_{ij}^k e_k$. Let $x = (x^1, x^2, \dots, x^d)$ be commutative variables. We obtain a Poisson algebra $(\mathbb{C}[x], \cdot, \{, \})$ by*

$$\{f, g\} := f\omega g := f \overleftarrow{\partial}_i \omega_{ij} \overrightarrow{\partial}_j g := (\partial_i f)\omega_{ij}(\partial_j g), \quad (6.4)$$

where $\partial_i = \frac{\partial}{\partial x^i}$ and $\omega_{ij} = f_{ij}^k x^k$.

The proof is given in [21] but for readers convenience we give the proof here.

Proof. $\{f, g\} = -\{g, f\}$ is followed from $f_{ij}^k = -f_{ji}^k$. The Leibniz's rule and the bilinearity are trivially satisfied by the definition. By direct calculations, the Jacobi identity is obtained as follows.

$$\begin{aligned} & \{\{f_1, f_2\}, f_3\} + \{\{f_2, f_3\}, f_1\} + \{\{f_3, f_1\}, f_2\} \\ &= (\partial_i f_1)(\partial_j f_2)(\partial_l f_3)\{(\partial_k \omega_{ij})\omega_{kl} + (\partial_k \omega_{jl})\omega_{ki} + (\partial_k \omega_{li})\omega_{kj}\}. \end{aligned}$$

By $\omega_{ij} = f_{ij}^k x^k$,

$$(\partial_k \omega_{ij})\omega_{kl} + (\partial_k \omega_{jl})\omega_{ki} + (\partial_k \omega_{li})\omega_{kj} = x^m (f_{ij}^k f_{kl}^m + f_{jl}^k f_{ki}^m + f_{li}^k f_{kj}^m) = 0$$

The last equality follows from the Jacobi identity of the Lie bracket $[e_i, e_j] = f_{ij}^k e_k$. \square

We denote this $(\mathbb{C}[x], \cdot, \{, \})$ by $A_{\mathfrak{g}}$. From the above discussions, we found that we can obtain a sequence of Lie algebras and a Poisson algebra from a fixed single Lie algebra.

Next, let us construct quantization maps from the Poisson algebra $A_{\mathfrak{g}}$ we have just obtained in Theorem 6.1 to the vector spaces in $\{T_{\mu}\}$ that generate original algebras. We define linear function $q_{\mu} : A_{\mathfrak{g}} \rightarrow T_{\mu}$ by

$$\sum_k f_{i_1, \dots, i_k} x^{i_1} \cdots x^{i_k} \mapsto \sum_k f_{i_1, \dots, i_k} e_{i_1}^{(\mu)} \cdots e_{i_k}^{(\mu)} = \sum_k f_{i_1, \dots, i_k} e_{(i_1, \dots, i_k)}^{(\mu)}, \quad (6.5)$$

where $f_{i_1, \dots, i_k} \in \mathbb{C}$ is completely symmetric. Then this correspondence q_{μ} satisfies the following.

Theorem 6.2. *Let $q_{\mu} : A_{\mathfrak{g}} \rightarrow T_{\mu} = \langle e^{(\mu)} \rangle$ be a linear function defined as (6.5). Then it satisfies*

$$[q_{\mu}(f), q_{\mu}(g)] = \hbar(\mu)q_{\mu}(\{f, g\}) + \tilde{O}(\hbar(\mu))$$

for $f, g \in A_{\mathfrak{g}}$. In other words, $q_{\mu} \in Q$.

Proof. We put $f = \sum_k f_{i_1, \dots, i_k} x^{i_1} \dots x^{i_k}$ and $g = \sum_l g_{j_1, \dots, j_l} x^{j_1} \dots x^{j_l}$.

$$\begin{aligned} [q_\mu(f), q_\mu(g)] &= \sum_{k,l}^{n_\mu} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} [e_{i_1}^{(\mu)} \dots e_{i_k}^{(\mu)}, e_{j_1}^{(\mu)} \dots e_{j_l}^{(\mu)}] \\ &= \sum_{k,l=1}^{n_\mu} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{n,m} e_{i_1}^{(\mu)} \dots [e_{i_n}^{(\mu)}, e_{j_m}^{(\mu)}] \dots e_{j_l}^{(\mu)}. \end{aligned}$$

Here using $[e_{i_n}^{(\mu)}, e_{j_m}^{(\mu)}] = \hbar(\mu) f_{i_n j_m}^k e_k^{(\mu)}$,

$$\begin{aligned} [q_\mu(f), q_\mu(g)] &= \hbar(\mu) \sum_{k,l=1}^{n_\mu} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{n,m} f_{i_n j_m}^p e_{i_1}^{(\mu)} \dots e_p^{(\mu)} \dots e_{j_l}^{(\mu)} \\ &= \hbar(\mu) \sum_{k,l=1}^{n_\mu} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{\substack{1 \leq n \leq k \\ 1 \leq m \leq l}} f_{i_n j_m}^p e_{(i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p)}^{(\mu)} + \tilde{O}(\hbar^{1+\epsilon}(\mu)). \end{aligned} \tag{6.6}$$

Here the index $\{i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p\}$ means $\{i_1, \dots, i_k, j_1, \dots, j_l, p\} - \{i_n, j_m\}$, so $e_{(i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p)}^{(\mu)}$ is a degree $k + l - 1$ polynomial. $\tilde{O}(\hbar^{1+\epsilon}(\mu))$ appeared when $e_{i_1}^{(\mu)} \dots e_p^{(\mu)} \dots e_{j_l}^{(\mu)}$ were sorted in the symmetric order. On the other hand,

$$\{f, g\} = \sum_{k,l=1} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{\substack{1 \leq n \leq k \\ 1 \leq m \leq l}} f_{i_n j_m}^p x^{i_1} \dots x^p \dots x^{j_l}.$$

Note that the degree of $x^{i_1} \dots x^p \dots x^{j_l}$ is $k + l - 1$. Then

$$q_\mu(\{f, g\}) = \sum_{\substack{1 \leq k,l \\ k+l \leq n_\mu+1}} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{\substack{1 \leq n \leq k \\ 1 \leq m \leq l}} f_{i_n j_m}^p e_{(i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p)}^{(\mu)}. \tag{6.7}$$

Let us subtract \hbar times (6.7) from (6.6):

$$\begin{aligned} [q_\mu(f), q_\mu(g)] - \hbar(\mu) q_\mu(\{f, g\}) &= \hbar(\mu) \sum_{\substack{2 \leq k,l \leq n_\mu \\ n_\mu+2 \leq k+l \leq 2n_\mu}} f_{i_1, \dots, i_k} g_{j_1, \dots, j_l} \sum_{\substack{1 \leq n \leq k \\ 1 \leq m \leq l}} f_{i_n j_m}^p e_{(i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p)}^{(\mu)} + \tilde{O}(\hbar^{1+\epsilon}(\mu)). \end{aligned}$$

Note that the degree of $e_{(i_1, \dots, \hat{i}_n, \dots, \hat{j}_m, \dots, j_l, p)}^{(\mu)}$ in the right hand side is bigger than or equal to $n_\mu + 1$. Recall that n_μ is determined as the highest degree of polynomials of $E_i (i = 1, \dots, D)$ that constitute basis of T_μ . So, any degree m polynomial of $e^{(\mu)}$ with $m > n_\mu$ is represented by some linear combination of polynomials whose degree is smaller than or equal to n_μ . From the definition (6.1), such linear combinations are $\tilde{O}(\hbar(\mu))$. Then we obtain

$$[q_\mu(f), q_\mu(g)] = \hbar(\mu) q_\mu(\{f, g\}) + \tilde{O}(\hbar(\mu)).$$

□

Thus, a sequence of quantization maps $\{q_\mu\} \subset Q$ is obtained, and we find $\{T_\mu\}$ is a sequence of objects in QW .

Other candidates for Poisson algebras exist besides $A_{\mathfrak{g}}$. Using I as an ideal of $A_{\mathfrak{g}}$ generated by relations that is invariants under acting Poisson brackets, we can introduce $A_{\mathfrak{g}}/I$. Even in this case, we expect to be able to construct $\{T_\mu\}$ and $\{q_\mu\}$ in the same way as in the $A_{\mathfrak{g}}$ case. In fact, q_μ is realized as t_k for a fuzzy sphere case in Appendix C.

Example 6.3. Let $e = \{e_1, e_2, e_3\}$ be a fixed base of $\mathfrak{su}(2)$ satisfying commutation relations $[e_i, e_j] = i\epsilon_{ijk}e_k$. Consider a sequence of spin j representation $\rho^\mu : \mathfrak{g} \rightarrow gl(V_\mu)$ ($\dim V_\mu = 2j + 1$). Then $e^{(\mu)}$ satisfies $[e_i^{(\mu)}, e_j^{(\mu)}] = i\hbar(\mu)\epsilon_{ijk}e_k^{(\mu)}$, and a Casimir relation

$$f^\mu(e^{(\mu)}, r, \hbar(\mu)) = (e_1^{(\mu)})^2 + (e_2^{(\mu)})^2 + (e_3^{(\mu)})^2 - r^2 Id$$

is imposed for each ρ^μ . Here $r/\hbar(\mu) = \sqrt{j(j+1)}$. From this sequence we obtain the sequence of T_k in Subsection 5.3 or in Subsection 5.5. The corresponding Poisson algebra of $\mathfrak{su}(2)$ determined by Theorem 6.1 is differ from \mathcal{A} because of the existence of the relation. However, quantization maps t_k in Subsection 5.3 or Subsection 5.5 is constructed in the same manner with q_μ in Theorem 6.2. Furthermore, as we saw in Subsection 5.3, this sequence gives the naive classical limit \mathcal{A} , when its diagram includes \mathcal{A} . And, in Subsection 5.5, it is shown that \mathcal{A} is isomorphic to the weak classical limit if there exists a weak classical limit with the sequence $\{Im(t_k)\}$ as the diagram.

As above, starting from a Lie algebra we obtain a sequence of corresponding objects $\{T_\mu\}$ in QW and some corresponding Poisson algebras. We already know that the diagram $\{T_\mu\} \cup A_{\mathfrak{g}}$ gives at least a naive classical limit from Theorem 5.2. This indicates that the above procedure may provide one new approach to the inverse problem.

Is it possible to derive a quantization family, a sequence like $\{T_\mu\}$ that gives a naive classical limit or a weak classical limit, from the principle of least action in terms of physics? Here we mention only one attempt to give such a quantization family. Let us consider a sequence of matrix models. Using $N \times N$ matrixes X_μ^N ($\mu = 1, \dots, D$, $N \in \mathbb{N}$) and a mass $\hbar(N)$, we consider the action

$$S_N(\hbar^2(N)) = \text{tr} \left(\frac{1}{4} [X_\mu^N, X_\nu^N]^2 + \hbar^2(N) \frac{1}{2} X^{N\mu} X_\mu^N \right) + \sum_i^k \langle \lambda_i^N, f_i^{(N)}(X^N, \nu) \rangle, \quad (6.8)$$

where we take contraction by the Killing metric with the structure constants $f_{\mu\rho}^\tau$ of \mathfrak{g} , i.e. $g_{\mu\nu} = f_{\mu\rho}^\tau f_{\nu\tau}^\rho$, and $\langle -, - \rangle$ represents the inner product with respect to this Killing metric. $f_i^{(N)}(X^N, \nu)$ ($i = 1, \dots, k \leq \text{rank } \mathfrak{g}$) are relations given by Casimir invariants like $f_i^\mu(e^{(\mu)}, \nu^{(\mu)}, \hbar(\mu))$ in (6.3) with some parameters ν . For simplicity, we assume the relations $\{f_i^{(N)}\}$ and parameters ν do not depend on N i.e. $f_i^{(N)}(x, \nu) = f_i^{(J)}(x, \nu)$ for all $N, J \in \mathbb{N}$ and $x \in \mathbb{C}^D$. $\sum_i \langle \lambda_i^N, f_i^{(N)}(X^N, \nu) \rangle$ is a gauge fixing term and λ_i^N is a Lagrange multiplier. The equation of motion with each λ_i^N is $f_i^{(N)}(X^N, \nu) = 0$.

Then the equation of motion obtained by the variation of X_ν^N with $\lambda_i^N = 0$ is given by

$$[X^{N\mu}, [X_\mu^N, X_\nu^N]] = \hbar^2(N)X_\nu^N.$$

If X_μ^N is a base of a representation of \mathfrak{g} such that $[X_\mu^N, X_\nu^N] = \hbar(N)f_{\mu\nu}^\tau X_\tau^N$, the left hand side of the equation of motion is calculated as

$$\begin{aligned} [X^{N\mu}, [X_\mu^N, X_\nu^N]] &= \hbar(N)f_{\mu\nu}^\rho g^{\mu\tau} [X_\tau^N, X_\rho^N] = \hbar^2(N)f_{\mu\nu}^\rho g^{\mu\tau} f_{\tau\rho}^\sigma X_\sigma^N \\ &= \hbar^2(N)\delta_\nu^\sigma X_\sigma^N = \hbar^2(N)X_\nu^N. \end{aligned}$$

Therefore, we found that such set of generators $\{X_\mu^N\}$ which consist a representation of \mathfrak{g} is a solution of the equation of motion of the action S_N . We denote these generators by $e^{(N)}$. A vector space V_N spanned by the solutions $e^{(N)}$ is determined. A subset of natural number $SN \subset \mathbb{N}$ determine a sequence of actions $\{S_N\} := \{S_N \mid N \in SN \subset \mathbb{N}\}$. Then we obtain sequence of V_N from this $\{S_N\}$ similarly. The sequence $\{V_N\}$ is a sequence of objects QW .

The way to get one Poisson algebra $A_{\mathfrak{g}}$ from generators $e^{(N)}$ is given in Theorem 6.1. For the case with relations $\{f_i^{(N)}(X^N, \nu^{(N)})\}$, a Poisson algebra with relations $\{f_i^{(N)}(x, \nu^{(N)})\}$ is also expected. (Recall that $f_i^{(N)}$ does not depend on N .) We assume the existence of the Poisson algebra and denote the Poisson algebra by $A_{\mathfrak{g}}/I$, where $I = \langle f_1^{(N)}(x, \nu), \dots, f_k^{(N)}(x, \nu) \rangle$. To obtain the classical limit, we need a sequence of quantization maps from the Poisson algebra to objects. Let us find the way to obtain the set of these quantization maps. From Theorem 6.2 quantization $q_N \in Q$ from $A_{\mathfrak{g}}/I$ to $\langle V_N \rangle$ is defined similar to (6.5). Using $\{q_N \mid N \in SN\}$, we define a new quantization sequence $\{q'_N\} = \{q'_N = q_N, q'_J = q_J, q'_K = q_K, \dots\} \subset Q$ as appropriate such that $\ker q'_N \supseteq \ker q'_J$ if $N < J$. (The similar condition appears in Fuzzy sphere case.)

Finally, using the objects $A_{\mathfrak{g}}/I$ and $q'_N(A_{\mathfrak{g}}/I)$ and the morphisms $\{q'_N\}$ in QW , as in Section 5.3, we can construct the index category and its diagram. We denote the diagram by $D_{\mathfrak{g}}$. The constructing process is parallel to the one in Section 5.3. Then, the diagram $D_{\mathfrak{g}}$ is obtained by (6.9) and the naive classical limit is given by $A_{\mathfrak{g}}/I$.

$$\left(\begin{array}{c} 1 \\ \downarrow \hat{N} \\ N \end{array} \begin{array}{c} \nearrow J \\ \searrow \hat{K} \end{array} \begin{array}{c} \dots \\ \nearrow J \\ \searrow \hat{K} \end{array} \dots \right) \xrightarrow{D_{\mathfrak{g}}} \left(\begin{array}{c} A_{\mathfrak{g}}/I \\ \downarrow q'_N \\ q'_N(A_{\mathfrak{g}}/I) \end{array} \begin{array}{c} \nearrow q'_J \\ \searrow q'_K \end{array} \begin{array}{c} \dots \\ \nearrow q'_J \\ \searrow q'_K \end{array} \dots \right). \quad (6.9)$$

Furthermore, as in the fuzzy sphere example in Subsection 5.4, if a weak limit exists, then $A_{\mathfrak{g}}/I$ is expected to be the weak limit.

Let us summarize the process to obtain the classical limit.

1. Prepare the sequence of actions $\{S_N\}$.
2. A sequence of the solution $e^{(N)} = \{e_1^{(N)}, \dots, e_d^{(N)}\}$ and the sequence of vector spaces $\{V_N\}$, where $e^{(N)}$ is a basis of V_N are obtained.

3. When $A_{\mathfrak{g}}/I$ is given as a Poisson algebra from $\{e^{(N)}\}$, a sequence of quantization $\{q'_N\} \subset Q$, and $q'_N(A_{\mathfrak{g}}/I)$ are constructed.
4. Objects $A_{\mathfrak{g}}/I$, $q'_N(A_{\mathfrak{g}}/I)$ and morphisms q'_N induce an index category and its diagram $D_{\mathfrak{g}}$.
5. The naive classical limit is determined as $A_{\mathfrak{g}}/I$. It is also expected to be in the weak classical limit.

It should be noted that, in contrast to conventional classical mechanics, the classical solution gives a basis for the vector space, and the set of basis (interpreted as the set of vector spaces) derives the sets of objects and morphisms in QW that may determine the classical limit.

The process discussed here is an example of approaching an inverse problem using the quantization world QW through a physics method. Since the method is based on a generalization of the fuzzy sphere, it works well for the case of fuzzy spheres, but it is not known at this time whether it will work for other manifolds. It is another future task to investigate this point.

7 Summary

We constructed a category QW that is composed of the all quantizations of all Poisson algebras. The characteristic of this category is that quantization is treated as a linear map not to an algebra, but to a module, which is a subset of some algebra. We defined what is required in this formulation and saw next that the definition works well. First, equivalence of quantization was defined. And it was shown that iff there is a pair of equivalent quantizations by the definition then there exists a pair of isomorphisms of Poisson algebras and the algebras generated by the images of the quantization maps. We also considered the category of quantizations for a fixed single Poisson algebra and discussed its classification of quantizations. Next, we defined the classical limits. Three types of classical limits were introduced: naive classical limits, weak classical limits and strong classical limits. The naive classical limit was defined in the context of category theory. It is a limit in category QW whose the vertex of cone is in Poisson algebras. As concrete examples, we introduced the naive classical limits for deformation quantizations of polynomials with the Moyal products and for matrix regularization of fuzzy spheres. The weak classical limit was defined by modifying the definition of the limit in the category theory. The diagrams for the limit were restricted them in $QP \subset QW$, and the cones to define the limit were restricted to only those with vertices in Poisson algebras. Like the limit of ordinary number sequences, the naive classical limit and the weak classical limit may not have a limit, depending on how the sequence of objects is chosen. On the other hand, the strong classical limit was defined for quantizations when a Poisson algebra is fixed. In contrast to the other classical limit, there is no flexibility in the choice of the classical limit, which is automatically determined when the Poisson algebra is chosen. Also, we found the fixed Poisson algebra is always the strong classical limit. Finally, we discussed the inverse problem of determining the classical limit from some noncommutative Lie algebra. From a fixed Lie algebra, we constructed a sequence of representations of the Lie algebra with relations, from which

we constructed a Poisson algebra. Next, by constructing a sequence of quantization maps from the Poisson algebra, we obtain the sequence of representations of the Lie algebra as a sequence of QW objects. A method to obtain this series of procedures from the principle of least action was proposed. The proposed method is just an example of how to approach to the inverse problem in the framework of QW , which is a generalization of the fuzzy sphere case. A more various discussions on how generalizations should be made are needed in the future. The challenge of obtaining classical manifolds from solutions of noncommutative manifolds of matrix models that actually describe string theory or M-theory is a future problem, too.

As for QW , it is important to investigate its properties from a purely mathematical point of view as well, since it is a noncommutative geometric object that naturally follows from the overall formulation of the quantization of all Poisson algebras. There are many interesting problems around QW to understand the whole picture of noncommutative geometry. These are issues to be clarified in the future.

Acknowledgements

A.S. was supported by JSPS KAKENHI Grant Number 21K03258. The author is grateful to A. Yoshioka for his detailed discussions and useful comments throughout the entire article. The author also thanks the participants in the workshop “Discrete Approaches to the Dynamics of Fields and Space-Time” for their useful comments.

A Definition of $\tilde{O}(z^{1+\epsilon})$

Since we have not defined a norm for objects of category QW in this paper, Landau symbol O does not make sense. So, we define an order \tilde{O} by $x \in \mathbb{R}$ with the Euclidean norm.

Definition A.1. Let \mathcal{M} be an R -module for a commutative algebra R over \mathbb{C} . Let f_i be a complex valued continuous function such that

$$\lim_{x \rightarrow 0} \frac{f_i(xz)}{x} = 0,$$

where $x \in \mathbb{R}$ and $z \in \mathbb{C}$. For $a_i \in \mathcal{M}$ which is independent of $z \in \mathbb{C}$, we denote the element described as $\sum_i f_i(z)a_i \in \mathcal{M}$ by $\tilde{O}(z^{1+\epsilon})$.

From this definition, the term of $\tilde{O}(\hbar^{1+\epsilon})$ in (2.3) is also 0 when $\hbar = 0$. Note that \hbar itself is not necessarily continuous.

B List of Symbols

In this paper, many symbols including new symbols are used. So, the list of them is useful to read through.

Table 1: List of Symbols

R	commutative ring over \mathbb{C}
$R\text{-Mod}$	category of R -module
$R\text{-alg}$	category of R -algebra
$\mathcal{P}oi\text{ss}$	category of Poisson algebras whose morphisms are surjective Poisson morphism
wkQ	set of weak quantization maps (including maps with $\hbar = 0$)
Qh_x	set of weak quantization maps with $ \hbar = x$
Q	set of weak quantization maps with $\hbar \neq 0$
$Q(\mathcal{P}oi\text{ss})$	$\{t(q) \mid q \in Q\}$
QP	subcategory of $R\text{-Mod}$: $ob(QP) := Q(\mathcal{P}oi\text{ss})$, $Mor(t(q_i), t(q_j))$ is a set of R -algebra homomorphisms restricted their domain into $t(q_i)$
$\mathcal{P}oi\text{ss} \sqcup QP$	subcategory of $R\text{-Mod}$: $ob(\mathcal{P}oi\text{ss} \sqcup QP) = ob(\mathcal{P}oi\text{ss}) \sqcup ob(QP)$, $Mor(\mathcal{P}oi\text{ss} \sqcup QP) = Mor(\mathcal{P}oi\text{ss}) \sqcup Mor(QP)$
QW	subcategory of $R\text{-Mod}$: $ob(QW) := ob(\mathcal{P}oi\text{ss}) \cup ob(QP)$, $Mor(QW) := Mor(\mathcal{P}oi\text{ss}) \sqcup Mor(QP) \sqcup Q$
U_P	forgetful functor from $\mathcal{P}oi\text{ss}$ to QW
I_{QP}	identically immersion functor from QP to QW
$(U_P \downarrow I_{QP})$	comma category of U_P and I_{QP}
F	functor from $(U_P \downarrow I_{QP})$ to $\mathcal{P}oi\text{ss} \times R\text{-alg}$
$P - QP$	$P - QP := F(U_P \downarrow I_{QP}) : \mathcal{P}oi\text{ss} - Q(\mathcal{P}oi\text{ss})$ quantization pair category
$(A \downarrow I_{QP})$	category of all quantizations of A
F_A	functor from $(A \downarrow I_{QP})$ to $R\text{-alg}$
$D(\mathcal{I})$	Quantization Family with an index category \mathcal{I}
J^\bullet	map from a category with total order to an index category
$\mathcal{I}_\hbar(A)$	$J^\bullet(A \downarrow I_{QP})$
D_\hbar	diagram $\mathcal{I}_\hbar(A) \rightarrow (A \downarrow I_{QP})$
A/QW_\hbar	category consists of $D_\hbar(\mathcal{I}_\hbar(A))$ and $\{(Id_A, A)\}$ with quantization maps whose source is A

C Fuzzy Sphere and Fuzzy Torus

The fuzzy sphere is considered in [21, 31]. See [21, 31, 2, 7] for details. Let x^a ($1 \leq a \leq 3$) be coordinates of three-dimensional Euclidean space and S^2 be the two-sphere given by

$$\delta_{ab}x^ax^b = 1. \quad (C.1)$$

Let $P[x^a]$ be the algebra of polynomials generated by x^a . For an ideal I which is generated by the relation (C.1), we consider a Poisson algebra $\mathcal{A} = P[x^a]/I$ with Poisson bracket

$$\{x^a, x^b\} = \epsilon^{abc}x^c. \quad (C.2)$$

For arbitrary $f \in \mathcal{A}$ is given as

$$f = f_0 + f_ax^a + \frac{1}{2}f_{ab}x^ax^b + \dots$$

where $f_{a_1 \dots a_i} \in \mathbb{C}$ is completely symmetric and trace-free. The morphism t_1 from \mathcal{A} to $T_1 = \mathbb{C}$ is defined by $t_1(f) := f_0$ and the morphism t_2 from \mathcal{A} to a subspace $T_2 := t_2(\mathcal{A})$ of a matrix algebra $M_2 = \text{Mat}_2$ is defined by

$$t_2(f) := f_0\mathbf{1}_2 + f_at_2(x^a), \quad t_2(x^a) := \frac{\hbar_2}{2}\sigma^a$$

where σ^a is Pauli matrix and $\mathbf{1}_k$ is a $k \times k$ unit matrix. t_2 means that the morphism gives a map from a polynomial to a 2×2 matrix. In the case of $k \geq 2$, morphisms $t_k : \mathcal{A} \rightarrow T_k := t_k(\mathcal{A}) \subset M_k = \text{Mat}_k$ are defined by

$$\begin{aligned} t_k(f) &:= f_0\mathbf{1}_k + f_{a_1}t_k(x^{a_1}) + \dots + f_{a_1 \dots a_{k-1}}t_k(x^{a_1} \dots x^{a_{k-1}}) \\ t_k(x^a) &:= \hbar_k J_k^a = X_k^a \end{aligned}$$

where J_k^a are generators for the k -dimensional irreducible representation of $\mathfrak{su}(2)$, and for $i \geq 2$ each $t_i(x^{a_1} \dots x^{a_{i-1}})$, which is fixed to be well-defined, is generated by J_i^a . The morphism t_k gives a map from a polynomial to a $k \times k$ matrix. From the Casimir relation

$$J_k^a J_{ka} = \frac{1}{4}(k^2 - 1)\mathbf{1}_k$$

and (C.1), the relationship between k and \hbar is given as

$$\frac{4}{k^2 - 1} = \hbar_k^2. \quad (C.3)$$

From the commutation relation of J_k^a

$$[J_k^a, J_k^b] = i\epsilon^{abc}J_k^c, \quad [t_k(x^a), t_k(x^b)] = i\hbar_k^2\epsilon^{abc}J_k^c = i\hbar_k\epsilon^{abc}t_k(x^c). \quad (C.4)$$

Using (C.4) we can show that $t_k \in Q$ ($k \geq 2$). Note that $\langle T_k \rangle = \text{Mat}_k =: M_k$.

Next, we consider a fuzzy torus in a similar way. (See for example [6].) Let (θ_1, θ_2) be a coordinate of $T^2 = S^1 \times S^1$, where $\theta_i \in [0, 2\pi)$. A algebra $\mathcal{B} = \{f(\theta_1, \theta_2) =$

$\sum_{l_1, l_2} f_{l_1, l_2} y_{l_1, l_2}$, where $y_{l_1, l_2} := \exp(\sum_i^2 \sqrt{-1} l_i \theta_i)$ ($l_i \in \mathbb{Z}_{\geq 0}$) are functions on T^2 . Its Poisson bracket is defined by

$$\{y_{l_1, l_2}, y_{m_1, m_2}\} = -\pi(l_1 m_2 - l_2 m_1) y_{l_1 + m_1, l_2 + m_2}.$$

The algebra of Fuzzy torus in $M_k = \text{Mat}_k$ is generated by the clock matrix U and the shift matrix V of size k determined by a complex number $q = e^{2\pi i/k}$, where

$$U = \begin{pmatrix} q^0 & & & \\ & q^1 & & \\ & & \ddots & \\ & & & q^{k-1} \end{pmatrix}, \quad V = \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ 1 & & & 0 \end{pmatrix}.$$

U, V satisfy $U^k = V^k = Id_k$ and $VU = qUV$. By using these U, V , we introduce

$$Y_{m, n} := U^m V^n.$$

Then they satisfy

$$[Y_{l_1, l_2}, Y_{m_1, m_2}] = (q^{l_2 m_1} - q^{l_1 m_2}) Y_{l_1 + m_1, l_2 + m_2}.$$

Note that $q^{l_2 m_1} - q^{l_1 m_2} = -\sqrt{-1} \frac{2\pi}{k} (l_1 m_2 - l_2 m_1) + O(1/k^2)$. The quantization $q_k : \mathcal{B} \rightarrow Y_k := q_k(\mathcal{B}) \subset M_k$ is defined as the same manner with the fuzzy sphere:

$$q_k \left(\sum_{l_1, l_2} f_{l_1, l_2} y_{l_1, l_2} \right) = \sum_{l_1, l_2} f_{l_1, l_2} Y_{l_1, l_2}.$$

Note that $\langle Y_k \rangle = M_k$.

References

- [1] A. Y. Alekseev and A. Z. Malkin. ‘‘Symplectic structures associated to Lie-Poisson groups.’’ *Communications in Mathematical Physics*, **162** (1994) 147-174 [arXiv:hep-th/9303038 [hep-th]].
- [2] J. Arnlind, M. Bordemann, L. Hofer, J. Hoppe. & H. Shimada. ‘‘Fuzzy Riemann surfaces.’’ *Journal of High Energy Physics*, **2009.06** (2009): 047.
- [3] J. Arnlind, J. Hoppe, and G. Huisken. ‘‘Multi-linear Formulation of Differential Geometry and Matrix Regularizations.’’ *Journal of differential geometry*, **91.1** (2012) 1-39.
- [4] T. Asakawa, G. Ishiki, T. Matsumoto, S. Matsuura, and H. Muraki. ‘‘Commutative geometry for non-commutative D-branes by tachyon condensation.’’ *Progress of Theoretical Physics*, (2018): 063B04
- [5] T. Banks, W. Fischler, S. Shenker, L. Susskind. ‘‘M Theory As A Matrix Model: A Conjecture.’’ *Physical Review D*, **55** (1997) 5112-5128
- [6] S. Bal, M. Hanada, H. Kawai and F. Kubo. ‘‘Fuzzy torus in matrix model.’’ *Nuclear Physics B*, **727** (2005) 196-217 [arXiv:hep-th/0412303 [hep-th]].

- [7] A. P. Balachandran, & S. Vaidya. “Lectures on fuzzy and fuzzy SUSY physics.” World Scientific. (2007).
- [8] S. Bates and A. Weinstein. “Lectures on the Geometry of Quantization.” vol. **8**. American Mathematical Society (1997).
- [9] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer. “Deformation theory and quantization. I. Deformations of symplectic structures.” *Annals of Physics*, **111**,(1978) 61-110.
- [10] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer. “Deformation theory and quantization. II. Physical applications.” *Annals of Physics*, **111** (1978) 111-151.
- [11] D. Berenstein and E. Dzienkowski. “Matrix embeddings on flat \mathbb{R}^3 and the geometry of membranes.” *Physical Review D*, **86** (2012): 086001
- [12] M. Bordemann, E. Meinrenken and M. Schlichenmaier. “Toeplitz quantization of Kahler manifolds and $gl(N)$, $N \rightarrow \infty$ limits.” *Communications in Mathematical Physics*, **165.2** (1994) 281-296.
- [13] M. Bertelson, P. Bieliavsky, S. Gutt. “Parametrizing equivalence classes of invariant star products.” *Letter of Mathematical Physics*, **46** (1998) 339345.
- [14] M. Bertelson, M. Cahen, S. Gutt. “Equivalence of star products.” *Classical Quantum Gravity*, **14** (1997) A93-A107
- [15] M. Bordemann, J. Hoppe, P. Schaller and M. Schlichenmaier. “ $gl(\infty)$ and geometric quantization.” *Communications in Mathematical Physics*, **138** no. 2,(1991) 209-244.
- [16] C. S. Chu, J. Madore and H. Steinacker. “Scaling limits of the fuzzy sphere at one loop,” *JHEP* **08** (2001), 038 [arXiv:hep-th/0106205 [hep-th]].
- [17] A. Connes, M.R. Douglas and A. Schwarz. “Noncommutative geometry and matrix theory.” *Journal of High Energy Physics* **1998.02** (1998): 003.
- [18] B. de Wit, J. Hoppe and H. Nicolai. “On the Quantum Mechanics of Supermembranes. *Nuclear Physics B*, **305.4** (1988) 545-581.
- [19] J. Gohara, Y. Hirota and A. Sako. “Categorical Perspective on Quantization of Poisson Algebra.” *Journal of Mathematical Physics* **61** (2020) no.7, 073506 [arXiv:1907.08665 [math-ph]].
- [20] G. Jumpei, H. Yuji and S. Akifumi. “A Generalization of the Quantization of Poisson Manifolds.” *Geometry, Integrability and Quantization*, **21** (2020) 138-148
- [21] J. Hoppe. “Quantum theory of a massless relativistic surface and a two-dimensional bound state problem.” *Soryushiron Kenkyu Electronics*, **80.3** (1989) 145-202.
- [22] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya. “A large- N reduced model as superstring.” *Nuclear Physics B*, **498.1-2** (1997) 467-491.
- [23] G. Ishiki, T. Matsumoto, and H. Muraki, “Kähler structure in the commutative limit of matrix geometry.” *JHEP*, **08** (2016): 042.
- [24] G. Ishiki, T. Matsumoto, and H. Muraki, “Information metric, Berry connection, and Berezin-Toeplitz quantization for matrix geometry.” *Physical Review D*, **98**(2018): 026002

- [25] B. Kostant. “Lie Group Representations on Polynomial Rings.” American Journal of Mathematics. vol. **85** no.3.(1963)324-404.
- [26] N. Landsman. “Strict deformation quantization of a particle in external gravitational and Yang-Mills fields.” Journal of Geometry and Physics, **12.2** (1993) 93-132.
- [27] T. Leinster. Basic category theory. vol. **143**. Cambridge University Press, (2014).
- [28] E. Lerman. “Geometric quantization: a crash course.” Contemporary Mathematics, **583** (2012) 147-174.
- [29] J.H. Lu, A. Weinstein. “Poisson-Lie groups, dressing transformations and Bruhat decompositions.” Journal of Differential Geometry, vol. **31**(1990)501-526.
- [30] S. Mac Lane. “Categories for the working mathematician.” vol. **5**. Springer Science & Business Media, (2013).
- [31] J. Madore. “The fuzzy sphere.” Classical and Quantum Gravity, **9.1** (1992): 69.
- [32] J. Madore. “ An Introduction to Noncommutative Differential Geometry and its Physical Applications.” (London Mathematical Society Lecture Note Series). Cambridge University Press, (1999).
- [33] S. Oh. “Poisson enveloping algebras.” Communications in Algebra, **27.5** (1999) 2181-2186.
- [34] S. Oh. “Hopf structure for Poisson enveloping algebras.” Contributions to Algebra and Geometry, **44.2** (2003) 567-574.
- [35] H. Omori, Y. Maeda , N. Miyazaki, and A. Yoshioka. “Deformation Quantization of Fréchet Poisson Algebras, - Convergence of the Moyal Product -.” In: Conference Moshé Flato 1999, Quantizations, Deformations and Symmetries vol III, Math. Phys. Studies 22, Kluwer, (2000): 233-246.
- [36] M. Rieffel. “Deformation quantization of Heisenberg manifolds.” Communications in mathematical physics, **122.4** (1989) 531-562.
- [37] M. Rieffel. “Quantization and C^* -algebras.” Contemporary Mathematics, **167** (1994) 67-97.
- [38] L. Schneiderbauer and H. Steinacker. “Measuring finite Quantum Geometries via Quasi-Coherent States.” Journal of Physics A, **49** no. 28(2016): 285301 [arXiv:1601.08007 [hep-th]].
- [39] M. Schlichenmaier. “Berezin-Toeplitz quantization and Berezin transform.” Long time behaviour of classical and quantum systems. World Scientific (2001): 271-287.
- [40] M.A. Semenov-Tian-Shansky, Dressing transformations and Poisson-Lie group actions.” Publ. RIMS, Kyoto University. **21**, no.6. (1985)1237-1260.
- [41] H. Shimada. “Membrane topology and matrix regularization.” Nuclear Physics B, 685 (2004) 297320
- [42] H. Steinacker. “Non-commutative geometry and matrix models.” PoS **QGGS2011** (2011), 004 [arXiv:1109.5521 [hep-th]].
- [43] T. Tomishita and A. Yoshioka. “Star Product and Star Exponential.” Geometry Integrability and Quantization, **11** (2011) 171 - 181,

- [44] U. Umirbaev. “Universal enveloping algebras and universal derivations of Poisson algebras.” *Journal of Algebra*, **354.1** (2012) 77-94.
- [45] I. Vaisman. “On the geometric quantization of Poisson manifolds.” *Journal of mathematical physics* **32.12**, (1991) 3339-3345.
- [46] C. Zhu and Y. Wang. “Realization of Poisson enveloping algebra.” *Frontiers of Mathematics in China*, **13.4** (2018): 999-1011.