

# Strongly nonexpansive mappings revisited: uniform monotonicity and operator splitting

Leon Liu\*, Walaa M. Moursi<sup>†</sup> and Jon Vanderwerff<sup>‡</sup>

May 18, 2022

## Abstract

The correspondence between the class of nonexpansive mappings and the class of maximally monotone operators via the reflected resolvents of the latter has played an instrumental role in the convergence analysis of the splitting methods. Indeed, the performance of some of these methods, e.g., Douglas–Rachford and Peaceman–Rachford methods hinges on iterating the so-called splitting operator associated with the individual operators. This splitting operator is a function of the composition of the reflected resolvents of the underlying operators. In this paper, we provide a comprehensive study of the class of uniformly monotone operators and their corresponding reflected resolvents. We show that the latter is closely related to the class of the strongly nonexpansive operators introduced by Bruck and Reich. Connections to duality via inverse operators are systematically studied. We provide applications to Douglas–Rachford and Peaceman–Rachford methods. Examples that illustrate and tighten our results are presented.

**2010 Mathematics Subject Classification:** 49M27, 65K10, 90C25; Secondary 47H14, 49M29.

**Keywords:** contraction mappings, Douglas–Rachford splitting, Peaceman–Rachford splitting, resolvent, reflected resolvent, strongly nonexpansive mapping, uniformly convex function, uniformly monotone operator.

## 1 Introduction

Throughout, we assume that

$$X \text{ is a real Hilbert space,} \tag{1}$$

---

\*Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada. E-mail: 13521iu@uwaterloo.ca.

<sup>†</sup>Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada. E-mail: walaa.moursi@uwaterloo.ca.

<sup>‡</sup>Department of Mathematics, La Sierra University, Riverside, CA 92515, USA. E-mail: jvanderw@lasierra.edu.

with inner product  $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{R}$  and induced norm  $\| \cdot \|$ . Let  $A : X \rightrightarrows X$  be a set-valued operator. The *graph* of  $A$  is  $\text{gra } A = \{(x, x^*) \in X \times X \mid x^* \in Ax\}$ . Recall that  $A$  is *monotone* if  $\{(x, x^*), (y, y^*)\} \subseteq \text{gra } A$  implies that  $\langle x - y, x^* - y^* \rangle \geq 0$ . A monotone operator  $A$  is *maximally monotone* if  $\text{gra } A$  does not admit a proper extension (in terms of set inclusion) to a graph of a monotone operator. The *resolvent* of  $A$  is  $J_A = (\text{Id} + A)^{-1}$  and the *reflected resolvent* of  $A$  is  $R_A = 2J_A - \text{Id}$ , where  $\text{Id} : X \rightarrow X : x \mapsto x$ .

The theory of monotone operators has been of significant interest in optimization: indeed a typical problem in convex optimization seeks finding a minimizer of the sum  $f + g$ , where both  $f$  and  $g$  are proper lower semicontinuous convex functions on  $X$ . Thanks to Rockafellar’s fundamental work (see [23, Theorem A]) the (possibly set-valued) *subdifferential* operators  $\partial f$  and  $\partial g$  of  $f$  and  $g$  respectively are maximally monotone. Assuming appropriate constraint qualifications the problem of minimizing  $f + g$  amounts to solving the monotone inclusion problem:

$$\text{Find } x \in X \text{ such that } x \in \text{zer}(A + B) = \{x \in X \mid 0 \in Ax + Bx\}. \quad (\text{P})$$

For a comprehensive discussion on (P) and its connection to optimization problems we refer the reader to [2], [10], [11], [13], [24], [25], [26], [29], [30] and the references therein. Splitting algorithms are potential candidates to solve (P). Many of these algorithms employ the resolvent and/or the reflected resolvent of the underlying operators  $A$  and  $B$ . The monotonicity of an operator  $A$  is reflected in the firm nonexpansiveness of its resolvents or, equivalently; the nonexpansiveness of its reflected resolvent. When a monotone operator  $A$  posses supplementary properties, e.g., strong monotonicity, Lipschitz continuity or cocoercivity its reflected resolvent enjoys refined notions of nonexpansiveness, see, e.g., [4], [3], [15], and [18]. However, none of these works studies the notion of *uniform monotonicity* and what the corresponding property in the reflected resolvent (if any) could be.

*The goal of this paper is to provide a systematic study of the class of uniformly monotone operators and their corresponding reflected resolvents. We show that the latter is closely related to the class of the strongly nonexpansive operators introduced by Bruck and Reich [12]. Connection to duality via the inverse operators is systematically studied. When the underlying operators are subdifferentials of proper lower semicontinuous convex functions better duality results hold. We provide applications to Douglas–Rachford, Peaceman–Rachford and forward-backward algorithms . Examples that illustrate and tighten our results are presented.*

## Organization and notation

The organization of this paper is as follows: [Section 2](#) contains a collection of auxiliary results and facts. Our main results appear in [Section 3–Section 7](#). In [Section 3](#) we provide key results concerning the correspondence between the class of uniformly monotone operators and the new class of super strongly nonexpansive mappings. In [Section 4](#) we prove the surjectivity of uniformly monotone operators. In [Section 5](#) and [Section 6](#) we demonstrate the power of self-dual properties and the connection to the class of contractions for large distances. [Section 7](#) is dedicated to the study of the compositions of the classes of nonexpansive mapping studied in the paper. Finally, [Section 8](#)

presents applications of our results to refine and strengthen known results in operator splitting methods.

The notation we adopt is standard and follows largely, e.g., [2] and [23].

## 2 Facts and auxiliary results

We start by recalling the following instrumental fact by Minty.

**Fact 2.1 (Minty's Theorem).** [17] (see also [2, Theorem 21.1]) *Let  $A: X \rightrightarrows X$  be monotone. Then*

$$\text{gra } A = \{(J_A x, (\text{Id} - J_A)x) \mid x \in \text{ran}(\text{Id} + A)\}. \quad (2)$$

Moreover,

$$A \text{ is maximally monotone} \Leftrightarrow \text{ran}(\text{Id} + A) = X. \quad (3)$$

Let  $A: X \rightrightarrows X$  be monotone and let  $(x, u) \in X \times X$ . In view of Fact 2.1, it is easy to check that

$$(x, u) \in \text{gra } J_A \Leftrightarrow (u, x - u) \in \text{gra } A, \quad (4)$$

and that

$$(x, u) \in \text{gra } R_A \Leftrightarrow \left(\frac{1}{2}(x + u), \frac{1}{2}(x - u)\right) \in \text{gra } A. \quad (5)$$

It is straightforward to verify that (see, e.g., [2, Proposition 23.38])

$$\text{zer } A = \text{Fix } J_A = \text{Fix } R_A, \quad (6)$$

and that

$$J_{A^{-1}} = \text{Id} - J_A \text{ and consequently } R_{A^{-1}} = -R_A. \quad (7)$$

**Example 2.2.** *Suppose that  $f: X \rightarrow ]-\infty, +\infty]$  is convex lower semicontinuous and proper. Then  $\partial f$  is maximally monotone. The resolvent  $J_A = J_{\partial f} = \text{Prox } f$  and the reflected resolvent is  $R_f = R_A = 2 \text{Prox } f - \text{Id}$  and hence, by (7),  $R_{f^*} = -R_A = \text{Id} - 2 \text{Prox } f$ .*

*Proof.* See [23] and [2, Example 23.3]. ■

Let  $\phi: \mathbb{R}_+ \rightarrow [0, +\infty]$  be an increasing function that vanishes only at 0 (such a function is called a modulus). Recall that  $f: X \rightarrow ]-\infty, +\infty]$  is *uniformly convex* with modulus  $\phi$  if  $(\forall x \in \text{dom } f) (\forall y \in \text{dom } f) (\forall \alpha \in ]0, 1[)$

$$f(\alpha x + (1 - \alpha)y) + \alpha(1 - \alpha)\phi(\|x - y\|) \leq \alpha f(x) + (1 - \alpha)f(y). \quad (8)$$

Recall also that  $A: X \rightrightarrows X$  is *uniformly monotone* with modulus  $\phi$  if  $\{(x, x^*), (y, y^*)\} \subseteq \text{gra } A$  implies that

$$\langle x - y, x^* - y^* \rangle \geq \phi(\|x - y\|). \quad (9)$$

The notion of uniform monotonicity is naturally motivated by properties of subdifferentials of uniformly convex functions. A comprehensive overview of uniformly convex functions is found in [28]. Some other results regarding uniformly convex functions can be found in [8, 9, 27].

**Fact 2.3.** *Suppose that  $f: X \rightarrow ]-\infty, +\infty]$  is uniformly convex with a modulus  $\phi$ . Then  $\partial f$  is uniformly monotone with a modulus  $2\phi$ .*

*Proof.* See [28, Theorem 3.5.10] and also [2, Example 22.4(iii)]. ■

We now turn to definitions of certain classes of mappings related to the notions of nonexpansiveness and Lipschitz continuity.

**Definition 2.4.** *Let  $T: X \rightarrow X$ , let  $(x, y) \in X \times X$  and let  $\alpha \in ]0, 1[$ .*

- (i)  *$T$  is nonexpansive if  $\|Tx - Ty\| \leq \|x - y\|$ .*
- (ii)  *$T$  is strongly nonexpansive if  $T$  is nonexpansive and we have the implication*

$$\left. \begin{array}{l} (x_n - y_n)_{n \in \mathbb{N}} \text{ is bounded} \\ \|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0 \end{array} \right\} \implies (x_n - y_n) - (Tx_n - Ty_n) \rightarrow 0. \quad (10)$$

- (iii)  *$T$  is  $\alpha$ -averaged if  $\alpha \in [0, 1[$  and there exists a nonexpansive operator  $N: X \rightarrow X$  such that  $T = (1 - \alpha)\text{Id} + \alpha N$ ; equivalently, we have (see [2, Proposition 4.35])*

$$(1 - \alpha)\|(\text{Id} - T)x - (\text{Id} - T)y\|^2 \leq \alpha(\|x - y\|^2 - \|Tx - Ty\|^2). \quad (11)$$

- (iv)  *$T$  is firmly nonexpansive if  $T$  is  $\frac{1}{2}$ -averaged; equivalently,*

$$\|Tx - Ty\|^2 + \|(\text{Id} - T)x - (\text{Id} - T)y\|^2 \leq \|x - y\|^2. \quad (12)$$

- (v)  *$T$  is Lipschitz for large distances (see [7, Proposition 1.11]) if for each  $\epsilon > 0$ , there exists  $K_\epsilon > 0$  so that  $\|Tx - Ty\| \leq K_\epsilon \|x - y\|$  whenever  $\|x - y\| \geq \epsilon$ .*

The following well-known fact summarizes the correspondences between the class of maximally monotone operators and the classes of firmly nonexpansive and nonexpansive mappings.

**Fact 2.5.** *Let  $T: X \rightarrow X$ , and set  $A = T^{-1} - \text{Id}$ . Then  $T = J_A$  and  $2T - \text{Id} = R_A$ . Moreover,*

$$A \text{ is maximally monotone} \Leftrightarrow T \text{ is firmly nonexpansive} \Leftrightarrow 2T - \text{Id} \text{ is nonexpansive.} \quad (13)$$

*Proof.* See, e.g., [14, Theorem 2] or [2, Corollary 23.11]. ■

We conclude this section with the following fact concerning the asymptotic behaviour of iterates of strongly nonexpansive mappings.

**Fact 2.6.** *Let  $T: X \rightarrow X$  be strongly nonexpansive. Suppose that  $\text{Fix } T \neq \emptyset$ . Let  $x_0 \in X$ . Then there exists  $\bar{x} \in \text{Fix } T$  such that  $(T^n x_0)_{n \in \mathbb{N}}$  converges weakly to  $\bar{x}$ .*

*Proof.* See [12, Corollary 1.1]. ■

### 3 Strongly nonexpansive and super strongly nonexpansive mappings

The main goal of this section is to show that the notion of uniform monotonicity of an operator corresponds to the notion of *super strong* nonexpansiveness (see [Definition 3.1](#) below) of its reflected resolvent. Throughout we assume that

$$A: X \rightrightarrows X \text{ and } B: X \rightrightarrows X \text{ are maximally monotone.} \quad (14)$$

We start by defining a new subclass of nonexpansive mappings.

**Definition 3.1.** *Let  $T: X \rightarrow X$ . We say that  $T$  is super strongly nonexpansive if  $T$  is nonexpansive and we have the implication*

$$\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0 \Rightarrow (x_n - y_n) - (Tx_n - Ty_n) \rightarrow 0. \quad (15)$$

**Proposition 3.2.** *Let  $T: X \rightarrow X$ . Suppose  $T$  is super strongly nonexpansive. Then  $T$  is strongly nonexpansive.*

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  be sequences in  $X$  such that  $(x_n - y_n)_{n \in \mathbb{N}}$  is bounded and suppose that  $\|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0$ . We claim that

$$\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0. \quad (16)$$

Indeed, the nonexpansiveness of  $T$  implies that  $(Tx_n - Ty_n)_{n \in \mathbb{N}}$  is bounded. Now  $\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 = (\|x_n - y_n\| - \|Tx_n - Ty_n\|)(\|x_n - y_n\| + \|Tx_n - Ty_n\|) \rightarrow 0$ . Since  $T$  is super strongly nonexpansive we conclude that  $(x_n - y_n) - (Tx_n - Ty_n) \rightarrow 0$ . Hence,  $T$  is strongly nonexpansive as claimed. ■

The converse of [Proposition 3.2](#) is not true in general as we illustrate in [Example 3.7](#) below. Nonetheless, when  $X = \mathbb{R}$  the converse of [Proposition 3.2](#) holds as we next illustrate in [Proposition 3.3](#). We will use the following simple observation. Let  $(a, b) \in X \times X$ . Then

$$(\|a\| - \|b\|)^2 \leq \|\|a\|^2 - \|b\|^2\|. \quad (17)$$

**Proposition 3.3 (the case of the real line).** *Suppose  $T: \mathbb{R} \rightarrow \mathbb{R}$  is strongly nonexpansive, then  $T$  is super strongly nonexpansive.*

*Proof.* Suppose for eventual contradiction that  $T$  is strongly nonexpansive, but  $T$  is not super strongly nonexpansive. Then we have sequences  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}$  so that

$$|x_n - y_n|^2 - |Tx_n - Ty_n|^2 \rightarrow 0 \quad (18)$$

but

$$(x_n - y_n) - (Tx_n - Ty_n) \not\rightarrow 0. \quad (19)$$

It follows from the nonexpansiveness of  $T$  and (17) applied with  $(a, b, X)$  replaced by  $(x_n - y_n, Tx_n - Ty_n, \mathbb{R})$  that  $(\forall n \in \mathbb{N}) 0 \leq (|x_n - y_n| - |Tx_n - Ty_n|)^2 \leq |x_n - y_n|^2 - |Tx_n - Ty_n|^2$ . Therefore (18) implies that

$$|x_n - y_n| - |Tx_n - Ty_n| \rightarrow 0, \quad (20)$$

and because  $T$  is strongly nonexpansive, (19) and (20) imply  $(x_n - y_n)$  is not bounded. Thus, without loss of generality, and swapping  $x_n$  and  $y_n$  as necessary, we may and do assume  $(\forall n \in \mathbb{N}) y_n - x_n > 1$ .

If  $(Ty_n - Tx_n)_{n \in \mathbb{N}}$  is eventually positive, that is, the same sign as  $(y_n - x_n)$  for all large  $n$ , then (20) would imply that  $(y_n - x_n) - (Ty_n - Tx_n) \rightarrow 0$  in contradiction with (19). Therefore, after passing to a subsequence and relabelling if necessary, we may and do assume that

$$(\forall n \in \mathbb{N}) Ty_n - Tx_n < 0. \quad (21)$$

Now consider  $z_n = x_n + 1$ , so  $x_n < z_n < y_n$ . Observe that

$$|x_n - y_n| = |x_n - z_n| + |z_n - y_n| = 1 + |z_n - y_n|. \quad (22)$$

Hence, because  $T$  is nonexpansive we learn, in view of the triangle inequality, (22) and (20), that

$$0 \leq |x_n - z_n| - |Tx_n - Tz_n| + |z_n - y_n| - |Tz_n - Ty_n| \quad (23a)$$

$$= |x_n - y_n| - |Tx_n - Tz_n| - |Tz_n - Ty_n| \leq |x_n - y_n| - |Tx_n - Ty_n| \rightarrow 0. \quad (23b)$$

It follows from the nonexpansiveness of  $T$  and (23) that  $|Tz_n - Tx_n| - |z_n - x_n| \rightarrow 0$ . Consequently, because  $T$  is strongly nonexpansive, we have  $(Tz_n - Tx_n) - 1 = (Tz_n - Tx_n) - (z_n - x_n) \rightarrow 0$ . After passing to a subsequence and relabelling if necessary, we may and do assume that

$$(\forall n \in \mathbb{N}) Tz_n \geq Tx_n. \quad (24)$$

Because  $T$  is nonexpansive, in view of (22) we have  $(\forall n \in \mathbb{N}) |Ty_n - Tz_n| \leq |y_n - z_n| = |y_n - x_n| - 1$ . This and (24) imply  $(\forall n \in \mathbb{N}) Ty_n \geq Tz_n - (|y_n - x_n| - 1) \geq Tx_n - |y_n - x_n| + 1$ . Therefore, by (20) we have  $(\forall n \in \mathbb{N}) -|y_n - x_n| + 1 \leq Ty_n - Tx_n < 0$ . That is  $(\forall n \in \mathbb{N}) |x_n - y_n| - |Tx_n - Ty_n| \geq 1$ , and this contradicts (20). This completes the proof.  $\blacksquare$

We now turn to the correspondence between uniformly monotone operators and super strongly nonexpansive operators. We start with the following lemma that provides a characterization of uniformly monotone operators via their reflected resolvents.

**Lemma 3.4.** *The following hold:*

(i) *Suppose that  $A$  is uniformly monotone with modulus  $\phi$ . Then  $(\forall (x, y) \in X \times X)$*

$$\|x - y\|^2 - \|R_A x - R_A y\|^2 \geq 4\phi\left(\frac{1}{2}\|(x - y) + (R_A x - R_A y)\|\right) = 4\phi(\|J_A x - J_A y\|). \quad (25)$$

(ii) *Suppose that there exists a modulus function  $\phi$  such that  $(\forall (x, y) \in X \times X)$*

$$\|x - y\|^2 - \|R_A x - R_A y\|^2 \geq \phi(\|(x - y) + (R_A x - R_A y)\|). \quad (26)$$

*Then  $A$  is uniformly monotone with a modulus  $\frac{1}{4}\phi \circ (2(\cdot))$ .*

*Proof.* (i): It follows from (5) that  $\text{gra } A = \{\frac{1}{2}(x + R_A x, x - R_A x) \mid x \in X\}$ . Now combine this with (9). (ii): Let  $\{(x, x^*), (y, y^*)\} \subseteq \text{gra } A$  and observe that (5) implies that

$$\{(x + x^*, x - x^*), (y + y^*, y - y^*)\} \subseteq \text{gra } R_A. \quad (27)$$

Now

$$\langle x - y, x^* - y^* \rangle \quad (28a)$$

$$= \frac{1}{4} \langle x + x^* - (y + y^*) + x - x^* - (y - y^*), x + x^* - (y + y^*) - (x - x^* - (y - y^*)) \rangle \quad (28b)$$

$$= \frac{1}{4} (\|x + x^* - (y + y^*)\|^2 - \|x - x^* - (y - y^*)\|^2) \geq \frac{1}{4} \phi(2\|x - y\|), \quad (28c)$$

where the inequality in (28c) follows from combining (26) applied with  $(x, y)$  replaced with  $(x + x^*, y + y^*)$  and (27). The proof is complete.  $\blacksquare$

**Proposition 3.5.** *Consider the following statements:*

- (i)  $A$  is uniformly monotone.
- (ii)  $-R_A$  is super strongly nonexpansive.
- (iii)  $-R_A$  is strongly nonexpansive.

Then (i)  $\Leftrightarrow$  (ii)  $\Rightarrow$  (iii).

*Proof.* (i)  $\Rightarrow$  (ii): Let  $\phi$  be a modulus function for  $A$ . Recalling (13) we have  $-R_A$ , as is  $R_A$ , is nonexpansive. Let  $(x_n - y_n)_{n \in \mathbb{N}}$  be a sequence in  $X$  and suppose that  $\|x_n - y_n\|^2 - \|R_A x_n - R_A y_n\|^2 \rightarrow 0$ . Combining this with Lemma 3.4, we learn that  $0 \leq \phi(\|(x_n + R_A x_n) - (y_n + R_A y_n)\|) \rightarrow 0$ . Since  $\phi$  is increasing and vanishes only at 0 we must have  $(x_n + R_A x_n) - (y_n + R_A y_n) \rightarrow 0$ , hence  $-R_A$  is super strongly nonexpansive.

(ii)  $\Rightarrow$  (i): Suppose  $A$  is not uniformly monotone but  $-R_A$  is super strongly nonexpansive. Because  $A$  is not uniformly monotone, there exist sequences  $(a_n, a_n^*)_{n \in \mathbb{N}}$  and  $(b_n, b_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  and  $\epsilon > 0$  such that

$$\langle a_n - b_n, a_n^* - b_n^* \rangle \rightarrow 0 \text{ but } (\forall n \in \mathbb{N}) \|a_n - b_n\| \geq \epsilon. \quad (29)$$

Set  $(\forall n \in \mathbb{N}) x_n = a_n + a_n^*$  and  $y_n = b_n + b_n^*$  and observe that Minty's parametrization of  $\text{gra } A$  implies that  $(\forall n \in \mathbb{N}) (a_n, a_n^*, b_n, b_n^*) = \frac{1}{2}(x_n + R_A x_n, x_n - R_A x_n, y_n + R_A y_n, y_n - R_A y_n)$ . Therefore

$$\begin{aligned} & \|x_n - y_n\|^2 - \| -R_A x_n - (-R_A y_n) \|^2 \\ &= \|x_n - y_n\|^2 - \|R_A x_n - R_A y_n\|^2 \end{aligned} \quad (30a)$$

$$= \langle x_n - y_n + (R_A x_n - R_A y_n), x_n - y_n - (R_A x_n - R_A y_n) \rangle \quad (30b)$$

$$= \langle x_n + R_A x_n - (y_n + R_A y_n), x_n - R_A x_n - (y_n - R_A y_n) \rangle \quad (30c)$$

$$= 4 \langle a_n - b_n, a_n^* - b_n^* \rangle \rightarrow 0, \quad (30d)$$

where the limit in (30d) follows from (29). Because  $-R_A$  is super strongly nonexpansive (30) implies

$$a_n - b_n = \frac{1}{2}((x_n + R_A x_n) - (y_n + R_A y_n)) = \frac{1}{2}((x_n - y_n) - (-R_A x_n - (-R_A y_n))) \rightarrow 0. \quad (31)$$

This contradicts (29), hence  $A$  is uniformly monotone. (ii) $\Rightarrow$ (iii): Apply Proposition 3.2 with  $T$  replaced by  $-R_A$ .  $\blacksquare$

**Corollary 3.6.** *Suppose that  $X = \mathbb{R}$ . The following are equivalent.*

- (i)  $A$  is uniformly monotone.
- (ii)  $-R_A$  is strongly nonexpansive.
- (iii)  $-R_A$  is super strongly nonexpansive.

*Proof.* Combine Proposition 3.5 and Proposition 3.3.  $\blacksquare$

**Example 3.7.** *Suppose that  $X = \mathbb{R}^2$ . Set  $a_0 = 0$  and set ( $\forall m \geq 1$ )*

$$a_m = 2^{m+1} - 2 \quad (32a)$$

$$w_m = \frac{1}{\sqrt{4^m+1}}(2^m, 1) \quad (32b)$$

$$K_m = (4^m - 4^{-m})^{1/2} \quad (32c)$$

$$\beta_m = \frac{1}{2^m} K_m \quad (32d)$$

$$D_m = \{(x, y) \mid x \leq a_m\}. \quad (32e)$$

Let

$$T(x, y): \mathbb{R}^2 \rightarrow \mathbb{R}^2: (x, y) \mapsto \begin{cases} (0, 0), & x \leq 0; \\ \sum_{j=0}^{m-1} K_j w_j + (\frac{x-a_{m-1}}{2^m}) K_m w_m, & x \in [a_{m-1}, a_m], m \geq 1. \end{cases} \quad (33)$$

Set  $\tilde{A} = (\frac{\text{Id}-T}{2})^{-1} - \text{Id}$ .

Then the following hold:

- (i) ( $\forall m \in \mathbb{N}$ )  $T|_{D_m}$  is a contraction with a constant  $\beta_m$ .
- (ii)  $T$  is strongly nonexpansive, hence nonexpansive.
- (iii) There exist sequences  $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}^2$  satisfying

$$\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0, \text{ nevertheless } (x_n - y_n) - (Tx_n - Ty_n) \not\rightarrow 0. \quad (34)$$

Consequently,  $T$  is not super strongly nonexpansive.

- (iv)  $T = -R_{\tilde{A}}$ .
- (v)  $\tilde{A}$  is maximally monotone.
- (vi)  $\tilde{A}$  is not uniformly monotone.

*Proof.* Observe that  $(w_m)_{m \in \mathbb{N}}$  is a sequence of unit vectors whose positive slopes are strictly decreasing to 0, that  $(\beta_m)_{m \in \mathbb{N}}$  is a sequence of strictly increasing real numbers in  $[0, 1[$  and that  $(K_m)_{m \in \mathbb{N}}$  is a sequence of strictly increasing real numbers with  $K_m \rightarrow +\infty$ .

(i): Let  $m \in \mathbb{N}$ . Observe that if  $m = 0$  the conclusion is obvious. Therefore, we assume  $m \geq 1$ . Let  $\{u, v\} \subseteq D_m$ , with  $u = (u_1, u_2)$  and  $v = (v_1, v_2)$  and let  $\{r, s\} \subseteq \{1, \dots, m\}$  be such that



$u_1 \in [a_{r-1}, a_r]$  and  $v_1 \in [a_{s-1}, a_s]$ . Without loss of generality we may and do assume that  $u_1 \leq v_1$  and hence  $r \leq s \leq m$ . If  $r = s$  then the definition of  $T$  implies that

$$Tu - Tv = \beta_r(u_1 - v_1)w_r. \quad (35)$$

Consequently, we have

$$\|Tu - Tv\| = \beta_r|u_1 - v_1| \leq \beta_r\|u - v\|. \quad (36)$$

Observe that  $(\forall (u_1, u_2) \in \mathbb{R}^2)$  we have  $T(u_1, u_2) = T(u_1, 0)$ . Moreover, let  $k \geq 1$ . Applying (35) with  $(u, v)$  replaced by  $((a_{k-1}, 0), (a_k, 0))$  yields:

$$T(a_k, 0) - T(a_{k-1}, 0) = \beta_k(a_k - a_{k-1})w_k = K_k w_k. \quad (37)$$

Using the triangle inequality, (35) and (37) we have

$$\|Tu - Tv\| = \|T(v_1, v_2) - T(u_1, u_2)\| = \|T(v_1, 0) - T(u_1, 0)\| \quad (38a)$$

$$\begin{aligned} &= \|T(v_1, 0) - T(a_{s-1}, 0) + T(a_{s-1}, 0) - T(a_{s-2}, 0) + \dots \\ &\quad - T(a_r, 0) + T(a_r, 0) - T(u_1, 0)\| \end{aligned} \quad (38b)$$

$$\begin{aligned} &\leq \|T(v_1, 0) - T(a_{s-1}, 0)\| + \|T(a_{s-1}, 0) - T(a_{s-2}, 0)\| + \dots \\ &\quad + \|T(a_r, 0) - T(u_1, 0)\| \end{aligned} \quad (38c)$$

$$= \beta_s(v_1 - a_{s-1}) + \beta_{s-1}(a_{s-1} - a_{s-2}) + \dots + \beta_{r+1}(a_{r+1} - a_r) + \beta_r(a_r - u_1) \quad (38d)$$

$$\leq \beta_m(v_1 - a_{s-1}) + \beta_m(a_{s-1} - a_{s-2}) + \dots + \beta_m(a_{r+1} - a_r) + \beta_m(a_r - u_1) \quad (38e)$$

$$= \beta_m(v_1 - u_1) \leq \beta_m\|u - v\|. \quad (38f)$$

(ii): Suppose  $(x_n)_{n \in \mathbb{N}}$ , and  $(y_n)_{n \in \mathbb{N}}$  are sequences in  $\mathbb{R}^2$  such that

$$\|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0 \text{ and } (x_n - y_n)_{n \in \mathbb{N}} \text{ is bounded.} \quad (39)$$

Let us denote  $x_n = (x_{n,1}, x_{n,2})$ ,  $y_n = (y_{n,1}, y_{n,2})$ . We proceed by proving the following claims:

CLAIM 1: The sequences  $(x_{n,1})_{n \in \mathbb{N}}$ , and  $(y_{n,1})_{n \in \mathbb{N}}$  are unbounded.

Indeed, suppose for eventual contradiction that one of the sequences  $(x_{n,1})_{n \in \mathbb{N}}$ , and  $(y_{n,1})_{n \in \mathbb{N}}$  is bounded. The boundedness of  $(x_n - y_n)_{n \in \mathbb{N}}$  implies that both sequences  $(x_{n,1})_{n \in \mathbb{N}}$  and  $(y_{n,1})_{n \in \mathbb{N}}$  must be bounded. Indeed, without loss of generality we may and do assume that  $(x_n - y_n) \not\rightarrow 0$ . Let  $m \geq 1$  be such that  $(\forall n \in \mathbb{N}) \max\{x_{n,1}, y_{n,1}\} \leq a_m$ . Observe that (i) implies that  $\|Tx_n - Ty_n\| \leq \beta_m\|x_n - y_n\|$ . Hence,

$$0 < (1 - \beta_m)\|x_n - y_n\| \leq \|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0. \quad (40)$$

That is,  $\|x_n - y_n\| \rightarrow 0$ , which is absurd. Therefore, the sequences  $(x_{n,1})_{n \in \mathbb{N}}$  and  $(y_{n,1})_{n \in \mathbb{N}}$  are unbounded as claimed. After passing to a subsequence and relabelling if necessary we conclude that

$$x_{n,1} \rightarrow \infty \text{ and } y_{n,1} \rightarrow \infty. \quad (41)$$

Because  $(x_n - y_n)_{n \in \mathbb{N}}$  is bounded and the slopes of the vectors  $(w_n)_{n \in \mathbb{N}}$  go to 0 as  $n \rightarrow \infty$ . CLAIM 2:

$$(\forall n \in \mathbb{N}) Tx_n - Ty_n = (c_n(x_{n,1} - y_{n,1}), d_n) \text{ where } c_n \rightarrow 1^- \text{ and } d_n \rightarrow 0. \quad (42)$$

To verify CLAIM 2, let  $M > 0$  be such that  $(\forall n \in \mathbb{N}) \|x_n - y_n\| \leq M$ . it follows from (35) that

$$Tu - Tv = \frac{K_r}{\sqrt{4^r+1}}(u_1 - v_1)\left(1, \frac{1}{2^r}\right). \quad (43)$$

In view of (43) and (36) we learn that for  $n$  and  $j$  such that  $\min\{x_{n,1}, y_{n,1}\} \geq a_{j-1}$ , it follows that

$$|d_n| \leq 2^{-j}M \quad \text{while} \quad c_n \geq \frac{K_j}{\sqrt{4^j+1}} = \left(\frac{4^j - 4^{-j}}{4^j + 1}\right)^{1/2}. \quad (44)$$

Note it is possible that the interval with endpoints  $x_{n,1}$  and  $y_{n,1}$  may intersect more than one interval  $[a_{r-1}, a_r]$ . However,  $2^{-n}$  decreases as  $n$  increases and  $K_n/\sqrt{4^n+1}$  increases as  $n$  increases in (43) and so (44) remains valid in this case as well since  $a_{j-1} \leq \min\{x_{n,1}, y_{n,1}\}$ . This verifies CLAIM 2.

CLAIM 3:

$$|x_{n,2} - y_{n,2}| \rightarrow 0. \quad (45)$$

Indeed, set  $(\forall n \in \mathbb{N}) \bar{x}_n = (x_{n,1}, 0)$  and  $\bar{y}_n = (y_{n,1}, 0)$  and note that  $\|\bar{x}_n - \bar{y}_n\| \leq \|x_n - y_n\|$  and that  $(T\bar{x}_n, T\bar{y}_n) = (Tx_n, Ty_n)$ . Therefore, the nonexpansiveness of  $T$  and (39) imply

$$0 \leq \|\bar{x}_n - \bar{y}_n\| - \|T\bar{x}_n - T\bar{y}_n\| = \|\bar{x}_n - \bar{y}_n\| - \|Tx_n - Ty_n\| \leq \|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0. \quad (46)$$

That is

$$\|\bar{x}_n - \bar{y}_n\| - \|Tx_n - Ty_n\| \rightarrow 0. \quad (47)$$

Subtracting (39) and (47) yields

$$\|x_n - y_n\| - \|\bar{x}_n - \bar{y}_n\| \rightarrow 0. \quad (48)$$

Consequently we have

$$|x_{n,2} - y_{n,2}|^2 = \|x_n - y_n\|^2 - \|\bar{x}_n - \bar{y}_n\|^2 \quad (49a)$$

$$= (\|x_n - y_n\| - \|\bar{x}_n - \bar{y}_n\|)(\|x_n - y_n\| + \|\bar{x}_n - \bar{y}_n\|) \rightarrow 0. \quad (49b)$$

Combining CLAIM 2 and CLAIM 3 we learn that

$$(x_n - y_n) - (Tx_n - Ty_n) \rightarrow (0, 0), \quad (50)$$

hence  $T$  is strongly nonexpansive.

(iii): Set  $(\forall n \in \mathbb{N}) x_n = (a_{n+1}, 0)$  and  $y_n = (a_n, 0)$ . Observe that  $x_n - y_n = (2^n, 0)$  while  $Tx_n - Ty_n = K_n w_n = \frac{K_n}{\sqrt{4^n+1}}(2^n, 1)$ . Therefore,  $\|x_n - y_n\|^2 = 4^n$  and  $\|Tx_n - Ty_n\|^2 = \|K_n w_n\|^2 = K_n^2 = 4^n - 4^{-n}$ . Hence

$$\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0 \quad (51)$$

However,

$$(x_n - y_n) - (Tx_n - Ty_n) = \left(1 - \frac{K_n}{\sqrt{4^n+1}}\right)2^n, -\frac{K_n}{\sqrt{4^n+1}} \rightarrow (0, -1) \neq (0, 0). \quad (52)$$

and so (34) holds.

(iv): This is clear.

(v): Combine (ii) and [2, Corollary 23.9 and Proposition 4.4].

(vi): Combine (iii), (iv) and Proposition 3.5. ■




Figure 1: A GeoGebra snapshot illustrating the operator  $T$  in Example 3.7. Here  $x(0) := T(x, y), x \leq 0, y \in \mathbb{R}$  and  $x(a_i) := T(a_i, y)$  where  $y \in \mathbb{R}, i \in \{1, 2, \dots, 7\}$ .

## 4 Properties of Uniformly Monotone Operators

The main goals of this section are to establish that uniformly monotone operators are surjective, have unique zeros and have uniformly continuous inverse operators.

The following result is motivated by Zălinescu's important result [28, Proposition 3.5.1] concerning the growth rate of the modulus of a uniformly convex function.

**Proposition 4.1.** *Let  $Y$  be a Banach space and suppose  $B: Y \rightrightarrows Y^*$  is uniformly monotone<sup>1</sup> with convex*

<sup>1</sup>Let  $Y$  be a Banach space. An operator  $B: Y \rightrightarrows Y^*$  is *uniformly monotone* with a modulus  $\phi: \mathbb{R}_+ \rightarrow [0, +\infty]$  if  $\phi$  is increasing, vanishes only at 0 and  $(\forall (x, x^*) \in \text{gra } B) (\forall (y, y^*) \in \text{gra } B) \langle x - y, x^* - y^* \rangle \geq \phi(\|x - y\|)$ .

domain. Then  $B$  has a supercoercive modulus  $\phi$  that satisfies the following property.

$$\text{For each } \epsilon > 0 \text{ } (\exists \beta_\epsilon > 0) \text{ so that } \phi(t) \geq \beta_\epsilon t^2 \text{ whenever } t \geq \epsilon, \text{ in particular } \liminf_{t \rightarrow \infty} \frac{\phi(t)}{t^2} > 0. \quad (53)$$

*Proof.* Because  $B$  is uniformly monotone we fix  $\alpha > 0$  so that  $(\forall (x, x^*) \in \text{gra } B) (\forall (y, y^*) \in \text{gra } B)$

$$\langle y - x, y^* - x^* \rangle \geq \alpha \text{ whenever } \|y - x\| \geq 1. \quad (54)$$

We claim that:

$$\langle y - x, y^* - x^* \rangle \geq \frac{\alpha}{4} \|x - y\|^2 \text{ whenever } \|y - x\| \geq 1, (x, x^*) \in \text{gra } B, (y, y^*) \in \text{gra } B. \quad (55)$$

To verify (55) it is sufficient to show that  $(\forall k \in \mathbb{N})$  we have

$$\langle y - x, y^* - x^* \rangle \geq 2^{2k} \alpha \text{ whenever } \|y - x\| \geq 2^k, (x, x^*) \in \text{gra } B, (y, y^*) \in \text{gra } B. \quad (56)$$

We proceed by induction on  $k$ . Indeed, (54) verifies the base case at  $k = 0$ . Now suppose that (56) is true for some  $k = n$ . Suppose that  $(x, x^*) \in \text{gra } B, (z, z^*) \in \text{gra } B$  satisfy  $\|z - x\| \geq 2^{n+1}$ . Let  $y = (x + z)/2$  and observe that  $y - x = z - y = (z - x)/2$ , hence  $\|y - x\| = \|z - y\| \geq 2^n$ . Because  $\text{dom } B$  is convex we have  $y \in \text{dom } B$  and so we can pick  $y^* \in B y$ . It follows from the inductive hypothesis that

$$2^{2n} \alpha \leq \langle y - x, y^* - x^* \rangle = \langle \frac{z-x}{2}, y^* - x^* \rangle \quad (57a)$$

$$2^{2n} \alpha \leq \langle z - y, z^* - y^* \rangle = \langle \frac{z-x}{2}, z^* - y^* \rangle \quad (57b)$$

Adding (57a) and (57b) yields  $\langle \frac{z-x}{2}, z^* - x^* \rangle \geq 2(2^{2n})\alpha$  and consequently

$$\langle z - x, z^* - x^* \rangle \geq 4(2^{2n})\alpha = 2^{2(n+1)}\alpha, \quad (58)$$

which proves (55). We now establish (53). Let  $\epsilon > 0$ . In view of (55) if  $\epsilon \geq 1$  we set  $\beta_\epsilon = \frac{\alpha}{4}$ . Now suppose that  $0 < \epsilon < 1$ . The uniform monotonicity of  $B$  implies that  $(\exists \alpha_\epsilon > 0)$  such that

$$\langle y - x, y^* - x^* \rangle \geq \alpha_\epsilon \text{ whenever } \|y - x\| \geq \epsilon, (x, x^*) \in \text{gra } B, (y, y^*) \in \text{gra } B. \quad (59)$$

In view of (55) the conclusion follows by setting  $\beta_\epsilon = \min\{\alpha/4, \alpha_\epsilon\}$  for  $0 < \epsilon < 1$ , and  $\beta_\epsilon = \alpha/4$  for  $\epsilon \geq 1$ . ■

Analogous to the concept of Lipschitz for large distances in [7], we introduce the following definition.

**Definition 4.2.** We say that  $T: X \rightarrow X$  is a contraction for large distances if for each  $\epsilon > 0$ ,  $(\exists K_\epsilon < 1)$  so that  $\|Tx - Ty\| \leq K_\epsilon \|x - y\|$  whenever  $(x, y) \in X \times X$  satisfy  $\|x - y\| \geq \epsilon$ .

**Remark 4.3.** Contractions for large distances are not a new concept. In fact, they coincide with an unnamed class of nonexpansive maps introduced by Rakotch in [20, Definition 2], and have been referred to as contractive by others, including, for example, Reich & Zaslavski in [21]. However, in [20] the term contractive referred to the more general class of strictly nonexpansive mappings.

**Lemma 4.4.** *Suppose  $A$  is uniformly monotone, then the following hold:*

- (i)  $J_{A^{-1}}$  is uniformly monotone with a supercoercive modulus.
- (ii) For each  $\epsilon > 0$  ( $\exists \beta_\epsilon \in ]0, 1[$ ) such that ( $\forall (x, y) \in X \times X$ ) satisfying  $\|x - y\| \geq \epsilon$  we have

$$\langle J_A x - J_A y, J_{A^{-1}} x - J_{A^{-1}} y \rangle + \|J_{A^{-1}} x - J_{A^{-1}} y\|^2 \geq \beta_\epsilon \|x - y\|^2. \quad (60)$$

- (iii)  $J_{A^{-1}}$  is surjective.
- (iv)  $J_A$  is a contraction for large distances.

*Proof.* Let  $\epsilon > 0$  and let  $\phi$  be a modulus function for  $A$ . Set

$$\alpha = \alpha(\epsilon) = \min\{\phi(\epsilon/2), \epsilon^2/4\}. \quad (61)$$

Then  $\alpha > 0$ . (i)&(ii): Let  $(x, y) \in X \times X$  be such that  $\|x - y\| \geq \epsilon$ . The triangle inequality and [Fact 2.1](#) imply that

$$\max\{\|J_A x - J_A y\|, \|J_{A^{-1}} x - J_{A^{-1}} y\|\} \geq \epsilon/2. \quad (62)$$

Therefore we obtain

$$\langle x - y, J_{A^{-1}} x - J_{A^{-1}} y \rangle = \langle J_A x - J_A y + J_{A^{-1}} x - J_{A^{-1}} y, J_{A^{-1}} x - J_{A^{-1}} y \rangle \quad (63a)$$

$$= \langle J_A x - J_A y, J_{A^{-1}} x - J_{A^{-1}} y \rangle + \|J_{A^{-1}} x - J_{A^{-1}} y\|^2 \quad (63b)$$

$$\geq \phi(\|J_A x - J_A y\|) + \|J_{A^{-1}} x - J_{A^{-1}} y\|^2 \geq \alpha. \quad (63c)$$

It follows from (63) that  $J_{A^{-1}}$  is uniformly monotone. Combining this with [Proposition 4.1](#) applied with  $(Y, A)$  replaced by  $(X, J_{A^{-1}})$  and the fact that  $\text{dom } J_{A^{-1}} = X$  we learn that  $J_{A^{-1}}$  is uniformly monotone with a supercoercive modulus that satisfies (53). The claim that  $\beta_\epsilon \leq 1$  is a direct consequence of the nonexpansiveness of  $J_{A^{-1}}$  and the monotonicity of  $A$  in view of (60).

(iii): Combine (i) and [2, Proposition 22.11(ii)]. (iv): Indeed, using (ii) and the firm nonexpansiveness of  $J_A$  there exists ( $\exists \beta_\epsilon \in ]0, 1[$ ) such that

$$0 \leq \|J_A x - J_A y\|^2 \leq \langle x - y, J_A x - J_A y \rangle = \|x - y\|^2 - \langle x - y, J_{A^{-1}} x - J_{A^{-1}} y \rangle \quad (64a)$$

$$\leq \|x - y\|^2 - \phi(\|x - y\|) \leq (1 - \beta_\epsilon) \|x - y\|^2, \quad (64b)$$

and the conclusion follows. ■

The previous results lead to nice consequences for uniformly monotone operators which we next state as the main result of this section.

**Theorem 4.5.** *Suppose  $A: X \rightrightarrows X$  is uniformly monotone. Then the following hold.*

- (i)  $A$  satisfies the growth condition

$$\lim_{\|x-y\| \rightarrow \infty} \inf_{\substack{(x, x^*) \in \text{gra } A \\ (y, y^*) \in \text{gra } A}} \frac{\|x^* - y^*\|}{\|x - y\|} > 0. \quad (65)$$

- (ii)  $A$  is surjective.
- (iii)  $A$  has a unique zero.
- (iv)  $A^{-1}$  is uniformly continuous.

*Proof. (i):* Suppose for eventual contradiction that there exist sequences  $(x_n, x_n^*)_{n \in \mathbb{N}}, (y_n, y_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  such that  $\|x_n - y_n\| \rightarrow \infty$  but

$$\lim_{n \rightarrow \infty} \frac{\|x_n^* - y_n^*\|}{\|x_n - y_n\|} = 0. \quad (66)$$

Indeed, write  $\|x_n^* - y_n^*\| = a_n \|x_n - y_n\|$  where  $a_n \rightarrow 0^+$ . Then using (60), we have  $\beta > 0$  such that

$$(\forall n \in \mathbb{N}) \quad a_n \|x_n - y_n\|^2 + a_n^2 \|x_n - y_n\|^2 \geq \beta (\|x_n - y_n\|^2 + a_n^2 \|x_n - y_n\|^2), \quad (67)$$

where on the right side, we drop the inner product from (60) because it is nonnegative by monotonicity of  $A$ . But the above inequality is impossible since  $a_n \rightarrow 0^+$  while  $\beta > 0$  is fixed. Hence, (65) holds.

(ii): Combine Lemma 4.4(iii) and the fact that  $\text{ran } A = \text{dom } A^{-1} = \text{dom}(\text{Id} + A^{-1}) = \text{ran}(\text{Id} + A^{-1})^{-1} = \text{ran } J_{A^{-1}}$ .

(iii): It follows from (ii) that  $\text{zer } A \neq \emptyset$ . Moreover,  $A$  is strictly monotone, hence  $\text{zer } A$  is at most a singleton by, e.g., [2, Proposition 23.35]. Altogether,  $A$  possess a unique zero.

(iv): Let  $\epsilon > 0$ . In view of (i) choose  $K > 0$  such that  $(\forall (x, x^*) \in \text{gra } A) (\forall (y, y^*) \in \text{gra } A)$

$$\|x - y\| \geq K \Rightarrow \|x^* - y^*\| > \epsilon. \quad (68)$$

Let  $\alpha = \phi(\epsilon)$  where  $\phi$  is a modulus function for  $A$ . Then  $\alpha > 0$  and  $(\forall (x, x^*) \in \text{gra } A) (\forall (y, y^*) \in \text{gra } A)$

$$\|x - y\| \geq \epsilon \Rightarrow \langle x - y, x^* - y^* \rangle \geq \alpha. \quad (69)$$

Now choose  $\delta = \min\{\alpha/K, \epsilon\}$ . Recalling (ii), let  $\{x^*, y^*\} \subseteq X = \text{dom } A^{-1}$ . Suppose  $\|x^* - y^*\| < \delta$  and let  $(x^*, x), (y^*, y)$  be points in  $\text{gra } A^{-1}$ , equivalently;  $(x, x^*), (y, y^*)$  are points in  $\text{gra } A$ . Because  $\|x^* - y^*\| < \delta \leq \epsilon$ , (68) implies  $\|x - y\| < K$ . Because  $\|x - y\| < K$ , using Cauchy-Schwarz we obtain

$$\langle x - y, x^* - y^* \rangle \leq \|x - y\| \|x^* - y^*\| < K\delta \leq \alpha. \quad (70)$$

Combining (70) and (69) we learn that  $\|x - y\| < \epsilon$ . Therefore  $A^{-1}$  is uniformly continuous as desired.  $\blacksquare$

**Remark 4.6.** In passing we point out that Theorem 4.5(ii)&(iii) relax the assumptions of [2, Proposition 22.11 and Corollary 23.37]. Indeed, Theorem 4.5(ii)&(iii) assume only the uniform monotonicity of  $A$  and do not require supercoercivity of the modulus.

Following [26, p. 160], we will say that  $S: X \rightrightarrows X$  is *coercive* provided that  $\inf \langle x, Sx \rangle / \|x\| \rightarrow \infty$  as  $\|x\| \rightarrow \infty$ , where we use the standard convention that the infimum of the empty set is  $+\infty$ . Or in other words, given any  $K > 0$  ( $\exists M > 0$ ) so that

$$\langle x, x^* \rangle \geq K \|x\| \text{ whenever } \|x\| \geq M, (x, x^*) \in \text{gra } S. \quad (71)$$

Neither the growth condition in [Theorem 4.5\(i\)](#) and nor coercivity implies the other for monotone operators as we illustrate in [Example 4.7](#) and [Example 4.8\(iii\)&\(v\)](#).

**Example 4.7.** Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined by

$$f(\zeta_1, \zeta_2) = \begin{cases} \zeta_1^2, & \text{if } 0 \leq \zeta_1, 0 \leq \zeta_2 \leq \zeta_1; \\ +\infty, & \text{otherwise.} \end{cases} \quad (72)$$

Set  $A = \partial f$ . Then

$$\liminf_{\|x\| \rightarrow \infty} \left\{ \frac{\langle x^*, x \rangle}{\|x\|^2} \mid x^* \in Ax \right\} > 0. \quad (73)$$

Hence,  $A$  is coercive. However,  $A$  does not satisfy the growth condition in [Theorem 4.5\(i\)](#).

*Proof.* Because  $f(x) \geq \frac{1}{2}\|x\|^2$  ( $\forall x \in \mathbb{R}^2$ ) it follows that [\(73\)](#) holds. To verify the second claim, set ( $\forall n \in \mathbb{N}$ )  $x_n = (n, 0)$ ,  $y_n = (n, n)$  and  $x_n^* = y_n^* = (2n, 0)$ . Observe that  $\{(x_n, x_n^*), (y_n, y_n^*)\} \subseteq \text{gra } A$ . Consequently, ( $\forall n \in \mathbb{N}$ )  $\frac{\|x_n^* - y_n^*\|}{\|x_n - y_n\|} = \frac{0}{n} = 0$ . The proof is complete. ■

**Example 4.8.** Suppose that  $S: \mathbb{R}^2 \rightarrow \mathbb{R}^2: (x_1, x_2) \rightarrow (-x_2, x_1)$  is the rotator by  $\pi/2$ . Then the following properties hold.

- (i)  $S$  and  $S^{-1} = -S$  are maximally monotone.
- (ii) Both  $S$  and  $S^{-1}$  are isometries, hence are Lipschitz continuous.
- (iii)  $\lim_{\|x-y\| \rightarrow \infty} \inf \frac{\|Sx - Sy\|}{\|x-y\|} = 1 > 0$ .
- (iv) Both  $S$  and  $S^{-1}$  are uniformly continuous.
- (v) Neither  $S$  nor  $S^{-1}$  are uniformly monotone, nor are they coercive.
- (vi) Both  $J_S$  and  $J_{S^{-1}}$  are strongly monotone.

*Proof.* [\(i\)](#): This is [\[2, Example 22.15\]](#). [\(ii\)](#): This is clear. [\(iii\)&\(iv\)](#): This is a direct consequence of [\(ii\)](#). [\(v\)](#): Indeed,  $S$  is neither uniformly monotone nor coercive because ( $\forall x \in \mathbb{R}^2$ )  $\langle Sx, x \rangle = 0$ . The same properties hold for  $S^{-1} = -S$ . [\(vi\)](#): Observe that  $J_S = (\text{Id} + S)^{-1}$  and so  $J_S = \frac{1}{2}(\text{Id} - S)$  and ( $\forall x \in \mathbb{R}^2$ )  $\langle J_S x, x \rangle = \frac{1}{2}\|x\|^2$ . Hence  $J_S$  is strongly monotone. Similarly, one verifies that ( $\forall x \in \mathbb{R}^2$ )  $\langle J_{S^{-1}} x, x \rangle = \frac{1}{2}\|x\|^2$ . ■

**Remark 4.9.**

- (i) In the convex function case,  $f$  is uniformly convex if and only if  $f^*$  has uniformly continuous derivative (see [\[28, Theorem 3.5.5 and Theorem 3.5.6\]](#)). [Example 4.8\(v\)](#) shows that this correspondence does not hold for general maximally monotone operators.
- (ii) [Example 4.8\(v\)&\(vi\)](#) also shows in a strong way that if  $J_{A^{-1}}$  is uniformly monotone (resp. coercive), it does not automatically follow that  $A$  is uniformly monotone (resp. coercive). This is in contrast to the situation for convex functions where the infimal convolution of  $\|\cdot\|^2$  and  $f$  is uniformly convex (resp. supercoercive) if and only if  $f$  is uniformly convex (resp. supercoercive). This follows by checking the conjugate of that infimal convolution is uniformly smooth if and only if  $f^*$  is. See [\[2, 8, 28\]](#) for more information on this.

## 5 Contractions for large distances

We start with the following lemma which provides a characterization of Banach contractions using averaged mappings.

**Lemma 5.1.** *Let  $T: X \rightarrow X$ . Then  $T$  is a Banach contraction if and only if [ $T$  is averaged and  $-T$  is averaged].*

*Proof.* ( $\Rightarrow$ ): Suppose that  $T$  is a Banach contraction. Observe that  $-T$  is also a Banach contraction. The conclusion follows from [2, Proposition 4.38].

( $\Leftarrow$ ): Suppose that both  $T$  and  $-T$  are averaged. It follows from [4, Proposition 4.3(ii)] that  $T = R_A$  and both  $A$  and  $A^{-1}$  are strongly monotone operators. Therefore, by [3, Corollary 4.7]  $T$  is a Banach contraction. ■

Before we proceed, we present the following useful result.

**Lemma 5.2.** *Let  $T: X \rightarrow X$ . Suppose that  $T$  and  $-T$  are strongly nonexpansive. Suppose that  $(x_n - y_n)_{n \in \mathbb{N}}$  is bounded and that  $\|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0$ . Then  $(x_n - y_n) \rightarrow 0$ .*

*Proof.* By assumption we have

$$(x_n - y_n) - (Tx_n - Ty_n) \rightarrow 0 \quad (74a)$$

$$(x_n - y_n) + (Tx_n - Ty_n) \rightarrow 0. \quad (74b)$$

Adding the above limits yields the desired result. ■

In an analogy to Lemma 5.1 we present the following result that characterizes contractions for large distances using either strongly nonexpansive mappings or super strongly nonexpansive mappings.

**Proposition 5.3.** *Suppose  $T: X \rightarrow X$  is a nonexpansive mapping. Then the following are equivalent:*

- (i) *Both  $T$  and  $-T$  are super strongly nonexpansive.*
- (ii) *Both  $T$  and  $-T$  are strongly nonexpansive.*
- (iii)  *$T$  is a contraction for large distances.*

*Proof.* (i)  $\Rightarrow$  (ii): Apply Proposition 3.2 to both  $T$  and  $-T$ .

(ii)  $\Rightarrow$  (iii): Fix  $\epsilon > 0$ , and define

$$\beta = \sup \left\{ \frac{\|Tx - Ty\|}{\|x - y\|} \mid \epsilon \leq \|x - y\| \leq 2\epsilon \right\}.$$

We claim that  $\beta < 1$ . Indeed, suppose for eventual contradiction that  $\beta = 1$ . Then there exist sequences  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  in  $X$  such that

$$(\forall n \in \mathbb{N}) \epsilon \leq \|x_n - y_n\| \leq 2\epsilon \quad \text{and} \quad \frac{\|Tx_n - Ty_n\|}{\|x_n - y_n\|} \rightarrow 1. \quad (75)$$



Because  $\epsilon \leq \|x_n - y_n\| \leq 2\epsilon$ , we have  $\|Tx_n - Ty_n\| - \|x_n - y_n\| \rightarrow 0$ . Therefore, it follows from [Lemma 5.2](#) that  $(x_n - y_n) \rightarrow 0$ , which is absurd since  $(\forall n \in \mathbb{N}) \|x_n - y_n\| \geq \epsilon$ . Therefore,  $0 \leq \beta < 1$ , and

$$\|Tx - Ty\| \leq \beta\|x - y\| \text{ whenever } \epsilon \leq \|x - y\| \leq 2\epsilon. \quad (76)$$

We next show  $(\forall k \in \{0, 1, 2, \dots\})$

$$\|Tx - Ty\| \leq \beta\|x - y\| \text{ whenever } 2^k\epsilon \leq \|x - y\| \leq 2^{k+1}\epsilon. \quad (77)$$

We proceed by induction. Indeed, (77) holds for  $k = 0$  by (76). Now suppose (77) holds for  $k = n$  where  $n \geq 0$ . Thus suppose  $\{x, y\} \subseteq X$  satisfies that  $2^{n+1}\epsilon \leq \|x - y\| \leq 2^{n+2}\epsilon$ . Now let  $z = (x + y)/2$ , then

$$x - z = \frac{x - y}{2} = z - y \text{ and so } 2^n\epsilon \leq \|x - z\| \leq 2^{n+1}\epsilon, 2^n\epsilon \leq \|z - y\| \leq 2^{n+1}\epsilon. \quad (78)$$

Because  $\|x - z\| = \|z - y\| = \frac{1}{2}\|x - y\|$ , the triangle inequality yields

$$\|Tx - Ty\| = \|Tx - Tz + Tz - Ty\| \leq \|Tx - Tz\| + \|Tz - Ty\| \quad (79a)$$

$$\leq \beta\|x - z\| + \beta\|z - y\| = \beta\|x - y\|. \quad (79b)$$

It follows by induction that (77) is true  $(\forall k \in \{0, 1, 2, \dots\})$ , and so  $T$  is a contraction for large distances.

(iii)  $\Rightarrow$  (i): Clearly  $T$  is a contraction for large distances if and only if  $-T$  is a contraction for large distances. Therefore it is sufficient to show the implication [ $T$  is a contraction for large distances  $\Rightarrow T$  is super strongly nonexpansive]. Suppose  $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$  are sequences in  $X$  such that

$$\|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0. \quad (80)$$

We claim that  $(x_n - y_n) \rightarrow 0$ . Indeed, suppose for eventual contradiction that  $\limsup \|x_n - y_n\| > 0$ . After passing to a subsequence and relabelling if necessary  $(\exists \epsilon > 0)$  such that  $(\forall n \in \mathbb{N}) \|x_n - y_n\| \geq \epsilon$ . This means  $(\exists \beta < 1)$  so that  $\|Tx_n - Ty_n\| \leq \beta\|x_n - y_n\|$ . Therefore,

$$(\forall n \in \mathbb{N}) \quad \|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \geq (1 - \beta^2)\|x_n - y_n\|^2 \geq (1 - \beta^2)\epsilon^2 \quad (81)$$

and this contradicts (80). Therefore  $\|x_n - y_n\| \rightarrow 0$  and consequently  $\|Tx_n - Ty_n\| \rightarrow 0$  which implies  $(x_n - y_n) - (Tx_n - Ty_n) \rightarrow 0$ . Thus  $T$  is super strongly nonexpansive.  $\blacksquare$

It is clear that every Banach contraction is a contraction for large distances. However, the opposite is not true as we illustrate in [Example 5.4](#) below.

**Example 5.4.** *Let*

$$T: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} 1, & x \geq \frac{\pi}{2}; \\ \sin x, & |x| < \frac{\pi}{2}; \\ -1, & \text{otherwise.} \end{cases} \quad (82)$$

*Then the following hold:*




Figure 2: A GeoGebra snapshot illustrating the mapping  $T$  in [Example 5.4](#).

- (i)  $T$  is nonexpansive.
- (ii)  $T$  is not a Banach contraction.
- (iii)  $T$  is a contraction for large distances.
- (iv) Both  $T$  and  $-T$  are super strongly nonexpansive.
- (v) Both  $T$  and  $-T$  are strongly nonexpansive.

*Proof.* Recall that (see, e.g., [6, Theorem 5.12]) if  $T: \mathbb{R} \rightarrow \mathbb{R}$  is differentiable then

$$T \text{ is Lipschitz continuous with a constant } K \geq 0 \text{ if and only if } (\forall x \in \mathbb{R}) |T'(x)| \leq K. \quad (83)$$

(i): One can directly verify that  $T$  is differentiable and that

$$T': \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} \cos x, & |x| < \frac{\pi}{2}; \\ 0, & \text{otherwise.} \end{cases} \quad (84)$$

Hence  $(\forall x \in \mathbb{R}) |T'(x)| \leq 1$ .

Consequently, by (83),  $T$  is Lipschitz continuous with a constant 1 and the conclusion follows.

(ii): Suppose for eventual contradiction that  $T$  is a Banach contraction. Then (83) implies that exists  $K \in [0, 1[$  such that  $(\forall x \in \mathbb{R}) |T'(x)| \leq K < 1$ . However,  $|T'(0)| = \cos 0 = 1 > K$ , which is absurd.

(iii): Let  $\epsilon > 0$ . Observe that if  $|t| \geq \epsilon/4$  then  $|T'(t)| \leq \alpha < 1$  where  $\alpha = |T'(\epsilon/4)|$ . We choose  $\beta = (\alpha + 3)/4$ . Now suppose  $|x - y| \geq \epsilon$ , where  $x < y$ . In the case  $|y| \geq |x|$ , we have  $y \geq |x - y|/2 \geq \epsilon/2$ . Therefore, by the Fundamental theorem of calculus, we write

$$|Tx - Ty| = \left| \int_x^y T'(t) dt \right| \leq \int_x^{|x-y|/4} 1 dt + \int_{|x-y|/4}^y \alpha \leq \beta|x - y|. \quad (85)$$

Similarly, if  $|x| \geq |y|$ , one has  $x \leq -|x - y|/2$ , and again,  $|Tx - Ty| \leq \beta|x - y|$ . Therefore,  $T$  is a contraction for large distances. (iv)–(v): Combine (iii) with [Proposition 5.3](#). ■

The next result and more general variations of it, are well-known in fixed point theory, see, for example, [20, Corollary, p. 463] and [1, Theorem 2.1]. Nevertheless, we include a simple proof based on [Theorem 4.5\(ii\)](#) for completeness.

**Proposition 5.5.** *Let  $T: X \rightarrow X$  be a contraction for large distances. Let  $x_0 \in X$  and set  $(\forall n \in \mathbb{N}) x_n = T^n x_0$ . Then  $(\exists \bar{x} \in X)$  such that the following hold:*

- (i)  $\text{Fix } T = \{\bar{x}\}$ .
- (ii)  $x_n \rightarrow \bar{x}$ .

*Proof.* (i): On the one hand because  $T$  is nonexpansive,  $T = R_A$  for some maximally monotone operator  $A: X \rightrightarrows X$  (see [2, Corollary 23.11, Proposition 4.4]). On the other hand, because  $-T = -R_A$  is a contraction for large distances it is super strongly nonexpansive by [Proposition 5.3](#). Therefore,  $A$  is uniformly monotone by [Proposition 3.5](#). Consequently, by [Theorem 4.5\(iii\)](#),  $A$  has a unique zero. Now combine this with (6).

(ii): Note that  $(\|x_n - \bar{x}\|)_{n \in \mathbb{N}}$  converges by, e.g., [2, Proposition 5.4(ii)]. Now, suppose by way of contradiction that  $(x_n)_{n \in \mathbb{N}}$  does not converge in norm to  $\bar{x}$ . Then  $\lim_{n \in \mathbb{N}} \|x_n - \bar{x}\| = \epsilon$  where  $\epsilon > 0$ . Thus we choose  $0 < \beta < 1$  so that

$$(\forall n \in \mathbb{N}) \quad \|x_{n+1} - \bar{x}\| = \|Tx_n - T\bar{x}\| \leq \beta \|x_n - \bar{x}\|. \quad (86)$$

Now for  $n$  sufficiently large, we have  $\|x_n - \bar{x}\| < \epsilon/\beta$ . Then

$$\|x_{n+1} - \bar{x}\| \leq \beta \|x_n - \bar{x}\| < \epsilon. \quad (87)$$

This contradiction completes the proof. Alternatively, use [Lemma 5.2](#) with  $(x_n, y_n)_{n \in \mathbb{N}}$  replaced by  $(T^n x_0, \bar{x})_{n \in \mathbb{N}}$  to conclude that  $T^n x_0 - \bar{x} \rightarrow 0$ . ■

## 6 Self-Dual Properties on Hilbert Spaces

**Lemma 6.1.** *Suppose  $C: X \rightarrow X$  is uniformly continuous. Then for each  $\epsilon > 0$  ( $\exists M > 0$ ) depending on  $\epsilon$  so that*

$$\|x - y\| \leq M \|u - v\| \quad \text{whenever} \quad \|x - y\| \geq \epsilon, \quad (u, v) \in J_C x \times J_C y. \quad (88)$$

*Proof.* Let  $\epsilon > 0$ . By the uniform continuity of  $C$  we choose  $0 < \delta < \epsilon/2$  so that

$$\|Cu - Cv\| < \frac{\epsilon}{2} \quad \text{whenever} \quad \|u - v\| < \delta. \quad (89)$$

Because  $C$  is uniformly continuous, it is Lipschitz for large distances (see [7, Proposition 1.11]). Thus we choose  $K > 0$  so that

$$\|Cu - Cv\| \leq K \|u - v\| \quad \text{whenever} \quad \|u - v\| \geq \delta. \quad (90)$$

Now let us suppose

$$\|x - y\| \geq \epsilon, \quad u \in J_C x, \quad v \in J_C y. \quad (91)$$

We will show that  $\|x - y\| \leq M\|u - v\|$  where  $M = K + 1$ . First, we verify that  $\|u - v\| \geq \delta$  where  $\delta$  is from (89) by way of a contradiction. So let us assume to the contrary that  $\|u - v\| < \delta$ . Then by (89) we have

$$\|Cu - Cv\| < \frac{\epsilon}{2} \quad (92)$$

Because  $u \in J_C x$  and  $v \in J_C y$ , this implies  $Cu = x - u$  and  $Cv = y - v$ . Then

$$\|x - u - (y - v)\| < \frac{\epsilon}{2} \Rightarrow \|x - y\| - \|u - v\| < \frac{\epsilon}{2} \Rightarrow \|x - y\| < \frac{\epsilon}{2} + \delta < \epsilon \quad (93)$$

This contradicts (91), and so  $\|u - v\| \geq \delta$ . Therefore, using (90), one has

$$\begin{aligned} \|x - y\| &= \|u + Cu - (v + Cv)\| \leq \|u - v\| + \|Cu - Cv\| \\ &\leq \|u - v\| + K\|u - v\| = M\|u - v\|, \end{aligned}$$

as desired. ■

**Theorem 6.2.** *The following hold:*

- (i) *Suppose  $A$  is uniformly monotone and uniformly continuous. Then  $R_A$  is a contraction for large distances.*
- (ii) *Suppose  $R_A$  is a contraction for large distances. Then  $A$  is uniformly monotone with a supercoercive modulus.*

*Proof. (i):* Let  $\phi$  be a modulus function for  $A$ , let  $\epsilon > 0$  and suppose  $\|x - y\| \geq \epsilon$ . On the one hand, it follows from Lemma 6.1 that  $(\exists K > 0)$  such that  $\|J_A x - J_A y\| \geq K\|x - y\| \geq K\epsilon$ . On the other hand, because  $\text{dom } A = X$ , Proposition 4.1 implies that  $(\exists \alpha > 0)$  such that  $(\forall t \geq K\epsilon) \phi(t) \geq \alpha t^2$ . Altogether, we learn that

$$\phi\left(\frac{1}{2}\|J_A x - J_A y\|\right) \geq \frac{\alpha K^2}{4}\|x - y\|^2. \quad (94)$$

Set  $\beta = \sqrt{\alpha}K$  and let  $(x, y) \in X \times X$ . Combining (94) and Lemma 3.4 we learn that  $\|x - y\| \geq \epsilon \Rightarrow \|R_A x - R_A y\|^2 + \beta^2\|x - y\|^2 \leq \|x - y\|^2$ ; equivalently,  $\|x - y\| \geq \epsilon \Rightarrow \|R_A x - R_A y\| \leq (1 - \beta)\|x - y\|$ . That is,  $R_A$  is a contraction for large distances. (ii): Let  $\epsilon > 0$  and let  $(x, x^*) \in \text{gra } A$ ,  $(y, y^*) \in \text{gra } A$ . Suppose that  $\|x - y\| \geq \epsilon$ . Set  $(u, v) = (x + x^*, y + y^*)$  and observe that (5) implies

$$(x, x^*) = \frac{1}{2}(u + R_A u, u - R_A u) \quad \text{and} \quad (y, y^*) = \frac{1}{2}(v + R_A v, v - R_A v). \quad (95)$$

It follows from (95), the nonexpansiveness of  $R_A$ , and the triangle inequality that

$$\|x - y\| = \frac{1}{2}\|u - v - (R_A u - R_A v)\| \leq \frac{1}{2}(\|u - v\| + \|R_A u - R_A v\|) \leq \|u - v\|. \quad (96)$$

Hence  $\|u - v\| \geq \epsilon$ . Consequently, because  $R_A$  is a contraction for large distances,  $(\exists \beta \in ]0, 1[)$  such that

$$\|R_A u - R_A v\| \leq \beta\|u - v\|. \quad (97)$$

Using (95) and (97) we learn that

$$\langle x - y, x^* - y^* \rangle = \frac{1}{4} \langle u - v + R_A u - R_A v, u - v - (R_A u - R_A v) \rangle \quad (98a)$$

$$= \frac{1}{4} (\|u - v\|^2 - \|R_A u - R_A v\|^2) \quad (98b)$$

$$\geq \frac{1}{4} (1 - \beta^2) \|u - v\|^2 \quad (98c)$$

$$\geq \frac{1}{4} (1 - \beta^2) \|x - y\|^2. \quad (98d)$$

Therefore, for  $t \geq \epsilon$ , we have

$$\inf \{ \langle x - y, x^* - y^* \rangle \mid (x, x^*) \in \text{gr}(A), (y, y^*) \in \text{gr}(A), \|x - y\| \geq t \} \geq \frac{1}{4} (1 - \beta^2) t^2. \quad (99)$$

That is,  $A$  has a modulus  $\phi$  satisfying  $\phi(t) \geq \frac{1}{4} (1 - \beta^2) t^2$  for  $t \geq \epsilon$ . ■

This brings us to our main duality result of this section.

**Theorem 6.3.** *The following are equivalent.*

- (i)  $A$  is uniformly monotone and uniformly continuous.
- (ii)  $R_A$  is a contraction for large distances.
- (iii) Both  $A$  and  $A^{-1}$  are uniformly monotone with supercoercive moduli.
- (iv) Both  $A$  and  $A^{-1}$  are uniformly monotone.
- (v)  $A^{-1}$  is uniformly monotone and uniformly continuous.
- (vi)  $R_A$  and  $R_{A^{-1}}$  are strongly nonexpansive.

*Proof.* (i)  $\Rightarrow$  (ii): [Theorem 6.2\(i\)](#).

(ii)  $\Rightarrow$  (iii): Since  $R_A$  is a contraction for large distances, so is  $R_{A^{-1}} = -R_A$ . Thus this follows by applying [Theorem 6.2\(ii\)](#) on  $R_A$  and on  $R_{A^{-1}}$ .

(iii)  $\Rightarrow$  (iv): is immediate, and (iv)  $\Rightarrow$  (v): follows from [Theorem 4.5](#) applied to  $A$  to deduce  $A^{-1}$  is uniformly continuous.

(v)  $\Rightarrow$  (i): The above implications show (i)  $\Rightarrow$  (v), so the reverse implication follows by applying (i)  $\Rightarrow$  (v) to the operator  $A^{-1}$ . (ii)  $\Leftrightarrow$  (vi): This is a direct consequence of [Proposition 5.3](#) applied with  $T$  replaced by  $R_A$ . ■

Let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous and proper. Recall that (see, e.g., [2, Corollary 16.30])

$$\partial f^* = (\partial f)^{-1}, \quad (100)$$

that<sup>2</sup> (see [28, Theorem 3.5.12])

$$f \text{ is uniformly convex} \Leftrightarrow f^* \text{ is uniformly smooth} \quad (101a)$$

$$\Leftrightarrow \partial f \text{ is uniformly monotone}, \quad (101b)$$

---

<sup>2</sup>Let  $f: X \rightarrow \mathbb{R}$  be convex. Then  $f$  is uniformly smooth if  $f$  is smooth and  $\nabla f$  is uniformly continuous.

and that (see, e.g., [2, Theorem 18.15] )

$$f \text{ is strongly convex} \Leftrightarrow f^* \text{ is differentiable and } \nabla f^* \text{ is Lipschitz continuous} \quad (102a)$$

$$\Leftrightarrow \partial f \text{ is monotone.} \quad (102b)$$

**Example 6.4.** *Let*

$$f: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} 4x^2 - 2, & x \leq -1; \\ 2x^4, & -1 < x < 0; \\ x^{3/2}, & 0 \leq x < 1; \\ \frac{3}{4}x^2 + \frac{1}{4}, & \text{otherwise.} \end{cases} \quad (103)$$

Then  $f$  is differentiable and  $f'$  is continuous and increasing. Moreover the following hold:

- (i) Both  $f$  and  $f^*$  are uniformly convex.
- (ii) Both  $f$  and  $f^*$  are uniformly smooth.
- (iii) Both  $f'$  and  $(f^*)'$  are uniformly monotone.
- (iv) Both  $f'$  and  $(f^*)'$  are uniformly continuous.
- (v) Neither  $f$  nor  $f^*$  is strongly convex.

*Proof.* It is straightforward to verify that  $f$  is differentiable and that

$$f': \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} 8x, & x \leq -1; \\ 8x^3, & -1 < x < 0; \\ \frac{3}{2}x^{1/2}, & 0 \leq x < 1; \\ \frac{3}{2}x, & \text{otherwise.} \end{cases} \quad (104)$$

Hence,  $f'$  is continuous and strictly increasing as claimed. (i): It follows from [27, Theorem 3.1] that  $f$  is uniformly convex on bounded sets. It follows from the strong (hence uniform) convexity of  $x^2$  that  $f$  is uniformly convex. We now prove the uniform convexity of  $f^*$ . In view of (101) it suffices to verify that  $f$  is uniformly smooth which can be easily deduced from (104).

(ii)&(iii): Combine (i) and (101). (iv): This follows from (ii).

(v): On the one hand  $2x^4$  is *not* strongly convex, hence  $f$  is *not* strongly convex. On the other hand,  $(x^{3/2})' = \frac{3}{2}\sqrt{x}$  is *not* Lipschitz continuous on  $]0, 1[$ . Therefore, in view of (102) applied with  $f$  replaced by  $f^*$  we conclude that  $f^*$  is *not* strongly convex. ■

**Example 6.5.** *Let*

$$T: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} 1, & x \geq \frac{\pi}{2}; \\ \sin x, & |x| < \frac{\pi}{2}; \\ -1, & \text{otherwise.} \end{cases} \quad (105)$$

Set  $A = \left(\frac{\text{Id}-T}{2}\right)^{-1} - \text{Id}$ . Then there exists a proper lower semicontinuous convex function  $f: \mathbb{R} \rightarrow ]-\infty, +\infty]$  such that

$$A = f'. \quad (106)$$

Moreover, the following hold:

- (i) Both  $A$  and  $A^{-1}$  are maximally monotone and uniformly monotone with supercoercive modulus.
- (ii) Both  $A$  and  $A^{-1}$  are uniformly continuous.
- (iii) Both  $A$  and  $A^{-1}$  are not strongly monotone.
- (iv) Both  $f$  and  $f^*$  are uniformly convex and uniformly smooth.
- (v) Both  $f$  and  $f^*$  are not strongly convex.

*Proof.* It is clear that  $T = -R_A$ . Because  $T$  is nonexpansive, from [Example 5.4\(i\)](#) we learn that  $A$  is maximally monotone. Therefore, by e.g., [\[2, Corollary 22.23\]](#) there exists a proper lower semicontinuous convex function  $f: \mathbb{R} \rightarrow ]-\infty, +\infty]$  such that  $A = \partial f$ . Finally, we show in [\(iv\)](#) below that  $f$  is uniformly smooth, hence  $A = f'$ .

[\(i\)&\(ii\)](#): Combine [Example 5.4\(iii\)](#), and [Theorem 6.3](#).

[\(iii\)](#): Suppose for eventual contradiction that  $A$  is strongly monotone. Then by [\[4, Proposition 4.3\(ii\)\]](#) ( $\exists \alpha \in ]0, 1[$ ) such that  $-R_A$  is  $\alpha$ -averaged. It follows from [\[2, Proposition 4.35\]](#) that  $(\forall (x, y) \in \mathbb{R} \times \mathbb{R}) \frac{1-\alpha}{\alpha} ((\text{Id}+T)x - (\text{Id}+T)y)^2 \leq (x-y)^2 - (Tx - Ty)^2$ . In particular, for  $y = 0$  the above inequality yields  $(\forall x \in ]0, \frac{\pi}{2}[$ )

$$\frac{1-\alpha}{\alpha} (x + \sin x)^2 \leq x^2 - \sin^2 x. \quad (107)$$

Simplifying and multiplying both sides of the above inequality by  $\alpha$  we obtain  $(1-2\alpha)x^2 + 2(1-\alpha)x \sin x + \sin^2 x \leq 0$ . Rearranging yields  $(x + \sin x)^2 \leq 2\alpha x(x + \sin x)$ . Because  $x \in ]0, \frac{\pi}{2}[$ , it follows that  $x + \sin x > 0$  and therefore the last inequality is equivalent to  $1 + \frac{\sin x}{x} \leq 2\alpha$ . Taking the limit as  $x \rightarrow 0^+$  we learn that  $2 \leftarrow 1 + \frac{\sin x}{x} \leq 2\alpha$  which is absurd. Hence,  $-R_A$  is not averaged; equivalently,  $A$  is not strongly monotone as claimed.

Using similar argument, one can show that  $R_A = -R_{A^{-1}}$  is not averaged; equivalently,  $A^{-1}$  is not strongly nonexpansive as claimed.

[\(iv\)](#): Combine [\(ii\)](#), [\(106\)](#) and [\[27, Theorem 3.5.10\]](#) in view of [\(100\)](#).

[\(v\)](#): Combine [\(iii\)](#), [\(106\)](#) and [\[2, Example 22.4\(iv\)\]](#). ■

**Remark 6.6.** In [Appendix A](#) we provide finer conclusions about the operator  $A$  and the function  $f$  introduced in [Example 6.5](#).

**Example 6.7.** Let  $f: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \frac{1}{4}x^4$  and let  $A = f'$ . Let  $a: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \sqrt[3]{108x + 12\sqrt{81x^2 + 12}}$  and set

$$T: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto x - 2 \left( \frac{a(x)}{6} - \frac{2}{a(x)} \right). \quad (108)$$

Then the following hold:

- (i)  $T = -R_A$ .
- (ii)  $A = x^3$  is maximally monotone but not uniformly continuous.
- (iii)  $A^{-1} = \sqrt[3]{x}$  is maximally monotone and uniformly continuous.
- (iv)  $A$  is uniformly monotone.

- (v)  $A^{-1}$  is not uniformly monotone.
- (vi)  $f$  is uniformly convex.
- (vii)  $f^*$  is not uniformly convex.
- (viii)  $T$  is super strongly nonexpansive and strongly nonexpansive.
- (ix)  $-T$  is neither super strongly nonexpansive nor strongly nonexpansive.

*Proof.* (i): It is enough to show that  $(\forall x \in \mathbb{R}) J_A(x) = \frac{a(x)}{6} - \frac{2}{a(x)}$ , equivalently, to show that  $(\text{Id} + A)\left(\frac{a(x)}{6} - \frac{2}{a(x)}\right) = x$ . Indeed,

$$(\text{Id} + A)\left(\frac{a(x)}{6} - \frac{2}{a(x)}\right) = \frac{a(x)}{6} - \frac{2}{a(x)} + \frac{(a(x))^3}{6^3} - \frac{a(x)}{6} + \frac{2}{a(x)} - \frac{2^3}{(a(x))^3} \quad (109a)$$

$$= \frac{(a(x))^3}{6^3} - \frac{2^3}{(a(x))^3} = \frac{(a(x))^6 - 12^3}{6^3(a(x))^3} = x. \quad (109b)$$

(ii)&(iii): This is clear.

(iv)–(vii): Combine (ii)&(iii) with [27, Theorem 3.5.10].

(viii): Combine (i), (iv) and Corollary 3.6.

(ix): Combine (i), (v) and Corollary 3.6. ■

## 7 Compositions

In this section we examine the behaviour of strongly nonexpansive mappings, super strongly nonexpansive mappings and contractions for large distances under structured compositions. The proof of the next result follows along the lines of the proof of [12, Proposition 1.1].

**Proposition 7.1.** *Let  $T_1: X \rightarrow X$  and  $T_2: X \rightarrow X$  be nonexpansive. Set  $T = T_2T_1$ . Then the following hold:*

- (i) *Suppose that  $T_1$  is strongly nonexpansive and  $T_2$  is strongly nonexpansive. Then  $T$  is strongly nonexpansive.*
- (ii) *Suppose that  $-T_1$  is strongly nonexpansive and  $-T_2$  is strongly nonexpansive. Then  $T$  is strongly nonexpansive.*
- (iii) *Suppose that  $-T_1$  is strongly nonexpansive and  $T_2$  is strongly nonexpansive. Then  $-T$  is strongly nonexpansive.*
- (iv) *Suppose that  $T_1$  is strongly nonexpansive and  $-T_2$  is strongly nonexpansive. Then  $-T$  is strongly nonexpansive.*

*Proof.* (i): This is [12, Proposition 1.1]. (ii): Clearly  $T$  is nonexpansive. Now suppose that  $(x_n - y_n)_{n \in \mathbb{N}}$  is bounded and that  $\|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0$ . Observe that the nonexpansiveness of



$T_1, T_2$  and, consequently,  $T$  implies that

$$0 \leq \|x_n - y_n\| - \|Tx_n - Ty_n\| = \underbrace{\|x_n - y_n\| - \|T_1x_n - T_1y_n\|}_{\geq 0} + \underbrace{\|T_1x_n - T_1y_n\| - \|Tx_n - Ty_n\|}_{\geq 0} \rightarrow 0. \quad (110)$$

Consequently, we learn that

$$\|x_n - y_n\| - \|(-T_1)x_n - (-T_1)y_n\| = \|x_n - y_n\| - \|T_1x_n - T_1y_n\| \rightarrow 0, \quad (111a)$$

$$\|T_1x_n - T_1y_n\| - \|(-T_2)(T_1x_n) - (-T_2)(T_1y_n)\| = \|T_1x_n - T_1y_n\| - \|Tx_n - Ty_n\| \rightarrow 0 \quad (111b)$$

The nonexpansiveness of  $T_1$  implies that  $\|T_1x_n - T_1y_n\|$  is bounded. Recalling that  $-T_1$  is strongly nonexpansive and  $-T_2$  is strongly nonexpansive we obtain

$$(x_n - y_n) + (T_1x_n - T_1y_n) \rightarrow 0 \quad (112a)$$

$$(T_1x_n - T_1y_n) + (T_2T_1x_n - T_2T_1y_n) \rightarrow 0. \quad (112b)$$

Hence,

$$(x_n - y_n) - (Tx_n - Ty_n) = (x_n - y_n) + (T_1x_n - T_1y_n) - ((T_1x_n - T_1y_n) + (T_2T_1x_n - T_2T_1y_n)) \rightarrow 0. \quad (113)$$

(iii): Proceeding similar to (ii) we learn that

$$(x_n - y_n) + (T_1x_n - T_1y_n) \rightarrow 0 \quad (114a)$$

$$(T_1x_n - T_1y_n) - (T_2T_1x_n - T_2T_1y_n) \rightarrow 0. \quad (114b)$$

Hence,

$$(x_n - y_n) + (Tx_n - Ty_n) = (x_n - y_n) + (T_1x_n - T_1y_n) - ((T_1x_n - T_1y_n) - (T_2T_1x_n - T_2T_1y_n)) \rightarrow 0. \quad (115)$$

(iv): Proceed similar to (iii). ■

The following analogous result holds for super strongly nonexpansive mappings.

**Proposition 7.2.** *Let  $T_1: X \rightarrow X$  and  $T_2: X \rightarrow X$  be nonexpansive. Set  $T = T_2T_1$ . Then the following hold:*

- (i) *Suppose that  $T_1$  is super strongly nonexpansive and  $T_2$  is super strongly nonexpansive. Then  $T$  is super strongly nonexpansive.*
- (ii) *Suppose that  $-T_1$  is super strongly nonexpansive and  $-T_2$  is super strongly nonexpansive. Then  $T$  is super strongly nonexpansive.*
- (iii) *Suppose that  $-T_1$  is super strongly nonexpansive and  $T_2$  is super strongly nonexpansive. Then  $-T$  is super strongly nonexpansive.*
- (iv) *Suppose that  $T_1$  is super strongly nonexpansive and  $-T_2$  is super strongly nonexpansive. Then  $-T$  is super strongly nonexpansive.*

*Proof. (i):* Let  $(x, y) \in X \times X$  and suppose that  $(x_n)_{n \in \mathbb{N}}$  and  $(y_n)_{n \in \mathbb{N}}$  are sequences in  $X$  such that  $0 \leq \|x_n - y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0$ . Rewrite the above limit as

$$0 \leq \|x_n - y_n\|^2 - \|T_1x_n - T_1y_n\|^2 + \|T_1x_n - T_1y_n\|^2 - \|Tx_n - Ty_n\|^2 \rightarrow 0, \quad (116)$$

and observe that the nonexpansiveness of  $T_1$  and  $T_2$  implies

$$\|x_n - y_n\|^2 - \|T_1 x_n - T_1 y_n\|^2 \rightarrow 0 \quad \text{and} \quad \|T_1 x_n - T_1 y_n\|^2 - \|T x_n - T y_n\|^2 \rightarrow 0. \quad (117)$$

Because  $T_1$  and  $T_2$  are super strongly nonexpansive we learn that

$$(x_n - y_n) - (T_1 x_n - T_1 y_n) \rightarrow 0 \quad (118a)$$

$$(T_1 x_n - T_1 y_n) - (T x_n - T y_n) \rightarrow 0. \quad (118b)$$

Adding (118a) and (118b) yields  $(x_n - y_n) - (T x_n - T y_n) \rightarrow 0$ , hence  $T$  is super strongly nonexpansive as claimed. (ii)–(iv): Proceed similar to the proof of Proposition 7.1(ii)–(iv). ■

We now turn to compositions of finitely many mappings each of which is either strongly nonexpansive or its negative is strongly nonexpansive.

**Theorem 7.3.** *Let  $m \geq 2$ , let  $I = \{1, \dots, m\}$ , let  $J \subseteq I$ , and let  $(T_i)_{i \in I}$  be a family of nonexpansive mappings from  $X$  to  $X$ . Suppose that  $(\forall i \in I \setminus J)$   $T_i$  is strongly nonexpansive and that  $(\forall j \in J)$   $-T_j$  is strongly nonexpansive. Set*

$$T = T_m \dots T_1. \quad (119)$$

Then  $(-1)^{|J|} T$  is strongly nonexpansive.

*Proof.* We proceed by induction on  $k \in \{2, \dots, m\}$ . To this end, let us set  $(\forall k \in \{2, \dots, m\})$   $J_k = \{j \in J \mid j \leq k\}$ . By Proposition 7.1 the claim is true for  $k = 2$ . Now assume that, for some  $k \in \{2, \dots, m\}$ , we have  $(-1)^{|J_k|} T_k \dots T_1$  is strongly nonexpansive. If  $T_{k+1}$  is strongly nonexpansive then  $|J_{k+1}| = |J_k|$  and the conclusion follows by applying Proposition 7.1(i) (respectively Proposition 7.1(iii)) with  $(T_1, T_2)$  replaced by  $(T_m \dots T_1, T_{m+1})$  in the case  $|J_k|$  is even (respectively  $|J_k|$  is odd). If  $-T_{k+1}$  is strongly nonexpansive then  $|J_{k+1}| = |J_k| + 1$  and the conclusion follows by applying Proposition 7.1(ii) (respectively Proposition 7.1(iv)) with  $(T_1, T_2)$  replaced by  $(T_m \dots T_1, T_{m+1})$  in the case  $|J_k|$  is even (respectively  $|J_k|$  is odd). ■

**Theorem 7.4.** *Let  $m \geq 2$ , let  $I = \{1, \dots, m\}$ , let  $J \subseteq I$ , and let  $(T_i)_{i \in I}$  be a family of nonexpansive mappings from  $X$  to  $X$ . Suppose that  $(\forall i \in I \setminus J)$   $T_i$  is super strongly nonexpansive and that  $(\forall j \in J)$   $-T_j$  is super strongly nonexpansive. Set*

$$T = T_m \dots T_1. \quad (120)$$

Then  $(-1)^{|J|} T$  is super strongly nonexpansive.

*Proof.* Proceed similar to the proof of Theorem 7.3 but use Proposition 7.2(i)–(iv) instead of Proposition 7.1(i)–(iv). ■

We conclude this section with the following result concerning compositions that involve contractions for large distances.

**Proposition 7.5.** *Let  $m \geq 2$ , let  $I = \{1, \dots, m\}$ , let  $\bar{m} \in I$ , and let  $(T_i)_{i \in I}$  be a family of nonexpansive mappings from  $X$  to  $X$  and suppose that  $T_{\bar{m}}$  is a contraction for large distances. Set*

$$T = T_m \dots T_1. \quad (121)$$

Then  $T$  is a contraction for large distances.

*Proof.* Let  $(x, y) \in X \times X$  and let  $\epsilon > 0$ . Suppose that  $\|x - y\| \geq \epsilon$ . We proceed by induction on  $k \in \{2, \dots, m\}$ . For  $m = 2$  we examine two cases: CASE 1:  $T_1$  is a contraction for large distances. Then  $(\exists \beta_\epsilon \in ]0, 1[)$  such that  $\|T_1x - T_1y\| \leq \beta_\epsilon \|x - y\|$ . Therefore, because  $T_2$  is nonexpansive we learn that  $\|Tx - Ty\| \leq \|T_1x - T_1y\| \leq \beta_\epsilon \|x - y\|$ . CASE 2:  $T_2$  is a contraction for large distances. If  $\|T_1x - T_1y\| \geq \frac{\epsilon}{2}$ . Then  $(\exists \alpha_\epsilon \in ]0, 1[)$  such that  $\|Tx - Ty\| \leq \alpha_{\epsilon/2} \|x - y\|$ . Now suppose that  $\|T_1x - T_1y\| < \frac{\epsilon}{2}$ . Then  $\|Tx - Ty\| \leq \|T_1x - T_1y\| < \frac{\epsilon}{2} \leq \frac{1}{2} \|x - y\|$ . Setting  $\beta_\epsilon = \max\{\alpha_{\epsilon/2}, \frac{1}{2}\}$  proves the claim for  $m = 2$ . Now assume that, for some  $k \in \{2, \dots, m\}$ , we have  $T_k \dots T_1$  is a contraction for large distances whenever  $T_{\bar{k}}$  is a contraction for large distances,  $\bar{k} \in \{1, \dots, k\}$ . Consider the composition  $T_{k+1}T_k \dots T_1$ . If  $(\exists \bar{k} \in \{1, \dots, k\})$  such that  $T_{\bar{k}}$  is a contraction for large distances then the inductive hypothesis implies that  $T_k \dots T_1$  is a contraction for large distances. Otherwise,  $T_{k+1}$  must be a contraction for large distances. In both cases the conclusion follows from applying the base case with  $(T_1, T_2)$  replaced by  $(T_k \dots T_1, T_{k+1})$ . The proof is complete. ■

## 8 Application to splitting algorithms

In this section we use our earlier conclusions to obtain stronger and more refined convergence results for some important splitting methods (see, e.g., [2, Chapter 26]); namely, Peaceman–Rachford algorithm (see [Theorem 8.1\(i\)\(c\)–\(i\)\(d\)&\(ii\)](#) below), Douglas–Rachford algorithm (see [Theorem 8.6\(ii\)](#) below) and forward-backward algorithm (see [Theorem 8.7\(ii\)](#) below).

Let  $C : X \rightrightarrows X$  be uniformly monotone and suppose that  $\text{zer } C \neq \emptyset$ . Then  $C$  is strictly monotone and it follows from, e.g., [2, Proposition 23.35] that

$$\text{zer } C \text{ is a singleton.} \quad (122)$$

**Theorem 8.1 (Peaceman–Rachford algorithm).** *Suppose that  $A$  is uniformly monotone. Set  $T = R_B R_A$ . Let  $x_0 \in X$  and set  $(\forall n \in \mathbb{N})$ :*

$$x_{n+1} = Tx_n, \quad (123a)$$

$$y_n = J_A x_n. \quad (123b)$$

*Then the following hold.*

- (i) *Suppose that  $\text{zer}(A + B) \neq \emptyset$ . Then we have:*
  - (a)  *$\text{zer}(A + B)$  is a singleton and  $\text{Fix } T \neq \emptyset$ .*
  - (b)  *$(\exists \bar{x} \in \text{Fix } T)$   $(y_n)_{n \in \mathbb{N}}$  converges strongly to  $J_A \bar{x}$  and  $\text{zer}(A + B) = \{J_A \bar{x}\}$ .*
- If, in addition,  $B$  is uniformly monotone then we also have:*
  - (c)  *$T$  is strongly nonexpansive.*
  - (d)  *$(x_n)_{n \in \mathbb{N}}$  converges weakly to  $\bar{x}$ .*
- (ii) *Suppose that  $A^{-1}$  is uniformly monotone. Then we have:*
  - (a)  *$T$  is a contraction for large distances and  $(\exists \bar{x} \in X)$  such that  $\text{Fix } T = \{\bar{x}\}$ .*
  - (b)  *$(x_n)_{n \in \mathbb{N}}$  converges strongly to  $\bar{x}$ .*

*Proof.* (i)(a): Applying (122) with  $C$  replaced by  $A + B$ , we conclude that  $\text{zer}(A + B)$  is a singleton. It follows from [2, Proposition 26.1(iii)(b)] that  $\text{Fix } T \neq \emptyset$ . (i)(b): This is [2, Proposition 26.13].

(i)(c): It follows from Proposition 3.5 applied to  $A$  (respectively  $B$ ) that  $-R_A$  and (respectively  $-R_B$ ) is strongly nonexpansive. Consequently,  $R_B R_A$  is strongly nonexpansive by Proposition 7.1(ii) applied with  $(T_1, T_2)$  replaced by  $(R_A, R_B)$ .

(i)(d): Combine (i)(c) with Fact 2.6.

(ii)(a)&(ii)(b): It follows from Theorem 6.3 that  $R_A$  is a contraction for large distances. Combining this with Proposition 7.5 applied with  $(m, T_1, T_2)$  replaced by  $(2, R_A, R_B)$  we learn that  $T$  is a contraction for large distances. Now combine this with Proposition 5.5(i)&(ii). ■

The assumption that  $B$  is uniformly monotone is critical in the conclusion of Theorem 8.1(i)(d) as we illustrate below.

**Example 8.2.** Suppose that  $X \neq \{0\}$ . Let  $A = N_{\{0\}}$  and let  $B \equiv 0$ . Then  $A$  is strongly monotone, hence uniformly monotone, and  $B$  is not uniformly monotone. Moreover,  $R_A = -\text{Id}$ ,  $R_B = \text{Id}$ . Consequently  $T = R_B R_A = -\text{Id}$ . Let  $x_0 \in X \setminus \{0\}$ . Then  $(\forall n \in \mathbb{N}) T^n x_0 = (-1)^n x_0$  and  $(T^n x_0)_{n \in \mathbb{N}}$  does not converge.

The assumption that  $A^{-1}$  is uniformly monotone is critical in the conclusion of Theorem 8.1(ii)(b) as we illustrate below.

**Example 8.3.** Suppose that  $X = \ell^2(\{1, 2, 3, \dots\})$  with the standard Schauder basis  $e_1 = (1, 0, \dots)$ ,  $e_2 = (0, 1, 0, \dots)$ , and so on. Let  $A = N_{\{0\}}$ , let  $R: X \rightarrow X: (x_1, x_2, \dots) \mapsto (0, x_1, x_2, \dots)$  (the right shift operator) and set  $B = \frac{1}{2}(\text{Id} - R)^{-1} - \text{Id}$ . Then  $A$  is strongly monotone, hence uniformly monotone. Moreover, because  $R_B = -R$  is nonexpansive, we conclude that  $B$  is maximally monotone by Fact 2.5. Observe that  $A^{-1} \equiv 0$  is not uniformly monotone. Moreover,  $R_A = -\text{Id}$ ,  $R_B = -R$ . Consequently  $T = R_B R_A = R$ . Let  $x_0 = e_1$ . Then  $(\forall n \in \mathbb{N}) T^n x_0 = e_{n+1}$  and  $(T^n x_0)_{n \in \mathbb{N}} = (e_1, e_2, \dots)$  converges weakly but not strongly to 0.

We now turn to Douglas–Rachford algorithm. We recall the following fact.

**Fact 8.4.** Let  $T_1: X \rightarrow X$ , let  $T_2: X \rightarrow X$  and let  $\lambda \in ]0, 1[$ . Set  $T = (1 - \lambda)T_1 + \lambda T_2$ . Suppose that  $T_1$  is strongly nonexpansive and that  $T_2$  is nonexpansive. Then  $T$  is strongly nonexpansive.

*Proof.* See [12, Proposition 1.3]. ■

**Proposition 8.5.** Let  $R: X \rightarrow X$  and let  $\lambda \in ]0, 1[$ . Set  $T = (1 - \lambda)\text{Id} + \lambda R$ . Suppose that  $-R$  is strongly nonexpansive. Then  $T$  is a contraction for large distances.

*Proof.* Clearly  $R$  is nonexpansive. Observe that Fact 8.4 applied with  $(T_1, T_2)$  replaced by  $(\text{Id}, R)$  implies that  $T$  is strongly nonexpansive. We claim that  $-T$  is strongly nonexpansive. Indeed, applying Fact 8.4 with  $(T_1, T_2, \lambda)$  replaced by  $(-R, -\text{Id}, 1 - \lambda)$  implies that  $-T = (1 - \lambda)(-R) + \lambda(-\text{Id})$  is strongly nonexpansive. Altogether, we conclude that  $T$  is a contraction for large distances in view of Proposition 5.3. ■

**Theorem 8.6 (Douglas–Rachford algorithm).** Suppose that  $\text{zer}(A + B) \neq \emptyset$ . Suppose that  $A$  is uniformly monotone. Set  $T = \frac{1}{2}(\text{Id} + R_B R_A)$ . Let  $x_0 \in X$  and set  $(\forall n \in \mathbb{N})$ :

$$x_{n+1} = T x_n, \quad (124a)$$

$$y_n = J_A x_n. \quad (124b)$$

Then  $\text{zer}(A + B)$  is a singleton,  $\text{Fix } T \neq \emptyset$  and  $(\exists \bar{x} \in \text{Fix } T)$  such that the following hold:

- (i)  $(y_n)_{n \in \mathbb{N}}$  converges strongly to  $\bar{y} := J_A \bar{x}$ .
- (ii) Suppose that  $(\exists C \in \{A^{-1}, B^{-1}\})$  such that  $C$  is uniformly monotone.

Then we additionally have:

- (a)  $T$  is a contraction for large distances and  $(\exists \bar{x} \in X)$  such that  $\text{Fix } T = \{\bar{x}\}$ .
- (b)  $(x_n)_{n \in \mathbb{N}}$  converges strongly to  $\bar{x}$ .

*Proof.* Applying (122) with  $C$  replaced by  $A + B$ , we conclude that  $\text{zer}(A + B) \neq \emptyset$  is a singleton. It follows from [2, Proposition 26.1(iii)(b)] that  $\text{Fix } T \neq \emptyset$ . (i): This is [2, Proposition 26.11(vi)(b)].

(ii)(a)&(ii)(b): Suppose that  $C = A^{-1}$ . It follows from Theorem 6.3 that  $R_A$  is a contraction for large distances. Consequently,  $T$  is a contraction for large distances. Now combine with Proposition 5.5(i)&(ii).

Now suppose that  $C = B^{-1}$ . It follows from Proposition 3.5 applied to  $A$  (respectively  $B^{-1}$ ) that  $-R_A$  and (respectively  $R_B = -R_{B^{-1}}$ ) is strongly nonexpansive. Consequently,  $-R_B R_A$  is strongly nonexpansive by Proposition 7.1(ii) applied with  $(T_1, T_2)$  replaced by  $(R_A, R_B)$ . Now combine with Proposition 8.5 applied with  $(\lambda, R)$  replaced by  $(\frac{1}{2}, R_B R_A)$ . This proves (i)(c). To show (i)(d), combine (ii)(a) with Fact 2.6. ■

We conclude this section with an application to the forward-backward algorithm.

**Theorem 8.7 (Forward-backward algorithm).** Let  $\beta > 0$ . Suppose that  $A$  is  $\beta$ -cocoercive and that  $B$  is uniformly monotone. Let  $\gamma \in ]0, 2\beta[$ . Set  $T = J_{\gamma B}(\text{Id} - \gamma A)$ . Let  $x_0 \in X$  and set  $(\forall n \in \mathbb{N})$ :

$$x_{n+1} = T x_n. \quad (125)$$

Then the following hold:

- (i)  $\text{zer}(A + B)$  is a singleton
- (ii)  $T$  is a contraction for large distances.
- (iii)  $(x_n)_{n \in \mathbb{N}}$  converges strongly to the unique point in  $\text{zer}(A + B)$ .

*Proof.* (i): Observe that  $A + B$  is maximally monotone by, e.g., [2, Corollary 25.5(i)], and uniformly monotone. Now combine this with Theorem 4.5(iii) applied with  $A$  replaced by  $A + B$ .

(ii): Observe that  $\text{Id} - \gamma A$  is  $\gamma/(2\beta)$  averaged, hence nonexpansive. Now combine this with Lemma 4.4(iv) (applied with  $A$  replaced by  $\gamma A$ ) and Proposition 7.5 applied with  $(m, T_1, T_2)$  replaced by  $(2, \text{Id} - \gamma A, J_{\gamma B})$ .

(iii): Combine (ii) and Proposition 5.5(ii). ■

## Acknowledgements

The research of WMM was partially supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant.

## References

- [1] Ya. I. Alber and S. Guerre-Delabriere, Principle of weakly contractive maps in Hilbert spaces, *New Results in Operator Theory and its Applications, Operator Theory: Advances and Applications* 98 (1997), 7–22.
- [2] H.H. Bauschke and P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, 2nd edition, Springer, 2017.
- [3] H.H. Bauschke, S. Moffat, and X. Wang, Firmly nonexpansive mappings and maximally monotone operators: correspondence and duality, *Set-valued and Variational Analysis* 20 (2012), 131–153.
- [4] H.H. Bauschke, W.M. Moursi, and X. Wang, Generalized monotone operators and their averaged resolvents, *Mathematical Programming (Series B)* 189 (2021) 55–74.
- [5] H.H. Bauschke, S.M. Moffat, and X. Wang, Firmly nonexpansive mappings and maximally monotone operators: correspondence and duality, *Set-valued and Variational Analysis* 20 (2012), 131–153.
- [6] A. Beck, *First-Order Methods in Optimization*, SIAM 2017. <https://doi.org/10.1137/1.9781611974997>
- [7] Y. Benyamini, J. Lindenstrauss, *Geometric Nonlinear Functional Analysis 1*, American Mathematical Society Colloquium Publications 48, American Mathematical Society, Providence, RI, 2000
- [8] J.M. Borwein and J. Vanderwerff, Constructions of uniformly convex functions, *Canadian Mathematical Bulletin* 55 (2012), 697–707.
- [9] J.M. Borwein and J. Vanderwerff, *Convex Functions: Constructions, Characterizations and Counterexamples*, Encyclopedia of Mathematics and Its Applications 109, Cambridge University Press, 2010.
- [10] J.M. Borwein, Fifty years of maximal monotonicity, *Optimization Letters* 4 (2010), 473–490.
- [11] H. Brezis, *Opérateurs Maximaux Monotones et Semi-Groupes de Contractions dans les Espaces de Hilbert*, North-Holland/Elsevier, 1973.
- [12] R.E. Bruck and S. Reich, Nonexpansive projections and resolvents of accretive operators in Banach spaces, *Houston Journal of Mathematics* 3 (1977), 459–470.
- [13] R.S. Burachik and A.N. Iusem, *Set-Valued Mappings and Enlargements of Monotone Operators*, Springer-Verlag, 2008.
- [14] J. Eckstein and D.P. Bertsekas, On the Douglas–Rachford splitting method and the proximal point algorithm for maximal monotone operators, *Mathematical Programming (Series A)* 55 (1992), 293–318.
- [15] P. Giselsson, Tight global linear convergence rate bounds for Douglas–Rachford splitting *Journal of Fixed Point Theory and Applications* 19 (2017), 2241–2270.
- [16] M.I. Garrido and J. Jaramillo, Lipschitz-type functions on metric spaces. *Journal of Mathematical Analysis and Applications* 340 (2008), 282–290.
- [17] G.J. Minty, Monotone (nonlinear) operators in Hilbert spaces, *Duke Mathematical Journal* 29 (1962), 341–346.
- [18] W.M. Moursi and L. Vandenbergh, Douglas–Rachford splitting for the sum of a Lipschitz continuous and a strongly monotone operator, *Journal of Optimization Theory and Applications* 183 (2019), 179–198
- [19] F. D. Parker, Integrals of Inverse Functions, *American Mathematical Monthly* 62, (1955) 439–440.
- [20] E. Rakotch, A note on contractive mappings, *Proceedings of the American Mathematical Society* 13 (1962), 459–465.
- [21] S. Reich and A.J. Zaslavski, Almost all nonexpansive mappings are contractive, *La Société Royale du Canada. L’Académie des Sciences. Comptes Rendus Mathématiques* 22 (2000), 118–124.

- [22] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, 1970.
- [23] R.T. Rockafellar, On the maximal monotonicity of subdifferential mappings, *Pacific Journal of Mathematics* 33, 209–216 (1970).
- [24] R.T. Rockafellar and R. J-B Wets, *Variational Analysis*, Springer-Verlag, corrected 3rd printing, 2009.
- [25] S. Simons, *Minimax and Monotonicity*, Springer-Verlag, 1998.
- [26] S. Simons, *From Hahn–Banach to Monotonicity*, Springer–Verlag, 2007.
- [27] C. Zălinescu, On uniformly convex functions, *Journal Mathematical Analysis and Applications* 95 (1983), 344–374.
- [28] C. Zălinescu, *Convex Analysis in General Vector Spaces*, World Scientific, 2002.
- [29] E. Zeidler, *Nonlinear Functional Analysis and Its Applications II/A: Linear Monotone Operators*, Springer-Verlag, 1990.
- [30] E. Zeidler, *Nonlinear Functional Analysis and Its Applications II/B: Nonlinear Monotone Operators*, Springer-Verlag, 1990.

## Appendix A

*Finer conclusions for Example 6.5.* Let  $g$  be the inverse function of the function  $x \mapsto x + \sin x$  over the interval  $]-\frac{\pi-2}{2}, \frac{\pi+2}{2}[$  and let  $h$  be the inverse function of the function  $x \mapsto x - \sin x$  over the interval  $]-\frac{\pi-2}{2}, \frac{\pi+2}{2}[$ . Set

$$A: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} x + 1, & x \leq -\frac{\pi-2}{4}; \\ g(2x) - x, & |x| < \frac{\pi+2}{4}; \\ x - 1, & \text{otherwise,} \end{cases} \quad (126)$$

and set

$$f: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \frac{1}{2} \begin{cases} x^2 + 2x, & x \leq -\frac{\pi-2}{4}; \\ 2xg(2x) - \frac{1}{2}(g(2x))^2 + \cos(g(2x)) - x^2 - \frac{\pi+1}{2}, & |x| < \frac{\pi+2}{4}; \\ x^2 - 2x, & \text{otherwise.} \end{cases} \quad (127)$$

Then

$$T = R_A, \quad (128)$$

$$A = f', \quad (129)$$

$$A^{-1}: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \begin{cases} x - 1, & x \leq -\frac{\pi+2}{4}; \\ h(2x) - x, & |x| < \frac{\pi-2}{4}; \\ x + 1, & \text{otherwise,} \end{cases} \quad (130)$$

and

$$f^*: \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \frac{1}{2} \begin{cases} x^2 - 2x, & x \leq -\frac{\pi+2}{4}; \\ 2xh(2x) - \frac{1}{2}(h(2x))^2 - \cos(g(2x)) - x^2 + \frac{\pi-1}{2}, & |x| < \frac{\pi-2}{4}; \\ x^2 + 2x, & \text{otherwise.} \end{cases} \quad (131)$$

Observe that  $T = R_A$  if and only if  $A = ((\text{Id} + T)/2)^{-1} - \text{Id}$ . Now

$$\frac{1}{2}(\text{Id} + T): \mathbb{R} \rightarrow \mathbb{R}: x \mapsto \frac{1}{2} \begin{cases} x - 1, & x \leq -\frac{\pi}{2}; \\ x + \sin x, & |x| < \frac{\pi}{2}; \\ x + 1, & \text{otherwise.} \end{cases} \quad (132)$$




Figure 3: A GeoGebra snapshot illustrating Example 6.5. Left: plot of  $A(x)$ . Right: plot of  $A^{-1}(x)$ .

Note that  $\frac{1}{2}(\text{Id}+T): \mathbb{R} \rightarrow \mathbb{R}$  is strictly increasing and differentiable, hence continuous. Therefore,  $(\frac{1}{2}(\text{Id}+T))^{-1}: \mathbb{R} \rightarrow \mathbb{R}$  is strictly increasing and continuous. To this end let  $(x, y) \in \mathbb{R} \times \mathbb{R}$ . Then  $y = (\frac{1}{2}(\text{Id}+T))^{-1}(x)$  if and only if  $x = \frac{1}{2}(y + Ty)$ . If  $y \leq -\frac{\pi}{2}$  then  $x = \frac{1}{2}(y - 1)$ . Hence,  $y = 2x + 1$  and  $x \leq \frac{-\pi-2}{4}$ . Similarly, if  $y \geq \frac{\pi}{2}$  then  $x = \frac{1}{2}(y + 1)$ . Hence,  $y = 2x - 1$  and  $x \geq \frac{\pi+2}{4}$ . If  $y < |\frac{\pi}{2}|$  then  $x = \frac{1}{2}(y + \sin y)$ ; equivalently,  $y = g(2x)$  and  $|x| \leq \frac{\pi+2}{4}$ . This proves (128).

We now show that  $f$  is an antiderivative of  $A$ . The formula for  $f$  over the intervals  $[\frac{\pi+2}{4}, +\infty[$  and  $] -\infty, -\frac{\pi+2}{4}]$  is straightforward. To compute an antiderivative of  $g(2x) - x$  we use [19] to learn that

$$\int (g(2x) - x) dx = \frac{1}{2} \left( 2xg(2x) - \frac{(g(x))^2}{2} + \cos(g(2x)) - x^2 \right) + C. \quad (133)$$

The continuity of  $A$  implies that  $g(\frac{\pi+2}{2}) = \frac{\pi}{2}$  and  $g(-\frac{\pi+2}{2}) = -\frac{\pi}{2}$ . This, together with the continuity of  $f$ , imply that  $C = -\frac{\pi+1}{4}$ . This proves (129).

To prove (130) proceed similar to the proof of (128) and observe that  $R_{A^{-1}} = -T$ . Analogously, the proof of (131) is similar to that of (129) by observing that  $A^{-1} = (f^*)'$ . ■






Figure 4: A GeoGebra snapshot illustrating [Example 6.5](#). Left: plot of  $f$ . Right: plot of  $f^*$ .