

Quantum Supermaps are Characterized by Locality

Matt Wilson^{1,2}, Giulio Chiribella^{1,2}, and Aleks Kissinger¹


¹Quantum Group, Department of Computer Science, University of Oxford

²HKU-Oxford Joint Laboratory for Quantum Information and Computation

We prove that the sequential and parallel composition rules for quantum channels are all that need to be referenced to axiomatise quantum supermaps from first principles. To do so we provide a simple definition of *locally-applicable transformation*, which can be stated for arbitrary symmetric monoidal categories, and so for arbitrary process theories and operational probabilistic theories. The definition can be rephrased in the language of category theory using the principle of naturality, and can be given an intuitive diagrammatic representation in terms of which all proofs are presented. In our main technical contribution we use this diagrammatic representation to show that locally-applicable transformations on quantum channels are in one-to-one correspondence with deterministic quantum supermaps. This re-characterization of quantum supermaps is proven to hold for supermaps on arbitrary convex subsets of channels, including as a special case the supermaps on non-signalling channels used in the study of quantum causal structure.

1 Introduction

Beyond the framework of standard quantum theory, in which states representing physical degrees of freedom incur changes over time, there is the framework of higher order quantum theory [1–9], in which dynamics themselves are transformed by higher order operations termed *supermaps* [1]. Quantum supermaps were originally defined to formalise the following intuitive picture:




In other words, supermaps were defined to formalize the notion of a higher order map S which may be applied to part of any bipartite process ϕ . Such supermaps can be generalised to act on constrained spaces, and are applied in the study of quantum information to analyze protocols in which quantum processes are treated as information-theoretic resources [10–25]. On top of providing a framework for formalising such protocols, supermaps are broad enough to incorporate higher order processes beyond those which can be interpreted as circuits with open holes [2, 26–33], the canonical example of such a supermap being the quantum switch [34]. Acting on the space of non-signalling channels,

Matt Wilson: matthew.wilson@cs.ox.ac.uk


Giulio Chiribella: giulio.chiribella@cs.ox.ac.uk

Aleks Kissinger: aleks.kissinger@cs.ox.ac.uk

the quantum switch uses a qubit to control the order in which wires are plugged together, and is often interpreted as a superposition of causal structures [34].



By being broad enough to incorporate examples such as the quantum switch, the supermap framework provides a way to study quantum causal structure as a resource in quantum information processing protocols [13–18, 22, 23, 35–38], and furthermore finds application in the study of quantum gravity. Indeed, the dependence of causal structure on mass distribution in general relativity and the possibility to superpose mass distributions in quantum theory, suggests the possibility of naturally occurring non-classical, or indefinite, causal structures [34, 39–49] such as those present in the quantum switch. To formalise this intuition supermaps (in this context often referred to as process matrices [40]) can be used, to model global spacetime structures as maps from interventions chosen in local quantum laboratories to probabilities [40].




This abstract approach to modeling global spacetime structures allows to study causal structures beyond those which are even switch-like. There are supermaps so causally indefinite that they can be used to break causal inequalities [40, 50–52], in analogy to the way which maximally entangled states are so non-local that they can be used to break Bell inequalities in standard quantum theory.

The motivating picture of a supermap appears to only reference the possibility to draw processes in quantum theory as boxes with multiple inputs and outputs, and yet, current constructions and definitions of quantum supermaps [1, 7, 40] rely on additional mathematical structures. These are typically structures of categories into which deterministic quantum theories embed, such as compact closure/Choi-Jamiołkowski isomorphism [1, 7, 53] and convexity/coarse graining [40]. This suggests that supermaps are in need of a more principled axiomatisation, so that the conceptual grounds on which supermaps are understood match the formal grounds on which they are defined, and so that the entire framework of higher order quantum theory may be more easily lifted to arbitrary physical theories.


In this paper we provide such an axiomatisation, by building on the process-theoretic/categorical approach to quantum theory [54–59] we discover that supermaps can indeed be axiomatised purely at the process-theoretic level, that is, with respect to symmetric monoidal structure [60]. From this result it follows that all of the operational physical principles such as compatibility with coarse-graining, convex combinations, linearity, and tensor extensions used in usual definitions of quantum supermaps can be viewed as consequences a simple principle:

A supermap is the kind of thing that can be applied locally.


The formalization of local applicability of supermaps that we use as our axiom can be understood in three consecutive steps. First, supermaps are functions on processes:



Second, supermaps are equipped with extensions to functions on all bipartite processes:



Third, localization is enforced by requiring that such functions commute with actions on their extensions:



We model these three principles for supermaps by a formal definition of *locally-applicable transformation*. The definition can be stated in entirely diagrammatic terms, and this diagrammatic phrasing is used throughout as a toolbox for proving our main theorem.

Theorem. *Quantum supermaps, including those on signalling-constrained channels, are in one-to-one correspondence with locally-applicable transformations.*

This characterisation is additionally noted to apply in the context of classical information theory. Given that supermaps are typically used to analyse causal structure, the above result can be viewed as complementary to the viewpoint of the causal decomposition program [61–63], in which causal structure in quantum theory is understood in terms of compositional structure. Indeed, we find that some of the very tools that are used to study quantum causal structure could have been defined purely at the compositional level all along.

Although we do not do so in the main text, many of the definitions and results (though not the proofs) can be neatly phrased in the language of category theory [60]. First, the presented correspondence between locally-applicable transformations and standard-definition quantum supermaps is compositional, meaning that it can be phrased as an equivalence of categories [60]. Second, the definition of locally-applicable transformation can itself be rephrased using the concept of natural transformation between functors. As a consequence of this observation our main theorem can be summarised concisely as a categorical characterisation of supermaps on quantum theory:

Quantum supermaps are equivalent to natural transformations.

Specialised instances of such natural transformations can in fact already be observed as playing a similar role to supermaps in the field of monoidal category theory, for instance

in the definition of a traced monoidal category [64,65] or similarly in the formalisation of closed time-like curves [66] where such curves are even termed super-operators.

The results presented in this paper show that the rich field of higher order quantum theory can be understood as the study of locally-applicable transformations in monoidal category theory, applied to the specific instance of quantum theory. In the long term the authors hope that the same viewpoint can be used to reconstruct all iterated layers of higher order quantum theory [1, 6, 7, 40] with only appeal to variations on the principle of formalisation of locality by naturality. We hope that this result furthermore allows for easier generalisation of higher order quantum transformations to non-monoidal physical frameworks. For instance the authors would like to see supermaps applied in decompositional approaches to physics [67,68] and to algebraic quantum field theories [69] where localizability of standard quantum transformations is of primary importance [70–73]. Ultimately, such generalisations would bring the recently developed higher-order approach to the study of quantum causal structure closer to being applicable to theories of quantum gravity, where such structures have been predicted to play a key role [39].

2 Preliminary Material

Here we review symmetric monoidal categories, an abstract model for theories with sequential and parallel composition, such as quantum circuits. We then review quantum supermaps, presenting them in terms of the graphical language for compact closed categories which captures the essence of the Choi-Jamiołkowski isomorphism. Throughout this paper we use purple-shaded boxes to represent standard quantum processes, and white boxes to indicate parts of a diagram which should be interpreted as higher-order maps to be applied to standard processes.

Symmetric Monoidal Categories To ease the following presentation we present symmetric monoidal categories which are “strict”, meaning that equalities between objects are written in place of natural isomorphisms. For a formal treatment of non-strict symmetric monoidal categories the reader is referred to [60], where it is noted that every symmetric monoidal category is suitably equivalent to a strict one. We from now on omit use of the word strict, leaving it as implicit for the remainder of the paper. Our working examples of symmetric monoidal categories will be the categories \mathbf{QC} , \mathbf{CP} of quantum channels and completely positive maps respectively. We use the term quantum channel to mean completely positive trace-preserving map.

The first formal ingredient in the definition of a symmetric monoidal category \mathbf{C} is the specification of objects of \mathbf{C} , which in quantum theory may be thought of as systems. In symmetric monoidal categories objects can be drawn as wires, for instance we draw an object A as:


$$\begin{array}{c} | \\ A \end{array}$$

Objects of each of \mathbf{QC} and \mathbf{CP} are given by finite dimensional Hilbert spaces, which are used to represent quantum degrees of freedom. The second formal ingredient of a symmetric monoidal category is the assignment of a set $\mathbf{C}(A, B)$ to each pair A, B of objects. The set $\mathbf{C}(A, B)$ will be referred to as the set of morphisms, and the elements $f \in \mathbf{C}(A, B)$ of this set are often denoted using the notation $f : A \rightarrow B$. Morphisms in symmetric monoidal categories can be drawn as boxes with input and output wires, used

to represent their input and output objects:



In **QC** the morphisms $f : H_A \rightarrow H_B$ are completely positive trace preserving maps from $\mathcal{L}(H_A)$ to $\mathcal{L}(H_B)$. Similarly, in **CP** the morphisms are taken to be completely positive maps. Symmetric monoidal categories are next equipped with two compatible notions, sequential composition and parallel composition. Sequential composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ can be denoted graphically by:




In quantum theory the sequential composition $g \circ f$ is typically interpreted as representing a process g occurring some time after a process f . In **QC** and **CP** the composition rule is inherited directly from the standard notion of sequential composition for linear maps.

Parallel composition addresses both objects and morphisms, for each pair of objects A, B a new object can be assigned called $A \otimes B$, typically interpreted as placing A next to B . For each pair of morphisms $f : A \rightarrow A'$ and $g : B \rightarrow B'$ one can assign a new morphism $f \otimes g : A \otimes B \rightarrow A' \otimes B'$ typically interpreted as the concurrent action of f and g . Diagrammatically $A \otimes B$ can be represented by placing wire A next to wire B :




further solidifying its interpretation as parallel composition. The expression $f \otimes g$ can then be represented by the following diagram:



which again displays concurrent action of f and g . In our working examples the parallel composition of objects H_A, H_B is given by the standard tensor product $H_A \otimes H_B$ of finite dimensional Hilbert spaces. A key feature of symmetric monoidal categories which allows them a diagrammatic calculus is the interchange law

$$(f \circ f') \otimes (g \circ g') = (f \otimes g) \circ (f' \otimes g')$$

which allows unambiguous interpretation of the following diagram:



In a monoidal category the parallel composition is required to be associative so that $(A \otimes B) \otimes C = A \otimes (B \otimes C)$, this is indeed true for the tensor product of Hilbert spaces up to natural isomorphism $H_A \otimes (H_B \otimes H_C) \cong (H_A \otimes H_B) \otimes H_C$ as addressed in the

introductory remarks. A monoidal category furthermore comes equipped with a notion of empty space given by an object I satisfying $A \otimes I = A = I \otimes A$, this object in our motivating examples is given by the Hilbert space \mathbb{C} satisfying $H_A \otimes \mathbb{C} \cong H_A \cong \mathbb{C} \otimes H_A$ for every Hilbert space H_A . Making use of the notion of empty space I , the morphisms of type $\rho : I \rightarrow A$ are often interpreted states A . This is well motivated when I -wires are omitted from diagrams so that morphisms of type $I \rightarrow A$ are drawn as having only output wires:



Finally, a symmetric monoidal category comes equipped with a swap-morphism $\beta : A \otimes B \Rightarrow B \otimes A$ depicted by:



satisfying a variety of natural properties [60]. In the course of this paper, we will need to address two refined monoidal structures which supermaps are typically defined with respect to, those monoidal structures which are additionally causal and those which are additionally compact closed.

Causality: The Trace Channel A symmetric monoidal category is causal [54, 74] if its unit object is terminal, meaning in concrete terms that for each A the set $\mathbf{C}(A, I)$ is a singleton. In a causal symmetric monoidal category we typically represent the unique effect of type $A \rightarrow I$ using the discard symbol:




Our working example **QC** is a causal category, with the unique morphism of type $H_A \rightarrow \mathbb{C}$ given by the trace quantum channel. When a category is causal a direction of time is essentially fixed as propagating up the page. Indeed, with no way to vary between effects of type $\mathbf{C}(A, I)$ then there is no way to send information in the opposite direction, from the top of the page to the bottom.

Compact Closure: Channel-State Duality A symmetric monoidal category is compact closed if it is equipped for each A with an object A^* and morphisms $\cup_A : I \rightarrow A^* \otimes A$ and $\cap_A : A \otimes A^* \rightarrow I$ which amongst other natural conditions [54] satisfy $(\cap \otimes id) \circ (id \otimes \cup) = id$, graphically this reads as:



When compact closure is present it offers significant flexibility by giving an internalised way to connect input and output ports of circuit diagrams together. When compact closure is not in some way present we find that we require a new externalised way to reason about operating on and plugging-together processes. Our working example **CP** is a compact closed category with H_A^* given by the dual Hilbert space to H_A and the cup $\cup_{H_A} : \mathbb{C} \rightarrow H_A^* \otimes H_A$ given by the maximally entangled Bell-state. Similarly the cap is given by the maximally entangled Bell-effect. The Choi-isomorphism [53], which provides an identification between completely positive maps and positive operators, is given by applying completely positive maps to Bell-states and so can be expressed graphically as:



Additive Structure The symmetric monoidal category \mathbf{CP} is equipped with a notion of closure under positive linear combination, for any pair $\phi_0, \phi_1 \in \mathbf{CP}(A, A')$ of completely positive maps and pair $a_0, a_1 \in \mathbb{R}_+$ of positive real numbers a new combined completely positive linear map $a_0\phi_0 + a_1\phi_1 \in \mathbf{CP}(A, A')$ can be constructed. The symmetric monoidal category \mathbf{QC} inherits a more restricted form of closure from \mathbf{CP} termed closure under convex combinations, meaning that for any pair $\phi_0, \phi_1 \in \mathbf{QC}(A, A')$ of quantum channels and any probability $p \in [0, 1]$ then the linear combination $p\phi_0 + (1-p)\phi_1 \in \mathbf{QC}(A, A')$ is itself a quantum channel.

Constrained Spaces In this paper, beyond supermaps on entire channels, we will aim to reconstruct supermaps which can be applied to channels satisfying signalling constraints [75, 76]. These quantum supermaps are the part of the higher-order toolbox typically used to study quantum causal structure. Let us begin with a simple example of a signalling constraint given by the specification of the set of bipartite one-way signaling channels, we denote the statement that ϕ be a one-way signaling channel by:

$$\begin{array}{c} | \\ | \\ \boxed{\Phi} \\ | \\ | \end{array} \in \mathcal{E}_{sig} \left(\begin{array}{c} \uparrow \quad \nearrow \quad \uparrow \\ \uparrow \quad \quad \uparrow \end{array} \right)$$

The depicted graph indicates that ϕ may not transmit information from the bottom right wire to the top left wire, this constraint is expressed concretely by the existence of a channel ϕ' satisfying the following equation:

$$\begin{array}{c} \text{---} \\ | \\ | \\ \boxed{\phi} \\ | \\ | \end{array} = \begin{array}{c} | \\ | \\ \boxed{\phi'} \\ | \\ | \end{array} \begin{array}{c} \text{---} \\ | \\ | \end{array}$$

In general more complex signalling constraints can be encoded by relations [61, 77]. For instance the statement

$$\begin{array}{c} | \\ | \\ | \\ \boxed{\Phi} \\ | \\ | \\ | \end{array} \in \mathcal{E}_{sig} \left(\begin{array}{c} \uparrow \nearrow \uparrow \nearrow \uparrow \nearrow \uparrow \\ \uparrow \nearrow \uparrow \nearrow \uparrow \end{array} \right)$$

can be used to encode the statements

$$\begin{array}{c} \text{---} \\ | \\ | \\ \boxed{\Phi} \\ | \\ | \end{array} = \begin{array}{c} \text{---} \\ | \\ | \\ \boxed{1} \\ | \\ | \end{array} \quad \text{and} \quad \begin{array}{c} \text{---} \\ | \\ | \\ \boxed{\Phi} \\ | \\ | \end{array} = \begin{array}{c} \text{---} \\ | \\ | \\ \boxed{2} \\ | \\ | \end{array} \begin{array}{c} \text{---} \\ | \\ | \end{array}$$

A formal definition for arbitrary relations is given in [77]. Useful for our reconstruction will be the observation that for any relation τ the set $\mathcal{E}_{sig}(\tau)$ of processes satisfying constraint τ has the convenient property of being closed under convex combinations. Another convenient property of signaling constraints in quantum theory for our reconstruction is their equivalence to localizability constraints, which we refer to here as pathing constraints so as to not confuse with our main definition of local applicability. As an example of a pathing constraint, the one-way signalling condition in quantum theory is equivalent to requiring the following decomposition:

$$\begin{array}{c} | \\ | \\ \boxed{\Phi} \\ | \\ | \end{array} = \begin{array}{c} | \\ | \\ \boxed{2} \\ | \\ \boxed{1} \\ | \\ | \end{array}$$

this decomposition abstracts the impossibility of signalling from the bottom right wire to the top left wire by forbidding the existence of a vertically directed path between them. Since in quantum theory pathing constraints are equivalent to signaling constraints [75], pathing constraints in quantum theory are also closed under convex combinations. Pathing constraints come with the advantage however that they can be phrased in arbitrary symmetric monoidal categories without reference to a prioritised effect or time-direction. In intuitive terms pathing constraints abstract from causality to compositionality, as an example of a more complex pathing constraint consider the statement:

$$\begin{array}{c} | \\ | \\ | \\ | \\ \hline \Phi \\ \hline | \\ | \\ | \\ | \end{array} \in \mathcal{E}_{path} \left(\begin{array}{c} \nearrow \nwarrow \nearrow \nwarrow \nearrow \nwarrow \nearrow \nwarrow \\ \nwarrow \nearrow \nwarrow \nearrow \nwarrow \nearrow \nwarrow \nearrow \end{array} \right)$$

this is a concise encoding of the statement that ϕ can be decomposed in such a way as to not provide a vertically directed path from the left-most input to the right-most output, or vice-versa, I.E

$$\begin{array}{c} | \\ | \\ | \\ | \\ \hline \Phi \\ \hline | \\ | \\ | \\ | \end{array} = \begin{array}{c} | \\ | \\ | \\ | \\ \hline 2 \\ \hline | \\ | \\ | \\ | \\ \hline 1 \\ \hline | \\ | \\ | \\ | \end{array} \quad \text{and} \quad \begin{array}{c} | \\ | \\ | \\ | \\ \hline \Phi \\ \hline | \\ | \\ | \\ | \end{array} = \begin{array}{c} | \\ | \\ | \\ | \\ \hline 2 \\ \hline | \\ | \\ | \\ | \\ \hline 1 \\ \hline | \\ | \\ | \\ | \end{array}$$

The equality $\mathcal{E}_{path}(\tau) = \mathcal{E}_{sig}(\tau)$ will allow us to reconstruct quantum supermaps on signaling constraints as locally-applicable transformations applied to their corresponding pathing constraints. Phrased in this way the quantum supermaps used to study quantum causal structure, are recovered from entirely compositional definitions with no reference to causality or chosen direction of the flow of time.

Extended Process Sets In this paper we will be often concerned with extending sets of processes to auxiliary systems. The principle is to define, for any set $K \subseteq \mathbf{C}(A, A')$ and pair X, X' , the extension by X, X' of K to be the set of all processes $\phi \in \mathbf{C}(A \otimes X, A' \otimes X')$ such that, whenever states and effects are applied to the X, X' part of ϕ the result is an element of K .

Definition 1. For each $K \subseteq \mathbf{C}(A, A')$ and pair (X, X') the set of extended channels $\text{Ext}_{X, X'}(K)$ is given by:


$$\Phi \in \text{Ext}_{X, X'}[K] \iff \forall \rho, \sigma : \begin{array}{c} | \\ | \\ | \\ | \\ \hline \sigma \\ \hline \Phi \\ \hline | \\ | \\ | \\ | \\ \hline \rho \\ \hline | \\ | \\ | \\ | \end{array} \in K$$

Note that for the example $K = \mathbf{C}(A, A')$ then the extension $\text{Ext}_{X, X'}(K)$ returns the entire set of bipartite morphisms $\mathbf{C}(A \otimes X, A' \otimes X')$. For a causal symmetric monoidal category the extended set can be rephrased in the following way:


$$\phi \in \text{Ext}_{X, X'}[K] \iff \forall \rho : \begin{array}{c} | \\ | \\ | \\ | \\ \hline \Phi \\ \hline | \\ | \\ | \\ | \\ \hline \rho \\ \hline | \\ | \\ | \\ | \end{array} \in K$$

Which is the form used in [34] to define extensions to subsets of quantum channels. In the appendix we note that such extended channel sets can be viewed in categorical language as functors into the category of sets. Whenever a subset $K \subseteq \mathbf{QC}(A, A')$ is closed under convex combinations then it follows that $\text{Ext}_{X, X'}(K)$ is closed under convex combinations, we will from now on rephrase the statement that a set K be closed under convex combinations as simply the statement that K be convex.

Quantum Supermaps: Standard Definition In this paper we use the category \mathbf{CP} of completely positive maps to express the definition of supermap, this is sufficient for our purposes since the sets we choose to work with are internal [3, 34]. We follow the presentation of [7] in which a completely positive linear map




is informally interpreted as a diagram with a hole, a function which accepts a channels of type $A \rightarrow A'$ and returns a channel of type $B \rightarrow B'$ by using compact closure or Choi-Jamiolkowski isomorphism.



This notation puts us in a position to concisely phrase the standard definition method for quantum supermaps.

Definition 2. Let $\mathbf{C} \subseteq \mathbf{P}$ be an inclusion of a symmetric monoidal category \mathbf{C} into a compact closed category \mathbf{P} . A \mathbf{P} -supermap on \mathbf{C} of type $S : K \rightarrow M$ with $K \subseteq \mathbf{C}(A, A')$ and $M \subseteq \mathbf{C}(B, B')$ is a morphism in \mathbf{P} of type $S : A^* \otimes A' \rightarrow B^* \otimes B'$ such that for every $\phi \in \mathbf{Ext}_{X, X'}[K]$ then




\mathbf{P} -supermaps can be composed in sequence, with this sequential composition inherited directly from \mathbf{P} , in categorical terms this means that the \mathbf{P} -supermaps define a category $\mathbf{Psup}[\mathbf{C}]$. For brevity we will use the term “quantum supermap” for \mathbf{CP} -supermap on \mathbf{QC} and denote $\mathbf{Psup}[\mathbf{C}]$ by \mathbf{QS} . Note that we are working with quantum supermaps applied to arbitrary subsets here, including those which satisfy signalling constraints. For the study of indefinite causal structure the most prevalent example is that of supermaps on non-signalling channels:



The lack of communication between wires in such channels allows them to be combined in a variety of ways without producing time-loops, so that the output may still be guaranteed to be a deterministic channel. Supermaps of this type which have been

the subject of considerable study are switches, which take channels and by reference to a control state, plug them together in a combination of orders:




Note that when the above diagrams are drawn in the category \mathbf{CP} they really represent classical mixtures of causal orders, a supermap which exhibits a quantum superposition of causal orders termed the quantum switch can be constructed in a similar way by including interference terms.

Summary In quantum information theory, supermaps on quantum channels are defined by using channel state duality (compact closure) of the theory of completely positive maps into which they embed. In this paper we ask the following: *What can we say when background compact closure cannot be assumed?* In other words: *Can supermaps be characterized in terms of sequential and parallel composition alone?* We answer positively, showing that quantum supermaps could have been defined all along by an abstract principle of local applicability (in category-theoretic language termed naturality). After motivating and defining locally-applicable transformations on arbitrary symmetric monoidal categories we show a one-to-one correspondence between them and quantum supermaps when applied to the symmetric monoidal category of quantum channels.


3 Formalisation of Local Applicability

Our goal is to find an axioms for quantum supermaps which can be applied to *any* symmetric monoidal category. In this section we show how to so, by defining higher order functions and requiring the existence of extensions for all auxiliary systems which further commute with actions on those auxiliary systems. We begin by warming up to the definition by observing a notion of local applicability present in monoidal categories, such as standard quantum theory, which we plan to abstract to higher order functions.


Local Applicability in Standard Quantum Theory The category \mathbf{QC} of quantum channels is a symmetric monoidal category, one consequence of the parallel composition rule $f \otimes g$ is that it gives a way to view any $f : A \rightarrow A'$ as locally applicable in an intuitive sense. We discuss this locality principle in the monoidal setting and then comment on how it can be abstracted to a statement about locality of functions built from such an f on state sets. To begin, consider a channel $f : A \rightarrow A'$ from A to A' :



Whenever A can be viewed as part of a larger system $N = A \otimes X$ then f can be locally applied to N by using $f \otimes id_X : N \rightarrow N$



Crucial to the interpretation of locality in $f \otimes id_X$ is that $f \otimes id_X$ commutes with all actions on X , this follows in this case by the interchange law for monoidal categories $(f \otimes id_{X'}) \circ (id_A \otimes g) = (id_{A'} \otimes g) \circ (f \otimes id_X)$:




In other words, a general monoidal category gives a collection of morphisms, all of which can be viewed as being locally applicable, in an informal sense.

A consequence of this local applicability is the possibility to construct from f a family of functions on states $l(f)_X : \mathbf{C}(I, A \otimes X) \rightarrow \mathbf{C}(I, A' \otimes X)$ which exhibit the local applicability of f . Explicitly, by using tensor extensions with the identity the function $l(f)_X(\rho_{AX}) := (f \otimes id_X)(\rho_{AX})$ can be defined for each X . The abstract functions $l(f)_X$ which represent the action of f on states indeed inherit a notion of local applicability from f . The functions $l(f)_X$ can be seen to leave the environment system X untouched in the sense that the action of any g on X commutes with the application of the function $l(f)_X$. The above sentence is captured in formal terms by the equation $l(f)_{X'}(id_A \otimes g(\rho_{AX})) = (id_{A'} \otimes g)(l_X(\rho_{AX}))$ which is guaranteed to hold for any $g : X \rightarrow X'$ since

$$\begin{aligned} l(f)_{X'}(id_A \otimes g(\rho_{AX})) &= (f \otimes id_{X'}) \circ (id_A \otimes g)(\rho_{AX}) \\ &= (id_{A'} \otimes g) \circ (f \otimes id_X)(\rho_{AX}) \\ &= (id_{A'} \otimes g)(l_X(\rho_{AX})) \end{aligned}$$


We now give an axiom which re-characterises quantum supermaps by generalizing this concept of local applicability of functions on states to local applicability of functions on processes. The only instances of locally-applicable transformations on quantum channels will turn out to be those which are simulated by the standard-definition quantum supermaps of [1]. We split the motivations for the definition of locally-applicable transformation into three consecutive principles.

Principle 1: Supermaps are Functions on Processes The kind of picture usually drawn with the aim of capturing diagrammatically the concept of a supermap from the space of processes $\mathbf{C}(A, A')$ to the space of processes $\mathbf{C}(B, B')$ is some variation of the following




As such our first step to characterising supermaps of type $\mathbf{C}(A, A') \rightarrow \mathbf{C}(B, B')$ is to consider functions of the same type $\mathbf{C}(A, A') \rightarrow \mathbf{C}(B, B')$. More generally for $K \subseteq \mathbf{C}(A, A')$ and $M \subseteq \mathbf{C}(B, B')$ the first step is to consider functions $S : K \rightarrow M$ from the set K to the set M .

Principle 2: Supermaps Can be Extended to Functions on All Bipartite Processes When we say that we wish for the map $S : \mathbf{C}(A, A') \rightarrow \mathbf{C}(B, B')$ to be locally applicable, we mean that we wish to formalise the following picture:



The next step toward such a formalisation is to specify for each X, X' the action of S when applied to the A, A' part of any morphism $\phi \in \mathbf{C}(A \otimes X, A' \otimes X')$. Consequently we say that a locally-applicable transformation must be equipped with a family of extended functions $S_{X, X'} : \mathbf{C}(A \otimes X, A' \otimes X') \rightarrow \mathbf{C}(B \otimes X, B' \otimes X')$ for every X, X' . For the generalised case of supermaps of type $S : K \rightarrow M$ we instead require the specification of a function $S_{X, X'} : \text{Ext}_{X, X'}(K) \rightarrow \text{Ext}_{X, X'}(M)$ for every X, X' . We now will need to find a way to enforce these extensions behave *as if* they are applied locally.

Principle 3: Supermaps Commute With Actions on Their Extensions A key feature of a local operation is commutation with operations applied to auxiliary spaces, we generalise this notion of locality to input-output operations, informally we aim to capture the equivalence of the following two pictures:



This can be formalised by the following condition:

$$S_{X, X'}((i \otimes g) \circ \phi \circ (i \otimes f)) = (i \otimes g) \circ S_{Y, Y'}(\phi) \circ (i \otimes f)$$

which can be required for functions on simple types $\mathbf{C}(A, A') \rightarrow \mathbf{C}(B, B')$ or on generalised types $K \rightarrow M$. The above considerations can be collected into a single concise definition of locally-applicable transformation, to which we will later assign a more intuitive entirely diagrammatic language.

Definition 3 (locally-applicable transformations). *A locally-applicable transformation of type $S : K \rightarrow M$ is a family of functions $S_{X X'} : \mathbf{Ext}_{X, X'}(K) \rightarrow \mathbf{Ext}_{X, X'}(M)$ such that for every $g : X' \rightarrow Y'$, $f : Y \rightarrow X$, and $\phi : A \otimes X \rightarrow A' \otimes X'$ then*


$$S_{Y Y'}((i \otimes g) \circ \phi \circ (i \otimes f)) = (i \otimes g) \circ S_{X X'}(\phi) \circ (i \otimes f).$$

In the appendix we note that locally-applicable transformations can be phrased in the language of category theory as natural transformations, and will consequently refer to the above notion of commutativity as naturality whenever it is used. We further note that locally-applicable transformations can be composed, given locally-applicable transformations $S : K \rightarrow M$ and $T : M \rightarrow N$ one can construct the locally-applicable transformation $(T \circ S) : K \rightarrow N$ by defining for each X, X' the functions $(S \circ T)_{X X'}(\phi) = S_{X X'}(T_{X X'}(\phi))$. This compositionality of locally-applicable transformations can be phrased in the language of category theory by stating that they form a category, which we denote by $\mathbf{Im}[\mathbf{C}]$. As a sanity check, we can check that every supermap constructed by the standard method indeed defines a locally-applicable transformation.


Example 1 (**P**-supermaps define locally-applicable transformations). *Every **P**-supermap of type $S : K \rightarrow M$ on \mathbf{C} defines a locally-applicable transformation $\mathcal{F}_{\mathbf{P}}(S) : K \rightarrow M$ of the same type on \mathbf{C} . Indeed one can define $\mathcal{F}_{\mathbf{P}}(S)$ by*

$$\mathcal{F}_{\mathbf{P}}(S)_{X, X'}(\phi) := \begin{array}{c} \begin{array}{c} B' \\ \boxed{S} \\ \text{---} \\ \boxed{\phi} \\ \text{---} \\ X \end{array} \end{array}$$

Naturality of the assigned family of functions is confirmed by noting that $\mathcal{F}_{\mathbf{P}}(S)_{X, X'}((id_{A'} \otimes g) \circ \phi \circ (id_A \otimes f))$ is given by:



which by the interchange law for monoidal categories is equal to:



which by definition is $(id_{B'} \otimes g) \circ \mathcal{F}_{\mathbf{P}}(S)_{Y,Y'}(S)(\phi) \circ (id_B \otimes f)$. The assignment $\mathcal{F}_{\mathbf{P}}$ is injective since if $S \neq T$ then this will be witnessed by the inequality $\mathcal{F}_{\mathbf{P}}(S)(\beta) \neq \mathcal{F}_{\mathbf{P}}(T)(\beta)$ of their actions on the swap process.

The embedding $\mathcal{F}_{\mathbf{P}}$ is in fact a type of structure-preserving map between categories called a functor $\mathcal{F}_{\mathbf{P}} : \mathbf{Psup}[\mathbf{C}] \rightarrow \mathbf{Lm}[\mathbf{C}]$ concretely this means that it preserves composition and identities. Injectivity is phrased in categorical language as faithfulness of the functor $\mathcal{F}_{\mathbf{P}}$. We will find that for quantum channels, any locally-applicable transformation S is implemented by a \mathbf{CP} -supermap in the above way, meaning that $\mathcal{F}_{\mathbf{CP}}$ can be inverted, to find for each S some T such that $S = \mathcal{F}_{\mathbf{CP}}(T)$. To aid us in proving this statement we will introduce an extension of the diagrammatic language for monoidal categories to include the defining rules of locally-applicable transformations.

Diagrammatic Language We will find that using a function-box notation simplifies the presentation of our proofs, the diagrammatic presentation given here can be traced back to [66]. For any locally-applicable transformation $S : K \rightarrow M$, any pair of objects X, X' and any $\phi \in \mathbf{Ext}_{X,X'}(K)$ we can denote the action of $S_{X,X'}$ by:

$$S_{X,X'}(\phi) := \begin{array}{c} \begin{array}{|c|} \hline X' \\ \hline \begin{array}{|c|} \hline S_{X,X'} \\ \hline \begin{array}{|c|} \hline \phi \\ \hline X \\ \hline \end{array} \\ \hline \end{array} \\ \hline X \end{array} \end{array}.$$

The defining condition for locally-applicable transformations is then easily introduced into function-box notation by including the following diagrammatic equation for *naturality*:

$$\begin{array}{c} \begin{array}{|c|} \hline g \\ \hline \begin{array}{|c|} \hline S_{Y,Y'} \\ \hline \begin{array}{|c|} \hline \phi \\ \hline \end{array} \\ \hline \end{array} \\ \hline f \end{array} = \begin{array}{c} \begin{array}{|c|} \hline \begin{array}{|c|} \hline S_{X,X'} \\ \hline \begin{array}{|c|} \hline \phi \\ \hline \end{array} \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline g \\ \hline f \end{array} \\ \hline \end{array} \end{array}.$$


In short, boxes can always be slid through dotted wires. The dotted lines are intended to model the notion that the action of S on the auxiliary systems is essentially that of an identity transformation. There is an additional property which can be phrased at the symmetric monoidal level, which if it wasn't true by other means for quantum theory, would be so primitive as to be justified as being an additional axiom for locally-applicable transformations. For other physical theories such as \mathbb{R} Real quantum theory we expect that we would in-fact need to state it as such.

Lemma 1 (Tensor separation). *Every locally-applicable transformation of type $K \rightarrow M$ on \mathbf{QC} commutes with products, meaning that for every $\phi : A \otimes X \rightarrow A' \otimes X'$ and $\psi : Y \rightarrow Y'$ then:*


$$S_{XY,X'Y'}(\phi \otimes \psi) = S_{X,X'}(\phi) \otimes \psi$$

This is indeed a property that would be expected of a function S which does not act on the environment Y, Y' .

Proof. For all causal ρ we have the following equation by naturality:




and following further equations, again by naturality:



Which together entail the result since quantum theory has enough causal states [7]. \square

When tensor separation is satisfied, it is again neatly captured by a diagrammatic rule which we term *box dragging*:



These two diagrammatic rules, will greatly ease the presentation of our derivations.

4 Characterisation of Standard Quantum Supermaps

We now prove that locally-applicable transformations on quantum channels are equivalent to quantum supermaps. We have already proven that all quantum supermaps define locally-applicable transformations, so what remains is to prove that this assignment can be inverted, that every locally-applicable transformation defines quantum supermap. We begin by identifying a key feature of quantum theory, the existence of control for convex sets. We then give our proof in three steps, proving inheritance of convex linearity for locally-applicable transformations on convex sets, proving their unique extension to action on completely positive maps, and finally proving their realisation in terms of standard quantum supermaps. We begin by addressing the property of control.

Definition 4 (Control). *A set K has control if for every pair $\phi_0, \phi_1 \in \text{Ext}_{X, X'}(K)$ there exists $\Phi \in \text{Ext}_{X \otimes Y, X' \otimes Y'}(S)$ and a pair of states ρ_0, ρ_1 such that:*



Conveniently convex sets in \mathbf{QC} are always controlled, in fact the existence of control is equivalent to asking for closure under convex combinations.

Lemma 2. *A set $K \subseteq \mathbf{QC}(A, A')$ has control if and only if it is convex.*

Proof. We begin by showing that whenever K is convex it has control. Note that whenever K is convex then $\mathbf{Ext}_{X, X'}(K)$ is convex for every choice of X and X' . Now choose a pair of states $\rho_0, \rho_1 \in \mathbf{QC}(I, Y)$ on an object Y which are distinguishable in the sense that there exist effects $e_0, e_1 \in \mathbf{CP}(Y, I)$ satisfying $e_i \circ \rho_j = \delta_{ij}$ and then construct the following process Φ

$$\begin{array}{c} | \\ \hline \boxed{\phi} \\ \hline | \end{array} := \begin{array}{c} | \\ \hline \boxed{\phi_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{e_0} \\ \hline | \end{array} + \begin{array}{c} | \\ \hline \boxed{\phi_1} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{e_1} \\ \hline | \end{array},$$

certainly by inserting ρ_0, ρ_1 into the rightmost wire the channels ϕ_0, ϕ_1 are recovered, what remains is to show that Φ is in $\mathbf{Ext}_{X, X'}(K)$. Indeed consider checking the reduction of Φ given by applying an arbitrary state and effect of \mathbf{QC} to its auxiliary wires, given that \mathbf{QC} is causal this is given by:

$$\begin{array}{c} | \\ \hline \boxed{\phi_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{e_0} \\ \hline | \end{array} + \begin{array}{c} | \\ \hline \boxed{\phi_1} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{e_1} \\ \hline | \end{array} \begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{\rho} \\ \hline | \end{array}$$

for some ρ . Now note that each of the post selected states is a normalised state equipped with a probability, for instance:

$$\begin{array}{c} | \\ \hline \boxed{e_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{\rho} \\ \hline | \end{array} = \begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{e_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{\rho} \\ \hline | \end{array} \times \frac{1}{\begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{e_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{\rho} \\ \hline | \end{array}}{\begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{e_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{\rho} \\ \hline | \end{array}} \times \begin{array}{c} | \\ \hline \boxed{e_0} \\ \hline | \end{array} \begin{array}{c} | \\ \hline \boxed{\rho} \\ \hline | \end{array} = p(e_0|\rho) \times \rho_{|0}$$

and similarly for e_1 . Since ϕ_0, ϕ_1 are elements of $\mathbf{Ext}_{X, X'}(K)$ it then follows from the above that the reduction of Φ is a convex combination of elements of K , explicitly the application of arbitrary state and effect to Φ returns:

$$p(e_0|\rho) \begin{array}{c} | \\ \hline \boxed{\phi_0} \\ \hline | \end{array} \begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{\rho_{|0}} \\ \hline | \end{array} + p(e_1|\rho) \begin{array}{c} | \\ \hline \boxed{\phi_1} \\ \hline | \end{array} \begin{array}{c} \bar{\vdash} \\ | \\ \hline \boxed{\rho_{|1}} \\ \hline | \end{array}$$

where $p(e_0|\rho) + p(e_1|\rho) = 1$. We now check the converse, that when K has control it is convex. Indeed, for any $\phi_0, \phi_1 \in K$ choose their control operation $\Phi \in \mathbf{Ext}_{Y, I}(K)$. Consider an arbitrary convex combination $p\phi_0 + (1-p)\phi_1$, this combination is given by inserting $\sigma := p\rho_0 + (1-p)\rho_1$ into the wire Y of Φ . Since Φ is in $\mathbf{Ext}_{Y, I}$ then insertion of σ into Φ must return an element of K and so it follows that the convex combination $p\phi_0 + (1-p)\phi_1$ is an element of K . \square

This equivalence is noted to furthermore hold for classical information theory in the appendix, by an identical proof method. We finish by stating a definition we will find convenient to reference later.

Definition 5. *The operational closure $K_{\mathbf{CP}}$ of a set $K \subseteq \mathbf{QC}(A, A')$ is given by the set of all:*

$$\begin{array}{c} | \\ \hline \boxed{\sigma} \\ \hline | \\ \hline \boxed{\Phi} \\ \hline | \\ \hline \boxed{\rho} \\ \hline | \end{array}$$

where $\Phi \in \mathbf{Ext}_{X, X'}(K)$, $\rho \in \mathbf{CP}(I, X)$, and the effect $\sigma \in \mathbf{CP}(X', I)$ for some X, X' .

The operational closure $K_{\mathbf{CP}}$ of K represents the set of all operations onto which locally-applicable transformations on K can be uniquely extended.

4.1 Proof of main theorem

In this section we prove that for any locally-applicable transformation S of type $\mathcal{E}_{path}(\tau) \rightarrow \mathcal{E}_{path}(\lambda)$ on \mathbf{QC} there exists a \mathbf{CP} -supermap S_Q of type $\mathcal{E}_{sig}(\tau) \rightarrow \mathcal{E}_{sig}(\lambda)$ which implements it, in the sense that such that $S = \mathcal{F}_{\mathbf{CP}}(S_Q)$ where $\mathcal{F}_{\mathbf{CP}}$ is the previously defined embedding from \mathbf{CP} -supermaps into locally-applicable transformations. We will note in passing that as a consequence we will have constructed an *equivalence of categories* between quantum supermaps on signalling constraints and locally-applicable transformations on pathing constraints, the latter definition is void of any reference to compact closure, linearity, coarse-graining, or even causal structure. The equivalence will in fact hold for all locally-applicable transformations $K \rightarrow M$ between convex sets K, M whether or not they be associated to signalling constraints. We begin by deriving convex linearity from locality in a general setting.

Lemma 3 (Convex linearity). *Let K, M be convex, then every locally-applicable transformation of type $S : K \rightarrow M$ on \mathbf{QC} preserves convex combinations.*

Proof. Consider a pair $\phi_i \in \mathbf{Ext}_{X, X'}(K)$ of channels, since K is convex it has control, there exists $\Phi \in \mathbf{Ext}_{X, X'}(K)$ such that

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_i \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{\phi_i} \\ | \\ \text{---} \end{array}$$

and so an arbitrary convex combination $p\phi_o + (1-p)\phi_1$ can be written as

$$p\phi_o + (1-p)\phi_1 = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_p \end{array}$$

With $\rho_p := p\rho_o + (1-p)\rho_1$. Now writing $p_1 = (1-p)$ then $S_{X, X'}(p_0\phi_0 + p_1\phi_1)$ is given using naturality by:

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{p\phi_o + (1-p)\phi_1} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_p \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_p \end{array}$$

Then rewriting ρ in terms of ρ_0, ρ_1 gives

$$= p_0 \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_0 \end{array} + p_1 \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_1 \end{array},$$

then using naturality again

$$= p_0 \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_0 \end{array} + p_1 \begin{array}{c} \text{---} \\ | \\ \text{---} \\ \boxed{S_{X, X'}} \\ | \\ \text{---} \\ \boxed{\phi} \\ | \\ \text{---} \\ \rho_1 \end{array}$$

and finally using the definition of control recovers the result

$$= p_0 \begin{array}{c} \boxed{S_{X,X'}} \\ \boxed{\phi_0} \end{array} + p_1 \begin{array}{c} \boxed{S_{X,X'}} \\ \boxed{\phi_1} \end{array} = p_0 S_{X,X'}(\phi_0) + p_1 S_{X,X'}(\phi_1)$$

and so $S_{X,X'}$ is indeed convex linear. \square

We note that S consequently has a unique extension to the real-linear span of K . We now show using this result that locally-applicable transformations on $\mathbf{QC}(A, A')$ can be uniquely extended to $\mathbf{CP}(A, A')$, more generally for any set K the map S can be extended to the operational closure $K_{\mathbf{CP}}$ of K .

Lemma 4 (Extension to operational closure). *Let K, M be convex, then every locally-applicable transformation of type $S : K \rightarrow M$ on \mathbf{QC} has a unique extension to a function $S_{\mathbf{CP}} : K_{\mathbf{CP}} \rightarrow M_{\mathbf{CP}}$.*

Proof. We give a candidate definition and then show that it is well-posed, consider some $\phi \in K_{\mathbf{CP}}$, for this ϕ there exists $\Phi \in \text{Ext}_{X,X'}(K)$ and ρ, σ such that:

$$\begin{array}{c} \boxed{\phi} \end{array} = \begin{array}{c} \boxed{\Phi} \\ \sigma \\ \rho \end{array}$$

With respect to this choice we can define

$$\begin{array}{c} \boxed{S_{\mathbf{CP}}} \\ \boxed{\Phi} \\ \sigma \\ \rho \end{array} := \begin{array}{c} \boxed{S_{X,X'}} \\ \boxed{\Phi} \\ \sigma \\ \rho \end{array}$$

We now show that this definition is well-posed by showing that for any other choice of Φ, ρ, σ the resulting outcome $S_{\mathbf{CP}}(\phi)$ would be the same. Indeed, let

$$\begin{array}{c} \boxed{\Phi} \\ \sigma \\ \rho \end{array} = \begin{array}{c} \boxed{\Phi'} \\ \sigma' \\ \rho' \end{array}$$

then consider the process Σ defined by

$$\begin{array}{c} \boxed{\Sigma} \end{array} = \begin{array}{c} \boxed{0} \\ \sigma \end{array} + \begin{array}{c} \boxed{1} \\ \ddagger \\ - \sigma \end{array}$$

where the output is taken to be at least 2-dimensional. A process Σ' can be defined similarly, note that both Σ, Σ' are trace preserving and so are members of \mathbf{QC} , meaning that they can be slid along dotted wires. We now consider the result of applying them to Φ, Φ' :

$$\begin{array}{c} \boxed{\Sigma} \\ \boxed{\Phi} \\ \rho \end{array} = \begin{array}{c} \boxed{0} \\ \sigma \\ \boxed{\Phi} \\ \rho \end{array} + \begin{array}{c} \boxed{1} \\ \ddagger \\ \boxed{\Phi} \\ \rho \end{array} - \begin{array}{c} \boxed{1} \\ \sigma \\ \boxed{\Phi} \\ \rho \end{array}$$

This in turn implies that

and so then by linearity

We begin by showing that σ, σ' can be safely pulled across dotted wires maintaining equality even though they are not quantum channels, Indeed consider the difference, and express in terms of Σ, Σ' :

Now since Σ, Σ' are quantum channels they can be pulled through dotted lines, after which we use equation (1):

Now the preparation $1 : I \rightarrow X'$ is a quantum channel it can be pulled back through dotted wires, after which orthogonality implies that the difference has to be 0.

Since the difference is 0 then it follows that

We can now consider the bottom side, which is easier to reason with since every $\rho \in \mathbf{CP}(I, X)$ is given by $a\eta$ with η a state in $\mathbf{QC}(I, X)$. Indeed, using the extension of S to \mathbb{R} -linearity (and so to multiplication by scalar α) gives

and similarly for the ϕ', ρ', σ' we find


All together then, using the separate deductions made for ρ and σ gives

and so $S_{\mathbf{CP}}$ is indeed well defined. \square


The above is the key to our result, we now are ready to construct a candidate \mathbf{CP} -supermap for simulating the action of our locally-applicable transformation $S_{X, X'}$ by tensor extension with identities on X, X' . To do so we apply our locally-applicable transformation S to the swap-morphism, the intuition being that the swap gives a way to noiselessly extract information about the input behaviour of a higher order map, by converting its input into a pair of lower-order objects.

Theorem 1 (Re-characterisation of supermaps). *Let K, M be convex subsets of channels of \mathbf{QC} , there is a one-to-one correspondence between the \mathbf{CP} -supermaps of type $K \rightarrow M$ and the locally-applicable transformations of type $K \rightarrow M$*


Proof. Given a locally-applicable transformation S of type $K \rightarrow M$ on \mathbf{QC} with $K \subseteq \mathbf{QC}(A, A')$ and $M \in \subseteq \mathbf{QC}(B, B')$ we define $S_Q : A^* \otimes A' \rightarrow B^* \otimes B'$ by:




In other words we apply S to the swap in \mathbf{QC} and then embed into \mathbf{CP} so that we may apply caps and cups. We now consider the application of arbitrary states and effects ρ, σ in \mathbf{CP} to the auxiliary wires, and use the tensor separation property:



We now use the well-posed definition of $S_{\mathbf{CP}}$ and then use compact closure to replace the cup and cap with the identity to give:



Since this is true for all ρ, σ in \mathbf{CP} it follows that:



and so there indeed exists a quantum supermap of type $S_Q : K \rightarrow M$ such that $\mathcal{F}_{\mathbf{CP}}(S_Q) = S$ where $\mathcal{F}_{\mathbf{CP}}$ is the previously defined embedding of \mathbf{CP} -supermaps on \mathbf{QC} into locally-applicable transformations on \mathbf{QC} . \square

Note that in \mathbf{QC} the set $\mathcal{E}_{sig}(\tau)$ is convex and that furthermore $\mathcal{E}_{sig}(\tau) = \mathcal{E}_{path}(\tau)$. It follows then that for arbitrary relations τ the locally-applicable transformations of type $\mathcal{E}_{path}(\tau) \rightarrow \mathcal{E}_{path}(\lambda)$ characterize the quantum supermaps of type $\mathcal{E}_{sig}(\tau) \rightarrow \mathcal{E}_{sig}(\lambda)$. As a stricter corollary the quantum supermaps on non-signalling channels, also referred to in the literature as process matrices have been characterised from principles of compositionality, without reference to causality or preferred time direction. To phrase this concisely we refer the morphisms in:

$$\mathcal{E}_{path}\left(\begin{array}{c} \uparrow \\ \uparrow \end{array}\right),$$

as non-pathing morphisms and refer to supermaps and locally-applicable transformations of type $K \rightarrow \mathbf{C}(B, B')$ for some B, B' as being *on* the set K .

Corollary 1. *There is a one-to-one correspondence between quantum supermaps on the set of non-signalling channels and locally-applicable transformations on the set of non-pathing channels.*

Other example of convex sets of interest for which supermaps are now consequently characterised in terms of locality are those convex sets which are specified by sectorial constraints [48, 63, 77, 78] which find use in the phrasing of more complex instances of pathing constraints [61] and in the analysis of fine-grained causal structures [47, 48]. Finally, we comment that by denoting the restriction of the category $\mathbf{CPsup}[\mathbf{QC}]$ to convex sets by $\mathbf{CPsup}[\mathbf{QC}]_{con} := \mathbf{QS}_{con}$ and similarly denoting restriction of the category $\mathbf{lm}[\mathbf{C}]$ to convex sets by $\mathbf{lm}[\mathbf{C}]_{con}$ we can state the one-to-one correspondence in concise categorical language.

Corollary 2. *There is an equivalence of categories $\mathbf{QS}_{con} \cong \mathbf{lm}[\mathbf{QC}]_{con}$*

This equivalence is confirmed by noting that the assignment $\mathcal{F}_{\mathbf{CP}}$ is functorial, surjective on objects, and full and faithful by the one-to-one correspondence observed in this section. In summary, quantum supermaps and equivalently process matrices, originally defined in terms of Choi-isomorphisms and probabilistic structure respectively, are instances of a purely compositional definition of higher order mapping of quantum channels.

5 Summary and Outlook

A definition of locally-applicable transformation is introduced which refers only to the circuit-theoretic structure of deterministic quantum information theory. By being purely compositional in nature it may be applied to arbitrary symmetric monoidal categories and stated concisely in the language of category theory using the notion of a natural transformation. When applied to quantum channels, including those equipped with signalling constraints, locally-applicable transformations are in one to one correspondence with quantum supermaps, so provide a re-axiomatisation for supermaps in terms of the principles of sequential composition, parallel composition, and locality.

The construction presented could in principle be a seed for a variety of new projects, including comparison of its infinite iteration with the construction of higher order causal categories, the free construction of various compositional features of quantum supermaps [6–8], and characterisation of locally-applicable transformations on infinite dimensional quantum theory. Less concretely and more broadly it is the authors hope that this definition of supermap is closer to allowing for a suitable generalisation to more elaborate notions of space and parallel composition [67, 68, 79] including but not limited to algebraic

quantum field theories [69], so that higher order quantum theories can be brought into closer contact with theories of quantum gravity where supermaps are hoped to provide insight by modelling of quantum causal structure [39].

Acknowledgements

MW is grateful to J Hefford, C Comfort, A Vanreitelde, N Ormrod, N Pinzani, and H Kristjánsson for useful conversations, and again to N Ormrod for reviewing an earlier draft of this manuscript. AK and GC are supported by Grant No. 61466 from the John Templeton Foundation as part of the QISS project. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. GC was supported by the Croucher Foundation and by the Hong Kong Research Grant Council (RGC) through the Senior Research Fellowship Scheme SRFS2021-7S02. MW was supported by University College London and the EPSRC Doctoral Training Centre for Delivering Quantum Technologies.

References

- [1] G. Chiribella, G. M. D’Ariano, and P. Perinotti. Transforming quantum operations: Quantum supermaps. *EPL (Europhysics Letters)*, 83(3):30004, 7 2008.
- [2] G. Chiribella, G. M. D’Ariano, and P. Perinotti. Quantum circuit architecture. *Physical Review Letters*, 101(6):060401, 8 2008. arxiv:0712.1325.
- [3] Giulio Chiribella, Giacomo Mauro D’Ariano, Paolo Perinotti, and Benoit Valiron. Quantum computations without definite causal structure. *Physical Review A - Atomic, Molecular, and Optical Physics*, 88(2):022318, 8 2013. arxiv:0912.0195.
- [4] Giulio Chiribella, Giacomo M. D’Ariano, and Paolo Perinotti. Theoretical framework for quantum networks. *Physical Review A - Atomic, Molecular, and Optical Physics*, 80(2), 4 2009. arxiv:0904.4483v2.
- [5] G. Chiribella, A. Toigo, and V. Umanità. Normal completely positive maps on the space of quantum operations. *Open Systems and Information Dynamics*, 20(1), 12 2010. arxiv:1012.3197v3.
- [6] Alessandro Bisio and Paolo Perinotti. Theoretical framework for higher-order quantum theory. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 475(2225):20180706, 5 2019.
- [7] Aleks Kissinger and Sander Uijlen. A categorical semantics for causal structure. volume 15. *Logical Methods in Computer Science*, 2019. arxiv:1701.04732.
- [8] Matt Wilson and Giulio Chiribella. A Mathematical Framework for Transformations of Physical Processes. 4 2022. arxiv:2204.04319.
- [9] Paolo Perinotti. Causal Structures and the Classification of Higher Order Quantum Computations. pages 103–127. Birkhäuser, Cham, 2017. arxiv:1612.05099.
- [10] Hlér Kristjánsson, Giulio Chiribella, Sina Salek, Daniel Ebler, and Matthew Wilson. Resource theories of communication. *New Journal of Physics*, 22(7):073014, 7 2020. arxiv:1910.08197.
- [11] Gilad Gour and Carlo Maria Scandolo. Dynamical Resources. arxiv:2101.01552v1.
- [12] Gilad Gour and Carlo Maria Scandolo. Entanglement of a bipartite channel. *Physical Review A*, 103(6):062422, 6 2021.

- [13] Daniel Ebler, Sina Salek, and Giulio Chiribella. Enhanced Communication with the Assistance of Indefinite Causal Order. *Physical Review Letters*, 120(12):120502, 3 2018.
- [14] Lorenzo M. Procopio, Francisco Delgado, Marco Enríquez, Nadia Belabas, and Juan Ariel Levenson. Communication Enhancement through Quantum Coherent Control of N Channels in an Indefinite Causal-Order Scenario. *Entropy*, 21(10):1012, 10 2019.
- [15] Lorenzo M. Procopio, Francisco Delgado, Marco Enríquez, Nadia Belabas, and Juan Ariel Levenson. Sending classical information via three noisy channels in superposition of causal orders. *Physical Review A*, 101(1):012346, 1 2020. arxiv:1910.11137.
- [16] Giulio Chiribella, Manik Banik, Some Sankar Bhattacharya, Tamal Guha, Mir Alimuddin, Arup Roy, Sutapa Saha, Sristy Agrawal, and Guruprasad Kar. Indefinite causal order enables perfect quantum communication with zero capacity channels. *New Journal of Physics*, 2 2021. arxiv:1810.10457v2.
- [17] Giulio Chiribella, Matthew Wilson, and H. F. Chau. Quantum and Classical Data Transmission Through Completely Depolarising Channels in a Superposition of Cyclic Orders. *arXiv*, 5 2020. arxiv:2005.00618.
- [18] Matthew Wilson and Giulio Chiribella. A Diagrammatic Approach to Information Transmission in Generalised Switches. 3 2020. arxiv:2003.08224.
- [19] Marco Túlio Quintino, Qingxiuxiong Dong, Atsushi Shimbo, Akihito Soeda, and Mio Muraō. Probabilistic exact universal quantum circuits for transforming unitary operations. *Physical Review A*, 100(6):062339, 12 2019. arxiv:1909.01366.
- [20] Marco Túlio Quintino, Qingxiuxiong Dong, Atsushi Shimbo, Akihito Soeda, and Mio Muraō. Reversing Unknown Quantum Transformations: Universal Quantum Circuit for Inverting General Unitary Operations. *Physical Review Letters*, 123(21):210502, 11 2019.
- [21] Qingxiuxiong Dong, Marco Túlio Quintino, Akihito Soeda, and Mio Muraō. Successor-Draw: A Strategy Allowing Repeat-Until-Success in Quantum Computation. *Physical Review Letters*, 126(15):150504, 4 2021.
- [22] Philippe Allard Guérin, Adrien Feix, Mateus Araújo, and Časlav Brukner. Exponential Communication Complexity Advantage from Quantum Superposition of the Direction of Communication. *Physical Review Letters*, 117(10):100502, 9 2016. arxiv:1605.07372.
- [23] Giulio Chiribella. Perfect discrimination of no-signalling channels via quantum superposition of causal structures. *Physical Review A - Atomic, Molecular, and Optical Physics*, 86(4):040301, 10 2012. arxiv:1109.5154.
- [24] Mateus Araújo, Fabio Costa, and Časlav Brukner. Computational advantage from quantum-controlled ordering of gates. *Physical Review Letters*, 113(25):250402, 12 2014. arxiv:1401.8127.
- [25] Adrien Feix, Mateus Araújo, and Časlav Brukner. Quantum superposition of the order of parties as a communication resource. *Physical Review A - Atomic, Molecular, and Optical Physics*, 92(5):052326, 11 2015.
- [26] Mario Román. Open Diagrams via Coend Calculus. pages 1–17, 4 2020. arxiv:2004.04526.

- [27] Mario Román. Comb Diagrams for Discrete-Time Feedback. Technical report, 2020. arxiv:2003.06214v1.
- [28] Guillaume Boisseau, Chad Nester, and Mario Roman. Cornering Optics. 5 2022. 10.48550/arxiv.2205.00842.
- [29] Jules Hedges. Coherence for lenses and open games. 4 2017. 10.48550/arxiv.1704.02230.
- [30] Mitchell Riley. Categories of Optics. 9 2018. arxiv:1809.00738.
- [31] Neil Ghani, Jules Hedges, Viktor Winschel, and Philipp Zahn. Compositional game theory. *Proceedings - Symposium on Logic in Computer Science*, pages 472–481, 3 2016. 10.48550/arxiv.1603.04641.
- [32] Felix A. Pollock, César Rodríguez-Rosario, Thomas Frauenheim, Mauro Paternostro, and Kavan Modi. Non-Markovian quantum processes: complete framework and efficient characterisation. *Physical Review A*, 97(1), 12 2015. 10.1103/PhysRevA.97.012127.
- [33] James Hefford and Cole Comfort. Coend Optics for Quantum Combs. 5 2022.
- [34] G. Chiribella, G. M. D’Ariano, P. Perinotti, and B. Valiron. Quantum computations without definite causal structure. *Physical Review A - Atomic, Molecular, and Optical Physics*, 88(2), 12 2009. arxiv:0912.0195.
- [35] Mateus Araújo, Fabio Costa, and Časlav Brukner. Computational advantage from quantum-controlled ordering of gates. *Physical Review Letters*, 113(25):250402, 12 2014. arxiv:1401.8127.
- [36] Mateus Araújo, Philippe Allard Guérin, and Ämin Baumeler. Quantum computation with indefinite causal structures. *Physical Review A*, 96(5), 6 2017. 10.1103/PhysRevA.96.052315.
- [37] Martin J. Renner and Časlav Brukner. Reassessing the computational advantage of quantum-controlled ordering of gates. *Physical Review Research*, 3(4), 2 2021. 10.1103/PhysRevResearch.3.043012.
- [38] David Felce and Vlatko Vedral. Quantum Refrigeration with Indefinite Causal Order. *Physical Review Letters*, 125(7):070603, 8 2020.
- [39] Lucien Hardy. Towards Quantum Gravity: A Framework for Probabilistic Theories with Non-Fixed Causal Structure. *Journal of Physics A: Mathematical and Theoretical*, 40(12):3081–3099, 8 2006. 10.1088/1751-8113/40/12/S12.
- [40] Ognjan Oreshkov, Fabio Costa, and Časlav Brukner. Quantum correlations with no causal order. *Nature Communications*, 3, 2012. arxiv:1105.4464.
- [41] Ämin Baumeler and Stefan Wolf. Perfect signaling among three parties violating predefined causal order. *IEEE International Symposium on Information Theory - Proceedings*, pages 526–530, 12 2013. 10.1109/ISIT.2014.6874888.
- [42] Ämin Baumeler and Stefan Wolf. The space of logically consistent classical processes without causal order. *New Journal of Physics*, 18(1), 7 2015. 10.1088/1367-2630/18/1/013036.
- [43] Esteban Castro-Ruiz, Flaminia Giacomini, and Časlav Brukner. Dynamics of Quantum Causal Structures. *Physical Review X*, 8(1):011047, 3 2018.

- [44] Veronika Baumann, Marius Krumm, Philippe Allard Guérin, and Časlav Brukner. Noncausal Page-Wootters circuits. *Physical Review Research*, 4(1), 5 2021. 10.1103/PhysRevResearch.4.013180.
- [45] Fabio Costa. A no-go theorem for superpositions of causal orders. *Quantum*, 6:663, 3 2022. 10.22331/q-2022-03-01-663.
- [46] Ralph Silva, Yelena Guryanova, Anthony J. Short, Paul Skrzypczyk, Nicolas Brunner, and Sandu Popescu. Connecting processes with indefinite causal order and multi-time quantum states. *New Journal of Physics*, 19(10), 1 2017. 10.1088/1367-2630/aa84fe.
- [47] V. Vilasini and Renato Renner. Embedding cyclic causal structures in acyclic space-times: no-go results for process matrices. 3 2022. 10.48550/arxiv.2203.11245.
- [48] Nick Ormrod, Augustin Vanrietvelde, and Jonathan Barrett. Causal structure in the presence of sectorial constraints, with application to the quantum switch. 4 2022. arxiv:2204.10273.
- [49] David Felce, Nicetu Tibau Vidal, Vlatko Vedral, and Eduardo O. Dias. Indefinite Causal Orders from Superpositions in Time. 7 2021. 10.48550/arxiv.2107.08076.
- [50] Ognjan Oreshkov and Christina Giarmatzi. Causal and causally separable processes. *New Journal of Physics*, 18(9):093020, 9 2016. 10.1088/1367-2630/18/9/093020.
- [51] Julian Wechs, Hippolyte Dourdent, Alastair A. Abbott, and Cyril Branciard. Quantum circuits with classical versus quantum control of causal order. *PRX Quantum*, 2(3), 1 2021. 10.1103/PRXQuantum.2.030335.
- [52] Tom Purves and Anthony J. Short. Quantum theory cannot violate a causal inequality. *Physical Review Letters*, 127(11), 1 2021. 10.1103/PhysRevLett.127.110402.
- [53] Man Duen Choi. Completely positive linear maps on complex matrices. *Linear Algebra and Its Applications*, 10(3):285–290, 1975.
- [54] Bob Coecke and Aleks Kissinger. *Picturing quantum processes: A first course in quantum theory and diagrammatic reasoning*. Cambridge University Press, 3 2017.
- [55] Chris Heunen and Jamie Vicary. Categories for Quantum Theory. *Categories for Quantum Theory*, 11 2019.
- [56] Samson Abramsky and Bob Coecke. A categorical semantics of quantum protocols. In *Proceedings - Symposium on Logic in Computer Science*, volume 19, pages 415–425, 2004. arxiv:quant-ph/0402130.
- [57] Bob Coecke. Kindergarten quantum mechanics: Lecture notes. In *AIP Conference Proceedings*, volume 810, pages 81–98, 1 2006.
- [58] Bob Coecke. Quantum pictorialism. *Contemporary Physics*, 51(1):59–83, 1 2010. arxiv:0908.1787.
- [59] John H. Selby, Carlo Maria Scandolo, and Bob Coecke. Reconstructing quantum theory from diagrammatic postulates. 2 2018. arxiv:1802.00367.
- [60] Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer New York, 1971. 10.1007/978-1-4612-9839-7.
- [61] Robin Lorenz and Jonathan Barrett. Causal and compositional structure of unitary transformations. *arXiv*, 1 2020. arxiv:2001.07774.
- [62] Jonathan Barrett, Robin Lorenz, and Ognjan Oreshkov. Cyclic Quantum Causal Models. Technical report. arxiv:2002.12157v2.

- [63] Augustin Vanrietvelde, Hlér Kristjánsson, and Jonathan Barrett. Routed quantum circuits. Technical report. arxiv:2001.07774.
- [64] Andre Joyal and Dominic Verity. Traced monoidal categories. *Math. Proc. Camb. Phil. Soc.*, 119:447–451, 2021.
- [65] Masahito Hasegawa. Recursion from Cyclic Sharing: Traced Monoidal Categories and Models of Cyclic Lambda Calculi. *Springer Verlag*, page 196213, 1997.
- [66] Nicola Pinzani, Stefano Gogioso, and Bob Coecke. Categorical Semantics for Time Travel. *Proceedings - Symposium on Logic in Computer Science*, 2019-June, 1 2019. arxiv:1902.00032.
- [67] Stefano Gogioso. A Process-Theoretic Church of the Larger Hilbert Space. 5 2019. arxiv:1905.13117.
- [68] Pablo Arrighi, Amélia Durbec, and Matt Wilson. Quantum networks theory. 10 2021. arxiv:2110.10587.
- [69] Rudolf Haag and Daniel Kastler. An Algebraic approach to quantum field theory. *J.Math.Phys.*, 5(7):848–861, 1964.
- [70] John E. Roberts. Localization in algebraic field theory. *Communications in Mathematical Physics*, 85(1):87–98, August 1982.
- [71] Sergio Doplichera. The principle of locality: Effectiveness, fate, and challenges. *Journal of Mathematical Physics*, 51(1):015218, 1 2010.
- [72] Romeo Brunetti, Klaus Fredenhagen, and Rainer Verch. The generally covariant locality principle – A new paradigm for local quantum physics. *Communications in Mathematical Physics*, 237(1-2):31–68, 12 2001. arxiv:math-ph/0112041v1.
- [73] F. Strocchi. Relativistic Quantum Mechanics and Field Theory. *Foundations of Physics*, 34(3):501–527, 1 2004. arxiv:hep-th/0401143v1.
- [74] Bob Coecke and Raymond Lal. Causal Categories: Relativistically Interacting Processes. *Foundations of Physics*, 43(4):458–501, 4 2013. arxiv:1107.6019.
- [75] T. Eggeling, D. Schlingemann, and R. F. Werner. Semicausal operations are semilocalizable. *Europhysics Letters*, 57(6):782–788, 2002. arxiv:quant-ph/0104027.
- [76] Paolo Perinotti. Causal influence in operational probabilistic theories. *Quantum*, 5:515, Aug 2021.
- [77] Matt Wilson and Augustin Vanrietvelde. Composable constraints. 12 2021. arxiv:2112.06818.
- [78] Augustin Vanrietvelde and Giulio Chiribella. Universal control of quantum processes using sector-preserving channels. *Quantum Information and Computation*, 21(15-16):1320–1352, dec 2021.
- [79] Pau Enrique Moliner, Chris Heunen, and Sean Tull. Space in Monoidal Categories. *Electronic Proceedings in Theoretical Computer Science, EPTCS*, 266:399–410, 4 2017. arxiv:1704.08086v3.

A Category Theory

Here we introduce three key concepts in abstract algebra, categories, functors, and natural transformations. Avoiding issues of size, a category is a set of objects A, B, \dots with for each pair A, B a set $\mathbf{C}(A, B)$ of morphisms often denoted as $f : A \rightarrow B$. A category is

furthermore equipped with, for each triple (A, B, C) of objects a composition function $\circ : \mathbf{C}(A, B) \times \mathbf{C}(B, C) \rightarrow \mathbf{C}(A, C)$ which is associative so that $f \circ (g \circ h) = (f \circ g) \circ h$, and for each object A a unit id_A so that for all $f : A \rightarrow B$ then $f \circ id_A = f = id_B \circ f$. In graphical terms a morphism $f : A \rightarrow A'$ may be written as a box labelled f with input wire A and output wire A' , sequential (\circ) composition represented by:

$$g \circ f := \begin{array}{c} \boxed{g} \\ \boxed{f} \end{array} \quad \text{and} \quad id = \begin{array}{c} | \\ | \\ | \end{array}$$

The identity box, can be diagrammatically represented as a wire. This notation has the convenient property that it absorbs the structural equations of a category. There is no graphical difference for instance between the diagram representing $(f \circ g) \circ h$ and the diagram representing $f \circ (g \circ h)$. A functor $\mathcal{F} : \mathbf{C} \rightarrow \mathbf{D}$ is a structure preserving map between categories, formally it is an assignment of an object $\mathcal{F}A$ to each object A along with for each pair A, B a function $\mathcal{F}_{AB} : \mathbf{C}(A, B) \rightarrow \mathbf{D}(\mathcal{F}A, \mathcal{F}B)$ which preserves composition in the sense that $\mathcal{F}(f \circ g) = \mathcal{F}(f) \circ \mathcal{F}(g)$ and $\mathcal{F}(i) = i$. Graphically a functor can be represented by a surrounding box, which satisfies box-merging and identity removal:

$$\begin{array}{c} \boxed{\mathcal{F}} \\ \boxed{f} \\ \boxed{\mathcal{F}} \\ \boxed{g} \end{array} = \begin{array}{c} \boxed{\mathcal{F}} \\ \boxed{f} \\ \boxed{g} \end{array} \quad \boxed{\mathcal{F}} = \begin{array}{c} | \\ | \\ | \end{array}$$

A natural transformation $\eta : \mathcal{F} \Rightarrow \mathcal{G}$ is for every object A a morphism $\eta_A : \mathcal{F}A \rightarrow \mathcal{G}A$ such that for every $f : A \rightarrow B$ then $\eta_B \circ \mathcal{F}(f) = \mathcal{G}(f) \circ \eta_A$. Graphically this reads as:

$$\begin{array}{c} \boxed{\eta_B} \\ \boxed{\mathcal{F}} \\ \boxed{f} \end{array} = \begin{array}{c} \boxed{\mathcal{G}} \\ \boxed{f} \\ \boxed{\eta_A} \end{array}$$

For any category \mathbf{C} the (*reverse*) opposite category \mathbf{C}^{op} can be defined in which $\mathbf{C}^{op}(A, B) := \mathbf{C}(B, A)$ with composition and identity inherited from \mathbf{C} so that $f \circ_{op} g := g \circ f$. Finally for any categories \mathbf{C}, \mathbf{D} the product category $\mathbf{C} \times \mathbf{D}$ can be defined in which objects are given by pairs (A, B) and morphisms given by pairs of morphisms (f, g) and composition rule inherited from \mathbf{C} and \mathbf{D} as $(f, g) \circ (f', g') := (f' \circ f, g' \circ g)$.


An example of a natural transformation of interest in this paper is given intuitively by the following: a family of functions $\eta_{A,B} : \mathbf{C}(A, B) \rightarrow \mathbf{C}(A, B)$ which commute with pre and post-composition by arbitrary processes, meaning that for all $f : A' \rightarrow A$ and $g : B \rightarrow B'$:

$$\eta_{A',B'}(g \circ \phi \circ f) = g \circ \eta_{A,B}(\phi) \circ f$$

In this paper we choose to represent such families graphically by introducing a function-box notation. We represent $\eta_{A,B}$ by

$$\eta_{A,B} \approx \begin{array}{c} B \\ \boxed{\eta_{A,B}} \\ A \end{array} \quad \eta_{A,B}(\phi) \approx \begin{array}{c} \boxed{\eta_{A,B}} \\ \boxed{\phi} \end{array}$$

The dotted wires are used to suggest the commutation condition satisfied by the family $\eta_{A,B}$:



Such a family of functions can be phrased as a natural transformation in the following way, we begin with definition of the “hom functor” $\mathbf{C}(-, =) : \mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{Set}$. The category $\mathbf{C}^{op} \times \mathbf{C}$ has as objects pairs (A, B) of objects of \mathbf{C} , for morphisms $(A, B) \rightarrow (A', B')$ it has pairs $(f : A' \rightarrow A, g : B \rightarrow B')$. A functor of type $\mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{Set}$ must send each pair (A, B) to a set, indeed the functor $\mathbf{C}(-, =)$ is defined as sending the object (A, B) to the set $\mathbf{C}(A, B)$. A functor of type $\mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{Set}$ must send each pair $(f : A' \rightarrow A, g : B \rightarrow B')$ to a function $\mathbf{C}(A, B) \rightarrow \mathbf{C}(A', B')$. Indeed the functor $\mathbf{C}(-, =)$ is defined by sending each pair (f, g) to the function

$$\begin{aligned} \mathbf{C}(f, g) : \mathbf{C}(A, B) &\rightarrow \mathbf{C}(A', B') \\ \mathbf{C}(f, g)(\phi) &= g \circ \phi \circ f \end{aligned}$$

Indeed this is a functor since $\mathbf{C}(i, i)(\phi) = \phi$ and $\mathbf{C}(f \circ f', g' \circ g)(\phi) = g' \circ g \circ \phi \circ f \circ f' = \mathbf{C}(f', g')(\mathbf{C}(f, g)(\phi)) = (\mathbf{C}(f', g') \circ \mathbf{C}(f, g))(\phi)$. A natural transformation of type $\eta : \mathbf{C}(-, =) \Rightarrow \mathbf{C}(-, =)$ is a family of functions $\eta_{A,B} : \mathbf{C}(A, B) \rightarrow \mathbf{C}(A, B)$ such that $\eta_{A,B} \circ \mathbf{C}(f, g) = \mathbf{C}(f, g) \circ \eta_{A,B}$, which when applied as functions reads $\forall \phi : \eta_{A',B'}(g \circ \phi \circ f) = g \circ \eta_{A,B}(\phi) \circ f$, our original commutativity condition.

B Phrasing of localizability as naturality

The previously introduced hom-functor $\mathbf{C}(-, =)$ can be generalised in monoidal categories to a functor

$$\mathbf{C}(A \otimes -, A' \otimes =) : \mathbf{C}^{op} \times \mathbf{C} \longrightarrow \mathbf{SET}$$

which assigns to each (X, X') the object $\mathbf{C}(A \otimes X, A' \otimes X')$ and to each morphism $f : Y \rightarrow X$ and each morphism $g : X' \rightarrow Y'$ the function

$$\begin{aligned} \mathbf{C}(A \otimes f, A' \otimes g) : \mathbf{C}(A \otimes X, A' \otimes X') &\rightarrow \mathbf{C}(A \otimes Y, A' \otimes Y') \\ \mathbf{C}(A \otimes f, A' \otimes g) :: \phi &\mapsto (i \otimes g) \circ \phi \circ (i \otimes f) \end{aligned}$$

This functor can be further generalised to a functor $\mathbf{Ext}(K)(-, =)$, which we now define.

Definition 6 (Extension functor). *For every $K \subseteq \mathbf{C}(A, A')$ in a symmetric monoidal category \mathbf{C} one can define a functor $\mathbf{Ext}(K)(-, =) : \mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{Set}$ given by*

- $\mathbf{Ext}(K)(X, X') := \mathbf{Ext}_{X, X'}(K)$
- $\mathbf{Ext}(K)(f, g) : \mathbf{Ext}_{X, X'}(K) \rightarrow \mathbf{Ext}_{Y, Y'}(K)$ defined by

$$\mathbf{Ext}(K)(f, g)(\phi) := \text{Diagram}$$

The functor $\mathbf{C}(A \otimes -, A' \otimes =)$ can be defined as the special case given by $\mathbf{Ext}(\mathbf{C}(A, A'))$. \mathbf{Ext} is well defined, whenever $\phi \in \mathbf{Ext}_{X, X'}(S)$ then $\mathbf{Ext}(S)(f, g)(\phi) \in \mathbf{Ext}_{Y, Y'}(S)$ since for each f, g and ρ, σ then

$$\begin{array}{c} \sigma \\ | \\ g \\ | \\ \Phi \\ | \\ f \\ | \\ \rho \end{array} = \begin{array}{c} \sigma' \\ | \\ \Phi \\ | \\ \rho' \end{array} \in K$$

since $\phi \in \mathbf{Ext}_{X, X'}(K)$. Furthermore the assignment $\mathbf{Ext}(S)(f, g)$ is functorial since

$$\mathbf{Ext}(S)(f', g')(\mathbf{Ext}(S)(f, g)(\phi)) = \begin{array}{c} g' \\ | \\ g \\ | \\ \Phi \\ | \\ f \\ | \\ f' \end{array} = \mathbf{Ext}(S)(f \circ f', g' \circ g)(\phi)$$

The definition of a locally-applicable transformation can be rephrased as the definition of a natural transformation respect to this functor. A natural transformation $S : \mathbf{C}(A \otimes -, A' \otimes =) \rightarrow \mathbf{C}(B \otimes -, B' \otimes =)$ will be any family of functions $S_{X, X'}$ making the following diagram commute for all f, g :

$$\begin{array}{ccc} \mathbf{C}(A \otimes X, A' \otimes X') & \xrightarrow{S_{X, X'}} & \mathbf{C}(B \otimes X, B' \otimes X') \\ \mathbf{C}(A \otimes f, A' \otimes g) \downarrow & & \downarrow \mathbf{C}(B \otimes f, B' \otimes g) \\ \mathbf{C}(A \otimes Y, A' \otimes Y') & \xrightarrow{S_{Y, Y'}} & \mathbf{C}(B \otimes Y, B' \otimes Y') \end{array}$$

In other words such that $S_{Y, Y'} \circ \mathbf{C}(B \otimes f, B' \otimes g) = \mathbf{C}(B \otimes f, B' \otimes g) \circ S_{X, X'}$. Evaluated on processes ϕ , this condition reads $S_{Y, Y'}(\mathbf{C}(B \otimes f, B' \otimes g)(\phi)) = \mathbf{C}(B \otimes f, B' \otimes g)(S_{X, X'}(\phi))$, which unpacking the definition of $\mathbf{C}(B \otimes f, B' \otimes g)$ is precisely the definition of a locally-applicable transformation. This observation extends to general subsets, so that a locally-applicable transformation of type $S : K \rightarrow M$ is exactly a natural transformation of type $S_{X, X'} : \mathbf{Ext}_{X, X'}(K) \rightarrow \mathbf{Ext}_{X, X'}(M)$.

C Characterisation of Classical Supermaps

The proof methods presented in the main text can also be used to characterise supermaps on finite dimensional classical information theory. We outline the story here, noting the key features common to quantum classical information theory that were used in our proof. First we must declare what would be aimed to be proven, and what we mean by classical information theory. We define deterministic classical maps to be stochastic matrices, just as with quantum channels it is easier to first state a compact closed category from which they are constructed.

Definition 7 (Positive real matrices). *The category $\mathbf{Mat}[\mathbb{R}_+]$ of positive real matrices is given by taking as objects the positive integers \mathbb{Z}_+ and as morphisms of type $f : n \rightarrow m$ the matrices of dimension $n \times m$. Sequential composition is given by matrix multiplication and identity morphism is given by the diagonal matrix of ones.*

The category $\mathbf{Mat}[\mathbb{R}_+]$ is symmetric monoidal and furthermore compact closed.

Definition 8 (Stochastic maps). *The category \mathbf{Stoch} of stochastic maps is given by the subcategory of $\mathbf{Mat}[\mathbb{R}_+]$ which contains only those matrices with column vectors which sum to 1.*

The category \mathbf{Stoch} is symmetric monoidal and furthermore causal, the unique effect of type $n \rightarrow 1$ is given by the row vector $(1, \dots, 1)$. Using the inclusion $\mathbf{Stoch} \subseteq \mathbf{Mat}[\mathbb{R}]$ one can immediately define the $\mathbf{Mat}[\mathbb{R}]$ -supermaps on \mathbf{Stoch} . We will now observe that these are precisely the locally-applicable transformations on \mathbf{Stoch} . First, we address the equivalence between convexity and control.

Lemma 5. *A subset $K \in \mathbf{Stoch}(A, A')$ is convex if and only if it has control*

Proof. All that was required to construct the proof was the existence of an object Y with a pair of distinguishable states in the sense that $e_i \circ \rho_j = \delta_{ij}$ and the possibility to take positive sums. Sums are taken care of by $\mathbf{Mat}[\mathbb{R}_+]$ and Y may be taken to be 2. Indeed one can define $\rho_i : 1 \rightarrow 2$ by taking the k^{th} component of the column vector ρ_i to be δ_{ik} and similarly for the effects $e_j : 2 \rightarrow 1$. \square

Lemma 6. *Let $K \in \mathbf{Stoch}(A, A')$ and $M \in \mathbf{Stoch}(B, B')$ be convex subsets, then every locally-applicable transformation $S : K \rightarrow M$ is convex linear.*

Proof. Follows directly from equivalence between convexity and control. \square

We take the operational closure $K_{\mathbf{Mat}[\mathbb{R}_+]}$ of a set $K \subseteq \mathbf{Mat}[\mathbb{R}_+]$ to be defined in the equivalence way as for subsets M of \mathbf{QC} , replacing the applying of effects from \mathbf{CP} with the applying of effects from $\mathbf{Mat}[\mathbb{R}_+]$

Lemma 7. *Every locally-applicable transformation $S : K \rightarrow M$ between convex sets in \mathbf{Stoch} extends to a function $S_{\mathbf{Mat}[\mathbb{R}_+]} : K_{\mathbf{Mat}[\mathbb{R}_+]} \rightarrow M_{\mathbf{Mat}[\mathbb{R}_+]}$.*

Proof. The required elements of the proof are that $\mathbf{Mat}[\mathbb{R}_+]$ embeds into $\mathbf{Mat}[\mathbb{R}]$ so that subtractions can be defined, and that for every effect $\sigma \in \mathbf{Mat}[\mathbb{R}_+](A, I)$ there exists $\lambda \in \mathbb{R}_+$ and $\sigma' \in \mathbf{Mat}[\mathbb{R}_+](A, I)$ such that $\lambda\sigma + \sigma' = (1, \dots, 1)$ (The discard). \square

Lemma 8 (Tensor separation). *Every locally-applicable transformation of type $S : K \rightarrow M$ on \mathbf{Stoch} tensor separates.*

Proof. Follows since $\mathbf{Mat}[\mathbb{R}_+]$ has enough causal states. \square

Theorem 2. *For K, M convex in \mathbf{Stoch} there is a one-to-one correspondence between the $\mathbf{Mat}[\mathbb{R}_+]$ -supermaps of type $K \rightarrow M$ on \mathbf{Stoch} and the locally-applicable transformations of type $K \rightarrow M$ on \mathbf{Stoch} .*

Proof. Here all that is required is compact closure of $\mathbf{Mat}[\mathbb{R}_+]$, and again that $\mathbf{Mat}[\mathbb{R}_+]$ has enough causal states. \square

Again as a corollary of this theorem, the classical supermaps on non-signalling channels are characterised as locally-applicable transformations on non-pathing channels in \mathbf{Stoch} .