

Strictly irreducible Markov shifts and ergodicity of skew-product extensions

FELIX POGORZELSKI, ELIAS ZIMMERMANN

ABSTRACT. We extend a criterion of Alexander I. Bufetov for the ergodicity of skew-product extensions of Markov shifts with finite state space to Markov shifts with countably infinite state space. We also show that the criterion is in fact a characterization. Finally, we discuss applications to pointwise ergodic theorems for Cesàro-type averages of transformations.

We consider a finite or countably infinite bundle $\{T_i : i \in I\}$ of measure preserving transformations on a common probability space (X, μ) , which are chosen randomly according to a stationary Markov chain. This setting can be described in terms of a step skew-product T with a Markov shift (Ω, ν, S) in the base and fibers T_i acting on X . In many situations, it is desirable to know whether ergodicity of the bundle implies ergodicity of the skew-product. For Markov shifts with finite state space Alexander I. Bufetov introduced in [3] the concept of strongly connected Markov measures, which arise from strictly irreducible matrices, in order to formulate a sufficient condition for the validity of the aforementioned implication. In the main theorem of this note, Theorem 2.3, we generalize this criterion to the case of countably infinite bundles. The proof is based on a characterization of the eigenfunctions of the Perron-Frobenius operator associated to the skew-product, which is due to Zbigniew S. Kowalski, cf. [14]. Moreover, we show that the criterion is in fact a characterization.

The note is organized as follows. In Section 1 we define and discuss the notion of strict irreducibility for Markov shifts with countable state space. Section 2 is devoted to the proof of the main theorem. In Section 3 we discuss several implications of this result. In particular, we explain how it can be applied for establishing pointwise ergodic theorems for Cesàro means of word-spherical averages.

1 Strict irreducibility of Markov shifts

Let I be a finite or countably infinite alphabet, i. e. $I = \{1, \dots, m\}$ for some $m \in \mathbb{N}$ or $I = \mathbb{N}$. A function $\pi : I \times I \rightarrow [0, 1]$ is called a *Markov kernel* if for all $i \in I$ we have $\sum_{j \in I} \pi(i, j) = 1$. It is not difficult to show that the product $\pi\kappa$ of two Markov kernels π and κ , which is defined by

$$(\pi\kappa)(i, j) := \sum_{k \in I} \pi(i, k)\kappa(k, j),$$

is again a Markov kernel. In case I is finite a Markov kernel is simply a row stochastic matrix Π , called the *transition matrix*, and the product of two Markov kernels coincides with the matrix product. We shall call a Markov kernel π *irreducible* if for all $i, j \in I$ there is an $n \in \mathbb{N}$ such that $\pi^n(i, j) > 0$. In the finite case this coincides with the irreducibility of the transition matrix Π . By an *initial distribution* we mean a sequence $\tau : I \rightarrow [0, 1]$ such that $\sum_{i \in I} \tau(i) = 1$. If $\tau(i) > 0$ for all $i \in I$, we call τ *strictly positive*. Given a Markov kernel π we say that τ is *invariant* with respect to π in case $\sum_{j \in I} \tau(j)\pi(j, i) = \tau(i)$ for all $i \in I$. In the finite case an invariant initial distribution corresponds to a stochastic eigenvector of the transition matrix Π .

Let I^* denote the set of finite words and Ω the set $I^{\mathbb{N}}$ of infinite sequences over I . We equip Ω with the product σ -algebra \mathcal{B} generated by cylinder sets of the form

$$[v] := \{ \omega \in \Omega : \omega_0 = v_0, \dots, \omega_{|v|-1} = v_{|v|-1} \}$$

for some word $v \in I^*$. Let π be a Markov kernel and τ be an initial distribution. Then, as a consequence of Kolmogorov's extension theorem, there is a unique probability measure ν on Ω such that

$$\nu(v) = \tau(v_0)\pi(v_0, v_1)\dots\pi(v_{n-2}, v_{n-1})$$

for every word $v = v_0\dots v_{n-1}$ over I , where $\nu(v)$ denotes the value of ν on the cylinder set $[v]$, see [8, Chapter 5]. The measure ν is called the *Markov measure* corresponding to π and τ . Let $S : \Omega \rightarrow \Omega$ denote the *shift map* on Ω , which is given by

$$S(\omega_0\omega_1\dots) = \omega_1\omega_2\dots$$

for $\omega \in \Omega$. Then S is \mathcal{B} -measurable. Furthermore, S is ν -invariant if and only if τ is invariant with respect to π . In this case we obtain a measure preserving dynamical system (MDS) (Ω, ν, S) , which we shall call a *Markov shift*. If τ is strictly positive, the MDS is ergodic if and only if π is irreducible, see [8, Chapter 7].

Given a Markov shift (Ω, ν, S) with Markov kernel π and strictly positive, invariant initial distribution τ , we may consider the *reverse Markov kernel* π^* , which is given by $\pi^*(i, j) := \tau(j)\pi(j, i)\tau(i)^{-1}$ for $i, j \in I$. We shall call the Markov chain *strictly irreducible* if $\pi\pi^*$ is irreducible and **-strictly irreducible* if $\pi^*\pi$ is irreducible. In the finite case these properties were introduced in [3] and [4] and correspond to the irreducibility of the matrices $\Pi\Pi^T$ and $\Pi^T\Pi$, where Π denotes the transition matrix of the Markov shift.

Strict irreducibility and strict *-irreducibility can be characterized in different ways, which we shall describe next. In the following a graph is a tuple (V, E) consisting of a set V of vertices and a set $E \subseteq V \times V$ of edges. By a finite

path we mean a finite sequence v_1, \dots, v_m of vertices such that $(v_i, v_{i+1}) \in E$ for $i = 1, \dots, m-1$. We shall say that a graph is *finitely connected* if any two vertices are contained in a finite path. Given a Markov kernel π we shall call to states $i, j \in I$ *crossing* if $i = j$ or $\pi(i, k) > 0$ and $\pi(j, k) > 0$ for some $k \in I$. The corresponding relation \sim on I is obviously reflexive and symmetric. Let \simeq be the transitive closure of \sim , i. e. we have $i \simeq j$ if and only if there are $k_1, \dots, k_n \in I$ such that

$$i \sim k_1, k_1 \sim k_2, \dots, k_{n-1} \sim k_n, k_n \sim j.$$

Then \simeq defines an equivalence relation. Using these notions we may give the following characterizations of strict irreducibility.

1.1 Proposition. *Let (Ω, ν, S) be a Markov shift with Markov kernel π and strictly positive initial distribution τ . Then the following assertions are equivalent:*

- (i) *The Markov kernel $\pi\pi^*$ is irreducible.*
- (ii) *The graph (I, \sim) is finitely connected.*
- (iii) *The relation \simeq has exactly one equivalence class.*

Proof: (ii) \Leftrightarrow (iii) is obvious. For (i) \Leftrightarrow (ii) set $\kappa = \pi\pi^*$ and observe that

$$\kappa(i, j) = \sum_{k \in I} \pi(i, k)\pi^*(k, j) = \sum_{k \in I} \pi(i, k) \frac{\tau(j)}{\tau(k)} \pi(j, k).$$

So we have $\kappa(i, j) > 0$ if and only if $i \sim j$. For the powers of κ we obtain

$$\kappa^{n+1}(i, j) = \sum_{k_1, \dots, k_n} \kappa(i, k_1)\kappa(k_1, k_2)\dots\kappa(k_{n-1}, k_n)\kappa(k_n, j),$$

so we have $\kappa^{n+1}(i, j) > 0$ if and only if there is a sequence $k_1, \dots, k_n \in I$ with $i \sim k_1, k_1 \sim k_2, \dots, k_n \sim j$. In particular κ is irreducible if and only if $i \simeq j$ for all $i, j \in I$. \square

A further characterization of strict irreducibility was given by Blackwell and Friedmann and independently by Grigorenko, see [2] and [11]. Given a Markov shift (Ω, ν, S) as above we may identify two sequences $\omega, \eta \in \Omega$ if they arise from each other by a finite permutation of symbols, i. e. there is some bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ with $\sigma(n) = n$ for all but finitely many n such that $\eta_{\sigma(n)} = \omega_n$ for all $n \in \mathbb{N}$. This defines an equivalence relation. The σ -algebra generated by the classes of this relation is called the *symmetric σ -algebra* of the Markov shift. In [2] and [11] it is shown that the Markov shift is strictly irreducible if and only if the symmetric σ -algebra is trivial.

For $*$ -strict irreducibility analogous characterizations can be given. To this end we have to consider the relation \sim on I , which relates two states $i, j \in I$ if $i = j$ or we find a state $k \in I$ such that $\pi(k, i) > 0$ and $\pi(k, j) > 0$. Considering the transitive closure \simeq of \sim we obtain the following equivalences. The proof is symmetric to that of Proposition 1.

1.2 Proposition. *Let (Ω, ν, S) be Markov shift with Markov kernel π and strictly positive initial distribution τ . Then the following assertions are equivalent:*

- (i) *The Markov kernel $\pi^* \pi$ is irreducible.*
- (ii) *The graph (I, \sim) is finitely connected.*
- (iii) *The relation \simeq has exactly one equivalence class.*




Figure 1: *Transition graph of an irreducible Markov shift. The equivalence classes of \simeq are $A_1 = \{0\}$, $A_2 = \{2\}$ and $A_3 = \{1, 3, 4\}$. The equivalence classes of \simeq are $C_1 = \{0, 1, 2\}$, $C_2 = \{3\}$ and $C_3 = \{4\}$.*

2 Ergodicity of skew-product extensions

Consider a probability space (X, μ) . Let $\{T_i : i \in I\}$ be a bundle of μ -preserving transformations on X . Then we may define a skew-product T on $\Omega \times X$ by

$$T(\omega, x) := (S\omega, T_{\omega_0} x)$$

for $\omega \in \Omega$ and $x \in X$. A skew-product of this form is sometimes called a *step skew-product*. It is not difficult to see that T is measurable with respect to the product σ -algebra on $\Omega \times X$ and preserves the product measure $\nu \otimes \mu$. The MDS $(\Omega \times X, \nu \otimes \mu, T)$ is an instance of a *bundle random dynamical system* or short *bundle RDS*. It is called *ergodic* in case the skew product T is an ergodic transformation.

Given a skew-product T as above we denote by $\mathcal{K}_T : L^\infty(\Omega \times X) \rightarrow L^\infty(\Omega \times X)$ the *Koopman operator* of T , which is given by $\mathcal{K}_T(\varphi) := \varphi \circ T$ for $\varphi \in L^\infty(\Omega \times X)$. As one can show, there is a unique operator $\mathcal{L}_T : L^1(\Omega \times X) \rightarrow L^1(\Omega \times X)$ such that \mathcal{K}_T is the adjoint of \mathcal{L}_T in the sense that

$$\langle \varphi, \mathcal{K}_T \psi \rangle = \langle \mathcal{L}_T \varphi, \psi \rangle, \quad \varphi \in L^1, \quad \psi \in L^\infty,$$

where $\langle \cdot, \cdot \rangle$ denotes the dual pairing between L^1 and L^∞ , see [14]. \mathcal{L}_T is called the *Perron-Frobenius operator* or short *P-F operator*. We will make extensive use of a characterization of the eigenvalues of the P-F operator of a step skew-product due to Kowalski. The following statement is a special case of [14, Theorem 3.1].

2.1 Theorem. *Let $(\Omega \times X, \nu \otimes \mu, T)$ be a step skew-product over a Markov shift (Ω, ν, S) with at most countable state space I . Let $\varphi \in L^1(\Omega \times X)$ be an eigenfunction of \mathcal{L}_T , i. e. $\mathcal{L}_T \varphi = \lambda \varphi$ for some $\lambda \in \mathbb{C}$ with $|\lambda| = 1$. Then there is a family $\{\varphi_i : i \in I\}$ of functions $\varphi_i \in L^1(X)$ such that*

$$\varphi(\omega, x) = \sum_{i \in I} \mathbb{1}_{[i]}(\omega) \varphi_i(x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in \Omega \times X$.

2.2 Corollary. *Let $(\Omega \times X, \nu \otimes \mu, T)$ be a step skew-product over a Markov shift (Ω, ν, S) with at most countable state space I and strictly positive initial distribution. Let $\varphi \in L^\infty(\Omega \times X)$ be a T -invariant function. Then there is a family $\{\varphi_i : i \in I\}$ of functions $\varphi_i \in L^\infty(X)$ such that*

$$\varphi(\omega, x) = \sum_{i \in I} \mathbb{1}_{[i]}(\omega) \varphi_i(x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in \Omega \times X$.

Proof: Let $\varphi \in L^\infty$ be T -invariant. Then we have

$$\begin{aligned} \langle \mathcal{L}_T \varphi, \psi \rangle &= \langle \varphi, \mathcal{K}_T \psi \rangle = \int \varphi \cdot \psi \circ T \, d\nu \otimes \mu \\ &= \int \varphi \circ T \cdot \psi \circ T \, d\nu \otimes \mu = \int \varphi \cdot \psi \, d\nu \otimes \mu \\ &= \langle \varphi, \psi \rangle \end{aligned}$$

for all $\psi \in L^1$. This shows that $\mathcal{L}_T \varphi = \varphi$, so the claimed representation of φ with $\varphi_i \in L^1$ follows from Theorem 2.1. Using the strict positivity of the initial distribution and the boundedness of φ it is not difficult to see that the φ_i have to

be L^∞ -functions. □

Assuming the setting of the above results let Σ_S and Σ_T denote the σ -algebras of S -invariant and T -invariant sets. Let furthermore Σ denote the σ -algebra of measurable sets $A \subseteq X$ such $T_i^{-1}(A) = A$ for all $i \in I$. In the following we shall characterize under which circumstances the ergodicity of the bundle $\{T_i : i \in I\}$, i. e. the triviality of Σ , implies the ergodicity of T , i. e. the triviality of Σ_T , in terms of the Markov shift (Ω, ν, S) . It is not difficult to see that the other direction is always true.

It is clear that if the Markov shift is not ergodic, T cannot be ergodic. To see this consider a set $B \in \Sigma_S$ with $0 < \nu(B) < 1$. Then $B \times X$ is a non trivial T -invariant set. Therefore we will assume the Markov shift to be ergodic, so Σ_S is the trivial σ -algebra Σ_{triv} on Ω . It is also clear that we have to require the initial distribution to be strictly positive. Otherwise we could construct a non-ergodic skew-product such that Σ is trivial by assigning an ergodic transformation to a symbol that is never reached.

We now state and prove the main theorem of this note. The theorem extends a criterion of Bufetov given in [3, Theorem 5] (see also [4]) for the validity of the implication discussed above to countable state Markov shifts and shows that it is in fact a characterization. This way we obtain a characterization of the $*$ -strict irreducibility of a Markov shift in terms of the ergodicity of suitable skew-product extensions.

2.3 Theorem. *Let (Ω, ν, S) be an ergodic Markov shift with at most countable state space I and strictly positive initial distribution. Then the following statements are equivalent:*

- (i) (Ω, ν, S) is $*$ -strictly irreducible.
- (ii) For every measure preserving step skew-product $(\Omega \times X, \nu \otimes \mu, T)$ over (Ω, ν, S) we have

$$\Sigma_T = \Sigma_S \otimes \Sigma = \Sigma_{\text{triv}} \otimes \Sigma.$$

- (iii) A measure preserving step skew-product $(\Omega \times X, \nu \otimes \mu, T)$ over (Ω, ν, S) is ergodic whenever the bundle $\{T_i : i \in I\}$ is ergodic.

Proof: (i) \Rightarrow (ii) Let $C = B \times A \in \Sigma_S \otimes \Sigma$. Then we have

$$T^{-1}(C) = \bigcup_{\omega \in S^{-1}(B)} \{\omega\} \times T_{\omega_0}^{-1}(A) = \bigcup_{\omega \in B} \{\omega\} \times A = B \times A = C,$$

so C is T -invariant. Since the sets of this form generate $\Sigma_S \otimes \Sigma$, this shows that $\Sigma_S \otimes \Sigma \subseteq \Sigma_T$.

To verify the other inclusion we will show that every bounded Σ_T -measurable, i. e. T -invariant function is also $\Sigma_S \otimes \Sigma$ -measurable. To this end let $\varphi \in L^\infty(\Omega \times X)$ be T -invariant. Then, by Corollary 2.2, we have

$$\varphi(\omega, x) = \sum_{i \in I} \mathbb{1}_{[i]}(\omega) \varphi_i(x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in \Omega \times X$ and a family $\{\varphi_i : i \in I\}$ of functions $\varphi_i \in L^\infty(X)$. Thus we obtain

$$\varphi \circ T(\omega, x) = \varphi(S\omega, T_{\omega_0}x) = \sum_{i \in I} \mathbb{1}_{[i]}(S\omega) \varphi_i(T_{\omega_0}x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in \Omega \times X$. This shows that $\varphi_i = \varphi_j \circ T_i$ μ -a.s. whenever $\pi(i, j) > 0$. Otherwise there would be a measurable set A with $\mu(A) > 0$, such that

$$\varphi(\omega, x) = \varphi_i(x) \neq \varphi_j \circ T_i(x) = \varphi \circ T(\omega, x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in [ij] \times A$. But since $\nu(ij) > 0$ this would contradict the T -invariance of φ .

Now consider $i, j \in I$ with $i \sim j$. Then we have a $k \in I$ such that $\pi(k, i) > 0$ and $\pi(k, j) > 0$. This implies

$$\varphi_i \circ T_k = \varphi_k = \varphi_j \circ T_k \quad \mu\text{-a.s.}$$

and since T_k is measure preserving we have $\varphi_i = \varphi_j$ μ -almost surely (Note that the Koopman operator is an isometry and therefore injective.) By assumption we find for all $i, j \in I$ a sequence $k_1, \dots, k_n \in I$ with $i \sim k_1, \dots, k_n \sim j$. Thus we obtain

$$\varphi_i = \varphi_{k_1} = \dots = \varphi_{k_n} = \varphi_j \quad \mu\text{-a. s.}$$

for all $i, j \in I$, which implies $\varphi = \mathbb{1} \otimes f$ for some $f \in L^\infty(X)$. The latter function has to be invariant under the bundle $\{T_i : i \in I\}$. Otherwise there would be some T_i and some measurable set A with $\mu(A) > 0$ such that $f \circ T_i \neq f$ on A . But this would imply that

$$\varphi \circ T(\omega, x) = f \circ T_i(x) \neq f(x) = \mathbb{1} \otimes f(\omega, x) = \varphi(\omega, x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in [i] \times A$. Since $\nu(i) > 0$ by assumption, this would again contradict the T -invariance of φ . So in particular, φ is a $\Sigma_S \otimes \Sigma$ -measurable function.

(ii) \Rightarrow (iii) Let T be ergodic and $A \in \Sigma$. Then we have $T^{-1}(\Omega \times A) = \Omega \times A$ by the same argument as above, which implies

$$\mu(A) = \nu(\Omega)\mu(A) = \nu \otimes \mu(\Omega \times A) \in \{0, 1\}.$$

On the other hand, assuming that $\Sigma_T = \Sigma_{\text{triv}} \otimes \Sigma$ and Σ is trivial, we obtain that Σ_T is trivial as well, so T is ergodic.

(iii) \Rightarrow (i) Assume that $(\Omega, \nu, \mathcal{S})$ is not $*$ -strictly irreducible. We shall construct a an ergodic bundle over I such that the corresponding skew-product is not ergodic. To this end consider the unit interval $[0, 1)$ equipped with its Borel σ -algebra and let λ denote the Lebesgue measure on $[0, 1)$. The dyadic odometer d on $[0, 1)$ is defined by

$$d(x) = \sum_{n=0}^{\infty} \mathbb{1}_{[s_n, s_{n+1})}(x) (x - s_n + q_{n+1})$$

for $x \in [0, 1)$, where $(q_n)_{n=1}^{\infty}$ denotes the geometric sequence given by $q_n := 2^{-n}$ and $s_n := \sum_{i=1}^n q_i$ denotes the n -th partial sum of the geometric series ($s_0 := 0$). The dyadic odometer is measure preserving and ergodic with respect to λ , see [16, Example 4.5.A]. Set $I_1 = [0, 1/2)$ and $I_2 = [1/2, 1)$. Then we have $d(I_1) = I_2$ and $d(I_2) = I_1$.

Let $\{C_r : 1 \leq r < N\}$ with $N \in \mathbb{N} \cup \{\infty\}$ denote the equivalence classes with respect to the relation \simeq . By Proposition 1.2 we have $N > 2$. For $i \in I$ let r_i denote the index of the equivalence class of i . Note that since the Markov shift is ergodic and the initial distribution is strictly positive, the kernel π is irreducible, so for all $i, j \in I$ there is some $n \in \mathbb{N}$ such that $\pi^{(n)}(i, j) > 0$. In particular, for all $i \in I$ there is some j such that $\pi(i, j) > 0$. But by definition of \simeq all j with $\pi(i, j) > 0$ are in the same equivalence class with respect to \simeq . So starting in i we may reach exactly one equivalence class C in one step. Denote the index of C by k_i .

Now we define a bundle $\{T_i : i \in I\}$ of λ -preserving transformations on $[0, 1)$ by setting $T_i := \text{Id}$ for $|r_i - k_i|$ even and $T_i := d$ for $|r_i - k_i|$ odd. Since $2\mathbb{N}$ is a semigroup, there must be at least one $i \in I$ with $|r_i - k_i|$ odd. Otherwise we would never reach an equivalence class with odd index when starting in an equivalence class with even index, which would contradict the ergodicity of the Markov shift. Thus $T_i = d$ for at least one $i \in I$. This ensures that Σ is trivial.

It remains to show that T is not ergodic. To see this consider the set

$$D := \bigcup_{r=1}^{N-1} \bigcup_{i \in C_r} [i] \times J_r,$$

where J_r denotes the interval $[0, 1/2)$ if r is odd and the interval $[1/2, 1)$ if r is even. Then D is obviously measurable. We claim that D is also T -invariant. To verify this fix $(\omega, x) \in D$, set $i := \omega_0$ and $j := \omega_1$ and assume that $\pi(i, j) > 0$. Then $k_i = r_j$ and therefore, if $|r_i - k_i| = |r_i - r_j|$ is even, we have $J_{r_i} = J_{r_j}$. In that case we have $T_{\omega_0}(x) = T_i(x) = \text{Id}(x) = x \in J_{r_j}$ and therefore $T(\omega, x) \in [j] \times J_{r_j} \subseteq D$. If on the other hand $|r_i - k_i| = |r_i - r_j|$ is odd, then $J_{r_i} = I_1$ and $J_{r_j} = I_2$ or vice versa. In both cases we obtain $T_{\omega_0}(x) = d(x) \in J_{r_j}$ and therefore again $T(\omega, x) \in [j] \times J_{r_j} \subseteq D$. Since the condition $\pi(\omega_0, \omega_1) > 0$ is satisfied for ν -almost all $\omega \in \Omega$ we may conclude that $(\omega, x) \in D$ implies $T(\omega, x) \in D$ for $\nu \otimes \lambda$ -almost all $(\omega, x) \in \Omega \times X$. Now the complement of D has the form

$$D^c = \bigcup_{r=1}^{N-1} \bigcup_{i \in C_r} [i] \times J_r^c,$$

so the same argument gives that $(\omega, x) \in D^c$ implies $T(\omega, x) \in D^c$ for $\nu \otimes \lambda$ -almost all (ω, x) . In sum we have shown that D is T -invariant.




Figure 2: The set D for the Markov shift in Fig. 1.

However, since $[i] \times J_i \subseteq D$ for arbitrary $i \in I$, we have $\nu \otimes \lambda(D) \geq \nu(i)\lambda(J_i) = \nu(i)/2 > 0$ by the strict positivity of the initial distribution. Since D^c has a subset of the same measure, we have $\nu \otimes \lambda(D^c) > 0$ and therefore $0 < \nu \otimes \lambda(D) < 1$. So T admits a non trivial T -invariant set, which implies that T is not ergodic. \square

As noted above, in the situation of the Theorem, the ergodicity of the skew-product implies always the ergodicity of the bundle. So the implication given in (iii) is in fact an equivalence. For the case of Bernoulli shifts (with arbitrary state space) the validity of this equivalence was already shown by Kakutani in [13]. In

the context of random walks on locally compact groups, Aaronson and Lemańczyk discuss this equivalence in [1], showing that it is tightly related to ergodicity and mixing properties of the group on its Poisson boundary (see also [12]).

In case of a finite state space I a further important characterization can be added to the statements of the above theorem. Let (Ω, ν, S) be an ergodic Markov shift with finite state space $I = \{1, \dots, m\}$, transition matrix Π and strictly positive initial distribution π . Let $\{T_i : i \in I\}$ be a bundle of measure preserving transformations on a common probability space (X, μ) as in the theorem. Denoting by $L_m^1(X) := L^1(X) \times \dots \times L^1(X)$ the m -fold cartesian product of the space $L^1(X)$ we may define bounded linear operators $A_j : L_m^1(X) \rightarrow L^1(X)$ for $j = 1, \dots, m$ and a bounded linear operator $A : L_m^1(X) \rightarrow L_m^1(X)$ by

$$A_j(\vec{\varphi}) := \sum_{i=1}^m \pi_{ji} \varphi_i \circ T_j$$

and

$$A(\vec{\varphi}) := (A_1(\vec{\varphi}), \dots, A_m(\vec{\varphi}))$$

for some vector $\vec{\varphi} = (\varphi_1, \dots, \varphi_m)$ of functions $\varphi_i \in L^1(X)$. Identifying $L_m^1(X)$ with $L^1(X \times I)$ one may consider A as an operator acting on L^1 -spaces. One can show that in this sense A is an instance of a Markov operator, see [5]. As is shown in [3, Theorem 4], the skew product T is ergodic if and only if A is an ergodic operator in the sense that the corresponding fixspace contains only constants. This observation plays an important role for obtaining ergodic theorems for so called strictly irreducible Markov groups, which we shall discuss below. However the identification of $L_m^1(X)$ with $L^1(X \times I)$ only works for finite I . Thus, in case of an infinite alphabet I , an analogous construction is not at hand.

3 Random and weighted ergodic theorems for bundles

The ergodicity of the skew-product extension of a Markov shift plays an important role for obtaining ergodic theorems for random dynamical systems. For a given bundle RDS $(\Omega \times X, \nu \otimes \mu, T)$ as above let $T_{\omega, n}$ denote the transformation $T_{\omega_{n-1}} \circ \dots \circ T_{\omega_0}$ for $n \geq 1$, and $T_{\omega, 0} := \text{Id}$. Considering a function $f \in L^1(X)$ we obtain μ -almost sure convergence of the fiberwise random averages

$$\frac{1}{n} \sum_{i=0}^{n-1} f \circ T_{\omega, i} = \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1} \otimes f \circ T^i$$

for ν -almost all $\omega \in \Omega$ by Birkhoff's ergodic theorem. However, it is not clear that the limit equals $\mathbb{E}[f | \Sigma]$. To see that this is not always the case consider the RDS

$(\{0, 1\}^{\mathbb{N}} \times \mathbb{T}, \nu \times \lambda, T)$, where \mathbb{T} denotes the torus, λ the Lebesgue-measure and ν the Markov measure given by the transition matrix

$$\Pi = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

and invariant initial distribution $\pi = (1/2, 1/2)^T$. The bundle of transformations is given by $T_0 = R_\alpha$ and $T_1 = R_\beta$, where R_α and R_β are rotations with irrational angles α, β such that $\alpha + \beta = 1$. Since rotations with irrational angles are ergodic, we obtain that Σ is trivial. However, since the measure ν is concentrated on the singletons $\{1010\dots\}$ and $\{0101\dots\}$ with equal probabilities, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{2n} \sum_{i=0}^{2n-1} \mathbb{1}_A \circ T_{\omega, i}(x) &\geq \lim_{n \rightarrow \infty} \frac{1}{2n} \sum_{j=0}^{n-1} \mathbb{1}_A \circ R_{\alpha+\beta}^j(x) \\ &= \frac{1}{2} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1}_A(x) = \frac{1}{2} > \nu(A) = \int_X \mathbb{1}_A d\lambda \end{aligned}$$

on the fiber $\omega = 01010101\dots$, every set A with $0 < \lambda(A) < \frac{1}{2}$ and every $x \in A$ and therefore on a set in $\Omega \times X$ of positive measure.

However, Theorem 2.3 allows us to give a sufficient criterion for the limit to be $\mathbb{E}[f|\Sigma]$ for every $f \in L^1(X)$ only in terms of the underlying Markov shift.

3.1 Theorem. *Let $(\Omega \times X, \nu \otimes \mu, T)$ be a finite bundle RDS over a $*$ -strictly irreducible ergodic Markov shift (Ω, ν, S) with at most countable state space I and strictly positive initial distribution. Then for all $f \in L^1(X)$ we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} f \circ T_{\omega, i}(x) = \mathbb{E}[f|\Sigma](x)$$

for $\nu \otimes \mu$ -almost all $(\omega, x) \in \Omega \times X$.

Proof: Since T is measure preserving we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} f \circ T_{\omega, i}(x) = \mathbb{E}[\mathbb{1} \otimes f | \Sigma_T](\omega, x)$$

$\nu \otimes \mu$ -almost surely by Birkhoff's ergodic theorem. The $*$ -strict irreducibility of (Ω, ν, S) implies $\Sigma_T = \Sigma_{\text{triv}} \otimes \Sigma$ by Theorem 2.3. Therefore we obtain

$$\mathbb{E}[\mathbb{1} \otimes f | \Sigma_T] = \mathbb{E}[\mathbb{1} | \Sigma_{\text{triv}}] \otimes \mathbb{E}[f | \Sigma] = \mathbb{E}[\mathbb{1}] \otimes \mathbb{E}[f | \Sigma] = \mathbb{1} \otimes \mathbb{E}[f | \Sigma],$$

which finishes the proof. \square

We may also ask for deterministic ergodic theorems along simultaneous averages over compositions of transformations in the bundle. More precisely we may consider the weighted spherical averages

$$S_n^v(f) := \sum_{|v|=n} v(v) f \circ T_v,$$

where $T_v := T_{v_0} \circ \dots \circ T_{v_{n-1}}$ denotes the composition of the T_i according to the word v . As we shall see below the above sum is well defined μ -a. s. for all $f \in L^1(X)$ since we may write it as the integral of the function $(\mathbb{1} \otimes f) \circ T^n(\cdot, x)$ against a suitable measure ν^* , the former being an L^1 -function on (Ω, ν^*) for μ -almost all $x \in X$. The following theorem shows that under reasonable conditions we obtain almost sure convergence of the Cesàro averages of $S_n^v(f)$ to the conditional expectation $\mathbb{E}[f|\Sigma]$. It extends results of Bufetov in [3] and [4] for the case of finite bundles, see also [6]. Recall the definition of the space $L \log L(X)$, which consists of all measurable functions f such that $\int |f| \log^+ |f| d\mu < \infty$.

3.2 Theorem. *Let (Ω, ν, S) be an ergodic Markov shift with at most countable state space I and strictly positive initial distribution. Let $\{T_i : i \in I\}$ be a bundle of measure preserving transformations on a probability space (X, μ) . Then for all $f \in L \log L(X)$ there is a function $\bar{f} \in L^1(X)$ such that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} S_i^v(f)(x) = \bar{f}(x)$$

for μ -almost all $x \in X$. If the Markov shift is strictly irreducible, then $\bar{f} = \mathbb{E}[f|\Sigma]$ μ -almost surely.

Proof: Let π be the Markov kernel and τ be the initial distribution of (Ω, ν, S) . We consider the Markov measure ν^* with reverse Markov kernel π^* and same initial distribution τ . It is not difficult to see that ν^* is also S -invariant, so (Ω, ν^*, S) is again a Markov shift. Accordingly, the skew-product T preserves the product measure $\nu^* \otimes \mu$.

Fix $f \in L \log L(X)$ and set $g := \mathbb{1} \otimes f$. For a pair $(\omega, x) \in \Omega \times X$ and a number $n \in \mathbb{N}$ consider the ergodic averages

$$M_n f(\omega, x) := \frac{1}{n} \sum_{i=0}^{n-1} g \circ T^i(\omega, x) = \frac{1}{n} \sum_{i=0}^{n-1} f \circ T_{\omega, i}(x).$$

Then by Birkhoff's ergodic theorem we have $\nu^* \otimes \mu$ -almost sure convergence of

$M_n f(\omega, x)$ to some limit \bar{g} in $L^1(\Omega \times X, \nu^* \otimes \mu)$. Furthermore, if the Markov shift (Ω, ν, S) is strictly irreducible, then the Markov shift (Ω, ν^*, S) is $*$ -strictly irreducible. In this case we may identify the limit as $\mathbb{1} \otimes \mathbb{E}[f|\Sigma]$ for $\nu^* \otimes \mu$ -almost all (ω, x) by Theorem 3.1.

Note that $f \in L \log L$ implies that f is integrable with respect to μ , which in turn gives that g and therefore $M_n f$ is integrable with respect to $\nu^* \otimes \mu$. The latter implies that $M_n f(\cdot, x)$ is in $L^1(\Omega, \nu^*)$ for μ -almost all $x \in X$. We may therefore integrate over Ω to obtain

$$\begin{aligned} \int_{\Omega} M_n f(\omega, x) d\nu^*(\omega) &= \frac{1}{n} \sum_{i=0}^{n-1} \int_{\Omega} f \circ T_{\omega, i}(x) d\nu^*(\omega) \\ &= \frac{1}{n} \sum_{i=0}^{n-1} \sum_{|v|=i} \nu^*(v) f \circ T_{v_{i-1}} \circ \dots \circ T_{v_0}(x) \end{aligned}$$

for μ -almost all $x \in X$, where the latter sum is absolutely convergent. Considering the bijections $\sigma: I^i \rightarrow I^i$ mapping a word $v_0 \dots v_{i-1}$ to the word $v_{i-1} \dots v_0$ and observing that $\nu^*(v_0 \dots v_{i-1}) = \nu(v_{i-1} \dots v_0)$ for all v of length i we obtain for the inner sums, which are absolutely and in particular unconditionally convergent, that

$$\begin{aligned} \sum_{|v|=i} \nu^*(v) f \circ T_{v_{i-1}} \circ \dots \circ T_{v_0}(x) &= \sum_{|v|=i} \nu(\sigma(v)) f \circ T_{\sigma(v)} \\ &= \sum_{|v|=i} \nu(v) f \circ T_v(x) \\ &= S_i^v(f)(x), \end{aligned}$$

so we have

$$\int_{\Omega} M_n f(\omega, x) d\nu^*(\omega) = \frac{1}{n} \sum_{i=0}^{n-1} S_i^v(f)(x)$$

for μ -almost all $x \in X$. Now set

$$M^* f(\omega, x) := \sup_{n \in \mathbb{N}} |M_n f(\omega, x)|$$

for $(\omega, x) \in \Omega \times X$. Since $f \in L \log L$, we have $g \in L \log L(\Omega \times X, \nu^* \otimes \mu)$. Therefore, by Wiener's dominated ergodic theorem, see [15, Chapter 1, Theorem 6.3], we have $M^* f \in L^1(\Omega \times X, \nu^* \otimes \mu)$. This implies that $M^* f(\cdot, x)$ is in $L^1(\Omega, \nu^*)$ for μ -almost all $x \in X$. In particular, for μ -almost all $x \in X$, the function $M^* f(\cdot, x)$ is an integrable majorant for the functions $M_n f(\cdot, x)$. This allows us

to use Lebesgue's theorem to obtain convergence of $\frac{1}{n} \sum_{i=0}^{n-1} S_i^v(f)(x)$ to

$$\bar{f}(x) := \int_{\Omega} \bar{g}(\omega, x) d\nu^*(\omega)$$

for μ -almost all $x \in X$. Since $g \in L^1(\Omega \times X, \nu^* \otimes \mu)$ we get $\bar{f} \in L^1(X, \mu)$. In case of a strictly irreducible Markov chain we have in addition

$$\bar{f}(x) = \int_{\Omega} \mathbb{1} \otimes \mathbb{E}[f|\Sigma](\omega, x) d\nu^*(\omega) = \mathbb{E}[f|\Sigma](x)$$

for μ -almost all $x \in X$. This finishes the proof. \square

In the case of a finite state space I one can rely on ergodic theoretic properties of the Markov operator A from above to obtain even convergence for functions in $L^1(X)$, see [3], [5] and [6]. Moreover, considering topological Markov chains and using for ν the Parry measure, one can replace the above averages by combinatorial averages in the sense that all points on a sphere get equal weight, see [5]. Using this one can e. g. obtain ergodic theorems for actions of finitely generated groups, which admit strictly irreducible Markov codings, such as the free (semi)group, see also [9] and [10], locally free (semi)groups, see [5], and a large class of Fuchsian groups, see [7]. However, since the approach relies on the ergodicity of the Markov operator defined above with respect to the reverse Markov chain, the characterization given in Theorem 2.3 determines the scope of this method in the sense that it is essentially restricted to such groups.

Acknowledgements. We would like to thank A. I. Bufetov for valuable remarks on a preliminary version of the manuscript. Both authors were partially supported by the grant I-1485-304.6/2019 by the German Israeli Foundation of Scientific Research and Development (GIF).

References

- [1] J. Aaronson and M. Lemańczyk: *Exactness of Rokhlin endomorphisms and weak mixing of Poisson boundaries*, Contemp. Math. 385, Amer. Math. Soc. Providence, RI, 77-87, 2005
- [2] D. Blackwell and D. Freedman: *The tail σ -field of a Markov chain and a theorem of Orey*, Ann. Math. Statist. 35, 1291-1295, 1964
- [3] A. I. Bufetov: *Skew products and ergodic theorems for group actions*, Zap. Nauchn. Semin. POMI, 266(5), 13–28, 2000. Translated in: J. Math. Sci. 113(4), 548–557, 2003

- [4] A. I. Bufetov: *Operator ergodic theorems for actions of free semigroups and groups*, Funktsional. Anal. i Prilozhen., 34(4), 1-17, 2000
- [5] A. I. Bufetov: *Markov averaging and ergodic theorems for several operators*, in: *Topology, ergodic theory, real algebraic geometry*, vol. 202 of Amer. Math. Soc. Transl. Ser. 2, Providence, 39-50, 2001
- [6] A. I. Bufetov and A. Klimenko: *On Markov operators and ergodic theorems for group actions*, Eur. J. Comb., 33, 1427 - 1443, 2012
- [7] A. I. Bufetov and C. Series: *A pointwise ergodic theorem for Fuchsian groups*, Math. Proc. Camb. Philos. Soc., 151(1), 145-159, 2011
- [8] R. Durrett: *Probability: Theory and examples*, 5th edition, Cambridge University Press, 2019
- [9] R. I. Grigorchuk: *Individual ergodic theorems for actions of free groups* (Russian), Proceedings of the Tambov Workshop in the Theory of Functions, 1986
- [10] R. I. Grigorchuk : *Ergodic theorems for actions of free groups and free semigroups*, Math. Notes, 65(5), 654–657, 1999
- [11] L. A. Grigorenko: *On the σ -algebra of symmetric events for a countable Markov chain*, Theory Probability Appl. 24, 199-204, 1979
- [12] V. Kaimanovich: *Double ergodicity of the Poisson boundary and applications to bounded cohomology*, Geom. Funct. Ana. 13, 852-861, 2003
- [13] S. Kakutani: *Random ergodic theorems and Markoff processes with a stable distribution*, in: *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, 1950, University of California Press, Berkeley and Los Angeles, 247-261, 1951
- [14] Z. S. Kowalski: *Frobenius-Perron operator description of Markov chains*, Prob. Math. Stat. 35, 325-341, 2015
- [15] U. Krengel: *Ergodic theorems*, de Gruyter, 1985
- [16] K. E. Petersen: *Ergodic theory*, Cambridge University Press, 1989

MATHEMATICAL INSTITUTE, LEIPZIG UNIVERSITY
AUGUSTUSPLATZ 10, 04109 LEIPZIG
felix.pogorzelski@math.uni-leipzig.de
elias.zimmermann@math.uni-leipzig.de