

Solving sparse polynomial systems using Gröbner bases and resultants

Matías R. Bender
 Technische Universität Berlin
 Berlin, Germany
 mbender@math.tu-berlin.de

Dedicated to the memory of Agnes Szanto

ABSTRACT

Solving systems of polynomial equations is a central problem in nonlinear and computational algebra. Since Buchberger’s algorithm for computing Gröbner bases in the 60s, there has been a lot of progress in this domain. Moreover, these equations have been employed to model and solve problems from diverse disciplines such as biology, cryptography, and robotics. Currently, we have a good understanding of how to solve generic systems from a theoretical and algorithmic point of view. However, polynomial equations encountered in practice are usually structured, and so many properties and results about generic systems do not apply to them. For this reason, a common trend in the last decades has been to develop mathematical and algorithmic frameworks to exploit specific structures of systems of polynomials.

Arguably, the most common structure is sparsity; that is, the polynomials of the systems only involve a few monomials. Since Bernstein, Khovanskii, and Kushnirenko’s work on the expected number of solutions of sparse systems, toric geometry has been the default mathematical framework to employ sparsity. In particular, it is the crux of the matter behind the extension of classical tools to systems, such as resultant computations, homotopy continuation methods, and most recently, Gröbner bases. In this work, we will review these classical tools, their extensions, and recent progress in exploiting sparsity for solving polynomial systems.

This manuscript complements its homonymous tutorial presented at the conference ISSAC 2022.

CCS CONCEPTS

• **Computing methodologies** → **Symbolic and algebraic algorithms**; *Hybrid symbolic-numeric methods*; *Symbolic calculus algorithms*; *Equation and inequality solving algorithms*.

KEYWORDS

Sparse polynomials, Gröbner bases, resultants, solving polynomial systems

1 INTRODUCTION

Systems of polynomial equations give us one of the simplest and general ways of dealing with non-linear objects. They are expressive enough to encode algebraic varieties and effective enough to allow us to compute with them. They generalize two ubiquitous kinds of equations appearing in Mathematics, linear equations and univariate polynomials. Solving polynomial equations, that is, to

find exact or approximate solutions for the equations of the systems, is one of the central problems in this setting and their applications span several domains in science and engineering [17, 33].

There are several tools to solve polynomial systems, e.g., geometric resolutions [64, 67, 68], Gröbner bases [56, 60][75, Chp. 2], homotopy continuation [18, 70, 86], normal form algorithms [2, 79, 88, 97], resultants [24, 49][34, Chp. 3], subdivisions [36, 89, 99], subresultants [43, 65, 95] and triangular decompositions [28, 101]. The aforementioned is a non-exhaustive list of methods and references; the interested reader can find the latest developments on many of these methods in the proceedings of the previous editions of ISSAC. Some nice introductions to the theory of solving systems of polynomial equations can be found in [34, 42, 75]. In this manuscript, and in its associated tutorial, we focus on Gröbner bases, resultants, and normal forms algorithms.

In practice, the polynomial systems that we encounter are structured, e.g., sparse polynomials in biology [62] and statistics [44], symmetric in cryptography [52], and determinantal in optimization [71]. However, the general-purpose strategies to solve polynomial systems does not exploit this structure. For this reason, in the last years, an important trend in polynomial system solving is to improve these techniques for specific structures. In this text, we will focus on the sparsity of the inputs. Besides sparsity, other kinds of structures include, e.g., symmetry [4, 54], determinantal [52, 77], black box evaluations [19], and degenerations [21].

1.1 Sparsity

Sparsity is arguably the most common structure appearing in practice. Before giving the definition of sparsity, let us introduce some notation.

Notation. Given $\alpha \in \mathbb{Z}^n$, we define $\mathbf{x}^\alpha := \prod_{i=1}^n x_i^{\alpha_i}$. The monomials $\{\mathbf{x}^\alpha : \alpha \in \mathbb{Z}^n\}$ generate the \mathbb{C} -algebra of *Laurent polynomials* $\mathbb{C}[x_1^\pm, \dots, x_n^\pm]$. We will write our polynomials in $\mathbb{C}[x_1^\pm, \dots, x_n^\pm]$ as a sum of terms, i.e., $\sum_{\alpha \in \mathbb{N}^n} c_\alpha \mathbf{x}^\alpha$, where $c_\alpha \in \mathbb{C}$ is the coefficient of the monomial \mathbf{x}^α and there are finite $\alpha \in \mathbb{Z}^n$ such that $c_\alpha \neq 0$. The *support* of a given polynomial f is the finite set of exponents with non-zero coefficients, i.e.,

$$\text{Supp} \left(\sum_{\alpha \in \mathbb{N}^n} c_\alpha \mathbf{x}^\alpha \right) := \{\alpha \in \mathbb{Z}^n : c_\alpha \neq 0\}.$$

We abuse notation and we also identify $\text{Supp}(f)$ with a set of the monomials with exponents in $\text{Supp}(f)$. In this text, we will only work with rational polytopes, i.e., its vertices are integer points. Given two polytopes P_1 and P_2 , we define its Minkowski sum $P_1 + P_2$ as their point-wise addition, i.e., $P_1 + P_2 := \{\alpha + \beta : \alpha \in P_1, \beta \in$

P_2 }. For each $i \in \mathbb{N}$, let $iP := P + (i-1)P$, and $0P = \{0\}$. We denote by Δ_n the n -dimensional standard simplex, i.e., the convex hull of $\{0, e_1, \dots, e_n\} \subset \mathbb{R}^n$, where $\{e_1, \dots, e_n\}$ is the canonical basis of \mathbb{R}^n .

Intuitively, a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is sparse when $\text{Supp}(f)$ is *small*, e.g., $\text{Supp}(f)$ has less elements than $\text{Supp}(g)$, where $g \in \mathbb{C}[x_1, \dots, x_n]$ is a generic polynomial of the same degree as f . However, in this text we will use a related, but different, notion of sparsity given by the *Newton polytope* of the polynomial.

Definition 1.1 (Newton polytope). Given a polynomial f , its *Newton polytope* $\text{NP}(f) \subset \mathbb{R}^n$ is the convex hull of $\text{Supp}(f)$ over \mathbb{R}^n .

Observe that the Newton polytope of a generic polynomial in $\mathbb{C}[x_1, \dots, x_n]$ of degree d is $d\Delta_n$, i.e., the d -dilation of the n -dimensional standard simplex.


Example 1.2 (Generalized eigenvalue problem). For most generalized eigenvalue problems in $\mathbb{C}^{2 \times 2}$, we can represent them as a system of polynomial equations in two variables λ, w , where $(1, \lambda)$ denotes the generalized eigenvalue and $v := (1, w)$, the eigenvector. For example, for the pencil (A, B) , $A := \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$, $B := \begin{pmatrix} 3 & 4 \\ -2 & -4 \end{pmatrix}$, we obtain the polynomial system $f_1 = f_2 = 0$,

$$(A + \lambda B)v = 0 \iff \begin{cases} f_1 := 1 + 3\lambda + 2w + 4\lambda w = 0 \\ f_2 := 3 - 2\lambda + 4w - 4\lambda w = 0 \end{cases}$$

The polynomials have degree 2 and their support are $\text{Supp}(f_1) = \text{Supp}(f_2) = \{1, \lambda, w, \lambda w\}$. The polynomials are sparse; written in terms of monomials of degree at most 2, we have that

$$f_1 = 1 + 3\lambda + 0\lambda^2 + 2w + 0w^2 + 4\lambda w.$$

The gray square in the following picture is their Newton polytope.



When we talk about sparsity, we look at Newton polytopes and not at the supports because of the next theorem, attributed to Bernstein [12], Khovanskii [73], and Kushnirenko [76].

THEOREM 1.3 (BBK BOUND). Fix a polynomial system (f_1, \dots, f_n) with a finite number of solutions over $(\mathbb{C}^*)^n$, where $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. Then, this number of solutions is upper bounded by the mixed volume of $\text{NP}(f_1), \dots, \text{NP}(f_n)$, i.e., $\text{MV}(\text{NP}(f_1), \dots, \text{NP}(f_n))$, where

$$\text{MV}(P_1, \dots, P_n) := (-1)^n + \sum_{k=1}^n (-1)^{n-k} \sum_{\substack{I \subset \{1, \dots, n\} \\ \#I=k}} \#((P_{I_1} + \dots + P_{I_k}) \cap \mathbb{Z}^n)$$

Moreover, consider finite subsets $A_1, \dots, A_n \subset \mathbb{Z}^n$ and a system of generic polynomials (f_1, \dots, f_n) supported on A_1, \dots, A_n , i.e., each $f_i = \sum_{\alpha \in A_i} c_{i,\alpha} x^\alpha$, where $(c_{i,\alpha})_{i \in \{1, \dots, n\}, \alpha \in A_i}$ is a generic vector. Then, the aforementioned bound for this generic system is an equality.

For a proof and extensions of this theorem, see [85]. The generic conditions of the previous theorem can be relaxed [15, 26, 29].

Example 1.4 (Cont. Ex. 1.2). Without considering the sparsity of the input, Bézout's theorem tells us that the system should have (at most) 2^2 solutions. However, it has only two, each corresponding to a different eigenspace. This number agrees with the mixed volume of the Newton polytopes defining the equations.

When working with sparse polynomials, we usually consider two class of systems. We say that a sparse polynomial system (f_1, \dots, f_r) is *unmixed* when there is an integer polytope $P \subset \mathbb{R}^n$ and integers d_1, \dots, d_r such that $\text{NP}(f_i) = d_i P$, i.e., each Newton polytope is a dilation of a common integer polytope. Otherwise, the system is *mixed*.

In this manuscript, we will discuss Gröbner bases and resultants in the context of sparse polynomials. We will employ the Newton polytopes of the input polynomials to speed up the computations and derive complexity bounds for our algorithms depending on the sparsity pattern of the inputs. These techniques rely on *toric geometry*. Toric geometry was also employed to solve sparse polynomial systems using other techniques such as homotopy continuation [50, 70, 82] or geometric resolutions [68].

Remark 1.5 (Other notions of sparsity). As the BKK bound shows, from a complex point of view, the solutions of the sparse systems are “determined” by the Newton polytopes of the input polynomials and not by their supports. However, from a real point of view, the support matters. In real algebraic geometry, a polynomial whose support is small is called a *fewnomial*. Khovanskii showed that the number of positive real solutions of a fewnomial system is upper bounded by a quantity singly exponential in the size of the supports of the fewnomials [74]. Determining the number of real solutions of a fewnomial system is an active area of research. For example, the bounds were improved for general [16] and particular systems [14]. Moreover, probabilistically, stronger bounds hold [20, 72].

A different kind of sparsity that has been studied is the *chordal structure*. This structure only cares about the variables appearing in each polynomial, regardless of the specific monomials. There are dedicated algorithms to compute Gröbner bases [30] and triangular sets [87] that exploit this notion.

1.2 Multiplication maps

An intermediate tool used to solve polynomial systems using the Gröbner bases and resultants is multiplication maps. To define them, let $f := \{f_1, \dots, f_r\} \subset \mathbb{C}[x_1, \dots, x_n]$ be an affine polynomial system with a finite number of solutions δ over \mathbb{C}^n , counting multiplicities. The quotient ring $\mathbb{C}[x_1, \dots, x_n]/\langle f \rangle$ is a finite dimensional \mathbb{C} -vector space whose dimension is δ [34, Chp. 4.2], so we will fix a basis b_1, \dots, b_δ to identify $\mathbb{C}[x_1, \dots, x_n]/\langle f \rangle$ with \mathbb{C}^δ . Given $g \in \mathbb{C}[x_1, \dots, x_n]/\langle f \rangle$, the *multiplication map* of g is the morphism

$$M_g : \mathbb{C}[x_1, \dots, x_n]/\langle f \rangle \rightarrow \mathbb{C}[x_1, \dots, x_n]/\langle f \rangle \\ h \mapsto M_g(h) := gh.$$

Using the basis $\{b_1, \dots, b_\delta\}$, we think about M_g as a matrix, which we call the *multiplication matrix*. Multiplication matrices are ubiquitous in algorithms for solving polynomial systems:

- On one hand, we can approximate the solutions by computing the eigenvalues of these maps. More precisely, the *eigenvalue theorem* [79] states that the eigenvalues of M_g correspond to the evaluations¹ of g at the δ solutions of the

¹The evaluation of $g \in \mathbb{C}[x_1, \dots, x_n]/\langle f \rangle$ at a point $p \in \mathbb{C}^n$ corresponds to $\bar{g}(p)$, where $\bar{g} \in \mathbb{C}[x_1, \dots, x_n]$ is such that \bar{g} belongs to the class of g in $\mathbb{C}[x_1, \dots, x_n]/\langle f \rangle$.

system (f_1, \dots, f_r) . See [32] for more details on this theorem and its history. Moreover, for each solution p of f , the vector given by the evaluations $(b_i(p))_{i \leq \delta}$ belongs to the eigenspace of M_g of eigenvalue $g(p)$ [2].

- On the other hand, multiplication maps are a standard tool to compute Gröbner bases for zero-dimension ideals. The algorithm FGLM [56, 57] allows us to derive, from the multiplication matrices, Gröbner bases of $\langle f \rangle$ with respect to any monomial order, e.g., lexicographical, or to find a rational univariate representation of the solutions [92].

To compute multiplication matrices, we use a *normal form*, which is, roughly speaking, a homomorphism $\mathbb{C}[x_1, \dots, x_n] \rightarrow \mathbb{C}^\delta$ whose kernel is $\langle f \rangle$. We can obtain such normal forms using, for example, Gröbner bases [75, Chp. 2], border bases [88][42, Chp. 4], or Sylvester formulas for the resultant [2, 97, 98]. We can compute these matrices by performing $d^{O(n)}$ arithmetic operations, where d is the maximal among the degrees of the polynomials in f [78].

The aforementioned strategies do not employ the sparsity of the inputs to speed up the computations (at least directly²). Moreover, their complexity bounds are independent of this structure and to achieve them, we must destroy the sparsity of the inputs, i.e., to perform random linear change of coordinates or to consider random combinations of the input polynomials. In the following sections, we discuss alternative approaches to avoid these issues.

2 THE SPARSE RESULTANT

In this section, we will assume that the reader is familiar with the classical (projective) resultant. We refer to [34, Chp. 3] and [22] for an introduction.

The sparse resultant is a generalization of the classical resultant that allow us to decide if a sparse affine polynomial system in n variables given by $n + 1$ polynomials has a common solution in $(\mathbb{C}^*)^n$. It is one of the first available tools to solve sparse polynomial systems and a subject of study since the 90's.

Given finite sets $A_0, \dots, A_n \subset \mathbb{Z}^n$, we can parameterize each different polynomial system supported in $A := (A_0, \dots, A_n)$ with points³ in $\mathbb{P}^A := \mathbb{P}^{\#A_0-1} \times \dots \times \mathbb{P}^{\#A_n-1}$,

$$(\mathbf{c}_0, \dots, \mathbf{c}_n) \in \mathbb{P}^{A_0} \times \dots \times \mathbb{P}^{A_n} \mapsto \left(\sum_{\alpha \in A_0} c_{0,\alpha} x^\alpha, \dots, \sum_{\alpha \in A_n} c_{n,\alpha} x^\alpha \right).$$

We use the *incidence variety* Ω to characterize all the sparse systems of polynomials supported on A_0, \dots, A_n with solutions in $(\mathbb{C}^*)^n$.

$$\Omega := \{(p, (F_0, \dots, F_n)) \in (\mathbb{C}^*)^n \times \mathbb{P}^A : (\forall i) F_i(p) = 0\},$$

If we consider the projection map $\pi : (\mathbb{C}^*)^n \times \mathbb{P}^A \rightarrow \mathbb{P}^A$, $\pi(\Omega) \subset \mathbb{P}^A$ determines the systems with solutions at $(\mathbb{C}^*)^n$. However, as $\pi(\Omega)$ is not an algebraic variety, we consider its algebraic closure $\overline{\pi(\Omega)}$ in \mathbb{P}^A . Under combinatorial assumptions on A [93, Cor. 1.1], $\overline{\pi(\Omega)}$ has co-dimension 1 on \mathbb{P}^A and it is an irreducible hypersurface defined by a multihomogeneous polynomial Res_A called

²Heuristically, implementations of the F4 algorithm to compute Gröbner bases [60] can benefit partially from the sparsity of the inputs; see, e.g., [13, 84].

³We identify the systems using the multi-projective space because multiplying the polynomials by non-zero constants does not change their solution set.

the *sparse resultant* of A^4 . For each i , the sparse resultant Res_A is homogeneous with respect to the coefficients of the polynomial F_i supported in A_i and its degree is $\text{MV}(P_0, \dots, P_{i-1}, P_{i+1}, \dots, P_n)$, where P_i is the convex hull of A_i . The sparse resultant is a generalization of the classical (projective) resultant. More precisely, if we have that $A_i = d_i \Delta_n \cap \mathbb{Z}^n$, for some $d_i \in \mathbb{N}$, for each i , the sparse and classical resultant agree. The interested reader can find more properties of the sparse resultant in, e.g., [34, Chp. 7], [63, Chp. 8], [40, 93].

2.1 Vanishing of the sparse resultant

As the classical resultant is a special case of the sparse resultant, it is easy to verify that having solutions on the torus $(\mathbb{C}^*)^n$ is a necessary but not sufficient condition for the vanishing of the sparse resultant. To understand when this vanishes, let us recall when the classical (projective) resultant vanishes. Given an affine polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$, we define its classical *homogenization*, $f^h \in \mathbb{C}[x_0, x_1, \dots, x_n]$, as the polynomial $f^h = x_0^{\deg(f)} f\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right)$. Given an affine polynomial system $f_0, \dots, f_n \in \mathbb{C}[x_1, \dots, x_n]$, the classical resultant of (f_0, \dots, f_n) will vanish if and only if the *homogenization* of the system, i.e. $f_0^h, \dots, f_n^h \in \mathbb{C}[x_0, x_1, \dots, x_n]$, has a common solution on \mathbb{P}^n . The homogenization does not change the solutions of the original system on \mathbb{C}^n , i.e., for $p \in \mathbb{C}^n$, if $f(p_1, \dots, p_n) = 0$, then $f^h(1, p_1, \dots, p_n) = 0$. However, it might introduce new solutions *at infinity*, i.e., $\mathbb{P}^n \setminus \mathbb{C}^n := \{[p_0 : \dots : p_n] \in \mathbb{P}^n : p_0 = 0\}$. In an analogous way, the sparse resultant Res_A of a sparse system will vanish if and only if a homogenization of the system vanishes over a normal projective toric variety X . This toric variety is a compact variety containing $(\mathbb{C}^*)^n$ which is constructed from the Minkowski sum $P := \sum_i P_i$; see, e.g., [35, Chp. 2.3] for more details on this construction. As in the classical setting, the homogenization of the system has the same solutions as the original system over $(\mathbb{C}^*)^n$, but might have extra zeros on $X \setminus (\mathbb{C}^*)^n$, which we will call *zeros at infinity*. More precisely, for each P_i , we can find a torus-invariant nef Cartier divisor D_i such that the global sections of its associated line bundle, denoted by $O_X(D_i)$, can be identified with the sparse polynomials whose Newton polytope is equal to (or contained in) P_i [35, Chp. 6], that is,

$$H^0(X_P, O_X(D_i)) = \{g \in \mathbb{C}[x_1^\pm, \dots, x_n^\pm] : \text{NP}(g) \subseteq P_i\}. \quad (1)$$

This identification corresponds to the homogenization. Somehow, if we have a global section $s \in H^0(X_P, O_X(D))$ identified with $g := \sum_{\alpha \in Q} c_\alpha x^\alpha$, then for every $p \in (\mathbb{C}^*)^n$, $s(p) = 0$ if and only if $g(p) = 0$. Using this intuition, we consider the incidence variety

$$\Omega^h := \{(p, (F_0, \dots, F_n)) \in X_P \times \mathbb{P}^A : (\forall i) \text{hom}(F_i)(p) = 0\},$$

where $\text{hom}(F_i)$ denotes the global section of $O_X(D)$ identified with F_i . If π now denotes the projection onto \mathbb{P}^A , $\pi(\Omega^h)$ is an algebraic variety (because X is projective) which we can prove agrees with $\overline{\pi(\Omega)}$. We refer the interested reader to [34, Chp. 7.3] for a more detailed explanation in the case of unmixed systems⁵.

⁴We follow Esterov's [51] and D'Andrea & Sombra's [40] definition of the sparse resultant. That is, the polynomial Res_A might not be irreducible, but it is the d -power of an irreducible polynomial, the *eliminant*, where d is the degree of π restricted to Ω ; see [40, Sec. 2]. Classically [63, 93], the sparse resultant was defined as the eliminant.

⁵Beware that the toric variety constructed in [34, Chp. 7.3] and X might not agree (X is its normalization; see [35, Chp. 3.A]), but the argument extends to X .

Remark 2.1. To define a homogenization in terms of coordinates, as we did over \mathbb{P}^n , we need to introduce a coordinate ring for X . However, such a ring depends on the way (embedding) that we use to think about X . For example, we can think about X as an *almost geometric quotient* and use its associated ring, the Cox ring [31], to homogenize the polynomials; see [35, Chp. 5]. Alternatively, we can use the so-called Cayley trick [63, Chp. 8], where we choose nef line bundles on X associated to each P_i and consider the embedding of X on the (weighted) multi-projective space defined by them. We will use the latter approach in Sec. 3.

2.2 Computing the sparse resultant

2.2.1 The determinant of a matrix. A classical way of computing the resultant is a factor of the determinant of a matrix. Sylvester [94] showed that we can compute the resultant of two univariate polynomials $f_0, f_1 \in \mathbb{C}[x_0, x_1]$ by considering the determinant of the Sylvester matrix, which linearizes the (Sylvester) map $(g_0, g_1) \mapsto g_0 f_0 + g_1 f_1$, where g_0 is a polynomial of degree at most $\deg(f_1) - 1$ and g_1 a polynomial of degree at most $\deg(f_0) - 1$. This construction was later generalized by Macaulay [81], who showed how to compute the classical (projective) resultant of $n + 1$ polynomials $f_0, \dots, f_n \in \mathbb{C}[x_1, \dots, x_n]$ of degree d_0, \dots, d_n . He also showed that there is a way of constructing monomial sets B_0, \dots, B_n such that each $x^\beta \in B_i$ has degree at most $\sum_{j \neq i} d_j - n$ and $\sum_i \#B_i$ agrees with the number of monomials of degree at most $\sum_j d_j - n$ in $\mathbb{C}[x_1, \dots, x_n]$. With these considerations, we can construct the *Macaulay matrix* M , whose columns we index with monomials in $\mathbb{C}[x_0, \dots, x_n]$ of degree at most $\sum_j d_j - n$ and the rows with pairs $\{(i, x^\beta) : i \in \{0, \dots, n\}, x^\beta \in B_i\}$. The element in M associated with the column indexed by x^α and row indexed by (i, x^β) corresponds to the coefficient of the monomial x^α in $x^\beta f_i$. Macaulay proved that the resultant of (f_0, \dots, f_n) is a factor of the determinant of M , i.e., $\det(M) = \text{Res}(f_0, \dots, f_n) \cdot E$. Moreover, he showed that the polynomial E , usually called the *extra factor*, is a specific minor of M . Observe that the matrix M is the transpose of a matrix representing the Sylvester map $(g_0, \dots, g_n) \mapsto \sum_i f_i g_i$, where the support of each g_i is contained in B_i . For this reason, we say that his construction is a *Sylvester formula* for the resultant. Besides Sylvester formulas, we observe that other maps can be used to compute resultants. We refer the reader to [48] and references therein.

Canny and Emiris [25] generalized Macaulay's construction to the sparse case. They presented a Sylvester formula for the resultant constructed out of mixed subdivisions of the Newton polytopes of the input polynomials. The interested reader can find more details and references in [34, Chp. 7.6]. However, Canny and Emiris did not characterize the extra factor appearing in their constructions. Different authors studied this question, e.g., [37, 38, 66], and very recently D'Andrea, Jeronimo, and Sombra [38] presented a complete characterization of the extra factor which allowed them to recover Macaulay's original results.

2.2.2 The determinant of a complex. A more general and elaborated way of computing the resultant is as the determinant of a complex of vector spaces. This approach is attributed to Cayley and was extended to study general kinds of resultants, such as the sparse one; the interested reader can find more details in [63, Chp. 3.4]. It turns out that the aforementioned Sylvester maps are

just the last maps appearing in certain Koszul complexes that we can use to compute resultants; see, e.g., [63, Chp. 13]. Moreover, we can use other kinds of complexes to construct smaller matrices from which to compute the resultant. In the best case scenario, the complex will involve only two non-zero modules and so its determinant will agree with the determinant of the maps between these modules. In this case, we will be able to construct *determinantal formulas*, that is, maps whose determinant is exactly the resultant, i.e., the extra factor E is a non-zero constant independent of the specific system.

Following [63], to construct the resultant as the determinant of a complex, we use sheaf cohomology. Let X be the normal projective toric variety constructed from the polytope P detailed before and consider the nef line bundles $O_X(D_0), \dots, O_X(D_n)$. Consider global sections $\tilde{f}_0, \dots, \tilde{f}_n$ such that $\tilde{f}_i \in H^0(X, O_X(D_i))$. If $(\tilde{f}_0, \dots, \tilde{f}_n)$ has no common zeros on X , the Koszul complex of sheaves \mathcal{K}_\bullet associated with the Sylvester map, defined locally at $U \subset X$ as $\text{SylV}(g_0, \dots, g_n) \mapsto \sum_i g_i \tilde{f}_i|_U$, is exact;

$$\begin{aligned} \mathcal{K}_\bullet : 0 \rightarrow O_X(-\sum_i D_i) \rightarrow \bigoplus_i O_X(-\sum_{j \neq i} D_j) \rightarrow \dots \\ \dots \rightarrow \bigoplus_{j \neq i} O_X(-D_i - D_j) \rightarrow \bigoplus_j O_X(-D_j) \xrightarrow{\text{SylV}} O_X \rightarrow 0, \end{aligned} \quad (2)$$

where $O_X(D-B) = O_X(D) \otimes O_X(B)^{-1}$, for any two Cartier divisors D and B . We will transform this complex of sheaves into a complex of vector spaces by considering its sheaf cohomology. Sheaf cohomology of line bundles on the toric variety X is a well-understood subject; for more details, see [35, Chp. 9] and [1].

Observe that, for any Cartier divisor D , the twisted complex $\mathcal{K}_\bullet \otimes O_X(D)$ is exact, because $O_X(D)$ is locally invertible. Hence, we twisted the exact complex $\mathcal{K}_\bullet \otimes O_X(\sum_i D_i)$. Every module in the twisted complex is a direct sum of sheaves of the form $O_X(\sum_{i \in I} D_i)$, for some $I \subset \{0, \dots, n\}$, so they are all nef. Hence, all the higher sheaf cohomologies of the modules in $\mathcal{K}_\bullet \otimes O_X(\sum_i D_i)$ vanish [35, Thm. 9.2.3] and so, the associated complex of global sections is exact [63, Lem. 2.2.4],

$$\begin{aligned} H^0(X, \mathcal{K}_\bullet \otimes O_X(\sum_i D_i)) : 0 \rightarrow H^0(X, O_X) \rightarrow \dots \\ \dots \rightarrow \bigoplus_j H^0(X, O_X(\sum_{i \neq j} D_i)) \xrightarrow{\text{SylV}} H^0(X, O_X(\sum_i D_i)) \rightarrow 0. \end{aligned} \quad (3)$$

The global sections of the line bundle $O_X(\sum_{i \in I} D_i)$, for $I \subset \{0, \dots, n\}$, correspond to the sparse polynomials with Newton polytope equal to (or contained in) $\sum_{i \in I} P_i$. Hence, the last map of the exact complex of global sections is the Sylvester map $(g_0, \dots, g_n) \mapsto \sum_i g_i \tilde{f}_i$, where g_i is a sparse polynomial such that $\text{NP}(g_i) \subset \sum_{j \neq i} P_j$. Canny and Emiris' matrix is a square submatrix of the matrix associated with this Sylvester map⁶. Using this construction, we can also recover a matrix related to Macaulay's construction. In this case, we consider $X = \mathbb{P}^n$ and so, its Picard group is \mathbb{Z} . We identify D_i with the degree of \tilde{f}_i and consider the complex $\mathcal{K}_\bullet \otimes O_X(\sum_i d_i - n)$. By Serre's vanishing theorem, the invertible sheaves appearing in this complex have no higher sheaf cohomologies and so, we can consider the exact complex of vector spaces $H^0(X, \mathcal{K}_\bullet \otimes O_X(\sum_i d_i - n))$. The last map of this complex is the Sylvester map $(g_0, \dots, g_n) \mapsto$

⁶However, it is not a submatrix of maximal size; see [83] for more details.

$\sum_i g_i f_i$, where g_i has degree $\sum_{j \neq i} \deg(f_j)$. Macaulay's matrix is a maximal square submatrix of this map.

We can also use this construction to understand why the Sylvester matrix gives us a determinantal formula. In this case, we have that $n = 1$ and $X = \mathbb{P}^1$. The aforementioned complex reduces to

$$0 \rightarrow \mathbb{C}[x_0, x_1]_{d_1-1} \oplus \mathbb{C}[x_0, x_1]_{d_0-1} \xrightarrow{\text{Sylv}} \mathbb{C}[x_0, x_1]_{d_0+d_1-1} \rightarrow 0,$$

where $d_i = \deg(f_i)$. The exactness of the Sylvester map determines the exactness of the full complex and agrees with Sylvester's construction. Hence, there is no extra-factor in his construction.⁷

In the general case, Weyman proposed an approach to derive from \mathcal{K}_\bullet an exact complex whose modules are direct sums of sheaves cohomologies of the form $H^i(X, \mathcal{O}_X(B))$ and such that the maps only depend on the coefficients of $(\tilde{f}_0, \dots, \tilde{f}_n)$ [102]. By twisting Weyman's complex by certain line bundles and studying the vanishing of the sheaf cohomologies appearing in it, several authors constructed smaller formulas for computing the sparse resultant, e.g. [39, 41, 103]. Moreover, by studying the case when the Weyman complex only involves two non-zero modules, this construction was used to derive determinantal formulas for the resultant of multihomogeneous systems, e.g., [6, 7, 23, 46, 47, 103]. In what follows, we present an example taken from [46, Ex. 5.3] of a determinantal formula obtained from Weyman's complex which is not a Sylvester formula.

Example 2.2 (Koszul-type formula). We consider bilinear forms $f_0, f_1, f_2 \in \mathbb{C}[x_0, x_1]_1 \otimes \mathbb{C}[y_0, y_1]_1$. Their resultant vanishes if and only if they have a common solution on $\mathbb{P}^1 \times \mathbb{P}^1$;

$$\begin{cases} f_0 = (a_{0,0} x_0 + a_{1,0} x_1) y_0 + (a_{0,1} x_0 + a_{1,1} x_1) y_1 \\ f_1 = (b_{0,0} x_0 + b_{1,0} x_1) y_0 + (b_{0,1} x_0 + b_{1,1} x_1) y_1 \\ f_2 = (c_{0,0} x_0 + c_{1,0} x_1) y_0 + (c_{0,1} x_0 + c_{1,1} x_1) y_1. \end{cases} \quad (4)$$

To construct a determinantal formula, we consider the bilinear map

$$\star : \mathbb{C}[y_0, y_1]_1 \times (\mathbb{C}[x_0, x_1]_1 \otimes \mathbb{C}[y_0, y_1]_1) \rightarrow \mathbb{C}[x_0, x_1]_1$$

$$y_i \star (x_k y_j) := \begin{cases} x_j & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

The determinant of $\phi : (\mathbb{C}[y_0, y_1]_1)^3 \rightarrow (\mathbb{C}[x_0, x_1]_1)^3$ is the resultant of the system;

$$y_i e_j \mapsto \phi(y_i e_j) := (y_i \star f_2) e_{J_1} - (y_i \star f_1) e_{J_2}, \quad (J = \{0, 1, 2\} \setminus \{j\})$$

This formula is not a Sylvester formula, but a Koszul one.

$$\begin{array}{c|cccccc} & x_0 e_0 & x_1 e_0 & x_1 e_2 & x_1 e_1 & x_0 e_2 & x_0 e_1 \\ \hline y_0 e_0 & 0 & 0 & b_{1,0} & -c_{1,0} & b_{0,0} & -c_{0,0} \\ y_1 e_0 & 0 & 0 & b_{1,1} & -c_{1,1} & b_{0,1} & -c_{0,1} \\ y_1 e_1 & -c_{0,1} & -c_{1,1} & a_{1,1} & 0 & a_{0,1} & 0 \\ y_1 e_2 & -b_{0,1} & -b_{1,1} & 0 & a_{1,1} & 0 & a_{0,1} \\ \hline y_0 e_1 & -c_{0,0} & -c_{1,0} & a_{1,0} & 0 & a_{0,0} & 0 \\ y_0 e_2 & -b_{0,0} & -b_{1,0} & 0 & a_{1,0} & 0 & a_{0,0} \end{array} \quad (5)$$

⁷Beware that the arguments at our exposition are not formal. To prove that the determinant of the complex is the resultant we need to consider the Koszul (or, more generally, the Weyman) complex not over X , but over $X \times \mathbb{P}^A$, i.e., the construction has to be done using systems with generic (symbolic) coefficients; see [63].

2.3 Solving sparse systems

In an analogous way to what we can do with the classical (projective) resultant, we can use the sparse resultant to compute all the solutions of a sparse square system. In this section, we will discuss eigenvalue methods to solve polynomial systems. Other solving strategies as the hidden-variable approach or U-resultants [34, Chp. 3.5] can be also extended; see, e.g., [40].

The following method was proposed by Emiris and Rege to recover multiplication maps using the sparse resultant [49]. Their approach generalizes ideas by Ausinger and Stetter for solving classical homogeneous systems [2]. Consider a sparse polynomial system f_1, \dots, f_n such that their number of solutions on $(\mathbb{C}^*)^n$ is the mixed volume of $\text{NP}(f_1), \dots, \text{NP}(f_n)$ and this volume is not zero. Let f_0 be a linear form in the variables x_1, \dots, x_n with Newton polytope Δ_n . We consider the matrix constructed by Canny and Emiris' algorithm for the sparse resultant of (f_0, \dots, f_n) [25]. This construction will generate monomial sets B_0, \dots, B_n, A such that $\sum \#B_i = \#A$, for any i and $x^\beta \in B_i$, for which $\text{Supp}(x^\beta f_i) \in A$, $\#B_0 = \text{MV}(P_1, \dots, P_n)$ and $B_0 \subset A$. From these sets, we consider the Macaulay matrix M whose rows are indexed by monomials in A and its columns by pairs (i, x^β) where $x^\beta \in B_i$, and where in the position corresponding to the row x^α and column (i, x^β) , the element is the coefficient of the monomial x^α in $x^\beta f_i$. We reorder M such that the bottom rows correspond to the pairs $(0, x^\beta)$ and the left-more columns corresponds to the monomials in B_0 , and we split the matrix accordingly:

$$\begin{array}{c} \left. \begin{array}{l} \{x^\beta f_i : i > 0, x^\beta \in B_i\} \\ \{x^\beta f_0 : x^\beta \in B_0\} \end{array} \right\} \begin{array}{|c|c|} \hline \begin{array}{c} A \setminus B_0 \\ \hline M_{1,1} \quad M_{1,2} \\ \hline M_{2,1} \quad M_{2,2} \end{array} \\ \hline \end{array} \quad (6)$$

Emiris and Rege showed that if the system (f_1, \dots, f_n) is generic enough then the matrix $M_{1,1}$ is invertible and so the Schur complement of $M_{2,2}$, i.e. the matrix $M_{2,2} - M_{1,2} M_{1,1}^{-1} M_{2,1}$, is the multiplication matrix of f_0 in $\mathbb{C}[x_1^\pm, \dots, x_n^\pm] / \langle f_1, \dots, f_n \rangle$. The latter holds because this matrix allows us to rewrite elements of the form $f_0 (\sum_{\beta \in B_0} c_\beta x^\beta)$ as $\sum_{\beta \in B_0} \bar{c}_\beta x^\beta + g$, where $g \in \langle f_1, \dots, f_n \rangle$. If the resultant of (f_0, \dots, f_n) does not vanish, then the map is invertible, and so B_0 is a monomial basis of $\mathbb{C}[x_1^\pm, \dots, x_n^\pm] / \langle f_1, \dots, f_n \rangle$ as its cardinal $\text{MV}(P_1, \dots, P_n)$ agrees with the dimension of the quotient ring [90]. The generic assumptions of the aforementioned construction can be relaxed [11, 91, 96] and the construction can be extended to solve overdetermined sparse systems [10, 83]. Recently, special attention has been given to the numerical aspects of this approach [10, 11, 96]. In general, we can use determinantal formulas to compute matrices which are similar to multiplication matrices, i.e., they share the same eigenvalues [6]. In contrast to the Sylvester formulas, the eigenvectors of the determinantal formulas are not well-understood, with the exception of the Koszul formulas [6].

Example 2.3 (Solving using determinantal formulas). Following our Example 2.2, we fix $f_0, f_1, f_2 \in \mathbb{C}[x_0, x_1]_1 \otimes \mathbb{C}[y_0, y_1]_1$ as in (4)

and partition the determinantal formula as in (5),

$$\left[\begin{array}{c|c} M_{1,1} & M_{1,2} \\ \hline M_{2,1} & M_{2,2} \end{array} \right]$$

We will show that, if the coordinates x_0 and y_0 of every solution of (f_1, f_2) are different to zero, then we can solve (f_1, f_2) by computing the eigenvalues of the Schur complement of $M_{2,2}$.

By assumption, the system $(x_0 y_0, f_1, f_2)$ has no solution and so the matrix $M_{1,1}$ is invertible as the resultant of $(x_0 y_0, f_1, f_2)$ is $\det(M_{1,1}) \neq 0$. We introduce a new variable λ and consider the resultant of the system $(f_0 - \lambda x_0, y_0, f_1, f_2)$. By the Poisson formula [40], the resultant is a polynomial in λ whose roots are of the form $\frac{f_0}{x_0 y_0}$ evaluated at the solutions of (f_1, f_2) . Hence, the determinantal formula for $(f_0 - \lambda x_0 y_0, f_1, f_2)$ given in (5), can be written as

$$\left[\begin{array}{c|c} M_{1,1} & M_{1,2} \\ \hline M_{2,1} & M_{2,2} - \lambda I \end{array} \right]$$

where I is the 2×2 identity matrix. Hence, the resultant of $(f_0 - \lambda x_0 y_0, f_1, f_2)$ agrees with the determinant of the Schur complement of $M_{2,2} - \lambda I$. Equivalently, the eigenvalues of the Schur complement of $M_{2,2}$ are the evaluations of $\frac{f_0}{x_0 y_0}$ at the solutions of (f_1, f_2) .

3 GRÖBNER BASES

The objective of this section is to compute Gröbner bases for sparse polynomial systems. Gröbner bases algorithms usually can incorporate a priori information of the system to speed up computations, e.g., syzygy module [45, 55] or Hilbert series [100]. The main issue when we compute with sparse polynomials is that they behave differently from dense systems, and we do not have much prior information on them. We will explain how toric geometry can help us understand the systems better and so, speed up computations.

3.1 Embedding of a toric variety

In this section, we will fix polytopes P_1, \dots, P_m and consider the projective normal toric variety X associated with $P := P_1 + \dots + P_m$ detailed in the previous section. We assume P is full-dimensional. Given an integer polytope Q , we say it is a \mathbb{N} -Minkowski summand of P if there is $k \in \mathbb{N}$ and an integer polytope S such that $Q + S = kP$. To each \mathbb{N} -Minkowski summand Q of P there is a nef Cartier divisor D_Q such that we can identify the global sections of the nef line bundle $O_X(D_Q)$ with the polynomials whose Newton polytope is equal to (or contained in) Q , as in (1) [35, Chp. 6]. Moreover, we have that for two \mathbb{N} -Minkowski summands Q, S , the global sections of $O_X(D_Q + D_S)$ correspond to the polynomials with Newton polytope $Q + S$. Given f_i with Newton polytope contained in P_i , we will homogenize f_i and write it as $\tilde{f}_i \in H^0(X, O_X(D_{P_i}))$. The common zeros of $\tilde{f}_1, \dots, \tilde{f}_m$ will determine a subscheme of X , that we denote by Y . The subscheme $Y \cap (\mathbb{C}^*)^n$ agrees with the subscheme defined by the original f_1, \dots, f_m on $(\mathbb{C}^*)^n$. Particularly, when $Y \subset (\mathbb{C}^*)^n$, Y is zero-dimensional and we can study the solutions of f_1, \dots, f_m on $(\mathbb{C}^*)^n$ by studying Y .

Example 3.1 (Intersection theory for toric varieties). If $m = n$ and Y is zero-dimensional, the mixed volume $MV(P_1, \dots, P_n)$ is actually the intersection number of D_{P_1}, \dots, D_{P_n} ; see [61, Chp. 5]. Hence, the number of solutions of f_1, \dots, f_n is bounded by the degree of

Y , and so by this mixed volume. Moreover, generically, the system $(\tilde{f}_1, \dots, \tilde{f}_n)$ has no solutions on $X \setminus (\mathbb{C}^*)^n$ and so the BKK bound is tight. In general, the BKK bound is tight when we count the solutions at infinity.

In what follows, we assume that there are \mathbb{N} -Minkowski summands Q_1, \dots, Q_r of P and vectors $d_1, \dots, d_m \in \mathbb{Z}^r$ such that $P_i = \sum_k d_{i,k} Q_k$. We consider the \mathbb{N}^r -graded algebra given by

$$R^h := \bigoplus_{b \in \mathbb{N}^r} H^0 \left(X, O_X \left(\sum_k b_k D_{Q_k} \right) \right) = \bigoplus_{b \in \mathbb{N}^r} \bigoplus_{\alpha \in (\sum_k b_k Q_k) \cap \mathbb{Z}^n} \mathbb{C} x^{(\alpha, b)}.$$

The Multi-Proj of R^h is the embedding of X in a (weighted) multi-projective space.

Hence, given polynomials f_1, \dots, f_m with Newton polytopes P_1, \dots, P_m , we can homogenize them as $(\tilde{f}_1, \dots, \tilde{f}_m)$ such that, for each i , $\tilde{f}_i \in S_{d_i}^h$. If the polynomials f_1, \dots, f_m are generic enough, we can predict the properties of $(\tilde{f}_1, \dots, \tilde{f}_m)$ on R^h . For this reason, we will compute Gröbner bases over R^h and use them to compute Gröbner bases over other rings. To formalize this, we introduce the notion of Gröbner bases over pointed affine semigroup algebras.

3.2 Gröbner bases for semigroup algebras

In this section, we will only consider affine semigroups with identity, that is, a semigroup $(S, +)$ with identity 0, isomorphic to a finitely-generated submonoid of \mathbb{Z}^k . We say that $(S, +)$ is pointed if $\alpha \in S \setminus \{0\}$, $-\alpha \notin S$, where $-\alpha$ is defined in the smallest group containing S . Given a semigroup, we define the semigroup algebra $\mathbb{C}[S]$ as the monomial \mathbb{C} -algebra generated by the monomials $\{x^\alpha : \alpha \in S\}$.

If $(S, +)$ is a pointed semigroup, it is possible to define a monomial ordering $<$ for $\mathbb{C}[S]$; see [53, Def. 3.1]. For example, if $S \subset \mathbb{N}^s$, for some $s \in \mathbb{N}$, then any monomial ordering for the standard algebra $\mathbb{C}[x_1, \dots, x_s]$ induces a monomial ordering on $\mathbb{C}[S]$. Hence, we can define Gröbner bases for ideals in $\mathbb{C}[S]$ using the standard definition, i.e., $G \subset \mathbb{C}[S]$ is a Gröbner basis for an ideal $I \subset \mathbb{C}[S]$ with respect to a monomial ordering $<$ if $G \subset I$ and, for every $f \in I$, there is $g \in G$ and $x^\alpha \in \mathbb{C}[S]$ such that $x^\alpha \text{LM}_<(f) = \text{LM}_<(g)$, where $\text{LM}_<(\cdot)$ is the leading monomial in $\mathbb{C}[S]$ with respect to $<$.

Given $\alpha \in \mathbb{R}^n \times \mathbb{R}^r$, let π_1, π_2 be its projections onto \mathbb{R}^n and \mathbb{R}^r , respectively. Clearly, as S is a semigroup, $\pi_1(S)$ and $\pi_2(S)$ are also semigroups. In what follows, we assume that, for the affine pointed semigroup S , it holds that $\pi_1(S)$ and $\pi_2(S)$ are also an affine pointed semigroups and the preimage of each $\gamma \in \pi_2(S)$ is finite. Somehow, we will think about $\pi_2(S)$ as a grading of S . In this case, we say that a monomial ordering $<$ for S is *graded* if there is another monomial ordering $<_1$ for $\pi_1(S)$ and $<_2$ for $\pi_2(S)$ such that

$$x^\alpha < x^\beta \iff \begin{cases} x^{\pi_2(\alpha)} <_2 x^{\pi_2(\beta)} \text{ or} \\ x^{\pi_2(\alpha)} = x^{\pi_2(\beta)} \text{ and } x^{\pi_1(\alpha)} <_1 x^{\pi_1(\beta)} \end{cases}$$

The algebra R^h defined in Subsection 3.1 corresponds to the semigroup algebra associated with the semigroup

$$S^h := \left\{ (\alpha, b) \in \mathbb{Z}^n \times \mathbb{Z}^r : \alpha \in \left(\sum_i b_i Q_i \right) \cap \mathbb{Z}^n \right\}.$$

If 0 is a vertex of each Q_i and of $\sum_i Q_i$, we can see that S^h satisfies the assumptions on the projections π_1 and π_2 detailed above. As the solution set of (f_1, \dots, f_m) over $(\mathbb{C}^*)^n$ does not change if we multiply the polynomials by monomials, equivalently, if we translate their Newton polytopes, we will assume with no loss of generality than 0 is a vertex of the aforementioned polytopes. Therefore, we consider the semigroup algebra

$$R := \mathbb{C}[\pi_1(S^h)] \subset \mathbb{C}[x_1^\pm, \dots, x_n^\pm].$$

We define π as the homomorphism between R^h and R taking $x^\alpha \mapsto \pi(x^\alpha) := x^{\pi_1(\alpha)}$. This morphism acts as a *dehomogenization* and it relates to graded orders as follow. Given a Gröbner basis G of an ideal of $I \subset R^h$ with respect to a graded order $<$, the set $\pi(G)$ is a Gröbner basis for $\pi(I) \subset R$ with respect to $<_1$ [53, Prop. 3.5].

3.3 Computing Gröbner bases

In this section we discuss an algorithm to compute Gröbner bases for an ideal $\langle f_1, \dots, f_r \rangle \subset R$ with respect to a monomial ordering $<_1$. For this, we will compute Gröbner bases for $\langle \tilde{f}_1, \dots, \tilde{f}_r \rangle \subset R^h$ with respect to a graded monomial ordering $<$ associated with $<_1$. In what follows, we ignore the subindex from the ordering $<_1$.

For each multidegree $b \in \mathbb{N}^r$, we will construct a Macaulay matrix M_b where the columns are indexed by the monomials of degree b in R^h , the rows by the pairs (i, x^β) , where x^β is a monomial of degree $b - d_i$. The rows are sorted in decreasing order with respect to $<$ and the columns are sorted in decreasing order in the following way: if $i > j$, or $i = j$ and $x^\alpha > x^\beta$, then $(i, x^\alpha) < (j, x^\beta)$. The element of the matrix M_b of column index x^α , and row (i, x^β) is the coefficient of the monomial x^α in $x^\beta \tilde{f}_i$. As Lazard realized [80], if we perform Gaussian elimination with no pivoting, we obtain new rows representing elements in the Gröbner basis of $\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle$ of degree b . As the Gröbner basis is finite, if we perform this computation for sufficiently many different degrees, we recover the complete Gröbner basis.

A classical bottleneck of Gröbner bases computations are reductions to zero. In the sketched algorithm, these reductions to zero correspond to the rows reducing to zero after performing Gaussian elimination. An established way of speeding up computations is to avoid these reductions using the F5 criterion [45, 55]. In this context, the F5 criterion is called Matrix-F5 [45, Sec. 3] and translates into skipping some rows from the construction of the matrix. The F5 criterion tells us that the row indexed by (i, x^β) will reduce to zero after performing Gaussian elimination if and only if x^β is the leading monomial of a polynomial of degree $b - d_i$ in the colon ideal $\langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle : f_i$, i.e., $x^\beta \in LM(\langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle : f_i)_{b-d_i}$. We can organize our computation in such a way that we can recover from it $LM(\langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle)_{b-d_i}$ [45, Sec. 3]. Hence, when these two sets agree, we can skip every reduction to zero, that is, the F5 criterion is *optimal*. A classical setting where this happens is when f_i is not a zero divisor in $R^h / \langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle$. Hence, when $(\tilde{f}_1, \dots, \tilde{f}_m)$ is a regular sequence, i.e., when the previous condition holds for every i , then we can avoid every reduction to zero. In the unmixed case, we have that $r = 1$ and the ring R^h is \mathbb{N} -graded. It can be shown that this is a Cohen-Macaulay ring and that, if $m \leq n$,

a generic unmixed sparse system $(\tilde{f}_1, \dots, \tilde{f}_m)$ forms a regular sequence on R^h [69]. This last fact was exploited by Faugère, Spaenlehauer, and Svartz [53] to compute Gröbner bases for unmixed systems. However, in the mixed case, this is not true anymore, that is, our system $(\tilde{f}_1, \dots, \tilde{f}_m)$ usually does not form a regular sequence on R^h . To solve this problem, together with Faugère and Tsigaridas [8, 9], we observed that the F5 criterion is optimal at *certain degree* b whenever $\langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle : f_i = \langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle_{b-d_i}$, for each i . This condition can be enforced by the vanishing of the first homology of the Koszul complex of $(\tilde{f}_1, \dots, \tilde{f}_i)$ at degree b , for each i . We can prove that, if the subscheme defined by $(\tilde{f}_1, \dots, \tilde{f}_i)$ on X is a complete intersection, then the first homology of the Koszul complex of $(\tilde{f}_1, \dots, \tilde{f}_i)$ vanishes at any degree $b \geq \sum_{j \leq i} d_j$. Moreover, this bound can be improved for special systems. For example, for multihomogeneous systems over $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_r}$, we have that $b \geq \sum_{j \leq i} d_j - (n_1, \dots, n_r)$ suffices [8]⁸. Hence, we can predict all the reductions to zero appearing at “big-enough” degree. Moreover, if our objective is to only compute Gröbner bases over R , we will just avoid the degrees at which we cannot predict every reduction.

Remark 3.2. This strategy generalizes partially previous work on computing Gröbner bases for special sparse systems such as bilinear [58] and weighted homogeneous systems [59].

To derive complexity bounds, we need to construct a bound of the maximal degree of an element in a Gröbner bases of $(\tilde{f}_1, \dots, \tilde{f}_m)$. However, the monomial orderings defined over semigroup algebras, in general, does not have the dehomogenization property of GRevLex [5, Ex. 7.2.1]. Moreover, we cannot talk about generic coordinates for the solutions, as random change of coordinates destroys the sparsity. Hence, we cannot use strategies as [3] to construct bounds for the maximal degrees of the elements in the Gröbner bases. As we will show in the next section, we can work around this problem in the zero-dimensional case, i.e., when the subscheme associated to $(\tilde{f}_1, \dots, \tilde{f}_m)$ has dimension zero.

3.4 Zero-dimensional systems

In this section, we will construct bounds for computing Gröbner bases of zero-dimensional ideals. Classically, to compute Gröbner bases for zero-dimensional systems, we first calculate a Gröbner basis with respect to GRevLex and then we use FGLM to recover the Gröbner basis with respect to the ordering we want. We do so because, with respect to GRevLex, the Gröbner basis involves elements of smaller degrees, for which we may find upper bounds. Generally, the maximal degree of an element in a Gröbner basis of a zero-dimensional ideal is given by the *Castelnuovo-Mumford regularity* of its homogenization [27, Cor. 3]. In the semigroup setting,

⁸A spectre is haunting this argument - a spectral sequence associated to the Koszul complex of sheaves mentioned in the previous section. Normal projective toric varieties are Cohen-Macaulay, so if $(\tilde{f}_1, \dots, \tilde{f}_i)$ defines a complete intersection Y on X , the Koszul complex gives a resolution for the structure sheaf \mathcal{O}_Y . In this case, we can transform this complex into a resolution of $R^h / \langle \tilde{f}_1, \dots, \tilde{f}_{i-1} \rangle$ at degree $b \in \mathbb{N}^r$, if for each $I \subseteq \{1, \dots, i\}$, $H^t(X, \mathcal{O}_X(\sum_j (b_j - \sum_{k \in I} d_{j,k}) D_{Q_j})) = 0$, for $t > 0$. It is easy to construct a priori bounds for b as $\sum_j d_j$ using vanishing theorems for the sheaf cohomology of line bundles over toric varieties; see, e.g., [11, Sec. 4]. In our papers [8, 9], we skipped this discussion by defining the concept of *Koszul regular systems* as the systems such that, for every subsystem, the first homology of its associated Koszul complex vanishes at certain degrees, which we can predict from the previous analysis.

where there usually does not exist a GRevLex ordering, the maximal degree of an element in the Gröbner basis could be higher than the regularity; see [5, Sec. 8.3.1]. To work around this problem, in [8, 9], we presented a way to recover the multiplication maps by truncating our computation of Gröbner bases at a certain degree. By doing so, we bounded the complexity of our computation. In what follows, we explain the idea behind this strategy in the general case of mixed sparse polynomial systems and refer the reader to [5, Chp. 8] for improvements in special cases.

Following the notation from the previous section, we consider an affine sparse system (f_1, \dots, f_m) such that the subscheme Y associated with $\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle \subset R^h$ has only solutions on $(\mathbb{C}^*)^n$ (in particular, Y is finite as X is compact). That is, it has no solutions at infinity. We assume without loss of generality that $1, x_1, \dots, x_n \in R$.⁹ In this case, we can prove [9, Lem. 4.11],

$$R/\pi(\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle) \simeq \mathbb{C}[x_1, \dots, x_n]/(\langle f_1, \dots, f_n \rangle : \langle \prod_i x_i \rangle^\infty).$$

Our strategy will be to perform FGLM using multiplication maps over $R/\pi(\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle)$ to recover the Gröbner basis of $\langle f_1, \dots, f_m \rangle : \langle \prod_i x_i \rangle^\infty$ in $\mathbb{C}[x_1, \dots, x_n]$. In what follows, we focus on computing these multiplication maps by adapting Emiris and Rege [49].

Consider $b = \sum_i d_i$, where d_i is the multidegree of \tilde{f}_i . Let d_0 be a multidegree such that $\Delta_n \cap \mathbb{N}^n \subset \pi(R_{d_0}^h)$, that is, after dehomogenization, we can obtain a linear form in R . At the degrees b and $b + d_0$, the ideal $R^h/\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle$ is a vector space of dimension δ equal to the number of points in Y , counting multiplicities [83, Thm. 3]. Let B_0 be the set of monomials of degree d which are not leading monomials in $LM(\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle)$. The cardinality of B_0 is δ . Moreover, also by [83, Thm. 3], as there are no solutions outside $(\mathbb{C}^*)^n$, for any monomial $x^\alpha \in R_{d_0}^h$, $R_{b+d_0}^h = \langle \tilde{f}_1, \dots, \tilde{f}_m, x^\alpha \rangle_{b+d_0}$, and using the F5 criterion, it is easy to see that the monomial set $\{x^\alpha x^\beta : x^\beta \in B_0\}$ forms a basis of the vector space $(R^h/\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle)_{b+d_0}$. Let x^{α_0} be the monomial of degree $b + d_0$ such that $\pi(x^{\alpha_0}) = 1$. Then, we can use our Macaulay matrix at degree $b + d_0$ related to $(\tilde{f}_1, \dots, \tilde{f}_m)$ to rewrite any product $x^\beta f_0$, where $f_0 \in R_{d_0}^h$ and $x^\beta \in B_0$, as a linear combination of monomials in $\{x^\alpha x^\beta : x^\beta \in B_0\}$. By dehomogenizing this relation, we recover the multiplication map of $\pi(f_0)$ over $R/\pi(\langle \tilde{f}_1, \dots, \tilde{f}_m \rangle)$, with respect to the basis $\pi(B_0)$. As the monomials $1, x_1, \dots, x_n$ belong to $\pi(B_0)$, we can recover every multiplication map in $\mathbb{C}[x_1, \dots, x_n]/(\langle f_1, \dots, f_n \rangle : \langle \prod_i x_i \rangle^\infty)$.

ACKNOWLEDGMENTS

This manuscript is dedicated to the memory of Agnes Szanto. Thank you very much, Agnes, for your selfless help and support.

I would like to thank Christopher Borger, Jean-Charles Faugère, Angelos Mantzaflaris, Pierre-Jean Spaenlehauer, Elias Tsigaridas, and Simon Telen, for our joint work where I learned a lot about what is detailed in this manuscript. Also, to Laurent Busé and Marc Chardin, for teaching me how to use local cohomology and spectral sequences to study the Castelnuovo-Mumford regularity. I am grateful to Fatima Abu Salem and Amir Hashemi for their several comments and corrections on this manuscript.

⁹If the monomials do not belong to R , we change the toric variety X by another one associated with $P + \Delta_n$. As $Y \subset (\mathbb{C}^*)^n$, this change does not affect Y and adds a degree d_0 to R^h as wanted; see [9, Sec. 4].

This work was funded by the ERC under the European's Horizon 2020 research and innovation programme (grant agreement 787840).

REFERENCES

- [1] Klaus Altmann and David Ploog. 2020. Displaying the cohomology of toric line bundles. *Izvestiya: Mathematics* 84, 4 (2020), 683.
- [2] Winfried Auzinger and Hans J. Stetter. 1988. An elimination algorithm for the computation of all zeros of a system of multivariate polynomial equations. In *Numerical Mathematics Singapore 1988: Proceedings of the International Conference on Numerical Mathematics held at the National University of Singapore, May 31–June 4, 1988*, Ravi P. Agarwal, Y. M. Chow, and S. J. Wilson (Eds.), 11–30.
- [3] David Bayer and Michael Stillman. 1987. A criterion for detecting m-regularity. *Inventiones Mathematicae* 87, 1 (Feb. 1987), 1–11.
- [4] Erick Rodriguez Bazan and Evelyne Hubert. 2020. Ideal interpolation, H-bases and symmetry. In *Proceedings of the 45th international symposium on symbolic and algebraic computation (Kalamata, Greece) (ISSAC '20)*, 402–409.
- [5] Matias R. Bender. 2019. *Algorithms for sparse polynomial systems: Groebner basis and resultants*. Ph. D. Dissertation. Sorbonne Université.
- [6] Matias R. Bender, Jean-Charles Faugère, Angelos Mantzaflaris, and Elias Tsigaridas. 2018. Bilinear systems with two supports: Koszul resultant matrices, eigenvalues, and eigenvectors. In *Proceedings of the 2018 ACM international symposium on symbolic and algebraic computation (ISSAC '18)*, 63–70.
- [7] Matias R. Bender, Jean-Charles Faugère, Angelos Mantzaflaris, and Elias Tsigaridas. 2021. Koszul-Type determinantal formulas for families of mixed multilinear systems. *SIAM Journal on Applied Algebra and Geometry* 5, 4 (2021), 589–619.
- [8] Matias R. Bender, Jean-Charles Faugère, and Elias Tsigaridas. 2018. Towards mixed gröbner basis algorithms: the multihomogeneous and sparse case. In *Proceedings of the 2018 ACM international symposium on symbolic and algebraic computation (ISSAC '18)*, 71–78.
- [9] Matias R. Bender, Jean-Charles Faugère, and Elias Tsigaridas. 2019. Groebner basis over semigroup algebras: Algorithms and applications for sparse polynomial systems. In *Proceedings of the 2019 on international symposium on symbolic and algebraic computation (ISSAC '19)*, 42–49.
- [10] Matias R. Bender and Simon Telen. 2021. Yet another eigenvalue algorithm for solving polynomial systems. *arXiv:2105.08472 [cs, math]* (May 2021).
- [11] Matias R. Bender and Simon Telen. 2022. Toric eigenvalue methods for solving sparse polynomial systems. *Math. Comp.* (2022). In press; arXiv: 2006.10654.
- [12] David N. Bernshtein. 1975. The number of roots of a system of equations. *Functional Analysis and Its Applications* 9, 3 (July 1975), 183–185.
- [13] Jérémy Berthomieu, Christian Eder, and Mohab Safey El Din. 2021. Msolve: a library for solving polynomial systems. In *Proceedings of the 2021 on international symposium on symbolic and algebraic computation (Virtual Event, Russian Federation) (ISSAC '21)*, 51–58.
- [14] Frédéric Bihan and Alicia Dickenstein. 2017. Descartes' rule of signs for polynomial systems supported on circuits. *International Mathematics Research Notices* 2017, 22 (2017), 6867–6893.
- [15] Frédéric Bihan and Ivan Soprunov. 2019. Criteria for strict monotonicity of the mixed volume of convex polytopes. *Advances in Geometry* 19, 4 (2019), 527–540.
- [16] Frédéric Bihan and Frank Sottile. 2007. New fewnomial upper bounds from Gale dual polynomial systems. *Moscow mathematical journal* 7, 3 (2007), 387–407.
- [17] Paul Breiding, Türkü Özlüm Çelik, Timothy Duff, Alexander Heaton, Aida Maraj, Anna-Laura Sattelberger, Lorenzo Venturello, and Oğuzhan Yürük. 2021. Nonlinear algebra and applications. *Numerical Algebra, Control and Optimization* (2021).
- [18] Peter Bürgisser and Felipe Cucker. 2013. *Condition: The geometry of numerical algorithms*.
- [19] Peter Bürgisser, Felipe Cucker, and Pierre Lairez. 2020. Rigid continuation paths II. Structured polynomial systems. *arXiv:2010.10997* (2020).
- [20] Peter Bürgisser, Alperen A Ergur, and Josué Tonelli-Cueto. 2019. On the number of real zeros of random fewnomials. *SIAM Journal on Applied Algebra and Geometry* 3, 4 (2019), 721–732.
- [21] Michael Burr, Frank Sottile, and Elise Walker. 2022. Numerical homotopies from Khovanskii bases. *Math. Comp.* (2022). In press; arXiv: 2008.13055.
- [22] Laurent Busé. 2006. *Elimination theory in codimension one and applications*. Research Report RR-5918. INRIA, 47 pages. Notes of lectures given at the CIMP-UNESCO-IRAN school in Zanjan, Iran, July 9-22 2005.
- [23] Laurent Busé, Angelos Mantzaflaris, and Elias Tsigaridas. 2020. Matrix formulæ for resultants and discriminants of bivariate tensor-product polynomials. *Journal of Symbolic Computation* 98 (May 2020), 65–83.
- [24] John Canny. 1990. Generalised characteristic polynomials. *Journal of Symbolic Computation* 9, 3 (March 1990), 241–250.

- [25] John Canny and Ioannis Emiris. 1993. An efficient algorithm for the sparse mixed resultant. In *Applied algebra, algebraic algorithms and error-correcting codes*, Gérard Cohen, Teo Mora, and Oscar Moreno (Eds.), 89–104.
- [26] John Canny and J. Maurice Rojas. 1991. An optimal condition for determining the exact number of roots of a polynomial system. In *Proceedings of the 1991 international symposium on Symbolic and algebraic computation (ISSAC '91)*, 96–102.
- [27] Marc Chardin. 2003. Bounds for Castelnuovo-Mumford regularity in terms of degrees of defining equations. In *Commutative algebra, singularities and computer algebra*, Jürgen Herzog and Victor Vuletescu (Eds.), 67–73.
- [28] Changbo Chen and Marc Moreno Maza. 2012. Algorithms for computing triangular decomposition of polynomial systems. *Journal of Symbolic Computation* 47, 6 (2012), 610–642.
- [29] Tianran Chen. 2019. Unmixing the mixed volume computation. *Discrete & Computational Geometry* 62, 1 (July 2019), 55–86.
- [30] Diego Cifuentes and Pablo A Parrilo. 2016. Exploiting chordal structure in polynomial ideals: A grobner bases approach. *SIAM Journal on Discrete Mathematics* 30, 3 (2016), 1534–1570.
- [31] David A. Cox. 1995. The homogeneous coordinate ring of a toric variety, Revised version. *Journal of Algebraic Geometry* 4 (1995), 17–50.
- [32] David A. Cox. 2021. Stickelberger and the eigenvalue theorem. In *Commutative algebra: Expository papers dedicated to David Eisenbud on the occasion of his 75th birthday*, Irena Peeva (Ed.), 283–298.
- [33] David A. Cox, Carlos D'Andrea, Alicia Dickenstein, Jonathan D. Hauenstein, Henry K. Schenck, and Jessica Sidman. 2020. *Applications of polynomial systems*. Number 134 in Conference Board of the Mathematical Sciences CBMS regional conference series in mathematics.
- [34] David A. Cox, John Little, and Donal O'Shea. 2005. *Using algebraic geometry* (2 ed.).
- [35] David A. Cox, John B. Little, and Henry K. Schenck. 2011. *Toric varieties*.
- [36] Felipe Cucker, Teresa Krick, Gregorio Malajovich, and Mario Wschebor. 2008. A numerical algorithm for zero counting. I: Complexity and accuracy. *Journal of Complexity* 24, 5-6 (2008), 2595–2629.
- [37] Carlos D'Andrea. 2002. Macaulay style formulas for sparse resultants. *Trans. Amer. Math. Soc.* 354, 7 (2002), 2595–2629.
- [38] Carlos D'Andrea, Gabriela Jeronimo, and Martín Sombra. 2022. The Canny-Emiris conjecture for the sparse resultant. *Foundations of Computational Mathematics* (2022), 1–61.
- [39] Carlos D'Andrea and Amit Khetan. 2005. Macaulay style formulas for toric residues. *Compositio Mathematica* 141, 03 (May 2005), 713–728.
- [40] Carlos D'Andrea and Martín Sombra. 2015. A Poisson formula for the sparse resultant. *Proceedings of the London Mathematical Society* 110, 4 (2015), 932–964.
- [41] Alicia Dickenstein and Ioannis Z. Emiris. 2003. Multihomogeneous resultant formulae by means of complexes. *Journal of Symbolic Computation* 36, 3 (2003), 317–342.
- [42] Alicia Dickenstein and Ioannis Z. Emiris (Eds.). 2005. *Solving polynomial equations*. Algorithms and Computation in Mathematics, Vol. 14.
- [43] Dimitrios I Diochnos, Ioannis Z Emiris, and Elias P Tsigaridas. 2009. On the asymptotic and practical complexity of solving bivariate systems over the reals. *Journal of Symbolic Computation* 44, 7 (2009), 818–835.
- [44] Mathias Drton, Bernd Sturmfels, and Seth Sullivant. 2009. *Lectures on algebraic statistics*.
- [45] Christian Eder and Jean-Charles Faugère. 2017. A survey on signature-based algorithms for computing Gröbner bases. *Journal of Symbolic Computation* 80 (2017), 719–784.
- [46] Ioannis Z Emiris and Angelos Mantzaflaris. 2012. Multihomogeneous resultant formulae for systems with scaled support. *Journal of Symbolic Computation* 47, 7 (2012), 820–842.
- [47] Ioannis Z. Emiris, Angelos Mantzaflaris, and Elias P. Tsigaridas. 2021. Multilinear polynomial systems: Root isolation and bit complexity. *Journal of Symbolic Computation* 105 (July 2021), 145–164.
- [48] Ioannis Z. Emiris and Bernard Mourrain. 1999. Matrices in elimination theory. *Journal of Symbolic Computation* 28, 1 (July 1999), 3–44.
- [49] Ioannis Z. Emiris and Ashutosh Rege. 1994. Monomial bases and polynomial system solving (extended abstract). In *Proceedings of the international symposium on Symbolic and algebraic computation - ISSAC '94*, 114–122.
- [50] Alperen A. Ergür and Timo de Wolff. 2021. A polyhedral homotopy algorithm for real zeros. *arXiv:1910.01957 [cs, math]* (Feb. 2021).
- [51] Alexander Esterov. 2010. Newton polyhedra of discriminants of projections. *Discrete & Computational Geometry* 44, 1 (2010), 96–148.
- [52] Jean-Charles Faugère, Mohab Safey El Din, and Pierre-Jean Spaenlehauer. 201. Computing loci of rank defects of linear matrices using Gröbner bases and applications to cryptology. In *Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation*, 257–264.
- [53] Jean-Charles Faugère, Pierre-Jean Spaenlehauer, and Jules Svartz. 2014. Sparse Gröbner bases: the unmixed case. *Proceedings of the 39th International Symposium on Symbolic and Algebraic Computation - ISSAC '14* (2014), 178–185.
- [54] Jean-Charles Faugère and Jules Svartz. 2013. Gröbner bases of ideals invariant under a commutative group: the non-modular case. In *Proceedings of the 38th international symposium on symbolic and algebraic computation*, 347–354.
- [55] Jean Charles Faugère. 2002. A new efficient algorithm for computing Gröbner bases without reduction to zero (F5). In *Proceedings of the 2002 international symposium on symbolic and algebraic computation (ISSAC '02)*, 75–83.
- [56] Jean-Charles Faugère, Patrizia Gianni, Daniel Lazard, and Teo Mora. 1993. Efficient computation of zero-dimensional Gröbner bases by change of ordering. *Journal of Symbolic Computation* 16, 4 (Oct. 1993), 329–344.
- [57] Jean-Charles Faugère and Chenqi Mou. 2017. Sparse FGLM algorithms. *Journal of Symbolic Computation* 80 (May 2017), 538–569.
- [58] Jean-Charles Faugère, Mohab Safey El Din, and Pierre-Jean Spaenlehauer. 2011. Gröbner bases of bihomogeneous ideals generated by polynomials of bidegree (1,1): Algorithms and complexity. *Journal of Symbolic Computation* 46, 4 (April 2011), 406–437.
- [59] Jean-Charles Faugère, Mohab Safey El Din, and Thibaut Verron. 2016. On the complexity of computing Gröbner bases for weighted homogeneous systems. *Journal of Symbolic Computation* 76 (Sept. 2016), 107–141.
- [60] Jean-Charles Faugère. 1999. A new efficient algorithm for computing Gröbner bases (F4). *Journal of Pure and Applied Algebra* 139, 1 (June 1999), 61–88.
- [61] William Fulton. 1993. *Introduction to toric varieties*.
- [62] Karin Gatermann and Birkett Huber. 2002. A family of sparse polynomial systems arising in chemical reaction systems. *Journal of Symbolic Computation* 33, 3 (2002), 275–305.
- [63] Israel M. Gelfand, Mikhail M. Kapranov, and Andrey V. Zelevinsky. 2008. *Discriminants, resultants, and multidimensional determinants*.
- [64] Marc Giusti, Grégoire Lecerf, and Bruno Salvy. 2001. A Gröbner free alternative for polynomial system solving. *Journal of Complexity* 17, 1 (2001), 154–211.
- [65] Laureano González-Vega. 1991. A subresultant theory for multivariate polynomials. In *Proceedings of the 1991 international symposium on symbolic and algebraic computation (ISSAC '91)*, 79–85. event-place: Bonn, West Germany.
- [66] Friedemann Groh. 2020. Subdivisions for Macaulay formulas of sparse systems. In *Proceedings of the 45th international symposium on symbolic and algebraic computation*, 218–225.
- [67] Joos Heintz, Teresa Krick, Susana Puddu, Juan Sabia, and Ariel Weissbein. 2000. Deformation techniques for efficient polynomial equation solving. *Journal of complexity* 16, 1 (2000), 70–109.
- [68] María Isabel Herrero, Gabriela Jeronimo, and Juan Sabia. 2013. Affine solution sets of sparse polynomial systems. *Journal of Symbolic Computation* 51 (2013), 34–54.
- [69] Melvin Hochster. 1972. Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes. *Annals of Mathematics* 96, 2 (1972), 318–337.
- [70] Birkett Huber and Bernd Sturmfels. 1995. A polyhedral method for solving sparse polynomial systems. *Math. Comp.* 64, 212 (1995), 1541–1555.
- [71] Birkett Huber and Jan Verschelde. 2000. Pieri homotopies for problems in enumerative geometry applied to pole placement in linear systems control. *SIAM Journal on Control and Optimization* 38, 4 (2000), 1265–1287.
- [72] Gorav Jindal, Anurag Pandey, Himanshu Shukla, and Charilaos Zisopoulos. 2020. How many zeros of a random sparse polynomial are real?. In *Proceedings of the 45th international symposium on symbolic and algebraic computation*, 273–280.
- [73] Askold G. Khovanskii. 1978. Newton polyhedra and the genus of complete intersections. *Functional Analysis and Its Applications* 12, 1 (Jan. 1978), 38–46.
- [74] Askold G. Khovanskii. 1991. *Fewnomials*. Vol. 88.
- [75] Martin Kreuzer and Lorenzo Robbiano. 2000. *Computational commutative algebra 1*.
- [76] Anatolii G. Kushnirenko. 1976. Newton polytopes and the Bézout theorem. *Functional Analysis and Its Applications* 10, 3 (July 1976), 233–235.
- [77] George Labahn, Mohab Safey El Din, Éric Schost, and Thi Xuan Vu. 2021. Homotopy techniques for solving sparse column support determinantal polynomial systems. *Journal of Complexity* 66 (2021), 101557.
- [78] Yagati N. Lakshman and Daniel Lazard. 1991. On the complexity of zero-dimensional algebraic systems. In *Effective methods in algebraic geometry*. Springer, 217–225.
- [79] Daniel Lazard. 1981. Resolution des systemes d'equations algebriques. *Theoretical Computer Science* 15, 1 (Jan. 1981), 77–110.
- [80] Daniel Lazard. 1983. Gröbner-Bases, gaussian elimination and resolution of systems of algebraic equations. In *Proceedings of the european computer algebra conference on computer algebra (EUROCAL '83)*, 146–156.
- [81] Francis Sowerby Macaulay. 1916. *The algebraic theory of modular systems*. Reprint in 1994.
- [82] Gregorio Malajovich. 2019. Complexity of sparse polynomial solving: homotopy on toric varieties and the condition metric. *Foundations of Computational Mathematics* 19, 1 (2019), 1–53.
- [83] César Massri. 2016. Solving a sparse system using linear algebra. *Journal of Symbolic Computation* 73 (March 2016), 157–174.

- [84] Michael Monagan and Roman Pearce. 2017. An algorithm for splitting polynomial systems based on F4. In *Proceedings of the international workshop on parallel symbolic computation*. 1–5.
- [85] Pinaki Mondal. 2021. *How many zeroes?: Counting solutions of systems of polynomials via toric geometry at infinity*. CMS/CAIMS Books in Mathematics, Vol. 2.
- [86] Alexander Morgan and Andrew Sommese. 1987. Computing all solutions to polynomial systems using homotopy continuation. *Appl. Math. Comput.* 24, 2 (1987), 115–138.
- [87] Chenqi Mou, Yang Bai, and Jiahua Lai. 2021. Chordal graphs in triangular decomposition in top-down style. *Journal of Symbolic Computation* 102 (2021), 108–131.
- [88] Bernard Mourrain. 1999. A new criterion for normal form algorithms. In *Applied algebra, algebraic algorithms and error-correcting codes (Lecture Notes in Computer Science)*, Marc Fossorier, Hideki Imai, Shu Lin, and Alain Poli (Eds.). 430–442.
- [89] Bernard Mourrain and Jean Pascal Pavone. 2009. Subdivision methods for solving polynomial equations. *Journal of Symbolic Computation* 44, 3 (2009), 292–306.
- [90] Paul Pedersen and Bernd Sturmfels. 1996. Mixed monomial bases. In *Algorithms in algebraic geometry and applications (Progress in Mathematics)*, Laureano González-Vega and Tomás Recio (Eds.). 307–316.
- [91] J. Maurice Rojas. 1999. Solving degenerate sparse polynomial systems faster. *Journal of Symbolic Computation* 28, 1 (July 1999), 155–186.
- [92] Fabrice Rouillier. 1999. Solving zero-dimensional systems through the rational univariate representation. *Applicable Algebra in Engineering, Communication and Computing* 9, 5 (1999), 433–461.
- [93] Bernd Sturmfels. 1994. On the Newton polytope of the resultant. *Journal of Algebraic Combinatorics* 3, 2 (April 1994), 207–236.
- [94] James Joseph Sylvester. 1840. XXIII. A method of determining by mere inspection the derivatives from two equations of any degree. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* 16, 101 (1840), 132–135.
- [95] Agnes Szanto. 2008. Solving over-determined systems by the subresultant method (with an appendix by Marc Chardin). *Journal of Symbolic Computation* 43, 1 (Jan. 2008), 46–74.
- [96] Simon Telen. 2020. Numerical root finding via Cox rings. *Journal of Pure and Applied Algebra* 224, 9 (2020), 106367.
- [97] Simon Telen. 2020. *Solving Systems of Polynomial Equations*. Ph. D. Dissertation. KU Leuven, Leuven, Belgium. Liras ID: LIRIAS1947881.
- [98] Simon Telen, Bernard Mourrain, and Marc van Barel. 2018. Solving polynomial systems via truncated normal forms. *SIAM J. Matrix Anal. Appl.* 39, 3 (Jan. 2018), 1421–1447.
- [99] Josué Tonelli-Cueto. 2019. *Condition and homology in semialgebraic geometry*. Doctoral Thesis. Technische Universität Berlin.
- [100] Carlo Traverso. 1996. Hilbert functions and the Buchberger algorithm. *Journal of Symbolic Computation* 22, 4 (1996), 355–376.
- [101] Dongming Wang, Rina Dong, and Chenqi Mou. 2020. Decomposition of polynomial sets into characteristic pairs. *Math. Comp.* 89, 324 (2020), 1993–2015.
- [102] Jerzy Weyman. 2003. *Cohomology of vector bundles and syzygies*.
- [103] Jerzy Weyman and Andrei Zelevinsky. 1994. Determinantal formulae for multi-graded resultants. *Journal of Algebraic Geometry* 3, 4 (1994), 569–597.