

Maximum zeroth-order general Randić index of orientations of cacti

Jiaxiang Yang¹, Hanyuan Deng^{1*}, Zikai Tang¹, Hechao Liu²

¹Key Laboratory of Computing and Stochastic Mathematics (Ministry of Education),

School of Mathematics and Statistics, Hunan Normal University,

Changsha, Hunan 410081, P. R. China.

²School of Mathematical Sciences, South China Normal University,

Guangzhou, 510631, P. R. China.

Abstract

The zeroth-order general Randić index R_{a+1}^0 of an n -vertices oriented graph D is equal to the sum of $(d_{u_i}^+)^a + (d_{u_j}^-)^a$ over all arcs $u_i u_j$ of D , where we denote by $d_{u_i}^+$ the out-degree of the vertex u_i and $d_{u_j}^-$ the in-degree of the vertex u_j , a is an arbitrary real number. In the paper, we determine the orientations of cacti with the maximum value of the zeroth-order general Randić index for $a \geq 1$.

Keywords: zeroth-order general Randić index; cactus; extremal graph.

1 Introduction

For a connected simple graph G with vertex set $V(G)$ and edge set $E(G)$, $N_G(u)$ denotes the set of all neighbors of u , and we denote by $d_G(u) = |N_G(u)|$ the degree of u in G . If $d_G(u) = 1$, then the vertex u is a pendent vertex, and $uv \in E(G)$ is a pendent edge. Topological index plays an important role in QSPR and QSAR; see [1, 2]. Li and Zheng [3] proposed first general Zagreb index. In the current study, most scholars call it the zeroth-order general Randić index which is defined as $R_a^0(G) = \sum_{u \in V(G)} (d_G(u))^a = \sum_{uv \in E(G)} [(d_G(u))^{a-1} + (d_G(v))^{a-1}]$, where a is an arbitrary real number. So, the zeroth-order general Randić index is also a VDB topological index. Over the past few decades, a great deal of study of researchers have shown many results of extremal zeroth-order general Randić index of a graph family; see [4–11]. Monsalve and Rada gave the definition of VDB topological indices over orientations of a graph in [13], and brought us some help to study extremal VDB topological indices of orientations of a graph in [12].

For $u, v \in V(G)$, we can add an edge $uv \notin E(G)$ to graph G and get the graph $G + uv$. For $u, v \in V(G)$, we can delete an edge $uv \in E(G)$ and get the graph $G - uv$. If we delete $u \in V(G)$ and all $e \in \{e | e = uv \in E(G), v \in V(G)\}$, then we get the graph $G - u$.

*Corresponding author: hydeng@hunnu.edu.cn

If a connected graph G only consists of blocks which are either edges or cycles, then G is called a cactus. Let $\delta(G) = \min\{d(u)|u \in V(G)\}$.

For a simple digraph D , the digraph D has vertex set $V(D)$ and arc set $A(D)$, where $uv \in A(D)$ is an arc that from vertex u to vertex v in the graph D . For vertex $u \in V(D)$, the out-degree $d_u^+ = |\{v|uv \in A(D), v \in V(D)\}|$ (resp. in-degree $d_u^- = |\{v|vu \in A(D), v \in V(D)\}|$).

For a vertex $w \in \{w|d_w^+ = 0, w \in V(D)\}$, we call w a sink vertex in D . For a vertex $w \in \{w|d_w^- = 0, w \in V(D)\}$, we call w a source vertex in D . For a vertex $w \in \{w|d_w^+ = d_w^- = 0, w \in V(D)\}$, we call w an isolated vertex in D . For each $uv \in E(G)$, we use $uv \in A$ or $vu \in A$ to replace $uv \in E(G)$ and get an oriented graph D (i.e. an orientation of G). An orientation of G is called a sink-source orientation of G if $d_w^+ = 0$ or $d_w^- = 0$ for any $w \in V(G)$. Let $\mathcal{O}(G) = \{D|D \text{ is a orientation of } G\}$.

The definition of the VDB topological indices of a digraph D is as [13]

$$\varphi(D) = \frac{1}{2} \sum_{uv \in A} \varphi(d_u^+, d_v^-),$$

where $\varphi(d_u^+, d_v^-)$ is an arbitrary real number. Specially, if $\varphi(i, j) = i^{a-1} + j^{a-1}$, then $\varphi(D)$ is the zeroth-order general Randić index of a digraph, denoted by $R_a^0(D)$, i.e.,

$$R_a^0(D) = \frac{1}{2} \sum_{uv \in A} [(d_u^+)^{a-1} + (d_v^-)^{a-1}].$$

Monsalve and Rada [13] determined sharp upper and lower bounds of the Randić index over the set of all digraphs in terms of the order, the set of all oriented trees in terms of the order, the set of all oriented paths and cycles in terms of the order, the set of all oriented hypercubes in terms of the dimension, respectively. Deng et al. [15] determined sharp upper and lower bounds of some VDB topological indices over the set of all digraphs in terms of the order, such as the harmonic index, the Atom-Bond-Connectivity index, the Geometric-Arithmetic index, the sum-connectivity index, the first and second Zagreb indices and the Randić index. The maximal values and minimal values of VDB topological indices over the set of all oriented trees in terms of the order and the set of all oriented graphs in terms of the order were obtained by Monsalve and Rada [12]. Soon after, the problem that is to determine a sharp upper bound of first Zagreb index over the set of all oriented unicyclic graphs in terms of the order and matching number was solved by Yang and Deng [14].

In this paper, we discuss orientations of cacti in terms of the order with maximum value of the zeroth-order general Randić index.

2 Some Lemmas

A lot of lemmas correlated with main result are given in the section. Let $\mathcal{G}(n, r)$ be the set of cacti of order n and r cycles. Let $G^0(n, r)$, $G^{(1)}(n, r)$, $G^{(2)}(n, r)$ be the graphs

depicted in Figure 1. $\tilde{\mathcal{G}}(n, r) = \{G^{(1)}(n, r), G^{(2)}(n, r)\}$. We denote by C_n the cycle which consists of n vertices, and $G^{(3)}(4, 1), G^{(4)}(4, 1)$ the sink-source orientations of C_4 . Let $\mathcal{G}^*(n, r) = \begin{cases} \tilde{\mathcal{G}}(n, r) \cup \{G^{(3)}(4, 1), G^{(4)}(4, 1)\}, & \text{if } a = 1 \text{ and } (n, r) = (4, 1) \\ \tilde{\mathcal{G}}(n, r), & \text{otherwise} \end{cases}$.




Figure 1: $G^0(n, r)$ and its two orientations

Lemma 2.1 [16] *If G is a connected bipartite graph, then G has two sink-source orientations.*

Lemma 2.2 *For any simple connected graph G , $D \in \mathcal{O}(G)$, $a \geq 1$, we have*

$$R_{a+1}^0(D) \leq \frac{R_{a+1}^0(G)}{2}$$

with equality if and only if D is a sink-source orientation of G .

Proof. $R_{a+1}^0(D) = \frac{1}{2} \sum_{uv \in A(D)} [(d_D^+(u))^a + (d_D^-(v))^a]$, $R_{a+1}^0(G) = \sum_{uv \in E(G)} [(d_G(u))^a + (d_G(v))^a]$.

And for any $u \in V(G)$, we have $d_G(u) \geq \max\{d_D^+(u), d_D^-(u)\}$. So,

$$R_{a+1}^0(D) = \frac{1}{2} \sum_{uv \in A(D)} [(d_D^+(u))^a + (d_D^-(v))^a] \leq \frac{1}{2} \sum_{uv \in E(G)} [(d_G(u))^a + (d_G(v))^a] = \frac{1}{2} R_{a+1}^0(G)$$

with equality if and only if $d_G(u) = \max\{d_D^+(u), d_D^-(u)\}$ for any $u \in V(G)$ which implies that D is a sink-source orientation of G . ■

Lemma 2.3 *Let $G \in \mathcal{G}(n, r)$, $n \geq 6$, $r \geq 2$, $\delta(G) \geq 2$. If there exists $u_0 u_1 \in E(G)$ such that $d_G(u_0) = d_G(u_1) = 2$, $\{u_2\} = N_G(u_0) \setminus \{u_1\}$ and $d_G(u_2) = d \geq 3$. $u_1 u_2 \notin E(G)$, $G' = G - u_0 + u_1 u_2$, $G'' = G - u_0$, $D \in \mathcal{O}(G)$, $D' \in \mathcal{O}(G')$, $D'' \in \mathcal{O}(G'')$, such that $A(D') \cap A(D) = A(D'')$, $a \geq 1$, then*

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2} [2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}]$$

with equality if and only if $u_1 u_0, u_2 u_0 \in A(D), u_1 u_2 \in A(D'), d_D^+(u_2) = d_G(u_2)$; or $u_0 u_1, u_0 u_2 \in A(D), u_2 u_1 \in A(D'), d_D^-(u_2) = d_G(u_2)$; or $u_0 u_1, u_2 u_0 \in A(D), u_1 u_2 \in A(D'), d_D^-(u_1) = d_G(u_1), d_D^+(u_2) = d_G(u_2)$; or $u_1 u_0, u_0 u_2 \in A(D), u_2 u_1 \in A(D'), d_D^+(u_1) = d_G(u_1), d_D^-(u_2) = d_G(u_2)$.

Proof. Let $N_G(u_1) \setminus \{u_0\} = \{w_1\}$, $N_G(u_2) \setminus \{u_0\} = \{v_1, v_2, \dots, v_{d_G(u_2)-1}\}$. It is straightforward to check that $d_D^+(w_1) = d_{D'}^+(w_1)$, $d_D^-(w_1) = d_{D'}^-(w_1)$; $d_D^+(v_i) = d_{D'}^+(v_i)$, $d_D^-(v_i) = d_{D'}^-(v_i)$ ($i = 1, 2, \dots, d_G(u_2) - 1$). Let $p_i = d_D^+(u_i)$, $i = 0, 1, 2$; $q_i = d_D^-(u_i)$, $i = 0, 1, 2$; $p'_i = d_{D'}^+(u_i)$, $i = 0, 1, 2$; $q'_i = d_{D'}^-(u_i)$, $i = 0, 1, 2$. We distinguish the following eight cases to prove the lemma.

Case 1. $u_0u_1, u_0u_2 \in A(D)$, $u_1u_2 \in A(D')$.

For $u_0u_1, u_0u_2 \in A(D)$, we have that $p_0 = p'_0 + 2 = 0 + 2 = 2$, $q_0 = q'_0 = 0$; For $u_0u_1 \in A(D)$, $u_1u_2 \in A(D')$, we have that $p_1 = p'_1 - 1$, $q_1 = q'_1 + 1$; For $u_0u_2 \in A(D)$, $u_1u_2 \in A(D')$, we have that $q_2 = q'_2$, $p_2 = p'_2$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[\varphi(p_0, q_1) + \varphi(p_0, q_2) - \varphi(p'_1, q'_2) + \sum_{i=1}^{q_1-1} [\varphi(q_1, d_D^+(w_i)) \\
&\quad - \varphi(q'_1, d_{D'}^+(w_i))] + \sum_{i=q_1}^{d_G(u_1)-1} [\varphi(p_1, d_D^-(w_i)) - \varphi(p'_1, d_{D'}^-(w_i))] \\
&\quad + \sum_{i=q_2}^{d_G(u_2)-1} [\varphi(p_2, d_D^-(v_i)) - \varphi(p'_2, d_{D'}^-(v_i))] + \sum_{i=1}^{q_2-1} [\varphi(q_2, d_D^+(v_i)) \\
&\quad - \varphi(q'_2, d_{D'}^+(v_i))] \\
&= \frac{1}{2}[(p_0)^a + (q_1)^a + (p_0)^a + (q_2)^a - (p_1 + 1)^a - (q_2)^a + (q_1 - 1) \\
&\quad [(q_1)^a - (q_1 - 1)^a] + (q_2 - 1)[(q_2)^a - (q_2)^a] \\
&\leq \frac{1}{2}[(p_0)^a + (d_G(u_1))^a + (p_0)^a - 1 + (d_G(u_1) - 1)[(d_G(u_1))^a \\
&\quad - (d_G(u_1) - 1)^a] \\
&\leq \frac{1}{2}[2^a + 2^a + 2^a - 1 + 2^a - 1] \\
&= 2^{a+1} - 1
\end{aligned}$$

Let $f(d) = \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}] - (2^{a+1} - 1) = \frac{1}{2}[d^{a+1} - (d-1)^{a+1} + 1 - 2^{a+1}]$, then $f'(d) = \frac{1}{2}[(a+1)d^a - (a+1)(d-1)^a] > 0$. Hence $\min_{d \geq 3} f(d) = f(3) = \frac{1}{2}[3^{a+1} - 2^{a+2} + 1] \geq \frac{1}{2}[9 \times 3^{a-1} - 8 \times 2^{a-1}] > 0$, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq 2^{a+1} - 1 < \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}].$$

Case 2. $u_0u_1, u_2u_0 \in A(D)$, $u_1u_2 \in A(D')$.

For $u_0u_1, u_2u_0 \in A(D)$, we have that $p_0 = p'_0 + 1 = 0 + 1 = 1$, $q_0 = q'_0 + 1 = 0 + 1 = 1$; For $u_0u_1 \in A(D)$, $u_1u_2 \in A(D')$, we have that $p_1 = p'_1 - 1$, $q_1 = q'_1 + 1$; For $u_2u_0 \in$

$A(D), u_1u_2 \in A(D')$, we have that $p_2 = p'_2 + 1, q_2 = q'_2 - 1$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_0)^a + (q_1)^a + (p_2)^a + (q_0)^a - (p_1 + 1)^a \\
&\quad - (q_2 + 1)^a + (q_1 - 1)[(q_1)^a - (q_1 - 1)^a] \\
&\quad + p_1[(p_1)^a - (p_1 + 1)^a] + (p_2 - 1)[(p_2)^a \\
&\quad - (p_2 - 1)^a] + q_2[(q_2)^a - (q_2 + 1)^a] \\
&\leq \frac{1}{2}[1 + (d_G(u_1))^a + (d_G(u_2))^a + 1 - 1 - 1 \\
&\quad + (d_G(u_1) - 1)[(d_G(u_1))^a - (d_G(u_1) - 1)^a] \\
&\quad + (d_G(u_2) - 1)[(d_G(u_2))^a - (d_G(u_2) - 1)^a]] \\
&= \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d - 1)^{a+1}]
\end{aligned}$$

with equality if and only if $q_1 = d_G(u_1), p_2 = d_G(u_2)$.

Case 3. $u_1u_0, u_0u_2 \in A(D), u_1u_2 \in A(D')$.

For $u_1u_0, u_0u_2 \in A(D)$, we have that $p_0 = p'_0 + 1 = 0 + 1 = 1, q_0 = q'_0 + 1 = 0 + 1 = 1$;
For $u_1u_0 \in A(D), u_1u_2 \in A(D')$, we have that $q_1 = q'_1, p_1 = p'_1$; For $u_0u_2 \in A(D), u_1u_2 \in A(D')$, we have that $p_2 = p'_2, q_2 = q'_2$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_1)^a + (q_0)^a + (p_0)^a + (q_2)^a - (p'_1)^a - (q'_2)^a] \\
&= \frac{1}{2}[(q_0)^a + (p_0)^a] \\
&= 1 < \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d - 1)^{a+1}]
\end{aligned}$$

Case 4. $u_1u_0, u_2u_0 \in A(D), u_1u_2 \in A(D')$.

For $u_1u_0, u_2u_0 \in A(D)$, we have that $p_0 = p'_0 = 0, q_0 = q'_0 + 2 = 0 + 2 = 2$;
For $u_1u_0 \in A(D), u_1u_2 \in A(D')$, we have that $p_1 = p'_1, q_1 = q'_1$; For $u_2u_0 \in A(D), u_1u_2 \in A(D')$, we have that $p_2 = p'_2 + 1, q_2 = q'_2 - 1$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_1)^a + (q_0)^a + (p_2)^a + (q_0)^a - (p_1)^a - (q_2 + 1)^a + (p_2 - 1)[(p_2)^a \\
&\quad - (p_2 - 1)^a] + q_2[(q_2)^a - (q_2 + 1)^a] \\
&\leq \frac{1}{2}[2^a + (d_G(u_2))^a + 2^a - 1 + (d_G(u_2) - 1)[(d_G(u_2))^a \\
&\quad - (d_G(u_2) - 1)^a] \\
&= \frac{1}{2}[2^a + d^a + 2^a - 1 + (d - 1)(d^a - (d - 1)^a)] \\
&= \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d - 1)^{a+1}]
\end{aligned}$$

with equality if and only if $p_2 = d_G(u_2)$.

Case 5. $u_0u_1, u_0u_2 \in A(D), u_2u_1 \in A(D')$.

Similarly to **Case 4**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d - 1)^{a+1}]$$

with equality if and only if $q_2 = d_G(u_2)$.

Case 6. $u_0u_1, u_2u_0 \in A(D)$, $u_2u_1 \in A(D')$.

Similarly to **Case 3**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}].$$

Case 7. $u_1u_0, u_0u_2 \in A(D)$, $u_2u_1 \in A(D')$.

Similarly to **Case 2**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}]$$

with equality if and only if $p_1 = d_G(u_1)$, $q_2 = d_G(u_2)$.

Case 8. $u_1u_0, u_2u_0 \in A(D)$, $u_2u_1 \in A(D')$.

Similarly to **Case 1**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}].$$

Consequently,

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} - 1 + d^{a+1} - (d-1)^{a+1}]$$

with equality if and only if $u_1u_0, u_2u_0 \in A(D)$, $u_1u_2 \in A(D')$, $d_D^+(u_2) = d_G(u_2)$; or $u_0u_1, u_0u_2 \in A(D)$, $u_2u_1 \in A(D')$, $d_D^-(u_2) = d_G(u_2)$; or $u_0u_1, u_2u_0 \in A(D)$, $u_1u_2 \in A(D')$, $d_D^-(u_1) = d_G(u_1)$, $d_D^+(u_2) = d_G(u_2)$; or $u_1u_0, u_0u_2 \in A(D)$, $u_2u_1 \in A(D')$, $d_D^+(u_1) = d_G(u_1)$, $d_D^-(u_2) = d_G(u_2)$. ■

Lemma 2.4 *Let $G \in \mathcal{G}(n, r)$ with $n \geq 6$ and $r \geq 2$, $\delta(G) \geq 2$. If there exists $u_0u_1 \in E(G)$ such that $d_G(u_0) = d_G(u_1) = 2$, $\{u_2\} = N_G(u_0) \setminus \{u_1\}$ and $d_G(u_2) = d \geq 3$. $u_1u_2 \in E(G)$, $G' = G - u_0 - u_1$, $D \in \mathcal{O}(G)$, $D' \in \mathcal{O}(G')$ such that $A(D') \cap A(D) = A(D')$, $a \geq 1$, then*

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}]$$

with equality if and only if $d_D^-(u_2) = d_G(u_2)$ or $d_D^+(u_2) = d_G(u_2)$.

Proof. Let $N_G(u_2) \setminus \{u_0, u_1\} = \{v_1, v_2, \dots, v_{d_G(u_2)-2}\}$. It is obvious that $d_D^+(v_i) = d_{D'}^+(v_i)$, $d_D^-(v_i) = d_{D'}^-(v_i)$ ($i = 1, 2, \dots, d_G(u_2) - 2$). Let $p_i = d_D^+(u_i)$, $i = 0, 1, 2$; $q_i = d_D^-(u_i)$, $i = 0, 1, 2$; $p'_i = d_{D'}^+(u_i)$, $i = 0, 1, 2$; $q'_i = d_{D'}^-(u_i)$, $i = 0, 1, 2$. According to u_0, u_1, u_2 , we have eight cases.

Case 1. $u_0u_1, u_0u_2 \in A(D)$, $u_1u_2 \in A(D)$.

For $u_0u_1, u_0u_2 \in A(D)$, we have that $p_0 = p'_0 + 2 = 2$, $q_0 = q'_0 = 0$; For $u_0u_1 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_1 = p'_1 + 1 = 0 + 1 = 1$, $q_1 = q'_1 + 1 = 0 + 1 = 1$; For $u_0u_2 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_2 = p'_2$, $q_2 = q'_2 + 2$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[\varphi(p_0, q_1) + \varphi(p_0, q_2) + \varphi(p_1, q_2) + \sum_{i=1}^{q_2-2} [\varphi(q_2, d_D^+(v_i)) \\
&\quad - \varphi(q_2', d_{D'}^+(v_i))] + \sum_{i=q_2-1}^{d_G(u_2)-2} [\varphi(p_2, d_D^-(v_i)) - \varphi(p_2', d_{D'}^-(v_i))] \\
&= \frac{1}{2}[(p_0)^a + (q_1)^a + (p_0)^a + (q_2)^a + (p_1)^a + (q_2)^a + (q_2 - 2) \\
&\quad [(q_2)^a - (q_2 - 2)^a]] \\
&\leq \frac{1}{2}[2^a + 1^a + 2^a + (d_G(u_2))^a + 1^a + (d_G(u_2))^a \\
&\quad + (d_G(u_2) - 2)[(d_G(u_2))^a - (d_G(u_2) - 2)^a]] \\
&= \frac{1}{2}[2^{a+1} + 2 + d^a + d^a + (d - 2)(d^a - (d - 2)^a)] \\
&= \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}]
\end{aligned}$$

with equality if and only if $q_2 = d_G(u_2)$.

Case 2. $u_0u_1, u_2u_0 \in A(D)$, $u_1u_2 \in A(D)$.

For $u_0u_1, u_2u_0 \in A(D)$, we have that $p_0 = p_0' + 1 = 0 + 1 = 1$, $q_0 = q_0' + 1 = 0 + 1 = 1$; For $u_0u_1 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_1 = p_1' + 1 = 0 + 1 = 1$, $q_1 = q_1' + 1 = 0 + 1 = 1$; For $u_2u_0 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_2 = p_2' + 1$, $q_2 = q_2' + 1$. Hence,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_0)^a + (q_1)^a + (p_2)^a + (q_0)^a + (p_1)^a + (q_2)^a + (q_2 - 1)[(q_2)^a \\
&\quad - (q_2')^a] + (p_2 - 1)[(p_2)^a - (p_2')^a] \\
&\leq \frac{1}{2}[4 + (q_2)^{a+1} - (q_2 - 1)^{a+1} + (p_2)^{a+1} - (p_2 - 1)^{a+1}]
\end{aligned}$$

Let $f(x) = \frac{1}{2}[4 + x^{a+1} - (x - 1)^{a+1} + (d - x)^{a+1} - (d - x - 1)^{a+1}]$ and $g(x) = (a + 1)x^a - (a + 1)(x - 1)^a$ ($1 \leq x \leq d - 1$), then $f'(x) = \frac{1}{2}[(a + 1)x^a - (a + 1)(x - 1)^a - [(a + 1)(d - x)^a - (a + 1)(d - x - 1)^a]] = \frac{1}{2}[g(x) - g(d - x)]$. By $g'(x) = a(a + 1)x^{a-1} - a(a + 1)(x - 1)^{a-1} \geq 0$, we have $f'(x) \geq 0$ for $x \geq d - x$, and $f'(x) \leq 0$ for $x \leq d - x$. It is obvious that $\max_{1 \leq x \leq d-1} f(x) = f(d - 1) = f(1) = \frac{1}{2}[4 + (d - 1)^{a+1} - (d - 2)^{a+1} + 1]$. Since

$$\begin{aligned}
&\frac{1}{2}[4 + (d - 1)^{a+1} - (d - 2)^{a+1} + 1] - \frac{1}{2}[d^{a+1} - (d - 2)^{a+1} + 2 + 2^{a+1}] \\
&= \frac{1}{2}[3 + (d - 1)^{a+1} - d^{a+1} - 2^{a+1}] \\
&< 0,
\end{aligned}$$

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}].$$

Case 3. $u_1u_0, u_0u_2 \in A(D)$, $u_1u_2 \in A(D)$.

For $u_1u_0, u_0u_2 \in A(D)$, we have that $p_0 = p_0' + 1 = 0 + 1 = 1$, $q_0 = q_0' + 1 = 0 + 1 = 1$; For $u_1u_0 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_1 = p_1' + 2 = 0 + 2 = 2$, $q_1 = q_1' = 0$; For

$u_0u_2 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_2 = p'_2$, $q_2 = q'_2 + 2$. Hence,

$$\begin{aligned} R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_1)^a + (q_0)^a + (p_0)^a + (q_2)^a + (p_1)^a + (q_2)^a + (q_2 - 2)[(q_2)^a \\ &\quad - (q_2 - 2)^a]] \\ &\leq \frac{1}{2}[2^a + 1^a + 1^a + (d_G(u_2))^a + 2^a + (d_G(u_2))^a + (d_G(u_2) - 2)[(d_G(u_2))^a \\ &\quad - (d_G(u_2) - 2)^a]] \\ &= \frac{1}{2}[2^a + 1 + 1 + d^a + 2^a + d^a + (d - 2)(d^a - (d - 2)^a)] \\ &= \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}] \end{aligned}$$

with equality if and only if $q_2 = d_G(u_2)$.

Case 4. $u_1u_0, u_2u_0 \in A(D)$, $u_1u_2 \in A(D)$.

For $u_1u_0, u_2u_0 \in A(D)$, we have that $p_0 = p'_0 = 0$, $q_0 = q'_0 + 2 = 0 + 2 = 2$; For $u_1u_0 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_1 = p'_1 + 2 = 0 + 2 = 2$, $q_1 = q'_1 = 0$; For $u_2u_0 \in A(D)$, $u_1u_2 \in A(D)$, we have that $p_2 = p'_2 + 1$, $q_2 = q'_2 + 1$. Hence, we have the inequality by the proof of **Case 2**.

$$\begin{aligned} R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2}[(p_1)^a + (q_0)^a + (q_0)^a + (p_2)^a + (p_1)^a + (q_2)^a + (q_2 - 1) \\ &\quad [(q_2)^a - (q_2 - 1)^a] + (p_2 - 1)[(p_2)^a - (p_2 - 1)^a]] \\ &= \frac{1}{2}[2^{a+2} + (q_2)^{a+1} - (q_2 - 1)^{a+1} + (p_2)^{a+1} - (p_2 - 1)^{a+1}] \\ &\leq \frac{1}{2}[2^{a+2} + (d - 1)^{a+1} - (d - 2)^{a+1} + 1] \end{aligned}$$

Let $h(d) = \frac{1}{2}[2^{a+2} + (d - 1)^{a+1} - (d - 2)^{a+1} + 1] - \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}] = \frac{1}{2}[2^{a+1} + (d - 1)^{a+1} - d^{a+1} - 1]$, we have $h'(d) = \frac{1}{2}[(a + 1)(d - 1)^a - (a + 1)d^a] < 0$, and $\max_{d \geq 3} h(d) = h(3) = \frac{1}{2}[2^{a+2} - 3^{a+1} - 1] = \frac{1}{2}[8 \times 2^{a-1} - 9 \times 3^{a-1} - 1] < 0$. Hence,

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}].$$

Case 5. $u_0u_1, u_0u_2 \in A(D)$, $u_2u_1 \in A(D)$.

Similarly to **Case 4**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}].$$

Case 6. $u_0u_1, u_2u_0 \in A(D)$, $u_2u_1 \in A(D)$.

Similarly to **Case 3**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d - 2)^{a+1}].$$

with equality if and only if $p_2 = d_G(u_2)$.

Case 7. $u_1u_0, u_0u_2 \in A(D)$, $u_2u_1 \in A(D)$.

Similarly to **Case 2**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') < \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}].$$

Case 8. $u_1u_0, u_2u_0 \in A(D), u_2u_1 \in A(D)$.

Similarly to **Case 1**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}]$$

with equality if and only if $p_2 = d_G(u_2)$.

Consequently,

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}]$$

with equality if and only if $d_D^-(u_2) = d_G(u_2)$ or $d_D^+(u_2) = d_G(u_2)$. ■

In the following, we determine the maximum zeroth-order general Randić index of trees, and the corresponding extreme graph.

Lemma 2.5 [4] *Let $G \in \mathcal{G}(n, 0)$ with $n \geq 2, a \geq 1$. Then*

$$R_{a+1}^0(G) \leq n - 1 + (n - 1)^{1+a}$$

with equality if and only if $G \cong G^0(n, 0)$.

From the maximum tree on the zeroth-order general Randić index and the corresponding extremum graph, we get the maximum oriented tree with respect to the zeroth-order general Randić index and the corresponding extremum digraph.

Lemma 2.6 *Let $G \in \mathcal{G}(n, 0)$ with $n \geq 2, D \in \mathcal{O}(G), a \geq 1$. Then*

$$R_{a+1}^0(D) \leq \frac{1}{2}[n - 1 + (n - 1)^{1+a}]$$

with equality if and only if $D \in \mathcal{G}^(n, 0)$.*

Proof. It is straightforward to check that G is a bipartite graph. By Lemma 2.1 and Lemma 2.2, $R_{a+1}^0(D) \leq \frac{1}{2}R_{a+1}^0(G)$ with equality if and only if D is a sink-source orientation of G .

We can obtain the following equality by Lemma 2.5,

$$\max\{R_{a+1}^0(D) | D \in \mathcal{O}(G), G \in \mathcal{G}(n, 0)\} = \max\{\frac{1}{2}R_{a+1}^0(G) | G \in \mathcal{G}(n, 0)\} = \frac{1}{2}R_{a+1}^0(G^0(n, 0))$$

Consequently,

$$R_{a+1}^0(D) \leq \frac{1}{2}[n - 1 + (n - 1)^{1+a}]$$

with equality if and only if $D \in \mathcal{G}^*(n, 0)$. ■

For a unicyclic graph of order 4 with a triangle, we have

Lemma 2.7 Let G_1 be the graph depicted in Figure 2, $D \in \mathcal{O}(G_1)$, $a \geq 1$. Then

$$R_{a+1}^0(D) \leq \frac{1}{2}[3^{a+1} + 2^{a+1} + 3]$$

with equality if and only if $D \in \{G^{(1)}(4, 1), G^{(2)}(4, 1)\}$.

Proof. It is straightforward to check that $\mathcal{O}(G_1) = \{D_1, D_2, \dots, D_6\}$, where D_1, D_2, \dots, D_6 are depicted in Figure 2. By direct computing, we have

$$R_{a+1}^0(D_5) = R_{a+1}^0(D_1) = \frac{1}{2}(3^{a+1} + 2^{a+1} + 3)$$

$$R_{a+1}^0(D_4) = R_{a+1}^0(D_2) = \frac{1}{2}(4 + 2^{a+2})$$

$$R_{a+1}^0(D_6) = R_{a+1}^0(D_3) = \frac{1}{2}(2 + 2^{a+2} + 2^{a+1}).$$

As the fact that

$$R_{a+1}^0(D_1) - R_{a+1}^0(D_2) = \frac{1}{2}(3^{a+1} + 2^{a+1} + 3) - \frac{1}{2}(4 + 2^{a+2}) = \frac{1}{2}(3^{a+1} - 2^{a+1} - 1) > 0,$$

$$R_{a+1}^0(D_1) - R_{a+1}^0(D_3) = \frac{1}{2}(3^{a+1} + 2^{a+1} + 3) - \frac{1}{2}(2 + 2^{a+2} + 2^{a+1}) = \frac{1}{2}(3^{a+1} - 2^{a+2} + 1) = \frac{1}{2}(9 \times 3^{a-1} - 8 \times 2^{a-1} + 1) > 0,$$

$$R_{a+1}^0(D) \leq \frac{1}{2}[3^{a+1} + 2^{a+1} + 3]$$

with equality if and only if $D \in \{D_1, D_5\} = \{G^{(1)}(4, 1), G^{(2)}(4, 1)\}$. ■




Figure 2: G_1 and its orientations in Lemma 2.7

Lemma 2.8 Let G be a simple connected graph, $d_G(u) = 1$ and $u \in V(G)$, $G' = G - u$, $D \in \mathcal{O}(G)$, $D' \in \mathcal{O}(G')$ such that $A(D') \cap A(D) = A(D')$, $v \in N_G(u)$, $a \geq 1$. Then

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2}[1 + (d_G(v))^{a+1} - (d_G(v) - 1)^{a+1}]$$

with equality if and only if $\max\{d_D^+(v), d_D^-(v)\} = d_G(v)$.

Proof. Let $N_G(v) = \{u = u_1, u_2, \dots, u_{d_G(v)}\}$ and $d_D^+(v) = p$, $d_D^-(v) = q$, $d_{D'}^+(v) = p'$, $d_{D'}^-(v) = q'$.

Case 1. $uv \in A(D)$

In this case, $q = q' + 1$, $p = p'$. It is straightforward to check that $d_D^+(u_i) = d_{D'}^+(u_i)$, $d_D^-(u_i) = d_{D'}^-(u_i)$, $i = 2, 3, \dots, d_G(v)$. Moreover,

$$\begin{aligned}
R_{a+1}^0(D) - R_{a+1}^0(D') &= \frac{1}{2} [R_{a+1}^0(d_D^+(u), q) + \sum_{i=2}^q [R_{a+1}^0(q, d_D^+(u_i)) - R_{a+1}^0(q', \\
&\quad , d_{D'}^+(u_i))] + \sum_{i=q+1}^{d_G(v)} [R_{a+1}^0(p, d_D^-(u_i)) - R_{a+1}^0(p', d_{D'}^-(u_i))] \\
&= \frac{1}{2} [(d_D^+(u))^a + (q)^a + (q-1)[(q)^a - (q-1)^a] \\
&\leq \frac{1}{2} [1 + (d_G(v))^a + (d_G(v) - 1)[(d_G(v))^a - (d_G(v) - 1)^a] \\
&= \frac{1}{2} [1 + (d_G(v))^{a+1} - (d_G(v) - 1)^{a+1}]
\end{aligned}$$

with equality if and only if $q = d_G(v)$.

Case 2. $vu \in A(D)$

Similarly to **Case 1**, we have

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2} [1 + (d_G(v))^{a+1} - (d_G(v) - 1)^{a+1}]$$

with equality if and only if $p = d_G(v)$.

Hence,

$$R_{a+1}^0(D) - R_{a+1}^0(D') \leq \frac{1}{2} [1 + (d_G(v))^{a+1} - (d_G(v) - 1)^{a+1}]$$

with equality if and only if $\max\{d_D^+(v), d_D^-(v)\} = d_G(v)$. ■

Next, we will get the oriented unicyclic graphs with the maximum of the zeroth-order general Randić index and the corresponding extreme graph, which will help to prove the main conclusions.

Lemma 2.9 *Let $G \in \mathcal{G}(n, 1)$, $n \geq 3$, $D \in \mathcal{O}(G)$, $a \geq 1$. Then*

$$R_{a+1}^0(D) \leq \frac{1}{2} [(n-1)^{a+1} + n - 1 + 2^{a+1}]$$

with equality if and only if $D \in \mathcal{G}^(n, 1)$.*

Proof. We proceed by applying induction on n .

If $n = 3$, then $G = C_3$, $D \in \mathcal{O}(C_3)$, it is straightforward to check that $R_{a+1}^0(D) \leq \frac{1}{2} [2^{a+2} + 2]$ with equality if and only if $D \in \{G^{(1)}(3, 1), G^{(2)}(3, 1)\} = \mathcal{G}^*(3, 1)$.

If $n = 4$, then $G = \{G_1, C_4\}$, where G_1 is shown in Figure 2. Let $f(x) = \frac{1}{2} [2^{x+1} + 3 + 3^{x+1}] - 2^{x+2}$. When $a \geq 2$, $f(a) = \frac{1}{2} [2^{a+1} + 3 + 3^{a+1}] - 2^{a+2} = \frac{1}{2} [3 + 3^{a+1} - 3 \times 2^{a+1}] = \frac{1}{2} [3 + 27 \times 3^{a-2} - 24 \times 2^{a-2}] > 0$; When $1 \leq a \leq 2$, $f'(a) = \frac{1}{2} [3^{a+1} \times \ln 3 - 3 \times 2^{a+1} \times \ln 2] > 0$ by MATLAB (see Figure 4 of the Appendix A). $f(a) \geq f(1) = 0$. From Lemma 2.2, if $D \in \mathcal{O}(C_4)$, we can get $R_{a+1}^0(D) \leq \frac{1}{2} R_{a+1}^0(C_4) = 2^{a+2} \leq \frac{1}{2} [2^{a+1} + 3 + 3^{a+1}]$ with equality if and only if $a = 1$ and D is a sink-source orientation of C_4 , which implies that

$D \in \{G^{(3)}(4, 1), G^{(4)}(4, 1)\}$. By Lemma 2.7, if $D \in \mathcal{O}(G_1)$, $M_1(D) \leq \frac{1}{2}[2^{a+1} + 3 + 3^{a+1}]$ with equality if and only if $D \in \{G^{(1)}(4, 1), G^{(2)}(4, 1)\}$. Thus the result holds for $n = 3, 4$.

If $n \geq 5$ and suppose that the lemma is true completely for all orientations of graphs in $\mathcal{G}(n-1, 1)$. Let $G \in \mathcal{G}(n, 1)$.

Case 1. $G = C_n$.

If $G = C_n$, $D \in \mathcal{O}(C_n)$, let $f(n) = \frac{1}{2}[2^{a+1} + n - 1 + (n-1)^{a+1}] - n \times 2^a$, then $f'(n) = \frac{1}{2}[1 + (a+1)(n-1)^a] - 2^a > 0$, and $f(n) \geq f(5) = \frac{1}{2}[2^{a+1} + 4 + 4^{a+1}] - 5 \times 2^a > 0$.

From Lemma 2.2, we can get that $R_{a+1}^0(D) \leq \frac{R_{a+1}^0(C_n)}{2} = n \times 2^a < \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2^{a+1}]$. The lemma holds.

Case 2. $G \neq C_n$.

Let $G' = G - \{u\}$, where $d_G(u) = 1$, then $G' \in \mathcal{G}(n-1, 1)$. Let $v \in V(G)$, $N_G(v) = \{u = u_1, u_2, \dots, u_s\}$, $1 \leq s = d_G(v) \leq n-1$, $D' \in \mathcal{O}(G')$ such that $A(D') \cap A(D) = A(D')$.

By the induction hypothesis,

$$R_{a+1}^0(D') \leq \frac{1}{2}[(n-2)^{a+1} + n - 2 + 2^{a+1}].$$

From Lemma 2.8, we can obtain that

$$\begin{aligned} R_{a+1}^0(D) &\leq R_{a+1}^0(D') + \frac{1}{2}[1 - (d_G(v) - 1)^{a+1} + (d_G(v))^{a+1}] \\ &\leq \frac{1}{2}[2^{a+1} + n - 2 + (n-2)^{a+1}] + \frac{1}{2}[1 + (n-1)^{a+1} - (n-2)^{a+1}] \\ &= \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2^{a+1}] \end{aligned}$$

with equality if and only if $R_{a+1}^0(D') = \frac{1}{2}[n - 2 + (n-2)^{a+1} + 2^{a+1}]$, and $\{d_D^+(u) = d_G(u), d_D^-(v) = d_G(v) = n-1\}$ or $\{d_D^+(v) = d_G(v) = n-1, d_D^-(u) = d_G(u)\}$, which implies that $D \in \mathcal{G}^*(n, 1)$.

The proof of the lemma is completed. ■

Lemma 2.10 *Let $D \in \mathcal{O}(G_2)$, $G_2 \in \mathcal{G}(5, 2)$, where G_2 is depicted in Figure 3, $a \geq 1$. Then*

$$R_{a+1}^0(D) \leq 2 \times 4^a + 2^{a+1} + 2$$

with equality if and only if $D \in \mathcal{G}^(5, 2)$.*

Proof. It is straightforward to check that $\mathcal{O}(G_2) = \{D_1, D_2, \dots, D_{10}\}$, where D_1, D_2, \dots, D_{10} are depicted in Figure 3. By direct computing, we obtain

$$\begin{aligned} R_{a+1}^0(D_1) &= \frac{1}{2}(8 + 2^{a+2}) \\ R_{a+1}^0(D_4) &= R_{a+1}^0(D_2) = \frac{1}{2}(7 + 3^{a+1} + 2^{a+1}) \\ R_{a+1}^0(D_9) &= R_{a+1}^0(D_3) = \frac{1}{2}(4 + 2^{a+3}) \\ R_{a+1}^0(D_7) &= R_{a+1}^0(D_5) = \frac{1}{2}(3 + 6 \times 2^a + 3^{a+1}) \\ R_{a+1}^0(D_6) &= \frac{1}{2}(12 \times 2^a) \\ R_{a+1}^0(D_{10}) &= R_{a+1}^0(D_8) = 2 \times 4^a + 2^{a+1} + 2. \end{aligned}$$




Figure 3: G_2 and its orientations

As the fact that

$$R_{a+1}^0(D_8) - R_{a+1}^0(D_1) = 2 \times 4^a + 2^{a+1} + 2 - \frac{1}{2}(8 + 2^{a+2}) = 2 \times 4^a - 2 > 0$$

$$R_{a+1}^0(D_8) - R_{a+1}^0(D_2) = 2 \times 4^a + 2^{a+1} + 2 - \frac{1}{2}(7 + 3^{a+1} + 2^{a+1}) = 2 \times 4^a - \frac{3}{2} \times 3^a + 2^a - 1.5 > 0$$

$$R_{a+1}^0(D_8) - R_{a+1}^0(D_3) = 2 \times 4^a + 2^{a+1} + 2 - \frac{1}{2}(4 + 2^{a+3}) = 2 \times 4^a - 2^{a+1} > 0$$

$$R_{a+1}^0(D_8) - R_{a+1}^0(D_5) = 2 \times 4^a + 2^{a+1} + 2 - \frac{1}{2}(3 + 6 \times 2^a + 3^{a+1}) = 2 \times 4^a - \frac{3}{2} \times 3^a - 2^a + \frac{1}{2} = 8 \times 4^{a-1} - \frac{9}{2} \times 3^{a-1} - 2 \times 2^{a-1} + \frac{1}{2} > 0$$

$$R_{a+1}^0(D_8) - R_{a+1}^0(D_6) = 2 \times 4^a + 2^{a+1} + 2 - \frac{1}{2}(12 \times 2^a) = (2^{a+1} + 2 - 6)2^a + 2 > 0$$

It is easy to get that

$$R_{a+1}^0(D) \leq 2 \times 4^a + 2^{a+1} + 2$$

with equality if and only if $D \in \{D_{10}, D_8\} = \mathcal{G}^*(5, 2)$. ■

3 Main Result

We will solve the problem which is to find the maximum zeroth-order general Randić index over the set of all oriented cacti in terms of the order in the section. The main result is as follows.

Theorem 3.1 *Let $G \in \mathcal{G}(n, r)$, $D \in \mathcal{O}(G)$, $a \geq 1$. Then*

$$R_{a+1}^0(D) \leq \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2r \times 2^a]$$

with equality if and only if $D \in \mathcal{G}^(n, r)$.*

Proof. We proceed by applying induction on n and r . By Lemma 2.6 and Lemma 2.9, the result holds for $r = 0$ and 1. If $r \geq 2$, then $n \geq 5$. If $n = 5$, $|\mathcal{G}(5, 2)| = 1$ and $G_2 \in \mathcal{G}(5, 2)$ (see Figure 3), then the result holds by Lemma 2.10.

If $n \geq 6$ and $r \geq 2$. We distinguish the following two cases according to $\delta(G)$.

Case 1. $\delta(G) = 1$.

Let $u, v \in V(G)$ and $d_G(u) = 1$, $v \in N_G(u)$, $G' = G - u$. Obviously, $G' \in \mathcal{G}(n-1, r)$, $d_G(v) \leq n-1$.

Let $D' \in \mathcal{O}(G')$ such that $A(D') \cap A(D) = A(D')$. By the induction hypothesis,

$$R_{a+1}^0(D') \leq \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a].$$

From Lemma 2.8, we have

$$\begin{aligned} R_{a+1}^0(D) &\leq R_{a+1}^0(D') + \frac{1}{2}[1 + (d_G(v))^{a+1} - (d_G(v) - 1)^{a+1}] \\ &\leq \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a + 1 + (n-1)^{a+1} - (n-2)^{a+1}] \\ &= \frac{1}{2}[(n-1)^{a+1} + n-1 + 2r \times 2^a] \end{aligned}$$

with equality if and only if $R_{a+1}^0(D') = \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a]$, $\max\{d_D^+(v), d_D^-(v)\} = d_G(v) = n-1$, which imply the fact that $D \in \mathcal{G}^*(n, r)$. The result holds.

Case 2. $\delta(G) \geq 2$.

It is obvious that there exists $u_0 u_1 \in E(G)$ in a terminal block of G such that $d_G(u_0) = d_G(u_1) = 2$, $N_G(u_0) = \{u_1, u_2\}$ and $d_G(u_2) = d \geq 3$. We will take two subcases into consideration.

Subcase 2.1. $u_1 u_2 \notin E(G)$.

Let $G' = G - u_0 + u_1 u_2$ and $G'' = G - u_0$. Obviously, $G' \in \mathcal{G}(n-1, r)$ and $d \leq n-2$. Let $D' \in \mathcal{O}(G')$, $D'' \in \mathcal{O}(G'')$ such that $A(D') \cap A(D) = A(D'')$. By the induction hypothesis, we can obtain the inequality that

$$R_{a+1}^0(D') \leq \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a].$$

From Lemma 2.3, we obtain

$$\begin{aligned} R_{a+1}^0(D) &\leq R_{a+1}^0(D') + \frac{1}{2}[2^{a+1} + d^{a+1} - (d-1)^{a+1} - 1] \\ &\leq \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a + 2^{a+1} + d^{a+1} - (d-1)^{a+1} - 1] \\ &\leq \frac{1}{2}[(n-2)^{a+1} + n-2 + 2r \times 2^a + 2^{a+1} + (n-2)^{a+1} - (n-3)^{a+1} - 1] \\ &= \frac{1}{2}[2(n-2)^{a+1} + n-3 + (r+1) \times 2^{a+1} - (n-3)^{a+1}] \end{aligned}$$

Let $f(n) = \frac{1}{2}[(n-1)^{a+1} + n-1 + 2r \times 2^a] - \frac{1}{2}[2(n-2)^{a+1} + n-3 + (r+1) \times 2^{a+1} - (n-3)^{a+1}]$ and $g(x) = x^a$ ($x > 0$), we have $g'(x) = ax^{a-1}$ and $g''(x) = a(a-1)x^{a-2} \geq 0$. Hence

$f'(n) = \frac{1}{2}[(a+1)(n-1)^a + 1] - \frac{1}{2}[2(a+1)(n-2)^a + 1 - (a+1)(n-3)^a] = \frac{1}{2}(a+1)[(n-1)^a - 2(n-2)^a + (n-3)^a] \geq 0$. It is obvious that $\min_{n \geq 6} f(n) = f(6)$.

If $a \geq 3$, then $f(6) = \frac{1}{2}[5^{a+1} + 5 + 2r \times 2^a] - \frac{1}{2}[2 \times 4^{a+1} + 3 + (r+1) \times 2^{a+1} - 3^{a+1}] = \frac{1}{2}[5^{a+1} + 2 - 2 \times 4^{a+1} + 3^{a+1} - 2^{a+1}] = \frac{1}{2}[625 \times 5^{a-3} + 2 - 512 \times 4^{a-3} + 3^{a+1} - 2^{a+1}] > 0$.

If $a = 1$, then $f(6) = \frac{1}{2}[5^{1+1} + 5 + 2r \times 2^1] - \frac{1}{2}[2 \times 4^{1+1} + 3 + (r+1) \times 2^{1+1} - 3^{1+1}] = 0$.

If $1 \leq a \leq 3$, let $h(a) = f(6)$, then $h'(a) = \frac{1}{2}[5^{a+1} \times \ln 5 - 2 \times 4^{a+1} \times \ln 4 + 3^{a+1} \times \ln 3 - 2^{a+1} \times \ln 2] > 0$ by direct computing of MATLAB (see Figure 5 of the Appendix A). Hence, $f(6) = h(a) \geq h(1) = 0$.

Consequently,

$$R_{a+1}^0(D) \leq \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2r \times 2^a]$$

with equality if and only if $a = 1$, $D' \in \mathcal{G}^*(n-1, r)$, $d = n-2$, $u_1 u_0, u_2 u_0 \in A(D)$, $u_1 u_2 \in A(D')$, $d_D^+(u_2) = d_G(u_2)$; or $u_0 u_1, u_0 u_2 \in A(D)$, $u_2 u_1 \in A(D')$, $d_D^-(u_2) = d_G(u_2)$; or $u_0 u_1, u_2 u_0 \in A(D)$, $u_1 u_2 \in A(D')$, $d_D^-(u_1) = d_G(u_1)$, $d_D^+(u_2) = d_G(u_2)$; or $u_1 u_0, u_0 u_2 \in A(D)$, $u_2 u_1 \in A(D')$, $d_D^+(u_1) = d_G(u_1)$, $d_D^-(u_2) = d_G(u_2)$. But if $D' \in \mathcal{G}^*(n-1, r)$, $u_1 u_2 \in A(D')$, we have $d_{D'}^+(u_2) = d_{G'}(u_2)$. It is a contradiction with $d_D^+(u_2) = d_G(u_2)$. If $D' \in \mathcal{G}^*(n-1, r)$, $u_2 u_1 \in A(D')$, we have $d_{D'}^+(u_2) = d_{G'}(u_2)$. It is a contradiction with $d_D^-(u_2) = d_G(u_2)$. Hence,

$$R_{a+1}^0(D) < \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2r \times 2^a].$$

Subcase 2.2. $u_1 u_2 \in E(G)$.

Let $G' = G - u_0 - u_1$. Obviously, $G' \in \mathcal{G}(n-2, r-1)$ and $d \leq n-1$. Let $D' \in \mathcal{O}(G')$ such that $A(D') \cap A(D) = A(D')$. By the induction hypothesis,

$$R_{a+1}^0(D') \leq \frac{1}{2}[(n-3)^{a+1} + n - 3 + 2(r-1) \times 2^a]$$

By Lemma 2.4, we can obtain

$$\begin{aligned} R_{a+1}^0(D) &\leq R_{a+1}^0(D') + \frac{1}{2}[2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}] \\ &\leq \frac{1}{2}[(n-3)^{a+1} + n - 3 + 2(r-1) \times 2^a + 2^{a+1} + 2 + d^{a+1} - (d-2)^{a+1}] \\ &\leq \frac{1}{2}[(n-3)^{a+1} + n - 3 + 2(r-1) \times 2^a + 2^{a+1} + 2 + (n-1)^{a+1} - (n-3)^{a+1}] \\ &= \frac{1}{2}[(n-1)^{a+1} + n - 1 + 2r \times 2^a] \end{aligned}$$

with equality if and only if $R_{a+1}^0(D') = \frac{1}{2}[(n-3)^{a+1} + n - 3 + 2(r-1) \times 2^a]$, $d = n-1$, or equivalently, $D' \in \mathcal{G}^*(n-2, r-1)$ and $\{d_D^-(u_2) = d_G(u_2) = n-1\}$ or $\{d_D^+(u_2) = d_G(u_2) = n-1\}$, i.e., $D \in \mathcal{G}^*(n, r)$. The result holds for n . \blacksquare

Acknowledgment. This work is supported by the Hunan Provincial Natural Science Foundation of China (2020JJ4423), the Department of Education of Hunan Province (19A318) and the National Natural Science Foundation of China (11971164).

Appendix A




Figure 4: $f1 = 2f'(a) = 3^{a+1} \times \ln 3 - 3 \times 2^{a+1} \times \ln 2 > 0$ ($1 \leq a \leq 2$)




Figure 5: $f1 = 2h'(a) = 5^{a+1} \times \ln 5 - 2 \times 4^{a+1} \times \ln 4 + 3^{a+1} \times \ln 3 - 2^{a+1} \times \ln 2 > 0$ ($1 \leq a \leq 3$)

References

- [1] M. V. Diudea, QSPR/QSAR Studies by Molecular Descriptors, Nova Science Publishers, New York, NY, USA, 2001.
- [2] A. T. Balaban, J. Devillers, Eds., Topological Indices and Related Descriptors in QSAR and QSPAR, CRC Press, Florida, FL, USA, 2014.
- [3] X. Li, J. Zheng, A unified approach to the extremal trees for different indices, MATCH Communications in Mathematical and in Computer Chemistry, 54 (2005) 195-208.
- [4] M. Liu, B. Liu, Some properties of the first general Zagreb index, Australasian Journal of Combinatorics, 47 (2010) 285.

- [5] J. M. Rodríguez, J. L. Sánchez, J. M. Sigarreta, CMMSE-on the first general Zagreb index, *Journal of Mathematical Chemistry*, 56 (2018) 1849-1864.
- [6] H. M. Awais, M. Javaid, A. Ali, First general Zagreb index of generalized F-sum graphs, *Discrete Dynamics in Nature and Society*, 2020 (2020) 1-16.
- [7] X. F. Pan, H. Liu, M. Liu, Sharp bounds on the zeroth-order general Randić index of unicyclic graphs with given diameter, *Applied Mathematics Letters*, 24 (2011) 687-691.
- [8] M. K. Jamil, I. Tomescu, Zeroth-order general Randić index of k-generalized quasi trees, (2018), <https://arxiv.org/abs/1801.03885>.
- [9] M. Azari, A. Iranmanesh, Generalized Zagreb index of graphs, *Studia Universitatis Babes-Bolyai Chemia*, 56 (2011) 59-70.
- [10] J. B. Liu, S. Javed, M. Javaid, K. Shabbir, Computing first general Zagreb index of operations on graphs, *IEEE access*, 7 (2019) 47494-47502.
- [11] L. Bedratyuk, O. Savenko, The star sequence and the general first Zagreb index, (2017), *MATCH Communications in Mathematical and in Computer Chemistry*, 79 (2018) 407-414.
- [12] J. Monsalve, J. Rada, Sharp upper and lower bounds of VDB topological indices of digraphs, *Symmetry*, 13 (2021) 1903.
- [13] J. Monsalve, J. Rada, Vertex-degree based topological indices of digraphs, *Discrete Applied Mathematics*, 295 (2021) 13-24.
- [14] J. Yang, H. Deng, maximum first Zagreb index of orientations of unicyclic graphs with given matching number, *Applied Mathematics and Computation*, 427 (2022) 127131.
- [15] H. Deng, Z. Tang, J. Yang, J. Yang, M. You, On the vertex-degree based invariants of digraphs, *Discrete Mathematics Letter*, 9 (2022) 2-9.
- [16] J. Monsalve, J. Rada, Oriented bipartite graphs with minimal trace norm, *Linear Multilinear Algebra*, 67 (2019) 1121-1131.