

Spherical tropicalization and Berkovich analytification

Desmond Coles

Abstract

Let X be a spherical variety. We show that Tevelev and Vogiannou's tropicalization map from X to its tropicalization factors through the Berkovich analytification X^{an} , as in the case for toric varieties. Furthermore we show that the tropicalization is a strong deformation retraction of X^{an} . We also give a strong deformation retraction of Thuillier's analytification X^{\square} onto a subspace described using the colored fan of X .

1 Introduction

Tropical geometry provides a set of tools for assigning combinatorial objects to algebraic varieties. One then studies the algebraic geometry of the original variety by looking at the combinatorics. One well known example is given by embedding a variety into a toric variety and then tropicalizing the toric variety [Pay09]. In [TV21] Tevelev and Vogiannou introduce tropicalization for spherical homogeneous spaces. Let k be an algebraically closed field. Recall that if G is a reductive group over k , and X is a normal G -variety, then X is **spherical** if there is a Borel subgroup of G with an open orbit in X . Spherical varieties can be considered the nonabelian analogue of toric varieties. Let $\overline{K} = k\{\{u\}\}$ be Puiseux series, the algebraic closure of Laurent series $K = k((u))$. Equip \overline{K} with the u -adic valuation $\|\cdot\|_u$. Given a spherical homogeneous space G/H , Tevelev and Vogiannou define a set theoretic tropicalization map $\text{trop}_G: G/H(\overline{K}) \rightarrow \mathcal{V}$, where \mathcal{V} is the set of $G(k)$ -invariant valuations on the function field $k(G/H)$. The set \mathcal{V} in fact has the structure of a finitely generated convex cone [Kno91, Corollary 5.3]. The map trop_G takes $x \in G/H(\overline{K})$ to the valuation $\|\cdot\|_x \in \mathcal{V}$ defined by the formula:

$$\|f\|_x = \|gf(x)\|_u \quad g \text{ in general position}$$

where general position means $g \in U_f$, $U_f \subseteq G(k)$ a nonempty Zariski open such that $\|gf(x)\|_u$ is constant for $g \in U_f$. For a subvariety $W \subseteq G/H$ one can then define $\text{Trop}_G(W)$ as the closure of $\text{trop}_G(W(\overline{K}))$. This construction agrees with the usual tropicalization map in the case when $G = T$, T a torus, and H trivial.

In [Nas17] Tevelev and Vogiannou's construction was extended to all spherical varieties, generalizing the extension of tropicalization of tori to toric varieties given in [Pay09]. Here the target of the tropicalization map, $\text{Trop}_G(X)$, consists of a gluing of the valuation cones associated to G -orbits in X . The tropicalization map is defined as a disjoint union of the maps given above. In [KM19] Kaveh and Manon develop tropical geometry for spherical varieties via Gröbner theory and it is shown that this agrees with the above notion.

Let X^{an} be the Berkovich analytification of X . In the case when X is a toric variety with dense torus T , $\text{Trop}_T(X)$ is canonically a subspace of X^{an} and the tropicalization map factors through a continuous retraction $X^{\text{an}} \rightarrow \text{Trop}_T(X)$. It was conjectured in [TV21] that this is true for tropicalizations of spherical homogeneous spaces. In [KM19] this is proved for affine spherical varieties. We show this is the case for all spherical varieties. Let G^{\square} denote the k -analytic space associated to G as defined in [Thu07, Section 1]. Let $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$ be the map given by $*$ -multiplication by the unique point in the Shilov boundary of G^{\square} .

Theorem A. *Let X be a spherical G -variety over an algebraically closed and trivially valued field k , and let \overline{K} denote Puiseux series in k . Then there is a canonical homeomorphism $\iota: \text{Trop}_G(X) \rightarrow \mathbf{p}(X^{\text{an}})$, and $\mathbf{p}: X^{\text{an}} \rightarrow \text{Trop}_G(X)$ is a retraction of topological spaces such that the following diagram commutes:*

$$\begin{array}{ccc} X(\overline{K}) & \longrightarrow & X^{\text{an}} \\ & \searrow \text{trop}_G & \downarrow \mathbf{p} \\ & & \text{Trop}_G(X). \end{array}$$

Corollary. For W a subvariety of X , there is a strong deformation retraction $H : [0, 1] \times W^{an} \rightarrow \text{Trop}_G(W)$ such that $H(1, p) = \mathbf{p}(p)$.

The construction of the map \mathbf{p} was inspired by the work of Thuillier in the toric case, [Thu07, Section 2]. The space G^\square is a compact analytic subdomain of the k -analytic space G^{an} , the Shilov boundary of this space is a point with important similarities to the generic point of G . The notion of $*$ -multiplication is an idea originally due to Berkovich which generalizes left-multiplication by points in $G^{an}(k)$.

Our work in Section 3 extends the notion of tropicalization to other nontrivially valued extensions of k besides \overline{K} . We show that for any field extension L/k with valuation $\|\cdot\|$ there is a well-defined tropicalization map $\text{trop}_{G,L} : X(L) \rightarrow \text{Trop}_G(X)$. For $x \in X(L)$, $\text{trop}_G(x) = \|\cdot\|_x$ where $\|\cdot\|_x$ is the valuation $\|\cdot\|_x \in \mathcal{V}$ given by the formula:

$$\|f\|_x = \|gf(x)\| \quad g \text{ in general position}$$

where general position means $g \in U_f$, $U_f \subseteq G(k)$ is a nonempty Zariski open such that $\|gf(x)\|$ is constant for $g \in U_f$. It follows from Theorem A that tropicalization is invariant under the choice of L .

Corollary. Let W be a subvariety of X , and L be a nontrivially valued and algebraically closed extension of k . Then the closure of $\text{trop}_{G,L}(W(L))$ in $\text{Trop}_G(X)$ is equal to $\mathbf{p}(W^{an})$, where W^{an} is considered as a subspace of X^{an} induced by the inclusion $W \subseteq X$.

Our second main result is on the restriction of the map \mathbf{p} to the subspace X^\square . This result generalizes Proposition 2.12 of [Thu07] on the toric case.

Theorem B. There is a strong deformation retraction from X^\square onto $\mathbf{p}(X^\square)$. If the colored fan $\mathfrak{F}(X)$ satisfies:

$$\bigcup_{(\sigma, \mathcal{F}) \in \mathfrak{F}(X)} \sigma \subseteq \mathcal{V}. \quad (\star)$$

Then $\mathbf{p}(X^\square)$ is homeomorphic to the canonical compactification of the fan $F(X) = \{(\sigma, \mathcal{F}) \in \mathfrak{F}(X)\}$.

Recall that just as toric varieties are classified by fans, spherical varieties are classified by colored fans. For each spherical homogeneous space G/H there is an associated vector space $N_{\mathbb{R}}$. A colored fan, defined precisely in Section 4, is a finite collection of pairs (σ, \mathcal{F}) where σ is a strictly convex cone in $N_{\mathbb{R}}$ and \mathcal{F} is a finite set. The **canonical compactification** of a cone σ is defined as $\text{hom}(\sigma^\vee, [0, \infty])$, where σ^\vee consists of positive linear functionals on σ and the homomorphisms are semigroup homomorphisms. The canonical compactification of a fan is the polyhedral complex given by gluing the canonical compactifications of the cones along shared faces. At the end of Section 6 we give a description of $\mathbf{p}(X^\square)$ in the case where \star does not hold.

Outline of paper. Section 2 introduces the necessary notions from Berkovich geometry. Section 3 introduces $*$ -multiplication and studies the map \mathbf{p} . Section 4 recalls the Luna-Vust classification of G/H -embeddings, and contains important notation for Sections 5 and 6. Section 5 recalls the work of [TV21] and [Nas17] and gives a proof of Theorem A. Section 6 recalls the work of [Thu07, Section 2] and gives a proof of Theorem B.

Convention Choice. Though somewhat nonstandard in the literature of spherical varieties we opt to use multiplicative notation for valuations to accommodate the conventions of Berkovich geometry. Our exact conventions are as follows. Let A be a ring. By a **seminorm** on A we mean a function $\|\cdot\|_A : A \rightarrow \mathbb{R}_{\geq 0}$ such that $\|1\|_A = 1$, $\|0\|_A = 0$, $\|fg\|_A \leq \|f\|_A \|g\|_A$, and $\|f + g\|_A \leq \max(\|f\|_A, \|g\|_A)$.¹ We say that $\|\cdot\|_A$ is a **norm** if $\|f\|_A \neq 0$ for all $f \neq 0$. A (semi)norm will be called a **(semi)valuation** if it is multiplicative: $\|fg\|_A = \|f\|_A \|g\|_A$.² If k is a field with valuation $\|\cdot\|_k$ and A is a k -algebra then a seminorm on A will always be assumed to extend the norm on k ; $\|\lambda f\|_A = \|\lambda\|_k \|f\|_A$ for all $\lambda \in k$. By a valued extension L/k we mean a field extension where L is equipped with a valuation (extending the valuation on k by our previous convention).

¹These are non-Archimedean seminorms, which are the only ones of interest because our base field will always be non-Archimedean.

²More commonly valuations given as functions ν into the additive semigroup $\mathbb{R} \sqcup \{\infty\}$, a multiplicative norm is equivalent to such a function ν , set $\nu = -\log(\|\cdot\|)$.

Acknowledgments. The author would like to thank their advisor, Sam Payne, for their support and guidance during this research. The paper also benefited greatly from conversations with Tom Gannon, Sam Raskin, Yixian Wu, Hernán Iriarte, Amy Li, and Logan White.

2 Overview of Berkovich geometry

The contents of this Section are an overview of the necessary notions in the theory of Berkovich spaces, in particular we need to work with analytifications of varieties and \square -spaces. Berkovich spaces provide a framework for doing analytic geometry over valued fields. Given a base field k with valuation, Berkovich spaces are called k -analytic spaces. These are locally ringed spaces equipped with a k -analytic atlas. We will not go into full detail of the theory as we are primarily concerned with the topology of these spaces, and will only introduce the notions we need. The content in this Section is laid out in detail in [Ber90], with the exception of \square -spaces for this we refer the reader to [Thu07].

Throughout this Section let k be a valued field, and let X be a separated locally finite type k -scheme.

2.1 Berkovich analytification

Definition 2.1. The **Berkovich analytification** of X , denoted X^{an} , has points given by pairs $(x, \|\cdot\|)$ where x is a scheme point of X and $\|\cdot\|$ is a valuation on $k(x)$. There is a map $\rho: X^{\text{an}} \rightarrow X$ given by $(x, \|\cdot\|) \mapsto x$. The topology on X^{an} is the coarsest such that ρ is continuous, and for any open $U \subseteq X$ and any $f \in \mathcal{O}_X(U)$, the map $\rho^{-1}(U) \rightarrow \mathbb{R}$ given by $(x, \|\cdot\|) \mapsto \|f\|$ is continuous.

Any valued extension L/k and map of schemes $x: \text{Spec } L \rightarrow X$ defines a point in X^{an} . The point is given by $(x, \|\cdot\|)$ where x is the point corresponding to the image of $\text{Spec } L$ and $\|\cdot\|$ is the valuation on L restricted to $k(x)$. There is also a sheaf of analytic functions on X^{an} , however we only need a few details. In particular for a point $p = (x, \|\cdot\|) \in X^{\text{an}}$ let $\mathcal{H}(p)$ be the *completed residue field*, it is the metric completion of $k(x)$ with respect to $\|\cdot\|$.

When $X = \text{Spec } A$, A a k -algebra, we can alternatively describe X^{an} as being the semivaluations on A . The topology on X^{an} in this case will be the coarsest such that evaluation at an element of A is continuous. One can see that this description of X^{an} is equivalent to the original as follows. Let the **kernel** of a semivaluation $\|\cdot\|$ on A , be $\ker(\|\cdot\|) := \{f \in A \mid \|f\| = 0\}$. Then $\|\cdot\|$ defines a valuation on the fraction field of $A/\ker(\|\cdot\|)$, which we also denote $\|\cdot\|$. This valuation is characterized by $\|f + \ker(\|\cdot\|)\| := \|f\|$.

Berkovich analytification is functorial. For any morphism of k -schemes $f: X \rightarrow Y$ there is a morphism of Berkovich spaces over k , $f^{\text{an}}: X^{\text{an}} \rightarrow Y^{\text{an}}$. In particular for a point $(x, \|\cdot\|) \in X^{\text{an}}$ one has that $f^{\text{an}}((x, \|\cdot\|)) = (f(x), \|f_x(\cdot)\|)$, where $f_x: k(f(x)) \rightarrow k(x)$ is the induced map on function fields.

2.2 The Berkovich spectrum and \square -spaces

We need to introduce one other type of k -analytic space. Let \mathcal{A} be a Banach k -algebra, i.e. \mathcal{A} is a k -algebra equipped with a norm $\|\cdot\|_{\mathcal{A}}$, and \mathcal{A} is complete with respect to this norm.

Definition 2.2. The **Berkovich spectrum**, $\mathcal{M}(\mathcal{A})$, is the set of bounded semivaluations on \mathcal{A} , i.e. semivaluations $\|\cdot\|$ such that where $\|\cdot\| \leq \|\cdot\|_{\mathcal{A}}$. The topology is the coarsest such that evaluation at an element of \mathcal{A} is continuous.

Example 2.1. The one-point spaces in the category of Berkovich spaces are given by $\mathcal{M}(L)$ where L is a complete field.

Let \mathcal{A} and \mathcal{B} be Banach k -algebras with norms $\|\cdot\|_{\mathcal{A}}$ and $\|\cdot\|_{\mathcal{B}}$. One defines the completed tensor product, $\mathcal{A} \hat{\otimes}_k \mathcal{B}$ to be the completion of $\mathcal{A} \otimes_k \mathcal{B}$, with respect to the norm:

$$\|f\|_{\mathcal{A}, \mathcal{B}} := \inf \left\{ \max_i \|g_i\|_{\mathcal{A}} \|f_i\|_{\mathcal{B}} \mid \sum_i f_i \otimes g_i \in \mathcal{A} \otimes_k \mathcal{B} \text{ such that } f = \sum_i f_i \otimes g_i \right\}.$$

The function $\|\cdot\|_{\mathcal{A} \otimes_k \mathcal{B}}$ is clearly a seminorm, that it is a norm and that $\mathcal{A} \otimes_k \mathcal{B}$ embeds isometrically into its completion follows from part 4 of Theorem 1 in [Gru66, Section 3].

Proposition 2.1. [Ber90, Chapter 3] The norm $\|\cdot\|_{\mathcal{A},\mathcal{B}}$ makes $\mathcal{M}(\mathcal{A}\hat{\otimes}_k\mathcal{B})$ the product $\mathcal{M}(\mathcal{A})\times\mathcal{M}(\mathcal{B})$ in the category of k -analytic spaces.

Notation. To avoid cluttered subscripts, when considering p in some analytic space will choose to write $\|\cdot\|_p$ for the norm on $\mathcal{H}(p)$ instead of $\|\cdot\|_{\mathcal{H}(p)}$. Similarly for a pair of points p and q we will write $\|\cdot\|_{p,q}$ for the norm on $\mathcal{H}(p)\hat{\otimes}_k\mathcal{H}(q)$ instead of $\|\cdot\|_{\mathcal{H}(p),\mathcal{H}(q)}$.

Consider when k is trivially valued. Let A be a finitely generated k -algebra, A equipped with the trivial valuation is complete. In this case we can identify $\mathcal{M}(A)$ with seminorms $\|\cdot\|$, that are bounded by 1. Recall that if X is a variety over k , L/k is a valued field extension, and R is the valuation ring of L , then the map of schemes $\text{Spec } L \rightarrow X$ **has center** if it factors through $\text{Spec } R \rightarrow X$. We say **the center** is the image of the closed point of $\text{Spec } R$. Furthermore if $X = \text{Spec } A$ and A is an integral domain, then a valuation $\|\cdot\|$ has center in X if and only if $\|\cdot\|$ is bounded by 1 on A . From this it follows that for any trivially valued finitely generated k -algebra A , $\mathcal{M}(A)$ can be taken to be pairs $(x, \|\cdot\|)$, x a scheme point of A , and $\|\cdot\|$ a valuation on $k(x)$ having center in $\text{Spec } A$. For a general locally finite type scheme X over a trivially valued field k , we define X^\square to be given by pairs of points $(x, \|\cdot\|)$ where $\|\cdot\|$ is a valuation on $k(x)$, and $\text{Spec } k(x) \rightarrow X$ has center in X . So $\mathcal{M}(A) = (\text{Spec } A)^\square$.

Proposition 2.2. [Thu07] The space X^\square is a compact analytic subdomain of X^{an} , and they are equal when X is proper.

2.3 The reduction map

Let $\mathcal{A}^\circ := \{f \in \mathcal{A} \mid \|f\|_{\mathcal{A}} \leq 1\}$ and $\mathcal{A}^{\circ\circ} := \{f \in \mathcal{A} \mid \|f\|_{\mathcal{A}} < 1\}$. Then the **reduction** of \mathcal{A} is said to be $\tilde{\mathcal{A}} := \mathcal{A}^\circ/\mathcal{A}^{\circ\circ}$. Any map of Banach k -algebras $\mathcal{A} \rightarrow \mathcal{B}$ induces a map of rings $\tilde{\mathcal{A}} \rightarrow \tilde{\mathcal{B}}$. In particular for $p \in \mathcal{M}(\mathcal{A})$ there is a map of rings $h_p: \tilde{\mathcal{A}} \rightarrow \widetilde{\mathcal{H}(p)}$. Then there is a map of sets called the **reduction map**, $r: \mathcal{M}(\mathcal{A}) \rightarrow \text{Spec}(\tilde{\mathcal{A}})$ defined by $p \mapsto \ker(h_p)$.

When A is a finitely generated algebra with the trivial norm we have $A^\circ = A$ and $A^{\circ\circ} = 0$, so reduction gives a map of sets $r: \mathcal{M}(A) \rightarrow \text{Spec } A$. More generally we have a map $r: X^\square \rightarrow X$. When X^\square is viewed as pairs $(x, \|\cdot\|)$, reduction maps $(x, \|\cdot\|)$ to its center. This map is anticontinuous [Ber90, Corollary 2.4.2], and surjective [Ber90, Proposition 2.4.4].

2.4 The Shilov boundary

A **boundary** of $\mathcal{M}(\mathcal{A})$ is a closed subset of $\mathcal{M}(\mathcal{A})$, such that for every $f \in \mathcal{A}$ the function $\text{ev}_f: \mathcal{M}(\mathcal{A}) \rightarrow \mathbb{R}$ given by $\|\cdot\| \mapsto \|f\|$, attains its maximum. A unique minimal boundary is said to be the **Shilov boundary**.

Proposition 2.3. [Ber90, Proposition 2.4.4] The reduction map $r: \mathcal{M}(\mathcal{A}) \rightarrow \text{Spec}(\tilde{\mathcal{A}})$ gives a bijection between the Shilov boundary of $\mathcal{M}(\mathcal{A})$ and the generic points of the components of $\text{Spec}(\tilde{\mathcal{A}})$.

3 The retraction map

In this Section we assume k is a trivially valued and algebraically closed field, G is a connected linear algebraic group over k , and X is a G -variety. Let m to denote the action map $m: G \times X \rightarrow X$. The objective is to introduce $*$ -multiplication, use it construct a retraction of topological spaces $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$, and give a description of the image of \mathbf{p} . Furthermore we show that when G is reductive we can construct a strong deformation from X^{an} onto $\mathbf{p}(X^{\text{an}})$ using $*$ -multiplication.

The notion of $*$ -multiplication is taken directly from [Ber90, Chapter 5]. Properties specific to $*$ -multiplication by the unique point in the Shilov boundary of G^\square are directly inspired by the results on tori from [Thu07, Section 2].

3.1 *-multiplication

Definition 3.1. We say that a commutative Banach k -algebra \mathcal{A} is **peaked** if for any valued extension L/k the norm on $L \hat{\otimes}_k \mathcal{A}$ is a valuation. We say a point p of a k -analytic space is peaked if the completed residue field $\mathcal{H}(p)$ is peaked.

Proposition 3.1. All the the points of G^{an} are peaked.

Proof. This is a particular case of [Poi13, Corollaire 3.14]. \square

Definition 3.2. Let $g \in G^{\text{an}}$, because g is peaked the norm $\|\cdot\|_{g,p}$ is a valuation and is thus an element of $\mathcal{M}(\mathcal{H}(g)) \times \mathcal{M}(\mathcal{H}(p))$. There is a map of sets $X^{\text{an}} \rightarrow X^{\text{an}}$ given by $p \mapsto g * p$; where $g * p$ is the image of $\|\cdot\|_{g,p}$ under the composition of maps of k -analytic spaces:

$$\mathcal{M}(\mathcal{H}(g)) \times \mathcal{M}(\mathcal{H}(p)) \hookrightarrow G^{\text{an}} \times X^{\text{an}} \xrightarrow{m^{\text{an}}} X^{\text{an}}.$$

This is called ***-multiplication by g** .

Example 3.1. Let $g \in G^{\text{an}}(k) = G(k)$. The map $X^{\text{an}} \rightarrow X^{\text{an}}$ given by $p \mapsto g * p$ is equal to the map induced by left multiplication by g .

Below are some general facts about *-multiplication.

Proposition 3.2. [Ber90, Proposition 5.2.8] Let G and H be algebraic groups acting on varieties X and Y , respectively. Let $\phi: G \rightarrow H$ be a morphism of algebraic groups, and $\psi: X \rightarrow Y$ a morphism of varieties.

1. The map $X^{\text{an}} \rightarrow X^{\text{an}}$ given by *-multiplication by a point $g \in G^{\text{an}}$ is continuous.
2. If g and h are points of G^{an} and H^{an} , respectively, and p a point of X^{an} , then $g * (h * p) = (g * h) * p$
3. Given a commutative diagram:

$$\begin{array}{ccc} G^{\text{an}} \times X^{\text{an}} & \longrightarrow & X^{\text{an}} \\ \downarrow \phi^{\text{an}} \times \psi^{\text{an}} & & \downarrow \psi^{\text{an}} \\ H^{\text{an}} \times Y^{\text{an}} & \longrightarrow & Y^{\text{an}} \end{array}$$

one has that $\psi^{\text{an}}(g * p) = \phi^{\text{an}}(g) * \psi^{\text{an}}(p)$.

Let $\mathbf{g} \in G^{\text{an}}$ be the point given by the trivial valuation. This is the unique point in the Shilov boundary of G^{an} . Let $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$ be given by $p \mapsto \mathbf{g} * p$. By Proposition 3.2 part 1, \mathbf{p} is a continuous map of topological spaces. The remainder of this section will be devoted to studying this map.

Remark 3.1. We can alternatively describe \mathbf{p} in the following way. We have that $\mathbf{p}(p)$ is the image of the unique point in the Shilov boundary of $G^{\text{an}} \times \mathcal{M}(\mathcal{H}(p))$ under the map of k -analytic spaces:

$$G^{\text{an}} \times \mathcal{M}(\mathcal{H}(p)) \hookrightarrow G^{\text{an}} \times X^{\text{an}} \rightarrow X^{\text{an}}.$$

So when $G = T$ is a torus, \mathbf{p} is the retraction map defined in [Thu07, Section 2].

Example 3.2. Let $X = T = \text{Spec } k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ be the n -dimensional torus, and equip $k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ with the trivial valuation. Let $\lambda \in [0, 1]$, and let \mathbf{g}_λ be the point of T^{an} corresponding to the semivaluation:

$$\left\| \sum_I a_I t^I \right\| = \max_I \lambda^{|I|}$$

where I denotes a multi-index and if $I = (I_1, \dots, I_n)$ then $|I| = I_1 + \dots + I_n$. Then one can see via computation that for any valued extension L , the norm $\|\cdot\|_{\mathbf{g}_\lambda, L}$ on $\mathcal{H}(\mathbf{g}_\lambda) \hat{\otimes}_k L$ is multiplicative. For $\lambda = 0$ this is because $\mathcal{H}(\mathbf{g}_\lambda) = k$. For $\lambda \in (0, 1]$ \mathbf{g}_λ defines a norm on $k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. For $f = \sum_I a_I t^I \in k[t_1^{\pm 1}, \dots, t_n^{\pm 1}] \otimes_k L = L[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ we compute that:

$$\|f\|_{\mathbf{g}_\lambda, L} = \max_I \|a_I\|_L \lambda^{|I|}$$

and this is a valuation.

Let $p \in T^{\text{an}}$ be given by $x = (x_1, \dots, x_n) \in T(L)$, where L is a valued extension of k , so $p = (x, \|\cdot\|_L)$. Let m_x the composition $T \times \text{Spec } k(x) \hookrightarrow T \times T \xrightarrow{m} T$. Then for $f = \sum_I a_I t^I \in \text{Spec } k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ seminorm $\mathbf{g}_\lambda * p$ is given by:

$$\|f\|_{\mathbf{g}_\lambda * p} = \|m_x^* f\|_{\mathbf{g}_\lambda, p} = \left\| \sum_I a_I t^I \otimes x^I \right\|_{\mathbf{g}_\lambda, p} = \max_I \|a_I x^I\|_L \lambda^{|I|}.$$

Corollary 3.1. The map: $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$ is idempotent, i.e. $\mathbf{g} * \mathbf{g} = \mathbf{g}$.

Proof. One has that $\mathcal{H}(\mathbf{g}) = k(G)$ and the norm on $k(G)$ defined by \mathbf{g} is the trivial valuation so we have that for $f \in \mathcal{H}(\mathbf{g}) \otimes_k \mathcal{H}(\mathbf{g})$:

$$\|f\|_{\mathbf{g}, \mathbf{g}} = \inf \left\{ \max_i \|g_i\|_{\mathbf{g}} \|f_i\|_{\mathbf{g}} \mid f = \sum_i g_i \otimes f_i \right\}$$

which is 0 or 1, and 0 if and only if $f = 0$. Therefore it is the trivial norm. so $\mathbf{g} * \mathbf{g} = \mathbf{g}$. So by Proposition 3.2 \mathbf{p} is idempotent. \square

Corollary 3.2. The map $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$ is a retraction of topological spaces.

Proposition 3.3. Let $p = (x, \|\cdot\|) \in X^{\text{an}}$, then $\mathbf{p}(p) = (\eta_Y, \overline{\|\cdot\|})$. Where η_Y is the generic point of Y , the Zariski closure of the image of the composition:

$$G \times \text{Spec}(k(x)) \hookrightarrow G \times X \xrightarrow{m} X$$

and $\overline{\|\cdot\|}$ is a $G(k)$ -invariant valuation on $k(Y)$.

The proof is similar to the proofs of [Thu07, Proposition 2.5 (2)] and [Thu07, Proposition 2.3 (ii)], which prove the above for $G = T$ a torus.

Proof. Recall that there is a map $\rho: X^{\text{an}} \rightarrow X$ given by $(x, \|\cdot\|) \mapsto x$. To see that $\rho(\mathbf{p}(p)) = \eta_Y$ notice that there is a commutative diagram:

$$\begin{array}{ccc} \mathcal{M}(\mathcal{H}(\mathbf{g})) \times \mathcal{M}(\mathcal{H}(p)) & \xrightarrow{\rho} & \text{Spec } k(\eta_G) \times \text{Spec } k(x) \\ \downarrow & & \downarrow \\ G^{\text{an}} \times X^{\text{an}} & \xrightarrow{\rho} & G \times X \\ \downarrow m^{\text{an}} & & \downarrow m \\ X^{\text{an}} & \xrightarrow{\rho} & X \end{array}$$

where η_G is the generic point of G . In this diagram the image of the generic point of $\text{Spec } k(\eta_G) \times \text{Spec } k(x)$ in X is η_Y . But then because $\rho(\|\cdot\|_{\mathbf{g}, p})$ is the generic point of $\text{Spec } k(\eta_G) \times \text{Spec } k(x)$ we have the first claim.

To see that $\overline{\|\cdot\|}$ is $G(k)$ -invariant notice that for any $g \in G(k)$, $g * \mathbf{g} = \mathbf{g}$. Thus by Proposition 3.2 part 2 we have that $g * (\mathbf{g} * p) = \mathbf{g} * p$. But $p \mapsto g * p$ is the map induced by left multiplication by g , which for points of the form $(\eta_Y, \|\cdot\|)$ is given by precomposing $\|\cdot\|$ with the left-multiplication by g . \square

Furthermore we have the following.

Proposition 3.4. The image of \mathbf{p} is exactly the set of points $(\eta_Y, \|\cdot\|)$, where Y is a G -invariant subvariety of X and $\|\cdot\|$ is $G(k)$ -invariant.

This proof is the same as the proof of [Thu07, Proposition 2.3 (ii)], which proves this fact for $G = T$ a torus. We include the argument, except for one lemma, for completeness.

Proof. By Proposition 3.3 we have one containment. For the other notice that points of the form $(\eta_Y, \|\cdot\|)$, where Y is a G -invariant subvariety and $\|\cdot\|$ is $G(k)$ -invariant, are exactly the points $(x, \|\cdot\|) \in X^{\text{an}}$ fixed by the action of $G(k)$. Let $p = (x, \|\cdot\|) \in X^{\text{an}}$ be a point fixed by the group action $G(k)$. The morphism:

$$G^{\square} \times \mathcal{M}(\mathcal{H}(p)) \hookrightarrow G^{\square} \times X^{\text{an}} \xrightarrow{m^{\text{an}}} X^{\text{an}}$$

maps the subset $G^{\square}(k) \times \mathcal{M}(\mathcal{H}(p))$ onto x . The map is continuous and x is closed as Berkovich spaces are locally hausdorff and thus T1, and so every point in the closure of $G^{\square}(k) \times \mathcal{M}(\mathcal{H}(p))$ will have image p . Because $\mathbf{p}(p)$ is the image in X^{an} of the unique Shilov boundary point of $G^{\square} \times \mathcal{M}(\mathcal{H}(p))$ under

$$G^{\square} \times \mathcal{M}(\mathcal{H}(p)) \hookrightarrow G^{\square} \times X^{\text{an}} \xrightarrow{m^{\text{an}}} X^{\text{an}}$$

we need to show that the unique point in the Shilov boundary of $\widehat{G^{\square} \times \mathcal{M}(\mathcal{H}(p))}$ is in the closure of $G^{\square}(k) \times \mathcal{M}(\mathcal{H}(p))$. The field k is algebraically closed so $G(k) \times \text{Spec } \widehat{\mathcal{H}(p)}$ is a dense subset of $G \times \text{Spec } \widehat{\mathcal{H}(p)}$. The following lemma will complete the proof. \square

Lemma 3.1. Let L/k be a nonarchimedean field extension, and let Σ be a subset of $G^{\square} \times_k \mathcal{M}(L)$ with $r(\Sigma)$ dense in $G \times_k \text{Spec}(\tilde{L})$. Then the Shilov boundary is in the closure of Σ .

Proof. The case when G is a torus is proved in [Thu07, lemma 2.4], and the proof proceeds the same for any connected linear algebraic group G . \square

3.2 The action of \mathbf{p} on homogeneous spaces

If G is reductive and X is a spherical G -variety, then any G -invariant subvariety is in fact the closure of the G -orbit of some closed point $y \in X$ ([Kno91]). Thus we will restrict our attention to when G acts on an orbit. In this case we will give a formula for \mathbf{p} in terms of valuations that will immediately relate it to the tropicalization map. We begin by recalling the construction of Luna and Vust which is used in [TV21] to define tropicalization.

Let $\overline{K} = k\{\{u\}\}$ denote Puiseux series over k , equipped with the u -adic valuation $\|\cdot\|_u$. In the case that G/H is spherical for any \overline{K} -point x , Luna and Vust define a $G(k)$ -invariant valuation $\|\cdot\|_x$ ([LV83, Section 4]). For any $f \in k(G/H)$ there is a nonempty open subset of $U \subseteq G(k)$ such that if $g \in U$ then $gf(x)$ is well defined. Luna and Vust show that for any $f \in k(G/H)$ there is a dense open $U_f \subseteq G(k)$ where $\|gf(x)\|_u$ is constant as g varies over U_f . The valuation $\|\cdot\|_x$ is given by $\|f\|_x = \|gf(x)\|_u$ for $g \in U_f$.

We will now explain how this generalizes to other valued extensions L/k . Let $x: \text{Spec } L \rightarrow G$ and denote the valuation on L by $\|\cdot\|$. The proof of the following lemma uses the same technique as the proof of [Kno91, Lemma 1.4].

Lemma 3.2. For each $f \in k(G)$ there is a nonempty Zariski open $U_f \subseteq U$ such that for any $g \in U_f$, $\|gf(x)\|$ is constant. For $f \in k[G]$ we have $\|gf(x)\| = \max_{h \in G(k)} \|hf(x)\|$ when $g \in U_f$.

Proof. Without loss of generality we can assume $f \in k[G]$. Let M be the G -module generated by f , this is a finite dimensional vector space and thus has basis $g_1 f, \dots, g_r f$. Let $v = \max_i \|g_i f(x)\|$. Let $V(v) = \{f' \in k[G] \mid \|f'(x)\| < v\}$. Then set $U_f = \{g \in G(k) \mid gf \notin V(v) \cap M\}$. \square

Let $\|\cdot\|_x : k(G) \rightarrow (0, \infty)$ be given by:

$$f \mapsto \|gf(x)\| \quad g \in U_f. \tag{1}$$

Corollary 3.3. The function $\|\cdot\|_x$ defines a $G(k)$ -invariant valuation on $k(G)$.

Lemma 3.3. Let $x: \text{Spec } L \rightarrow G$ be as above and let $p = (x, \|\cdot\|)$ be the corresponding point in G^{an} . Then if $\mathbf{p}(p) = (\eta_G, \overline{\|\cdot\|})$, we have that $\overline{\|\cdot\|} = \|\cdot\|_x$.

Proof. We will show that $\|\cdot\|_x$ and $\overline{\|\cdot\|}$ agree for a regular functions $f \in k[G]$. Let m_x be the composition $G \times \text{Spec } k(x) \hookrightarrow G \times G \xrightarrow{m} G$. Then if $m_x^* f = \sum_i g_i \otimes f_i(x) \in k[G] \otimes k(x)$ we have $\|f\|_x = \|gf(x)\| = \|\sum_i g_i(g) \otimes f_i(x)\| \leq \max_i (\|f_i(x)\|)$, where $g \in U_f$, U_f the open subset of $G(k)$ from Lemma 3.2. But then $\overline{\|f\|} = \|m_x^* f\|_{\mathbf{g}, p}$ and

$$\|m_x^* f\|_{\mathbf{g}, p} = \inf \left\{ \max_i \|g_i\|_{\mathbf{g}} \|f_i(x)\|_p \mid \sum_i g_i \otimes f_i(x) = m_x^* f \right\}.$$

Because $\|\cdot\|_{\mathbf{g}}$ is trivial we have that $\|f\|_x \leq \overline{\|f\|}$.

Conversely, to show that $\|f\|_x \geq \overline{\|f\|}$ it suffices to show that there is one expression $m_x^* f = \sum_i g_i \otimes f_i(x)$ such that $\|f\|_x = \max_i \|f_i(x)\|$. Let $m_x^* f = \sum_i g_i \otimes f_i(x)$, by manipulating the sum we can assume all the g_i and $f_i(x)$ are nonzero and the functions g_i are k -linearly independent. We can also assume that $\max_i (\|f_i(x)\|) = \|f_1(x)\|$. Let V be the k -span in L of $f_i(x)$. Fix $\alpha > 1$. It follows from [BGR84, Subsection 2.6.2, Proposition 3]³ there is a k -basis of V , v_1, \dots, v_r such that v_1 is a multiple of $f_1(x)$, and for any $\lambda_1, \dots, \lambda_r \in k$

$$\left\| \sum_j \lambda_j v_j \right\| \geq \alpha^{-1} \max_j (\|v_j\|).$$

Then we have for each i , $f_i(x) = \sum_j \lambda_j^i v_j$. So then $m_x^* f = \sum_j (\sum_i \lambda_j^i g_i) \otimes v_j$, by assumption for each j , the function $\sum_i \lambda_j^i g_i \in k[G]$ is not identically 0. Take $g \in U_f$ and such that $\sum_i \lambda_j^i g_i(g) \neq 0$, which exists because U_f is nonempty and open and so is the set of points such that $\sum_i \lambda_j^i g_i(g) \neq 0$. Now we have that:

$$\|f\|_x = \left\| \sum_j (\sum_i \lambda_j^i g_i(g)) \otimes v_j \right\| \geq \alpha^{-1} \max_j (\|v_j\|) = \alpha^{-1} \|f_1(x)\|.$$

But this holds for all $\alpha > 1$. So $\|f\|_x = \|f_1(x)\| = \max_i (\|f_i(x)\|)$. So $\|f\|_x \geq \overline{\|f\|}$, and thus the valuations are equal on $k[G]$, and thus are equal on all of $k(G)$. \square

Let Y be the G -orbit in X of some closed point y . The morphism $G \rightarrow Y$ given by $g \mapsto gy$ defines an inclusion $k(Y) \rightarrow k(G)$. Given some valued extension L/k and $x \in G(L)$, one can restrict $\|\cdot\|_x$ to $k(Y)$. Let x' be the image of x under the map $G(L) \rightarrow Y(L)$. Then $\|\cdot\|_x$ is in fact given by

$$\|f\|_x = \|gf(x')\| \quad g \in U_f. \quad (2)$$

We denote the restriction of $\|\cdot\|_x$ to $k(Y)$ by $\|\cdot\|_{x'}$.

Corollary 3.4. The map $\mathbf{p}: Y^{\text{an}} \rightarrow Y^{\text{an}}$ is given by $(x, \|\cdot\|) \mapsto (\eta_Y, \|\cdot\|_{x'})$. Furthermore all the $G(k)$ -invariant valuations on $k(Y)$ have the form $\|\cdot\|_{x'}$ for some $x' \in Y$.

Proof. We have a surjection of sets $G^{\text{an}} \rightarrow Y^{\text{an}}$ and a commutative diagram of maps of analytic spaces:

$$\begin{array}{ccc} G^{\text{an}} \times G^{\text{an}} & \longrightarrow & G^{\text{an}} \\ \downarrow & & \downarrow \\ G^{\text{an}} \times Y^{\text{an}} & \longrightarrow & Y^{\text{an}} \end{array}$$

Thus by Proposition 3.2 part 3 we see that $\mathbf{p}((x, \|\cdot\|)) = (\eta_Y, \|\cdot\|_{x'})$.

To see that this is all $G(k)$ -invariant valuations on $k(Y)$, notice that it follows from Proposition 3.4 that all $G(k)$ -invariant valuations on $k(G)$ are of the form $\|\cdot\|_x$ for some $x \in X(L)$ and it follows from [Kno91, Corollary 1.5] that any $G(k)$ -invariant valuation on $k(Y)$ is the restriction of one on $k(G)$. \square

3.3 A strong deformation retraction when G is reductive

Assume G is a reductive group. We construct a strong deformation retraction from X^{an} onto $\mathbf{p}(X^{\text{an}})$, using the same method as the proof of [Ber90, Theorem 6.2.1]. Fix a maximal torus T in G , Borel subgroup B_+ containing T , and let B_- be the opposite Borel i.e. the unique Borel subgroup of G such that $B_+ \cap B_- = T$. Let U_+ be the unipotent radical of B_+ , and let U_- be the unipotent radical of B_- . Then the map:

$$U_- \times T \times U_+ \rightarrow G$$

given by the multiplication map, is in fact an open immersion [GP11, Exposé XXII]. Denote the image by Ω . The groups U_- and U_+ are isomorphic affine N -space for some N . Let $U_+ = \text{Spec } k[u_1, \dots, u_N]$, and

³While [BGR84] works with k nontrivially valued, this result holds for trivially valued k as well, the proof is effectively the same. Such bases are called α -Cartesian bases.

$U_- = \text{Spec } k[u_{-1}, \dots, u_{-N}]$, and let $T = \text{Spec}(k[t_1^{\pm 1}, \dots, t_n^{\pm 1}])$. Then the coordinate ring of Ω , $k[\Omega]$, is given by $\text{Spec } k[t_i^{\pm 1}, u_j]$. Then set $T_i = t_i - 1$. For each $\lambda \in [0, 1]$ there is a point $\mathbf{g}_\lambda \in k[\Omega]^{\square}$ given by the semivaluation $\|\cdot\|_{\mathbf{g}_\lambda}$, which is defined by:

$$\left\| \sum_{I,J} \alpha_{I,J} T^I u^J \right\|_{\mathbf{g}_\lambda} = \min_{I,J} \|\alpha_{I,J}\| \lambda^{|I|+|J|}$$

where I and J denote multi-indices. Then $\mathbf{g}_1 = \mathbf{g}$ and $\mathbf{g}_0 = e$, the identity of G . Thus we have a map of sets:

$$H: [0, 1] \times X^{\text{an}} \rightarrow X^{\text{an}}$$

$(\lambda, p) \mapsto \mathbf{g}_\lambda * p$. Furthermore $H(0, p) = p$ and $H(1, p) = \mathbf{p}(X^{\text{an}})$. Notice that because $\mathbf{g}_\lambda \in k[\Omega]^{\square}$ for any $\lambda \in [0, 1]$, we can restrict H to a map $[0, 1] \times X^{\square} \rightarrow X^{\square}$, with $H(0, p) = p$ and $H(1, p) = \mathbf{p}(X^{\square})$.

Proposition 3.5. The map H is a homotopy, and in fact a strong deformation retraction of X^{an} onto $\mathbf{p}(X^{\text{an}})$. Similarly H defines a strong deformation retraction from X^{\square} onto $\mathbf{p}(X^{\square})$.

Proof. That H is continuous follows directly from [Ber90, Corollary 6.1.2]. It follows from Proposition 3.2 part 2 that H is strong deformation retract because we have that $\mathbf{g}_\lambda * \mathbf{g}_1 = \mathbf{g}_1$. Indeed, for $f \in k[\Omega]$ we will show that $\|m^* f\|_{\mathbf{g}_\lambda, \mathbf{g}_1} = 1$. Let $m^* f = \sum a_{I,J}^{L,M} t^I u^J \otimes t^L u^M$. Without loss of generality we can assume that $a_{I,J}^{L,M} = 0$ whenever I or L is less than 0, because otherwise we can factor out a power of $t^R \otimes t^S$, and $\|t^R \otimes t^S\|_{\mathbf{g}_\lambda, \mathbf{g}_1} = 1$. But then we can rewrite the sum by substituting T_i for $t_i - 1$, and we have that $m^* f = \sum b_{I,J}^{L,M} T^I u^J \otimes T^L u^M$. Note that T^I and u^J both evaluate to 0 on the identity. If $f \neq 0$ then $e^* \times \text{id} \circ m^* f \neq 0$, so there must be some term $b_{0,0}^{L,M} \neq 0$. But then:

$$\|m^* f\|_{\mathbf{g}_\lambda, \mathbf{g}_1} = \left\| \sum b_{I,J}^{L,M} T^I u^J \otimes T^L u^M \right\|_{\mathbf{g}_\lambda, \mathbf{g}_1} = \max_{I,J,L,M} (\|b_{I,J}^{L,M}\|) (\lambda^{|I|+|J|}) = 1$$

as the maximum is achieved when $|I| + |J| = 0$. □

4 The Luna-Vust Classification of Spherical varieties

Here we review the classification of embeddings of spherical homogeneous spaces via colored fans. The classification was first given by Luna and Vust in [LV83], though we will follow the exposition of [Kno91]. Much of the notation for spherical varieties is different than that of toric varieties, we will attempt to align our notation with that of toric varieties, similar to the notation in [Nas17].

Let k be an algebraically closed and trivially valued field⁴ and let G be a reductive group over k . Let H be a closed subgroup such that G/H has an open B -orbit, for some Borel subgroup $B \subseteq G$. A G/H -**embedding** is a normal G -variety X with a G -equivariant open embedding $G/H \hookrightarrow X$. Note that if X is a spherical variety with open B -orbit B_0 , then it is a G_0 -embedding where G_0 is the open G -orbit GB_0 . The Luna-Vust theory is a generalization of the toric case which puts embeddings of a torus in bijection with fans. Here G/H -embeddings are put in bijection with *colored fans*. To define colored fans and state the classification we will need to introduce some data associated to G/H , primarily a lattice M , its dual lattice N , a convex cone \mathcal{V} in $N \otimes_{\mathbb{Z}} \mathbb{R}$, and a finite set of *colors* $\mathcal{D}(G/H)$.

4.1 Data associated to G/H

Let $\mathfrak{X}(B)$ be the character lattice of B and define:

$$k(G/H)^{(B)} = \{f \in k(G/H)^{\times} \mid \text{there exists } \chi_f \in \mathfrak{X}(B) \text{ such that for any } b \in B(k), b \cdot f = \chi_f(b)f\}.$$

⁴In the literature on spherical varieties one does not generally specify a valuation on k , but it is assumed the valuations on $k(X)$ are trivial when restricted to k .

We say elements of $k(G/H)^{(B)}$ are B semi-invariant. There is a map $k(G/H)^{(B)} \rightarrow \mathfrak{X}(B)$ given by:

$$f \mapsto \chi_f$$

and the kernel is exactly k^\times . Let the image of this map be M , set $N = \text{hom}(M, \mathbb{Z})$, $M_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$, and $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. We say M is the **weight lattice** of G/H . Notice that every valuation $\|\cdot\|$ on $k(G/H)$ defines an element $\varrho(\|\cdot\|) \in N_{\mathbb{R}}$ by:

$$\varrho(\|\cdot\|)(\chi_f) = -\log(\|f\|).$$

In particular if \mathcal{V} is the set of $G(k)$ -invariant valuations on $k(G/H)$ then the map ϱ restricted to \mathcal{V} is an injection by [Kno91, Corollary 1.8] and the image is a finitely generated convex cone [Kno91, Corollary 5.3]. In general though the map ϱ is *not* injective. We say \mathcal{V} is the **valuation cone** of G/H and identify it with its image in $N_{\mathbb{R}}$.

Now let X be a G/H -embedding, and let Y be a G -orbit. Then define:

$$\begin{aligned} \mathcal{D}(X) &:= \{D \mid D \text{ a } B\text{-stable prime divisor in } X\} \\ \mathcal{D}_Y(X) &:= \{D \mid D \text{ a } B\text{-stable prime divisor in } X \text{ and } D \supseteq Y\} \end{aligned}$$

For any prime divisor D in X let $\|\cdot\|_D$ be the valuation D induces on $k(G/H)$. For convenience we may write $\varrho(D)$ when we mean $\varrho(\|\cdot\|_D)$.

4.2 Colored fans

We can now define colored fans and colored cones.

Definition 4.1. A **colored cone** is a pair (σ, \mathcal{F}) where σ is a strictly convex cone in $N_{\mathbb{R}}$, and \mathcal{F} is a subset of $\mathcal{D}(G/H)$ (referred to as the colors), such that the following are satisfied:

CC1 σ is generated by $\varrho(\mathcal{F})$ and finitely many elements of \mathcal{V} .

CC2 The relative interior of σ intersects \mathcal{V} .

CC3 $0 \notin \varrho(\mathcal{F})$.

We say (σ', \mathcal{F}') is a **colored face** of a colored cone (σ, \mathcal{F}) if σ' is a face of the cone σ , the relative interior of σ' intersects \mathcal{V} , and $\mathcal{F}' = \mathcal{F} \cap \varrho^{-1}(\sigma')$. A **colored fan**, \mathfrak{F} , is a finite set of colored cones containing all the colored faces of any colored cone in \mathfrak{F} , and for any $\|\cdot\| \in \mathcal{V}$ there is at most one colored cone $(\sigma, \mathcal{F}) \in \mathfrak{F}$ with $\|\cdot\|$ in the relative interior of σ .

We associate a colored cone to each G -orbit in X, Y , as follows. First define:

$$\begin{aligned} \mathcal{B}_Y(X) &= \{D \in \mathcal{D}_Y(X) \mid D \text{ is } G\text{-stable}\} \\ \mathcal{F}_Y(X) &= \{D \in \mathcal{D}_Y(X) \mid D \text{ is not } G\text{-stable}\} \end{aligned}$$

and let $\sigma_Y(X)$ be the cone in $N_{\mathbb{R}}$ generated by $\rho(\mathcal{B}_Y(X))$ and $\rho(\mathcal{F}_Y(X))$. We identify the elements of $\mathcal{F}_Y(X)$ with elements of $\mathcal{D}(G/H)$ by intersecting each element D with G/H , as $D \cap G/H$ is dense in D when D isn't G -stable. Then we get a colored cone $(\sigma_Y(X), \mathcal{F}_Y(X))$. Just as affine toric varieties correspond to cones, we have a particular type of G/H -embedding corresponding to colored cones. A G/H -embedding is **simple** if it has a unique closed G -orbit.

Theorem 4.1. [Kno91, Theorem 3.1] The map

$$X \mapsto (\sigma_Y(X), \mathcal{F}_Y(X))$$

gives a bijection between isomorphism classes of simple G/H -embeddings and colored cones.

We can construct all simple subembeddings of X in the following way. Let Y be a G -orbit in X , then define:

$$X_0 := X \setminus \left(\bigcup_{D \in \mathcal{D}(X) \setminus \mathcal{D}_Y(X)} D \right). \quad (3)$$

Note that $\mathcal{D}(X) \setminus \mathcal{D}_Y(X)$ consists of all divisors D such that $\|\cdot\|_D$ is *not* an element of $\sigma_Y(X)$.

Theorem 4.2. [Kno91, Theorem 2.1] The open subvariety X_0 is a B -stable, affine open subset of X . The intersection $Y \cap X_0$ is an open B -orbit, and GX_0 is a simple embedding with closed orbit Y .

When X is simple we have that $GX_0 = X$. The following theorem describes the cone $\sigma_Y(X)$ in this case.

Theorem 4.3. [Kno91, Theorem 2.5] Let X be a simple embedding with closed orbit Y , and let X_0 be as in 3.

1. $k[X_0]^{(B)} = \{f \in k(G/H)^{(B)} \mid \chi_f \in (\sigma_Y(X))^\vee\}$.
2. The center of $\|\cdot\|$ exists if and only if $\|\cdot\| \in \sigma_Y(X)$.
3. The center of $\|\cdot\|$ is Y if and only if $\|\cdot\|$ is in the relative interior of $\sigma_Y(X)$.

Furthermore we have the following about the faces of $\sigma_Y(X)$ and the orbits of X .

Lemma 4.1. [Kno91, Lemma 3.2] The map

$$Z \mapsto (\sigma_Z(X), \mathcal{F}_Z(X))$$

gives a bijection between orbits Z whose closure contains Y and the colored faces of $(\sigma_Y(X), \mathcal{F}_Y(X))$.

Due to the above, and the fact that there are finitely many G -orbits in X ([Kno91]), we have that the set $\mathfrak{F}(X) := \{(\sigma_Y(X), \mathcal{F}_Y(X)) \mid Y \text{ is a } G\text{-orbit}\}$ is a colored fan. We can now state the main theorem in the classification.

Theorem 4.4. [Kno91, Theorem 3.3] The map:

$$X \mapsto \mathfrak{F}(X)$$

is a bijection between isomorphism classes of G/H -embeddings and colored fans.

Example 4.1. The classification of toric varieties by fans is a special case of the Luna-Vust theory. Let $G = T$ be a torus, let H be trivial. Then $\mathcal{D}(T) = \emptyset$, N is the cocharacter lattice of T , and T -embeddings are classified by fans in $N_{\mathbb{R}}$.

Example 4.2. Let $G = \mathrm{SL}_2$ and let H be the subgroup of upper triangular matrices with 1s on the diagonal. Let O denote the origin of \mathbb{A}^2 , there is an isomorphism $G/H \rightarrow \mathbb{A}_k^2 \setminus O$ given by mapping a matrix to its first column. The complement of the x -axis is an open orbit of the Borel subgroup of upper triangular matrices, B . We have that $\mathfrak{X}(B) \cong \mathbb{Z}$, where the generator is the character χ given by:

$$\begin{bmatrix} a & b \\ 0 & a^{-1} \end{bmatrix} \mapsto a.$$

Furthermore $k(G/H)^{(B)}/k^\times = \{y^n \mid n \in \mathbb{Z}\}$ and $\chi_{y^n} = \chi^{-n}$, so $M = \mathfrak{X}(B)$. Thus $N_{\mathbb{R}} \cong \mathbb{R}$. Let D be the x -axis in \mathbb{A}^2 . Consider the embedding $\mathbb{A}^2 \hookrightarrow \mathbb{P}^2$ given by $(a, c) \mapsto [a : c : 1]$. Extend the SL_2 -action to \mathbb{P}^2 by letting SL_2 act on the first two homogeneous coordinates while preserving the third.

Let L be the line at infinity in \mathbb{P}^2 . Then both $\|\cdot\|_D$ and $\|\cdot\|_L$ are $G(k)$ -invariant and furthermore $\varrho(\|\cdot\|_D) = 1$ and $\varrho(\|\cdot\|_L) = -1$, thus $\mathcal{V} = N$. There is one color, given by the x -axis. The table below consists of a complete list of all G/H -embeddings. The hollow circle denotes the color D , and E denotes the exceptional divisor in the blow ups at O : $\mathrm{Bl}_O \mathbb{A}^2$ and $\mathrm{Bl}_O \mathbb{P}^2$.

X	Closed G -orbits	$\mathfrak{F}(X)$
$\mathbb{A}^2 \setminus O$		-----●-----
$\mathrm{Bl}_O \mathbb{A}^2$	E	-----●----- ----->
$\mathbb{P}^2 \setminus O$	L	<-----●-----
$\mathrm{Bl}_O \mathbb{P}^2$	L, E	<-----●----- ----->
\mathbb{A}^2	O	-----●○----->
\mathbb{P}^2	L, O	<-----●○----->

4.3 Orbits and Orbit Closures

Let Z be the G -orbit of a closed point in X . Notice that from Theorem 4.2 it follows that Z is spherical. Let \overline{Z} be the closure of Z in X , so \overline{Z} is a Z -embedding. Let $M(Z)$ be the weight lattice of Z , $N_{\mathbb{R}}(Z) = \text{hom}(M(Z), \mathbb{R})$, $\mathcal{V}(Z)$ be the valuation cone, and $\mathcal{D}(Z)$ be the B -stable prime divisors. We will relate the above to M , $N_{\mathbb{R}}$, \mathcal{V} , and $\mathcal{D}(G/H)$. Furthermore we will give the relationship between the colored fan of X and that of \overline{Z} . See [Kno91, Section 4] for further discussion of the following.

Given $z \in Z$ there is a map $\varphi: G/H \rightarrow Z$ given by $gH \mapsto gz$. This induces a map of function fields $k(Z) \hookrightarrow k(G/H)$. The map $k(Z) \hookrightarrow k(G/H)$ gives an inclusion $M(Z) \hookrightarrow M$ and after dualizing, a surjection $\varphi_*: N_{\mathbb{R}} \rightarrow N_{\mathbb{R}}(Z)$. Let $\tau = \sigma_Z(X)$, $N(\tau) = N_{\mathbb{R}}/\text{span}(\tau)$, and let $M(\tau)$ be the dual vector space to $N(\tau)$. Consider the following subset of \mathcal{D} :

$$\mathcal{F}_{\varphi} = \{D \in \mathcal{D} \mid D \text{ maps dominantly to } Z\}.$$

Let $\mathcal{F}_{\phi}^c = \mathcal{D} \setminus \mathcal{F}_{\phi}$.

Theorem 4.5. [Kno91, Theorem 4.4] We have that $N(\tau) = N(Z)$, the map φ_* is the projection $N \rightarrow N(\tau)$, $M(\tau) = M(Z)$, $\varphi_*(\mathcal{V}) = \mathcal{V}(Z)$, and there is a bijection $\mathcal{F}_{\varphi}^c \rightarrow \mathcal{D}(Z)$ given by φ .

To state the relationship between $\mathfrak{F}(X)$ and $\mathfrak{F}(\overline{Z})$ define the following colored fan:

$$\text{Star}(\tau) := \{(\varphi_*(\sigma), \varphi_*(\mathcal{F} \cap \mathcal{F}_{\varphi}^c)) \mid (\sigma, \mathcal{F}) \in \mathfrak{F}(X) \text{ and } (\tau, \mathcal{F}_Z(X)) \text{ is a colored face of } (\sigma, \mathcal{F})\}.$$

Then as a result of [Kno91, Theorem 4.5] we have that.

Theorem 4.6. Let X be a G/H -embedding, and let $Z \subseteq X$ be a G -orbit. Let the Zariski closure of Z in X be \overline{Z} . Then $\text{Star}(\sigma_Z(X)) = \mathfrak{F}(\overline{Z})$.

5 Spherical Tropicalization

In this Section we review tropicalization for toric varieties, tropicalization of spherical varieties, and prove Theorem A. We also discuss applications of Theorem A. Let k be an algebraically closed and trivially valued field, let G be a reductive group over k , let X be a spherical G -variety with open G -orbit G/H .

5.1 Tropicalization of toric varieties

Let $\overline{K} = k\{\{u\}\}$ be the field of Puiseux series, equipped with the u -adic valuation which we denote $\|\cdot\|_u$. Let $T = \text{Spec } k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ and let be W a subvariety of T . There is a map of sets $\text{trop}_T: W(\overline{K}) \rightarrow \mathbb{R}^n$ given by:

$$(t_1, \dots, t_n) \mapsto (-\log(\|t_1\|_u), \dots, -\log(\|t_n\|_u)).$$

We say the closure of $\text{trop}_T(W(\overline{K}))$ is the **tropicalization of W** , denoted $\text{Trop}_T(W)$.

This construction extends to subvarieties of toric varieties, as follows. Let X be a toric variety, with dense torus T . Let N be the cocharacter lattice of T , and let $F(X)$ be the fan in $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$ associated to X . Given a cone $\sigma \in F(X)$, consider the vector space $N(\sigma) = N_{\mathbb{R}}/\text{span}(\sigma)$. As a set we define we the tropicalization of X to be:

$$\text{Trop}_T(X) := \bigsqcup_{\sigma \in F(X)} N(\sigma).$$

To define a topology on $\text{Trop}_T(X)$ we first consider the case when X is affine; so X corresponds to a cone $\sigma \in F(X)$. Let $M_{\mathbb{R}}$ be the dual vector space to $N_{\mathbb{R}}$; let $\sigma^{\vee} \subseteq M_{\mathbb{R}}$ denote linear functions on $N_{\mathbb{R}}$ that are nonnegative on σ . Let $\overline{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$. If $\text{hom}(\sigma^{\vee}, \overline{\mathbb{R}})$ denotes semigroup homomorphisms $\sigma^{\vee} \rightarrow \overline{\mathbb{R}}$, then there is a natural bijection $\text{Trop}_T(X) \rightarrow \text{hom}(\sigma^{\vee}, \overline{\mathbb{R}})$ defined as follows. The fan $F(X)$ consists of the faces of σ . Let τ be a face of σ and let $M(\tau)$ be the dual of $N(\tau)$. Recall that $M(\tau) = \tau^{\perp} \subseteq M_{\mathbb{R}}$. So $\alpha \in N(\tau)$ defines a semigroup homomorphism $\tilde{\alpha}: \sigma^{\vee} \rightarrow \overline{\mathbb{R}}$

$$\tilde{\alpha}(v) = \begin{cases} \alpha(v), & v \in \tau^{\perp} \cap \sigma^{\vee} \\ \infty, & \text{else} \end{cases}$$

This then defines a bijection $\text{Trop}_T(X) \rightarrow \text{hom}(\sigma^\vee, \overline{\mathbb{R}})$ and we give $\text{Trop}_T(X)$ the topology $\text{hom}(\sigma^\vee, \overline{\mathbb{R}})$ inherits from $\overline{\mathbb{R}}^{\sigma^\vee}$. For a general toric variety X we glue the tropicalizations of affine toric subvarieties according to the shared faces of their respective cones in $F(X)$. For further discussion of tropicalization of toric varieties and their subvarieties see [Pay09].

Recall that X is a disjoint union of tori which are in bijection with the cones of $F(X)$. Thus there is a tropicalization map $X(\overline{K}) \rightarrow \text{Trop}_T(X)$ which is defined by taking a disjoint union of the tropicalization maps of the tori $T(\sigma) \rightarrow N(\sigma)$, where $T(\sigma)$ is the torus in X corresponding to $\sigma \in F(X)$. Thus for any subvariety W of a toric variety X we have a tropicalization map from $W(\overline{K})$ into $\text{Trop}_T(X)$.

5.2 Tropicalization of spherical varieties

In [TV21] Tevelev and Vogianou define a tropicalization map $\text{trop}_G: G/H(\overline{K}) \rightarrow \mathcal{V}$. The map is given as follows. For each $x \in G/H(\overline{K})$ define $\|\cdot\|_x$ to be the valuation given by:

$$\|f\|_x = \|gf(x)\|_u \quad g \in U_f$$

as in equation 1. Then define $\text{trop}_G(x) = \|\cdot\|_x$, this gives a map $G/H(\overline{K}) \rightarrow \mathcal{V}$. For a subvariety $W \subseteq G/H$ the **tropicalization** is the closure of $\text{trop}_G(W(\overline{K}))$, in particular the tropicalization of G/H is \mathcal{V} .

In [Nas17] the above is extended to construct a tropicalization for any G/H -embedding X , this is done similarly to the toric case. As a set the **tropicalization**, denoted $\text{Trop}_G(X)$, is defined to be:

$$\text{Trop}_G(X) := \bigsqcup_{(\tau, \mathcal{F}) \in \mathfrak{F}(X)} \mathcal{V}(\tau)$$

where $\mathcal{V}(\tau)$ is the valuation cone of the G -orbit associated to τ .

Remark 5.1. If G is a torus and X is a toric variety then the cones $\mathcal{V}(\tau)$ are exactly the vector spaces $N(\tau)$ as in the previous subsection.

We now put a topology on $\text{Trop}_G(X)$. Consider first the case when X is simple with closed orbit Y ; so X corresponds to the colored cone $(\sigma_Y(X), \mathcal{F}_Y(X))$. Let $\sigma = \sigma_Y(X)$. Recall that $\text{hom}(\sigma^\vee, \overline{\mathbb{R}})$ is in bijection with the disjoint union of the vector spaces $N(\tau) = N_{\mathbb{R}}/\text{span}(\tau)$, where τ ranges over the faces of σ . The bijection is given by $\alpha \mapsto \tilde{\alpha}$ where

$$\tilde{\alpha}(v) = \begin{cases} \alpha(v), & v \in \tau^\perp \cap \sigma^\vee \\ \infty, & \text{else} \end{cases}$$

The colored faces of $(\sigma, \mathcal{F}_Y(X))$ are exactly the faces of σ whose relative interior intersects \mathcal{V} . Given such a face τ , $N(\tau)$ contains the valuation cone $\mathcal{V}(\tau)$. Thus we have that $\text{Trop}_G(X) \subseteq \text{hom}(\sigma^\vee, \overline{\mathbb{R}}) \subseteq \overline{\mathbb{R}}^{\sigma^\vee}$ and we give $\text{Trop}_G(X)$ the induced topology. For X not necessarily simple, $\text{Trop}_G(X)$ is obtained by tropicalizing the simple spherical subvarieties and gluing the valuation cones of shared orbits.

Remark 5.2. Let X be simple with closed orbit Y , and $\sigma = \sigma_Y(X)$. In general $\text{Trop}_G(X)$ is not equal to $\text{hom}(\sigma^\vee, \overline{\mathbb{R}})$ as we do not necessarily have that $\sigma \subseteq \mathcal{V}$ or that $\mathcal{V}(\tau) = N(\tau)$, for a given colored face (τ, \mathcal{F}) .

Example 5.1. Let $X = \text{Bl}_O \mathbb{P}^2$ be the embedding of SL_2/H as described in example 4.2. Then the tropicalization is given by



The left most point representing the valuation cone of L , the right most of E , and the interior representing the valuation cone of SL_2/H . One can find further illustrated examples of this in Sections 4 and 5 of [Nas17].

We have a tropicalization map $\text{Trop}_G: X(\overline{K}) \rightarrow \text{Trop}_G(X)$ extending the one defined in [TV21]. This is because all G -orbits of X are spherical homogeneous spaces so we can take the disjoint union of the tropicalization maps. Explicitly, for $x \in X(\overline{K})$ if x is in the G -orbit Y then set $\text{Trop}_G(x)$ to be the valuation $\|\cdot\|_x$ on $k(Y)$. Then for a subvariety $W \subseteq X$ we can again define $\text{Trop}_G(W) := \text{trop}_G(W(\overline{K}))$

Remark 5.3. In [Nas17] Nash actually defines a *colored tropicalization*. This consists of attaching the finite set of colors associated to a colored face, to the corresponding valuation cone. However we do not deal with the colors here so this was not discussed.

We now prove Theorem A. Recall that in Section 3 we showed that we have a retraction of topological spaces $\mathbf{p}: X^{\text{an}} \rightarrow X^{\text{an}}$, given by $*$ -multiplication by the unique point of the Shilov boundary of G^{\square} (Corollary 3.2).

Theorem A. *Let X be a spherical G -variety over an algebraically closed and trivially valued field k , and let \overline{K} denote Puiseux series in k . Then there is a canonical homeomorphism $\iota: \text{Trop}_G(X) \rightarrow \mathbf{p}(X^{\text{an}})$, and $\mathbf{p}: X^{\text{an}} \rightarrow \text{Trop}_G(X)$ is a retraction of topological spaces such that the following diagram commutes:*

$$\begin{array}{ccc} X(\overline{K}) & \longrightarrow & X^{\text{an}} \\ & \searrow \text{trop}_G & \downarrow \mathbf{p} \\ & & \text{Trop}_G(X). \end{array}$$

Proof. First observe that as sets we have an inclusion $\iota: \text{Trop}_G(X) \hookrightarrow X^{\text{an}}$ given as follows. For each orbit Z of X , and $\|\cdot\| \in \mathcal{V}(Z)$, $\|\cdot\|$ defines an element of X^{an} by:

$$\|\cdot\| \mapsto (\eta_Z, \|\cdot\|)$$

where η_Z is the generic point of Z . All the G -invariant subvarieties of a spherical variety are closures of G -orbits ([Kno91]); thus by Proposition 3.4 we have that $\mathbf{p}(X^{\text{an}}) = \iota(\text{Trop}_G(X))$ as sets. That the diagram commutes as maps of sets follows from Corollary 3.4. Finally, $\mathbf{p}: X^{\text{an}} \rightarrow \mathbf{p}(X^{\text{an}})$ is continuous by Proposition 3.2. So it remains to show that the topology on $\text{Trop}_G(X)$ agrees with the topology of $\mathbf{p}(X^{\text{an}})$, i.e. that ι is a homeomorphism onto its image. We will need to temporarily emphasize the difference between $\text{Trop}_G(X)$ and its image in X^{an} . Let \mathcal{I} be the set $\iota(\text{Trop}_G(X))$, and give $\mathcal{I} \subseteq X^{\text{an}}$ the subspace topology.

Consider the case when X is simple with closed orbit Y . Let X_0 be the B -stable affine open defined by equation 3. The open X_0 meets every G -orbit in X by Theorem 4.2. Thus X_0^{an} contains \mathcal{I} . The topology on X_0^{an} is generated by the functions $\text{ev}_f: k[X_0]^{\text{an}} \rightarrow \overline{\mathbb{R}}$, where $\text{ev}_f(\|\cdot\|) = \|f\|$, $f \in k[X_0]$. The functions ev_f for $f \in k[X_0]^{(B)}$ define the topology on $\text{Trop}_G(X)$ by part 1 of Theorem 4.3. So the inclusion $\iota: \text{Trop}_G(X) \rightarrow \mathcal{I}$ is open. Given that for a general G/H -embedding $\text{Trop}_G(X)$ is given by gluing the tropicalizations of simple subembeddings we have an open bijection $\iota: \text{Trop}_G(X) \rightarrow \mathcal{I}$ for all X .

Consider the case then when X is proper. Then X^{an} is compact, and $\mathcal{I} = \mathbf{p}(X^{\square})$ is a closed subset, and therefore \mathcal{I} is compact. Thus $\iota: \text{Trop}_G(X) \rightarrow \mathcal{I}$ is a homeomorphism onto its image. Then for a general X there exists a proper spherical G -variety \overline{X} and an open G -equivariant embedding $X \hookrightarrow \overline{X}$ by [Sum74, Theorem 3]. But then we have a commutative diagram:

$$\begin{array}{ccc} X^{\text{an}} & \hookrightarrow & \overline{X}^{\text{an}} \\ \uparrow & & \uparrow \\ \text{Trop}_G(X) & \hookrightarrow & \text{Trop}_G(\overline{X}) \end{array}$$

where the composition $\text{Trop}_G(X) \hookrightarrow \text{Trop}_G(\overline{X}) \hookrightarrow \overline{X}^{\text{an}}$ is a homeomorphism onto its image by the above. Furthermore the map $X^{\text{an}} \hookrightarrow \overline{X}^{\text{an}}$ is an open embedding so the inclusion $\text{Trop}_G(X) \hookrightarrow X^{\text{an}}$ is a homeomorphism onto \mathcal{I} . \square

Remark 5.4. In [KM19] it is shown that tropicalization factors through X^{an} when X is affine. It follows from Corollary 3.4 that \mathbf{p} and the map $X^{\text{an}} \rightarrow \text{Trop}_G(X)$ given in Definition 5.18 of [KM19] agree.

Corollary 5.1. For W a subvariety of X , $\text{Trop}_G(W)$ is a strong deformation retract of W^{an} .

Proof. We have that $\text{Trop}_G(W) = \mathbf{p}(W^{\text{an}})$ so this follows from Proposition 3.5. \square

Remark 5.5. Notice that as a result of Proposition 3.2 one has that Trop_G defines a functor from the category of spherical G -varieties to the category of topological spaces. One can describe the morphism as follows. If $X \rightarrow X'$ is a morphism of G -varieties, then for a $Z \subseteq X$ a G -orbit, Z maps dominantly onto a G -orbit $Z' \subseteq X'$. Then $\|\cdot\| \in \mathcal{V}(Z)$ is mapped to its restriction to $k(Z')$ under the inclusion $k(Z') \hookrightarrow k(Z)$. This is the same as the construction of tropicalization of morphisms given in Section 2 of [Nas20].

From Theorem A we can see that tropicalization did not depend on our choice to use Puiseux series. Let L/k be an algebraically closed, valued extension, with nontrivial valuation. Define $\text{trop}_{G,L}: X(L) \rightarrow \mathcal{V}$ to be the map given $x \mapsto \|\cdot\|_x$, where x is contained in some G -orbit Y and $\|\cdot\|_x$ is the valuation on $k(Y)$ defined in 2.

Corollary 5.2. The closure of $\text{trop}_{G,L}(W(L))$ in $\text{Trop}_G(X)$ is equal to $\mathbf{p}(W^{\text{an}}) = \text{Trop}_G(W)$, where W^{an} is considered as a subspace of X^{an} induced by the inclusion $W \subseteq X$.

Proof. For a subvariety $V \subseteq X$, $W(L)$ is a dense subset of W^{an} , and we have a commutative diagram

$$\begin{array}{ccc} X(L) & \longrightarrow & X^{\text{an}} \\ & \searrow \text{trop}_{G,L} & \downarrow \mathbf{p} \\ & & \text{Trop}_G(X). \end{array}$$

So the closure of $\text{trop}_G(W(L))$ in $\text{Trop}_G(X)$ for any subvariety $W \subseteq X$ must be equal to $\mathbf{p}(W^{\text{an}})$. \square

In [Nas20] Nash gives a description of tropicalization by embedding X into a toric variety and then applying the usual tropicalization process for toric varieties. See [Nas20, Section 6] for details. In particular this process is used to prove the following result, for which we can give an alternate proof.

Theorem 5.1. [Nas20, Theorem 6.7] If $W \subseteq G/H$ is a subvariety, and \overline{W} is the closure of W in X . Then we have that:

$$\text{Trop}_G(\overline{W}) = \overline{\text{Trop}_G(W)}$$

where $\overline{\text{Trop}_G(W)}$ is the closure taken in $\text{Trop}_G(X)$.

Proof. This follows from Theorem A, as \mathbf{p} is a retraction. \square

6 Compactification of the colored fan

Recall from subsection 2.2 that when working over a trivially valued field X^{an} has a compact analytic subdomain X^{\triangleright} . The space X^{\triangleright} is defined as the subspace whose points are pairs $(x, \|\cdot\|) \in X^{\text{an}}$, where $\|\cdot\|$ has center in X . In this Section we study $\mathbf{p}(X^{\triangleright})$ and prove Theorem B. In the case of toric varieties it is shown in [Thu07, Section 2] that $\mathbf{p}(X^{\triangleright})$ is the canonical compactification of the fan associated to X . We begin by reviewing the work of Thuillier on the toric case.

Let X be a toric variety over k with dense torus T . Let $F(X)$ denote the fan associated to X . Let σ be a cone in $F(X)$ and U_σ the T -stable open affine associated to σ . We will assume all the cones of $F(X)$ are strictly convex. In [Thu07] Thuillier defines a map of topological spaces $X^{\triangleright} \rightarrow X^{\triangleright}$; which is equal to the map \mathbf{p} given by $*$ -multiplication by the unique point in the Shilov boundary of T^{\triangleright} . Let σ^\vee denote positive linear functionals on σ . Thuillier shows that $\rho^{-1}(T) \cap \mathbf{p}(U_\sigma^{\triangleright})$ is naturally homeomorphic to $\text{hom}(\sigma^\vee, [0, \infty))$, where hom denotes semigroup homomorphisms and $[0, \infty)$ has the additive semigroup structure. The space $\text{hom}(\sigma^\vee, [0, \infty))$ is naturally isomorphic to σ , as σ is strictly convex. Furthermore $\mathbf{p}(U_\sigma^{\triangleright})$ is compact, and in fact is canonically homeomorphic to $\text{hom}(\sigma^\vee, [0, \infty))$, the **canonical compactification** of σ . We will denote the canonical compactification of a strictly convex cone by $\overline{\sigma}$. In this way $\mathbf{p}(X^{\triangleright})$ is the canonical compactification of the fan $F(X)$.

Let X be a spherical G -variety with dense G -orbit G/H . First observe that \mathbf{p} has image consisting only of G -invariant valuations, and colored fans may contain points of Q lying outside \mathcal{V} . As an example consider

[Nas17, Example 4.4], where $G = \mathrm{GL}_2 \times \mathrm{GL}_2$, and H is the diagonal so $G/H \cong \mathrm{GL}_2$. This is depicted in Figure 1. Because of this we will assume the following:

$$\bigcup_{(\sigma, \mathcal{F}) \in \mathfrak{F}(X)} \sigma \subseteq \mathcal{V} \quad (\star)$$

Horospherical varieties, for example, satisfy this condition [Kno91, Corollary 6.2]; symmetric varieties often may not [Vus90].




Figure 1: To the left is the vector space $N_{\mathbb{R}}$, the valuation cone \mathcal{V} , and the one color D , for the homogeneous space GL_2 . The blue circle depicts $\varrho(\|\cdot\|_D)$, and the shaded triangle depicts \mathcal{V} . To the right is the colored cone for the G/H -embedding \mathbb{A}_k^4 . The cone is generated by $(1, 0)$ and $\varrho(\|\cdot\|_D)$.

Lemma 6.1. Let X be a G/H -embedding satisfying \star . Let Z be a G -orbit of X , and \bar{Z} be the closure of Z in X . Then the Z -embedding \bar{Z} satisfies \star .

Proof. Without loss of generality we may assume X is simple. Denote the closed orbit by Y . Let $\tau = \sigma_Z(X)$, the face of $\sigma_Y(X)$ corresponding to Z . For $z \in Z$ let $\varphi: G/H \rightarrow Z$ be the map given by $gH \mapsto gz$, and let $\varphi_*: N_{\mathbb{R}} \rightarrow N_{\mathbb{R}}(\tau)$ be the induced map. It follows from Theorem 4.6 that $\varphi_*(\sigma_Y(X)) = \sigma_Y(\bar{Z})$, and we know from Theorem 4.5 that $\varphi_*(\mathcal{V}) = \mathcal{V}(Z)$. So $\sigma_Y(\bar{Z}) \subseteq \mathcal{V}(\tau)$. \square

Because colored cones correspond to simple embeddings, we will first consider simple embeddings. Let X be simple with closed G -orbit Y . Let $\sigma = \sigma_Y(X)$. We begin by defining a map $\Phi: \mathbf{p}(X^{\triangleright}) \rightarrow \bar{\sigma}$. The set $\rho^{-1}(G/H) \cap \mathbf{p}(X^{\triangleright})$ consists of $G(k)$ -invariant valuations on $k(G/H)$ having center in X . Given such a valuation $\|\cdot\|$ we know it uniquely defines a semigroup homomorphism $\varrho(\|\cdot\|): M \rightarrow \mathbb{R}$ by:

$$\varrho(\|\cdot\|)(\chi_f) = -\log(\|f\|).$$

If $p = (\eta_{G/H}, \|\cdot\|) \in X^{\triangleright}$ then denote the restriction of $\varrho(\|\cdot\|)$ to σ^{\vee} by ϕ_p . By Theorem 4.3 part 2, we have that ϕ_p takes values in $[0, \infty)$. Thus we have a map of sets $\rho^{-1}(G/H) \cap \mathbf{p}(X^{\triangleright}) \rightarrow \mathrm{hom}(\sigma^{\vee}, [0, \infty))$ given by $p \mapsto \phi_p$.

A general point of $\mathbf{p}(X^{\triangleright})$ is a point $p = (\eta_Z, \|\cdot\|)$ where η_Z is the generic point of a G -orbit Z and $\|\cdot\|$ is a $G(k)$ -invariant valuation having center in \bar{Z} , where \bar{Z} is the Zariski closure of Z in X . Let τ be the face of σ corresponding to Z . Then $\|\cdot\|$ defines an element of $\mathrm{hom}(M(\tau), \mathbb{R}) = \mathrm{hom}(\tau^{\perp} \cap \sigma^{\vee}, \mathbb{R})$. This extends uniquely to a morphism of semigroups $\phi_p: \sigma^{\vee} \rightarrow \bar{\mathbb{R}}$ by setting $\phi_p(u) = \infty$ when $u \notin \tau^{\perp}$. But by Theorem 4.5 and 4.6 we have that $\tau^{\perp} \cap \sigma^{\vee} = \sigma_Y(\bar{Z})$, thus by Theorem 4.3 part 2 we see that ϕ_p only takes values in $[0, \infty]$. So we have a map $\Phi: \mathbf{p}(X^{\triangleright}) \rightarrow \bar{\sigma}$, given by $\Phi(p) = \phi_p$.

Proposition 6.1. Let X be a simple G/H -embedding corresponding to the colored cone (σ, \mathcal{F}) , and satisfying \star . Then there is a commutative diagram:

$$\begin{array}{ccc} \mathbf{p}(X^{\triangleright}) & \xrightarrow{\sim} & \bar{\sigma} \\ \uparrow & & \uparrow \\ \mathbf{p}(X^{\triangleright}) \cap \rho^{-1}(G/H) & \xrightarrow{\sim} & \sigma \end{array}$$

where the horizontal arrows are homeomorphisms and are given by Φ .

Proof. Notice that

$$\bar{\sigma} = \bigsqcup_{\tau} \text{hom}(\tau^{\perp} \cap \sigma^{\vee}, [0, \infty))$$

where τ runs over the faces of σ . But then because X satisfies \star we have that:

$$\bigsqcup_{\tau} \text{hom}(\tau^{\perp} \cap \sigma^{\vee}, [0, \infty)) = \bigsqcup_Z \text{hom}(\sigma_Y(\bar{Z})^{\vee}, [0, \infty))$$

where Z runs over the orbits of X , and \bar{Z} is the Zariski closure of Z in X . Because X satisfies \star all elements of $\text{hom}(\sigma_Y(\bar{Z})^{\vee}, [0, \infty))$ are uniquely given by $G(k)$ -invariant valuations on $k(Z)$, and by Theorem 4.3 part 2, $\text{hom}(\sigma_Y(\bar{Z})^{\vee}, [0, \infty))$ is exactly those $G(k)$ -invariant valuations with center in \bar{Z} . Thus it follows that Φ defines a bijection $\mathbf{p}(X^{\triangleright}) \rightarrow \bar{\sigma}$ and $\mathbf{p}(X^{\triangleright}) \cap \rho^{-1}(G/H) \rightarrow \sigma$.

It remains to show that Φ is a homeomorphism, and because $\bar{\sigma}$ is compact it suffices to show Φ is open. Let X_0 be the B -stable affine open from equation 3, note that X_0 intersects all G -orbits of X . So $\mathbf{p}(X^{\triangleright}) \subseteq X_0^{\triangleright}$. Recall from Theorem 4.3 part 1 that $k[X_0]^{(B)} = \{f \in k(G/H)^{(B)} \mid \chi_f \in (\sigma_Y(X))^{\vee}\}$. But then evaluation at the functions $f \in k[X_0]^{(B)}$ generates the topology on $\bar{\sigma}$, so Φ is open. \square

As a corollary we have the following.

Theorem B. *There is a strong deformation retraction from X^{\triangleright} onto $\mathbf{p}(X^{\triangleright})$. If the colored fan $\mathfrak{F}(X)$ satisfies:*

$$\bigcup_{(\sigma, \mathcal{F}) \in \mathfrak{F}(X)} \sigma \subseteq \mathcal{V}. \quad (\star)$$

Then $\mathbf{p}(X^{\triangleright})$ is homeomorphic to the canonical compactification of the fan $F(X) = \{\sigma \mid (\sigma, \mathcal{F}) \in \mathfrak{F}(X)\}$.

Proof. That X^{\triangleright} strong deformation retracts onto $\mathbf{p}(X^{\triangleright})$ space follows from Corollary 3.5. By Proposition 6.1 if X satisfies \star then $\mathbf{p}(X^{\triangleright})$ is homeomorphic to the canonical compactification of the fan $F(X) = \{\sigma \mid (\sigma, \mathcal{F}) \in \mathfrak{F}(X)\}$. \square

Because the map \mathbf{p} maps into the closure of \mathcal{V} in X^{\triangleright} by construction, one will not have the above for a general spherical variety. This is because a colored cone need not be contained in \mathcal{V} . However given a simple embedding X , one will have that $\mathbf{p}(X^{\triangleright}) = \overline{\sigma \cap \mathcal{V}}$, where the closure is taken in $\text{hom}(\sigma^{\vee}, [0, \infty])$. Consider for example the GL_2 -embedding $\mathbf{p}(X^{\triangleright})$ as in Figure 1, one can see $\mathbf{p}(X^{\triangleright})$ depicted in Figure 2.




Figure 2: The canonical compactification of the cone σ associated to X , along with $\mathbf{p}(X^{\triangleright})$ depicted as the pink shaded region to the right.

Remark 6.1. Because of this difference between spherical varieties that satisfy \star and those that do not, it may be interesting to describe \star -multiplication by \mathbf{b} , the unique point in the Shilov boundary of B^\square . All of the results described in Section 3 regarding \mathbf{g} apply to \mathbf{b} , including the construction of the homotopy from X^\square to the image of \star -multiplication by \mathbf{b} .

References

- [Ber90] Vladimir Berkovich. *Spectral theory and analytic geometry over non-Archimedean fields*. Vol. 33. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1990, pp. x+169. ISBN: 0-8218-1534-2. DOI: [10.1090/surv/033](https://doi.org/10.1090/surv/033). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1090/surv/033>.
- [BGR84] Siegfried Bosch, Ulrich Güntzer, and Reinhold Remmert. *Non-Archimedean analysis*. Vol. 261. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. A systematic approach to rigid analytic geometry. Springer-Verlag, Berlin, 1984, pp. xii+436. ISBN: 3-540-12546-9. DOI: [10.1007/978-3-642-52229-1](https://doi.org/10.1007/978-3-642-52229-1). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/978-3-642-52229-1>.
- [BP87] Michel Brion and Franz Pauer. “Valuations des espaces homogènes sphériques”. In: *Comment. Math. Helv.* 62.2 (1987), pp. 265–285. ISSN: 0010-2571. DOI: [10.1007/BF02564447](https://doi.org/10.1007/BF02564447).
- [CLS11] David Cox, John Little, and Henry Schenck. *Toric varieties*. Vol. 124. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011, pp. xxiv+841. ISBN: 978-0-8218-4819-7. DOI: [10.1090/gsm/124](https://doi.org/10.1090/gsm/124). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1090/gsm/124>.
- [Ful93] William Fulton. *Introduction to toric varieties*. Vol. 131. Annals of Mathematics Studies. The William H. Roever Lectures in Geometry. Princeton University Press, Princeton, NJ, 1993, pp. xii+157. ISBN: 0-691-00049-2. DOI: [10.1515/9781400882526](https://doi.org/10.1515/9781400882526). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1515/9781400882526>.
- [GP11] Philippe Gille and Patrick Polo, eds. *Schémas en groupes (SGA 3). Tome III. Structure des schémas en groupes réductifs*. Vol. 8. Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by Michel Demazure and Alexander Grothendieck with the collaboration of Michael Artin, Bertin, Jean-Étienne Bertin, Pierre Gabriel, Michèle Raynaud and Jean-Pierre Serre, Revised and annotated edition of the 1970 French original. Société Mathématique de France, Paris, 2011, pp. lvi+337. ISBN: 978-2-85629-324-9.
- [Gru66] Laurent Gruson. “Théorie de Fredholm p -adique”. In: *Bull. Soc. Math. France* 94 (1966), pp. 67–95. ISSN: 0037-9484. URL: http://www.numdam.org/item?id=BSMF_1966__94__67_0.
- [KM19] Kiumars Kaveh and Christopher Manon. “Gröbner theory and tropical geometry on spherical varieties”. In: *Transform. Groups* 24.4 (2019), pp. 1095–1145. ISSN: 1083-4362. DOI: [10.1007/s00031-019-09536-5](https://doi.org/10.1007/s00031-019-09536-5). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/s00031-019-09536-5>.
- [Kno91] Friedrich Knop. “The Luna-Vust theory of spherical embeddings”. In: *Proc. of the Hyderabad Conf. on Algebraic Groups*, Manoj Prakashan. 1991, pp. 225–249.
- [LV83] Domingo Luna and Thierry Vust. “Plongements d’espaces homogènes”. In: *Comment. Math. Helv.* 58.2 (1983), pp. 186–245. ISSN: 0010-2571. DOI: [10.1007/BF02564633](https://doi.org/10.1007/BF02564633). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/BF02564633>.
- [Mil17] James Milne. *Algebraic groups*. Vol. 170. Cambridge Studies in Advanced Mathematics. The theory of group schemes of finite type over a field. Cambridge University Press, Cambridge, 2017, pp. xvi+644. ISBN: 978-1-107-16748-3. DOI: [10.1017/9781316711736](https://doi.org/10.1017/9781316711736). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1017/9781316711736>.
- [MS15] Diane Maclagan and Bernd Sturmfels. *Introduction to tropical geometry*. Vol. 161. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2015, pp. xii+363. ISBN: 978-0-8218-5198-2. DOI: [10.1090/gsm/161](https://doi.org/10.1090/gsm/161). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1090/gsm/161>.
- [Nas17] Evan Nash. *Tropicalizing Spherical Embeddings*. 2017. arXiv: [1609.07455](https://arxiv.org/abs/1609.07455) [math.AG].
- [Nas20] Evan Nash. “Global spherical tropicalization via toric embeddings”. In: *Math. Z.* 294.1-2 (2020), pp. 615–637. ISSN: 0025-5874. DOI: [10.1007/s00209-019-02284-y](https://doi.org/10.1007/s00209-019-02284-y). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/s00209-019-02284-y>.
- [Pas] Boris Pasquier. *Introduction to spherical varieties and description of special classes of spherical varieties*. <http://www-math.sp2mi.univ-poitiers.fr/~bpasquie/KIAS.pdf>.

- [Pay09] Sam Payne. “Analytification is the limit of all tropicalizations”. In: *Math. Res. Lett.* 16.3 (2009), pp. 543–556. ISSN: 1073-2780. DOI: [10.4310/MRL.2009.v16.n3.a13](https://doi.org/10.4310/MRL.2009.v16.n3.a13). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.4310/MRL.2009.v16.n3.a13>.
- [Poi13] Jérôme Poineau. “Les espaces de Berkovich sont angéliques”. In: *Bull. Soc. Math. France* 141.2 (2013), pp. 267–297. ISSN: 0037-9484. DOI: [10.24033/bsmf.2648](https://doi.org/10.24033/bsmf.2648). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.24033/bsmf.2648>.
- [Sum74] Hideyasu Sumihiro. “Equivariant completion”. In: *Journal of Mathematics of Kyoto University* 14.1 (1974), pp. 1–28. DOI: [10.1215/kjm/1250523277](https://doi.org/10.1215/kjm/1250523277). URL: <https://doi.org/10.1215/kjm/1250523277>.
- [Thu07] Amaury Thuillier. “Géométrie toroïdale et géométrie analytique non archimédienne. Application au type d’homotopie de certains schémas formels”. In: *Manuscripta Math.* 123.4 (2007), pp. 381–451. ISSN: 0025-2611. DOI: [10.1007/s00229-007-0094-2](https://doi.org/10.1007/s00229-007-0094-2). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/s00229-007-0094-2>.
- [Tim11] Dmitry Timashev. *Homogeneous spaces and equivariant embeddings*. Vol. 138. Encyclopaedia of Mathematical Sciences. Invariant Theory and Algebraic Transformation Groups, 8. Springer, Heidelberg, 2011, pp. xxii+253. ISBN: 978-3-642-18398-0. DOI: [10.1007/978-3-642-18399-7](https://doi.org/10.1007/978-3-642-18399-7). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/978-3-642-18399-7>.
- [TV21] Jenia Tevelev and Tassos Voggiannou. “Spherical tropicalization”. In: *Transform. Groups* 26.2 (2021), pp. 691–718. ISSN: 1083-4362. DOI: [10.1007/s00031-021-09641-4](https://doi.org/10.1007/s00031-021-09641-4). URL: <https://doi-org.ezproxy.lib.utexas.edu/10.1007/s00031-021-09641-4>.
- [Vus90] Thierry Vust. “Plongements d’espaces symétriques algébriques: une classification”. In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 17.2 (1990), pp. 165–195. ISSN: 0391-173X. URL: http://www.numdam.org/item?id=ASNSP_1990__17_2_165_0.