

PRETRIANGULATED 2-REPRESENTATIONS VIA DG ALGEBRA 1-MORPHISMS

ROBERT LAUGWITZ AND VANESSA MIEMIETZ

ABSTRACT. This paper develops a theory of pretriangulated 2-representations of dg 2-categories. We characterize cyclic pretriangulated 2-representations, under certain compactness assumptions, in terms of modules over dg algebra 1-morphisms internal to associated dg 2-categories of compact dg modules. Further, we investigate the Morita theory and quasi-equivalences for such dg 2-representations. We relate this theory to various classes of examples of dg categorifications from the literature.

CONTENTS

1. Introduction	2
Acknowledgements	4
2. Generalities	5
2.1. Dg categories	5
2.2. pretriangulated categories	5
2.3. Compact dg modules	7
2.4. Compactness and adjunctions	9
2.5. The homotopy category of compact semi-free modules	11
3. Dg 2-categories and 2-representations	13
3.1. Dg 2-categories	13
3.2. Dg 2-representations	15
3.3. Compact pretriangulated 2-representations	18
3.4. Dg ideals of 2-representations and dg 2-subrepresentations	20
3.5. Cyclic and quotient-simple dg 2-representations	20
3.6. Homotopy 2-representations	21
4. Dg algebra 1-morphisms	22
4.1. Compact modules over dg algebra 1-morphisms	22
4.2. The algebra structure on internal homs $[X, X]$	24
4.3. An equivalence of dg 2-representations	25
4.4. Morphisms of dg 2-representations and dg algebra 1-morphisms	28

Date: May 23, 2022.

2020 Mathematics Subject Classification. 18N10, 18N25, 16E45.

Key words and phrases. 2-category, 2-representation, differential graded category, pretriangulated category, homotopy category.

4.5. Morita equivalence	30
4.6. Morita quasi-equivalence	31
4.7. Restriction and pushforward	32
4.8. Local quasi-equivalences and dg 2-representations	34
5. Examples	37
5.1. The dg 2-category \mathcal{C}_A	37
5.2. Pretriangulated hulls of finitary 2-categories and 2-representations	39
5.3. Pretriangulated hulls of finite multitensor categories	39
5.4. Categorification of $\mathbb{Z}[i]$	39
5.5. Braid group categorification	40
References	41

1. INTRODUCTION

Categorification aims to lift important algebraic structures to a higher categorical level, where elements are replaced by functors (or 1-morphisms) and equations are upgraded to natural isomorphisms (or 2-morphisms). The underlying classical structure can then be recovered through a process of passing to Grothendieck groups. Categorifications of Hecke algebras, through Soergel bimodules, or quantum groups have been among the most celebrated achievements of the field and featured in solutions to long-standing conjectures in representation theory, see e.g. [CR, EW, W]. A natural framework for categorification is that of 2-categories containing monoidal or tensor categories as the one-object case.

Studying categorical representations was proposed in [Ro3]. A systematic theory for categorical representations of tensor categories (such as fusion categories) was developed by Etingof–Ostrik and others [EGNO]. These categories are usually abelian while several categorifications are based on non-abelian additive categories. A systematic theory of 2-representations tailored to the latter, categorifying in some sense the theory of representations of finite-dimensional algebras, has been developed in a programme starting with [MM1], see also [MM2, MM3, MM5, MMT, MMMZ] among others.

Many of the important categorifications, such as categorified braid groups or categorical braid group actions [ST, KS, Ro1, Ro2], only emerge when passing from additive categories to triangulated categories. This is manifest, for example, in the fact that braid group relations only become isomorphisms on the level of homotopy or derived categories. This observation motivates the need for a theory of 2-representations that allows working with homotopy categories. In fact, the literature contains several constructions of categorifications of 2-representations of specific categorified algebras on the level of triangulated categories, including, e.g., [BFK, RZ, Ca, CL, Sa, JY] to list a few examples.

Working directly with triangulated categories involves technical obstacles to a formal treatment of 2-representations. For a formal theory of 2-representations, it is important to account for coherence conditions of the structural isomorphisms involved. The definition of triangulated categories, in addition to being very involved, lacks fundamental properties such as functoriality of cones. A solution to these technical issues was

provided by Bondal–Kapranov [BK] by working with *pretriangulated* categories. Pretriangulated categories are *differentially graded (dg)* categories closed under taking cones and shifts, thus ensuring that the associated homotopy or derived categories are triangulated. In this case, the dg category is an enhancement of the triangulated category [LO]. This approach solves key technical issues associated with triangulated categories such as functoriality of cones, taking duals and tensor products [LO, BLL]. Dg enhancements have been used in numerous constructions in categorification, for example, in the theory of spherical twists underlying categorical braid group actions [AL].

The present paper proposes a theory of 2-representations suitable for working with pretriangulated categories. We extend the setup used in [MM1, MM2, MMT] to working with pretriangulated categories. In particular, this setup allows to consider pretriangulated hulls of finitary 2-categories, \mathbb{Z} -gradings, and relaxes the strict finiteness requirements on spaces of 2-morphisms for some of the constructions of finitary 2-representation theory. This first paper in this direction focuses on representing cyclic 2-representations through modules over dg algebra 1-morphisms generalizing results of [EGNO] and [MMT].

The theory of pretriangulated 2-representations proposed in this paper applies to several classes of 2-categories, some of which we start to explore here.

- Pretriangulated hulls of finitary 2-categories. This allows us to consider 2-representations acting on (bounded) complexes of (projective) modules over finite-dimensional \mathbb{k} -algebras. These 2-representations descend to triangulated categories through acting on the associated homotopy categories.
- Categorifications involving dg 2-categories such as [Ti1, Ti2, Ti3, KT]. As a first example, we explore the categorification of $\mathbb{Z}[\sqrt{-1}]$ and its natural action on $\mathbb{Z}^{\oplus 2}$ of [Ti1] in detail in Section 5.4.
- Dg 2-categories \mathcal{C}_A build from projective bimodules over finite-dimensional dg \mathbb{k} -algebras. In finitary 2-representation theory, these 2-categories can be used to classify simple transitive 2-representation of categorified finite-dimensional Lie algebras, see [MM5, Section 7.2]. The first example of such dg 2-categories is given by $\mathcal{C}_{\mathbb{k}}$, the 2-category of bounded complexes of finite-dimensional \mathbb{k} -vector spaces. Using results of Orlov [Or2], we prove in Proposition 5.3 that $\mathcal{C}_{\mathbb{k}}$ has a unique non-acyclic quotient-simple pretriangulated 2-representation.
- Finally, we explain how the theory developed in this paper can be applied to categorical braid group actions of [KS, Ro1, Ro2] in Section 5.5. We relate 2-representations of categorified braid groups in the approach of [KS] to dg 2-representations of categories \mathcal{C}_A , where A is a zigzag algebra with trivial differential. More generally, categorified braid group representations of [Ro1] are related to pretriangulated 2-representations of pretriangulated hulls of 2-categories of Soergel bimodules.

Given a dg 2-category \mathcal{C} , we define the dg 2-category of *pretriangulated 2-representations* $\mathcal{C}\text{-mod}^{\text{pre}}$ in Section 3.2. Given a pretriangulated 2-representation \mathbf{M} , one obtains the *homotopy 2-representation* \mathbf{KM} by passing to the associated homotopy categories $\mathbf{KM}(\mathbf{i})$ at each object \mathbf{i} of \mathcal{C} , see Section 3.6.

The main results of this paper involve an assignment of internal dg algebra 1-morphisms to pretriangulated 2-representations which admit a generator under the action of the dg 2-category \mathcal{C} and taking thick closures. We obtain the following results:

- We define the *internal dg 2-representation* \mathbf{M}_A consisting of certain modules over an internal dg algebra 1-morphism A which lives in the dg 2-category $\vec{\mathcal{C}}$ of compact dg modules introduced in this paper. This construction of $\vec{\mathcal{C}}$ provides a generalization and dg enhancement of the completion of a finitary additive 2-category under cokernels.
- In Corollary 4.11 we prove that if X is a \mathcal{C} -generator of a *compact* pretriangulated 2-representation \mathbf{M} , then the dg idempotent completion \mathbf{M}° is dg equivalent to the internal dg 2-representation \mathbf{M}_{A_X} , where $A_X = [X, X]$ is the internal endomorphism dg algebra 1-morphism of X .
A version of this result, where X is only a \mathcal{C} -quasi-generator, i.e., X generates the associated homotopy 2-representation \mathbf{KM} , is given in Corollary 4.22.
- We define a possible notion of “simple” dg 2-representations of dg 2-categories called *quotient-simple pretriangulated 2-representations*. We show in Corollary 4.18 that a cyclic pretriangulated 2-representation \mathbf{M} is quotient-simple if and only if A_X is simple as a dg algebra 1-morphism.
- Using results by Keller [Ke1], we characterize equivalences (and quasi-equivalences) between internal dg 2-representations \mathbf{M}_A and \mathbf{M}_B over \mathcal{C} in terms of Morita equivalence, see Sections 4.5 and 4.6.
- In Section 4.8 we explain how a dg 2-functor $\mathbf{F}: \mathcal{C} \rightarrow \mathcal{D}$ which is a local quasi-equivalence and essentially surjective on objects induces a correspondence between internal dg 2-representations of \mathcal{C} and \mathcal{D} up to quasi-equivalence, see Proposition 4.26, 4.27, and 4.29 as well as Corollary 4.30 for precise statements.

The definition of quotient-simple pretriangulated 2-representations was inspired by an extension of the theory of cell 2-representations of [MM1] to the pretriangulated setup that will appear in a forthcoming paper in preparation.

The results of this paper refine and extend some of the constructions given in the setup of p -dg 2-representations from [LM]. It would be possible to adapt the constructions of algebra 1-morphisms of the present paper to the p -dg setup (for $\text{char } \mathbb{k} = p$) in order to be applied to the categorifications of quantum groups at p -th roots of unity of [EQ1, EQ2]. Further, the setup presented here could be adapted to incorporate $\mathbb{Z}/2\mathbb{Z}$ -gradings as used, e.g., in [EIQ, EL].

The paper is organized as follows. Section 2 introduces the technical constructions for dg categories required in this paper, including a description of compact dg modules as a dg enriched generalization of the projective abelianization of a finitary category of [Fr] by adding cokernels. Next, Section 3 introduces dg 2-categories and pretriangulated 2-representations, including the concepts of *cyclic* and *quotient-simple* pretriangulated 2-representations and explains how pretriangulated 2-representations induce triangulated homotopy 2-representations. The main results of the paper are found in Section 4 revolving around associating dg algebra 1-morphisms to pretriangulated 2-representations. Finally, Section 5 discusses various classes of examples.

Acknowledgements. The authors would like to thank Ben Elias, Gustavo Jasso, and Marco Mackaay for helpful conversations related to this work. V. M. is supported by EPSRC grant EP/S017216/1 and R. L. is supported by a Nottingham Research Fellowship.

2. GENERALITIES

2.1. Dg categories. We denote by $\mathbb{k}\text{-mod}^{\text{dg}}$ the category of **dg \mathbb{k} -modules**. Its objects are \mathbb{Z} -graded \mathbb{k} -vector spaces equipped with a differential ∂ of degree $+1$ such that $\partial^2 = 0$. A morphism in $f: V \rightarrow W$ in $\mathbb{k}\text{-mod}^{\text{dg}}$ of degree n is a \mathbb{k} -linear map satisfying $f(V^k) \subset W^{k+n}$ on the k -th graded piece. The category $\mathbb{k}\text{-mod}^{\text{dg}}$ is symmetric monoidal (see e.g. [Ke2, Section 2.1]) and enriched over itself. It is equipped with a **shift functor** defined by $(V\langle 1 \rangle)^k = V^{k+1}$ on the k -th graded piece.

A **dg category** \mathcal{C} is a category enriched over the category $\mathbb{k}\text{-mod}^{\text{dg}}$. We assume that all dg categories considered are small.

Given a dg category, we denote by $\mathcal{Z}(\mathcal{C})$ the additive category of **dg morphisms** given by the same objects as \mathcal{C} with morphisms given by the \mathbb{k} -vector spaces

$$\text{Hom}_{\mathcal{Z}(\mathcal{C})}(X, Y) = \{f \in \text{Hom}_{\mathcal{C}}(X, Y) \mid \deg f = 0 \text{ and } \partial f = 0\}.$$

Let \mathcal{C} be a dg category. We define the dg category $\mathcal{C}\text{-mod}^{\text{dg}}$ of **dg modules over \mathcal{C}** to be the dg category of dg functors $\mathcal{C}^{\text{op}} \rightarrow \mathbb{k}\text{-mod}^{\text{dg}}$ (cf. [Ke2, Section 1.2], [Or1, Section 2.2]). Given a dg category \mathcal{C} , the Yoneda lemma gives a fully faithful dg functor

$$\mathcal{C} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}, \quad X \mapsto X^\vee := \text{Hom}_{\mathcal{C}}(-, X).$$

We refer to X^\vee as a **free dg \mathcal{C} -module**. It has the property that given any dg module M over \mathcal{C} , there is a dg isomorphism

$$\text{Hom}_{\mathcal{C}\text{-mod}^{\text{dg}}}(X^\vee, M) \cong M(X).$$

We require the following notion of idempotent completion to obtain dg categories closed under taking dg direct summands. Given a dg category \mathcal{C} , the **dg idempotent completion** \mathcal{C}° is defined as having objects X_e for any dg idempotent $e \in \text{End}_{\mathcal{C}}(X)$, i.e., $\partial(e) = 0$, and morphism spaces

$$\text{Hom}_{\mathcal{C}^\circ}(X_e, Y_f) = f \text{Hom}_{\mathcal{C}}(X, Y) e.$$

Then \mathcal{C}° is a dg category and the embedding $\mathcal{C} \hookrightarrow \mathcal{C}^\circ$ is a dg functor. Furthermore, $(\mathcal{C}^\circ)^\circ$ is dg equivalent to \mathcal{C}° .

Given a dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$, we obtain an induced dg functor

$$(2.1) \quad F^\circ: \mathcal{C}^\circ \rightarrow \mathcal{D}^\circ, \quad F(X_e) = F(X)_{F(e)}.$$

Further, given a natural transformation of dg functors $\phi: F \rightarrow G$,

$$(2.2) \quad \phi_{X_e}^\circ := G(e) \circ \phi_X \circ F(e)$$

defines a natural transformation $\phi^\circ: F^\circ \rightarrow G^\circ$.

2.2. pretriangulated categories. A dg category \mathcal{C} is **pretriangulated** if its Yoneda embedding into $\mathcal{C}\text{-mod}^{\text{dg}}$ via $X \mapsto X^\vee = \text{Hom}_{\mathcal{C}}(-, X)$ is closed under shifts and cones. Note that we do not require \mathcal{C} to be closed under dg direct summands. We say that a full dg subcategory $\mathcal{D} \subseteq \mathcal{C}$ is a **pretriangulated subcategory** if it is closed under shifts, cones and dg isomorphisms, and say that \mathcal{D} is **thick** if it is, in addition, closed under all direct summands which exist in \mathcal{C} .

Let \mathcal{C} be a pretriangulated category. Given a pretriangulated category \mathcal{C} , and a set X of objects in \mathcal{C} , we define the **thick closure** of X in \mathcal{C} to be the full subcategory \widehat{X} of \mathcal{C} generated by all objects in X under shifts, cones, direct sums, dg direct summands, and closed under dg isomorphisms. That is, \widehat{X} is the smallest thick subcategory of \mathcal{C}

containing X . An object X in \mathcal{C} is a **(classical) generator** for \mathcal{C} if the thick closure $\widehat{\{X\}}$ of X is all of \mathcal{C} .

Given a dg category \mathcal{C} , we denote by $\overline{\mathcal{C}}$ the dg category of (one-sided) twisted complexes in \mathcal{C} . Explicitly, we define $\overline{\mathcal{C}}$ as the dg category whose

- objects are given by pairs $(\bigoplus_{m=1}^s F_m, \alpha = (\alpha_{k,l})_{k,l})$ where the F_m are shifts of objects in \mathcal{C} and $\alpha_{k,l} \in \text{Hom}_{\mathcal{C}}(F_l, F_k)$, $\alpha_{k,l} = 0$ for all $k \geq l$ such that the matrix $\partial \cdot \mathbb{I}_s + ((\alpha_{kl})_{k,l})_{kl}$ acts as a differential on $\bigoplus_{m=1}^s F_m^\vee$ in $\mathcal{C}\text{-mod}^{\text{dg}}$ (here \mathbb{I}_s is the identity matrix);
- morphisms are matrices of morphisms between the corresponding objects, with the differential of a homogeneous morphism

$$\gamma = (\gamma_{n,m})_{n,m}: \left(\bigoplus_{m=1}^s F_m, \alpha = (\alpha_{k,l})_{k,l} \right) \longrightarrow \left(\bigoplus_{n=1}^t G_n, \beta = (\beta_{k,l})_{k,l} \right)$$

defined as

$$\partial((\gamma_{n,m})_{n,m}) := (\partial\gamma_{n,m} + (\beta\gamma)_{n,m} - (-1)^{\deg \gamma} (\gamma\alpha)_{n,m})_{n,m}.$$

The notation $\bigoplus_i X_i$ denotes an ordered list of objects rather than a direct sum internal to \mathcal{C} . We give $\overline{\mathcal{C}}$ the explicit additive structure

$$(X, \alpha) \oplus (Y, \beta) := \left(X \oplus Y, \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \right),$$

where $X \oplus Y$ is the concatenation of ordered lists of objects. This additive structure on $\overline{\mathcal{C}}$ is strict, in the sense that $(X \oplus Y) \oplus Z = X \oplus (Y \oplus Z)$.

Note that $\overline{\mathcal{C}}$ is a pretriangulated category, and the smallest pretriangulated category containing \mathcal{C} , cf. [LO, Section 1], [AL, Section 3.2].

Let $f: X \rightarrow Y$ be a dg morphism in $\overline{\mathcal{C}}$, where $X = (\bigoplus_{i=1}^t X_i, \alpha)$, $Y = (\bigoplus_{i=1}^t Y_i, \beta)$. The **cone** C_f of f is the object

$$C_f = \left(Y \oplus X\langle 1 \rangle, \begin{pmatrix} \beta & -f \\ 0 & \alpha \end{pmatrix} \right).$$

The cone C_f comes equipped with the dg morphisms

$$C_f\langle -1 \rangle \rightarrow X, \quad Y \rightarrow C_f,$$

such that pre- (respectively, post-) composition with f yields a null-homotopic morphism.

Given a dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$, we obtain an induced dg functor

$$(2.3) \quad \overline{F}: \overline{\mathcal{C}} \rightarrow \overline{\mathcal{D}}, \quad \overline{F}\left(\bigoplus_{i=1}^t X_i, \alpha\right) = \left(\bigoplus_{i=1}^t F(X_i), F(\alpha)\right),$$

by applying F component-wise to α and to morphisms in $\overline{\mathcal{C}}$. It follows directly that $\overline{G \circ F} = \overline{G} \circ \overline{F}$ for compatible dg functors $F: \mathcal{C} \rightarrow \mathcal{D}$, $G: \mathcal{D} \rightarrow \mathcal{E}$. Moreover, a natural transformation $\tau: F \rightarrow G$ induces a natural transformation $\overline{\tau}: \overline{F} \rightarrow \overline{G}$ using the diagonal matrix

$$(2.4) \quad \tau_{(\bigoplus_{i=1}^t X_i, \alpha)} = \text{diag}(\tau_{X_1}, \dots, \tau_{X_t}).$$

This way, we obtain a dg 2-functor $\overline{(-)}$ that sends dg categories \mathcal{C} to pretriangulated categories $\overline{\mathcal{C}}$.

For future reference, we record the following lemmas.

Lemma 2.1. *Let \mathcal{C} be a full subcategory of \mathcal{C}' , and \mathcal{I} a dg ideal in \mathcal{C} . Then the restriction to \mathcal{C} of the dg ideal generated by \mathcal{I} in \mathcal{C}' is equal to \mathcal{I} .*

Lemma 2.2. *Suppose \mathcal{C} is a dg category equivalent to $\widehat{\{X\}}$ for some $X \in \mathcal{C}$ and let \mathcal{I} be a dg ideal in \mathcal{C} . Then \mathcal{I} is completely determined on $\widehat{\{X\}}$. Thus, the subset $\mathcal{I} \cap \text{End}_{\mathcal{C}}(X)$ generates \mathcal{I} .*

Proof. Let $Y, Z \in \widehat{\{X\}}$. Then there exist objects $Y', Z' \in \widehat{\{X\}}$ and dg idempotents $e_Y \in \text{End}_{\widehat{\{X\}}}(Y')$ and $e_Z \in \text{End}_{\widehat{\{X\}}}(Z')$ such that $\text{Hom}_{\widehat{\{X\}}}(Y, Z) = e_Z \text{Hom}_{\widehat{\{X\}}}(Y', Z') e_Y$. We claim that $\text{Hom}_{\mathcal{I}}(Y, Z) \cong e_Z \text{Hom}_{\mathcal{I}}(Y', Z') e_Y$.

Indeed, the inclusion $\text{End}_{\mathcal{I}}(Y, Z) \supseteq e_Z \text{End}_{\mathcal{I}}(Y', Z') e_Y$ is trivial from \mathcal{I} being an ideal. An element of $\text{Hom}_{\mathcal{I}}(Y, Z)$ is of the form $e_Z \circ f \circ e_Y = e_Z \circ (e_Z \circ f \circ e_Y) \circ e_Y$ and $f = e_Z \circ f \circ e_Y \in \text{Hom}_{\mathcal{I}}(Y', Z')$, so the inclusion $\text{End}_{\mathcal{I}}(Y, Z) \subseteq e_Z \text{End}_{\mathcal{I}}(Y', Z') e_Y$ also holds. This proves the lemma. \square

Given a dg ideal \mathcal{I} in \mathcal{C} we denote by $\overline{\mathcal{I}}$ the dg ideal of morphisms in $\overline{\mathcal{C}}$ with components contained in \mathcal{I} .

Lemma 2.3. *Let \mathcal{C} be a dg category and \mathcal{I} a dg ideal in \mathcal{C} . Then there is an equivalence of pretriangulated categories $(\mathcal{C}/\mathcal{I}) \simeq \overline{\mathcal{C}}/\overline{\mathcal{I}}$. Moreover, if \mathcal{J} is a dg ideal in $\overline{\mathcal{C}}$, then $\mathcal{J} = \overline{\mathcal{J}|_{\mathcal{C}}}$, where $\mathcal{J}|_{\mathcal{C}}$ is the restriction of \mathcal{J} to \mathcal{C} .*

Proof. The quotient dg functor $P: \mathcal{C} \rightarrow \mathcal{C}/\mathcal{I}$ induces a dg functor $\overline{P}: \overline{\mathcal{C}} \rightarrow \overline{(\mathcal{C}/\mathcal{I})}$. On morphisms in $\overline{\mathcal{C}}$, this functor is given by component-wise application of π . Thus, the kernel of \overline{P} is precisely the ideal $\overline{\mathcal{I}}$, proving the first claim.

For the second part, denote $\overline{\mathcal{I}} = \overline{\mathcal{J}|_{\mathcal{C}}}$. Let $f = (f_{ij}) \in \mathcal{J}$, then using the injection ι_i and projection π_j , $f_{ij} = \pi_j f \iota_i \in \mathcal{J}$. Thus, $f \in \overline{\mathcal{I}}$. Conversely, $\overline{\mathcal{I}} \subseteq \mathcal{J}$ implies $\overline{\mathcal{I}} \subseteq \overline{\mathcal{J}}$. \square

It follows that if \mathcal{C} has a generator X , then \mathcal{C}/\mathcal{I} has a generator, which is the image of X under the projection functor.

Lemma 2.4. *If \mathcal{C} is a pretriangulated category, then the dg idempotent completion \mathcal{C}° is also pretriangulated and $\iota: \mathcal{C} \rightarrow \mathcal{C}^\circ$ displays \mathcal{C} as a full pretriangulated subcategory. In particular, this implies $\overline{\mathcal{C}^\circ} \simeq \overline{\mathcal{C}}^\circ$ for any dg category \mathcal{C} .*

Proof. It is clear that $X_e \langle 1 \rangle = (X \langle 1 \rangle)_e$. To show that \mathcal{C}° is closed under cones, we observe that for a dg morphism $g = fge: X_e \rightarrow Y_f$ in \mathcal{C}° , the object $Z \begin{pmatrix} f & 0 \\ 0 & e \end{pmatrix}$, for $Z = (Y \oplus X \langle 1 \rangle, \begin{pmatrix} 0 & g \\ 0 & 0 \end{pmatrix})$ is the cone of g in \mathcal{C}° . This shows \mathcal{C}° is pretriangulated. \square

2.3. Compact dg modules. We now give a diagrammatic description of the dg category of compact dg modules. If \mathcal{C} is a dg category, let $\overline{\mathcal{C}}$ denote the category

- whose objects are diagrams of the form $X_1 \xrightarrow{x} X_0$ for $X_0, X_1 \in \overline{\mathcal{C}}$ and x a dg morphism in $\overline{\mathcal{C}}$;

- whose morphism are pairs (ϕ_0, ϕ_1) of morphisms in $\overline{\mathcal{C}}$ producing solid commutative diagrams of the form

$$(2.5) \quad \begin{array}{ccc} X_1 & \xrightarrow{x} & X_0 \\ \downarrow \phi_1 & \dashrightarrow \eta & \downarrow \phi_0 \\ Y_1 & \xrightarrow{y} & Y_0 \end{array}$$

modulo the subgroup generated by diagrams where there exists a morphism η , as indicated by the dashed arrow, such that $\phi_0 = y\eta$;

- the differential of such a pair (ϕ_0, ϕ_1) is simply the componentwise differential in $\overline{\mathcal{C}}$.

We note that $\overline{\mathcal{C}}$ is a full dg subcategory of $\vec{\mathcal{C}}$ by mapping X to $0 \rightarrow X$ and that $\vec{\mathcal{C}}$ is a pretriangulated category, with shifts and cones inherited from $\overline{\mathcal{C}}$.

Any dg functor $F: \overline{\mathcal{C}} \rightarrow \overline{\mathcal{D}}$ extends to a dg functor $\vec{F}: \vec{\mathcal{C}} \rightarrow \vec{\mathcal{D}}$ component-wise, that is,

$$\vec{F}(X_1 \xrightarrow{x} X_0) = F(X_1) \xrightarrow{F(x)} F(X_0).$$

In particular, consider the Yoneda embedding

$$Y: \mathcal{C} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}, \quad X \mapsto X^\vee = \text{Hom}_{\mathcal{C}}(-, X).$$

This dg functor extends to a dg functor $Y: \vec{\mathcal{C}} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}$, where an object $X_1 \xrightarrow{x} X_0$ is mapped to the dg module given by the cokernel of the dg morphism $Y(x) = x \circ (-)$. By construction, the functor Y is fully faithful on morphism spaces.

An object X in a dg category \mathcal{C} is **compact** if the *corepresentable* functor

$$\text{Hom}_{\mathcal{C}}(X, -): \mathcal{C} \rightarrow \mathbb{k}\text{-mod}^{\text{dg}}$$

commutes with filtered colimits. More precisely, for a directed set I and a diagram $(X_i)_{i \in I}$ of objects in \mathcal{C} with dg morphisms between them, there is a canonical dg isomorphism

$$\text{Hom}_{\mathcal{C}}(X, \underset{i \in I}{\text{colim}} X_i) \cong \underset{i \in I}{\text{colim}} \text{Hom}_{\mathcal{C}}(X, X_i).$$

provided that the colimit $\underset{i \in I}{\text{colim}} X_i$ exists in \mathcal{C} (cf. [LO, Definition 1.6] for the related concept for triangulated categories). We denote the full dg subcategory of $\mathcal{C}\text{-mod}^{\text{dg}}$ on compact dg modules over \mathcal{C} by $\mathcal{C}\text{-mod}^{\text{dg}, \text{cp}}$.

Lemma 2.5. *Every object in $\mathcal{C}\text{-mod}^{\text{dg}}$ is dg isomorphic to a filtered colimit of objects in the image of $Y: \vec{\mathcal{C}} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}$.*

Proof. Let M be a dg module over \mathcal{C} . By [Ke2, Section 3.1] we find a semi-free dg module N (with a possibly infinite filtration by free dg modules) together with a surjective dg morphism $N \xrightarrow{\pi} M$. Denote $K := \ker \pi$. As N is semi-free, there exists a filtration $0 \subseteq F^1 N \subseteq F^2 N \subseteq \dots$ of N . We denote by \mathcal{S} the set of pairs (i, U) where $i \geq 0$ and U is a dg submodule of $K \cap F^i N$ such that there exists a compact semifree dg \mathcal{C} -module surjecting onto U . It follows that \mathcal{S} is a directed set, where $(i, U) \leq (j, V)$ if and only if $i \leq j$ and $U \subseteq V$. The meet operation is given by the sum of dg submodules. A standard argument now shows that M is dg isomorphic to the filtered colimit over quotients $F^i N/U$, where (i, U) is an index from \mathcal{S} . We note that all quotients $F^i N/U$ are in the image of the functor Y since $F^i N$ is a compact semi-free dg \mathcal{C} -module. \square

Lemma 2.6. *The full subcategory $\mathcal{C}\text{-mod}^{\text{dg, cp}}$ of $\mathcal{C}\text{-mod}^{\text{dg}}$ given by compact objects is dg equivalent to the full subcategory on the image of $\vec{\mathcal{C}}$ under Y .*

Proof. Let X be an object of \mathcal{C} . By the dg Yoneda embedding, we have that $\text{Hom}_{\mathcal{C}\text{-mod}^{\text{dg}}}(X^\vee, M) \cong M(X)$ for any dg \mathcal{C} -module M . This implies directly that X^\vee is compact in $\mathcal{C}\text{-mod}^{\text{dg}}$ as colimits in $\mathcal{C}\text{-mod}^{\text{dg}}$ are computed object-wise. More generally, for any object $(\bigoplus_{i=1}^t X_i, \alpha)$ in $\vec{\mathcal{C}}$, as a \mathbb{k} -linear functor, $Y(\bigoplus_{i=1}^t X_i, \alpha)$ is the direct sum $\bigoplus_{i=1}^t X_i^\vee$, with differential induced by α . Therefore,

$$\text{Hom}_{\mathcal{C}\text{-mod}^{\text{dg}}}(Y(\bigoplus_{i=1}^t X_i, \alpha), M) \cong M(\bigoplus_{i=1}^t X) \cong \bigoplus_{i=1}^t M(X),$$

which also commutes with filtered colimits.

Now assume given an object $X := (X_1 \xrightarrow{x} X_0)$ in $\vec{\mathcal{C}}$. By construction, its image under Y is the cokernel of the dg morphism $Y(X_1) \xrightarrow{Y(x)} Y(X_0)$ in $\mathcal{C}\text{-mod}^{\text{dg}}$. Thus, we have an exact sequence

$$Y(X_1) \xrightarrow{Y(x)} Y(X_0) \rightarrow Y(X) \rightarrow 0$$

in $\mathcal{Z}(\mathcal{C}\text{-mod}^{\text{dg}})$, which is an abelian category. Thus, by the five lemma, using that $Y(X_1)$ and $Y(X_0)$ are compact, $Y(X)$ is also compact.

Conversely, let M be a compact object in $\mathcal{C}\text{-mod}^{\text{dg}}$. By Lemma 2.5, M is dg isomorphic to a filtered colimit $\text{colim}_{i \in I} M_i$, with M_i in the image of Y . Since M is compact, $\text{Hom}_{\mathcal{C}\text{-mod}^{\text{dg}}}(M, M) \cong \text{colim}_{i \in I} \text{Hom}_{\mathcal{C}\text{-mod}^{\text{dg}}}(M, M_i)$. Thus, id_M factors through $M_i \rightarrow M$ for some i via some dg morphism. Therefore, M is a dg direct summand of M_i and hence contained in the image of Y itself. \square

Corollary 2.7. *Let \mathcal{C} be a dg category. Then the category $\mathcal{Z}(\vec{\mathcal{C}})$ has cokernels.*

Proof. By Lemma 2.6, $\mathcal{Z}(\vec{\mathcal{C}})$ is equivalence to the full subcategory on compact objects of the abelian category $\mathcal{Z}(\mathcal{C}\text{-mod}^{\text{dg}})$. As compact objects are closed under cokernels, the claim follows. Explicitly, the cokernel of a morphism (f_1, f_0) from $X_1 \xrightarrow{x} X_0$ to $Y_1 \xrightarrow{y} Y_0$ is given by

$$\begin{array}{ccc} Y_1 & \xrightarrow{y} & Y_0 \\ \downarrow \begin{pmatrix} \text{id} \\ 0 \end{pmatrix} & & \downarrow \text{id} \\ Y_1 \oplus X_0 & \xrightarrow{(y, f_0)} & Y_0. \end{array} \quad \square$$

Recall that, for a dg category \mathcal{C} , a **compact semifree dg \mathcal{C} -module** is a dg functor from \mathcal{C}^{op} to $\mathbb{k}\text{-mod}^{\text{dg}}$ which has a finite filtration by free dg \mathcal{C} -modules, cf. [Or1, Section 2.2] where such dg modules are called *finitely-generated semifree*.

Lemma 2.8 ([Or2, Section 2.3]). *The dg category $\vec{\mathcal{C}}$ is dg equivalent to $\mathcal{C}\text{-csf}$, the dg category of compact semi-free dg \mathcal{C} -modules.*

2.4. Compactness and adjunctions. Let \mathcal{C} and \mathcal{D} be pretriangulated categories. Assume given a dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$. We now establish compactness conditions to ensure that $\vec{F}: \vec{\mathcal{C}} \rightarrow \vec{\mathcal{D}}$, obtained by extending F as in Section 2.3, has a right adjoint $G: \vec{\mathcal{D}} \rightarrow \vec{\mathcal{C}}$.

Recall the Yoneda embedding $Y: \mathcal{C} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}$, $X \mapsto X^\vee = \text{Hom}_{\mathcal{C}}(-, X)$. The dg functor F induces a dg functor

$$F^*: \mathcal{D}\text{-mod}^{\text{dg}} \rightarrow \mathcal{C}\text{-mod}^{\text{dg}}, \quad M \mapsto M \circ F.$$

Given a \mathcal{D} -module M , we refer to F^*M as the **pullback** of M along F . By [Ke1, Section 6.1], the dg functor F^* is right adjoint to the dg functor $F_*: \mathcal{C}\text{-mod}^{\text{dg}} \rightarrow \mathcal{D}\text{-mod}^{\text{dg}}$ defined by $M \mapsto F_*M$, where for $D \in \mathcal{D}$, $F_*M(D)$ is the cokernel of the dg morphism

$$\bigoplus_{C_1, C_2 \in \mathcal{C}} M(C_2) \otimes \text{Hom}_{\mathcal{C}}(C_1, C_2) \otimes \text{Hom}_{\mathcal{D}}(D, FC_1) \xrightarrow{\phi} \bigoplus_C M(C) \otimes \text{Hom}_{\mathcal{D}}(D, FC),$$

$$\phi(m \otimes f \otimes g) = M(f)(m) - m \otimes F(f) \circ g.$$

The dg functor F_* extends F under the Yoneda embedding, i.e. $F_* \circ Y = Y \circ F$. In particular, F_* restricts to the dg functor $\vec{F}: \vec{\mathcal{C}} \rightarrow \vec{\mathcal{D}}$. The following lemma characterizes when this restriction has a right adjoint.

Lemma 2.9. *The functor F^* restricts to a dg functor from $\vec{\mathcal{D}}$ to $\vec{\mathcal{C}}$ if and only if for any object D of \mathcal{D} , the dg \mathcal{C} -module $F^*(D^\vee) = \text{Hom}_{\mathcal{D}}(F(-), D)$ is compact.*

Proof. Using Lemma 2.6, the right adjoint dg functor F^* restricts to a dg functor $R: \vec{\mathcal{D}} \rightarrow \vec{\mathcal{C}}$ if and only if $F^*(Y(D))$ is a compact dg \mathcal{C} -module, for any object $D = (D_1 \xrightarrow{d} D_0)$ in $\vec{\mathcal{D}}$. If the latter condition is satisfied for all $D \in \vec{\mathcal{D}}$, then it holds, in particular, for all $D^\vee = Y(D)$ for $D \in \mathcal{D} \subseteq \vec{\mathcal{D}}$. Conversely, assume that $F^*(D^\vee)$ is a compact dg \mathcal{C} -module for any $D \in \mathcal{D}$. Using a similar argument as in the proof of Lemma 2.6, the dg \mathcal{C} -module

$$F^*\left(Y(D_1 \xrightarrow{d} D_0)\right) = \text{coker}\left(F^*D_1^\vee \xrightarrow{F^*Y(d)} F^*D_0^\vee\right)$$

is compact, as the cokernel of a dg morphism between two compact dg \mathcal{C} -modules, cf. Lemma 2.7. \square

If a dg functor F satisfies the equivalent conditions of Lemma 2.9, we say that F **has a compact right adjoint**. In the presence of generators for \mathcal{C} and \mathcal{D} , we have alternative characterizations of the existence of a right adjoint for \vec{F} , i.e., for F having a compact right adjoint.

If \mathcal{C} has a compact generator X , then by Lemma 2.8, the category $\vec{\mathcal{C}}$ is dg equivalent to that of right compact semi-free modules over the dg algebra $A := \text{End}_{\mathcal{C}}(X)$. Here, the product in A is $fg := f \circ g$.

Lemma 2.10. *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a dg functor.*

- (a) *Assume \mathcal{D} has a generator D . Then \vec{F} has a right adjoint if and only if $F^*(D^\vee) = \text{Hom}_{\mathcal{C}}(F(-), D)$ is compact as a dg \mathcal{C} -module.*
- (b) *Assume \mathcal{D} has a compact generator D , \mathcal{C} has a generator C , and denote by A the dg algebra $\text{End}_{\mathcal{C}}(C)$. Then \vec{F} has a right adjoint if and only if $M_F := \text{Hom}_{\mathcal{C}}(F(C), D)$ is compact as a right dg A -module. The action of A on M_F is given by $m \otimes a := m \circ F(a)$, for $m \in M_F$ and $a \in A$.*

Proof. Part (a) follows from the fact that if $F^*(D^\vee)$ is compact, then for any object X in the thick closure of D inside of $\vec{\mathcal{D}}$, $F^*(X^\vee)$ is compact.

Part (b) follows from Part (a) under use of the dg equivalence between $\mathcal{C}\text{-mod}^{\text{dg}}$ and the category of right dg A -modules, which is given by sending X^\vee to $X^\vee(C) = \text{Hom}_{\mathcal{C}}(C, X)$, for $X \in \mathcal{C}$. \square

2.5. The homotopy category of compact semi-free modules. In this section, we discuss the passage from dg categories to triangulated categories in the setup used for this paper.

Let \mathcal{C} be a dg category. We denote the **homotopy category** of \mathcal{C} by $\mathcal{K}(\mathcal{C})$ and recall that $\mathcal{K}(\mathcal{C})$ has the same objects as \mathcal{C} but morphism spaces are given by 0-th cohomology, i.e.

$$\text{Hom}_{\mathcal{K}(\mathcal{C})}(X, Y) = H^0(\text{Hom}_{\mathcal{C}}(X, Y)).$$

It follows that if \mathcal{C} is pretriangulated, then $\mathcal{K}(\mathcal{C})$ is a triangulated category, see [BK, §1, Proposition 2], [Ke1, Section 2.2]. A dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$ induces a functor $\mathcal{K}(F): \mathcal{K}(\mathcal{C}) \rightarrow \mathcal{K}(\mathcal{D})$. The functor $\mathcal{K}(F)$ is a *triangle functor* (also called exact functor of triangulated categories) [BK, §3]. Moreover, a dg natural transformation $\tau: F \rightarrow G$ between two dg functors descends to a natural transformation $\mathcal{K}(\tau): \mathcal{K}(F) \rightarrow \mathcal{K}(G)$.

We say that a morphism $f: X \rightarrow Y$ is a **homotopy isomorphism** if it descends to an isomorphism in $\mathcal{K}(\mathcal{C})$.

A dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is (part of) a **quasi-equivalence** if it induces quasi-isomorphisms (i.e., isomorphisms in all cohomological degrees) between $\text{Hom}_{\mathcal{C}}(X, Y)$ and $\text{Hom}_{\mathcal{D}}(FX, FY)$ for any pair of objects X, Y in \mathcal{C} and $\mathcal{K}(F)$ is an equivalence of categories [Ke2, Section 2.3]. Note that if \mathcal{C} and \mathcal{D} are both pretriangulated, closure under shifts implies that $\text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}}(FX, FY)$ being a quasi-isomorphism for all $X, Y \in \mathcal{C}$ is equivalent to $\mathcal{K}(F)$ being fully faithful.

Given a dg category \mathcal{C} , a morphism $f: M \rightarrow N$ of dg \mathcal{C} -modules is a **quasi-isomorphism** if $\partial(f) = 0$ and f induces a quasi-isomorphism $f(X): N(X) \rightarrow M(X)$ for each object X of \mathcal{C} . The **derived category** $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})$ is the localization of the category $\mathcal{Z}(\mathcal{C}\text{-mod}^{\text{dg}})$ of dg morphisms (or, equivalently, of $\mathcal{K}(\mathcal{C}\text{-mod}^{\text{dg}})$), see Section 2.1, by the set of quasi-isomorphisms [Ke2, Section 3.2]. The derived category can also be defined as the Verdier quotient of $\mathcal{K}(\mathcal{C}\text{-mod}^{\text{dg}})$ by the full subcategory of acyclic modules. Here, a \mathcal{C} -module M is **acyclic** if the complex $M(X)$ is acyclic for any object X of \mathcal{C} [Or1, Section 2.2].

The full subcategory of **perfect objects** of $\mathcal{C}\text{-mod}^{\text{dg}}$ is given by those dg modules whose images in $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})$ are in the idempotent completion of compact semi-free modules. The full subcategory of perfect objects yields the subcategory $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})^{\text{cp}}$ of compact objects in $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})$ [Ke2, Corollary 3.7]. The following well-known result justifies that we will mostly be working with $\mathcal{K}(\overline{\mathcal{C}})$ as a model for the compact derived category of dg modules over \mathcal{C} .

Proposition 2.11. *The functor $\mathcal{K}(\overline{\mathcal{C}})^\circ \hookrightarrow \mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})^{\text{cp}}$ forms part of an equivalence of triangulated categories.*

Proof. By [Ke2, Corollary 3.7], every compact object in $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})$ is a direct summand of an object in the image of $\mathcal{K}(\overline{\mathcal{C}})$ in the derived category. As the derived category is idempotent complete [BN, Proposition 3.2], this implies that the idempotent completion $\mathcal{K}(\overline{\mathcal{C}})^\circ$ (see [BS]) admits a triangle functor into $\mathcal{D}(\mathcal{C}\text{-mod}^{\text{dg}})^{\text{cp}}$. This functor is fully faithful by [Ke2, Section 3.2] and hence gives an equivalence of categories. \square

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a dg functor and recall the adjoint dg functors

$$F_*: \mathcal{C}\text{-mod}^{\text{dg}} \rightleftarrows \mathcal{D}\text{-mod}^{\text{dg}} : F^*$$

from Section 2.4.

Lemma 2.12. *If $F: \mathcal{C} \rightarrow \mathcal{D}$ is a quasi-equivalence, then so is $\vec{F}: \vec{\mathcal{C}} \rightarrow \vec{\mathcal{D}}$.*

Proof. Recall that if F is a dg functor that has a compact right adjoint (cf. Section 2.4), then both dg functors F^* , F_* restrict to compact objects to a pair of adjoint dg functors

$$F_*: \mathcal{C}\text{-mod}^{\text{dg, cp}} \rightleftarrows \mathcal{D}\text{-mod}^{\text{dg, cp}} : F^*.$$

We first consider the unit η of this adjunction. For a representable dg \mathcal{C} -module X^\vee , we have $F^*F_*(X^\vee) = \text{Hom}_{\mathcal{D}}(F(-), FX)$. Then $\eta_{X^\vee}: \text{Hom}_{\mathcal{C}}(-, X) \rightarrow \text{Hom}_{\mathcal{D}}(F(-), FX)$ is simply given by applying F to morphism spaces, and is a quasi-isomorphism by assumption. By Lemma 2.6, any compact dg \mathcal{C} -module M is dg isomorphic to a cokernel of a morphism between dg modules in the pretriangulated closure of representable dg \mathcal{C} -modules. By right exactness of F_* and exactness of F^* , we have a commutative diagram

$$\begin{array}{ccccc} X_1^\vee & \longrightarrow & X_0^\vee & \longrightarrow & M \\ \eta_{X_1^\vee} \downarrow & & \eta_{X_0^\vee} \downarrow & & \downarrow \eta_M \\ F^*F_*(X_1^\vee) & \longrightarrow & F^*F_*(X_0^\vee) & \longrightarrow & F^*F_*M \end{array}$$

or dg morphisms in $\mathcal{C}\text{-mod}^{\text{dg, cp}}$. Since $\eta_{X_1^\vee}, \eta_{X_0^\vee}$ are quasi-isomorphisms, so is the induced morphism η_M on the cokernel. Thus η is a quasi-isomorphism on all compact dg \mathcal{C} -modules.

Now consider the counit of the adjunction. Let $X \in \mathcal{D}$ and consider the dg \mathcal{D} -module $F_*F^*(X^\vee)$. This is given as the cokernel of

$$\begin{array}{ccc} \bigoplus_{C_1, C_2 \in \mathcal{C}} \text{Hom}_{\mathcal{D}}(FC_2, X) \otimes \text{Hom}_{\mathcal{C}}(C_1, C_2) \otimes \text{Hom}_{\mathcal{D}}(-, FC_1) & & h \otimes f \otimes g \\ \downarrow \phi & & \downarrow \\ \bigoplus_{C \in \mathcal{C}} \text{Hom}_{\mathcal{D}}(FC, X) \otimes \text{Hom}_{\mathcal{D}}(-, FC) & & hF(f) \otimes g - h \otimes F(f)g. \end{array}$$

There is a natural map $\psi: \bigoplus_{C \in \mathcal{C}} \text{Hom}_{\mathcal{D}}(FC, X) \otimes \text{Hom}_{\mathcal{D}}(-, FC) \rightarrow \text{Hom}_{\mathcal{D}}(-, X)$ induced by composition, which annihilates the image of ϕ and hence factors over the cokernel, producing the component $\epsilon_{X^\vee}: F_*F^*(X^\vee) \rightarrow X^\vee$ of the counit.

Next, we show that in the special case $X = FY$, for $Y \in \mathcal{C}$, $\epsilon_{(FY)^\vee}$ is a quasi-isomorphism. In fact, we can compute the cokernel of ϕ by evaluating the maps ϕ_Z and c_Z for any object Z in \mathcal{D} . Note that ψ_Z is surjective, with $\text{id}_Z \otimes f$ being a preimage of $f \in \text{Hom}_{\mathcal{D}}(Z, FY)$. Moreover, if $h \in \text{Hom}_{\mathcal{D}}(FC, FY)$, $g \in \text{Hom}_{\mathcal{D}}(Z, FC)$ are dg morphisms such that $\psi_Z(h \otimes g) = hg = 0$, there exists a morphism $f: \mathcal{C} \rightarrow Y$ such that $F(f) = h$ in $H^0(\text{Hom}_{\mathcal{D}}(FC, FY))$, i.e. $F(f) = h + \partial(a)$ for some morphism a .

Thus, we compute

$$\begin{aligned}
\phi(\text{id} \otimes f \otimes g) &= F(f) \otimes g - \text{id} \otimes F(f)g \\
&= (h + \partial(a)) \otimes g - \text{id} \otimes (h + \partial(a))g \\
&= h \otimes g + \partial(a) \otimes g - \text{id} \otimes hg - \text{id} \otimes \partial(a)g \\
&= h \otimes g + \partial(a) \otimes g - \text{id} \otimes \partial(a)g \\
&= h \otimes g + \partial(a \otimes g - \text{id} \otimes ag),
\end{aligned}$$

where in the last step we use that $\partial(g) = 0$. This shows that, after passing to cohomology, $\ker \phi_Z = \text{Im } \psi_Z$ and hence the cokernel of ϕ is given by the representable functor associated to FY . In other words, $\epsilon_{(FY)^\vee}$ gives an isomorphism in $\mathcal{K}(\mathcal{D}\text{-mod}^{\text{dg, cp}})$.

Now consider a general compact dg module $X_1^\vee \xrightarrow{x} X_0^\vee \rightarrow M$, with X_1, X_0 in $\overline{\mathcal{D}}$. Then by right exactness of F_* and exactness of F^* , we have a commutative diagram

$$\begin{array}{ccccc}
F_*F^*(X_1^\vee) & \longrightarrow & F_*F^*(X_0^\vee) & \twoheadrightarrow & F_*F^*M \\
\epsilon_{X_1^\vee} \downarrow & & \downarrow \epsilon_{X_0^\vee} & & \downarrow \epsilon_M \\
X_1^\vee & \longrightarrow & X_0^\vee & \twoheadrightarrow & M
\end{array}$$

of dg morphisms in $\mathcal{D}\text{-mod}^{\text{dg, cp}}$. Arguing similarly to before, the morphism ϵ_M is a quasi-isomorphism as both $\epsilon_{X_1^\vee}$ and $\epsilon_{X_0^\vee}$ are quasi-isomorphisms. Thus ϵ is a quasi-isomorphism on all compact dg \mathcal{D} -modules which shows that $\mathcal{K}(F^*)$ and $\mathcal{K}(F_*)$ are equivalences. Thus, as categories of compact dg modules are closed under shift, F^* , F_* induce quasi-equivalences on compact dg modules as claimed. Since \overline{F} corresponds to F_* under the dg equivalence of Lemma 2.6, it is also a quasi-equivalence. \square

3. DG 2-CATEGORIES AND 2-REPRESENTATIONS

In this section, we collect generalities on pretriangulated 2-categories and define different classes of pretriangulated 2-representations.

3.1. Dg 2-categories. We call a 2-category \mathcal{C} a **dg 2-category** if the categories $\mathcal{C}(i, j)$ are dg categories for any pair of objects $i, j \in \mathcal{C}$, and horizontal composition is a dg functor.

Given a dg 2-category \mathcal{C} , we can associate a new dg 2-category of **(one-sided) twisted complexes**, denoted by $\overline{\mathcal{C}}$. It consists of

- the same objects as \mathcal{C} ;
- 1-morphism categories $\overline{\mathcal{C}(i, j)}$;
- with horizontal composition of two 1-morphisms $X = (\bigoplus_{m=1}^s F_m, \alpha) \in \overline{\mathcal{C}(k, l)}$ and $X' = (\bigoplus_{n=1}^t F'_n, \alpha') \in \overline{\mathcal{C}(j, k)}$ given by

$$(3.1) \quad X \circ X' = \left(\bigoplus_{(m, n)} F_m F'_n, (\delta_{k', l'} \alpha_{k, l} \circ_0 \text{id}_{F'_{k'}} + \delta_{k, l} \text{id}_{F_k} \circ_0 \alpha'_{k', l'})_{(k, k'), (l, l')} \right),$$

where pairs (m, n) are ordered lexicographically;

The above composition of 1-morphism gives a strict operation (for details, see [LM, Proposition 3.5]). We say that a dg 2-category is **pretriangulated** if the embedding $\mathcal{C} \hookrightarrow \overline{\mathcal{C}}$ is part of a dg biequivalence.

We can further extend the dg 2-category $\overline{\mathcal{C}}$ by formally adding dg quotients to obtain a dg 2-category $\vec{\mathcal{C}}$.

Lemma 3.1. *There exists a dg 2-category $\vec{\mathcal{C}}$ containing $\overline{\mathcal{C}}$ as a dg 2-subcategory. A given dg 2-functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ extends naturally to a dg 2-functor $\vec{\Phi}: \vec{\mathcal{C}} \rightarrow \vec{\mathcal{D}}$.*

Proof. We define $\vec{\mathcal{C}}(i, j) := \overline{\mathcal{C}(i, j)}$. The composition in $\vec{\mathcal{C}}$ is defined as

$$(G_1 \xrightarrow{g} G_0)(H_1 \xrightarrow{h} H_0) := (G_1 H_0 \oplus G_0 H_1 \xrightarrow{(g \circ_0 \text{id}_{H_0} \quad \text{id}_{G_0} \circ_0 h)} G_0 H_0),$$

for objects $(G_1 \xrightarrow{g} G_0)$ in $\vec{\mathcal{C}}(j, k)$ and $(H_1 \xrightarrow{h} H_0)$ in $\vec{\mathcal{C}}(i, j)$. Here, we crucially use the explicit additive structure on $\vec{\mathcal{C}}$ making composition of 1-morphisms in $\vec{\mathcal{C}}$ strictly associative.

Given morphisms

$$(\phi_0, \phi_1): (G_1 \xrightarrow{g} G_0) \rightarrow (G'_1 \xrightarrow{g'} G'_0), \quad \text{and} \quad (\psi_0, \psi_1): (H_1 \xrightarrow{h} H_0) \rightarrow (H'_1 \xrightarrow{h'} H'_0)$$

in $\vec{\mathcal{C}}$, we define

$$(\phi_0, \phi_1) \circ_0 (\psi_0, \psi_1) := \left(\phi_0 \circ_0 \psi_0, \begin{pmatrix} \phi_1 \circ_0 \psi_0 & 0 \\ 0 & \phi_0 \circ_0 \psi_1 \end{pmatrix} \right).$$

Note that given a homotopy $\eta: G_0 \rightarrow G'_1$ rendering the morphism (ϕ_0, ϕ_1) zero in $\vec{\mathcal{C}}(j, k)$, the homotopy $\begin{pmatrix} \eta \circ_0 \psi_0 \\ 0 \end{pmatrix}$ shows that the morphism $(\phi_0, \phi_1) \circ_0 (\psi_0, \psi_1)$ is zero in $\vec{\mathcal{C}}(i, k)$. Similarly, if $\epsilon: H_0 \rightarrow H'_1$ is a homotopy rendering the morphism (ψ_0, ψ_1) zero, use the homotopy $\begin{pmatrix} 0 \\ \phi_0 \circ_0 \epsilon \end{pmatrix}$. Thus, horizontal composition in $\vec{\mathcal{C}}$ is well-defined.

Next, we define vertical composition \circ_1 and the differential on morphism spaces component-wise. One directly verifies that this way, $\vec{\mathcal{C}}$ obtains the structure of dg 2-category.

Given a dg 2-functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$, the underlying dg functors $\Phi_{i,j}: \mathcal{C}(i, j) \rightarrow \mathcal{D}(\Phi i, \Phi j)$ extend by component-wise application to give dg functors to give dg functors

$$\vec{\Phi}_{i,j}: \vec{\mathcal{C}}(i, j) \rightarrow \vec{\mathcal{D}}(\Phi i, \Phi j),$$

which assemble into a dg 2-functor as required. \square

Corollary 3.2. *If the dg 2-functor Φ in Lemma 3.1 is a local quasi-equivalence, then so is $\vec{\Phi}$.*

Proof. This follows directly from Lemma 2.12. \square

Definition 3.3. Let \mathcal{C} be a dg 2-category.

- (a) We say that a dg 2-category \mathcal{C} **has generators** if each category $\overline{\mathcal{C}(i, j)}$ has a generator.
- (b) We say that a dg 2-category \mathcal{C} **has compact left adjoints** if for any F in $\overline{\mathcal{C}(i, j)}$, there exists a morphism $*F \in \vec{\mathcal{C}}(j, i)$ together with dg morphisms

$$u_F: \mathbb{1}_i \rightarrow F^*F \in \vec{\mathcal{C}}(i, i), \quad c_F: *FF \rightarrow \mathbb{1}_j \in \vec{\mathcal{C}}(j, j)$$

satisfying the usual identities of an adjunction internal to $\vec{\mathcal{C}}$.

Given a dg 2-category \mathcal{C} , the **dg idempotent completion** \mathcal{C}° is the dg 2-category with the same objects as \mathcal{C} and $\mathcal{C}^\circ(i, j) = \mathcal{C}(i, j)^\circ$, the closure under dg idempotents, for two objects i, j . We note that by Lemma 2.4, if \mathcal{C} is pretriangulated, then \mathcal{C}° is pretriangulated.

To a dg 2-category \mathcal{C} , we can associated the **2-category $\mathcal{Z}\mathcal{C}$ of dg 2-morphisms** which has the same objects and 1-morphisms as \mathcal{C} but only those 2-morphisms which are of degree zero and annihilated by the differential.

The **homotopy 2-category $\mathcal{K}\mathcal{C}$** associated to \mathcal{C} is defined by having the same objects as \mathcal{C} and categories of 1-morphisms given by the homotopy categories

$$\mathcal{K}\mathcal{C}(i, j) := \mathcal{K}(\mathcal{C}(i, j)).$$

If \mathcal{C} is pretriangulated, each $\mathcal{K}\mathcal{C}(i, j)$ is triangulated, and we will hence call $\mathcal{K}\mathcal{C}$ a triangulated 2-category. Note that horizontal composition with 1-morphisms is, in particular, a triangle functor.

3.2. Dg 2-representations. For the purpose of giving targets for dg 2-representations in this section, we define the dg 2-category \mathfrak{M}^{dg} as the 2-category whose

- objects are pretriangulated categories \mathcal{C} ;
- 1-morphisms are dg functors between such categories;
- 2-morphisms are all morphisms of such dg functors.

The dg 2-subcategory $\mathfrak{M}_g^{\text{dg}}$ of \mathfrak{M}^{dg} consists of those dg categories which have a generator.

A **pretriangulated 2-representation** of a dg 2-category \mathcal{C} is a 2-functor $\mathbf{M}: \mathcal{C} \rightarrow \mathfrak{M}^{\text{dg}}$ such that locally the functors from $\mathcal{C}(i, j)$ to $\mathfrak{M}^{\text{dg}}(\mathbf{M}(i), \mathbf{M}(j))$ are dg functors. Explicitly, \mathbf{M} sends

- an object $i \in \mathcal{C}$ to a pretriangulated category $\mathbf{M}(i)$,
- a 1-morphism $G \in \mathcal{C}(i, j)$ to a dg functor $\mathbf{M}(G): \mathbf{M}(i) \rightarrow \mathbf{M}(j)$,
- a 2-morphism $\alpha: G \rightarrow H \in \mathcal{C}(i, j)$ to a morphism of dg functors $\mathbf{M}(\alpha): \mathbf{M}(G) \rightarrow \mathbf{M}(H)$.

A **morphisms of dg 2-representations** $\Phi: \mathbf{M} \rightarrow \mathbf{N}$ consists of

- dg functor $\Phi_i: \mathbf{M}(i) \rightarrow \mathbf{N}(i)$ for each $i \in \mathcal{C}$ and
- natural dg isomorphisms $\eta_F: \Phi_j \circ \mathbf{M}(F) \rightarrow \mathbf{N}(F) \circ \Phi_i$ for each $F \in \mathcal{C}(i, j)$ such that for composable F, G

$$\eta_{FG} = (\text{id}_{\mathbf{N}(F)} \circ_0 \eta_G) \circ_1 (\eta_F \circ_0 \text{id}_{\mathbf{M}(G)}).$$

The collection of pretriangulated dg 2-representations of \mathcal{C} , together with morphisms of dg 2-representations and modifications satisfying the same condition as in [MM3, Section 2.3], form a dg 2-category, which we denote by $\mathcal{C}\text{-mod}^{\text{pre}}$.

We say a pretriangulated 2-representation **has generators** if its target is $\mathfrak{M}_g^{\text{dg}}$, that is, if $\mathbf{M}(i)$ has a generator for any object i . The 1-full and 2-full dg 2-subcategory of $\mathcal{C}\text{-mod}^{\text{pre}}$ consisting of pretriangulated 2-representations with generators is denoted by $\mathcal{C}\text{-mod}_g^{\text{pre}}$.

A pretriangulated 2-representation \mathbf{M} is **acyclic** if, for any object i , any $X \in \mathbf{M}(i)$ is acyclic, i.e. there exists a morphism f such that $\partial(f) = \text{id}_X$.

Note that a pretriangulated 2-representation \mathbf{M} of a dg 2-category \mathcal{C} extends to a pretriangulated 2-representation $\overline{\mathbf{M}}$ of $\overline{\mathcal{C}}$.

Lemma 3.4. *Extending from \mathcal{C} to $\overline{\mathcal{C}}$ and restricting from $\overline{\mathcal{C}}$ to \mathcal{C} define mutually inverse dg biequivalences between $\mathcal{C}\text{-mod}^{\text{pre}}$ and $\overline{\mathcal{C}}\text{-mod}^{\text{pre}}$.*

The following lemma describes how to extend pretriangulated 2-representations from \mathcal{C} to $\overline{\mathcal{C}}$.

Lemma 3.5. *Let \mathcal{C} be a dg 2-category*

- (a) *Given a pretriangulated 2-representation \mathbf{M} of \mathcal{C} , we can define a pretriangulated 2-representation $\overline{\mathbf{M}}$ by $\overline{\mathbf{M}}(\mathbf{i}) = \overline{\mathbf{M}}(\mathbf{i})$. Moreover, $\overline{\mathbf{M}}$ extends to a pretriangulated 2-representation of $\overline{\mathcal{C}}$.*
- (b) *Given a morphism of 2-representations $\Phi: \mathbf{M} \rightarrow \mathbf{N}$ of \mathcal{C} , we obtain a morphism of dg 2-representations $\overline{\Phi}: \overline{\mathbf{M}} \rightarrow \overline{\mathbf{N}}$ of $\overline{\mathcal{C}}$.*
- (c) *Given morphisms of dg 2-representations $\Phi, \Psi: \mathbf{M} \rightarrow \mathbf{N}$, a modification $\theta: \Phi \rightarrow \Psi$ extends to a modification $\overline{\theta}: \overline{\Phi} \rightarrow \overline{\Psi}$. This assignment yields a faithful dg morphism*

$$\text{Hom}_{\text{Hom}_{\mathcal{C}}(\mathbf{M}, \mathbf{N})}(\Phi, \Psi) \rightarrow \text{Hom}_{\text{Hom}_{\overline{\mathcal{C}}}(\overline{\mathbf{M}}, \overline{\mathbf{N}})}(\overline{\Phi}, \overline{\Psi}).$$

Proof. We first prove (a). The fact that $\overline{\mathbf{M}}(\mathbf{i}) := \overline{\mathbf{M}}(\mathbf{i})$ defines a dg 2-representation $\overline{\mathbf{M}}$ of $\overline{\mathcal{C}}$ follows from the observation that for $G \in \mathcal{C}(\mathbf{i}, \mathbf{j})$ and H in $\mathcal{C}(\mathbf{j}, \mathbf{k})$, the induced dg functor satisfy the equality $\overline{\mathbf{M}}(\overline{HG}) = \overline{\mathbf{M}}(\overline{H})\overline{\mathbf{M}}(\overline{G})$, since these dg functors are defined by component-wise application of \mathbf{M} .

Given an object $(G_1 \xrightarrow{g} G_0)$ in $\overline{\mathcal{C}}(\mathbf{i}, \mathbf{j})$, we define

$$\overline{\mathbf{M}}(G_1 \xrightarrow{g} G_0)(X_1 \xrightarrow{x} X_0) = (\overline{\mathbf{M}}(G_1)X_0 \oplus \overline{\mathbf{M}}(G_0)X_1 \xrightarrow{(\overline{\mathbf{M}}(g)(X_0) \ \overline{\mathbf{M}}(G_0)(x))} \overline{\mathbf{M}}(G_0)X_0),$$

for any $(X_1 \xrightarrow{x} X_0)$ in $\overline{\mathbf{M}}(\mathbf{i})$, and extend to morphisms accordingly. It is readily verified that this extends $\overline{\mathbf{M}}$ to a dg 2-representation of $\overline{\mathcal{C}}$ proceeding similarly to the proof of Lemma 3.1. We note that the dg 2-representation is pretriangulated as $\overline{\mathbf{M}}(\mathbf{i})$ is pretriangulated for any object \mathbf{i} , cf. Section 2.3.

To prove (b), we denote by $\eta_G: \Phi_j \circ \mathbf{M}(G) \rightarrow \mathbf{N}(G) \circ \Phi_i$ the dg natural isomorphism given as part of the data of a morphism of dg 2-representation, which is also natural in $G \in \mathcal{C}(\mathbf{i}, \mathbf{j})$. Given an object $(G_1 \xrightarrow{g} G_0)$ in $\overline{\mathcal{C}}(\mathbf{i}, \mathbf{j})$ and $(X_1 \xrightarrow{x} X_0)$ in $\overline{\mathbf{M}}(\mathbf{i})$, we define the dg morphism

$$\left(\overline{\eta}_{(G_1 \xrightarrow{g} G_0)} \right)_{(X_1 \xrightarrow{x} X_0)} : \overline{\Phi}_j \circ \overline{\mathbf{M}}(G_1 \xrightarrow{g} G_0)(X_1 \xrightarrow{x} X_0) \rightarrow \overline{\mathbf{N}}(G_1 \xrightarrow{g} G_0) \circ \overline{\Phi}_i(X_1 \xrightarrow{x} X_0)$$

by the diagram

$$\begin{array}{ccc} \Phi_j \mathbf{M}(G_1)(X_0) \oplus \Phi_j \mathbf{M}(G_0)(X_1) & \xrightarrow{(\Phi_j \mathbf{M}(g)_{X_0} \ \Phi_j \mathbf{M}(G_0)(x))} & \Phi_j \mathbf{M}(G_0)(X_0) \\ \downarrow \left(\begin{array}{cc} (\eta_{G_1})_{X_0} & 0 \\ 0 & (\eta_{G_0})_{X_1} \end{array} \right) & & \downarrow (\eta_{G_0})_{X_0} \\ \mathbf{N}(G_1) \Phi_i(X_0) \oplus \mathbf{N}(G_0) \Phi_i(X_1) & \xrightarrow{(\mathbf{N}(g)_{\Phi_i(X_0)} \ \mathbf{N}(G_0) \Phi_i(x))} & \mathbf{N}(G_0) \Phi_i(X_0). \end{array}$$

This diagram commutes since

$$\mathbf{N}(g)_{\Phi_i(X_0)} \circ (\eta_{G_1})_{X_0} = (\eta_{G_0})_{X_0} \circ \Phi_j \mathbf{M}(g)_{X_0}$$

by naturality of $(\eta_G)_{X_0}$ in G applied to the morphism g , and

$$\mathbf{N}(G_0)_{\Phi_i(x)} \circ (\eta_{G_0})_{X_1} = (\eta_{G_0})_{X_0} \circ \Phi_j \mathbf{M}(G_0)(x)$$

by naturality of η_{G_0} applied to the morphism x .

The morphisms $(\vec{\eta}_{(G_1 \xrightarrow{g} G_0)})_{(X_1 \xrightarrow{x} X_0)}$ thus defined are natural in $(G_1 \xrightarrow{g} G_0)$ and $(X_1 \xrightarrow{x} X_0)$ because all their components are instances of morphisms $(\eta_G)_X$ which are natural in G and X . As the differential in $\vec{\mathbf{N}}(\vec{j})$ is defined component-wise, we see that $\vec{\eta}$ consists of dg morphisms. The equation

$$\eta_{HG} = (\text{id}_{\mathbf{N}(H)} \circ_0 \eta_G) \circ_1 (\eta_H \circ_0 \text{id}_{\mathbf{M}(H)})$$

implies the corresponding equation for $\vec{\eta}$ since it holds component-wise.

We leave the verification of (c) to the reader. \square

For any dg 2-category \mathcal{C} and i one of its objects, we define the **i -th principal dg 2-representation \mathbf{P}_i** , which sends

- an object j to $\overline{\mathcal{C}(i, j)}$,
- a 1-morphism G in $\mathcal{C}(j, k)$ to the dg functor $\overline{\mathcal{C}(i, j)} \rightarrow \overline{\mathcal{C}(i, k)}$ induced by composition with G ,
- a 2-morphism to the induced dg morphism of functors

pretriangulated 2-representations can also be extended to the dg idempotent completion \mathcal{C}° of \mathcal{C} . For this, given $\mathbf{M} \in \mathcal{C}\text{-mod}^{\text{pre}}$, consider $\mathbf{M}^\circ \in \mathcal{C}^\circ\text{-mod}^{\text{pre}}$ given by $\mathbf{M}^\circ(i) = \mathbf{M}(i)^\circ$ and extending the dg functors $\mathbf{M}(F)$ to $\mathbf{M}^\circ(F)$ via (2.1). The dg 2-representation \mathbf{M}° extends to a pretriangulated 2-representation of \mathcal{C}° where F_e , for $e: F \rightarrow F$ an idempotent in $\mathcal{C}(i, j)$, is sent to the dg functor $\mathbf{M}^\circ(F_e): \mathbf{M}^\circ(i) \rightarrow \mathbf{M}^\circ(j)$ defined by

$$X \mapsto \mathbf{M}^\circ(F_e)(X) = (\mathbf{M}(F)(X))_{\mathbf{M}(e)_X}, \quad f \mapsto \mathbf{M}(e)_Y \circ \mathbf{M}(F)(f) \circ \mathbf{M}(e)_X,$$

where X, Y are objects and $f: X \rightarrow Y$ is a morphism in $\mathbf{M}^\circ(i)$. Functoriality of $\mathbf{M}(F_e)$ follows using naturality of $\mathbf{M}(e)$.

Lemma 3.6. *There is a dg 2-functor*

$$(-)^\circ: \mathcal{C}\text{-mod}^{\text{pre}} \rightarrow \mathcal{C}^\circ\text{-mod}^{\text{pre}},$$

where the assignments on dg 2-representations, morphisms, and modifications extend the given structures from \mathcal{C} to \mathcal{C}° .

Proof. A morphism of dg 2-representations $\Phi: \mathbf{M} \rightarrow \mathbf{N}$ naturally extends to a morphism of dg 2-representations $\Phi^\circ: \mathbf{M}^\circ \rightarrow \mathbf{N}^\circ$ of \mathcal{C}° using (2.1), which is a functorial assignment. For a 1-morphism F and a dg idempotent $e: F \rightarrow F$, one defines

$$(\eta_{F_e})_X := \mathbf{N}(e)_{\Phi^\circ(X)} \circ (\eta_F)_X \circ \Phi^\circ \mathbf{M}(e)_X$$

and extends η_{F_e} via (2.2). This way, it follows that Φ° indeed commutes with the action of \mathcal{C}° defined above. Modifications of dg morphisms of dg 2-representations are naturally extended using (2.2). \square

The dg 2-functor $(-)^{\circ}: \mathcal{C}\text{-mod}^{\text{pre}} \rightarrow \mathcal{C}^{\circ}\text{-mod}^{\text{pre}}$ is, in a certain sense, left dg 2-adjoint to restriction $\mathbf{N} \mapsto \mathbf{N}|_{\mathcal{C}}$ along the inclusion of \mathcal{C} into \mathcal{C}° .

Lemma 3.7. *There are dg equivalences*

$$E: \text{Hom}_{\mathcal{C}}(\mathbf{M}, \mathbf{N}|_{\mathcal{C}}) \rightleftarrows \text{Hom}_{\mathcal{C}^{\circ}}(\mathbf{M}^{\circ}, \mathbf{N}) : R,$$

where \mathbf{N} is a dg idempotent complete pretriangulated 2-representation of \mathcal{C}° and $\mathbf{M} \in \mathcal{C}\text{-mod}^{\text{pre}}$.

In particular, the dg 2-functor $(-)^{\circ}$ restricts to a dg biequivalence on the full 2-subcategories of dg idempotent complete pretriangulated 2-representations.

Proof. The dg functor E is defined by extending a morphism of 2-representations $\Phi: \mathbf{M} \rightarrow \mathbf{N}|_{\mathcal{C}}$ of \mathcal{C} by setting

$$E(\Phi)_i = \Phi_i^{\circ}.$$

The target of this dg functor is $\mathbf{N}(i) \simeq \mathbf{N}(i)^{\circ}$ since \mathbf{N} is dg idempotent complete. The dg functor R is simply given by restricting to $\mathbf{N}(i) \subset \mathbf{N}(i)^{\circ}$. Given a dg morphism $\Psi: \mathbf{M}^{\circ} \rightarrow \mathbf{N}$ and an object (X, e) in $\mathbf{N}(i)$, it is clear that $E(\Psi)_i(X, e) = (\Psi_i(X), \Psi_i(e))$ gives a splitting for the idempotent $\Psi_i(e)$ and is hence isomorphic to $\Psi_i(X, e)$. \square

3.3. Compact pretriangulated 2-representations. Given a pretriangulated 2-representation \mathbf{M} and an object X of $\mathbf{M}(i)$, we define, for any object j of \mathcal{C} , **evaluation at X** to be the dg functor

$$\text{Ev}_X: \overline{\mathcal{C}}(i, j) \rightarrow \mathbf{M}(j), \quad F \mapsto \mathbf{M}(F)X.$$

We say that a pretriangulated 2-representation \mathbf{M} is **compact** if for any objects i, j of \mathcal{C} and any $X \in \mathbf{M}(i)$, the functors Ev_X satisfy the equivalent conditions of Lemma 2.9 (see also Lemma 2.10), i.e., the extended dg functors $\overline{\text{Ev}}_X$ have right adjoints. In this situation, we will also say that Ev_X has a compact right adjoint.

Lemma 3.8. *Let \mathbf{M} be a pretriangulated 2-representation of a dg 2-category \mathcal{C} and $X = (X' \oplus X'', \begin{pmatrix} 0 & \alpha \\ 0 & 0 \end{pmatrix}) \in \mathbf{M}(i)$. Assume that $\text{Ev}_{X'}$, $\text{Ev}_{X''}$ have compact right adjoints. Then Ev_X has a compact right adjoint.*

Proof. For $G \in \overline{\mathcal{C}}(i, j)$, $Y \in \mathbf{M}(j)$, we compute

$$\begin{aligned} \text{Hom}_{\mathbf{M}(j)}(\text{Ev}_X(G), Y) &\cong \text{Hom}_{\mathbf{M}(j)}((GX' \oplus GX'', \begin{pmatrix} 0 & G\alpha \\ 0 & 0 \end{pmatrix}), Y) \\ &\cong (\text{Hom}_{\mathbf{M}(j)}(GX'', Y) \oplus \text{Hom}_{\mathbf{M}(j)}(GX', Y), \begin{pmatrix} 0 & -\circ G\alpha \\ 0 & 0 \end{pmatrix}) \\ &\cong \left(\text{Hom}_{\overline{\mathcal{C}}(i, j)}(G, \text{Ev}_{X''}^* Y) \oplus \text{Hom}_{\mathbf{M}(i)}(G, \text{Ev}_{X'}^* Y), \begin{pmatrix} 0 & \text{Hom}_{\overline{\mathcal{C}}(i, j)}(G, (\text{Ev}_{\alpha}^*)^* Y) \\ 0 & 0 \end{pmatrix} \right) \\ &\cong \text{Hom}_{\overline{\mathcal{C}}(i, j)} \left(G, \left(\text{Ev}_{X''}^* \oplus \text{Ev}_{X'}^*, \begin{pmatrix} 0 & \text{Ev}_{\alpha}^* \\ 0 & 0 \end{pmatrix} \right) Y \right), \end{aligned}$$

where in the third isomorphism we have used that Ev is in fact a bifunctor by definition of a 2-representation. This computation shows that Ev_X has the compact right adjoint $(\text{Ev}_{X''}^* \oplus \text{Ev}_{X'}^*, \begin{pmatrix} 0 & \text{Ev}_{\alpha}^* \\ 0 & 0 \end{pmatrix})$. \square

We have the following dg 2-subcategories of $\mathcal{C}\text{-mod}^{\text{pre}}$.

- The 2-subcategory of *compact* pretriangulated 2-representations is denoted by $\mathcal{C}\text{-mod}_c^{\text{pre}}$.

- The 2-subcategory of *compact* pretriangulated 2-representations *with generators* is denoted by $\mathcal{C}\text{-mod}_{\mathbb{C}\mathbb{g}}^{\text{pre}}$.

We observe that \mathbf{P}_i is in $\mathcal{C}\text{-mod}_{\mathbb{C}\mathbb{g}}^{\text{pre}}$ and in $\mathcal{C}\text{-mod}_{\mathbb{C}\mathbb{g}}^{\text{pre}}$ provided that \mathcal{C} has generators (see Section 3.1). The following lemma establishes when \mathbf{P}_i belongs to $\mathcal{C}\text{-mod}_{\mathbb{C}\mathbb{g}}^{\text{pre}}$.

Lemma 3.9. *Let \mathcal{C} be a dg 2-category. Then all principal 2-representations \mathbf{P}_i are compact if and only if \mathcal{C} has compact left adjoints.*

Proof. First assume that \mathcal{C} has compact left adjoints. Let $F \in \mathcal{C}(i, j)$ be a 1-morphism. Using Lemma 2.9, we need to show that the $\mathcal{C}(j, k)$ -module $\text{Ev}_F^*(G^\vee) = \text{Hom}_{\vec{\mathcal{C}}(i, k)}((-)F, G)$ is compact for any 1-morphism G in $\mathcal{C}(i, k)$.

For the left adjoint $*F \in \vec{\mathcal{C}}(j, i)$ of F , there is a natural dg isomorphism

$$\text{Hom}_{\vec{\mathcal{C}}(i, k)}(HF, G) \xrightarrow{\sim} \text{Hom}_{\vec{\mathcal{C}}(j, k)}(H, G^*F), \quad \alpha \mapsto (\alpha \circ_0 \text{id}_{*F}) \circ_1 (\text{id}_H \circ_0 u_F),$$

for any $H \in \mathcal{C}(j, k)$. This provides a dg isomorphism of dg $\mathcal{C}(j, k)$ -modules between $\text{Hom}_{\vec{\mathcal{C}}(i, k)}((-)F, G)$ and the image under Y of G^*F , which is, by Lemma 2.6, a compact dg $\mathcal{C}(j, k)$ -module.

Conversely, assume \mathbf{P}_i is compact for any object i of \mathcal{C} . Then, in particular, for any $F \in \mathcal{C}(i, j)$ the functor $\text{Hom}_{\vec{\mathcal{C}}(j, j)}((-)F, \mathbb{1}_j)$ is compact. Again using Lemma 2.6, there exists a 1-morphism $*F$ in $\vec{\mathcal{C}}(j, i)$ such that $\text{Hom}_{\vec{\mathcal{C}}(j, j)}((-)F, \mathbb{1}_j)$ is dg isomorphic to $Y(*F)$ as a dg $\mathcal{C}(j, i)$ -module. Thus, there are dg isomorphisms

$$\text{Hom}_{\vec{\mathcal{C}}(j, j)}(GF, \mathbb{1}_j) \cong \text{Hom}_{\vec{\mathcal{C}}(j, i)}(G, *F),$$

natural in $G \in \mathcal{C}(j, i)$. These dg functors and natural dg isomorphisms extend to inputs G from $\vec{\mathcal{C}}(j, i)$. In particular, there are natural dg isomorphisms

$$\text{Hom}_{\vec{\mathcal{C}}(j, j)}(*FF, \mathbb{1}_j) \cong \text{Hom}_{\vec{\mathcal{C}}(j, j)}(*F, *F).$$

We define c_F to be the element corresponding to id_{*F} under this isomorphism. Similarly, $\text{Hom}_{\vec{\mathcal{C}}(i, j)}((-)F, F)$ is compact. Applying this to the identity 1-morphism $\mathbb{1}_i$, there exists a natural dg isomorphism

$$\text{Hom}_{\vec{\mathcal{C}}(i, j)}(\mathbb{1}_i F, F) \cong \text{Hom}_{\vec{\mathcal{C}}(i, i)}(\mathbb{1}_i, F^*F).$$

The image of the identity id_F gives the element u_F . The adjunction axioms for u_F, c_F now follow from the fact that the isomorphisms $\text{Hom}_{\vec{\mathcal{C}}(i, k)}(GF, H) \cong \text{Hom}_{\vec{\mathcal{C}}(j, k)}(G, H^*F)$ are natural in G, H adapting the argument of [McL, Section IV.1]. \square

Example 3.10. Let $R = \mathbb{k}[x]$ considered as a dg algebra concentrated in degree zero and define the dg 2-category \mathcal{C}_R to have one object \bullet and morphism category $\mathcal{C}_R(\bullet, \bullet)$ given by a strictification of $\{R \oplus \widehat{R \otimes_{\mathbb{k}} R}\}$ with horizontal composition induced by the tensor product over R , and $\mathbb{1}$ corresponding to R .

The dg 2-category \mathcal{C}_R has its natural dg 2-representation \mathbf{N} on $\mathbf{N}(\bullet) = \overline{\{R\}}$ induced by viewing the dg bimodules in $\mathcal{C}_R(\bullet, \bullet)$ as acting by tensor functors. This pretriangulated 2-representation is not compact. Indeed, since

$$\text{Hom}_{\mathbf{N}(\bullet)}(R \otimes_{\mathbb{k}} R \otimes_R R, R) \cong \text{Hom}_{\mathbb{k}\text{-mod}^{\text{dgs}}}(R, R)$$

is not a compact dg R - R -bimodule, the requirement

$$\mathrm{Hom}_{\mathbf{N}(\bullet)}(R \otimes_{\mathbb{k}} R \otimes_R R, R) \cong \mathrm{Hom}_{\vec{\mathcal{C}}(\bullet, \bullet)}(R \otimes_{\mathbb{k}} R, [R, R])$$

cannot be satisfied.

On the other hand, we have the trivial dg 2-representation \mathbf{M} with $\mathbf{M}(\mathbf{i}) = \mathbb{k}\text{-mod}^{\mathrm{dg}}$, on which $R \otimes_{\mathbb{k}} R$ acts by 0, R acts as the identity functor, with the endomorphism x of R acting by zero. This is a compact dg 2-representation, since

$$\mathrm{Hom}_{\mathbf{M}(\bullet)}(\mathbf{M}(F)\mathbb{k}, \mathbb{k}) \cong \mathrm{Hom}_{\vec{\mathcal{C}}(\bullet, \bullet)}(F, [\mathbb{k}, \mathbb{k}])$$

is satisfied for any F when taking $[\mathbb{k}, \mathbb{k}]$ to be the diagram $\mathbb{1} \xrightarrow{x} \mathbb{1}$ in $\vec{\mathcal{C}}(\bullet, \bullet)$.

3.4. Dg ideals of 2-representations and dg 2-subrepresentations. In this section we collect facts about dg ideals used later in the paper.

A **dg ideal** of a pretriangulated 2-representation \mathbf{M} of a dg 2-category \mathcal{C} is a collection of subcategories $\mathbf{I}(\mathbf{i}) \subset \mathbf{M}(\mathbf{i})$ such that $\mathbf{I}(\mathbf{i})$ is an ideal in the category $\mathbf{M}(\mathbf{i})$, which is closed under differentials and the \mathcal{C} -action.

Definition 3.11. Let \mathbf{M} be a pretriangulated 2-representation of \mathcal{C} , $\mathbf{i} \in \mathcal{C}$, and f a morphism in $\mathbf{M}(\mathbf{i})$. Define the **dg ideal $\mathbf{I}_{\mathbf{M}}(f)$ generated by f** to be the smallest dg ideal of \mathbf{M} containing f .

If \mathbf{I} is a dg ideal in a pretriangulated 2-representation \mathbf{M} , we can form the quotient \mathbf{M}/\mathbf{I} acting on

$$(\mathbf{M}/\mathbf{I})(\mathbf{i}) = \mathbf{M}(\mathbf{i})/\mathbf{I}(\mathbf{i}).$$

In particular, given a dg 2-subrepresentation \mathbf{N} of \mathbf{M} , we define the **quotient \mathbf{M}/\mathbf{N}** to be the quotient of \mathbf{M} by the dg ideal generated by \mathbf{N} .

Let \mathbf{M} be a pretriangulated 2-representation of \mathcal{C} . A pretriangulated **2-subrepresentation \mathbf{N}** of \mathbf{M} is a collection of thick subcategories $\mathbf{N}(\mathbf{i}) \subseteq \mathbf{M}(\mathbf{i})$ for all objects \mathbf{i} of \mathcal{C} such that for all 1-morphisms G and any 2-morphism α in $\mathcal{C}(\mathbf{i}, \mathbf{j})$, we have that

$$\mathbf{N}G(N) = \mathbf{M}G(N) \in \mathbf{N}(\mathbf{j}), \quad \mathbf{N}G(f) = \mathbf{M}G(f) \in \mathbf{N}(\mathbf{j}), \quad \mathbf{N}(\alpha)_N = \mathbf{M}(\alpha)_N,$$

for all objects N and morphisms f in $\mathbf{N}(\mathbf{i})$. In particular, the thick subcategories $\mathbf{N}(\mathbf{i})$ are closed under the \mathcal{C} -action, which is the restriction of the \mathcal{C} -action given by the dg 2-functor \mathbf{M} . Note that we require that $\mathbf{N}(\mathbf{i})$ is closed under forming biproducts and taking dg direct summands that exist in $\mathbf{M}(\mathbf{i})$.

Given a collection of pretriangulated 2-subrepresentations $\{\mathbf{N}^\nu\}_{\nu \in I}$ of a fixed pretriangulated 2-representation \mathbf{M} , we define the **sum $\sum_{\nu \in I} \mathbf{N}^\nu$** , as the smallest pretriangulated 2-subrepresentation of \mathbf{M} containing all \mathbf{N}^ν .

3.5. Cyclic and quotient-simple dg 2-representations.

Definition 3.12. Given a pretriangulated 2-representation \mathbf{M} of \mathcal{C} and an object $X \in \mathbf{M}(\mathbf{i})$, for some $\mathbf{i} \in \mathcal{C}$, we denote by $\mathbf{G}_{\mathbf{M}}(X)$ the 2-representation on the thick closure of $\{\mathbf{M}(G)X \mid G \in \mathcal{C}(\mathbf{i}, \mathbf{j}), \mathbf{j} \in \mathcal{C}\}$ inside $\coprod_{\mathbf{j} \in \mathcal{C}} \mathbf{M}(\mathbf{j})$. We call $\mathbf{G}_{\mathbf{M}}(X)$ the dg 2-subrepresentation **\mathcal{C} -generated by X** . If $\mathbf{G}_{\mathbf{M}}(X) = \mathbf{M}$, we say X **\mathcal{C} -generates \mathbf{M}** . We call \mathbf{M} **cyclic** if there exists an $X \in \mathbf{M}(\mathbf{i})$, for some $\mathbf{i} \in \mathcal{C}$, which \mathcal{C} -generates \mathbf{M} .

Lemma 3.13. *Let \mathcal{C} be a dg 2-category and \mathbf{M} a cyclic dg 2-representation \mathcal{C} -generated by $X \in \mathbf{M}(\mathbf{i})$.*

- (a) If \mathcal{C} has generators, then $\mathbf{M} \in \mathcal{C}\text{-mod}_{\mathbf{g}}^{\text{pre}}$.
- (b) If \mathcal{C} has compact left adjoints, and Ev_X has a compact right adjoint, then $\mathbf{M} \in \mathcal{C}\text{-mod}_{\mathbf{c}}^{\text{pre}}$.

Proof. To prove (a), we first note that by assumption that X \mathcal{C} -generates \mathbf{M} , the thick closure of $\{\mathbf{M}(\mathbf{F})X\}$, where \mathbf{F} ranges over the 1-morphisms in $\mathcal{C}(\mathbf{i}, \mathbf{j})$, is $\mathbf{M}(\mathbf{j})$. If $\mathcal{C}(\mathbf{i}, \mathbf{j})$ is the thick closure of $\mathbf{F}_{\mathbf{i}, \mathbf{j}}$, then every 1-morphism is a direct summand of an object in $\overline{\{\mathbf{F}_{\mathbf{i}, \mathbf{j}}\}}$. Hence, every object in $\mathbf{M}(\mathbf{j})$ is a direct summand of an object in $\overline{\{\mathbf{M}(\mathbf{F}_{\mathbf{i}, \mathbf{j}})X\}}$ since the dg 2-functor \mathbf{M} preserves shifts and cones, so $\mathbf{M} \in \mathcal{C}\text{-mod}_{\mathbf{g}}^{\text{pre}}$.

To prove (b), we first argue that, given the assumptions, for any $\mathbf{F} \in \mathcal{C}(\mathbf{i}, \mathbf{j})$, $\text{Ev}_{\mathbf{M}(\mathbf{F})X}$ also has a compact right adjoint. For this, we show that for any object $Y \in \mathbf{M}(\mathbf{k})$, $\text{Ev}_{\mathbf{M}(\mathbf{F})X}^*(Y)$ is dg isomorphic to $\text{Ev}_X^*(Y)^*F$ and hence compact. This follows from the following computation

$$\begin{aligned} \text{Hom}_{\mathbf{M}(\mathbf{k})}(\text{Ev}_{\mathbf{M}(\mathbf{F})X}(G), Y) &= \text{Hom}_{\mathbf{M}(\mathbf{k})}(\text{Ev}_X(GF), Y) \\ &\cong \text{Hom}_{\mathcal{C}(\mathbf{i}, \mathbf{k})}(GF, \text{Ev}_X^*(Y)) \\ &\cong \text{Hom}_{\mathcal{C}(\mathbf{j}, \mathbf{k})}(G, \text{Ev}_X^*(Y)^*F) \end{aligned}$$

of the right adjoint of $\text{Ev}_{\mathbf{M}(\mathbf{F})X}$, for $G \in \mathcal{C}(\mathbf{j}, \mathbf{k})$. Here, the first equality follows using $\mathbf{M}(GF) = \mathbf{M}(G)\mathbf{M}(\mathbf{F})$, the second dg isomorphism uses the right adjoint of Ev_X , and the last dg isomorphism uses Lemma 3.9. Thus, by Lemma 2.9, $\text{Ev}_{\mathbf{M}(\mathbf{F})X}$ has a compact right adjoint. For X' in the thick closure of the $\{\mathbf{M}(\mathbf{F})X\}$, the right adjoint of $\text{Ev}_{X'}$ has compact right adjoint by Lemma 3.8 and additivity. Therefore $\text{Ev}_{X'}$ has a compact right adjoint for any $X' \in \mathbf{M}(\mathbf{i})$ and \mathbf{M} is compact. \square

Given a dg 2-subrepresentation \mathbf{N} of a pretriangulated 2-representation \mathbf{M} we define the **weak closure** \mathbf{N}^\diamond as the dg 2-subrepresentation defined on the full dg subcategories $\mathbf{N}^\diamond(\mathbf{i}) \subseteq \mathbf{M}(\mathbf{i})$ on objects whose identities are contained in the ideal generated by \mathbf{N} . A dg 2-subrepresentation \mathbf{N} is **weakly closed** in \mathbf{M} if $\mathbf{N} = \mathbf{N}^\diamond$.

We say that $X \in \mathbf{M}(\mathbf{i})$, for some \mathbf{i} in \mathcal{C} , **weakly \mathcal{C} -generates** \mathbf{M} if $\mathbf{G}_{\mathbf{M}}(X)^\diamond = \mathbf{M}$.

We say \mathbf{M} is **weakly transitive** if the weak closure of any dg 2-subrepresentation of \mathbf{M} equals \mathbf{M} . Equivalently, \mathbf{M} is weakly transitive if it is \mathcal{C} -generated by any non-zero object X in any $\mathbf{M}(\mathbf{i})$.

We call a pretriangulated 2-representation \mathbf{M} **quotient-simple** provided it has no proper nonzero dg ideals. Note that, by definition, if \mathbf{M} is quotient simple, it is also weakly transitive.

3.6. Homotopy 2-representations. Let \mathcal{C} be a dg 2-category. In this section, we describe how a pretriangulated 2-representation \mathbf{M} descends to a 2-representations \mathbf{KM} on the corresponding triangulated categories. For this, we define $\mathbf{KM}(\mathbf{i}) := \mathcal{K}(\mathbf{M}(\mathbf{i}))$ for any object \mathbf{i} in \mathcal{C} and

$$\mathbf{KM}(\mathbf{F}) := \mathcal{K}(\mathbf{M}(\mathbf{F})): \mathbf{KM}(\mathbf{i}) \rightarrow \mathbf{KM}(\mathbf{j}),$$

for any 1-morphism in $\mathcal{C}(\mathbf{i}, \mathbf{j})$. If $\alpha: \mathbf{F} \rightarrow \mathbf{G}$ is a dg 2-morphism, then α induces a natural transformation $\mathbf{KM}(\alpha): \mathbf{KM}(\mathbf{F}) \rightarrow \mathbf{KM}(\mathbf{G})$. By construction, \mathbf{KM} is an additive 2-representation of $\mathcal{Z}\mathcal{C}$, cf. Section 3.1.

Lemma 3.14. *The 2-representation \mathbf{KM} is a 2-representation of the homotopy 2-category $\mathcal{H}\mathcal{C}$.*

Proof. If $\alpha: F \rightarrow G$ is null-homotopic, i.e., $\partial(\beta) = \alpha$ for some morphism $\beta: F \rightarrow G$, then

$$\mathbf{KM}(\alpha) = \mathbf{KM}(\partial\beta) = \partial(\mathbf{KM}(\beta)),$$

and hence $\mathbf{KM}(\alpha) = 0$ as a natural transformation from $\mathbf{KM}(F)$ to $\mathbf{KM}(G)$. Thus, \mathbf{KM} descends to the quotient \mathcal{HC} of \mathcal{LC} . \square

We call \mathbf{KM} the **homotopy 2-representation** derived from \mathbf{M} . We refrain from giving a general definition of a triangulated 2-representation of \mathcal{HC} but note that the homotopy 2-representations \mathbf{KM} defined above are triangulated in the sense that they define 2-functors to the target 2-category \mathfrak{T}^{dg} whose

- objects are triangulated categories of the form $\mathcal{K}(\overline{\mathcal{C}})$;
- 1-morphisms are functors between such categories that are induced from dg functors and hence are functors of triangulated categories;
- 2-morphisms are natural transformations of such functors coming from dg natural transformations of the corresponding dg categories.

We will restrict ourselves to triangulated 2-representations that have dg enhancements as described above. We note that the idempotent completion $(\mathbf{KM})^\circ$ is again triangulated by [BS].

The dg 2-subcategory $\mathfrak{T}_g^{\text{dg}}$ of \mathfrak{T}^{dg} consists of such triangulated categories which have a generator (also called a classical generator, see e.g. [LO, Definition 1.10]), i.e., are equal to the thick closure of a single object. We say a 2-representation of the form \mathbf{KM} **has generators** if its target is $\mathfrak{T}_g^{\text{dg}}$, that is, if $\mathbf{KM}(i)$ has a generator for any object i . We note that if \mathbf{M} has generators as a pretriangulated 2-representation, then, in particular, \mathbf{KM} has generators as triangulated 2-representation.

We will use the following weaker notion of equivalence of dg 2-representations. Let \mathbf{M} and \mathbf{N} be pretriangulated 2-representations. A morphism of dg 2-representations $\Phi: \mathbf{M} \rightarrow \mathbf{N}$ is a **quasi-equivalence** if the dg functors

$$\Phi_i: \mathbf{M}(i) \rightarrow \mathbf{N}(i)$$

are quasi-equivalences (see Section 2.5). In this case, we say that \mathbf{M} and \mathbf{N} are **quasi-equivalent**. In particular, a dg morphism Φ induces an equivalence $\mathbf{K}\Phi$ of triangulated 2-representations if and only if Φ is a quasi-equivalence.

We say that X is a \mathcal{C} -**quasi-generator** for a pretriangulated 2-representation \mathbf{M} if the inclusion of $\mathbf{G}_M(X)$ into \mathbf{M} is a quasi-equivalence (and hence $\mathbf{K}\mathbf{G}_M(X)$ and \mathbf{KM} are equivalent).

For instance, if \mathbf{M} is acyclic, the zero object is a \mathcal{C} -quasi-generator for \mathbf{M} .

4. DG ALGEBRA 1-MORPHISMS

Throughout this section, assume that \mathcal{C} is a dg 2-category and \mathbf{M} a pretriangulated 2-representation in $\mathcal{C}\text{-mod}^{\text{pre}}$.

4.1. Compact modules over dg algebra 1-morphisms. We say that A is a **dg algebra 1-morphism** in $\mathcal{C}(i, i)$ if it comes with dg morphisms $u_A: \mathbb{1}_i \rightarrow A$, $m_A: AA \rightarrow A$ satisfying the usual axioms of a unitary product internal to \mathcal{C} . Similarly, a morphism of dg algebra 1-morphisms is just a 2-morphism commuting with the multiplication and unit morphisms. Given A , define $\text{mod}_{\mathcal{C}(i, j)}^{\text{dg}}\text{-}A$ to be the category of **right dg A-modules** in $\mathcal{C}(i, j)$.

Lemma 4.1. For 1-morphisms G in $\mathcal{C}(i, j)$ and $Y \in \text{mod}_{\mathcal{C}(i, j)}^{\text{dg}}\text{-}A$, there is a dg isomorphism

$$\text{Hom}_{\text{mod}_{\mathcal{C}(i, j)}^{\text{dg}}\text{-}A}(GA, Y) \cong \text{Hom}_{\mathcal{C}(i, j)}(G, Y),$$

natural in G and Y .

Proof. The proof is an adaptation of the purely formal argument in [EGNO, 7.8.12], noting that the mutually inverse morphisms

$$f \mapsto (f \circ (\text{id}_G \circ_0 u_A)) \quad \text{and} \quad (\rho_Y \circ (g \circ_0 \text{id}_A)) \leftarrow g$$

are dg morphisms since u_A and ρ_Y , the right A -action on Y , are. \square

We now construct pretriangulated 2-representations of modules over a dg algebra 1-morphism A in $\overrightarrow{\mathcal{C}}(i, i)$.

Definition 4.2 ($\overrightarrow{\mathbf{M}}_A, \mathbf{M}_A$). Let A be a dg algebra 1-morphism in $\overrightarrow{\mathcal{C}}(i, i)$.

- (1) Define $\overrightarrow{\mathbf{M}}_A(j) = \text{mod}_{\mathcal{C}(i, j)}^{\text{dg}}\text{-}A$, for any object j of \mathcal{C} . Left multiplication by 1- and 2-morphisms in \mathcal{C} induces a natural structure of a pretriangulated 2-representation of \mathcal{C} on the $\overrightarrow{\mathbf{M}}_A(j)$ which we denote by $\overrightarrow{\mathbf{M}}_A$.
- (2) Define $\mathbf{M}_A(j)$ to be the thick closure of the set $\{GA \mid G \in \mathcal{C}(i, j)\}$ inside $\text{mod}_{\mathcal{C}(i, j)}^{\text{dg}}\text{-}A$. The collection of dg subcategories $\mathbf{M}_A(j)$ forms a pretriangulated 2-subrepresentation of $\overrightarrow{\mathbf{M}}_A$ of \mathcal{C} , which we denote by \mathbf{M}_A .

Lemma 4.3. The dg 2-representation $\overrightarrow{\mathbf{M}}_A$ is dg equivalent to $\overrightarrow{\mathbf{M}}_A$.

Proof. For any j , the dg category $\overrightarrow{\mathbf{M}}_A(j)$ is closed under taking cokernels of dg morphisms. Thus, the embedding $I: \mathbf{M}_A(j) \rightarrow \overrightarrow{\mathbf{M}}_A(j)$ extends to a fully faithful dg functor $\overrightarrow{I}: \overrightarrow{\mathbf{M}}_A(j) \rightarrow \overrightarrow{\mathbf{M}}_A(j)$.

Now, any object $X \in \overrightarrow{\mathbf{M}}_A$ is dg isomorphic to the cokernel of the dg morphism

$$\text{id}_X \circ_0 m - a \circ_0 \text{id}_A: XAA \rightarrow XA.$$

where $a: XA \rightarrow A$ denotes the right A -action on X . Both modules XAA, XA are in the image of \overrightarrow{I} . But the image of \overrightarrow{I} is closed under taking cokernels of dg morphisms. This follows as cokernels of dg morphisms exist in $\overrightarrow{\mathbf{M}}_A(j)$ by Lemma 2.7 and these cokernels are preserved by \overrightarrow{I} . Thus, X is in the essential image of \overrightarrow{I} . This proves the claim. \square

Lemma 4.4. Let A be a dg algebra 1-morphism in $\overrightarrow{\mathcal{C}}(i, i)$.

- (a) If \mathcal{C} has generators, then $\mathbf{M}_A \in \mathcal{C}\text{-mod}_{\mathbf{g}}^{\text{pre}}$.
- (b) If \mathcal{C} has compact left adjoints, then $\mathbf{M}_A \in \mathcal{C}\text{-mod}_{\mathbf{c}}^{\text{pre}}$.

Proof. If $\mathcal{C}(i, j)$ has a generator $F_{i, j}$, then, by definition, $\mathbf{M}_A(j)$ is the thick closure of $F_{i, j}A$ and thus \mathbf{M}_A has generators, i.e. $\mathbf{M}_A \in \mathcal{C}\text{-mod}_{\mathbf{g}}^{\text{pre}}$. This proves (a).

To prove (b), we use Lemma 3.13(b) to see that it suffices to show that Ev_A has a compact right adjoint. By Lemma 4.1, for any $G \in \mathcal{C}(j, k), Y \in \overrightarrow{\mathbf{M}}_A(k)$, we obtain natural isomorphisms

$$\begin{aligned} \text{Hom}_{\overrightarrow{\mathbf{M}}_A(k)}(\overrightarrow{\text{Ev}}_A(G), Y) &= \text{Hom}_{\overrightarrow{\mathbf{M}}_A(k)}(GA, Y) \\ &\cong \text{Hom}_{\mathcal{C}(i, k)}(G, Y) \end{aligned}$$

thus Ev_A has the forgetful functor on $\overrightarrow{\mathbf{M}}_A \rightarrow \overrightarrow{\mathcal{C}}$ as right adjoint, which is a compact right adjoint by construction. \square

Similarly to dg modules one also defines dg bimodules in a general dg 2-category. Working with $\overrightarrow{\mathcal{C}}$, the existence of cokernels ensures that these dg bimodules can be composed. Indeed, let A, B, C be dg algebra 1-morphisms in $\overrightarrow{\mathcal{C}}$ and let M be a dg A - B -bimodule 1-morphism with (commuting) left and right actions λ_M and ρ_M and let N be a dg B - C -bimodule 1-morphism with (commuting) left and right actions λ_N and ρ_N . We define the **relative composition** $M \circ_B N$ as the cokernel of the dg morphism

$$\text{MBN} \xrightarrow{\rho_M \circ_0 \text{id}_N - \text{id}_M \circ_0 \lambda_N} \text{MN},$$

which exists in $\overrightarrow{\mathcal{C}}$ by Lemma 2.7. Note that $M \circ_B N$ is now a dg A - C -bimodule.

4.2. The algebra structure on internal homs $[X, X]$. Let $X \in \mathbf{M}(i)$ be nonzero. We consider the evaluation dg functors

$$\text{Ev}_X: \mathcal{C}(i, j) \rightarrow \mathbf{M}(j), \quad F \mapsto \mathbf{M}(F)X,$$

which we may extend to $\overrightarrow{\mathcal{C}}(i, j)$ defined at the beginning of Section 3.3.

If Ev_X has a compact right adjoint (i.e., satisfies the equivalent conditions of Lemma 2.9), then we denote the right adjoint dg functor of $\overrightarrow{\text{Ev}}_X$ by

$$[X, -]: \overrightarrow{\mathbf{M}}(j) \rightarrow \overrightarrow{\mathcal{C}}(i, j),$$

called the **internal hom**. We note that $[X, -]$ is natural in X . Furthermore, for the restriction of $[X, -]$ to $\overrightarrow{\mathbf{M}}(j)$, and an object $(\bigoplus_i Y_i, \alpha)$ in $\overrightarrow{\mathbf{M}}(j)$, we have

$$\left[X, \left(\bigoplus_i Y_i, \alpha \right) \right] \cong \left(\bigoplus_i [X, Y_i], [X, \alpha] \right).$$

For $X \in \mathbf{M}(i), Y \in \overrightarrow{\mathbf{M}}(j)$, we define

$$\text{ev}_{X,Y}: \overrightarrow{\mathbf{M}}([X, Y])X \rightarrow Y$$

as the image of the identity morphism under the dg isomorphism

$$\text{Hom}_{\overrightarrow{\mathcal{C}}(i,j)}([X, Y], [X, Y]) \cong \text{Hom}_{\overrightarrow{\mathbf{M}}(j)}(\overrightarrow{\mathbf{M}}([X, Y])X, Y).$$

Note that, as the image of a dg morphism under a dg isomorphism, $\text{ev}_{X,Y}$ is also a dg morphism.

Lemma 4.5. *For nonzero $X \in \mathbf{M}(i)$ such that Ev_X has a compact right adjoint, the object $A_X := [X, X] \in \overrightarrow{\mathcal{C}}(i, i)$ carries the structure of a dg algebra 1-morphism.*

Proof. We define the unit morphism u_{A_X} as the image of the identity on X under the dg isomorphism

$$\text{Hom}_{\overrightarrow{\mathbf{M}}(i)}(X, X) \cong \text{Hom}_{\overrightarrow{\mathcal{C}}(i,i)}(\mathbb{1}_i, A_X).$$

We define the multiplication m_{A_X} as the image of the dg morphism

$$\text{ev}_{X,X} \circ \overrightarrow{\mathbf{M}}(A_X)(\text{ev}_{X,X}): \overrightarrow{\mathbf{M}}(A_X)\overrightarrow{\mathbf{M}}(A_X)X \rightarrow \overrightarrow{\mathbf{M}}(A_X)X \rightarrow X$$

under the dg isomorphism

$$\text{Hom}_{\overrightarrow{\mathbf{M}}(i)}(\overrightarrow{\mathbf{M}}(A_X A_X)X, X) \cong \text{Hom}_{\overrightarrow{\mathcal{C}}(i,i)}(A_X A_X, A_X).$$

It is a routine exercise to check the algebra axioms. \square

Note that if \mathbf{M} is compact, i.e., $\mathbf{M} \in \mathcal{C}\text{-mod}_c^{\text{pre}}$, then we can associate an algebra 1-morphism A_X to any nonzero object $X \in \mathbf{M}(i)$.

Lemma 4.6. *For any $Y \in \overrightarrow{\mathbf{M}}(j)$, the object $[X, Y]$ in $\overrightarrow{\mathcal{C}}(i, j)$ has the structure of a right dg module over A_X . Moreover, for any morphism $f: Y_1 \rightarrow Y_2 \in \overrightarrow{\mathbf{M}}(j)$, the morphism $[X, f]$ commutes with the right action of A_X and satisfies $[X, \partial f] = \partial[X, f]$.*

Proof. We define the right action $\rho_{[X, Y]}: [X, Y]A_X \rightarrow [X, Y]$ as the image of

$$\text{ev}_{X, Y} \circ \overrightarrow{\mathbf{M}}([X, Y])(\text{ev}_{X, X}): \overrightarrow{\mathbf{M}}([X, Y])\overrightarrow{\mathbf{M}}(A_X)X \rightarrow \overrightarrow{\mathbf{M}}([X, Y])X \rightarrow Y$$

under the dg isomorphism

$$\text{Hom}_{\overrightarrow{\mathbf{M}}(j)}(\overrightarrow{\mathbf{M}}([X, Y]A_X)X, Y) \cong \text{Hom}_{\overrightarrow{\mathcal{C}}(i, j)}([X, Y]A_X, [X, Y]).$$

It is, again, a routine exercise to check that this morphism satisfies the axioms of a right action. Compatibility with differentials follows from $\text{ev}_{X, Y}$ being a dg morphism. \square

Note that Lemma 4.6 can be reformulated as saying that $[X, -]$ is a dg functor from $\overrightarrow{\mathbf{M}}(j)$ to $\overrightarrow{\mathbf{M}}_{A_X}(j)$ for any $j \in \mathcal{C}$.

4.3. An equivalence of dg 2-representations. Throughout this subsection, we will assume that \mathcal{C} has compact left adjoints, see Definition 3.3(b). We additionally assume that \mathbf{M} is compact though, strictly speaking, we could fix X and only require that Ev_X has a compact right adjoint.

Lemma 4.7. *For any $G \in \overrightarrow{\mathcal{C}}(i, j)$, $X \in \mathbf{M}(i)$, and $Y \in \mathbf{M}(j)$, we have a dg isomorphism*

$$[X, \mathbf{M}(G)Y] \cong G[X, Y]$$

in $\overrightarrow{\mathcal{C}}(i, j)$, natural in all variables.

Proof. For any $H \in \overrightarrow{\mathcal{C}}(i, j)$, there is a sequence of dg isomorphisms, natural in all variables,

$$\begin{aligned} \text{Hom}_{\overrightarrow{\mathcal{C}}(i, j)}(H, [X, \mathbf{M}(G)Y]) &\cong \text{Hom}_{\mathbf{M}(j)}(\mathbf{M}(H)X, \mathbf{M}(G)Y) \\ &\cong \text{Hom}_{\overrightarrow{\mathbf{M}}(i)}((\overrightarrow{\mathbf{M}}(G))\overrightarrow{\mathbf{M}}(H)X, Y) \\ &\cong \text{Hom}_{\overrightarrow{\mathbf{M}}(i)}(\overrightarrow{\mathbf{M}}((\overrightarrow{*}G)H)X, Y) \\ &\cong \text{Hom}_{\overrightarrow{\mathcal{C}}(i, j)}((\overrightarrow{*}G)H, [X, Y]) \\ &\cong \text{Hom}_{\overrightarrow{\mathcal{C}}(i, j)}(H, G[X, Y]), \end{aligned}$$

where the second and third dg isomorphisms follow from 2-functoriality of $\overrightarrow{\mathbf{M}}$, which, in particular, implies that $\overrightarrow{\mathbf{M}}(\overrightarrow{*}G)$ is left adjoint to $\overrightarrow{\mathbf{M}}(G)$. \square

Proposition 4.8. *There is a morphism of dg 2-representations from $\overrightarrow{\mathbf{M}}$ to $\overrightarrow{\mathbf{M}}_{A_X}$ induced by the functor $[X, -]$.*

Proof. For notational simplicity, we denote the dg functor $[X, -]: \overrightarrow{\mathbf{M}}(j) \rightarrow \overrightarrow{\mathbf{M}}_{A_X}(j)$, which indeed has the latter target by Lemma 4.6, by Φ . By Lemma 4.7, for each 1-morphism G , there is a dg isomorphism of functors $\eta_G: \Phi \circ \overrightarrow{\mathbf{M}}(G) \rightarrow \overrightarrow{\mathbf{M}}_{A_X}(G) \circ \Phi$, natural in G .

To prove that the dg functor Φ together with the natural dg isomorphism η indeed induces a morphism of 2-representations, it hence suffices to check that

$$\eta_{GH} = (\text{id}_{\overline{\mathbf{M}}_{A_X}(G)} \circ_0 \eta_H) \circ_1 (\eta_G \circ_0 \text{id}_{\overline{\mathbf{M}}(H)}),$$

for any objects $k, 1$ and 1-morphism $H \in \overline{\mathcal{C}}(j, k)$, $G \in \overline{\mathcal{C}}(k, 1)$, $K \in \overline{\mathcal{C}}(i, 1)$.

This follows from commutativity of

$$\begin{array}{ccc} \text{Hom}_{\overline{\mathbf{M}}(1)}(\overline{\mathbf{M}}(K)X, \overline{\mathbf{M}}(GH)Y) & \xrightarrow{\sim} & \text{Hom}_{\overline{\mathbf{M}}(k)}(\overline{\mathbf{M}}((*G)K)X, \overline{\mathbf{M}}(H)Y) \\ \downarrow \sim & & \downarrow \sim \\ \text{Hom}_{\overline{\mathbf{M}}(j)}(\overline{\mathbf{M}}(* (GH)K)X, Y) & & \text{Hom}_{\overline{\mathbf{M}}(j)}(\overline{\mathbf{M}}((*H)(*G)K)X, Y) \\ \downarrow \sim & & \downarrow \sim \\ \text{Hom}_{\overline{\mathcal{C}}(i,j)}(* (GH)K, [X, Y]) & & \text{Hom}_{\overline{\mathcal{C}}(i,j)}((*H)(*G)K, [X, Y]) \\ \downarrow \sim & & \downarrow \sim \\ \text{Hom}_{\overline{\mathcal{C}}(i,1)}(K, GH[X, Y]) & \xleftarrow{\sim} & \text{Hom}_{\overline{\mathcal{C}}(i,k)}((*G)K, H[X, Y]) \end{array}$$

in exactly the same vein as in [MMMT, Lemma 6]. \square

Proposition 4.9. *The morphism of dg 2-representations constructed in Proposition 4.8 is fully faithful when restricted to $\mathbf{G}_{\mathbf{M}}(X)$, the pretriangulated 2-subrepresentation \mathcal{C} -generated by X .*

Proof. First note that

$$\begin{aligned} \text{Hom}_{\overline{\mathbf{M}}_{A_X}(j)}([X, \mathbf{M}(G)X], [X, \mathbf{M}(H)X]) &\cong \text{Hom}_{\overline{\mathbf{M}}_{A_X}(j)}(\mathbf{G}A_X, [X, \mathbf{M}(H)X]) \\ &\cong \text{Hom}_{\overline{\mathcal{C}}(i,j)}(\mathbf{G}, [X, \mathbf{M}(H)X]) \\ &\cong \text{Hom}_{\overline{\mathbf{M}}(j)}(\mathbf{M}(G)X, \mathbf{M}(H)X) \\ &\cong \text{Hom}_{\mathbf{M}(j)}(\mathbf{M}(G)X, \mathbf{M}(H)X) \end{aligned}$$

where the second isomorphism is given by Lemma 4.1. This shows that the functor $[X, -]$ is full and faithful for objects of the form $\mathbf{M}(G)X$.

From the definition of morphism spaces in $\mathbf{M}(j) \simeq \overline{\mathbf{M}}(j)$, it is immediate that

$$\text{Hom}_{\overline{\mathbf{M}}_{A_X}(j)}([X, Y], [X, Z]) \cong \text{Hom}_{\mathbf{M}(j)}(Y, Z),$$

for all Y, Z in the thick closure of $\{\mathbf{M}(G)X \mid G \in \mathcal{C}(i, j)\}$, completing the proof. \square

The dg 2-representation $\mathbf{G}_{\mathbf{M}}(X)$ may not be dg equivalent to \mathbf{M}_{A_X} as the latter is closed under dg idempotents but the former is not. However, when replacing \mathbf{M} by the dg idempotent completion \mathbf{M}° , we obtain an equivalence of dg 2-representation between $\mathbf{G}_{\mathbf{M}^\circ}(X)$ and \mathbf{M}_{A_X} .

Proposition 4.10. *The morphism of dg 2-representations from Proposition 4.8 induces an equivalence of dg 2-representation $\mathbf{G}_{\mathbf{M}^\circ}(X) \rightarrow \mathbf{M}_{A_X}$.*

Proof. Note that $\mathbf{F}A \cong [X, \mathbf{F}X]$ and $[X, -]$ commutes with taking dg direct summands. Hence, the full image of $[X, -]$ is the thick closure of the set $\{\mathbf{F}A_X \mid \mathbf{F}\}$. \square

As X \mathcal{C} -generating \mathbf{M} implies that X also \mathcal{C} -generates \mathbf{M}° , we obtain the following result.

Corollary 4.11. *If the object X \mathcal{C} -generates \mathbf{M} , then the dg 2-representations \mathbf{M}_{A_X} and \mathbf{M}° are dg equivalent.*

Observe that, if $\mathbf{M}(j) = \widehat{\{X_j\}}$, then $\mathbf{M}_{A_X}(j) \simeq \widehat{\{[X, X_j]\}}$ and $\mathbf{M}_{A_X}(j)$ is hence in $\mathfrak{M}_g^{\text{dg}}$. Hence, if such X_j exists for any j , then $\mathbf{M}_{A_X} \in \mathcal{C}\text{-mod}_g^{\text{pre}}$.

Corollary 4.12. *Provided that X \mathcal{C} -generates \mathbf{M} , the morphism of dg 2-representations $\overline{\mathbf{M}} \rightarrow \overline{\mathbf{M}}_{A_X}$ constructed in Proposition 4.8 is a dg equivalence.*

Proof. We note that the equivalences from Corollary 4.11 and Lemma 4.3 give an equivalence of $\overline{\mathbf{M}^\circ} \simeq \overline{\mathbf{M}}$ and $\overline{\mathbf{M}}_{A_X} \simeq \overline{\mathbf{M}}_{A_X}$. \square

Corollary 4.13. *If \mathbf{M} is cyclic, then there exists a dg algebra 1-morphism A in \mathcal{C} , such that \mathbf{M}° is dg equivalent to \mathbf{M}_A .*

Proof. This is seen by choosing a \mathcal{C} -generator X of \mathbf{M} , and setting $A = A_X$. \square

If \mathbf{M} only has a weak \mathcal{C} -generator, the following weaker statement still holds true.

Proposition 4.14. *Let X be a weak \mathcal{C} -generator for \mathbf{M} . Then $[X, -]$ restricts to a full and faithful morphism of pretriangulated 2-representations $\mathbf{M}^\diamond \rightarrow \mathbf{M}_{A_X}^\diamond$. Here, the target is the weak closure taken in $\overline{\mathbf{M}}_{A_X}$.*

Proof. Assume given an object $Y \in \mathbf{M}^\diamond(i)$, we find an object $Y' \in \mathbf{M}(i)$ together with (non-dg) morphisms $\iota: Y \rightarrow Y', p: Y' \rightarrow Y$ such that $\text{id}_Y = p\iota$. As $[X, -]$ is a functor, $[X, p][X, \iota]$ is the identity on $[X, Y]$ and hence the latter is contained in the closure $\mathbf{M}_{A_X}^\diamond$. The arguments in the proof of Proposition 4.9 can again be used to prove that the functor $[X, -]$ to the closure $\mathbf{M}^\diamond(i)$ is fully faithful. For this, observe that, for 1-morphisms $G, G' \in \overline{\mathcal{C}}(i, j)$, and objects $Y, Y' \in \mathbf{M}(j)$ together with morphisms $\text{id}_Y = Y \xrightarrow{\iota} \mathbf{M}(G)X \xrightarrow{p} Y$ and $\text{id}_{Y'} = Y' \xrightarrow{\iota'} \mathbf{M}(G')X \xrightarrow{p'} Y'$, the diagram

$$\begin{array}{ccc} \text{Hom}(\mathbf{M}(G)X, \mathbf{M}(G')X) & \xrightarrow{[X, -]} & \text{Hom}([X, \mathbf{M}(G)X], [X, \mathbf{M}(G')X]) \\ \begin{array}{c} \uparrow [X, \iota'](-) \circ [X, p] \\ \downarrow [X, p'] \circ (-) \circ [X, \iota] \end{array} & & \begin{array}{c} \uparrow \iota' \circ (-) \circ p \\ \downarrow p' \circ (-) \circ \iota \end{array} \\ \text{Hom}(Y, \mathbf{M}(G')X) & \xrightarrow{[X, -]} & \text{Hom}([X, Y], [X, \mathbf{M}(G')X]). \end{array}$$

commutes. The horizontal maps are the application of a dg functor and hence commute with differentials. Although the vertical maps are not necessarily dg morphisms, they display the lower row as direct summands of the upper row. Hence, the lower horizontal map is the restriction of an isomorphism and hence a dg isomorphism. \square

The morphism $[X, -]$ of dg 2-representations, when extended to the weak closures, is not necessarily an equivalence. For example, take \mathcal{C} to be the trivial dg 2-category with single object \bullet , 1-morphism $\mathbb{1} = \mathbb{1}_\bullet$, and $\text{End}_{\mathcal{C}(\bullet, \bullet)}(\mathbb{1}) = \mathbb{k}$. Let $\mathbf{M}(\bullet)$ consist of acyclic objects in $\mathbb{k}\text{-mod}^{\text{dg}}$. Then any generator of \mathbf{M} yields $\mathbf{M}_{A_X}^\diamond(\bullet) \simeq \mathbb{k}\text{-mod}^{\text{dg}}$, that is, $\mathbf{M}_{A_X}^\diamond$ is the natural pretriangulated 2-representation, into which $\mathbf{M} = \mathbf{M}^\diamond$ embeds as a proper pretriangulated 2-subrepresentation.

4.4. Morphisms of dg 2-representations and dg algebra 1-morphisms. In this section, we again assume that the dg 2-category \mathcal{C} has compact left adjoints, see Definition 3.3(b), and that \mathbf{M}, \mathbf{N} are compact pretriangulated 2-representations.

Lemma 4.15. *Let A, B be dg algebra 1-morphisms in $\vec{\mathcal{C}}$ and assume given a morphism $\Phi: \mathbf{M}_A \rightarrow \mathbf{M}_B$ of dg 2-representations. Then there exists a dg A-B-bimodule X and a dg isomorphism $\Phi \cong (-) \circ_A X$.*

Proof. Given A, B , let i, j denote the objects such that $A \in \vec{\mathcal{C}}(i, i)$, $B \in \vec{\mathcal{C}}(j, j)$. Assume that we have a morphism of dg 2-representations $\Phi: \mathbf{M}_A \rightarrow \mathbf{M}_B$. We note that Φ induces a morphism $\vec{\mathbf{M}}_A \rightarrow \vec{\mathbf{M}}_B$ of dg 2-representations of $\vec{\mathcal{C}}$ by Lemma 3.5(b), which we will also simply denote by Φ . Let Φ_k denote the dg functor $\vec{\mathbf{M}}_A(k) \rightarrow \vec{\mathbf{M}}_B(k)$ underlying this morphism of dg 2-representations. Define $X := \Phi_i(A) \in \vec{\mathcal{C}}(j, i)$. As $A \in \mathbf{M}_A(i)$, X is a right B-module in $\mathbf{M}_B(i)$. Furthermore, X carries a left A-module structure via the composition

$$A\Phi_i(A) = \vec{\mathbf{M}}_B(A)\Phi_i(A) \cong \Phi_i(\vec{\mathbf{M}}_A(A)A) = \Phi_i(AA) \xrightarrow{\Phi_i(\mu_A)} \Phi_i(A),$$

where the dg isomorphism is given by the natural transformation η included in the data of a morphism of 2-representations. Thus, X is an A-B-bimodule in $\vec{\mathcal{C}}(j, i)$, with $X_B \in \mathbf{M}_B(i)$.

Next, we show that there exists a dg isomorphism $\Phi \cong (-) \circ_A X$. Indeed, for any object Y in $\mathbf{M}_A(k)$, we have

$$\begin{aligned} Y \circ_A X &= \text{coker} \left(YAX \xrightarrow{\rho_Y \circ_0 \text{id}_X - \text{id}_Y \circ_0 \lambda_X} YX \right) \\ &\cong \text{coker} \left(\Phi_k(YAA) \xrightarrow{\Phi_k(\rho_Y) \circ_0 \text{id}_A - \text{id}_Y \circ_0 \Phi_k(\mu_A)} \Phi_k(YA) \right) \\ &\cong \Phi_k \left(\text{coker} \left(YAA \xrightarrow{\rho_Y \circ_0 \text{id}_A - \text{id}_Y \circ_0 \mu_A} YA \right) \right) \\ &\cong \Phi_k(Y). \end{aligned}$$

Here, the second step uses that Φ commutes with the $\vec{\mathcal{C}}$ -action and the definition of the left action on $X = \Phi_i(A)$, the third step uses that the dg functor Φ_k preserves cokernels of dg morphisms (as it is induced by a dg functor from $\mathbf{M}_A(k)$ to $\mathbf{M}_B(k)$), and the final step uses that Y and $Y \circ_A A$ are dg isomorphic. The dg isomorphisms used above are all natural with respect to morphisms of right A-modules in $\vec{\mathcal{C}}$. In fact, a morphism of such modules induces morphisms of the diagrams of the displayed cokernels, and the dg isomorphism η used in the third step is natural in Y . \square

Denoting by $\mathcal{C}\text{-mod}_{\text{int}}^{\text{pre}}$ the dg 2-category of internal dg 2-representations, it follows from Lemma 4.13 and Lemma 4.15 that $\mathcal{C}\text{-mod}_{\text{int}}^{\text{pre}}$ is dg biequivalent to the 2-full dg 2-subcategory of $\mathcal{C}\text{-mod}_{\text{dg}}^{\text{pre}}$ consisting of pretriangulated 2-representations which are dg idempotent complete and cyclic.

Proposition 4.16. *Let $\Phi: \mathbf{M} \rightarrow \mathbf{N}$ be a morphism of dg 2-representations and $\alpha: A \rightarrow B$ be a dg morphism of dg algebra 1-morphisms in $\vec{\mathcal{C}}(i, i)$.*

(a) *For any object X in $\mathbf{M}(i)$, Φ induces a dg morphism of dg algebra 1-morphisms $\alpha_\Phi: A_X \rightarrow A_{\Phi_i X}$.*

- (b) The morphism α induces an essentially surjective morphism of dg 2-representations $\Phi^\alpha: \mathbf{M}_A \rightarrow \mathbf{M}_B$.
- (c) We have $\alpha = \alpha_{\Phi^\alpha}$.
- (d) Φ is full if and only if α_Φ is an epimorphism in $\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})$.
- (e) Φ^α is full if and only if α is an epimorphism in $\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})$.

Proof. (a) For simplicity of notation, we set $Y = \Phi_{\mathfrak{i}} X$. Then, for any $G \in \mathcal{C}(\mathfrak{i}, \mathfrak{i})$, the functor $\Phi_{\mathfrak{i}}$ induces a morphism

$$\mathrm{Hom}_{\mathbf{M}(\mathfrak{i})}(\mathbf{M}(G)X, X) \rightarrow \mathrm{Hom}_{\mathbf{N}(\mathfrak{i})}(\mathbf{N}(G)Y, Y),$$

natural in G . Transferring this via the adjunction isomorphism, we obtain a morphism

$$\mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, [X, X]) \rightarrow \mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, [Y, Y]),$$

again natural in G . This implies that there is a dg morphism $\alpha_\Phi: [X, X] \rightarrow [Y, Y]$, such that the second morphism of morphism spaces is given by postcomposition with α_Φ . Thanks to naturality of all constructions, α_Φ preserves the algebra structure.

- (b) Let $M \in \mathbf{M}_A(\mathfrak{i})$. We define $\Phi^\alpha(M) = M \circ_A B$, noting the natural structure on B as a left dg A -module. It is immediate that this assignment induces a morphism of dg 2-representations. Since \mathbf{M}_B is thick closure of the GB , for all 1-morphisms G in \mathcal{C} , essential surjectivity follows from $GB \cong GA \circ_A B$.
- (c) To prove $\alpha = \alpha_{\Phi^\alpha}$, consider the diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, A) & \xrightarrow{\alpha_{\Phi^\alpha} \circ -} & \mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, B) \\ \parallel \sim & & \parallel \sim \\ \mathrm{Hom}_{\mathbf{M}_A(\mathfrak{i})}(GA, A) & \xrightarrow{\Phi^\alpha} \mathrm{Hom}_{\mathbf{M}_B(\mathfrak{i})}(GA \circ_A B, A \circ_A B) \xrightarrow{\sim} & \mathrm{Hom}_{\mathbf{M}_B(\mathfrak{i})}(GB, B) \end{array}$$

and trace $g \in \mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, A)$ along the path going down, to the right, and then up again. The image of g going down is $\mu_A \circ (g \circ_0 \mathrm{id}_A)$, which is mapped to $(\mu_A \circ_A \mathrm{id}_B) \circ (g \circ_0 \mathrm{id}_A \circ_A \mathrm{id}_B)$ by Φ^α , and then in the natural isomorphism is identified with $\mu_B \circ (\alpha \circ_0 \mathrm{id}_B) \circ (g \circ_0 \mathrm{id}_B) = \mu_B \circ ((\alpha \circ_1 g) \circ_0 \mathrm{id}_B)$. The latter is then sent to $\alpha \circ_1 g$ going back up, and hence $\alpha_{\Phi^\alpha} = \alpha$, as claimed.

- (d) Assume Φ is full. Then for any G in $\mathcal{C}(\mathfrak{i}, \mathfrak{i})$, the morphism

$$\Phi_{\mathfrak{i}}: \mathrm{Hom}_{\mathbf{M}(\mathfrak{i})}(\mathbf{M}(G)X, X) \rightarrow \mathrm{Hom}_{\mathbf{N}(\mathfrak{i})}(\mathbf{N}(G)Y, Y)$$

is an epimorphism, implying that the morphism

$$\mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, [X, X]) \rightarrow \mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, [Y, Y]),$$

is also an epimorphism. This implies that α_Φ is an epimorphism. Conversely, notice that for G in $\mathcal{C}(\mathfrak{i}, \mathfrak{i})$, under the Yoneda embedding, $\mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, -)$ is sent to $\mathrm{Hom}_{\mathcal{C}(\mathfrak{i}, \mathfrak{i})\text{-mod}^{\mathrm{dg}}}(G^\vee, -)$, which is exact and thus $\mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, -)$ is exact. Thus, if α_Φ is an epimorphism, so is $\mathrm{Hom}_{\vec{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(G, \alpha_\Phi)$ as well as the induced morphism

$$\mathrm{Hom}_{\mathbf{M}(\mathfrak{i})}(\mathbf{M}(G)X, X) \rightarrow \mathrm{Hom}_{\mathbf{N}(\mathfrak{i})}(\mathbf{N}(G)Y, Y).$$

- (e) This follows from $\alpha = \alpha_{\Phi^\alpha}$ and (d). \square

We call a dg algebra 1-morphism A **simple** if any dg morphisms of algebra 1-morphisms $A \rightarrow B$ in $\vec{\mathcal{C}}$ which is an epimorphism in $\vec{\mathcal{C}}$ is necessarily an isomorphism.

The following is an analogue of [MMMZ, Corollary 12].

Lemma 4.17. *Let A be a dg algebra 1-morphism in $\vec{\mathcal{C}}$. Then A is simple if and only if \mathbf{M}_A is quotient-simple.*

Proof. Assume A is simple and assume there is a full an essentially surjective morphism of dg 2-representations $\Phi: \mathbf{M}_A \rightarrow \mathbf{N}$. Let $X = \Phi(A)$, then $\mathbf{N} \cong \mathbf{M}_B$ for $B = [X, X]$. By Lemma 4.16(a), this induces a morphisms of dg algebra 1-morphisms $A \rightarrow B$, which is an epimorphism in $\vec{\mathcal{C}}$ by Lemma 4.16(d), since Φ is full. By assumption, this implies that this epimorphism is an isomorphism, and Φ is an equivalence.

Conversely, assume that \mathbf{M}_A is quotient-simple and we have a morphisms of dg algebra 1-morphisms $\alpha: A \rightarrow B$, which is an epimorphism in $\vec{\mathcal{C}}$. Then we obtain an essentially surjective morphism of dg 2-representations $\Phi^\alpha: \mathbf{M}_A \rightarrow \mathbf{M}_B$ by Lemma 4.16(b), which is full by Lemma 4.16(e), since α is an epimorphism. This implies that \mathbf{M}_B is dg equivalent to \mathbf{M}_A/\mathbf{I} for some dg ideal \mathbf{I} in \mathbf{M}_A . Hence, by quotient-simplicity of \mathbf{M}_A , we see that \mathbf{I} is trivial and thus Φ^α is a dg equivalence, which implies that $\alpha = \alpha_{\Phi^\alpha}$ is a dg isomorphism. \square

Denote by \mathcal{C}_i the 2-category on the single object i with morphism category $\mathcal{C}(i, i)$. We have the following corollary, cf. [MMMT, Corollary 4.10].

Corollary 4.18. *There is a bijection between equivalence classes of cyclic dg 2-representations $\mathbf{M} \in \mathcal{C}\text{-mod}_c^{\text{pre}}$ which are \mathcal{C} -generated by some object $X \in \mathbf{M}(i)$ and equivalence classes of cyclic dg 2-representations in $\mathcal{C}_i\text{-mod}_c^{\text{pre}}$.*

This bijection preserves the class of quotient-simple dg 2-representations.

Proof. Let $\mathbf{M} \in \mathcal{C}\text{-mod}_c^{\text{pre}}$ be \mathcal{C} -generated by $X \in \mathbf{M}(i)$. Then $A_X \in \vec{\mathcal{C}}(i, i)$ and hence A_X gives rise to a dg 2-representation of \mathcal{C}_i . Similarly, any dg algebra 1-morphism in $\vec{\mathcal{C}}_i$ is also a dg algebra 1-morphism in $\vec{\mathcal{C}}$, giving the converse map. This yields the first statement.

Since simplicity of A_X is a property formulated inside of the dg category $\vec{\mathcal{C}}(i, i)$, A_X is a simple dg algebra 1-morphism in $\vec{\mathcal{C}}_i$ if and only if it is a simple dg algebra 1-morphism in $\vec{\mathcal{C}}$. Thus, the second statement follows from Lemma 4.17. \square

4.5. Morita equivalence. Two dg algebra 1-morphisms will be called **dg Morita equivalent** provided that \mathbf{M}_A and \mathbf{M}_B are dg equivalent as pretriangulated 2-representations of \mathcal{C} .

Proposition 4.19. *Let $A \in \vec{\mathcal{C}}(i, i)$ and $B \in \vec{\mathcal{C}}(j, j)$ be two dg algebra 1-morphisms. The following are equivalent:*

- (a) *A and B are dg Morita equivalent;*
- (b) *There exist a dg A-B-bimodule $X \in \vec{\mathcal{C}}(j, i)$ and a dg B-A-bimodule $Y \in \vec{\mathcal{C}}(i, j)$ with $Y_A \in \mathbf{M}_A(j)$ and $X_B \in \mathbf{M}_B(i)$ such that $X \circ_B Y \cong A$ and $Y \circ_A X \cong B$.*

Proof. The implication (b) \Rightarrow (a) is obvious, since $- \circ_A X$ and $- \circ_B Y$ provide mutually inverse dg equivalences between \mathbf{M}_A and \mathbf{M}_B .

For the implication (a) \Rightarrow (b) assume that we have a dg equivalence $\Phi: \vec{\mathbf{M}}_A \rightarrow \vec{\mathbf{M}}_B$. Let Φ_k be the dg functor $\mathbf{M}_A(k) \rightarrow \mathbf{M}_B(k)$ underlying this morphism of dg 2-representations. Define $X := \Phi_i(A)$ be the dg A-B-bimodule inducing Φ by Lemma 4.15. We similarly obtain a dg B-A-bimodule Y from the dg functor $\Psi_j: \mathbf{M}_B(j) \rightarrow \mathbf{M}_A(j)$ which is part of the given dg equivalence. The given dg isomorphisms $\Psi_k \circ \Phi_k \cong \text{Id}_{\mathbf{M}_A(k)}$ and $\Phi_k \circ \Psi_k \cong \text{Id}_{\mathbf{M}_B(k)}$ provide dg isomorphisms

$$A \cong \Psi_i \circ \Phi_i(A) \cong \Psi_i(X) \cong X \circ_B Y, \quad \text{and} \quad B \cong \Phi_j \circ \Psi_j(B) \cong \Phi_j(Y) \cong Y \circ_A X$$

as required. \square

Corollary 4.20. *Let \mathcal{C} be a pretriangulated 2-category with compact left adjoints. Further, let \mathbf{M}, \mathbf{N} be two compact pretriangulated 2-representations \mathcal{C} -generated by X and Y , respectively. Then the dg 2-representations \mathbf{M}° and \mathbf{N}° are dg equivalent if and only if the dg algebra 1-morphisms A_X and A_Y are dg Morita equivalent.*

4.6. Morita quasi-equivalence. Let $A \in \vec{\mathcal{C}}(i, i)$ and $B \in \vec{\mathcal{C}}(j, j)$ be dg algebra 1-morphisms and $\Phi: \vec{\mathbf{M}}_A \rightarrow \vec{\mathbf{M}}_B$ a morphism of dg 2-representations. Let $X \in \vec{\mathcal{C}}(j, i)$ be the dg A-B bimodule inducing Φ by Lemma 4.15.

Proposition 4.21. *The morphism of dg 2-representations $\Phi: \vec{\mathbf{M}}_A \rightarrow \vec{\mathbf{M}}_B$ is a quasi-equivalence if and only if the following conditions hold:*

- (a) *the natural maps $\text{Hom}_{\mathbf{M}_A(k)}(\text{FA}, \text{GA}) \rightarrow \text{End}_{\mathbf{M}_B(k)}(\text{FX}, \text{GX})$ are quasi-isomorphisms for all 1-morphisms $F, G \in \mathcal{C}(i, k)$ for any $k \in \mathcal{C}$;*
- (b) *X cyclically generates the homotopy 2-representation \mathbf{KM}_B .*

If we have generators $F_{i,k}$ of $\mathcal{C}(i, k)$, for $k \in \mathcal{C}$, we can simplify (a) to requiring a quasi-isomorphism $\text{End}_{\mathbf{M}_A(k)}(F_{i,k}A) \rightarrow \text{End}_{\mathbf{M}_B(k)}(F_{i,k}X)$.

Proof. First assume that Φ is a quasi-equivalence. As Φ_k sends FA to $\text{FA} \circ_A X \cong \text{FX}$, condition (a) follows. Condition (b) follows since A cyclically generates \mathbf{KM}_A .

For the converse, first note that since Φ_k is a dg functor, it descends to a collection of triangle functors $\mathbf{K}\Phi_k: \mathbf{KM}_A(k) \rightarrow \mathbf{KM}_B(k)$. Then (a) implies that each $\mathbf{K}\Phi_k$ is fully faithful, by [Ke1, Lemma 4.2 (a)], since every object in $\mathbf{KM}_A(k)$ is in the thick closure of the the FA , for $F \in \mathcal{C}(i, k)$. Moreover, $\mathbf{K}\Phi_k$ is dense by (b). \square

We also obtain the following analogue of Corollary 4.11 for \mathcal{C} -quasi-generators.

Corollary 4.22. *Let \mathbf{M} be a pretriangulated 2-representation with a \mathcal{C} -quasi-generator X . Then the morphism of dg 2-representations $[X, -]: \mathbf{M} \rightarrow \vec{\mathbf{M}}_{A_X}$ induces an equivalence between $(\mathbf{KM})^\circ$ and $(\mathbf{KM}_{A_X})^\circ$.*

Proof. By assumption that X is a \mathcal{C} -quasi-generator for \mathbf{M} , it follows that the inclusion of $\mathbf{KG}_M(X)$ into \mathbf{KM} is an equivalence of homotopy 2-representations.

We claim that $(\mathbf{KG}_M(X))^\circ$ is equivalent to $(\mathbf{KM}_{A_X})^\circ$. Note that the functors $\mathcal{K}[X, -]: \mathbf{KM}(i) \rightarrow \mathbf{KM}_{A_X}(i)$ restrict to fully faithful functors $\mathbf{KG}_M(X)(i) \rightarrow \mathbf{KM}_{A_X}(i)$ by Proposition 4.9. The dg isomorphisms $[X, \text{GX}] \cong \text{GA}_X$ are natural in the second component and descend to the homotopy categories. Clearly, all objects GA_X are in the image of these functors. Using fully faithfulness, passing to the idempotent completions implies the claim. Thus, we conclude that the homotopy 2-representations $(\mathbf{KM})^\circ$ and $(\mathbf{KM}_{A_X})^\circ$ are equivalent. \square

Remark 4.23. If the categories $\mathbf{KM}(\mathfrak{i})$ have a bounded t-structure, then they are idempotent complete by [LC] and hence $\mathbf{KM} \simeq (\mathbf{KM})^\circ$. In this case, assuming the setup of Corollary 4.22, consider the fully faithful functors $\mathcal{K}[X, -]: \mathbf{KG}_M(X)(\mathfrak{i}) \rightarrow \mathbf{KM}_{A_X}(\mathfrak{i})$. Given an idempotent e in the image, $e = [X, f]$ for a morphism f in $\mathbf{KM}(\mathfrak{i})$ which is also an idempotent and hence splits off a dg direct summand Y in $\mathbf{KG}_M(X)(\mathfrak{i}) \simeq \mathbf{KM}(\mathfrak{i})$. Thus, e splits off the object $[X, Y]$ as a dg direct summand and, hence, $\mathbf{KM}_{A_X}(\mathfrak{i})$ is idempotent complete. Hence, if \mathbf{M} has a bounded t-structure, the morphism of dg 2-representations $[X, -]: \mathbf{M} \rightarrow \mathbf{M}_{A_X}$ is a quasi-equivalence provided that X is a \mathcal{C} -quasi-generator.

Algebraic conditions on the categories $\mathbf{M}(\mathfrak{i})$ which imply the existence of a bounded t-structure are provided in [Sch, Theorem 1]. Namely, if A is a positively graded dg algebra with A_0 semisimple such that $\partial(A_0) = 0$, then $\mathcal{K}(A\text{-csf})$ is closed under taking direct summands.

4.7. Restriction and pushforward. Let \mathcal{C}, \mathcal{D} be dg 2-categories and $\mathbf{F}: \mathcal{C} \rightarrow \mathcal{D}$ a dg 2-functor.

We obtain a dg 2-functor $\mathbf{R} = \mathbf{R}_F: \mathcal{D}\text{-mod}_{\text{int}}^{\text{pre}} \rightarrow \mathcal{C}\text{-mod}^{\text{pre}}$ by precomposition, i.e. $\mathbf{R}(\mathbf{M}) = \mathbf{M} \circ \mathbf{F}$.

On the other hand, we can consider a dg 2-functor $\mathbf{P} = \mathbf{P}_F: \mathcal{C}\text{-mod}_{\text{int}}^{\text{pre}} \rightarrow \mathcal{D}\text{-mod}_{\text{int}}^{\text{pre}}$, by defining $\mathbf{P}(\mathbf{M}_A) = \mathbf{M}_{F(A)}$.

Lemma 4.24. *Given $\mathbf{M}_A \in \mathcal{C}\text{-mod}_{\text{int}}^{\text{pre}}$, there is a morphism of dg 2-representations $\Phi: \mathbf{M} \rightarrow \mathbf{RPM}$ induced by \mathbf{F} .*

Proof. Writing $\mathbf{RPM} = \mathbf{RM}_{F(A)}$, we define Φ_j by mapping an object $X \in \mathbf{M}(j)$ to $\mathbf{F}(X)$ and similarly on morphisms. It is straightforward to check that this defines a morphism of dg 2-representations. \square

Let $\mathbf{M}_B \in \mathcal{D}\text{-mod}_{\text{int}}^{\text{pre}}$. Assuming that $B \in \mathbf{M}(\mathfrak{i})$ is a \mathcal{C} -generator for \mathbf{RM}_B , we have a canonical choice of internal dg 2-representation of \mathcal{C} dg equivalent to \mathbf{RM}_B defined by $\mathbf{M}_{C'}$, for $C' = \mathcal{C}[B, B]$, where we distinguish internal homs in \mathcal{C} and \mathcal{D} by denoting them by $\mathcal{C}[-, -]$ and $\mathcal{D}[-, -]$, respectively. We can then define \mathbf{PRM}_B as $\mathbf{PM}_{C'}$.

Lemma 4.25. *Let $\mathbf{M} = \mathbf{M}_B \in \mathcal{D}\text{-mod}_{\text{int}}^{\text{pre}}$ and assume that $B \in \mathbf{M}(\mathfrak{i})$ is a \mathcal{C} -generator for \mathbf{RM} . Then there is an essentially surjective morphism of dg 2-representations $\Psi: \mathbf{PRM}_B \rightarrow \mathbf{M}_B$.*

Proof. Set $C' = \mathcal{C}[B, B]$, the dg algebra 1-morphism associated to B , and $C = \mathbf{F}(C')$. By assumption that B is a \mathcal{C} -generator for \mathbf{RM} , the latter is dg equivalent to $\mathbf{M}_{C'}$ by Corollary 4.11 as the restriction is dg idempotent complete. Then, by construction, we have an equivalence of dg 2-representations $\mathbf{PRM}_B \rightarrow \mathbf{M}_{C'}$. We define the map $\varphi: C \rightarrow B$ as the image of $\text{id}_{C'}$ under the chain of isomorphisms

$$\begin{aligned} \text{Hom}_{\overline{\mathcal{C}}(\mathfrak{i}, \mathfrak{i})}(C', C') &\cong \text{Hom}_{\overline{\mathbf{RM}}(\mathfrak{i})}(\mathbf{RM}(C')B, B) \\ &= \text{Hom}_{\overline{\mathbf{M}}(\mathfrak{i})}(\mathbf{F}(C')B, B) = \text{Hom}_{\overline{\mathbf{M}}(\mathfrak{i})}(CB, B) \\ &\cong \text{Hom}_{\overline{\mathcal{D}}(\mathfrak{i}, \mathfrak{i})}(C, B). \end{aligned}$$

Recalling that the inverse dg isomorphisms

$$\text{Hom}_{\overline{\mathbf{M}}(\mathfrak{i})}(\mathbf{F}B, B) \cong \text{Hom}_{\overline{\mathcal{D}}(\mathfrak{i}, \mathfrak{i})}(\mathbf{F}, B)$$

for $F \in \mathcal{C}(\mathbf{i}, \mathbf{i})$ are given by

$$(4.1) \quad \alpha: \begin{array}{ccc} g & \mapsto & g \circ_1 (\text{id}_F \circ_0 u_B) \\ m_B \circ_1 (f \circ_0 \text{id}_B) & \leftarrow & f \end{array} : \beta,$$

we see that, explicitly, $\varphi = \alpha(\text{ev}_{BB}^{\mathcal{C}}) = \text{ev}_{BB}^{\mathcal{C}} \circ_1 (\text{id}_C \circ_0 u_B)$.

We claim that this is an algebra homomorphism, i.e. $\varphi \circ_1 m_C = m_B \circ_1 (\varphi \circ_0 \varphi)$. In order to prove this, we consider the images of both sides of the claimed equality under the dg isomorphism

$$\text{Hom}_{\overline{\mathcal{D}}(\mathbf{i}, \mathbf{i})}(\text{CC}, B) \cong \text{Hom}_{\overline{\mathbf{M}}(\mathbf{i})}(\text{CCB}, B).$$

The strategy is to show that the morphisms on both sides of the claimed equation correspond to $\text{ev}_{BB}^{\mathcal{C}} \circ_1 (\text{id}_C \circ_0 \text{ev}_{BB}^{\mathcal{C}})$.

First, observe that

$$(4.2) \quad m_B \circ_1 (\varphi \circ_0 \text{id}_B) = \beta(\varphi) = \beta\alpha(\text{ev}_{BB}^{\mathcal{C}}) = \text{ev}_{BB}^{\mathcal{C}}.$$

On the one hand, $m_B \circ_1 (\varphi \circ_0 \varphi)$ corresponds to

$$m_B \circ_1 ((m_B \circ_1 (\varphi \circ_0 \varphi)) \circ_0 \text{id}_B).$$

Consider the diagram

$$\begin{array}{ccccc} \text{CCB} & \xrightarrow{\varphi \circ_0 \text{id}_{CB}} & \text{BCB} & \xrightarrow{\text{id}_B \circ_0 \varphi \circ_0 \text{id}_B} & \text{BBB} & \xrightarrow{m_B \circ_0 \text{id}_B} & \text{BB} \\ & \searrow \text{id}_C \circ_0 \text{ev}_{BB}^{\mathcal{C}} & & \searrow \text{id}_B \circ_0 \text{ev}_{BB}^{\mathcal{C}} & \downarrow \text{id}_B \circ_0 m_B & & \downarrow m_B \\ & & \text{CB} & \xrightarrow{\varphi \circ_0 \text{id}_B} & \text{BB} & \xrightarrow{m_B} & \text{BB} \\ & & & \searrow \text{ev}_{BB}^{\mathcal{C}} & & & \downarrow m_B \\ & & & & & & B \end{array}$$

where the two triangles commute by (4.2), the leftmost square commutes by the interchange law, and the rightmost square commutes by associativity of m_B . This shows that $m_B \circ_1 (\varphi \circ_0 \varphi)$ corresponds to $\text{ev}_{BB}^{\mathcal{C}} \circ_1 (\text{id}_C \circ_0 \text{ev}_{BB}^{\mathcal{C}})$ as claimed.

On the other hand, $\varphi \circ_1 m_C$ corresponds to

$$\begin{aligned} m_B \circ_1 ((\varphi \circ_1 m_C) \circ_0 \text{id}_B) &= m_B \circ_1 (\varphi \circ_0 \text{id}_B) \circ_1 (m_C \circ_0 \text{id}_B) \\ &= \text{ev}_{BB}^{\mathcal{C}} \circ_1 (m_C \circ_0 \text{id}_B), \end{aligned}$$

where the second equality uses (4.2). Now consider the definition of $m_C = \mathbf{F}(m_{C'})$. Then the map $m_C \circ_0 \text{id}_B: \text{CCB} \rightarrow \text{CB}$ in $\overline{\mathbf{M}}(\mathbf{i})$ is really the map $\mathbf{F}(m_{C'}) \circ_0 \text{id}_B: \mathbf{F}(C'C)B \rightarrow \mathbf{F}(C')B$ in $\overline{\mathbf{M}}(\mathbf{i})$, which, by definition of \mathbf{R} , describes the action of $m_{C'}$ via $\mathbf{R}m_B$. By associativity of the action of $\mathbf{R}m_B(C')$ via $\text{ev}_{BB}^{\mathcal{C}}$, we have

$$\text{ev}_{BB}^{\mathcal{C}} \circ_1 (\mathbf{F}(m_{C'})) \circ_0 \text{id}_B = \text{ev}_{BB}^{\mathcal{C}} \circ_1 (\mathbf{F}(\text{id}_{C'}) \circ_0 \text{ev}_{BB}^{\mathcal{C}}) = \text{ev}_{BB}^{\mathcal{C}} \circ_1 (\text{id}_C \circ_0 \text{ev}_{BB}^{\mathcal{C}})$$

as claimed.

Since φ defines a left dg C-module structure on B, we obtain a morphism of dg 2-representations $\mathbf{M}_C \rightarrow \mathbf{M}_B$ given by $- \circ_C B$. As this morphism sends C to B, it is essentially surjective. \square

4.8. Local quasi-equivalences and dg 2-representations. Throughout this subsection, we use the the same setup as in the previous one, and additionally assume that the dg 2-functor $\mathbf{F}: \mathcal{C} \rightarrow \mathcal{D}$ is a local quasi-equivalence and, moreover, essentially surjective on objects. That is, for any object j in \mathcal{D} , there exists an object i in \mathcal{C} such that $\mathbf{P}_k = \mathcal{D}(k, -)$ is dg equivalent to $\mathbf{P}_{\mathbf{F}(i)} = \mathcal{D}(\mathbf{F}(i), -)$ through composing with 1-morphisms in \mathcal{D} .

Proposition 4.26. *Under the assumptions on \mathbf{F} as above, $\Phi: \mathbf{M}_A \rightarrow \mathbf{RPM}_A$ is a quasi-equivalence of dg 2-representations.*

Proof. Recall that, if $A \in \mathcal{C}(i, i)$, $\mathbf{M}_A(j)$ is the thick closure of the GA with $G \in \mathcal{C}(i, j)$ and $\mathbf{RPM}_A(j)$ is the thick closure of the $\mathbf{HF}(A)$ for $H \in \mathcal{D}(\mathbf{F}(i), \mathbf{F}(j))$. Since $\mathcal{K}\mathbf{F}_{i,j}$ is essentially surjective, we can find $G \in \mathcal{C}(i, j)$ such that $H \in \mathcal{D}(\mathbf{F}(i), \mathbf{F}(j))$ is quasi-isomorphic to $\mathbf{F}(G)$ and hence $\mathbf{HF}(A)$ is quasi-isomorphic to $\mathbf{F}(G)\mathbf{F}(A) = \mathbf{F}(GA)$, so $\mathcal{K}\Phi(i)$ is also essentially surjective. Moreover, we have a commutative diagram

$$(4.3) \quad \begin{array}{ccc} \mathrm{Hom}_{\mathbf{M}_A(j)}(GA, G'A) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathcal{D}(i,j)}(G, G'A) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_{\mathbf{RPM}_A(j)}(\mathbf{F}(GA), \mathbf{F}(G'A)) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathcal{D}(i,j)}(\mathbf{F}(G), \mathbf{F}(G'A)). \end{array}$$

Since the right vertical arrow is a quasi-isomorphism by assumption, so is the left vertical arrow. This proves the lemma. \square

Proposition 4.27. *Assume, in addition, that B is a \mathcal{C} -generator for \mathbf{RM}_B . Then $\Psi: \mathbf{PRM}_B \rightarrow \mathbf{M}_B$ is a quasi-equivalence.*

Proof. It suffices to prove that $- \circ_C B$ is a quasi-equivalence. By Lemma 4.25, this morphism of dg 2-representation is essentially surjective. Thus, it suffices to show that it induces quasi-isomorphisms of morphism spaces. Recall that $\varphi = \mathrm{ev}_{\mathbf{BB}}^{\mathcal{C}} \circ_1 (\mathrm{id}_C \circ_0 u_B)$.

We first claim that the diagram

$$(4.4) \quad \begin{array}{ccc} \mathrm{Hom}_{\mathbf{M}_C(j)}(GC, HC) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathcal{D}(j)}(G, HC) \\ - \circ_C \mathrm{id}_B \downarrow & & \downarrow (\mathrm{id}_H \circ_0 \varphi) \circ_1 - \\ \mathrm{Hom}_{\mathbf{M}_B(j)}(GB, HB) & \xrightarrow{\cong} & \mathrm{Hom}_{\mathcal{D}(j)}(G, HB) \end{array}$$

commutes. Indeed, using $\mathrm{ev}_{\mathbf{CC}}^{\mathcal{C}} = m_C$, a morphism $f \in \mathrm{Hom}_{\mathcal{D}(j)}(G, HC)$ corresponds to

$$GC \xrightarrow{f \circ_0 \mathrm{id}_C} HCC \xrightarrow{\mathrm{id}_H \circ_0 m_C} HC$$

in $\mathrm{Hom}_{\mathbf{M}_C(j)}(GC, HC)$, which, under $- \circ_C \mathrm{id}_B$ maps to

$$GB \xrightarrow{f \circ_0 \mathrm{id}_B} HCB \xrightarrow{\mathrm{id}_H \circ_0 \mathrm{ev}_{\mathbf{BB}}^{\mathcal{C}}} HB.$$

Here we have used that $m_C \circ_C \mathrm{id}_B = \mathrm{ev}_{\mathbf{BB}}^{\mathcal{C}}$ since both, by construction, correspond to the action on B when viewed as a left C -module.

The above morphism then corresponds to

$$G \xrightarrow{\mathrm{id}_G \circ_0 u_B} GB \xrightarrow{f \circ_0 \mathrm{id}_B} HCB \xrightarrow{\mathrm{id}_H \circ_0 \mathrm{ev}_{\mathbf{BB}}^{\mathcal{C}}} HB$$

in $\text{Hom}_{\vec{\mathcal{D}}(j)}(G, \text{HB})$. Given

$$(f \circ_0 \text{id}_B) \circ_1 (\text{id}_G \circ_0 u_B) = (\text{id}_{\text{HC}} \circ_0 u_B) \circ_1 f,$$

we see that

$$\begin{aligned} (\text{id}_H \circ_0 \text{ev}_{\text{BB}}^{\mathcal{C}}) \circ_1 (f \circ_0 \text{id}_B) \circ_1 (\text{id}_G \circ_0 u_B) &= (\text{id}_H \circ_0 (\text{ev}_{\text{BB}}^{\mathcal{C}} \circ_1 (\text{id}_G \circ_0 u_B))) \circ_1 f \\ &= (\text{id}_H \circ_0 \varphi) \circ_1 f, \end{aligned}$$

proving that the diagram in (4.4) commutes.

It thus suffices to show that the right vertical map in (4.4) is a quasi-isomorphism. To see this, first assume $G = \mathbf{F}(G')$ and $H = \mathbf{F}(H')$ and consider the diagram

$$\begin{array}{ccc} \text{Hom}_{\vec{\mathcal{C}}(i,j)}(G', H'C') & \xrightarrow{\mathbf{F}(-)} & \text{Hom}_{\vec{\mathcal{D}}(j)}(G, \text{HC}) \\ \downarrow \cong & & \uparrow \\ \text{Hom}_{\mathbf{M}_B(j)}(\text{GB}, \text{HB}) & & \\ \downarrow \cong & & \swarrow (\text{id}_H \circ_0 \varphi) \circ_1 - \\ \text{Hom}_{\vec{\mathcal{D}}(j)}(G, \text{HB}) & & \end{array}$$

Under the vertical isomorphisms, $f \in \text{Hom}_{\vec{\mathcal{C}}(i,j)}(G', H'C)$ is first mapped to $(\text{id}_H \circ_0 \text{ev}_{\text{BB}}^{\mathcal{C}}) \circ_1 (\mathbf{F}(f) \circ_0 \text{id}_B)$ and subsequently, using the same arguments as before, to

$$(\text{id}_H \circ_0 \text{ev}_{\text{BB}}^{\mathcal{C}}) \circ_1 (\mathbf{F}(f) \circ_0 \text{id}_B) \circ_1 (\text{id}_G \circ_0 u_B) = (\text{id}_H \circ_0 \varphi) \circ_1 \mathbf{F}(f),$$

and hence the diagram commutes. Given that the vertical maps are isomorphisms, and the horizontal map is a quasi-isomorphism, the map $(\text{id}_H \circ_0 \varphi) \circ_1 -$ is also a quasi-isomorphism as desired in this case.

For general G, H , there exist G', H' and zigzags of quasi-isomorphisms between G and $\mathbf{F}(G')$ as well as H and $\mathbf{F}(H')$. The quasi-isomorphism

$$(\text{id}_{\mathbf{F}(H')} \circ_0 \varphi) \circ_1 - : \text{Hom}_{\vec{\mathcal{D}}(j)}(\mathbf{F}(G'), \mathbf{F}(H')C) \rightarrow \text{Hom}_{\vec{\mathcal{D}}(j)}(\mathbf{F}(G'), \mathbf{F}(H')B)$$

propagates along these to imply that

$$(\text{id}_H \circ_0 \varphi) \circ_1 - : \text{Hom}_{\vec{\mathcal{D}}(j)}(G, \text{HC}) \rightarrow \text{Hom}_{\vec{\mathcal{D}}(j)}(G, \text{HB})$$

is also a quasi-isomorphism. This completes the proof. \square

Even if B does not \mathcal{C} -generate \mathbf{RM}_B as assumed in Lemma 4.25 and Proposition 4.27, \mathbf{F} being a local quasi-equivalence and essentially surject on objects implies that it is equivalent to a \mathcal{C} -quasi-generator.

Lemma 4.28. *The dg algebra 1-morphism B corresponds to a \mathcal{C} -quasi-generator for \mathbf{RM}_B .*

Proof. By assumption that \mathbf{F} is essentially surjective on objects, we can assume w.l.o.g. that $B \in \mathcal{D}(\mathbf{F}(k), \mathbf{F}(k))$, for an object k of \mathcal{C} . We need to show that the inclusion $I: \mathbf{GRM}_B(B) \hookrightarrow \mathbf{RM}_B$ is a quasi-equivalence. As I is fully faithful, it remains to show that \mathbf{KI} is essentially surjective. That is, we claim that every object X in $\mathbf{KRM}_B(j)$ is isomorphic to an object in the thick closure of $\{\mathbf{F}(G)B \mid G \in \mathcal{C}(j, k)\}$ in the homotopy category $\mathbf{KRM}_B(j)$. Note that $X \in \mathbf{RM}_B(j) = \mathbf{M}_B(\mathbf{F}(j))$ is dg isomorphic to an object in the thick closure of $\{\text{HB} \mid H \in \mathcal{D}(\mathbf{F}(j), \mathbf{F}(k))\}$. Since any 1-morphism H of \mathcal{D} is homotopy isomorphic to $\mathbf{F}(G)$ for some 1-morphism G of \mathcal{C} , the claim follows. \square

Pullback and pushforward preserve quasi-equivalences of internal dg 2-representations, see Section 4.6.

Proposition 4.29. *Retain the assumptions on $\mathbf{F}: \mathcal{C} \rightarrow \mathcal{D}$ of this section.*

- (a) *For any quasi-equivalence $\Phi: \mathbf{M}_{A_1} \rightarrow \mathbf{M}_{A_2}$ of internal dg 2-representations of \mathcal{C} , $\mathbf{P}(\Phi)$ is a quasi-equivalence.*
- (b) *For any quasi-equivalence $\Psi: \mathbf{M}_{B_1} \rightarrow \mathbf{M}_{B_2}$ of internal dg 2-representations of \mathcal{D} , $\mathbf{R}(\Psi)$ is a quasi-equivalence.*

Proof. To prove Part (a), recall Proposition 4.21 and denote by X a dg bimodule giving the quasi-equivalence $\Phi: \mathbf{M}_{A_1} \rightarrow \mathbf{M}_{A_2}$. That is, $\Phi(A_1) \cong X$, which implies that $\mathbf{P}(\Phi)(\mathbf{F}(A_1)) \cong \mathbf{F}(X)$. By Proposition 4.21, A_2 appears in the thick closure of $\{GX|G\}$ in \mathbf{KM}_{A_2} , for G varying over 1-morphisms of \mathcal{C} . Thus, $\mathbf{F}(A_2)$ appears in the thick closure of $\{\mathbf{F}(G)\mathbf{F}(X)|G\}$ in $\mathbf{KM}_{\mathbf{F}(A_2)}$. As $\mathbf{F}(A_2)$ generates \mathbf{PM}_{A_2} by definition, we see that the second condition in Proposition 4.21 holds for $\mathbf{F}(X)$.

As in the diagram (4.3), we have quasi-isomorphisms

$$\mathrm{Hom}_{\mathbf{M}_{A_1}(k)}(\mathbf{F}A_1, \mathbf{G}A_1) \longrightarrow \mathrm{Hom}_{\mathbf{M}_{\mathbf{F}(A_1)}(\mathbf{F}k)}(\mathbf{F}(\mathbf{F})\mathbf{F}(A_1), \mathbf{F}(\mathbf{G})\mathbf{F}(A_1)).$$

Thus, we have the following commutative diagram of dg morphisms

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbf{M}_{A_1}(k)}(\mathbf{F}A_1, \mathbf{G}A_1) & \longrightarrow & \mathrm{End}_{\mathbf{M}_{A_2}(k)}(\mathbf{F}X, \mathbf{G}X) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_{\mathbf{M}_{\mathbf{F}(A_1)}(\mathbf{F}k)}(\mathbf{F}(\mathbf{F})\mathbf{F}(A_1), \mathbf{F}(\mathbf{G})\mathbf{F}(A_1)) & \longrightarrow & \mathrm{End}_{\mathbf{M}_{\mathbf{F}(A_2)}(\mathbf{F}k)}(\mathbf{F}(\mathbf{F})\mathbf{F}(X), \mathbf{F}(\mathbf{G})\mathbf{F}(X)), \end{array}$$

where the vertical morphisms are quasi-isomorphisms, and the top vertical morphism is a quasi-isomorphism as $\Phi: \mathbf{M}_{A_1} \rightarrow \mathbf{M}_{A_2}$ is a quasi-equivalence. Thus, the bottom vertical morphism is a quasi-isomorphism. For general 1-morphisms H, K in \mathcal{D} , we have homotopy isomorphisms $\mathbf{F}(F) \cong H$ and $\mathbf{F}(G) \cong K$. Composing with these homotopy isomorphisms we obtain the commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbf{PM}_{A_1}(k)}(\mathbf{F}(\mathbf{F})\mathbf{F}(A_1), \mathbf{F}(\mathbf{G})\mathbf{F}(A_1)) & \longrightarrow & \mathrm{End}_{\mathbf{PM}_{A_2}(k)}(\mathbf{F}(\mathbf{F})\mathbf{F}(X), \mathbf{F}(\mathbf{G})\mathbf{F}(X)) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_{\mathbf{PM}_{A_1}(k)}(\mathbf{H}\mathbf{F}(A_1), \mathbf{K}\mathbf{F}(A_1)) & \longrightarrow & \mathrm{End}_{\mathbf{PM}_{A_2}(k)}(\mathbf{H}\mathbf{F}(X), \mathbf{K}\mathbf{F}(X)), \end{array}$$

where the vertical morphisms are quasi-isomorphism by the above. As the top horizontal morphisms is a quasi-isomorphism, so is the bottom morphism.

Hence, both conditions of Proposition 4.21 are satisfied for the dg bimodule $\mathbf{F}(X)$, which implies that the morphism $\mathbf{P}(\Phi)$ from $\mathbf{PM}_{A_1} \cong \mathbf{M}_{\mathbf{F}(A_1)}$ to $\mathbf{PM}_{A_2} \cong \mathbf{M}_{\mathbf{F}(A_2)}$ is a quasi-equivalence and Part (a) follows.

To prove Part (b), assume given a quasi-equivalence $\Psi: \mathbf{M}_{B_1} \rightarrow \mathbf{M}_{B_2}$ of internal dg 2-representations of \mathcal{D} . Since \mathbf{F} is essentially surjective on objects, w.l.o.g., $B_1 \in \mathcal{D}(\mathbf{F}(i), \mathbf{F}(i))$ and $B_2 \in \mathcal{D}(\mathbf{F}(j), \mathbf{F}(j))$ for objects i, j in \mathcal{C} . By Proposition 4.21, Ψ corresponds to composing with a dg B_1 - B_2 -bimodule Y in $\vec{\mathcal{D}}$ which descends to a cyclic generator for \mathbf{KM}_{B_2} and induces quasi-isomorphisms on morphism spaces. We observe that

$$\Psi(B_1) \cong Y \in (\mathbf{G}_{\mathbf{M}_{B_2}}(B_2))(\mathbf{F}(j)) = (\mathbf{G}_{\mathbf{RM}_{B_2}}(B_2))(j).$$

By Lemma 4.28, B_2 cyclically generates \mathbf{KRM}_{B_2} . Now, as \mathbf{F} is a local quasi-equivalence, B_2 is in the thick closure of $\{\mathbf{F}(G)Y|G\}$ in \mathbf{KRM}_{B_2} . Thus, Y cyclically generates \mathbf{KRM}_{B_2} . Moreover, the quasi-isomorphisms

$$\mathrm{Hom}_{\mathbf{M}_{B_1}(\mathbf{k})}(\mathbf{H}B_1, \mathbf{K}B_1) \xrightarrow{\Psi} \mathrm{Hom}_{\mathbf{M}_{B_2}(\mathbf{k})}(\mathbf{H}Y, \mathbf{K}Y),$$

for 1-morphisms H, K in \mathcal{D} , induced by Ψ provide quasi-isomorphisms

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbf{RM}_{B_1}(\mathbf{h})}(\mathbf{F}(\mathbf{F})B_1, \mathbf{F}(\mathbf{G})B_1) & \xrightarrow{\mathbf{R}\Psi} & \mathrm{Hom}_{\mathbf{RM}_{B_2}(\mathbf{h})}(\mathbf{F}(\mathbf{F})Y, \mathbf{F}(\mathbf{G})Y) \\ \parallel & & \parallel \\ \mathrm{Hom}_{\mathbf{M}_{B_1}(\mathbf{Fh})}(\mathbf{F}(\mathbf{F})B_1, \mathbf{F}(\mathbf{G})B_1) & \xrightarrow{\Psi} & \mathrm{Hom}_{\mathbf{M}_{B_2}(\mathbf{Fh})}(\mathbf{F}(\mathbf{F})Y, \mathbf{F}(\mathbf{G})Y) \end{array}$$

when restricting to acting by 1-morphisms of the form $\mathbf{F}(\mathbf{F})$, $\mathbf{F}(\mathbf{G})$, for 1-morphisms F, G in \mathcal{C} , and \mathbf{h} an object of \mathcal{C} . Thus, $\mathbf{R}(\Psi)$ is a quasi-equivalence of dg 2-representations of \mathcal{C} . \square

The results of this section imply the following corollary.

Corollary 4.30. *Assume given a dg 2-functor $\mathbf{F}: \mathcal{C} \rightarrow \mathcal{D}$ that is a local quasi-equivalence and essentially surjective on objects.*

- (a) *For every internal dg 2-representation \mathbf{M}_B of \mathcal{D} there exists an internal dg 2-representation \mathbf{M}_A of \mathcal{C} and a quasi-equivalence of dg 2-representations of \mathcal{D} from \mathbf{PM}_A to \mathbf{M}_B .*
- (b) *For every internal dg 2-representation \mathbf{M}_A of \mathcal{C} there is a quasi-equivalence of dg 2-representations of \mathcal{C} from \mathbf{M}_A to $\mathbf{RM}_{\mathbf{F}(A)}$.*

Proof. We first proof Part (a). Assume given an internal dg 2-representation \mathbf{M}_B of \mathcal{D} and, w.l.o.g., $B \in \mathcal{D}(\mathbf{F}(\mathbf{k}), \mathbf{F}(\mathbf{k}))$. Set $C' = {}_{\mathcal{C}}[B, B]$ and $C = \mathbf{F}(C')$, then $\mathbf{PM}_{C'} = \mathbf{M}_C$. The proof of Lemma 4.25 shows that, even if B is not a \mathcal{C} -generator for \mathbf{RM}_B , there is a natural morphism of dg 2-representations $\Psi: \mathbf{M}_C \rightarrow \mathbf{M}_B$ which, by the proof of Proposition 4.27, is quasi-equivalence. Setting $A = C'$, Part (a) follows. Part (b) directly follows from Proposition 4.26. \square

5. EXAMPLES

5.1. The dg 2-category \mathcal{C}_A . Let $A = A_1 \times \cdots \times A_n$ be a finite-dimensional dg algebra, where each A_i is indecomposable as a \mathbb{k} -algebra.

Let \mathcal{A}_i be a small dg category dg equivalent to $\widehat{\{A_i\}}$ inside $A_i\text{-mod}^{\mathrm{dg}}$, and set $\mathcal{A} = \coprod_{i=1}^n \mathcal{A}_i$. Note that, by definition, \mathcal{A} is a pretriangulated category.

Definition 5.1. We define \mathcal{C}_A as the dg 2-category with

- objects $1, \dots, n$ where we identify i with \mathcal{A}_i ;
- 1-morphisms in $\mathcal{C}_A(i, j)$ are all functors dg isomorphic to tensoring with dg A_j - A_i -bimodules in the thick closure of $A_j \otimes_{\mathbb{k}} A_i$, if $i \neq j$, and in the thick closure of $A_i \oplus A_i \otimes_{\mathbb{k}} A_i$, if $i = j$;
- 2-morphisms all natural transformations of such functors.

We define the **natural 2-representation** \mathbf{N} of \mathcal{C}_A as its defining action on \mathcal{A} , that is, $\mathbf{N}(i) = \mathcal{A}_i$, $\mathbf{N}(F) = F$ for a 1-morphism F , and $\mathbf{N}(\alpha) = \alpha$ for a 2-morphism α . By

construction, \mathcal{C}_A is a pretriangulated 2-category, and \mathbf{N} is in $\mathcal{C}_A\text{-mod}^{\text{pre}}$. It is cyclic with any $A_i \in \mathbf{N}(i) = \mathcal{A}_i$ as a \mathcal{C}_A -generator.

We further denote by $F_{j,i}$ the functor given by tensoring with $A_j \otimes_{\mathbb{k}} A_i$.

Lemma 5.2. *Choosing A_i as a \mathcal{C}_A -generator for \mathbf{N} , the corresponding algebra dg 1-morphism A_{A_i} is given by (tensoring with) the dg bimodule $A_i^* \otimes_{\mathbb{k}} A_i$.*

We remark that A_{A_i} is indeed a dg 1-morphism in $\overrightarrow{\mathcal{C}_A}$ by our assumption of finite-dimensionality of A_i . In fact, \mathcal{C}_A has compact left adjoints and \mathbf{N} is compact as required in Section 4.3.

Proof. First notice that since $A_i \in \mathbf{N}(i)$, the dg algebra 1-morphism A_{A_i} is thus in $\overrightarrow{\mathcal{C}_A}(i, i)$ and we only need to apply Ev_{A_i} to the generator of this dg category, which is given by $\mathbb{1}_i \oplus F_{i,i}$. We thus verify that

$$\begin{aligned} \text{Hom}_{\overrightarrow{\mathcal{C}_A}(i,i)}(\mathbb{1}_i \oplus F_{i,i}, A_{A_i}) &\cong \text{Hom}_{A_i\text{-mod}^{\text{dg}}-A_i}(A_i \oplus A_i \otimes_{\mathbb{k}} A_i, A_i^* \otimes_{\mathbb{k}} A_i) \\ &\cong \text{Hom}_{A_i\text{-mod}^{\text{dg}}-A_i}(A_i \oplus A_i \otimes_{\mathbb{k}} A_i, \text{Hom}_{\mathbb{k}}(A_i^{\mathbb{k}}) \otimes_{\mathbb{k}} A_i) \\ &\cong \text{Hom}_{A_i\text{-mod}^{\text{dg}}-A_i}(A_i \oplus A_i \otimes_{\mathbb{k}} A_i, \text{Hom}_{\mathbb{k}}(A_i, A_i)) \\ &\cong \text{Hom}_{A_i\text{-mod}^{\text{dg}}}(A_i \oplus A_i \otimes_{\mathbb{k}} A_i, A_i) \\ &\cong \text{Hom}_{\mathbf{N}(i)}(\text{Ev}_{A_i}(\mathbb{1}_i \oplus F_{i,i}), A_i), \end{aligned}$$

which proves the claim. \square

Hence, as $\mathbf{N}(j)$ is dg idempotent complete for any object j , Corollary 4.11 implies that \mathbf{N} is dg equivalent to the internal dg 2-representation $\mathbf{M}_{A_{A_i}}$, for any i .

As a first example, we can consider the dg 2-category $\mathcal{C}_{\mathbb{k}}$. In this case, $\overrightarrow{\mathcal{C}_{\mathbb{k}}}$ and $\mathcal{C}_{\mathbb{k}}$ are dg biequivalent and a dg algebra 1-morphism in $\mathcal{C}_{\mathbb{k}}$ corresponds to a finite-dimensional dg algebra over \mathbb{k} . Results of [Or2] imply the following corollary.

Proposition 5.3. *There is a unique non-acyclic quotient-simple pretriangulated 2-representations of $\mathcal{C}_{\mathbb{k}}$ up to dg equivalence.*

Proof. By Lemma 4.17, a quotient simple 2-representation \mathbf{M} corresponds to a dg \mathbb{k} -algebra A that does not have any proper dg ideals. Using [Or2], this in particular implies that the dg ideal denoted by J_+ in loc. cit. is zero, or $J_+ = A$, in which case the dg algebra and the hence the dg 2-representation are acyclic. Thus, if \mathbf{M} is not acyclic, the (ungraded) \mathbb{k} -algebra underlying A is a product of matrix rings by [Or2, Proposition 2.16]. It is remarked in the proof of loc. cit. that the central idempotents defining the individual factors are annihilated by ∂ , and hence simplicity of A implies that the \mathbb{k} -algebra underlying A is isomorphic to $M_n(\mathbb{k})$ for some n .

Given a (non-zero) dg \mathbb{k} -module V , $V^* \otimes V$ is a dg \mathbb{k} -algebra with underlying \mathbb{k} -algebra isomorphic to $M_n(\mathbb{k})$. As a dg algebra 1-morphism in $\mathcal{C}_{\mathbb{k}}$, $V^* \otimes V$ appears as the internal hom $[V, V]$ obtained from \mathbf{N} with V as a (weak) $\mathcal{C}_{\mathbb{k}}$ -generator similarly to Lemma 5.2. In fact, all differentials on matrix rings are of this form by [Or2, Proposition 2.15]. Thus, A is isomorphic to $V^* \otimes V$ as a dg algebra, for some dg \mathbb{k} -module V .

It remains to show that the pretriangulated 2-representations $\mathbf{M}_{V^* \otimes V}$ are dg equivalent for any choice of V . For this, we apply Proposition 4.19 to V viewed as a \mathbb{k} - $V^* \otimes V$ -bimodule and V^* viewed as a $V^* \otimes V$ - \mathbb{k} -bimodule. Since $V \otimes_{V^* \otimes V} V^* \cong \mathbb{k}$ is an isomorphism of dg \mathbb{k} - \mathbb{k} -bimodules, it follows that $V^* \otimes V$ and \mathbb{k} are dg Morita equivalent

and hence $\mathbf{M}_{V^* \otimes V}$ is dg equivalent to the natural 2-representation $\mathbf{N} \cong \mathbf{M}_{\mathbb{k}}$ of $\mathcal{C}_{\mathbb{k}}$. This proves the claim. \square

Example 5.4. Consider the dg \mathbb{k} -algebra $D = \mathbb{k}[x]/(x^2)$ with $\partial(x) = 1$. Then D has no proper dg ideals. The *internal dg ideal* I_- of D is equal to zero, while the *external dg ideal* $I_+ = D$. This implies that D is acyclic [Or2, Section 2.1]. In particular D has no proper dg ideals, but is not given by a matrix algebra.

5.2. Pretriangulated hulls of finitary 2-categories and 2-representations. Let \mathcal{D} be a \mathbb{k} -finitary 2-category in the sense of [MM1]. We can view \mathcal{D} a dg 2-category with zero differential. Let $\mathcal{C} = \overline{\mathcal{D}}$ be the pretriangulated 2-category associated to it. Any finitary 2-representation \mathbf{M} of \mathcal{D} extends naturally to a pretriangulated 2-representation $\overline{\mathbf{M}}$ of \mathcal{C} . It is easy to see that given an object $X \in \mathbf{M}(\mathbf{i})$ and viewing it as an object in $\overline{\mathbf{M}}(\mathbf{i})$ by placing it in degree 0, the associated dg algebra 1-morphism A_X in $\overrightarrow{\mathcal{C}}(\mathbf{i}, \mathbf{i})$ is simply the algebra 1-morphism obtained from the finitary 2-representation \mathbf{M} , again interpreted as an object in $\overrightarrow{\mathcal{C}}(\mathbf{i}, \mathbf{i})$ by placing it in degree zero with zero differential.

For example, if $A = A_1 \times \dots \times A_n$, for A_i finite-dimensional connected \mathbb{k} -algebras, one defines the \mathbb{k} -finitary 2-category \mathcal{C}_A following [MM3, Section 4.5]. Then the pretriangulated hull of \mathcal{C}_A recovers the dg category \mathcal{C}_{A_0} defined in Section 5.1, for the dg algebra A_0 given by A with zero differential. The dg category $\mathcal{C}_{A_0}(\mathbf{i}, \mathbf{j})$ is dg equivalent to that of bounded complexes of A_j - A_i -bimodules that are projective (if $\mathbf{i} \neq \mathbf{j}$) or a direct sum of projective A_i - A_i -bimodules and direct sums of copies of A_i , if $\mathbf{i} = \mathbf{j}$.

Question 5.5. Are there quotient-simple dg 2-representations of \mathcal{C} which are not dg equivalent to ones of the form $\overline{\mathbf{M}}$ for a simple transitive 2-representation \mathbf{M} of \mathcal{D} ?

We remark that this question has a negative answer in case $\mathcal{D} = \mathcal{C}_{\mathbb{k}}$ by Proposition 5.3.

5.3. Pretriangulated hulls of finite multitensor categories. We can consider a (strict) finite multitensor category \mathcal{D} as in [EGNO], viewed as a 2-category with one object \bullet , and consider its pretriangulated hull $\mathcal{C} = \overline{\mathcal{D}}$ as in the preceding section. For any 2-representation \mathbf{N} of (in [EGNO] called a (strict) module category over) \mathcal{D} gives rise to a pretriangulated 2-representation $\mathbf{M} = \overline{\mathbf{N}}$ of \mathcal{C} .

Now assume that \mathcal{D} is a strict finite multitensor category. In particular, \mathcal{D} is abelian, locally finite, and every 1-morphism has left and right adjoints. These structures extend to the pretriangulated hull \mathcal{C} . Therefore, the evaluation dg functor $\text{Ev}_X: \mathcal{C}(\bullet, \bullet) \rightarrow \mathbf{M}(\bullet)$ from Section 4.2 has a right adjoint $[X, -]: \mathbf{M}(\bullet) \rightarrow \mathcal{C}(\bullet, \bullet)$ without extending to $\overrightarrow{\mathcal{C}}$ (see [EGNO, Sections 4.2, 7.3]). Our results in Section 4 generalize aspects of [EGNO, Section 7]. Similarly to Section 5.2, algebra 1-morphisms associated to $\mathbf{M} = \overline{\mathbf{N}}$ can be chosen inside \mathcal{D} (i.e., concentrated in degree zero).

5.4. Categorification of $\mathbb{Z}[i]$. In [Ti1], Tian constructs an explicit monoidal dg category \mathcal{I} which decategorifies to $\mathbb{Z}[i]$, and its action on a dg category $DGP(R)$, which categorifies the natural action of $\mathbb{Z}[i]$ on $\mathbb{Z}^{\oplus 2}$. Here, we explicitly compute a dg algebra 1-morphism associated to this dg 2-representation. We adopt the notation in loc. cit.

Let $X = P(x)$, the projective R -module at vertex x , which generates the dg 2-representation. Using the fact that $\mathcal{I} = \overline{\{Q^0, Q^1 = Q\}}$ (see [Ti1, Definition 2.2] together with the isomorphism $Q^2 \cong Q^0[1]$) it suffices to compute morphism spaces to

$[X, X]$ from Q^0 and Q^1 . We compute

$$\mathrm{Hom}_{\mathcal{I}}(Q^0, [X, X]) \cong \mathrm{Hom}_{\mathcal{I}}(Q^0 \otimes_R P(x), P(x)) \cong \mathrm{Hom}_{\mathcal{I}}(P(x), P(x)) \cong \mathbb{k}[d]/(d^2)$$

$$\mathrm{Hom}_{\mathcal{I}}(Q^1, [X, X]) \cong \mathrm{Hom}_{\mathcal{I}}(Q^1 \otimes_R P(x), P(x)) \cong \mathrm{Hom}_{\mathcal{I}}(P(y), P(x)) \cong \mathbb{k}[-1],$$

where d is the degree 1-endomorphism of $P(x)$. This implies that the dg algebra 1-morphism in $\vec{\mathcal{I}}$ is given by $Q^1[-2] \xrightarrow{\alpha_0 \circ \overline{hf}} Q^0$. The multiplication is given by

$$\begin{array}{ccc} Q^1[-2] \oplus Q^1[-2] & \xrightarrow{(\alpha_0 \circ \overline{hf}, \alpha_0 \circ \overline{hf})} & Q^0 \\ \downarrow (\mathrm{id}, \mathrm{id}) & & \downarrow \mathrm{id} \\ Q^1[-2] & \xrightarrow{\alpha_0 \circ \overline{hf}} & Q^0. \end{array}$$

As multiplication is induced by the identity on Q_0 , there is a dg algebra epimorphism given by

$$\begin{array}{ccc} Q^1[-2] & \xrightarrow{\alpha_0 \circ \overline{hf}} & Q^0 \\ \downarrow \alpha_1 & & \downarrow \mathrm{id} \\ Q^1[-1] & \xrightarrow{\overline{hf}} & Q^0 \end{array}$$

(note that $\alpha_0 \circ \overline{hf} = \overline{hf} \circ \alpha_1$). The given dg 2-representation is thus not quotient-simple. Indeed, it has a quotient given by factoring out the radicals of R and M , i.e. acting on $DGP(R')$ where $R' = \mathbb{k}e_x \times \mathbb{k}e_y$ and the action of Q_1 given by the R' - R' -bimodule $M' = \mathbb{k}e_y \otimes e_x \oplus \mathbb{k}e_x \otimes e_y[1]$.

5.5. Braid group categorification. The setup developed in this paper can be applied to existing categorified braid group actions from the literature. Firstly, in [KS], the authors consider a categorification of the braid group of type A_n using complexes of bimodules over the symmetric zigzag algebra $Z = Z_n$ on n -vertices. We denote the resulting dg 2-category by \mathcal{B}_n with generating 1-morphisms $T_i = Ze_i \otimes e_i Z \rightarrow Z$, $T'_i = Z \rightarrow Ze_i \otimes e_i Z$, as well as the identity bimodule.

Note that any pretriangulated 2-representation of \mathcal{B}_n is necessarily closed under cones, and hence extends to a dg 2-representation of \mathcal{C}_Z , see Section 5.1. In fact, \mathcal{C}_Z is the dg idempotent completion of the $\overline{\mathcal{B}_n}$.

Given that any dg ideal in a dg 2-representation of \mathcal{B}_n gives rise to a dg ideal in the extended dg 2-representation of \mathcal{C}_Z and vice versa, there is a one-to-one correspondence between quotient-simple pretriangulated 2-representations of \mathcal{B}_n and \mathcal{C}_Z . Thus, it follows from Lemma 5.2 that the dg algebra 1-morphism of the defining action corresponds to the dg algebra 1-morphism $Z \otimes_{\mathbb{k}} Z$ in $\overline{\mathcal{B}_n}$.

More generally, Rouquier [Ro1] considers a categorification of the braid group associated to an arbitrary finitely generated Coxeter system (W, S, V) in the homotopy category of Soergel bimodules. One could alternatively consider the dg 2-category \mathcal{B}_W generated by the complexes denoted in loc. cit. by F_s , which has a natural action on the category of bounded chain complexes of graded $\mathbb{k}[V]$ -modules. The homotopy category of this dg 2-category gives rise to Rouquier's categorification of the braid group, which he conjecturally strictifies in [Ro2].

As before, we can describe the dg idempotent completion of $\overline{\mathcal{B}_W}$ as the monoidal category of chain complexes of Soergel bimodules \mathcal{S}_W , and we again restrict or extend

pretriangulated 2-representations to pass from one to the other. Note that in both cases the dg ideal generated by the total invariants $\mathbb{k}[V]^W$ acts by zero on every quotient-simple dg 2-representation, so we can pass to the quotient dg 2-categories defined in the same way, but over the coinvariant algebra $C = \mathbb{k}[V]/(\mathbb{k}[V]_+^W)$ instead of $\mathbb{k}[V]$, which, provided W is finite, is a finite-dimensional symmetric algebra, see e.g. [MMMTZ], and references therein, for details. Then, for finite W , the natural (or defining) dg 2-representation of \mathcal{B}_W on chain complexes of C -modules simply corresponds to the pretriangulated hull of the cell 2-representation of \mathcal{S}_W associated to the longest element w_0 of the (finite) Coxeter group whose Kazhdan Lusztig basis element is categorified by $C \otimes C$, see [E, Sections 2.1,2.2] for more explanations, where this bimodule is described as $C \otimes_{C^S} C$ with C^S denoting the total invariants in the coinvariants and hence is isomorphic to \mathbb{k} (more generally for $J \subseteq S$, $C \otimes_{C^J} C$ decategorifies to the Kazhdan–Lusztig basis element corresponding to the longest element of the parabolic subgroup determined by J — this uses the Decategorification Theorem [EW]). The corresponding dg algebra 1-morphism is the simply given by $C \otimes C$, which is indeed a 1-morphism in $\overrightarrow{\mathcal{B}}_W = \overrightarrow{\mathcal{S}}_W$.

REFERENCES

- [AL] R. Anno, T. Logvinenko. Spherical DG-functors. *J. Eur. Math. Soc. (JEMS)* **19** (2017), no. 9, 2577–2656.
- [BS] P. Balmer, M. Schlichting. Idempotent completion of triangulated categories. *J. Algebra* **236** (2001), no. 2, 819–834.
- [BFK] J. Bernstein, I. Frenkel, M. Khovanov. A categorification of the Temperley-Lieb algebra and Schur quotients of $U(\mathfrak{sl}_2)$ via projective and Zuckerman functors. *Selecta Math. (N.S.)* **5** (1999), no. 2, 199–241.
- [BN] M. Bökstedt, A. Neeman. Homotopy limits in triangulated categories. *Compositio Math.* **86** (1993), no. 2, 209–234.
- [BLL] A.I. Bondal, M. Larsen, V. A. Lunts. Grothendieck ring of pretriangulated categories. *Int. Math. Res. Not.* 2004, no. 29, 1461–1495.
- [BK] A.I. Bondal, M.M. Kapranov. Enhanced triangulated categories. *Mat. Sb.* **181** (1990), no. 5, 669–683; transl. in *Math. USSR-Sb.* **70** (1991), no. 1, 93–107.
- [Ca] S. Cautis. Clasp technology to knot homology via the affine Grassmannian. *Math. Ann.* **363** (2015), no. 3-4, 1053–1115.
- [CL] S. Cautis, A. Licata. Heisenberg categorification and Hilbert schemes. *Duke Math. J.* **161** (2012), no. 13, 2469–2547.
- [LC] J. Le, X.-W. Chen. Karoubianness of a triangulated category. *J. Algebra* **310** (2007), no. 1, 452–457.
- [CR] J. Chuang, R. Rouquier. Derived equivalences for symmetric groups and \mathfrak{sl}_2 -categorification. *Ann. of Math. (2)* **167** (2008), no. 1, 245–298.
- [EL] I. Egilmez, A. D. Lauda. DG structures on odd categorified quantum $sl(2)$. *Quantum Topol.* **11** (2020), no. 2, 227–294.
- [E] B. Elias. Thicker Soergel calculus in type A. *Proc. Lond. Math. Soc. (3)* **112** (2016), no. 5, 924–978.
- [EQ1] B. Elias, Y. Qi. An approach to categorification of some small quantum groups II. *Advances in Mathematics* **288** (2016), 81–151.
- [EQ2] B. Elias, Y. Qi. A categorification of quantum $sl(2)$ at prime roots of unity. *Advances in Mathematics* **299** (2016), 863–930.
- [EW] B. Elias, G. Williamson. The Hodge theory of Soergel bimodules. *Ann. of Math. (2)* **180** (2014), no. 3, 1089–1136.
- [EIQ] A. P. Ellis, Y. Qi. The differential graded odd nilHecke algebra. *Comm. Math. Phys.* **344** (2016), no. 1, 275–331.
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik. Tensor categories. *Mathematical Surveys and Monographs*, 205. American Mathematical Society, Providence, RI, 2015.

- [Fr] P. Freyd. Representations in abelian categories. 1966 Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965) pp. 95–120 Springer, New York, 1966
- [JY] P. Jorgensen, M. Yakimov. Green groupoids of 2-Calabi–Yau categories, derived Picard actions, and hyperplane arrangements. Preprint arXiv:1910.05560 (2019).
- [Ke1] B. Keller. Deriving DG categories. Ann. Sci. École Norm. Sup. (4) **27** (1994), no. 1, 63–102.
- [Ke2] B. Keller. On differential graded categories. International Congress of Mathematicians. Vol. II, 151–190, Eur. Math. Soc., Zürich, 2006.
- [KS] M. Khovanov, P. Seidel. Quivers, Floer cohomology, and braid group actions. J. Amer. Math. Soc. **15** (2002), no. 1, 203–271.
- [KT] M. Khovanov, Y. Tian. How to categorify the ring of integers localized at two. Quantum Topol. **10** (2019), no. 4, 723–775.
- [LM] R. Laugwitz, V. Miemietz. Cell 2-representations and categorification at prime roots of unity. Adv. Math. **361** (2020), 106937, 66 pp.
- [LO] V.A. Lunts, D.O. Orlov. Uniqueness of enhancement for triangulated categories. J. Amer. Math. Soc. **23** (2010), no. 3, 853–908.
- [McL] S. Mac Lane. Categories for the working mathematician. Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998.
- [MMMT] M. MacKaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer. Simple transitive 2-representations via (co-)algebra 1-morphisms. Indiana Univ. Math. J. **68** (2019), no. 1, 1–33.
- [MMMTZ] M. MacKaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, X. Zhang. Simple transitive 2-representations of Soergel bimodules for finite Coxeter types. Preprint, arXiv:1906.11468.
- [MM1] V. Mazorchuk, V. Miemietz. Cell 2-representations of finitary 2-categories. Compositio Math. **147** (2011), 1519–1545.
- [MM2] V. Mazorchuk, V. Miemietz. Additive versus abelian 2-representations of fiat 2-categories. Moscow Math. J. **14** (2014), no. 3, 595–615.
- [MM3] V. Mazorchuk, V. Miemietz. Endomorphisms of cell 2-representations. Int. Math. Res. Not. (2016), no. 24, 7471–7498.
- [MM5] V. Mazorchuk, V. Miemietz. Transitive representations of finitary 2-categories. Trans. Amer. Math. Soc. **368** (2016), no. 11, 7623–7644.
- [MMMZ] M. Mackaay, V. Mazorchuk, V. Miemietz, X. Zhang. Analogues of centralizer subalgebras for fiat 2-categories and their 2-representations. J. Inst. Math. Jussieu **19** (2020), no. 6, 1793–1829.
- [Or1] D. Orlov. Smooth and proper noncommutative schemes and gluing of dg categories. Adv. Math. **302** (2016), 59–105.
- [Or2] D. Orlov. Finite-dimensional differential graded algebras and their geometric realizations. Adv. Math. **366** (2020), 107096, 33 pp.
- [Ro1] R. Rouquier. Categorification of the braid groups, Preprint, arXiv:math/0409593 (2004).
- [Ro2] R. Rouquier. Categorification of \mathfrak{sl}_2 and braid groups. Trends in representation theory of algebras and related topics, 137–167, Contemp. Math. 406, Amer. Math. Soc., Providence, RI, 2006.
- [Ro3] R. Rouquier. 2-Kac-Moody algebras. Preprint, arXiv:0812.5023 (2008).
- [RZ] R. Rouquier, A. Zimmermann. Picard groups for derived module categories. Proc. London Math. Soc. (3) **87** (2003), no. 1, 197–225.
- [Sa] A. Sartori. Categorification of tensor powers of the vector representation of $U_q(\mathfrak{gl}(1|1))$. Selecta Math. (N.S.) **22** (2016), no. 2, 669–734.
- [Sch] O.M. Schnürer. Perfect derived categories of positively graded DG algebras, Appl. Categ. Structures **19** (2011), no. 5, 757–782.
- [ST] P. Seidel, R. Thomas. Braid group actions on derived categories of coherent sheaves. Duke Math. J. **108** (2001), no. 1, 37–108.
- [Ti1] Y. Tian. A categorification of the square root of -1 . Fund. Math. **232** (2016), no. 1, 89–98.
- [Ti2] Y. Tian. A diagrammatic categorification of a Clifford algebra. Int. Math. Res. Not. (2015), no. 21, 10872–10928
- [Ti3] Y. Tian. A categorification of $U_T(\mathfrak{sl}(1|1))$ and its tensor product representations. Geom. Topol. **18** (2014), no. 3, 1635–1717.
- [W] G. Williamson. Schubert calculus and torsion explosion. With a joint appendix with Alex Kontorovich and Peter J. McNamara. J. Amer. Math. Soc. **30** (2017), no. 4, 1023–1046.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM, NG7 2RD, UK

Email address: `robert.laugwitz@nottingham.ac.uk`

SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH NR4 7TJ, UK

Email address: `v.miemietz@uea.ac.uk`

URL: <https://www.uea.ac.uk/~byr09xgu/>