

# PISTA: PRECONDITIONED ITERATIVE SOFT THRESHOLDING ALGORITHM FOR GRAPHICAL LASSO

GAL SHALOM\*, ERAN TREISTER<sup>†</sup>, AND IRAD YAVNEH\*

**Abstract.** We propose a novel quasi-Newton method for solving the sparse inverse covariance estimation problem also known as the graphical least absolute shrinkage and selection operator (GLASSO). This problem is often solved using a second order quadratic approximation. However, in such algorithms the Hessian term is complex and computationally expensive to handle. Therefore, our method uses the inverse of the Hessian as a preconditioner to simplify and approximate the quadratic element at the cost of a more complex  $\ell_1$  element. The variables of the resulting preconditioned problem are coupled only by the  $\ell_1$  sub-derivative of each other, which can be guessed with minimal cost using the gradient itself, allowing the algorithm to be parallelized and implemented efficiently on GPU hardware accelerators. Numerical results on synthetic and real data demonstrate that our method is competitive with other state-of-the-art approaches.

**Key words.** Graphical LASSO, Sparse Precision Matrix Estimation, Proximal Methods, Preconditioning.

**AMS subject classifications.** 90C25, 65D18, 65K10, 65F08

**1. Introduction.** Inverse covariance estimation is a fundamental problem in modern statistics. Specifically, the inverse covariance matrices of multivariate normal distributions are used in numerous applications. One of the most common uses of the inverse covariance, also known as the *Precision Matrix*, is to describe statistical models. That is, the graph structure of Gaussian graphical models can be inferred from the Precision Matrix [19]. This inferred graph can be used for analyzing gene networks [8], financial assets and stocks dependencies [10, 30], social networks [13] and other inter-dependent data [5, 18].

The most straightforward approach to estimating the Precision Matrix would be to use the inverse of the empirical covariance matrix. Evidently, inverting the empirical covariance matrix requires a number of samples that equals or exceeds the dimension of the matrix. However, many problems of interest are high dimensional [8, 17], to the extent that they are significantly larger than the number of available samples, so the empirical covariance matrix is rank-deficient. In such cases, some kind of regularization is essential. One of the most common ways to estimate the Precision Matrix is to solve a  $\ell_1$ -regularized maximum likelihood estimation problem, known as the graphical least absolute shrinkage and selection operator (Graphical LASSO, or GLASSO) problem. The use of  $\ell_1$ -regularization aims to achieve a sparse estimation while keeping the problem convex. Sparsity of the Precision Matrix may be interpreted as simplicity—a sparse Precision Matrix implies a simple inferred graph structure [15]. Thus, when a genuine empirical Precision Matrix cannot be computed, a sparse valid estimation is often the preferred choice [7, 15, 31]. We refer the reader to the first chapter of [15] and to [7] for additional motivation.

As the problem is well studied, many algorithms and methods for its solution have been presented over the years [1, 2, 6, 9, 12, 14, 16, 17, 21, 22, 24, 26, 27]. Some of these

---

\*Faculty of Computer Science, Technion — Israel Institute of Technology, Israel (gal-shalom@cs.technion.ac.il, irad@cs.technion.ac.il). Supported in part by the Israel Science Foundation, grant No. 1639/19.

<sup>†</sup>Department of Computer Science, Ben-Gurion University of the Negev, Israel (erant@cs.bgu.ac.il). This work was supported in part by the Israeli Council for Higher Education (CHE) via the Data Science Research Center, Ben-Gurion University of the Negev, Israel.

methods are computationally intensive or difficult to parallelize, rendering parallel computation accelerators such as GPUs ineffective. As far as we know, none of the algorithms in the literature were designed specifically with an efficient GPU deployment in mind. Nevertheless, various existing algorithms are able to take advantage of GPU processing power with a simple implementation, e.g., G-ISTA [14], VSM [21], PSM [9], ALM [24], Newton-Lasso [22] and Orthant-Based Newton [22]. Each of these has its strengths and weaknesses. Other algorithms, such as BCD-IC [26], can be deployed on a GPU as well, however, the method and the authors' implementation are complex, and the extensive usage of scalar operations may hinder its performance.

In this work we introduce a preconditioned Iterative Soft Thresholding Algorithm (pISTA) for solving the graphical LASSO problem. pISTA is a quasi-Newton algorithm designed to be highly parallel and suitable for using GPU capabilities to its benefit by exploiting the problem structure effectively. A traditional second order quadratic approximation includes a complex Hessian term which is computationally expensive to handle. Moreover, all the elements of the Hessian term are coupled, making efficient GPU deployment challenging. To this end, pISTA uses the inverse of the Hessian as a preconditioner to simplify and approximate the quadratic element, at the cost of a more complex  $\ell_1$  element. This is highly beneficial since the Hessian inverse is easily obtained for the GLASSO problem. The variables of the resulting preconditioned problem are coupled only by the  $\ell_1$  sub-derivative of each other, which can be guessed with minimal cost using the gradient itself. The resulting pISTA algorithm uses only matrix operations, making it easier for exploiting the GPU efficiently.

This paper is organized as follows. In section 2 we refer to related work, and in section 3 we formulate the problem. The pISTA algorithm is introduced in section 4, and its convergence is proved in section 5. Experimental results are presented in section 6, and conclusions follow in section 7.

**2. Related Work.** Many numerical algorithms have been developed, which are specifically designed for solving the sparse inverse covariance estimation problem. G-ISTA [14] uses a proximal gradient method which estimates the inverse covariance iteratively. GLASSO [12] splits the problem into smaller LASSO problems and updates the estimation by solving them separately. QUIC [16] uses a proximal Newton method on the objective function. BCD-IC [26] uses a block coordinate descent method. ALM [24] uses an alternating linearization technique by splitting the objective function into two linear approximations and minimizing them alternately. PSM [9] employs a projected gradient method. VSM [21] uses Nesterov's smooth optimization technique. Newton-Lasso [22] and Orthant-Based Newton [22] use FISTA [3] and Conjugate Gradient methods, respectively, to solve reduced second order approximations. Although some algorithms are designed to be run in parallel using the multi-core model of the CPU, none of them take into specific consideration the parallelism model of the GPU and the usage of GPUs as a computation accelerator.

Additionally, some algorithm were developed specifically for large scale matrices where the matrix can only reside in memory in sparse format. Big&Quic and SQUIC [4, 17] extend QUIC [16] to large scales. Big&Quic uses on-demand computation of the Hessian's columns and a special procedure of the linesearch conditions using Schur complements. SQUIC utilizes sparse Cholesky factors of the iterations matrix. ML-BCD [27] takes BCD-IC [26], which is already suitable to large scale problems, and accelerates it using a generic multilevel framework, which was originally suggested in [28] for LASSO. This acceleration can in principle be used with our proposed method as well. Large scale algorithms may benefit from the GPU processing power,

however memory constraints and usage of sparse format make this task non-trivial.

Graphical LASSO can also be used in a mixture model setup when multiple sparse inverse covariances are utilized [11]. This allows a richer statistical model, but requires the estimation of multiple sparse Precision Matrices. An efficient GPU-based solver is highly beneficial in such scenarios as well.

### 3. Background.

**3.1. Sparse Inverse Covariance Estimation.** Estimating the parameters of multivariate Gaussian distributions is a fundamental problem in statistics. Given  $m$  independent samples  $\{y_i\}_{i=1}^m \in \mathbb{R}^n$ , where  $y_i \sim \mathcal{N}(\mu, \Sigma)$ , one would like to estimate the mean  $\mu \in \mathbb{R}^n$  and either the covariance  $\Sigma$ , or its inverse  $\Sigma^{-1}$ , which is also called the *Precision Matrix*. Both the mean  $\mu$  and the covariance  $\Sigma$  are often estimated by the maximum likelihood estimator (MLE). The MLE is given by the parameters  $\mu$  and  $\Sigma$  that maximize the probability to sample the observed data  $\{y_i\}_{i=1}^m$ :

$$\begin{aligned} \hat{\mu}, \hat{\Sigma} &= \arg \max_{\mu, \Sigma} \prod_{i=1}^m \mathbb{P}(y_i | \Sigma, \mu) \\ (3.1) \quad &= \arg \max_{\mu, \Sigma} \prod_{i=1}^m \frac{1}{\sqrt{(2\pi)^m \det(\Sigma)}} \exp\left(-\frac{1}{2}(y_i - \mu)^T \Sigma^{-1} (y_i - \mu)\right). \end{aligned}$$

MLE has an analytical solution:

$$(3.2) \quad \hat{\mu} = \frac{1}{m} \sum_{i=1}^m y_i, \quad \hat{\Sigma} = S \triangleq \frac{1}{m} \sum_{i=1}^m (y_i - \hat{\mu})(y_i - \hat{\mu})^T,$$

where  $\hat{\mu}$  and  $\hat{\Sigma}$  are called the empirical mean and empirical covariance, respectively.

Usually, one considers the log-likelihood objective (negative *log* of (3.1)), minimized over the inverse covariance matrix, yielding the inverse covariance MLE:

$$(3.3) \quad \hat{\Sigma}^{-1} = \arg \min_{A \succ 0} f(A) \triangleq -\log(\det(A)) + \text{Tr}(SA).$$

The solution is indeed the inverse of (3.2). However, in cases where the number of available samples is smaller than the dimension of  $y_i$  ( $m < n$ ), the matrix  $S$  is rank-deficient and thus non-invertible, whereas the true  $\Sigma$  is assumed to have full rank and is positive definite. In other words, we cannot estimate  $\Sigma^{-1}$  by inverting  $S$ , and further assumptions should be considered. A common choice in this case is to assume that  $\Sigma^{-1}$  is sparse.  $\Sigma^{-1}$  can be interpreted as a conditional dependence matrix, that is, its off-diagonal entries indicate the dependence between the row and column variables, given all the remaining variables [23].

A common approach is to regularize the log-likelihood objective (3.3) with a sparsity promoting  $\ell_1$ -penalty [1, 2, 6]:

$$(3.4) \quad \hat{\Sigma}^{-1} = \arg \min_{A \succ 0} F(A) \triangleq \arg \min_{A \succ 0} f(A) + \alpha \|A\|_1,$$

where  $\alpha > 0$  is a scalar and  $\|A\|_1 = \sum_{i,j} |A_{i,j}|$ . Equation (3.4) is known as the *Graphical Lasso* problem.

As seen from (3.4), the objective function is convex and is composed of two parts—a smooth convex part  $f(A)$  and a non-smooth convex part  $\alpha \|A\|_1$ . Although the objective is convex, the non-smooth term makes traditional algorithms ineffective, and more specialized solvers are needed.

**3.2. Proximal Methods for Sparse Inverse Covariance Estimation.** A common approach to solving convex problems comprised of smooth and non-smooth parts is known as *proximal methods*, where we approximate the objective function at each iteration as follows: the smooth part is approximated by a quadratic function while the non-smooth part is kept unchanged. This approach is especially attractive when the non-smooth part is a separable function (e.g., point-wise).

Specifically, applying proximal methods to the objective function  $F(A)$  of (3.4), the smooth term  $f(A)$  is approximated by a quadratic function while the term  $\alpha\|A\|_1$  remains unchanged. This approximation yields a linear LASSO [25] problem ( $\ell_1$ -regularized quadratic objective). Once defined, we minimize it approximately using a LASSO solver. Denoting the descent direction at the  $k$ -th iteration by  $D^{(k)}$ , then at each iteration we solve:

$$(3.5) \quad \begin{aligned} D^{(k)} &= \arg \min_D \tilde{F}(A^{(k)} + D) \\ &= \arg \min_D f(A^{(k)}) + \langle g^{(k)}, D \rangle + \frac{1}{2} \langle D, H^{(k)} D \rangle + \alpha \|A^{(k)} + D\|_1, \end{aligned}$$

where  $g^{(k)} = \nabla f(A^{(k)})$  is the gradient at the  $k$ th iteration, and  $H^{(k)}$  depends on the particular method. For  $H^{(k)} = I$  we get the ‘‘Proximal Gradient’’ method, which is called G-ISTA [14]. At the other extreme, for  $H^{(k)} = \nabla^2 f(A^{(k)})$ , the Hessian at the  $k$ th iteration, we get the ‘‘Proximal Newton’’ method known as QUIC [16].

The gradient and the Hessian are given by:

$$(3.6) \quad \nabla f(A) = S - A^{-1} \quad , \quad \nabla^2 f(A) = A^{-1} \otimes A^{-1} ,$$

where  $\otimes$  is the Kronecker product. We note that once the gradient has been computed, we can compute Hessian-vector products at relatively low cost by using Kronecker product properties. However, solving (3.5) with  $H^{(k)} = \nabla^2 f(A^{(k)})$  is highly complex and it is typically done by coordinate descent iterations [16]. This makes it hard to use computation accelerators (such as GPUs) efficiently.

#### 4. The preconditioned Iterative Soft Thresholding Algorithm (pISTA).

In this section we develop the *preconditioned Iterative Soft Thresholding Algorithm* (pISTA). In [16], the Proximal Newton method for our problem, each coordinate computation depends on the values of all the other coordinates due to the Hessian in the quadratic element. In pISTA, we aim for a simpler and easier to solve quadratic element at the cost of a more complex non-smooth  $\ell_1$  part.

First, in subsection 4.1, we restrict our problem to a smaller set of variables as done in other algorithms [16, 22, 27]. In subsection 4.2, we simplify and relax the quadratic part using a preconditioner, at the cost of making the non-smooth part  $\alpha\|A\|_1$  more complex. Lastly, in subsection 4.3, we solve the resulting problem, completing the development of pISTA. In contrast to second order methods like QUIC, for example, in pISTA each coordinate depends on the other variables only through the sub-gradient of the non-smooth part ( $\ell_1$  norm), which is more complex. However, the sub-gradient of the  $\ell_1$  term can be approximated easily, and relatively well, by the sign of the elements of the current iterate (if it is nonzero) or of its gradient (where the elements of the iterate vanish), allowing our quasi-Newton method to be efficient.

**4.1. Restricting the Updates to the Free-set.** As introduced in [16] and used in [22, 27], we restrict the descent direction  $D$  to the *free-set* of  $A^{(k)}$ . That is, an element in  $D$  which is not in the *free-set* of  $A^{(k)}$  is set to zero. Denote by  $\mathcal{S}_A^{(k)}$  the

*free-set* at iteration  $k$ , defined as follows:

$$(4.1) \quad \mathcal{S}_A^{(k)} = \left\{ (i, j) \mid A_{i,j}^{(k)} \neq 0 \right\} \cup \left\{ (i, j) \mid |\nabla f(A^{(k)})|_{i,j} > \alpha \right\}.$$

According to Lemma 7 of [16], solving (3.5) restricted to the variables that are not in the *free-set* of  $A^{(k)}$  will result in a zero value in every element in the direction  $D$ . Thus, restricting  $D$  to the *free-set* is equivalent to solving (3.5) in an alternating two-block manner: the first step restricted to the variables not in the *free-set* (which makes no change in our approximation), and the second step restricted to the *free-set*. The free set is recomputed before the next iteration.

Define the restriction mask at the  $k$ -th iteration:

$$(4.2) \quad [\mathcal{M}_A^{(k)}]_{i,j} = \begin{cases} 1 & (i, j) \in \mathcal{S}_A^{(k)} \\ 0 & (i, j) \notin \mathcal{S}_A^{(k)} \end{cases}.$$

Then, the restricted problem we solve at the  $k$ th iteration can be written as:

$$(4.3) \quad D^{(k)} = \arg \min_D \tilde{F}(A^{(k)} + (\mathcal{M}_A^{(k)} \odot D)),$$

where  $\odot$  is the Hadamard product. Note that, in the solution of (4.3), the elements of  $D^{(k)}$  that are not in the *free-set* of  $A^{(k)}$  may have nonzero values. However, we restrict the updates to the indices in the *free-set* of  $A^{(k)}$  only, therefore, our descent direction is  $\mathcal{M}_A^{(k)} \odot D^{(k)}$ , which is equivalent to setting the values of  $D^{(k)}$  that are not in the free-set to zero.

**4.2. Preconditioning of the Descent Direction.** Note the following property of the Hessian in (3.6):

$$(4.4) \quad (\nabla^2 f(A))^{-1} = A \otimes A,$$

where  $\otimes$  is the Kronecker product. This means that the Hessian inverse can be obtained without any computational overhead, using the typically sparse  $A$  instead of the typically dense  $A^{-1}$ . Here we are interested in using this appealing property to accelerate the solution of (3.5), where  $H^{(k)} = \nabla^2 f(A^{(k)})$ . To this end, we use it as a preconditioner to  $D$ :

$$(4.5) \quad \text{vec}(D) = (\nabla^2 f(A^{(k)}))^{-1} \text{vec}(\Delta) \Rightarrow D = A^{(k)} \Delta A^{(k)},$$

where  $\text{vec}$  denotes the column-stacking of a matrix into a vector.

Denote  $W^{(k)} = (A^{(k)})^{-1}$ . Using the preconditioned descent direction (4.5) in the restricted problem (4.3), with  $H^{(k)} = \nabla^2 f(A^{(k)}) = W^{(k)} \otimes W^{(k)}$ , results in the following equation:

$$(4.6) \quad \begin{aligned} \Delta^{(k)} &= \arg \min_{\Delta} \tilde{F} \left( A^{(k)} + \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right) \\ &= \arg \min_{\Delta} f(A^{(k)}) + \left\langle g^{(k)}, \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right\rangle \\ &\quad + \frac{1}{2} \left\langle \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}), W^{(k)} \left( \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right) W^{(k)} \right\rangle \\ &\quad + \alpha \left\| A^{(k)} + \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right\|_1. \end{aligned}$$

After applying the mask restriction  $\mathcal{M}$  and the preconditioning to (3.5), the quadratic part (the second element in (4.6)) is still complex. We relax it by removing the mask restriction  $\mathcal{M}$  and replacing it with a multiplicative scalar:

$$(4.7) \quad \frac{1}{2} \left\langle \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}), W^{(k)} \left( \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right) W^{(k)} \right\rangle \\ \approx \frac{1}{2t} \left\langle A^{(k)} \Delta A^{(k)}, W^{(k)} (A^{(k)} \Delta A^{(k)}) W^{(k)} \right\rangle = \frac{1}{2t} \left\langle A^{(k)} \Delta A^{(k)}, \Delta \right\rangle,$$

where  $t$  is a scalar that is computed using linesearch. Since we do not change the first order terms (the gradient and  $\ell_1$ ), this can still result in a monotonically convergent method. Denote the resulting quasi-Newton approximation by  $P(\Delta; t)$ , then the  $k$ -th iterate is given by:

$$(4.8) \quad \Delta^{(k)} = \arg \min_{\Delta} P(\Delta; t) \\ = \arg \min_{\Delta} f(A^{(k)}) + \left\langle g^{(k)}, \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right\rangle + \frac{1}{2t} \left\langle A^{(k)} \Delta A^{(k)}, \Delta \right\rangle \\ + \alpha \left\| A^{(k)} + \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right\|_1.$$

**4.3. pISTA.** Our pISTA algorithm solves the problem defined by (4.8) in each iteration, with an appropriately chosen  $t$ . First, we find a solution  $\Delta^{(k)}$  to (4.8). To obtain this, we need to formulate the sub-differential of the non-smooth  $\ell_1$  term  $\|A + D\|_1$ . First, denote the sub-differential of  $\|A + D\|_1$  by:

$$(4.9) \quad G^{(k)}(D) \triangleq \frac{\partial \|A^{(k)} + D\|_1}{\partial D} \\ = \left\{ \mathcal{G}^{(k)} \left| \begin{array}{ll} \mathcal{G}_{i,j}^{(k)} = [\text{sign}(A^{(k)} + D)]_{i,j} & [A^{(k)} + D]_{i,j} \neq 0 \\ \mathcal{G}_{i,j}^{(k)} \in [-1, 1] & [A^{(k)} + D]_{i,j} = 0 \end{array} \right. \right\}.$$

Next, using the chain rule and some known derivatives with subsection 4.3, we formulate the sub-differential of the quasi-Newton approximation:

$$(4.10) \quad \frac{\partial P(\Delta; t)}{\partial \Delta} = \left\{ \frac{1}{t} A^{(k)} \Delta A^{(k)} + A^{(k)} \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} + \alpha A^{(k)} \left( \mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} \right. \\ \left. \left| \mathcal{G}^{(k)} \in G^{(k)} \left( \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta A^{(k)}) \right) \right. \right\},$$

where  $\mathcal{G}^{(k)}$  represents a sub-gradient of the  $\ell_1$  term.

The desired  $\Delta^{(k)}$  that will be used to compute the descent direction is one which includes the matrix 0 in its sub-gradients:

$$(4.11) \quad \Delta^{(k)} : \quad 0 \in \frac{\partial P(\Delta^{(k)}; t)}{\partial \Delta}.$$

Notice that after finding  $\Delta^{(k)}$ , it will be used to compute the descent direction  $\mathcal{M}_A^{(k)} \odot D^{(k)} = \mathcal{M}_A^{(k)} \odot (A^{(k)} \Delta^{(k)} A^{(k)})$ . Thus, we can equivalently find  $D^{(k)} = A^{(k)} \Delta^{(k)} A^{(k)}$  instead of  $\Delta^{(k)}$ . In other words, we shall find  $D^{(k)}$  which has 0 as one

of its sub-gradients:

$$(4.12) \quad D^{(k)} : 0 \in \left\{ \frac{1}{t} D^{(k)} + A^{(k)} \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} + \alpha A^{(k)} \left( \mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} \right. \\ \left. \left| \mathcal{G}^{(k)} \in G^{(k)} \left( \mathcal{M}_A^{(k)} \odot D^{(k)} \right) \right\}.$$

Note that the change of variables has made the first (Hessian) term trivial. To find  $D^{(k)}$ , we split the sub-gradient to  $n^2$  equations where the  $(i, j)$ -th equation is composed by the  $(i, j)$  element in the sub-gradient. For each equation  $(i, j)$ , we define the equation variable as  $D_{i,j}^{(k)}$ . The approximate solution (developed in Appendix A) is:

$$(4.13) \quad D_{i,j}^{(k)} = -A_{i,j}^{(k)} + \text{SoftThreshold} \left( A^{(k)} - t \cdot B_{i,j}^{(k)}, t \cdot C_{i,j}^{(k)} \right),$$

where

$$(4.14) \quad C_{i,j}^{(k)} = \begin{cases} \alpha \cdot \left( A_{i,i}^{(k)} \cdot A_{j,j}^{(k)} \right), & i = j \\ \alpha \cdot \left( A_{i,i}^{(k)} \cdot A_{j,j}^{(k)} + A_{i,j}^{(k)} \cdot A_{j,i}^{(k)} \right), & i \neq j \end{cases},$$

and

$$(4.15) \quad B^{(k)} = A^{(k)} \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} + \alpha A^{(k)} \left( \mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} - C^{(k)} \odot \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right).$$

So far, we developed a second order approximation for the GLASSO using the Hessian inverse and a quadratic relaxation. We found a closed form direction  $D^{(k)}$  for each  $(i, j)$  entry, assuming that the rest of the entries in  $\mathcal{G}^{(k)}$  are given. In the following subsection we show how we approximate those entries.

Algorithm 4.1 describes the pISTA method using (4.13). One part that is still missing from the description so far is the computation of  $\mathcal{G}^{(k)}$  and the linesearch procedure for finding an appropriate  $t$ , which we describe next. It is important to note that all the steps and operations in Algorithm 4.1 can be computed using matrix operations. This property allows us to easily benefit from GPU computation acceleration and easy GPU deployment.

---

**Algorithm 4.1** pISTA( $S, \alpha, A^{(0)}$ )

---

- 1: **Result:**  $A$
  - 2: **Initialization:**  $k = 0$ ;
  - 3: **while** stop criteria not met **do**
  - 4:   Compute  $\mathcal{S}_A^{(k)}$  according to (4.1);
  - 5:   Compute  $\mathcal{M}_A^{(k)}$  according to (4.2);
  - 6:   Compute  $\mathcal{G}^{(k)}$  according to (4.16);
  - 7:   Find an appropriate  $t$  and compute  $D^{(K)}$  according to (4.13);
  - 8:    $A^{(k+1)} = A^{(K)} + \mathcal{M}_A^{(k)} \odot D^{(K)}$ ;
  - 9: **end while**
-

**4.3.1. The Approximation of  $\mathcal{G}$ .** The algorithm depends on a good approximation of  $\mathcal{G}^{(k)}$ , especially at the beginning of the solution process. Recall that  $\mathcal{G}_{i,j}^{(k)}$  is the sub-derivative of  $|A_{i,j}^{(k)} + D_{i,j}^{(k)}|$ , and although the sub-derivative cannot be computed, it can be approximated by  $\text{sign}(A_{i,j}^{(k)} + D_{i,j}^{(k)})$ . Eventually, at the late iterations  $\mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)}$  will converge to  $\text{sign}(A^{(k)})$ , as all zero elements in  $A^{(k)}$  will be out of the free-set  $\mathcal{S}_A^{(k)}$ . Therefore, their elements in  $\mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)}$  will be zero, and all non-zero elements in  $A^{(k)}$  will have a constant  $\mathcal{G}^{(k)}$ , which is their sign. Moreover, at those later iterations, where  $\text{sign}(A^{(k)})$  is not expected to change at all, the above formula (4.13) solves (4.12) completely as all the terms except  $D^{(k)}$  are constant. On the other hand, at the initial iterations  $\mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)}$  cannot be easily computed because some elements in the free set may change sign or be zero in the final solution. To this end, we approximate  $\mathcal{G}^{(k)}$  by predicting the “next” sign of the elements. That is, we take the sign of the non-zero entries, and if an entry is zero, then we take the opposite sign of its gradient. Thus, a good approximation of  $\mathcal{G}^{(k)}$  is:

$$(4.16) \quad \mathcal{G}_{i,j}^{(k)} = \begin{cases} \text{sign}(A_{i,j}^{(k)}) & A_{i,j}^{(k)} \neq 0 \\ -\text{sign}(g_{i,j}^{(k)}) & A_{i,j}^{(k)} = 0 \end{cases},$$

where  $g^{(k)} = \nabla f(A^{(k)})$  as defined before. This means that we choose the next signs according to a small step of proximal gradient descent. The above  $\mathcal{G}^{(k)}$  together with the free-set restriction mask  $\mathcal{M}^{(k)}$  define the minimum sub-gradient of  $F(A^{(k)})$  with respect to  $\ell_2$ -norm.

**4.3.2. Performing Linesearch to find  $t$ .** At each iteration we perform linesearch to obtain a parameter  $t$  satisfying:

$$(4.17) \quad A^{(k)} + \mathcal{M}_A^{(k)} \odot D^{(k)} \succ 0,$$

$$(4.18) \quad F(A^{(k)} + \mathcal{M}_A^{(k)} \odot D^{(k)}) < F(A^{(k)}).$$

**5. Convergence Analysis.** Throughout this section, unless stated otherwise, all the matrices are assumed to be symmetric and real. The definitions of  $f(A)$ ,  $F(A)$ ,  $P(\Delta; t)$ ,  $g$ ,  $\mathcal{M}$ ,  $G$  and  $D$  are as above. We will denote  $D_t^{*(k)}$  as the direction  $D^{(k)}$  which satisfies equation (4.12) for a given  $t$ :

$$(5.1) \quad 0 \in \left\{ \frac{1}{t} D_t^{*(k)} + A^{(k)} \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} + \alpha A^{(k)} \left( \mathcal{G}^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} \right. \\ \left. \left| \mathcal{G}^{(k)} \in G^{(k)} \left( \mathcal{M}_A^{(k)} \odot D_t^{*(k)} \right) \right\}.$$

Also, denote by  $\mathcal{G}_t^{*(k)}$  the  $\mathcal{G}^{(k)}$  for which  $D_t^{*(k)}$  attains 0:

$$(5.2) \quad 0 = \frac{1}{t} D_t^{*(k)} + A^{(k)} \left( g^{(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)} + \alpha A^{(k)} \left( \mathcal{G}_t^{*(k)} \odot \mathcal{M}_A^{(k)} \right) A^{(k)}.$$

In the proofs only, we will omit the iteration symbol when the meaning is evident.

This section is organized as follows. In subsection 5.1 we prove that if our pISTA iteration does not update  $A^{(k)}$ , then  $A^{(k)} = \arg \min_{A \succ 0} F(A)$ . In subsection 5.2 we prove that for every iteration there exists a  $t \geq t_{pd} > 0$  which satisfies (4.17) and



keeps the matrix positive definite. Lastly, in subsection 5.3 we prove that for every iteration there exists such a  $t \geq t_{dc} > 0$  that also satisfies (4.18) and decreases our function value. Combining all the theorems and lemmas we get the following:

**COROLLARY 5.1.** *By Lemma 5.2, Theorem 5.4, Corollary 5.5, Lemma 5.10, Lemma 5.12 and Corollary 5.14 - pISTA converges to  $\arg \min_{A \succ 0} F(A)$ .*

**5.1. Fixed point iteration at minimum.** The following Lemma 5.2 states that if a pISTA iteration ends with no update to  $A^{(k)}$ , then  $A^{(k)} = \arg \min_{A \succ 0} F(A)$ .

**LEMMA 5.2.**  $D_t^{*(k)} = 0 \iff A^{(k)} = \arg \min_{A \succ 0} F(A)$ .

*Proof.* First, we will prove that if  $A$  is the minimum of  $F(\cdot)$ , then pISTA will result in no update. Assume  $A^{(k)} = \arg \min_{A \succ 0} F(A)$ , therefore 0 is one of the sub-gradients of  $F(A)$  at  $A^{(k)}$ :

$$(5.3) \quad 0 \in \frac{\partial F(A^{(k)})}{\partial A} = \left\{ g + \alpha \mathcal{G} \left| \begin{array}{ll} \mathcal{G}_{i,j} = \text{sign}(A_{i,j}) & A_{i,j} \neq 0 \\ \mathcal{G}_{i,j} \in [-1, 1] & A_{i,j} = 0 \end{array} \right. \right\} = \left\{ g + \alpha \mathcal{G} \left| \mathcal{G} \in G(0) \right. \right\}.$$

Let  $\mathcal{G}^*$  be the sub-gradient  $\mathcal{G}$  for which  $g + \alpha \mathcal{G}^* = 0$ . Thus:

$$\begin{aligned} g + \alpha \mathcal{G}^* &= 0 \\ \Rightarrow (g + \alpha \mathcal{G}^*) \odot \mathcal{M}_A &= 0 \\ \Rightarrow A((g + \alpha \mathcal{G}^*) \odot \mathcal{M}_A) A &= 0 \\ \Rightarrow A(g \odot \mathcal{M}_A) A + \alpha A(\mathcal{G}^* \odot \mathcal{M}_A) A &= 0. \end{aligned}$$

If we write this in a sub-differential form, we get:

$$0 \in \left\{ A(g \odot \mathcal{M}_A) A + \alpha A(\mathcal{G} \odot \mathcal{M}_A) A \left| \mathcal{G} \in G(0) \right. \right\},$$

which, according to (5.1), means that  $D_t^* = 0$  has a 0 in its sub-gradients.

Now, we will prove that if a pISTA iteration ends with no update, then  $A$  is the  $F(\cdot)$  minimum. Assume  $D_t^* = 0$ , then, according to (5.2),  $\mathcal{G}_t^*$  satisfies:

$$\begin{aligned} A(g \odot \mathcal{M}_A) A + \alpha A(\mathcal{G}_t^* \odot \mathcal{M}_A) A &= 0 \\ \Rightarrow A((g + \alpha \mathcal{G}_t^*) \odot \mathcal{M}_A) A &= 0 \quad / \cdot A^{-1} \\ \Rightarrow (g + \alpha \mathcal{G}_t^*) \odot \mathcal{M}_A &= 0. \end{aligned}$$

Let us now define  $\hat{\mathcal{G}}$  so that  $g + \alpha \hat{\mathcal{G}} \in \frac{\partial F}{\partial A}$ :

$$\hat{\mathcal{G}}_{i,j} = \begin{cases} [\mathcal{G}_t^*]_{i,j} & [\mathcal{M}_A]_{i,j} \neq 0 \\ [g/\alpha]_{i,j} & [\mathcal{M}_A]_{i,j} = 0 \end{cases}.$$

It is clear that  $g + \alpha \hat{\mathcal{G}} = 0$ , because if  $[\mathcal{M}_A]_{i,j} = 0$ , then by definition  $A_{i,j} = 0$  and  $|g| \leq \alpha$ . This implies that  $A^{(k)} = \arg \min_{A \succ 0} F(A)$ .  $\square$

In the proof of Lemma 5.2, no assumptions on  $t$  were made, therefore we have the following:

**COROLLARY 5.3.**  $\exists t > 0 : D_t^{*(k)} = 0 \iff \forall t > 0 : D_t^{*(k)} = 0$ .

**5.2. Satisfying positive definiteness.** The following Theorem 5.4 states that if  $A^{(k)} \neq \arg \min_{A \succ 0} F(A)$  then there exists a  $t_0$  such that every step-size  $t_0 > t > 0$  keeps the matrix positive definite. Thus, as stated in Corollary 5.5, the linesearch parameter can be bounded away from zero.

**THEOREM 5.4.** *For every iteration  $k$ , there exists a  $t_0$  such that for every  $t_0 > t > 0$ :  $A^{(k)} + \mathcal{M}_A^{(k)} \odot D_t^{*(k)} \succ 0$ .*

*Proof.* We first define a new constant  $\eta$ :

$$(5.4) \quad \eta = \max_{\mathcal{G}_{i,j} \in [-1,1]} \|A(g \odot \mathcal{M}_A)A + \alpha A(\mathcal{G} \odot \mathcal{M}_A)A\|_F^2.$$

Using  $\eta$  we can upper-bound the Frobenius norm of the descent step  $\mathcal{M}_A \odot D_t^*$ :

$$\begin{aligned} \|\mathcal{M}_A \odot D_t^*\|_F^2 &\leq \|D_t^*\|_F^2 \\ &= t^2 \|A(g \odot \mathcal{M}_A)A + \alpha A(\mathcal{G}_t^* \odot \mathcal{M}_A)A\|_F^2 \leq t^2 \eta. \end{aligned}$$

Let  $t_0 = \frac{\lambda_{\min}(A)}{\sqrt{\eta}}$ , then, for  $0 < t < t_0$ :

$$|\lambda_{\min}(\mathcal{M}_A \odot D_t^*)| \leq \|\mathcal{M}_A \odot D_t^*\|_F \leq t\sqrt{\eta} < \lambda_{\min}(A),$$

and by Weyl's inequality:

$$\begin{aligned} \lambda_{\min}(A + \mathcal{M}_A \odot D_t^*) &\geq \lambda_{\min}(A) + \lambda_{\min}(\mathcal{M}_A \odot D_t^*) > 0 \\ &\Rightarrow A + \mathcal{M}_A \odot D_t^* \succ 0. \end{aligned} \quad \square$$

**COROLLARY 5.5.** *The linesearch parameter  $t$  can be chosen to be bounded away from zero while keeping the matrix positive definite, i.e.,  $t \geq \frac{t_0}{2} > 0$ .*

**5.3. Decrease in function objective.** In this section we show that under suitable conditions the sequence  $\{F(A^{(k)})\}$  converges to  $\min_{A \succ 0} F(A)$  which means that the sequence  $\{A^{(k)}\}$  converges to  $\arg \min_{A \succ 0} F(A)$ . Then, we prove that our pISTA iteration satisfies those conditions, which completes the convergence proof. In this section we denote the space of symmetric positive definite matrices by  $S_{++}$ .

Denote:

$$\begin{aligned} T(\Delta; t) &= P(\mathcal{M}_A^{(k)} \odot \Delta; t) = f(A^{(k)}) \\ &\quad + \left\langle g^{(k)}, \mathcal{M}_A^{(k)} \odot \left( A^{(k)}(\mathcal{M}_A^{(k)} \odot \Delta)A^{(k)} \right) \right\rangle \\ &\quad + \frac{1}{2t} \left\langle A^{(k)}(\mathcal{M}_A^{(k)} \odot \Delta)A^{(k)}, (\mathcal{M}_A^{(k)} \odot \Delta) \right\rangle \\ &\quad + \alpha \left\| A^{(k)} + \mathcal{M}_A^{(k)} \odot \left( A^{(k)}(\mathcal{M}_A^{(k)} \odot \Delta)A^{(k)} \right) \right\|_1. \end{aligned}$$

We state a relation between  $\arg \min_{\Delta} T(\Delta; t)$  and the descent direction  $\mathcal{M}_A^{(k)} \odot D_t^{*(k)}$ :

**LEMMA 5.6.** *Denote  $\Delta_t^{*(k)} \triangleq \arg \min_{\Delta} T(\Delta; t)$ , then*

$$A^{(k)} \left( \mathcal{M}_A^{(k)} \odot \Delta_t^{*(k)} \right) A^{(k)} = \mathcal{M}_A^{(k)} \odot D_t^{*(k)}.$$

*Proof.* The sub-differential of  $T(\Delta; t)$  is defined as:

$$\frac{\partial T(\Delta; t)}{\partial \Delta} = \left\{ \mathcal{M}_A \odot \left( \frac{1}{t} A (\mathcal{M}_A \odot \Delta) A + A (g \odot \mathcal{M}_A) A + \alpha A (\mathcal{G} \odot \mathcal{M}_A) A \right) \right. \\ \left. \left| \mathcal{G} \in G(\mathcal{M}_A \odot (A(\mathcal{M}_A \odot \Delta)A)) \right. \right\}.$$

Denote  $Z = A(\mathcal{M}_A \odot \Delta)A$ , then we get:

$$\frac{\partial T(\Delta; t)}{\partial \Delta} = \left\{ \mathcal{M}_A \odot \left( \frac{1}{t} Z + A (g \odot \mathcal{M}_A) A + \alpha A (\mathcal{G} \odot \mathcal{M}_A) A \right) \right. \\ \left. \left| \mathcal{G} \in G(\mathcal{M}_A \odot Z) \right. \right\}.$$

We notice that, for  $Z_t^* = \mathcal{M}_A \odot D_t^*$ , 0 is in the sub-differential:

$$0 \in \left\{ \mathcal{M}_A \odot \left( \frac{1}{t} Z_t^* + A (g \odot \mathcal{M}_A) A + \alpha A (\mathcal{G} \odot \mathcal{M}_A) A \right) \right. \\ \left. \left| \mathcal{G} \in G(\mathcal{M}_A \odot Z_t^{*(k)}) \right. \right\}.$$

Thus, if  $\Delta_t^* = \arg \min_{\Delta} T(\Delta; t)$ , then  $A(\mathcal{M}_A \odot \Delta_t^*)A = \mathcal{M}_A \odot D_t^*$ .  $\square$

We next prove that  $f(A)$  has a Lipschitz continuous gradient under suitable conditions and state an important property for this kind of functions.

LEMMA 5.7. *For any square matrices  $A \succ 0$  and  $B$ :  $\|AB\|_F \leq \|A\|_2 \|B\|_F$ .*

*Proof.* Denote by  $b_i$  the  $i$ -th column of matrix  $B$ , then:

$$\|AB\|_F^2 = \sum_i \|Ab_i\|_2^2 \leq \|A\|_2^2 \sum_i \|b_i\|_2^2 = \|A\|_2^2 \|B\|_F^2. \quad \square$$

LEMMA 5.8. *Let  $\beta$  be a positive constant, then the function  $f(A)$  has a Lipschitz continuous gradient over the domain  $B = \{\beta I \preceq X \preceq \gamma I \mid X \in S_{++}\}$  with a Lipschitz constant of  $\frac{1}{\beta^2}$ .*

*Proof.* To prove that the function  $f(A)$  has a Lipschitz continuous gradient with a Lipschitz constant of  $\frac{1}{\beta^2}$ , we need to prove that  $\|\nabla f(X) - \nabla f(Y)\|_F \leq \frac{1}{\beta^2} \|X - Y\|_F$  for every  $X, Y \in B$ . The gradient is given by  $\nabla f(A) = S - A^{-1}$ . Thus, for every  $X, Y \in B$ :

$$\begin{aligned} \|\nabla f(X) - \nabla f(Y)\|_F &= \|Y^{-1} - X^{-1}\|_F \\ &= \|Y^{-1}(X - Y)X^{-1}\|_F \\ &\leq \|Y^{-1}\|_2 \|X - Y\|_F \|X^{-1}\|_2 \\ &= \frac{1}{\lambda_{\min}(Y)} \frac{1}{\lambda_{\min}(X)} \|X - Y\|_F \\ &\leq \frac{1}{\beta^2} \|X - Y\|_F. \quad \square \end{aligned}$$

COROLLARY 5.9. *[Properties of a function with Lipschitz continuous gradient] Let  $\beta$  be a positive constant and  $B = \{\beta I \preceq X \preceq \gamma I \mid X \in S_{++}\}$ , then for every  $Y, X \in B$ :*

$$f(Y) \leq f(X) + \langle \nabla f(X), (Y - X) \rangle + \frac{1}{2\beta^2} \langle Y - X, Y - X \rangle .$$

The rest of the proofs are inspired by [27], [16] and [29], following the proofs of [27] with slight modifications.

LEMMA 5.10 (corresponds to lemma 4.1 in [27]). *Let  $\beta$  be a positive constant and  $B = \{\beta I \preceq X \preceq \gamma I \mid X \in S_{++}\}$ . Assume that for every iteration  $k$ ,  $A^{(k)} \in B$ , then:*

$$(5.5) \quad F(A^{(k)}) - F(A^{(k+1)}) \geq L \cdot \|A^{(k)} - A^{(k+1)}\|_F^2 .$$

Where  $L > 0$ . Furthermore, the linesearch parameter  $t$  can be chosen to be bounded away from zero, i.e.,  $t \geq t_{min} > 0$ .

*Proof.* According to pISTA iteration,  $A^{(k+1)} = A^{(k)} + \mathcal{M}_A^{(k)} \odot D_t^{*(k)}$ . Therefore,

$$F(A^{(k)}) - F(A^{(k+1)}) = F(A^{(k)}) - F(A^{(k)} + \mathcal{M}_A^{(k)} \odot D_t^{*(k)}) ,$$

and from now on, we shall omit the iteration symbol  $k$ :

$$\begin{aligned} F(A^{(k)}) - F(A^{(k+1)}) &= F(A) - F(A + \mathcal{M}_A \odot D_t^*) \\ &= f(A) + \alpha \|A\|_1 - f(A + \mathcal{M}_A \odot D_t^*) - \alpha \|A + \mathcal{M}_A \odot D_t^*\|_1 . \end{aligned}$$

Using Corollary 5.9 on  $f(A + \mathcal{M}_A \odot D_t^*)$ :

$$\begin{aligned} F(A^{(k)}) - F(A^{(k+1)}) &\geq \alpha \|A\|_1 - \left( \langle g, \mathcal{M}_A \odot D_t^* \rangle \right. \\ &\quad \left. + \frac{1}{2\beta^2} \langle \mathcal{M}_A \odot D_t^*, \mathcal{M}_A \odot D_t^* \rangle + \alpha \|A + \mathcal{M}_A \odot D_t^*\|_1 \right) . \end{aligned}$$

By the definition of  $T(\Delta; t)$  and Lemma 5.6:

$$\langle g, \mathcal{M}_A \odot D_t^* \rangle = T(\Delta_t^*; t) - \frac{1}{2t} \langle \mathcal{M}_A \odot D_t^*, A^{-1} (\mathcal{M}_A \odot D_t^*) A^{-1} \rangle - f(A) ,$$

therefore,

$$\begin{aligned} F(A^{(k)}) - F(A^{(k+1)}) &\geq f(A) + \alpha \|A\|_1 - \frac{1}{2\beta^2} \langle \mathcal{M}_A \odot D_t^*, \mathcal{M}_A \odot D_t^* \rangle \\ &\quad + \frac{1}{2t} \langle \mathcal{M}_A \odot D_t^*, A^{-1} (\mathcal{M}_A \odot D_t^*) A^{-1} \rangle - T(\Delta_t^*; t) . \end{aligned}$$

Because  $\Delta_t^* = \arg \min_{\Delta} T(\Delta; t)$ , we know that  $T(\Delta_t^*; t) \leq T(0; t) = f(A) + \alpha \|A\|_1$ :

$$\begin{aligned} F(A^{(k)}) - F(A^{(k+1)}) &\geq f(A) + \alpha \|A\|_1 - \frac{1}{2\beta^2} \langle \mathcal{M}_A \odot D_t^*, \mathcal{M}_A \odot D_t^* \rangle \\ &\quad + \frac{1}{2t} \langle \mathcal{M}_A \odot D_t^*, A^{-1} (\mathcal{M}_A \odot D_t^*) A^{-1} \rangle - T(0; t) \\ &\geq -\frac{1}{2\beta^2} \langle \mathcal{M}_A \odot D_t^*, \mathcal{M}_A \odot D_t^* \rangle \\ &\quad + \frac{1}{2t} \langle \mathcal{M}_A \odot D_t^*, A^{-1} (\mathcal{M}_A \odot D_t^*) A^{-1} \rangle \\ &\geq \left( \frac{1}{2t\gamma^2} - \frac{1}{2\beta^2} \right) \langle \mathcal{M}_A \odot D_t^*, \mathcal{M}_A \odot D_t^* \rangle . \end{aligned}$$

Denote  $L = \frac{1}{2t\gamma^2} - \frac{1}{2\beta^2}$ , then  $L$  is positive as long as  $t < (\frac{\beta}{\gamma})^2$ . Furthermore,  $t$  can be chosen to be bounded away from zero, i.e.,  $t \geq t_{min} = \frac{1}{2} \cdot (\frac{\beta}{\gamma})^2 > 0$ .  $\square$

LEMMA 5.11 (corresponds to lemma 4.2 on [27]). *Let  $\beta$  be a positive constant and  $B = \{\beta I \preceq X \preceq \gamma I | X \in S_{++}\}$ , and assume that for every iteration  $k$ ,  $A^{(k)} \in B$ . Let  $\{A^{(k_j)}\}$  be any infinite and converging sub-series of  $\{A^{(k)}\}$  and let  $\bar{A}$  denote its limit. Then  $\bar{A} = \arg \min_{A \succ 0} F(A)$ .*

*Proof.* Since the sub-series  $\{A^{(k_j)}\}$  converges to  $\bar{A}$ , then  $\{F(A^{(k_j)})\}$  converges to  $F(\bar{A})$ . According to Lemma 5.10, the full series  $\{F(A^{(k)})\}$  is monotone and hence convergent because  $\{F(A^{(k_j)})\}$  is convergent. Therefore,  $\{F(A^{(k_{j+1})}) - F(A^{(k_j)})\} \rightarrow 0$ , which implies following (5.5) that  $\|A^{(k_j)} - A^{(k_{j+1})}\|_F^2 \rightarrow 0$  and  $\lim_{j \rightarrow \infty} A^{(k_{j+1})} = \bar{A}$ . According to pISTA,  $A^{(k+1)} = A^{(k)} + \mathcal{M}_A^{(k)} \odot D_t^{*(k)}$  and because  $\mathcal{M}_A^{(k)} \neq 0$ ,  $\|A^{(k_j)} - A^{(k_{j+1})}\|_F^2 \rightarrow 0$  implies  $\|D_t^{*(k)}\|_F^2 \rightarrow 0$ . We know that  $D_t^{*(k)}$  satisfies (5.1), by taking the limit as  $j \rightarrow \infty$  and using Lemma 5.2 we get that  $\bar{A} = \arg \min_{A \succ 0} F(A)$ .  $\square$

LEMMA 5.12. [corresponds to lemma 4.3 on [27]] *Let  $\beta$  be a positive constant and  $B = \{\beta I \preceq X \preceq \gamma I | X \in S_{++}\}$ . Assume that for every iteration  $k$ ,  $A^{(k)} \in B$ . Then the series  $\{A^{(k)}\}$  has a limit and it is given by  $\arg \min_{A \succ 0} F(A)$ .*

*Proof.* The domain  $B$  is compact, the rest of the proof follows the proof of lemma 4.3 in [27] exactly.  $\square$

We have shown that pISTA converges under the condition of bounded eigenvalues. It remains to show that pISTA satisfies it.

LEMMA 5.13 (corresponds to lemma 2 on [16]). *Let  $U = \{A | F(A) \leq F(A_0) \text{ and } A \in S_{++}\}$ . Then,  $U \subseteq B = \{\beta I \preceq X \preceq \gamma I | X \in S_{++}\}$  where  $\beta > 0$ .*

*Proof.* See lemma 2 in [16].  $\square$

COROLLARY 5.14. *The series  $\{A^{(k)}\}$  created by the pISTA algorithm satisfies (4.18), hence,  $A$  is contained in  $U$  and in  $B$ .*

**6. Numerical Results.** We compare the performance of our pISTA algorithm to the performance of GISTA [14], Orthant-Based Newton (OBN) [22] and ALM [24] (without skipping step<sup>1</sup>), all of which can be implemented efficiently and easily on GPU based on matrix operations. We note that Newton-Lasso [22], VSM [21] and PSM [9] can be implemented in the same manner on GPU too, but all of them have worse performance than OBN [22] and ALM [24] according to the authors' measurements. Other algorithms, such as BIG&QUIC [17] and BCD-IC [26], can be implemented on GPU as well. However, their implementations might be highly complex due to the differences between the CPU parallelism model and the GPU parallelism model, or their implementations might have worse results than the CPU counterparts due to the numerous scalar operations. The algorithms are all implemented in Python as recommended by their authors using 32bit floating point precision. Each algorithm is implemented using the GPU as efficiently as possible with the Cupy library.

We initialize all methods using GISTA [14] initialization where:

$$(6.1) \quad A_{i,j}^{(0)} = \begin{cases} (S_{i,i} + \alpha)^{-1} & i = j \\ 0 & i \neq j \end{cases}.$$

<sup>1</sup>As implemented by the authors: <https://www.math.ucdavis.edu/~sqma/ALM-SICS.html>

For ALM only, we consider any number which is less than or equal to  $10 \cdot \epsilon_{machine}$  as zero and set those numbers to zero at end of each iteration. For OBN we use 10 inner iterations, and for ALM we use the hyperparameters used by [24] except that we set  $\mu_0 = \frac{1}{\alpha}$  if  $\alpha < 0.5$ . For our pISTA algorithm, we limit our linesearch over  $t$  by stopping if  $t$  is less than  $10^{-4}$ . In that case, we use a step of  $t \leq \left(\frac{0.9}{cond(A^{(k)})}\right)^2$  which keeps the matrix positive definite. As proved in subsection 5.3, we can use a  $t < \left(\frac{\beta}{\gamma}\right)^2$  where  $\beta$  and  $\gamma$  are lower and upper bounds of the eigenvalues, respectively. However, we cannot compute those bounds, so we assume that  $\left(\frac{0.9}{cond(A^{(k)})}\right)^2$  is sufficient and it should satisfy the linesearch criteria.

As a stopping criterion for all the method, we follow [17, 26, 27] and use  $\min_z \|\partial F(A^{(k)})\|_1 < 10^{-2} \|A^{(k)}\|_1$ , where  $\min_z \|\partial F(A^{(k)})\|_1$  is the minimum sub-gradient norm.

All experiments were run on a machine with *Intel(R) Xeon(R) Gold 6230 2.10GHz* processor with 4 cores, 20GB RAM, *GeForce RTX 2080 Ti* GPU and *Ubuntu 18.04.5* operating system. We also use *Python3.9.2*, *CUDA11.0*, *Numpy1.20.1* and *Cupy9.0*. Our full python framework and code can be found at

[https://github.com/GalSha/GLASSO\\_Framework](https://github.com/GalSha/GLASSO_Framework).

**6.1. Synthetic Experiments.** First, we evaluate the algorithms on synthetic data. We use three different types of matrices as our ground truth:

- Chain graphs: as described in [16]. The ground truth matrix  $\Sigma^{-1}$  is set to be  $\Sigma_{i,i}^{-1} = 1$  and  $\Sigma_{i,i+1}^{-1} = \Sigma_{i,i-1}^{-1} = -0.5$ .
- Graphs with random sparsity structures: as described in [20]. We generate a sparse matrix  $U$  with non-zero elements set to be +1 or -1. We define the ground truth as  $\Sigma^{-1} = U^T U$  where all off diagonal elements are in  $[-1, 1]$ . We control the number of non-zeros in  $U$  and tune it such that  $\Sigma^{-1}$  has approximately 0.5% non-zeros.
- Planar graphs: as described in [27]. First we create our graph  $G(V, E)$ . We generate  $n$  random points on the unit square to be our vertices  $V$ . We use Delaunay triangulation to generate our edges  $E$ . Given the graph connectivity, we define  $\Sigma^{-1}$  as its graph Laplacian, i.e.,  $\Sigma_{i,j}^{-1} = -1$  if  $(i, j) \in E$  and  $\Sigma_{i,i}^{-1} = d_i$  where  $d_i$  is the degree of vertex  $i$ .

To ensure that the matrices are positive definite, we add a predefined diagonal term of  $\max\{-1.2\lambda_{min}, 10^{-1}\} \cdot I$ .

We do two sets of experiments, with  $n = 1,000$  and with  $n = 10,000$ . In each set we draw  $3\% \cdot n$  of samples and run the algorithms with two different values of  $\alpha$ . We repeat each experiment five times and show the average results in Table 1 and Table 2 for the first set and second set, respectively. Also, we compare the results of a CPU only implementation of our pISTA algorithm for both sets of experiments.

In the tables, we show the average time and iterations it took for each algorithm to reach the stopping criterion. We also show the number of non-zeros in the output matrix, and the minimum sub-gradient Frobenius norm.

From Table 1, we see that pISTA outperforms every other algorithm with respect to running time on all the low-dimension matrices except one. We notice that pISTA requires less iterations than GISTA and more iterations than OBN. This is expected as GISTA is a first order method, OBN is a second order method, and the pISTA algorithm is a quasi-Newton method. Thus, pISTA convergence rate should be between

Problem Parameters		pISTA (GPU)	pISTA (CPU)	GISTA	OBN	ALM
n	$\Sigma^{-1}$ type	time (iter)				
$m/n$	$ \Sigma^{-1} _0$	$\min_z \ \partial F(\tilde{\Sigma}^{-1})\ _F$				
$\alpha$		$\ \tilde{\Sigma}^{-1}\ _0$				
1,000	<i>Chain</i> 2998	<b>0.04s (2.0)</b>	0.35s (2.0)	0.05s (3.0)	0.05s (2.0)	0.55s (11.0)
3%		<b>0.0017</b>	0.0017	0.0275	0.0020	0.0312
0.6		<b>2959.2</b>	2959.2	2970.4	<b>2959.2</b>	2966.0
1,000	<i>Chain</i> 2998	<b>0.10s (6.6)</b>	1.40s (6.6)	0.15s (10.6)	0.11s ( <b>4.0</b> )	2.47s (53.8)
3%		0.0142	0.0142	0.0336	<b>0.0003</b>	0.5966
0.4		25307.2	25307.2	25302.0	<b>25246.0</b>	25246.8
1,000	<i>Random</i> 5936	<b>0.04s (2.2)</b>	0.32s (2.2)	0.06s (3.6)	0.06s (2.2)	0.61s (14.0)
3%		<b>0.0265</b>	0.0265	0.0462	0.0824	0.8561
0.6		<b>2184.0</b>	2184.0	2191.2	2185.2	2209.0
1,000	<i>Random</i> 5936	<b>0.10s (6.4)</b>	1.05s (6.4)	0.21s (15.2)	0.13s ( <b>4.6</b> )	2.41s (53.4)
3%		0.0666	0.0666	0.0286	<b>0.0005</b>	2.5377
0.4		26335.2	26335.2	26400.4	<b>26246.4</b>	26256.8
1,000	<i>Planar</i> 6958	<b>0.03s (2.0)</b>	0.31s (2.0)	0.04s (2.4)	0.05s (2.0)	0.47s (10.2)
3%		<b>0.0008</b>	0.0008	0.0269	0.0028	0.3156
0.6		<b>2995.6</b>	2995.6	3011.6	2997.6	3039.6
1,000	<i>Planar</i> 6958	0.24s (15.4)	3.05s (15.6)	0.28s (20.2)	<b>0.15s (5.6)</b>	3.13s (69.6)
3%		0.0244	0.0244	0.0241	<b>0.0130</b>	1.8171
0.4		28495.2	28471.6	28623.2	<b>28290.4</b>	28312.8

Table 1: Results for  $1,000 \times 1,000$  Precision Matrix  
 Bold text marks the best results

Problem Parameters		pISTA (GPU)	pISTA (CPU)	GISTA	OBN	ALM
n	$\Sigma^{-1}$ type	time (iter)				
$m/n$	$ \Sigma^{-1} _0$	$\min_z \ \partial F(\tilde{\Sigma}^{-1})\ _F$				
$\alpha$		$\ \tilde{\Sigma}^{-1}\ _0$				
10,000	<i>Chain</i> 29,998	<b>5.69s (3.0)</b>	249.18s (3.0)	10.80s (6.0)	13.85s (3.0)	103.44s (28.6)
3%		0.0101	0.0101	0.1478	<b>0.0060</b>	28.5176
0.4		<b>34685.2</b>	34685.2	35104.0	34839.2	34879.6
10,000	<i>Chain</i> 29,998	<b>12.89s (7.0)</b>	578.37s (7.0)	27.53s (16.6)	20.02s (4.0)	168.46s (48.0)
3%		0.0951	0.0951	0.8418	<b>0.0129</b>	24.0052
0.2		<b>97765.6</b>	97765.6	98432.8	97881.2	98339.2
10,000	<i>Random</i> 508,787	<b>1.56s (0.6)</b>	68.74s (0.6)	1.85s (0.6)	1.60s (0.6)	767.79s (237.4)
3%		<b>2.1558</b>	2.1558	3.0901	<b>2.1558</b>	44.2140
0.4		<b>10823.6</b>	<b>10823.6</b>	<b>10823.6</b>	<b>10823.6</b>	11072.0
10,000	<i>Random</i> 508,787	<b>5.67s (3.0)</b>	263.86s (3.0)	15.95s (9.0)	13.97s (3.0)	153.16s (44.2)
3%		4.3416	4.3416	<b>0.8136</b>	1.0047	21.1711
0.2		<b>140699.6</b>	140699.6	140869.6	140715.6	141009.8
10,000	<i>Planar</i> 69,949	<b>3.92s (2.0)</b>	174.70s (2.0)	8.84s (5.0)	13.73s (3.0)	100.34s (28.2)
3%		1.2172	1.2172	<b>0.0477</b>	0.1113	6.6973
0.4		<b>63249.6</b>	63249.6	63446.8	63324.0	63572.8
10,000	<i>Planar</i> 69,949	<b>20.91s (10.2)</b>	943.43s (10.2)	63.17s (35.8)	41.54s (7.6)	269.97s (79.4)
3%		1.4050	1.4049	<b>0.5368</b>	1.3284	12.4953
0.2		<b>194261.6</b>	194261.2	195714.0	192740.0	193075.0

Table 2: Results for  $10,000 \times 10,000$  Precision Matrix  
 Bold text marks the best results

GISTA and OBN. For small  $\alpha$ , OBN achieves a better minimum sub-gradient and better sparsity, however, the difference is negligible. Also, the GPU implementation of pISTA is up to  $\times 14$  times faster than its CPU counterpart, proving the desirability of GPU centric algorithms.

In Table 2, we see similar results. pISTA outperforms every other algorithm with respect to running time on all matrices. We notice that pISTA also achieves better sparsity patterns than the other algorithms. As expected, pISTA requires less iterations than GISTA and more iterations than OBN. For high-dimension matrices, the speedup of the GPU over CPU is up to  $\times 44$ , making it extremely valuable in high dimensions.




Fig. 1: Semilog plot of  $F(\tilde{\Sigma}^{-1}) - \min F(A)$  as function of iteration<sup>a</sup> for  $10,000 \times 10,000$  Precision Matrix with  $\alpha = 0.2$

<sup>a</sup>The iterations are not equal in complexity or time between the algorithms

In Figure 1, we show a semi-log plot of  $F(\tilde{\Sigma}^{-1}) - \min F(A)$  as a function of the iteration, however, it is important to note that the iterations are not equal in complexity or required time. We only present the plots of the  $10,000 \times 10,000$  Precision Matrices for  $\alpha = 0.2$  because of space consideration. We define  $\min F(A)$  as the minimum value achieved for  $F$  among all the algorithms and iterations in that experiment. The plots show what we expect, pISTA, as a quasi-Newton method, achieves a convergence rate between linear (GISTA) and quadratic (OBN). Moreover, we see that the convergence rate is quadratic in the first few iterations.

**6.2. Real World Data Experiments.** For real world data we use gene expression data sets that are available at the Gene Expression Omnibus <http://www.ncbi.nlm.nih.gov/geo/>. We preprocess the data to have zero mean and unit variance for each variable, i.e.,  $\text{diag}(S) = I$ . Table 3 shows the results for data sets of various sizes including the name codes of the data sets used.

In Table 3, we see that there is no silver bullet. There are some problems where



Problem Parameters		pISTA	GISTA	OBN	ALM
Data set	n m/n $\alpha$	time (iter) $\min_z \ \partial F(\tilde{\Sigma}^{-1})\ _F$ $\ \tilde{\Sigma}^{-1}\ _0$			
GSE-3016	1322 4.77% (63) 0.85	<b>0.09s (3)</b> 0.1915 <b>3478</b>	0.11s (3) <b>0.0378</b> 3560	0.16s (3) 0.2185 3490	1.01s (18) 0.2921 3596
GSE-3016	1322 4.77% (63) 0.75	<b>0.14s (5)</b> 0.0092 <b>10796</b>	0.35s (14) 0.0049 10890	0.32s (6) <b>0.0025</b> 10806	2.06s (39) 8.5051 10809
GSE-3016	1322 4.77% (63) 0.65	1.02s (33) 0.0134 20682	1.55s (66) 0.0158 20660	<b>0.66s (11)</b> 0.0016 <b>20386</b>	3.17s (59) <b>0.0011</b> 20440
GSE-26242	1536 6.25% (96) 0.85	<b>0.08s (2)</b> 0.0118 4876	0.15s (4) 0.0186 4962	0.14s (2) 0.0103 <b>4862</b>	1.67s (26) <b>0.0081</b> 4912
GSE-26242	1536 6.25% (96) 0.75	<b>0.45s (14)</b> 0.0003 13896	0.70s (24) 0.0056 13900	0.47s (7) <b>0.0001</b> <b>13798</b>	2.39s (40) 3.5156 13810
GSE-26242	1536 6.25% (96) 0.65	1.81s (32) 0.0096 28494	2.43s (83) 0.0148 28942	<b>1.01s (13)</b> <b>0.0087</b> <b>28148</b>	4.28s (66) 6.7166 28182
GSE-7039	6264 3.99% (250) 0.9	<b>2.31s (4)</b> 0.0397 34286	8.52s (15) 0.0798 34364	5.92s (4) <b>0.0014</b> <b>34250</b>	74.19s (77) 33.9381 34266
GSE-7039	6264 3.99% (250) 0.8	10.42s (14) 0.0566 49200	39.91s (69) 0.1218 50020	<b>9.39s (6)</b> <b>0.0449</b> <b>49054</b>	*s (>1000) * *
GSE-7039	6264 3.99% (250) 0.7	41.87s (41) 0.0206 70700	101.95s (166) 0.1302 70820	<b>16.39s (10)</b> <b>0.0012</b> <b>70572</b>	*s (>1000) * *
GSE-52076	11064 5.39% (596) 0.9	<b>9.78s (4)</b> <b>0.0027</b> <b>32840</b>	30.65s (13) 0.1908 32850	Out of GPU memory	103.10s (23) 23.4233 32990
GSE-52076	11064 5.39% (596) 0.8	<b>10.05s (4)</b> <b>0.1162</b> <b>42852</b>	55.54s (23) 0.2827 43086	Out of GPU memory	168.27s (39) 71.0796 43471
GSE-52076	11064 5.39% (596) 0.7	<b>17.34s (7)</b> 0.3950 <b>49612</b>	167.93s (65) <b>0.3866</b> 49644	Out of GPU memory	317.99s (75) 58.4818 49840

Table 3: Results for real world data set  
Bold text marks the best results

pISTA will outshine and there are some problems where OBN will outshine. However, OBN requires much more memory than pISTA, making it hard to use in high dimension on GPUs.

**7. Conclusions.** In this work we presented a quasi-Newton method for solving sparse inverse covariance estimation problem. Our method creates simplified approximate second order optimization where the different variables dependency can be guessed in a simple manner. Moreover, the method is designed to be implemented using matrix operations which can be done on a GPU, allowing us to solve the problem efficiently. We showed the desirability of GPU implementations and that our pISTA method has better results for various problems structures.

## Appendix A.

**A.1. General solution for sign derivative equation.** Define  $\mathcal{T}(x)$ :

$$\begin{aligned} x \neq 0: & \quad \mathcal{T}(x) = \text{sign}(x) \\ x = 0: & \quad \mathcal{T}(x) \in [-1, 1]. \end{aligned}$$

LEMMA A.1. *The solution for:*

$$(A.1) \quad x + b + c\mathcal{T}(x + a) = 0, \quad c > 0$$

is given by:

$$(A.2) \quad x = \begin{cases} -b + c & c < b - a \\ -a & c > |b - a| \\ -b - c & -c > b - a \end{cases} = -a + \text{SoftThreshold}(a - b, c).$$

*Proof.* First, consider the case where  $x > -a \Rightarrow \mathcal{T}(x + a) = 1$ :

$$\begin{cases} x + b + c = 0 \\ x > -a \end{cases} \Rightarrow \begin{cases} x = -b - c \\ x > -a \end{cases} \Rightarrow \begin{cases} x = -b - c \\ -b - c > -a \end{cases} \Rightarrow \begin{cases} x = -b - c \\ -c > b - a \end{cases}.$$

In a similar way, for  $x < -a \Rightarrow \mathcal{T}(x + a) = -1$  and we get:

$$\begin{cases} x = -b + c \\ c < b - a \end{cases}.$$

Lastly, consider the case that  $x = -a$ , then since  $c > 0$  we get:

$$0 \in -a + b + c \cdot t, \quad t \in [-1, 1]$$

which leads to

$$a - b \in [-c, c] \Rightarrow c \leq |a - b| \Rightarrow c \leq |b - a|. \quad \square$$

**A.2. Approximate solution for the pISTA sub-gradient equation.** Consider the following problem:

$$(A.3) \quad D: \quad 0 \in \left\{ \frac{1}{t}D + A(g \odot \mathcal{M}_A)A + \alpha A(\mathcal{G} \odot \mathcal{M}_A)A \right. \\ \left. \begin{array}{l} \mathcal{G}_{i,j} = [\text{sign}(A + \mathcal{M}_A \odot D)]_{i,j} \quad [A + \mathcal{M}_A \odot D]_{i,j} \neq 0 \\ \mathcal{G}_{i,j} \in [-1, 1] \quad [A + \mathcal{M}_A \odot D]_{i,j} = 0 \end{array} \right\}.$$

Let us split the equation into  $n^2$  scalar equations where the  $(i, j)$ -th equation is composed of the elements with indices  $(i, j)$ . Denote the equation variable  $D_{i,j} = x$ . The  $(i, j)$  equation is:

$$(A.4) \quad \frac{1}{t}x + [A(g \odot \mathcal{M}_A)A]_{i,j} + \alpha [A(\mathcal{G} \odot \mathcal{M}_A)A]_{i,j} = 0,$$

where  $\mathcal{G}_{i,j} = \mathcal{T}(A_{i,j} + [\mathcal{M}_A]_{i,j} \cdot x)$ . As we consider equations where  $[\mathcal{M}_A]_{i,j} \neq 0$  (the rest are zeros), we can use  $\mathcal{G}_{i,j} = \mathcal{T}(A_{i,j} + x)$ . Define:

$$(A.5) \quad \mathcal{G}_{k,l}^{-i,-j} = \begin{cases} 0 & k = i, l = j \\ 0 & k = j, l = i \\ \mathcal{G}_{k,l} & \text{otherwise} \end{cases}.$$

We can write (A.4) as:

$$(A.6) \quad \frac{1}{t}x + [A(g \odot \mathcal{M}_A)A]_{i,j} + \alpha [A(\mathcal{G}^{-i,-j} \odot \mathcal{M}_A)A]_{i,j} + C_{i,j}\mathcal{T}(A_{i,j} + x) = 0,$$

where

$$(A.7) \quad C_{i,j} = \begin{cases} \alpha \cdot (A_{i,i} \cdot A_{j,j}), & i = j \\ \alpha \cdot (A_{i,i} \cdot A_{j,j} + A_{i,j} \cdot A_{j,i}), & i \neq j \end{cases}$$

is the diagonal entry of the Kronecker matrix  $\alpha \cdot A \otimes A$  corresponding to the entry  $(i, j)$ . Note that  $C_{i,j} > 0$  since  $\alpha > 0$  and  $A^{(k)}$  is symmetric positive definite, and its diagonal is strictly positive. According to Lemma A.1, the solution to (A.6) is:

$$x = -A_{i,j} + \text{SoftThreshold}\left(A_{i,j} - t \cdot \left([A(g \odot \mathcal{M}_A)A]_{i,j} + \alpha [A(\mathcal{G}^{-i,-j} \odot \mathcal{M}_A)A]_{i,j}\right), t \cdot C_{i,j}\right).$$

Notice that:

$$(A.8) \quad \alpha [A(\mathcal{G}^{-i,-j} \odot \mathcal{M}_A)A]_{i,j} = \alpha [A(\mathcal{G} \odot \mathcal{M}_A)A]_{i,j} - C_{i,j} \cdot \mathcal{G}_{i,j} \cdot [\mathcal{M}_A]_{i,j} \\ = [\alpha A(\mathcal{G} \odot \mathcal{M}_A)A - C \odot (\mathcal{G} \odot \mathcal{M}_A)]_{i,j},$$

where  $\odot$  is the Hadmard product. To write (A.2) more compactly, first define:

$$B = A(g \odot \mathcal{M}_A)A + \alpha A(\mathcal{G} \odot \mathcal{M}_A)A - C \odot (\mathcal{G} \odot \mathcal{M}_A),$$

then we get

$$x = -A_{i,j} + \text{SoftThreshold}(A_{i,j} - t \cdot B_{i,j}, t \cdot C_{i,j}) \\ \Rightarrow D = -A + \text{SoftThreshold}(A - t \cdot B, t \cdot C).$$

#### REFERENCES

- [1] O. BANERJEE, L. EL GHAOU, AND A. D'ASPREMONT, *Model selection through sparse maximum likelihood estimation for multivariate gaussian or binary data*, J. Mach. Learn. Res., (2008).
- [2] O. BANERJEE, L. E. GHAOU, A. D'ASPREMONT, AND G. NATSOULIS, *Convex optimization techniques for fitting sparse gaussian graphical models*, in Proceedings of the 23rd International Conference on Machine Learning, 2006, p. 89–96.
- [3] A. BECK AND M. TEOULLE, *A fast iterative shrinkage-thresholding algorithm for linear inverse problems*, SIAM Journal on Imaging Sciences, (2009), pp. 183–202.
- [4] M. BOLLHOFFER, A. EFTEKHARI, S. SCHEIDEGGER, AND O. SCHENK, *Large-scale sparse inverse covariance matrix estimation*, SIAM Journal on Scientific Computing, (2019), pp. A380–A401.
- [5] X. CHEN, Y. LIU, H. LIU, AND J. CARBONELL, *Learning spatial-temporal varying graphs with applications to climate data analysis*, AAAI Conference on Artificial Intelligence, (2010).
- [6] A. D'ASPREMONT, O. BANERJEE, AND L. GHAOU, *First-order methods for sparse covariance selection*, SIAM Journal on Matrix Analysis and Applications, (2006).
- [7] A. P. DEMPSTER, *Covariance selection*, Biometrics, (1972), pp. 157–175.
- [8] A. DOBRA, C. HANS, M. JONES, J. NEVINS, G. YAO, AND M. WEST, *Sparse graphical models for exploring gene expression data*, Journal of Multivariate Analysis, (2004), pp. 196–212.
- [9] J. DUCHI, S. GOULD, AND D. KOLLER, *Projected subgradient methods for learning sparse gaussians*, in Uncertainty in Artificial Intelligence, 2008, p. 153–160.
- [10] J. FAN, J. ZHANG, AND K. YU, *Vast portfolio selection with gross-exposure constraints*, Journal of the American Statistical Association, (2012), pp. 592–606.
- [11] S. E. FINDER, E. TREISTER, AND O. FREIFELD, *Effective learning of a GMRF mixture model*, IEEE Access, (2022).

- [12] J. FRIEDMAN, T. HASTIE, AND R. TIBSHIRANI, *Sparse inverse covariance estimation with the graphical lasso*, *Biostatistics*, (2007), pp. 432–441.
- [13] A. GOLDENBERG AND A. W. MOORE, *Bayes net graphs to understand co-authorship networks?*, in *Proceedings of the 3rd International Workshop on Link Discovery*, 2005, p. 1–8.
- [14] D. GUILLOT, B. RAJARATNAM, B. ROLFS, A. MALEKI, AND I. WONG, *Iterative thresholding algorithm for sparse inverse covariance estimation*, *Advances in Neural Information Processing Systems*, (2012).
- [15] T. HASTIE, R. TIBSHIRANI, AND M. WAINWRIGHT, *Statistical Learning with Sparsity: The Lasso and Generalizations*, Chapman & Hall/CRC, 2015.
- [16] C.-J. HSIEH, M. A. SUSTIK, I. S. DHILLON, AND P. RAVIKUMAR, *Quic: Quadratic approximation for sparse inverse covariance estimation*, *Journal of Machine Learning Research*, (2014).
- [17] C.-J. HSIEH, M. A. SUSTIK, I. S. DHILLON, P. K. RAVIKUMAR, AND R. POLDRACK, *Big & quic: Sparse inverse covariance estimation for a million variables*, in *Neural Information Processing Systems*, 2013, pp. 3165–3173.
- [18] T. IDÉ, A. C. LOZANO, N. ABE, AND Y. LIU, *Proximity-Based Anomaly Detection using Sparse Structure Learning*, 2009, pp. 97–108.
- [19] S. LAURITZEN, *Graphical models*, Oxford Statistical Science Series, Clarendon Press, 1996.
- [20] L. LI AND K.-C. TOH, *An inexact interior point method for  $l_1$ -regularized sparse covariance selection*, *Mathematical Programming Computation*, (2010), p. 291–315.
- [21] Z. LU, *Smooth optimization approach for sparse covariance selection*, *SIAM Journal on Optimization*, (2009).
- [22] P. A. OLSEN, F. OZTOPRAK, J. NOCEDAL, AND S. J. RENNIE, *Newton-like methods for sparse inverse covariance estimation*, in *Neural Information Processing Systems*, 2012, p. 755–763.
- [23] H. RUE AND L. HELD, *Gaussian Markov random fields: theory and applications*, CRC press, 2005.
- [24] K. SCHEINBERG, S. MA, AND D. GOLDFARB, *Sparse inverse covariance selection via alternating linearization methods*, in *Neural Information Processing Systems*, 2010, p. 2101–2109.
- [25] R. TIBSHIRANI, *Regression shrinkage and selection via the lasso*, *Journal of the Royal Statistical Society. Series B (Methodological)*, (1996), pp. 267–288.
- [26] E. TREISTER AND J. S. TUREK, *A block-coordinate descent approach for large-scale sparse inverse covariance estimation*, in *Neural Information Processing Systems (NIPS)*, 2014.
- [27] E. TREISTER, J. S. TUREK, AND I. YAVNEH, *A multilevel framework for sparse optimization with application to inverse covariance estimation and logistic regression*, *SIAM Journal on Scientific Computing*, 38 (2016), pp. S566–S592.
- [28] E. TREISTER AND I. YAVNEH, *A multilevel iterated-shrinkage approach to  $l_1$  penalized least-squares minimization*, *IEEE transactions on signal processing*, 60 (2012), pp. 6319–6329.
- [29] S. J. WRIGHT, R. D. NOWAK, AND M. A. T. FIGUEIREDO, *Sparse reconstruction by separable approximation*, *IEEE Transactions on Signal Processing*, (2009), pp. 2479–2493.
- [30] X. XUAN AND K. MURPHY, *Modeling changing dependency structure in multivariate time series*, in *Proceedings of the 24th International Conference on Machine Learning*, 2007, p. 1055–1062.
- [31] M. YUAN AND Y. LIN, *Model selection and estimation in the gaussian graphical model*, *Biometrika*, (2007), pp. 19–35.