

Sharp estimates for screened Vlasov-Poisson system around Penrose-stable equilibria in \mathbb{R}^d , $d \geq 3$.

Lingjia Huang*, Quoc-Hung Nguyen[†] and Yiran Xu[‡]

Abstract

In this paper, we study the asymptotic stability of Penrose-stable equilibria among solutions of the screened Vlasov-Poisson system in \mathbb{R}^d with $d \geq 3$ that was first established by Bedrossian, Masmoudi, and Mouhot in [6] with smooth initial data. More precisely, we prove the sharp decay estimates for the density of the perturbed system, exactly like the free transport with only Hölder (i.e. C^a for $0 < a < 1$) perturbed initial data. This improves the recent works in [17] by Han-Kwan, Nguyen, and Rousset for lower derivatives of the density and in [29] by T. Nguyen for higher derivatives with a logarithmic correction in time. We establish new estimates and cancellations of the kernel to the linearized problem to obtain this result. Furthermore, we also prove this result for the Vlasov-Poisson system in which the electric field obeys a general nonlinear Poisson equation containing electrons mass/ions case.

1 Introduction

This paper is devoted to the study the asymptotic stability of equilibria for the Vlasov-Poisson system of the form:

$$\begin{cases} \partial_t f_i + v \cdot \nabla_x f_i + E \cdot \nabla_v f_i = 0 \\ E = -\nabla_x U, -\Delta U + U = \rho_i - 1 + A(U), \quad \rho_i(t, x) = \int_{\mathbb{R}^d} f_i(t, x, v) dv \end{cases} \quad (1.1)$$

on the whole space $x \in \mathbb{R}^d$, $v \in \mathbb{R}^d$, $d \geq 3$, where $f_i = f_i(t, x, v) \geq 0$ is the probability distribution of charged particles in plasma, $\rho_i(t, x)$ is the electric charge density, and $E = E(t, x)$ is electric field and $A : \mathbb{R} \rightarrow \mathbb{R}$ is smooth and satisfies $A(r) = o(r)$ as $r \rightarrow 0$. In particular, massless electrons/ ions case if $A(r) = r + 1 - e^r$; screened case if $A(r) = 0$; and unscreened case if $A(r) = r$.

This system describes a hot, unconfined, electrostatic plasma, ions in d dimensions of electrons on a uniform, static, background of ions with $d \geq 3$. This system has been extensively studied ([1–4, 8, 11, 13–16, 19, 20, 26, 30, 31]), focusing on the global existence, regularity results and longtime behavior of solutions. The first paper on global existence of weak solutions with $A(r) = r$ is due to Arsenév in [1]. Then Batt in [4] established global existence for spherically symmetric data, and this was extended by Horst in [19] to global classical solvability with cylindrically symmetric data. After that in [2], Bardos and Degond obtained global existence for small data with $d \geq 3$ by using Lagrangian approach. Next, in [30] Pfaffelmoser proved the global existence of a smooth solution with large (unrestricted size) data. Later, simpler proofs of the same were published by Schaeffer [31], Horst [20], and Lions and Pertharne [26]. Moreover, the system with massless electrons (i.e. $A(r) = r + 1 - e^r$) in \mathbb{R}^d with $d \leq 3$ was also studied by Bouchut in [8], (see also [13, 16]). In recent years, in [22] they indicated the sharp faster decay of derivatives. Later on, in [10] they extended the results of (1.1) to $d \geq 2$ with better decay of the electric field in the case of screened interactions. Vector fields approach in [32] and Fourier analysis approach in [23, 33] were also developed in the last few years.

In this paper, we study the stability and the long time behavior of solutions f_i to (1.1) in the form

$$f_i(t, x, v) = \mu(v) + f(t, x, v)$$

where $\mu(v)$ is a stable equilibrium with $\int_{\mathbb{R}^d} \mu(v) dv = 1$ and $f_i(t = 0, x, v)$ closes to $\mu(v)$. So, f solves the following perturbed system

$$\begin{cases} \partial_t f + v \cdot \nabla_x f + E \cdot \nabla_v \mu = -E \cdot \nabla_v f, \\ E = -\nabla_x U, -\Delta U + U = \rho + A(U), \quad \rho(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv, \\ f|_{t=0} = f_0. \end{cases} \quad (1.2)$$

Here are our assumptions in this paper.

*E-mail address: ljhuang20@fudan.edu.cn, Fudan University, 220 Handan Road, Yangpu, Shanghai, 200433, China.

[†]E-mail address: qhnguyen@amss.ac.cn, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, PR China.

[‡]E-mail address: yrxu20@fudan.edu.cn, Fudan University, 220 Handan Road, Yangpu, Shanghai, 200433, China.

Assumption 1 μ satisfies the Penrose stability condition:

$$\inf_{\tau \in \mathbb{R}, \xi \in \mathbb{R}^d} \left| 1 - \int_0^{+\infty} e^{-i\tau s} \frac{1}{1 + |\xi|^2} i\xi \cdot \widehat{\nabla_v \mu}(s\xi) ds \right| \geq \hat{c}, \quad (1.3)$$

for some constant $\hat{c} > 0$, $\widehat{\nabla_v \mu}$ is the Fourier transform of $\nabla_v \mu$ in \mathbb{R}^d .

Assumption 2 $\mu \in L^1$ satisfies

$$\|\langle \cdot \rangle^N \nabla_v \mu(\cdot)\|_{W^{2,\infty}} + \|\langle \cdot \rangle^{d+5} \nabla \mu(\cdot)\|_{W^{2d+7,1}} + \frac{1}{N-d} \leq M^*,$$

for some $N > d$ and $M^* < \infty$

Assumption 3 $A : \mathbb{R} \rightarrow \mathbb{R}$ is C^2 and satisfies

$$\sup_{|r| \leq 1} \left(\left| \frac{A(r)}{r^2} \right| + \left| \frac{A'(r)}{r} \right| + |A''(r)| \right) \leq C_A,$$

for some constant $C_A > 0$.

Note that a particular example for the assumption 3 is $A(r) = r + 1 - e^r$ corresponding to the Vlasov–Poisson system with electrons mass, see [8].

It is well-known that the free transport equation $\partial_t f + v \nabla_x f = 0$ exhibits phase mixing which the spatial density $\rho(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv$ decays in time. It was an observation of Landau in [25] that the linearized Vlasov-Poisson equations near homogeneous Penrose stable equilibria also have the spatial density decaying in time. Under the Penrose condition (1.3), nonlinear Landau damping was proved on $\mathbb{T}^d \times \mathbb{R}^d$ in the pioneering work [27] by Mouhot and Villani for data with Gevrey regularity (see also [7, 29] for refinements and simplifications). In the whole space $\mathbb{R}^d \times \mathbb{R}^d$ with $d \geq 3$, it was established for the screened Vlasov-Poisson system in [6] by Bedrossian, Masmoudi and Mouhot for data with finite Sobolev regularity. Their proof relies on the dispersive mechanism in Fourier space to control the plasma echo resonance. However, a decay in time of their result is far from being optimal, as the dispersion in the physical space was not taken into account. In the recent work [17], Han-Kwan, Nguyen, and Rousset revisited the asymptotic stability of Penrose stable equilibria. They obtained the decay estimates for the density ρ as follows:

$$\|\rho(t)\|_{L^1} + (1+t)\|\nabla \rho(t)\|_{L^1} + (1+t)^d \|\rho(t)\|_{L^\infty} + (1+t)^{d+1} \|\nabla \rho(t)\|_{L^\infty} \lesssim \varepsilon_0 \log(t+2) \quad \forall t > 0. \quad (1.4)$$

This is achieved by a pointwise dispersive estimate, directly on the resolvent kernel for the linearized system around Penrose stable equilibria μ . Moreover, (1.4) is optimal up to a logarithmic correction. Very recently, in [24] Ionescu, Pausader, Wang and Widmayer proved the first asymptotic stability result for the unscreened Vlasov-Poisson system in \mathbb{R}^3 around the poisson equilibrium. The unscreened case is open for the general equilibria. However, in [5, 18] they studied the linearized unscreened Vlasov-Poisson equation around suitably stable homogeneous equilibria in $\mathbb{R}_x^d \times \mathbb{R}_v^d$.

Our goal in this paper is to prove (1.4) for (1.2) *without* $\log(t+2)$ (in RHS). We recall the Besov norm and Triebel-Lizorkin norm as

$$\|g\|_{\dot{B}_{p,\infty}^s} := \sup_{\alpha} \frac{\|\delta_{\alpha} g(x)\|_{L_x^p}}{|\alpha|^s}, \quad \|g\|_{\dot{F}_{p,\infty}^s} := \left\| \sup_{\alpha} \frac{|\delta_{\alpha} g(x)|}{|\alpha|^s} \right\|_{L_x^p},$$

and for $\kappa \in (0, 1)$ we define a norm weighted in time:

$$\|g\|_{\kappa} = \sum_{p=1,\infty} \sup_{s \in [0,\infty)} \left(\langle s \rangle^{\frac{d(p-1)}{p}} \|g(s)\|_{L^p} + \langle s \rangle^{\kappa + \frac{d(p-1)}{p}} \|g(s)\|_{\dot{B}_{p,\infty}^{\kappa}} \right), \quad \|g\|_{l,\kappa} = \sum_{j=0}^l \|\langle s \rangle^j \nabla^j g\|_{\kappa}.$$

By Gagliardo-Nirenberg interpolation inequality, we know that

$$\|g\|_{l,\kappa} \sim \|g\|_{\kappa} + \|\langle s \rangle^l \nabla^l g\|_{\kappa} \sim \sum_{p=1,\infty} \sup_{s \in [0,\infty)} \left(\langle s \rangle^{\frac{d(p-1)}{p}} \|g(s)\|_{L^p} + \langle s \rangle^{l+\kappa + \frac{d(p-1)}{p}} \|\nabla^l g(s)\|_{\dot{B}_{p,\infty}^{\kappa}} \right).$$

Let us introduce some notations.

Notation 1.1 Let $g : \mathbb{R}_x^d \times \mathbb{R}_v^d \rightarrow \mathbb{R}^m$. We define, for $(x, v) \in \mathbb{R}_x^d \times \mathbb{R}_v^d$ and $a \in (0, 1)$

$$\begin{aligned} \dot{D}_1^a g(x, v) &= \sup_{z \in \mathbb{R}^d} \frac{|g(x, v) - g(x - z, v)|}{|z|^a}, & \dot{D}_2^a g(x, v) &= \sup_{z \in \mathbb{R}^d} \frac{|g(x, v) - g(x, v - z)|}{|z|^a}, \\ \dot{D}^a g &= \dot{D}_1^a g + \dot{D}_2^a g, & D^a g &= |g| + \dot{D}^a g, & D_i^a g &= |g| + \dot{D}_i^a g \quad i = 1, 2. \end{aligned}$$

Our main result is as follows.

Theorem 1.2 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$. There exist $C_0 > 0$, $\tilde{\varepsilon} \in (0, 1)$ such that for $0 < \varepsilon \leq \tilde{\varepsilon}$ if*

$$\sum_{p=1, \infty} \sum_{i=0, 1} \|\mathcal{D}^a(\nabla_{x,v}^i f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} \leq \frac{1}{C_0} \varepsilon, \quad (1.5)$$

then the problem (1.2) has a unique global solution f with

$$\|\rho\|_a + \tilde{\varepsilon}^{\frac{1}{3}} \|U\|_a \leq \varepsilon \quad \text{and} \quad \|\rho\|_{1,a} + \|U\|_{1,a} \lesssim_{\varepsilon, a, M^*} 1.$$

In addition, with $(\varrho_1(t, x), \mathcal{U}_1(t, x)) = (\int f_0(x - tv, v) dv, 0)$ we have the estimate that

$$\|\rho - \varrho_1\|_a + \tilde{\varepsilon}^{\frac{1}{3}} \|U - \mathcal{U}_1\|_a \lesssim_{\varepsilon, M^*, a} \varepsilon^{\frac{4}{3}}. \quad (1.6)$$

Moreover, for $m \in \mathbb{N}^+$ and $b \in (0, 1)$ if

$$\sum_{j=2}^{m+1} \sup_{|r| \leq 1} |A^{(j)}(r)| \leq C_{A,m}, \quad \sum_{j=0}^m \sum_{p=1, \infty} \|\mathcal{D}^a(\nabla_{x,v}^j f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} \leq \mathbf{c}, \quad \sum_{j=1}^{m+3} \sup_{v \in \mathbb{R}^d} \langle v \rangle^N |\nabla^j \mu(v)| \leq \mathbf{c}',$$

for some $N > d$, then

$$\|\rho\|_{m,b} + \|U\|_{m,b} \lesssim_{\varepsilon, a, M^*, \mathbf{c}, \mathbf{c}'} 1.$$

Remark 1.3 *Thanks to (1.6), we obtain that our estimate $\langle t \rangle^{\frac{d(p-1)}{p}} \|\rho(t)\|_{L_x^p} \lesssim \varepsilon$ is optimal. Indeed, consider $f_0(x, v) = \varepsilon(1 + |x|^2)^{-d}(1 + |v|^2)^{-d}$, then we have $\langle t \rangle^{\frac{d(p-1)}{p}} \|\varrho_1(t)\|_{L_x^p} \gtrsim \varepsilon$ which implies $\langle t \rangle^{\frac{d(p-1)}{p}} \|\rho(t)\|_{L_x^p} \gtrsim \varepsilon$ for ε small enough.*

Remark 1.4 *If we have initial data f_0 with higher-order derivatives, we can get more regularity of the solution ρ . It is different from the condition in [29], which made smallness assumptions on higher derivatives of f_0 .*

Remark 1.5 *The crucial decay estimates of $E(t, x)$ depend on the fact that we work in dimension $d \geq 3$. And the decay will be stronger when d becomes higher. The decay estimates is insufficient in the case $d \leq 2$ which can be seen in the proof of Proposition 4.1, where we need to estimate $\sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{a-\delta}}$. That is the reason why we only consider $d \geq 3$ in our paper. We deal with the 2d case in [21].*

Remark 1.6 *While finishing our article, we learned that Toan Nguyen was working on this problem and had a similar result to ours. But the two works are independent.*

As [17, Corollary 1.1], thanks to Theorem 1.2, we also obtain the following scattering property for the solution to (1.2). The proof is omitted as it is very analogous to [17, Proof of Corollary 1.1].

Corollary 1.7 *With the same assumptions and notations as in Theorem 1.2, there is a function $f_{\infty} \in W^{1, \infty}$ given by*

$$f_{\infty}(x, v) = f_0(x + Y_{\infty}(x, v), v + W_{\infty}(x, v)) + \mu(v + W_{\infty}(x, v)) - \mu(v),$$

such that

$$\|f(t, x + tv, v) - f_{\infty}(x, v)\|_{L_{x,v}^{\infty}} \lesssim_{a, M^*} \frac{\varepsilon}{\langle t \rangle^{d+a-1}}, \quad t \geq 0,$$

and $\|Y_{\infty}\|_{L_{x,v}^{\infty}} + \|W_{\infty}\|_{L_{x,v}^{\infty}} \lesssim \varepsilon$.

Let us discuss the main idea in this paper. First, we give the equivalence of (ρ, U) and (f, ρ) satisfies the system

$$\begin{aligned} (\rho, U) &= (\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U)), \\ \mathcal{F}_1(\rho, U) &= G *_{(t,x)} ((\mathcal{I} + \mathcal{R})(\rho, U)) + (\mathcal{I} + \mathcal{R})(\rho, U), \\ \mathcal{F}_2(\rho, U) &= (1 - \Delta)^{-1}(\rho + A(U)), \end{aligned}$$

where the kernel G is defined in (2.6) and \mathcal{I}, \mathcal{R} are defined in (2.3). This means, (ρ, U) is a fixed point of $\mathcal{F} = (\mathcal{F}_1, \mathcal{F}_2)$. We establish the boundedness of the mapping $\mathcal{G}(\cdot) = G *_{(t,x)} \cdot$ under the norm $\|\cdot\|_a$. To do this, we need to estimate some pointwise decay estimates of the kernel $G(t, x)$, which gives a new proof to be different from the scaling method introduced by Han-Kwan, Nguyen, and Rousset in [17]. It should be emphasized that the cancellation of $G(t, x)$, i.e. $\int_{\mathbb{R}^d} G(t, x) dx = 0$, $\forall t \geq 0$, and the estimate of $\| |x|^a |\delta_{\alpha} G(t, x)| \|_{L_x^p}$ play a very important role in proving the boundedness of the mapping \mathcal{G} . Using the decay estimates for the characteristics

to prove maps $\rho \mapsto \mathcal{I}(\rho)$, $\rho \mapsto \mathcal{R}(\rho)$ be compressed mappings with small contracting constant. Finally, we apply the contracting mapping principle to get a fixed point of \mathcal{F} .

The paper is organized as follows. In section 2, we establish the equivalence of (ρ, U) and f , then we focus on the estimate of (ρ, U) in the following sections. In section 3, we give the pointwise decay estimates of the kernel $G(t, x)$ and its derivatives. Then, we prove the bound: $\|\mathcal{G}(\mathbf{f})\|_a \leq M\|\mathbf{f}\|_a$, for some constant $M = M(\check{c}, d, M_1)$. In section 4, we establish the pointwise estimate of the characteristics $(X_{s,t}(x, v), V_{s,t}(x, v))$ and their derivatives. In section 5, we estimate the contribution of the initial data and explain that why we need the norm $L_x^1 L_v^1 \cap L_v^1 L_x^1 \cap L_x^1 L_v^\infty \cap L_v^1 L_x^\infty$ of the initial data f_0 . In Section 6, we estimate the reaction term $\mathcal{R}(\rho)$ and its higher derivatives. To handle it, we shall introduce a general map

$$\mathcal{T}[F, \eta](t, x) = - \int_0^t \int_{\mathbb{R}^d} F(s, X_{s,t}(x, v)) \eta(V_{s,t}(x, v)) dv ds + \int_0^t \int_{\mathbb{R}^d} F(s, x - (t-s)v) \eta(v) dv ds,$$

and give the estimate in Proposition 6.1. In Section 7, we prove the main theorem.

The following are some notations in our paper.

Notation 1.8 *Throughout this paper, $A \lesssim B$ means that there exists constant C only depending dimension d such that $A \leq CB$. $A \lesssim_M B$ means that there exists constant $C = C(d, M)$ such that $A \leq CB$. The same convention is adopted for $A \gtrsim B$ and $A \sim B$.*

Notation 1.9 *In this paper, $\tilde{\cdot}$ is the "space-time" Fourier transform on \mathbb{R}^{d+1} as follows*

$$\tilde{g}(\tau, \xi) = \int_{\mathbb{R}} \int_{\mathbb{R}^d} e^{-i\tau t} e^{-ix \cdot \xi} g(t, x) dx dt.$$

And we define inverse Fourier transform

$$\mathcal{F}^{-1}(h)(t, x) = \frac{1}{(2\pi)^{d+1}} \int_{\mathbb{R}} \int_{\mathbb{R}^d} e^{i\tau t} e^{ix \cdot \xi} h(\tau, \xi) d\tau d\xi.$$

Notation 1.10 *For multi-index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$, we denote $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_d$.*

Notation 1.11 *For a vector $x \in \mathbb{R}^d$, and $k \in \mathbb{N}$, we define $x^{\otimes k} = \underbrace{x \otimes x \dots \otimes x}_k$.*

Acknowledgements: Q.H.N. is supported by the Academy of Mathematics and Systems Science, Chinese Academy of Sciences startup fund, and the National Natural Science Foundation of China (No. 12050410257 and No. 12288201) and the National Key R&D Program of China under grant 2021YFA1000800. Q.H.N. also wants to thank Benoit Pausader for his stimulating comments and suggestion to consider the Vlasov-Poisson system with massless electrons.

2 Equivalence of (ρ, U) and f

In this section, we will show that the system (1.2) is equivalent to an equation in terms of (ρ, U) . Let $(X_{s,t}(x, v), V_{s,t}(x, v))$ be the flow associated to the vector field $(v, E(t, x))$, i.e. $(X_{s,t}(x, v), V_{s,t}(x, v))$ satisfies the ODE system:

$$\begin{cases} \frac{d}{ds} X_{s,t}(x, v) = V_{s,t}(x, v), & X_{t,t}(x, v) = x, \\ \frac{d}{ds} V_{s,t}(x, v) = E(s, X_{s,t}(x, v)), & V_{t,t}(x, v) = v, \end{cases} \quad (2.1)$$

for any $0 \leq s \leq t$, $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$. Then, the solution f of the equation (1.2) is given by

$$f(t, x, v) = f_0(X_{0,t}(x, v), V_{0,t}(x, v)) - \int_0^t E(s, X_{s,t}(x, v)) \cdot \nabla_v \mu(V_{s,t}(x, v)) ds, \quad (2.2)$$

(see (1.3) in [28] with $\mathbf{B}(t, x, v) = (v, E(t, x))$). We denote

$$\begin{aligned} \mathcal{I}(\rho, U)(t, x) &= \int_{\mathbb{R}^d} f_0(X_{0,t}(x, v), V_{0,t}(x, v)) dv, \\ \mathcal{R}(\rho, U)(t, x) &= \int_0^t \int_{\mathbb{R}^d} (E(s, x - (t-s)v) \cdot \nabla_v \mu(v) - E(s, X_{s,t}(x, v)) \cdot \nabla_v \mu(V_{s,t}(x, v))) dv ds. \end{aligned} \quad (2.3)$$

Hence, $\rho(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv$ is the solution of the following equation:

$$\begin{aligned}
\rho(t, x) &= \int_0^t \int_{\mathbb{R}^d} [\nabla_x (1 - \Delta_x)^{-1} (\rho + A \circ U)](s, x - (t - s)v) \cdot \nabla_v \mu(v) dv ds + \mathcal{I}(\rho, U)(t, x) + \mathcal{R}(\rho, U)(t, x) \\
&= \int_0^t \int_{\mathbb{R}^d} [\nabla_x (1 - \Delta_x)^{-1} (\rho + A \circ U)](s, w) \cdot \nabla_v \mu\left(\frac{x - w}{t - s}\right) \frac{1}{(t - s)^d} dw ds + \mathcal{I}(\rho, U)(t, x) + \mathcal{R}(\rho, U)(t, x) \\
&= \int_0^t \int_{\mathbb{R}^d} (\rho(s, w) + A \circ U(s, w)) \left(1 - \frac{1}{(t - s)^2} \Delta_v\right)^{-1} \Delta_v \mu\left(\frac{x - w}{t - s}\right) \frac{1}{(t - s)^{d+1}} dw ds \\
&\quad + \mathcal{I}(\rho, U)(t, x) + \mathcal{R}(\rho, U)(t, x).
\end{aligned} \tag{2.4}$$

Define $K(t, x) = \frac{1}{t^{d+1}} \left((1 - \frac{1}{t^2} \Delta_v)^{-1} \Delta_v \mu \right) \left(\frac{x}{t} \right) \mathbf{1}_{t>0}$, then ρ can be expressed in the following way

$$\rho = (K *_{(t,x)} (\rho + A(U))) + \mathcal{I}(\rho, U) + \mathcal{R}(\rho, U).$$

Next, taking Fourier transform in (t, x) , we have

$$\tilde{\rho} = \tilde{K} \tilde{\rho} + \tilde{K} \widetilde{A(U)} + \widetilde{\mathcal{I}(\rho, U)} + \widetilde{\mathcal{R}(\rho, U)}.$$

Then we have

$$\rho = G *_{(t,x)} (\mathcal{I}(\rho) + \mathcal{R}(\rho) + A(U)) + \mathcal{I}(\rho) + \mathcal{R}(\rho), \tag{2.5}$$

where G is the kernel satisfying:

$$\tilde{G} = \frac{\tilde{K}}{1 - \tilde{K}}, \tag{2.6}$$

and

$$\tilde{K}(\tau, \xi) = \int_0^{+\infty} e^{-i\tau t} \frac{1}{1 + |\xi|^2} i\xi \cdot \widehat{\nabla_v \mu}(t\xi) dt.$$

Thanks to the Penrose condition, one has $|1 - \tilde{K}| \geq \mathring{c} > 0$, which implies that \tilde{G} is well defined.

It is clear that if we have f , we can obtain the formula of ρ and U . On the other hand, if we have ρ and U , we can get E , then we recover the characteristics $(X_{s,t}(x, v), V_{s,t}(x, v))$ to finally get f from (2.2).

Remark 2.1 We will use $\|\langle \cdot \rangle^N \nabla_v \mu(\cdot)\|_{W^{2,\infty}}$ to estimate the reaction term $\mathcal{R}(\rho, U)$, and $\|\langle \cdot \rangle^{d+5} \nabla \mu(\cdot)\|_{W^{2d+7,1}}$ will be used to deal with the boundedness of the operator \mathcal{G} .

We define the operator \mathcal{F} as:

$$\mathcal{F}(\rho, U) = (\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U)), \tag{2.7}$$

where

$$\begin{aligned}
\mathcal{F}_1(\rho, U) &= G *_{(t,x)} (\mathcal{I}(\rho, U) + \mathcal{R}(\rho, U) + A(U)) + \mathcal{I}(\rho, U) + \mathcal{R}(\rho, U), \\
\mathcal{F}_2(\rho, U) &= (1 - \Delta)^{-1} (\rho + A(U)).
\end{aligned}$$

We need to prove that \mathcal{F} has a fixed point i.e. $\mathcal{F}(\rho, U) = (\rho, U)$.

We define the norm

$$\|(g_1, g_2)\|_{S_\kappa^\varepsilon} := \|g_1\|_\kappa + \varepsilon^{\frac{1}{3}} \|g_2\|_\kappa, \quad \|(g_1, g_2)\|_{l,\kappa} := \|g_1\|_{l,\kappa} + \|g_2\|_{l,\kappa}.$$

The main Theorem 1.2 is a consequence of the following theorems:

Theorem 2.2 Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$. There exist constants $C_0 > 0$ and $\tilde{\varepsilon} > 0$ such that for any $\varepsilon \in (0, \tilde{\varepsilon}]$ we have

$$\|\mathcal{F}(\rho, U)\|_{S_\varepsilon^\varepsilon} \leq \varepsilon, \quad \|\mathcal{F}(\rho_1, U_1) - \mathcal{F}(\rho_2, U_2)\|_{S_\varepsilon^\varepsilon} \leq \frac{1}{2} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_\varepsilon^\varepsilon},$$

for any $(\rho, U), (\rho_1, U_1), (\rho_2, U_2) \in \mathfrak{B}(\varepsilon) := \{(\rho, U) \in L^\infty((0, \infty); L^1 \cap L^\infty(\mathbb{R}^d)) : \|(\rho, U)\|_{S_\varepsilon^\varepsilon} \leq \varepsilon\}$, provided that

$$\sum_{i=0,1} \sum_{p=1,\infty} \|\mathcal{D}^a(\nabla_{x,v}^i f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} \leq \frac{1}{C_0} \varepsilon. \tag{2.8}$$

Under condition (2.8), \mathcal{F} has a unique fixed point (ρ, U) in $\mathfrak{B}(\varepsilon)$ with $\|(\rho, U)\|_{1,a} \lesssim_{\varepsilon,a,M^*} 1$. Moreover, set $(\varrho_1(t, x), \mathcal{U}_1(t, x)) = (\int f_0(x - tv, v) dv, 0)$, we have

$$\|(\rho - \varrho_1, U - \mathcal{U}_1)\|_{S_\varepsilon^\varepsilon} \lesssim_{\varepsilon, M^*, a} \varepsilon^{\frac{4}{3}}.$$

Theorem 2.3 Let ρ be in Theorem 2.2. Then for $b \in (0, 1)$ and $m \geq 2$, we have

$$\|(\rho, U)\|_{m,b} \lesssim_{\check{c}, a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} 1,$$

provided the initial data f_0 , the stationary state μ and function A with

$$\sum_{j=2}^{m+1} \sup_{|r| \leq 1} |A^{(j)}(r)| \leq C_{A,m}, \quad \sum_{j=0}^m \sum_{p=1, \infty} \|\mathcal{D}^a(\nabla_{x,v}^j f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} \leq \mathbf{c}', \quad \langle v \rangle^N \sum_{j=0}^{m+3} |\nabla^j \mu(v)| \leq \mathbf{c}'. \quad (2.9)$$

3 Estimate of the kernel \mathbf{G} and boundedness of operator \mathcal{G}

In this section, we prove the boundedness of the operator $\mathcal{G} : \mathbf{f} \mapsto G *_{(t,x)} \mathbf{f}$ under the norm $\|\cdot\|_a$. The main result is

$$\|\mathcal{G}(\mathbf{f})\|_a \leq M \|\mathbf{f}\|_a,$$

for some constant $M = M(\check{c}, d, M_1)$.

To do this, we need the following Lemmas:

Lemma 3.1 Define

$$Q_0(u, \tau, \xi) = \int_0^{+\infty} e^{-i\tau t} u(t\xi) dt.$$

Then,

$$|Q_0(u, \tau, \xi)| \lesssim \frac{\sup_x (|u(x)| \langle x \rangle^2)}{|\xi|}, \quad (3.1)$$

$$|Q_0(u, \tau, \xi)| \lesssim \sum_{j=1}^{m-1} \frac{|\xi|^j}{|\tau|^{j+1}} |\nabla^j u(0)| + \frac{|\xi|^{m-1}}{|\tau|^m} \sup_{x \in \mathbb{R}} (|\nabla^m u(x)| \langle x \rangle^2). \quad (3.2)$$

Proof. Firstly, we can get the result (3.1) directly as follows,

$$|Q_0(u, \tau, \xi)| \leq \int_0^{+\infty} |u(t\xi)| dt \leq \sup_{z \in \mathbb{R}^d} (|u(z)| \langle z \rangle^2) \int_0^{+\infty} \frac{1}{\langle t\xi \rangle^2} dt \sim \sup_{z \in \mathbb{R}^d} (|u(z)| \langle z \rangle^2) \frac{1}{|\xi|}.$$

Moreover, we integrate by parts m -times to obtain that for any nonnegative integer m ,

$$Q_0(u, \tau, \xi) = - \left(\sum_{j=0}^{m-1} \frac{1}{(i\tau)^{j+1}} \xi^{\otimes j} : \nabla^j u(0) \right) + \frac{1}{(i\tau)^m} Q(\xi^{\otimes m} : \nabla^m u(t\xi), \tau, \xi).$$

Combining the above equality with the first inequality, it is easy to check the conclusion (3.2). ■

Lemma 3.2 For $k \in \mathbb{Z}^+$, define

$$Q_k(u, \tau, \xi) = \int_0^{+\infty} t^k e^{-i\tau t} u(t\xi) dt. \quad (3.3)$$

Then,

$$|Q_k(u, \tau, \xi)| \lesssim \frac{\sup_x (|u(x)| \langle x \rangle^{2+k})}{|\xi|^{k+1}}, \quad (3.4)$$

$$|Q_k(u, \tau, \xi)| \lesssim \frac{1}{|\tau|^{k+1}} \left(|u(0)| + \sum_{l=0}^k \sup_{x \in \mathbb{R}} (|\nabla^{k-l+1} u(x)| \langle x \rangle^{2+k-l}) \right), \quad (3.5)$$

$$|Q_k(u, \tau, \xi)| \lesssim \frac{1}{(|\tau| + |\xi|)^{k+1}} \left(\sup_x (|u(x)| \langle x \rangle^{2+k}) + \sum_{l=0}^k \sup_{x \in \mathbb{R}} (|\nabla^{k-l+1} u(x)| \langle x \rangle^{2+k-l}) \right). \quad (3.6)$$

Moreover, if $u(0) = 0$, we have

$$|Q_k(u, \tau, \xi)| \lesssim \frac{|\xi|}{|\tau|^{k+2}} \sum_{l=0}^{k+1} \sup_{x \in \mathbb{R}} (|\nabla^{k-l+2} u(x)| \langle x \rangle^{3+k-l}), \quad (3.7)$$

$$|Q_k(u, \tau, \xi)| \lesssim \frac{|\xi|}{(|\tau| + |\xi|)^{k+2}} \left(\sup_x (|u(x)| \langle x \rangle^{2+k}) + \sum_{l=0}^{k+1} \sup_{x \in \mathbb{R}} (|\nabla^{k-l+2} u(x)| \langle x \rangle^{3+k-l}) \right). \quad (3.8)$$

Proof. First of all, $\forall j \in \{1, 2, \dots, d\}$, we obtain

$$\xi_j^k Q_k(u, \tau, \xi) = \int_0^{+\infty} (t\xi_j)^k e^{-i\tau t} u(t\xi) dt = Q_0(u_1, \tau, \xi),$$

where $u_1(\xi) = \xi_j^k u(\xi)$. By (3.1), one has,

$$|\xi_j^k Q_k(u, \tau, \xi)| \lesssim \frac{\sup_x (|u(x)| \langle x \rangle^{2+k})}{|\xi|},$$

which implies (3.4).

Note that

$$\nabla^l u_1(0) = 0, \quad \forall l \leq k-1, \quad |\nabla^l u_1(0)| \lesssim |\nabla^{l-k} u(0)|, \quad \forall l \geq k.$$

Then, we know from (3.2) with $m = k+1$ that

$$\begin{aligned} |\xi_j^k Q_k(u, \tau, \xi)| &= Q_0(u_1, \tau, \xi) \lesssim \frac{|\xi|^k}{|\tau|^{k+1}} |\nabla^k u_1(0)| + \frac{|\xi|^k}{|\tau|^{k+1}} \sup_{x \in \mathbb{R}} (|\nabla^{k+1} u_1(x)| \langle x \rangle^2) \\ &\lesssim \frac{|\xi|^k}{|\tau|^{k+1}} |u(0)| + \frac{|\xi|^k}{|\tau|^{k+1}} \sum_{l=0}^k \sup_{x \in \mathbb{R}} (|\nabla^{k-l+1} u(x)| \langle x \rangle^{2+k-l}). \end{aligned}$$

Then,

$$|\xi|^k |Q_k(u, \tau, \xi)| \lesssim \frac{|\xi|^k}{|\tau|^{k+1}} |u(0)| + \frac{|\xi|^k}{|\tau|^{k+1}} \sum_{l=0}^k \sup_{x \in \mathbb{R}} (|\nabla^{k-l+1} u(x)| \langle x \rangle^{2+k-l}),$$

which implies (3.5).

Then (3.4) and (3.5) imply (3.6) directly.

Moreover, if $u(0) = 0$, thanks to (3.2) with $m = k+2$, we have

$$\begin{aligned} |\xi_j^k Q_k(u, \tau, \xi)| &\lesssim \frac{|\xi|^{k+1}}{|\tau|^{k+2}} |\nabla^{k+1} u_1(0)| + \frac{|\xi|^{k+1}}{|\tau|^{k+2}} \sup_{x \in \mathbb{R}} (|\nabla^{k+2} u_1(x)| \langle x \rangle^2) \\ &\lesssim \frac{|\xi|^{k+1}}{|\tau|^{k+2}} |\nabla u(0)| + \frac{|\xi|^{k+1}}{|\tau|^{k+2}} \sum_{l=0}^{k+1} \sup_{x \in \mathbb{R}} (|\nabla^{k-l+2} u(x)| \langle x \rangle^{3+k-l}) \\ &\lesssim \frac{|\xi|^{k+1}}{|\tau|^{k+2}} \sum_{l=0}^{k+1} \sup_{x \in \mathbb{R}} (|\nabla^{k-l+2} u(x)| \langle x \rangle^{3+k-l}), \end{aligned}$$

which implies (3.7). Finally, (3.4) and (3.7) imply (3.8). ■

We define

$$\|F\|_{\mathcal{W}_\alpha^\beta} := \sum_{j=0}^{\beta} \int_{\mathbb{R}^d} \langle x \rangle^\alpha |\nabla^j F(x)| dx,$$

and use this norm in the following lemmas.

Lemma 3.3 For integer $\alpha \geq 0$ and $\beta \geq 2$, we have following estimates:

$$\left| \nabla_\xi \tilde{K}(\tau, \xi) \right| \lesssim \frac{|\xi|}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} \|\nabla \mu\|_{\mathcal{W}_3^\beta}, \quad (3.9)$$

$$\left| \nabla_\xi^\beta \tilde{K}(\tau, \xi) \right| \lesssim \left(\frac{1}{\langle \xi \rangle^\beta} \frac{1}{(|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2} \frac{1}{(|\tau| + |\xi|)^\beta} \right) \|\nabla \mu\|_{\mathcal{W}_{\beta+2}^{2\beta+1}}, \quad (3.10)$$

$$\left| \partial_\tau^\alpha \tilde{K}(\tau, \xi) \right| \lesssim \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{\alpha+2}} \|\nabla \mu\|_{\mathcal{W}_{\alpha+3}^{\alpha+2}}. \quad (3.11)$$

Proof. Recall that

$$\tilde{K}(\tau, \xi) = \int_0^{+\infty} e^{-i\tau t} \frac{1}{1 + |\xi|^2} i\xi \cdot \widehat{\nabla}_v \mu(t\xi) dt. \quad (3.12)$$

Then for $\alpha \geq 0$, we have

$$\partial_\tau^\alpha \tilde{K}(\tau, \xi) = \sum_{j=1}^d \frac{(-i)^\alpha \xi_j}{\langle \xi \rangle^2} \int_0^{+\infty} t^\alpha e^{-i\tau t} P_j(t\xi) dt,$$

where

$$P(\xi) = (P_1, P_2, \dots, P_d)(\xi) = \widehat{\nabla_v \mu}(\xi), \quad P(0) = 0.$$

Thanks to (3.8), we have

$$\begin{aligned} \left| \int_0^{+\infty} t^\alpha e^{-i\tau t} P_j(t\xi) dt \right| &\lesssim \frac{|\xi|}{(|\tau| + |\xi|)^{\alpha+2}} \left(\sup_x (|P(x)| \langle x \rangle^{2+\alpha}) + \sum_{l=0}^{\alpha+1} \sup_{x \in \mathbb{R}} (|\nabla^{\alpha-l+2} P(x)| \langle x \rangle^{3+\alpha-l}) \right) \\ &\lesssim \frac{|\xi|}{(|\tau| + |\xi|)^{\alpha+2}} \|\nabla \mu\|_{\mathcal{W}_{\alpha+3}^{\alpha+2}}. \end{aligned}$$

Thus we obtain that

$$\left| \partial_\tau^\alpha \tilde{K}(\tau, \xi) \right| \lesssim \sum_{j=1}^d \frac{|\xi|}{\langle \xi \rangle^2} \left| \int_0^{+\infty} t^\alpha e^{-i\tau t} P_j(t\xi) dt \right| \lesssim \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{\alpha+2}} \|\nabla \mu\|_{\mathcal{W}_{\alpha+3}^{\alpha+2}}.$$

Thanks to (3.6) with $k = \beta_1$, one has

$$\begin{aligned} &\left| \int_0^{+\infty} t^{\beta_1} e^{-i\tau t} (\nabla^{\beta_1} P_j)(t\xi) dt \right| \\ &\lesssim \frac{1}{(|\tau| + |\xi|)^{\beta_1+1}} \left(\sup_x (|\nabla^{\beta_1} P(x)| \langle x \rangle^{2+\beta_1}) + \sum_{l=0}^{\beta_1} \sup_{x \in \mathbb{R}} (|\nabla^{2\beta_1-l+1} P(x)| \langle x \rangle^{2+\beta_1-l}) \right) \\ &\lesssim \frac{1}{(|\tau| + |\xi|)^{\beta_1+1}} \|\nabla \mu\|_{\mathcal{W}_{\beta_1+2}^{2\beta_1+1}}. \end{aligned}$$

Therefore,

$$\left| \nabla_\xi^\beta \tilde{K}(\tau, \xi) \right| \lesssim \sum_{j=1}^d \frac{|\xi|}{\langle \xi \rangle^2} \left| \int_0^{+\infty} t e^{-i\tau t} (\nabla P_j)(t\xi) dt \right| + \sum_{j=1}^d \frac{1}{\langle \xi \rangle^2} \left| \int_0^{+\infty} e^{-i\tau t} P_j(t\xi) dt \right| \lesssim \frac{|\xi|}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} \|\nabla \mu\|_{\mathcal{W}_3^3}.$$

For $\beta \geq 2$, we have

$$\begin{aligned} \left| \nabla_\xi^\beta \tilde{K}(\tau, \xi) \right| &\lesssim \sum_{j=1}^d \sum_{\beta_1=0}^{\beta} \left| \nabla_\xi^{\beta-\beta_1} \left(\frac{\xi_j}{\langle \xi \rangle^2} \right) \right| \left| \int_0^{+\infty} t^{\beta_1} e^{-i\tau t} (\nabla^{\beta_1} P_j)(t\xi) dt \right| \\ &\lesssim \sum_{j=1}^d \sum_{\beta_1=1}^{\beta-1} \frac{1}{\langle \xi \rangle^{\beta-\beta_1+1}} \left| \int_0^{+\infty} t^{\beta_1} e^{-i\tau t} (\nabla^{\beta_1} P_j)(t\xi) dt \right| \\ &\quad + \sum_{j=1}^d \frac{|\xi|}{\langle \xi \rangle^2} \left| \int_0^{+\infty} t^\beta e^{-i\tau t} (\nabla^\beta P_j)(t\xi) dt \right| + \sum_{j=1}^d \frac{1}{\langle \xi \rangle^{\beta+1}} \left| \int_0^{+\infty} e^{-i\tau t} P_j(t\xi) dt \right| \\ &\lesssim \|\nabla \mu\|_{\mathcal{W}_{\beta+2}^{2\beta+1}} \left(\frac{1}{\langle \xi \rangle^\beta} \frac{1}{(|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2} \frac{1}{(|\tau| + |\xi|)^\beta} \right). \end{aligned}$$

Thus, the proof is complete. \blacksquare

Lemma 3.4 For any integer $j \geq 1$, there holds

$$\left| \nabla^j \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \right| \lesssim M_j \left(\frac{1}{\langle \xi \rangle^j (|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2 (|\tau| + |\xi|)^j} \right), \quad (3.13)$$

$$\left| \partial_\tau^j \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \right| \lesssim M_j \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{j+2}}, \quad (3.14)$$

where

$$M_j = \|\nabla \mu\|_{\mathcal{W}_{j+2}^{2j+1}} \left(1 + \|\nabla \mu\|_{\mathcal{W}_{j+2}^{2j+1}} \right)^{j-1}.$$

Proof. Thanks to the Penrose condition, one has $|1 - \tilde{K}(\tau, \xi)| \geq \hat{c}$. Firstly, we have

$$\left| \nabla \frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right| = \left| \nabla \left(\frac{1}{1 - \tilde{K}(\tau, \xi)} \right) \right| \lesssim \hat{c} |\nabla \tilde{K}(\tau, \xi)| \stackrel{(3.9)}{\lesssim} \frac{|\xi|}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} \|\nabla \mu\|_{\mathcal{W}_3^3}.$$

Note that we get the following lines for any $j \geq 1$,

$$\begin{aligned}
\left| \nabla^j \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \right| &= \left| \nabla^j \left(\frac{1}{1 - \tilde{K}(\tau, \xi)} \right) \right| \lesssim_{\tilde{c}} \sum_{i=1}^j |\nabla^i \tilde{K}(\tau, \xi)|^{\frac{j}{i}} \\
&\lesssim_{\tilde{c}} \sum_{i=2}^j \left(\frac{1}{\langle \xi \rangle^j} \frac{1}{(|\tau| + |\xi|)^{\frac{2j}{i}}} + \frac{1}{\langle \xi \rangle^{\frac{2j}{i}}} \frac{1}{(|\tau| + |\xi|)^j} \right) \|\nabla \mu\|_{\mathcal{W}_3^{2i+1}}^{\frac{j}{i}} + \frac{|\xi|^j}{\langle \xi \rangle^{2j} (|\tau| + |\xi|)^{2j}} \|\nabla \mu\|_{\mathcal{W}_3^3}^j \\
&\lesssim_{\tilde{c}} \left(\frac{1}{\langle \xi \rangle^j} \frac{1}{(|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2} \frac{1}{(|\tau| + |\xi|)^j} \right) \left(\|\nabla \mu\|_{\mathcal{W}_3^{2j+1}} + \|\nabla \mu\|_{\mathcal{W}_3^{2j+1}}^j \right) + \frac{|\xi|^j}{\langle \xi \rangle^{2j} (|\tau| + |\xi|)^{2j}} \|\nabla \mu\|_{\mathcal{W}_3^3}^j \\
&\lesssim_{\tilde{c}} M_j \left(\frac{1}{\langle \xi \rangle^j (|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2 (|\tau| + |\xi|)^j} \right).
\end{aligned}$$

Then, we consider the derivative of τ to obtain

$$\begin{aligned}
\left| \partial_\tau^j \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \right| &= \left| \partial_\tau^j \left(\frac{1}{1 - \tilde{K}(\tau, \xi)} \right) \right| \lesssim_{\tilde{c}} \sum_{j_1=1}^j \left| \partial_\tau^{j_1} \tilde{K}(\tau, \xi) \right|^{\frac{j}{j_1}} \stackrel{(3.11)}{\lesssim_{\tilde{c}}} \sum_{j_1=1}^j \left| \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{j_1+2}} \right|^{\frac{j}{j_1}} \|\nabla \mu\|_{\mathcal{W}_3^{j_1+3}}^{\frac{j}{j_1}} \\
&\lesssim_{\tilde{c}} \frac{1}{(|\tau| + |\xi|)^j} \left(\frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} + \frac{|\xi|^{2j}}{\langle \xi \rangle^{2j} (|\tau| + |\xi|)^{2j}} \right) \left(\|\nabla \mu\|_{\mathcal{W}_3^{j+3}}^j + \|\nabla \mu\|_{\mathcal{W}_3^{j+3}}^{j+2} \right) \\
&\lesssim_{\tilde{c}} M_j \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{j+2}}.
\end{aligned}$$

This completes the proof. \blacksquare

Lemma 3.5 Define $G_n = \mathcal{F}^{-1} \left(\tilde{G} \chi_n \right)$, where $\chi \in [0, 1]$ is a smooth compactly supported function in the annulus $1/4 \leq |\xi| \leq 4$ and $\chi_n(\xi) = \chi(\frac{\xi}{2^n})$. Then, we have for any integer $\gamma \geq 0$

$$|\nabla^\gamma G_n(t, x)| \lesssim_{\tilde{c}} 2^{(d+\gamma+1)n-2n^+} M_{d+\gamma+2} \min \left\{ 1, \frac{1}{2^n(t+|x|)} \right\}^{d+\gamma+2},$$

where $n^+ = \max\{0, n\}$.

Proof. Firstly, we estimate $|\nabla^\gamma G_n(t, x)|$ directly.

$$\begin{aligned}
|\nabla^\gamma G_n(t, x)| &= \left| \mathcal{F}^{-1} \left(\xi^{\otimes \gamma} \frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \chi_n(\xi) \right) \right| \lesssim \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \left| \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \xi^{\otimes \gamma} \chi_n(\xi) \right| d\xi d\tau \\
&\lesssim_{\tilde{c}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} 2^{\gamma n} |\tilde{K}(\tau, \xi)| d\tau d\xi.
\end{aligned}$$

Using (3.11), one has

$$|\nabla^\gamma G_n(t, x)| \lesssim_{\tilde{c}} \|\nabla \mu\|_{\mathcal{W}_3^2} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} 2^{\gamma n} \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} d\tau d\xi \lesssim_{\tilde{c}} 2^{(d+\gamma+1)n-2n^+} \|\nabla \mu\|_{\mathcal{W}_3^2}. \quad (3.15)$$

Then we integrate by parts in τ to get that

$$\begin{aligned}
|\nabla^\gamma G_n(t, x)| &\sim \left| \int_{\mathbb{R} \times \mathbb{R}^d} e^{i\tau t} e^{i\xi \cdot x} \frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \xi^{\otimes \gamma} \chi_n(\xi) d\xi d\tau \right| \\
&\lesssim \frac{1}{|t|^{m_1}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \left| \partial_\tau^{m_1} \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right) \xi^{\otimes \gamma} \chi_n(\xi) \right| d\xi d\tau.
\end{aligned}$$

Using (3.14), we know that

$$\begin{aligned}
|\nabla^\gamma G_n(t, x)| &\lesssim_{\tilde{c}} M_{m_1} \frac{1}{|t|^{m_1}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \frac{|\xi|^{2+\gamma}}{\langle \xi \rangle^2 (|\tau| + |\xi|)^{m_1+2}} d\xi d\tau \\
&\lesssim_{\tilde{c}} \frac{1}{(|t|2^n)^{m_1}} 2^{(d+\gamma+1)n-2n^+} M_{m_1}.
\end{aligned} \quad (3.16)$$

Finally, we integrate by parts in ξ to get that

$$\begin{aligned}
|\nabla^\gamma G_n(t, x)| &\sim \left| \int_{\mathbb{R} \times \mathbb{R}^d} e^{i\tau t} e^{i\xi \cdot x} \frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \xi^{\otimes \gamma} \chi_n(\xi) d\xi d\tau \right| \\
&\lesssim \frac{1}{|x|^{m_2}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \left| \nabla^{m_2} \left(\frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \xi^{\otimes \gamma} \chi_n(\xi) \right) \right| d\xi d\tau \\
&\lesssim_{\dot{c}} \frac{1}{|x|^{m_2}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} |\tilde{K}(\tau, \xi)| |\nabla^{m_2} (\xi^{\otimes \gamma} \chi_n(\xi))| d\xi d\tau \\
&\quad + \frac{1}{|x|^{m_2}} \sum_{j=1}^{m_2} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \left| \nabla^j \frac{\tilde{K}(\tau, \xi)}{1 - \tilde{K}(\tau, \xi)} \right| |\nabla^{m_2-j} (\xi^{\otimes \gamma} \chi_n(\xi))| d\xi d\tau.
\end{aligned}$$

Using (3.9) and (3.13), we know that

$$\begin{aligned}
|\nabla^\gamma G_n(t, x)| &\lesssim_{\dot{c}} \|\nabla \mu\|_{W_3^2} \frac{1}{|x|^{m_2}} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} \frac{|\xi|^2}{\langle \xi \rangle^2 (|\tau| + |\xi|)^2} 2^{(\gamma-m_2)n} d\xi d\tau \\
&\quad + \frac{1}{|x|^{m_2}} \sum_{j=1}^{m_2} \int_{\tau \in \mathbb{R}, |\xi| \sim 2^n} M_j \left(\frac{1}{\langle \xi \rangle^j} \frac{1}{(|\tau| + |\xi|)^2} + \frac{1}{\langle \xi \rangle^2} \frac{1}{(|\tau| + |\xi|)^j} \right) 2^{(\gamma-m_2+j)n} d\xi d\tau \\
&\lesssim_{\dot{c}} \frac{M_{m_2}}{|x|^{m_2}} \sum_{j=1}^{m_2} \int_{|\xi| \sim 2^n} \left[\frac{|\xi|}{\langle \xi \rangle^2} 2^{(\gamma-m_2)n} + \left(\frac{1}{\langle \xi \rangle^j} \frac{1}{|\xi|} + \frac{1}{\langle \xi \rangle^2} \frac{1}{|\xi|^{j-1}} \right) 2^{(\gamma-m_2+j)n} \right] d\xi \\
&\lesssim_{\dot{c}} \frac{1}{(|x|2^n)^{m_2}} 2^{(d+\gamma+1)n-2n^+} M_{m_2}.
\end{aligned} \tag{3.17}$$

Combining (3.15), (3.16) with (3.17) and taking $m_1 = m_2 = d + \gamma + 2$, we obtain

$$|\nabla^\gamma G_n(t, x)| \lesssim_{\dot{c}} 2^{(d+\gamma+1)n-2n^+} M_{d+\gamma+2} \min \left\{ 1, \frac{1}{2^n(t+|x|)} \right\}^{d+\gamma+2}.$$

This gives the result. ■

Theorem 3.6 *Let $\gamma \geq 0$, $0 \leq a_0 < 1$ and $0 < b_0 < 1$. Then,*

$$|\nabla^\gamma G(t, x)| \lesssim_{\dot{c}} \frac{\tilde{M}_\gamma}{(t+|x|)^{d+\gamma-1} (1+t+|x|)^2}. \tag{3.18}$$

In particular,

$$\|\nabla^\gamma G(t, x)\|_{L_x^\infty} \lesssim_{\dot{c}} \frac{\tilde{M}_\gamma}{t^{d+\gamma-1} (1+t)^2}, \tag{3.19}$$

$$\||x|^{a_0} \nabla^\gamma G(t, x)\|_{L_x^1} \lesssim_{\dot{c}} \frac{\tilde{M}_\gamma}{t^{\gamma-1-a_0} (1+t)^2} + \tilde{M}_\gamma \mathbf{1}_{t \leq 1} + \tilde{M}_\gamma \mathbf{1}_{\gamma=1+a_0} \log^+ \left(\frac{1}{t} \right), \tag{3.20}$$

$$\||x|^{b_0} \delta_\alpha G(t, x)\|_{L_x^1} \lesssim_{\dot{c}} \tilde{M}_1 \frac{|\alpha|}{(1+t+|\alpha|)^{2-b_0}} + \tilde{M}_1 \mathbf{1}_{|\alpha| > t} \frac{|\alpha|^{b_0}}{1+t} + \tilde{M}_1 \mathbf{1}_{|\alpha| \leq t} \frac{|\alpha|^{d+b_0}}{t^{d-1} (1+t)^2}, \tag{3.21}$$

where

$$\tilde{M}_\gamma = \overline{M}_\gamma (1 + \overline{M}_\gamma)^{d+\gamma+1}, \quad \text{and} \quad \overline{M}_\gamma = \|\langle \cdot \rangle^{d+\gamma+4} \nabla \mu(\cdot)\|_{W^{2d+2\gamma+5,1}}.$$

Remark 3.7 *To obtain the results of this paper, we use these estimates with $\gamma = 0, 1$. But we would like to give the estimate for all $\gamma \geq 0$, which will be used in our future work.*

Proof. Firstly, we prove (3.18). Note that

$$\|\nabla \mu\|_{W_\alpha^\beta} \lesssim \|\langle \cdot \rangle^\alpha \nabla \mu\|_{W^{\beta,1}},$$

thus

$$M_{d+\gamma+2} \lesssim \overline{M}_\gamma (1 + \overline{M}_\gamma)^{d+\gamma+1} = \tilde{M}_\gamma.$$

Since

$$|\nabla^\gamma G(t, x)| \leq \sum_{n \leq 0} |\nabla^\gamma G_n(t, x)| + \sum_{n > 0} |\nabla^\gamma G_n(t, x)|,$$

for $n > 0$, combining this with Lemma 3.5, we can yield that

$$\begin{aligned}
\sum_{n>0} |\nabla^\gamma G_n(t, x)| &\lesssim_{\dot{c}} \widetilde{M}_\gamma \sum_{n>0} 2^{(d+\gamma-1)n} \min \left\{ 1, \frac{1}{2^n(t+|x|)} \right\}^{d+\gamma+2} \\
&\sim_{\dot{c}} \widetilde{M}_\gamma \sum_{n>0} \left(\mathbf{1}_{t+|x|\leq 1} \sum_{1\leq 2^n\leq (t+|x|)^{-1}} 2^{(d+\gamma-1)n} + \mathbf{1}_{t+|x|\leq 1} \sum_{2^n>(t+|x|)^{-1}} 2^{-3n} \frac{1}{(t+|x|)^{d+\gamma+2}} \right. \\
&\quad \left. + \mathbf{1}_{t+|x|>1} \frac{1}{2^{3n}(t+|x|)^{d+\gamma+2}} \right) \\
&\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\mathbf{1}_{t+|x|\leq 1} \frac{1}{(t+|x|)^{d+\gamma-1}} + \mathbf{1}_{t+|x|>1} \frac{1}{(t+|x|)^{d+\gamma+2}} \right).
\end{aligned}$$

Similarly, we have for $n \leq 0$

$$\begin{aligned}
\sum_{n\leq 0} |\nabla^\gamma G_n(t, x)| &\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\mathbf{1}_{t+|x|\leq 1} \sum_{n\leq 0} 2^{(d+\gamma+1)n} + \mathbf{1}_{t+|x|>1} \sum_{2^n\leq (t+|x|)^{-1}} 2^{(d+\gamma+1)n} \right. \\
&\quad \left. + \mathbf{1}_{t+|x|>1} \sum_{(t+|x|)^{-1}<2^n\leq 1} \frac{2^{-n}}{(t+|x|)^{d+\gamma+2}} \right) \\
&\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\mathbf{1}_{t+|x|\leq 1} + \mathbf{1}_{t+|x|>1} \frac{1}{(t+|x|)^{d+\gamma+1}} \right).
\end{aligned}$$

Finally, we yield that

$$\begin{aligned}
|\nabla^\gamma G(t, x)| &\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\mathbf{1}_{t+|x|\leq 1} \frac{1}{(t+|x|)^{d+\gamma-1}} + \mathbf{1}_{t+|x|>1} \frac{1}{(t+|x|)^{d+\gamma+1}} \right) \\
&\lesssim_{\dot{c}} \frac{\widetilde{M}_\gamma}{(t+|x|)^{d+\gamma-1}(1+t+|x|)^2}.
\end{aligned}$$

(1) Now we prove (3.19), we consider the L^∞ norm of G as follows

$$\|\nabla^\gamma G(t, x)\|_{L_x^\infty} \lesssim_{\dot{c}} \sup_{x\in\mathbb{R}^d} \frac{\widetilde{M}_\gamma}{(t+|x|)^{d+\gamma-1}(1+t+|x|)^2} \lesssim_{\dot{c}} \frac{\widetilde{M}_\gamma}{t^{d+\gamma-1}(1+t)^2}.$$

(2) Now we prove (3.20), for the L^1 norm, we have

$$\||x|^{a_0} \nabla^\gamma G(t, x)\|_{L_x^1} \lesssim_{\dot{c}} \widetilde{M}_\gamma \int \frac{|x|^{a_0}}{(t+|x|)^{d+\gamma-1}(1+t+|x|)^2} dx.$$

Now we consider the case $t \geq 1$, and changing variables to obtain

$$\||x|^{a_0} \nabla^\gamma G(t, x)\|_{L_x^1} \lesssim_{\dot{c}} \widetilde{M}_\gamma \int \frac{|x|^{a_0}}{(t+|x|)^{d+\gamma+1}} dx \sim_{\dot{c}} \frac{\widetilde{M}_\gamma}{t^{\gamma+1-a_0}} \int \frac{|x|^{a_0}}{(1+|x|)^{d+\gamma+1}} dx \sim_{\dot{c}} \frac{\widetilde{M}_\gamma}{t^{\gamma+1-a_0}}.$$

For the case $0 \leq t \leq 1$, we know that

$$\begin{aligned}
\||x|^{a_0} \nabla^\gamma G(t, x)\|_{L_x^1} &\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\int_{|x|\leq t} + \int_{t<|x|<1} + \int_{|x|\geq 1} \right) \frac{|x|^{a_0}}{(t+|x|)^{d+\gamma-1}(1+t+|x|)^2} dx \\
&\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\int_{|x|\leq t} \frac{|x|^{a_0}}{t^{d+\gamma-1}} dx + \int_{t<|x|<1} \frac{|x|^{a_0}}{|x|^{d+\gamma-1}} dx + \int_{|x|\geq 1} \frac{|x|^{a_0}}{|x|^{d+\gamma+1}} dx \right) \\
&\lesssim_{\dot{c}} \widetilde{M}_\gamma \left(\frac{1}{t^{\gamma-1-a_0}} + 1 + \mathbf{1}_{\gamma<1+a_0} + \mathbf{1}_{\gamma=1+a_0} \log \left(\frac{1}{t} \right) + \frac{\mathbf{1}_{\gamma>1+a_0}}{t^{\gamma-a_0-1}} \right) \\
&\sim_{\dot{c}} \widetilde{M}_\gamma \left(\frac{1}{t^{\gamma-1-a_0}} + 1 + \mathbf{1}_{\gamma=1+a_0} \log \left(\frac{1}{t} \right) \right).
\end{aligned}$$

Combining these two cases, we deduce that

$$\begin{aligned}
\||x|^{a_0} \nabla^\gamma G(t, x)\|_{L_x^1} &\lesssim_{\dot{c}} \mathbf{1}_{t\geq 1} \frac{\widetilde{M}_\gamma}{t^{\gamma+1-a_0}} + \mathbf{1}_{t\leq 1} \widetilde{M}_\gamma \left(\frac{1}{t^{\gamma-1-a_0}} + 1 + \mathbf{1}_{\gamma=1+a_0} \log \left(\frac{1}{t} \right) \right) \\
&\sim_{\dot{c}} \frac{\widetilde{M}_\gamma}{t^{\gamma-1-a_0}(1+t)^2} + \widetilde{M}_\gamma \mathbf{1}_{t\leq 1} + \mathbf{1}_{\gamma=1+a_0} \log^+ \left(\frac{1}{t} \right).
\end{aligned}$$

(3) Now we prove (3.21), we focus on $\| |x|^{b_0} \delta_\alpha G(t, x) \|_{L_x^1}$.

$$\| |x|^{b_0} \delta_\alpha G(t, x) \|_{L_x^1} = \left(\int_{|\alpha| \leq |x|/2} + \int_{|\alpha| \geq |x|/2} \right) |x|^{b_0} |\delta_\alpha G(t, x)| dx =: I_{G, b_0}^1 + I_{G, b_0}^2.$$

For the term I_{G, b_0}^1 ,

$$\begin{aligned} I_{G, b_0}^1 &\leq \int_{|\alpha| \leq |x|/2} \int_0^1 |x|^{b_0} |\nabla G(t, x - \tau\alpha)| |\alpha| d\tau dx \\ &\lesssim \varepsilon |\alpha| \int_0^1 \int_{|\alpha| \leq |x|/2} |x|^{b_0} \frac{\widetilde{M}_1}{(t + |x - \tau\alpha|)^d (1 + t + |x - \tau\alpha|)^2} dx d\tau \\ &\lesssim \varepsilon |\alpha| \int_{|\alpha| \leq |x|/2} |x|^{b_0} \frac{\widetilde{M}_1}{(t + |x|/2)^d (1 + t + |x|/2)^2} dx \\ &\lesssim \varepsilon \mathbf{1}_{|\alpha| \geq t/2} |\alpha| \int_{|\alpha| \leq |x|/2} \frac{\widetilde{M}_1}{|x|^{d-b_0} (1 + |x|)^2} dx + \mathbf{1}_{|\alpha| < t/2} |\alpha| \int_{|\alpha| \leq |x|/2} |x|^{b_0} \frac{\widetilde{M}_1}{(t + |x|/2)^d (1 + t + |x|/2)^2} dx \\ &=: I_{G, b_0}^{1,1} + I_{G, b_0}^{1,2}. \end{aligned}$$

Then, we estimate the first one and obtain

$$\begin{aligned} I_{G, b_0}^{1,1} &\lesssim \mathbf{1}_{|\alpha| \geq t/2} \mathbf{1}_{|\alpha| \geq 1} |\alpha| \int_{|\alpha| \leq |x|/2} \frac{\widetilde{M}_1}{|x|^{d+2-b_0}} dx \\ &\quad + \mathbf{1}_{|\alpha| \geq t/2} \mathbf{1}_{|\alpha| < 1} |\alpha| \left(\int_{|\alpha| \leq |x| \leq 2} \frac{\widetilde{M}_1}{|x|^{d-b_0}} dx + \int_{|x| \geq 2} \frac{\widetilde{M}_1}{|x|^{d+2-b_0}} dx \right) \\ &\lesssim \widetilde{M}_1 |\alpha| \left(\mathbf{1}_{|\alpha| \geq t/2} \mathbf{1}_{|\alpha| \geq 1} \frac{1}{|\alpha|^{2-b_0}} + \mathbf{1}_{|\alpha| \geq t/2} \mathbf{1}_{|\alpha| < 1} \right) \sim \widetilde{M}_1 \mathbf{1}_{|\alpha| \geq t/2} \frac{|\alpha|}{(1 + |\alpha|)^{2-b_0}}. \end{aligned} \quad (3.22)$$

For the other one, we have

$$\begin{aligned} I_{G, b_0}^{1,2} &\lesssim \mathbf{1}_{|\alpha| < t/2} |\alpha| \int_{|x| \geq t} \frac{\widetilde{M}_1}{|x|^{d-b_0} (1 + |x|)^2} dx + \mathbf{1}_{|\alpha| < t/2} |\alpha| \int_{2|\alpha| \leq |x| \leq t} \frac{\widetilde{M}_1 |x|^{b_0}}{t^d (1 + t)^2} dx \\ &\lesssim \mathbf{1}_{|\alpha| < t/2} \mathbf{1}_{t \geq 1} |\alpha| \int_{|x| \geq t} \frac{\widetilde{M}_1}{|x|^{d-b_0+2}} dx + \mathbf{1}_{|\alpha| < t/2} \mathbf{1}_{t \leq 1} |\alpha| \left(\int_{t \leq |x| \leq 1} \frac{\widetilde{M}_1}{|x|^{d-b_0}} dx + \int_{|x| \geq 1} \frac{\widetilde{M}_1}{|x|^{d-b_0+2}} dx \right) \\ &\quad + \mathbf{1}_{|\alpha| < t/2} |\alpha| \int_{|\alpha| \leq |x| \leq 2t} \frac{\widetilde{M}_1 |x|^{b_0}}{t^d (1 + t)^2} dx \\ &\sim \widetilde{M}_1 |\alpha| \mathbf{1}_{|\alpha| < t/2} \left(\mathbf{1}_{t \geq 1} \frac{1}{t^{2-b_0}} + \mathbf{1}_{t \leq 1} + \frac{t^{b_0}}{(1 + t)^2} \right) \sim \widetilde{M}_1 \mathbf{1}_{|\alpha| < t} \frac{|\alpha|}{(1 + t)^{2-b_0}}. \end{aligned} \quad (3.23)$$

Combining (3.22) with (3.23), we can yield

$$I_{G, b_0}^1 \lesssim \varepsilon \widetilde{M}_1 \mathbf{1}_{|\alpha| \geq t} \frac{|\alpha|}{(1 + |\alpha|)^{2-b_0}} + \widetilde{M}_1 \mathbf{1}_{|\alpha| < t} \frac{|\alpha|}{(1 + t)^{2-b_0}} \sim \varepsilon \widetilde{M}_1 \frac{|\alpha|}{(1 + t + |\alpha|)^{2-b_0}}. \quad (3.24)$$

For the term I_{G, b_0}^2 , one has

$$\begin{aligned} I_{G, b_0}^2 &\lesssim \varepsilon \widetilde{M}_0 |\alpha|^{b_0} \int_{|x| \lesssim |\alpha|} \frac{1}{(t + |x|)^{d-1} (1 + t + |x|)^2} dx \sim \varepsilon \widetilde{M}_0 |\alpha|^{b_0} \int_0^{|\alpha|} \frac{r^{d-1}}{(t + r)^{d-1} (1 + t + r)^2} dr \\ &\sim \varepsilon \mathbf{1}_{|\alpha| > t} \widetilde{M}_0 |\alpha|^{b_0} \left(\int_0^t \frac{r^{d-1}}{t^{d-1} (1 + t)^2} dr + \int_t^{|\alpha|} \frac{r^{d-1}}{r^{d-1} (1 + r)^2} dr \right) + \mathbf{1}_{|\alpha| \leq t} \widetilde{M}_0 |\alpha|^{b_0} \int_0^{|\alpha|} \frac{r^{d-1}}{t^{d-1} (1 + t)^2} dr \\ &\lesssim \varepsilon \mathbf{1}_{|\alpha| > t} \widetilde{M}_0 |\alpha|^{b_0} \left(\frac{t}{(1 + t)^2} + \frac{1}{1 + t} \right) + \mathbf{1}_{|\alpha| \leq t} \frac{\widetilde{M}_0 |\alpha|^{d+b_0}}{t^{d-1} (1 + t)^2} \\ &\lesssim \varepsilon \widetilde{M}_0 \mathbf{1}_{|\alpha| > t} \frac{|\alpha|^{b_0}}{1 + t} + \widetilde{M}_0 \mathbf{1}_{|\alpha| \leq t} \frac{|\alpha|^{d+b_0}}{t^{d-1} (1 + t)^2}. \end{aligned}$$

Combining this with (3.24) and note that $\widetilde{M}_0 \leq \widetilde{M}_1$, one has

$$\| |x|^{b_0} \delta_\alpha G(t, x) \|_{L_x^1} \lesssim \varepsilon \widetilde{M}_1 \frac{|\alpha|}{(1 + t + |\alpha|)^{2-b_0}} + \widetilde{M}_1 \mathbf{1}_{|\alpha| > t} \frac{|\alpha|^{b_0}}{1 + t} + \widetilde{M}_1 \mathbf{1}_{|\alpha| \leq t} \frac{|\alpha|^{d+b_0}}{t^{d-1} (1 + t)^2}$$

which implies (3.21). ■

Lemma 3.8 *For the kernel G defined in (2.6), there holds*

$$\int_{\mathbb{R}^d} G(t, x) dx = 0, \quad \forall t \geq 0. \quad (3.25)$$

Proof. By (3.11) with $\alpha = 0$, one has $\tilde{K}(\tau, 0) = 0$. Then from (2.6), we obtain that $\tilde{G}(\tau, 0) = 0$, for all $\tau \in \mathbb{R}$, which implies the conclusion (3.25). ■

Then we can prove the main result in this section.

Theorem 3.9 *There holds*

$$\|\mathcal{G}(\mathbf{f})\|_a \leq M \|\mathbf{f}\|_a,$$

with $M = M(\dot{c}, d, M_1)$.

It is easy to check that for any $t \in [0, 1]$

$$\sum_{p=1, \infty} \left(\|G *_{(t,x)} \mathbf{f}(t)\|_{L^p} + \|G *_{(t,x)} \mathbf{f}(t)\|_{\dot{B}_{p,\infty}^a} \right) \leq \|\mathbf{f}\|_a \int_0^1 \|G(s)\|_{L_x^1} ds \stackrel{(3.20)}{\lesssim} \dot{c}, d, M_1 \|\mathbf{f}\|_a.$$

To prove the theorem 3.9, we need to have the following Lemmas.

Lemma 3.10 *Let $t \geq 1$, and $p = 1, \infty$, we have*

$$\langle t \rangle^{\frac{d(p-1)}{p}} \|G *_{(t,x)} \mathbf{f}(t)\|_{L^p} \lesssim_{\dot{c}, M_0} \sup_{s \in [0, \infty)} \left(\|\mathbf{f}(s)\|_{L^1} + \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right).$$

Proof. We use the idea in [9]. Since we have the cancellation showed in the (3.25), we rewrite lines as follows when changing of variable $z = x - y$ in the third term,

$$\begin{aligned} \|G *_{(t,x)} \mathbf{f}\|_{L^p} &\leq \left\| \int_0^{t/2} \int_{\mathbb{R}^d} G(t-s, x-y) \mathbf{f}(s, y) dy ds \right\|_{L_x^p} \\ &\quad + \left\| \int_{t/2}^t \int_{\mathbb{R}^d} G(t-s, z) (\mathbf{f}(s, x) - \mathbf{f}(s, x+z)) dz ds \right\|_{L_x^p} := I_1 + I_2. \end{aligned}$$

First, we consider the term

$$I_1 \leq \int_0^{t/2} \|G(t-s)\|_{L^p} \|\mathbf{f}(s)\|_{L^1} ds.$$

Using (3.19) and (3.20) in Theorem 3.6, we obtain

$$I_1 \lesssim_{\dot{c}, M_0} \int_0^{t/2} \frac{1}{\langle t-s \rangle^{1 + \frac{d(p-1)}{p}}} \|\mathbf{f}(s)\|_{L^1} ds \lesssim_{\dot{c}, M_0} \widetilde{M} \langle t \rangle^{-\frac{d(p-1)}{p}} \sup_{s \in [0, \infty)} \|\mathbf{f}(s)\|_{L^1}. \quad (3.26)$$

Similarly, we have

$$\begin{aligned} I_2 &= \left\| \int_{t/2}^t \int_{\mathbb{R}^d} G(t-s, z) |z|^a \frac{\mathbf{f}(s, x) - \mathbf{f}(s, x+z)}{|z|^a} dz ds \right\|_{L_x^p} \\ &\leq \int_{t/2}^t \int_{\mathbb{R}^d} |G(t-s, z)| |z|^a \frac{\|\mathbf{f}(s, x) - \mathbf{f}(s, x+z)\|_{L_x^p}}{|z|^a} dz ds \\ &\leq \int_{t/2}^t \sup_z \frac{\|\delta_z \mathbf{f}(s, x)\|_{L_x^p}}{|z|^a} \|G(t-s, z)\|_{L_x^1} |z|^a ds. \end{aligned} \quad (3.27)$$

Using (3.20) in Theorem 3.6, we have,

$$\begin{aligned} I_2 &\lesssim_{\dot{c}, M_0} \widetilde{M} \int_{t/2}^t \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \frac{1}{\langle t-s \rangle^{1-a}} ds \\ &\lesssim_{\dot{c}, M_0} \widetilde{M} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right] \langle t \rangle^{-\frac{d(p-1)}{p} - a} \int_{t/2}^t \frac{1}{\langle t-s \rangle^{1-a}} ds \\ &\lesssim_{\dot{c}, M_0} \widetilde{M} \langle t \rangle^{-\frac{d(p-1)}{p}} \sup_{s \in [0, \infty)} \left(\langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right). \end{aligned}$$

Combining (3.26) and (3.27) we obtain that

$$\langle t \rangle^{\frac{d(p-1)}{p}} \|G^{*(t,x)} \mathbf{f}(t)\|_{L^p} \lesssim_{\dot{c}, M_0} \widetilde{M} \sup_{s \in [0, \infty)} \left(\|\mathbf{f}(s)\|_{L^1} + \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}\|_{\dot{B}_{p,\infty}^a} \right).$$

This implies the result. ■

Lemma 3.11 *Let $t \geq 1$, and $p = 1, \infty$, we have estimates as follows,*

$$\langle t \rangle^{a + \frac{d(p-1)}{p}} \|G^{*(t,x)} \mathbf{f}(t)\|_{\dot{B}_{p,\infty}^a} \lesssim_{\dot{c}, M_1} \widetilde{M} \sup_{s \in [0, \infty)} \left(\|\mathbf{f}(s)\|_{L^1} + \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right).$$

Proof. When referring to $\|G^{*(t,x)} \mathbf{f}(t)\|_{\dot{B}_{p,\infty}^a}$, we consider $\|\delta_\alpha(G^{*(t,x)} \mathbf{f}(t))\|_{L^p}$ at first. Set

$$\begin{aligned} u_1(t, x) &= \int_0^{t/2} \int_{\mathbb{R}^d} G(t-s, y) \mathbf{f}(s, x-y) dy ds, \\ u_2(t, x) &= \int_{t/2}^t \int_{\mathbb{R}^d} G(t-s, y) \mathbf{f}(s, x-y) dy ds. \end{aligned}$$

Thus,

$$\|\delta_\alpha(G^{*(t,x)} \mathbf{f}(t))\|_{L^p} \leq \|\delta_\alpha u_1(t, x)\|_{L_x^p} + \|\delta_\alpha u_2(t, x)\|_{L_x^p}.$$

Firstly, we focus on the second term. Thanks to (3.20) and (3.25), we have for $|\alpha| > t/2 \geq \frac{1}{2}$

$$\begin{aligned} \frac{\|\delta_\alpha u_2(t, x)\|_{L_x^p}}{|\alpha|^a} &\leq \frac{2}{|\alpha|^a} \|u_2(t, x)\|_{L_x^p} = \frac{2}{|\alpha|^a} \left\| \int_{t/2}^t \int_{\mathbb{R}^d} G(t-s, y) \mathbf{f}(s, x-y) dy ds \right\|_{L_x^p} \\ &\lesssim \frac{2}{|\alpha|^a} \int_{t/2}^t \| |y|^a G(t-s, y) \|_{L_y^1} \sup_y \left(\frac{\|\delta_y \mathbf{f}(s)\|_{L^p}}{|y|^a} \right) ds \\ &\lesssim_{\dot{c}} \frac{1}{|\alpha|^a} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}\|_{\dot{B}_{p,\infty}^a} \right] \langle t \rangle^{-\frac{d(p-1)}{p} - a} \int_{t/2}^t \frac{\widetilde{M}}{\langle t-s \rangle^{1-a}} ds \\ &\lesssim_{\dot{c}} \widetilde{M} \frac{\langle t \rangle^a}{|\alpha|^a} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}\|_{\dot{B}_{p,\infty}^a} \right] \langle t \rangle^{-\frac{d(p-1)}{p} - a} \\ &\lesssim_{\dot{c}} \widetilde{M} \langle t \rangle^{-\frac{d(p-1)}{p} - a} \sup_{s \in [0, \infty)} \left(\langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}\|_{\dot{B}_{p,\infty}^a} \right). \end{aligned}$$

For the case $|\alpha| \leq t/2$, we know that

$$\begin{aligned} \frac{\|\delta_\alpha u_2(t, x)\|_{L_x^p}}{|\alpha|^a} &\lesssim \frac{1}{|\alpha|^a} \left\| \int_{t/2}^t \int_{\mathbb{R}^d} \mathbf{1}_{|t-s| \leq |\alpha|} \delta_\alpha G(t-s, y) \delta_y \mathbf{f}(s, x) dy ds \right\|_{L_x^p} \\ &\quad + \frac{1}{|\alpha|^a} \left\| \int_{t/2}^t \int_{\mathbb{R}^d} \mathbf{1}_{|t-s| > |\alpha|} \delta_\alpha G(t-s, y) \delta_y \mathbf{f}(s, x) dy ds \right\|_{L_x^p} =: \mathcal{J}_1 + \mathcal{J}_2. \end{aligned}$$

For the term \mathcal{J}_1 , one has

$$\begin{aligned} \mathcal{J}_1 &\leq \frac{1}{|\alpha|^a} \left\| \int_{t/2}^t \int_{\mathbb{R}^d} \mathbf{1}_{|t-s| \leq |\alpha|} G(t-s, y) \delta_y \mathbf{f}(s, x) dy ds \right\|_{L_x^p} \\ &\leq \frac{1}{|\alpha|^a} \int_{t/2}^t \mathbf{1}_{|t-s| \leq |\alpha|} \| |y|^a G(t-s, y) \|_{L^1} \sup_y \frac{\|\delta_y \mathbf{f}(s)\|_{L^p}}{|y|^a} ds \\ &\stackrel{(3.20)}{\lesssim_{\dot{c}}} \frac{1}{|\alpha|^a} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right] \langle t \rangle^{-\frac{d(p-1)}{p} - a} \int_0^{t/2} \| |y|^a G(s, y) \|_{L_y^1} \mathbf{1}_{|s| \leq |\alpha|} ds. \end{aligned}$$

Since we have

$$\int_0^{t/2} \| |y|^a G(s, y) \|_{L_y^1} \mathbf{1}_{|s| \leq |\alpha|} ds \stackrel{(3.20)}{\lesssim_{\dot{c}}} \widetilde{M} \int_0^{|\alpha|} \frac{1}{s^{1-a}} ds \leq \widetilde{M} |\alpha|^a,$$

then

$$\mathcal{J}_1 \lesssim_{\dot{c}, M_0} \widetilde{M} \langle t \rangle^{-\frac{d(p-1)}{p} - a} \sup_{s \in [0, \infty)} \left(\langle s \rangle^{\frac{d(p-1)}{p} + a} \|\mathbf{f}(s)\|_{\dot{B}_{p,\infty}^a} \right).$$

Now we deal with the term \mathcal{J}_2 ,

$$\begin{aligned}
\mathcal{J}_2 &\leq \frac{1}{|\alpha|^a} \int_{t/2}^t \mathbf{1}_{|t-s|>|\alpha|} \left(\int_{\mathbb{R}^d} |y|^a |\delta_\alpha G(t-s, y)| dy \right) \sup_y \frac{\|\delta_y \mathbf{f}(s, x)\|_{L_x^p}}{|y|^a} ds \\
&\leq \frac{1}{|\alpha|^a} \langle t \rangle^{-a - \frac{d(p-1)}{p}} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p, \infty}^a} \right] \int_{|\alpha|}^{t/2} \| |y|^a \delta_\alpha G(s, y) \|_{L_y^1} ds \\
&\stackrel{(3.21)}{\lesssim_{\dot{c}, M_0}} \frac{1}{|\alpha|^a} \langle t \rangle^{-a - \frac{d(p-1)}{p}} \left[\sup_{s \in [0, \infty)} \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p, \infty}^a} \right] \int_{|\alpha|}^{t/2} \frac{|\alpha|}{(1+s)^{2-a}} ds \\
&\lesssim_{\dot{c}, M_0} \langle t \rangle^{-a - \frac{d(p-1)}{p}} \sup_{s \in [0, \infty)} \left(\langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p, \infty}^a} \right).
\end{aligned}$$

Combining the estimation of \mathcal{J}_1 and \mathcal{J}_2 , we have following estimation

$$\frac{\|\delta_\alpha u_2(t, x)\|_{L_x^p}}{|\alpha|^a} \lesssim_{\dot{c}, M_0} \langle t \rangle^{-a - \frac{d(p-1)}{p}} \sup_{s \in [0, \infty)} \left(\langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p, \infty}^a} \right).$$

By the same method, we obtain similar conclusion about $u_1(t, x)$. We firstly consider the case $t \geq 1$,

$$\begin{aligned}
\|\delta_\alpha u_1(t, x)\|_{L_x^p} &= \left\| \int_0^{t/2} \int_{\mathbb{R}^d} \delta_\alpha G(t-s, x-y) \mathbf{f}(s, y) dy ds \right\|_{L_x^p} \\
&\leq \int_{t/2}^t \|\delta_\alpha G(s)\|_{L_x^p} ds \sup_{s \in [0, \infty)} \|\mathbf{f}(s)\|_{L^1}.
\end{aligned}$$

Note that for any $s \gtrsim 1$, we have

$$\begin{aligned}
\|\delta_\alpha G(s)\|_{L_x^p} &\leq \min\{\|G(s)\|_{L_x^p}, |\alpha| \|\nabla G(s)\|_{L_x^p}\} \\
&\stackrel{(3.19), (3.20)}{\lesssim_{\dot{c}, M_1}} \min \left\{ \frac{1}{\langle s \rangle^{1 + \frac{d(p-1)}{p}}}, \frac{|\alpha|}{\langle s \rangle^{2 + \frac{d(p-1)}{p}}} \right\} \lesssim_{\dot{c}, M_1} \frac{|\alpha|^a}{\langle s \rangle^{1+a + \frac{d(p-1)}{p}}}.
\end{aligned}$$

Then for $t \geq 1$,

$$\int_{t/2}^t \|\delta_\alpha G(s)\|_{L_x^p} ds \lesssim_{\dot{c}, M_1} \frac{|\alpha|^a}{\langle t \rangle^{1+a + \frac{d(p-1)}{p}}} \langle t \rangle \lesssim_{\dot{c}, M_1} |\alpha|^a \langle t \rangle^{-a - \frac{d(p-1)}{p}}.$$

Thus we obtain

$$\frac{\|\delta_\alpha u_1(t, x)\|_{L_x^p}}{|\alpha|^a} \lesssim_{\dot{c}, M_1} \langle t \rangle^{-a - \frac{d(p-1)}{p}} \sup_{s \in [0, \infty)} \|\mathbf{f}(s)\|_{L^1}.$$

By the arbitrariness of α and the fact $M_0 \leq M_1$, we have

$$\langle t \rangle^{a + \frac{d(p-1)}{p}} \|G *_{(t, x)} \mathbf{f}(t)\|_{\dot{B}_{p, \infty}^a} \lesssim_{\dot{c}, M_1} \sup_{s \in [0, \infty)} \left(\|\mathbf{f}(s)\|_{L^1} + \langle s \rangle^{a + \frac{d(p-1)}{p}} \|\mathbf{f}(s)\|_{\dot{B}_{p, \infty}^a} \right).$$

Until now we finish the proof of Theorem 3.9. ■

4 Estimates for Characteristics

In this section, we learn about the flow $(X_{s,t}(x, v), V_{s,t}(x, v))$ defined in (2.1), from which we have for any $0 \leq s \leq t < \infty$,

$$\begin{aligned}
X_{s,t}(x, v) &= x - v(t-s) + \int_s^t (\tau-s) E(\tau, X_{\tau,t}(x, v)) d\tau, \\
V_{s,t}(x, v) &= v - \int_s^t E(\tau, X_{\tau,t}(x, v)) d\tau.
\end{aligned} \tag{4.1}$$

Set

$$\begin{aligned}
Y_{s,t}(x, v) &= \int_s^t (\tau-s) E(\tau, x + \tau v + Y_{\tau,t}(x, v)) d\tau, \\
W_{s,t}(x, v) &= - \int_s^t E(\tau, x + \tau v + Y_{\tau,t}(x, v)) d\tau.
\end{aligned} \tag{4.2}$$

Hence,

$$X_{s,t}(x, v) = x - (t - s)v + Y_{s,t}(x - vt, v), \quad V_{s,t}(x, v) = v + W_{s,t}(x - vt, v). \quad (4.3)$$

We obtain the bounds on characteristics as follows.

Proposition 4.1 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, then there exists $\tilde{\varepsilon}_0 \in (0, 1)$ such that for any $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$ and $0 < \delta < a$, we have following estimates:*

$$\begin{aligned} & \sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \left(\|Y_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_x Y_{s,t}\|_{L_{x,v}^\infty} + \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \right) \\ & + \sup_{0 \leq s \leq t} \langle s \rangle^{d-3+a} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} + \delta \sup_{0 \leq s \leq t} \langle s \rangle^{d-3+\delta} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^{a-\delta}} \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}, \end{aligned} \quad (4.4)$$

and

$$\begin{aligned} & \sup_{0 \leq s \leq t} \langle s \rangle^{d-1+a} \left(\|W_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_x W_{s,t}\|_{L_{x,v}^\infty} + \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x W_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \right) \\ & + \sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \|\nabla_v W_{s,t}(x, v)\|_{L_{x,v}^\infty} + \sup_{0 \leq s \leq t} \langle s \rangle^{d-2} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v W_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \end{aligned} \quad (4.5)$$

Here we use the notation

$$\delta_\alpha^x Y_{s,t}(x, v) = Y_{s,t}(x, v) - Y_{s,t}(x - \alpha, v), \quad \delta_\alpha^v Y_{s,t}(x, v) = Y_{s,t}(x, v) - Y_{s,t}(x, v - \alpha).$$

Before we prove this Proposition, we state the following Lemma:

Lemma 4.2 *For $p = 1, \infty$ and $\kappa, \kappa_1 \in (0, 1)$, one has*

$$\begin{aligned} & \sum_{j=0}^1 \|\nabla^j (1 - \Delta)^{-1} \psi\|_{L^p} + \|\nabla (1 - \Delta)^{-1} \psi\|_{\dot{B}_{p,\infty}^\kappa} \leq c_0 \|\psi\|_{L^p}, \\ & \sum_{j=1}^2 \|\nabla^j (1 - \Delta)^{-1} \psi\|_{L^p} + [2 + \kappa - (\kappa_1 + j)] \sum_{\substack{j=0,1,2 \\ \kappa_1 + j < 2 + \kappa}} \|\nabla^j (1 - \Delta)^{-1} \psi\|_{\dot{B}_{p,\infty}^{\kappa_1} \cap \dot{F}_{\infty,\infty}^\kappa} \leq c_0 \|\psi\|_{\dot{B}_{p,\infty}^\kappa}, \\ & \|\nabla^2 (1 - \Delta)^{-1} \psi\|_{\dot{B}_{\infty,\infty}^\kappa} \leq c_0 \|\psi\|_{\dot{B}_{\infty,\infty}^\kappa}, \end{aligned}$$

for some $c_0 = c_0(\kappa)$.

Proof of Theorem 4.1. Note that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0$ implies $\|U\|_a \leq \varepsilon_0^{\frac{2}{3}}$, then by Assumption 3, we know that

$$\begin{aligned} \|A(U)\|_{\dot{B}_{p,\infty}^a} & \leq \max_{|r| \leq \varepsilon_0^{\frac{2}{3}}} |A'(r)| \|U\|_{\dot{B}_{p,\infty}^a} \leq C_A \varepsilon_0^{\frac{2}{3}} \|U\|_{\dot{B}_{p,\infty}^a}, \\ \|A(U)\|_{L^p} & \leq C_A \|U^2\|_{L^p} \leq C_A \varepsilon_0^{\frac{2}{3}} \|U\|_{L^p}. \end{aligned} \quad (4.6)$$

Thus taking $\varepsilon_0 \leq C_A^{-3}$, we have

$$\|\rho + A(U)\|_a \leq \|\rho\|_a + C_A \varepsilon_0^{\frac{2}{3}} \|U\|_a \leq \|\rho\|_a + \varepsilon_0^{\frac{1}{3}} \|U\|_a = \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \quad (4.7)$$

Thanks to Lemma 4.2 we have

$$\|E(\tau)\|_{L^\infty} + \|\nabla_x E(\tau)\|_{L^\infty} + \|\nabla_x E(\tau)\|_{\dot{B}_{\infty,\infty}^a} \leq c_0(a) \|\rho + A(U)\|_{\dot{B}_{\infty,\infty}^a} \leq c_0(a) \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d+a}}, \quad (4.8)$$

and there exists a constant $c_1 > 0$ such that

$$\int_s^t \frac{1}{\langle \tau \rangle^{d-i+a}} d\tau \leq \frac{c_1}{\langle s \rangle^{d-i+a-1}}, \quad i = 0, 1, 2, \quad (4.9)$$

where c_1 only depend on d .

(1) By (4.8) one has

$$\begin{aligned} \|Y_{s,t}\|_{L_{x,v}^\infty} &\leq \int_s^t (\tau-s) \|E(\tau, X_{\tau,t}(x,v))\|_{L_{x,v}^\infty} d\tau \leq \int_s^t \frac{c_0 \|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-1+a}} d\tau \leq c_0 c_1 \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}}, \\ \|\nabla_x Y_{s,t}\|_{L_{x,v}^\infty} &\leq (1 + \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}) \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{L_{x,v}^\infty} d\tau \\ &\leq c_0 \|(\rho, U)\|_{S_a^{\varepsilon_0}} (1 + \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}) \int_s^t \frac{1}{\langle \tau \rangle^{d-1+a}} d\tau \\ &\leq \frac{c_0 c_1 \|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} (1 + \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}). \end{aligned}$$

Then, the right hand side can be absorbed by the left provided that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \frac{1}{400c_0c_1d^2}$.
By the same method, we have

$$\begin{aligned} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} &\leq \int_s^t |\tau-s| \|\nabla_x E(\tau)\|_{L^\infty} \left(|\tau| + \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right) d\tau \\ &\leq c_0 c_1 \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\frac{1}{\langle s \rangle^{d-3+a}} + \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-3+a} \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right). \end{aligned}$$

Hence,

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \left(\|Y_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_x Y_{s,t}\|_{L_{x,v}^\infty} \right) + \sup_{0 \leq s \leq t} \langle s \rangle^{d-3+a} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} \leq 2c_0 c_1 \|(\rho, U)\|_{S_a^{\varepsilon_0}}, \quad (4.10)$$

provided $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \frac{1}{400c_0c_1d^2}$.

(2) By (4.8), (4.10), one obtains

$$\begin{aligned} \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} &\leq \sup_\alpha \frac{1}{|\alpha|^a} \left[\int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{L^\infty} \|\delta_\alpha^x \nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty} d\tau \right. \\ &\quad \left. + \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{\dot{B}_{\infty,\infty}^a} \left(|\alpha| + \|\delta_\alpha^x Y_{\tau,t}\|_{L_{x,v}^\infty} \right)^a \left(1 + \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty} \right) d\tau \right] \\ &\leq \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{L^\infty} d\tau \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \\ &\quad + \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{\dot{B}_{\infty,\infty}^a} \left(1 + \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty} \right)^{1+a} d\tau \\ &\leq c_0 c_1 \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} + c_0 c_1 \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}}. \end{aligned}$$

Similarly,

$$\begin{aligned} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^{a-\delta}} &\leq \sup_\alpha \frac{1}{|\alpha|^{a-\delta}} \left(\int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{L^\infty} \|\delta_\alpha^v \nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} d\tau \right. \\ &\quad \left. + \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{\dot{B}_{\infty,\infty}^{a-\delta}} \left(|\alpha| |\tau| + \|\delta_\alpha^v Y_{\tau,t}\|_{L_{x,v}^\infty} \right)^{a-\delta} \left(|\tau| + \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right) d\tau \right) \\ &\leq c_0 c_1 \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d+a-2}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-3+\delta} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^{a-\delta}} + c_0 c_1 \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\delta \langle s \rangle^{d-3+\delta}}. \end{aligned} \quad (4.11)$$

These implies

$$\sup_{0 \leq s \leq t} \left(\langle s \rangle^{d-2+a} \sup_\alpha \frac{\|\delta_\alpha^x \nabla_x Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} + \delta \langle s \rangle^{d-3+\delta} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^{a-\delta}} \right) \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \quad (4.12)$$

provided $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \frac{1}{400c_0c_1d^2}$.

Combining this with (4.10), one gets (4.4). By the same argument, we also obtain (4.5). Thus we choose $\tilde{\varepsilon}_0 = \min\{C_A^{-3}, \frac{1}{400c_0c_1d^2}\}$ to finish the proof. \blacksquare

Remark 4.3 For $d \geq 4$, using the same method in (4.11), we obtain

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-3} \sup_\alpha \frac{\|\delta_\alpha^v \nabla_v Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \quad (4.13)$$

Proposition 4.4 Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, $\tilde{\varepsilon}_0$ be as in Proposition 4.1. Assume that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$ and $\|(\rho, U)\|_{\gamma, b} \leq \mathbf{c}$ for some $\gamma > 0$ and $b \in (0, 1)$. Then, for $j = 1, \dots, \gamma$ and $\delta \in (0, b)$, we have

$$\begin{aligned} & \sup_{0 \leq s \leq t} \langle s \rangle^{d-2+b} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} + \langle s \rangle^{d-1+b} \|\nabla_v W_{s,t}\|_{L_{x,v}^\infty} \lesssim_{\mathbf{c}, b} 1, \\ & \sum_{j=1}^{\gamma} \sup_{0 \leq s \leq t} \langle s \rangle^{d-2} \sup_{\alpha} \frac{\|\delta_\alpha^v \nabla_v^j Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} + \langle s \rangle^{d-1} \sup_{\alpha} \frac{\|\delta_\alpha^v \nabla_v^j W_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} \lesssim_{\mathbf{c}, b} 1, \\ & \sup_{0 \leq s \leq t} \delta \langle s \rangle^{d-3+\delta} \sup_{\alpha} \frac{\|\delta_\alpha^v \nabla_v^{\gamma+1} Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^{b-\delta}} + \langle s \rangle^{d-2} \sup_{\alpha} \frac{a \delta_\alpha^v \nabla_v^{\gamma+1} W_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} \lesssim_{\mathbf{c}, b} 1, \\ & \sum_{k=1}^{\gamma+1} \sup_{0 \leq s \leq t} \langle s \rangle^{d-1+b} \sup_{\alpha} \frac{\|\delta_\alpha^x \nabla_x^k Y_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} + \langle s \rangle^{d+b} \sup_{\alpha} \frac{\|\delta_\alpha^x \nabla_x^k W_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} \lesssim_{\mathbf{c}, b} 1, \end{aligned}$$

provided that

$$\sum_{j=2}^{\gamma+1} \sup_{|r| \leq 1} |A^{(j)}(r)| \leq C_{A, \gamma}. \quad (4.14)$$

Proof. Note that for $0 \leq j \leq \gamma$ and $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0$, we have

$$\begin{aligned} \|\nabla^j(A(U))\|_{\dot{B}_{p,\infty}^a} & \leq \sum_{k=1}^j |A^{(k)}(U)|_{L^\infty} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = j}} \left\| \prod_{i=1}^k \nabla^{m_i} U \right\|_{\dot{B}_{p,\infty}^a} + \sum_{k=1}^j |A^{(k)}(U)|_{\dot{B}_{p,\infty}^a} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = j}} \left\| \prod_{i=1}^k \nabla^{m_i} U \right\|_{L^\infty} \\ & \leq C_{A, \gamma} \left(\sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = j}} \left\| \prod_{i=1}^k \nabla^{m_i} U \right\|_{\dot{B}_{p,\infty}^a} + \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = j}} \left\| \prod_{i=1}^k \nabla^{m_i} U \right\|_{L^\infty} \right) \lesssim_{\mathbf{c}, C_{A, \gamma}} 1. \end{aligned}$$

Combining this with Lemma 4.2, one has there exists a constant $c_3 = c_3(d, b, b', C_{A, \gamma}, \mathbf{c})$ such that

$$\sup_{\tau \geq 0} \sum_{j=0}^{\gamma+1} \langle \tau \rangle^{d+\min\{j+1, \gamma+b\}} \|\nabla_x^j E(\tau)\|_{L^\infty} + \langle \tau \rangle^{d+\min\{j+b'+1, \gamma+b\}} \|\nabla_x^j E(\tau)\|_{\dot{B}_{\infty,\infty}^{b'}} \lesssim_{b, b'} \|\rho + A(U)\|_{\gamma, b} \leq c_3, \quad (4.15)$$

for any $b' \in (0, b]$.

(1) Let c_1 be in (4.9), as the proof of (4.10), we have

$$\begin{aligned} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} & \leq \int_s^t |\tau - s| \|\nabla_x E(\tau)\|_{L^\infty} |\tau| d\tau + \int_s^t |\tau - s| \|\nabla_x E(\tau)\|_{L^\infty} \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} d\tau \\ & \leq c_3 \int_s^t \frac{|\tau|^2}{\langle \tau \rangle^{d+b+1}} d\tau + c_0 \left(\sup_{s \leq \tau \leq t} \langle \tau \rangle^{d-2+b} \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right) \int_s^t \frac{|\tau| \|\rho, U\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-2+b} \langle \tau \rangle^{d+a}} d\tau \\ & \leq c_3 c_1 \frac{1}{\langle s \rangle^{d+b-2}} + c_0 c_1 \|\rho, U\|_{S_a^{\varepsilon_0}} \left(\sup_{s \leq \tau \leq t} \langle \tau \rangle^{d-2+b} \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right) \frac{1}{\langle s \rangle^{d+b-2}}. \end{aligned} \quad (4.16)$$

Since $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \frac{1}{400c_1 d^2}$, then we have

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-2+b} \|\nabla_v Y_{s,t}\|_{L_{x,v}^\infty} \leq 2c_3 c_1 \mathbf{c} \lesssim_{\mathbf{c}, b, C_{A, \gamma}} 1.$$

Similarly, we obtain

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-1+b} \|\nabla_v W_{s,t}\|_{L_{x,v}^\infty} \lesssim_{\mathbf{c}, b, C_{A, \gamma}} 1.$$

(2) Using the method in (4.16) inductively, we know that

$$\sum_{j=0}^{\gamma+1} \|\nabla_v^j(Y_{s,t}, W_{s,t})\|_{L_{x,v}^\infty} + \sum_{j=0}^{\gamma} \sup_{\alpha} \frac{\|\delta_\alpha^v \nabla_v^j(Y_{s,t}, W_{s,t})\|_{L_{x,v}^\infty}}{|\alpha|^b} \lesssim_{\mathbf{c}, b, C_{A, \gamma}} 1, \quad (4.17)$$

for any $0 \leq s \leq t < \infty$.

By (4.17), one has

$$\begin{aligned}
\sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}} &\leq \int_s^t (\tau-s) \|\nabla E(\tau)\|_{L^{\infty}} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}} d\tau \\
&\quad + \int_s^t (\tau-s) \|\nabla E(\tau)\|_{B_{\infty,\infty}^{b-\delta}} \left(|\tau| + \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^{\infty}} \right)^{b-\delta} \|\nabla_v^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}} d\tau \\
&\quad + C(\mathbf{c}) \sum_{j=1}^k \int_s^t (\tau-s) \|\nabla^{j+1} E(\tau)\|_{B_{\infty,\infty}^{b-\delta}} (|\tau|+1)^{j+1+b-\delta} d\tau \\
&\quad + C(\mathbf{c}) \sum_{j=1}^k \int_s^t (\tau-s) \|\nabla^{j+1} E(\tau)\|_{L^{\infty}} (|\tau|+1)^{j+1} d\tau \\
&\stackrel{(4.8),(4.15)}{\leq} c_0 \int_s^t \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-1+a}} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}} d\tau + \frac{C(\mathbf{c}, b, C_{A,\gamma})}{\langle s \rangle^{d-2+\delta}} + \frac{C(\mathbf{c}, b, C_{A,\gamma}) \mathbf{1}_{k=\gamma}}{\delta \langle s \rangle^{d-3+\delta}}.
\end{aligned}$$

One derives that

$$\omega(s) \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}} \leq \frac{1}{4} B(t) + C(\mathbf{c}, b, C_{A,\gamma}),$$

where

$$B(t) = \sup_{0 \leq \tau \leq t} \omega(\tau) \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}}, \quad \omega(s) = \begin{cases} \langle s \rangle^{d-2} & \text{if } k \leq \gamma - 1, \\ \delta \langle s \rangle^{d-3+\delta} & \text{if } k = \gamma. \end{cases}$$

This implies

$$B(t) \lesssim_{\mathbf{c}, b, C_{A,\gamma}} 1.$$

In particular,

$$\sum_{k=0}^{\gamma-1} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} + \delta \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-3+\delta} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{\gamma+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^{b-\delta}} \lesssim_{\mathbf{c}, b, C_{A,\gamma}} 1. \quad (4.18)$$

Similarly, one also has,

$$\begin{aligned}
\sup_{\alpha} \frac{\|\delta_{\alpha}^x \nabla_x^{k+1} Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} &\leq \int_s^t (\tau-s) \|\nabla E(\tau)\|_{L^{\infty}} \sup_{\alpha} \frac{\|\delta_{\alpha}^x \nabla_x^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} d\tau \\
&\quad + \int_s^t (\tau-s) \|\nabla E(\tau)\|_{B_{\infty,\infty}^a} \|\nabla_x^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}} d\tau \\
&\quad + C(\mathbf{c}) \sum_{j=1}^k \int_s^t (\tau-s) \left(\|\nabla^{j+1} E(\tau)\|_{B_{\infty,\infty}^b} + \|\nabla^{j+1} E(\tau)\|_{L^{\infty}} \right) d\tau \\
&\stackrel{(4.8),(4.15)}{\leq} c_0 \int_s^t \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-1+a}} \sup_{\alpha} \frac{\|\delta_{\alpha}^x \nabla_x^{k+1} Y_{\tau,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} + \frac{C(\mathbf{c}, b, C_{A,\gamma})}{\langle s \rangle^{d-1+b}}.
\end{aligned}$$

This gives

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-1+b} \sup_{\alpha} \frac{\|\delta_{\alpha}^x \nabla_x^{k+1} Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} \lesssim_{\mathbf{c}, b, C_{A,\gamma}} 1. \quad (4.19)$$

By the same argument, we obtain

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-1-\mathbf{1}_{k=\gamma}} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{k+1} W_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^a} + \sup_{0 \leq s \leq t} \langle s \rangle^{d+a} \sup_{\alpha} \frac{\|\delta_{\alpha}^x \nabla_x^{k+1} W_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^a} \lesssim_{\mathbf{c}, b, C_{A,\gamma}} 1.$$

Therefore, the proof is complete. \blacksquare

Remark 4.5 For $d \geq 4$, we have

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-3} \sup_{\alpha} \frac{\|\delta_{\alpha}^v \nabla_v^{\gamma+1} Y_{s,t}\|_{L_{x,v}^{\infty}}}{|\alpha|^b} \lesssim_{\mathbf{c}, b, C_{A,\gamma}} 1.$$

Proposition 4.6 Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, $\tilde{\varepsilon}_0$ be as in Proposition 4.1, ρ_1, ρ_2 be such that $\|(\rho_1, U_1)\|_{S_a^{\varepsilon_0}}, \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. $(X_{s,t}^1, V_{s,t}^1), (X_{s,t}^2, V_{s,t}^2)$ are solutions to (2.1) associated to $E^1(s, x) = \nabla_x(1 - \Delta)^{-1}(\rho_1 + A(U_1))(x)$ and $E^2(s, x) = \nabla_x(1 - \Delta)^{-1}(\rho_2 + A(U_2))(x)$. Let $(Y_{s,t}^1, W_{s,t}^1), (Y_{s,t}^2, W_{s,t}^2)$ be such that

$$X_{s,t}^i(x, v) = x - (t-s)v + Y_{s,t}^i(x - vt, v), V_{s,t}^i(x, v) = v + W_{s,t}^i(x - vt, v), \quad i = 1, 2. \quad (4.20)$$

Then for $\mathbf{Y}_{s,t} = Y_{s,t}^1 - Y_{s,t}^2, \mathbf{W}_{s,t} = W_{s,t}^1 - W_{s,t}^2$, we have

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \left(\|\mathbf{Y}_{s,t}\|_{L_{x,v}^\infty} + \sup_\alpha \frac{\|\delta_\alpha^x \mathbf{Y}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \right) + \sup_{0 \leq s \leq t} \langle s \rangle^{d-2} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{Y}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}, \quad (4.21)$$

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-1+a} \left(\|\mathbf{W}_{s,t}\|_{L_{x,v}^\infty} + \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{W}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \right) + \sup_{0 \leq s \leq t} \langle s \rangle^{d-1} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{W}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (4.22)$$

Proof. Let c_0, c_1 be the constant mentioned in the proof of Theorem 4.1, then we have

$$Y_{s,t}^k(x, v) = \int_s^t (\tau - s) E^k(\tau, x + \tau v + Y_{\tau,t}^k(x, v)) d\tau, \\ W_{s,t}^k(x, v) = - \int_s^t E^k(\tau, x + \tau v + Y_{\tau,t}^k(x, v)) d\tau, \quad k = 1, 2.$$

From $\|(\rho_k, U_k)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0$, one has $\|U_k\|_a \leq \varepsilon_0^{\frac{2}{3}}$, for $k = 1, 2$. Combining with Assumption 3, we obtain that

$$\begin{aligned} \|A(U_1) - A(U_2)\|_{\dot{B}_{\infty,\infty}^a} &= \left\| (U_1 - U_2) \int_0^1 A'(\varpi U_1 + (1 - \varpi)U_2) d\varpi \right\|_{\dot{B}_{\infty,\infty}^a} \\ &\leq \|U_1 - U_2\|_{L^\infty} \left\| \int_0^1 A'(\varpi U_1 + (1 - \varpi)U_2) d\varpi \right\|_{\dot{B}_{\infty,\infty}^a} \\ &+ \|U_1 - U_2\|_{\dot{B}_{\infty,\infty}^a} \left\| \int_0^1 A'(\varpi U_1 + (1 - \varpi)U_2) d\varpi \right\|_{L^\infty} \\ &\leq \|U_1 - U_2\|_{L^\infty} \left(\max_{|r| \leq \varepsilon_0^{2/3}} |A'(r)| \right)^{1-a} \left(\varepsilon_0^{\frac{2}{3}} \max_{|r| \leq \varepsilon_0^{2/3}} |A''(r)| \right)^a + \|U_1 - U_2\|_{\dot{B}_{\infty,\infty}^a} \max_{|r| \leq \varepsilon_0^{2/3}} |A'(r)| \\ &\leq C_A \varepsilon_0^{\frac{2}{3}} \left(\|U_1 - U_2\|_{L^\infty} + \|U_1 - U_2\|_{\dot{B}_{\infty,\infty}^a} \right). \end{aligned}$$

Therefore,

$$\begin{aligned} \|(E^1 - E^2)(\tau)\|_{L^\infty} + \|\nabla_x(E^1 - E^2)(\tau)\|_{L^\infty} + \|\nabla_x(E^1 - E^2)(\tau)\|_{\dot{B}_{\infty,\infty}^a} &\leq c_0(a) \|\rho_1 - \rho_2 + A(U_1) - A(U_2)\|_{\dot{B}_{\infty,\infty}^a} \\ &\leq c_0(a) \left(\|\rho_1 - \rho_2\|_a + C_A \varepsilon_0^{\frac{2}{3}} \left(\|U_1 - U_2\|_{L^\infty} + \|U_1 - U_2\|_{\dot{B}_{\infty,\infty}^a} \right) \right) \leq \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d+a}}. \end{aligned} \quad (4.23)$$

These imply

$$\begin{aligned} \mathbf{Y}_{s,t}(x, v) &= \int_s^t (\tau - s) (E^1 - E^2)(\tau, x + \tau v + Y_{\tau,t}^1(x, v)) d\tau \\ &+ \int_s^t (\tau - s) (E^2(\tau, x + \tau v + Y_{\tau,t}^1(x, v)) - E^2(\tau, x + \tau v + Y_{\tau,t}^2(x, v))) d\tau. \end{aligned}$$

Hence we know that

$$\begin{aligned} \|\mathbf{Y}_{s,t}\|_{L_{x,v}^\infty} &\leq \int_s^t (\tau - s) \|(E^1 - E^2)(\tau)\|_{L^\infty} d\tau + \int_s^t (\tau - s) \|\nabla E^2(\tau)\|_{L^\infty} \|\mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty} d\tau \\ &\leq c_0 \int_s^t \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-1+a}} d\tau + c_0 \int_s^t \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle \tau \rangle^{d-1+a}} \|\mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty} d\tau \\ &\leq c_0 c_1 \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} + c_0 c_1 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{2d-4+2a}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \|\mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}. \end{aligned}$$

Since $\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \frac{1}{400_0 c_1 d^2}$, we can absorb the last term by the left handside to obtain

$$\sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \|\mathbf{Y}_{s,t}\|_{L_{x,v}^\infty} \leq 2c_0 c_1 \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}.$$

For the next term, we have

$$\begin{aligned} \sup_\alpha \frac{\|\delta_\alpha^x \mathbf{Y}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} &\leq \int_s^t (\tau - s) \|(E^1 - E^2)(\tau)\|_{B_{\infty,\infty}^a} d\tau + \int_s^t (\tau - s) \|\nabla E^2(\tau)\|_{B_{\infty,\infty}^a} \|\mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty} d\tau \\ &\quad + \int_s^t (\tau - s) \|\nabla E^2(\tau)\|_{L^\infty} \sup_\alpha \frac{\|\delta_\alpha^x \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} d\tau \\ &\leq c_0 c_1 \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} + 2c_0^2 c_1^2 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} \\ &\quad + c_0 c_1 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{2(d-2+a)}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \sup_\alpha \frac{\|\delta_\alpha^x \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \\ &\leq c_0 c_1 \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} + c_0 c_1 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{2(d-2+a)}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2+a} \sup_\alpha \frac{\|\delta_\alpha^x \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a}, \end{aligned}$$

and

$$\begin{aligned} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{Y}_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} &\leq \int_s^t (\tau - s) \|(E^1 - E^2)(\tau)\|_{B_{\infty,\infty}^a} \langle \tau \rangle^a d\tau + \int_s^t (\tau - s) \|\nabla E^2(\tau)\|_{B_{\infty,\infty}^a} \|\mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty} \langle \tau \rangle^a d\tau \\ &\quad + \int_s^t (\tau - s) \|\nabla E^2(\tau)\|_{L^\infty} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} d\tau \\ &\leq c_0 c_1 \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2}} + 2c_0^2 c_1^2 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2}} \\ &\quad + c_0 c_1 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{2(d-2)+a}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a} \\ &\leq c_0 c_1 \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2}} + c_0 c_1 \frac{\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{2(d-2)+a}} \sup_{0 \leq \tau \leq t} \langle \tau \rangle^{d-2} \sup_\alpha \frac{\|\delta_\alpha^v \mathbf{Y}_{\tau,t}\|_{L_{x,v}^\infty}}{|\alpha|^a}. \end{aligned}$$

As above, we can get the estimations of $\mathbf{W}_{s,t}$. Therefore, we get (4.21) provided $\|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \frac{1}{400_0 c_1 d^2}$. Similarly, we also get (4.22). The proof is complete. \blacksquare

Proposition 4.7 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, $\tilde{\varepsilon}_0$ be as in Proposition 4.1, (ρ, U) be such that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. Then, for any $0 \leq s \leq t < \infty$, we have a C^1 map $(x, v) \mapsto \Psi_{s,t}(x, v)$, which satisfies for all $x, v \in \mathbb{R}^d$:*

$$X_{s,t}(x, \Psi_{s,t}(x, v)) = x - (t - s)v, \quad (4.24)$$

and

$$\langle s \rangle^{d-1+a} (|\Psi_{s,t}(x, v) - v| + |\nabla_x \Psi_{s,t}(x, v)|) + \langle s \rangle^{d-2+a} |\nabla_v (\Psi_{s,t}(x, v) - v)| \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \quad (4.25)$$

Proof. Define

$$\Phi_{s,t}(x, v) = -\frac{1}{t-s} \int_s^t (\tau - s) E(\tau, x - (t - \tau)v + Y_{\tau,t}(x - vt, v)) d\tau.$$

Hence, we know from (4.1) that

$$X_{s,t}(x, v) = x - v(t - s) + Y_{s,t}(x - vt, v) = x - (t - s)(v + \Phi_{s,t}(x, v)). \quad (4.26)$$

For the first estimate, since we know from (4.10) that $\|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}$ is bounded, according to

$$\|\Phi_{s,t}\|_{L_{x,v}^\infty} \lesssim \int_s^t \|E(\tau)\|_{L_x^\infty} d\tau \lesssim_a \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d+a-1}} \quad (4.27)$$

and

$$\|\nabla_x \Phi_{s,t}\|_{L_{x,v}^\infty} \leq \frac{1}{t-s} \int_s^t (\tau - s) \|\nabla_x E(\tau)\|_{L_x^\infty} (1 + \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty}) d\tau \lesssim_a \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d+a-1}}, \quad (4.28)$$

we have

$$\sup_{0 \leq s, t \leq \infty} \langle s \rangle^{d+a-1} \left(\|\Phi_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_x \Phi_{s,t}\|_{L_{x,v}^\infty} \right) \lesssim_a \|(\rho, U)\|_{S_a^{\varepsilon_0}}. \quad (4.29)$$

Now one has

$$\begin{aligned} \|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} &\leq \frac{1}{t-s} \int_s^t (\tau-s) \|\nabla_x E(\tau)\|_{L^\infty} \left(t-\tau + t \|\nabla_x Y_{\tau,t}\|_{L_{x,v}^\infty} + \|\nabla_v Y_{\tau,t}\|_{L_{x,v}^\infty} \right) d\tau \\ &\leq \frac{c_0 \|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t-s} \int_s^t \frac{\tau-s}{\langle \tau \rangle^{d+a}} \left((t-\tau) + (t-\tau) + \tau + 1 \right) d\tau \\ &\leq 4c_0 \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_s^t \frac{1}{\langle \tau \rangle^{d+a-1}} d\tau. \end{aligned}$$

Let c_0, c_1 be in (4.8) and (4.9) by (4.4) and (4.8), one obtains that

$$\|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} \leq \frac{4c_0 c_1 \|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} \leq \frac{1}{100d^2},$$

provided $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \frac{1}{400c_0 c_1 d^2}$, and it is clear that under the assumption $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \frac{1}{400c_0 c_1 d^2}$ the map $(x, v) \mapsto (x, v + \Phi_{s,t}(x, v))$ is a C^1 diffeomorphism. Thus, there exists a map $(x, v) \mapsto \Psi_{s,t}(x, v)$ satisfying (4.24). Moreover, combining (4.24) and (4.26), we have $\Psi_{s,t}(x, v) + \Phi_{s,t}(x, \Psi_{s,t}(x, v)) = v$. Then,

$$\begin{aligned} |\Psi_{s,t}(x, v) - v| &\leq \|\Phi_{s,t}\|_{L_{x,v}^\infty} \lesssim_a \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d+a-1}}, \\ |\nabla_x \Psi_{s,t}(x, v)| &\leq \|\nabla_x \Phi_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} |\nabla_x \Psi_{s,t}(x, v)|, \\ |\nabla_v (\Psi_{s,t}(x, v) - v)| &\leq \|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} + \|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} (|\nabla_v (\Psi_{s,t}(x, v) - v)| + 1). \end{aligned}$$

Then, the right hand side can be absorbed by the left,

$$\begin{aligned} \|\nabla_x \Psi_{s,t}(x, v)\|_{L_{x,v}^\infty} &\leq 2\|\nabla_x \Phi_{s,t}\|_{L_{x,v}^\infty} \lesssim_a \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d+a-1}}, \\ |\nabla_v (\Psi_{s,t}(x, v) - v)| &\leq 3\|\nabla_v \Phi_{s,t}\|_{L_{x,v}^\infty} \lesssim_a \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}}. \end{aligned}$$

These follow (4.25). ■

5 Contribution of the initial data

For $h = h(x, v) : \mathbb{R}_x^d \times \mathbb{R}_v^d \rightarrow \mathbb{R}$, define

$$\mathcal{I}_h(\rho, U) = \int_{\mathbb{R}^d} h(X_{0,t}(x, v), V_{0,t}(x, v)) dv. \quad (5.1)$$

Proposition 5.1 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, $\tilde{\varepsilon}_0$ be as in Proposition 4.1 (ρ, U) such that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. Then there holds*

$$\|\mathcal{I}_h(\rho, U)\|_\sigma \lesssim_a \sum_{p=1, \infty} \|\mathcal{D}^\sigma(h)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p}, \quad (5.2)$$

for any $\sigma \in [a, 1)$. Moreover, for $0 < \delta_0 < 2 - \sqrt{3}$, if $\|(\rho, U)\|_{l, 1-\delta_0} \leq \mathbf{c}$ for some $l \geq 0$, then for any $b \in (0, 1 - \delta_0)$, we have

$$\|\mathcal{I}_h(\rho, U)\|_{l+1, b} \lesssim_{\mathbf{c}, a, b, C_{A, l}} \frac{1}{1 - \delta_0 - b} \sum_{i=1}^{l+1} \|\mathcal{D}^b(\nabla_{x,v}^i h)\|_{L_x^1 L_v^b \cap L_v^1 L_x^b}, \quad (5.3)$$

provided that $\sum_{j=2}^{l+1} \sup_{|r| \leq 1} |A^{(j)}(r)| \leq C_{A, l}$.

Remark 5.2 *By (4.13), for $d \geq 4$, we can obtain better result as follows*

$$\|\mathcal{I}_h(\rho, U)\|_{l+1, 1-\delta_0} \lesssim_{\mathbf{c}, a, b, C_{A, l}} \sum_{i=1}^{l+1} \|\mathcal{D}^{1-\delta_0}(\nabla_{x,v}^i h)\|_{L_x^1 L_v^b \cap L_v^1 L_x^b}. \quad (5.4)$$

Remark 5.3 We can not only use $\|h\|_{L_x^1 L_v^1 \cap L_x^1 L_v^\infty}$ to control $\|I_h(\rho, U)(t)\|_{L^\infty}$ for $0 \leq t \leq 1$. For example we take

$$h(x, v) = \frac{1}{|x|^\theta (|x|^2 + |v|^2 + 1)^{2d}}, \quad X_{0,t}(x, v) = x - tv, \quad V_{0,t}(x, v) = v, \quad U = 0,$$

for $0 < \theta < d$. And it is easy to check that for $t \in [0, 1]$

$$\|h\|_{L_x^1 L_v^1 \cap L_x^1 L_v^\infty} < \infty, \quad \|h\|_{L_v^1 L_x^\infty} = \infty, \quad \text{and} \quad \|I_h(\rho, U)(t)\|_{L^\infty} \sim \frac{1}{t^\theta}.$$

Proof. (1) Firstly, we deal with $\|\mathcal{I}_h(\rho, U)(t, x)\|_{L_x^p}$. Note that $(x, v) \mapsto (X_{0,t}(x, v), V_{0,t}(x, v))$ corresponding to the transport matrix

$$\mathfrak{A}(t, x, v) := \begin{bmatrix} I_d + \nabla_x Y_{0,t}(x - vt, v) & -tI_d + \nabla_v Y_{0,t}(x - vt, v) \\ \nabla_x W_{0,t}(x - vt, v) & I_d + \nabla_v W_{0,t}(x - vt, v) \end{bmatrix}.$$

Since we have

$$\|\nabla_{x,v} Y_{0,t}(x - vt, v)\|_{L_{x,v}^\infty} + \|\nabla_{x,v} W_{0,t}(x - vt, v)\|_{L_{x,v}^\infty} \leq 2c_0 c_1 \|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq 2c_0 c_1 \varepsilon_0 \leq \frac{1}{400d^2},$$

thus, $\|\det \mathfrak{A}^{-1}(t, x, v)\|_{L_{x,v}^\infty} \lesssim 1$. Then we directly obtain the inequality that

$$\begin{aligned} \|\mathcal{I}_h(\rho, U)(t, x)\|_{L_x^1} &= \left\| \int_{\mathbb{R}^d} h(x - vt + Y_{0,t}(x - vt, v), v + W_{0,t}(x - vt, v)) dv \right\|_{L_x^1} \\ &= \left\| \int_{\mathbb{R}^d} h(x, v) |\det \mathfrak{A}^{-1}(t, x, v)| dv \right\|_{L_x^1} \lesssim \|h\|_{L_v^1 L_x^1}. \end{aligned} \quad (5.5)$$

For the $\|\mathcal{I}_h(\rho, U)(t, x)\|_{L_x^\infty}$ with $t \leq 1$, changing variables as $\tilde{v} = v + W_{0,t}(x - vt, v)$, we have

$$\begin{aligned} \|\mathcal{I}_h(\rho, U)(t, x)\|_{L_x^\infty} &= \left\| \int_{\mathbb{R}^d} h(X_{0,t}(x, v), \tilde{v}) \det(I_d + \nabla_v W_{0,t}(x - vt, v))^{-1} d\tilde{v} \right\|_{L_x^\infty} \\ &\lesssim \left\| \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(X_{0,t}(x, v), \tilde{v}) d\tilde{v} \right\|_{L_x^\infty} \lesssim_a \int_{\mathbb{R}^d} \|h(\cdot, \tilde{v})\|_{L^\infty} d\tilde{v} = \|h\|_{L_v^1 L_x^\infty}. \end{aligned} \quad (5.6)$$

For $t \geq 1$, we change variable $w = x - vt$ to get more decay:

$$\mathcal{I}_h(\rho, U)(t, x) = \int_{\mathbb{R}^d} h\left(w + Y_{0,t}\left(w, \frac{x-w}{t}\right), \frac{x-w}{t} + W_{0,t}\left(w, \frac{x-w}{t}\right)\right) \frac{dw}{t^d},$$

Thus

$$\begin{aligned} \|\mathcal{I}_h(\rho, U)(t, x)\|_{L_x^\infty} &= \left\| \int_{\mathbb{R}^d} h(w, V_{0,t}(x, \Psi_{0,t}(x, \frac{x-w}{t}))) \det((\nabla_v \Psi_{0,t})(x, \frac{x-w}{t})) \frac{dw}{t^d} \right\|_{L_x^\infty} \\ &\lesssim \left\| \int_{\mathbb{R}^d} h\left(w, V_{0,t}\left(x, \Psi_{0,t}\left(x, \frac{x-w}{t}\right)\right)\right) \frac{dw}{t^d} \right\|_{L_{x,w}^\infty} \left\| \det\left((\nabla_v \Psi_{0,t})(x, \frac{x-w}{t})\right) \right\|_{L_{x,w}^\infty} \\ &\lesssim_a \frac{1}{t^d} \int_{\mathbb{R}^d} \|h(w, \cdot)\|_{L^\infty} dw = \frac{1}{t^d} \|h\|_{L_x^1 L_v^\infty}. \end{aligned} \quad (5.7)$$

Combine (5.5) with (5.6) and (5.7), we have

$$\langle t \rangle^{\frac{d(p-1)}{p}} \|\mathcal{I}_h(\rho, U)(t)\|_{L^p} \lesssim_a \|h\|_{L_x^1 L_v^p \cap L_x^1 L_v^\infty}, \quad p = 1, \infty. \quad (5.8)$$

(2) Now we deal with the term $\|\mathcal{I}_h(\rho, U)(t)\|_{B_{p,\infty}^\sigma}$. From Proposition 4.1, we have

$$\begin{aligned} \|\nabla_x X(x, v)\|_{L_{x,v}^\infty} + \|\nabla_x V(x, v)\|_{L_{x,v}^\infty} &\lesssim_a 1, \\ \sup_{\alpha/t} \frac{|\delta_{\alpha/t} Y_{0,t}(w, \cdot)(\frac{x-w}{t})|}{|\alpha/t|} + \sup_{\alpha/t} \frac{|\delta_{\alpha/t} W_{0,t}(w, \cdot)(\frac{x-w}{t})|}{|\alpha/t|} &\lesssim_a 1. \end{aligned}$$

Thus

$$\frac{|\delta_\alpha \mathcal{I}_h(\rho, U)(t, x)|}{|\alpha|^\sigma} \lesssim_a \sum_{z=x, x-\alpha} \int_{\mathbb{R}^d} \dot{D}^\sigma(h)(X_{0,t}(z, v), V_{0,t}(z, v)) dv,$$

and

$$\begin{aligned} \frac{|\delta_\alpha \mathcal{I}_h(\rho, U)(t, x)|}{|\alpha|^\sigma} &\lesssim_a \int_{\mathbb{R}^d} \dot{\mathcal{D}}_1^\sigma(h) \left(w + Y_{0,t} \left(w, \frac{x-w}{t} \right), \frac{x-w}{t} + W_{0,t} \left(w, \frac{x-w}{t} \right) \right) \frac{dw}{t^{d+\sigma}} \\ &\quad + \int_{\mathbb{R}^d} \dot{\mathcal{D}}_2^\sigma(h) \left(w + Y_{0,t} \left(w, \frac{x-\alpha-w}{t} \right), \frac{x-\alpha-w}{t} + W_{0,t} \left(w, \frac{x-\alpha-w}{t} \right) \right) \frac{dw}{t^{d+\sigma}} \\ &\lesssim_a \frac{1}{t^\sigma} \sum_{z=v, v-\frac{\alpha}{t}} \int_{\mathbb{R}^d} \dot{\mathcal{D}}^\sigma(h)(X_{0,t}(x, z), V_{0,t}(x, z)) dv. \end{aligned}$$

Thanks to (5.8) with $h = \dot{\mathcal{D}}^\sigma(h)$, we get

$$\langle t \rangle^{\sigma + \frac{d(p-1)}{p}} \|\mathcal{I}_h(\rho, U)(t)\|_{B_{p,\infty}^\sigma} \lesssim_a \|\dot{\mathcal{D}}^\sigma(h)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p}, \quad p = 1, \infty. \quad (5.9)$$

Until now we finish the proof of (5.2).

(3) Finally we consider the term $\|\mathcal{I}_h(\rho, U)\|_{l+1, b}$. Thanks to Proposition 4.4, one has

$$\sum_{k=1}^{l+1} (1 - \delta_0 - b) \sup_{0 \leq s \leq t} \sup_{\alpha} \frac{\|\delta_\alpha^x \nabla_x^k(Y, W)_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} + \sum_{k=1}^{l+1} \sup_{0 \leq s \leq t} \|\nabla_x^k(Y, W)_{s,t}\|_{L_{x,v}^\infty} \lesssim_{\mathbf{c}, b, C_{A,l}} 1.$$

Therefore

$$\frac{|\delta_\alpha \nabla^l \mathcal{I}_h(\rho, U)(t, x)|}{|\alpha|^b} \lesssim_{\mathbf{c}, b, C_{A,l}} \frac{1}{1 - \delta_0 - b} \sum_{j=1}^l \sum_{z=x, x-\alpha} \int_{\mathbb{R}^d} \mathcal{D}^b(\nabla_{x,v}^j h)(X_{0,t}(z, v), V_{0,t}(z, v)) dw.$$

Thus,

$$\|\nabla^l \mathcal{I}_h(\rho, U)(t)\|_{\dot{B}_{p,\infty}^b} \lesssim_{\mathbf{c}, a, b, C_{A,l}} \frac{1}{1 - \delta_0 - b} \sum_{j=1}^l \|\mathcal{D}^b(\nabla_{x,v}^j h)\|_{L_x^1 L_v^p}. \quad (5.10)$$

One the other hand, changing of variables as $w = x - vt$ and using the estimation

$$\sum_{k=1}^{l+1} (1 - \delta_0 - b) \sup_{0 \leq s \leq t} \sup_{\alpha} \frac{\|\delta_\alpha^x \nabla_v^k(Y, W)_{s,t}\|_{L_{x,v}^\infty}}{|\alpha|^b} + \sum_{k=1}^{l+1} \sup_{0 \leq s \leq t} \|\nabla_v^k(Y, W)_{s,t}\|_{L_{x,v}^\infty} \lesssim_{\mathbf{c}, b, C_{A,l}} 1,$$

we have

$$\begin{aligned} &\frac{|\delta_\alpha \nabla^l \mathcal{I}_h(\rho, U)(t, x)|}{|\alpha|^b} \\ &\lesssim_{\mathbf{c}, b} \frac{1}{1 - \delta_0 - b} \frac{1}{t^{d+l+b}} \left[\sum_{j=1}^l \int_{\mathbb{R}^d} \mathcal{D}_1^b(\nabla_{x,v}^j h) \left(w + Y_{0,t} \left(w, \frac{x-w}{t} \right), \frac{x-w}{t} + W_{0,t} \left(w, \frac{x-w}{t} \right) \right) dw \right. \\ &\quad \left. + \int_{\mathbb{R}^d} \mathcal{D}_2^b(\nabla_{x,v}^j h) \left(w + Y_{0,t} \left(w, \frac{x-\alpha-w}{t} \right), \frac{x-\alpha-w}{t} + W_{0,t} \left(w, \frac{x-\alpha-w}{t} \right) \right) dw \right] \\ &\lesssim_{\mathbf{c}, b} \frac{1}{(1 - \delta_0 - b)t^{l+b}} \sum_{j=1}^l \sum_{z=v, v-\alpha/t} \int_{\mathbb{R}^d} (\mathcal{D}^b(\nabla_{x,v}^j h))(X_{0,t}(x, z), V_{0,t}(x, z)) dv. \end{aligned}$$

Thanks to (5.8), we get for $p = 1, \infty$

$$t^{l+b-\delta + \frac{d(p-1)}{p}} \|\nabla^l \mathcal{I}_h(\rho, U)(t)\|_{B_{p,\infty}^{b-\delta}} \lesssim_{\mathbf{c}, a, b, C_{A,l}} \frac{1}{1 - \delta_0 - b} \sum_{j=1}^l \|\mathcal{D}^{b'}(\nabla_{x,v}^j h)\|_{L_x^1 L_v^p}. \quad (5.11)$$

Therefore, we obtain (5.3). ■

Proposition 5.4 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$, $\tilde{\varepsilon}_0$ be as in Proposition 4.1. Suppose that $\|(\rho_1, U_1)\|_{S_a^{\varepsilon_0}}, \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. There holds,*

$$\|\mathcal{I}_h(\rho_1, U_1) - \mathcal{I}_h(\rho_2, U_2)\|_a \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \sum_{p=1, \infty} \|\mathcal{D}^a(\nabla_{x,v} h)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p}. \quad (5.12)$$

Proof. Let $X_{s,t}^1, X_{s,t}^2, Y_{s,t}^1, Y_{s,t}^2, V_{s,t}^1, V_{s,t}^2, \mathbf{Y}_{s,t}, \mathbf{W}_{s,t}$ be in Proposition 4.6. Then we have

$$\begin{aligned} X_{s,t}^1(x, v) - X_{s,t}^2(x, v) &= Y_{s,t}^1(x - vt, v) - Y_{s,t}^2(x - vt, v) = \mathbf{Y}_{s,t}(x - vt, v), \\ V_{s,t}^1(x, v) - V_{s,t}^2(x, v) &= W_{s,t}^1(x - vt, v) - W_{s,t}^2(x - vt, v) = \mathbf{W}_{s,t}(x - vt, v). \end{aligned}$$

Meanwhile, denote

$$\begin{aligned} X_{s,t}(x, v, \varpi) &= (1 - \varpi)X_{s,t}^1(x, v) + \varpi X_{s,t}^2(x, v), \quad V_{s,t}(x, v, \varpi) = (1 - \varpi)V_{s,t}^1(x, v) + \varpi V_{s,t}^2(x, v), \\ Y_{s,t}(x, v, \varpi) &= (1 - \varpi)Y_{s,t}^1(x, v) + \varpi Y_{s,t}^2(x, v), \quad W_{s,t}(x, v, \varpi) = (1 - \varpi)W_{s,t}^1(x, v) + \varpi W_{s,t}^2(x, v). \end{aligned} \quad (5.13)$$

(1) For the term $\|\mathcal{I}_h(\rho_1, U_1)(t) - \mathcal{I}_h(\rho_2, U_2)(t)\|_{L^p}$, we have

$$\begin{aligned} |\mathcal{I}_h(\rho_1, U_1)(t, x) - \mathcal{I}_h(\rho_2, U_2)(t, x)| &\leq \left(\|\mathbf{Y}_{0,t}\|_{L_{x,v}^\infty} + \|\mathbf{W}_{0,t}\|_{L_{x,v}^\infty} \right) \int_{\mathbb{R}^d} \int_0^1 |(\nabla_{x,v} h)(X_{0,t}(x, v, \varpi), V_{0,t}(x, v, \varpi))| d\varpi dv \\ &\stackrel{(4.22)(4.21)}{\lesssim_a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_{\mathbb{R}^d} \sup_{0 \leq \varpi \leq 1} |(\nabla_{x,v} h)(X_{0,t}(x, v, \varpi), V_{0,t}(x, v, \varpi))| dv. \end{aligned}$$

As the proof of (5.8), one obtains

$$\int_{\mathbb{R}^d} \sup_{0 \leq \varpi \leq 1} |(\nabla_{x,v} h)(X_{0,t}(x, v, \varpi), V_{0,t}(x, v, \varpi))| dv \lesssim_a \sum_{p=1, \infty} \|\nabla_{x,v} h\|_{L_x^1 L_v^p \cap L_v^1 L_x^p}.$$

Thus

$$\sum_{p=1, \infty} \langle t \rangle^{\frac{d(p-1)}{p}} \|\mathcal{I}_h(\rho_1)(t) - \mathcal{I}_h(\rho_2)(t)\|_{L^p} \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \sum_{p=1, \infty} \|\nabla_{x,v} h\|_{L_x^1 L_v^p \cap L_v^1 L_x^p}. \quad (5.14)$$

(2) Now we consider the term $\|\mathcal{I}_h(\rho_1, U_1)(t) - \mathcal{I}_h(\rho_2, U_2)(t)\|_{\dot{B}_{p,\infty}^a}$. We compute directly to get

$$\begin{aligned} &\frac{|\delta_\alpha (\mathcal{I}_h(\rho_1, U_1) - \mathcal{I}_h(\rho_2, U_2))(t, x)|}{|\alpha|^a} \\ &\leq \left(\sup_z \frac{\|\delta_z^x \mathbf{Y}_{0,t}\|_{L_{x,v}^\infty}}{|z|^a} + \sup_z \frac{\|\delta_z^x \mathbf{W}_{0,t}\|_{L_{x,v}^\infty}}{|z|^a} \right) \int_{\mathbb{R}^d} \int_0^1 |(\nabla_{x,v} h)(X_{0,t}(x, v, \varpi), V_{0,t}(x, v, \varpi))| d\varpi dv \\ &\quad + \left(\|\mathbf{Y}_{0,t}\|_{L_{x,v}^\infty} + \|\mathbf{W}_{0,t}\|_{L_{x,v}^\infty} \right) \int_{\mathbb{R}^d} \int_0^1 \sum_{z=x, x-\alpha} \dot{D}^a (\nabla_{x,v} h)(X_{0,t}(z, v, \varpi), V_{0,t}(z, v, \varpi)) d\varpi dv \\ &\stackrel{(4.22)(4.21)}{\lesssim_a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \sum_{z=x, x-\alpha} \int_{\mathbb{R}^d} \sup_{0 \leq \varpi \leq 1} \mathcal{D}^a (\nabla_{x,v} h)(X_{0,t}(z, v, \varpi), V_{0,t}(z, v, \varpi)) d\varpi dv. \end{aligned} \quad (5.15)$$

On the other hand, we first change of variables as $w = x - vt$, then taking δ_α to get the decay of t^{-a} , after this we change $v = \frac{x-w}{t}$ to obtain

$$\begin{aligned} &\frac{|\delta_\alpha (\mathcal{I}_h(\rho_1, U_1) - \mathcal{I}_h(\rho_2, U_2))(t, x)|}{|\alpha|^a} \\ &\leq \frac{1}{t^a} \left(\sup_z \frac{\|\delta_z^v \mathbf{Y}_{0,t}\|_{L_{x,v}^\infty}}{|z|^a} + \sup_z \frac{\|\delta_z^v \mathbf{W}_{0,t}\|_{L_{x,v}^\infty}}{|z|^a} \right) \int_{\mathbb{R}^d} \int_0^1 |(\nabla_{x,v} h)(X_{0,t}(x, v, \varpi), V_{0,t}(x, v, \varpi))| d\varpi dv \\ &\quad + \frac{1}{t^a} \left(\|\mathbf{Y}_{0,t}\|_{L_{x,v}^\infty} + \|\mathbf{W}_{0,t}\|_{L_{x,v}^\infty} \right) \int_{\mathbb{R}^d} \int_0^1 \sum_{z=x, x-\alpha} \dot{D}^a (\nabla_{x,v} h)(X_{0,t}(z, v, \varpi), V_{0,t}(z, v, \varpi)) d\varpi dv \\ &\stackrel{(4.22)(4.21)}{\lesssim_a} \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{t^a} \sum_{z=x, x-\alpha} \int_{\mathbb{R}^d} \sup_{0 \leq \varpi \leq 1} \mathcal{D}^a (\nabla_{x,v} h)(X_{0,t}(z, v, \varpi), V_{0,t}(z, v, \varpi)) d\varpi dv. \end{aligned} \quad (5.16)$$

Combining (5.15) and (5.16), we have

$$\begin{aligned} &\frac{|\delta_\alpha (\mathcal{I}_h(\rho_1, U_1) - \mathcal{I}_h(\rho_2, U_2))(t, x)|}{|\alpha|^a} \\ &\lesssim_a \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{\langle t \rangle^a} \sum_{z=x, x-\alpha} \int_{\mathbb{R}^d} \sup_{0 \leq \varpi \leq 1} \mathcal{D}^a (\nabla_{x,v} h)(X_{0,t}(z, v, \varpi), V_{0,t}(z, v, \varpi)) d\varpi dv. \end{aligned}$$

Therefore, applying the the same method in (5.8), we obtain

$$\sup_{t>0} \langle t \rangle^{a+\frac{d(p-1)}{p}} \|(\mathcal{I}_h(\rho_1, U_1) - \mathcal{I}_h(\rho_2, U_2))(t)\|_{\dot{B}_{p,\infty}^a} \lesssim_a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \sum_{p=1, \infty} \|\mathcal{D}^a (\nabla_{x,v} h)\|_{L_x^1 L_v^p}. \quad (5.17)$$

Now the proof is complete. \blacksquare

6 Contribution Of The Reaction Term

In this section, we focus on estimate of the reaction term:

$$\mathcal{R}(\rho, U)(t, x) = \int_0^t \int_{\mathbb{R}^d} E(s, x - (t-s)v) \cdot \nabla_v \mu(v) dv ds - \int_0^t \int_{\mathbb{R}^d} E(s, X_{s,t}(x, v)) \cdot \nabla_v \mu(V_{s,t}(x, v)) dv ds. \quad (6.1)$$

Denote that for any given $F : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\mathcal{T}[F, \eta](t, x) = -\mathcal{T}_{NL}[F, \eta](t, x) + \mathcal{T}_L[F, \eta](t, x),$$

with

$$\begin{aligned} \mathcal{T}_L[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, x - (t-s)v) \eta(v) dv ds, \\ \mathcal{T}_{NL}[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, X_{s,t}(x, v)) \eta(V_{s,t}(x, v)) dv ds. \end{aligned}$$

Hence, it is clear that

$$\mathcal{R}(\rho) = \sum_{i=1}^d \mathcal{T}[E_i, \partial_{x_i} \mu].$$

By changing variable, we reformuate as follows:

$$\begin{aligned} \mathcal{T}_L[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, w + \frac{s}{t}(x-w)) \eta\left(\frac{x-w}{t}\right) \frac{dw ds}{t^d}, \\ \mathcal{T}_{NL}[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F\left(s, Y_{s,t}(w, \frac{x-w}{t}) + w + \frac{s(x-w)}{t}\right) \eta\left(W_{s,t}(w, \frac{x-w}{t}) + \frac{x-w}{t}\right) \frac{dw ds}{t^d}. \end{aligned}$$

Then, we have

Lemma 6.1 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$ and $\tilde{\varepsilon}_0 > 0$ be as in Proposition 4.1, ρ be such that $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. $(X_{s,t}, V_{s,t})$ are solutions to (2.1) associated to $E(s, x) = \nabla_x(1 - \Delta)^{-1}(\rho(s, x))$. Let $Y_{s,t}, W_{s,t}$ be such that*

$$X_{s,t}(x, v) = x - (t-s)v + Y_{s,t}(x - vt, v), V_{s,t}(x, v) = v + W_{s,t}(x - vt, v). \quad (6.2)$$

Suppose η be such that

$$\sum_{j=0}^2 \langle v \rangle^N |\nabla^j \eta(v)| \leq \mathbf{c}_1 \quad (6.3)$$

for $N > d$. Then,
(1)

$$|\mathcal{T}[F, \eta](t, x)| \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| \frac{1}{\langle s \rangle^{d-2+a}} \frac{dv ds}{\langle v \rangle^N}. \quad (6.4)$$

In particular, for $p = 1, \infty$

$$\langle t \rangle^{\frac{d(p-1)}{p}} \|\mathcal{T}[F, \eta](t)\|_{L^p} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} + \sup_{s \in [0, t]} \|F(s)\|_{L^1} \right). \quad (6.5)$$

(2) For any $\sigma \in [a, 1)$

$$\sum_{p=1, \infty} \langle t \rangle^{\frac{d(p-1)}{p} + \sigma} \|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p, \infty}^{\sigma}} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^{\sigma} \sum_{p=1, \infty} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p, \infty}^{\sigma}} \right). \quad (6.6)$$

Remark 6.2 *To estimate $\|\mathcal{T}[F, \eta](t)\|_{L^p}$, we only need η satisfies $\sum_{j=0,1} \langle v \rangle^N |\nabla^j \eta(v)| \leq \mathbf{c}_1$. Moreover, if η satisfies $\sum_{j=0,1} (\langle v \rangle^N + \langle v - \frac{\alpha}{t} \rangle^N) |\nabla^j \eta(v)| \leq \mathbf{c}_1$, we can also prove (6.5) by the same method.*

Proof. (1) First of all, we change variable and obtain

$$\begin{aligned}
|\mathcal{T}[F, \eta](t, x)| &= \left| - \int_0^t \int_{\mathbb{R}^d} F(s, x - (t-s)v) \eta(V_{s,t}(x, \Psi_{s,t}(x, v))) |\det(\nabla_v \Psi_{s,t}(x, v))| dv ds \right. \\
&\quad \left. + \int_0^t \int_{\mathbb{R}^d} F(s, x - (t-s)v) \eta(v) dv ds \right| \\
&\leq \int_0^t \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| |\eta(V_{s,t}(x, \Psi_{s,t}(x, v))) - \eta(v)| dv ds \\
&\quad + \int_0^t \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| \left| |\det(\nabla_v \Psi_{s,t}(x, v))| - 1 \right| |\eta(V_{s,t}(x, \Psi_{s,t}(x, v)))| dv ds.
\end{aligned}$$

Hence, since we have

$$\begin{aligned}
|\eta(V_{s,t}(x, \Psi_{s,t}(x, v))) - \eta(v)| &\leq \int_0^1 |\nabla \eta(\varpi V_{s,t}(x, v) + (1-\varpi)v)| |V_{s,t}(x, v) - v| d\varpi \\
&\lesssim_{\mathbf{c}_1} \sup_{\varpi \in [0,1]} \frac{|V_{s,t}(x, v) - v|}{\langle \varpi W_{s,t}(x - vt, v) + v \rangle^N} \\
&\sim_{\mathbf{c}_1} \frac{|W_{s,t}(x - vt, v)|}{\langle v \rangle^N} \stackrel{(4.5), (4.25)}{\lesssim_{a, \mathbf{c}_1}} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-1+a} \langle v \rangle^N},
\end{aligned}$$

and $|\eta(V_{s,t}(x, \Psi_{s,t}(t, x)))| \lesssim_{\mathbf{c}_1} \langle W_{s,t}(x - t\Psi_{s,t}(t, x), \Psi_{s,t}(t, x)) + v \rangle^{-N} \lesssim_{\mathbf{c}_1} \langle v \rangle^{-N}$, where we use the fact that $\sup_{0 \leq s \leq t} \|W_{s,t}\|_{L_{x,v}^\infty} \leq 1$ in (4.5). Hence

$$\begin{aligned}
|\mathcal{T}[F, \eta](t, x)| &\stackrel{(4.25)}{\lesssim_{a, \mathbf{c}_1}} \int_0^t \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| \left(\frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-1+a}} + \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{\langle s \rangle^{d-2+a}} \right) \frac{dv ds}{\langle v \rangle^N} \\
&\lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| \frac{1}{\langle s \rangle^{d-2+a} \langle v \rangle^N} dv ds.
\end{aligned}$$

This gives (6.4). Moreover,

$$\|\mathcal{T}[F, \eta](t)\|_{L^p} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \left\| \int_{\mathbb{R}^d} |F(s, x - (t-s)v)| \frac{dv}{\langle v \rangle^N} \right\|_{L_x^p} \frac{ds}{\langle s \rangle^{d-2+a}}.$$

Then, we apply Lemma 8.1 with $\mathcal{H}_1 = F$ and $\mathcal{H}_2 = 0$ to obtain

$$\begin{aligned}
\|\mathcal{T}[F, \eta](t)\|_{L^p} &\lesssim_{a, \mathbf{c}_1} t^{-\frac{d(p-1)}{p}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^{t/2} \|F(s)\|_{L^1} \frac{ds}{\langle s \rangle^{d-2+a}} + \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_{t/2}^t \|F(s)\|_{L^p} \frac{ds}{\langle s \rangle^{d-2+a}} \\
&\lesssim_{a, \mathbf{c}_1} t^{-\frac{d(p-1)}{p}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \underbrace{\sup_{s \in [0, t]} \|F(s)\|_{L^1} \int_0^{t/2} \frac{ds}{\langle s \rangle^{d-2+a}}}_{\text{Thanks to (8.2)}} \\
&\quad + \underbrace{\|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} \right) \int_{t/2}^t \frac{ds}{\langle s \rangle^{d-2+a} \langle s \rangle^{\frac{d(p-1)}{p}}}}_{\text{Thanks to (8.1)}} \\
&\lesssim_{a, \mathbf{c}_1} \frac{1}{t^{\frac{d(p-1)}{p}}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|F(s)\|_{L^1} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} \right).
\end{aligned}$$

Combining this with

$$\|\mathcal{T}[F, \eta](t)\|_{L^p} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \|F(s)\|_{L^p} \frac{1}{\langle s \rangle^{d-2+a}} ds \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \sup_{s \in [0, t]} \|F(s)\|_{L^p},$$

to obtain (6.5).

(2) Now we estimate $\|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p, \infty}^\sigma}$.

2.1) Estimate $\|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p, \infty}^\sigma}$ for $0 \leq t < 1$.

We divide $\delta_\alpha \mathcal{T}[F, \eta](t, x)$ into three terms:

$$\delta_\alpha \mathcal{T}[F, \eta](t, x) = \overline{\mathcal{T}}_\alpha^1[F, \eta](t, x) + \overline{\mathcal{T}}_\alpha^2[F, \eta](t, x) + \overline{\mathcal{T}}_\alpha^3[F, \eta](t, x) + \overline{\mathcal{T}}_\alpha^4[F, \eta](t, x),$$

where

$$\begin{aligned}\overline{\mathcal{T}}_\alpha^1[F, \eta](t, x) &= - \int_0^t \int_{\mathbb{R}^d} \delta_\alpha F(s, \cdot)(X_{s,t}(x, v)) \eta(V_{s,t}(x, v)) dv ds + \int_0^t \int_{\mathbb{R}^d} \delta_\alpha F(s, \cdot)(x - (t-s)v) \eta(v) dv ds, \\ \overline{\mathcal{T}}_\alpha^2[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} [F(s, X_{s,t}(x - \alpha, v)) - F(s, X_{s,t}(x, v) - \alpha)] \eta(V_{s,t}(x, v)) dv ds, \\ \overline{\mathcal{T}}_\alpha^3[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, X_{s,t}(x - \alpha, v)) [\eta(V_{s,t}(x - \alpha, v)) - \eta(V_{s,t}(x, v))] dv ds.\end{aligned}$$

For the first term, using (6.3) and (6.4) we have

$$\begin{aligned}\|\overline{\mathcal{T}}_\alpha^1[F, \eta](t)\|_{L^p} &= \|\mathcal{T}[\delta_\alpha F, \eta](t)\|_{L^p} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|\delta_\alpha F(s)\|_{L^1} + \sup_{s \in [0, t]} \|\delta_\alpha F(s)\|_{L^p} \right) \\ &\lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|F(s)\|_{\dot{B}_{1, \infty}^\sigma} + \sup_{s \in [0, t]} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma} \right).\end{aligned}$$

Since

$$|X_{s,t}(x - \alpha, v) - (X_{s,t}(x - \alpha, v) - \alpha)| = |Y_{s,t}(x - tv, v) - Y_{s,t}(x - tv, v)| \leq |\alpha| \|\nabla_x Y_{s,t}\|_{L^\infty},$$

we can estimate $\overline{\mathcal{T}}_\alpha^2[F, \eta](t, x)$ as follows

$$\begin{aligned}\|\overline{\mathcal{T}}_\alpha^2[F, \eta](t, x)\|_{L_x^p} &\lesssim_{\mathbf{c}_1} |\alpha|^\sigma \int_0^t \int_{\mathbb{R}^d} \|\sup_z |z|^{-\sigma} |\delta_z F(s, \cdot)|\|_{L^p} \|\nabla_x Y_{s,t}\|_{L_{x,v}^\infty}^\sigma \frac{dv}{\langle v \rangle^N} ds \\ &\lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma} \right) \int_0^t \langle s \rangle^{-(d+a-2)\sigma} ds \lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \sup_{s \in [0, t]} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma},\end{aligned}$$

where in the last inequality, we used $(d+a-2)\sigma \geq (a+1)a > 1$.

Note that by (4.5), one has $\sup_{s,t} \|W_{s,t}\|_{L_{x,v}^\infty} \leq 1$. Then,

$$\langle \varpi V_{s,t}(x, v) + (1 - \varpi) V_{s,t}(x - \alpha, v) \rangle = \langle v + \varpi W_{s,t}(x - tv, v) + (1 - \varpi) W_{s,t}(x - \alpha - tv, v) \rangle \sim \langle v \rangle. \quad (6.7)$$

We can estimate $\overline{\mathcal{T}}_\alpha^3[F, \eta](t)$.

$$\begin{aligned}\|\overline{\mathcal{T}}_\alpha^3[F, \eta](t, x)\|_{L^p} &\leq \int_0^t \int_{\mathbb{R}^d} \int_0^1 \|F(s)\|_{L^p} |\nabla \eta(\cdot)| (\varpi V_{s,t}(x, v) + (1 - \varpi) V_{s,t}(x - \alpha, v)) |\delta_\alpha^x V_{s,t}(x, v)| dv ds d\varpi \\ &\lesssim_{\mathbf{c}_1} |\alpha|^\sigma \int_0^t \int_{\mathbb{R}^d} \|F(s)\|_{L^p} \frac{1}{\langle v \rangle^N} \sup_\alpha \frac{|\delta_\alpha^x V_{s,t}(x, v)|}{|\alpha|^\sigma} dv ds \\ &\lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|F(s)\|_{L^p} \right) \int_0^t \frac{ds}{\langle s \rangle^{d-1+a}} \lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \sup_{s \in [0, t]} \|F(s)\|_{L^p},\end{aligned}$$

where we apply (4.5).

In conclusion, for $0 < t < 1$

$$\|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p, \infty}^\sigma} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma} + \sup_{s \in [0, t]} \|F(s)\|_{L^p} \right). \quad (6.8)$$

2.2) Estimate $\|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p, \infty}^\sigma}$ for $t \geq 1$.

Set

$$\begin{aligned}Z_1(x) &= Y_{s,t}(w, \frac{x-w}{t}) + w + \frac{s(x-w)}{t}, \quad Z_2(x) = W_{s,t}(w, \frac{x-w}{t}) + \frac{x-w}{t}, \\ Z_3(x) &= W_{s,t}(w, \frac{x-\alpha-w}{t}) + \frac{x-w}{t}, \quad Z_4(x) = Y_{s,t}(w, \frac{x-w}{t}) + w + \frac{s(x-\alpha-w)}{t}.\end{aligned} \quad (6.9)$$

We have

$$\delta_\alpha \mathcal{T}[F, \eta](t, x) = \mathcal{T}_\alpha^1[F, \eta](t, x) + \mathcal{T}_\alpha^2[F, \eta](t, x) + \mathcal{T}_\alpha^3[F, \eta](t, x) + \mathcal{T}_\alpha^4[F, \eta](t, x),$$

where

$$\begin{aligned}\mathcal{T}_\alpha^1[F, \eta](t, x) &= - \int_0^t \int_{\mathbb{R}^d} \delta_{\frac{s\alpha}{t}} F(s, \cdot) (Z_1(x)) \eta(Z_2(x)) \frac{dw ds}{t^d} + \int_0^t \int_{\mathbb{R}^d} \delta_{\frac{s\alpha}{t}} F(s, \cdot) (w + \frac{s}{t}(x-w)) \eta(\frac{x-w}{t}) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^2[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, Z_1(x-\alpha)) (\delta_{\frac{\alpha}{t}} \eta)(Z_2(x)) \frac{dw ds}{t^d} - \int_0^t \int_{\mathbb{R}^d} F(s, w + \frac{s}{t}(x-w-\alpha)) (\delta_{\frac{\alpha}{t}} \eta)(\frac{x-w}{t}) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^3[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} (F(s, Z_1(x-\alpha)) - F(s, Z_4(x))) \eta(Z_2(x)) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^4[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, Z_1(x-\alpha)) (\eta(Z_3(x)) - \eta(Z_2(x))) \frac{dw ds}{t^d}.\end{aligned}$$

Going back to the variable $v = \frac{x-w}{t}$, and denote

$$Z_5(x) = Y_{s,t}(x-tv, v - \frac{\alpha}{t}) + x - (t-s)v - \frac{s\alpha}{t}, \quad Z_6(x) = W_{s,t}(x-tv, v - \frac{\alpha}{t}) + v.$$

we reformulate as follows:

$$\begin{aligned}\mathcal{T}_\alpha^1[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} (-\delta_{\frac{s\alpha}{t}} F(s, \cdot) (X_{s,t}(x, v)) \eta(V_{s,t}(x, v)) + \delta_{\frac{s\alpha}{t}} F(s, \cdot) (x - (t-s)v) \eta(v)) dv ds, \\ \mathcal{T}_\alpha^2[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} (F(s, X_{s,t}(x, v)) (\delta_{\frac{\alpha}{t}} \eta)(V_{s,t}(x, v)) - F(s, x - (t-s)v) (\delta_{\frac{\alpha}{t}} \eta)(v)) dv ds, \\ \mathcal{T}_\alpha^3[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} (F(s, X_{s,t}(x, v) - \frac{s\alpha}{t}) - F(s, Z_5(x))) \eta(V_{s,t}(x, v)) dv ds, \\ \mathcal{T}_\alpha^4[F, \eta](t, x) &= \int_0^t \int_{\mathbb{R}^d} F(s, Z_5(x)) (\eta(V_{s,t}(x, v)) - \eta(Z_6(x))) dv ds.\end{aligned}$$

2.2.1) Estimate $\mathcal{T}_\alpha^1[F, \eta](t, x)$.

Applying (6.5) with $\delta_{\frac{s\alpha}{t}} F$, we have

$$\begin{aligned}\|\mathcal{T}_\alpha^1[F, \eta](t)\|_{L^p} &= \|\mathcal{T}[\delta_{\frac{s\alpha}{t}} F, \eta](t)\|_{L^p} \lesssim_{a, \mathbf{c}_1} t^{-\frac{d(p-1)}{p}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|\delta_{\frac{s\alpha}{t}} F(s)\|_{L^1} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|\delta_{\frac{s\alpha}{t}} F(s)\|_{L^p} \right) \\ &\lesssim_{a, \mathbf{c}_1} |\alpha|^\sigma t^{-\frac{d(p-1)}{p} - \sigma} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{B}_{1, \infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma} \right).\end{aligned}$$

Thus, we can yield

$$t^{\frac{d(p-1)}{p} + \sigma} \sup_\alpha \frac{\|\mathcal{T}_\alpha^1[F, \eta](t)\|_{L^p}}{|\alpha|^\sigma} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{B}_{1, \infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{B}_{p, \infty}^\sigma} \right). \quad (6.10)$$

2.2.2) Estimate $\mathcal{T}_\alpha^2[F, \eta](t, x)$.

Note that

$$|\delta_{\frac{\alpha}{t}} \eta(v)| + |\nabla_v (\delta_{\frac{\alpha}{t}} \eta)(v)| \lesssim_{\mathbf{c}_1} \min \left\{ \frac{|\alpha|}{t}, 1 \right\} \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right) \lesssim_{\mathbf{c}_1} \frac{|\alpha|^\sigma}{t^\sigma} \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right).$$

Thus, thanks to Remark 6.2, we have

$$\begin{aligned}\|\mathcal{T}_\alpha^2[F, \eta](t)\|_{L^p} &= \frac{|\alpha|^\sigma}{t^\sigma} \left\| \mathcal{T} \left[F, \frac{t^\sigma}{|\alpha|^\sigma} \delta_{\frac{\alpha}{t}} \eta(v) \right] \right\|_{L^p} \\ &\lesssim_{a, \mathbf{c}_1} \frac{|\alpha|^\sigma}{t^{\sigma + \frac{d(p-1)}{p}}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \|F(s)\|_{L^1} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} \right),\end{aligned}$$

which implies

$$t^{\frac{d(p-1)}{p} + \sigma} \sup_\alpha \frac{\|\mathcal{T}_\alpha^2[F, \eta](t)\|_{L^p}}{|\alpha|^\sigma} \lesssim_{a, \mathbf{c}_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} + \sup_{s \in [0, t]} \|F(s)\|_{L^1} \right).$$

2.2.3 Estimate $\mathcal{T}_\alpha^3[F, \eta](t, x)$.

$$\begin{aligned}
& |\mathcal{T}_\alpha^3[F, \eta](t, x)| \\
& \lesssim_{c_1} \int_0^t \int_{\mathbb{R}^d} \left| F\left(s, X_{s,t}(x, v) - \frac{s\alpha}{t}\right) - F\left(s, Z_5(x)\right) \right| \frac{dv ds}{\langle v \rangle^N} \\
& \lesssim_{c_1} \int_0^t \int_{\mathbb{R}^d} \left(\sup_z |z|^{-\sigma} \left| \delta_z F(s, \cdot) \left(Y_{s,t}(x - tv, v) + x - (t-s)v - \frac{s\alpha}{t} \right) \right| \left| Y_{s,t}(x - tv, v) - Y_{s,t}\left(x - tv, v - \frac{\alpha}{t}\right) \right|^\sigma \frac{dv ds}{\langle v \rangle^N} \right) \\
& \lesssim_{a, c_1} \frac{|\alpha|^\sigma}{t^\sigma} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \int_0^t \int_{\mathbb{R}^d} \sup_z |z|^{-\sigma} \left| \delta_z F(s, \cdot) \left(Y_{s,t}(x - tv, v) + x - (t-s)v - \frac{s\alpha}{t} \right) \right| \frac{ds}{\langle s \rangle^{\sigma(d-3+a)}} \frac{dv}{\langle v \rangle^N}. \tag{6.11}
\end{aligned}$$

Thus applying Lemma 8.1 with $\mathcal{H} = \sup_z |z|^{-\sigma} |\delta_z F(s, \cdot)|$ and $\varphi(x, v) = Y_{s,t}(x - tv, v - \frac{\alpha}{t}) - \frac{s\alpha}{t}$,

$$\begin{aligned}
& \|\mathcal{T}_\alpha^3[F, \eta](t)\|_{L^p} \\
& \lesssim_{a, c_1} \underbrace{\frac{|\alpha|^\sigma}{t^{\frac{d(p-1)}{p} + \sigma}} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \int_0^{t/2} \|F(s)\|_{\dot{F}_{1,\infty}^\sigma} \frac{1}{\langle s \rangle^{\sigma(d-3+a)}} ds}_{\text{Thanks to (8.2)}} + \underbrace{\frac{|\alpha|^\sigma}{t^\sigma} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \int_{t/2}^t \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \frac{1}{\langle s \rangle^{\sigma(d-3+a)}} ds}_{\text{Thanks to (8.1)}} \\
& \lesssim_{a, c_1} \frac{|\alpha|^\sigma}{t^{\frac{d(p-1)}{p} + \sigma}} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{F}_{1,\infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \right) \int_0^t \frac{1}{\langle s \rangle^{\sigma(d-2+a)}} ds \\
& \lesssim_{a, c_1} \frac{|\alpha|^\sigma}{t^{\frac{d(p-1)}{p} + \sigma}} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{F}_{1,\infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \right),
\end{aligned}$$

where in the last line, we use the fact that $a^2 + a > 1$. Thus,

$$t^{\frac{d(p-1)}{p} + \sigma} \sup_\alpha \frac{\|\mathcal{T}_\alpha^3[F, \eta](t)\|_{L^p}}{|\alpha|^\sigma} \lesssim_{a, c_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{F}_{1,\infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \right).$$

2.2.4 Estimate $\mathcal{T}_\alpha^4[F, \eta](t, x)$.

One has

$$\begin{aligned}
& |\mathcal{T}_\alpha^4[F, \eta](t, x)| \lesssim_{c_1} \int_0^t \int_{\mathbb{R}^d} |F(s, Z_5(x))| \left| W_{s,t}(x - tv, v) - W_{s,t}\left(x - tv, v - \frac{\alpha}{t}\right) \right| \frac{dv ds}{\langle v \rangle^N} \\
& \lesssim_{a, c_1} \min \left\{ 1, \frac{|\alpha|}{t} \right\} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \int_{\mathbb{R}^d} \left| F\left(s, Y_{s,t}\left(x - tv, v - \frac{\alpha}{t}\right) + x - (t-s)v - \frac{s\alpha}{t}\right) \right| \frac{1}{\langle s \rangle^{d-2+a}} \frac{dv ds}{\langle v \rangle^N} \\
& \lesssim_{a, c_1} \frac{|\alpha|^\sigma}{t^\sigma} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^t \int_{\mathbb{R}^d} \left| F\left(s, Y_{s,t}\left(x - tv, v - \frac{\alpha}{t}\right) + x - (t-s)v - \frac{s\alpha}{t}\right) \right| \frac{dv}{\langle v \rangle^N} \frac{ds}{\langle s \rangle^{d-2+a}}. \tag{6.12}
\end{aligned}$$

Then, applying Lemma 8.1 with $\mathcal{H} = F$ and $\varphi(x, v) = Y_{s,t}(x - tv, v - \frac{\alpha}{t}) - \frac{s\alpha}{t}$, one gets

$$\begin{aligned}
& \|\mathcal{T}_\alpha^4[F, \eta](t)\|_{L^p} \lesssim_{a, c_1} \underbrace{\frac{|\alpha|^\sigma}{t^{\sigma + \frac{d(p-1)}{p}}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_0^{t/2} \|F(s)\|_{L^1} \frac{ds}{\langle s \rangle^{d-2+a}}}_{\text{Thanks to (8.2)}} + \underbrace{\frac{|\alpha|^\sigma}{t^\sigma} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \int_{t/2}^t \|F(s)\|_{L^p} \frac{ds}{\langle s \rangle^{d-2+a}}}_{\text{Thanks to (8.1)}} \\
& \lesssim_{a, c_1} \frac{|\alpha|^\sigma}{t^{\frac{d(p-1)}{p} + \sigma}} \|(\rho, U)\|_{S_a^{\varepsilon_0}} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} + \sup_{s \in [0, t]} \|F(s)\|_{L^1} \right).
\end{aligned}$$

Hence, we obtain

$$t^{\frac{d(p-1)}{p} + \sigma} \sup_\alpha \frac{\|\mathcal{T}_\alpha^4[F, \eta](t)\|_{L^p}}{|\alpha|^\sigma} \lesssim_{a, c_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\sup_{s \in [0, t]} \langle s \rangle^\sigma \|F(s)\|_{\dot{F}_{1,\infty}^\sigma} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \right).$$

Summing up, we conclude that for $t \geq 1$

$$\sum_{p=1, \infty} t^{\frac{d(p-1)}{p} + \sigma} \|\mathcal{T}[F, \eta](t)\|_{\dot{B}_{p,\infty}^\sigma} \lesssim_{a, c_1} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \sum_{p=1, \infty} \left(\sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p}} \|F(s)\|_{L^p} + \sup_{s \in [0, t]} \langle s \rangle^{\frac{d(p-1)}{p} + \sigma} \|F(s)\|_{\dot{F}_{p,\infty}^\sigma} \right). \tag{6.13}$$

Hence, we proved the Lemma with (6.8) and (6.13). \blacksquare

Proposition 6.3 *Let a, σ and ρ satisfy the same condition in Lemma 6.1, then we have*

$$\|\mathcal{R}(\rho, U)\|_\sigma \lesssim_{a, M^*} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^{1+\sigma}. \quad (6.14)$$

Proof. Since $\langle v \rangle^N \sum_{j=0}^2 |\nabla^j (\nabla \mu(v))| \leq M^*$ thus, by Lemma 6.1, we have

$$\|\mathcal{R}(\rho, U)\|_\sigma \lesssim_{a, M^*} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^\sigma \left(\|E\|_\sigma + \sup_{s \in [0, t]} \langle s \rangle^\sigma \|E(s)\|_{\dot{F}_{1, \infty}^\sigma} \right) \lesssim_{a, M^*} \|(\rho, U)\|_{S_a^{\varepsilon_0}}^{1+\sigma}.$$

This implies the result. \blacksquare

Proposition 6.4 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$ and $\tilde{\varepsilon}_0 > 0$ be as in Proposition (4.1). Assume $\|(\rho_1, U_1)\|_{S_a^{\varepsilon_0}}, \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$. There holds,*

$$\|\mathcal{R}(\rho_1, U_1) - \mathcal{R}(\rho_2, U_2)\|_a \lesssim_{a, M^*} \left(\|(\rho_1, U_1)\|_{S_a^{\varepsilon_0}}^a + \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}^a \right) \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (6.15)$$

Proof. In this proof, we recall the notations $X_{s,t}^1, X_{s,t}^2, Y_{s,t}^1, Y_{s,t}^2, V_{s,t}^1, V_{s,t}^2, \mathbf{Y}_{s,t}, \mathbf{W}_{s,t}$ in Proposition 4.6, and $X_{s,t}(x, v, \varpi), Y_{s,t}(x, v, \varpi), V_{s,t}(x, v, \varpi), W_{s,t}(x, v, \varpi)$ in (5.13). Then we have

$$\begin{aligned} \mathbf{R}(\rho_1, \rho_2, U_1, U_2)(t, x) &:= \mathcal{R}(\rho_1, U_1)(t, x) - \mathcal{R}(\rho_2, U_2)(t, x) \\ &= \int_0^t \int_{\mathbb{R}^d} (E^1 - E^2)(s, x - (t-s)v) \cdot \nabla_v \mu(v) dv ds - \int_0^t \int_{\mathbb{R}^d} (E^1 - E^2)(s, X_{s,t}^1(x, v)) \cdot \nabla_v \mu(V_{s,t}^1(x, v)) dv ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d} (E^2(s, X_{s,t}^1(x, v)) - E^2(s, X_{s,t}^2(x, v))) \cdot \nabla \mu(V_{s,t}^1(x, v)) \\ &\quad + \int_0^t \int_{\mathbb{R}^d} E^2(s, X_{s,t}^2(x, v)) \cdot (\nabla \mu(V_{s,t}^1(x, v)) - \nabla \mu(V_{s,t}^2(x, v))) \\ &=: \sum_{i=1}^d \mathcal{T}[(E^1 - E^2)_i, \partial_i \mu](t, x) + \mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x) + \mathbf{R}^2(\rho_1, \rho_2, U_1, U_2)(t, x). \end{aligned}$$

Thanks to Lemma 6.1 with $F = (E^1 - E^2)_i, \eta = \partial_i \mu$ and $\sigma = a$, we get

$$\sum_{i=1}^d \|\mathcal{T}[(E^1 - E^2)_i, \partial_i \mu]\|_a \lesssim_{a, M^*} \|(\rho_1, U_1)\|_{S_a^{\varepsilon_0}}^a \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (6.16)$$

(1) Estimate on $\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)$, one has

$$\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x) = - \sum_{i=1}^d \int_0^1 \int_0^t \int_{\mathbb{R}^d} \mathbf{Y}_{s,t}(x - vt, v) \nabla E_i^2(s, X_{s,t}(x, v, \varpi)) \partial_i \mu(V_{s,t}^1(x, v)) d\varpi ds dv.$$

Hence,

$$\begin{aligned} |\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)| &\leq \sup_{0 \leq s \leq t} \langle s \rangle^{d-2+a} \|\mathbf{Y}_{s,t}\|_{L^\infty} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \frac{|\nabla E^2(s, X_{s,t}(x, v, \varpi))|}{\langle s \rangle^{d-2+a}} |\nabla \mu(V_{s,t}^1(x, v))| ds dv d\varpi \\ &\stackrel{(4.21)}{\lesssim_{a, M^*}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2-a} |\nabla E^2(s, X_{s,t}(x, v, \varpi))| \frac{1}{\langle v \rangle^N} ds dv d\varpi. \end{aligned}$$

Note that $X_{s,t}(x, v, \varpi) = x - (t-s)v + Y_{s,t}(x - tv, v, \varpi)$, then we apply Lemma 8.2 with $\mathcal{H} = \nabla E^2$ and $\varphi(x, v) = Y_{s,t}(x - tv, v, \varpi)$ to obtain that for $t > 1$

$$\begin{aligned} &\|\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)\|_{L^p} \\ &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_a \int_0^1 \int_0^t \langle s \rangle^{-d+2-a} \left\| \int_{\mathbb{R}^d} \nabla E^2(s, X_{s,t}(x, v, \varpi)) \frac{1}{\langle v \rangle^N} dv \right\|_{L_x^p} ds d\varpi \\ &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \left(\int_0^{t/2} \langle s \rangle^{-d+2-a} t^{-\frac{d(p-1)}{p}} \|\nabla E^2(s)\|_{L_x^1} ds + \int_{t/2}^t \langle s \rangle^{-d+2-a} \|\nabla E^2(s)\|_{L_x^p} ds \right) \\ &\lesssim_{a, M^*} \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{t^{\frac{d(p-1)}{p}}} \left(\int_0^{t/2} + \int_{t/2}^t \right) \frac{\langle s \rangle^{2-a}}{\langle s \rangle^{d+a}} ds \lesssim \frac{\|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}}{t^{\frac{d(p-1)}{p}}}. \end{aligned} \quad (6.17)$$

Meanwhile, for $0 \leq t \leq 1$

$$\begin{aligned} \|\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)\|_{L^p} &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^t \langle s \rangle^{-d+2-a} \|\nabla E^2(s)\|_{L^p} ds \\ &\lesssim_{a, M^*} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \end{aligned} \quad (6.18)$$

Then, with (6.18) and (6.17), we can obtain

$$\sum_{p=1, \infty} \langle t \rangle^{\frac{d(p-1)}{p}} \|\mathbf{R}^1(\rho_1, \rho_2)\|_{L^p} \lesssim_{a, M^*} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (6.19)$$

Now we deal with $\frac{|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2)(t, x)|}{|\alpha|^a}$, for the case $t \leq 1$ we have

$$\begin{aligned} \frac{|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2)(t, x)|}{|\alpha|^a} &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2+a} |\nabla E^2(s, X_{s,t}(x, v, \varpi))| \frac{dv}{\langle v \rangle^N} ds d\varpi \\ &+ \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2} \sup_z \frac{|\delta_z \nabla E^2(s, \cdot)(X_{s,t}(x, v, \varpi))|}{|z|^a} \frac{dv}{\langle v \rangle^N} ds d\varpi \\ &+ \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2-a} |\nabla E^2(s, X_{s,t}(x - \alpha, v, \varpi))| \frac{dv}{\langle v \rangle^N} ds d\varpi. \end{aligned}$$

Then thanks to (8.3), we deduce that

$$\begin{aligned} \frac{\|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2)\|_{L^p}}{|\alpha|^a} &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^t \left(\langle s \rangle^{-d+2-a} \|\nabla E^2(s)\|_{L^p} + \langle s \rangle^{-d+2} \|\nabla E^2(s)\|_{\dot{F}_{p, \infty}^a} \right) ds \\ &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \int_0^t \left(\frac{\langle s \rangle^{-d+2-a}}{\langle s \rangle^{\frac{d(p-1)}{p}+a}} + \frac{\langle s \rangle^{-d+2}}{\langle s \rangle^{\frac{d(p-1)}{p}+a}} \right) ds \\ &\lesssim_{a, M^*} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}. \end{aligned}$$

For the case $t > 1$, $\frac{|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)|}{|\alpha|^a}$ can be recast as

$$\begin{aligned} \frac{|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)(t, x)|}{|\alpha|^a} &\lesssim_{a, M^*} \frac{1}{t^a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2} |\nabla E^2(s, X_{s,t}(x, v, \varpi))| \frac{dv}{\langle v \rangle^N} ds d\varpi \\ &+ \frac{1}{t^a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2} \sup_z \frac{|\delta_z \nabla E^2(s, \cdot)(X_{s,t}(x, v, \varpi))|}{|z|^a} \frac{dv}{\langle v \rangle^N} ds d\varpi \\ &+ \frac{1}{t^a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^1 \int_0^t \int_{\mathbb{R}^d} \langle s \rangle^{-d+2-a} |\nabla E_i^2(s, X_{s,t}(x - \alpha, v, \varpi))| \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right) dv d\varpi ds. \end{aligned}$$

Thus, it holds that for $t > 1$

$$\begin{aligned} &\frac{\|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)\|_{L^p}}{|\alpha|^a} \\ &\lesssim_{a, M^*} \frac{1}{t^a} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_0^{t/2} \left(\langle s \rangle^{-d+2-a} \|\nabla E^2(s)\|_{L^1} + \langle s \rangle^{-d+2} \|\nabla E^2(s)\|_{\dot{F}_{1, \infty}^a} \right) ds \\ &+ \frac{1}{t^{a+\frac{d(p-1)}{p}}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \int_{t/2}^t \left(\langle s \rangle^{-d+2-a} \|\nabla E^2(s)\|_{L^p} + \langle s \rangle^{-d+2} \|\nabla E^2(s)\|_{\dot{F}_{p, \infty}^a} \right) ds \\ &\lesssim_{a, M^*} \frac{1}{t^{a+\frac{d(p-1)}{p}}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}}. \end{aligned}$$

Then,

$$\sum_{p=1, \infty} \langle t \rangle^{a+\frac{d(p-1)}{p}} \frac{\|\delta_\alpha \mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)\|_{L^p}}{|\alpha|^a} \lesssim_{a, M^*} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (6.20)$$

Combining this with (6.19), one gets

$$\|\mathbf{R}^1(\rho_1, \rho_2, U_1, U_2)\|_{S_a^{\varepsilon_0}} \lesssim_{a, M^*} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}. \quad (6.21)$$

(2) Estimate on $\mathbf{R}^2(\rho_1, \rho_2, U_1, U_2)(t, x)$. We infer that

$$\begin{aligned} & \mathbf{R}^2(\rho_1, \rho_2, U_1, U_2)(t, x) \\ &= - \sum_{i=1}^d \int_0^1 \int_0^t \int_{\mathbb{R}^d} \mathbf{W}_{s,t}(x-vt, v) E_i^2(s, X_{s,t}^2(x, v)) (\nabla \partial_i \mu)(V_{s,t}(x, v, \varpi)) dv ds d\varpi. \end{aligned}$$

Analogously, we obtain

$$\|\mathbf{R}^2(\rho_1, \rho_2, U_1, U_2)\|_{S_a^{\varepsilon_0}} \lesssim_{a, M^*} \|(\rho_2, U_2)\|_{S_a^{\varepsilon_0}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^{\varepsilon_0}}, \quad (6.22)$$

which complete the proof. \blacksquare

Lemma 6.5 *Let $H_0 : \mathbb{R}^d \rightarrow \mathbb{R}$ and $A_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $\lambda \in \mathbb{R}$. Then, for any multi-index $\alpha \neq 0$*

$$\begin{aligned} & |\partial_x^\alpha [H_0(\lambda x + A_0(x))] - (\partial_x^\alpha H_0)(\lambda x + A_0(x)) \lambda^{|\alpha|}| \lesssim |(\nabla^{|\alpha|} H_0)(\lambda x + A_0(x))| |\nabla A_0(x)| (|\lambda| + |\nabla A_0(x)|)^{|\alpha|-1} \\ & + \sum_{k=1}^{|\alpha|-1} \sum_{j=0}^{k-1} \sum_{i=2}^{|\alpha|-j} |(\nabla^k H_0)(\lambda x + A_0(x))| (|\lambda| + |\nabla A_0(x)|)^j |\nabla^i A_0(x)|^{\frac{|\alpha|-j}{i}}. \end{aligned}$$

Proof. Assume $\partial_x^\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_d}^{\alpha_d}$. We know that, for $\tilde{A}_0(x) = \lambda x + A_0(x)$

$$|\partial_x^\alpha [H_0(\lambda x + A_0(x))] - \mathbf{Q}| \lesssim \sum_{k=1}^{|\alpha|-1} |(\nabla^k H_0)(\tilde{A}_0(x))| \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = |\alpha|}} \prod_{j=1}^k |\nabla^{m_j} \tilde{A}_0(x)|$$

with

$$\mathbf{Q} = (\nabla^{|\alpha|} H_0)(\lambda x + A_0(x)) : [(\partial_{x_1} \tilde{A}_0(x))^{\otimes \alpha_1} \otimes (\partial_{x_2} \tilde{A}_0(x))^{\otimes \alpha_2} \otimes \dots \otimes (\partial_{x_d} \tilde{A}_0(x))^{\otimes \alpha_d}]. \quad (6.23)$$

It is easy to check that

$$\begin{aligned} & \sum_{k=1}^{|\alpha|-1} |(\nabla^k H_0)(\tilde{A}_0(x))| \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = |\alpha|}} \prod_{j=1}^k |\nabla^{m_j} \tilde{A}_0(x)| \\ & \lesssim \sum_{k=1}^{|\alpha|-1} \sum_{j=0}^{k-1} \sum_{i=2}^{|\alpha|-j} |(\nabla^k H_0)(\lambda x + A_0(x))| (|\lambda| + |\nabla A_0(x)|)^j |\nabla^i A_0(x)|^{\frac{|\alpha|-j}{i}} \end{aligned}$$

and

$$\left| \mathbf{Q} - (\partial_x^\alpha H_0)(\lambda x + A_0(x)) \lambda^{|\alpha|} \right| \lesssim |(\nabla^{|\alpha|} H_0)(\lambda x + A_0(x))| |\nabla A_0(x)| (|\lambda| + |\nabla A_0(x)|)^{|\alpha|-1}. \quad (6.24)$$

Combining (6.23) and (6.24), we finish the proof. \blacksquare

Proposition 6.6 *Let $a \in (\frac{\sqrt{5}-1}{2}, 1)$ and assume $\|(\rho, U)\|_{S_a^{\varepsilon_0}} \leq \varepsilon_0 \leq \tilde{\varepsilon}_0$ with $\tilde{\varepsilon}_0$ mentioned in Proposition 4.1. Then we have for $b \in (0, 1 - \delta_0)$,*

$$\|\mathcal{R}(\rho, U)\|_{l+1, b} \lesssim_{a, b, \delta_0, \mathbf{c}, \mathbf{c}', C_{A, l}} 1, \quad (6.25)$$

provided $\|(\rho, U)\|_{l, 1-\delta_0} \leq \mathbf{c}$, $\sum_{j=2}^{l+1} |A^{(j)}(r)| \leq C_{A, l}$, $\forall |r| \leq 1$, and μ satisfies

$$\sum_{i=1}^{l+3} \langle v \rangle^N |\nabla^i \mu(v)| \leq \mathbf{c}', \quad (6.26)$$

where $l \geq 0$, $0 < \delta_0 < 2 - \sqrt{3}$ and $N > d$.

Proof. In this proof, we denote $C = C(a, b, \delta_0, \mathbf{c}, \mathbf{c}', C_{A, l})$.

(1) We firstly consider $t \geq 1$. Set

$$\begin{aligned} F_{1,1}(x) &= \delta_{\frac{s}{t}} E_i(s, \cdot)(Z_1(x)), \quad \eta_{1,1}(x) = \partial_i \mu(Z_2), \\ F_{1,2}(x) &= \delta_{\frac{s}{t}} E_i(s, \cdot)(w + \frac{s}{t}(x-w)), \quad \eta_{1,2}(x) = \partial_i \mu(\frac{x-w}{t}), \\ F_{2,1}(x) &= E_i(s, Z_1(x)), \quad \eta_{2,1}(x) = \delta_{\frac{s}{t}} \partial_i \mu(Z_2(x)), \\ F_{2,2}(x) &= E_i(s, w + \frac{s}{t}(x-w)), \quad \eta_{2,2}(x) = \delta_{\frac{s}{t}} \partial_i \mu(\frac{x-w}{t}), \\ F_3(x) &= E_i(s, Z_1(x-\alpha)) - E_i(s, Z_4(x)), \quad \eta_3(x) = \partial_i \mu(Z_3(x)) - \partial_i \mu(Z_2(x)), \end{aligned}$$

where Z_1, Z_2, Z_3, Z_4 are defined in (6.9) and $w = x - vt$, then we have,

$$\delta_\alpha \mathcal{T}[E_i, \partial_i \mu](t, x) = \sum_{j=1}^4 \mathcal{T}_\alpha^j[E_i, \partial_i \mu](t, x),$$

in which

$$\begin{aligned} \mathcal{T}_\alpha^1[E_i, \partial_i \mu](t, x) &= - \int_0^t \int_{\mathbb{R}^d} (F_{1,1}(x)\eta_{1,1}(x) - F_{1,2}(x)\eta_{1,2}(x)) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^2[E_i, \partial_i \mu](t, x) &= \int_0^t \int_{\mathbb{R}^d} (F_{2,1}(x)\eta_{2,1}(x) - F_{2,2}(x)\eta_{2,2}(x)) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^3[E_i, \partial_i \mu](t, x) &= \int_0^t \int_{\mathbb{R}^d} F_3(x)\eta_{1,1}(x) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^4[E_i, \partial_i \mu](t, x) &= \int_0^t \int_{\mathbb{R}^d} F_{2,1}(x - \alpha)\eta_3(x) \frac{dw ds}{t^d}. \end{aligned}$$

Hence, we obtain

$$|\nabla_x^{l+1} \delta_\alpha \mathcal{T}^j[E_i, \partial_i \mu](t, x)| \lesssim \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_\alpha^j[E_i, \partial_i \mu, \beta_1, \beta_2](t, x)|,$$

where

$$\begin{aligned} \mathcal{T}_\alpha^1[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= - \int_0^t \int_{\mathbb{R}^d} (\partial_x^{\beta_1} F_{1,1}(x) \partial_x^{\beta_2} \eta_{1,1}(x) - \partial_x^{\beta_1} F_{1,2}(x) \partial_x^{\beta_2} \eta_{1,2}(x)) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^2[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= \int_0^t \int_{\mathbb{R}^d} (\partial_x^{\beta_1} F_{2,1}(x) \partial_x^{\beta_2} \eta_{2,1}(x) - \partial_x^{\beta_1} F_{2,2}(x) \partial_x^{\beta_2} \eta_{2,2}(x)) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^3[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= \int_0^t \int_{\mathbb{R}^d} \partial_x^{\beta_1} F_3(x) \partial_x^{\beta_2} \eta_{1,1}(x) \frac{dw ds}{t^d}, \\ \mathcal{T}_\alpha^4[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= \int_0^t \int_{\mathbb{R}^d} \partial_x^{\beta_1} F_{2,1}(x - \alpha) \partial_x^{\beta_2} \eta_3(x) \frac{dw ds}{t^d}. \end{aligned}$$

1.1) Estimate of terms in $\nabla_x^{l+1} \mathcal{T}_\alpha^1[E_i, \partial_i \mu]$.

Note that

$$\begin{aligned} & \partial_x^{\beta_1} F_{1,1}(x) \partial_x^{\beta_2} \eta_{1,1}(x) - \partial_x^{\beta_1} F_{1,2}(x) \partial_x^{\beta_2} \eta_{1,2}(x) \\ &= \left(\partial_x^{\beta_1} F_{1,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1) \right) \left(\partial_x^{\beta_2} \eta_{1,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} (\partial_x^{\beta_2} \partial_i \mu)(Z_2) \right) \\ &+ \left(\partial_x^{\beta_1} F_{1,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1) \right) \left(\frac{1}{t}\right)^{|\beta_2|} (\partial_x^{\beta_2} \partial_i \mu)(Z_2) \\ &+ \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1) \left(\partial_x^{\beta_2} \eta_{1,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} (\partial_x^{\beta_2} \partial_i \mu)(Z_2) \right) \\ &+ \left(\left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1) \left(\frac{1}{t}\right)^{|\beta_2|} (\partial_x^{\beta_2} \partial_i \mu)(Z_2) - \partial_x^{\beta_1} F_{1,2}(x) \partial_x^{\beta_2} \eta_{1,2}(x) \right) \\ &:= \sum_{k=1}^4 \mathcal{T}_{\alpha,k}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t}). \end{aligned}$$

We first deal with the fourth term directly.

$$\begin{aligned} & \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \int_0^t \int_{\mathbb{R}^d} \mathcal{T}_{\alpha,4}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t}) \frac{dw ds}{t^d} \right\|_{L_x^p} = \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \int_0^t \int_{\mathbb{R}^d} \mathcal{T}_{\alpha,4}^{1,\beta_1,\beta_2}(t, x, v) dv ds \right\|_{L_x^p} \\ &= \frac{1}{t^{l+1}} \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \mathcal{T}_\alpha^1[s^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i, \partial_x^{\beta_2} \partial_i \mu] \right\|_{L_x^p} \\ &\stackrel{(6.10)}{\lesssim} \frac{1}{a, c} \frac{|\alpha|^b}{t^{d+l+1+b}} \sum_{|\beta_1|=0}^{l+1} \left(\sup_{0 \leq s \leq t} |s|^{|\beta_1|+b} \|\nabla^{|\beta_1|} E_i(s)\|_{\dot{F}_{1,\infty}^b} + \sup_{0 \leq s \leq t} |s|^{|\beta_1|+b+d} \|\nabla^{|\beta_1|} E_i(s)\|_{\dot{F}_{\infty,\infty}^b} \right) \\ &\stackrel{(4.15)}{\lesssim} C \frac{|\alpha|^b}{t^{d+l+1+b}} \sum_{|\beta_1|=0}^{l+1} \sup_{0 \leq s \leq t} \frac{|s|^{|\beta_1|+b}}{\langle s \rangle^{|\beta_1|+1-\delta_0+d}} \lesssim C \frac{|\alpha|^b}{t^{d+l+1+b}}. \end{aligned} \tag{6.27}$$

Then we start to estimate $\mathcal{T}_{\alpha,k}^{1,\beta_1,\beta_2}$, where $k = 1, 2, 3$.

Applying Lemma 6.5 and Proposition 4.4, 4.4, we get

$$\begin{aligned}
& \left| \partial_x^{\beta_1} F_{1,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1) \right| \lesssim \frac{\mathbf{1}_{|\beta_1| \geq 1}}{t^{|\beta_1|}} |\nabla^{|\beta_1|} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| |\nabla_v Y_{s,t}| (s + |\nabla_v Y_{s,t}|)^{|\beta_1|-1} \\
& + \frac{\mathbf{1}_{|\beta_1| \geq 1}}{t^{|\beta_1|}} \sum_{m=1}^{|\beta_1|-1} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_1|-j} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| (|s| + |\nabla_v Y_{s,t}|)^j |\nabla_v^i Y_{s,t}|^{\frac{|\beta_1|-j}{i}} \\
& \lesssim_C \mathbf{1}_{|\beta_1| \geq 1} \left(\frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} |\nabla^{|\beta_1|} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{|\beta_1|-d+\delta_0} + \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} \sum_{m=1}^{|\beta_1|-1} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \right) \\
& \lesssim_C \mathbf{1}_{|\beta_1| \geq 1} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} \sum_{m=1}^{|\beta_1|} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1},
\end{aligned}$$

with $\nabla_v^i Y_{s,t} := (\nabla_v^i Y_{s,t})(w, \frac{x-w}{t})$. Similarly, we also can estimate

$$\begin{aligned}
& \left| \partial_x^{\beta_2} \eta_{1,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} (\partial_x^{\beta_2} \partial_i \mu)(Z_2) \right| \lesssim \frac{\mathbf{1}_{|\beta_2| \geq 1}}{t^{|\beta_2|}} |\nabla^{|\beta_2|} \partial_i \mu(Z_2)| |\nabla_v W_{s,t}| (1 + |\nabla_v W_{s,t}|)^{|\beta_2|-1} \\
& + \frac{\mathbf{1}_{|\beta_2| \geq 1}}{t^{|\beta_2|}} \sum_{m=1}^{|\beta_2|-1} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_2|-j} |\nabla^m \partial_i \mu(Z_2)| (1 + |\nabla_v W_{s,t}|)^j |\nabla_v^i W_{s,t}|^{\frac{|\beta_2|-j}{i}} \\
& \stackrel{(4.4),(4.4)}{\lesssim_C} \mathbf{1}_{|\beta_2| \geq 1} \left(\frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_2|}} |\nabla^{|\beta_2|} \partial_i \mu(Z_2)| \langle s \rangle^{-d+\delta_0} + \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_2|}} \sum_{m=1}^{|\beta_2|} |\nabla^m \partial_i \mu(Z_2)| \langle s \rangle^{-d+1} \right) \\
& \lesssim_C \mathbf{1}_{|\beta_2| \geq 1} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_2|}} \frac{1}{\langle \frac{x-w}{t} \rangle^N} \langle s \rangle^{-d+1}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,1}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \lesssim_C \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}^2}{t^{l+1}} \sum_{|\beta_1|=1}^l \sum_{m=1}^{|\beta_1|} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-2d+2} \frac{1}{\langle \frac{x-w}{t} \rangle^N}, \\
& \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,2}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \lesssim_C \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}^2}{t^{l+1}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \frac{1}{\langle \frac{x-w}{t} \rangle^N}, \\
& \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,3}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \lesssim_C \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}^2}{t^{l+1}} \sum_{|\beta_1|=0}^l |\partial_x^{\beta_1} \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{|\beta_1|-d+1} \frac{1}{\langle \frac{x-w}{t} \rangle^N}.
\end{aligned}$$

Hence, we get

$$\sum_{k=1}^3 \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,k}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \lesssim_C \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}^2}{t^{l+1}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \frac{1}{\langle \frac{x-w}{t} \rangle^N}. \quad (6.28)$$

Note that we have (8.1) and (8.2) in the Appendix with $\mathcal{H} = \nabla^m \delta_{\frac{s\alpha}{t}} E_i$ and $\varphi = Y_{s,t}(x, v)$, we firstly change of variables $v = \frac{x-w}{t}$, then applying (4.15) to obtain

$$\begin{aligned}
& \left\| \int_0^t \int_{\mathbb{R}^d} \sum_{k=1}^3 \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,k}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \frac{dw ds}{t^d} \right\|_{L_x^p} \\
& \lesssim_C \frac{1}{t^{l+1}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_0^t \langle s \rangle^{m-d+1} \left\| \int_{\mathbb{R}^d} |\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s, \cdot)(X_{s,t}(x, v))| \frac{1}{\langle v \rangle^N} dv \right\|_{L_x^p} ds \\
& \lesssim_C \frac{1}{t^{l+1+\frac{d(p-1)}{p}}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_0^{t/2} \langle s \rangle^{m-d+1} \|\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s)\|_{L^1} ds + \frac{1}{t^{l+1}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_{t/2}^t \langle s \rangle^{m-d+1} \|\nabla^m \delta_{\frac{s\alpha}{t}} E_i(s)\|_{L^p} ds \\
& \lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \left(\int_0^{t/2} + \int_{t/2}^t \right) \frac{|s|^b \langle s \rangle^{m-d+1}}{\langle s \rangle^{m+1-\delta_0}} ds \lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}}.
\end{aligned} \quad (6.29)$$

Thanks to (6.27) and (6.29), we have

$$t^{\frac{d(p-1)}{p}+l+1+b} \sup_{\alpha} \frac{\|\nabla_x^{l+1} \mathcal{T}_{\alpha}^1[E_i, \partial_i \mu](t)\|_{L_x^p}}{|\alpha|^b} \lesssim_{a,b,\delta_0,c,c',C_{A,l}} 1. \quad (6.30)$$

1.2) Estimate of terms in $\nabla_x^{l+1} \mathcal{T}_{\alpha}^2[E_i, \partial_i \mu]$. Now, we consider the other term. It is clear that

$$\begin{aligned} & \partial_x^{\beta_1} F_{2,1}(x) \partial_x^{\beta_2} \eta_{2,1}(x) - \partial_x^{\beta_1} F_{2,2}(x) \partial_x^{\beta_2} \eta_{2,2}(x) \\ &= \left(\partial_x^{\beta_1} F_{2,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} E_i(s, \cdot)(Z_1) \right) \left(\partial_x^{\beta_2} \eta_{2,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} \partial_x^{\beta_2} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2) \right) \\ &+ \left(\partial_x^{\beta_1} F_{2,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} E_i(s, \cdot)(Z_1) \right) \left(\frac{1}{t}\right)^{|\beta_2|} \partial_x^{\beta_2} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2) \\ &+ \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} E_i(s, \cdot)(Z_1) \left(\partial_x^{\beta_2} \eta_{2,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} \partial_x^{\beta_2} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2) \right) \\ &+ \left(\left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} E_i(s, \cdot)(Z_1) \left(\frac{1}{t}\right)^{|\beta_2|} \partial_x^{\beta_2} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2) - \partial_x^{\beta_1} F_{2,2}(x) \partial_x^{\beta_2} \eta_{2,2}(x) \right) \\ &:= \sum_{k=1}^4 \mathcal{T}_{\alpha,k}^{2,\beta_1,\beta_2} \left(t, x, \frac{x-w}{t} \right). \end{aligned}$$

Then, we also compute the last term first as follows

$$\begin{aligned} & \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \int_0^t \int_{\mathbb{R}^d} \mathcal{T}_{\alpha,4}^{2,\beta_1,\beta_2} \left(t, x, \frac{x-w}{t} \right) \frac{dw ds}{t^d} \right\|_{L_x^p} = \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \int_0^t \int_{\mathbb{R}^d} \mathcal{T}_{\alpha,4}^{2,\beta_1,\beta_2} (t, x, v) dv ds \right\|_{L_x^p} \\ &= \frac{1}{t^{l+1}} \sum_{|\beta_1|+|\beta_2|=l+1} \left\| \mathcal{T}_{\alpha}^2 [s^{|\beta_1|} \partial^{\beta_1} E_i, \partial^{\beta_2} \partial_i \mu] \right\|_{L_x^p} \\ &\stackrel{(6.10)}{\lesssim_{a,c}} \frac{|\alpha|^b}{t^{d+l+1+b}} \sum_{|\beta_1|=0}^{l+1} \left(\sup_{0 \leq s \leq t} |s|^{|\beta_1|} \|\nabla^{|\beta_1|} E_i(s)\|_{L^1} + \sup_{0 \leq s \leq t} |s|^{|\beta_1|+d} \|\nabla^{|\beta_1|} E_i(s)\|_{L^\infty} \right) \\ &\stackrel{(4.15)}{\lesssim_C} \frac{|\alpha|^b}{t^{d+l+1+b}} \sum_{|\beta_1|=0}^{l+1} \sup_{0 \leq s \leq t} \frac{|s|^{|\beta_1|}}{\langle s \rangle^{|\beta_1|+1-\delta_0+d}} \lesssim_C \frac{|\alpha|^b}{t^{d+l+1+b}}. \end{aligned} \quad (6.31)$$

Now, we turn to $\mathcal{T}_{\alpha,k}^{2,\beta_1,\beta_2}$, where $k = 1, 2, 3$. Applying Proposition 4.4, we can get

$$\begin{aligned} & \left| \partial_x^{\beta_1} F_{2,1}(x) - \left(\frac{s}{t}\right)^{|\beta_1|} \partial_x^{\beta_1} E_i(s, \cdot)(Z_1) \right| \lesssim \frac{\mathbf{1}_{|\beta_1| \geq 1}}{t^{|\beta_1|}} |\nabla^{|\beta_1|} E_i(s, \cdot)(Z_1)| |(\nabla_v Y_{s,t})| (s + |\nabla_v Y_{s,t}|)^{|\beta_1|-1} \\ &+ \frac{\mathbf{1}_{|\beta_1| \geq 1}}{t^{|\beta_1|}} \sum_{m=1}^{|\beta_1|-1} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_1|-j} |\nabla^m E_i(s, \cdot)(Z_1)| (|s| + |\nabla_v Y_{s,t}|)^j |\nabla_v^i Y_{s,t}|^{\frac{|\beta_1|-j}{i}} \\ &\stackrel{(4.4),(4.4)}{\lesssim_C} \mathbf{1}_{|\beta_1| \geq 1} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} \left(|\nabla^{|\beta_1|} E_i(s, \cdot)(Z_1)| \langle s \rangle^{|\beta_1|-d+\delta_0} + \sum_{m=1}^{|\beta_1|-1} |\nabla^m E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \right) \\ &\lesssim_C \mathbf{1}_{|\beta_1| \geq 1} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} \sum_{m=1}^{|\beta_1|} |\nabla^m E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \end{aligned}$$

and

$$\begin{aligned} & \left| \partial_x^{\beta_2} \eta_{2,1}(x) - \left(\frac{1}{t}\right)^{|\beta_2|} \partial_x^{\beta_2} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2) \right| \lesssim \frac{\mathbf{1}_{|\beta_2| \geq 1}}{t^{|\beta_2|}} |\nabla^{|\beta_2|} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2)| |\nabla_v W_{s,t}| (1 + |\nabla_v W_{s,t}|)^{|\beta_2|-1} \\ &+ \frac{\mathbf{1}_{|\beta_2| \geq 1}}{t^{|\beta_2|}} \sum_{m=1}^{|\beta_2|} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_2|-j} |\nabla^m (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2)| (1 + |\nabla_v W_{s,t}|)^j |\nabla_v^i W_{s,t}|^{\frac{|\beta_2|-j}{i}} \\ &\stackrel{(4.4)}{\lesssim_C} \mathbf{1}_{|\beta_2| \geq 1} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_2|}} \left(|\nabla^{|\beta_2|} (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2)| \langle s \rangle^{-d+\delta_0} + \sum_{m=1}^{|\beta_2|-1} |\nabla^m (\delta_{\frac{\alpha}{t}} \partial_i \mu)(Z_2)| \langle s \rangle^{-d+1} \right) \\ &\lesssim_C \mathbf{1}_{|\beta_2| \geq 1} \frac{|\alpha|^b \|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_2|+b}} \left(\frac{1}{\langle \frac{x-w}{t} \rangle^N} + \frac{1}{\langle \frac{x-w-\alpha}{t} \rangle^N} \right) \langle s \rangle^{-d+1}. \end{aligned}$$

Define

$$\Xi(x, w, \alpha, t) = \frac{1}{\langle \frac{x-w}{t} \rangle^N} + \frac{1}{\langle \frac{x-w-\alpha}{t} \rangle^N}.$$

Thus, we have

$$\begin{aligned} \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,1}^{2,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| &\lesssim_C \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=1}^l \sum_{m=1}^{|\beta_1|} |\nabla^m E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-2d+2} \Xi(x, w, \alpha, t), \\ \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,2}^{2,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| &\lesssim_C \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} |\nabla^m E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \Xi(x, w, \alpha, t), \\ \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,3}^{2,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| &\lesssim_C \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=0}^l |\partial_x^{\beta_1} E_i(s, \cdot)(Z_1)| \langle s \rangle^{|\beta_1|-d+1} \Xi(x, w, \alpha, t). \end{aligned}$$

We observe to get

$$\sum_{k=1}^3 \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,k}^{2,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \lesssim_C \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} |\nabla^m E_i(s, \cdot)(Z_1)| \langle s \rangle^{m-d+1} \Xi(x, w, \alpha, t). \quad (6.32)$$

Since we have Remark 8.2 in the Appendix with $\mathcal{H} = \nabla^m E_i$ and $\varphi = Y_{s,t}(x, v)$, we firstly change of variables $v = \frac{x-w}{t}$, then applying (4.15) to obtain

$$\begin{aligned} &\left\| \int_0^t \int_{\mathbb{R}^d} \sum_{k=1}^3 \sum_{|\beta_1|+|\beta_2|=l+1} |\mathcal{T}_{\alpha,k}^{1,\beta_1,\beta_2}(t, x, \frac{x-w}{t})| \frac{dw ds}{t^d} \right\|_{L_x^p} \\ &\lesssim_C \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_0^t \langle s \rangle^{m-d+1} \left\| \int_{\mathbb{R}^d} |\nabla^m E_i(s, \cdot)(X_{s,t}(x, v))| \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right) dv \right\|_{L_x^p} ds \\ &\lesssim_C \frac{|\alpha|^b}{t^{l+1+b+\frac{d(p-1)}{p}}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_0^{t/2} \langle s \rangle^{m-d+1} \|\nabla^m E_i(s)\|_{L^1} ds \\ &\quad + \frac{|\alpha|^b}{t^{l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \int_{t/2}^t \langle s \rangle^{m-d+1} \|\nabla^m E_i(s)\|_{L^p} ds \\ &\lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}} \sum_{|\beta_1|=1}^{l+1} \sum_{m=1}^{|\beta_1|} \left(\int_0^{t/2} + \int_{t/2}^t \right) \frac{|s|^b \langle s \rangle^{m-d+1}}{\langle s \rangle^{m+1-\delta_0}} ds \lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}}. \end{aligned} \quad (6.33)$$

Thanks to (6.31) and (6.33), we have

$$t^{\frac{d(p-1)}{p}+l+1+b} \sup_{\alpha} \frac{\|\nabla_x^{l+1} \mathcal{T}_{\alpha}^2[E_i, \partial_i \mu](t)\|_{L_x^p}}{|\alpha|^b} \lesssim_C 1. \quad (6.34)$$

1.3) Estimate of terms in $\mathcal{T}_{\alpha}^3[E_i, \partial_i \mu]$.

Firstly, we have

$$\begin{aligned} |\partial_x^{\beta_1} F_3(x)| &\lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_1|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_4(x)) (\partial^{\beta_3} E_i)(s, Z_4(x)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_4(x-\alpha)) (\partial^{\beta_3} E_i)(s, Z_1(x-\alpha)) \right| \\ &\leq \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_1|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_4(x)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_4(x-\alpha)) \right| |(\nabla^{|\beta_3|} E_i)(s, Z_4(x))| \\ &\quad + \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} Z_4(x-\alpha)| \left| (\nabla^{|\beta_3|} E_i)(s, Z_4(x)) - (\nabla^{|\beta_3|} E_i)(s, Z_1(x-\alpha)) \right|. \end{aligned}$$

Then, by Proposition 4.1 and 4.4, for $|\beta_1| \geq 1$

$$|\partial_x^{\beta_1} F_3(x)| \lesssim_{b,\delta_0,c} \frac{|\alpha|^b}{t^{|\beta_1|+b}} \sum_{1 \leq |\beta_3| \leq \beta_1}^{|\beta_3|} \left(\left[\langle s \rangle^{|\beta_3|-d+1} \mathbf{1}_{(|\beta_3|,|\beta_1|) \neq (1,l+1)} + \langle s \rangle^{-d+2+\delta_0+b} \mathbf{1}_{(|\beta_3|,|\beta_1|) = (1,l+1)} \right] |(\nabla^{|\beta_3|} E_i)(s, Z_4)| \right. \\ \left. + \langle s \rangle^{|\beta_3|-(d-1-\delta_0)b-1} \mathbf{1}_{|\beta_3| < |\beta_1|} \sup_z \frac{|(\delta_z \nabla^{|\beta_3|} E_i)(s, Z_4)|}{|z|^b} \right).$$

And if $|\beta_1| = 0$, we can obtain

$$|F_3(x)| \leq \left| \frac{\alpha}{t} \right|^b \frac{|\delta_z E_i(s, \cdot)|}{|z|^b} \|\nabla_v Y_{s,t}\|_{L^\infty}^b \lesssim_{b,\delta_0,c} \left| \frac{\alpha}{t} \right|^b \sup_z \frac{|\delta_z E_i(s, \cdot)|}{|z|^b} \langle s \rangle^{-b(d-1-\delta_0)}.$$

Now we estimate the L^1 and L^∞ norm of $\partial_x^{\beta_1} F_3(x)$ as follows

$$\|\partial_x^{\beta_1} F_3(x)\|_{L_x^p} \lesssim_{b,\delta_0,c} \frac{|\alpha|^b}{t^{|\beta_1|+b}} \sum_{1 \leq |\beta_3| \leq \beta_1} \left[\langle s \rangle^{|\beta_3|-d+1} \mathbf{1}_{(|\beta_3|,|\beta_1|) \neq (1,l+1)} + \langle s \rangle^{-d+2+\delta_0+b} \mathbf{1}_{(|\beta_3|,|\beta_1|) = (1,l+1)} \right] \|(\nabla^{|\beta_3|} E_i)(s)\|_{L^p} \\ + \frac{|\alpha|^b}{t^{|\beta_1|+b}} \left(\sum_{1 \leq |\beta_3| \leq \beta_1} \langle s \rangle^{|\beta_3|-(d-1-\delta_0)b-1} \mathbf{1}_{|\beta_3| < |\beta_1|} \|\nabla^{|\beta_1|} E_i(s)\|_{\dot{F}_{p,\infty}^b} + \|E_i(s)\|_{\dot{F}_{p,\infty}^b} \langle s \rangle^{-b(d-1-\delta_0)} \right) \\ \lesssim_C \frac{|\alpha|^b}{t^{|\beta_1|+b}} \left(\frac{1}{\langle s \rangle^{d-\delta_0+\frac{d(p-1)}{p}}} + \frac{1}{\langle s \rangle^{d-b-2\delta_0+\frac{d(p-1)}{p}}} + \frac{1}{\langle s \rangle^{-\delta_0+(d-1-\delta_0)b+\frac{d(p-1)}{p}}} \right) \\ \lesssim_C \frac{|\alpha|^b}{t^{|\beta_1|+b}} \langle s \rangle^{\delta_0-(d-1-\delta_0)b-\frac{d(p-1)}{p}}. \quad (6.35)$$

On the other hand, for $|\beta_2| \geq 0$ we have

$$|\partial_x^{\beta_2} \eta_{1,1}(x)| \lesssim_{t^{|\beta_2|}} \sum_{\substack{1 \leq |\beta_3| \leq |\beta_2|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|} = |\beta_2|}} \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} Z_2(x)| |\partial^{\beta_3} \partial_i \mu|(Z_2(x)) \lesssim_C \frac{1}{t^{|\beta_2|}} \frac{1}{\langle \frac{x-w}{t} \rangle^N}.$$

Since $(d-1-\delta_0)b-\delta_0 > 1$, $\delta_0 \leq 2-\sqrt{3}$, we apply (6.35) to get

$$\|\mathcal{T}_\alpha^3[E_i, \partial_i \mu](t)\|_{L^p} \lesssim_{b,\delta_0,c,c'} \int_0^{t/2} \frac{1}{t^{|\beta_2|}} \|\partial_x^{\beta_1} F_3(x)\|_{L_x^1} \frac{ds}{(t-s)^{\frac{d(p-1)}{p}}} + \int_{t/2}^t \int_{\mathbb{R}^d} \frac{1}{t^{|\beta_2|}} \|\partial_x^{\beta_1} F_3(x)\|_{L_x^p} \frac{1}{\langle v \rangle^N} dv ds \\ \lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}} \left(\int_0^{t/2} + \int_{t/2}^t \right) \frac{1}{\langle s \rangle^{-\delta_0+(d-1-\delta_0)b}} ds \lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}}.$$

Thus, we have

$$t^{\frac{d(p-1)}{p}+l+1+b} \sup_\alpha \frac{\|\mathcal{T}_\alpha^3[E_i, \partial_i \mu](t)\|_{L_x^p}}{|\alpha|^b} \lesssim_C 1. \quad (6.36)$$

1.4) Estimate of terms in $\mathcal{T}_\alpha^4[E_i, \partial_i \mu]$.

Firstly, we have for $|\beta_1| \geq 1$

$$|\partial_x^{\beta_1} F_{2,1}(x-\alpha)| \lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|-1, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|} = |\beta_1|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_1(x-\alpha)) (\partial^{\beta_3} E_i)(s, Z_1(x-\alpha)) \right| \\ + \sum_{|\vartheta_1| + \dots + |\vartheta_{|\beta_1|} = |\beta_1|, |\vartheta_i| > 0}^{|\beta_1|} \left| \prod_{j=1}^{|\beta_1|} (\partial_x^{\vartheta_j} Z_1(x-\alpha)) (\partial^{\beta_1} E_i)(s, Z_1(x-\alpha)) \right| \\ \lesssim_{b,\delta_0,c} \frac{\|(\rho, U)\|_{S_a^{\varepsilon_0}}}{t^{|\beta_1|}} \sum_{1 \leq |\beta_3| \leq |\beta_1|} \langle s \rangle^{|\beta_3|-1} \mathbf{1}_{|\beta_3| < |\beta_1|} |(\partial^{\beta_3} E_i)(s, Z_1(x-\alpha))|,$$

and for $|\beta_1| = 0$, we obtain

$$|F_{2,1}(x-\alpha)| = |E_i(s, Z_1(x-\alpha))|.$$

Then, we estimate the L^1 and L^∞ norm of $\partial_x^{\beta_1} F_{2,1}$

$$\begin{aligned} \|\partial_x^{\beta_1} F_{2,1}\|_{L^p} &\lesssim_{b,\delta_0,c} \frac{\mathbf{1}_{|\beta_1|\geq 1}}{t^{|\beta_1|}} \sum_{\substack{1\leq|\beta_3|\leq|\beta_1| \\ |\beta_3|-1\leq|\beta_3|<|\beta_1|}} \langle s \rangle^{|\beta_3|-1} \|(\partial^{\beta_3} E_i)(s, Z_1(x-\alpha))\|_{L^p} + \mathbf{1}_{|\beta_1|=0} \|E_i(s, Z_1(x-\alpha))\|_{L^p} \\ &\lesssim_C \frac{1}{t^{|\beta_1|}} \langle s \rangle^{\delta_0-1-\frac{d(p-1)}{p}} \lesssim_C \frac{1}{t^{|\beta_1|}} \langle s \rangle^{\delta_0-1-\frac{d(p-1)}{p}}. \end{aligned} \quad (6.37)$$

Meanwhile, for $|\beta_2| \geq 1$ we have

$$\begin{aligned} |\partial_x^{\beta_2} \eta_3(x)| &\lesssim \sum_{\substack{1\leq|\beta_3|\leq|\beta_2|, |\vartheta_i|>0 \\ |\vartheta_1|+\dots+|\vartheta_{|\beta_3|}|=|\beta_2|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_2(x)) (\partial^{\beta_3} \partial_i \mu)(s, Z_2(x)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_2(x-\alpha)) (\partial^{\beta_3} \partial_i \mu)(s, Z_3(x)) \right| \\ &\leq \sum_{\substack{1\leq|\beta_3|\leq|\beta_2|, |\vartheta_i|>0 \\ |\vartheta_1|+\dots+|\vartheta_{|\beta_3|}|=|\beta_2|}} \left(\left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_2(x)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} Z_2(x-\alpha)) \right| |(\nabla^{|\beta_3|} \partial_i \mu)(s, Z_2(x))| \right. \\ &\quad \left. + \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} Z_2(x-\alpha)| |(\nabla^{|\beta_3|} \partial_i \mu)(s, Z_2(x)) - (\nabla^{|\beta_3|} \partial_i \mu)(s, Z_3(x))| \right) \\ &\lesssim_C \frac{|\alpha|^b \|(\rho, U)\|_{S_{a^0}^{\varepsilon_0}} \langle s \rangle^{-d+2}}{t^{|\beta_2|+b} \langle \frac{x-w}{t} \rangle^N} + \frac{\min\left\{\frac{|\alpha|}{t}, 1\right\} \langle s \rangle^{-d+2}}{t^{|\beta_2|} \langle \frac{x-w}{t} \rangle^N} \lesssim_C \frac{|\alpha|^b \langle s \rangle^{-d+2}}{t^{|\beta_2|+b} \langle \frac{x-w}{t} \rangle^N}. \end{aligned}$$

For $|\beta_2| = 0$, we have

$$\begin{aligned} |\eta_3(x)| &= |(\partial_i \mu)(Z_2(x)) - (\partial_i \mu)(Z_3(x))| = \frac{|\alpha|^b |\partial_i \mu(Z_2(x)) - \partial_i \mu(Z_3(x))|}{t^b |Z_2(x) - Z_3(x)|^b} \left| \frac{Z_2(x) - Z_3(x)}{|\alpha|/t} \right|^b \\ &\lesssim_C \frac{|\alpha|^b \|(\rho, U)\|_{S_{a^0}^{\varepsilon_0}}^b \langle s \rangle^{(\delta_0-d)b}}{t^b \langle \frac{x-w}{t} \rangle^N} \lesssim_C \frac{|\alpha|^b \langle s \rangle^{(\delta_0-d)b}}{t^b \langle \frac{x-w}{t} \rangle^N}. \end{aligned}$$

Thus, with (6.37), we have

$$\begin{aligned} \|\mathcal{T}_\alpha^4[E_i, \partial_i \mu](t, x)\|_{L^p} &\lesssim_C \sum_{|\beta_1|+|\beta_2|=l+1} \sum_{|\beta_1|=0}^l \int_0^{t/2} \frac{|\alpha|^b \langle s \rangle^{-d+2} \|\partial_x^{\beta_1} F_{2,1}(x)\|_{L_x^1}}{t^{|\beta_2|+b} \langle s \rangle^{-d+2}} \frac{ds}{(t-s)^{\frac{d(p-1)}{p}}} \\ &\quad + \int_0^{t/2} \frac{|\alpha|^b \langle s \rangle^{(\delta_0-d)b}}{t^b} \|\partial_x^{l+1} F_{2,1}(x)\|_{L_x^1} \frac{ds}{(t-s)^{\frac{d(p-1)}{p}}} \\ &\quad + \sum_{|\beta_1|+|\beta_2|=l+1} \sum_{|\beta_1|=0}^l \int_0^{t/2} \int_{\mathbb{R}^d} \frac{|\alpha|^b \langle s \rangle^{-d+2}}{t^{|\beta_2|+b} \langle v \rangle^N} \|\nabla_x^{\beta_1} F_{2,1}(x)\|_{L_x^p} dv ds \\ &\quad + \int_0^{t/2} \int_{\mathbb{R}^d} \frac{|\alpha|^b \langle s \rangle^{(\delta_0-d)b}}{t^b \langle v \rangle^N} \|\partial_x^{l+1} F_{2,1}(x)\|_{L_x^p} dv ds \\ &\lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}} \left(\int_0^{t/2} + \int_{t/2}^t \right) \left(\frac{1}{\langle s \rangle^{d-1-\delta_0}} + \frac{1}{\langle s \rangle^{(d-\delta_0)b+1-\delta_0}} \right) ds \\ &\lesssim_C \frac{|\alpha|^b}{t^{\frac{d(p-1)}{p}+l+1+b}}. \end{aligned}$$

This implies

$$t^{\frac{d(p-1)}{p}+l+1+b} \sup_\alpha \frac{\|\mathcal{T}_\alpha^4[E_i, \partial_i \mu](t)\|_{L_x^p}}{|\alpha|^b} \lesssim_C 1. \quad (6.38)$$

Hence, combining (6.30), (6.34), (6.36) with (6.38) we obtain

$$t^{\frac{d(p-1)}{p}+l+1+b} \|\nabla_x^{l+1} \mathcal{T}[E_i, \partial_i \mu](t)\|_{\dot{B}_{p,\infty}^b} \lesssim_C 1. \quad (6.39)$$

(2) We now consider $t \leq 1$. Denote

$$\begin{aligned}\bar{F}_{1,1}(x) &= \delta_\alpha E_i(s, \cdot)(X_{s,t}(x, v)), \quad \bar{\eta}_{1,1}(x) = \partial_i \mu(V_{s,t}(x, v)), \\ \bar{F}_{1,2}(x) &= \delta_\alpha E_i(s, \cdot)(x - (t-s)v), \quad \bar{\eta}_{1,2}(x) = \partial_i \mu(v), \\ \bar{F}_{2,1}(x) &= E_i(s, X_{s,t}(x, v)), \quad \bar{F}_3(x) = E_i(s, X_{s,t}(x - \alpha, v)) - E_i(s, X_{s,t}(x, v) - \alpha), \\ \bar{\eta}_3(x) &= \partial_i \mu(V_{s,t}(x - \alpha, v)) - \partial_i \mu(V_{s,t}(x, v)).\end{aligned}$$

Thus, then we have,

$$\delta_\alpha \mathcal{T}[E_i, \partial_i \mu](t, x) = \sum_{j=1}^3 \bar{\mathcal{T}}_\alpha^j[E_i, \partial_i \mu](t, x),$$

with

$$\begin{aligned}\bar{\mathcal{T}}_\alpha^1[E_i, \partial_i \mu](t, x) &= - \int_0^t \int_{\mathbb{R}^d} (\bar{F}_{1,1}(x) \bar{\eta}_{1,1}(x) - \bar{F}_{1,2}(x) \bar{\eta}_{1,2}(x)) dv ds, \\ \bar{\mathcal{T}}_\alpha^2[E_i, \partial_i \mu](t, x) &= \int_0^t \int_{\mathbb{R}^d} \bar{F}_3(x) \bar{\eta}_{1,1}(x) dv ds, \\ \bar{\mathcal{T}}_\alpha^3[E_i, \partial_i \mu](t, x) &= \int_0^t \int_{\mathbb{R}^d} \bar{F}_{2,1}(x - \alpha) \bar{\eta}_3(x) dv ds.\end{aligned}$$

Hence, we obtain

$$|\nabla_x^{l+1} \delta_\alpha \bar{\mathcal{T}}^j[E_i, \partial_i \mu](t, x)| \lesssim \sum_{|\beta_1| + |\beta_2| = l+1} \left| \bar{\mathcal{T}}_\alpha^j[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) \right|,$$

where

$$\begin{aligned}\bar{\mathcal{T}}_\alpha^1[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= - \int_0^t \int_{\mathbb{R}^d} (\partial_x^{\beta_1} \bar{F}_{1,1}(x) \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) - \partial_x^{\beta_1} \bar{F}_{1,2}(x) \partial_x^{\beta_2} \bar{\eta}_{1,2}(x)) dv ds, \\ \bar{\mathcal{T}}_\alpha^2[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= \int_0^t \int_{\mathbb{R}^d} \partial_x^{\beta_1} \bar{F}_3(x) \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) dv ds, \\ \bar{\mathcal{T}}_\alpha^3[E_i, \partial_i \mu, \beta_1, \beta_2](t, x) &= \int_0^t \int_{\mathbb{R}^d} \partial_x^{\beta_1} \bar{F}_{2,1}(x - \alpha) \partial_x^{\beta_2} \bar{\eta}_3(x) dv ds.\end{aligned}$$

2.1) Estimate of $\bar{\mathcal{T}}_\alpha^1[E_i, \partial_i \mu]$.

we consider the following line

$$\begin{aligned}& \left| \partial_x^{\beta_1} \bar{F}_{1,1}(x) \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) - \partial_x^{\beta_1} \bar{F}_{1,2}(x) \partial_x^{\beta_2} \bar{\eta}_{1,2}(x) \right| \\ & \leq \left| \partial_x^{\beta_1} \bar{F}_{1,1}(x) - \partial_x^{\beta_1} \delta_\alpha E_i(s, \cdot)(X_{s,t}(x, v)) \right| \left| \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) - (\partial_x^{\beta_2} \partial_i \mu)(V_{s,t}(x, v)) \right| \\ & \quad + \left| \partial_x^{\beta_1} \bar{F}_{1,1}(x) - \partial_x^{\beta_1} \delta_\alpha E_i(s, \cdot)(X_{s,t}(x, v)) \right| \left| (\partial_x^{\beta_2} \partial_i \mu)(V_{s,t}(x, v)) \right| \\ & \quad + \left| \partial_x^{\beta_1} \delta_\alpha E_i(s, \cdot)(X_{s,t}(x, v)) \right| \left| \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) - (\partial_x^{\beta_2} \partial_i \mu)(V_{s,t}(x, v)) \right| \\ & \quad + \left| \partial_x^{\beta_1} \delta_\alpha E_i(s, \cdot)(X_{s,t}(x, v)) (\partial_x^{\beta_2} \partial_i \mu)(V_{s,t}(x, v)) - \bar{F}_{1,2}(x) \bar{\eta}_{1,2}(x) \right|.\end{aligned}$$

Thanks to Lemma 6.5 we have

$$\begin{aligned}& \left| \partial_x^{\beta_1} \bar{F}_{1,1}(x) - \partial_x^{\beta_1} \delta_\alpha F(s, \cdot)(X_{s,t}(x, v)) \right| \\ & \lesssim \mathbf{1}_{|\beta_1| \geq 1} |\nabla^{|\beta_1|} \delta_\alpha F(s, \cdot)(X_{s,t}(x, v))| |(\nabla_x Y_{s,t})| (1 + |\nabla_x Y_{s,t}|)^{|\beta_1| - 1} \\ & \quad + \mathbf{1}_{|\beta_1| \geq 1} \sum_{m=1}^{|\beta_1| - 1} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_1| - j} |\nabla^m \delta_\alpha F(s, \cdot)(X_{s,t}(x, v))| (1 + |\nabla_x Y_{s,t}|)^j |\nabla_x^i Y_{s,t}|^{\frac{|\beta_1| - j}{i}} \\ & \lesssim C \mathbf{1}_{|\beta_1| \geq 1} \sum_{m=1}^{|\beta_1|} |\nabla^m \delta_\alpha F(s, \cdot)(X_{s,t}(x, v))| \langle s \rangle^{-d + \delta_0}.\end{aligned}$$

Then, for $|\beta_2| \geq 1$

$$\begin{aligned}& \left| \partial_x^{\beta_2} \bar{\eta}_{1,1}(x) - (\partial_x^{\beta_2} \partial_i \mu)(V_{s,t}(x, v)) \right| \\ & \lesssim \mathbf{1}_{|\beta_2| \geq 1} \left(|\nabla^{|\beta_2|} \partial_i \mu(V_{s,t}(x, v))| |\nabla_x W_{s,t}|^{|\beta_2|} + \sum_{m=1}^{|\beta_2|} \sum_{j=0}^{m-1} \sum_{i=2}^{|\beta_2| - j} |\nabla^m \partial_i \mu(V_{s,t}(x, v))| |\nabla_x W_{s,t}|^j |\nabla_x^i W_{s,t}|^{\frac{|\beta_2| - j}{i}} \right) \\ & \stackrel{(4.4), (4.4)}{\lesssim} C \mathbf{1}_{|\beta_2| \geq 1} \sum_{m=1}^{|\beta_2|} |\nabla^m \partial_i \mu(V_{s,t}(x, v))| \langle s \rangle^{-d-1+\delta_0} \|(\rho, U)\|_{S_a^\varepsilon} \lesssim C \frac{\mathbf{1}_{|\beta_2| \geq 1}}{\langle v \rangle^N} \langle s \rangle^{-1-d+\delta_0}.\end{aligned}$$

Then, we have for $p = 1, \infty$

$$\begin{aligned} \left\| \overline{\mathcal{T}}_\alpha^1[E_i, \partial_i \mu] \right\|_{L_x^p} &\lesssim_C \sum_{|\beta_1|=0}^{l+1} \sum_{m=0}^{|\beta_1|} \int_0^t \int_{\mathbb{R}^d} \|\nabla^m \delta_\alpha E_i(s)\|_{L^p} \langle s \rangle^{-d+\delta_0} \frac{dv ds}{\langle v \rangle^N} + \sum_{|\beta_1|+|\beta_2|=l+1} \|\mathcal{T}[\partial_x^{\beta_1} \delta_\alpha E_i, \partial_x^{\beta_2} \partial_i \mu]\|_{L_x^p} \\ &\lesssim_C |\alpha|^b \sum_{|\beta_1|+1}^{l+1} \sum_{m=0}^{|\beta_1|} \left(\int_0^t \frac{1}{\langle s \rangle^{m+1-\delta_0}} \langle s \rangle^{-d+\delta_0} ds + \|\nabla^{|\beta_1|} E_i(s)\|_{\dot{B}_{p,\infty}^b} \right) \lesssim_C |\alpha|^b. \end{aligned} \quad (6.40)$$

2.2) Estimate of $\overline{\mathcal{T}}_\alpha^2[E_i, \partial_i \mu]$.

Firstly, we consider $\partial_x^{\beta_1} \overline{F}_3(x)$ as follows. If $|\beta_1| \geq 1$, we get

$$\begin{aligned} |\partial_x^{\beta_1} \overline{F}_3(x)| &\lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_1|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} X_{s,t}(x, v)) (\partial^{\beta_3} E_i)(s, X_{s,t}(x, v) - \alpha) \right. \\ &\quad \left. - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} X_{s,t}(x - \alpha, v)) (\partial^{\beta_3} E_i)(s, X_{s,t}(x - \alpha, v)) \right| \\ &\leq \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_1|}} \left(\left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} X_{s,t}(x, v)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} X_{s,t}(x - \alpha, v)) \right| \left| (\nabla^{|\beta_3|} E_i)(s, X_{s,t}(x, v) - \alpha) \right| \right. \\ &\quad \left. + \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} X_{s,t}(x - \alpha, v)| \left| (\nabla^{|\beta_3|} E_i)(s, X_{s,t}(x, v) - \alpha) - (\nabla^{|\beta_3|} E_i)(s, X_{s,t}(x - \alpha, v)) \right| \right) \\ &\lesssim_C |\alpha|^b \sum_{1 \leq |\beta_3| \leq |\beta_1|} \left(\langle s \rangle^{-d+\delta_0} |\nabla^{|\beta_3|} E_i|(s, X_{s,t}(x, v) - \alpha) + \langle s \rangle^{-(d-\delta_0)b} \sup_z \frac{|\delta_z \nabla^{|\beta_3|} E_i(s, \cdot)(X_{s,t}(x - \alpha, v))|}{|z|^b} \right). \end{aligned}$$

If $|\beta_1| = 0$, we can yield

$$|\overline{F}_3(x)| \leq \frac{|\overline{F}_3(x)|}{|\delta_\alpha Y_{s,t}(x, v)|^b} \|\nabla_x Y_{s,t}\|_{L^\infty}^b \lesssim_{b, \delta_0, c} |\alpha|^b \sup_z \frac{|\delta_z E_i(s, \cdot)(X_{s,t}(x - \alpha, v))|}{|z|^b} \langle s \rangle^{-(d-\delta_0)b}.$$

Then, we consider another term. For $|\beta_2| \geq 0$,

$$|\partial_x^{\beta_2} \overline{\eta}_{1,1}(x)| \lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_2|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_2|}} \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} W_{s,t}(x - vt, v)| |\partial^{\beta_3} \partial_i \mu|(V_{s,t}(x, v)) \lesssim_{b, \delta_0, c, c'} \frac{1}{\langle v \rangle^N}.$$

Hence, we obtain

$$\begin{aligned} \|\overline{\mathcal{T}}_\alpha^2[E_i, \partial_i \mu](t)\|_{L^p} &\lesssim_C |\alpha|^b \int_0^t \int_{\mathbb{R}^d} \sum_{\substack{1 \leq |\beta_1| \leq l+1 \\ 1 \leq |\beta_3| \leq |\beta_1|}} \left(\langle s \rangle^{-d+\delta_0} \|(\nabla^{|\beta_3|} E_i)(s)\|_{L^p} + \langle s \rangle^{-(d-\delta_0)b} \|\nabla^{|\beta_3|} E_i\|_{\dot{F}_{p,\infty}^b} \right) \frac{dv ds}{\langle v \rangle^N} \\ &\quad + |\alpha|^b \int_0^t \int_{\mathbb{R}^d} \|E_i(s)\|_{\dot{F}_{p,\infty}^b} \langle s \rangle^{-(d-\delta_0)b} \frac{1}{\langle v \rangle^N} dv ds \\ &\lesssim_C |\alpha|^b \int_0^t \left(\frac{1}{\langle s \rangle^{d-2\delta_0+2}} + \frac{1}{\langle s \rangle^{(d-\delta_0)b+2-\delta_0}} \right) ds \lesssim_C |\alpha|^b. \end{aligned} \quad (6.41)$$

2.3) Estimate of $\overline{\mathcal{T}}_\alpha^3[E_i, \partial_i \mu]$.

Firstly, we consider $\partial_x^{\beta_1} \overline{F}_{2,1}(x - \alpha)$, and when $|\beta_1| \geq 1$,

$$\begin{aligned} |\partial_x^{\beta_1} \overline{F}_{2,1}(x - \alpha)| &\lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_1|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_1|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} X_{s,t}(x - \alpha, v)) (\partial^{\beta_3} E_i)(s, X_{s,t}(x - \alpha, v)) \right| \\ &\lesssim_C \sum_{1 \leq |\beta_3| \leq |\beta_1|} \left| (\partial^{\beta_3} E_i)(s, X_{s,t}(x - \alpha, v)) \right|. \end{aligned}$$

Combining above with the definition of $\partial_x^{\beta_1} \overline{F}_{2,1}(x - \alpha)$, we get, for any $|\beta_1| \geq 0$

$$|\partial_x^{\beta_1} \overline{F}_{2,1}(x - \alpha)| \lesssim_C \sum_{0 \leq |\beta_3| \leq |\beta_1|} |(\partial^{\beta_3} E_i)(s, X_{s,t}(x - \alpha, v))|.$$

Then, we compute the other term. For $|\beta_2| \geq 1$,

$$\begin{aligned} |\partial_x^{\beta_2} \overline{\eta}_3(x)| &\lesssim \sum_{\substack{1 \leq |\beta_3| \leq |\beta_2|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_2|}} \left| \prod_{j=1}^{|\beta_3|} \partial_x^{\vartheta_j} W_{s,t}(x - vt, v) (\partial^{\beta_3} \partial_i \mu)(V_{s,t}(x, v)) \right. \\ &\quad \left. - \prod_{j=1}^{|\beta_3|} \partial_x^{\vartheta_j} W_{s,t}(x - \alpha - vt, v) (\partial^{\beta_3} \partial_i \mu)(V_{s,t}(x - \alpha, v)) \right| \\ &\leq \sum_{\substack{1 \leq |\beta_3| \leq |\beta_2|, |\vartheta_i| > 0 \\ |\vartheta_1| + \dots + |\vartheta_{|\beta_3|}| = |\beta_2|}} \left| \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} W_{s,t}(x - vt, v)) - \prod_{j=1}^{|\beta_3|} (\partial_x^{\vartheta_j} W_{s,t}(x - \alpha - vt, v)) \right| |(\nabla^{|\beta_3|} \partial_i \mu)(V_{s,t}(x, v))| \\ &\quad + \prod_{j=1}^{|\beta_3|} |\partial_x^{\vartheta_j} W_{s,t}(x - \alpha - vt, v)| \left| (\nabla^{|\beta_3|} \partial_i \mu)(V_{s,t}(x, v)) - (\nabla^{|\beta_3|} \partial_i \mu)(V_{s,t}(x - \alpha, v)) \right| \\ &\lesssim_C |\alpha|^b \frac{\langle s \rangle^{-d+\delta_0-1}}{\langle v \rangle^N} + \min\{|\alpha|, 1\} \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \alpha \rangle^N} \right) \langle s \rangle^{-d+\delta_0-1} \\ &\lesssim_C |\alpha|^b \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \alpha \rangle^N} \right) \langle s \rangle^{-d+\delta_0-1}. \end{aligned}$$

If $|\beta_2| = 0$, we have

$$|\overline{\eta}_3(x)| \lesssim_C \frac{|\alpha|^b}{\langle v \rangle^N} \langle s \rangle^{-d+\delta_0-1}.$$

Hence, we know that

$$\begin{aligned} \left\| \overline{\mathcal{T}}_\alpha^3[E_i, \partial_i \mu](t, x) \right\|_{L^p} &\lesssim_{b, \delta_0, c, c'} |\alpha|^b \int_0^t \int_{\mathbb{R}^d} \sum_{\substack{0 \leq |\beta_1| \leq l \\ 0 \leq |\beta_3| \leq |\beta_1|}} \|(\partial^{\beta_3} E_i)(s)\|_{L^p} \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \alpha \rangle^N} \right) \langle s \rangle^{-d+\delta_0-1} dv ds \\ &\quad + |\alpha|^b \int_0^t \int_{\mathbb{R}^d} \sum_{1 \leq |\beta_3| \leq l+1} \|(\partial^{\beta_3} E_i)(s)\|_{L^p} \langle s \rangle^{\delta_0-d-1} \frac{1}{\langle v \rangle^N} dv ds \\ &\lesssim_C |\alpha|^b \int_0^t \langle s \rangle^{-d+\delta_0-1} ds \lesssim_C |\alpha|^b. \end{aligned} \tag{6.42}$$

Thus, combining (6.40), (6.41) and (6.42), we have for $0 \leq t \leq 1$

$$\sup_\alpha \frac{\|\overline{\mathcal{T}}_\alpha[E_i, \partial_i \mu](t)\|_{L^\infty}}{|\alpha|^b} \lesssim_C 1, \tag{6.43}$$

In conclusion, thanks to (6.39) and (6.43), we finish the proof. \blacksquare

7 Proof of the Theorem 2.2 and Theorem 2.3

Recall the operator \mathcal{F} as:

$$\mathcal{F}(\rho, U) = \left(\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U) \right),$$

where $\mathcal{F}_1(\rho, U) = G *_{(t,x)} (\mathcal{I}(\rho, U) + \mathcal{R}(\rho, U) + A(U)) + \mathcal{I}(\rho, U) + \mathcal{R}(\rho, U)$, $\mathcal{F}_2(\rho, U) = (1 - \Delta)^{-1}(\rho + A(U))$. Now we start to prove Theorem 2.2 and 2.3.

Proof of Theorem 2.2. Let $\tilde{\varepsilon}_0$ be in Proposition 4.1 and take $\tilde{\varepsilon} \leq \tilde{\varepsilon}_0$. Applying Theorem 3.9 with $\mathbf{f} = \mathcal{I}_{f_0}(\rho, U) + \mathcal{R}(\rho, U)$ and $\mathbf{f} = \mathcal{I}_{f_0}(\rho_1, U_1) - \mathcal{I}_{f_0}(\rho_2, U_2) + \mathcal{R}(\rho_1, U_1) - \mathcal{R}(\rho_2, U_2)$, it is clear that for any

$\|(\rho, U)\|_{S_a^\varepsilon}, \|(\rho_1, U_1)\|_{S_a^\varepsilon}, \|(\rho_2, U_2)\|_{S_a^\varepsilon} \leq \varepsilon \leq \tilde{\varepsilon}$, there exists a constant $M = M(\mathring{c}, d, M^*)$ such that

$$\|\mathcal{F}_1(\rho, U)\|_a \leq M \left(\|(\mathcal{I}_{f_0} + \mathcal{R})(\rho, U)\|_a + \|A(U)\|_a \right), \quad (7.1)$$

$$\|\mathcal{F}_1(\rho_1, U_1) - \mathcal{F}_1(\rho_2, U_2)\|_a \leq M \left(\|(\mathcal{I}_{f_0} + \mathcal{R})(\rho_1, U_1) - (\mathcal{I}_{f_0} + \mathcal{R})(\rho_2, U_2)\|_a + \|A(U_1) - A(U_2)\|_a \right). \quad (7.2)$$

Thanks to Lemma 4.2, Proposition 5.1, Proposition 6.3 and (4.6), we know that there exists a constant $C_{3,1} = C_{3,1}(a, \mathring{c}, d, M^*) > 0$ and $C_{3,2} = C_{3,2}(a, d) > 0$ such that

$$\begin{aligned} \|\mathcal{F}_1(\rho, U)\|_a &\leq C_{3,1} \left(\sum_{p=1, \infty} \|\mathcal{D}^a(f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} + \|(\rho, U)\|_{S_a^\varepsilon}^{1+a} + C_A \varepsilon^{\frac{1}{3}} \|(\rho, U)\|_{S_a^\varepsilon} \right) \\ &\leq C_{3,1} \left(\frac{\varepsilon}{C_2} + \varepsilon^{1+a} + \varepsilon \varepsilon^{\frac{1}{3}} \right) \leq \left(\frac{C_{3,1}}{C_2} + C_{3,1}(1 + C_A) \tilde{\varepsilon}^{\frac{1}{3}} \right) \varepsilon. \end{aligned} \quad (7.3)$$

and

$$\|\mathcal{F}_2(\rho, U)\|_a \leq C_{3,2} \|\rho + A(U)\|_a \leq C_{3,2} \|(\rho, U)\|_{S_a^\varepsilon} \leq C_{3,2} \varepsilon. \quad (7.4)$$

$$\|\mathcal{F}_2(\rho_1, U_1) - \mathcal{F}_2(\rho_2, U_2)\|_a \leq C_{3,2} \|\rho_1 - \rho_2 + A(U_1) - A(U_2)\|_a \leq C_{3,2} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^\varepsilon} \quad (7.5)$$

Moreover, with Proposition 5.4 and Proposition 6.4, there exists a constant $C_4 = C_4(a, \mathring{c}, d, M^*) > 0$ such that

$$\begin{aligned} \|\mathcal{F}_1(\rho_1, U_1) - \mathcal{F}_1(\rho_2, U_2)\|_a &\leq C_4 \|A(U_1) - A(U_2)\|_a \\ &+ C_4 \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^\varepsilon} \left(\sum_{p=1, \infty} \|\mathcal{D}^a(\nabla_{x,v} f_0)\|_{L_x^1 L_v^p \cap L_v^1 L_x^p} + \|(\rho_1, U_1)\|_{S_a^\varepsilon}^a + \|(\rho_2, U_2)\|_{S_a^\varepsilon} \right) \\ &\leq \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^\varepsilon} \left(\frac{C_4}{C_2} + 2C_4 \right) \varepsilon^a + C_4 C_A \varepsilon^{\frac{1}{3}} \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^\varepsilon} \\ &\leq \|(\rho_1 - \rho_2, U_1 - U_2)\|_{S_a^\varepsilon} \left(\frac{C_4}{C_2} + (2 + C_A) C_4 \right) \tilde{\varepsilon}^{\frac{1}{3}}. \end{aligned} \quad (7.6)$$

Combining (7.3), (7.6), (7.5) with (7.4), we obtain that

$$\begin{aligned} \|\mathcal{F}(\rho, U)\|_{S_a^\varepsilon} &= \|\mathcal{F}_1(\rho, U)\|_a + \tilde{\varepsilon}^{\frac{1}{3}} \|\mathcal{F}_2(\rho, U)\|_a \leq \left(\frac{C_{3,1}}{C_2} + (C_{3,1}(1 + C_A) + C_{3,2}) \tilde{\varepsilon}^{\frac{1}{3}} \right) \varepsilon \\ \|\mathcal{F}(\rho_1, U_1) - \mathcal{F}(\rho_2, U_2)\|_{S_a^\varepsilon} &= \|\mathcal{F}_1(\rho_1, U_1) - \mathcal{F}_1(\rho_2, U_2)\|_a + \tilde{\varepsilon}^{\frac{1}{3}} \|\mathcal{F}_2(\rho_1, U_1) - \mathcal{F}_2(\rho_2, U_2)\|_a \\ &\leq \|(\rho_1 - \rho_2, U_1 - U_2)\|_a \left(\frac{C_4}{C_2} + (2 + C_A) C_4 + C_{3,2} \right) \tilde{\varepsilon}^{\frac{1}{3}}. \end{aligned} \quad (7.7)$$

Hence, we choose $C_2 = \max\{1, 8C_{3,1}, 8C_4\}$, and $\tilde{\varepsilon} = \min \left\{ \left(\frac{1}{4(C_{3,1}(1+C_A)+C_{3,2})} \right)^3, \left(\frac{1}{4((2+C_A)C_4+C_{3,2})} \right)^3, \tilde{\varepsilon}_0 \right\}$, we can yield that

$$\|\mathcal{F}(\rho, U)\|_a \leq \varepsilon, \quad \|\mathcal{F}(\rho_1, U_1) - \mathcal{F}(\rho_2, U_2)\|_a \leq \frac{1}{2} \|(\rho_1 - \rho_2, U_1 - U_2)\|_a. \quad (7.8)$$

Therefore, we conclude that \mathcal{F} is a compressed mapping in $\mathfrak{B}(\varepsilon)$, for $\varepsilon \in (0, \tilde{\varepsilon}]$. Now we consider the iterative sequence $(\varrho_n, \mathcal{U}_n) = \mathcal{F}(\varrho_{n-1}, \mathcal{U}_{n-1})$ with $\varrho_1(t, x) = \int f_0(x - tv, v) dv$. and $\mathcal{U}_1 = 0$. From (7.7) we obtain that for $n \in \mathbb{N}$

$$\begin{aligned} \|\mathcal{F}(\varrho_{n+1}, \mathcal{U}_{n+1}) - \mathcal{F}(\varrho_n, \mathcal{U}_n)\|_a &\leq \frac{1}{2} \|(\varrho_n - \varrho_{n-1}, \mathcal{U}_n - \mathcal{U}_{n-1})\|_a \\ &\leq \frac{1}{2^n} \|(\mathcal{F}_1(\varrho_1, \mathcal{U}_1) - \varrho_1, \mathcal{F}_2(\varrho_1, \mathcal{U}_1) - \mathcal{U}_1)\|_a \leq \frac{\varepsilon}{2^{n-1}}. \end{aligned} \quad (7.9)$$

Thus $\{(\varrho_n, \mathcal{U}_n)\}_{n=1}^\infty$ is a Cauchy sequence. We denote

$$(\rho, U) = \left(\sum_{n=2}^\infty (\varrho_n - \varrho_{n-1}) + \varrho_1, \sum_{n=2}^\infty (\mathcal{U}_n - \mathcal{U}_{n-1}) + \mathcal{U}_1 \right),$$

it is clear that $(\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U)) = (\rho, U) \in \mathfrak{B}(\varepsilon)$. Moreover by (7.9), we have

$$\|(\rho - \varrho_1, U - \mathcal{U}_1)\|_a \leq \sum_{n=2}^\infty \|(\varrho_n - \varrho_{n-1}, \mathcal{U}_n - \mathcal{U}_{n-1})\|_a \lesssim_{\mathring{c}, M^*, a} \varepsilon \sum_{n=2}^\infty \varepsilon^{\frac{n-1}{3}} \lesssim_{\mathring{c}, M^*, a} \varepsilon_2^{\frac{4}{3}},$$

Recall the definition of $\mathcal{F}(\rho)$ in (2.7), we know that the fixed point is exactly the solution of (2.5). Now we consider $\|(\rho, U)\|_{1,a}$, note that

$$\begin{aligned}\|\langle s \rangle \nabla U\|_a &= \|\langle s \rangle \nabla \mathcal{F}_2(\rho)\|_a \leq \|\langle s \rangle \nabla \rho\|_a + C_A \|U\|_a \|\langle s \rangle \nabla U\|_a \leq \|\langle s \rangle \nabla \rho\|_a + \varepsilon^{\frac{1}{3}} \|\langle s \rangle \nabla U\|_a, \\ \|\langle s \rangle \nabla \rho\|_a &= \|\langle s \rangle \nabla \mathcal{F}_1(\rho, U)\|_a = \left\| \langle s \rangle G_{*(t,x)} \nabla \left(\mathcal{I}_{f_0}(\rho, U) + \mathcal{R}(\rho, U) + A(U) \right) + \langle s \rangle \nabla \left(\mathcal{I}_{f_0}(\rho, U) + \mathcal{R}(\rho, U) \right) \right\|_a \\ &\leq M \left(\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_a + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_a + \|\langle s \rangle \nabla (A(U))\|_a \right) \\ &\leq M \left(\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_a + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_a + \varepsilon^{\frac{1}{3}} \|\langle s \rangle \nabla U\|_a \right).\end{aligned}$$

Therefore we have

$$\begin{aligned}\|\langle s \rangle \nabla \rho\|_a + \frac{1}{4} \|\langle s \rangle \nabla U\|_a \\ \leq M \left(\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_a + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_a \right) + M \varepsilon^{\frac{1}{3}} \|\langle s \rangle \nabla U\|_a + \frac{1}{4} \|\langle s \rangle \nabla \rho\|_a + \varepsilon^{\frac{1}{3}} \|\langle s \rangle \nabla U\|_a,\end{aligned}\quad (7.10)$$

which implies that

$$\|\langle s \rangle \nabla \rho\|_a + \|\langle s \rangle \nabla U\|_a \leq 8M \left(\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_a + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_a \right).$$

Thanks to (5.2) and (6.14), we get

$$\|\mathcal{I}_{f_0}(\rho, U)\|_{1-\delta_1} + \|\mathcal{R}(\rho, U)\|_{1-\delta_1} \lesssim_{\hat{c}, a, M^*} 1,$$

where $\delta_1 = \min\{\frac{2-\sqrt{3}}{2}, \frac{1-a}{2}\}$. Then we use Proposition 6.25 and (5.3) to obtain

$$\|\mathcal{I}_{f_0}(\rho, U)\|_{1,a} + \|\mathcal{R}(\rho, U)\|_{1,a} \lesssim_{\hat{c}, a, M^*} 1.$$

This implies $\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_a + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_a \lesssim_{\hat{c}, a, M^*} 1$. Consequently, it gives that

$$\|(\rho, U)\|_{1,a} \leq \|\rho\|_a + \|U\|_a + \|\langle s \rangle \nabla \rho\|_a + \|\langle s \rangle \nabla U\|_a \lesssim_{\hat{c}, a, M^*} 1.$$

Proof of Theorem 2.3. From Theorem 2.2 we know that $\|(\rho, U)\|_{S_a^{\tilde{\varepsilon}}} \leq \tilde{\varepsilon}_0$ and $\|(\rho, U)\|_{1,a} \lesssim_{\hat{c}, a, M^*} 1$. Applying the Gagliardo-Nirenberg interpolation inequality, we get $\|(\rho, U)\|_{1-\delta_0} \lesssim_{a, M^*} 1$, where $\delta_0 \in (0, \min\{2-\sqrt{3}, \frac{1-b}{m+1}\})$. Combining Theorem 3.9 and Proposition 5.12 with Proposition 6.6, one has

$$\begin{aligned}\|\langle s \rangle \nabla \rho\|_{1-2\delta_0} &= \|\langle s \rangle \nabla \mathcal{F}_1(\rho, U)\|_{1-2\delta_0} \leq M \left(\|\langle s \rangle \nabla \mathcal{I}_{f_0}(\rho, U)\|_{1-2\delta_0} + \|\langle s \rangle \nabla \mathcal{R}(\rho, U)\|_{1-2\delta_0} + \|\langle s \rangle \nabla A(U)\|_{1-2\delta_0} \right), \\ \|\langle s \rangle \nabla U\|_{1-2\delta_0} &= \|\langle s \rangle \nabla \mathcal{F}_2(\rho, U)\|_{1-2\delta_0} \leq \|\langle s \rangle \nabla \rho\|_a + \|\langle s \rangle \nabla (A(U))\|_{1-2\delta_0} \leq \|\langle s \rangle \nabla \rho\|_a + \varepsilon^{\frac{1}{3}} \|\langle s \rangle \nabla U\|_a, \\ \|\mathcal{I}_{f_0}(\rho)\|_{1,1-2\delta_0} + \|\mathcal{R}(\rho)\|_{1,1-2\delta_0} &\lesssim_{a, M^*, \mathbf{c}, \mathbf{c}'} 1.\end{aligned}$$

Using the same method in (7.10) we have $\|(\rho, U)\|_{1,1-2\delta_0} = \|(\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U))\|_{1,1-2\delta_0} \lesssim_{a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} 1$. Note that

$$\|\langle s \rangle^2 \nabla^2 (A(U))\|_{\dot{B}_{p,\infty}^{1-3\delta_0}} \leq C_{A,m} \|U\|_{1,1-3\delta_0} + C_A \varepsilon^{\frac{2}{3}} \|\langle s \rangle^2 \nabla^2 U\|_{\dot{B}_{p,\infty}^{1-3\delta_0}}$$

We do one more step to obtain

$$\begin{aligned}\|(\rho, U)\|_{2,1-3\delta_0} &\leq \|(\rho, U)\|_a + \|\nabla^2(\rho, U)\|_{1-3\delta_0} = \|(\rho, U)\|_a + \|\nabla^2(\mathcal{F}_1(\rho, U), \mathcal{F}_2(\rho, U))\|_{1-3\delta_0} \\ &\lesssim_{\hat{c}, a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} \|(\rho, U)\|_a + \|\mathcal{I}_{f_0}(\rho, U)\|_{2,1-3\delta_0} + \|\mathcal{R}(\rho, U)\|_{2,1-3\delta_0} + \|(\rho, U)\|_{2,1-3\delta_0} + \|U\|_{1,1-3\delta_0} \\ &\lesssim_{\hat{c}, a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} 1.\end{aligned}$$

Then, with m times iteration, we have

$$\|(\rho, U)\|_{m,1-(m+1)\delta_0} \lesssim_{\hat{c}, a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} 1.$$

Note that $1 - (m+1)\delta_0 > b$, thus by the Gagliardo-Nirenberg interpolation inequality again to get

$$\|(\rho, U)\|_{m,b} \lesssim_{\hat{c}, a, M^*, \mathbf{c}, \mathbf{c}', C_{A,m}} 1,$$

which completes the proof. \blacksquare

8 Appendix

Lemma 8.1 Assume $\mathcal{H} \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ and $\varphi(x, v)$ satisfies $\|\nabla_{x,v}\varphi(x, v)\|_{L^\infty_{x,v}} \leq \frac{1}{2}$, then for $p = 1, \infty$, $0 \leq s \leq t$ and $t \geq 0$ we have

$$\left\| \int_{\mathbb{R}^d} \mathcal{H}(\varphi(x, v) + x - (t-s)v) \frac{dv}{\langle v \rangle^N} \right\|_{L^p_x} \lesssim \|\mathcal{H}\|_{L^p}. \quad (8.1)$$

Moreover, for $0 \leq s \leq \frac{t}{2}$ and $t \geq 1$, we also get

$$\left\| \int_{\mathbb{R}^d} \mathcal{H}(\varphi(x, v) + x - (t-s)v) \frac{dv}{\langle v \rangle^N} \right\|_{L^p_x} \lesssim \frac{1}{t^{\frac{d(p-1)}{p}}} \|\mathcal{H}\|_{L^1}. \quad (8.2)$$

Proof. We can directly prove the first inequality as follows

$$\left\| \int_{\mathbb{R}^d} \mathcal{H}(\varphi(x, v) + x - (t-s)v) \frac{dv}{\langle v \rangle^N} \right\|_{L^p_x} \leq \|\mathcal{H}\|_{L^p} \left(1 + \|\nabla_{x,v}\varphi(x, v)\|_{L^\infty_{x,v}}\right) \lesssim \|\mathcal{H}\|_{L^p}.$$

Furthermore, we change variable $\zeta = x - (t-s)v$ and obtain for $0 \leq s \leq \frac{t}{2}$,

$$\begin{aligned} \left\| \int_{\mathbb{R}^d} \mathcal{H}(\varphi(x, v) + x - (t-s)v) \frac{dv}{\langle v \rangle^N} \right\|_{L^\infty_x} &= \frac{1}{(t-s)^d} \left\| \int_{\mathbb{R}^d} \mathcal{H}\left(\varphi\left(x, \frac{x-\zeta}{t-s}\right) + \zeta\right) \frac{d\zeta}{\langle \frac{x-\zeta}{t-s} \rangle^N} \right\|_{L^\infty_x} \\ &\lesssim \frac{1}{t^d} \left[\sup_{x, \zeta} \left(\left| (\nabla_v \varphi)\left(x, \frac{x-\zeta}{t-s}\right) \right| (t-s)^{-d} + 1 \right)^{-1} \right] \|\mathcal{H}\|_{L^1} \lesssim \frac{1}{t^d} \|\mathcal{H}\|_{L^1}. \end{aligned}$$

■

Remark 8.2 Similarly, under the same assumption of \mathcal{H} and φ one has

$$\left\| \int_{\mathbb{R}^d} |\mathcal{H}(\varphi(x, v) + x - (t-s)v)| \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right) dv \right\|_{L^p_x} \lesssim \|\mathcal{H}\|_{L^p}, \quad (8.3)$$

and for $0 \leq s \leq \frac{t}{2}$, $t \geq 1$,

$$\left\| \int_{\mathbb{R}^d} |\mathcal{H}(\varphi(x, v) + x - (t-s)v)| \left(\frac{1}{\langle v \rangle^N} + \frac{1}{\langle v - \frac{\alpha}{t} \rangle^N} \right) dv \right\|_{L^p_x} \lesssim \frac{1}{t^{\frac{d(p-1)}{p}}} \|\mathcal{H}\|_{L^1}. \quad (8.4)$$

References

- [1] A. Arsenév. Global Existence of a Weak Solution of Vlasov system of equations, *U.S.S.R. Comp. Math. Math. Phys.*, 15 (1975), pp. 131-143.
- [2] C. Bardos and P. Degond. Global existence for the Vlasov-Poisson equation in 3 space variables with small initial data. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 2(2):101-118, 1985.
- [3] C. Bardos, F. Golse, T. Nguyen, and R. Sentis. The Maxwell-Boltzmann approximation for ion kinetic modeling. *Physics. D*, 376/377:94-107, 2018.
- [4] J. Batt. Global Symmetric Solutions of the Initial Value Problem of Stellar Dynamics, *Journal of Differential Equations*, 25 (1977), pp. 342-364.
- [5] J. Bedrossian, N. Masmoudi and C. Mouhot. Linearized wave-damping structure of Vlasov-Poisson in \mathbb{R}^3 . *arXiv:2007.08580*, 2020.
- [6] J. Bedrossian, N. Masmoudi and C. Mouhot. Landau damping in finite regularity for unconfined systems with screened interactions. *Communications on Pure and Applied Mathematics*, 71(3): 537-576, 2018.
- [7] J. Bedrossian, N. Masmoudi and C. Mouhot. Landau damping: paraproducts and Gevrey regularity. *Annals of PDE*, 2(1): 1-71, 2016.
- [8] F. Bouchut. Global weak solution of the Vlasov-Poisson system for small electrons mass. *Comm. Partial Differential Equations*, 16(8-9):1337-1365, 1991.
- [9] K. Chen and Q.H. Nguyen. The Peskin Problem with $B^\infty_{\infty, \infty}$ initial data, submitted, *arXiv:2107.13854*.

- [10] S.-H. Choi, S.-Y. Ha and H. Lee. Dispersion estimates for the two-dimensional Vlasov–Yukawa system with small data. *Journal of Differential Equations*, 250(1):515–550, 2011.
- [11] R. T. Glassey. The Cauchy problem in kinetic theory. *Society for Industrial and Applied Mathematics, Philadelphia, PA*, 1996.
- [12] E. Grenier, T. T. Nguyen and I. Rodnianski. Landau damping for analytic and Gevrey data. *arXiv:2004.05979*, 2020.
- [13] M. Griffin-Pickering and M. Iacobelli. Global well-posedness in 3-dimensions for the Vlasov–Poisson system with massless electrons. *arXiv:1810.06928*.
- [14] Y. Guo and B. Pausader. Global Smooth Ion Dynamics in the Euler-Poisson System. *Commun. Math. Phys.* 303, 89–125 (2011). <https://doi.org/10.1007/s00220-011-1193-1>.
- [15] D. Han-Kwan. Quasineutral limit of the Vlasov-Poisson system with massless electrons. *Communications in Partial Differential Equations*, 36(8):1385-1425, 2011.
- [16] D. Han-Kwan and M. Iacobelli. The quasineutral limit of the Vlasov-Poisson equation in Wasserstein metric. *Commun. Math. Sci.*, 15(2):481–509, 2 2017.
- [17] D. Han-Kwan, T. T. Nguyen and F. Rousset. Asymptotic stability of equilibria for screened Vlasov–Poisson systems via pointwise dispersive estimates. *Annals of PDE*, 7(2): 1-37, 2021.
- [18] D. Han-Kwan, T. Nguyen and F. Rousset. On the linearized Vlasov–Poisson system on the whole space around stable homogeneous equilibria. *Communications in Mathematical Physics*, 387(3): 1405-1440, 2021.
- [19] E. Horst. On the classical solutions of the initial value problem for the unmodified non-linear Vlasov equation (Parts I and II), *Math. Meth. Appl. Sci.* 3 (1981),pp. 229-248 and 4 (1982), pp. 19-32.
- [20] E. Horst. On the asymptotic growth of the solutions of the Vlasov-Poisson system. *Mathematical Methods in the Applied Sciences*, 16(2):75–86, 1993.
- [21] L. Huang, Q.H. Nguyen and Y. Xu. Sharp estimates for screened Vlasov-Poisson system around Penrose-stable equilibria in \mathbb{R}^2 . *In preparation*.
- [22] H.-J. Hwang, A. Rendall, and J.-L. Velázquez. Optimal gradient estimates and asymptotic behaviour for the Vlasov-Poisson system with small initial data. *Archive for Rational Mechanics and Analysis*, 200(1):313–360, 2011.
- [23] A. Ionescu, B. Pausader, X. Wang and K. Widmayer. On the Asymptotic Behavior of Solutions to the Vlasov–Poisson System. *International Mathematics Research Notices*, 2021, rnab155, <https://doi.org/10.1093/imrn/rnab155>
- [24] A. Ionescu, B. Pausader, X. Wang and K. Widmayer. Nonlinear Landau damping for the Vlasov-Poisson system in \mathbb{R}^3 : the Poisson equilibrium. *arXiv:2205.04540*, 2022.
- [25] L. Landau. On the vibration of the electronic plasma. *J. Phys. USSR*, 10(25), 1946.
- [26] P. L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system. *Inventiones mathematicae*, 105(1): 415-430, 1991.
- [27] C. Mouhot and C. Villani. On Landau damping. *Acta Mathematica*, 207(1):29–201, 2011.
- [28] Q. H. Nguyen. Quantitative estimates for regular Lagrangian flows with BV vector fields. *Communications on Pure and Applied Mathematics*, 74(6): 1129-1192, 2021.
- [29] T. T. Nguyen . Derivative estimates for screened Vlasov-Poisson system around Penrose-stable equilibria. *ArXiv:2004.05546*, 2020.
- [30] K. Pfaffelmoser. Global classical solutions of the Vlasov-Poisson system in three dimensions for general initial data. *Journal of Differential Equations*, 95(2):281–303, 1992.
- [31] J. Schaeffer. Global existence of smooth solutions to the Vlasov-Poisson system in three dimensions. *Communications in Partial Differential Equations*, 16(8-9):1313–1335, 1991.
- [32] J. Smulevici. Small data solutions of the Vlasov-Poisson system and the vector field method. *Annals of PDE*, 2(2): 11-55, 2016.
- [33] X. Wang. Decay estimates for the 3D relativistic and non-relativistic Vlasov-Poisson systems. *arXiv:1805.10837*, 2018.