

Fermion dynamics in presence of non abelian monopoles

Alejandro Morano ^{*}and Osvaldo P. Santillán [†]

Abstract

In the present letter, the dynamics of a spin one-half particle with non abelian charge, interacting with a non abelian monopole like configuration, is studied. In the non spinning case, these equations correspond to the Wong ones [1], and the associated dynamics has been extensively studied in [2]-[3]. The classical limit of a spinning particle in an abelian was studied in [4], where an interpretation of the celebrated Callan-Rubakov effect [5]-[6] was obtained in purely classical terms. The present work studies an interpolating situation, in which spin and non abelian charges are turned on. The corresponding equations are obtained by taking into account some earlier works about spinning particles [7]-[9]. The conservation laws are studied in this context, and it is found that energy and some generalization of angular momentum are conserved just in particular limits. This is a reflection that the non abelian interaction contains gauge field quartic interaction, which spoils the symmetries leading to these conservation laws. The precession of the particle is analyzed and compared with its abelian counterpart in this limit.

1. Introduction

Non abelian gauge theories are one of the most important subjects in modern theoretical physics, as they may explain phenomena such quark confinement and several aspects of hadron physics. One of the aspects that makes them harder to understand is that quarks seemingly are not part of the asymptotic states of the theory. It was suggested at some point that non abelian monopoles [10]-[11] may play an important role for confinement, since they may help for realizing a dual Meissner effect in which the quarks become the analog of confined magnets. In this scenario, the monopoles are decoupled at low energy due to their large mass [12]-[14]. As far as the authors know, this idea has not yet successfully implemented. However, the study of flux lines, confinement and dualities has been proved to be fruitful in the supersymmetric context. Supersymmetry has not been discovered yet, nevertheless, these studies are an interesting theoretical laboratory for understanding the possible phases that a gauge theory may realize [15]-[30].

The present work is focused on the classical dynamics of a particle with one-half spin in presence of non abelian fields, in particular monopole configurations. At first sight, one may wonder about the possible applications of such studies, as quarks are not free and the quantum effects are believed to play a significant role in the dynamics. However, some motivations may be given. An example is the study of a non abelian plasma [31]-[42] at very high densities or temperatures. These studies may have applications in the very early universe, the dynamics of neutron stars or supernova explosions. In addition, the study of a electroweak plasma may have interesting applications for modelling baryon number violation at these cosmological stages.

There exist excellent books such as [43]-[45] which covers topics about non abelian monopoles. However, the authors did not found yet a discussion focused on classical aspects of particles in these non abelian configurations in full detail. This is the motivation for writing the present short work.

^{*}Departamento de Física, FCEyN, Universidad de Buenos Aires, Buenos Aires, Argentina a.g.morano@gmail.com and osantil@dm.uba.ar.

[†]IMAS, FCEyN, Universidad de Buenos Aires, Buenos Aires, Argentina firenzecita@hotmail.com and osantil@dm.uba.ar.

We should however open the possibility that this analysis has been done before. If this is the case, we are simply not acquainted with that reference.

The present work is organized as follows. In section 2 the equations of motion are derived, and the abelian limit and the spinless case are also discussed. In section 3 and 4 the energy and generalized angular momentum are discussed, and it is shown that these quantities are conserved only in specific situations. An example is presented in section 5, where the spin and isotopic charge precession are studied in some detail. This section also contains a discussion of the presented results.

2. Derivation of the equations of motion

2.1 An abelian probe with spin in a monopole field

Before deriving the equations of motion of a spinning non abelian particle in the field of a non abelian monopole, it may be advisable to study some simpler situations. In an abelian electromagnetic field, and in the non relativistic limit, the equations of motion of a charged particle are given by

$$m\ddot{\vec{r}} = Qe(\vec{E} + \vec{v} \times \vec{B}) + \frac{\mu e}{2m} \nabla(\vec{S} \cdot \vec{B}), \quad (2.1)$$

$$\frac{d\vec{S}}{dt} = \frac{\mu e}{2m} (\vec{S} \times \vec{B}), \quad (2.2)$$

where Qe is the particle charge, m its mass and μ its anomalous magnetic moment. The back reaction of the particle on the source is neglected, which means that the gauge fields are not affected by the dynamics. The second term of (2.1) represents the interaction of the spin and the magnetic field and the equation (2.2) represents Larmor spin precession. These equations are well known. It is also known that the energy

$$E = \frac{m}{2} [\dot{r}^2 + r^2(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2)] - \frac{\mu e}{2m} \vec{B} \cdot \vec{S}, \quad (2.3)$$

is conserved for a magnetic field independent on time. Such is the case for the abelian monopole field

$$\vec{B} = \frac{g\vec{r}}{r^3}, \quad (2.4)$$

with g the monopole magnetic charge, which satisfies the Dirac quantization $2eg = 1$.

The issues related to angular momentum conservation are more tricky. Even taking into account the rotational symmetry of the monopole magnetic field, the angular momentum is not conserved as the spin-magnetic field coupling spoils this symmetry. However, the following quantity

$$\vec{J} = mr^2[\dot{\theta}\hat{\phi} - \dot{\phi}\sin\theta\hat{\theta}] + \vec{S} - Qeg\hat{r}, \quad (2.5)$$

has dimensions of an angular momentum and is conserved. Here the standard spherical coordinates were employed, in which the axis \hat{x} corresponds to $2\theta = \pi$ and $\phi = 0$. If a particle incident to the monopole is considered, with impact parameter d and velocity v in the negative \hat{x} direction, then the value of \vec{J} is specified as

$$\vec{J} = mdv\hat{z} - Qeg\hat{x} + \vec{S}_0, \quad (2.6)$$

with \vec{S}_0 the initial spin direction. On the other hand, (2.5) implies that

$$m^2r^4(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) = j^2 + s^2 - 2\vec{J} \cdot \vec{S} - Q^2e^2g^2,$$

where j and s indicates the modulus of their capital letter counterparts. By use of the value (2.6) it follows that

$$m^2r^4(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) = m^2d^2v^2 - 2\vec{S} \cdot [\vec{S}_0 - \vec{S}] + 2mdv[(S_0)_z - S_z] - 2Qeg[(S_0)_x - S_x]$$

By eliminating, by use of the last expression, the angular part in the energy $E = mv^2/2$, it is arrived to

$$\dot{r}^2 = \sqrt{v^2 - \frac{m^2 d^2 v^2 + 2(mdv\hat{z} - Qeg\hat{x} - \bar{S}) \cdot (\bar{S}_0 - \bar{S})}{mr^2}}, \quad (2.7)$$

The spin equations are

$$\begin{aligned} \frac{dS_x}{dt} &= \frac{\mu ge}{2mr^2} (S_y \cos \theta - S_z \sin \theta \sin \phi), \\ \frac{dS_y}{dt} &= \frac{\mu ge}{2mr^2} (S_z \sin \theta \cos \phi - S_x \cos \theta), \\ \frac{dS_z}{dt} &= \frac{\mu ge}{2mr^2} (S_x \sin \theta \sin \phi - S_y \sin \theta \cos \phi). \end{aligned} \quad (2.8)$$

The angular momentum in spherical coordinates reads

$$\bar{J} = mr^2[\dot{\theta}\hat{\phi} - \dot{\phi}\sin\theta\hat{\theta}] + \bar{S} - Qeg\hat{r}, \quad (2.9)$$

From here and the initial value, it is found that

$$\frac{d\phi}{dt} = \frac{mdv + Qeg \cos \theta + S_{0z} - S_z}{mr^2 \sin^2 \theta}, \quad (2.10)$$

$$\frac{d\theta}{dt} = \frac{Qeg \sin \theta \sin \phi + S_{0y} - S_y}{mr^2 \cos \phi} + \cot \theta \tan \phi \left[\frac{mdv + Qeg \cos \theta + S_{0z} - S_z}{mr^2} \right] \quad (2.11)$$

From (2.7) it is seen that there is a spin dependent radius of return when

$$mv^2 = \frac{m^2 d^2 v^2 + 2(mdv\hat{z} - Qeg\hat{x} - \bar{S}) \cdot (\bar{S}_0 - \bar{S})}{mr^2}.$$

The extremal values of the quantity of the right hand side with respect to \bar{S} are found to be

$$r_e^2 = \frac{m^2 d^2 v^2 + (mdv\hat{z} - Qeg\hat{x}) \cdot \bar{S}_0}{m^2 v^2}.$$

The term $(mdv\hat{z} - Qeg\hat{x}) \cdot \bar{S}_0$ takes the minimum when the vectors are anti parallel, the result is

$$r_e^2 = \frac{m^2 d^2 v^2 - s\sqrt{m^2 d^2 v^2 + s^2}}{m^2 v^2}.$$

This potential will be repulsive if

$$mdv \geq \text{Max}(s, \mu ge, Qge).$$

The equality defines an approximate critical value d_c for which the potential can be attractive depending on the spin direction. Thus, the corresponding scattering cross section may be approximated by

$$\sigma \sim \pi d_c^2 \sim \pi \left(\frac{Qeg}{mv} \right)^2,$$

which, for $3v \sim 1$ is of the order of the Callan-Rubakov estimation [5]-[6]. These classical considerations were employed in [4], although with slightly different methodology.

2.2 A non abelian spinless charge in a non abelian field

On the other hand, the relativistic dynamics of an spinless particle in a non abelian gauge field is described by the Wong equations [1]

$$m\dot{u}_\mu = gF_{\mu\nu}^a u^\nu I^a, \quad (2.12)$$

$$\frac{dI^a}{dt} = gf^{abc}I^b[A_0^c + \bar{v} \cdot \bar{A}^c]. \quad (2.13)$$

Here the signature convention $(1, -1, -1, -1)$ is employed and u^μ is the relativistic four velocity $u^\mu = dx^\mu/d\tau$, with τ the proper time. The non abelian charge is characterized by the vector I^a , which may depend on τ during the particle evolution. The external non abelian gauge field is denoted by A_μ , and the non abelian stress tensor is calculated by means of the formula

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig[A_\mu, A_\nu].$$

By employing generators T^a of the gauge group such that $[T^a, T^b] = if_{abc}T^c$ with f_{abc} real constants, the stress tensor may be expressed as

$$F_{\mu\nu} = F_{\mu\nu}^a T^a, \quad F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf_{abc}A_\mu^b A_\nu^c.$$

In particular, in the the $SU(2)$ case considered below $a = 1, 2, 3$, $f_{abc} = \epsilon_{abc}$ and $2T^a = \sigma^a$, with σ^a the standard Pauli matrices. In general, the covariant derivative of given non abelian field φ is expressed as

$$D_\mu \varphi = \partial_\mu \varphi - ig[A_\mu, \varphi].$$

In these terms, the equations of motion (2.12)-(2.13) can be shown to be equivalent to

$$\frac{d}{dt} \left(\frac{m\bar{v}}{\sqrt{1-v^2}} \right) = g(\bar{E}^a + \bar{v} \times \bar{B}^a)I^a, \quad (2.14)$$

$$\frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} \right) = g\bar{v} \cdot \bar{E}^a I^a, \quad (2.15)$$

$$\frac{dI^a}{dt} = gf^{abc}I^b[A_0^c + \bar{v} \cdot \bar{A}^c]. \quad (2.16)$$

Here the external non abelian electric and magnetic fields \bar{E}^a and \bar{B}^a are given by

$$E_i^a = F_{0i}^a = \partial_0 A_i^a - \partial_i A_0^a + gf_{abc}A_0^b A_i^c, \quad B_i^a = \frac{\epsilon_{ijk}}{2} F_{jk}^a. \quad (2.17)$$

The interpretation of these equations is clear. The first are the relativistic Newton equations in a non abelian gauge field. The second is the kinetic energy variation, as a consequence of the work of the electric force on the particle. The last ones represent the evolution of the charge vector I^a due to the particle dynamics. Classical solutions of this equations were studied, for instance, in [2]-[3].

2.3 A non abelian charge with spin in a non abelian field

The equations of motion for an spinning probe particle in a general electromagnetic field will be taken, following the references [7]-[9], as follows

$$\frac{d}{dt} \left(\frac{m\bar{v}}{\sqrt{1-v^2}} \right) = g(\bar{E}^a + \bar{v} \times \bar{B}^a)I^a + \frac{\mu g}{2m} I^a S_l [\nabla B_l^a - gf^{abc}B_l^b \bar{A}^c], \quad (2.18)$$

$$\frac{d\bar{S}}{dt} = \frac{\mu g}{2m} I^a (\bar{S} \times \bar{B}^a), \quad (2.19)$$

$$\frac{dI^a}{dt} = g f^{abc} I^b [A_0^c + \bar{v} \cdot \bar{A}^c] + \frac{\mu g}{2m} \epsilon^{abc} I^b \bar{S} \cdot \bar{B}^c. \quad (2.20)$$

The last system interpolate between the two situations described above namely, the abelian case and the non abelian Wong equations. The first limit follows by choosing the structure constants $f^{abc} = 0$ and the second by putting $\bar{S} = 0$ in the last equations. The next issue is to understand in which situations there are conserved quantities for the given system of equations. The non abelianity of the fields may alter the intuition about the conserved quantities, as there are quartic gauge field interactions which are absent in the abelian case.

3. The energy balance equation

By multiplying (2.18) by $v = \dot{\bar{r}}$, the following expression for the time variation of the kinetic energy of the particle is obtained

$$\frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} \right) = g \bar{v} \cdot \bar{E}^a I^a + \frac{\mu g}{2m} I^a S_l [\bar{v} \cdot \nabla B_l^a - g f^{abc} B_l^b (\bar{v} \cdot \bar{A}^c)]. \quad (3.21)$$

On the other hand, the generalization of the potential energy in an electric field for the non abelian case will be $I^a A_0^a$. Its variation is

$$\frac{d(I^a A_0^a)}{dt} = \frac{dI^a}{dt} A_0^a + I^a [\partial_0 A_0^a + \bar{v} \cdot \nabla \bar{A}_0^a].$$

By using (2.17) in order to eliminate the term $\nabla \bar{A}_0^a$ and also by taking into account (2.20) it is found that

$$\begin{aligned} \frac{d(I^a A_0^a)}{dt} &= g f^{abc} A_0^a I^b [A_0^c + \bar{v} \cdot \bar{A}^c] + \frac{\mu g}{2m} \epsilon^{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a \\ &\quad + I^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a - \bar{E}^a + g f_{abc} A_0^b \bar{A}^c)]. \end{aligned}$$

In the last expression, the first term is zero due to the anti-symmetry of f_{abc} with respect of its indices. The second term cancels the last one. Thus

$$\frac{d(I^a A_0^a)}{dt} = \frac{\mu g}{2m} \epsilon^{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a + I^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a - \bar{E}^a)].$$

The sum of the last expression multiplied with g with (3.21) then leads to

$$\begin{aligned} \frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} + g I^a A_0^a \right) &= \frac{\mu g}{2m} I^a S_l [\bar{v} \cdot \nabla B_l^a - g f^{abc} B_l^b (\bar{v} \cdot \bar{A}^c)] \\ &\quad + \frac{\mu g^2}{2m} \epsilon^{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a + g I^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a)]. \end{aligned}$$

Note that the terms proportional to the non abelian electric field E_i^a cancelled each other. The last formula may be expressed as

$$\begin{aligned} \frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} + g I^a A_0^a \right) &= -\frac{\mu g^2}{2m} f_{abc} I^a S_l B_l^b (\bar{v} \cdot \bar{A}^c) + \frac{\mu g^2}{2m} \epsilon_{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a \\ &\quad + \frac{\mu g}{2m} I^a S_l \frac{dB_l^a}{dt} + g I^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a)]. \end{aligned}$$

On the other hand, (2.19) shows that $\dot{\bar{S}} \cdot \bar{B} = 0$ and therefore

$$\frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} + g I^a A_0^a - \frac{\mu g}{2m} I^a B_l^a S_l \right) = -\frac{\mu g^2}{2m} f_{abc} I^a S_l B_l^b (\bar{v} \cdot \bar{A}^c) + \frac{\mu g^2}{2m} \epsilon_{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a$$

$$-\frac{\mu g}{2m} B_l^a S_l \frac{dI^a}{dt} + gI^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a)].$$

By further taking into account (2.20) and the anti-symmetry of f_{abc} the last expression becomes

$$\begin{aligned} \frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} + gI^a A_0^a - \frac{\mu g}{2m} I^a B_l^a S_l \right) &= -\frac{\mu g^2}{2m} f_{abc} I^a S_l B_l^b (\bar{v} \cdot \bar{A}^c) + \frac{\mu g^2}{2m} \epsilon_{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a \\ &\quad - \frac{\mu g^2}{2m} f_{abc} B_l^a S_l I^b [A_0^c + \bar{v} \cdot \bar{A}^c] + gI^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a)]. \end{aligned}$$

The first and the fourth term cancel out, and the second and the third are proportional. Thus

$$\begin{aligned} \frac{d}{dt} \left(\frac{m}{\sqrt{1-v^2}} + gI^a A_0^a - \frac{\mu g}{2m} I^a B_l^a S_l \right) &= \frac{\mu g^2}{m} \epsilon_{abc} I^b \bar{S} \cdot \bar{B}^c A_0^a \\ &\quad + gI^a [\partial_0 A_0^a + \bar{v} \cdot (\partial_0 \bar{A}^a)]. \end{aligned} \quad (3.22)$$

This formula has important consequences. One may define the full energy of the particle as

$$E = \frac{m}{\sqrt{1-v^2}} + gI^a A_0^a - \frac{\mu g}{2m} \bar{S} \cdot \bar{B}^a I^a. \quad (3.23)$$

This is the sum of the kinetic energy, the generalization of the electric potential, and the Larmor one. If there is a gauge for which A_μ^a is time independent, the last two terms cancel. However, this energy is not conserved due to the first term in (3.22). This may be interpreted as an effect due to the non linear interaction of the gauge field of the particle and the external one.

The results given above show that energy is not conserved in general for non abelian probe particles moving in an abelian field. However, as it will be shown below, it is conserved for a monopole field in a $SU(2)$ gauge theory, in certain limit.

4. The angular momentum balance equation

The angular momentum is not expected to be conserved either, even for a static field configuration. In the present section, the attention will be restricted to the class of static $SU(2)$ non abelian fields given by

$$A_0^a = \frac{f}{gr} n^a, \quad A_i^a = \frac{(1-a)}{gr} \epsilon_{aij} n^j. \quad (4.24)$$

Here $a = a(r)$ and $f = f(r)$ are generic functions of the distance r between the observation point and the non abelian source, whose explicit profile depends on the model of consideration. In the last expression, the unit vector

$$n^a = \frac{x^a}{r}, \quad \dot{n}^a \cdot n^a = 0,$$

has been introduced, with x^a a system of cartesian coordinates in R^3 describing the position of the probe particle. In the present case, the structure constants $f_{abc} = \epsilon_{abc}$. For the calculation to be performed below, it is convenient to write (2.18)-(2.20) in terms of (4.24), and the corresponding color electromagnetic fields. From (4.24) and (2.17) it is found out that

$$E_i^a = -\frac{1}{g} \left[\left(\frac{f}{r} \right)' n^a n^i + \frac{af}{r^2} (\delta^{ai} - n^a n^i) \right], \quad (4.25)$$

$$B_i^a = -\frac{1}{g} \left[\frac{a^2 - 1}{r^2} n^a n^i + \frac{a'}{r} (\delta^{ai} - n^a n^i) \right]. \quad (4.26)$$

By use of the last expressions, (4.24) and (2.19) it is found the following equation describing spin precession

$$\frac{d\bar{S}}{dt} = \frac{\mu}{2m} \left[\frac{1-a^2}{r^2} (\bar{I} \cdot \bar{n}) \bar{S} \times \bar{n} - \frac{a'}{r} \bar{S} \times \bar{I} + \frac{a'}{r} (\bar{I} \cdot \bar{n}) \bar{S} \times \bar{n} \right]. \quad (4.27)$$

On the other hand, (2.20) and the identity $\bar{v} = \dot{r}\bar{n} + r\dot{\bar{n}}$ imply that

$$\begin{aligned} \frac{d\bar{I}}{dt} &= \frac{f}{r} \bar{I} \times \bar{n} + (1-a)[(\bar{I} \cdot \bar{n})\dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}})\bar{n}] \\ &+ \frac{\mu}{2m} \left[\frac{1-a^2}{r^2} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} - \frac{a'}{r} \bar{S} \times \bar{I} + \frac{a'}{r} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} \right]. \end{aligned} \quad (4.28)$$

This, in particular, leads to the following formula

$$\frac{d\bar{I} \cdot \bar{n}}{dt} = a(\bar{I} \cdot \dot{\bar{n}}) - \frac{\mu}{2m} \frac{a'}{r} (\bar{S} \times \bar{I}) \cdot \bar{n}, \quad (4.29)$$

which will be employed below. Now, vector multiplication of (2.18) with respect to \bar{v} leads to the following equation for the angular momentum of the particle

$$\frac{d\bar{L}}{dt} = g(\bar{r} \times \bar{E}^a + \bar{r} \times \bar{v} \times \bar{B}^a) I^a + \frac{\mu g}{2m} I^a S_l [(\bar{r} \times \nabla) B_l^a - g f^{abc} B_l^b (\bar{r} \times \bar{A}^c)]. \quad (4.30)$$

Here

$$\bar{L} = \frac{m\bar{r} \times \bar{v}}{\sqrt{1-v^2}}.$$

There are several terms to be calculated. First, from (4.25)-(4.26) and by use of $\bar{v} = \dot{r}\bar{n} + r\dot{\bar{n}}$, it is found that

$$g(\bar{r} \times \bar{E}^a + \bar{r} \times \bar{v} \times \bar{B}^a) I^a = \frac{a f}{r} \bar{I} \times \bar{n} + (1-a^2)(\bar{I} \cdot \bar{n})\dot{\bar{n}} + \frac{da}{dt} (\bar{I} - (\bar{I} \cdot \bar{n})\bar{n}). \quad (4.31)$$

On the other hand, expression for the term ∇B_l^a is a bit cumbersome, but the expression for $(\bar{r} \times \nabla) B_l^a$ is significantly simpler. A direct calculation shows the validity of this identity for the present configuration

$$\begin{aligned} \frac{\mu g}{2m} I^a S_l (\bar{r} \times \nabla) B_l^a &= \frac{\mu}{2m} \left[\frac{1-a^2}{r^2} (\bar{S} \cdot \bar{n}) (\bar{n} \times \bar{I}) + \frac{1-a^2}{r^2} (\bar{I} \cdot \bar{n}) (\bar{n} \times \bar{S}) \right. \\ &\left. + \frac{a'}{r} (\bar{S} \cdot \bar{n}) (\bar{n} \times \bar{I}) + \frac{a'}{r} (\bar{I} \cdot \bar{n}) (\bar{n} \times \bar{S}) \right]. \end{aligned}$$

This expression describes the third term in (4.30). Comparison of the last terms of this formula with (4.27) and (4.28) allows to find a simpler expression for the third term, namely

$$\frac{\mu g}{2m} I^a S_l (\bar{r} \times \nabla) B_l^a = -\frac{d\bar{S}}{dt} - \frac{d\bar{I}}{dt} + \frac{f}{r} \bar{I} \times \bar{n} + (1-a)[(\bar{I} \cdot \bar{n})\dot{\bar{n}} + (\bar{I} \cdot \dot{\bar{n}})\bar{n}] - \frac{\mu}{m} \frac{a'}{r} \bar{S} \times \bar{I}. \quad (4.32)$$

Finally, the last term in (4.30) can be calculated to give

$$\begin{aligned} -\frac{\mu g^2}{2m} I^a S_l \epsilon^{abc} B_l^b (\bar{r} \times \bar{A}^c) &= \frac{\mu}{2m} \frac{(1-a)a'}{r} \bar{I} \cdot (\bar{S} \times \bar{n}) \bar{n} \\ &+ \frac{\mu}{2m} (1-a) \left[\frac{1-a^2}{r^2} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} - \frac{a'}{r} \bar{S} \times \bar{I} + \frac{a'}{r} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} \right]. \end{aligned}$$

The use of (4.27) and (4.28) again shows that the last formula is equivalent to the following one

$$\begin{aligned} -\frac{\mu g^2}{2m} I^a S_l \epsilon^{abc} B_l^b (\bar{r} \times \bar{A}^c) &= \frac{\mu}{2m} \frac{(1-a)a'}{r} \bar{I} \cdot (\bar{S} \times \bar{n}) \bar{n} + (1-a) \frac{d\bar{I}}{dt} \\ &\quad - (1-a) \frac{f}{r} \bar{I} \times \bar{n} - (1-a)^2 [(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}}) \bar{n}]. \end{aligned} \quad (4.33)$$

After these calculations, it is found that sum of the formulas (4.31)-(4.33) imply that

$$\begin{aligned} \frac{d\bar{L}}{dt} + \frac{d\bar{S}}{dt} &= \frac{af}{r} \bar{I} \times \bar{n} + (1-a^2)(\bar{I} \cdot \bar{n}) \dot{\bar{n}} + \frac{da}{dt} (\bar{I} - (\bar{I} \cdot \bar{n}) \bar{n}) \\ -a \frac{d\bar{I}}{dt} + \frac{af}{r} \bar{I} \times \bar{n} + a(1-a)[(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}}) \bar{n}] &+ \frac{\mu}{m} \frac{a'}{r} \bar{S} \times \bar{I} + \frac{\mu}{2m} \frac{(1-a)a'}{r} \bar{I} \cdot (\bar{S} \times \bar{n}) \bar{n}. \end{aligned}$$

It can not be concluded from this expression that the sum $\bar{J} = \bar{L} + \bar{S}$ is conserved. This is not peculiar however, as this quantity is not conserved in the abelian case either. Nevertheless, the use of (4.29) allows to put the last expression in the following form

$$\begin{aligned} \frac{d}{dt} (\bar{L} + \bar{S} - a\bar{I} - (1-a)(\bar{I} \cdot \bar{n}) \bar{n}) &= \frac{2af}{r} \bar{I} \times \bar{n} + (1-a^2)(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (1-a)(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (1-a)[a(\bar{I} \cdot \dot{\bar{n}}) - \frac{\mu}{2m} \frac{a'}{r} (\bar{S} \times \bar{I}) \cdot \bar{n}] \bar{n} \\ &\quad - 2a \frac{d\bar{I}}{dt} + a(1-a)[(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}}) \bar{n}] + \frac{\mu}{m} \frac{a'}{r} \bar{S} \times \bar{I} + \frac{\mu}{2m} \frac{(1-a)a'}{r} \bar{I} \cdot (\bar{S} \times \bar{n}) \bar{n}. \end{aligned}$$

The use of (4.28) leads to

$$\begin{aligned} \frac{d}{dt} (\bar{L} + \bar{S} - a\bar{I} - (1-a)(\bar{I} \cdot \bar{n}) \bar{n}) &= (1-a^2)(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (1-a)(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (1-a)a(\bar{I} \cdot \dot{\bar{n}}) \bar{n} \\ &\quad + a(1-a)[(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}}) \bar{n}] - 2a(1-a)[(\bar{I} \cdot \bar{n}) \dot{\bar{n}} - (\bar{I} \cdot \dot{\bar{n}}) \bar{n}] \\ &+ \frac{\mu}{m} \frac{a'}{r} \bar{S} \times \bar{I} + \frac{\mu}{2m} \frac{(1-a)a'}{r} (\bar{I} \cdot (\bar{S} \times \bar{n})) \bar{n} + \frac{\mu(1-a)a'}{2m} \frac{a'}{r} ((\bar{S} \times \bar{I}) \cdot \bar{n}) \bar{n} - \frac{\mu a}{2m} \left[\frac{1-a^2}{r^2} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} \right. \\ &\quad \left. - \frac{a'}{r} \bar{S} \times \bar{I} + \frac{a'}{r} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} \right]. \end{aligned}$$

Several terms cancel each other out, and this formula simplify to

$$\begin{aligned} \frac{d}{dt} (\bar{L} + \bar{S} - a\bar{I} - (1-a)(\bar{I} \cdot \bar{n}) \bar{n}) &= \frac{\mu}{m} \frac{a'(1-a)}{r} \bar{S} \times \bar{I} + \frac{\mu}{2m} \frac{(1-a)a'}{r} ((\bar{S} \times \bar{n}) \cdot \bar{I}) \bar{n} \\ &\quad + \frac{\mu(1-a)a'}{2m} \frac{a'}{r} ((\bar{S} \times \bar{I}) \cdot \bar{n}) \bar{n} - \frac{\mu a}{2m} \left[\frac{1-a^2}{r^2} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} + \frac{a'}{r} (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n} \right]. \end{aligned}$$

By use of algebra vector identities, this becomes

$$\frac{d}{dt} (\bar{L} + \bar{S} - a\bar{I} - (1-a)(\bar{I} \cdot \bar{n}) \bar{n}) = \frac{\mu}{m} \frac{a'(1-a)}{r} \bar{S} \times \bar{I} - \frac{\mu a}{2m} \left[\frac{1-a^2}{r^2} + \frac{a'}{r} \right] (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n}.$$

This means that

$$\frac{d\bar{M}}{dt} = \frac{\mu}{m} \frac{a'(1-a)}{r} \bar{S} \times \bar{I} - \frac{\mu a}{2m} \left[\frac{1-a^2}{r^2} + \frac{a'}{r} \right] (\bar{S} \cdot \bar{n}) \bar{I} \times \bar{n},$$

where

$$\bar{M} = \bar{L} + \bar{S} - a\bar{I} - (1-a)(\bar{I} \cdot \bar{n}) \bar{n}.$$

Therefore the quantity \bar{M} is conserved in this case only if $a = 0$ or $a = 1$. The case $a = 0$ corresponds approximately to the asymptotic region, while the case $a = 1$ describes the core of the object. As the core is very small, the solution may be approximated with $a = 0$ in almost all the region of the space. Note that, in the abelian limit $\bar{I} \cdot \bar{n} = Qe$, the resulting conserved quantity coincides with (2.5).

5. Specific calculation

In the following, the approximation $a = 0$ will be employed. In this limit, the formula (4.29) leads to

$$\bar{\mathbf{I}} \cdot \bar{\mathbf{n}} = \bar{\mathbf{I}}_0 \cdot \bar{\mathbf{n}}_0 \quad (5.34)$$

where the 0 represent the quantities at the beginning of the motion. The last formula implies that the projection of the abelian charge on the direction joining the probe particle and the source is conserved. In addition, in this case, the energy balance equation implies that (3.22) that

$$E = \frac{m}{\sqrt{1-v^2}} + gI^a A_0^a - \frac{\mu g}{2m} \bar{\mathbf{S}} \cdot \bar{\mathbf{B}}^a I^a. \quad (5.35)$$

is conserved, as the term $\epsilon_{abc} I^b \bar{\mathbf{S}} \cdot \bar{\mathbf{B}}^c A_0^a$ is zero in this limit. In fact, for $a = 0$ the gauge fields are given by

$$A_0^a = \frac{f}{gr} n^a, \quad A_i^a = \frac{1}{gr} \epsilon_{aij} n^j.$$

$$E_i^a = -\frac{1}{g} \left(\frac{f}{r} \right)' n^a n^i, \quad B_i^a = \frac{1}{gr^2} n^a n^i.$$

These formulas imply that $\epsilon_{abc} I^b \bar{\mathbf{S}} \cdot \bar{\mathbf{B}}^c A_0^a$ is proportional to $\epsilon_{abc} n^b n^c$, which clearly vanishes. In addition, it follows that

$$\partial_j B_i^a = \frac{1}{gr^3} (\delta^{aj} n^i + \delta^{ai} n^j) - \frac{4n^i n^a n^j}{gr^3}.$$

The equations of motion simplify to

$$\frac{d}{dt} \left(\frac{m\bar{\mathbf{v}}}{\sqrt{1-v^2}} \right) = - \left(\frac{f}{r} \right)' (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) \bar{\mathbf{n}} + \frac{1}{r^2} (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) (\bar{\mathbf{v}} \times \bar{\mathbf{n}}) + \frac{\mu}{2mr^3} [(\bar{\mathbf{I}} \cdot \bar{\mathbf{S}}) \bar{\mathbf{n}} - 3(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) (\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}) \bar{\mathbf{n}}],$$

$$\frac{d\bar{\mathbf{S}}}{dt} = \frac{\mu}{2mr^2} (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) (\bar{\mathbf{S}} \times \bar{\mathbf{n}}),$$

$$\frac{d\bar{\mathbf{I}}}{dt} = \frac{f}{r} \bar{\mathbf{I}} \times \bar{\mathbf{n}} + \frac{\bar{\mathbf{I}} \times \bar{\mathbf{v}} \times \bar{\mathbf{n}}}{r} + \frac{\mu}{2mr^2} (\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}) (\bar{\mathbf{I}} \times \bar{\mathbf{n}}).$$

The conserved quantities are

$$E = \frac{m}{\sqrt{1-v^2}} + \frac{f}{r} (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) - \frac{\mu}{2mr^2} (\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}) (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}).$$

$$\frac{m\bar{\mathbf{v}} \times \bar{\mathbf{n}}}{\sqrt{1-v^2}} = \frac{\bar{\mathbf{S}}}{r} - \frac{\bar{\mathbf{M}}}{r} + \frac{(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) \bar{\mathbf{n}}}{r}.$$

This implies that

$$(\bar{\mathbf{S}} + \bar{\mathbf{I}}) \cdot \bar{\mathbf{n}} = \bar{\mathbf{M}} \cdot \bar{\mathbf{n}}.$$

Even taking into account (5.34), note that the last equation does not imply that $\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}$ is conserved. The point is that $\bar{\mathbf{M}}$ is a constant of motion, but its multiplication with $\bar{\mathbf{n}}$ is not. The conserved quantities may be written as

$$E = \frac{m}{\sqrt{1-v^2}} + \left[\frac{f}{r} - \frac{\mu}{2mr^2} (\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}) \right] (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}).$$

$$\frac{m\bar{\mathbf{v}} \times \bar{\mathbf{n}}}{\sqrt{1-v^2}} = \frac{\bar{\mathbf{S}}}{r} - \frac{(\bar{\mathbf{S}} \cdot \bar{\mathbf{n}}) \bar{\mathbf{n}}}{r} - \frac{\bar{\mathbf{M}}}{r} + \frac{(\bar{\mathbf{M}} \cdot \bar{\mathbf{n}}) \bar{\mathbf{n}}}{r}.$$

Explicitly, for the non relativistic case, the energy is

$$E = \frac{m}{2}(\dot{r}^2 + r^2\dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2) + \left[\frac{f}{r} - \frac{\mu}{2mr^2}(S_x \sin \theta \cos \phi + S_y \sin \theta \sin \phi + S_z \cos \theta) \right] (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}).$$

By taking this into account, the spin equations and the conservation of M_i it is obtained the following linear system

$$\begin{aligned} \frac{dS_x}{dt} &= \frac{\mu(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}})}{2mr^2}(S_z \sin \theta \sin \phi - S_y \cos \theta), \\ \frac{dS_y}{dt} &= \frac{\mu(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}})}{2mr^2}(S_x \cos \theta - S_z \sin \theta \cos \phi), \\ \frac{dS_z}{dt} &= \frac{\mu(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}})}{2mr^2}(S_y \sin \theta \cos \phi - S_x \sin \theta \sin \phi), \\ \dot{\theta} &= \frac{1}{mr^2} \left[(M_x - S_x) \sin \phi - (M_y - S_y) \cos \phi \right], \\ \dot{\phi} &= \frac{1}{mr^2 \sin \theta} \left[(M_x - S_x) \cos \theta \cos \phi + (M_y - S_y) \cos \theta \sin \phi - (M_z - S_z) \sin \theta \right], \end{aligned}$$

$$\dot{r} = \pm \sqrt{\frac{2}{m} \left\{ E - \frac{m}{2}(r^2\dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2) - \left[\frac{f}{r} - \frac{\mu}{2mr^2}(S_x \sin \theta \cos \phi + S_y \sin \theta \sin \phi + S_z \cos \theta) \right] (\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}) \right\}},$$

where the fourth and the fifth equations have to be replaced in the last one for eliminating $\dot{\theta}$ and $\dot{\phi}$. The last are a system of equation of first order for the quantities describing the dynamics namely, S_i , θ , ϕ and r . This is a closed system. It is clear that this system describes a particle which moves as an abelian particle with charge $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}$, which is constant during the whole evolution. This does not implies however, that non abelian and abelian particles can not be distinguished by the dynamics. This is discussed below.

6. Discussion

It is seen by comparing the obtained system with the equations (2.1)-(2.11) that the dynamics are equivalent, by interpreting the constant $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}$ as an effective charge, analogous to Qe for the abelian case. However, the sign of the charge depends on the initial conditions, and the force may be attractive or repulsive depending on the initial choice. In addition, there exist a regime $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}} = 0$ for which the charge does not experience the force of the monopole and just pass through with a straight line. This does not imply that the charges do not evolve in time. In fact, in the limit $a = 0$ the equations (4.28) are given by

$$\frac{d\bar{\mathbf{I}}}{dt} = \frac{f}{r}\bar{\mathbf{I}} \times \bar{\mathbf{n}} + [(\bar{\mathbf{I}} \cdot \bar{\mathbf{n}})\dot{\bar{\mathbf{n}}} - (\bar{\mathbf{I}} \cdot \dot{\bar{\mathbf{n}}})\bar{\mathbf{n}}] + \frac{\mu}{2mr^2}(\bar{\mathbf{S}} \cdot \bar{\mathbf{n}})\bar{\mathbf{I}} \times \bar{\mathbf{n}}. \quad (6.36)$$

It is seen that even if $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}} = 0$ and $\bar{\mathbf{n}}$ describes a straight line, this time derivative is not zero. Thus, there is charge precession even in this almost trivial case.

The discussion given above may suggest that the dynamics of a non abelian charge is not distinguishable from an abelian one. This not necessarily the case, and there are several effects that can be considerably different. For a single charge, as discussed above, the effective charge $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}$ is a constant. However, consider a beam of particles with the same initial value of $\bar{\mathbf{I}}$. The value of $\bar{\mathbf{I}} \cdot \bar{\mathbf{n}}$ will be clearly different for any different particle composing the beam. In particular, consider a line composed by a beam an initial value of $\bar{\mathbf{I}}$, which is parallel to the direction $\bar{\mathbf{n}}$ pointing to an angle 0.25π with respect to the direction joining the monopole with the nearest point of the beam. The particles corresponding

to this direction will interact maximally with the monopole. Those with angle -0.25π will not interact at all, as $\vec{I} \cdot \vec{n} = 0$. Those with angles $\alpha < -0.25\pi$ will have opposite interaction with those with $\alpha > -0.25\pi$, that is, repulsion will change into attraction or viceversa. This does not happen for an abelian beam, as all the particles will have charges Qe independently on the direction \vec{n} . The cross section will depend on an average on $\vec{I} \cdot \vec{n}$, and the result will be of the order of Callan-Rubakov, despite those differences in the dynamics.

There is a possible additional difference, although the following argument is not completely rigorous. Consider a charge with spin in the middle of two identical abelian monopoles. The formula (2.1) shows that the force will be zero. However, if a direction dependent charge $\vec{I} \cdot \vec{n}$ is in the middle of two identical non abelian monopoles, there is no cancellation, as this seems to act as a positive charge for one monopole and negative for the other. This argument is dubious however, as the conservation of $\vec{I} \cdot \vec{n}$ was obtained only for the limit $a = 0$ in a single monopole configuration and may be violated for a two monopole field. In any case, we suggest that the dynamics of a non abelian charge in the field of two or more monopoles may be of academic interest. We leave this issue for a future publication.

Acknowledgments

O. S is supported by CONICET, Argentina.

References

- [1] S. Wong Il Nuovo Cimento LXV, 4 (1970) 689.
- [2] I. Alekseev Theoret. and Math. Phys., 77 (1988) 1273.
- [3] L. Gy. Feher Acta Physica Polonica 15 (1984) 920.
- [4] A. Davis and W. Perkins Phys. Lett. B 228 (1989) 37.
- [5] C.G. Callan, Phys. Rev. D 26 (1982) 2058; V.A. Rubakov, Nucl. Phys. B 203 (1982) 311.
- [6] V.A. Rubakov and M.S. Serebryakov, Nucl. Phys. B 218 (1983) 240; V.A. Rubakov, JETP Len. 33 (1981) 644.
- [7] N. Linden, A. Macfarlane and J van Holten Czechoslovak J. Phys. 46 (1996) 209.
- [8] A. Barducci, R. Casalbuoni and L. Lusanna Nuclear Physics B 124 (1977) 93
- [9] A. P. Balachandran Per Salomson,Bo-Sture Skagerstam, and Jan-Olov Winnberg Phys. Rev. D 15 (1977) 2308.
- [10] G. 't Hooft Nuclear Physics B. 79 (1975)276.
- [11] A. Polyakov JETP Letters. 20 (1974) 194.
- [12] G. 't Hooft Nucl. Phys. B 190 (1981) 455.
- [13] Y. Nambu, Phys. Rev. D 10 (1974) 4262.
- [14] S. Mandelstam Phys. Rept. 23 (1976) 245.
- [15] N. Seiberg and E. Witten, Nucl. Phys. B 426 (1994) 19.
- [16] N. Seiberg and E. Witten Nucl. Phys. B 431 (1994) 484.

- [17] M. Shifman and A. Yung Phys.Rev. D 66 (2002) 045012.
- [18] M. Shifman and A. Yung Phys.Rev.D 70 (2004) 045004.
- [19] M. Shifman and A. Yung Phys.Rev.D 72 (2005) 085017.
- [20] M. Shifman and A. Yung Phys.Rev.D 76 (2007) 045005.
- [21] R. Auzzi, S. Bolognesi and M. Shifman Phys.Rev.D 81 (2010) 085011.
- [22] A. Gorsky, M. Shifman and A. Yung Phys. Rev. D 83 (2011) 085027.
- [23] M. Shifman and A. Yung Phys. Rev. D 83 (2011) 105021.
- [24] S. Chen, Y. Li, Y. Yang Phys. Rev. D 86 (2012) 085030.
- [25] M. Shifman and A. Yung Phys. Rev. D 90 (2014) 065014.
- [26] N. Seiberg, Nucl. Phys. B 435 (1995) 129.
- [27] N. Seiberg, Phys. Rev. D 49 (1994) 6857.
- [28] M. A. Shifman, Prog. Part. Nucl. Phys. 39 (1997) 1.
- [29] K. A. Intriligator and N. Seiberg, Nucl. Phys. Proc. Suppl. 45BC (1996) 1.
- [30] K. Intriligator, N. Seiberg and D. Shih, JHEP 0604 (2006) 021.
- [31] D. Litim and C. Manuel Phys.Rept. 364 (2002) 451.
- [32] D. F. Litim, Semi-classical transport theory for non-Abelian plasmas, habilitation thesis (University of Heidelberg, May 2000).
- [33] D. F. Litim, Aspects of semi-classical transport theory for QCD, hep-ph/0010259.
- [34] D. F. Litim and C. Manuel Phys. Rev. Lett. 82 (1999) 4981.
- [35] D. F. Litim and C. Manuel Nucl. Phys. B562 (1999) 237.
- [36] D. F. Litim and C. Manuel Phys. Rev. D61 (2000) 125004.
- [37] D. F. Litim and C. Manuel, Deriving effective transport equations for non-Abelian plasmas, hep-ph/0003302.
- [38] D. F. Litim and C. Manuel Phys. Rev. Lett. 87 (2001) 052002.
- [39] D. F. Litim and C. Manuel Phys. Rev. D64 (2001) 094013.
- [40] C. Manuel Phys. Rev. D53 (1996) 5866.
- [41] C. Manuel Annals Phys. 263 (1998) 238 [hep-ph/9612494].
- [42] Y. A. Markov and M. A. Markova, Theor. Math. Phys. 103 (1995) 444.
- [43] M. Shifman and A. Yung Supersymmetric Solitons (2009) Cambridge University Press. ISBN: 0521516382.
- [44] M. Shifman Advanced Topics in Quantum Field Theory. (2012) Cambridge University Press. ISBN 978-0521190848.
- [45] Y. Shnir Magnetic Monopoles (2008) Springer. SBN-10: 3540808841.