

HECKE ORBITS ON SHIMURA VARIETIES OF HODGE TYPE

MARCO D'ADDEZIO AND POL VAN HOF TEN

ABSTRACT. We prove the Hecke orbit conjecture of Chai–Oort for Shimura varieties of Hodge type at primes of good reduction, under a mild assumption on the size of the prime. Our proof uses a new generalisation of Serre–Tate coordinates for deformation spaces of central leaves in such Shimura varieties, constructed using work of Caraiani–Scholze and Kim. We use these coordinates to give a new interpretation of Chai–Oort’s notion of strongly Tate-linear subspaces of these deformation spaces. This lets us prove upper bounds on the local monodromy of these subspaces using the Cartier–Witt stacks of Bhatt–Lurie. We also prove a rigidity result in the style of Chai–Oort for strongly Tate-linear subspaces. Another main ingredient of our proof is a new result on the local monodromy groups of F -isocrystals “coming from geometry”, which should be of independent interest.

CONTENTS

1. Introduction	1
2. Background	5
3. Local monodromy of F -isocrystals	11
4. Automorphism groups of p -divisible groups and Dieudonné–Lie algebras	16
5. Deformation theory of central leaves	26
6. Rigidity	37
7. Proof of the main theorem and some variants	46
Appendix A. Complete restricted perfection, after Chai–Oort	52
References	59

1. INTRODUCTION

1.1. The Hecke orbit conjecture. Let p be a prime number and g a positive integer. Problem 15 on Oort’s 1995 list of open problems in algebraic geometry, [61], is the following conjecture.

Conjecture 1. *Let $x = (A_x, \lambda)$ be an $\overline{\mathbb{F}}_p$ -point of the moduli space \mathcal{A}_g of principally polarised abelian varieties of dimension g over $\overline{\mathbb{F}}_p$. The Hecke orbit of x , consisting of all points $y \in \mathcal{A}_g(\overline{\mathbb{F}}_p)$ corresponding to principally polarised abelian varieties related to (A_x, λ) by symplectic isogenies, is Zariski dense in the Newton stratum of \mathcal{A}_g containing x .*

Date: May 23, 2022.

2010 Mathematics Subject Classification. Primary 11G18; Secondary 14G35.

More generally, for the special fibre of a Shimura variety of Hodge type at a prime of good reduction one expects that the isogeny classes are Zariski dense in the Newton strata containing them. This article contains a proof of this expectation under the assumption that p is not too small with respect to the given Shimura datum.

1.1.1. There is a refined version of Conjecture 1, also due to Oort, which considers instead the prime-to- p Hecke-orbit of x , consisting of all $y \in \mathcal{A}_g(\overline{\mathbb{F}}_p)$ related to x by prime-to- p symplectic isogenies. In this case, the quasi-polarised p -divisible group $(A_x[p^\infty], \lambda)$ is constant on prime-to- p Hecke orbits (not simply constant up to isogeny). Therefore, the prime-to- p Hecke orbit of x is contained in the central leaf

$$C(x) = \{y \in \mathcal{A}_g(\overline{\mathbb{F}}_p) \mid A_y[p^\infty] \simeq_\lambda A_x[p^\infty]\},$$

where \simeq_λ denotes a symplectic isomorphism. Oort proved in [60] that $C(x)$ is a smooth closed subvariety of the Newton stratum of \mathcal{A}_g containing x . He also conjectured that the prime-to- p Hecke orbit of x was Zariski dense in the central leaf $C(x)$. This conjecture is known as the *Hecke orbit conjecture* (for \mathcal{A}_g). Thanks to Mantovan–Oort product formula, [52, 60], the Hecke orbit conjecture implies Conjecture 1.

Central leaves and prime-to- p Hecke orbits can be also defined for the special fibres of Shimura varieties of Hodge type at primes of good reduction. The Hecke orbit conjecture for Shimura varieties of Hodge type then predicts that the prime-to- p Hecke orbits of points are Zariski dense in the central leaves containing them (see question 8.2.1 of [51] and Conjecture 3.2 of [14]).

The Hecke orbit conjecture naturally splits up into a discrete part and a continuous part. The discrete part states that the prime-to- p Hecke orbit of x intersects each connected component of $C(x)$, whereas the continuous part states that the Zariski closure of the prime-to- p Hecke orbit of x is of the same dimension as $C(x)$. The discrete part of the conjecture is Theorem C of [51] (see [39] for related results). In this paper, we will focus instead on the continuous part of the conjecture.

1.2. Main result. Let (G, X) be a Shimura datum of Hodge type with reflex field E , and assume for simplicity that G^{ad} is \mathbb{Q} -simple throughout this introduction. Let p be a prime such that $G_{\mathbb{Q}_p}$ is quasi-split and split over an unramified extension, let $U_p \subseteq G(\mathbb{Q}_p)$ be a hyperspecial subgroup, and let $U^p \subseteq G(\mathbb{A}_f^p)$ be a sufficiently small compact open subgroup. We choose a place v of E dividing p and we write Sh_G for the geometric special fibre of the canonical integral model over $\mathcal{O}_{E,v}$ of the Shimura variety $\mathbf{Sh}_{U^p U_p}$ of level over E . The canonical integral model exists by work of Kisin [49]. Let $C \subseteq \text{Sh}_G$ be a central leaf as constructed in [32] (c.f. [45]).

Theorem I (Theorem 7.3.2). *If $Z \subseteq C$ is a nonempty reduced closed subvariety that is stable under the prime-to- p Hecke operators and $p \geq h(G)$, then $Z = C$.*

When $\text{Sh}_G = \mathcal{A}_g$ this result, without any conditions on p , is due to Chai–Oort, see their forthcoming book [17] for the continuous part and [19] for the discrete part. Their proofs do not generalise to more general Shimura varieties because they rely on the existence of hypersymmetric points in Newton strata, which is usually false for Shimura varieties of Hodge type. Moreover, their proof of the continuous part of the conjecture relies on the fact that any point $x \in \mathcal{A}_g(\overline{\mathbb{F}}_p)$ is contained in a large Hilbert modular variety, and they use work of Chai–Oort–Yu on the Hecke orbit conjecture for Hilbert modular varieties at (possibly ramified) primes. There are many other partial results e.g. for prime-to- p Hecke orbits of hypersymmetric points in the PEL case, [70], or for prime-to- p Hecke orbits of ordinary points, [10, 38, 55, 64, 72].

We also prove that isogeny classes are dense in the Newton strata containing them, see Theorem 7.4.1. Moreover, we prove results about the ℓ -adic Hecke orbits for primes $\ell \neq p$ generalising work of Chai, [13], in the Siegel case, see Theorem 7.6.1.

Remark 1.2.1. The condition that $p \geq h(G)$, as well as the assumption that G^{ad} is \mathbb{Q} -simple, can be relaxed at the expense of introducing more notation, see Theorem 7.3.2 for a precise statement.

Remark 1.2.2. Theorem 8.10 of [8], which gives a potentially good reduction criterion for K3 surfaces, is conditional on the Hecke orbit conjecture for certain orthogonal Shimura varieties, see Conjecture 8.2 of [*ibid.*]. In Section 7.5 we explain that our results can be used to prove this conjecture under mild assumptions on p .

1.3. Local monodromy of F -isocrystals. One of the main tools of the proof is the theory of *monodromy groups of F -isocrystals*, defined in [21]. We will prove a new result about the local monodromy groups of F -isocrystals “coming from geometry”, which should be of independent interest. This result is used to prove that the local monodromy groups of the crystalline Dieudonné modules of the universal p -divisible groups over $Z \subseteq C$ (notation as in Theorem I) are “big”. We explain it in a more general setting.

Let X be a smooth irreducible variety over a perfect field with a rational point x (for simplicity) and let $(\mathcal{M}^\dagger, \Phi_{\mathcal{M}^\dagger})$ be a semi-simple overconvergent F -isocrystal over X with constant Newton polygon. Since the Newton polygon is constant, the associated F -isocrystal $(\mathcal{M}, \Phi_{\mathcal{M}})$ admits the slope filtration. Let $X^{/x} := \text{Spf } \hat{\mathcal{O}}_{X,x}$ be the formal spectrum of the complete local ring of X at x . The main result of [23] tells us that the *monodromy group* of \mathcal{M} (Definition 2.3.4), denoted by $G(\mathcal{M}, x)$, is the parabolic subgroup $P \subseteq G(\mathcal{M}^\dagger, x)$ associated to the slope filtration. We refine that result for the local monodromy group at x .

Theorem II. *The monodromy group $G(\mathcal{M}_{X^{/x}}, x)$ of the restriction of \mathcal{M} to $X^{/x}$ is the unipotent radical of the monodromy group $G(\mathcal{M}, x)$ of \mathcal{M} over X .*

When \mathcal{M} is the crystalline Dieudonné-module of an ordinary p -divisible group, then this result is proved by Chai in [11], by doing explicit computations with Serre–Tate coordinates. Our proof builds instead on the techniques developed in [23] and uses new descent results for isocrystals from [27, 53]. Since $X^{/x}$ is geometrically simply connected, each isocrystal underlying an isoclinic F -isocrystal over $X^{/x}$ is trivial by Theorem 6 of [2]. This already implies that $G(\mathcal{M}_{X^{/x}}, x)$ is unipotent. To relate $G(\mathcal{M}_{X^{/x}}, x)$ and the unipotent radical of $G(\mathcal{M}, x)$ we pass through the respective generic points. Write k for the function field of X and k_x for the function field of $X^{/x}$. We first prove in Theorem 3.2.5 that when we pass to $\text{Spec } k$ we get the same monodromy group as the global one over X . Then we also show that if we pass instead to the separable closure of k we obtain the unipotent radical of $G(\mathcal{M}, x)$ (Proposition 3.4.2). This means that passing to k^{sep} we kill precisely a Levi subgroup of $G(\mathcal{M}, x)$. Subsequently, using the fact that the field extension $k \subseteq k_x$ is a separable extension such that k and k_x share a common (absolute) p -basis (Lemma 3.4.1), we show that when we pass from k^{sep} to k_x^{sep} we do not get extra splitting of the slope filtration (Proposition 3.3.5). This is enough to prove that the local monodromy group $G(\mathcal{M}_{X^{/x}}, x)$ is the same as the monodromy group over k^{sep} , which in turn is the unipotent radical of the monodromy group $G(\mathcal{M}, x)$.

1.4. An overview of the proof of Theorem I. The overall structure of the proof of Theorem I is similar to the proof of the ordinary Hecke orbit conjecture in [38] and is based on a strategy implicit in the work of Chai–Oort and sketched to us by Chai in a letter.

To explain the proof we first need to establish some notation. Let $Z \subseteq C \subseteq \text{Sh}_{G,b}$ be a reduced closed subvariety that is stable under the prime-to- p Hecke operators as in the statement of Theorem I. The Newton stratum $\text{Sh}_{G,b} \subseteq \text{Sh}_G$ corresponds to an element $b \in B(G_{\mathbb{Q}_p})$ which has an associated

Newton (fractional) cocharacter ν_b . Attached to this cocharacter is a parabolic subgroup P_{ν_b} with unipotent radical U_{ν_b} .

Let Z^{sm} be the smooth locus of Z . Then Corollary 2.2.1 of [38] tells us that the monodromy of the crystalline Dieudonné module \mathcal{M} of the universal p -divisible group over Z^{sm} is isomorphic to P_{ν_b} . Theorem II then tells us that for $x \in Z^{\text{sm}}(\overline{\mathbb{F}}_p)$, the monodromy of \mathcal{M} over $Z/x := \text{Spf } \widehat{\mathcal{O}}_{Z,x}$ is equal to U_{ν_b} . We are going to leverage this fact to show that $Z/x = C/x$, which will allow us to conclude that Z^{sm} and hence Z is equidimensional of the same dimension as C .

For this we will construct generalised Serre–Tate coordinates on the formal completion $C/x := \text{Spf } \widehat{\mathcal{O}}_{C,x}$. To be precise, we will show that there is a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra¹ \mathfrak{a}^+ over $\check{\mathbb{Z}}_p$ governing the structure of C/x . For example if C is the ordinary locus in $\mathcal{A}_{g,N}$ then C/x is a p -divisible formal group by the classical theory of Serre–Tate coordinates, and $\mathfrak{a}^+ = \mathbb{D}(C/x)$ is its Dieudonné module, equipped with the trivial Lie bracket.

More generally the perfection of C/x admits the structure of a (functor to) nilpotent Lie \mathbb{Q}_p -algebra(s) whose Dieudonné module is $\mathfrak{a} := \mathfrak{a}^+[\frac{1}{p}]$. More canonically, the perfection of C/x is a trivial torsor for a unipotent formal group $\tilde{\Pi}(\mathfrak{a})$ related to the aforementioned nilpotent Lie algebra by the Baker–Campbell–Hausdorff formula. In the Siegel case, this unipotent formal group is the identity component of the group of self-quasi isogenies (compatible with the polarisation up to a scalar) of the p -divisible group $A_x[p^\infty]$. These results come from the perspective of Caraiani–Scholze, [9], on C and the perspective of Kim, [45], on C/x .

We note that our generalisation of Serre–Tate coordinates is different from the notion of a p -divisible cascade given by Moonen in [58]. Our alternative definition is necessary in our work since in the Hodge type case the deformation spaces we consider do not have in general a cascade structure. Already in the μ -ordinary case, all we can hope for is a *shifted subcascade* in the sense of [65] (c.f. [40]) and, as far as we can see, there is no way to run our arguments with shifted subcascades.

The Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra \mathfrak{a} turns out to be isomorphic to $\text{Lie } U_{\nu_b}$ equipped with a natural F -structure coming from b . If $\mathfrak{b} \subseteq \mathfrak{a} = \text{Lie } U_{\nu_b}$ is an F -stable Lie subalgebra, we construct a formally smooth closed formal subscheme $Z(\mathfrak{b}^+) \subseteq C/x$ (Definition 4.3.14), which is strongly Tate-linear in the sense of Chai–Oort, [16]. The construction is such that when $\mathfrak{b} = \mathfrak{a}$, the formal subscheme $Z(\mathfrak{b}^+)$ is C/x itself. It turns out that Z/x admits such a description.

Theorem 1.4.1 (Theorem 6.1.1). *There is an F -stable Lie subalgebra $\mathfrak{b} \subseteq \mathfrak{a}$ such that $Z/x = Z(\mathfrak{b}^+)$.*

To prove Theorem I, we are then reduced to show that the formal subschemes $Z(\mathfrak{b}^+)$ for $\mathfrak{b} \subsetneq \mathfrak{a}$ have small monodromy. In this direction we prove the following result:

Theorem 1.4.2 (Theorem 5.5.3). *The Lie algebra of the monodromy group of \mathcal{M} over $Z(\mathfrak{b}^+)$ is contained in \mathfrak{b} .*

By the previous discussion, we know that the Lie algebra of the monodromy group of \mathcal{M} over Z/x is equal to \mathfrak{a} . Therefore, if $Z/x = Z(\mathfrak{b}^+)$ for some \mathfrak{b} , then $\mathfrak{a} \subseteq \mathfrak{b} \subseteq \mathfrak{a}$, so that $Z/x = Z(\mathfrak{b}^+) = Z(\mathfrak{a}^+) = C/x$.

The proof of Theorem 1.4.2 uses the Cartier–Witt stacks of Bhatt–Lurie [4] in combination with the interpretation of C/x as a formal deformation space of the trivial torsor for a certain group scheme $\Pi(\mathfrak{a}^+)$, due to Chai–Oort [16] in the PEL case. In particular, we show that the closed formal

¹See Definition 4.3.2.

subscheme $Z(\mathfrak{b}^+) \subseteq C^{/x}$ can be identified with the formal deformation space of the trivial torsor for a certain closed subgroup $\Pi(\mathfrak{b}^+) \subseteq \Pi(\mathfrak{a}^+)$. Our argument with Cartier–Witt stacks happens in Section 5.5.

Theorem 1.4.1 is related to rigidity results for p -divisible formal groups of Chai, [15], and to rigidity results for biextensions of p -divisible formal groups of Chai–Oort, [20]. We would also like to mention unpublished work of Tao Song which proves rigidity results in the case of p -divisible 4-cascades.

The proof of Theorem 1.4.1, which happens in Section 6, was inspired by the proof of the rigidity result for biextensions of [20] and uses some of the result of [*ibid.*] on the topological commutative algebra of completions of perfection of power series rings over $\overline{\mathbb{F}}_p$, which are summarised in Appendix A. However, it is not entirely straightforward to translate the proofs of [20], which happen with biextensions, to the setting of unipotent groups and Dieudonné–Lie algebras.

Remark 1.4.3. The unipotent formal group $\tilde{\Pi}(\mathfrak{a})$ for $\mathfrak{a} = \text{Lie } U_{\nu_b}$ is closely related to the “unipotent group diamond” $\tilde{G}_b^{>0}$ introduced in Chapter III.5 of Fargues–Scholze, [30].

1.5. Structure of the article. In Section 2 we cover some background theory on p -divisible groups, isocrystals, formal schemes and Cartier–Witt stacks. We prove Theorem II in Section 3. In Section 4 we discuss internal hom p -divisible groups and groups of (quasi-)automorphisms of p -divisible groups, and introduce Dieudonné–Lie algebras. In Section 5 we discuss central leaves for Shimura varieties of Hodge type and describe their deformation theory in terms of Dieudonné–Lie algebras using work of Kim, [45]. We then relate our perspective on the deformation central leaves to the one of Chai–Oort, [16], and prove Theorem 1.4.2. Section 6 is devoted entirely to the proof of Theorem 1.4.1.

1.6. Acknowledgements. We are indebted to Ching-Li Chai and Frans Oort for the ideas they developed and we used for our work on the Hecke orbit conjecture. The first named author would also like to thank Ching-Li Chai for his letter on the relation between the parabolicity conjecture and the Hecke orbit conjecture. We are grateful to Brian Conrad, Sean Cotner, Andrew Graham, and Richard Taylor for many helpful discussions about this work. We also thank H el ene Esnault and Matteo Tamiozzo for their comments on an earlier version of this article.

The first named author was funded by the Deutsche Forschungsgemeinschaft (DA-2534/1-1, project ID: 461915680).

2. BACKGROUND

2.1. Formal schemes over a field. In this text we will repeatedly make use of the theory of formal schemes, as introduced in [26, §10]. The admissible topological rings we will consider will be *adic rings* with a finitely generated ideal of definition.

Definition 2.1.1. A *pre-adic* ring* is a topological ring R endowed with the I -adic topology for some finitely generated ideal I . An *adic* ring* is an I -adically complete pre-adic* ring.

If R is an adic* ring over a field κ (not asked to be perfect) we write $h_{\text{Spf } R} : \text{Alg}_{\kappa}^{\text{op}} \rightarrow \text{Sets}$ for the associated functor of point, where $\text{Alg}_{\kappa}^{\text{op}}$ is the opposite of the category of κ -algebras endowed with the fpqc topology. If S is another adic* ring, by [68, Lemma 0AN0], a homomorphism of functors $f : h_{\text{Spf } S} \rightarrow h_{\text{Spf } R}$ comes uniquely from a continuous homomorphism $f : R \rightarrow S$.

2.1.2. *Scheme-theoretic images.* Let $f : R \rightarrow S$ be a morphism of adic* rings with kernel K (a closed ideal of R by the completeness of S). If I is a finitely generated ideal of definition of R , then we can form the ring

$$T := \varprojlim_n R/(I^n + K),$$

which is just the J -adic completion of R/K where $J := I/K$. The ring T is J -adically complete because J is finitely generated. By Lemma 0APT, the morphism $R \rightarrow T$ is surjective. This implies that $T = R/K$, or in other words, that R/K is J -adically complete. Note that this implies that T is a subring of S , so that if S is a domain, the same is true for T .

Definition 2.1.3. The closed immersion $\mathrm{Spf} T \hookrightarrow \mathrm{Spf} R$ is called the *scheme-theoretic image* of $f : \mathrm{Spf} S \rightarrow \mathrm{Spf} R$.

We present now some technical lemmas on formal schemes over κ that we will use later on. We would like to thank Brian Conrad and Sean Cotner for suggesting the proofs we propose here (although any errors are due to the authors).

Lemma 2.1.4. *Suppose that $\mathrm{Spf} R$ and $\mathrm{Spf} S$ are endowed with an action of a group scheme G/κ and let $f : \mathrm{Spf} S \rightarrow \mathrm{Spf} R$ be a G -equivariant morphism of formal schemes over κ . There is a unique action of G on the scheme-theoretic image $\mathrm{Spf} T$ such that the induced maps $\mathrm{Spf} T \rightarrow \mathrm{Spf} R$ and $\mathrm{Spf} S \rightarrow \mathrm{Spf} T$ are G -equivariant.*

Proof. Let $C = \Gamma(S)$ be the coordinate ring of G , then there is a commutative diagram (all the tensor products are over κ)

$$\begin{array}{ccccc} R & \longrightarrow & R/K & \longrightarrow & S \\ \downarrow & & \downarrow & & \downarrow \\ R \widehat{\otimes} C & \longrightarrow & R/K \widehat{\otimes} C & \longrightarrow & S \widehat{\otimes} C, \end{array}$$

where K is the kernel of $f : R \rightarrow S$. We are trying to show that there is a unique way to fill in the dotted arrow, which will follow from the following claim.

Claim 2.1.5. *The map $R/K \widehat{\otimes} C \rightarrow S \widehat{\otimes} C$ is injective.*

Showing that the induced map $\mathrm{Spf} R/K = T \rightarrow G \times T$ satisfies the axioms of a group action is straightforward. \square

Proof of Claim 2.1.5. The completed tensor product of R/K with C is the I -adic completion of the usual tensor product $R/K \otimes C$, where I is an ideal of definition of R (and thus also an ideal of definition of R/K). As a module over κ we know that $C = \kappa^{\oplus \mathcal{A}}$ for some index set \mathcal{A} and since tensor products commute with direct sums we see that $R/K \otimes C = (R/K)^{\oplus \mathcal{A}}$. The \mathfrak{m}_R -adic completion of this tensor product is the subset of the product of R/K over \mathcal{A} , consisting of those sequences where for all $m \geq 0$ almost all elements of the sequence lie in I^m .

Let J be an ideal of definition of S containing $f(I)$, then the completion $S \widehat{\otimes} C$ is the J -adic completion of $S \otimes C$. This can be identified with the subset of the product of S over \mathcal{A} , consisting of those sequences where for all $m \geq 0$ almost all elements of the sequence lie in J^m . Since $R/K \rightarrow S$ is injective we see that the map

$$\prod_{\mathcal{A}} R/K \rightarrow \prod_{\mathcal{A}} S$$

is injective and hence the map $R/K\widehat{\otimes}C \rightarrow S\widehat{\otimes}C$ is injective. \square

Lemma 2.1.6. *Let $(A, \mathfrak{m}), (B, \mathfrak{n})$ be Noetherian complete local rings over κ , let C, D over κ be adic rings and let $A \rightarrow C$ and $B \rightarrow D$ be adic and injective maps of κ -algebras. Then the induced map (where the tensor products are over κ)*

$$A\widehat{\otimes}B \rightarrow C\widehat{\otimes}D$$

is injective.

Proof. The map of A -modules

$$A \otimes B \rightarrow C \otimes D$$

is injective because we are working over a field. For $j \geq 0$ let $I_j = \mathfrak{m}_A^j C$, then the topology on C is defined by the chain of ideals I_j and since the topology on C is Hausdorff by assumption we find that $\bigcap_j I_j = (0)$. This means that we similarly have $\bigcap_j (I_j \cap A) = 0$ and because A is Noetherian, Chevalley's theorem (see exercise 8.7 of [54] and its solution on page 290) says that for each j there is an $n(j)$ such that $(I_{n(j)} \cap A) \subseteq \mathfrak{m}_A^j$. By construction $\mathfrak{m}_A^j \subseteq (I_j \cap A)$ and so we have shown that the \mathfrak{m}_A -adic topology on A is equal to the linear topology induced by the chain of ideals $(I_1 \cap A) \supseteq (I_2 \cap A) \supseteq \dots$. In other words,

$$A = \varprojlim_j A/(I_j \cap A),$$

as a topological κ -vector space, and similarly

$$B = \varprojlim_j B/(J_j \cap B)$$

where $J_j = \mathfrak{m}_B^k D$. By construction the maps $A/(I_j \cap A) \rightarrow C/I_j$ and $B/(J_j \cap B) \rightarrow D/J_j$ are injective. It follows that for each (j, j') the map

$$A/(I_j \cap A) \otimes B/(J_{j'} \cap B) \rightarrow C/I_j \otimes D/J_{j'}$$

is injective because it factors as a composition of the maps

$$A/(I_j \cap A) \otimes B/(J_{j'} \cap B) \rightarrow A/I_j \otimes D/J_{j'} \rightarrow C/I_j \otimes D/J_{j'}$$

which are injective because we are over a field. By the definition of the completed tensor product the map that we are trying to show is injective is

$$\varprojlim_{j, j'} (A/(I_j \cap A) \otimes B/(J_{j'} \cap B)) \rightarrow \varprojlim_{j, j'} (C/I_j \otimes D/J_{j'}).$$

This is an inverse limit of injective maps and thus injective. \square

Lemma 2.1.7. *Let (A, \mathfrak{m}) and (B, \mathfrak{n}) be complete (Noetherian) local rings over κ with residue fields isomorphic to κ . Let $f : \mathrm{Spf} B \rightarrow \mathrm{Spf} A$ be a monomorphism of formal schemes over κ , then it is a closed immersion.*

Proof. We have a local homomorphism of κ -algebras $f : (A, \mathfrak{m}) \rightarrow (B, \mathfrak{n})$ that induces an isomorphism on residue fields and such that for every Artinian local ring R the induced map $\mathrm{Hom}(B, R) \rightarrow \mathrm{Hom}(A, R)$ is injective.

Let $R = B/(\mathfrak{m}B + \mathfrak{n}^j)$ and assume that $\mathfrak{m}B + \mathfrak{n}^j$ is strictly contained in \mathfrak{n} . Then the natural map $B \rightarrow R$ does not factor through B/\mathfrak{n} . Note that there is another natural map $B \rightarrow R$ given by

$$B \rightarrow B/\mathfrak{n} = A/\mathfrak{m} \rightarrow B/\mathfrak{m}B \rightarrow R,$$

and these two natural maps are different by our assumption. But they agree after precomposition with A , so this contradicts the injectivity of $\mathrm{Hom}(B, R) \rightarrow \mathrm{Hom}(A, R)$, and thus $\mathfrak{m}_B + \mathfrak{n}^j = \mathfrak{n}$.

Lemma 0AMS of [68] then tells us that the closure of the ideal $\mathfrak{m}B$ is equal to \mathfrak{n} and it follows that the closure of the ideal $\mathfrak{m}^j B$ is equal to \mathfrak{n}^j . Therefore the map f is taut and thus by Lemma 0APU of [68] the topology on B is given by the \mathfrak{m} -adic topology. Thus B/\mathfrak{m}_B^j has the discrete topology and it follows that the following diagram is Cartesian

$$\begin{array}{ccc} \mathrm{Spec} B/\mathfrak{m}_B^j & \longrightarrow & \mathrm{Spf} B \\ \downarrow & & \downarrow \\ \mathrm{Spec} A/\mathfrak{m}^j & \longrightarrow & \mathrm{Spf} A. \end{array}$$

This tells us that $f : \mathrm{Spf} B \rightarrow \mathrm{Spf} A$ is representable, and Lemma 0GHZ of [68] allows us to conclude that f is a closed immersion. \square

2.2. Cartier–Witt stacks. In this section we want to briefly recall the main properties of the *Cartier–Witt stacks*. To simplify the exposition, we will deal only with *quasi-syntomic schemes*. Let p be a prime number and X a scheme over \mathbb{F}_p .

Definition 2.2.1. We say that X is *quasi-syntomic* if the relative cotangent complex $L\Omega_{X/\mathbb{F}_p}^1$ has Tor-amplitude in $[-1, 0]$.

Recall that if X is of finite type over a perfect field then X is quasi-syntomic if and only if X is a local complete intersection. Moreover, any regular Noetherian scheme is quasi-syntomic by Example C.8 of [3].

2.2.2. Write $\mathrm{Nil}_{\mathbb{Z}_p}^{\mathrm{op}} \subseteq \mathrm{Alg}_{\mathbb{Z}_p}^{\mathrm{op}}$ for the full subcategory of p -nilpotent algebras with the fpqc topology. A *p -adic formal stack* is a groupoid valued functor \mathcal{F} on $\mathrm{Nil}_{\mathbb{Z}_p}^{\mathrm{op}}$ whose diagonal is representable by formal algebraic spaces and which admits an fpqc cover $\mathcal{X} \rightarrow \mathcal{F}$, where \mathcal{X} is a p -adic formal algebraic space over $\mathrm{Spf} \mathbb{Z}_p$ (cf. [68, Definition 0AIM]).

Bhatt and Lurie define a functor

$$X \mapsto X^\Delta$$

which goes from the category of quasi-syntomic \mathbb{F}_p -schemes to the category of p -adic formal stacks endowed with an endomorphism $F : X^\Delta \rightarrow X^\Delta$, lifting the Frobenius on the special fibre².

2.2.3. For every quasi-syntomic scheme X , Proposition 8.15 of [4] tells us that there is an equivalence between the category of crystals in quasi-coherent \mathcal{O} -modules on the absolute prismatic site of X (or the absolute crystalline site by Example 4.7 of [6]) and quasi-coherent \mathcal{O} -modules on the Zariski site of X^Δ .

There are a few important properties of this functor that we will use.

- If $X = \mathrm{Spec} R$ is a semiperfect quasi-syntomic scheme, then X^Δ is simply $\mathrm{Spf} A_{\mathrm{cris}}(R)$, the formal spectrum of Fontaine’s ring of crystalline periods (this is Lemma 6.1 of [4], see Proposition 4.1.3 of [63] for a definition of $A_{\mathrm{cris}}(R)$). For example $(\mathrm{Spec} \mathbb{F}_p)^\Delta = \mathrm{Spf} \mathbb{Z}_p$.

²They also define a derived version of this functor, which we will not use in this text.

- If $f : X \rightarrow Y$ is a quasi-syntomic cover in the sense of Section 4 of [5], then $f : X^\Delta \rightarrow Y^\Delta$ is an fpqc cover (this is Proposition 7.5 of [4]). For example, this means that $X^\Delta \rightarrow (\mathrm{Spec} \mathbb{F}_p)^\Delta = \mathrm{Spf} \mathbb{Z}_p$ is automatically flat.
- The functor commutes with products and with fibre products in the case that the structure maps are flat and quasi-syntomic by Remark 8.9 of [4] and Proposition 7.5 of [4].

2.3. Frobenius-smooth schemes and the Tannakian category of isocrystals. Among all quasi-syntomic schemes, there is a class of them where the prismaticisation has better properties.

Definition 2.3.1. We say that a scheme X over \mathbb{F}_p is *Frobenius-smooth* if $F : X \rightarrow X$ is syntomic.

By Lemma 2.1.1.(2) of [27] this is equivalent to X being Zariski locally of the form $\mathrm{Spec} B$ where B has a finite (absolute) p -basis. In other words there exist elements x_1, \dots, x_n such that every element $b \in B$ can be uniquely written as

$$b = \sum b_\alpha^p x_1^{\alpha_1} \cdots x_n^{\alpha_n},$$

where $0 \leq \alpha_i \leq p-1$ and where $b_\alpha \in B$. The main examples of Frobenius-smooth schemes we will encounter are smooth schemes over perfect fields and power series rings over perfect fields. If X is a Noetherian Frobenius-smooth scheme, then it is regular by a result of Kunz (see Lemma 0EC0 of [68]), thus in particular quasi-syntomic.

Definition 2.3.2. If X is a scheme over \mathbb{F}_p we denote by $\mathrm{Isoc}(X)$ the category of crystals of finitely generated quasi-coherent \mathcal{O} -modules on the absolute crystalline site of X with Hom-sets tensored by \mathbb{Q} .

Following [27], we define

$$\kappa := \bigcap_{i=1}^{\infty} \Gamma(X, \mathcal{O}_X)^{p^i}.$$

to be the *ring of constants* of X and we write K for $W(\kappa)[\frac{1}{p}]$. Note that if X is a geometrically connected scheme of finite type over a perfect field then κ coincides with the base field.

Proposition 2.3.3 ([27, Corollary 3.3.3]). *If X is an irreducible Noetherian Frobenius-smooth scheme, then $\mathrm{Isoc}(X)$ is a K -linear Tannakian category.*

This allows us to define the monodromy groups of isocrystals in this situation.

Definition 2.3.4. Let X be a scheme satisfying the assumptions in Proposition 2.3.3 and let \mathcal{M} be an isocrystal over X . We define $\langle \mathcal{M} \rangle$ to be the Tannakian subcategory of $\mathrm{Isoc}(X)$ generated by \mathcal{M} . If η is an Ω -point for some perfect field Ω , we define $G(\mathcal{M}, \eta)$ to be the Tannaka group of $\langle \mathcal{M} \rangle$ with respect to the fibre functor induced by η . We call it the *monodromy group of \mathcal{M}* with respect to η .

2.4. p -divisible groups and Dieudonné-theory. Let R be a semiperfect \mathbb{F}_p -algebra and let $A_{\mathrm{cris}}(R)$ be Fontaine's ring of crystalline periods with $\varphi : A_{\mathrm{cris}}(R) \rightarrow A_{\mathrm{cris}}(R)$ induced by the absolute Frobenius on R .

Definition 2.4.1. A *Dieudonné module* over R is a pair (M^+, φ_{M^+}) , where M^+ is a finite locally free $A_{\text{cris}}(R)$ -module and where

$$\varphi_{M^+} : \varphi^* M[\frac{1}{p}] \rightarrow M[\frac{1}{p}]$$

is an isomorphism such that

$$M^+ \subseteq \varphi_{M^+}(M^+) \subseteq \frac{1}{p}M^+.$$

Remark 2.4.2. Usually one instead asks that

$$pM^+ \subseteq \varphi_{M^+}(M^+) \subseteq M^+;$$

our conventions agree, for example, with the ones in [9]. A p -divisible group X over R has a covariant Dieudonné-module $\mathbb{D}(X), \varphi_X$, normalised as in [9]. In particular this means that the Dieudonné-module of $\mathbb{Q}_p/\mathbb{Z}_p$ over R is $A_{\text{cris}}(R)$ equipped with the trivial Frobenius, and the Dieudonné-module of μ_{p^∞} is $A_{\text{cris}}(R)$ equipped with Frobenius given by $1/p$.

If X and X' are p -divisible groups over R then there is a natural map

$$\text{Hom}_R(X, X') \rightarrow \text{Hom}_\varphi(\mathbb{D}(X), \mathbb{D}(X')),$$

where the right hand side denotes homomorphisms of $A_{\text{cris}}(R)$ -modules that intertwine φ_X and $\varphi_{X'}$. Theorem 4.8.5 of [1] tells us that this natural map is an isomorphism if R is quasi-syntomic and semiperfect. Note that the theorem is stated for contravariant Dieudonné-theory with the usual normalisation, but this does not matter.

Definition 2.4.3 (c.f. Definition 1.1.4 of [1]). We call an \mathbb{F}_p -scheme *qrs*p (quasi-regular semiperfect) if it is quasi-syntomic and semiperfect.

The main example is

$$\text{Spec } \kappa[X_1^{1/p^\infty}, \dots, X_m^{1/p^\infty}]/(X_1, \dots, X_m)$$

where κ is a perfect field.

2.4.4. We recall that a connected p -divisible group Y over $\overline{\mathbb{F}}_p$, considered as a functor on $\overline{\mathbb{F}}_p$ -algebras, is representable by the formal spectrum of a power series ring over $\overline{\mathbb{F}}_p$ (see e.g. Lemma 3.1.1 of [63] for a more general result). The p -adic Tate-module

$$T_p Y := \varprojlim_n Y[p^n]$$

where the transition maps are given by multiplication by p , is representable by an affine scheme. Proposition 4.12.(4) of [9] tells us that this affine scheme is isomorphic to the spectrum of

$$\overline{\mathbb{F}}_p[X_1^{1/p^\infty}, \dots, X_m^{1/p^\infty}]/(X_1, \dots, X_m)$$

for some m ; note that this ring is qrs. If Y is connected then the universal cover

$$\tilde{Y} := \varprojlim Y,$$

where the transition maps are given by multiplication by p , is representable by the formal spectrum of the completed perfection of a power series ring by Proposition 3.1.3.(iii) of [63]. Therefore \tilde{Y} is a filtered colimit of spectra of semiperfect rings and so it is determined by its values on semiperfect rings.

2.4.5. Both \tilde{Y} and Y are determined by their restriction to the category of semiperfect $\overline{\mathbb{F}}_p$ -algebras, and we can describe them explicitly on the category of qrsp $\overline{\mathbb{F}}_p$ -algebras as follows.

Lemma 2.4.6. *There is a commutative diagram of natural transformation of functors on the category of qrsp $\overline{\mathbb{F}}_p$ -algebras, which evaluated at an object R gives*

$$\begin{array}{ccc} T_p Y(R) & \xrightarrow{\cong} & \left(A_{\text{cris}}(R) \otimes_{\mathbb{Z}_p} \mathbb{D}(Y) \right)^{\varphi=1} \\ \downarrow & & \downarrow \\ \tilde{Y}(R) & \xrightarrow{\cong} & \left(B_{\text{cris}}^+(R) \otimes_{\mathbb{Z}_p} \mathbb{D}(Y)[\frac{1}{p}] \right)^{\varphi=1}, \end{array}$$

where φ is given by the diagonal Frobenius.

Proof. Let R be qrsp, then Theorem 4.8.5 of [1] tells us that Dieudonné-module functor gives us a natural isomorphism

$$\begin{aligned} T_p Y(R) &= \text{Hom}_R((\mathbb{Q}_p/\mathbb{Z}_p)_R, Y_R) \\ &\rightarrow \text{Hom}_{A_{\text{cris}}, F}(A_{\text{cris}}(R), A_{\text{cris}}(R) \otimes_{\mathbb{Z}_p} \mathbb{D}(Y)) \\ &\simeq \left(A_{\text{cris}}(R) \otimes_{\mathbb{Z}_p} \mathbb{D}(Y) \right)^{\varphi=1} \end{aligned}$$

where the latter bijection is induced by evaluation at 1. Similarly after inverting p we get a natural isomorphism

$$\begin{aligned} \tilde{Y}(R) &= \text{Hom}_R((\mathbb{Q}_p/\mathbb{Z}_p)_R, Y_R[\frac{1}{p}]) \\ &\rightarrow \left(B_{\text{cris}}^+(R) \otimes_{\mathbb{Q}_p} \mathbb{D}(Y)[\frac{1}{p}] \right)^{\varphi=1} \end{aligned}$$

and the diagram commutes by construction. \square

3. LOCAL MONODROMY OF F -ISOCRYSTALS

This section is an interlude on monodromy groups of isocrystals in which we prove Theorem II. We first get some general results on Tannakian categories and isocrystals and then we use them in Section 3.4 to prove the main theorem.

3.1. A Tannakian criterion. Let K be a field and let V be a finite-dimensional K -vector space.

Lemma 3.1.1. *If U, U' are unipotent subgroups of $\text{GL}(V)$, then the following two properties are equivalent.*

- (i) $U' \subseteq U$.
- (ii) $W^U \subseteq W^{U'}$ for every algebraic representation W of $\text{GL}(V)$.

Proof. The implication (i) \Rightarrow (ii) is obvious. To prove (ii) \Rightarrow (i) just note that by Chevalley's theorem there is a representation W of $\text{GL}(V)$ such that U is the stabiliser of a line $L \subseteq W$. Since U does not admit non-trivial characters, we deduce that $L \subseteq W^U$. Thanks to (ii), this implies that U' fixes L and this yields the desired result. \square

Thanks to this lemma, we get a Tannakian criterion to prove that a unipotent subgroup of an algebraic group coincides with the unipotent radical. This will be the criterion that we will use to prove Theorem II.

Proposition 3.1.2. *Let $U \subseteq G \subseteq \mathrm{GL}(V)$ be a chain of subgroups of $\mathrm{GL}(V)$ and suppose that there exists a cocharacter $\nu : \mathbb{G}_m \rightarrow \mathrm{GL}(V)$ such that both U and $R_u(G)$ are in U_ν . The following two properties are equivalent.*

- (i) $U = R_u(G)$.
- (ii) *For every representation W of $\mathrm{GL}(V)$, the group G stabilises W^U and the induced representation factors through $G/R_u(G)$.*

Proof. (1) \Rightarrow (2) follows from the observation that $R_u(G)$ is normal in G . For (2) \Rightarrow (1) first note that by the assumptions $U \subseteq R_u(G)$ since $R_u(G) = G \cap U_\nu$. For the other inclusion, thanks to (2) we deduce that for every representation W we have that $W^U \subseteq W^{R_u(G)}$. Thus by Lemma 3.1.1, we conclude that $R_u(G) \subseteq U$. \square

3.2. Passing to the generic point. Another ingredient that we will need for Theorem II is a way to pass from varieties to fields. This is provided by Theorem 3.2.5, that we want to prove here.

Let A_0 be an \mathbb{F}_p -algebra and let $A_\infty = \varinjlim_{i \in I} A_i$ be a filtered colimit of A_0 -algebras, where A_0 is the initial object of the system. We write I' for $I \cup \{\infty\}$ and for a crystal in quasi-coherent modules \mathcal{M}_0 over $\mathrm{Spec} A_0/\mathbb{Z}_p$ we consider for every $i \in I'$ the base changes of \mathcal{M}_0 to $\mathrm{Cris}(\mathrm{Spec} A_i/\mathbb{Z}_p)$, denoted by \mathcal{M}_i .

Lemma 3.2.1. *For every $j \geq 0$, we have*

$$H^j(\mathrm{Cris}(\mathrm{Spec} A_\infty/\mathbb{Z}_p), \mathcal{M}_\infty) = \left(\varinjlim_{i \in I} H^j(\mathrm{Cris}(\mathrm{Spec} A_i/\mathbb{Z}_p), \mathcal{M}_i) \right)^\wedge.$$

Proof. For $i \in I'$, let P_i be the free commutative \mathbb{Z}_p -algebra associated to the set underlying A_i , so that $P_\infty = \varinjlim_{i \in I} P_i$. We write $P_i(n)$ (resp. $A_i(n)$) for the tensor product of $n+1$ copies of P_i (resp. A_i) over \mathbb{Z}_p (resp. \mathbb{F}_p), and $J_i(n)$ for the kernel of $P_i(n) \rightarrow A_i(n)$. We also write $D_i(n)$ for the p -adic completion of the divided power envelope of $P_i(n)$ with respect to the ideal $J_i(n)$ and for every $i \in I'$, $n \geq 0$, and $e > 0$ we write $M_{i,e}(n)$ for

$$\mathcal{M}_i(A_i(n), D_i(n)/p^e, \gamma_{i,e}(n)),$$

where $\gamma_{i,e}(n)$ is the natural divided power structure on $J_i(n)/p^e$. The cohomology of \mathcal{M}_i is computed as the cohomology of the complex $\varprojlim_e M_{i,e}(\bullet)$. Since $D_\infty(n)/p^e = \varinjlim_{i \in I} D_i(n)/p^e$, we deduce that for each $e > 0$ we have that $M_{\infty,e}(\bullet) = \varinjlim M_{i,e}(\bullet)$. This yields the desired result. \square

Proposition 3.2.2. *Let X be an irreducible Noetherian Frobenius-smooth scheme over \mathbb{F}_p with generic point η . For every isocrystal \mathcal{M} over X , we have*

$$H^0(\eta, \mathcal{M}_\eta) = \varinjlim_{U \subseteq X} H^0(U, \mathcal{M}_U),$$

where the colimit is over the dense open subschemes of X .

Proof. By Lemma 3.2.1, we have that $H^0(\eta, \mathcal{M}_\eta) = \left(\varinjlim_{U \subseteq X} H^0(U, \mathcal{M}_U) \right)^\wedge$. Let κ be the field of constants of X . We note that κ is the field of constants of every $U \subseteq X$ and η . Therefore, if K is the fraction field of the ring of Witt vectors of κ , each group $H^0(U, \mathcal{M}_U)$ is a K -linear subspace of $H^0(\eta, \mathcal{M}_\eta)$. Since η is Frobenius-smooth, this implies that $\varinjlim_{U \subseteq X} H^0(U, \mathcal{M}_U)$ is a finite dimensional K -vector space as well. In particular, it is already p -adically complete. This ends the proof. \square

Suppose now that X is a smooth irreducible variety over a perfect field with generic point η .

Proposition 3.2.3. *If \mathcal{N} is a subquotient of an isocrystal \mathcal{M} over X such that $F^*\mathcal{M} \simeq \mathcal{M}$, then $H^0(\eta, \mathcal{N}_\eta) = H^0(X, \mathcal{N})$.*

Proof. By Theorem 5.10 of [29], we know that \mathcal{N} is a subobject of some isocrystal \mathcal{M}' such that $F^*\mathcal{M}' \simeq \mathcal{M}'$. Thanks to [*ibid.*, Lemma 5.6] it is enough to prove the result for \mathcal{M}' . By [28, Theorem 2.2.3], for every dense open $U \subseteq X$, we have that $H^0(X, \mathcal{M}) = H^0(U, \mathcal{M}_U)$. Therefore, thanks to Proposition 3.2.2 we deduce that $H^0(\eta, \mathcal{M}'_\eta) = H^0(X, \mathcal{M}')$. \square

Proposition 3.2.4. *If $(\mathcal{N}_\eta, \Phi_{\mathcal{N}_\eta})$ is an isoclinic subobject of $(\mathcal{M}_\eta, \Phi_{\mathcal{M}_\eta})$, then $\mathcal{N}_\eta \subseteq \mathcal{M}_\eta$ extends to a subobject of \mathcal{M} .*

Proof. It is enough to prove the result after replacing $\Phi_{\mathcal{M}}$ with some power and after twisting $(\mathcal{M}, \Phi_{\mathcal{M}})$ by a rank 1 F -isocrystal. Therefore we may assume that $(\mathcal{N}_\eta, \Phi_{\mathcal{N}_\eta})$ is a unit-root F^n -isocrystal. We have that $(\mathcal{N}_\eta, \Phi_{\mathcal{N}_\eta})$ is a subobject of the unit-root part of the graded object associated to the slope filtration of $(\mathcal{M}_\eta, \Phi_{\mathcal{M}_\eta})$, denoted by $(\mathcal{N}'_\eta, \Phi_{\mathcal{N}'_\eta})$. By [2, Theorem 6], the F^n -isocrystals $(\mathcal{N}_\eta, \Phi_{\mathcal{N}_\eta})$ and $(\mathcal{N}'_\eta, \Phi_{\mathcal{N}'_\eta})$ correspond to lisse \mathbb{Q}_p^n -sheaves V and V' over η . Since $(\mathcal{M}, \Phi_{\mathcal{M}})$ admits the slope filtration, V' extends to a lisse sheaf over X . The lisse sheaf V , being a subobject of V' , extends as well over X . This implies that $(\mathcal{N}_\eta, \Phi_{\mathcal{N}_\eta})$ extends to an F^n -isocrystal $(\mathcal{N}, \Phi_{\mathcal{N}})$ over X . Write \mathcal{H} for the isocrystal $\underline{\text{Hom}}(\mathcal{N}, \mathcal{M})$. \square

Theorem 3.2.5. *For every F -isocrystal $(\mathcal{M}, \Phi_{\mathcal{M}})$ over X with constant Newton polygon we have that $\langle \mathcal{M} \rangle \rightarrow \langle \mathcal{M}_\eta \rangle$ is an equivalence of Tannakian categories. In other words, $G(\mathcal{M}, \eta^{\text{perf}}) = G(\mathcal{M}_\eta, \eta^{\text{perf}})$.*

Proof. The natural functor $\langle \mathcal{M} \rangle \rightarrow \langle \mathcal{M}_\eta \rangle$ is a functor between Tannakian categories. Thanks to Proposition A.3 and A.4.1 of [29], to show that it is an equivalence we have to prove that it is fully faithful and that every rank 1 object $\mathcal{L}_\eta \in \langle \mathcal{M}_\eta \rangle$ is a direct summand of a semi-simple isocrystal coming from $\langle \mathcal{M} \rangle$. For the first part it is enough to prove that the functor induces isomorphisms at the level of global sections, which follows from Proposition 3.2.3. To prove the second part we note that a rank 1 isocrystal $\mathcal{L}_\eta \in \langle \mathcal{M}_\eta \rangle$ is the subquotient of some isocrystal $\mathcal{M}'_\eta \in \langle \mathcal{M}_\eta \rangle$ coming from $\langle \mathcal{M} \rangle$ which can be endowed with a Frobenius structure with constant Newton polygon. Also, taking the subquotient of \mathcal{M}'_η associated to some slope, we may assume that \mathcal{L}_η is actually a subobject of \mathcal{M}'_η and that the latter isocrystal can be endowed with an isoclinic Frobenius structure. Write $\mathcal{N}_\eta \subseteq \mathcal{M}'_\eta$ for the sum of rank 1 subobjects of \mathcal{M}'_η . By construction, $\mathcal{N}_\eta \subseteq \mathcal{M}'_\eta$ is kept stable by the Frobenius structure of \mathcal{M}'_η . By virtue of Proposition 3.2.4, this implies that isocrystal \mathcal{N}_η comes from an isocrystal $\mathcal{N} \in \langle \mathcal{M} \rangle$. This ends the proof. \square

3.3. Descent for isocrystals. We prove now various descent results that we will need in the next section. Let $f : Y \rightarrow X$ be a pro-étale Π -cover of Noetherian Frobenius-smooth schemes over \mathbb{F}_p , where Π is a profinite group and let $y \in Y(\Omega)$ be an Ω -point of Y with Ω a perfect field. Write K for the fraction field of the ring of Witt vectors of Ω .

Lemma 3.3.1. *For every isocrystal \mathcal{M} over X , the maximal trivial subobject of $f^*\mathcal{M}$ descends to a subobject $\mathcal{N} \subseteq \mathcal{M}$. Moreover, if \mathcal{M} is endowed with a Frobenius structure $\Phi_{\mathcal{M}}$, the inclusion $\mathcal{N} \subseteq \mathcal{M}$ upgrades to an inclusion $(\mathcal{N}, \Phi_{\mathcal{N}}) \subseteq (\mathcal{M}, \Phi_{\mathcal{M}})$ of F -isocrystals and $(\mathcal{N}, \Phi_{\mathcal{N}})$ is a direct sum of isoclinic objects.*

Proof. Since the cover $Y \rightarrow X$ is a quasi-syntomic cover, it satisfies descent for isocrystals by the descent result in Proposition 3.5.4 of [27] (see also [53] or Section 2 of [6]). By the assumption,

$$Y \times_X Y \simeq \varinjlim_{U \subseteq \Pi} (Y \times_X Y)^U$$

where the limit runs over all the open normal subgroups of Π and $(Y \times_X Y)^U := \coprod_{[\gamma] \in \Pi/U} Y_{[\gamma]}$ is a disjoint union of copies of Y . The group Π acts on $Y \times_X Y$ in the obvious way. Since $f^* \mathcal{M}$ descends to X , it is endowed with a descent datum with respect to the cover $Y \rightarrow X$. This datum consists of isomorphisms $\gamma^* \mathcal{M}_{(Y \times_X Y)^U} \simeq \mathcal{M}_{(Y \times_X Y)^U}$ for each $U \subseteq \Pi$ and $\gamma \in \Pi$. The functor γ^* sends trivial objects to trivial objects, which implies that the descent datum restricts to a descent datum for \mathcal{T} , the maximal trivial subobject of $f^* \mathcal{M}$. Therefore, \mathcal{T} descends to a subobject $\mathcal{N} \subseteq \mathcal{M}$, as we wanted. If \mathcal{M} is endowed with a Frobenius structure, then it induces a Frobenius structure on each isocrystal $\mathcal{M}_{(Y \times_X Y)^U}$ and this has to preserve each maximal trivial subobject of given slope. This implies that also the descended object $\mathcal{N} \subseteq \mathcal{M}$ is stabilised by the Frobenius and the induced Frobenius structure satisfies the desired property. \square

Proposition 3.3.2. *Let $(\mathcal{M}, \Phi_{\mathcal{M}})$ be an F -isocrystal which admits the slope filtration and write ν for the associated Newton cocharacter. If $R_u(G(\mathcal{M}, f(y))) \subseteq U_{\nu}$ and $\mathrm{Gr}_{S_{\bullet}}(f^* \mathcal{M})$ is trivial, then $G(f^* \mathcal{M}, y) = R_u(G(\mathcal{M}, f(y)))$.*

Proof. Since $\mathrm{Gr}_{S_{\bullet}}(f^* \mathcal{M})$ is trivial, the group $G(f^* \mathcal{M}, y)$ is a unipotent subgroup of $G(\mathcal{M}, f(y)) \otimes_{K'} K$ sitting inside U_{ν} . Therefore, we are in the situation of Proposition 3.1.2 and we have to prove that (ii) is satisfied. This amounts to show that for every $m, n \geq 0$, the maximal trivial subobject $\mathcal{T} \subseteq f^*(\mathcal{M}^{\otimes m} \otimes (\mathcal{M}^{\vee})^{\otimes n})$ descends to a semi-simple isocrystal over X . By Lemma 3.3.1, we know that \mathcal{T} descends to an isocrystal \mathcal{N} which is the direct sum of isocrystals which can be endowed with an isoclinic Frobenius structure. Since $R_u(G(\mathcal{M}, f(y)))$ is contained in U_{ν} , we deduce that \mathcal{N} is semi-simple, as we wanted. \square

Lemma 3.3.3. *If k'/k is a separable field extension and k' admits a finite p -basis, then $k' \otimes_k k'$ admits a finite p -basis as well.*

Proof. Thanks to [54, Theorem 26.6], the field k admits a finite p -basis t_1, \dots, t_d which extends to a finite p -basis $t_1, \dots, t_d, u_1, \dots, u_e$ of k' . We claim that $\Gamma := \{t_i \otimes 1\}_i \cup \{u_i \otimes 1\}_i \cup \{1 \otimes u_i\}_i$ is a finite p -basis of $k' \otimes_k k'$. It is clear from the construction that the elements of Γ generate $k' \otimes_k k'$ over $(k' \otimes_k k')^p$. On the other hand, the exact sequence

$$0 \rightarrow \Omega_{k/\mathbb{F}_p}^1 \otimes_k (k' \otimes_k k') \rightarrow \Omega_{k' \otimes_k k'/\mathbb{F}_p}^1 \rightarrow (\Omega_{k'/k}^1 \otimes_k k') \oplus (k' \otimes_k \Omega_{k'/k}^1) \rightarrow 0$$

shows that the elements $d\gamma$ with $\gamma \in \Gamma$ form a basis of the free module $\Omega_{k' \otimes_k k'/\mathbb{F}_p}^1$. We deduce the p -independence of the elements of Γ by arguing as in Lemma 07P2 of [68]. \square

Lemma 3.3.4. *If X is a Frobenius-smooth scheme over \mathbb{F}_p , every locally free F -isocrystal with constant Newton polygon which does not admit slope 0 has no global sections fixed by the Frobenius structure.*

Proof. Since X is Frobenius-smooth, the global sections of any isocrystal over X embed into the global sections of the base change to X^{perf} . Over X^{perf} one can argue as in the proof of [44, Theorem 2.4.2]. We reduce to the case when $X^{\mathrm{perf}} = \mathrm{Spec} A$ with A perfect. Then we can embed A into a product of perfect fields. This reduces the problem to the case of perfect fields where the result is well-known. \square

Proposition 3.3.5. *Let $k \subseteq k'$ a separable extension of characteristic p fields with finite p -basis and let $(\mathcal{M}, \Phi_{\mathcal{M}})$ be a free F -isocrystal over k with slope filtration S_{\bullet} of length n . If $\mathcal{M}_{k'}$ admits a Frobenius-stable splitting of the form $\mathcal{N}_{k'} \oplus S_{n-1}(\mathcal{M}_{k'})$ with $\mathcal{N}_{k'}$ some subobject of $\mathcal{M}_{k'}$, the same is true for \mathcal{M} .*

Proof. Since $\mathrm{Spec} k' \rightarrow \mathrm{Spec} k$ is a quasi-syntomic cover, it satisfies descent for isocrystals. Indeed, this follows from the descent results of Drinfeld and Mathew in [27, 53] (see Theorem 2.2 of [6]). Therefore, in order to descend $\mathcal{N}_{k'}$ to k it is enough to show that the splitting $\mathcal{N}'_{k' \otimes_k k'} \oplus S_{n-1}(\mathcal{M}'_{k' \otimes_k k'})$ is unique. Suppose that $\mathcal{N}'_{k' \otimes_k k'} \oplus S_{n-1}(\mathcal{M}'_{k' \otimes_k k'})$ was a different splitting. Then there would exist a non-trivial Frobenius-equivariant morphism $\mathcal{N}'_{k' \otimes_k k'} \rightarrow S_{n-1}(\mathcal{M}'_{k' \otimes_k k'})$. In other words, the F -isocrystal $\underline{\mathrm{Hom}}(\mathcal{N}'_{k' \otimes_k k'}, S_{n-1}(\mathcal{M}'_{k' \otimes_k k'}))$ would have a non-trivial Frobenius-invariant global section. Since the slopes of $\underline{\mathrm{Hom}}(\mathcal{N}'_{k' \otimes_k k'}, S_{n-1}(\mathcal{M}'_{k' \otimes_k k'}))$ are all negative by definition and $k' \otimes_k k'$ admits a finite p -basis by 3.3.3, this would contradict Lemma 3.3.4. \square

3.4. The local monodromy theorem. We are ready to put all the previous results together and prove Theorem II. Let X be a smooth irreducible variety over a perfect field and let x be a closed point of X . We denote by k the function field of X and by k_x the function field of X^x . We also write η^{sep} (resp. $\bar{\eta}$) for the points over the generic point of X associated to a separable (resp. algebraic) closure of k .

Lemma 3.4.1. *The fields k and k_x have a common finite p -basis. In particular, $k \subseteq k_x$ is a separable field extension.*

Proof. By [54, Theorem 26.7], it is enough to show that $\Omega_{k/\mathbb{F}_p}^1 \otimes_k k_x = \Omega_{k_x/\mathbb{F}_p}^1$. Write A for the local ring of X at x and A_x^\wedge for the completion with respect to the maximal ideal \mathfrak{m}_x . Since A is regular, thanks to Theorem 30.5 and Theorem 30.9 of [54], we deduce that $\Omega_{A/\mathbb{F}_p}^1 \otimes_A A_x^\wedge = \Omega_{A_x^\wedge/\mathbb{F}_p}^1$. We get the desired result after inverting $\mathfrak{m}_x - \{0\}$. \square

Proposition 3.4.2. *If $(\mathcal{M}, \Phi_{\mathcal{M}})$ is an F -isocrystal over X such that $R_u(G(\mathcal{M}, \bar{\eta})) \subseteq U_\nu$, then $G(\mathcal{M}_{\eta^{\mathrm{sep}}}, \bar{\eta}) = R_u(G(\mathcal{M}, \bar{\eta}))$.*

Proof. By Theorem 3.2.5 we have that $G(\mathcal{M}, \bar{\eta}) = G(\mathcal{M}_\eta, \bar{\eta})$, so that we are reduced to prove the statement for $G(\mathcal{M}_\eta, \bar{\eta})$. Note that the cover $f : \eta^{\mathrm{sep}} \rightarrow \eta$ is a pro-étale $\mathrm{Gal}(k^{\mathrm{sep}}/k)$ -cover and $\mathrm{Gr}_{S_{\bullet}}(f^* \mathcal{M}_\eta)$ is trivial because η^{sep} is simply connected. This shows that we can apply Proposition 3.3.2 and deduce the desired result. \square

Proposition 3.4.3. *If $(\mathcal{M}, \Phi_{\mathcal{M}})$ is an F -isocrystal over X coming from an irreducible overconvergent F -isocrystal with constant Newton polygon, then $H^0(X^x, (S_1(\mathcal{M}))^x) = H^0(X^x, \mathcal{M}^x)$.*

Proof. By Galois descent we may assume that the ring of constants of X is an algebraically closed field. The inclusion $H^0(X^x, (S_1(\mathcal{M}))^x) \subseteq H^0(X^x, \mathcal{M}^x)$ is an inclusion of F -isocrystals over κ . We suppose by contradiction that this is not an equality. Let $s_r > s_1$ be the greatest slope appearing in $H^0(X^x, \mathcal{M}^x)$ and let v be a non-zero vector such that $\Phi_{\mathcal{M}^x}^n(v) = p^{s_r n} v$ for $n \gg 0$. Write $(\tilde{\mathcal{M}}, \Phi_{\tilde{\mathcal{M}}})$ for the base change of $(\mathcal{M}, \Phi_{\mathcal{M}})$ to η^{sep} .

By the parabolicity conjecture, [23], $R_u(G(\mathcal{M}, \bar{\eta}))$ is contained in U_ν because $(\mathcal{M}, \Phi_{\mathcal{M}})$ comes from an irreducible overconvergent F -isocrystal. Proposition 3.4.2 then implies that the monodromy group $G(\tilde{\mathcal{M}}, \bar{\eta})$ is equal to $G(\mathcal{M}, \bar{\eta}) \cap U_\nu$. Therefore, the line spanned by v determines a rank 1

subobject $\tilde{\mathcal{L}} \subseteq S_r(\tilde{\mathcal{M}})/S_{r-1}(\tilde{\mathcal{M}})$ stabilised by the Frobenius. The preimage of this isocrystal in $S_r(\tilde{\mathcal{M}})$, denoted by $\tilde{\mathcal{N}}$, is kept invariant by the Frobenius and sits in an exact sequence

$$0 \rightarrow S_{r-1}(\tilde{\mathcal{M}}) \rightarrow \tilde{\mathcal{N}} \rightarrow \tilde{\mathcal{L}} \rightarrow 0.$$

Since $(\mathcal{M}, \Phi_{\mathcal{M}})$ comes from an irreducible overconvergent F -isocrystal, the sequence does not admit a Frobenius-equivariant splitting by [23, Theorem 4.1.3]. By Proposition 3.3.5, the base change of this extension to k_x^{sep} does not split as well. This leads to a contradiction since v is a vector in $H^0(X/x, \mathcal{M}/x)$ which produces a non-trivial global section of $\tilde{\mathcal{N}} \subseteq \tilde{\mathcal{M}}$. \square

We write η_x for the generic point of X/x and $G(\mathcal{M}/x, \eta_x^{\text{perf}})$ for the monodromy group³ of \mathcal{M}/x with respect to the perfection of η_x .

Theorem 3.4.4. *If $(\mathcal{M}, \Phi_{\mathcal{M}})$ comes from a semi-simple overconvergent F -isocrystal with constant Newton polygon, then*

$$G(\mathcal{M}/x, \eta_x^{\text{perf}}) = R_u(G(\mathcal{M}, \eta_x^{\text{perf}})).$$

Proof. Write G for the group $G(\mathcal{M}, \eta)$ and V for the induced G -representation. By [23] we have that $R_u(G)$ is contained in U_{ν} where ν is the Newton cocharacter. Since X/x is geometrically simply connected we deduce that $\text{Gr}_{S_{\bullet}}(\mathcal{M})/x$ is trivial. This implies that $G(\mathcal{M}/x, \eta_x^{\text{perf}}) \subseteq R_u(G) \subseteq U_{\nu}$. Therefore, in order to apply the criterion of Proposition 3.1.2 it is enough to show that for every $\mathcal{N} \in \langle \mathcal{M} \rangle$, the space of global sections of \mathcal{N}/x is the same as the fibre at x of some direct sum of isoclinic subobjects of \mathcal{N} . To prove this, we may assume that \mathcal{N} can be endowed with a Frobenius structure $\Phi_{\mathcal{N}}$ and $(\mathcal{N}, \Phi_{\mathcal{N}})$ is irreducible. Thanks to Proposition 3.4.3, we deduce that the fibre of $S_1(\mathcal{N})$ at x is the same as $H^0(X/x, \mathcal{N}/x)$. This yields the desired result. \square

4. AUTOMORPHISM GROUPS OF p -DIVISIBLE GROUPS AND DIEUDONNÉ–LIE ALGEBRAS

The goal of this section is to define various groups of tensor preserving automorphisms and endomorphisms of p -divisible groups with G -structure, correcting some definitions from [45]. Once this is done, we will axiomatise these groups in terms of Dieudonné–Lie $\tilde{\mathbb{Z}}_p$ -algebras and prove various properties of these groups using this point of view.

4.1. Endomorphisms and automorphisms. We closely follow Section 3 of [16] and also Section 4 of [9] and [45]. For p -divisible groups Y and Z over $\overline{\mathbb{F}}_p$, Chai and Oort construct finite group schemes

$$\underline{\text{Hom}}^{\text{st}}(Y[p^n], Z[p^n]) \subseteq \underline{\text{Hom}}(Y[p^n], Z[p^n])$$

together with maps

$$\pi_n : \underline{\text{Hom}}^{\text{st}}(Y[p^n], Z[p^n]) \rightarrow \underline{\text{Hom}}^{\text{st}}(Y[p^{n+1}], Z[p^{n+1}])$$

such that (the additive group underlying)

$$\varinjlim_n \underline{\text{Hom}}^{\text{st}}(Y[p^n], Z[p^n])$$

is a p -divisible group $\mathcal{H}_{Y,Z}$, called the *internal Hom p -divisible group*, with p -adic Tate module isomorphic to the scheme

$$\underline{\text{Hom}}(Y, Z)$$

³Arguing as in [24, Proposition 2.4.8] one can show that this monodromy group is the same whether you consider isocrystals over the formal scheme X/x or over the associated scheme.

of homomorphisms from Y to Z . Lemma 4.1.7 of [9] tells us that there is an isomorphism

$$\mathbb{D}(\mathcal{H}_{Y,Z})[\frac{1}{p}] = \underline{\mathbf{Hom}}(\mathbb{D}(Y)[\frac{1}{p}], \mathbb{D}(Z)[\frac{1}{p}])^{\leq 0},$$

where $(-)^{\leq 0}$ denotes the operation of taking the subspace of slope ≤ 0 and where

$$\underline{\mathbf{Hom}}(\mathbb{D}(Y)[\frac{1}{p}], \mathbb{D}(Z)[\frac{1}{p}]).$$

denotes the internal hom in the category of F -isocrystals. By Lemma 4.1.8 of [9] there is an isomorphism of formal group schemes

$$\tilde{\mathcal{H}}_{Y,Z} \simeq \underline{\mathbf{Hom}}(Y, Z)[\frac{1}{p}],$$

where $\tilde{\mathcal{H}}_{Y,Z}$ is the universal cover of $\mathcal{H}_{Y,Z}$ in the sense of Scholze–Weinstein (see Section 2.4).

We will mostly be interested in $\mathcal{H}_{Y,Y} =: \mathcal{H}_Y$ for a p -divisible group Y in which case $T_p \mathcal{H}_Y$ and $\mathbb{D}(\mathcal{H}_Y)$ have an algebra structure, and the same for their rational variants. Up to isogeny we can write $Y = \bigoplus_i Y_i$ as a direct sum of isoclinic p -divisible groups with slopes in increasing order and then we can write endomorphisms of \tilde{Y} in matrix form to get

$$\tilde{\mathcal{H}}_Y = \bigoplus_{i,j} \tilde{\mathcal{H}}_{Y_i, Y_j}.$$

Corollary 4.1.10 of [9] tells us that the p -divisible groups H_{Y_i, Y_j} are zero when $i > j$, they are étale p -divisible groups when $i = j$ and connected p -divisible groups when $i < j$. This means that we get a lower triangular matrix form (see the proof of Proposition 4.2.11 of [9]), and that the connected part

$$\tilde{\mathcal{H}}_Y^\circ = \bigoplus_{i < j} \tilde{\mathcal{H}}_{Y_i, Y_j}$$

consists of nilpotent endomorphisms. The étale part is precisely the locally profinite group scheme associated to the \mathbb{Q}_p -algebra given by

$$\underline{\mathbf{End}}(Y)(\overline{\mathbb{F}}_p)[\frac{1}{p}].$$

In order to generalise away from GL_n , it will be more fruitful to use the commutator bracket on $\underline{\mathbf{Hom}}(Y, Y)[\frac{1}{p}]$ to equip $\tilde{\mathcal{H}}_Y$ with the structure of a Lie algebra. To precise it is a Lie algebra over the locally profinite ring-scheme $\underline{\mathbb{Q}}_p$ associated to the topological ring \mathbb{Q}_p . Here if V is a topological space we use the notation \underline{V} for the functor on $\overline{\mathbb{F}}_p$ -schemes sending $T \mapsto \mathrm{Hom}_{\mathrm{cont}}(|T|, V)$, where $|T|$ is the topological space underlying the scheme T . The functor \underline{V} is representable by a finite scheme if V is finite and discrete, and therefore also representable if T is profinite or locally profinite.

There is a descending sequence of Lie $\underline{\mathbb{Q}}_p$ -algebra ideals

$$\tilde{\mathcal{H}}_Y^\circ = \mathrm{Fil}^0 \supseteq \mathrm{Fil}^1 \supseteq \mathrm{Fil}^2 \supseteq \dots$$

where Fil^k consists of those endomorphisms that lie in

$$\bigoplus_{j \geq i+k} \tilde{\mathcal{H}}_{Y_i, Y_j} \subseteq \bigoplus_{i,j} \tilde{\mathcal{H}}_{Y_i, Y_j}.$$

Note that the graded quotients are isomorphic to the abelian Lie $\underline{\mathbb{Q}}_p$ -algebra

$$\mathrm{Fil}^k / \mathrm{Fil}^{k+1} \simeq \bigoplus_i \tilde{\mathcal{H}}_{Y_i, Y_{i+k}}.$$

If we recall that $\mathbb{D}(H_Y^\circ)[\frac{1}{p}]$ is an isocrystal over $\check{\mathbb{Q}}_p$ equipped with a Lie bracket preserving F , then the filtration $\mathrm{Fil}_{\mathbb{D}}^\bullet$ given by

$$\mathrm{Fil}_{\mathbb{D}}^k = \mathbb{D}(\mathrm{Fil}^k)$$

is its lower central series.

4.1.1. *Automorphisms.* Let $\mathbf{Aut}(\tilde{Y})$ be functor of automorphisms of \tilde{Y} , then there is a monomorphism

$$\begin{aligned} \alpha : \mathbf{Aut}(\tilde{Y}) &\rightarrow \left(\tilde{\mathcal{H}}_Y\right)^{\oplus 2} \\ \gamma &\mapsto (\gamma, \gamma^{-1}) \end{aligned}$$

and the image consists of those pairs (γ, γ') such that $\gamma \circ \gamma' = 1 = \gamma' \circ \gamma$. It follows from this that α is representable by closed immersions and therefore $\mathbf{Aut}(\tilde{Y})$ is a formal algebraic space. If we intersect with the subscheme $\underline{\mathbf{Hom}}(Y, Y)^{\oplus 2} = T_p \mathcal{H}_Y^{\oplus 2}$ we recover the group scheme $\mathbf{Aut}(Y)$. The projection to the diagonal map

$$\mathbf{Aut}(\tilde{Y}) \rightarrow \underline{\mathbf{Aut}(\tilde{Y})(\overline{\mathbb{F}}_p)},$$

has connected kernel $\mathbf{Aut}(\tilde{Y})^\circ$. This implies that we have a semi-direct product decomposition

$$\mathbf{Aut}(\tilde{Y}) \simeq \mathbf{Aut}(\tilde{Y})^\circ \rtimes \underline{\mathbf{Aut}(\tilde{Y})(\overline{\mathbb{F}}_p)}.$$

4.1.2. Our matrix description implies that all elements γ of $\mathbf{Aut}(\tilde{Y})^\circ$ are unipotent automorphisms of \tilde{Y} . This means that the logarithm map

$$\begin{aligned} L : \mathbf{Aut}(\tilde{Y})^\circ &\rightarrow \tilde{\mathcal{H}}_Y^\circ \\ X &\mapsto \sum_{i=1}^{\infty} (-1)^{i+1} \frac{(X-1)^i}{i} \end{aligned}$$

and the exponential map

$$\begin{aligned} E : \tilde{\mathcal{H}}_Y^\circ &\rightarrow \mathbf{Aut}(\tilde{Y})^\circ \\ X &\mapsto \sum_{i=0}^{\infty} \frac{X^i}{i!} \end{aligned}$$

are well defined. We know that $E \circ L = 1$ and $L \circ E = 1$ because we can check this in $\mathbb{Q}[[X]]$ by universality and there the result follows from complex analysis.

The Baker–Campbell–Hausdorff formula gives an expression for the group structure on $\mathbf{Aut}(\tilde{Y})^\circ$ in terms of the Lie bracket on $\tilde{\mathcal{H}}_Y^\circ$. In fact if V is a (possibly infinite dimensional) vector space over a field of characteristic zero and X, Y are two nilpotent endomorphisms of V , then the BCH formula expresses $\exp(X)\exp(Y)$ in terms of X and Y and the Lie bracket. This is explained in Chapter XVI of [35].

It follows from this that the filtration Fil^\bullet induces a filtration $\mathrm{Fil}_{\mathbf{Aut}}^\bullet = E(\mathrm{Fil}^\bullet)$ of $\mathbf{Aut}(\tilde{Y})^\circ$ by normal subgroups, with graded quotients isomorphic to

$$\bigoplus_i \tilde{\mathcal{H}}_{Y_i, Y_{i+k}}.$$

Indeed, we can identify the graded pieces of the two filtrations via the exponential map, and the Baker–Campbell–Hausdorff formula tells us that the exponential map of an abelian Lie algebra is an isomorphism.

4.2. Automorphism groups for Shimura varieties of Hodge type. In order to generalise the previous section to Shimura varieties of Hodge type, we let $b \in \mathrm{GL}_n(\check{\mathbb{Q}}_p)$ be the Frobenius of the F -isocrystal $\mathbb{D}(Y)[\frac{1}{p}]$ associated to our p -divisible group Y . Then the internal hom isocrystal

$$\underline{\mathrm{Hom}}(\mathbb{D}(Y)[\frac{1}{p}], \mathbb{D}(Y)[\frac{1}{p}])$$

is isomorphic to the isocrystal

$$(\mathfrak{gl}_n, \mathrm{Ad} \sigma b).$$

If we are given a reductive group $G \subseteq \mathrm{GL}_n(\mathbb{Q}_p)$ such that $b \in G(\check{\mathbb{Q}}_p)$, then we get an inclusion of F -isocrystals

$$(\mathfrak{g}, \mathrm{Ad} \sigma b) \subseteq (\mathfrak{gl}_n, \mathrm{Ad} \sigma b).$$

The slope filtration of this F -isocrystal is described in Section 3 of [45]: The slope t part is given by

$$\bigoplus_{\substack{\alpha_0 \in \Phi_0^+ \\ \langle \alpha_0, \nu_b \rangle = t}} \mathfrak{u}_{\alpha_0},$$

where α_0 runs over the relative roots of G with respect a fixed maximal split torus contained a Borel B with respect to which the Newton cocharacter ν_b of b is dominant. We see that the slope ≤ 0 part corresponds precisely to Lie algebra of the standard parabolic subgroup $P_b = P_{\nu_b}$ associated to ν_b and that the slope 0 part corresponds to the Lie algebra of the Levi M_b .

Taking non-positive slope parts we get an F -stable sub-isocrystal

$$\mathrm{Lie} P_{\nu_b} \subseteq \mathbb{D}(\mathcal{H}_Y)[\frac{1}{p}],$$

and intersecting with $\mathbb{D}(\mathcal{H}_Y)$ we get the Dieudonné module $\mathbb{D}(\mathcal{H}_Y^G)$ of a p -divisible group \mathcal{H}_Y^G .

Corollary 4.2.1. *The dimension of \mathcal{H}_Y^G is equal to $\langle 2\rho, \nu_b \rangle$.*

Proof. Its dimension can be computed as

$$\sum_{\alpha_0 \in \Phi_0^+} \langle \alpha_0, \nu_b \rangle \cdot \dim \mathfrak{u}_{\alpha_0} = \sum_{\alpha \in \Phi^+} \langle \alpha, \nu_b \rangle = \langle 2\rho, \nu_b \rangle.$$

□

It follows from Lemma 4.3.5 that

$$\tilde{\mathcal{H}}_Y^G \subseteq \tilde{\mathcal{H}}_Y = \underline{\mathrm{Hom}}(Y, Y)[\frac{1}{p}]$$

is closed under the Lie bracket (the commutator) of $\mathrm{Hom}(Y, Y)[\frac{1}{p}]$. From the description of the isocrystal above we see that the étale part of \mathcal{H}_Y^G is the locally profinite \mathbb{Q}_p -Lie algebra \mathfrak{J}_b corresponding to the Lie algebra of the twisted Levi J_b of G . Moreover the Dieudonné module

$$\mathbb{D}(\mathcal{H}_Y^{G, \circ})[\frac{1}{p}]$$

is stable under the Lie bracket and consists precisely of the strictly negative slope part of the isocrystal $\mathrm{Lie} P_{\nu_b}$, which can be identified with

$$\mathrm{Lie} U_{\nu_b} \subseteq \mathrm{Lie} P_{\nu_b}.$$

4.2.2. If $G \subseteq \mathrm{GL}_n$ is the stabiliser of a finite collection of tensors $\{s_\alpha\}_\alpha$, then $\mathfrak{g} \subseteq \mathfrak{gl}_n$ is also the stabiliser of these tensors as a Lie subalgebra. In other words, it consists of those endomorphisms g satisfying $g^*s_\alpha = 0$, let us call such endomorphisms *tensor annihilating endomorphisms*. It follows that

$$\mathrm{Lie} P_{\nu_b} \subseteq \mathbb{D}(\mathcal{H}_Y) \left[\frac{1}{p} \right]$$

consists precisely of the tensor annihilating endomorphisms. Therefore by Remark 4.3.6 it follows that for qrsp $\overline{\mathbb{F}}_p$ -algebras R we have that

$$\mathcal{H}_Y^G(R) \subseteq \mathcal{H}_Y(R)$$

precisely consists of the tensor annihilating endomorphisms. There is also the notion of *tensor preserving automorphism* which is an automorphism g such that $g^*s_\alpha = s_\alpha$. We claim that there is a closed subgroup

$$\mathbf{Aut}_G(\tilde{Y}) \subseteq \mathbf{Aut}(\tilde{Y})$$

such that on qrsp $\overline{\mathbb{F}}_p$ -algebras R the subgroup

$$\mathbf{Aut}_G(\tilde{Y})(R) \subseteq \mathbf{Aut}(\tilde{Y})(R)$$

consists precisely of those quasi-automorphisms of Y_R that induce tensor preserving automorphisms on rational Dieudonné-modules. Indeed, we know that $\mathbf{Aut}(\tilde{Y})$ is a filtered colimit of spectra of qrsp algebras and therefore we can define a subgroup as above.

Our definition does *not* agree with definition 2.3.1 of [45], which tries to define $\mathbf{Aut}_G(\tilde{Y})$ as the intersection of

$$\left(\tilde{\mathcal{H}}_Y^G \times \tilde{\mathcal{H}}_Y^G \right) \cap \mathbf{Aut}(\tilde{Y})$$

inside $\tilde{\mathcal{H}}_Y \times \tilde{\mathcal{H}}_Y$. However this cannot be a subgroup as $\tilde{\mathcal{H}}_Y^G$ does not contain the identity endomorphism unless $G = \mathrm{GL}_n$.

However it is clear from the definition that the exponential of a nilpotent tensor annihilating endomorphism is a unipotent tensor preserving automorphism, and conversely the logarithm of a unipotent tensor preserving automorphism is a nilpotent tensor annihilating endomorphism. Thus the exponential map defines an isomorphism of formal schemes

$$\mathcal{H}_Y^{G,\circ} \simeq \mathbf{Aut}_G(\tilde{Y})^\circ$$

and as before the (convergent) BCH formula expresses the group operation on the right hand side in terms of the Lie bracket on the left hand side. Similarly we can prove that

$$\mathbf{Aut}_G(\tilde{Y}) \simeq \mathbf{Aut}_G(\tilde{Y})^\circ \rtimes J_b(\mathbb{Q}_p)$$

and therefore Proposition 3.2.4. of [45] is still correct. To get a filtration on $\mathcal{H}_Y^{G,\circ}$ we will not take the filtration induced by Fil^k , but rather we will take the filtration coming from the lower central series filtration on the Lie algebra

$$\mathbb{D}(\mathcal{H}_Y^{G,\circ}) \left[\frac{1}{p} \right].$$

We will need these filtrations in more generality, so we will give an axiomatic treatment in the next section. We end this section by defining

$$\mathbf{Aut}_G(Y) = \mathbf{Aut}_G(\tilde{Y}) \cap \mathbf{Aut}(Y),$$

and noting that the image of its identity component $\mathbf{Aut}_G(Y)^\circ$ under the logarithm map is given by $T_p \mathcal{H}_Y^{G,\circ}$ if $p \gg 0$. Indeed if $p \gg 0$ then in the formula for the logarithm map we do not need to divide by p . We will make the condition on p precise in the next section as well.

4.3. Dieudonné–Lie algebras. We start by recalling the definition of a Dieudonné module over $\overline{\mathbb{F}}_p$ (see Definition 2.4.1)

Definition 4.3.1. A *Dieudonné module* over $\overline{\mathbb{F}}_p$ is a pair (M^+, φ_{M^+}) , where M^+ is a finite free $W(\overline{\mathbb{F}}_p)$ -module and where

$$\varphi_{M^+} : \varphi^* M^+[\frac{1}{p}] \rightarrow M^+[\frac{1}{p}]$$

is an isomorphism such that

$$M^+ \subseteq \varphi_{M^+}(M^+) \subseteq \frac{1}{p}M^+.$$

Definition 4.3.2. A *Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra* is a triple $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+}, [-, -])$ where $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+})$ is a Dieudonné module and where

$$[-, -] : \mathfrak{a}^+ \times \mathfrak{a}^+ \rightarrow \mathfrak{a}^+$$

is a Lie bracket, such that the following diagram commutes

$$\begin{array}{ccc} \varphi^* \mathfrak{a}^+ \times \varphi^* \mathfrak{a}^+ & \xrightarrow{\varphi_{\mathfrak{a}^+} \times \varphi_{\mathfrak{a}^+}} & \varphi^* \mathfrak{a}^+ \times \mathfrak{a}^+ \\ \downarrow \varphi^*[-, -] & & \downarrow [-, -] \\ \varphi^* \mathfrak{a}^+ & \xrightarrow{\varphi_{\mathfrak{a}^+}} & \mathfrak{a}^+. \end{array}$$

A morphism of Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebras is a $\check{\mathbb{Z}}_p$ -linear map $f : \mathfrak{a}^+ \rightarrow \mathfrak{b}^+$ that respects the Lie brackets and induces a homomorphism of Dieudonné modules. We refer to the isogeny category of Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebras as *Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebras*. We write $(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$ for the Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra associated to $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+}, [-, -])$.

The main example that we will see is

$$\mathfrak{a}^+ = \mathbb{D}(\mathcal{H}_Y^{G, \circ})$$

equipped with the Frobenius $\varphi_{\mathfrak{a}^+}$ coming from the Dieudonné module structure and the Lie-bracket coming from the algebra structure on

$$\mathbb{D}(\mathcal{H}_Y).$$

We moreover note that the F -isocrystal $(\mathfrak{a}, \varphi_{\mathfrak{a}})$ underlying a Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra has slopes contained in $[-1, 0]$. We say that a Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra $(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$ is nilpotent if the underlying Lie algebra $(\mathfrak{a}, [-, -])$ is nilpotent, which is the case in our example. Note that it follows from the definition that the lower central series

$$\mathfrak{a} \supseteq [\mathfrak{a}, \mathfrak{a}] \supseteq [\mathfrak{a}, [\mathfrak{a}, \mathfrak{a}]] \supseteq \dots$$

is a filtration by F -stable Lie subalgebras, which we will write as

$$\mathfrak{a} = \mathfrak{a}_0 \supseteq \mathfrak{a}_1 \supseteq \dots \mathfrak{a}_{n(\mathfrak{a})} = 0.$$

The integer $n(\mathfrak{a})$ is called the *nilpotency class* of \mathfrak{a} . We will write

$$\mathfrak{a}_i^+ = \mathfrak{a}^+ \cap \mathfrak{a}_i,$$

which gives us a filtration of \mathfrak{a}^+ by Dieudonné–Lie $\check{\mathbb{Z}}_p$ -subalgebras which satisfy

$$[\mathfrak{a}^+, \mathfrak{a}_i^+] \subseteq \mathfrak{a}_{i+1}^+.$$

Moreover the graded quotients $\mathfrak{a}_i^+ / \mathfrak{a}_{i+1}^+$ again have the structure of Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebras.

Lemma 4.3.3. *Let $(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$ be a Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra where all the slopes are strictly smaller than 0. Let μ_1 be the smallest slope of \mathfrak{a} and let $\mathfrak{b} \subseteq \mathfrak{a}$ be the maximal Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra that is isoclinic of that slope. Then \mathfrak{b} is contained in the centre of \mathfrak{a} .*

Proof. There are no nonzero F -equivariant maps

$$\mathfrak{b} \otimes \mathfrak{a} \rightarrow \mathfrak{a}$$

because all the slopes of $\mathfrak{b} \otimes \mathfrak{a}$ are strictly smaller than the slopes of \mathfrak{a} since \mathfrak{a} has strictly negative slopes. Hence the restriction of the Lie-bracket to $\mathfrak{b} \times \mathfrak{a}$ is trivial. \square

4.3.4. *Dieudonné–theory and bilinear maps.* Let $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+}, [-, -])$ be a Dieudonné–Lie algebra and let $\mathbb{X}(\mathfrak{a}^+)$ be the unique p -divisible group over $\overline{\mathbb{F}}_p$ with (covariant) Dieudonné module $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+})$. We would like to equip its universal cover and its Tate-module with a bilinear map, coming from the Lie bracket on \mathfrak{a}^+ . For this we record a result that tells us how Dieudonné–theory for universal covers of p -divisible groups interacts with \mathbb{Q}_p -bilinear maps. The analogous result for bihomomorphisms of finite flat group schemes is Corollary 1.2.5 of [34]; we’ve decided to include the below proof for the benefit of the reader because it is quite short.

Lemma 4.3.5. *Let Y_1, Y_2, Y_3 be p -divisible groups over $\overline{\mathbb{F}}_p$, then there is a functorial and \mathbb{Q}_p -linear bijection between the space of $\check{\mathbb{Q}}_p$ -bilinear maps*

$$g : \mathbb{D}(Y_1)[\frac{1}{p}] \times \mathbb{D}(Y_2)[\frac{1}{p}] \rightarrow \mathbb{D}(Y_3)[\frac{1}{p}]$$

that satisfy $(g(\varphi_{Y_1}x, \varphi_{Y_2}y) = \varphi_{Y_3}(g(x, y)))$ and the space of bilinear maps

$$f : \tilde{Y}_1 \times \tilde{Y}_2 \rightarrow \tilde{Y}_3.$$

Moreover if $Y_1 = Y_2 = Y_3$ then f satisfies the Jacobi identity if and only if g does.

Proof. Recall that the internal hom p -divisible group \mathcal{H}_{Y_2, Y_3} satisfies

$$\tilde{\mathcal{H}}_{Y_2, Y_3} = \underline{\mathbf{Hom}}(Y_2, Y_3)[\frac{1}{p}] = \underline{\mathbf{Hom}}(\tilde{Y}_2, \tilde{Y}_3).$$

It follows from the usual tensor-hom adjunction for \mathbb{Q}_p -vector spaces that bilinear maps

$$\tilde{Y}_1 \times \tilde{Y}_2 \rightarrow \tilde{Y}_3$$

are in bijection with homomorphisms

$$\tilde{Y}_1 \rightarrow \tilde{\mathcal{H}}_{Y_2, Y_3}.$$

It also follows from the tensor-hom adjunction in the category of F -isocrystals that $\check{\mathbb{Q}}_p$ -bilinear maps

$$\mathbb{D}(Y_1)[\frac{1}{p}] \times \mathbb{D}(Y_2)[\frac{1}{p}] \rightarrow \mathbb{D}(Y_3)[\frac{1}{p}]$$

that ‘commute with the Frobenius’ as above are in bijection with morphisms of F -isocrystals

$$\mathbb{D}(Y_1)[\frac{1}{p}] \rightarrow \underline{\mathbf{Hom}}(\mathbb{D}(Y_2)[\frac{1}{p}], \mathbb{D}(Y_3)[\frac{1}{p}]),$$

where $\underline{\mathbf{Hom}}$ denotes the internal hom in the category of F -isocrystals. Because the slope of $\mathbb{D}(Y_1)[\frac{1}{p}]$ is bounded above by 0, these are also in bijection with morphisms of F -crystals

$$\mathbb{D}(Y_1)[\frac{1}{p}] \rightarrow \underline{\mathbf{Hom}}(\mathbb{D}(Y_2)[\frac{1}{p}], \mathbb{D}(Y_3)[\frac{1}{p}])^{\leq 0},$$

where $()^{\leq 0}$ denotes taking the slope ≤ 0 subspace. Lemma 4.1.7 of [9] tells us that there is an isomorphism

$$\mathbb{D}(\mathcal{H}_{Y_2, Y_3})[\frac{1}{p}] = \underline{\mathbf{Hom}}(\mathbb{D}(Y_2)[\frac{1}{p}], \mathbb{D}(Y_3)[\frac{1}{p}])^{\leq 0}$$

and Dieudonné-theory over $\overline{\mathbb{F}}_p$ tells us that homomorphisms

$$\tilde{Y}_1 \rightarrow \tilde{\mathcal{H}}_{Y_2, Y_3}$$

are in bijection with morphisms of F -isocrystals

$$\mathbb{D}(Y_1)[\frac{1}{p}] \rightarrow \mathbb{D}(\mathcal{H}_{Y_2, Y_3})[\frac{1}{p}],$$

which is what we wanted to prove. We can similarly prove a statement about trilinear maps and use this to deduce that the Jacobi identity, which is just the vanishing of a certain trilinear map, is the same condition on both sides. \square

Remark 4.3.6. The map from bilinear maps of rational Dieudonné-modules to bilinear maps of universal covers of p -divisible groups can be described explicitly on R -points for qrsp R using the isomorphism

$$\tilde{Y}_i(R) = \left(B_{\text{cris}}^+(R) \otimes_{\check{\mathbb{Q}}_p} \mathbb{D}(Y_i)[\frac{1}{p}] \right)^{\varphi=1}$$

of Lemma 2.4.6. Indeed, there is an induced $B_{\text{cris}}^+(R)$ -bilinear map

$$B_{\text{cris}}^+(R) \otimes_{\check{\mathbb{Q}}_p} \mathbb{D}(Y_1)[\frac{1}{p}] \times B_{\text{cris}}^+(R) \otimes_{\check{\mathbb{Q}}_p} \mathbb{D}(Y_2)[\frac{1}{p}] \rightarrow B_{\text{cris}}^+(R) \otimes_{\check{\mathbb{Q}}_p} \mathbb{D}(Y_3)[\frac{1}{p}]$$

which induces a \mathbb{Q}_p -bilinear map on the $\varphi = 1$ subspaces.

Corollary 4.3.7. *Suppose that we are given a $\check{\mathbb{Z}}_p$ -bilinear map*

$$g^+ : \mathbb{D}(Y_1) \times \mathbb{D}(Y_2) \rightarrow \mathbb{D}(Y_3)$$

satisfying $g^+(\varphi_{Y_1}(x), \varphi_{Y_2}(y)) = \varphi_{Y_3}(g^+(x, y))$. The induced map $f : \tilde{Y}_1 \times \tilde{Y}_2 \rightarrow \tilde{Y}_3$ restricts to a \mathbb{Z}_p -bilinear Lie bracket

$$f^+ : T_p Y_1 \times T_p Y_2 \rightarrow T_p Y_3.$$

Proof. It suffices to show this on R -valued points for semiperfect R and in fact by Yoneda it suffices to check it in the universal case that R is the ring underlying $T_p Y_1 \times T_p Y_2$. But this R is qrsp and so we can use Lemma 2.4.6 which tells us that

$$T_p Y_i(R) = \left(A_{\text{cris}}(R) \otimes_{\check{\mathbb{Q}}_p} \mathbb{D}(Y_i) \right)^{\varphi=1},$$

and we see that f^+ sends $T_p Y_1(R) \times T_p Y_2(R)$ to $T_p Y_3(R)$ because g^+ sends $\mathbb{D}(Y_1) \times \mathbb{D}(Y_2)$ to $\mathbb{D}(Y_3)$. \square

4.3.8. Now let \mathfrak{a}^+ be a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra. Thanks to Corollary 4.3.7, the Lie bracket on \mathfrak{a}^+ induces a $\underline{\mathbb{Z}}_p$ -linear Lie bracket

$$T_p \mathbb{X}(\mathfrak{a}^+) \times T_p \mathbb{X}(\mathfrak{a}^+) \rightarrow T_p \mathbb{X}(\mathfrak{a}^+)$$

and after inverting p a $\underline{\mathbb{Q}}_p$ -linear Lie bracket

$$\tilde{\mathbb{X}}(\mathfrak{a}) \times \tilde{\mathbb{X}}(\mathfrak{a}) \rightarrow \tilde{\mathbb{X}}(\mathfrak{a}).$$

The lower central series of \mathfrak{a} induces a filtration

$$\tilde{\mathbb{X}}(\mathfrak{a}) = \tilde{\mathbb{X}}(\mathfrak{a}_0) \supseteq \tilde{\mathbb{X}}(\mathfrak{a}_1) \supseteq \cdots \supseteq \tilde{\mathbb{X}}(\mathfrak{a}_{n(\mathfrak{a})-1})$$

by $\underline{\mathbb{Q}}_p$ -Lie algebra ideals satisfying

$$[\tilde{\mathbb{X}}(\mathfrak{a}), \tilde{\mathbb{X}}(\mathfrak{a}_i)] \subseteq \tilde{\mathbb{X}}(\mathfrak{a}_{i+1}).$$

Moreover the associated graded quotient \mathbb{Q}_p -Lie algebras satisfy (note that $\mathfrak{a} \mapsto \tilde{\mathcal{H}}_{\mathfrak{a}}$ is functorial in morphisms of Dieudonné–Lie \mathbb{Q}_p -algebras)

$$\frac{\tilde{\mathbb{X}}(\mathfrak{a}_i)}{\tilde{\mathbb{X}}(\mathfrak{a}_{i+j})} \simeq \tilde{\mathbb{X}}\left(\frac{\mathfrak{a}_i}{\mathfrak{a}_{i+j}}\right).$$

Its intersection with $T_p\mathbb{X}(\mathfrak{a})$ induces a filtration by \mathbb{Z}_p -Lie algebra ideals

$$T_p\mathbb{X}(\mathfrak{a}) = T_p\mathbb{X}(\mathfrak{a}_0) \supseteq T_p\mathbb{X}(\mathfrak{a}_1) \supseteq \cdots \supseteq T_p\mathbb{X}(\mathfrak{a}_{n(\mathfrak{a})-1}),$$

whose graded quotients are isomorphic to

$$T_p\mathbb{X}\left(\frac{\mathfrak{a}_i^+}{\mathfrak{a}_{i+j}^+}\right).$$

Definition 4.3.9. Let $n(\mathfrak{a})$ be the nilpotency class of the Lie algebra \mathfrak{a} , i.e., the length of its lower central series. We define $\tilde{\Pi}(\mathfrak{a})$ to be the group scheme obtained by endowing $\tilde{\mathbb{X}}(\mathfrak{a})$ with the group structure given by the Baker–Campbell–Hausdorff formula. If $p > n(\mathfrak{a})$, then the Baker–Campbell–Hausdorff formula has coefficients in $\mathbb{Z}_{(p)}$ by Theorem 1 of [36]. Therefore, the group structure on $\tilde{\mathbb{X}}(\mathfrak{a})$ given by the BCH formula restricts to a group structure on $T_p\mathbb{X}(\mathfrak{a}^+)$, and we let $\Pi(\mathfrak{a}^+)$ denote this group scheme.

Hypothesis 4.3.10. From now on we will assume that $p > n(\mathfrak{a})$.

Lemma 4.3.11. *The group scheme $\Pi(\mathfrak{a}^+)$ is quasi-syntomic and semiperfect.*

Proof. It follows from Proposition 4.1.2.(4) of [9] that for any connected p -divisible group $Y/\overline{\mathbb{F}}_p$ the Tate-module $T_p Y$ is representable by a scheme of the form

$$\overline{\mathbb{F}}_p[[x_1^{1/p^\infty}, \dots, x_n^{1/p^\infty}]]/(x_1, \dots, x_n),$$

which is qrsp. Since $\Pi(\mathfrak{a}^+) \simeq T_p\mathbb{X}(\mathfrak{a}^+)$ as schemes, we are done. \square

4.3.12. Let us record the following representability result.

Proposition 4.3.13. *Assume Hypothesis 4.3.10. The quotient of $\tilde{\Pi}(\mathfrak{a})$ by the subgroup $\Pi(\mathfrak{a}^+)$ is representable by a formally smooth formal scheme of dimension equal to $\dim \mathbb{X}(\mathfrak{a}^+)$.*

Proof. What we have proved above implies that

$$\frac{\tilde{\Pi}(\mathfrak{a}_i)/\Pi(\mathfrak{a}_i^+)}{\tilde{\Pi}(\mathfrak{a}_{i+1})/\Pi(\mathfrak{a}_{i+1}^+)} = \frac{\tilde{\mathbb{X}}\left(\frac{\mathfrak{a}_i}{\mathfrak{a}_{i+1}}\right)}{T_p\mathbb{X}\left(\frac{\mathfrak{a}_i^+}{\mathfrak{a}_{i+1}^+}\right)} = \mathbb{X}\left(\frac{\mathfrak{a}_i^+}{\mathfrak{a}_{i+1}^+}\right),$$

which is a p -divisible group and hence representable by a formally smooth formal scheme. Now consider the tower

$$\frac{\tilde{\Pi}(\mathfrak{a})}{\Pi(\mathfrak{a}^+)} \rightarrow \frac{\tilde{\Pi}(\mathfrak{a})/\Pi(\mathfrak{a}^+)}{\tilde{\Pi}(\mathfrak{a}_{n(\mathfrak{a})-1})/\Pi(\mathfrak{a}_{n(\mathfrak{a})-1}^+)} \rightarrow \cdots \rightarrow \frac{\tilde{\Pi}(\mathfrak{a})/\Pi(\mathfrak{a}^+)}{\tilde{\Pi}(\mathfrak{a}_1)/\Pi(\mathfrak{a}_1^+)}.$$

By the discussion in 4.3.8, the rightmost object of the tower is representable by a p -divisible group. The representability of $\frac{\tilde{\Pi}(\mathfrak{a})}{\Pi(\mathfrak{a}^+)}$ then follows by induction because the k -th object of the tower is a torsor under a p -divisible group over the $(k-1)$ -th object, and torsors under p -divisible groups are formally smooth and representable by formal schemes. \square

Definition 4.3.14. We denote by $Z(\mathfrak{a}^+)$ the formally smooth formal scheme $\frac{\tilde{\Pi}(\mathfrak{a})}{\Pi(\mathfrak{a}^+)}$.

4.3.15. Let $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+}, [-, -])$ be a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra with associated Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra $(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$. Let $\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}})$ be the automorphism group of the underlying isocrystal, considered as an algebraic group over \mathbb{Q}_p as in §1.11 of [62]. Then the Lie algebra of $\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}})$ can be identified with the endomorphism ring of the isocrystal, equipped with the commutator bracket.

There is a closed subgroup

$$\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -]) \subseteq \mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}})$$

consisting of those automorphisms preserving the Lie bracket.

Definition 4.3.16. Let Q be an algebraic group over \mathbb{Q}_p equipped with a group homomorphism

$$Q \rightarrow \mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -]).$$

We call such a group homomorphism (or action) *strongly non-trivial* if the induced linear representation of Q on \mathfrak{a} has no non-trivial subquotients.

In the situation of Definition 4.3.16, the elements of $Q(\mathbb{Q}_p)$ act via F -equivariant $\check{\mathbb{Q}}_p$ -linear Lie algebra automorphisms on $(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$ and thus via Lie $\check{\mathbb{Q}}_p$ -algebra automorphisms on $\tilde{\mathbb{X}}(\mathfrak{a})$ by functoriality. There is a compact open subgroup $\Gamma \subseteq Q(\mathbb{Q}_p)$ preserving \mathfrak{a}^+ . By construction, the action of Γ on $\tilde{\mathbb{X}}(\mathfrak{a})$ preserves $T_p X(\mathfrak{a}^+)$, therefore there is an induced action of Γ on the p -divisible group $\mathbb{X}(\mathfrak{a}^+)$. Moreover, there is an action of Γ on the formally smooth formal scheme $Z(\mathfrak{a}^+)$.

Example 4.3.17. If $\mathfrak{a} = \mathbb{D}(\mathcal{H}_Y^{G,\circ})[\frac{1}{p}]$ then the algebraic group J_b has a natural and strongly non-trivial action on \mathfrak{a} . Moreover, the restriction of this action to a maximal torus $T \subseteq J_b$ is still strongly non-trivial, as explained in the proof of Proposition 3.2.1 of [38].

The map $Q \rightarrow \mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}})$ induces a Lie algebra homomorphism

$$\mathrm{Lie} Q \rightarrow \mathrm{Lie} \mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -]) \rightarrow \mathrm{Lie} \mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}) = \mathrm{End}(\mathfrak{a}, \varphi_{\mathfrak{a}})$$

and a routine computation shows that the Lie algebra of $\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])$ consist of all those homomorphism of F -isocrystals $g : \mathfrak{a} \rightarrow \mathfrak{a}$ such that for all $x, y \in \mathfrak{a}$ we have

$$[gx, gy] + [x, gy] + [gx, y] = 0.$$

If U is as above this means that elements of the Lie algebra (in the sense of p -adic Lie groups) of U induce endomorphisms of the Dieudonné module \mathfrak{a}^+ satisfying this relation, and by functoriality these will induce endomorphisms of $\tilde{\mathbb{X}}(\mathfrak{a})$ preserving $T_p \mathbb{X}(\mathfrak{a}^+)$.

Lemma 4.3.18 (Continuity). *The action of the (abstract group) $\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])(\mathbb{Q}_p)$ on $\tilde{\mathbb{X}}(\mathfrak{a})$ can be upgraded uniquely to an action of the locally profinite group scheme*

$$\underline{\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])}(\mathbb{Q}_p).$$

Proof. By the definition of the locally profinite topology on $\mathbf{Aut}(\mathfrak{a}, \varphi_{\mathfrak{a}}, [-, -])(\mathbb{Q}_p)$ it suffices to show that for sufficiently small compact open subgroups U , the action of U on \mathfrak{a} extends to an action of the profinite group scheme \underline{U} . In particular we may assume that U preserves \mathfrak{a}^+ and therefore its action on $\tilde{\mathbb{X}}(\mathfrak{a})$ preserves the Tate module $T_p \mathbb{X}(\mathfrak{a}^+)$ of $\mathbb{X}(\mathfrak{a}^+)$.

Since the action of U on $T_p\mathbb{X}(\mathfrak{a}^+)$ is by group homomorphisms, there are induced actions of U on the quotients

$$T_p\mathbb{X}(\mathfrak{a}^+)/p^n T_p\mathbb{X}(\mathfrak{a}^+) = \mathbb{X}(\mathfrak{a}^+)[p^n]$$

of $T_p\mathbb{X}(\mathfrak{a}^+)$.

Claim 4.3.19. *The action of U on these quotients factors through a finite quotient of U .*

Proof. We can check this after shrinking U and then U is a pro- p group and equal to the exponential of its p -adic Lie algebra. But the action of $\mathrm{Lie} U$ on $T_p\mathbb{X}(\mathfrak{a}^+)/p^n T_p\mathbb{X}(\mathfrak{a}^+)$ factors through $\mathrm{Lie} U/p^n \mathrm{Lie} U$ by linearity and $\mathrm{Lie} U/p^n \mathrm{Lie} U$ is finite. \square

It follows from the claim that the action of U induces a unique action of \underline{U} on $\mathbb{X}(\mathfrak{a}^+)[p^n]$. Since

$$T_p\mathbb{X}(\mathfrak{a}^+) = \varprojlim_n \mathbb{X}(\mathfrak{a}^+)[p^n]$$

it follows that the action of U on $T_p\mathbb{X}(\mathfrak{a}^+)$ upgrades uniquely to an action of \underline{U} . Because the action of U commutes with multiplication by p it follows that the action of U on $\frac{1}{p^n} T_p\mathbb{X}(\mathfrak{a}^+)$ upgrades uniquely to an action of \underline{U} and since

$$\tilde{\mathbb{X}}(\mathfrak{a}) = \varinjlim_n \frac{1}{p^n} T_p\mathbb{X}(\mathfrak{a}^+)$$

we deduce that the action of U on $\tilde{\mathbb{X}}(\mathfrak{a})$ upgrades uniquely to an action of \underline{U} . \square

5. DEFORMATION THEORY OF CENTRAL LEAVES

Let (G, X) be a Shimura datum of Hodge type with reflex field E and let p be a prime such that $G_{\mathbb{Q}_p}$ is quasi-split and split over an unramified extension. Let $U_p \subseteq G(\mathbb{Q}_p)$ be a hyperspecial subgroup and let $U^p \subseteq G(\mathbb{A}_f^p)$ be a sufficiently small compact open subgroup. Let \mathbf{Sh}_G/E be the Shimura variety of level $U = U^p U_p$ and for a prime $v|p$ of E let $\mathcal{S}_G/\mathcal{O}_{E,v}$ be the canonical integral model of \mathbf{Sh}_G constructed in [48] (see [46] for the $p = 2$ case). Let Sh_G be the base change to $\overline{\mathbb{F}}_p$ of this integral canonical model for some choice of map $\mathcal{O}_{E,v} \rightarrow \overline{\mathbb{F}}_p$.

A choice of Hodge embedding will give us a closed immersion $\mathrm{Sh}_G \rightarrow \mathcal{A}_{g,N}$ for some g and N (this uses the main result of [71]) and therefore a family of abelian varieties A/Sh_G with associated p -divisible group $X = A[p^\infty]$. In this section we will recall the definitions of central leaves C and Igusa varieties from [32, 33, 45]. The main goal of the section is to study the structure of the formal completions $C^{/x}$ of central leaves at point $x \in C(\overline{\mathbb{F}}_p)$. Using the Hodge embedding, these formal schemes admit closed immersions to deformation spaces $\mathbf{Def}(Y)$ of p -divisible groups Y over $\overline{\mathbb{F}}_p$ and in fact they admit closed immersions to central leaves $\mathbf{Def}_{\mathrm{sus}}(Y) \subseteq \mathbf{Def}(Y)$ inside these deformation spaces⁴. If the corresponding p -divisible group is completely slope divisible, then $\mathbf{Def}_{\mathrm{sus}}(Y)$ has the structure of a p -divisible *cascade* in the sense of [58]. For example if Y has two slopes then $\mathbf{Def}_{\mathrm{sus}}(Y)$ has the structure of a p -divisible formal group. As observed already in [65] (c.f. [40]), one cannot expect that $C^{/x} \subseteq \mathbf{Def}_{\mathrm{sus}}(Y)$ is a sub-cascade but only that it is a *shifted subcascade*. We will not make use of p -divisible cascades or shifted subcascades in this paper and instead show, using the work of Caraiani–Scholze and Kim, that the perfection of the formal schemes $C^{/x}$ and $\mathbf{Def}_{\mathrm{sus}}(Y)$ canonically have a (non-commutative) group structure. These groups will be of the form $\tilde{\Pi}(\mathfrak{a})$ for a Dieudonné–Lie $\check{\mathbb{Q}}_p$ -algebra as introduced in Section 4. As a

⁴The notation $\mathbf{Def}_{\mathrm{sus}}$ denotes the *sustained deformation space* in the sense of Chai–Oort, this will be explained in Section 5.3.

consequence C/x and $\mathbf{Def}_{\text{sus}}(Y)$ have the structure of an *iterated BT-fibration*. In other words, it is an iterated fibration of p -divisible formal groups.

Let $\text{Sh}_{G,b}$ be the Newton stratum containing the central leaf C and let ν_b be the Newton cocharacter of b . As before we consider the parabolic subgroup P_{ν_b} of G with unipotent radical U_{ν_b} . Then we will show that \mathfrak{a} introduced above has underlying Lie algebra isomorphic to $\text{Lie } U_{\nu_b}$.

5.1. Central leaves. Let $A/\mathcal{A}_{g,N}$ denote the universal abelian variety and let λ be the universal polarisation. The moduli space $\mathcal{A}_{g,N}$ is equipped with a Newton stratification; the Newton stratum that a point $x \in \mathcal{A}_{g,N}(\overline{\mathbb{F}}_p)$ lies in is determined by the isogeny class of the (quasi-polarised) p -divisible group $(A_x[p^\infty], \lambda_x)$. Central leaves are defined similarly using instead the isomorphism class of $(A_x[p^\infty], \lambda_x)$. To be more precise, given a point $x \in \mathcal{A}_{g,N}(\overline{\mathbb{F}}_p)$, the central leaf C_λ passing through x is defined as

$$C_\lambda = \{y \in \mathcal{A}_{g,N}(\overline{\mathbb{F}}_p) \mid (A_y[p^\infty], \lambda_y) \simeq_\lambda (A_x[p^\infty], \lambda_x)\},$$

where the notation \simeq_λ means that there is an isomorphism preserving the polarisation up to a scalar in \mathbb{Z}_p^\times . It follows from results of Oort, [60], that C_λ is equal to the set of $\overline{\mathbb{F}}_p$ -point of a smooth equidimensional variety that is closed inside the Newton stratum containing x , and thus locally closed inside $\mathcal{A}_{g,N}$.

Hypothesis 5.1.1. From now on we will write $Y = A_x[p^\infty]$ and we will assume that Y is completely slope divisible. This implies that $X = A[p^\infty]$ over C_λ admits a slope filtration and we will denote the associated graded pieces for the slope filtration by X_i .

We now define Igusa towers (M for Mantovan and CS for Caraiani–Scholze)

$$\text{Ig}_{\text{CS},\lambda} \rightarrow \text{Ig}_{\text{M},\lambda} \rightarrow C_\lambda,$$

where $\text{Ig}_{\text{M},\lambda}$ to C_λ is the moduli space of isomorphisms (compatible with the polarisations up to a \mathbb{Z}_p^\times -scalar)

$$X_i \simeq Y_{i,C_\lambda}$$

and $\text{Ig}_{\text{CS},\lambda} \rightarrow C_\lambda$ is the moduli space of isomorphisms (compatible with the polarisations up to a \mathbb{Z}_p^\times -scalar)

$$X \simeq Y_{C_\lambda}.$$

Let $\mathbf{Aut}_\lambda(Y)$ be the group scheme of automorphisms of Y that preserve the polarisation λ up to a \mathbb{Z}_p^\times -scalar (this is $\mathbf{Aut}_G(Y)$ for $G = \text{GSp}$) and let $\underline{\mathbf{Aut}}_\lambda(Y)(\overline{\mathbb{F}}_p)$ be its pro-étale group scheme of connected components.

It follows from work of Mantovan, [52], that C_λ is smooth and that $\text{Ig}_{\text{M},\lambda} \rightarrow C_\lambda$ is a pro-étale torsor for $\underline{\mathbf{Aut}}_\lambda(Y)(\overline{\mathbb{F}}_p)$. Caraiani and Scholze, [9], prove that $\text{Ig}_{\text{CS},\lambda}$ is perfect and that the map $\text{Ig}_{\text{CS},\lambda} \rightarrow \text{Ig}_{\text{M},\lambda}$ identifies it with the perfection of $\text{Ig}_{\text{M},\lambda}$. This implies that $\text{Ig}_{\text{CS},\lambda} \rightarrow C_\lambda$ is faithfully flat and thus a torsor for $\mathbf{Aut}_\lambda(Y)$. Moreover they prove that the action of $\mathbf{Aut}_\lambda(Y)$ on $\text{Ig}_{\text{CS},\lambda}$ extends to an action of $\mathbf{Aut}_\lambda(\tilde{Y})$, the group of automorphisms of \tilde{Y} preserving the polarisation up to a \mathbb{Q}_p^\times -scalar.

We note that the following diagram is Cartesian

$$(5.1.1) \quad \begin{array}{ccc} \text{Ig}_{\text{CS},\lambda} & \longrightarrow & \text{Ig}_{\text{M},\lambda} \\ \downarrow & & \downarrow \\ C_\lambda^{\text{perf}} & \longrightarrow & C_\lambda. \end{array}$$

5.2. Central leaves for Shimura varieties of Hodge type. Let $\mathfrak{a}^+ = \mathbb{D}(\mathcal{H}_Y^{G,\circ})$ and $\mathfrak{a} = \mathfrak{a}^+ \otimes \check{\mathbb{Q}}_p$ as above. We assume that $p > n(\mathfrak{a})$, where $n(\mathfrak{a})$ is the nilpotency class of the Lie algebra \mathfrak{a} . We have seen in Section 4.2 that $\mathfrak{a} = \text{Lie } U_{\nu_b}$, so that this condition only depends on the Newton stratum that we are in. We have that $\mathbf{Aut}_G(\tilde{Y})^\circ \simeq \tilde{\Pi}(\mathfrak{a})$ and $\mathbf{Aut}_G(Y)^\circ \simeq \Pi(\mathfrak{a}^+)$.

By the main result of [71], a choice of a Hodge embedding gives us a closed immersion $\text{Sh}_G \hookrightarrow \mathcal{A}_{g,N}$. In this setting, Hamacher and Kim defined central leaves

$$C \subseteq (\text{Sh}_G \cap C_\lambda)$$

which are, roughly speaking, the locally closed subsets where the p -divisible group with G -structure is constant. These are locally closed smooth subschemes of $\mathcal{A}_{g,N}$ of dimension $\langle 2\rho, \nu_b \rangle$, by Corollary 5.3.1 of [45].

Remark 5.2.1. The reader might worry that $\text{Sh}_G \cap C_\lambda$ might be always empty because we have assumed that the universal p -divisible group over C_λ is completely slope divisible. Fortunately, Proposition 2.4.5 of [45] tells us that for every Newton stratum $\text{Sh}_{G,b} \subseteq \text{Sh}_G$ we can always find a central leaf $C \subseteq \text{Sh}_{G,b}$ such that the universal p -divisible group over it (coming from our choice of a Hodge embedding) is completely slope divisible.

Hamacher and Kim in [33] define an Igusa variety $\text{Ig}_{\text{CS}} \rightarrow C^{\text{perf}}$ by pulling back $\text{Ig}_{\text{CS},\lambda} \rightarrow C_\lambda^{\text{perf}}$ and then taking the closed subset where the universal isomorphism

$$X \simeq Y_{C_\lambda}$$

preserves the tensors on geometric points. They then prove that $\text{Ig}_{\text{CS}} \rightarrow C^{\text{perf}}$ is a pro-étale torsor for $\mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p) \subseteq \mathbf{Aut}_\lambda(Y)(\overline{\mathbb{F}}_p)$ and that Ig_{CS} is stable under the action of $\mathbf{Aut}_G(\tilde{Y}) \subseteq \mathbf{Aut}(\tilde{Y})$. Using the invariance of the étale site under perfection, we get a Cartesian diagram

$$\begin{array}{ccc} \text{Ig}_{\text{CS}} & \longrightarrow & \text{Ig}_{\text{M}} \\ \downarrow & & \downarrow \\ C^{\text{perf}} & \longrightarrow & C \end{array}$$

which fits in a commutative cube with (5.1.1).

The map $\text{Ig}_{\text{CS}} \rightarrow C$ is faithfully flat since $\text{Ig}_{\text{CS}} \rightarrow C^{\text{perf}}$ is a pro-étale cover and C is smooth. We want to prove that it is actually an fpqc torsor under $\mathbf{Aut}_G(Y)$. We first prove it at the level of infinitesimal neighbourhoods (defined as in Section 5.2 of [33]).

Lemma 5.2.2. *If $y \in \text{Ig}_{\text{CS}}(\overline{\mathbb{F}}_p)$ is an $\overline{\mathbb{F}}_p$ -point over $x \in C(\overline{\mathbb{F}}_p)$, then $\text{Ig}_{\text{CS}}^{/y} \rightarrow C^{/x}$ is a torsor under the formal group $\mathbf{Aut}_G^\circ(Y)$.*

Proof. In Section 5.2 of [33], Hamacher and Kim show that the formal completion at a point of Ig_{CS} is isomorphic to $\mathbf{Aut}_G^\circ(\tilde{Y})$ and that the natural action of $\mathbf{Aut}_G^\circ(\tilde{Y})$ on Ig_{CS} corresponds to the multiplication map under this isomorphism. Moreover, the morphism $\text{Ig}_{\text{CS}}^{/y} \rightarrow C^{/x}$ corresponds to the restriction of the action map

$$\mathbf{Aut}_G^\circ(\tilde{Y}) \times \text{Sh}_{G,b}^{/x} \rightarrow \text{Sh}_{G,b}^{/x}$$

to the closed point $x \in \text{Sh}_{G,b}^{/x}$. Theorem 5.1.3 of [45] tells us that the scheme-theoretic image of this restriction

$$\mathbf{Aut}_G^\circ(\tilde{Y}) \rightarrow \text{Sh}_{G,b}^{/x}$$

is C'/x and in fact this identifies $\mathbf{Aut}_G^\circ(\tilde{Y})$ with the formal completion of the perfection of C'/x . If we apply this when $G = \mathrm{GSp}$, the fact that

$$\mathrm{Ig}_{\mathrm{CS},\lambda} \rightarrow \mathrm{Ig}_{\mathrm{M},\lambda}$$

is a torsor for $\mathbf{Aut}_\lambda^\circ(Y)$ (plus the fact that $\mathrm{Ig}_{\mathrm{M},\lambda} \rightarrow C_\lambda$ is pro-étale) tells us that

$$C_\lambda'/x \simeq \frac{\mathbf{Aut}_\lambda^\circ(\tilde{Y})}{\mathbf{Aut}_\lambda^\circ(Y)}.$$

Since $\mathbf{Aut}_G^\circ(Y)$ is the intersection of $\mathbf{Aut}_\lambda^\circ(Y)$ with $\mathbf{Aut}_G(\tilde{Y})$, it follows similarly that $\mathbf{Aut}_G^\circ(Y) \subseteq \mathbf{Aut}_G^\circ(\tilde{Y})$ is the stabiliser of $x \in C'/x$. Therefore, we get a monomorphism of formal schemes

$$\frac{\mathbf{Aut}_G^\circ(\tilde{Y})}{\mathbf{Aut}_G^\circ(Y)} \rightarrow C'/x,$$

both of which are formally smooth of dimension $\langle 2\rho, \nu_b \rangle$. Indeed for the left hand side this is Proposition 4.3.13, while for the right hand side it is Corollary 5.3.1 of [45]. Lemma 2.1.7 tells us that the map is a closed immersion and therefore we deduce that is an isomorphism. \square

Proposition 5.2.3. *The map $\mathrm{Ig}_{\mathrm{CS}} \rightarrow C$ is an fpqc torsor under $\mathbf{Aut}_G(Y)$.*

Proof. Since $\mathrm{Ig}_{\mathrm{CS}} \rightarrow C$ is faithfully flat, it suffices to prove that it is a quasi-torsor under $\mathbf{Aut}_G(Y)$. In other words, we want to show that the action map

$$(5.2.1) \quad \mathbf{Aut}_G(Y) \times \mathrm{Ig}_{\mathrm{CS}} \rightarrow \mathrm{Ig}_{\mathrm{CS}} \times_C \mathrm{Ig}_{\mathrm{CS}}$$

is an isomorphism. This map is clearly a homeomorphism because $\mathrm{Ig}_{\mathrm{M}} \rightarrow C$ is an $\mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p)$ -torsor and both $\mathrm{Ig}_{\mathrm{CS}} \rightarrow \mathrm{Ig}_{\mathrm{M}}$ and $\mathbf{Aut}_G(\tilde{Y}) \rightarrow \mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p)$ are universal homeomorphisms.

It follows from Lemma 5.2.3 of [33] that when Z is either the source or the target of (5.2.1) and $z \in Z(\overline{\mathbb{F}}_p)$ is an $\overline{\mathbb{F}}_p$ -point over x , then Z/z is pro-represented by the formal spectrum of the I -adic completion of $\mathcal{O}_{Z,x}$, where I is the maximal ideal of $\mathcal{O}_{C,x}$. Moreover, they prove that $\mathcal{O}_{Z,x} \rightarrow \mathcal{O}_{Z,x}^{\wedge,I}$ is faithfully flat.

It follows from Lemma 5.2.2 that the action map (5.2.1) is flat at all closed points and therefore it is flat by (7) of Lemma 00HT of [68]. We moreover know that the action map is a closed immersion because this is true in the Siegel case, so the action map is a surjective flat closed immersion and therefore an isomorphism (see Lemma 04PW of [68]). \square

5.3. Deformation theory of central leaves. If we let the notation be as above then Chai and Oort prove that the deformation theory of C_λ is completely determined by the $\mathbf{Aut}_\lambda(Y)$ -torsor $\mathrm{Ig}_{\mathrm{CS},\lambda} \rightarrow C_\lambda$. Let $\mathbf{Def}_{\mathrm{sus}}(Y, \lambda)$ denote the deformation space (considered as a functor on the category of Artin local rings over $\overline{\mathbb{F}}_p$ with residue field $\overline{\mathbb{F}}_p$) of the trivial $\mathbf{Aut}_\lambda(Y)$ -torsor over $\overline{\mathbb{F}}_p$. Then Theorem 4.3 of [16] states that the natural map

$$C_\lambda'/x \rightarrow \mathbf{Def}_{\mathrm{sus}}(Y, \lambda)$$

is an isomorphism of formal schemes. They prove this by showing that $\mathbf{Def}_{\mathrm{sus}}(Y, \lambda)$ is formally smooth of the same dimension as C_λ'/x . We will need a version of their argument later, so we'll give an axiomatic treatment of it. Note that the deformation space of the trivial $\mathbf{Aut}_\lambda(Y)$ -torsor over $\overline{\mathbb{F}}_p$ is (non-canonically) isomorphic to the deformation space of the trivial $\mathbf{Aut}_\lambda^\circ(Y)$ -torsor over $\overline{\mathbb{F}}_p$ because torsors for pro-étale group schemes have trivial deformation theory.

Proposition 5.3.1. *Let $P/\overline{\mathbb{F}}_p$ be a group scheme that is an inverse limit of finite flat group schemes $\{P_i\}_{i \in \mathbb{Z}_{\geq 1}}$, equipped with a filtration Fil_P^\bullet . Assume that for each k there is a p -divisible group \mathbb{X}_k such that*

$$\frac{\text{Fil}_P^{k+1}}{\text{Fil}_P^k} \simeq T_p \mathbb{X}_k,$$

Let $\text{Fil}_{P_i}^\bullet$ be the induced filtration of P_i , and assume that there is an isomorphism of inverse system

$$\left\{ \frac{\text{Fil}_{P_i}^{k+1}}{\text{Fil}_{P_i}^k} \right\}_{i \in \mathbb{Z}_{\geq 1}} \simeq \{\mathbb{X}_k[p^i]\}_{i \in \mathbb{Z}_{\geq 1}}.$$

Then the formal deformation space of the trivial P -torsor is formally smooth of dimension

$$\sum_k \dim \mathbb{X}_k.$$

Proof. Let $R' \rightarrow R$ be a square-zero extension of Artinian local rings with over $\overline{\mathbb{F}}_p$ with kernel I , and write $S = \text{Spec } R, S' = \text{Spec } R'$. Let T/S be a P -torsor together with a trivialisation $\alpha : T \times_{S_0} \text{Spec } \overline{\mathbb{F}}_p \simeq P$. We need to show that there is a P -torsor T' over S' lifting T . Since T is the inverse limit of the P_i -torsors T_i induced from T via $P \rightarrow P_i$, it suffices to show that the T_i lift compatibly to S' .

For this, we will use the deformation theory for torsors under flat group schemes locally of finite presentation from Section 2.4 of Chapter VII of [41]. It follows from loc. cit. that there is a complex $\ell_{T_n/S}$ of \mathcal{O}_S -modules of amplitude contained in $[-1, 0]$, called the co-Lie complex. Theorem 2.4.4 of loc. cit. tells us that there is an obstruction element

$$o(T_i, R' \rightarrow R) \in H^2(\text{Spec } \overline{\mathbb{F}}_p, \ell_{T_i/S}^\vee \otimes_{\overline{\mathbb{F}}_p} I),$$

which vanishes precisely when T_n lifts to S' . Moreover the set of all such lifts is a torsor under

$$H^1(\text{Spec } \overline{\mathbb{F}}_p, \ell_{P_i/\overline{\mathbb{F}}_p}^\vee \otimes_{\overline{\mathbb{F}}_p} I) =: \nu_{P_i}$$

The obstruction element vanishes for cohomological degree reasons, and so the torsor T_i lifts to S' . In order to show that our deformation space is formally smooth, we now need to show that

$$H^1(\text{Spec } \overline{\mathbb{F}}_p, \ell_{P_{i+1}/\overline{\mathbb{F}}_p}^\vee \otimes_{\overline{\mathbb{F}}_p} I) \rightarrow H^1(\text{Spec } \overline{\mathbb{F}}_p, \ell_{P_i/\overline{\mathbb{F}}_p}^\vee \otimes_{\overline{\mathbb{F}}_p} I)$$

is surjective for all i , so that we can pick a compatible sequence of lifts of the T_i to S' . In fact we will show that this is an isomorphism for all i , so that we can compute the dimension in the case of $i = 1$.

To show that this cohomology group does not depend on i , we will show that ν_{P_i} does not depend on i . Proposition 3.3.1 of Chapter VII of [41] tells us that a short exact sequence of flat locally of finite presentation group schemes leads to a distinguished triangle between their co-Lie complexes. Using the filtration $\text{Fil}_{P_i}^k$, we see that it suffices to show that $\nu_{\mathbb{X}_k[p^i]}$ does not depend on i , but this is Proposition 2.2.1.c of [42].

To compute the dimension of our deformation functor, we note that Proposition 2.2.1.c of [42] also tells us that the dimension of

$$H^1(\text{Spec } \overline{\mathbb{F}}_p, \ell_{\mathbb{X}[p]}^\vee)$$

is equal to the dimension of the tangent space of X , and therefore equal to the dimension of X ; the proposition follows. \square

5.3.2. There is a canonical $\mathbf{Aut}_\lambda(Y)$ -torsor over $C_\lambda^{/x}$ given by the base-change of $\mathrm{Ig}_{\mathrm{CS},\lambda}$.

Lemma 5.3.3. *The induced map*

$$C_\lambda^{/x} \rightarrow \mathbf{Def}_{\mathrm{sus}}(Y, \lambda).$$

is an isomorphism.

Proof. It suffices to show that it is a closed immersion because both sides are formally smooth of the same dimension. Indeed, the dimension of the right hand side can be computed using Proposition 5.3.1 and this agrees with the dimension of the left hand side. There is a natural closed immersion

$$C_\lambda^{/x} \rightarrow \mathbf{Def}(Y)$$

to the deformation space of Y , given by Serre–Tate theory. There is a subfunctor of the deformation space of Y , given by those deformations that are fpqc locally isomorphic to Y . This subfunctor can be identified with $\mathbf{Def}_{\mathrm{sus}}(Y, \lambda)$ because the universal $\mathbf{Aut}_\lambda(Y)$ -torsor determines the universal p -divisible group. Moreover the subfunctor is automatically closed by Lemma 5.3.4. \square

In the Hodge type case we define $\mathbf{Def}_{\mathrm{sus}}(Y, G)$ to be the deformation space of the trivial $\mathbf{Aut}_G(Y)$ -torsor, then there is a commutative diagram

$$(5.3.1) \quad \begin{array}{ccc} C^{/x} & \longrightarrow & \mathbf{Def}_{\mathrm{sus}}(Y, G) \\ \downarrow & & \downarrow i \\ C_\lambda^{/x} & \hookrightarrow & \mathbf{Def}_{\mathrm{sus}}(Y, \lambda). \end{array}$$

Lemma 5.3.4. *The map $C^{/x} \rightarrow \mathbf{Def}_{\mathrm{sus}}(Y, G)$ is an isomorphism.*

Proof. We start by proving that i is a closed immersion. For this, we note that by Lemma 2.1.7 it suffices to prove that it is a monomorphism. This means that given an Artin local ring R together with a $\mathbf{Aut}_\lambda(Y)$ -torsor \mathcal{P} over R , the set of reductions of structure group of P to $\mathbf{Aut}_G(Y)$, compatible with the identification

$$P \otimes_R \overline{\mathbb{F}}_p = \mathbf{Aut}_\lambda(Y),$$

is either empty or consists of a single element. This set is either empty or it has a transitive action of $\mathbf{Aut}_\lambda(Y)^\circ(R)$ which is the trivial group because

$$\mathbf{Aut}_\lambda^\circ(Y) \simeq \mathrm{Spec} \overline{\mathbb{F}}_p[X_1^{1/p^\infty}, \dots, X_n^{1/p^\infty}]/(X_1, \dots, X_n)$$

has no non-trivial sections over Noetherian test objects.

Proposition 5.3.1 together with Lemma 4.2.1 tells us that $\mathbf{Def}_{\mathrm{sus}}(Y, G)$ is formally smooth of dimension $\langle 2\rho, \nu_b \rangle$, which is equal to the dimension of C by Corollary 5.3.1 of [45].

It follows from the diagram (5.3.1) that the map $C^{/x} \rightarrow \mathbf{Def}_{\mathrm{sus}}(Y, G)$ is a closed immersion between two formally smooth formal schemes of the same dimension, and therefore it is an isomorphism. \square

Remark 5.3.5. If $Y/\overline{\mathbb{F}}_p$ is a completely slope divisible p -divisible group, then one can prove that the deformation space $\mathbf{Def}_{\mathrm{sus}}(Y)$ of the trivial $\mathbf{Aut}(Y)^\circ$ -torsor admits a closed immersion to the deformation space $\mathbf{Def}(Y)$ of Y . Its image is identified with the subspace of deformations of Y

that are fpqc locally isomorphic to the constant deformation of Y . It follows from Section 5 of [45] that there is an action of $\mathbf{Aut}(\tilde{Y})^\circ$ on $\mathbf{Def}_{\text{sus}}(Y)$ which gives an identification

$$\mathbf{Def}_{\text{sus}}(Y) \simeq \frac{\mathbf{Aut}(\tilde{Y})^\circ}{\mathbf{Aut}(Y)^\circ}.$$

Remark 5.3.6. If $Y = Y_1 \oplus Y_2$ then $\mathbf{Aut}(\tilde{Y})^\circ$ is isomorphic to $\tilde{\mathcal{H}}_{Y_1, Y_2}$ and $\mathbf{Aut}(Y)^\circ \simeq T_p \mathcal{H}_{Y_1, Y_2}$ so that $\mathbf{Def}_{\text{sus}}(Y) \simeq \mathcal{H}_{Y_1, Y_2}$. This gives $\mathbf{Def}_{\text{sus}}(Y)$ the structure of a p -divisible formal group. There is another group structure on $\mathbf{Def}_{\text{sus}}(Y)$ because it is the deformation space of the trivial $\mathbf{Aut}(Y)^\circ$ -torsor and $\mathbf{Aut}(Y)^\circ \simeq T_p \mathcal{H}_{Y_1, Y_2}$ is abelian. This second group structure coincides with the group structure we get from thinking of $\mathbf{Def}_{\text{sus}}(Y)$ as parametrising certain extensions of Y_1 by Y_2 .

It is not obvious to us that the first group structure agrees with the second group structure and it is not something that we will need in the rest of the paper. However we would like to point out that there is a preprint of Chai, [12], where he computes the Dieudonné module of the second group structure and it seems to be the same as the Dieudonné-module of the first group structure.

5.4. Strongly Tate-linear subspaces. Let $Y/\overline{\mathbb{F}}_p$ be a complete slope divisible p -divisible group and let X be the universal p -divisible group over the sustained deformation space $\mathbf{Def}_{\text{sus}}(Y)$. Let $Z \subseteq \mathbf{Def}_{\text{sus}}(Y)$ be a formally smooth closed subscheme, then the monodromy group $G(\mathcal{M}_Z)$ of the isocrystal $\mathcal{M} = \mathbb{D}(X)[\frac{1}{p}]$ restricted to Z with respect to the closed point of Z has a natural inclusion

$$\text{Lie } G(\mathcal{M}_Z) \subseteq \mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}] \subseteq \mathbb{D}(\mathcal{H}_Y)[\frac{1}{p}] = \text{Lie } \text{GL}(\mathbb{D}(Y))[\frac{1}{p}].$$

because X admits a slope filtration with constant graded parts. Since \mathcal{M} has the structure of an F -isocrystal we get, by Section 2.2 of [21], an isomorphism

$$F^* G(\mathcal{M}_Z) \rightarrow G(\mathcal{M}_Z),$$

which induces an isomorphism

$$F^* \text{Lie } G(\mathcal{M}_Z) \rightarrow \text{Lie } G(\mathcal{M}_Z)$$

compatible with the F -structure on $\mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}]$. In particular

$$\mathfrak{b} := \text{Lie } G(\mathcal{M}_Z) \subseteq \mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}] =: \mathfrak{a}_Y$$

is a sub- F -isocrystal stable under the Lie bracket. Note that

$$\mathfrak{a}_Y^+ =: \mathbb{D}(\mathcal{H}_Y^\circ)$$

is a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra, thus we get a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra

$$\mathfrak{b}^+ = \mathfrak{a}_Y^+ \cap \mathfrak{b}.$$

5.4.1. Let $\mathfrak{b} \subseteq \mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}]$ be an F -stable Lie subalgebra as above. There is a closed immersion of Lie \mathbb{Q}_p -algebras

$$\tilde{\mathbb{X}}(\mathfrak{b}) \hookrightarrow \tilde{\mathbb{X}}(\mathfrak{a}_Y) = \tilde{\mathcal{H}}_Y^\circ.$$

Using the BCH-formula we get a closed group homomorphism

$$\tilde{\Pi}(\mathfrak{b}) \hookrightarrow \tilde{\Pi}(\mathfrak{a}_Y) = \mathbf{Aut}(\tilde{Y})^\circ$$

and similarly if $p > n$ we have

$$\Pi(\mathfrak{b}^+) \subseteq \Pi(\mathfrak{a}_Y^+) = \mathbf{Aut}(Y)^\circ.$$

This means that we get a formally smooth closed formal subscheme

$$Z(\mathfrak{b}^+) = \frac{\tilde{\Pi}(\mathfrak{b})}{\Pi(\mathfrak{b}^+)} \hookrightarrow \mathbf{Def}_{\text{sus}}(Y)$$

and the same proof as in the previous section shows that $Z(\mathfrak{b}^+)$ can be identified with the deformation space of the trivial $\Pi(\mathfrak{b}^+)$ -torsor. To be precise Lemma 5.3.4 tells us that the natural map to the deformation space of the trivial $\Pi(\mathfrak{b}^+)$ is a closed immersion, and then it is an isomorphism because source and target are formally smooth of the same dimension.

Therefore the subspace $Z(\mathfrak{b}^+)$ is a strongly Tate-linear subspace of $\mathbf{Def}_{\text{sus}}(Y)$ in the sense of Chai–Oort, see Definition 6.2 of [16]. This requires a bit of translation since they work with projective systems of torsors for a projective system of finite flat group schemes and we are instead working with torsors for the inverse limit of the projective system of finite flat group schemes.

Conjecture 5.4.2. *Let $Z \hookrightarrow \mathbf{Def}_{\text{sus}}(Y)$ be a closed immersion. Then for each F -stable Lie subalgebra $\mathfrak{b} \subseteq \mathfrak{a}_Y$ there is an inclusion $Z \subseteq Z(\mathfrak{b}^+)$ if and only if $\text{Lie } G(\mathcal{M}_Z) \subseteq \mathfrak{b}$. In particular $\text{Lie } G(\mathcal{M}_{Z(\mathfrak{b}^+)}) = \mathfrak{b}$.*

Remark 5.4.3. If $\mathfrak{b} = \mathbb{D}(\mathcal{H}_Y^{G,\circ})$ for a Shimura variety of Hodge type (G, X) with G^{ad} simple over \mathbb{Q} , then the conjecture is true. Indeed the global monodromy of C is equal to P_{ν_b} by Corollary 2.2.1 of [38], which uses [22, 23], and then Theorem 3.4.4 allows us to conclude.

As a special case of this if Y has height h and dimension d , then $\mathbf{Def}_{\text{sus}}(Y)$ can be realised as the complete local ring of a central leaf in a PEL type unitary Shimura variety of signature $(h - d, d)$ associated to an imaginary quadratic field E in which p splits. In particular we know that the monodromy group of $\mathcal{M}/\mathbf{Def}_{\text{sus}}(Y)$ is isomorphic to the unipotent group corresponding to the nilpotent Lie algebra $\mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}] = \mathfrak{a}_Y$.

5.4.4. There is a $\check{\mathbb{Z}}_p$ -algebra structure on $\mathbb{D}(\mathcal{H}_Y)$ and $1 + \mathbb{D}(\mathcal{H}_Y^\circ)$ is a subgroup, defining a unipotent algebraic group over $\check{\mathbb{Z}}_p$. We can also consider $\mathbb{D}(\mathcal{H}_Y^\circ) \subseteq \mathbb{D}(\mathcal{H}_Y)$ as a Lie subalgebra, and consider the associated unipotent algebraic group. If Y is completely slope divisible and $p > n \left(\mathbb{D}(\mathcal{H}_Y^\circ)[\frac{1}{p}] \right)$, or equivalently if p is greater than the number of slopes of Y minus one, then these two constructions are the same.

Similarly given $\mathfrak{b} \subseteq \mathfrak{a}_Y$ we get a nilpotent $\check{\mathbb{Z}}_p$ -Lie algebra $\mathfrak{b}^+ = \mathfrak{a}_Y \cap \mathfrak{b}$ whose exponential defines a unipotent algebraic group $U(\mathfrak{b}^+)$ over $\check{\mathbb{Z}}_p$. Its generic fibre is a unipotent group $U(\mathfrak{b})$ which only depends on \mathfrak{b} and comes with a closed immersion

$$U(\mathfrak{b}) \subseteq 1 + \mathbb{D}(\mathcal{H}_Y^\circ).$$

5.5. Local monodromy of strongly Tate-linear subvarieties.

5.5.1. *Algebraisation.* Let $Z = Z(\mathfrak{b}) \subseteq \mathbf{Def}_{\text{sus}}(Y)$ be a strongly Tate-linear formal closed subscheme of $\mathbf{Def}_{\text{sus}}(Y)$ corresponding to a Dieudonné–Lie subalgebra $\mathfrak{b} \subseteq \mathfrak{a}_Y$. The formal scheme Z is equal to $\text{Spf } A$ for a complete Noetherian local ring A . Proposition 2.4.8 of [24] tells us that the category of Dieudonné isocrystals over $\text{Spf } A$ is equivalent to the category of Dieudonné isocrystals over $\text{Spec } A$. Moreover, we have that the $\Pi(\mathfrak{b}^+)$ -torsor P_Z over $Z = \text{Spf } A$ comes from a $\Pi(\mathfrak{b}^+)$ -torsor P_Z^{alg} over $\text{Spec } A$. Indeed P is an inverse limit of torsors under finite flat group schemes over Z , and those all algebraise because finite modules do functorially.

Notation 5.5.2. In Section 5.5 we will treat Z as the affine scheme $\text{Spec } A$ rather than the formal scheme $\text{Spf}(A)$ and we will simply write $P_Z \rightarrow Z$ for the algebraic torsor P_Z^{alg} defined above.

Let \mathcal{M}^+ be the F -crystal over $\mathbf{Def}_{\text{sus}}(Y)$ attached to the universal p -divisible group. We write \mathcal{M} for the induced F -isocrystal and \mathcal{M}_Z the restriction of \mathcal{M} to Z . Write $i : z \hookrightarrow Z$ for the closed embedding of the closed point of Z .

Theorem 5.5.3. *The monodromy group $G(\mathcal{M}_Z)$ of the restriction of \mathcal{M} to Z is contained in $U(\mathfrak{b})$.*

Proof. Let $P \rightarrow Z$ be the restriction of the universal $\mathbf{Aut}(Y)$ -torsor and let $P_Z \rightarrow Z$ be the reduction of P to a $\Pi(\mathfrak{b}^+)$ -torsor. The basic idea of the proof is to use descent of isocrystals along $P_Z \rightarrow Z$ to describe \mathcal{M}_{P_Z} as a constant isocrystal equipped with a descent datum (or equivalently an $\Pi(\mathfrak{b}^+)$ -equivariant structure). However it seems quite hard to compare the group scheme $\Pi(\mathfrak{b}^+)$ over $\overline{\mathbb{F}}_p$ with the monodromy group of \mathcal{M} , which is an algebraic group over $\check{\mathbb{Q}}_p$. This is where the Cartier–Witt stacks of [4] come in.

5.5.4. *Cartier–Witt stacks.* The Dieudonné module $\mathbb{D}(Y)$ of Y is a trivial vector bundle on $\overline{\mathbb{F}}_p^\Delta = \text{Spf}(\check{\mathbb{Z}}_p)$ endowed with a Frobenius. We denote by $\text{GL}(\mathbb{D}(Y))$ the p -adic formal group over $\text{Spf}(\check{\mathbb{Z}}_p)$ of $\check{\mathbb{Z}}_p$ -linear automorphisms of $\mathbb{D}(Y)$ (thus forgetting the F -structure). Let $U(\mathfrak{b}^+) \subset \text{GL}(\mathbb{D}(Y))$ denote the inclusion of the p -adic completion of the unipotent group $U(\mathfrak{b}^+)$.

By the formalism of Cartier–Witt stacks, the crystal \mathcal{M}^+ corresponds to a vector bundle \mathcal{V}^+ on Z^Δ . In turns, this defines a formal $\text{GL}(\mathbb{D}(Y))$ -torsor

$$\text{Isom}(\mathcal{V}^+, \mathbb{D}(Y)_{Z^\Delta}) \rightarrow Z^\Delta$$

over Z^Δ which we denote by $\Omega \rightarrow Z^\Delta$. On the other hand, $P_Z \rightarrow Z$ induces a $\Pi(\mathfrak{b}^+)^\Delta$ -torsor $P_Z^\Delta \rightarrow Z^\Delta$ of p -adic formal stacks by the following lemma.

Lemma 5.5.5. *Let T be a torsor over a quasi-syntomic scheme S over \mathbb{F}_p under a qrsp group scheme G . The prismaticisation G^Δ of G is a formal group scheme and $T^\Delta \rightarrow S^\Delta$ is a G^Δ -torsor of formal stacks.*

Proof. Since G is qrsp, the prismaticisation is a p -adic formal scheme. In addition, since prismaticisation of \mathbb{F}_p -schemes commutes with products, it follows that G^Δ is a formal group scheme. The fact that prismaticisation sends quasi-syntomic covers to fpqc covers and commutes with fibre products when the structure maps are quasi-syntomic covers tells us that $T^\Delta \rightarrow S^\Delta$ is a torsor for G^Δ . \square

5.5.6. We may apply Lemma 5.5.5 in our situation since the group scheme $\Pi(\mathfrak{b}^+)$ over $\overline{\mathbb{F}}_p$ is qrsp by Lemma 4.3.11. Consider the element $g_{\text{univ}} \in \Pi(\mathfrak{b}^+)(R)$ corresponding to the identity map $R \rightarrow R$. This element corresponds to an automorphism

$$g_{\text{univ}} : Y_R \rightarrow Y_R,$$

and this induces an automorphism of Dieudonné-modules

$$\mathbb{D}(g_{\text{univ}}) : \mathbb{D}(Y) \otimes_{\check{\mathbb{Z}}_p} A_{\text{cris}}(R) \rightarrow \mathbb{D}(Y) \otimes_{\check{\mathbb{Z}}_p} A_{\text{cris}}(R).$$

This corresponds precisely to an $\text{Spf } A_{\text{cris}}(R) = \Pi(\mathfrak{b}^+)^\Delta$ -point of $\text{GL}(\mathbb{D}(Y))$, in other words, it corresponds to a map

$$\rho : \Pi(\mathfrak{b}^+)^\Delta \rightarrow \text{GL}(\mathbb{D}(Y)).$$

Lemma 5.5.7. *The image of ρ lands in the closed subgroup $U(\mathfrak{b}^+) \subseteq \text{GL}(\mathbb{D}(Y))$. Moreover, the morphism ρ is a group homomorphism.*

Proof. The definition of $\Pi(\mathfrak{b}^+)$ tells us g_{univ} has logarithm in $\tilde{X}(\mathfrak{b})(R)$. Since the logarithm map commutes with the Dieudonné-module functor (because the functor commutes with composition and is additive) we see that $\mathbb{D}(g_{\text{univ}})$ has logarithm contained in

$$1 + \mathfrak{b} \otimes_{\mathbb{Q}_p} B_{\text{cris}}^+(R) \subset \text{End}(\mathbb{D}(Y) \otimes_{\mathbb{Z}_p} B_{\text{cris}}^+(R)).$$

Our assumption that $p > n(\mathfrak{a}) \geq n(\mathfrak{b})$ implies that the logarithm of $\mathbb{D}(g_{\text{univ}})$ in fact lies in

$$1 + \mathfrak{b}^+ \otimes_{\mathbb{Z}_p} A_{\text{cris}}(R) \subset \text{End}(\mathbb{D}(Y) \otimes_{\mathbb{Z}_p} A_{\text{cris}}(R)),$$

therefore ρ factors through the unipotent group associated to \mathfrak{b}^+ . The second claim of the lemma is that the following diagram commutes:

$$\begin{array}{ccc} \Pi(\mathfrak{b}^+)^{\Delta} \times \Pi(\mathfrak{b}^+)^{\Delta} & \xrightarrow{\rho} & \text{GL}(\mathbb{D}(Y)) \times \text{GL}(\mathbb{D}(Y)) \\ \downarrow & & \downarrow \\ \Pi(\mathfrak{b}^+)^{\Delta} & \longrightarrow & \text{GL}(\mathbb{D}(Y)). \end{array}$$

For $i = 1, 2$ let $p_{i,\text{GL}} : \text{GL}(\mathbb{D}(Y)) \times \text{GL}(\mathbb{D}(Y)) \rightarrow \text{GL}(\mathbb{D}(Y))$ and $p_{i,\Pi} : \Pi(\mathfrak{b}^+)^{\Delta} \times \Pi(\mathfrak{b}^+)^{\Delta} \rightarrow \Pi(\mathfrak{b}^+)^{\Delta}$ be the projection maps. Using the Yoneda lemma, it suffices to show the equality

$$p_{1,\text{GL}}^* \mathbb{D}(g_{\text{univ}}) \circ p_{2,\text{GL}}^* \mathbb{D}(g_{\text{univ}}) = \mathbb{D}(p_{1,\Pi}^* g_{\text{univ}} \circ p_{2,\Pi}^* g_{\text{univ}})$$

as elements of

$$\text{GL}(\mathbb{D}(Y)) \left(\Pi(\mathfrak{b}^+)^{\Delta} \times \Pi(\mathfrak{b}^+)^{\Delta} \right) = \mathbf{Aut}(\mathbb{D}(Y) \otimes_{\mathbb{Z}_p} (A_{\text{cris}}(R) \otimes_{\mathbb{Z}_p} A_{\text{cris}}(R))).$$

But functoriality of Dieudonné-theory tells us that

$$\begin{aligned} \mathbb{D}(p_{1,\Pi}^* g_{\text{univ}} \circ p_{2,\Pi}^* g_{\text{univ}}) &= \mathbb{D}(p_{1,\Pi}^* g_{\text{univ}}) \circ \mathbb{D}(p_{2,\Pi}^* g_{\text{univ}}) \\ &= p_{1,\text{GL}}^* \mathbb{D}(g_{\text{univ}}) \circ p_{2,\text{GL}}^* \mathbb{D}(g_{\text{univ}}). \end{aligned}$$

□

5.5.8. There is an isomorphism

$$h_{\text{univ}} : X_{P_Z} \rightarrow Y_{P_Z}$$

corresponding to the identity map $P_Z \rightarrow P_Z$ and applying the Dieudonné-theory functor we get an isomorphism

$$\mathcal{V}_{P_Z^{\Delta}}^+ \rightarrow \mathbb{D}(Y)_{P_Z^{\Delta}},$$

which corresponds to a morphism

$$\sigma : P_Z^{\Delta} \rightarrow \Omega.$$

Lemma 5.5.9. *The map σ is $\Pi(\mathfrak{b}^+)^{\Delta}$ -equivariant, where $\Pi(\mathfrak{b}^+)^{\Delta}$ acts on Ω via ρ .*

Proof. We are trying to show that the following diagram commutes

$$\begin{array}{ccc} \Pi(\mathfrak{b}^+)^{\Delta} \times P_Z^{\Delta} & \xrightarrow{(\rho, \sigma)} & \text{GL}(\mathbb{D}(Y)) \times \Omega \\ \downarrow & & \downarrow \\ P_Z^{\Delta} & \xrightarrow{\sigma} & \Omega, \end{array}$$

where the vertical maps are given by the respective action maps. It suffices to prove that the diagram commutes on $\Pi(\mathfrak{b}^+)^{\Delta} \times P_Z^{\Delta}$ -points, which we will write as $\text{Spf } A_{\text{cris}}(R) \times \text{Spf } A_{\text{cris}}(S)$. The identity

map $R \otimes S \rightarrow R \otimes S$ corresponds to $(g_{\text{univ}}^{\Delta}, h_{\text{univ}}^{\Delta})$ and (ρ, σ) corresponds to $(\mathbb{D}(g_{\text{univ}}), \mathbb{D}(h_{\text{univ}}))$. The map to P_Z^{Δ} corresponds to the composition $g_{\text{univ}} \circ h_{\text{univ}}$. The commutativity of the diagram is equivalent to the equality

$$\mathbb{D}(g_{\text{univ}} \circ h_{\text{univ}}) = \mathbb{D}(g_{\text{univ}}) \circ \mathbb{D}(h_{\text{univ}}),$$

which follows from functoriality of Dieudonné-theory. \square

Since ρ factors through $U(\mathfrak{b}^+)$, we get a reduction of the $\text{GL}(\mathbb{D}(Y))$ -torsor $\mathfrak{Q} \rightarrow Z^{\Delta}$ to a $U(\mathfrak{b}^+)$ -torsor $\mathfrak{R} \rightarrow Z^{\Delta}$ sitting between P_Z^{Δ} and \mathfrak{Q} . We can associate to this the symmetric tensor functor

$$\Psi : \text{Rep}_{\check{\mathbb{Z}}_p}(U(\mathfrak{b}^+)) \rightarrow \text{Vect}(Z^{\Delta})$$

which sends $V \in \text{Rep}_{\check{\mathbb{Z}}_p}(U(\mathfrak{b}^+))$ to

$$\mathfrak{R} \times^{U(\mathfrak{b}^+)} (V \otimes_{\check{\mathbb{Z}}_p} Z^{\Delta}).$$

The tautological representation $U(\mathfrak{b}^+) \hookrightarrow \text{GL}(\mathbb{D}(Y))$ is sent by Ψ to the vector bundle \mathcal{V}^+ . Note that since Z is Frobenius-smooth, the vector bundle

$$\mathfrak{R} \times^{U(\mathfrak{b}^+)} (V \otimes_{\check{\mathbb{Z}}_p} Z^{\Delta})$$

over the formal stack Z^{Δ} is simply the vector bundle

$$\mathfrak{R}_{W(Z^{\text{perf}})} \times^{U(\mathfrak{b}^+)} (V \otimes_{\check{\mathbb{Z}}_p} W(Z^{\text{perf}}))$$

over the formal scheme $W(Z^{\text{perf}})$ over $\check{\mathbb{Z}}_p$, together with a descent datum with respect to the fpqc cover $W(Z^{\text{perf}}) \rightarrow Z^{\Delta}$.

In order to pass to the generic fibre, we need the following result, which is a special case of the proposition stated in Section 6.4 of [69].

Lemma 5.5.10. *Let \mathfrak{G} be a smooth group scheme over $\check{\mathbb{Z}}_p$ with generic fibre \mathfrak{G}_{η} . Then every representation $\rho : \mathfrak{G}_{\eta} \rightarrow \text{GL}(V)$, where V is a finite dimensional $\check{\mathbb{Q}}_p$ -vector space, extends to a representation $\mathfrak{G} \rightarrow \text{GL}(\Lambda)$ for some $\check{\mathbb{Z}}_p$ -lattice $\Lambda \subseteq V$.*

Applying the lemma and passing to isogeny categories, we get an exact tensor functor

$$\Psi_{\check{\mathbb{Q}}_p} : \text{Rep}_{\check{\mathbb{Q}}_p}(U(\mathfrak{b})) \rightarrow \text{Vect}(Z^{\Delta})[\frac{1}{p}]$$

sending the defining representation of $U(\mathfrak{b})$ to \mathcal{V} . We can compose this with the natural inclusion

$$\text{Vect}(Z^{\Delta})[\frac{1}{p}] \hookrightarrow \text{Isoc}(Z)$$

and apply Tannaka duality to get a morphism of group schemes

$$G(\mathcal{M}_Z) \rightarrow U(\mathfrak{b}).$$

This is a closed immersion because the constructed functor

$$\Psi_{\check{\mathbb{Q}}_p} : \text{Rep}_{\check{\mathbb{Q}}_p}(U(\mathfrak{b})) \rightarrow \text{Isoc}(Z)$$

between Tannakian categories commutes with the $\check{\mathbb{Q}}_p$ -linear fibre functor obtained by restricting the objects to the closed point of Z (see Proposition 2.21.(b) of [25]). \square

6. RIGIDITY

6.1. Statement. Let $(\mathfrak{a}^+, \varphi_{\mathfrak{a}^+}, [-, -])$ be a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebra equipped with a strongly non-trivial action of an algebraic group Q (see Definition 4.3.16) and let $\Gamma \subseteq Q(\mathbb{Q}_p)$ be a compact open subgroup preserving \mathfrak{a}^+ . We assume as always that $p > n(\mathfrak{a})$.

Recall that for any F -stable Lie subalgebra $\mathfrak{b} \subseteq \mathfrak{a}$ there is a Dieudonné–Lie $\check{\mathbb{Z}}_p$ -subalgebra $\mathfrak{b}^+ \subseteq \mathfrak{a}^+$ defined by $\mathfrak{b}^+ = \mathfrak{b} \cap \mathfrak{a}^+$. In particular, such a \mathfrak{b} defines a subspace

$$Z(\mathfrak{b}^+) \subseteq Z(\mathfrak{a}^+) = \frac{\tilde{\Pi}(\mathfrak{a})}{\Pi(\mathfrak{a}^+)}.$$

Theorem 6.1.1 (Rigidity). *Let $Z \subseteq Z(\mathfrak{a}^+)$ be a Γ -stable integral closed formal subscheme. Then there is an F -stable Lie subalgebra $\mathfrak{b}_Z \subseteq \mathfrak{a}$ such that $Z = Z(\mathfrak{b}_Z^+)$.*

In other words, every subspace $Z \subseteq Z(\mathfrak{a}^+)$ that is stable under a strongly non-trivial action of a p -adic Lie group is strongly Tate-linear. This confirms an expectation of Chai–Oort, see Section 6.3 of [16]. Our proof is inspired by the proof of the rigidity result for biextensions by Chai–Oort in [20]. We will make a consistent effort throughout to point out the similarities between their proof and our proof, because our proof uses a different language than theirs (Dieudonné–Lie algebras vs biextensions). We start the proof by noticing that the statement is invariant under isogenies.

Lemma 6.1.2. *If $(\mathfrak{a}_1^+, \varphi_{\mathfrak{a}_1^+}, [-, -])$ and $(\mathfrak{a}_2^+, \varphi_{\mathfrak{a}_2^+}, [-, -])$ are isogenous Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebras and Theorem 6.1.1 is true for $Z(\mathfrak{a}_1^+)$ for every choice of Γ , then the same is true for $Z(\mathfrak{a}_2^+)$.*

Proof. Suppose that $\mathfrak{a}_1^+ \rightarrow \mathfrak{a}_2^+$ is a morphism of Dieudonné–Lie $\check{\mathbb{Z}}_p$ -algebras that is an isomorphism after inverting p . Then there is a closed immersion of Lie $\underline{\mathbb{Z}}_p$ -algebras

$$T_p \mathbb{X}(\mathfrak{a}_1^+) \hookrightarrow T_p \mathbb{X}(\mathfrak{a}_2^+),$$

with finite flat cokernel. In particular, there is a closed immersion of group schemes

$$\Pi(\mathfrak{a}_1^+) \hookrightarrow \Pi(\mathfrak{a}_2^+)$$

that induces a finite flat surjective map (note the reversal of the order)

$$(6.1.1) \quad Z(\mathfrak{a}_2^+) \rightarrow Z(\mathfrak{a}_1^+).$$

For $i = 1, 2$, let Γ_i be a compact open subgroup of $Q(\mathbb{Q}_p)$ preserving \mathfrak{a}_i^+ . Without loss of generality, we may assume that $\Gamma_1 \subseteq \Gamma_2$. Let $Z_2 \subseteq Z(\mathfrak{a}_2^+)$ be a Γ_2 -stable integral closed formal subscheme as in the statement of Theorem 6.1.1.

Let Z_1 be the scheme-theoretic image of Z_2 under (6.1.1). The formal subscheme $Z_1 \subseteq Z(\mathfrak{a}_1^+)$ is Γ_1 -stable by Lemma 2.1.4 and integral by construction. This means that the assumptions of Theorem 6.1.1 are also satisfied for Z_1 . If we assume that Theorem 6.1.1 is true for $Z(\mathfrak{a}_1^+)$, then there is an F -stable Lie subalgebra $\mathfrak{b} \subseteq \mathfrak{a} = \mathfrak{a}_1^+[\frac{1}{p}]$ with associated $\check{\mathbb{Z}}_p$ -lattice $\mathfrak{b}_1^+ \subseteq \mathfrak{a}_1^+$ such that

$$Z_1 = Z(\mathfrak{b}_1^+).$$

If we define $\mathfrak{b}_2^+ := \mathfrak{a}_2^+ \cap \mathfrak{b}$, we claim that $Z_2 = Z(\mathfrak{b}_2^+)$. It is clear from the construction that $Z(\mathfrak{b}_2^+)$ is the inverse image under (6.1.1) of $Z(\mathfrak{b}_1^+)$ and therefore we find that $Z_2 \subseteq Z(\mathfrak{b}_2^+)$. But both formal schemes are formal spectra of complete local domains of the same dimension and therefore this inclusion has to be an isomorphism. \square

Hypothesis 6.1.3. Thanks to Lemma 6.1.2, during the proof of Theorem 6.1.1 throughout Section 6 we may (and will) assume that $\mathbb{X}(\mathfrak{b}^+)$ is completely slope divisible. In addition, we will assume that Z is not contained in $Z(\mathfrak{e}^+)$ for any F -stable Lie subalgebra $\mathfrak{e} \subsetneq \mathfrak{a}$. This can be done without loss of generality by replacing \mathfrak{a} by the smallest F -stable Lie subalgebra $\mathfrak{d} \subset \mathfrak{a}$ such that $Z \subset Z(\mathfrak{d}^+)$.

6.2. An approximation lemma. Let μ_1 be the smallest slope of the Dieudonné–Lie algebra \mathfrak{a} and let $\mathfrak{b} \subseteq \mathfrak{a}$ be the largest F -stable subalgebra that is isoclinic of slope μ_1 . Lemma 4.3.3 tell us that \mathfrak{b} is contained in the center of \mathfrak{a} and therefore, if we equipped it with the trivial Lie bracket, it becomes an F -stable Lie subalgebra of \mathfrak{a} . We then define $\mathfrak{c} := \frac{\mathfrak{a}}{\mathfrak{b}}$ and $\mathfrak{c}^+ := \frac{\mathfrak{a}^+}{\mathfrak{b}^+}$.

The short exact sequence

$$0 \rightarrow \mathfrak{b} \rightarrow \mathfrak{a} \rightarrow \mathfrak{c} \rightarrow 0$$

has a unique F -equivariant $\check{\mathbb{Q}}_p$ -linear splitting for slope reasons. Using Lemma 6.1.2 we may assume without loss of generality that this is induced by a (unique) F -equivariant $\check{\mathbb{Z}}_p$ -linear splitting of

$$0 \rightarrow \mathfrak{b}^+ \rightarrow \mathfrak{a}^+ \rightarrow \mathfrak{c}^+ \rightarrow 0.$$

We let $\sigma : \mathfrak{c}^+ \rightarrow \mathfrak{a}^+$ be the (unique) F -equivariant section of the natural map, and similarly we let $\rho : \mathfrak{a}^+ \rightarrow \mathfrak{b}^+$ be the (unique) F -equivariant of the natural map. These maps are automatically equivariant for the action of Γ because Γ preserves slope filtrations⁵, but they are generally *not* compatible with the Lie brackets. We also let ρ and σ denote the induced maps on p -divisible groups and their Tate-modules and universal covers.

Lemma 6.2.1. *Consider the morphism*

$$\beta : \tilde{\mathbb{X}}(\mathfrak{a}) \xrightarrow{\rho} \tilde{\mathbb{X}}(\mathfrak{b}) \longrightarrow \mathbb{X}(\mathfrak{b}^+).$$

Then the induced map

$$\Gamma(\mathbb{X}(\mathfrak{b}^+), \mathcal{O}) \rightarrow \Gamma(\tilde{\mathbb{X}}(\mathfrak{a}), \mathcal{O}) = \Gamma(Z(\mathfrak{a}^+), \mathcal{O})^{\text{perf}, \wedge}$$

factors through a complete restricted perfection of $\Gamma(Z(\mathfrak{a}^+), \mathcal{O})$.

Proof. Consider the Lie $\underline{\mathbb{Z}}_p$ -subalgebra

$$\frac{1}{p^n} T_p \mathbb{X}(\mathfrak{c}^+) \subseteq \tilde{\mathbb{X}}(\mathfrak{c}).$$

Using the BCH formula, we get a closed subgroup

$$\frac{1}{p^n} \Pi(\mathfrak{c}^+) \subseteq \tilde{\Pi}(\mathfrak{c}),$$

containing $\Pi(\mathfrak{c}^+)$. Therefore, it is stable under the translation action of $\Pi(\mathfrak{c}^+)$ and hence descends to a closed subscheme⁶

$$Z(\mathfrak{c}^+) [p^n] \subseteq Z(\mathfrak{c}^+).$$

We let $Z(\mathfrak{a}^+)_n$ denote the inverse image of $Z(\mathfrak{c}^+) [p^n]$ in $Z(\mathfrak{a}^+)$. Its inverse image under $\tilde{\Pi}(\mathfrak{a}) \rightarrow Z(\mathfrak{a}^+)$ can be described as the subgroup of $\tilde{\Pi}(\mathfrak{a})$ corresponding to the Lie $\underline{\mathbb{Z}}_p$ -subalgebra

$$\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^n} T_p \mathbb{X}(\sigma(\mathfrak{c}^+)) \subseteq \tilde{\mathbb{X}}(\mathfrak{a})$$

⁵Indeed, if we conjugate our splitting by the action of an element of Γ , then we get another F -equivariant splitting which equals our original splitting by uniqueness

⁶If \mathfrak{c} is an abelian Lie algebra then $Z(\mathfrak{c}^+)$ is a p -divisible group and the closed subscheme $Z(\mathfrak{c}^+) [p^n]$ defined above is indeed its p^n -torsion.

where σ is the unique F -equivariant $\tilde{\mathbb{Z}}_p$ -linear section of $\mathfrak{a}^+ \rightarrow \mathfrak{c}^+$ as before. We will often omit the σ by abuse of notation because it is unique.

Claim 6.2.2. *There exists a morphism $\eta_n : Z(\mathfrak{a}^+)_n \rightarrow \mathbb{X}(\mathfrak{b}^+)$ such that the following diagram commutes*

$$\begin{array}{ccc} \mathbb{X}(\mathfrak{b}^+) & \hookrightarrow & Z(\mathfrak{a}^+)_n \\ & \downarrow p^n & \swarrow \eta_n \\ \mathbb{X}(\mathfrak{b}^+) & & \end{array}$$

Proof. Consider the diagram

$$\begin{array}{ccc} \tilde{\mathbb{X}}(\mathfrak{b}) & \xrightarrow{\rho} & \tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^n} T_p \mathbb{X}(\mathfrak{c}^+) \\ \downarrow & & \downarrow \\ \mathbb{X}(\mathfrak{b}^+) & \xrightarrow{\quad} & Z(\mathfrak{a}^+)_n. \end{array}$$

To prove the claim, it suffices to prove that the map given by ρ , followed by the natural projection $\tilde{\mathbb{X}}(\mathfrak{b}) \rightarrow \mathbb{X}(\mathfrak{b}^+)$ and by the multiplication by p^n is invariant under the action of $\Pi(\mathfrak{a}^+)$ on the source; this action is given by multiplication inside the group $\tilde{\Pi}(\mathfrak{a})$ as governed by the BCH formula. Let (v, w) be an R -point of $\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^n} T_p \mathbb{X}(\mathfrak{c}^+)$ and let $(v', w') \in \Pi(\mathfrak{a}^+)$. Using the assumption that $p > n(\mathfrak{a})$ and [36], the BCH formula tells us that

$$\rho((v', w') \cdot (v, w)) = v + v' + \rho(\mathbb{Z}_{(p)}\text{-linear combinations of iterated Lie brackets of } w \text{ and } w').$$

Since iterated Lie brackets of w and w' (and $\mathbb{Z}_{(p)}$ -linear combinations of them) are all in

$$\frac{1}{p^n} T_p \mathbb{X}(\mathfrak{a}^+)(R)$$

by the integrality of the Lie-bracket, and since ρ sends $T_p \mathbb{X}(\mathfrak{a}^+)$ to $T_p \mathbb{X}(\mathfrak{b}^+)$, we see that the difference between $\rho((v', w') \cdot (v, w))$ and v lands in

$$\frac{1}{p^n} T_p \mathbb{X}(\mathfrak{b}^+)(R).$$

This is precisely the kernel of

$$\tilde{\mathbb{X}}(\mathfrak{b})(R) \longrightarrow \mathbb{X}(\mathfrak{b}^+)(R) \xrightarrow{p^n} \mathbb{X}(\mathfrak{b}^+)(R)$$

and so we win. \square

We define now $Z(\mathfrak{a}^+)[F^n] \subseteq Z(\mathfrak{a}^+)$ to be closed subscheme defined by $\varphi_{\text{rel}}^n(\mathfrak{m})$, where \mathfrak{m} is the maximal ideal of the coordinate ring of $Z(\mathfrak{a}^+)$ and where φ_{rel} is the Frobenius relative to $\overline{\mathbb{F}}_p$ on the coordinate ring. We will use this notation later also for other formal spectra of complete local Noetherian rings over $\overline{\mathbb{F}}_p$ with residue field $\overline{\mathbb{F}}_p$.

Let a and r be sufficiently divisible integers such that $a/r = \mu_1$ is the slope of $\mathbb{X}(\mathfrak{b}^+)$. Since $\mathbb{X}(\mathfrak{b}^+)$ is completely slope divisible this means that

$$\mathbb{X}(\mathfrak{b}^+)[p^{na}] = \mathbb{X}(\mathfrak{b}^+)[F^{nr}].$$

Let μ_0 be a rational number that is smaller than μ_1 but larger than all the slopes of the Dieudonné-Lie algebra $\mathfrak{a}/\mathfrak{b}$. We can write $\mu_0 = a/s$ for $s > r$, possibly after multiplying both a and r by suitable positive integers.

Claim 6.2.3. *There is an integer n_0 such that for all $n \geq n_0$ we have $Z(\mathfrak{a}^+)[F^{ns}] \subseteq Z(\mathfrak{a}^+)_{na}$. Moreover, the restriction of η_{na} to $Z(\mathfrak{a}^+)[F^{ns}]$ factors through $\mathbb{X}(\mathfrak{b}^+)[F^{ns}]$.*

Proof. For the first claim it suffices to show that there exists an n_0 such that $Z(\mathfrak{c}^+)[F^{ns}] \subseteq Z(\mathfrak{c}^+)[p^{na}]$ for all $n \geq n_0$. If the Lie algebra \mathfrak{c} is abelian, then $Z(\mathfrak{c}^+)[p^n]$ is the actual p^n -torsion of the p -divisible group $\mathbb{X}(\mathfrak{c}^+)$ and the result follows from general properties of slopes. In general the result is obtained by induction because $Z(\mathfrak{c}^+)[p^n]$ is a tower of torsors for $H_i[p^n]$ where H_i is a p -divisible group of slope less than μ_0 . The second part of the claim follows from the fact that maps between formal spectra of complete local rings induce local homomorphisms on coordinate rings. \square

Now write $\mathbb{X}(\mathfrak{b}^+) = \mathrm{Spf} \overline{\mathbb{F}}_p[[u_1, \dots, u_d]]$ with

$$[p^a]^* u_j = u_j^{p^r}.$$

This is possible because $\mathbb{X}(\mathfrak{b}^+)$ is completely slope divisible. The map $\beta : \tilde{\mathbb{X}}(\mathfrak{a}) \rightarrow \mathbb{X}(\mathfrak{b}^+)$ corresponds to a d -tuple of topologically nilpotent elements a_1, \dots, a_d in

$$\Gamma(\tilde{\mathbb{X}}(\mathfrak{a}), \mathcal{O}) = \Gamma(Z(\mathfrak{a}^+), \mathcal{O})^{\mathrm{perf}, \wedge}.$$

The geometric discussion above tells us that the image of $a_j^{p^{nr}}$ in

$$\Gamma\left(\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^{na}} T_p \mathbb{X}(\mathfrak{c}^+), \mathcal{O}\right)$$

lies in the image of

$$R = \Gamma(Z(\mathfrak{a}^+)_{na}, \mathcal{O}) \rightarrow \Gamma\left(\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^{na}} T_p \mathbb{X}(\mathfrak{c}^+), \mathcal{O}\right)$$

Since $Z(\mathfrak{a}^+)_{na} \supseteq Z(\mathfrak{a}^+)[F^{ns}]$ this implies that for all $n \geq n_0$ the elements $a_i^{p^{nr}}$ are congruent to elements $c_{i,n}$ of R modulo $\varphi_{\mathrm{rel}}^{ns}(\mathfrak{m}_R)$ for all $n \geq n_0$. By definition, this means that for all i we have

$$a_i \in \Gamma(Z(\mathfrak{a}^+), \mathcal{O})_{r,s \geq n_0}^{\mathrm{perf}, \wedge}.$$

\square

Remark 6.2.4. In [20] Chai and Oort construct the morphisms η_n for all n by explicitly working with biextensions. They then define $a_{n,j} := \eta_n^* u_j \in R/\varphi_{\mathrm{rel}}^{ns}(\mathfrak{m}_R)$ which satisfy $a_{n,j}^{p^r} = a_{n+1,j}$ in $R/\varphi_{\mathrm{rel}}^{ns+r}(\mathfrak{m}_R)$, and use this to motivate the definitions of certain complete restricted perfection. They then use these complete restricted perfections to define (a variant of) ρ . In our proof, we are doing the opposite. We first constructed ρ and then used it to define the morphisms η_n .

6.3. Proof of the Theorem 6.1.1. We want to prove the result by induction on the dimension of $Z(\mathfrak{a}^+)$. The base case when $Z(\mathfrak{a}^+)$ has dimension zero is vacuous because $Z(\mathfrak{a}^+)$ is a point. Let $\tilde{Z} \hookrightarrow \tilde{\mathbb{X}}(\mathfrak{a})$ be the perfection of the closed immersion $Z \hookrightarrow Z(\mathfrak{a}^+)$ and consider the Γ -equivariant morphism

$$\alpha : \tilde{Z} \hookrightarrow \tilde{\mathbb{X}}(\mathfrak{a}) \xrightarrow{\rho} \tilde{\mathbb{X}}(\mathfrak{b}) \longrightarrow \mathbb{X}(\mathfrak{b}^+)$$

where \mathfrak{b} is as in Section 6.2. We denote by $T \subseteq \mathbb{X}(\mathfrak{b}^+)$ the scheme-theoretic image of α . The formal scheme T is integral and Γ -invariant by Lemma 2.1.4, thus the main result of [15] tells us that T is a p -divisible subgroup. In other words, $T = Z(\mathfrak{b}_1^+)$ for some Dieudonné–Lie subalgebra $\mathfrak{b}_1 \subseteq \mathfrak{b}$. Note that $\tilde{\mathbb{X}}(\mathfrak{b})$ acts on $\tilde{\mathbb{X}}(\mathfrak{a})$ by translation and that this action induces an action of the p -divisible group $\mathbb{X}(\mathfrak{b}^+) = Z(\mathfrak{b}^+)$ on $Z(\mathfrak{a}^+)$.

We need now the following two results.

Proposition 6.3.1. *The closed formal subscheme $Z \subseteq Z(\mathfrak{a}^+)$ is stable under the action of $Z(\mathfrak{b}_1^+)$.*

Proposition 6.3.2. *The subgroup $Z(\mathfrak{b}_1^+) \subseteq Z(\mathfrak{b}^+)$ is non-trivial.*

If we assume both propositions, we deduce that there is an inclusion

$$\frac{Z}{Z(\mathfrak{b}_1^+)} \subseteq \frac{Z(\mathfrak{a}^+)}{Z(\mathfrak{b}_1^+)} = Z\left(\frac{\mathfrak{a}^+}{\mathfrak{b}_1^+}\right),$$

and that the dimension of $Z\left(\frac{\mathfrak{a}^+}{\mathfrak{b}_1^+}\right)$ is smaller than the dimension of $Z(\mathfrak{a}^+)$. Therefore, by the inductive hypothesis, we find that

$$\frac{Z}{Z(\mathfrak{b}_1^+)} = Z(\mathfrak{d}^+)$$

for some F -stable Lie subalgebra $\mathfrak{d} \subseteq \frac{\mathfrak{a}}{\mathfrak{b}_1}$. However, if $\mathfrak{d} \neq \frac{\mathfrak{a}}{\mathfrak{b}_1}$ then Z is contained in $Z(\pi^{-1}(\mathfrak{d}^+))$, where

$$\pi : \mathfrak{a} \rightarrow \frac{\mathfrak{a}}{\mathfrak{b}_1}$$

is the natural projection. This contradicts our assumption that Z is not contained in $Z(\mathfrak{e}^+)$ for any F -stable Lie subalgebra $\mathfrak{e} \subsetneq \mathfrak{a}$.

We end Section 6 with a proof of Proposition 6.3.1 and Proposition 6.3.2. Note that Proposition 6.3.1 is an analogue of Proposition 7.3 of [20] and Proposition 6.3.2 is a generalisation of Proposition 6.2 of [ibid.], as explained in Remark 6.5.2.

6.4. Proof of Proposition 6.3.1. Let $X \in \text{Lie } \Gamma$ and choose $n \gg 0$ such that $g = \exp(p^n X)$ converges in Γ . Write $X = (B, C)$ where B is the induced endomorphism of \mathfrak{b}^+ and C is the induced endomorphism of \mathfrak{c}^+ and consider the following commutative diagram (compare it with the diagram on page 64 of [20])

$$(6.4.1) \quad \begin{array}{ccccc} Z & \hookrightarrow & Z(\mathfrak{a}^+) & \xlongequal{\quad} & Z(\mathfrak{a}^+) \\ & & \star \uparrow & & \star \uparrow \\ & & \mathbb{X}(\mathfrak{b}_1^+) \times Z & \longrightarrow & \mathbb{X}(\mathfrak{b}^+) \times Z(\mathfrak{a}^+) \\ & & B \times 1 \uparrow & & B \times 1 \uparrow \\ & & \mathbb{X}(\mathfrak{b}_1^+) \times Z & \hookrightarrow & \mathbb{X}(\mathfrak{b}^+) \times Z(\mathfrak{a}^+) & \swarrow & \tilde{\Pi}(\mathfrak{a}) \\ & & \uparrow & & \uparrow & & \star \uparrow \\ & & \tilde{\mathbb{X}}(\mathfrak{b}_1) \times \tilde{Z} & \hookrightarrow & \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) & \xrightarrow{B \times 1} & \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) \\ & & \alpha \times 1 \uparrow & & \rho \times 1 \uparrow & & \\ & & \tilde{Z} \times \tilde{Z} & \hookrightarrow & \tilde{\Pi}(\mathfrak{a}) \times \tilde{\Pi}(\mathfrak{a}) & & \\ & & (\varphi_{\text{rel}^1}^{nr}) \uparrow & & (\varphi_{\text{rel}^1}^{nr}) \uparrow & & \\ & & \tilde{Z} & \longrightarrow & \tilde{\Pi}(\mathfrak{a}). & & \end{array}$$

Lemma 6.4.1. *The dotted arrow exists.*

Proof. We start by studying the automorphism of $\frac{1}{p^n}T_p\mathbb{X}(\mathfrak{c}^+)$ induced by $g_n = \text{Exp}(p^n X)$ for $n \gg 0$. It follows from the Taylor series for the exponential map that g_n is congruent to the identity modulo $T_p\mathbb{X}(\mathfrak{c}^+)$, thus g_n induces the identity on $Z(\mathfrak{c}^+)[p^n]$. Similarly we can write the automorphism of $\tilde{\mathbb{X}}(\mathfrak{b})$ induced by g_n as

$$1 + p^n \circ B + \sum_{j \geq 2} \frac{p^{jn}}{j!} B^j,$$

and this shows that the automorphism of $\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^n}T_p\mathbb{X}(\mathfrak{c}^+)$ induced by g_n is congruent modulo $T_p\mathbb{X}(\mathfrak{b}^+) \oplus T_p\mathbb{X}(\mathfrak{c}^+)$ to

$$\left(B \circ [p^n] \circ \rho + \sum_{j \geq 2} \frac{p^{jn}}{j!} B^j \circ \rho \right) \star 1,$$

where 1 is the identity map and \star denotes the translation action of $\tilde{\mathbb{X}}(\mathfrak{b})$ on $\tilde{\mathbb{X}}(\mathfrak{b}) \oplus \frac{1}{p^n}T_p\mathbb{X}(\mathfrak{c}^+)$.⁷ We know that g_n induces an automorphism of the quotient $Z(\mathfrak{a}^+)_n$ and there g_n acts via the following formula

$$g_n = \left(B \circ \eta_n + \sum_{j \geq 2} \frac{p^{(j-1)n}}{j!} B^j \circ \eta_n \right) \star 1,$$

where 1 denotes the identity on $Z(\mathfrak{a}^+)_n$, where \star denotes the translation action of $\mathbb{X}(\mathfrak{b}^+)$ on $Z(\mathfrak{a}^+)_n$, and where B denotes the endomorphism of $\mathbb{X}(\mathfrak{a}^+)$ induced by B . If we now look at the action of $\exp(p^{na} X)$ on $Z(\mathfrak{a}^+)_{na}$ then it is given by the following formula

$$g_{na} = \left(B \circ \eta_{na} + \sum_{j \geq 2} \frac{p^{(j-1)na}}{j!} B^j \circ \eta_{na} \right) \star 1$$

Claim 6.4.2. *There exists a constant c such that for all sufficiently large n there is an equality of endomorphisms*

$$\left(B \circ \eta_{na} + \sum_{j \geq 2} \frac{p^{(j-1)na}}{j!} B^j \circ \eta_{na} \right) \star 1 = (B \circ \eta_{na}) \star 1$$

of the scheme $Z(\mathfrak{a}^+)_{na}[F^{2nr-c}]$.

Proof. This just comes down to showing that

$$\frac{p^{(j-1)na}}{j!} B^j \circ \eta_{na} \equiv 0 \pmod{\varphi_{\text{rel}}^{2nr-c}(\mathfrak{m})}.$$

It is not so hard to see that η_{na} itself is zero modulo $\varphi_{\text{rel}}^{nr}(\mathfrak{m})$ because $p^{na} = \varphi_{\text{rel}}^{nr}$ on $\mathbb{X}(\mathfrak{b}^+)$. Thus we just need to remark that the p -adic valuation of

$$\frac{p^{(j-1)na}}{j!}$$

is at least $na - c$ for $n \gg 0$ to conclude, which is an elementary calculation. \square

⁷The same formula appears in Proposition 5.3.2 of [20] and inspired ours.

Let $Q \in R = \Gamma(Z(\mathfrak{a}^+), \mathcal{O})$ be a function that vanishes on Z , to prove Lemma 6.4.1 we have to show that the pullback

$$(B \times 1)^* \circ \star^* Q$$

vanishes. We first note that the map $\tilde{Z} \times Z \rightarrow \mathbb{X}(\mathfrak{b}_1^+) \times Z$ given by composing $\alpha \times 1$ with the natural map $\tilde{X}(\mathfrak{b}) \rightarrow \mathbb{X}(\mathfrak{b}_1^+)$ is injective on coordinate rings. Indeed this is true for $\tilde{X}(\mathfrak{b}) \rightarrow \mathbb{X}(\mathfrak{b}_1^+)$ by Lemma A.1.6, it is true for $\alpha : \tilde{Z} \rightarrow X(\mathfrak{b}_1)$ by the definition of scheme-theoretic image and by Lemma 2.1.6, the induced map on completed tensor products is also injective. Thanks to this observation, it suffices to show that the further pullback

$$(6.4.2) \quad (\alpha \times 1)^* \circ (B \times 1)^* \circ \star^* Q$$

vanishes. By Proposition 2.1 of [15], there is an injective local homomorphism

$$\Gamma(Z, \mathcal{O}) \rightarrow \overline{\mathbb{F}}_p[[X_1, \dots, X_m]] =: S.$$

If we write W for $\mathrm{Spf} S$, we have the following commutative diagram

$$\begin{array}{ccc} \tilde{W} \times \tilde{W} & \longrightarrow & \tilde{Z} \times \tilde{Z} \\ (\varphi_{\mathrm{rel}, 1}^{nr}) \uparrow & & (\varphi_{\mathrm{rel}, 1}^{nr}) \uparrow \\ \tilde{W} & \longrightarrow & \tilde{Z}. \end{array}$$

Note that the induced map

$$\Gamma(\tilde{Z} \times \tilde{Z}, \mathcal{O}) \rightarrow \Gamma(\tilde{W} \times \tilde{W}, \mathcal{O})$$

is again injective because the formation of completed perfection commutes with completed tensor products by Lemma A.1.12 and completed perfection of injective maps are injective by Lemma A.1.9. Thus we want to prove that the image of (6.4.2) in

$$\Gamma(\tilde{W} \times \tilde{W}, \mathcal{O}) = \overline{\mathbb{F}}_p[[X_1, \dots, X_m, Y_1, \dots, Y_m]]^{\mathrm{perf}, \wedge}$$

is zero. Choose coordinates u_1, \dots, u_d for $X(\mathfrak{b})$ and coordinates v_1, \dots, v_e for $Z(\mathfrak{a}^+)$. The pullbacks of u_1, \dots, u_d under

$$\tilde{W} \longrightarrow \tilde{Z} \xrightarrow{\alpha} X(\mathfrak{b})$$

are topologically nilpotent elements g_1, \dots, g_d contained, by Lemma 6.2.1, in $\overline{\mathbb{F}}_p[[X_1, \dots, X_m]]_{r, s, \geq n_0}^{\mathrm{perf}, \wedge}$. Similarly the pullbacks of v_1, \dots, v_e under

$$\tilde{W} \longrightarrow \tilde{Z} \longrightarrow \tilde{\Pi}(\mathfrak{a}) \longrightarrow Z(\mathfrak{a}^+)$$

are given by topologically nilpotent elements

$$h_1, \dots, h_e \in \overline{\mathbb{F}}_p[[X_1, \dots, X_m]].$$

If we write the pullback of a to $\mathbb{X}(\mathfrak{b}^+) \times Z(\mathfrak{a}^+)$ as a power series $f(u_1, \dots, u_d, v_1, \dots, v_e)$, then the image of (6.4.2) in

$$\overline{\mathbb{F}}_p[[X_1, \dots, X_m, Y_1, \dots, Y_m]]^{\mathrm{perf}, \wedge}$$

is given by

$$f(g_1, \dots, g_d, h_1, \dots, h_e).$$

We note that in this case the composition of completed perfected power series makes sense. Lemma 6.4.1 follows then from the following claim. \square

Claim 6.4.3. *The function $f(g_1, \dots, g_d, h_1, \dots, h_e)$ vanishes.*

Proof. Proposition 6.4 of [20] (stated in Appendix A as Theorem A.2.4) tells us that to show that $f(g_1, \dots, g_d, h_1, \dots, h_e) = 0$ it suffices to find an integer r and a sequence of integers d_i such that

$$\lim_{n \rightarrow \infty} \frac{p^{rn}}{d_i} = 0$$

and such that (with $q = p^r$)

$$f(g_1(\underline{X})^{q^n}, \dots, g_e(\underline{X})^{q^n}, h_1(\underline{X}), \dots, h_d(\underline{X})) \equiv 0 \pmod{(X_1, \dots, X_m)^{d^n}}.$$

Note that

$$f(g_1(\underline{X})^{q^n}, \dots, g_d(\underline{X})^{q^n}, h_1(\underline{X}), \dots, h_e(\underline{X}))$$

is precisely the pullback of $f(g_1, \dots, g_d, h_1, \dots, h_e)$ under $(\varphi_{\text{rel}}^{nr}, 1)$. We will apply this theorem with r as above, where we recall that $a/r = \mu_1$ is the slope of $\mathbb{X}(\mathfrak{b}_1^+)$, and we will take $d^n = p^{2mnr-c'}$ where c' is some constant. Proving the congruence comes down to chasing the diagram in (6.4.1) and applying the congruence from Claim 6.4.2.

Indeed, our assumption that Z is stable under the action of Γ tells us that the pullback of Q under the automorphism induced by $g_{na} = \exp(p^{na}X)$ of $Z(\mathfrak{a}^+)$ maps to zero in $\Gamma(Z, \mathcal{O})$, and so further pullback to $\Gamma(\tilde{\Pi}(\mathfrak{a}), \mathcal{O})$ maps to zero in $\Gamma(\tilde{Z}, \mathcal{O})$. The congruence

$$\exp(p^n X) \equiv (B \circ \eta_{na}) \star 1 \pmod{\varphi_{\text{rel}}^{2nr-c}(\mathfrak{m})}$$

tells us that the pullback of Q via $(B \circ p^{na} \circ \rho) \star 1$, maps to zero in $\Gamma(Z, \mathcal{O})$ modulo $\varphi_{\text{rel}}^{2nr-c}(\mathfrak{m})$. Since $\eta_{na} = [p^n] \circ \rho_{na}$ the further pullback to $\Gamma(\tilde{\Pi}(\mathfrak{a}), \mathcal{O})$ is the same as the pullback of Q via

$$\tilde{\Pi}(\mathfrak{a}) \xrightarrow{(\rho, 1)} \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) \xrightarrow{(B \circ p^{na}, 1)} \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) \xrightarrow{\star} \tilde{\Pi}(\mathfrak{a}) \longrightarrow Z(\mathfrak{a}^+).$$

Since $p^{na} = \varphi_{\text{rel}}^{nr}$ on $\mathbb{X}(\mathfrak{b}^+)$ we can identify this morphism instead with

$$\tilde{\Pi}(\mathfrak{a}) \xrightarrow{(\varphi_{\text{rel}}^{nr}, 1)} \tilde{\Pi}(\mathfrak{a}) \times \tilde{\Pi}(\mathfrak{a}) \xrightarrow{\rho \times 1} \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) \xrightarrow{(B, 1)} \tilde{\mathbb{X}}(\mathfrak{b}) \times \tilde{\Pi}(\mathfrak{a}) \xrightarrow{\star} \tilde{\Pi}(\mathfrak{a}) \longrightarrow Z(\mathfrak{a}^+).$$

To summarise, the pullback of Q along the composition of this morphism with $\tilde{Z} \rightarrow \tilde{\Pi}(\mathfrak{a})$ vanishes modulo $\varphi_{\text{rel}}^{2nr-c}$. By the commutativity of the big diagram (6.4.1) this is precisely saying that the pullback of Q along the five vertical maps in the second column vanishes modulo $\varphi_{\text{rel}}^{2nr-c}(\mathfrak{m})$, and so the same is true for $f(g_1(\underline{X})^{q^n}, \dots, g_d(\underline{X})^{q^n}, h_1(\underline{X}), \dots, h_e(\underline{X}))$. To be precise, it is zero modulo

$$(X_1^{p^{2nr-c}}, \dots, X_m^{p^{2nr-c}}) \supseteq (X_1, \dots, X_n)^{p^{2mnr-cm}},$$

and so Proposition 6.4 of [20] applies and we win. \square

Finally, Proposition 6.3.1 follows from Lemma 6.4.1 by applying Lemma 4.1.1 of [15], as in the proof of Theorem 7.2 of [20] or the proof of Theorem 4.3 of [15]. The assumptions of this lemma are satisfied because the action of Γ is strongly non-trivial.

6.5. Proof of Proposition 6.3.2. Suppose that the scheme-theoretic image of α is trivial and let σ be the (unique) F -equivariant section of $\mathfrak{a} \rightarrow \mathfrak{c}$. Our assumption implies that \tilde{Z} is contained in the image of this section. In other words, it is contained in

$$\tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \subseteq \tilde{\mathbb{X}}(\mathfrak{a}) = \tilde{\Pi}(\mathfrak{a}).$$

We are going to show that $\tilde{\mathbb{X}}(\sigma(\mathfrak{c}))$ is a Lie subalgebra of $\tilde{\mathbb{X}}(\mathfrak{a})$, contradicting our assumption that Z is not contained in $Z(\mathfrak{d}^+)$ for an F -stable Lie-subalgebra $\mathfrak{d} \subseteq \mathfrak{a}$. This contradiction uses the following claim.

Claim 6.5.1. *The \mathbb{Q}_p -subspace*

$$\tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \subseteq \tilde{\mathbb{X}}(\mathfrak{a})$$

is stable under the Lie bracket if and only if the F -stable \mathbb{Q}_p -subspace

$$\sigma(\mathfrak{c}) \subseteq \mathfrak{a}$$

is stable under the Lie bracket.

Proof. This is an immediate consequence of Lemma 4.3.5. □

Let Z' be the inverse image of Z in $\tilde{\Pi}(\mathfrak{a})$, and note that

$$Z' \rightarrow Z$$

is a torsor for $\Pi(\mathfrak{a}^+)$. The restriction of ρ to Z' factors through $T_p\mathbb{X}(\mathfrak{b}^+) \subseteq \tilde{\mathbb{X}}(\mathfrak{b})$ by the assumption that the scheme-theoretic image of \tilde{Z} under α is trivial. In fact Z' surjects onto $T_p\mathbb{X}(\mathfrak{b}^+)$ because ρ is equivariant for the translation action of $T_p\mathbb{X}(\mathfrak{b}^+)$ via $T_p\mathbb{X}(\mathfrak{b}^+) \subseteq \Pi(\mathfrak{a}^+)$. Moreover since ρ is a retraction of the inclusion of $\tilde{\mathbb{X}}(\mathfrak{b})$ it follows that

$$\rho_R : Z'(R) \rightarrow T_p\mathbb{X}(\mathfrak{b}^+)(R)$$

is surjective for every $\overline{\mathbb{F}}_p$ -algebra R .

Let $x = (v, w) \in Z'(R)$ with v denoting $\rho(x)$ and w denoting the image of x in $\tilde{\Pi}(\mathfrak{c})$ and similarly let $x' = (v', w') \in \Pi(\mathfrak{a}^+)$. Then the action of x' on x can be described as

$$(v + v' + \rho([w, w']), w \cdot w').$$

where \cdot denotes the multiplication in $\tilde{\Pi}(\mathfrak{c})$. Since

$$\rho(Z'(R)) = T_p\mathbb{X}(R)$$

is a subgroup containing v, v' and $v + v' + \rho([w, w'])$ we see that it must contain $\rho([w, w'])$. The existence of the section σ shows that

$$Z'(R) \rightarrow \tilde{\Pi}(\mathfrak{c})(R)$$

is surjective for all R . So we can choose w arbitrarily and similarly we can choose w' arbitrarily. In other words

$$\rho(Z'(R)) = T_p\mathbb{X}(R)$$

contains the image of the map

$$(6.5.1) \quad T_p\mathbb{X}(\sigma(\mathfrak{c}^+))(R) \times \tilde{\mathbb{X}}(\sigma(\mathfrak{c}))(R) \rightarrow \tilde{\mathbb{X}}(\mathfrak{a})(R) \rightarrow \tilde{\mathbb{X}}(\mathfrak{b})(R),$$

where the last map is given by ρ and the first by the Lie bracket. Since the pairing is bilinear and $\tilde{\mathbb{X}}(\sigma(\mathfrak{c}))(R)$ is a \mathbb{Q}_p -vector space we see that the image of (6.5.1) is equal to the image of

$$\tilde{\mathbb{X}}(\sigma(\mathfrak{c}))(R) \times \tilde{\mathbb{X}}(\sigma(\mathfrak{c}))(R) \rightarrow \tilde{\mathbb{X}}(\mathfrak{a})(R) \rightarrow \tilde{\mathbb{X}}(\mathfrak{b})(R).$$

But this means that the morphism

$$\tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \times \tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \rightarrow \tilde{\mathbb{X}}(\mathfrak{a}) \rightarrow \tilde{\mathbb{X}}(\mathfrak{b}).$$

factors through the inclusion

$$T_p\mathbb{X}(\sigma(\mathfrak{b}^+)) \hookrightarrow \tilde{\mathbb{X}}(\sigma(\mathfrak{b}))$$

and hence is constant. Indeed the source is perfect and the coordinate ring of the target has no non-trivial maps to perfect rings because its maximal ideal coincides with the nilpotent radical. Therefore, we find that the Lie bracket

$$\tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \times \tilde{\mathbb{X}}(\sigma(\mathfrak{c})) \rightarrow \tilde{\mathbb{X}}(\mathfrak{a})$$

lands in the kernel of ρ which is given by $\tilde{\mathbb{X}}(\sigma(\mathfrak{c}))$. This leads to a contradiction using Claim 6.5.1, and so we are done. \square

Remark 6.5.2. Proposition 6.3.2 can be thought of as a generalisation of Proposition 6.2 of [20]. Indeed, this proposition states (when \mathfrak{a} has a biextension structure) that the Lie bracket on

$$\frac{\tilde{\mathbb{X}}(\mathfrak{a})}{\tilde{\mathbb{X}}(\mathfrak{b}_1)}$$

is trivial, or equivalently that the biextension is split. This is what suggested to us that Proposition 6.3.2 might be true.

7. PROOF OF THE MAIN THEOREM AND SOME VARIANTS

7.1. Preliminaries on Hecke operators. Let (G, X) be a Shimura datum of abelian type with reflex field E and let p be a prime such that $G_{\mathbb{Q}_p}$ is quasi-split and split over an unramified extension. Let $U_p \subset G(\mathbb{Q}_p)$ be a hyperspecial subgroup and let $U^p \subset G(\mathbb{A}_f^p)$ be a sufficiently small compact open subgroup. Let $\mathbf{Sh}_{G,U}/E$ be the Shimura variety of level $U = U^p U_p$ and for a prime $v|p$ of E let $\mathcal{S}_{G,U}/\mathcal{O}_{E,v}$ be the canonical integral model of $\mathbf{Sh}_{G,U}$ constructed in [48] (resp [46] for $p = 2$).

Let Sh_G be the base change to $\overline{\mathbb{F}}_p$ of this integral canonical model for some choice of map $\mathcal{O}_{E,v} \rightarrow \overline{\mathbb{F}}_p$. Let $G^{\mathrm{sc}} \rightarrow G^{\mathrm{der}}$ be the simply-connected cover of the derived group of G ; we will almost always identify groups like $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ and $G^{\mathrm{sc}}(\mathbb{Q}_\ell)$ with their images in $G(\mathbb{A}_f^p)$ and $G(\mathbb{Q}_\ell)$. Let $G \rightarrow G^c$ be the quotient of G obtained by killing the part of the centre that splits over \mathbb{R} but not over \mathbb{Q} . Let $U^{p,c}$ be the image of U^p in $G^c(\mathbb{A}_f^p)$.

There is a pro-étale $\underline{U}^{p,c}$ -torsor

$$\pi : \varprojlim_{K^p \subseteq U^p} \mathrm{Sh}_{G,K^p} \rightarrow \mathrm{Sh}_G$$

over Sh_G such that the action of U^p via $U^p \rightarrow U^{p,c}$ extends to an action of $G(\mathbb{A}_f^p)$. Let $Z \subseteq \mathrm{Sh}_G$ be a locally closed subvariety and let \tilde{Z} be the inverse image of Z under π . We say that Z is stable under the prime-to- p -Hecke operators, or that Z is $G(\mathbb{A}_f^p)$ -stable, if \tilde{Z} is $G(\mathbb{A}_f^p)$ -stable. Similarly we will call Z stable under $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ if \tilde{Z} is.

Remark 7.1.1. For $g \in G(\mathbb{A}_f^p)$ and $K^p \subset U^p$ there is a finite étale correspondence

$$\begin{array}{ccc} & \mathrm{Sh}_{G,K^p \cap gK^p g^{-1}} & \\ p_1 \swarrow & & \searrow p_2 \\ \mathrm{Sh}_{G,K^p} & & \mathrm{Sh}_{G,gK^p g^{-1}} \xrightarrow{g} \mathrm{Sh}_{G,K^p} \end{array}$$

and the Hecke operator attached to g is $g \circ p_2 \circ p_1^{-1}$. A locally closed subvariety $Z \subset \mathrm{Sh}_{G,K^p}$ is stable under the Hecke operator attached to g if and only if \tilde{Z} is stable under the action of g considered as an element of $G(\mathbb{A}_f^p)$.

7.2. Local stabiliser principle. Now suppose that (G, X) is of Hodge type and choose a Hodge embedding which leads to a closed immersion (by the main result of [71])

$$\mathrm{Sh}_G \hookrightarrow \mathcal{A}_{g,N}$$

for some $N > 0$. Let $C \subset \mathrm{Sh}_{G,b}$ be a central leaf with completely slope divisible p -divisible group y , and let $x \in C(\overline{\mathbb{F}}_p)$. Then we have seen that the profinite group $\mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p)$ acts on C/x .

For $x \in \mathrm{Sh}_G(\overline{\mathbb{F}}_p)$ we let I_x be the algebraic group over \mathbb{Q} consisting of tensor preserving self-quasi-isogenies of the abelian variety A_x , introduced by Kisin in Section 2.1.2 of [49]. By definition it is a closed subgroup of the algebraic group over \mathbb{Q} whose R -points are given by

$$\left(\mathrm{End}_{\overline{\mathbb{F}}_p}(A_x) \otimes_{\mathbb{Z}} R \right)^\times.$$

There are natural maps

$$(7.2.1) \quad \begin{aligned} I_{x, \mathbb{A}_f^p} &\rightarrow G_{\mathbb{A}_f^p} \\ I_{x, \mathbb{Q}_p} &\rightarrow J_b. \end{aligned}$$

Let $X_\mu(b)(\overline{\mathbb{F}}_p)$ be the affine Deligne-Lusztig set associated to the Newton stratum $\mathrm{Sh}_{G,b}$. This set has an action of $J_b(\mathbb{Q}_p)$ and is equipped with a point z such corresponding to our p -divisible group with G -structure Y such that the stabiliser of z under the $J_b(\mathbb{Q}_p)$ -action can be identified with the compact open subgroup

$$\mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p) \subset J_b(\mathbb{Q}_p).$$

Kisin proves (Proposition 2.1.3 of [49]) that the isogeny class \mathcal{S}_x of x can be $G(\mathbb{A}_f^p)$ -equivariantly identified with

$$I_x(\mathbb{Q}) \backslash G(\mathbb{A}_f^p) \times X_\mu(b)(\overline{\mathbb{F}}_p) / U^p$$

where $I_x(\mathbb{Q})$ acts diagonally on the product via the maps of (7.2.1). Moreover, the point $(1, y)$ is mapped to x under this identification. Define

$$I_x(\mathbb{Z}_{(p)}) = I_x(\mathbb{Q}) \cap \mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p).$$

This is precisely the stabiliser of the isogeny class \mathcal{S}_x in the inverse limit over all prime-to- p levels.

Corollary 2.3.5 of [49] tells us that I_x is an inner form of a subgroup of G , and therefore we can define I'_x as the inner form corresponding to the intersection of this subgroup with G^{sc} . Similarly we can define $J'_b \subset J_b$ because J_b is an inner form of a Levi of G . Then there are natural maps

$$\begin{aligned} I_{x, \mathbb{A}_f^p} &\rightarrow G_{\mathbb{A}_f^p}^{\mathrm{sc}} \\ I_{x, \mathbb{Q}_p} &\rightarrow J'_b. \end{aligned}$$

We define $I'_x(\mathbb{Z}_{(p)})$ to be the intersection of $I_x(\mathbb{Z}_{(p)})$ with $I'_x(\mathbb{Q})$.

Proposition 7.2.1 (Local stabiliser principle). *If $Z \subset C$ be any $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -stable closed reduced subset containing x , then $Z/x \subset C/x$ is stable under the action of*

$$I'_x(\mathbb{Z}_{(p)}) \subset \mathbf{Aut}_G(Y)(\overline{\mathbb{F}}_p).$$

Proof. This is Theorem 9.5 of [18] in the Siegel case, and the same proof goes through in our case. The main point is that the Hecke correspondences are finite étale and therefore induce isomorphisms on complete local rings. \square

7.3. Proof of Theorem I. Let (G, X) be a Shimura variety of abelian type with reflex field E , let p be a prime such that $G_{\mathbb{Q}_p}$ is quasi-split and split over an unramified extension, let $U_p \subseteq G(\mathbb{Q}_p)$ be a hyperspecial subgroup and let $U^p \subseteq G(\mathbb{A}_f^p)$ be a sufficiently small compact open subgroup. Let $v|p$ be a place of E and let Sh_G be the geometric special fibre of the canonical integral model over $\mathcal{O}_{E,v}$ of the Shimura variety \mathbf{Sh} of level $U^p U_p$ over E . Note that the canonical integral model exists by work of Kisin, [49], see [46] for the $p = 2$ case. Let $B(G, X) \subseteq B(G_{\mathbb{Q}_p})$ be the finite set of admissible σ -conjugacy classes and let $b \in B(G, X)$ with associated Newton stratum $\mathrm{Sh}_{G,b} \subseteq \mathrm{Sh}_G$ (see [32]).

Let $C \subseteq \mathrm{Sh}_{G,b}$ be a canonical central leaf as in Section 5.4.5 of [67], let ν_b be the Newton cocharacter of b and let P_{ν_b} be the associated parabolic subgroup with unipotent radical U_{ν_b} . Let $G^{\mathrm{ad}} = G_1 \times \cdots \times G_n$ be a decomposition of G^{ad} into a product of \mathbb{Q} -simple groups.

Definition 7.3.1 (Definition 5.3.2 of [51]). The element b is called \mathbb{Q} -non-basic if the image of b in $B(G_{i,\mathbb{Q}_p})$ is non-basic for all i .

Theorem 7.3.2. *Suppose that p is greater than the nilpotency class of the nilpotent Lie algebra $\mathrm{Lie} U_{\nu_b}$. Then any nonempty reduced closed subscheme $Z \subseteq C$ that is stable under $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ is equal to a union of connected components of C . If we suppose moreover that (G, X) is of Hodge type, that the element $b \in B(G, X)$ is \mathbb{Q} -non-basic and that Z is $G(\mathbb{A}_f^p)$ -stable, then $Z = C$.*

The assumption that the Coxeter number $h(G) \leq p$ in Theorem I implies that p is greater than the nilpotency class of $\mathrm{Lie} U_{\nu_b}$ by Claim 2.1.1 of [43] together with Proposition 31 in Section VI.1.11 of [7]. Therefore Theorem 7.3.2 implies Theorem I.

The discussion in Section 1.6 of [51] implies that the theorem is also true when b is \mathbb{Q} -basic, that is, when b_i is basic for all i . In this case the central leaves are finite and the claim is that $G(\mathbb{A}_f^p)$ acts transitively on them. We expect Theorem 7.3.2 to be true for arbitrary b , but we don't know how to prove the discrete part.

Proof. We reduce immediately to the case that Z is the Zariski closure of the $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -orbit of a point.

Both the assumption that Z is a closed subvariety of $\mathrm{Sh}_{G,b}$ given by the Zariski-closure of the $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -orbit of a point and the conclusion that Z is equidimensional of dimension $\langle 2\rho, \nu_b \rangle$ behave well with respect to passing from $(G, X) \rightarrow (G^{\mathrm{ad}}, X^{\mathrm{ad}})$ and behaves well with respect to products of Shimura varieties. Therefore, it suffices to prove the first part of the theorem in the case that (G, X) is of Hodge type with G^{ad} simple over \mathbb{Q} and that $Z \subseteq C$ is stable under the action of $G^{\mathrm{sc}}(\mathbb{A}_f^p)$; we will assume that we are in this case from now on.

We will also assume that b is non-basic, because if b is basic then C is a finite set and the first part of the theorem is automatically true. Proposition 2.4.5 of [45] tells us that there is a central leaf $C' \subseteq \mathrm{Sh}_{G,b}$ such that the universal p -divisible group over C' is completely slope divisible. Since C and C' share a $G(\mathbb{A}_f^p)$ -equivariant finite étale cover, it suffices to prove the theorem for C' and therefore we will assume without loss of generality that the universal p -divisible group over C is completely slope divisible.

It follows from Lemma 2.1.1 of [38] that the smooth locus Z^{sm} of Z is also $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -stable. Moreover if we choose a Hodge embedding $\mathrm{Sh}_G \hookrightarrow \mathcal{A}_{g,N}$ then we get an isocrystal \mathcal{M} associated to the universal abelian variety A/Sh_G . It follows from Corollary 2.2.1 of [38] that the monodromy group of \mathcal{M} over Z^{sm} is equal to P_{ν_b} (strictly speaking the corollary is stated for $G(\mathbb{A}_f^p)$ -stable

subvarieties but the proof goes through for $G^{\text{sc}}(\mathbb{A}_f^p)$ -stable subvarieties). Theorem II then tells us that for $x \in Z^{\text{sm}}(\overline{\mathbb{F}}_p)$ the monodromy of the isocrystal \mathcal{M} over $\text{Spec } \widehat{\mathcal{O}}_{Z,x}$ is equal to U_{ν_b} .

The assumption that the universal p -divisible group over C is completely slope divisible tells us that $Z^{/x} \subseteq \text{Def}_{\text{sus}}(Y)$ for a completely slope divisible group Y . Theorem 5.5.3 tells us that that $Z^{/x}$ is not contained in $Z(\mathfrak{b}^+)$ for any F -stable Lie algebra $\mathfrak{b} \subsetneq \mathfrak{a} = \mathbb{D}(\mathcal{H}_Y^{G,\circ})[\frac{1}{p}] = \text{Lie } U_{\nu_b}$.

Proposition 7.2.1 tells us that $Z^{/x}$ is stable under the action of

$$I'_x(\mathbb{Z}_{(p)}) \subset \mathbf{Aut}_G(\overline{\mathbb{F}}_p).$$

By continuity it is also stable under its closure $\Gamma \subset \mathbf{Aut}_G(\overline{\mathbb{F}}_p)$. It follows as in the proof of Proposition 3.2.1 of [38] that Γ acts strongly non-trivially on $C^{/x} = Z(\mathfrak{a}^+)$. Therefore Theorem 6.1.1 tells us that $Z^{/x} = Z(\mathfrak{b}^+)$ for some F -stable Lie algebra $\mathfrak{b} \subseteq \mathfrak{a}$, and the previous paragraph tells us that $\mathfrak{a} = \mathfrak{b}$. In other words, $Z^{/x} = C^{/x}$ for all points $x \in Z^{\text{sm}}(\overline{\mathbb{F}}_p)$. Since $Z^{\text{sm}} \subseteq Z$ is dense because Z is reduced, it follows that Z is equidimensional of the same dimension as C and therefore it is a union of connected components of C .

The further claim that if (G, X) is of Hodge type, the element $b \in B(G, X)$ is assumed to be \mathbb{Q} -non-basic and Z is $G(\mathbb{A}_f^p)$ -stable that then $Z = C$ follows from this in combination with Theorem C of [51], which states that under these assumptions any $G(\mathbb{A}_f^p)$ -stable subvariety $Z \subseteq C$ intersects each connected component of C non-trivially. \square

7.4. Isogeny classes are dense in Newton strata. Let (G, X) be a Shimura variety of Hodge type, let $x \in \text{Sh}_{G,b}(\overline{\mathbb{F}}_p)$ and let $\mathcal{I}_x \subseteq \text{Sh}_{G,b}(\overline{\mathbb{F}}_p)$ be the isogeny class of x in the sense of [49].

Theorem 7.4.1. *Suppose that p is greater than the nilpotency class of the nilpotent Lie algebra $\text{Lie } U_{\nu_b}$, then \mathcal{I}_x is dense in the Newton stratum $\text{Sh}_{G,b}$.*

Proof. The isogeny class $\mathcal{I}_x \subseteq \text{Sh}_{G,b}(\overline{\mathbb{F}}_p)$ intersects every central leaf by the Rapoport–Zink uniformisation of isogeny classes (which follows from the main result of [49], see Section 1.4 of op. cit.). Thus if $W \subseteq \text{Sh}_{G,b}$ is a $G(\mathbb{A}_f^p)$ -stable closed subset containing $\mathcal{I}_x \subseteq \text{Sh}_{G,b}(\overline{\mathbb{F}}_p)$, then W intersects every central leaf C in a $G(\mathbb{A}_f^p)$ -stable *nonempty* closed subset W_C . Theorem 7.3.2 now tells us that $W_C = C$ and since $\text{Sh}_{G,b}$ is the (set-theoretic) union of all the central leaves it follows that $W = \text{Sh}_{G,b}$. \square

7.5. Orthogonal Shimura varieties. Conjecture 8.2 of [8] predicts that prime-to- p Hecke orbits are Zariski dense in certain Newton strata of the *orthogonal* Shimura varieties. These are the Shimura varieties for the group $\text{SO}(M)$ where M has signature $(2, m-2)$; they are of abelian type by Appendix B of [56].

In general one does *not* expect that prime-to- p Hecke orbits are Zariski dense in Newton strata, but when the Shimura datum is fully Hodge–Newton decomposable at p , then Newton strata are equal to central leaves. This is indeed the case for orthogonal Shimura varieties by Theorem D of [31] and Theorem E.(2) of [66].

It follows from the results of [51] and the proof of Proposition 6.2.4 of [39] that

$$\pi_0(\text{Sh}_{G,b}) \rightarrow \pi_0(\text{Sh}_G)$$

is a bijection for \mathbb{Q} -non-basic b for Shimura varieties of Hodge type. Since Newton strata behave well with respect to the devissage from Hodge type to abelian type, this results also hold for Shimura varieties of abelian type (see Section 2 of [67]). Thus Conjecture 8.2 of [8] comes down to showing

that the Zariski closure of Hecke orbits have the correct dimensions, and this can be reduced to the Hodge type case and then to Theorem 7.3.2. The assumption on the Coxeter number comes down to the assumption that $p \geq 2(m-1)$. In the required applications to K3 surfaces it suffices to assume that the number of slopes of the K3-crystal is less than or equal to p .

Remark 7.5.1. This line of reasoning shows that the Hecke orbit conjecture holds for fully Hodge–Newton decomposable Shimura varieties of abelian type, at primes of hyperspecial good reduction that satisfy the condition $p > n(\mathrm{Lie} U_{\nu_b})$, for central leaves in Newton strata corresponding to \mathbb{Q} -non-basic b . In fact these central leaves will be equal to Newton strata which makes it possible to reduce the discrete part of the Hecke orbit conjecture to the Hodge type case.

7.6. ℓ -power Hecke orbits. In this section we study the Zariski closures of ℓ -adic Hecke orbits of points for primes $\ell \neq p$. Since the ℓ -adic Hecke operators do not, generally, act transitively on $\pi_0(\mathrm{Sh}_G)$, all we can hope to prove is that ℓ -adic Hecke orbits are dense in a union of connected components of a central leaf. However, this cannot be true if $G^{\mathrm{sc}}/\mathbb{Q}_\ell$ is totally anisotropic because then there are no non-trivial ℓ -power Hecke operators. Let $C \subset \mathrm{Sh}_{G,b}$ be a central leaf as before.

Theorem 7.6.1. *Assume that Sh_G is proper, that $G^{\mathrm{sc}}/\mathbb{Q}_\ell$ is totally isotropic and that p is greater than the nilpotency class of the nilpotent Lie algebra $\mathrm{Lie} U_{\nu_b}$. Then any $G^{\mathrm{sc}}(\mathbb{Q}_\ell)$ -stable reduced closed subscheme $Z \subset C$ is a union of connected components of C*

Proof. The theorem follows at once from Theorem 7.3.2 and the Lemma 7.6.2 below. \square

Lemma 7.6.2. *If $Z \subseteq \mathrm{Sh}_{G,b}$ is irreducible and stable under the action of $G^{\mathrm{sc}}(\mathbb{Q}_\ell)$, then Z is stable under the action of $G^{\mathrm{sc}}(\mathbb{A}_f^p)$.*

Proof. Since the proof is almost exactly the same as the proof of Proposition 4.6 of [13], we will only give a sketch: A standard argument using the quasi-affineness of the Ekedahl–Oort stratification and the properness of Sh_G will show that the Zariski closure \overline{Z} of Z in Sh_G will contain a point $x \in \mathrm{Sh}_G(\overline{\mathbb{F}}_p)$ with a finite ℓ -power Hecke orbit. This Hecke orbit can be written as

$$I_x^{\mathrm{sc}}(\mathbb{Z}[\frac{1}{\ell}]) \backslash G^{\mathrm{sc}}(\mathbb{Q}_\ell) / U_\ell^{\mathrm{sc}},$$

where $I_x^{\mathrm{sc}}(\mathbb{Z}[\frac{1}{\ell}])$ is the intersection of the local stabiliser group with G^{sc} and U_ℓ^{sc} is the intersection of the level at ℓ with G^{sc} .

Claim 7.6.3. *The group I_x^{sc} is an inner form of G^{sc} and therefore x is contained in the basic locus of Sh_G .*

Proof. If I_x^{sc} is not an inner form of G^{sc} then there is a unipotent subgroup $N \subseteq G_{\mathbb{Q}_\ell}^{\mathrm{sc}}$ that intersects trivially with $I_{x, \mathbb{Q}_\ell}^{\mathrm{sc}}$. But then it is clear that

$$I_x^{\mathrm{sc}}(\mathbb{Z}[\frac{1}{\ell}]) \backslash G^{\mathrm{sc}}(\mathbb{Q}_\ell) / U_\ell^{\mathrm{sc}}$$

cannot be finite because the quotient of $N(\mathbb{Q}_\ell)$ by the compact open subgroup $N(\mathbb{Q}_\ell) \cap U_\ell^{\mathrm{sc}}$ is not finite. \square

The prime-to- p Hecke orbit of x has the form

$$I_x^{\mathrm{sc}}(\mathbb{Z}_{(p)}) \backslash G^{\mathrm{sc}}(\mathbb{A}_f^p) / U^{\mathrm{sc}, p}$$

and strong approximation away from ∞, ℓ tells us that

$$I_x^{\mathrm{sc}}(\mathbb{Z}_{(p)}) \subseteq G^{\mathrm{sc}}(\mathbb{A}_f^{p, \ell})$$

is dense, so that the prime-to- p Hecke orbit of x is equal to the ℓ -power Hecke orbit of x . Now the proof of Proposition 4.6 of [13] shows that the formal scheme $\overline{Z}^x \subseteq \mathrm{Sh}_G^x$ is stable under the action of the prime-to- p Hecke-operators. Moreover the same is true for \overline{Z}^y for any y in the prime-to- p Hecke orbit of x .

Now take a λ -adic Hecke operator $g_\lambda \in G^{\mathrm{sc}}(\mathbb{Q}_\lambda)$ for $\lambda \neq p, \ell$ and let W be the image of \overline{Z} under g_λ . Then every irreducible component W_i of W must contain a point y_i in the prime-to- p Hecke orbit of x , and since \overline{Z} is irreducible and contains y_i we know that W_i and \overline{Z} intersect in y_i . But then we get formal subschemes

$$W_i^{/y_i}, \overline{Z}^{/y_i} \subseteq \mathrm{Sh}_G^{/y_i}$$

where $W_i^{/y_i}$ is the Hecke orbit of $\overline{Z}^{/y_i}$ under g_λ . But we have shown that $\overline{Z}^{/y_i}$ is stable under prime-to- p Hecke orbits and so $W_i^{/y_i} \subseteq \overline{Z}^{/y_i}$ and for dimension reasons they must be equal. It follows that $W_i = \overline{Z}$ for all i and so \overline{Z} was already stable under the action of g_λ . We conclude that \overline{Z} is stable under the action of $G^{\mathrm{sc}}(\mathbb{A}_f^p)$.

We know that Z is the intersection of \overline{Z} with $\mathrm{Sh}_{G,b}$ and as the intersection of two $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -stable subschemes is itself $G^{\mathrm{sc}}(\mathbb{A}_f^p)$ -stable. \square

7.7. Further questions of Chai–Oort. Let $Z \subseteq C_\lambda$ be an irreducible smooth closed subvariety and let $x \in Z(\overline{\mathbb{F}}_p)$. We call Z *strongly Tate-linear at x* if $Z^z \subseteq C_\lambda^x$ is a strongly Tate-linear subvariety.

Question 7.7.1. Suppose that Z is strongly Tate-linear at some closed point $x_0 \in Z(\overline{\mathbb{F}}_p)$, is Z then strongly Tate-linear at all closed point $x \in Z(\overline{\mathbb{F}}_p)$?

It follows from Theorem II that the monodromy group of \mathcal{M} over $\mathrm{Spec} \widehat{\mathcal{O}}_{Z,x}$ does not depend on x . Now the validity of Conjecture 5.4.2 would imply that the question above has an affirmative answer.

Question 7.7.2. Suppose that Z is strongly Tate-linear at some closed point $x_0 \in Z(\overline{\mathbb{F}}_p)$, must Z then be an irreducible component of a central leaf in the mod p reduction of a Shimura variety of Hodge type?

The stronger assertion that Z must itself be an irreducible component of a Shimura variety of Hodge type is false in general, because only finitely many central leaves in a given Newton stratum contain the mod p reductions of special points by Theorem 1.3 of [50]. We note that if C_λ is the ordinary locus and Z is proper, then this stronger assertion is true by work of Moonen [57].

7.8. Results at ramified primes and parahoric level. The statement of the results make sense for central leaves in special fibres of the Kisin–Pappas [47] integral models of Shimura varieties of parahoric level at tamely ramified primes.

7.8.1. If the group is unramified and the level is parahoric, then the Hecke orbit conjecture for central leaves at parahoric level follows immediately from Theorem 7.3.2. The main observation is that the forgetful map

$$C' \rightarrow C,$$

where C' is a central leaf at Iwahori level and C is a central leaf at hyperspecial level, is $G(\mathbb{A}_f^p)$ -equivariant and induces a bijection on π_0 . This last statement can be proven using the surjectivity

of $C' \rightarrow C^8$ and the explicit description of connected components of Igusa varieties in [39, 51]. Rapoport–Zink uniformisation of isogeny classes at parahoric level⁸, implies as before that isogeny classes are dense in the Newton strata containing them.

7.8.2. If the group is ramified, then it is not always true that $G(\mathbb{A}_f^p)$ acts transitively on $\pi_0(\mathrm{Sh}_G)$ (see [59] for explicit counterexamples) and therefore it is not necessarily true that $G(\mathbb{A}_f^p)$ acts transitively on $\pi_0(C)$ either because $\pi_0(C) \rightarrow \pi_0(\mathrm{Sh}_G)$ is surjective. Nevertheless, we expect that the continuous part of the Hecke orbit conjecture is true for ramified groups and can be proven with more or less the same proofs. However one has to be careful for example with the proof of Corollary 2.2.1 of [38] because one does not know that EKOR strata are quasi-affine in this generality.

APPENDIX A. COMPLETE RESTRICTED PERFECTIONS, AFTER CHAI–OORT

In this section we will collect some results about completed perfection and complete restricted perfection of complete Noetherian local rings over a perfect field κ of characteristic p . For the latter part we will follow Chai–Oort [20].

A.1. Completed perfection. Throughout this section we will consider complete Noetherian local κ -algebras with maximal ideal \mathfrak{m} and residue field $R/\mathfrak{m} = \kappa$, the main example being the ring of power series $R = \kappa[[X_1, \dots, X_n]]$. We let $\varphi_{\mathrm{rel}} : R \rightarrow R$ denote the relative Frobenius homomorphism over κ .

Definition A.1.1. The (relative) perfection R^{perf} of R over κ is the (filtered) colimit

$$\varinjlim_{i \in \mathbb{Z}} R_i,$$

where $R_i = R$ and the transition maps $R_i \rightarrow R_{i+1}$ are given by φ_{rel} .

Example A.1.2. When $R = \kappa[[X_1, \dots, X_n]]$, we get the ring

$$\bigcup_m \kappa[[X_1^{1/p^m}, \dots, X_n^{1/p^m}]].$$

When $n = 1$ we note that this ring does not contain the element

$$\sum_{i=1}^{\infty} X^i \cdot X^{1/p^i},$$

while the partial sums do converge in the \mathfrak{m} -adic topology. In other words, R^{perf} is not \mathfrak{m} -adically complete.

Definition A.1.3. Let (R, \mathfrak{m}) be a complete Noetherian local κ -algebra with residue field κ . Then we define $R^{\mathrm{perf}, \wedge}$ to be the $I = \mathfrak{m} \cdot R^{\mathrm{perf}}$ -adic completion of R^{perf} .

Lemma A.1.4. *The ring $R^{\mathrm{perf}, \wedge}$ is an I -adically complete local κ -algebra with residue field κ .*

Proof. Lemma 05GG of [68] tells us that it is complete, because I is finitely generated. To show that it is a local ring it suffices to show that $R^{\mathrm{perf}}/I = R^{\mathrm{perf}, \wedge}/IR^{\mathrm{perf}, \wedge}$ is a local ring because every maximal ideal of $R^{\mathrm{perf}, \wedge}$ contains $IR^{\mathrm{perf}, \wedge}$ by Lemma 05GI of [68], and R^{perf}/I is a local ring because R^{perf} is. \square

⁸ This surjectivity is axiom 4c of the He–Rapoport axioms, see [66], and follows from Rapoport–Zink uniformisation at parahoric level, which is Theorem A of [37].

Lemma A.1.5. *Let (R, \mathfrak{m}) be a complete Noetherian local κ -algebra with residue field κ . Then there is a natural isomorphism of topological rings*

$$\varprojlim_{\varphi_{\text{rel}}} \left(R^{\text{perf}} / \mathfrak{m} R^{\text{perf}} \right) \simeq R^{\text{perf}, \wedge}.$$

In particular $R^{\text{perf}, \wedge}$ is perfect.

Proof. There is an isomorphism of inverse systems

$$\begin{array}{ccccc} \dots & \xrightarrow{\varphi_{\text{rel}}} & R^{\text{perf}} / \mathfrak{m} R^{\text{perf}} & \dots & \xrightarrow{\varphi_{\text{rel}}} & R^{\text{perf}} / \mathfrak{m} R^{\text{perf}} & \xrightarrow{\varphi_{\text{rel}}} & R^{\text{perf}} / \mathfrak{m} R^{\text{perf}} \\ & & g_2 \uparrow & & & g_1 \uparrow & & g_0 \uparrow \\ \dots & \longrightarrow & R^{\text{perf}} / \varphi_{\text{rel}}^2(\mathfrak{m}) R^{\text{perf}} & \dots & \longrightarrow & R^{\text{perf}} / \varphi_{\text{rel}}(\mathfrak{m}) R^{\text{perf}} & \longrightarrow & R^{\text{perf}} / \mathfrak{m} R^{\text{perf}}, \end{array}$$

where the transition maps on the bottom are the natural reduction maps. The maps g_i are constructed by precomposing the natural surjection $R^{\text{perf}} \rightarrow R^{\text{perf}} / \mathfrak{m} R^{\text{perf}}$ with $\varphi_{\text{rel}}^{-i}$ and noticing that this map is surjective with kernel $\varphi_{\text{rel}}^i(\mathfrak{m})$, giving the isomorphisms g_i .

Because the sequence of ideals $\varphi_{\text{rel}}^i(\mathfrak{m}) R^{\text{perf}}$ and $\mathfrak{m}^i R^{\text{perf}} = (\mathfrak{m} R^{\text{perf}})^i$ are cofinal, the inverse limit over the bottom inverse system is canonically isomorphic to $R^{\text{perf}, \wedge}$. \square

Lemma A.1.6 (c.f. Corollary 4.2.3 of [20]). *Let (R, \mathfrak{m}) be a complete Noetherian local κ -algebra with residue field κ . Then the natural map $R^{\text{perf}} \rightarrow R^{\text{perf}, \wedge}$ is injective.*

Proof. The lemma comes down to showing that the \mathfrak{m} -adic topology on R^{perf} is separated, in other words that

$$\bigcap_k (\mathfrak{m}^k R^{\text{perf}}) = (0).$$

For this we use the following result of Chai–Oort:

Claim A.1.7 (Proposition 4.2.2 of [20]). *Let R be a Noetherian local domain whose normalisation in its fraction field is finite over R . Then there exists a positive integer n_0 such that for all integers $a \geq 0$ and for all $n \geq a \cdot n_0$*

$$\{x \in R : x^a \in \mathfrak{m}^n\} \subseteq \mathfrak{m}^{\lfloor \frac{n}{a} - n_0 \rfloor}.$$

Now y be an element in $\bigcap_k (\mathfrak{m}^k R^{\text{perf}})$, and choose m such that $x = y^{p^m} \in R$. Then for all k we can write

$$y = \sum_{|I|=k} x_I r_I^{1/p^{n(k)+m}},$$

with $r_I \in R$. Here x_1, \dots, x_j are generators of \mathfrak{m} and for a multiset I of integers between 1 and j we write $x_I = \prod_{i \in I} x_i$. Then clearly

$$x^{p^{n(k)}} = y^{p^{n(k)+m}} = \sum_{|I|=k} (x_I)^{p^{n(k)+m}} r_I \in \mathfrak{m}^{k \cdot (p^{n(k)+m})} R.$$

By the claim for k large enough (larger than n_0 , say) we have

$$x \in \mathfrak{m}^{\lfloor k \cdot p^m - k \rfloor}.$$

Since this holds for all sufficiently large k , Krull's intersection theorem tells us that $x = 0$ and therefore $y = 0$. \square

A.1.8. If $\alpha : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ is a local homomorphism of complete local Noetherian κ -algebras, then there is an induced homomorphism

$$\alpha^{\text{perf}} : R^{\text{perf}} \rightarrow S^{\text{perf}}$$

by functoriality of the perfection construction and then similarly a morphism

$$\alpha^{\text{perf}, \wedge} : R^{\text{perf}, \wedge} \rightarrow S^{\text{perf}, \wedge}.$$

Lemma A.1.9 (c.f. Corollary 4.2.4 of [20]). *Suppose that S is a domain. Then if α is injective then so are α^{perf} and $\alpha^{\text{perf}, \wedge}$. In addition, $R^{\text{perf}} \subseteq S^{\text{perf}}$ has the subspace topology.*

Proof. Lemma A.1.6 tells us that we get a commutative diagram with injective horizontal maps

$$\begin{array}{ccccc} R & \hookrightarrow & R^{\text{perf}} & \hookrightarrow & R^{\text{perf}, \wedge} \\ \downarrow \alpha & & \downarrow \alpha^{\text{perf}} & & \downarrow \alpha^{\text{perf}, \wedge} \\ S & \hookrightarrow & S^{\text{perf}} & \hookrightarrow & S^{\text{perf}, \wedge}. \end{array}$$

The map α^{perf} is injective because S^{perf} is reduced and since elements in R^{perf} are $\overline{\mathbb{F}}_p$ -linear combinations of p^n -th roots of elements of R .

Because $R \rightarrow S$ is a local homomorphism we find that $\mathfrak{m}^k \subseteq \mathfrak{n}^k \cap R$ and Chevalley's theorem tells us that for each $k \geq 0$ there exists an $n(k)$ such that $\mathfrak{n}^{n(k)} \cap R \subseteq \mathfrak{m}^k$. Therefore the \mathfrak{m} -adic topology on R is equal to the subspace topology coming from S . If we can show that R^{perf} has the subspace topology coming from S^{perf} then Lemma 0ARZ of [68] tells us that the induced map on completions is injective.

In other words, we want to show that for each $k \gg 0$ there exists an $n'(k)$ such that

$$(\mathfrak{n}^{n'(k)} S^{\text{perf}}) \cap R^{\text{perf}} \subseteq \mathfrak{m}^k R^{\text{perf}}.$$

Now fix k and let $y \in (\mathfrak{n}^n S^{\text{perf}}) \cap R^{\text{perf}}$, then we are going to choose $n = n'(k)$ such that the above inclusion holds. If we choose m such that $x = y^{p^m} \in R$, then we can write

$$y = \sum_{|I|=n} x_I r_I^{1/p^{t+m}},$$

for some t and some $r_I \in S$. Here x_1, \dots, x_j is a basis of \mathfrak{n} and for a multiset I of integers between 1 and j we write $x_I = \prod_{i \in I} x_i$. Then clearly

$$x^{p^t} = y^{p^{t+m}} = \sum_{|I|=n} x_I^{p^{t+m}} r_I \in \mathfrak{n}^{n \cdot (p^{t+m})} S.$$

And so x^{p^t} is contained in

$$\mathfrak{n}^{n \cdot (p^{t+m})} \cap R.$$

If we can prove the following claim then the lemma follows:

Claim A.1.10. *For $k \gg 0$ there is a choice of $n = n'(k)$, independent of t and m , such that*

$$\mathfrak{n}^{n \cdot (p^{t+m})} \cap R \subseteq \mathfrak{m}^{(k+1) \cdot p^{t+m}}.$$

Indeed Claim A.1.7 says that if $k \gg 0$ we have

$$x \in \mathfrak{n}^{\lfloor (k+1) \cdot p^m - n_0 \rfloor}.$$

and so if we choose m such that $p^m \gg n_0$ then we find that y is in

$$\mathfrak{m}^k R^{\text{perf}}.$$

□

Proof of Claim A.1.10. We can certainly choose n such that

$$\mathfrak{n}^n \cap R \subseteq \mathfrak{m}^{k+1}$$

and then

$$\mathfrak{n}^{n \cdot (p^{t+m})} \subseteq \varphi_{\text{rel}}^{t+m-c}(\mathfrak{n}^n)$$

where c_0 is a constant only depending on the characteristic p and the number of generators of \mathfrak{n} . Therefore

$$\begin{aligned} (\mathfrak{n}^{n \cdot (p^{t+m})} \cap R) &\subseteq \varphi_{\text{rel}}^{t+m-c_0}(\mathfrak{n}^n) \cap R \\ &\subseteq \varphi_{\text{rel}}^{t+m-c_1}(\mathfrak{m}^{k+1}) \\ &\subseteq \mathfrak{m}^{(k+1) \cdot p^{t+m-c_2}} \end{aligned}$$

which is contained in $\mathfrak{m}^{k \cdot p^{t+m}}$ for $k \gg 0$. The constant c_1 only depends on c_0 and $R \rightarrow S$ (in particular on the biggest j for which there exists an $x \in \mathfrak{m}$ with $x^{1/p^j} \in S$). □

A.1.11. Let (R, \mathfrak{m}) and (S, \mathfrak{n}) be complete Noetherian local κ -algebras with residue field κ , then their completed tensor product

$$T = R \widehat{\otimes}_{\kappa} S$$

is again a complete Noetherian local κ -algebra with residue field k .

Lemma A.1.12. *There is a natural continuous isomorphism*

$$T^{\text{perf}, \wedge} \rightarrow R^{\text{perf}, \wedge} \widehat{\otimes}_k S^{\text{perf}, \wedge}.$$

Proof. It suffices to show that there is a natural isomorphism of functors

$$\text{Spf } T^{\text{perf}, \wedge} = \text{Spf } R^{\text{perf}, \wedge} \times \text{Spf } S^{\text{perf}, \wedge},$$

which is a straightforward consequence of Lemma A.1.17 below. Indeed, limits commute with products in categories of sheaves and so the perfection of the product of two formal schemes is naturally isomorphic to the product of the perfections. □

A.1.13. If $R = \kappa[[X_1, \dots, X_n]]$ then

$$R^{\text{perf}} = \bigcup_m \kappa[[X_1^{1/p^m}, \dots, X_n^{1/p^m}]].$$

and $R^{\text{perf}, \wedge}$ is its completion, which we will now describe explicitly. Let S be the ring of formal expressions

$$\sum_{I \in (\mathbb{Z}_{\geq 0}[1/p])^n} a_I x_I$$

with $a_I \in \kappa$ and such that for each $N \in \mathbb{Z}_{\geq 0}$ the set

$$\{I \in (\mathbb{Z}_{\geq 0}[1/p])^n : a_I \neq 0\} \cap [0, N]^n$$

is finite (thus for example the element $x_1^{1/p} + x_1^{1/p^2} + x_1^{1/p^3} + \dots$ is not contained in S). The ring structure on S is given by the usual ring structure on power series, which makes sense because of

the finiteness condition. It is easy to see that S is an integral domain for example by looking at leading coefficients. It is also clear that S is a perfect κ -algebra, because we are allowed to divide by p in the exponents and this does not change the finiteness condition.

There is a natural injective ring homomorphism $R \rightarrow S$ sending X_i to x_i and since S is perfect this extends to an injective ring homomorphism $R^{\text{perf}} \rightarrow S$.

Lemma A.1.14. *The ring S is I -adically complete, where $I = (x_1, \dots, x_n)$.*

Proof. If we are given a sequence f_1, f_2, \dots of elements in S such that f_{k+m} is congruent to f_k mod I^k for all k, m , then the coefficients of the f_i form a stabilising sequence which leads to an element $f \in S$ which is congruent to f_k mod I^k for all k . To be precise the element f_k determines the coefficients $a_{f,i}$ for $i \in [0, k]^n$ and so we get a well defined limit f . \square

The natural map $R^{\text{perf}} \rightarrow S$ clearly induces isomorphisms

$$S/I^k S \simeq R^{\text{perf}}/I^k R^{\text{perf}}$$

for all k and therefore $R^{\text{perf}} \rightarrow S$ can be identified with the natural map $R^{\text{perf}} \rightarrow R^{\text{perf}, \wedge}$.

Corollary A.1.15. *When $R = \kappa[[X_1, \dots, X_n]]$, the completed perfection $R^{\text{perf}, \wedge}$ is an integral domain.*

We do not know whether $R^{\text{perf}, \wedge}$ is always an integral domain when R is.

A.1.16. Recall our conventions on formal schemes from Section 2.1: Let R be a κ -algebra that is complete with respect to a finitely generated ideal $I \subseteq R$. We write $\text{Spf } R$ for functor

$$\varinjlim_j \text{Spec } R/I^j R$$

on the category of κ -algebras, where the transition maps are the ones induced by the natural quotient maps $R/I^{j+1} \rightarrow R/I^j$.

Lemma A.1.17. *Let (R, \mathfrak{m}) be a complete Noetherian local κ -algebra with residue field κ , and consider the inverse system*

$$\dots \longrightarrow \text{Spf } R \xrightarrow{\varphi_{\text{rel}}} \text{Spf } R \longrightarrow \dots$$

Then the inverse limit of this diagram (in the category of functors) is isomorphic to $\text{Spf } R^{\text{perf}, \wedge}$.

Proof. Let us write $\text{Spf } R = \varinjlim \text{Spec } R/\varphi_{\text{rel}}^n(\mathfrak{m})$, then the following diagram is Cartesian

$$\begin{array}{ccc} \text{Spec } R/\varphi_{\text{rel}}^{n+1}(\mathfrak{m}) & \hookrightarrow & \text{Spf } R \\ \downarrow & & \downarrow \varphi \\ \text{Spec } R/\varphi_{\text{rel}}^n(\mathfrak{m}) & \hookrightarrow & \text{Spf } R, \end{array}$$

and the left vertical map corresponds to the map

$$R/\varphi_{\text{rel}}^n(\mathfrak{m}) \rightarrow R/\varphi_{\text{rel}}^{n+1}(\mathfrak{m})$$

induced by φ , which is injective if R is reduced. The diagram

$$\varinjlim_m R/\varphi_{\text{rel}}^{n+m}(\mathfrak{m})$$

where the transition maps are the maps induced by φ can be identified with the diagram

$$\varinjlim_m R_m / \varphi_{\text{rel}}^n(\mathfrak{m}) R_m$$

where R is the ring R equipped with an R -algebra structure via the map $\varphi_{\text{rel}}^m : R \rightarrow R$. It is clear that the limit of this diagram of R -algebras is given by $R^{\text{perf}} / \varphi_{\text{rel}}^n(\mathfrak{m}) R^{\text{perf}}$.

So if \mathcal{F} is the limit of the diagram in the statement of the Lemma, then there is a Cartesian diagram

$$\begin{array}{ccc} \text{Spec } R^{\text{perf}} / \varphi_{\text{rel}}^n(\mathfrak{m} R^{\text{perf}}) & \hookrightarrow & \mathcal{F} \\ \downarrow & & \downarrow \\ \text{Spec } R / \varphi^n(\mathfrak{m}) & \hookrightarrow & \text{Spf } R. \end{array}$$

It follows from this (c.f. Lemma 0AJH of [68]) that

$$\mathcal{F} \simeq \varinjlim_n \text{Spec } R^{\text{perf}} / \varphi_{\text{rel}}^n(\mathfrak{m} R^{\text{perf}})$$

which proves the lemma. \square

A.2. Complete restricted perfections. In this section we follow Section 3.5 of [20]. Let (R, \mathfrak{m}) be a complete Noetherian local κ -algebra with residue field κ . Let r, s, n_0 be natural numbers with $0 < r < s$ and $n_0 \geq 0$ and consider the following subset of $R^{\text{perf}, \wedge}$:

$$R_{s,r,\geq n_0}^{\text{perf}, \wedge} := \left\{ x \in R^{\text{perf}, \wedge} \mid \text{For } n \geq n_0 \text{ there is an } y \in R \text{ such that } x^{p^{nr}} \equiv y \pmod{\varphi^{ns}(\mathfrak{m}) R^{\text{perf}, \wedge}} \right\}$$

This is clearly a subring because the image of

$$R / \varphi_{\text{rel}}^{ns}(\mathfrak{m}) \rightarrow R^{\text{perf}, \wedge} / \varphi_{\text{rel}}^{ns}(\mathfrak{m}) R^{\text{perf}, \wedge}$$

is a subring. It is moreover closed in the \mathfrak{m} -adic topology on $R^{\text{perf}, \wedge}$ because it is defined by congruence conditions. Furthermore if $R \rightarrow S$ is a local homomorphism then there is clearly an induced ring homomorphism

$$R_{s,r,\geq n_0}^{\text{perf}, \wedge} \rightarrow S_{s,r,\geq n_0}^{\text{perf}}$$

which is injective if $R \rightarrow S$ is injective and S is a domain, by Lemma A.1.9. These rings are called complete restricted perfections because they are complete and sit between R and $R^{\text{perf}, \wedge}$, hence the *restricted* perfection.

A.2.1. If $R = \kappa[[X_1, \dots, X_n]]$ then

$$R_{s,r,\geq n_0}^{\text{perf}, \wedge} \subseteq R^{\text{perf}, \wedge}$$

can be described as a subring of S as follows. It consists of those formal series

$$\sum_{I \in (\mathbb{Z}_{\geq 0}[1/p])^n} a_I x_I$$

such that for all $n \geq n_0$ the truncated series

$$\sum_{\substack{I \in (\mathbb{Z}_{\geq 0}[1/p])^n \\ |p^{nr} I|_{\infty} \leq p^{ns}}} a_I x_{p^{nr} I}$$

is an actual power series. Here the norm $|\cdot|_\infty$ on \mathbb{R}^n is the maximum of the norms of the coordinates. An elementary argument shows that this condition is equivalent to the condition that

$$-\text{ord}_p(I) \leq \max \left\{ n_0, s \cdot \left\lfloor \frac{\log_p |I|_\infty}{s-r} \right\rfloor + 1 \right\},$$

where $\text{ord}_p(I)$ is the minimum of the p -adic valuations of the coordinates.

A.2.2. Chai and Oort also define a variant

$$\kappa[[X_1, \dots, X_n]]_{E,C;d}^{\text{perf},\wedge} \subseteq \kappa[[X_1, \dots, X_n]]^{\text{perf},\wedge}$$

for real numbers $C > 0, d \geq 0, E > 0$ consisting of those series

$$\sum_{I \in (\mathbb{Z}_{\geq 0}[1/p])^n} a_I x_I$$

satisfying

$$p^{-\text{ord}_p(I)} \leq \max(C \cdot (|I|_\infty + D)^E, 1).$$

Remark A.2.3. Chai and Oort use the notation

$$\kappa\langle\langle X_1^{p^{-\infty}}, \dots, X_n^{p^{-\infty}} \rangle\rangle_{C;d}^{E,\sharp}$$

for these rings; the superscript \sharp is used to distinguish it from a \flat -variant

$$\kappa\langle\langle X_1^{p^{-\infty}}, \dots, X_n^{p^{-\infty}} \rangle\rangle_{C;d}^{E,\flat} \supseteq \kappa\langle\langle X_1^{p^{-\infty}}, \dots, X_n^{p^{-\infty}} \rangle\rangle_{C;d}^{E,\sharp}.$$

We have opted not to use this notation because it could cause confusion with the \sharp and \flat notation introduced by Scholze to denote tilts and untilts of (semi-)perfect rings.

Chai and Oort prove (Lemma 4.12 of [20]) that for each r, s, n_0 there are choices of C, d, E such that

$$\kappa[[X_1, \dots, X_n]]_{s,r;\geq n_0}^{\text{perf}} \subseteq \kappa[[X_1, \dots, X_n]]_{E,C;d}^{\text{perf},\wedge}.$$

These rings are then used to state Proposition of 6.4 of [20], where we note that composition of perfected power series is well defined.

Theorem A.2.4 (Proposition 6.4 of [20]). *Let $f(\underline{u}, \underline{v}) \in \kappa[[u_1, \dots, u_a, v_1, \dots, v_b]]$ be a formal power series. Let $(g_1(\underline{x}), \dots, g_a(\underline{x}))$ be an a -tuple of elements in*

$$\kappa[[X_1, \dots, X_n]]_{E,C;d}^{\text{perf},\wedge},$$

whose degree zero terms vanish, and similarly let $(h_1(\underline{y}), \dots, h_b(\underline{y}))$ be an b -tuple of elements in

$$\kappa[[Y_1, \dots, Y_n]]_{E,C;d}^{\text{perf},\wedge},$$

whose degree zero terms vanish. Suppose that there is a sequence of nonnegative integers d_n for $n \in \mathbb{Z}_{\geq 0}$ such that

$$\lim_{n \rightarrow \infty} \frac{q^n}{d_n} = 0$$

where $q = p^r$ is some power of p and such that for all n the element

$$f(g_1(\underline{X}), \dots, g_a(\underline{X}), h_1(\underline{X})^{q^n}, \dots, h_b(\underline{X})^{q^n})$$

is congruent to zero modulo $(X_1, \dots, X_n)^{d_n}$. Then

$$f(g_1(\underline{X}), \dots, g_a(\underline{X}), h_1(\underline{Y}), \dots, h_b(\underline{Y})) = 0.$$

Proof. This is a special case of Proposition 6.4 of [20] (c.f. the remark after its statement); strictly speaking this proposition is stated for the \flat -versions of the complete restricted perfection of Chai–Oort, but those contain the \sharp -versions (see A.2.3). \square

Remark A.2.5. Let us recall from Section 6.1 of [20] some intuition behind Theorem A.2.4 in the case that the g_i and h_i are ordinary power series. Define

$$U = \mathrm{Spf} \kappa[[u_1, \dots, u_a]], \quad V = \mathrm{Spf} \kappa[[v_1, \dots, v_b]], \quad W = \mathrm{Spf} \kappa[[x_1, \dots, x_n]].$$

Then $f(u, v)$ corresponds to a global function f on $U \times V$, and $(g_1, \dots, g_a), (h_1, \dots, h_b)$ correspond to morphisms

$$\begin{aligned} g &: W \rightarrow U \\ h &: W \rightarrow V. \end{aligned}$$

The theorem can then be interpreted as saying that $(g, h)^*(f)$ is trivial if and only if for all n the global function

$$(1, \varphi_{\mathrm{rel}}^n)^* \circ (g, h)^*(f)$$

is trivial mod $(X_1, \dots, X_n)^{d_n}$ for a sequence of integers d_n satisfying

$$\lim_{n \rightarrow \infty} \frac{q^n}{d_n} = 0.$$

REFERENCES

- [1] Johannes Anshütz and Arthur-César Le Bras, *Prismatic Dieudonné theory*, arXiv e-prints (July 2019), available at 1907.10525.
- [2] Pierre Berthelot and William Messing, *Théorie de Dieudonné cristalline. I*, Journées de Géométrie Algébrique de Rennes (Rennes, 1978), Vol. I, 1979, pp. 17–37. MR563458
- [3] Bhargav Bhatt and Jacob Lurie, *Absolute prismatic cohomology*, arXiv e-prints (January 2022), available at 2201.06120.
- [4] ———, *The prismatization of p -adic formal schemes*, arXiv e-prints (January 2022), available at 2201.06124.
- [5] Bhargav Bhatt, Matthew Morrow, and Peter Scholze, *Topological Hochschild homology and integral p -adic Hodge theory*, Publ. Math. Inst. Hautes Études Sci. **129** (2019), 199–310. MR3949030
- [6] Bhargav Bhatt and Peter Scholze, *Prismatic F -crystals and crystalline Galois representations*, arXiv e-prints (June 2021), available at 2106.14735.
- [7] Nicolas Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley. MR1890629
- [8] Daniel Bragg and Ziquan Yang, *Twisted Derived Equivalences and Isogenies between $K3$ Surfaces in Positive Characteristic*, arXiv e-prints (February 2021), available at 2102.01193.
- [9] Ana Caraiani and Peter Scholze, *On the generic part of the cohomology of compact unitary Shimura varieties*, Ann. of Math. (2) **186** (2017), no. 3, 649–766. MR3702677
- [10] Ching-Li Chai, *Every ordinary symplectic isogeny class in positive characteristic is dense in the moduli*, Invent. Math. **121** (1995), no. 3, 439–479. MR1353306
- [11] ———, *Families of ordinary abelian varieties: canonical coordinates, p -adic monodromy, Tate-linear subvarieties and hecke orbits*, 2003. preprint available at https://www2.math.upenn.edu/~chai/papers_pdf/fam_ord_av.pdf.
- [12] ———, *Canonical coordinates on leaves of p -divisible groups: The two-slope case* (2005). Available at https://www2.math.upenn.edu/~chai/papers_pdf/2slope_v14.pdf.
- [13] ———, *Monodromy of Hecke-invariant subvarieties*, Pure Appl. Math. Q. **1** (2005), no. 2, Special Issue: In memory of Armand Borel. Part 1, 291–303. MR2194726
- [14] ———, *Hecke orbits as Shimura varieties in positive characteristic*, International Congress of Mathematicians. Vol. II, 2006, pp. 295–312. MR2275599
- [15] ———, *A rigidity result for p -divisible formal groups*, Asian J. Math. **12** (2008), no. 2, 193–202. MR2439259
- [16] ———, *Sustained p -divisible groups and a foliation on moduli spaces of abelian varieties*, Proceedings of the International Consortium of Chinese Mathematicians 2018, [2020] ©2020, pp. 3–30. MR4251135
- [17] Ching-Li Chai and Frans Oort, *Hecke orbits*. book project in preparation.
- [18] ———, *Moduli of abelian varieties and p -divisible groups*, Arithmetic geometry, 2009, pp. 441–536. MR2498069

- [19] ———, *Monodromy and irreducibility of leaves*, Ann. of Math. (2) **173** (2011), no. 3, 1359–1396. MR2800716
- [20] ———, *Rigidity for biextensions of formal groups* (2017). Available at https://www2.math.upenn.edu/~chai/papers_pdf/rig_biext_v4d.pdf.
- [21] Richard Crew, *F-isocrystals and their monodromy groups*, Ann. Sci. École Norm. Sup. (4) **25** (1992), no. 4, 429–464. MR1186910
- [22] Marco D'Addezio, *The monodromy groups of lisse sheaves and overconvergent F-isocrystals*, Selecta Math. (N.S.) **26** (2020), no. 3.
- [23] Marco D'Addezio, *Parabolicity conjecture of F-isocrystals*, arXiv e-prints (December 2020), available at 2012.12879.
- [24] Aise Johan de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*, Inst. Hautes Études Sci. Publ. Math. **82** (1995), 5–96 (1996). MR1383213
- [25] P. Deligne and J. S. Milne, *Tannakian categories*, Hodge cycles, motives, and shimura varieties, 1982, pp. 101–228.
- [26] Jean Dieudonné and Alexander Grothendieck, *Éléments de géométrie algébrique: I. Le langage des schémas*, Inst. Hautes Études Sci. Publ. Math. **4** (1961).
- [27] Vladimir Drinfeld, *A stacky approach to crystals*, arXiv e-prints (October 2018), arXiv:1810.11853, available at 1810.11853.
- [28] Vladimir Drinfeld and Kiran Kedlaya, *Slopes of indecomposable F-isocrystals*, Pure Appl. Math. Quart. **13** (2017), 131–192.
- [29] Marco D'Addezio and Hélène Esnault, *On the Universal Extensions in Tannakian Categories*, International Mathematics Research Notices (202105), available at <https://academic.oup.com/imrn/advance-article-pdf/doi/10.1093/imrn/rnab107/38377494/rnab107.pdf>.
- [30] Laurent Fargues and Peter Scholze, *Geometrization of the local Langlands correspondence*, arXiv e-prints (February 2021), available at 2102.13459.
- [31] Ulrich Görtz, Xuhua He, and Sian Nie, *Fully Hodge–Newton Decomposable Shimura Varieties*, Peking Math. J. **2** (2019), no. 2, 99–154. MR4060001
- [32] Paul Hamacher, *The product structure of Newton strata in the good reduction of Shimura varieties of Hodge type*, J. Algebraic Geom. **28** (2019), no. 4, 721–749. MR3994311
- [33] Paul Hamacher and Wansu Kim, *l-adic étale cohomology of Shimura varieties of Hodge type with non-trivial coefficients*, Math. Ann. **375** (2019), no. 3-4, 973–1044. MR4023369
- [34] S. Mohammad Hadi Hedayatzadeh, *Exterior powers of π -divisible modules over fields*, J. Number Theory **138** (2014), 119–174. MR3168925
- [35] Gerhard P. Hochschild, *Basic theory of algebraic groups and Lie algebras*, Graduate Texts in Mathematics, vol. 75, Springer-Verlag, New York-Berlin, 1981. MR620024
- [36] Harald Hofstätter, *Denominators of coefficients of the Baker–Campbell–Hausdorff series*, arXiv e-prints (October 2020), available at 2010.03440.
- [37] Pol van Hoftten, *Mod p points on Shimura varieties of parahoric level (with an appendix by Rong Zhou)*, arXiv e-prints (October 2020), available at 2010.10496.
- [38] ———, *On the ordinary Hecke orbit conjecture*, arXiv e-prints (December 2021), available at 2112.12422.
- [39] Pol van Hoftten and Luciena Xiao Xiao, *Monodromy and Irreducibility of Igusa Varieties*, arXiv e-prints (February 2021), available at 2102.09870.
- [40] Serin Hong, *On the Hodge–Newton filtration for p -divisible groups of Hodge type*, Math. Z. **291** (2019), no. 1-2, 473–497. MR3936079
- [41] Luc Illusie, *Complexe cotangent et déformations. II*, Lecture Notes in Mathematics, Vol. 283, Springer-Verlag, Berlin-New York, 1972. MR0491681
- [42] ———, *Déformations de groupes de Barsotti–Tate (d'après A. Grothendieck)*, 1985, pp. 151–198. Seminar on arithmetic bundles: the Mordell conjecture (Paris, 1983/84). MR801922
- [43] Guy Kapon and Lior Hadassy, *Characterising Semi-Simple Lie Algebras by Their Borel Nilpotent Radical*, arXiv e-prints (November 2021), available at 2111.03323.
- [44] Nicholas M. Katz, *Slope filtration of F-crystals*, Journées de Géométrie Algébrique de Rennes (Rennes, 1978), Vol. I, 1979, pp. 113–163. MR563463
- [45] Wansu Kim, *On central leaves of Hodge-type Shimura varieties with parahoric level structure*, Math. Z. **291** (2019), no. 1-2, 329–363. MR3936073
- [46] Wansu Kim and Keerthi Madapusi Pera, *2-adic integral canonical models*, Forum Math. Sigma **4** (2016), Paper No. e28, 34. MR3569319
- [47] M. Kisin and G. Pappas, *Integral models of Shimura varieties with parahoric level structure*, Publ. Math. Inst. Hautes Études Sci. **128** (2018), 121–218. MR3905466
- [48] Mark Kisin, *Integral models for Shimura varieties of abelian type*, J. Amer. Math. Soc. **23** (2010), no. 4, 967–1012. MR2669706

- [49] ———, *mod p points on Shimura varieties of abelian type*, J. Amer. Math. Soc. **30** (2017), no. 3, 819–914. MR3630089
- [50] Mark Kisin, Yeuk Hay Joshua Lam, Ananth N. Shankar, and Padmavathi Srinivasan, *Finiteness of reductions of Hecke orbits*, arXiv e-prints (September 2021), available at 2109.05147.
- [51] Arno Kret and Sug Woo Shin, *H^0 of Igusa varieties via automorphic forms*, arXiv e-prints (February 2021), available at 2102.10690.
- [52] Elena Mantovan, *On the cohomology of certain PEL-type Shimura varieties*, Duke Math. J. **129** (2005), no. 3, 573–610. MR2169874
- [53] Akhil Mathew, *Faithfully flat descent of almost perfect complexes in rigid geometry*, arXiv e-prints (December 2019), available at 1912.10968.
- [54] Hideyuki Matsumura, *Commutative algebra*, W. A. Benjamin, Inc., New York, 1970. MR0266911
- [55] Davesh Maulik, Ananth N. Shankar, and Yunqing Tang, *Picard ranks of K3 surfaces over function fields and the Hecke orbit conjecture*, arXiv e-prints (November 2020), available at 2011.08887.
- [56] J. S. Milne, *Introduction to Shimura varieties*, Harmonic analysis, the trace formula, and Shimura varieties, 2005, pp. 265–378. MR2192012
- [57] Ben Moonen, *Linearity properties of Shimura varieties. II*, Compositio Math. **114** (1998), no. 1, 3–35. MR1648527
- [58] ———, *Serre-Tate theory for moduli spaces of PEL type*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 2, 223–269. MR2061781
- [59] Yasuhiro Oki, *On the connected components of Shimura varieties for CM unitary groups in odd variables*, arXiv e-prints (April 2021), available at 2104.13086.
- [60] Frans Oort, *Foliations in moduli spaces of abelian varieties*, J. Amer. Math. Soc. **17** (2004), no. 2, 267–296. MR2051612
- [61] ———, *Appendix 1: Some questions in algebraic geometry*, Open problems in arithmetic algebraic geometry, [2019] ©2019, pp. 263–283. Reprint of 1995 original. MR3971187
- [62] M. Rapoport and M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, Compositio Math. **103** (1996), no. 2, 153–181. MR1411570
- [63] Peter Scholze and Jared Weinstein, *Moduli of p -divisible groups*, Camb. J. Math. **1** (2013), no. 2, 145–237. MR3272049
- [64] Ananth N. Shankar, *The Hecke-orbit conjecture for “modèles étranges”*, 2016. preprint.
- [65] Ananth N. Shankar and Rong Zhou, *Serre-Tate theory for Shimura varieties of Hodge type*, Math. Z. **297** (2021), no. 3-4, 1249–1271. MR4229601
- [66] Xu Shen, Chia-Fu Yu, and Chao Zhang, *Ekor strata for Shimura varieties with parahoric level structure*, Duke Math. J. (2021). To appear, preprint available at arxiv.org/abs/1910.07785.
- [67] Xu Shen and Chao Zhang, *Stratifications in good reductions of Shimura varieties of abelian type*, arXiv e-prints (July 2017), available at 1707.00439.
- [68] The Stacks project authors, *The stacks project*, 2022. <https://stacks.math.columbia.edu>.
- [69] Torsten Wedhorn, *On Tannakian duality over valuation rings*, J. Algebra **282** (2004), no. 2, 575–609. MR2101076
- [70] Luciena Xiao Xiao, *On The Hecke Orbit Conjecture for PEL Type Shimura Varieties*, arXiv e-prints (June 2020), available at 2006.06859.
- [71] Yujie Xu, *Normalization in integral models of Shimura varieties of Hodge type*, arXiv e-prints (July 2020), available at 2007.01275.
- [72] Rong Zhou, *Motivic cohomology of quaternionic Shimura varieties and level raising*, arXiv e-prints (January 2019), available at 1901.01954.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, SU - 4 PLACE JUSSIEU, CASE 247, 75005 PARIS

Email address: daddezio@imj-prg.fr

STANFORD MATHEMATICS DEPARTMENT, 450 JANE STANFORD WAY, BUILDING 380, STANFORD, CA 94305, USA

Email address: pol.van.hoften@stanford.edu