

ABUNDANCE THEOREM FOR MINIMAL COMPACT KÄHLER MANIFOLDS WITH VANISHING SECOND CHERN CLASS

MASATAKA IWAI AND SHIN-ICHI MATSUMURA

ABSTRACT. In this paper, for compact Kähler manifolds with nef cotangent bundle, we study the abundance conjecture and the associated Iitaka fibrations. We show that, for a minimal compact Kähler manifold, the second Chern class vanishes if and only if the cotangent bundle is nef and the canonical bundle has the numerical dimension 0 or 1. Additionally, in this case, we prove that the canonical bundle is semi-ample. Furthermore, we give a relation between the variation of the fibers of the Iitaka fibration and a certain semipositivity of the cotangent bundle.

CONTENTS

1. Introduction	2
Acknowledgments	4
2. Preliminaries	5
2.1. Notation	5
2.2. Positivity, flatness, and slope of torsion-free sheaves.	5
3. Abundance Theorem and Structure Theorem	8
3.1. Abundance theorem	8
3.2. Structure theorem	10
4. On Variation of the Fibers of Iitaka Fibrations	11
4.1. On the Fujita decomposition of nef cotangent bundles	11
4.2. Hermitian flatness of nef cotangent bundles	13
5. Generically Nef Vector Bundles with Vanishing Second Chern Class	18
5.1. Case of smooth projective varieties	18
5.2. Case of compact Kähler manifolds	21
6. Appendix	22
6.1. Proof of Theorem 5.7	22
6.2. Related open problems	27
References	27

Date: May 24, 2022, version 0.01.

2020 Mathematics Subject Classification. Primary 14E30, Secondary 14D06, 32M25, 32Q26.

Key words and phrases. Abundance conjecture, Projectively numerically flat vector bundles, Hermitian flat vector bundles, Generically ample vector bundles, Slope stabilities, Fujita decomposition,

1. INTRODUCTION

The following conjecture, the so-called *abundance conjecture*, is one of the most important open problems in algebraic geometry. The abundance conjecture has been solved for projective varieties of dimension at most 3 (see [Miy87, Miy88a, Miy88b, Kaw92]). However, for higher-dimensional varieties, it remains generally unsolved, and even partial results are scarce (see [DHP13, GM17, LP18] for some approaches).

Conjecture 1.1 (Abundance conjecture for smooth projective varieties). *Let X be a smooth projective variety. If the canonical bundle K_X is nef, then K_X is semi-ample.*

In this paper, we study the abundance conjecture and the associated Iitaka fibration for a smooth projective variety X with the nef cotangent bundle Ω_X , motivated by the following problems.

- (a) To solve the abundance conjecture for the smooth projective variety X with the nef cotangent bundle Ω_X .
- (b) To find a relation between the Fujita decomposition of Ω_X and the Iitaka fibration $f : X \rightarrow Y$ of X . Here, the Fujita decomposition decomposes nef vector bundles into numerically flat vector bundles and generically ample vector bundles (see Proposition 4.2 for the precise statement).
- (c) To understand a mechanism by which the variation of the fibers of the Iitaka fibration $f : X \rightarrow Y$ behaves differently depending on a certain semipositivity of curvatures of Ω_X .

The first main result of this paper is to solve the abundance conjecture for a compact Kähler manifold X such that Ω_X is nef and $\nu(K_X)$ is 0 or 1, equivalently, the second Chern class $c_2(X)$ vanishes (see the theorems below). Here $\nu(K_X)$ denotes the numerical dimension of the canonical bundle K_X . Additionally, in this case, we reveal a detailed structure of the Iitaka fibration $f : X \rightarrow Y$ (see Theorem 3.5, cf. [Hör13]). We emphasize that Theorem 1.3 is valid not only for smooth projective varieties but also compact Kähler manifolds.

Theorem 1.2. *Let X be a compact Kähler manifold with the nef canonical bundle K_X . Then, the following conditions are equivalent:*

- (1) *The second Chern class $c_2(X)$ vanishes in $H^{2,2}(X, \mathbb{R})$.*
- (2) *Ω_X is nef, and $\nu(K_X)$ is 0 or 1.*

Theorem 1.3. *Let X be a compact Kähler manifold with the nef cotangent bundle Ω_X . If $\nu(K_X) = 0$ or $\nu(K_X) = 1$, then the canonical bundle K_X is semi-ample.*

Let X be a smooth projective variety such that K_X is nef and $\nu(K_X) = 1$. Theorem 1.3 should be compared to the result of [LP18], which states that $\kappa(K_X) \geq 0$ if $\chi(\mathcal{O}_X) \neq 0$. Here $\chi(\mathcal{O}_X)$ is the holomorphic Euler characteristic. In contrast to the assumption of this result, the Euler characteristic $\chi(\mathcal{O}_X)$ always vanishes when Ω_X is nef and $\nu(K_X) < \dim X$. Further, when X is a surface, our assumption $c_2(X) = 0$ is equivalent to $\chi(\mathcal{O}_X) = 0$. In this sense, Theorem 1.3 partially complements the result of [LP18].

Wu-Zheng and Liu in [WZ02, Liu14] proved that K_X is semi-ample under the assumption that X admits a Kähler metric with nonpositive holomorphic bisectional curvature. This assumption is much stronger than the condition of Ω_X being nef.

In fact, under this assumption, the Iitaka fibration $f : X \rightarrow Y$ is locally trivial and actually gives a product structure $X \cong F \times Y$ with the fiber F , up to finite étale covers of X . However, when Ω_X is merely nef, the Iitaka fibration $f : X \rightarrow Y$ (even if it exists) is not necessarily locally trivial (see [Hör13]). This phenomenon constitutes a difference from the nonpositive holomorphic bisectional curvature and presents difficulty in accommodating nef cotangent bundles. We focus on the Fujita decomposition to understand the reason why such a phenomenon occurs.

The second main result gives a characterization for the Iitaka fibration $f : X \rightarrow Y$ to be locally trivial in terms of the Fujita decomposition and Hermitian flatness (see Theorem 1.4).

Let us explain a detailed motivation for this result. In the case where X has nonpositive holomorphic bisectional curvature, the Ricci kernel foliation \mathcal{R} can be constructed, and further \mathcal{R} determines the Iitaka fibration $f : X \rightarrow Y$, which is the core of the arguments in [WZ02, Liu14]. The Ricci kernel foliation \mathcal{R} is a certain “flat part” of the cotangent bundle Ω_X . Then, we can show that

$$\mathcal{R} = \mathcal{F} = \Omega_{X/Y},$$

where \mathcal{F} is the flat part of the Fujita decomposition of Ω_X and $\Omega_{X/Y}$ is the relative cotangent bundle of the Iitaka fibration $f : X \rightarrow Y$. Unfortunately, when only the nefness of Ω_X is assumed, the Ricci kernel foliation is no longer constructed. Nevertheless, the Fujita decomposition always exists for the nef cotangent bundle Ω_X . Problem (b) asks when the flat part \mathcal{F} of Ω_X might determine the Iitaka fibration $f : X \rightarrow Y$ with $\mathcal{F} = \Omega_{X/Y}$. The second main result (Theorem 1.4) answers this question.

Theorem 1.4. *Let X be a smooth projective variety with the nef cotangent bundle Ω_X . Assuming the abundance conjecture, we consider the Iitaka fibration $f : X \rightarrow Y$ of X . Then, the following conditions are equivalent:*

- (1) *The flat part \mathcal{F} of the Fujita decomposition of Ω_X coincides with $\Omega_{X/Y}$.*
- (2) *The relative cotangent bundle $\Omega_{X/Y}$ is Hermitian flat.*
- (3) *The Iitaka fibration $f : X \rightarrow Y$ is locally trivial.*

Assuming the abundance conjecture for a smooth projective variety X with nef cotangent bundle, Höring in [Hör13] proved that $f : X \rightarrow Y$ is an abelian group scheme (up to finite étale covers), i.e., that there exists a fibration $g : Y \rightarrow Z$ onto a subvariety Z in the fine moduli of polarized abelian varieties with level structure so that $f : X \rightarrow Y$ is recovered by the fiber product of the universal family via $g : Y \rightarrow Z$. From this viewpoint, the local triviality of $f : X \rightarrow Y$ can be rephrased as Z being the one point. Hence, the dimension $\dim Z$ can be interpreted as measuring the variation of the fibers. Theorem 1.4 relates the condition of $\dim Z = 0$ to a certain flatness of $\Omega_{X/Y}$. Problem (c) asks a more detailed mechanism of the variation of the fibers of the Iitaka fibration $f : X \rightarrow Y$. The third main result answers this question in terms of the Fujita decomposition.

Theorem 1.5. *Under the same situation as in Theorem 1.4, we consider the Fujita decomposition of the nef cotangent bundle Ω_X :*

$$0 \rightarrow \mathcal{H} \rightarrow \Omega_X \rightarrow \mathcal{F} \rightarrow 0,$$

where \mathcal{F} is a Hermitian flat vector bundle and \mathcal{H} is a generically ample vector bundle.

(1) Let l be the rank of maximal Hermitian flat subbundles of $\mathcal{H}|_{X_y}$, where $\mathcal{H}|_{X_y}$ is the restriction of \mathcal{H} to the fiber $X_y := f^{-1}(y)$ at a general point $y \in Y$. Then, we have $\dim Z = \text{rk}\mathcal{H} - l$.

(2) The Iitaka fibration $f : X \rightarrow Y$ is locally trivial if Ω_X is semipositive in the following sense: the hyperplane bundle $\mathcal{O}_{\mathbb{P}(\Omega_X)}(1)$ admits a smooth Hermitian metric with semipositive Chern curvature on the projective space bundle $\mathbb{P}(\Omega_X)$.

We emphasize that the conditions of differential geometry (e.g., Hermitian flatness or semipositivity) are related to the variation of the fibers, which is an interesting point. As one of benchmarks, Theorem 1.5 (2) should be compared to [Hör13, 1.3 Theorem], which states that $f : X \rightarrow Y$ is locally trivial if Ω_X is semi-ample. The semipositivity of $\mathcal{O}_{\mathbb{P}(\Omega_X)}(1)$ in Theorem 1.5 (2) is weaker than the condition of X admitting nonpositive bisectional curvature or Ω_X being semi-ample. Therefore, under the abundance conjecture, Theorem 1.5 can be seen as a generalization of [WZ02, Liu14] and [Hör13, 1.3. Theorem] to the semipositive case.

We finally give a technical remark on the condition $c_2(X) = 0$, which enables us to relax the assumption of Theorem 1.2. Note that the cotangent bundle Ω_X is always generically nef when K_X is nef by [Miy87]. As an application, we obtain a structure theorem for a compact Kähler manifold X such that $-K_X$ is nef and $c_2(X) = 0$. When X is projective, Corollary 1.7 was already proved in [Cao13] and [Ou17] by using the theory of extremal rays. Our method is different from their proof.

Theorem 1.6. *Let (X, ω) be a compact Kähler manifold of dimension n , and \mathcal{E} be a reflexive coherent sheaf on X . Assume that \mathcal{E} is generically nef and $c_1(\mathcal{E})$ is nef.*

(1) *There exists a positive number ε_0 such that \mathcal{E} is a nef vector bundle and $c_2(\mathcal{E}) = 0$, if $c_2(\mathcal{E})(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} = 0$ holds for some $\varepsilon_0 > \varepsilon > 0$.*

(2) *Assume that X is projective. The sheaf \mathcal{E} is a nef vector bundle and $c_2(\mathcal{E}) = 0$ if $c_2(\mathcal{E})A^{n-2} = 0$ holds for some ample line bundle A .*

Corollary 1.7. *Let X be a compact Kähler manifold with the nef anti-canonical bundle $-K_X$. If $c_2(X) = 0 \in H^{2,2}(X, \mathbb{R})$, then the tangent bundle T_X is nef. Moreover, there exists a finite étale cover $\pi : X' \rightarrow X$ such that X' is a torus or a \mathbb{P}^1 -fiber bundle over a torus.*

This paper is organized as follows: In Section 2, we summarize various positivity conditions for torsion-free sheaves. In Section 3, we prove Theorems 1.2, 1.3 and a structure theorem of the Iitaka fibration $f : X \rightarrow Y$ (Theorem 3.5). In Section 4, we prove Theorems 1.4 and 1.5. In Section 5, we prove Theorem 1.5. Section 6 is an appendix where we provide small remarks.

Acknowledgments. The authors thank Prof. Wenhao Ou for some helpful comments about second Chern classes of torsion-free coherent sheaves and giving the reference [Gri10]. M. I. thanks Dr. Masaru Nagaoka for discussions and helpful comments. He was partially supported by Osaka City University, Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics, JPMXP0619217849) and Grant-in-Aid for Early Career Scientists #22K13907. He also thanks the members in RACMaS -Research Alliance Center

for Mathematical Sciences-, Tohoku University. S.M. thanks Prof. Xiaojun Wu for discussions about special varieties. This paper was written during his stay at the University of Bayreuth. He also thanks the members of the University of Bayreuth for their hospitality. He was partially supported by Grant-in-Aid for Scientific Research (B) #21H00976 and Fostering Joint International Research (A) #19KK0342 from JSPS.

2. PRELIMINARIES

2.1. Notation. Throughout this paper, we work over the field of complex numbers, and basically follow the notation of [Har77, KM98]. We interchangeably use the words “Cartier divisors,” “invertible sheaves,” and “line bundles,” and we also identify “locally free sheaves of finite rank” with “vector bundles.” We assume that all the sheaves are coherent without explicit mention. For a torsion-free sheaf \mathcal{F} on a normal variety X , we define the dual sheaf \mathcal{F}^* and the determinant sheaf $\det \mathcal{F}$ by

$$\mathcal{F}^* := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_X) \quad \text{and} \quad \det \mathcal{F} := (\Lambda^{\text{rk} \mathcal{F}} \mathcal{F})^{**}.$$

2.2. Positivity, flatness, and slope of torsion-free sheaves. In this subsection, we summarize some properties defined for vector bundles or torsion-free sheaves. Throughout this subsection, let X be a compact Kähler manifold of dimension n , and let E be a vector bundle of rank r on X .

Definition 2.1 ([DPS94, Proposition 1.11, Definitions 1.7, 1.9], [Kob87, Propositions 2.5, 4.21, 4.22, Corollary 2.7], [Pet12, Definition 2.1]). A vector bundle E is said to be

- (1) *Griffith semipositive* if E admits a smooth Hermitian metric with Griffith semipositive curvature;
- (2) *nef* if the hyperplane bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ is a nef line bundle on the projective space bundle $\mathbb{P}(E)$;
- (3) *semipositive* if $\mathcal{O}_{\mathbb{P}(E)}(1)$ admits a smooth Hermitian metric with a semipositive Chern curvature on the projective space bundle $\mathbb{P}(E)$;
- (4) *Hermitian flat* if E is Griffiths semipositive and $c_1(E) = 0$;
- (5) *numerically flat* if E is nef and $c_1(E) = 0$;
- (6) *flat* if E admits a connection ∇ whose curvature vanishes;
- (7) *projectively Hermitian flat* if E admits a smooth Hermitian metric h such that the Chern curvature tensor $\Theta_{E,h}$ satisfies $\Theta_{E,h} = \alpha \text{Id}_E$ for some 2-form α ;
- (8) *projectively flat* if E admits a connection ∇ such that $\nabla^2 = \alpha \text{Id}_E$ for some 2-form α , equivalently, there exists a representation

$$\rho : \pi_1(X) \rightarrow \mathbb{P}GL(r, \mathbb{C})$$

such that $\mathbb{P}(E) \cong X_{\text{univ}} \times_{\rho} \mathbb{P}^{r-1}$, where X_{univ} is the universal cover of X .

We further assume that X is a smooth projective variety. Let A_1, \dots, A_{n-1} be ample line bundles on X . Then E is said to be

- (9) (A_1, \dots, A_{n-1}) -generically ample (resp. (A_1, \dots, A_{n-1}) -generically nef) if $E|_C$ is ample (resp. nef) for a curve $C := D_1 \cap \dots \cap D_{n-1}$ defined by general members $D_i \in |m_i A_i|$ and $m_i \gg 0$.

The vector bundle E is simply called *generically ample* (resp. *generically nef*) if E is (A_1, \dots, A_{n-1}) -generically ample (resp. (A_1, \dots, A_{n-1}) -generically nef) for any ample line bundles $\{A_i\}_{i=1}^{n-1}$.

We now summarize relations between a certain flatness and stabilities after we review slopes of torsion-free sheaves. Let \mathcal{E} be a torsion-free sheaf of rank r on X and $\alpha \in H^{n-1, n-1}(X, \mathbb{R})$ be a movable class on X . Then, the *slope* $\mu_\alpha(\mathcal{E})$ of \mathcal{E} with respect to α is defined by

$$\mu_\alpha(\mathcal{E}) := \frac{c_1(\mathcal{E})\alpha}{r} \quad \text{and} \quad c_1(\mathcal{E}) := c_1(\det \mathcal{E}).$$

The maximal slope $\mu_\alpha^{\max}(\mathcal{E})$ (resp. the minimal slope $\mu_\alpha^{\min}(\mathcal{E})$) is defined by the supremum of $\mu_\alpha(\mathcal{S})$ for any nonzero subsheaf $\mathcal{S} \subset \mathcal{E}$ (resp. the infimum of $\mu_\alpha(\mathcal{Q})$ for any nonzero torsion-free quotient sheaf $\mathcal{E} \rightarrow \mathcal{Q} \rightarrow 0$). The sheaf \mathcal{E} is said to be *α -semistable with respect to α* if $\mathcal{E} = \mathcal{E}_{\max}$. Here $\mathcal{E}_{\max} \subset \mathcal{E}$, which is called the *maximal destabilizing sheaf*, is the maximal subsheaf with respect to the inclusion such that $\mu_\alpha(\mathcal{E}_{\max}) = \mu_\alpha^{\max}(\mathcal{E})$, which uniquely exists (see [Kob87, Lemma 7.17]).

Theorem-Definition 2.2 ([Nak04, Chapter 4. Theorem 4.1], [Wu20, Proposition 13, Corollary 10], [LOY20, Definition 4.1]). A torsion-free sheaf \mathcal{E} on X is said to be *projectively numerically flat* if it satisfies one of the equivalent following conditions:

- (1) \mathcal{E} is locally free, and the \mathbb{Q} -twisted vector bundle $\mathcal{E}\langle \frac{\det \mathcal{E}^*}{r} \rangle$ is nef.
- (2) \mathcal{E} is $\{\omega\}^{n-1}$ -semistable and satisfies that

$$\left(c_2(\mathcal{E}) - \frac{r-1}{2r} c_1(\mathcal{E})^2 \right) \{\omega\}^{n-2} = 0$$


for some Kähler form ω .

- (3) \mathcal{E} is locally free, and there exists a filtration of subbundles:

$$0 =: E_0 \subset E_1 \subset \dots \subset E_l := \mathcal{E}$$

such that $G_i := E_i/E_{i-1}$ is a projectively Hermitian flat vector bundle and $c_1(G_i)/\text{rk}G_i = c_1(E)/r \in H^{1,1}(X, \mathbb{R})$ holds for any $i = 1, \dots, l$.

These relations can be summarized by the following table.



The notion of generically ample vector bundles with respect to ample line bundles can be characterized by the slope, which enables us to extend the definition to any movable classes on compact Kähler manifolds.

Lemma 2.3. *Let X be a smooth projective variety of dimension n . Let A_1, \dots, A_{n-1} be ample line bundles on X . Then, a torsion-free sheaf \mathcal{E} is (A_1, \dots, A_{n-1}) -generically ample (resp. (A_1, \dots, A_{n-1}) -generically nef) if and only if $\mu_{A_1 \dots A_{n-1}}^{\min}(\mathcal{E}) > 0$ (resp. $\mu_{A_1 \dots A_{n-1}}^{\min}(\mathcal{E}) \geq 0$).*

Definition 2.4. Let α be a movable class on a compact Kähler manifold X . Then, a torsion-free sheaf \mathcal{E} is said to be α -generically ample (resp. α -generically nef) if $\mu_{\alpha}^{\min}(\mathcal{E}) > 0$ (resp. $\mu_{\alpha}^{\min}(\mathcal{E}) \geq 0$).

Proof of Lemma 2.3. Assume that \mathcal{E} is (A_1, \dots, A_{n-1}) -generically ample. Let

$$0 =: \mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots \subset \mathcal{E}_l := \mathcal{E}.$$

be the Harder-Narasimhan filtration with respect to this polarization. Then, straightforward computation yields $\mu_{A_1 \dots A_{n-1}}^{\min}(\mathcal{E}) = \mu_{A_1 \dots A_{n-1}}(\mathcal{E}_l/\mathcal{E}_{l-1})$. By definition, we have

$$\mu_{A_1 \dots A_{n-1}}(\mathcal{E}_l/\mathcal{E}_{l-1}) = \frac{1}{m_1 \dots m_{n-1} \cdot \text{rank}(\mathcal{E}_l/\mathcal{E}_{l-1})} \deg_C((\mathcal{E}_l/\mathcal{E}_{l-1})|_C),$$

where $C := D_1 \cap \dots \cap D_{n-1}$ is a curve defined by general members D_i in $|m_i A_i|$ and $m_i \gg 0$. Then, the right-hand side is positive since $\mathcal{E}|_C$ is ample.

The converse implication follows from Mehta-Ramanathan's theorem. The restriction of the Harder-Narasimhan filtration to C constructed above also gives a Harder-Narasimhan filtration of $\mathcal{E}|_C$. Hence, we obtain

$$0 < \mu_{A_1 \dots A_{n-1}}^{\min}(\mathcal{E}) = \mu_{A_1 \dots A_{n-1}}(\mathcal{E}_l/\mathcal{E}_{l-1}) = \mu(\mathcal{E}_l/\mathcal{E}_{l-1}|_C) = \mu^{\min}(\mathcal{E}|_C),$$

which finishes the proof. The proof for the nef case is almost the same, and thus we omit it. \square

The following lemma is easily obtained, but useful in this paper.

Lemma 2.5. *Let X be a compact Kähler manifold and E be a vector bundle on X .*

- (1) *If E is pseudoeffective, then E is α -generically nef for any movable class α .*
- (2) *If E is pseudoeffective and $\mu_{\beta}^{\min}(E) > 0$ for some movable class β , then E is α -generically ample for any class α in the interior of the movable cone.*

Proof. (1). For any torsion-free quotient $E \rightarrow Q$, the Chern class $c_1(Q)$ is pseudoeffective, and thus $\mu_{\alpha}^{\min}(E) \geq 0$ holds for any movable class α .

(2). Assume that there exists a class α in the interior of the movable cone such that $\mu_{\alpha}^{\min}(E) \leq 0$. By definition, we can find a torsion-free quotient $E \rightarrow Q$ such that $\mu_{\alpha}(Q) \leq 0$. Then, we obtain $c_1(Q) = 0$ since $\det Q$ is pseudoeffective (for example, see [Mat13, Lemma 2.1]). On the other hand, by assumption, we obtain $\mu_{\beta}^{\min}(E) \leq \mu_{\beta}(Q) = 0$, which is a contradiction. \square

Remark 2.6. In this paper, we say E is *pseudoeffective* if the nonnef locus of $\mathcal{O}_{\mathbb{P}(E)}(1)$ is not dominant over X . This definition requires a stronger condition than the pseudoeffectivity of $\mathcal{O}_{\mathbb{P}(E)}(1)$. The converse of (1) in Lemma 2.5 does not hold. Indeed, by [CPe11], the cotangent bundle Ω_X is generically nef when K_X is nef. Therefore, the cotangent bundle of a (strict) Calabi-Yau manifold is generically nef, but it is not pseudoeffective by [DPS01, Corollary 6.12].

3. ABUNDANCE THEOREM AND STRUCTURE THEOREM

3.1. Abundance theorem. This subsection is devoted to the proofs of Theorems 1.2 and 1.3. We first prove Theorem 1.2, assuming Theorem 1.6. Theorem 1.6 is proved in Section 5.

Proof of Theorem 1.2. Assume Condition (2). Then, from [DPS94, Corollary 2.6], we obtain

$$0 \leq c_2(X)\omega^{n-2} \leq c_1(X)^2\omega^{n-2} = 0$$

for any Kähler form ω . This implies that $c_2(X) = 0$ by Theorem 1.6 (1).

Conversely, we assume Condition (1). Since K_X is nef, the cotangent bundle Ω_X is $\{\omega\}^{n-1}$ -generically nef for any Kähler form ω by [Eno93, Theorem 1.4] and [Cao13, Theorem 1.2]. Then, by Theorem 1.6 (1) and $c_2(X) = 0$, we can see that Ω_X is nef. It remains to show that $\nu(K_X) \leq 1$. If $\nu(K_X) > 1$, there exists a nef line bundle $L \subset \Omega_X$ such that $c_1(L) = c_1(K_X)$. This follows from [Cao13] in the Kähler case and [Ou17] in the projective case (see Theorems 5.7 and 5.3). Then, the Bogomolov-Sommese vanishing theorem implies $H^0(X, \Omega_X \otimes L^*) = 0$. This contradicts the fact that L is a subbundle of Ω_X . Hence, we conclude that $\nu(K_X) = 0$ or 1. \square

For the proof of Theorem 1.3, we focus on the notion of special manifolds introduced by Campana in [Cam04]. A compact Kähler manifold X is said to be *special* if any $p \in \mathbb{Z}_+$ and any saturated subsheaf $F \subset \Omega_X^p$ of rank 1 satisfies $\kappa(F) < p$, where $\kappa(F)$ denotes the Kodaira dimension of F . In general, it is quite difficult to determine whether a given manifold is special or nonspecial. Nevertheless, the following theorem gives a criteria of manifolds to be nonspecial.

Theorem 3.1. *Let X be a compact Kähler manifold. If there exists a projectively flat subsheaf $E \subset \Omega_X$ with $\nu(c_1(E)) = 1$, then X is not special.*

Proof. Any finite étale cover of special manifolds is special by [Cam04, Theorem 5.12], and thus we can freely replace X with its finite étale covers. By the definition of the projective flatness, there exists a representation

$$\rho : \pi_1(X) \rightarrow \mathbb{P}GL(r, \mathbb{C})$$

such that $\mathbb{P}(E) \cong X_{\text{univ}} \times_{\rho} \mathbb{P}^{r-1}$, where $r := \text{rk} E$.

In the case where $\text{Im}(\rho)$ is finite, the Kernel $\text{Ker}(\rho) \subset \pi_1(X)$ is of finite index. Hence, after replacing X with the finite étale cover corresponding to $\text{Ker}(\rho)$, we may assume that $\text{Im}(\rho)$ is the one point. This implies that $E \cong L^{\oplus r}$ holds for some line bundle L . We have $\nu(L) = \nu(c_1(E)) = 1$ by assumption, and hence X is not special by [PRT21, Theorem A].

From now on, we assume that $\text{Im}(\rho)$ is infinite. Let H be the algebraic Zariski closure of $\text{Im}(\rho)$. Note H has finitely many connected components since H is a linear algebraic group. After replacing X with its finite étale cover, we may assume that H is connected. For the solvable radical $\text{Rad}(H)$ of H , we consider

$$\rho' : \pi_1(X) \xrightarrow{\rho} H \twoheadrightarrow H/\text{Rad}(H).$$

and the kernel $K := \text{Ker}(\rho')$. By Selberg's Lemma, we may assume that the image of ρ' is torsion-free. We now take a K -Shafarevich map $\text{sha}^K : X \dashrightarrow \text{Sha}^K(X)$ (see [Kol95, Theorem 3.6] or [Cam94, Chapter 3] for Shafarevich maps.)

In the case of $\dim \text{Sha}^K(X) = 0$, after replacing X with its finite étale cover, we may assume that the image of ρ is contained in $\text{Rad}(H)$. Then, by the argument of [GKP21, Proposition 3.3], we can conclude that

$$E \cong A \otimes G,$$

where A is a line bundle and G is a flat vector bundle admitting a filtration

$$0 = G_0 \subset G_1 \subset \cdots \subset G_r = G$$

such that G_i is a flat vector bundle and $G_1 \cong \mathcal{O}_X$. From $G_1 \cong \mathcal{O}_X$, we obtain

$$A = G_1 \otimes A \subset G \otimes A = E \subset \Omega_X.$$

Hence, from $\nu(A) = \nu(c_1(E)) = 1$, we can conclude that X is not special by using [PRT21, Theorem A] again.

When $\dim \text{Sha}^K(X) > 0$, the base variety $\text{Sha}^K(X)$ of the K -Shafarevich map is of general type since $H/\text{Rad}(H)$ is semi-simple and the image of ρ' is torsion-free, which follows from [CCE15, Théorème 1]. Then, the manifold X is not special by the definition of special manifolds. \square

The theorem below ensures that nonspecial manifolds admit a fibration of general type in the sense of Campana (see [Cam04, Cam11] for the details).

Theorem 3.2 ([Cam04, Theorems 2.25, 2.26 and Corollaries 1.13, 4.7]). *Let X be a compact Kähler manifold. Assume X is not special. Then, there exists an almost holomorphic map $f : X \dashrightarrow Y$, called the core fibration, onto a compact Kähler manifold Y with $0 < \dim Y \leq \dim X$ satisfying the following diagram and properties:*

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \pi_X \downarrow & & \downarrow \pi_Y \\ X & \dashrightarrow & Y, \end{array}$$

- (1) Both $\pi_X : X' \rightarrow X$ and $\pi_Y : Y' \rightarrow Y$ are bimeromorphic maps between compact Kähler manifolds.
- (2) A general fiber F' of $f' : X' \rightarrow Y'$ is special.
- (3) $\kappa(K_{Y'} + \Delta(f')) = \dim Y'$ holds, where $\Delta(f')$ is the multiplicity divisor of f' .
- (4) $\kappa(K_{X'}) = \kappa(K_{F'}) + \dim Y'$ holds for a general fiber F' of $f' : X' \rightarrow Y'$.

Under the above preparation, we prove Theorem 1.3.

Proof of Theorem 1.3. When $\nu(K_X) = 0$, there exists a finite étale cover $T \rightarrow X$ from a torus T . This implies that K_X is semi-ample.

In the case of $\nu(K_X) = 1$, we prove the semi-ampleness of K_X by induction on $n := \dim X$. In this case, we have $c_2(X) = 0$ by Theorem 1.2. Hence, by applying Theorem 5.7 (see Theorem 5.3 in the projective case), we can find a projectively numerically flat subbundle $E \subset \Omega_X$ with $\nu(c_1(E)) = 1$. The subbundle E is projectively flat by [LOY20, Theorem 4.6], and thus X is not special by Theorem 3.1. Then, by Theorem 3.2, there exists a nontrivial core fibration $f : X \dashrightarrow Y$. Note

that $\kappa(K_{X'}) = \kappa(K_X)$ and $\kappa(K_{F'}) = \kappa(K_F)$ since X and F are smooth, where F is a general fiber of $f : X \dashrightarrow Y$. This implies that

$$\kappa(K_X) = \kappa(K_F) + \dim Y.$$

It is sufficient to show that $\kappa(K_F) \neq -\infty$.

The cotangent bundle Ω_F is nef, and $\nu(K_F) = \nu(K_X|_F) \leq \nu(K_X) = 1$ holds. Hence, by the induction hypothesis, the canonical bundle K_F is semi-ample. In particular, we have $\kappa(K_F) \geq 0$. This implies that

$$1 = \nu(K_X) \geq \kappa(K_X) = \kappa(K_F) + \dim Y \geq \dim Y \geq 1.$$

Hence, the canonical divisor K_X is semi-ample. \square

3.2. Structure theorem. In this subsection, with specific examples, we examine properties of the Iitaka fibration $f : X \rightarrow Y$ of X with nef cotangent bundle. When X is a smooth projective variety, Höring in [Hör13] showed that, up to finite étale covers, the Iitaka fibration $f : X \rightarrow Y$ (if it exists) is an abelian group scheme onto a smooth projective variety Y . In particular, the Iitaka fibration $f : X \rightarrow Y$ always admits a section $s : Y \rightarrow X$. Furthermore, the base variety Y has the nef cotangent bundle and the ample canonical bundle K_Y . However, by Example 3.3, we see that the Iitaka fibration does not necessarily have a section when we assume only that X is Kähler. This comes from the lack of fine moduli of complex tori. Further, by Example 3.4, we see that the Iitaka fibration $f : X \rightarrow Y$ is not smooth and K_Y is not ample, without taking finite étale covers even if X is projective.

Example 3.3. This example was given by Donu Arapura in Mathoverflow [Math]. Let C be a smooth curve of the genus $g \geq 2$ curve and E be an elliptic curve. Fixing a nontorsion point $p \in E$ and the isomorphism $H_1(C, \mathbb{Z}) \cong \mathbb{Z}^{2g}$, we define a homomorphism

$$\begin{aligned} H_1(C, \mathbb{Z}) \cong \mathbb{Z}^{2g} &\rightarrow E \\ (x_1, \dots, x_{2g}) &\mapsto \left(\sum_{i=1}^{2g} x_i\right)p. \end{aligned}$$

By composing the natural morphism $\pi_1(C) \rightarrow H_1(C, \mathbb{Z})$ and \tilde{h} , we obtain a homomorphism $h : \pi_1(C) \rightarrow E$. Note that the image of h is dense in E since $p \in E$ is a nontorsion point.

Let C_{univ} be a universal cover of C . We define an action of $\pi_1(C)$ on $C_{\text{univ}} \times E$ to be $\gamma(x, y) = (\gamma^{-1}x, y + h(\gamma))$ for $\gamma \in \pi_1(C)$. Let X be the quotient of $C_{\text{univ}} \times E$ by this action. Then the induced morphism $f : X \rightarrow C$ by projection gives the Iitaka fibration of X . The Iitaka fibration $f : X \rightarrow C$ does not admit a section even if we replace X with finite étale covers. Indeed, if the Iitaka fibration $X' \rightarrow C'$ of a finite étale cover $X' \rightarrow X$ admits a section, the image of the section gives a relatively ample divisor on X . Therefore, we can conclude that X' (and thus X) is projective. However, the elliptic curve E does not have a nontrivial line bundle that is invariant by the action of $h : \pi_1(C) \rightarrow E$. This implies that X is not projective.

Example 3.4 ([Cam11, Example 2.12]). This example shows that the Iitaka fibration $f : X \rightarrow Y$ is not smooth and that the base variety Y is not of general type without taking finite étale covers of X .

Let E be an elliptic curve with a translation τ of order 2 and C be a hyperelliptic curve of genus $g \geq 2$ with an involution $i : C \rightarrow C$. Then, the group $G := \mathbb{Z}/2\mathbb{Z}$ acts

on $C \times E$ by i and τ . Let us consider the quotient map $p : C \times E \rightarrow (C \times E)/G$. Then, since p is a finite étale cover, the cotangent bundle Ω_X is nef and $\kappa(K_X) = \nu(K_X) = 1$ holds. The morphism $f : X \rightarrow C/\langle i \rangle \cong \mathbb{P}^1$ induced by the first projection is the Iitaka fibration of X . Note that the base variety \mathbb{P}^1 is, of course, not of general type, but $f : X \rightarrow C/\langle i \rangle \cong \mathbb{P}^1$ is the core fibration of X and a morphism of general type.

We prove a structure theorem in the case of $\nu(K_X) = 1$.

Theorem 3.5. *Let X be a compact Kähler manifold such that Ω_X is nef and $\nu(K_X) = 1$. Then, there exists a finite étale cover $X' \rightarrow X$ such that the Iitaka fibration $f' : X' \rightarrow Y'$ is a smooth fibration onto a smooth curve Y' of genus ≥ 2 whose fibers are complex tori.*

Proof. The Iitaka fibration $f : X \rightarrow Y$ is equidimensional from $\dim Y = 1$, and the tangent bundle T_X is f -numerically flat. Hence, by [Hör13, Lemma 2.1 (2)], there exists a finite étale cover $\pi : X' \rightarrow X$ such that f' is a smooth morphism and all fibers of f' are complex tori, where f' and π' are the morphisms that give the Stein factorization of $f \circ \pi$.

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \pi \downarrow & & \downarrow \pi' \\ X & \xrightarrow{f} & Y \end{array}$$

We show that $K_{Y'}$ is big. Let $X' \dashrightarrow Z$ be the core fibration of X . From the proof of Theorem 1.3, the core fibration of X' coincides with the Iitaka fibration of X' and the base variety Z is a curve. Furthermore, since the fibers of f' are complex tori (and thus special), the morphism $f' : X' \rightarrow Y'$ gives the core fibration of X' (and thus the Iitaka fibration). By the definition of the core fibration, the divisor $K_{Y'} + \Delta(f')$ is big. In our case, since f' is smooth, we have $\Delta(f') = 0$, which implies that $K_{Y'}$ is big. \square

4. ON VARIATION OF THE FIBERS OF IITAKA FIBRATIONS

4.1. On the Fujita decomposition of nef cotangent bundles. In this subsection, using the Fujita decomposition, we study the abundance conjecture for smooth projective varieties with nef cotangent bundles. To motivate this study, we first observe the Fujita decomposition of the cotangent bundle in the case of [WZ02, Liu14].

Observation 4.1. (Fujita decomposition in the case of nonpositive holomorphic bisectional curvature). Let X be a smooth projective variety admitting a Kähler metric g with nonpositive holomorphic bisectional curvature. Wu-Zheng and Liu in [WZ02, Liu14] showed that a holomorphic Ricci kernel foliation \mathcal{R} induced by the metric g determines a locally trivial fibration $f : X \rightarrow Y$, which actually coincides with the Iitaka fibration. Then, we can show that the standard exact sequence of tangent bundles

$$0 \rightarrow f^*\Omega_Y \rightarrow \Omega_X \rightarrow \Omega_{X/Y} = \mathcal{R} \rightarrow 0$$

corresponds to the Fujita decomposition of Ω_X .

Under the weaker assumption that the cotangent bundle Ω_X is nef, we cannot construct a holomorphic Ricci kernel foliation \mathcal{R} and not expect that the Iitaka fibration $f : X \rightarrow Y$ is locally trivial. However, we can still consider the flat part of the Fujita decomposition of Ω_X . Therefore, it is natural to ask how the Fujita decomposition of Ω_X is related to the Iitaka fibration of X . Specifically, in this paper, we study whether the Fujita decomposition determines the Iitaka fibration and what relation exists between the flat part and the local triviality.

We first begin with the definition of the Fujita decomposition.

Proposition 4.2 (cf. [Iwa21, Section 3]). *Let E be a nef vector bundle on a smooth projective variety X .*

(a) *There exists a numerically flat vector bundle F and a generically ample vector bundle H such that the sequence $0 \rightarrow H \rightarrow E \rightarrow F \rightarrow 0$ is exact.*

(b) *In the case of $E = \Omega_X$, we further have:*

- (1) *F is Hermitian flat, $\det F$ is a torsion line bundle, and $F^* \subset T_X$ is a regular foliation.*
- (2) *$0 \rightarrow H \rightarrow E \rightarrow F \rightarrow 0$ splits.*

Proof. Conclusion (a) has been proven in [Iwa21, Lemma 3.1]. We prove only Conclusion (b) by reviewing [Iwa21, Lemma 3.1].

Let A be an ample line bundle on X . When $\mu_{A^{n-1}}^{\min}(\Omega_X) > 0$, the cotangent bundle Ω_X is generically ample by Lemma 2.5. Then, we see that $F := 0$ and $H := \Omega_X$ satisfy the desired conclusions. Hence, we may assume that $\mu_{A^{n-1}}^{\min}(\Omega_X) = 0$.

In this case, we define \mathcal{F} by the maximal destabilizing subsheaf of T_X with respect to A^{n-1} . Then, we have

$$\mu_{A^{n-1}}(\mathcal{F}) = \mu_{A^{n-1}}^{\max}(T_X) = -\mu_{A^{n-1}}^{\min}(\Omega_X) = 0.$$

The sheaf \mathcal{F} is pseudoeffective, as is $\det \mathcal{F}^*$ since it is a quotient of the nef cotangent bundle. We obtain $c_1(\mathcal{F}) = 0 \in H^{1,1}(X, \mathbb{R})$ from $c_1(\mathcal{F}^*)A^{n-1} = 0$. Hence, the sheaf \mathcal{F} is a numerically flat vector bundle (see [GKP16, Theorem 1.20] or [HIM21, Theorem 1.2]). On the other hand, the canonical bundle K_X is nef, and \mathcal{F} is a saturated subsheaf with $c_1(\mathcal{F}) = 0$. Thus, by [LPT18, Theorem 5.2] and [PeT13, Lemma 2.1], we see that

- $\mathcal{F} \subset T_X$ is a regular foliation,
- \mathcal{F} is polystable with respect to A ,
- $K_{\mathcal{F}} = \det \mathcal{F}^*$ is a torsion line bundle,
- $T_X = \mathcal{F} \oplus \mathcal{G}$ holds for some regular foliation $\mathcal{G} \subset T_X$.

The vector bundle \mathcal{F} is numerically flat and polystable, and thus F is Hermitian flat. It remains to show that \mathcal{G}^* is generically ample. If $\mu_{A^{n-1}}^{\max}(\mathcal{G}) \geq 0$, we can find a nonzero subsheaf $\mathcal{N} \subset \mathcal{G}$ with $\mu_{A^{n-1}}(\mathcal{N}) \geq 0$. This contradicts the maximality of \mathcal{F} by $\mu_{A^{n-1}}(\mathcal{N} \oplus \mathcal{F}) \geq 0$. Hence, we obtain $\mu_{A^{n-1}}^{\max}(\mathcal{G}) < 0$, and thus \mathcal{G}^* is generically ample with respect to A . By Lemma 2.5, we see that \mathcal{G}^* is generically ample since \mathcal{G}^* is nef. \square

Example 4.3. Conclusion (b) in Proposition 4.2 does not hold for a nef vector bundle $E \neq \Omega_X$. A numerically flat vector bundle is not necessarily Hermitian flat (for example, see [DPS94]). Hence, Property (1) does not hold in general. When X

admits an ample line bundle L with $H^1(X, L) \neq 0$, we obtain a nonsplitting exact sequence:

$$0 \rightarrow L \rightarrow E \rightarrow \mathcal{O}_X \rightarrow 0$$

from $H^1(X, L) \cong \text{Ext}^1(\mathcal{O}_X, L) \neq 0$. This sequence gives the Fujita decomposition of E but does not split by construction.

The Fujita decomposition shows that Conjecture 4.4 can be reduced to the case where Ω_X is nef and generically ample.

Conjecture 4.4. _{$\leq n$} *Let X be a smooth projective variety of dimension n with a nef cotangent bundle. Then, the canonical bundle K_X is semi-ample.*

Proposition 4.5. *Let X be a smooth projective variety of dimension n with the nef cotangent bundle Ω_X . The flat part of the Fujita decomposition of Ω_X is nonzero. Then, if Conjecture 4.4 _{$\leq n-1$} holds, then K_X is semi-ample.*

Proof. Let $\Omega_X = F \oplus H$ be the Fujita decomposition as in Lemma 4.2. Since F is a nonzero Hermitian flat vector bundle by assumption, there exists a representation

$$\rho : \pi_1(X) \rightarrow U(r, \mathbb{C})$$

such that $F \cong X_{\text{univ}} \times_{\rho} \mathbb{C}^r$, where $r := \text{rk} F > 0$ and X_{univ} is the universal cover of X .

In the case where $\text{Im}(\rho)$ is finite, we may assume that F is a trivial vector bundle after replacing X with a finite étale cover. Then, we have $H^0(X, T_X) \neq 0$, and thus, up to finite étale covers, we can find an abelian variety A and a smooth projective variety Y with $H^0(Y, T_Y) = 0$ such that $X \cong A \times Y$ by a theorem of Lieberman [Lie78] (cf. [AMN12, Theorem 3.2]). The canonical bundle K_Y is semi-ample by Conjecture 4.4 _{$\leq n-1$} , and thus so is K_X .

We may assume that $\text{Im}(\rho)$ is infinite. Let G be the algebraic Zariski closure of $\text{Im}(\rho)$. Up to finite étale covers, we may assume that G is connected. Let K be the kernel of $\bar{\rho}$ defined by

$$\bar{\rho} : \pi_1(X) \xrightarrow{\rho} G \twoheadrightarrow G/\text{Rad}(G),$$

where $\text{Rad}(G)$ is the solvable radical of G . By Selberg's Lemma, we may assume that the image of $\bar{\rho}$ is torsion-free. We consider the K -Shafarevich map $\text{sha}^K : X \dashrightarrow \text{Sha}^K(X)$. Since $\text{Im}(\rho)$ is not virtually solvable by [PeT13, Corollary 2.7], we obtain $\dim \text{Sha}^K(X) > 0$. Since $G/\text{Rad}(G)$ is semi-simple and the image of $\bar{\rho}$ is torsion-free, the base variety $\text{Sha}^K(X)$ is of general type by [CCE15, Théorème 1]. The canonical bundle K_X is abundant by Conjecture 4.4 _{$\leq n-1$} and [GKP21, Proposition 2.10]. \square

4.2. Hermitian flatness of nef cotangent bundles. In this subsection, assuming the abundance conjecture, we study the associated Iitaka fibration $f : X \rightarrow Y$ of a smooth projective variety X with the nef cotangent bundle Ω_X . In particular, we study a relation among the Fujita decomposition of Ω_X , a certain flatness of $\Omega_{X/Y}$, and a variation of the fibers.

Compared to Observation 4.1, when Ω_X is merely nef, the Iitaka fibration $f : X \rightarrow Y$ is not necessarily locally trivial. This difference can be captured by the moduli of the abelian varieties. Precisely, assuming the abundance conjecture, Höring in

[Hör13] proved that, up to finite étale covers of X , the Iitaka fibration $f : X \rightarrow Y$ is an abelian group scheme, i.e., there exists a fibration $g : Y \rightarrow Z$ onto a subvariety Z in the fine moduli of polarized abelian varieties with level structure so that $f : X \rightarrow Y$ is recovered from the fiber product via $g : Y \rightarrow Z$. From this viewpoint, the local triviality of $f : X \rightarrow Y$ can be rephrased as Z being the one point. In this subsection, we prove that the dimension $\dim Z$, which measures how far $f : X \rightarrow Y$ is from the local triviality of f , is related to a certain flatness.

Keeping the above situation, we prove Theorems 1.4 and 1.5 in this subsection.

We consider the Fujita decomposition of the nef cotangent bundle Ω_X :

$$(4.1) \quad 0 \rightarrow \mathcal{H} \rightarrow \Omega_X \rightarrow \mathcal{F} \rightarrow 0,$$

where \mathcal{F} is a Hermitian flat vector bundle and \mathcal{H} is a generically ample vector bundle. The relative cotangent bundle $\Omega_{X/Y}$ is nef by the standard exact sequence of tangent bundles

$$(4.2) \quad 0 \rightarrow f^*\Omega_Y \rightarrow \Omega_X \rightarrow \Omega_{X/Y} \rightarrow 0.$$

Hence, we can consider the Fujita decomposition of $\Omega_{X/Y}$:

$$(4.3) \quad 0 \rightarrow \mathcal{G} \rightarrow \Omega_{X/Y} \rightarrow \mathcal{E} \rightarrow 0,$$

where \mathcal{E} is a numerically flat vector bundle and \mathcal{G} is a generically ample vector bundle. Then, we have:

Proposition 4.6. *The flat part of \mathcal{F} of Ω_X coincides with the flat part \mathcal{E} of $\Omega_{X/Y}$.*

Proof. We first prove that $f^*\Omega_Y$ is a generically ample vector bundle on X . For this purpose, we confirm that Ω_Y (and thus $f^*\Omega_Y$) is a nef vector bundle on X . There exists a section $s : Y \rightarrow X$ of $f : X \rightarrow Y$ since it is an abelian group scheme, and thus we obtain a surjective bundle morphism $\Omega_X|_{s(Y)} \rightarrow \Omega_{s(Y)}$ on the image $s(Y) \subset X$. This implies that Ω_Y is nef since $\Omega_{s(Y)}$ can be identified with Ω_Y via the isomorphism $s : Y \cong s(Y)$.

We now show that $f^*\Omega_Y$ is a generically ample vector bundle. The canonical bundle K_Y is ample by [Hör13]; hence, Y admits a Kähler-Einstein metric. This implies that Ω_Y is semistable. Let us consider a curve $C := A_1 \cap A_2 \cap \cdots \cap A_{\dim Y - 1}$ defined by general members $A_i \in |mK_Y|$, where $m \gg 1$. Then, by Mehta-Ranamathan's theorem, the restriction $\Omega_Y|_C$ is also stable. Miyaoka's characterization of semistabilities shows that the \mathbb{Q} -vector bundle

$$\Omega_Y|_C \otimes \det(\Omega_Y|_C)^{1/\dim Y - 1}$$

is nef. This implies that $\Omega_Y|_C$ is ample since K_Y is ample. We take a curve $C' \subset X$ such that the restriction $f|_{C'} : C' \rightarrow C$ is a surjective morphism. Then, we see that $(f^*\Omega_Y)|_{C'} = (f|_{C'})^*(\Omega_Y|_C)$ is an ample vector bundle on C' since $f|_{C'} : C' \rightarrow C$ is a finite morphism. However, for the flat part \mathcal{F}' of the nef vector bundle $f^*\Omega_Y$, we have the surjective morphism $(f^*\Omega_Y)|_{C'} \rightarrow \mathcal{F}'|_{C'}$ by restricting $f^*\Omega_Y \rightarrow \mathcal{F}'$ to C' . The vector bundle $(f^*\Omega_Y)|_{C'}$ is ample by the choice of C' , but the restriction $\mathcal{F}'|_{C'}$ is still flat. This is a contradiction. Hence, we conclude that $\mathcal{F}' = 0$ (i.e., $f^*\Omega_Y$ is generically ample).

By using exact sequences (4.1), (4.2), and (4.3), the snake lemma yields the following commutative diagram.

$$(4.4) \quad \begin{array}{ccccccc} & & & 0 & & 0 & \\ & & & \uparrow & & \uparrow & \\ & & & T_X/\mathcal{E}^* & \longrightarrow & f^*T_Y & \longrightarrow 0 \\ & 0 \longrightarrow & \mathcal{G}^* & \longrightarrow & & & \\ & & & \uparrow & & \uparrow & \\ & & & T_X & \xlongequal{\quad} & T_X & \\ & & & \uparrow & & \uparrow & \\ & 0 \longrightarrow & \mathcal{E}^* & \longrightarrow & T_{X/Y} & \longrightarrow & \mathcal{G}^* \longrightarrow 0 \\ & & & \uparrow & & \uparrow & \\ & & & 0 & & 0 & \end{array}$$

The dual bundle $(T_X/\mathcal{E}^*)^*$ is generically ample since \mathcal{G} and $f^*\Omega_X$ are as well. This implies that the (dual of) the left column in the above diagram actually corresponds to the Fujita decomposition of Ω_X . Hence, the flat part \mathcal{F} of Ω_X is the flat part \mathcal{E} of $\Omega_{X/Y}$. \square

Theorem 1.4 immediately follows from Proposition 4.6.

Proof of Theorem 1.4. The moduli part M of $f : X \rightarrow Y$ defined by $f^*M := K_{X/Y}$ is semi-ample by [Fuj03, Theorem 1.2] and [Uen78]. The semi-ample fibration $h : Y \rightarrow W$ of M coincides with $g : Y \rightarrow Z$, and thus $M = g^*A$ holds for some ample line bundle A on Z . Indeed, for a point $z \in Z$, the restriction $f|_{X_z} : X_z := (g \circ f)^{-1}(z) \rightarrow Y_z := g^{-1}(z)$ is a locally trivial fibration, and thus $M|_{Y_z}$ is trivial. Conversely, for a point $w \in W$, the relative canonical bundle K_{X_w/Y_w} is trivial, and thus $f|_{X_w} : X_w := (h \circ f)^{-1}(w) \rightarrow Y_w := g^{-1}(w)$ is locally trivial. This implies that $g(Y_w)$ comprises one point.

If $f : X \rightarrow Y$ is locally trivial (i.e., $\dim Z = 0$), we have $c_1(\Omega_{X/Y}) = 0$ by $K_{X/Y} = f^*M = f^*g^*A$. Hence, since $\Omega_{X/Y}$ is nef, we see that $\Omega_{X/Y}$ is numerically flat, and thus the vector bundle $\Omega_{X/Y}$ coincides with the flat part of Ω_X , which is Hermitian flat. Hence, Condition (3) leads to Conditions (1) and (2).

It is obvious that Condition (1) is equivalent to Condition (2). Therefore, it remains to show that Condition (1) leads to Condition (3). The Fujita decomposition of cotangent bundles always splits, and the sequence (4.2) of tangent bundles also splits under Condition (1). This implies that the Kodaira-Spencer map vanishes everywhere. Hence, the Iitaka fibration $f : X \rightarrow Y$ is locally trivial. \square

We finally prove Theorem 1.5.

Proof of Theorem 1.5. By pushing forward (4.4) by f , we obtain:

$$(4.5) \quad \begin{array}{ccccccc} & & R^1 f_* \mathcal{E}^* & \longrightarrow & R^1 f_* T_{X/Y} & & \\ & & \uparrow & & \uparrow \varphi & & \\ 0 & \longrightarrow & f_* \mathcal{G}^* & \longrightarrow & f_* \mathcal{H}^* = f_*(T_X/\mathcal{E}^*) & \xrightarrow{\psi} & T_Y \longrightarrow \cdots \\ & & \uparrow & & \uparrow & & \\ & & f_* T_X & \xlongequal{\quad\quad\quad} & f_* T_X & & \\ & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & f_* \mathcal{E}^* & \longrightarrow & f_* T_{X/Y} & \longrightarrow & \cdots \\ & & \uparrow & & \uparrow & & \\ & & 0 & & 0, & & \end{array}$$

where ψ and φ are the induced morphisms. Note that φ is the Kodaira-Spencer map. The Fujita decomposition (4.1) splits, and thus the induced morphism $f_* \mathcal{H}^* \rightarrow R^1 f_* \mathcal{E}^*$ is the zero map. Hence, for a general point $y \in Y$, we obtain

$$\text{Ker}(\varphi_y) = \text{Im}(\psi_y) \cong (f_* \mathcal{H}^*)_y / (f_* \mathcal{G}^*)_y = H^0(X_y, \mathcal{H}^*|_{X_y}) / H^0(X_y, \mathcal{G}^*|_{X_y}).$$

Furthermore, we have $T_{Y/Z, y} = \text{Ker}(\varphi_y)$ since $g : Y \rightarrow Z$ is the morphism into the moduli of the abelian varieties as in the proof of Theorem 1.4. Hence, we obtain

$$\dim Y - \dim Z = h^0(X_y, \mathcal{H}^*|_{X_y}) - h^0(X_y, \mathcal{G}^*|_{X_y})$$

We now have the exact sequence

$$0 \rightarrow \mathcal{E}^*|_{X_y} \rightarrow T_{X/Y}|_{X_y} \rightarrow \mathcal{G}^*|_{X_y} \rightarrow 0$$

from (4.1). The vector bundle $\mathcal{E}^*|_{X_y}$ is numerically flat, and $T_{X/Y}|_{X_y}$ is a trivial vector bundle. This implies that this sequence splits, and thus that $\mathcal{G}^*|_{X_y}$ is numerically flat and globally generated. Hence, we see that $\mathcal{G}^*|_{X_y}$ is a trivial vector bundle by [DPS94, Proposition 1.16]. This implies that $h^0(X_y, \mathcal{G}^*|_{X_y}) = \dim X - \dim Y - \text{rk} \mathcal{E}$, and thus, we conclude that

$$(4.6) \quad \dim Z = \text{rk} \mathcal{H} - h^0(X_y, \mathcal{H}^*|_{X_y}).$$

For the proof of Theorem 1.5, we investigate $h^0(X_y, \mathcal{H}^*|_{X_y})$ in detail. Let us consider the induced commutative diagram:

$$(4.7) \quad \begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & \mathcal{G}|_{X_y} \cong T_{\text{rk}\mathcal{G}} & \longrightarrow & \Omega_{X/Y}|_{X_y} \cong T_{n-m} & \longrightarrow & \mathcal{E}|_{X_y} \cong T_{\text{rk}\mathcal{E}} \longrightarrow 0 \\ & & \uparrow & & \uparrow p & & \parallel \\ 0 & \longrightarrow & \mathcal{H}|_{X_y} & \longrightarrow & \Omega_X|_{X_y} & \longrightarrow & \mathcal{E}|_{X_y} \cong T_{\text{rk}\mathcal{E}} \longrightarrow 0 \\ & & & & \uparrow i & & \\ & & & & f^*\Omega_Y|_{X_y} \cong T_m & & \\ & & & & \uparrow & & \\ & & & & 0, & & \end{array}$$

where $n = \dim X$ and $m := \dim Y$. Here, the notation T_k denotes the trivial vector bundle on X_y of rank k . Note that $\Omega_X|_{X_y}$ is numerically flat from the middle column.

The following lemma will be applied to $H := \mathcal{H}|_{X_y}$.

Lemma 4.7. *Let A be a vector bundle on a compact complex manifold. Assume that A is an extension of a trivial vector bundle by a trivial vector bundle. Then, any Hermitian flat subbundle $F \subset A$ is a trivial vector bundle.*

Proof. By assumption, we have the exact sequence:

$$0 \longrightarrow T_a \xrightarrow{i} A \xrightarrow{p} T_b \longrightarrow 0,$$

where T_k denotes the trivial vector bundle of rank k . This implies that A is numerically flat. This sequence and $F \subset A$ lead to the following commutative diagram:

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & p(F) & \longrightarrow & T_b & \longrightarrow & T_b/p(F) \longrightarrow 0 \\ & & \uparrow & & \uparrow p & & \uparrow \\ 0 & \longrightarrow & F & \longrightarrow & A & \longrightarrow & A/F \longrightarrow 0 \\ & & \uparrow & & \uparrow i & & \uparrow \\ 0 & \longrightarrow & i^{-1}(F) & \longrightarrow & T_a & \longrightarrow & T_a/i^{-1}(F) \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & 0 & & 0 & & 0. \end{array}$$

Considering the middle row, the quotient bundle A/F is nef and $c_1(A/F) = c_1(A) - c_1(F) = 0$, and thus A/F is numerically flat. This implies that the dual bundle $(T_a/i^{-1}(F))^*$ is nef. On the other hand, the quotient $T_a/i^{-1}(F)$ is globally generated from the bottom column, and thus $T_a/i^{-1}(F)$ is a trivial vector bundle. The quotient

$T_b/p(F)$ of A/F is nef and satisfies $c_1(T_b/p(F)) = c_1(A/F) - c_1(T_a/i^{-1}(F)) = 0$. Hence, we see that $T_b/p(F)$ is numerically and globally generated from the top column. Therefore $T_b/p(F)$ is also a trivial vector bundle. Then, we can conclude that the rows in the top and bottom split, and thus $p(F)$ and $i^{-1}(F)$ are trivial vector bundles. The left column splits since F is Hermitian flat by assumption. This leads to the desired conclusion. \square

We take a filtration of the numerically flat vector bundle $H := \mathcal{H}|_{X_y}$ by subbundles $\{F_k\}_{k=0}^p$, i.e.,

$$0 =: F_0 \subset F_1 \subset \cdots \subset F_{p-1} \subset F_p := H,$$

such that each quotient bundle F_k/F_{k-1} is a Hermitian flat vector bundle. By construction, the vector bundle F_k is inductively obtained as follows: F_k is the inverse image of \overline{F}_k by $H \rightarrow H/F_{k-1}$, where \overline{F}_k is a subbundle of H/F_{k-1} of the minimal rank that satisfies $c_1(\overline{F}_k) \cdot c_1(A)^{\dim X-1} \geq 0$.

The vector bundle $H = \mathcal{H}|_{X_y}$ satisfies the assumption of Lemma 4.7, and F_1 is Hermitian flat, and thus F_1 is a trivial vector bundle. This implies that F_1 is a trivial line bundle since F_1 has the minimal rank. The quotient H/F_1 also satisfies the assumption of Lemma 4.7 by the same argument as in Lemma 4.7. Hence, it follows that \overline{F}_2 is a trivial line bundle since \overline{F}_2 is Hermitian flat and has the minimal rank. Therefore, we can conclude that $p = \text{rk}H$ and F_k/F_{k-1} is a trivial line bundle.

We confirm Theorem 1.5 (1); i.e., $\dim Z = \text{rk}H - l$ holds. Here l is the maximal integer such that F_l is Hermitian flat. Let us consider

$$0 \rightarrow F_{k-1} \rightarrow F_k \rightarrow F_k/F_{k-1} \rightarrow 0$$

for every $k \in \mathbb{Z}_+$. Then, by $F_k/F_{k-1} = T_1$, the above sequence splits if and only if $h^0(X_y, F_k) = h^0(X_y, F_{k-1}) + 1$. This implies that $h^0(X_y, H) = l$. Theorem 1.5 (1) follows from (4.6).

We finally confirm Theorem 1.5 (2). Let $p : \mathbb{P}(\Omega_X) \rightarrow X$ be the projective space bundle over X . Then, we have

$$p_*(K_{\mathbb{P}(\Omega_X)/X} + \mathcal{O}_{\mathbb{P}(\Omega_X)}(n+1)) = \Omega_X \otimes \det \Omega_X = \Omega_X \otimes K_X.$$

By assumption, the hyperplane bundle $\mathcal{O}_{\mathbb{P}(\Omega_X)}(1)$ is semipositive, and thus, we can conclude that $\Omega_X \otimes K_X$ is Griffiths semipositive by applying the theory of positivity of direct images (see [PăT18, HPS18, BP08]). The restriction $K_X|_{X_y}$ is numerically trivial since $f : X \rightarrow Y$ is the Iitaka fibration. This implies that $\Omega_X|_{X_y}$ is Griffiths semipositive and that $H = \mathcal{H}|_{X_y}$ is Hermitian flat. Then, Theorem 1.5 (2) follows from (1). \square

5. GENERICALLY NEF VECTOR BUNDLES WITH VANISHING SECOND CHERN CLASS

This section describes the proof of Theorem 1.6.

5.1. Case of smooth projective varieties. We first begin with the following lemma. This lemma follows from the proof of [Nak04, Proposition 4.6] or [Wu20, Proposition 13], but we give a proof for reader's convenience.

Lemma 5.1. *Let X be a complex manifold and let*

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{E} \rightarrow \mathcal{G} \rightarrow 0$$

*be an exact sequence of torsion-free sheaves on X . Assume that \mathcal{F} is locally free, \mathcal{E} is reflexive, \mathcal{G}^{**} is locally free, and $\text{codim}(\text{Supp}(\mathcal{G}^{**}/\mathcal{G})) \geq 3$. Then, both \mathcal{E} and \mathcal{G} are locally free on X .*

Proof. Take a Zariski open set $X_0 \subset X$ with $\text{codim}(X \setminus X_0) \geq 3$ such that $\mathcal{G}|_{X_0} \cong \mathcal{G}^{**}|_{X_0}$ and $\mathcal{E}|_{X_0}$ are locally free. Let us consider the following exact sequence:

$$(5.1) \quad 0 \rightarrow \mathcal{F}|_{X_0} \rightarrow \mathcal{E}|_{X_0} \rightarrow \mathcal{G}^{**}|_{X_0} \rightarrow 0.$$

Since $\mathcal{G}^{**}|_{X_0}$ is locally free, we have $\text{Tor}_{\mathcal{O}_{X,x}}^1(\mathcal{O}_{X,x}/\mathfrak{m}_x, \mathcal{G}_x^{**}) = 0$ for any $x \in X_0$, where \mathfrak{m}_x is the maximal ideal of $\mathcal{O}_{X,x}$. Hence, the sequence (5.1) is exact as a sequence of vector bundles, and thus $\mathcal{E}|_{X_0}$ is determined by an extension class in $H^1(X_0, \mathcal{F}|_{X_0} \otimes (\mathcal{G}^{**}|_{X_0})^*)$. By $\text{codim}(X \setminus X_0) \geq 3$ and [Wu20, Lemma 4], we have

$$H^1(X_0, \mathcal{F}|_{X_0} \otimes (\mathcal{G}^{**}|_{X_0})^*) \cong H^1(X, \mathcal{F} \otimes \mathcal{G}^*).$$

This implies that $\mathcal{E}|_{X_0}$ can be extended to a vector bundle \mathcal{E}' on X . We obtain $\mathcal{E} = \mathcal{E}'$ by reflexivity. \square

We prove Theorem 1.6 (2) by using Theorems 5.2, 5.3.

Theorem 5.2 ([Miy87, Theorem 6.1]). *Let X be a smooth projective variety of dimension $n \geq 2$, let \mathcal{E} be a torsion-free coherent sheaf, and let A_1, \dots, A_{n-2} be ample line bundles on X . Assume that $c_1(\mathcal{E})$ is nef and $\mu_{A_1 \cdots A_{n-2} L}^{\min}(\mathcal{E}) \geq 0$ holds for any nef line bundle L . Then, we have*

$$c_2(\mathcal{E})A_1 \cdots A_{n-2} \geq 0.$$

Theorem 5.3 ([Ou17, Chapter 6]). *Let X be a smooth projective variety of dimension $n \geq 2$, let \mathcal{E} be a torsion-free sheaf, and let A be an ample line bundle. Assume that $c_1(\mathcal{E})$ is nef, \mathcal{E} is generically nef, and $c_2(\mathcal{E})A^{n-2} = 0$. Then, we have:*

- (1) *In the case of $\nu(c_1(\mathcal{E})) \geq 2$, there exists a line bundle $\mathcal{E}_1 \subset \mathcal{E}$ such that $\mathcal{E}/\mathcal{E}_1$ is torsion-free, $c_1(\mathcal{E}_1) = c_1(\mathcal{E})$, $c_1(\mathcal{E}/\mathcal{E}_1) = 0$, and $c_2(\mathcal{E}/\mathcal{E}_1)A^{n-2} = 0$.*
- (2) *In the case of $\nu(c_1(\mathcal{E})) = 1$, there exists a filtration of subsheaves:*

$$0 =: \mathcal{E}_0 \subset \mathcal{E}_1 \subset \cdots \subset \mathcal{E}_l := \mathcal{E}$$

satisfying the following conditions:

- *The above filtration is the $(c_1(\mathcal{E}) + \varepsilon A)^{n-1}$ -Harder-Narasimhan filtration for any $0 < \varepsilon \ll 1$.*
 - *$c_1(\mathcal{E}_k/\mathcal{E}_{k-1})$ is numerically proportional to $c_1(\mathcal{E})$ for any $k = 1, \dots, l$.*
 - *$c_2(\mathcal{E}_k/\mathcal{E}_{k-1})(c_1(\mathcal{E}) + \varepsilon A)^{n-2} = 0$ for any $k = 1, \dots, l$ and any $0 < \varepsilon \ll 1$.*
- (3) *In the case of $\nu(c_1(\mathcal{E})) = 0$, the reflexive hull \mathcal{E}^{**} is a numerically flat vector bundle.*

Remark 5.4. We should pay attention to the choice of polarizations when considering generically nef sheaves in Theorems 5.3 and 5.7. For example, we assume that \mathcal{E} is generically nef (with respect to any ample polarizations) in Theorem 5.3, and thus \mathcal{E} is generically (H_1, \dots, H_{n-2}) -semipositive by Theorem 5.2, which was an assumption in [Ou17, Proposition 6.1]. On the other hand, in Theorem 5.7, we only assume that

\mathcal{E} is $\{\eta\}^{n-1}$ -generically nef for a Kähler form η , but we consider only the polarization of the form on $(c_1(\mathcal{E}) + \varepsilon\{\omega\})$ in Theorems 5.7 or 1.6 (1). See Subsection 6.2 for related open problems.

Proof of Theorem 1.6 (2). We first remark that F is nef if a vector bundle F is projectively numerically flat and $c_1(F)$ is nef. This because $F\langle\frac{\det F^*}{\text{rk}F}\rangle$ is nef. The proof is divided by the cases according to $\nu(c_1(\mathcal{E}))$.

Case of $\nu(c_1(\mathcal{E})) \geq 2$. By Theorem 5.3, there exists a nef line bundle \mathcal{E}_1 such that $\mathcal{E}_2 := \mathcal{E}/\mathcal{E}_1$ is torsion-free, $c_1(\mathcal{E}_2) = 0$, and $c_2(\mathcal{E}_2)A^{n-2} = 0$. Then, from Theorem 5.2, we obtain $c_2(\mathcal{E}_2^{**})A^{n-2} \geq 0$ since \mathcal{E}_2^{**} is generically nef. This implies that

$$0 = c_2(\mathcal{E}_2)A^{n-2} \geq c_2(\mathcal{E}_2^{**})A^{n-2} \geq 0,$$

and thus $c_2(\mathcal{E}_2^{**})A^{n-2} = 0$. Hence, we see that \mathcal{E}_2^{**} is a projectively numerically flat vector bundle by Theorem-Definition 2.2. By [Cao13, Remark 8] or [Ou17, Lemma 6.9], we have $\text{codim}(\text{Supp}(\mathcal{E}_2^{**}/\mathcal{E}_2)) \geq 3$. By Lemma 5.1, the sheaf \mathcal{E} is locally free and satisfies the exact sequence

$$0 \rightarrow \mathcal{E}_1 \rightarrow \mathcal{E} \rightarrow \mathcal{E}_2^{**} \rightarrow 0.$$

This implies that

$$c_2(\mathcal{E}) = c_1(\mathcal{E}_1)c_1(\mathcal{E}_2^{**}) + c_2(\mathcal{E}_2^{**}) + c_2(\mathcal{E}_1) = 0.$$

Case of $\nu(c_1(\mathcal{E})) = 1$. We use induction on the rank of \mathcal{E} . We show that the subsheaf $\mathcal{E}_1 \subset \mathcal{E}$ appearing in Theorem 5.3 is actually a nef vector bundle. Note that \mathcal{E}_1 is reflexive since \mathcal{E}_1 is saturated in \mathcal{E} . The sheaf \mathcal{E}_1 is $(c_1(\mathcal{E}) + \varepsilon A)^{n-1}$ -semistable and satisfies $c_2(\mathcal{E}_1)(c_1(\mathcal{E}) + \varepsilon A)^{n-2} = 0$. Furthermore, we have $c_1(\mathcal{E}_1)^2 = 0$ since $c_1(\mathcal{E}_1)$ is numerically proportional to $c_1(\mathcal{E})$. Hence, the sheaf \mathcal{E}_1 is a projectively numerically flat vector bundle and satisfies

$$c_2(\mathcal{E}_1) = \frac{\text{rk}\mathcal{E}_1 - 1}{2\text{rk}\mathcal{E}_1} c_1(\mathcal{E}_1)^2 = 0.$$

We conclude that \mathcal{E}_1 is a nef vector bundle.

The quotient sheaf $\mathcal{E}_2 := \mathcal{E}/\mathcal{E}_1$ is a generically nef sheaf that satisfies that $c_1(\mathcal{E}_2)$ is nef, $c_1(\mathcal{E}_2)^2 = 0$, and $c_2(\mathcal{E}_2)(c_1(\mathcal{E}) + \varepsilon A)^{n-2} = 0$. By the same argument as above, we have $\text{codim}(\text{Supp}(\mathcal{E}_2^{**}/\mathcal{E}_2)) \geq 3$. The induction hypothesis shows that \mathcal{E}_2^{**} is also a nef vector bundle with $c_2(\mathcal{E}_2^{**}) = 0$.

The sheaf \mathcal{E} is locally free by Lemma 5.1. Furthermore, we have the following exact sequence:

$$0 \rightarrow \mathcal{E}_1 \rightarrow \mathcal{E} \rightarrow \mathcal{E}_2^{**} \rightarrow 0.$$

Therefore, \mathcal{E} is nef since both \mathcal{E}_1 and \mathcal{E}_2 are nef. Since both $c_1(\mathcal{E}_1)$ and $c_1(\mathcal{E}_2^{**})$ are numerically proportional to $c_1(\mathcal{E})$, we have

$$c_2(\mathcal{E}) = c_2(\mathcal{E}_1) + c_1(\mathcal{E}_1)c_1(\mathcal{E}_2^{**}) + c_2(\mathcal{E}_2^{**}) = 0.$$

Case of $\nu(c_1(\mathcal{E})) = 0$. The sheaf $\mathcal{E} = \mathcal{E}^{**}$ is a numerically flat vector bundle by Theorem 5.3, which finishes the proof. \square

Remark 5.5. By the proof of Theorem 1.6 (2), if \mathcal{E} is reflexive, $c_1(\mathcal{E})$ is nef, if \mathcal{E} is generically nef, and if $c_2(\mathcal{E})A^{n-2} = 0$, then there exists a filtration of subbundles:

$$0 =: E_0 \subset E_1 \subset \cdots \subset E_l := \mathcal{E}$$

such that E_i/E_{i-1} is projectively numerically flat and $c_1(E_i/E_{i-1})$ is numerically proportional to $c_1(\mathcal{E})$. In particular, each quotient E_i/E_{i-1} is nef, and E is an extension of projectively numerically flat bundles.

Theorem 1.6 (2) leads to the following corollary.

Corollary 5.6. *Let X be a smooth projective variety, and let A be an ample line bundle on X .*

- (1) *If K_X is nef and $c_2(X)A^{n-2} = 0$, then Ω_X is nef and $c_2(X) = 0$ in $H^{2,2}(X, \mathbb{R})$.*
- (2) *If $-K_X$ is nef and $c_2(X)A^{n-2} = 0$, then T_X is nef and $c_2(X) = 0$ in $H^{2,2}(X, \mathbb{R})$.*

Proof. By [Miy87] (cf. [CPe11, Theorem 0.3] or [CPă19, Theorem 1.3]), the cotangent bundle Ω_X is generically nef. Hence, Conclusion (1) follows from Theorem 1.6. By [Ou17, Theorem 1.4], the tangent bundle T_X is also generically nef. Hence, Conclusion (2) follows from Theorem 1.6. \square

5.2. Case of compact Kähler manifolds. Theorems 5.2 and 5.3 can be generalized to compact Kähler manifolds. See Section 6 for some remarks for the Chern classes of coherent sheaves on compact Kähler manifolds. The proof is essentially the same as in [Cao13]; hence, we put the proof in Appendix 6. By using Theorem 5.7 instead of Theorem 5.3, we can repeat the same argument as in Theorem 1.6 (2) for compact Kähler manifolds, which proves Theorem 1.6 (1). Hence, we omit the proof.

Theorem 5.7. (cf. [Cao13, Proposition 4.4 and Lemma 4.5]) *Let (X, ω) be a compact Kähler manifold of dimension $n \geq 2$, and let \mathcal{E} be a torsion-free coherent sheaf. Assume that $c_1(\mathcal{E})$ is nef and \mathcal{E} is $\{\eta\}^{n-1}$ -generically nef for any Kähler form η . Then, $c_2(\mathcal{E})(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} \geq 0$ holds for any $0 < \varepsilon \ll 1$.*

Moreover, there exists ε_0 depending on (X, ω) and \mathcal{E} such that if $c_2(\mathcal{E})(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} = 0$ for some $\varepsilon_0 > \varepsilon > 0$, then the following holds:

- (1) *In the case of $\nu(c_1(\mathcal{E})) \geq 2$, there exists a line bundle $\mathcal{E}_1 \subset \mathcal{E}$ such that $\mathcal{E}/\mathcal{E}_1$ is torsion-free, $c_1(\mathcal{E}_1) = c_1(\mathcal{E})$, $c_1(\mathcal{E}/\mathcal{E}_1) = 0$, and $c_2(\mathcal{E}/\mathcal{E}_1)(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} = 0$.*
- (2) *In the case of $\nu(c_1(\mathcal{E})) = 1$, there exists a filtration of torsion-free coherent sheaves:*

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \cdots \subset \mathcal{E}_l = \mathcal{E}$$

which satisfies the following conditions.

- *This filtration is the $(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-1}$ -Harder-Narasimhan filtration.*
 - *$c_1(\mathcal{E}_k/\mathcal{E}_{k-1})$ is nef and numerically proportional to $c_1(\mathcal{E})$ for any $k = 1, \dots, l$.*
 - *$c_2(\mathcal{E}_k/\mathcal{E}_{k-1})(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} = 0$ for any $k = 1, \dots, l$.*
- (3) *In the case of $\nu(c_1(\mathcal{E})) = 0$, the reflexive hull \mathcal{E}^{**} is a numerically flat vector bundle.*

Corollary 5.8. *Let X be a compact Kähler manifold and let ω be a Kähler form. We take $\varepsilon_0 > 0$ as in Theorem 5.7.*

- (1) *If K_X is nef and $c_2(X)(c_1(K_X) + \varepsilon\{\omega\})^{n-2} = 0$ for some $\varepsilon_0 > \varepsilon > 0$, then Ω_X^1 is nef and $c_2(X) = 0$ in $H^{2,2}(X, \mathbb{R})$.*
- (2) *If $-K_X$ is nef and $c_2(X)(c_1(-K_X) + \varepsilon\{\omega\})^{n-2} = 0$ for some $\varepsilon_0 > \varepsilon > 0$, then T_X is nef and $c_2(X) = 0$ in $H^{2,2}(X, \mathbb{R})$.*

Proof. By [Cao13, Theorem 1.2] or [Eno93, Theorem 1.4], the cotangent bundle Ω_X is $\{\eta\}^{n-1}$ -generically nef for any Kähler form η . Hence, Conclusion (1) follows from Theorem 1.6. By [Cao13, Theorem 1.2], the tangent bundle T_X is $\{\eta\}^{n-1}$ -generically nef for any Kähler form η . Hence, Conclusion (2) follows from Theorem 1.6. \square

We prove Corollary 1.7 by using the following proposition:

Proposition 5.9. *If X is a Fano manifold of dimension $n \geq 2$, then $c_2(X) \neq 0$ in $H^{2,2}(X, \mathbb{R})$.*

Proof. Assume that $c_2(X) = 0$ in $H^{2,2}(X, \mathbb{R})$. Then, we have $c_2(X)c_1(-K_X)^{n-2} = 0$ and $\nu(c_1(-K_X)) = n \geq 2$. By the proof of Theorem 1.6 (2), there exists a quotient bundle $T_X \rightarrow E_2 \rightarrow 0$ of rank $\dim X - 1$ such that E_2 is a numerically flat vector bundle. The vector bundle E_2 is trivial since X is simply connected. This implies that $h^0(X, \Omega_X) \geq \dim X - 1 > 0$, which contradicts the rational connectedness of X . \square

Proof of Corollary 1.7. Since T_X is nef by Corollary 5.8, there exists a finite étale morphism $\pi : X' \rightarrow X$ and a smooth morphism $\alpha : X' \rightarrow Y'$ such that Y' is a torus and any fiber F of f is Fano by [DPS94, Main Theorem]. By considering the restriction of the standard exact sequence of cotangent bundles to F , we obtain $c_2(F) = 0$ by $c_2(X') = 0$. Since the fiber F is Fano, we conclude that $\dim F \leq 1$ by Proposition 5.9. If $\dim F = 0$, then $X' \cong Y'$ is a complex torus. If $\dim F = 1$, then $F \cong \mathbb{P}^1$. Then $\alpha : X' \rightarrow Y$ is a \mathbb{P}^1 -fiber bundle. \square

If Ω_X is nef and $c_2(X) = 0$, then $K_X = \det \Omega_X$ is semi-ample by Theorem 1.3. However, the following example shows that this is not true when replacing Ω_X with T_X .

Example 5.10. Let C be an elliptic curve, and let D be a divisor on C such that $\deg D = 0$ and D is not a torsion element in $\text{Cl}(X)$. Set $X := \mathbb{P}(\mathcal{O}_C \oplus \mathcal{O}_C(-D))$. We denote by $\phi : X \rightarrow C$ the natural projection. Then, the anticanonical bundle $-K_X = -K_{X/C}$ is nef but not semi-ample (for example, see [EIM21, Example 6.2]). From $c_1(\phi^*K_C) = 0$, we obtain $c_2(X) = c_1(K_{X/C})c_1(\phi^*K_C) = 0$.

6. APPENDIX

6.1. Proof of Theorem 5.7. In this subsection, we prove Theorem 5.7. The proof is essentially the same as in [Cao13], but we provide the details for the reader's convenience.

We first give some remarks for the Chern classes of coherent sheaves on compact Kähler manifolds. By [Gri10, Theorem 1 and Corollary 1], for a coherent sheaf \mathcal{F} on a compact Kähler manifold X , we can define the Chern class $c_i(\mathcal{F}) \in H^{2i}(X, \mathbb{Q})$,

which is compatible with the rational topological Chern class. Thus, as in [CHP16, Definition 7.1], we can define the intersection number $c_2(\mathcal{F})\{\omega\}^{n-2}$ for a Kähler form ω on X . Further we can use this intersection number in the same way as in the case of projective manifolds thanks to results proved in [CHP16, Chapter 7].

We now check an elementary result used in [Ou17, Lemma 6.7] and [Cao13].

Lemma 6.1. *Let m be a positive integer. We define the quadratic form by*

$$q(x, y) := x_0y_0 - (x_1y_1 + \cdots + x_my_m)$$

for $x = (x_0, x_1, \dots, x_m), y = (y_0, y_1, \dots, y_m) \in \mathbb{R}^{m+1}$. Then, we have:

- (1) If $q(x, x) \geq 0$, then $q(x, x)q(y, y) \leq q(x, y)^2$.
- (2) If $q(x, x) > 0$ and $q(x, y) = q(y, y) = 0$, then $y = 0$.
- (3) If $q(x, x) = q(x, y) = q(y, y) = 0$, then x, y are linearly dependent.
- (4) If $x \neq 0$ and $q(x, y) = q(x, x) = 0$, then $q(y, y) \leq 0$.

Proof. Set $F(t) := q(x + ty, x + ty) = q(x, x) + 2tq(x, y) + t^2q(y, y)$.

(1). We may assume that $q(x, x) > 0$ and $q(y, y) > 0$. From $y_0 \neq 0$, we have $F(0) > 0$ and $F(-\frac{x_0}{y_0}) \leq 0$. Thus, $q(x, y)^2 \geq q(x, x)q(y, y)$ since $F(t) = 0$ has a real solution.

(2). If $y_0 \neq 0$, then $0 \geq F(-\frac{x_0}{y_0}) = F(0) > 0$, which is impossible. Thus, $y_0 = 0$. From $q(y, y) = 0$, we have $y = 0$.

(3). We may assume that $y \neq 0$. From $F(-\frac{x_0}{y_0}) = 0$, we have $x - \frac{x_0}{y_0}y = 0$.

(4). The proof follows from $q(y, y) = q(y - \frac{y_0}{x_0}x, y - \frac{y_0}{x_0}x) \leq 0$. \square

Let X be a compact Kähler manifold of dimension $n \geq 2$, and let ω be a Kähler form. We define $q(\eta_1, \eta_2) = \{\eta_1\}\{\eta_2\}\{\omega\}^{n-2}$ for $\eta_1, \eta_2 \in H^{1,1}(X, \mathbb{R})$. Then, the signature of q is $(1, h^{1,1}(X, \mathbb{R}) - 1)$ (for example, see [Voi02]). Hence, we can apply Lemma 6.1 to $q(\eta_1, \eta_2)$.

Proof of Theorem 5.7. The conclusion is obvious in the case of $\text{rk}\mathcal{E} = 1$. Hence, we may assume that $\text{rk}\mathcal{E} \geq 2$.

Set $\nu := \nu(c_1(\mathcal{E}))$ and $\alpha_\varepsilon := c_1(\mathcal{E}) + \varepsilon\{\omega\}$ for any $\varepsilon > 0$. By [Cao13, Proposition 2.3], if $\varepsilon > 0$ is small enough, the α_ε^{n-1} -Harder Narasimhan filtration

$$0 =: \mathcal{E}_0 \subset \mathcal{E}_1 \subset \cdots \subset \mathcal{E}_l := \mathcal{E}$$

is independent of ε . Set $\mathcal{G}_i := \mathcal{E}_i/\mathcal{E}_{i-1}$ and $r_i := \text{rk}(\mathcal{G}_i)$. Since \mathcal{E} is $\{\eta\}^{n-1}$ -generically nef for any Kähler form η , we have $\mu_{\alpha_\varepsilon}(\mathcal{G}_i) \geq 0$. The sheaf \mathcal{G}_i is an α_ε^{n-1} -semistable sheaf, and thus the Bogomolov-Gieseker inequality yields

$$\begin{aligned} 2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} &= \left(c_1(\mathcal{E})^2 + \sum_{1 \leq i \leq l} (2c_2(\mathcal{G}_i) - c_1(\mathcal{G}_i)^2) \right) \alpha_\varepsilon^{n-2} \\ (6.1) \quad &\geq \left(c_1(\mathcal{E})^2 - \sum_{1 \leq i \leq l} \frac{c_1(\mathcal{G}_i)^2}{r_i} \right) \alpha_\varepsilon^{n-2}. \end{aligned}$$

Claim 6.2. $c_1(\mathcal{G}_i)c_1(\mathcal{E})^\nu\{\omega\}^{n-1-\nu} = 0$ holds if $\nu \leq n - 1$.

Proof of Claim 6.2. We have $c_1(\mathcal{G}_i)c_1(\mathcal{E})^\nu\{\omega\}^{n-1-\nu} \geq 0$ from

$$0 \leq \mu_{\alpha_\varepsilon}(\mathcal{G}_i) = \binom{n-1}{\nu} (c_1(\mathcal{G}_i)c_1(\mathcal{E})^\nu\{\omega\}^{n-1-\nu}) \varepsilon^{n-1-\nu} + O(\varepsilon^{n-\nu}).$$

The sum of them can be computed as follows:

$$\sum_{1 \leq i \leq l} c_1(\mathcal{G}_i)c_1(\mathcal{E})^\nu\{\omega\}^{n-1-\nu} = \sum_{1 \leq i \leq l} (c_1(\mathcal{E}_i) - c_1(\mathcal{E}_{i-1}))c_1(\mathcal{E})^\nu\{\omega\}^{n-1-\nu} = c_1(\mathcal{E})^{\nu+1}\{\omega\}^{n-1-\nu}.$$

The right-hand side is zero by the definition of the numerical dimension. This completes the proof. \square

The proof is divided into three cases: $\nu \geq 2$, $\nu = 1$, and $\nu = 0$.

Case of $\nu \geq 2$. (6.1) shows that $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} > 0$ holds if $l = 1$. Hence, we may assume that $l \geq 2$. Set $a_i := c_1(\mathcal{G}_i)c_1(\mathcal{E})^{\nu-1}\{\omega\}^{n-\nu}$. Then, we have

$$\mu_{\alpha_\varepsilon}(\mathcal{G}_i) = \binom{n-1}{\nu-1} \frac{a_i}{r_i} \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}).$$

From $\mu_{\alpha_\varepsilon}(\mathcal{G}_1) > \cdots > \mu_{\alpha_\varepsilon}(\mathcal{G}_l) \geq 0$, we obtain $a_1/r_1 \geq a_2/r_2 \geq \cdots \geq a_l/r_l \geq 0$ for sufficiently small $\varepsilon > 0$.

Claim 6.3. The following estimate holds:

$$2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq \binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right)^{-1} \left(\sum_{1 \leq i \leq l} \frac{(r_i-1)a_i^2}{r_i} + \sum_{1 \leq i < j \leq l} 2a_i a_j \right) \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}).$$

In particular, if there exists $a > 0$ such that

$$\sum_{1 \leq i \leq l} \frac{(r_i-1)a_i^2}{r_i} + \sum_{1 \leq i < j \leq l} 2a_i a_j \geq a + O(\varepsilon),$$

then $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} > 0$ for sufficiently small $\varepsilon > 0$.

Proof of Claim 6.3. By Lemma 6.1, we have

$$(6.2) \quad c_1(\mathcal{G}_i)^2 \alpha_\varepsilon^{n-2} \leq \frac{(c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2})^2}{c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2}}.$$

Hence, we obtain the following equalities:

$$\begin{aligned}
 c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} &= \binom{n-2}{\nu-2} a_i \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}), \\
 c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2} &= \sum_{1 \leq k \leq l} c_1(\mathcal{G}_k)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} \\
 &= \binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right) \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}), \\
 (c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2})^{-1} &= \left(\binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right) \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}) \right)^{-1} \\
 &= \frac{\varepsilon^{-n+\nu}}{\binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right)} + O(\varepsilon^{-n+\nu+1}).
 \end{aligned}$$

From (6.1) and (6.2), we obtain

$$\begin{aligned}
 2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} &\geq c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2} - \sum_{1 \leq i \leq l} \frac{(c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2})^2}{r_i c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2}} \\
 &= \binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right)^{-1} \left(\sum_{1 \leq i \leq l} a_i \sum_{1 \leq j \leq l} a_j - \sum_{1 \leq i \leq l} \frac{a_i^2}{r_i} \right) \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}) \\
 &= \binom{n-2}{\nu-2} \left(\sum_{1 \leq k \leq l} a_k \right)^{-1} \left(\sum_{1 \leq i \leq l} \left(1 - \frac{1}{r_i} \right) a_i^2 + \sum_{1 \leq i < j \leq l} 2a_i a_j \right) \varepsilon^{n-\nu} + O(\varepsilon^{n-\nu+1}).
 \end{aligned}$$

□

If $a_2 > 0$ or $r_1 \neq 1$, we can find that there exists $a > 0$ such that

$$\sum_{1 \leq i \leq l} \frac{(r_i - 1)a_i^2}{r_i} + \sum_{1 \leq i < j \leq l} 2a_i a_j \geq a + O(\varepsilon)$$

from $a_1/r_1 \geq a_2/r_2 \geq \dots \geq a_l/r_l \geq 0$. Hence, by Claim 6.3, we can conclude that $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} > 0$ for sufficiently small $\varepsilon > 0$.

From now on, we may assume that $a_2 = 0$ and $r_1 = 1$. Then, from (6.1), we obtain

$$\begin{aligned}
 (6.3) \quad 2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} &\geq 2 \sum_{2 \leq i \leq l} c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} - \left(\sum_{2 \leq i \leq l} c_1(\mathcal{G}_i) \right)^2 \alpha_\varepsilon^{n-2} - \sum_{2 \leq i \leq l} \frac{c_1(\mathcal{G}_i)^2 \alpha_\varepsilon^{n-2}}{r_i}.
 \end{aligned}$$

Let us consider the case where $c_1(\mathcal{G}_2)c_1(\mathcal{E})^{\nu-t}\{\omega\}^{n-1-\nu+t} \neq 0$ holds for some $t \in \{2, \dots, \nu-1\}$. Then, we take the minimal number $s \in \{2, \dots, \nu-1\}$ such that $c_1(\mathcal{G}_2)c_1(\mathcal{E})^{\nu-s}\{\omega\}^{n-1-\nu+s} \neq 0$, and set $b_i := c_1(\mathcal{G}_i)c_1(\mathcal{E})^{\nu-s}\{\omega\}^{n-1-\nu+s}$ for any $i =$

$2, \dots, l$. Then, since we have

$$c_1(\mathcal{G}_i)\alpha_\varepsilon^{n-1} = \binom{n-1}{\nu-s} b_i \varepsilon^{n-\nu+s-1} + O(\varepsilon^{n-\nu+s}) \geq 0,$$

we obtain $b_2 > 0$. By Lemma 6.1, we obtain the following equalities:

$$\begin{aligned} c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} &= \binom{n-2}{\nu-s-1} b_i \varepsilon^{n-\nu+s-1} + O(\varepsilon^{n-\nu+s}), \\ \left(\sum_{2 \leq i \leq l} c_1(\mathcal{G}_i) \right)^2 \alpha_\varepsilon^{n-2} &\leq \frac{(\sum_{2 \leq i \leq l} c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2})^2}{c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2}} \\ &= \frac{\binom{n-1}{\nu-s-1}^2 (\sum_{2 \leq i \leq l} b_i)^2}{\binom{n-2}{\nu-2} a_1} \varepsilon^{n-\nu+2s-2} + O(\varepsilon^{n-\nu+2s-1}), \\ c_1(\mathcal{G}_i)^2 \alpha_\varepsilon^{n-2} &\leq \frac{(c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2})^2}{c_1(\mathcal{E})^2 \alpha_\varepsilon^{n-2}} = \frac{\binom{n-2}{\nu-s-1}^2 b_i^2}{\binom{n-2}{\nu-2} a_1} \varepsilon^{n-\nu+2s-2} + O(\varepsilon^{n-\nu+2s-1}). \end{aligned}$$

From $s-1 > 0$ and (6.3),

$$2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq 2 \binom{n-2}{\nu-s-1} \left(\sum_{2 \leq i \leq l} b_i \right) \varepsilon^{n-\nu+s-1} + O(\varepsilon^{n-\nu+s}).$$

This implies that $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} > 0$ for sufficiently small $\varepsilon > 0$.

From now on, we may assume that $c_1(\mathcal{G}_2)c_1(\mathcal{E})\nu^{-t}\{\omega\}^{n-1-\nu+t} = 0$ for any $t = 2, \dots, \nu-1$. Since $c_1(\mathcal{G}_i)c_1(\mathcal{E})^t\{\omega\}^{n-1-t} = 0$ for any $t = 1, \dots, n-1$ and $i = 2, \dots, l$, we have

$$(6.4) \quad c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} = c_1(\mathcal{G}_i)c_1(\mathcal{E})(c_1(\mathcal{E}) + \varepsilon\{\omega\})^{n-2} = 0.$$

From $c_1(\mathcal{E})^2\alpha_\varepsilon^{n-2} > 0$, we obtain $c_1(\mathcal{G}_i)^2\alpha_\varepsilon^{n-2} \leq 0$ by Lemma 6.1. By the same argument as above, we obtain $(\sum_{2 \leq i \leq l} c_1(\mathcal{G}_i))^2\alpha_\varepsilon^{n-2} \leq 0$, and finally $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq 0$ for sufficiently small $\varepsilon > 0$ from (6.3). Hence, we can take $\varepsilon_0 > 0$ such that $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq 0$ for any $\varepsilon_0 > \varepsilon > 0$.

If $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} = 0$ for some $\varepsilon_0 > \varepsilon > 0$, then $c_1(\mathcal{G})^2\alpha_\varepsilon^{n-2} = 0$, where $\mathcal{G} := \mathcal{E}/\mathcal{E}_1$. By Lemma 6.1, $c_1(\mathcal{G}) = 0$ and $c_1(\mathcal{E}_1) = c_1(\mathcal{E})$. Furthermore, \mathcal{E}_1 is a nef line bundle by $r_1 = 1$. Hence, we obtain that

$$c_2(\mathcal{G})\alpha_\varepsilon^{n-2} = c_2(\mathcal{E})\alpha_\varepsilon^{n-2} - c_1(\mathcal{G})c_1(\mathcal{E}_1)\alpha_\varepsilon^{n-2} = 0.$$

Case of $\nu = 1$. By Claim 6.2, we have

$$(6.5) \quad c_1(\mathcal{G}_i)c_1(\mathcal{E})\alpha_\varepsilon^{n-2} = c_1(\mathcal{E})^2\alpha_\varepsilon^{n-2} = 0.$$

Furthermore, from $c_1(\mathcal{E}) \neq 0$ and Lemma 6.1, we have $c_1(\mathcal{G}_i)^2\alpha_\varepsilon^{n-2} \leq 0$. Together with (6.1), we obtain

$$2c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq - \sum_{1 \leq i \leq l} \frac{c_1(\mathcal{G}_i)^2}{r_i} \alpha_\varepsilon^{n-2} \geq 0.$$

If $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} = 0$ holds for some $\varepsilon > 0$, then $c_1(\mathcal{G}_i)^2\alpha_\varepsilon^{n-2} = 0$. By Lemma 6.1 and (6.5), $c_1(\mathcal{G}_i)$ and $c_1(\mathcal{E})$ are linearly dependent. This implies that $c_1(\mathcal{G}_i)c_1(\mathcal{G}_j)\alpha_\varepsilon^{n-2} = 0$ for any $1 \leq i \leq j \leq l$. Since \mathcal{G}_i is α_ε^{n-1} -semistable, we obtain $c_2(\mathcal{G}_i)\alpha_\varepsilon^{n-2} \geq 0$ from the Bogomolov-Gieseker inequality. Furthermore, from

$$c_2(\mathcal{E})\alpha_\varepsilon^{n-2} = \sum_{1 \leq i \leq l} c_2(\mathcal{G}_i)\alpha_\varepsilon^{n-2} = 0,$$

we obtain $c_2(\mathcal{G}_i)\alpha_\varepsilon^{n-2} = 0$ any $i = 1, \dots, l$.

Case of $\nu = 0$. From $c_1(\mathcal{E}) = 0$, \mathcal{E} is α_ε^{n-1} -semistable. By the Bogomolov-Gieseker inequality, we obtain $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} \geq 0$. If $c_2(\mathcal{E})\alpha_\varepsilon^{n-2} = 0$, then $c_2(\mathcal{E}^{**})\alpha_\varepsilon^{n-2} = 0$. By Theorem-Definition 2.2, the reflexive hull \mathcal{E}^{**} is a numerically projectively flat vector bundle; thus, \mathcal{E}^{**} is numerically flat by $c_1(\mathcal{E}^{**}) = 0$. \square

6.2. Related open problems. In this subsection, we collect some problems related to generically nef vector bundles.

Conjecture 6.4. *Let X be a compact Kähler manifold, let \mathcal{E} be a torsion-free coherent sheaf, and let $\alpha_1, \dots, \alpha_{n-2}$ be Kähler classes on X .*

(1) *If $c_1(\mathcal{E})$ is nef and \mathcal{E} is generically nef, then the inequality $c_2(\mathcal{E})\alpha_1 \cdots \alpha_{n-2} \geq 0$ holds.*

(2) *Let β be a nef class. We further assume that \mathcal{E} is $\alpha_1 \cdots \alpha_{n-2}\beta$ -semistable. Then, the following inequality holds:*

$$\left(c_2(\mathcal{E}) - \frac{r-1}{2r} c_1(\mathcal{E})^2 \right) \alpha_1 \cdots \alpha_{n-2} \geq 0.$$

Note that Conjecture 6.4 (2) leads to Conjecture 6.4 (1) by the argument of [Miy87, Theorem 6.1]. Conjecture 6.4 has been affirmatively solved when X is projective and both α_i and β are in the Néron-Severi group of X .

The following conjecture has also been affirmatively solved in the projective case but is still open in the compact Kähler case. This conjecture is more difficult than we expected due to the lack of the foliation theory developed in [CPă19] on compact Kähler manifolds.

Conjecture 6.5 (cf. [Cao13, Conjecture 4.3]). *Let X be a compact Kähler manifold, and let $\alpha_1, \dots, \alpha_{n-1}$ be Kähler classes.*

(1) *If K_X is nef, then Ω_X is $\alpha_1 \cdots \alpha_{n-1}$ -generically nef.*

(2) *If $-K_X$ is nef, then T_X is $\alpha_1 \cdots \alpha_{n-1}$ -generically nef.*

REFERENCES

- [AMN12] J. Amorós, M. Manjarín, M. Nicolau. *Deformations of Kähler manifolds with nonvanishing holomorphic vector fields*. J. Eur. Math. Soc. (JEMS) **14** (2012), no. 4, 997–1040.
- [BP08] B. Berndtsson, M. Păun. *Bergman kernels and the pseudoeffectivity of relative canonical bundles*. Duke Math. J. **145** (2008), no. 2, 341–378.
- [CCE15] F. Campana, B. Claudon, P. Eyssidieux. *Représentations linéaires des groupes kählériens: factorisations et conjecture de Shafarevich linéaire*. (French) *Linear representations of Kähler groups: factorizations and linear Shafarevich conjecture*. Compos. Math. **151** (2015), no. 2, 35–376.
- [CHP16] F. Campana, A. Höring, T. Peternell. *Abundance for Kähler threefolds*. Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), no. 4, 971–1025.

- [CPă19] F. Campana, M. Păun. *Foliations with positive slopes and birational stability of orbifold cotangent bundles*. Publ. Math. Inst. Hautes Études Sci. **129** (2019), 1–49.
- [CPe11] F. Campana, T. Peternell. *Geometric stability of the cotangent bundle and the universal cover of a projective manifold*. Bull. Soc. Math. France **139** (2011), no. 1, 41–74.
- [Cam94] F. Campana. *Remarques sur le revêtement universel des variétés kählériennes compactes*. Bulletin de la Société Mathématique de France, Tome **122** (1994) no. 2, pp. 255–284.
- [Cam04] F. Campana. *Orbifolds, special varieties and classification theory*. Ann. Inst. Fourier (Grenoble) **54** (2004), no. 3, 499–630.
- [Cam11] F. Campana. *Special orbifolds and birational classification: a survey*. Classification of algebraic varieties, 123–170, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, (2011).
- [Cao13] J. Cao. *A remark on compact Kähler manifolds with nef anticanonical bundles and its applications*. Preprint, arXiv:1305.4397v2.
- [DHP13] J.-P. Demailly, C. D. Hacon, M. Păun. *Extension theorems, non-vanishing and the existence of good minimal models*. Acta Math. **210** (2013), no. 2, 203–259.
- [DPS94] J.-P. Demailly, T. Peternell, M. Schneider. *Compact complex manifolds with numerically effective tangent bundles*. J. Algebraic Geom., **3**, (1994), no.2, 295–345.
- [DPS01] J.-P. Demailly, T. Peternell, M. Schneider. *Pseudo-effective line bundles on compact Kähler manifolds*. Internat. J. Math. **12** (2001), no. 6, 689–741.
- [EIM21] S. Ejiri, M. Iwai, S. Matsumura. *On asymptotic base loci of relative anti-canonical divisors of algebraic fiber spaces*. Preprint, arXiv:2005.04566v1.
- [Eno93] I. Enoki. *Kawamata-Viehweg vanishing theorem for compact Kähler manifolds*. Einstein metrics and Yang-Mills connections (Sanda, 1990), 59–68, Lecture Notes in Pure and Appl. Math., **145**, Dekker, New York, (1993).
- [Fuj03] O. Fujino. *A canonical bundle formula for certain algebraic fiber spaces and its applications*. Nagoya Math. J. **172** (2003), 129–171.
- [Gri10] J. Grivaux. *Chern classes in Deligne cohomology for coherent analytic sheaves*. Math. Ann. **347** (2010), no. 2, 249–284.
- [GM17] Y. Gongyo, S. Matsumura. *Versions of injectivity and extension theorems*. Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 2, 479–502.
- [GKP16] D. Greb, S. Kebekus, T. Peternell. *Étale fundamental groups of Kawamata log terminal spaces, flat sheaves, and quotients of abelian varieties*. Duke Math. J. **165** (2016), no. 10, 1965–2004.
- [GKP21] D. Greb, S. Kebekus, T. Peternell. *Projectively flat KLT varieties*. Journal de l’Ecole Polytechnique - Mathématiques **8** (2021), 1005–1036.
- [HPS18] C. Hacon, M. Popa, C. Schnell. *Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Păun*. Local and global methods in algebraic geometry, 143–195, Contemp. Math., **712**, Amer. Math. Soc., Providence, RI, 2018.
- [Har77] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics **52** (1977).
- [Hör13] A. Höring. *Manifolds with nef cotangent bundle*. Asian J. Math. **17** (2013), no. 3, 561–568.
- [HIM21] G. Hosono, M. Iwai, S. Matsumura. *On projective manifolds with pseudo-effective tangent bundle*. to appear in J. Inst. Math. Jussieu, DOI: <https://doi.org/10.1017/S1474748020000754>.
- [Iwa21] M. Iwai. *Almost nef regular foliations and Fujita’s decomposition of reflexive sheaves*. To appear in Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), DOI: https://doi.org/10.2422/2036-2145.202010_055.
- [Kaw92] Y. Kawamata. *Abundance theorem for minimal threefolds*. Invent. Math. **108** (1992), no. 2, 229–246.
- [Kob87] S. Kobayashi. *Differential geometry of complex vector bundles*. Publications of the Mathematical Society of Japan, **15** Kano Memorial Lectures, **5**. Princeton University Press, Princeton, NJ; Princeton University Press, Princeton, NJ, (1987). xii+305.
- [Kol95] J. Kollár. *Shafarevich maps and automorphic forms*. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, (1995). x+201.

- [KM98] J. Kollár, S. Mori. *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. Cambridge Tracts in Math., Vol. 134. Cambridge: Cambridge Univ. Press, 1998.
- [LP18] V. Lazić, T. Peternell. *Abundance for varieties with many differential forms*. *Épjournal Géom. Algébrique* 2 (2018), Art. 1, 35 pp.
- [Lie78] D. Lieberman. *Compactness of the Chow scheme: applications to automorphisms and deformations of Kähler manifolds*. *Fonctions de plusieurs variables complexes, III (Sém. Francois Norguet, 1975–1977)*, pp. 140–186, Lecture Notes in Math., **670**, Springer, Berlin, (1978).
- [Liu14] G. Liu. *Compact Kähler manifolds with nonpositive bisectional curvature*. *Geom. Funct. Anal.* **24** (2014), no. 5, 1591–1607.
- [LOY20] J. Liu, W. Ou, X. Yang. *Projective manifolds whose tangent bundle contains a strictly nef subsheaf*. Preprint, arXiv:2004.08507
- [LPT18] F. Loray, J. V. Pereira, F. Touzet. *Singular foliations with trivial canonical class*. *Invent. Math.* **213** (2018), no. 3, 1327–1380.
- [Math] Mathoverflow: *Are most Kähler manifolds non-projective?* Available at <https://mathoverflow.net/questions/257147/are-most-k%C3%A4hler-manifolds-non-projective>
- [Mat13] S. Matsumura. *An ampleness criterion with the extendability of singular positive metrics*. *Math. Z.* 273 (2013), no. 1-2, 43–54.
- [Miy87] Y. Miyaoka. *The Chern classes and Kodaira dimension of a minimal variety*. *Algebraic geometry, Sendai, (1985)*, 449-476, *Adv. Stud. Pure Math.*, **10**, North-Holland, Amsterdam, (1987).
- [Miy88a] Y. Miyaoka. *Abundance conjecture for 3-folds: case $\nu=1$* . *Compositio Math.* **68** (1988), no. 2, 203–220.
- [Miy88b] Y. Miyaoka. *On the Kodaira dimension of minimal threefolds*. *Math. Ann.* **281** (1988), no. 2, 325–332.
- [Nak04] N. Nakayama. *Zariski-decomposition and abundance*. *MSJ Memoirs*, **14**, Mathematical Society of Japan, Tokyo, 2004. xiv+277.
- [Ou17] W. Ou. *On generic nefness of tangent sheaves*. Preprint, arXiv:1703.03175.
- [PäT18] M. Păun, S. Takayama. *Positivity of twisted relative pluricanonical divisors and their direct images*. *J. Algebraic Geom.* **27** (2018), 211–272.
- [PeT13] J. V. Pereira, F. Touzet. *Foliations with vanishing Chern classes*. *Bull. Braz. Math. Soc. (N.S.)* **44** (2013), no. 4, 731–754.
- [PRT21] J. V. Pereira, E. Rousseau, F. Touzet. *Numerically nonspecial varieties*. Preprint, arXiv:2106.12275v1.
- [Pet12] T. Peternell. *Varieties with generically nef tangent bundles*. *J. Eur. Math. Soc. (JEMS)* **14** (2012), no. 2, 571–603. MR 2881306
- [Uen78] K. Ueno. *On algebraic fibre spaces of abelian varieties*. *Math. Ann.* **237** (1978), no. 1, 1–22.
- [Voi02] C. Voisin. *Hodge theory and complex algebraic geometry. I*. Translated from the French original by Leila Schneps. *Cambridge Studies in Advanced Mathematics*, 76. Cambridge University Press, Cambridge, 2002. x+322 pp. ISBN: 0-521-80260-1.
- [WZ02] H.-H. Wu, F. Zheng. *Compact Kähler manifolds with nonpositive bisectional curvature*. *J. Differential Geom.* **61** (2002), no. 2, 263–287.
- [Wu20] X. Wu. *Pseudo-effective and numerically flat reflexive sheaves*. Preprint, arXiv:2004.14676v2, to appear in the *Journal of Geometric Analysis*.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, 1-1,
MACHIKANNEYAMA-CHO, TOYONAKA, OSAKA 560-0043, JAPAN.

Email address: masataka@math.sci.osaka-u.ac.jp

Email address: masataka.math@gmail.com

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, 6-3, ARAMAKI AZA-AOBA, AOBA-KU,
SENDAI 980-8578, JAPAN.

Email address: mshinichi-math@tohoku.ac.jp

Email address: mshinichi0@gmail.com