

FAILURE OF STRONG ARENS IRREGULARITY IN GENERALIZED GROUP ALGEBRAS

LAV KUMAR SINGH

ABSTRACT. In this short note, we show that the generalized group algebra $L^1(G, \mathcal{A})$ is not strongly Arens irregular for any infinite discrete group G and any Banach algebra \mathcal{A} with an orthonormal Schauder basis equivalent to the canonical Schauder basis of $\ell^1(\mathbb{N})$. This is interestingly in contrast to the celebrated result by Losert and Lau that the group algebra $L^1(G)$ is strongly Arens irregular for any locally compact group G .

1. INTRODUCTION

For any Banach algebra \mathcal{A} , Richard Arens (in [1]) defined two products \square and \diamond on its bidual space \mathcal{A}^{**} such that each product makes \mathcal{A}^{**} into a Banach algebra and the canonical isometric inclusion $J : \mathcal{A} \rightarrow \mathcal{A}^{**}$ becomes a homomorphism with respect to both the products. A Banach algebra \mathcal{A} is said to be *Arens regular* if the two products \square and \diamond agree on \mathcal{A}^{**} , i.e. $f \square g = f \diamond g$ for all $f, g \in \mathcal{A}^{**}$. Even when a Banach algebra is not Arens regular, the size of its topological center carry a great deal of information. Dales and Lau introduced the notion of strong Arens irregularity to distinguish between non-Arens regular Banach algebras.

Briefly speaking, a Banach algebra \mathcal{A} is said to be *left* (resp., *right*) *strongly Arens irregular* if its so called *left topological center* $Z_t^{(l)}(\mathcal{A}^{**})$ (resp., *right topological center* $Z_t^{(r)}(\mathcal{A}^{**})$) equals \mathcal{A} . And, \mathcal{A} is said to be *strongly Arens irregular* (in short, S.A.I.) if it is both left and right strongly Arens irregular. Interestingly, on the other extreme, it is known that \mathcal{A} is Arens regular if and only if $Z_t^{(l)}(\mathcal{A}^{**}) = \mathcal{A}^{**} = Z_t^{(r)}(\mathcal{A}^{**})$. Reflexive spaces turn out to be at the junction of Arens regularity and strong Arens irregularity because $\mathcal{A} = \mathcal{A}^{**} = Z^{(l)}(\mathcal{A}^{**}) = Z^{(t)}(\mathcal{A}^{**})$.

After the pioneering paper [1] by Richard Arens, people started investigating the Arens regularity of various Banach algebras that appear in abstract Harmonic analysis. The group algebra $L^1(G)$ has been a center of attention for quite some time. Arens himself proved that the semi-group algebra $\ell^1(\mathbb{N})$ (with respect to convolution) is not Arens regular. Graham, in [3], proved very elegantly that the group algebra $L^1(G)$ is not Arens regular for all infinite discrete amenable groups and non-discrete locally compact groups. Finally, Losert and Lau in [5] proved that $L^1(G)$ is strongly Arens irregular for all infinite locally compact groups.

It is then quite natural to analyze the strong Arens irregularity of generalized group algebras $L^1(G, \mathcal{A})$ for infinite locally compact groups G and Banach algebras \mathcal{A} ; and, this short article is just a report on some progress made in this direction. A lot still remains to be explored. Since $L^1(G)$ is not Arens regular, and because $L^1(G, \mathcal{A}) \cong L^1(G) \otimes^\gamma \mathcal{A}$, we know from [9] that $L^1(G, \mathcal{A})$ is also not Arens regular. We wish to study under what conditions on \mathcal{A} , the Banach algebra $L^1(G, \mathcal{A})$ is strongly Arens irregular. Needless to mention, if $\mathcal{A} = \mathbb{C}^n$ with point-wise multiplication and any p -norm and G is infinite, then $L^1(G, \mathcal{A})$ is strongly Arens irregular (due

2010 *Mathematics Subject Classification.* 43A20, 46B28, 46M05, 46H20.

Key words and phrases. Banach algebras, Arens regularity, projective tensor product, Group algebra.

This research work was carried out with the financial support from the Council of Scientific and Industrial Research (Government of India) through a Senior Research Fellowship with No. **09/263(1133)/2017-EMR-I**.

to [5]). Further, since there is an isometric algebra isomorphism from $L^1(G) \otimes^\gamma L^1(H)$ onto $L^1(G \times H)$ (w.r.t. convolution on both the legs) - see [4], it follows again from [5] that $L^1(G, \mathcal{A})$ is strongly Arens irregular for $\mathcal{A} = L^1(H)$, where H is an infinite group. However, on the other extreme, we give a class of Banach algebras \mathcal{A} for which $L^1(G, \mathcal{A})$ is not S.A.I. for any infinite discrete group G . This is interestingly in contrast to the above mentioned fact that the (infinite) group algebra $L^1(G)$ is always S.A.I. - see [5]. Our approach relies on chasing the annihilators in the bidual of $L^1(G, \mathcal{A})$ using appropriate functionals associated to the non-principal ultra-filters on \mathbb{N} .

2. PRELIMINARIES

2.1. Arens regularity and strong Arens irregularity. We quickly recall the definitions of the two products \square and \diamond . Let \mathcal{A} be a Banach algebra. For $a \in \mathcal{A}$, $\omega \in \mathcal{A}^*$, $f \in \mathcal{A}^{**}$, consider the functionals $\omega_a, {}_a\omega \in \mathcal{A}^*$, $\omega_f, {}_f\omega \in \mathcal{A}^{**}$ given by

$$w_a = (L_a)^*(\omega), {}_a\omega = (R_a)^*(\omega); \omega_f(a) = f({}_a\omega) \text{ and } {}_f\omega(a) = f(\omega_a).$$

Then, for $f, g \in \mathcal{A}^{**}$ the operations \square and \diamond are given by

$$(f \square g)(\omega) = f({}_g\omega) \text{ and } (f \diamond g)(\omega) = g(\omega_f)$$

for all $\omega \in \mathcal{A}^*$. And, \mathcal{A} is said to be *Arens regular* if the products \square and \diamond agree.

Further, the left and the right topological centers of \mathcal{A} are defined, respectively, as

$$\begin{aligned} Z_t^{(l)}(\mathcal{A}^{**}) &= \{\varphi \in \mathcal{A}^{**} : \varphi \square \psi = \varphi \diamond \psi \text{ for all } \psi \in \mathcal{A}^{**}\}; \text{ and} \\ Z_t^{(r)}(\mathcal{A}^{**}) &= \{\psi \in \mathcal{A}^{**} : \psi \square \phi = \psi \diamond \phi \text{ for all } \phi \in \mathcal{A}^{**}\}. \end{aligned}$$

Identifying \mathcal{A} with $J(\mathcal{A})$, where $J : \mathcal{A} \rightarrow \mathcal{A}^{**}$ is the canonical isometry, it is known that $\mathcal{A} \subseteq Z_t^{(l)}(\mathcal{A}^{**}) \cap Z_t^{(r)}(\mathcal{A}^{**})$. It is easily seen that \mathcal{A} is Arens regular if and only if $Z_t^{(l)}(\mathcal{A}^{**}) = \mathcal{A}^{**} = Z_t^{(r)}(\mathcal{A}^{**})$. A Banach algebra \mathcal{A} is said to be *left* (resp., *right*) *strongly Arens irregular* if $Z_t^{(l)}(\mathcal{A}^{**}) = \mathcal{A}$ (resp., $Z_t^{(r)}(\mathcal{A}^{**}) = \mathcal{A}$). And, \mathcal{A} is said to be *strongly Arens irregular* if it is both left and right strongly Arens irregular. We refer the reader to [2] for further discussion on topological centers.

2.2. Projective tensor product. Let X and Y be Banach spaces. Then, there are various ways of imposing a normed space structure on their algebraic tensor product $X \otimes Y$. For instance, for each $u \in X \otimes Y$, its projective norm is given by

$$\|u\|_\gamma = \inf \left\{ \sum_{i=1}^n \|x_i\| \|y_i\| : u = \sum_{i=1}^n x_i \otimes y_i \right\}.$$

This norm turns out to be a *cross norm*, i.e., $\|x \otimes y\|_\gamma = \|x\| \|y\|$ for all $(x, y) \in X \times Y$; and, the completion of the normed space $X \otimes Y$ with respect to this norm is usually denoted by $X \otimes^\gamma Y$.

Let \mathcal{A} and \mathcal{B} be Banach algebras. Then, there is a natural multiplication structure on their algebraic tensor product $\mathcal{A} \otimes \mathcal{B}$ satisfying $(a_1 \otimes b_1)(a_2 \otimes b_2) = a_1 a_2 \otimes b_1 b_2$ for all $a_i \in \mathcal{A}$ and $b_i \in \mathcal{B}$, $i = 1, 2$. And, there are various ways of imposing a normed algebra structure on $\mathcal{A} \otimes \mathcal{B}$. For instance, completing $\mathcal{A} \otimes \mathcal{B}$ with respect to the projective norm gives a Banach algebra structure on $\mathcal{A} \otimes^\gamma \mathcal{B}$, which occupies a prominent space in the theory of tensor products of Banach algebras.

2.2.1. *Generalized group algebras.* Let G be a locally compact group and A be a Banach algebra. Then, recall that, $L^1(G, \mathcal{A})$ denotes the space of all Bochner integrable functions from G to \mathcal{A} with the norm $\|f\|_1 := \int_G \|f\| d\mu$ and forms a Banach algebra with respect to the convolution product defined as $f \star g(x) = \int_G f(s)g(s^{-1}x)ds$ - see [4] for further details on generalized group algebras. When G is a discrete group, then $L^1(G, \mathcal{A})$ is nothing but the collection of all absolutely summable \mathcal{A} -valued functions and $\|f\|_1 = \sum_{g \in G} \|f(g)\|$.

The following natural identifications are well known and will be used ahead - see, for instance, [4, 7].

Theorem 2.1. (1) *For any two Banach spaces X and Y , there is a canonical surjective isometry $\theta : B(X, Y^*) \rightarrow (X \otimes^\gamma Y)^*$ satisfying*

$$\theta(f)(x \otimes^\gamma y) = f(x)(y) \text{ for all } f \in B(X, Y^*), x \in X, y \in Y.$$

(2) *For any locally compact group G and Banach algebra A , there exists an isometric algebra isomorphism $T : L^1(G) \otimes^\gamma A \rightarrow L^1(G, A)$ satisfying*

$$T(f \otimes a)(g) = f(g)a \text{ for all } f \in L^1(G), a \in A, g \in G.$$

The following elementary observations will be used ahead. We include their proofs just for the sake of convenience.

Lemma 2.2. *Let G be an infinite discrete group and \mathcal{A} be a Banach algebra. Then,*

- (1) *if $x = \sum_{n \in F} \delta_{g_n} \otimes a_n \in L^1(G) \otimes^\gamma \mathcal{A}$ for some $F \subseteq \mathbb{N}$, a collection $\{g_n : n \in F\} \subseteq G$ with no repetitions and a collection $\{a_n : n \in F\} \subset \mathcal{A}$, then $\|x\|_\gamma = \sum_n \|a_n\|$; and,*
- (2) *for each $x \in L^1(G) \otimes^\gamma \mathcal{A}$, there exists an $F \subset \mathbb{N}$, a collection $\{h_n : n \in F\} \subset G$ with no repetitions and a collection $\{b_n : n \in F\} \subset \mathcal{A} \setminus \{0\}$ such that $x = \sum_{n \in F} \delta_{h_n} \otimes b_n$.*

Proof. By Theorem 2.1(2), we know that $L^1(G) \otimes^\gamma \mathcal{A} \cong L^1(G, \mathcal{A})$ isometrically algebraically.

(1): Identifying x with its corresponding element in $L^1(G, \mathcal{A})$, we see that $x(g) = a_n$ if $g = g_n$ for some n and 0 otherwise; so,

$$\|x\|_\gamma = \|x\|_1 = \sum_{g \in G} \|x(g)\| = \sum_{n \in F} \|x(g_n)\| = \sum_{n \in F} \|a_n\|.$$

(2): Again, identifying x with its corresponding element in $L^1(G, \mathcal{A})$, it is a standard fact that $x : G \rightarrow \mathcal{A}$ has a (possibly finite) countable support, say, $\{h_n : n \in F\}$ for some $F \subset \mathbb{N}$. Let $b_n := x(h_n) \in \mathcal{A}$, $n \in F$. Next, note that, $\|x\|_\gamma = \|x\|_1 = \sum_{g \in G} \|x(g)\| = \sum_{n \in F} \|x(h_n)\|$. Hence, $\{b_n : n \in F\}$ converges absolutely in \mathcal{A} and we also have $x = \sum_{n \in F} \delta_{h_n} \otimes b_n$. \square

3. TOPOLOGICAL CENTER OF GENERALIZED GROUP ALGEBRAS

Throughout this section, G will denote an infinite discrete group.

Recall that, for a Banach space Z with a normalized Schauder basis $\{z_n\}$, the dual functionals $\{z_n^*\}$ are given by $z_n^*(\sum_m \alpha_m z_m) = \alpha_n$ and it is easily seen that each z_n^* is a bounded linear functional on Z . In fact, the collection $\{z_n^*\}$ is uniformly bounded - see [6, Chapter 4].

The following observation is the crux of this article.

Proposition 3.1. *Let \mathcal{A} be a Banach algebra with a (normalized) Schauder basis and let $X := L^1(G, \mathcal{A})$. Then, for each non-principal ultrafilter \mathcal{U} on \mathbb{N} , there exists a $\varphi_{\mathcal{U}} \in X^{**} \setminus J(X)$.*

Proof. Let \mathcal{U} be a non-principal ultrafilter on \mathbb{N} . Fix a countable set $S = \{h_n\}_{n=1}^\infty$ consisting of distinct elements of G .

Notice that, for each $\tau \in X^*$, via the identifications illustrated in Theorem 2.1, we have

$$|\tau(\delta_{h_n} \otimes a_n)| \leq \|\tau\| \text{ for all } n \in \mathbb{N}.$$

Thus, since limit along an ultrafilter in a compact set always exists, we have a map $\varphi_{\mathcal{U}} : X^* \rightarrow \mathbb{C}$ given by

$$(3.1) \quad \varphi_{\mathcal{U}}(\tau) = \lim_{\mathcal{U}} \tau(\delta_{h_n} \otimes a_n) \text{ for all } \tau \in X^*.$$

Clearly, $\varphi_{\mathcal{U}}$ is a bounded linear functional on X^* . Also, $\varphi_{\mathcal{U}} = \lim_{\mathcal{U}}^w J(\delta_{h_n} \otimes a_n)$.

We first assert that $\varphi_{\mathcal{U}} \neq 0$. Consider $\eta : L^1(G) \rightarrow \mathcal{A}^*$ defined as

$$\eta\left(\sum_{g \in G} \alpha_g \delta_g\right) = \sum_{i=1}^{\infty} \alpha_{h_i} a_i.$$

One can easily verify $\eta \in B(L^1(G), \mathcal{A}^*)$ and that $\|\eta\| \leq 1$. Clearly, $\varphi_{\mathcal{U}}(\eta) = 1$. Hence, $\varphi_{\mathcal{U}} \neq 0$.

Next, consider the subset $\Delta := \text{span}\{\mathcal{T}_{g,i} \mid g \in G, i \in \mathbb{N}\}$ of X^* , where $\mathcal{T}_{g,i} : L^1(G) \rightarrow \mathcal{A}^*$ is defined as

$$\mathcal{T}_{g,i}\left(\sum_{h \in G} c_h \delta_h\right) = c_g a_i^*.$$

(One can easily see that $\mathcal{T}_{g,i} \in B(L^1(G), \mathcal{A}^*)$ and are uniformly bounded and, hence, in view of Theorem 2.1, Δ is a well defined non-trivial set in X^* .)

We now claim that J_x does not vanish on $\overline{\Delta}$ for any $0 \neq x \in X$, while $\varphi_{\mathcal{U}}$ vanishes (on Δ and hence) on $\overline{\Delta}$. Since $\varphi_{\mathcal{U}} \neq 0$, this will then imply that $\varphi_{\mathcal{U}} \notin J(X)$.

The latter claim is evident from the fact that $\varphi_{\mathcal{U}}(\mathcal{T}_{g,i}) = 0$ for each $g \in G$ and $i \in \mathbb{N}$.

To prove the first part of the claim, suppose $x \in X \setminus \{0\}$. Then, as observed in Lemma 2.2, there exist countable sets $F \subset \mathbb{N}$, $\{u_j : j \in F\} \subset G$ and $\{b_j : j \in F\} \subset \mathcal{A} \setminus \{0\}$ such that $x = \sum_{j=1}^{\infty} \delta_{u_j} \otimes b_j$.

Note that, each b_j can be written as $b_j = \sum_{n=1}^{\infty} c_n^{(j)} a_n$. So, for each $j \in \mathbb{N}$, we can choose an n_j such that $c_{n_j}^{(j)} \neq 0$. Consider $\mathcal{T} \in X^*$ given by $\mathcal{T} = \sum_{j=1}^{\infty} \frac{1}{j^2} \beta_j \mathcal{T}_{u_j, n_j}$, where $\beta_j = \frac{\overline{c_{n_j}^{(j)}}}{|c_{n_j}^{(j)}|}$. (Note that $\mathcal{T} \in X^*$ because its defining series is absolutely convergent in the norm of X^* .) Clearly, $\mathcal{T} \in \overline{\Delta}$.

Now, notice that

$$\begin{aligned} J_x(\mathcal{T}) &= \mathcal{T}(x) \\ &= \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} \frac{1}{j^2} \beta_j \mathcal{T}_{u_j, n_j}(\delta_{u_i})(b_i) \\ &= \sum_{j=1}^{\infty} \frac{1}{j^2} \beta_j a_{n_j}^*(b_j) \\ &= \sum_{j=1}^{\infty} \frac{1}{j^2} \beta_j c_{n_j}^{(j)} \\ &= \sum_{j=1}^{\infty} \frac{1}{j^2} |c_{n_j}^{(j)}| \neq 0. \end{aligned}$$

Thus, $\varphi_{\mathcal{U}} \in X^{**} \setminus J(X)$. □

The following definition is well known to the experts - see, for instance, [6, Chapter 4].

Definition 3.2. A (normalized) Schauder basis $\{z_n\}$ of a Banach space Z is said to be equivalent to the canonical Schauder basis of $\ell^1(\mathbb{N})$ if there exists a positive scalar C such that

$$\left\| \sum_{i=1}^N \lambda_i z_i \right\| \leq \sum_{i=1}^N |\lambda_i| \leq C \left\| \sum_{i=1}^N \lambda_i z_i \right\|$$

for any set of scalars $\{\lambda_i\}_{i=1}^N$ and $N \in \mathbb{N}$.

Notice that the first inequality in the preceding definition is naturally true for any normalized Schauder basis in any Banach space. It is worth noticing that $C \geq 1$ always.

Lemma 3.3. Let G be an infinite discrete group and Z be a Banach space with a normalized Schauder basis $\{z_n\}_{n=1}^\infty$ which is equivalent to the canonical Schauder basis of $\ell^1(\mathbb{N})$ with equivalence constant C (as in Definition 3.2). Then,

$$\frac{1}{C} \sum_{i=1}^N |\lambda_i| \leq \left\| \sum_{i=1}^N \lambda_i \delta_{g_i} \otimes z_{n_i} \right\|_\gamma$$

for any $\{g_1, g_2, \dots, g_N\} \subset G$, $\{n_1, n_2, \dots, n_N\} \subset \mathbb{N}$ and $N \in \mathbb{N}$.

Proof. We prove this by induction on N . Notice that the statement is true for $N = 2$ because

$$\|\lambda_1 \delta_{g_1} \otimes z_{n_1} + \lambda_2 \delta_{g_2} \otimes z_{n_2}\|_\gamma = \begin{cases} |\lambda_1| + |\lambda_2| & \text{if } g_1 \neq g_2; \text{ and} \\ \|\lambda_1 z_{n_1} + \lambda_2 z_{n_2}\| & \text{if } g_1 = g_2, \end{cases}$$

where the first equality follows from Lemma 2.2(1). Thus, since $C \leq 1$, we have

$$\frac{1}{C} \sum_{i=1}^2 |\lambda_i| \leq \left\| \sum_{i=1}^2 \lambda_i \delta_{g_i} \otimes z_{n_i} \right\|_\gamma.$$

Now, let the statement be true for $N = r > 2$. Then, for any given collections $\{g_1, \dots, g_{r+1}\} \subset G$ and $\{n_1, \dots, n_{r+1}\} \subset \mathbb{N}$, consider

$$f := \sum_{i=1}^{r+1} \lambda_i \delta_{g_i} \otimes z_{n_i} \in L^1(G) \otimes^\gamma Z.$$

If all the g_i 's are same then $\|f\|_\gamma = \|\sum_{i=1}^{r+1} \lambda_i z_{n_i}\| \geq \frac{1}{C} \sum_{i=1}^{r+1} |\lambda_i|$, because $\|\cdot\|_\gamma$ is a cross norm and the Schauder basis $\{z_n\}$ is normalized and equivalent with the canonical Schauder basis of $\ell^1(\mathbb{N})$.

On the other hand, if some of the g_i 's are distinct, we can choose a g_j such that it has less than $r + 1$ (say k) repetitions (we may even have $k = 1$); and, suppose that $\{n_t : 1 \leq t \leq k\} = \{m : g_j = g_m\}$. Then,

$$\begin{aligned} \|f\|_\gamma &= \|\delta_{g_j} \otimes (\sum_{t=1}^k \lambda_{j_t} z_{n_{j_t}})\|_\gamma + \left\| \sum_{i \neq j_t \text{ for any } t} \lambda_i \delta_{g_i} \otimes z_{n_i} \right\|_\gamma \\ &\geq \left\| \sum_{t=1}^k \lambda_{j_t} z_{n_{j_t}} \right\| + \frac{1}{C} \sum_{i \neq j_t \text{ for any } t} |\lambda_i| && \text{(by induction hypothesis)} \\ &\geq \frac{1}{C} \sum_{t=1}^k |\lambda_{j_t}| + \frac{1}{C} \sum_{i \neq j_t \text{ for any } t} |\lambda_i| && \text{(by equivalence of bases)} \end{aligned}$$

$$= \frac{1}{C} \sum_{i=1}^{r+1} |\lambda_i|.$$

Thus, by the principle of induction, the statement is true for all $N \in \mathbb{N}$. \square

Recall that a Schauder basis $\{a_n\}$ of a Banach algebra \mathcal{A} is said to be orthogonal if $a_n \cdot a_m = \delta_{mn} a_n$ for all $m, n \in \mathbb{N}$. We suggest the reader to see [8] for more on orthogonal Schauder bases of Banach algebras.

Theorem 3.4. *Let G be an infinite discrete group, \mathcal{A} be a Banach algebra with an orthonormal Schauder basis equivalent to the canonical basis of $\ell^1(\mathbb{N})$ and \mathcal{U} be a non-principal ultrafilter on \mathbb{N} . Then, $\varphi_{\mathcal{U}} \in Z_t^{(l)}((L^1(G, \mathcal{A}))^{**})$.*

In particular, the generalized group algebra $L^1(G, \mathcal{A})$ is not strongly Arens irregular.

Proof. Let $X := L^1(G, \mathcal{A})$ and $\{a_n\}$ denote an orthonormal Schauder basis of \mathcal{A} which is equivalent to the standard Schauder basis of $\ell^1(\mathbb{N})$. Consider $S := \{h_n\} \subseteq G$ as in Proposition 3.1. Then, for any $\psi \in X^{**}$ and $\mathcal{T} \in X^*$, we have

$$\begin{aligned} (\varphi_{\mathcal{U}} \square \psi)(\mathcal{T}) &= \lim_{\mathcal{U}} \psi(\mathcal{T}(\delta_{h_n} \otimes a_n)) \\ &= \lim_{\mathcal{U}} \psi(\mathcal{T}_{\delta_{h_n} \otimes a_n}). \end{aligned}$$

We claim that $\lim_n \psi(\mathcal{T}_{\delta_{h_n} \otimes a_n}) = 0$.

To prove this we assume the contrary. Then, there exists an $\epsilon > 0$ and an increasing sequence $\{n_t\}_{t \in \mathbb{N}} \subset \mathbb{N}$ such that $|\psi(\mathcal{T}_{\delta_{h_{n_t}} \otimes a_{n_t}})| \geq \epsilon$ for all $t \in \mathbb{N}$.

Consider the functionals $h_N := \sum_{t=1}^N c_{n_t} \mathcal{T}_{\delta_{h_{n_t}} \otimes a_{n_t}}$ for $N \in \mathbb{N}$, where $c_{n_t} := \frac{\psi(\mathcal{T}_{\delta_{h_{n_t}} \otimes a_{n_t}})}{|\psi(\mathcal{T}_{\delta_{h_{n_t}} \otimes a_{n_t}})|}$. Then, for any two finite sets $\{g_i : 1 \leq i \leq k\} \subset G$ and $\{s_i : 1 \leq i \leq k\} \subset \mathbb{N}$ (with possible repetitions), if $x := \sum_{i=1}^k \lambda_i \delta_{g_i} \otimes a_{s_i} \in L^1(G) \otimes \mathcal{A}$, then we have

$$\begin{aligned} \|h_N(x)\| &= \left| \mathcal{T} \left(\sum_{t=1}^N \sum_{i=1}^k \lambda_i c_{n_t} \delta_{h_{n_t}} \star \delta_{g_i} \otimes a_{n_t} \cdot a_{s_i} \right) \right| \\ &= \left| \mathcal{T} \left(\sum_{i=1}^k \lambda_i c_{s_i} \delta_{h_{s_i} + g_i} \otimes a_{s_i} \right) \right| && (\because a_i \cdot a_j = \delta_{i,j} a_i) \\ &\leq \|\mathcal{T}\| \cdot \left\| \sum_{i=1}^k \lambda_i c_{s_i} \delta_{h_{s_i} + g_i} \otimes a_{s_i} \right\|_{\gamma} \\ &\leq \|\mathcal{T}\| \cdot \sum_{i=1}^k |\lambda_i| && (\because |c_{s_i}| = 1) \\ &\leq C \|\mathcal{T}\| \|x\|_{\gamma}. && (\text{due to Lemma 3.3}) \end{aligned}$$

Since such x 's form a dense subspace of $L^1(G) \otimes_{\gamma} \mathcal{A}$, it follows that $\|h_N\| \leq C \|\mathcal{T}\|$ for all N . Now notice that

$$\psi(h_N) = \sum_{t=1}^N |\psi(\mathcal{T}_{\delta_{h_{n_t}} \otimes a_{n_t}})| \geq N\epsilon$$

for all $N \in \mathbb{N}$. But this is absurd because ψ is a bounded linear functional and $\{h_N\}$ is a bounded sequence. Thus, our assumption is wrong, and we must have

$$\lim_n \psi(\mathcal{T}_{\delta_{h_n} \otimes a_n}) = 0.$$

Thus, $(\varphi_{\mathcal{U}} \square \psi)(\mathcal{T}) = 0$ (because the limit along an ultrafilter in a compact Hausdorff space is unique and every non-principal ultrafilter contains the cofinite filter). Since ψ and \mathcal{T} were arbitrary, we conclude that $\varphi_{\mathcal{U}} \square \psi = 0$ for all $\psi \in X^{**}$.

Next, we turn to the second Arens product. We have $(\varphi_{\mathcal{U}} \diamond \psi)(\mathcal{T}) = \psi(\mathcal{T}_{\varphi_{\mathcal{U}}})$ for all $\psi \in X^{**}$, $\mathcal{T} \in X^*$. Notice that for any $u \in L^1(G) \otimes^{\gamma} \mathcal{A}$, we have

$$\begin{aligned} \mathcal{T}_{\varphi_{\mathcal{U}}}(u) &= \varphi_{\mathcal{U}}(u\mathcal{T}) \\ &= \lim_{\mathcal{U}} u\mathcal{T}(\delta_{h_n} \otimes a_n) \\ &= \lim_{\mathcal{U}} \mathcal{T}_{\delta_{h_n} \otimes a_n}(u). \end{aligned}$$

Thus, $\mathcal{T}_{\varphi_{\mathcal{U}}} = \lim_{\mathcal{U}} \mathcal{T}_{\delta_{h_n} \otimes a_n}^{w^*}$ and hence $\varphi_{\mathcal{U}} \diamond \psi(\mathcal{T}) = \psi(\lim_{\mathcal{U}} \mathcal{T}_{\delta_{h_n} \otimes a_n}^{w^*})$ for all $\psi \in X^{**}$, $\mathcal{T} \in X^*$. Further, for an arbitrary $u = \sum_{i=1}^{\infty} \lambda_i f_i \otimes b_i \in L^1(G, \mathcal{A})$, we have

$$\begin{aligned} \lim_n |\mathcal{T}_{\delta_{h_n} \otimes a_n}(u)| &= \lim_n \left| \mathcal{T} \left(\sum_{i=1}^{\infty} \lambda_i \delta_{h_n} \star f_i \otimes a_n^*(b_i) \right) \right| \quad (\because a_n \cdot b_i = a_n^*(b_i) \text{ due to orthonormality}) \\ &= 0. \quad (\because \lim_n a_n^*(b_i) = 0 \forall i) \end{aligned}$$

Thus, $\lim_{\mathcal{U}} \mathcal{T}_{\delta_{g_n} \otimes e_n}^{w^*} = 0$. Hence, $\varphi_{\mathcal{U}} \diamond \psi(\mathcal{T}) = 0$ for all $\psi \in X^{**}$ and all $\mathcal{T} \in X^*$.

Thus, we have shown that $\varphi_{\mathcal{U}} \square \psi = 0 = \varphi_{\mathcal{U}} \diamond \psi$ for all $\psi \in X^{**}$. In particular, $\varphi_{\mathcal{U}} \in Z_t^{(l)}(X^{**})$.

By Proposition 3.1, we know that $\varphi_{\mathcal{U}} \notin J(L^1(G, \mathcal{A}))$. Hence, $L^1(G, \mathcal{A})$ is not strongly Arens irregular. \square

Remark 3.5. A Banach algebra \mathcal{A} satisfying the hypothesis of Theorem 3.4 is in fact forced to be algebraically isomorphic (not necessarily isometric) to $\ell^1(\mathbb{N})$ (with respect to pointwise multiplication). Indeed, the map $\theta : \ell^1(\mathbb{N}) \rightarrow \mathcal{A}$ defined as $\theta(\sum_{n=1}^{\infty} c_n e_n) = \sum_{n=1}^{\infty} c_n a_n$ defines an algebra isomorphism. Thus, in other words, such an \mathcal{A} can always be thought of as $\ell^1(\mathbb{N})$ with some equivalent Banach algebra norm.

REFERENCES

- [1] R. Arens, The adjoint of a bilinear operation, Proc. Amer. Math. Soc. **26** (1951), 839-848.
- [2] H. G. Dales and A. T.-M. Lau, The second duals of Beurling algebras, Mem. Amer. Math. Soc. **177** (2005), no. 836.
- [3] C. C. Graham, A B.A.I. proof of the non-Arens regularity of $L^1(G)$, Proc. Amer. Math. Soc. **133** (2005), no. 1, 163-165.
- [4] E. Kaniuth, A Course in Commutative Banach Algebras, Springer New York, 2009.
- [5] A. T.-M. Lau and V. Losert, On the second conjugate algebra of $L^1(G)$ of a locally compact group, J. London Math. Soc. **37** (1988), no. 2, 464-470.
- [6] V. Montesinos, M. Fabian, P. Hájek, P. Habala and V. Zizler, Banach Space Theory: The Basis for Linear and Nonlinear Analysis, Springer New York, 2011.
- [7] R. A. Ryan, Introduction to Tensor Products of Banach Spaces, Springer London, 2013.
- [8] T. Husain and S. Watson, Topological algebras with orthogonal Schauder basis, Pacific J. Math. **91** (1980), no. 2, 339-347.
- [9] A. Ülger, Arens regularity of the algebra $A \otimes^{\gamma} B$, Trans. Amer. Math. Soc. **305** (1988), no. 2, 623-639.
- [10] A. Ülger, Erratum to: "Arens regularity of the algebra $A \otimes^{\gamma} B$ " [Trans. Amer. Math. Soc. 305 (1988), no. 2, 623-639], Trans. Amer. Math. Soc. **355** (2003), no. 9, 3839.

SCHOOL OF PHYSICAL SCIENCES, JAWAHARLAL NEHRU UNIVERSITY, NEW DELHI
Email address: lavksingh@hotmail.com