

On point estimators for Gamma and Beta distributions

Nickos Papadatos*

*National and Kapodistrian University of Athens, Department of Mathematics,
Section of Statistics and Operations Research, Panepistemiopolis, 157 84 Athens, Greece.*

Abstract

Let X_1, \dots, X_n be a random sample from the Gamma distribution with density $f(x) = \lambda^\alpha x^{\alpha-1} e^{-\lambda x} / \Gamma(\alpha)$, $x > 0$, where both $\alpha > 0$ (the shape parameter) and $\lambda > 0$ (the reciprocal scale parameter) are unknown. The main result shows that the *uniformly minimum variance unbiased estimator* (UMVUE) of the shape parameter, α , exists if and only if $n \geq 4$; moreover, it has finite variance if and only if $n \geq 6$. More precisely, the form of the UMVUE is given for all parametric functions α , λ , $1/\alpha$ and $1/\lambda$. Furthermore, a highly efficient estimating procedure for the two-parameter Beta distribution is also given. This is based on a Stein-type covariance identity for the Beta distribution, followed by an application of the theory of U -statistics and the delta-method.

MSC: Primary 62F10; 62F12; Secondary 62E15.

Key words and phrases: unbiased estimation; Gamma distribution; Beta distribution; Ye-Chen-type closed-form estimators; asymptotic efficiency; U -statistics; Stein-type covariance identity; delta-method.

1 Introduction

Let $\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$, $\alpha > 0$, be the Gamma function. The Gamma distribution, $\mathcal{G}(\alpha, \lambda)$, with density

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x > 0,$$

and parameters $\alpha > 0$ (the shape) and $\lambda > 0$ (the reciprocal scale), generates one of the most useful statistical models, especially when the data are nonnegative. This distribution belongs to the Pearson family, and in particular, to the Integrated Pearson family of distributions, see, e.g., Afendras and Papadatos (2015). Therefore, estimating procedures for the parameters α , λ , and/or $1/\alpha$ and $1/\lambda$, are of fundamental importance in applications. It is well-known that the maximum likelihood estimators (MLEs) are not tractable and, consequently, one has to apply numerical procedures for their evaluation. Moreover, to the best of our knowledge, it is not even known whether an unbiased estimator for α (and λ) exists, for some sample size n . In the contrary, unbiased estimators for the reciprocal parameters $1/\alpha$ and $1/\lambda$ do exist; recently, Ye and Chen (2017) obtained closed-form unbiased estimators for the reciprocal parameters with very high asymptotic efficiency. It should be noted that an augmented-sample (simulation) technique for obtaining exact confidence intervals for α is introduced by Iliopoulos (2016).

*e-mail: npapadat@math.uoa.gr, url: users.uoa.gr/~npapadat/

In the present work we show that an unbiased estimator for α exists if and only if the sample size n is at least 4, and the same is true for λ (provided that α is not too small). More precisely, in Section 2 we obtain the uniformly minimum variance unbiased estimators (UMVUEs) of α , λ , $1/\alpha$, $1/\lambda$. In particular, the UMVUE of $1/\lambda$, based on two observations, can be written as a positive symmetric kernel, whence, the corresponding U -statistic generates the Ye-Chen (2017) estimator; see Remark 2.2.

Let $B(\alpha, \beta) = \Gamma(\alpha)\Gamma(\beta)/\Gamma(\alpha+\beta)$, $\alpha > 0, \beta > 0$, be the Beta function. The Beta distribution has density

$$f(x) = \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}, \quad 0 < x < 1. \quad (1.1)$$

This distribution belongs to the Integrated Pearson family, and, at least to our knowledge, it is not known whether the parameters can be unbiasedly estimated. The MLEs do not have closed forms, and one has to overcome similar difficulties as for the Gamma case. In Section 3 we obtain closed-form, Ye-Chen-type, estimators for the parameters. Our method is different than that of Ye and Chen (2017), since it is based on a Stein-type covariance identity for the Beta distribution, followed by an application of the theory of U -statistics. The covariance identity generates an unbiased bivariate symmetric kernel for the parameter $\theta = 1/(\alpha + \beta)$. Then, an application of the classical theory of U -statistics produces an unbiased estimator of θ , which, in turn, yields the desired estimators of α and β . The asymptotic efficiency of the proposed estimators is studied in some detail, and it is shown to be very high (see Table 1). The Beta model is useful not only for analyzing data with bounded support, but also when positive data are mapped to the interval $(0, 1)$ by means of a transformation, e.g., $x \mapsto x/(1+x)$.

2 Unbiased estimation of the parameters of Gamma distribution

Let $\mathbf{X} = (X_1, \dots, X_n)$ ($n \geq 2$) be a random sample from $\Gamma(\alpha, \lambda)$, where the unknown parameter $\theta = (\alpha, \lambda) \in \Theta = (0, \infty)^2$. It is well-known that the complete, sufficient statistic can be written in the form $\mathbf{T}(\mathbf{X}) = (X, Y)$, where $X = \sum_{i=1}^n X_i$ and $Y^{1/n}$ is the ratio of the geometric to the arithmetic mean of the data. The random variables X, Y are independent, as follows, e.g., from Basu's theorem, because the distribution of Y is free of λ , while X is complete for λ when $\alpha > 0$ is known. The random variable X follows a $\mathcal{G}(n\alpha, \lambda)$ distribution, while Y is distributed like $\prod_{i=1}^{n-1} B_\alpha(i/n)$, where the random variables $B_\alpha(i/n)$, $i = 1, \dots, n-1$, are independent, and $B_\alpha(i/n)$ has a Beta distribution (1.1) with parameters α and $\beta = i/n$.

By using the Beta product representation, it follows inductively that the random variable Y has density

$$f_Y(y; \alpha) = K(\alpha) y^{\alpha-1} (1-y)^{(n-3)/2} G(y), \quad 0 < y < 1, \quad (2.1)$$

where $K(\alpha) = \Gamma(\alpha)^{-n} \prod_{i=1}^{n-1} \Gamma(\alpha + i/n)/\Gamma(i/n)$, and the (power-series) function G is given by

$$G(y) = \sum_{m=0}^{\infty} d_m (1-y)^m, \quad 0 < y < 1. \quad (2.2)$$

Notice that our notation suspends the dependence of $K(\alpha)$ and $G(y)$ on n .

In (2.2) the coefficients $(d_m)_{m \geq 0}$ are defined recurrently as follows:

$$\begin{aligned} \gamma_1(m) &= I(m=0) = \begin{cases} 1, & \text{if } m=0, \\ 0, & \text{if } m=1, 2, \dots, \end{cases} \\ \gamma_{i+1}(m) &= \frac{\Gamma(m+i(i+1)/(2n))}{\Gamma(m+(i+1)(i+2)/(2n))} \sum_{k=0}^m \frac{\Gamma(m-k+(i+1)/n)}{(m-k)!} \gamma_i(k), \quad i=1, \dots, n-2, \\ d_m &:= \gamma_{n-1}(m), \quad m=0, 1, \dots \end{aligned} \quad (2.3)$$

Most of the preceding results can be found in Glaser (1976), Nandi (1980), and Tang and Gupta (1984). Obviously, $G(y) \equiv 1$ when $n=2$. Moreover, for $n \geq 3$, G is strictly decreasing and positive; more precisely, $d_m > 0$ for all m , $G'(y) < 0$ and $G(y) > G(1-) = d_0 > 0$, $0 < y < 1$. It should be noted at this point that, when $n \geq 3$, the auxiliary random variable $Y_1 = \prod_{i=1}^{n-2} B_\alpha(i/n)$ has density (cf. Tang and Gupta, 1984)

$$f_{Y_1}(y; \alpha) = K_1(\alpha) y^{\alpha-1} (1-y)^{(n-1)(n-2)/(2n)-1} G_1(y), \quad 0 < y < 1, \quad (2.4)$$

where $K_1(\alpha) = \Gamma(\alpha)^{-(n-1)} \prod_{i=1}^{n-2} \Gamma(\alpha+i/n)/\Gamma(i/n)$ and

$$G_1(y) = \sum_{k=0}^{\infty} \gamma_{n-2}(k) (1-y)^k, \quad 0 < y < 1. \quad (2.5)$$

The random variable Y_1 will be used in the proof of Theorem 2.1, below.

Definition 2.1. An estimator $T(X)$ will be called *uniformly minimum variance unbiased estimator* (UMVUE) for the parametric function $h(\alpha, \lambda)$ if $T(X) = u(X, Y)$ is a (Borel) function of the complete, sufficient statistic (X, Y) , and $\mathbf{E}_{(\alpha, \lambda)} u(X, Y) = h(\alpha, \lambda)$, for all $\alpha > 0$, $\lambda > 0$, where the subscript (α, λ) denotes expectation w.r.t. the joint density of (X_1, \dots, X_n) ,

$$f(x_1, \dots, x_n) = \frac{\lambda^{n\alpha}}{\Gamma(\alpha)^n} \left(\prod_{i=1}^n x_i \right)^{\alpha-1} \exp \left(-\lambda \sum_{i=1}^n x_i \right), \quad (x_1, \dots, x_n) \in (0, \infty)^n.$$

Accordingly, an UMVUE may, or may not have finite variance. For completeness of the presentation, the term UMVUE will be also used in a wider sense, namely, in the case where the equality $\mathbf{E}_{(\alpha, \lambda)} u(X, Y) = h(\alpha, \lambda)$ is merely satisfied for all (α, λ) in a subset of $(0, \infty)^2$ with nonempty interior.

In the sequel we shall make use of the following simple, but useful, observation, saying that if two functions have identical Laplace transforms in an arbitrarily small interval then they coincide.

Lemma 2.1. Let $I = (\lambda_1, \lambda_2)$ where $0 \leq \lambda_1 < \lambda_2 \leq \infty$. Suppose that for the Borel functions $g_i : (0, \infty) \rightarrow \mathbb{R}$, $i=1, 2$, it is true that $\int_0^\infty e^{-\lambda x} |g_i(x)| dx < \infty$ for all $\lambda \in I$. If

$$\int_0^\infty e^{-\lambda x} g_1(x) dx = \int_0^\infty e^{-\lambda x} g_2(x) dx, \quad \lambda \in I,$$

then $g_1(x) = g_2(x)$ for almost all $x \in (0, \infty)$.

Proof: For $i = 1, 2$ write $g_i = g_i^+ - g_i^-$ where $g_i^+(x) = \max\{g_i(x), 0\}$, $g_i^-(x) = \max\{-g_i(x), 0\}$. By assumption,

$$0 \leq \int_0^\infty e^{-\lambda x} w_1(x) dx = \int_0^\infty e^{-\lambda x} w_2(x) dx < \infty, \quad \lambda \in I,$$

where $w_1 = g_1^+ + g_2^- \geq 0$, $w_2 = g_1^- + g_2^+ \geq 0$. Fix $\lambda_0 \in I$ and set $\nu_0 := \int_0^\infty e^{-\lambda_0 x} w_1(x) dx = \int_0^\infty e^{-\lambda_0 x} w_2(x) dx \geq 0$. Since $w_i \geq 0$, it is clear that $\nu_0 = 0$ implies that $w_1 = 0$ and $w_2 = 0$ a.e.; then, $w_1 = w_2$ a.e. and, thus, $g_1 = g_2$ a.e. If $\nu_0 > 0$, we may define the probability densities $f_i(x) = e^{-\lambda_0 x} w_i(x) / \nu_0$, $i = 1, 2$; then, for $\lambda \in I$,

$$\int_0^\infty e^{(\lambda_0 - \lambda)x} f_1(x) dx = \frac{1}{\nu_0} \int_0^\infty e^{-\lambda x} w_1(x) dx = \frac{1}{\nu_0} \int_0^\infty e^{-\lambda x} w_2(x) dx = \int_0^\infty e^{(\lambda_0 - \lambda)x} f_2(x) dx.$$

If M_i is the moment generating function of f_i , it follows that $M_1(t) = M_2(t) < \infty$ for all $t \in J := (\lambda_0 - \lambda_2, \lambda_0 - \lambda_1)$, and since J contains the origin, $f_1 = f_2$ a.e.; this implies that $w_1 = w_2$ a.e., and consequently, $g_1 = g_2$ a.e.

Proposition 2.1. Let $s(x, y) : (0, \infty)^2 \rightarrow \mathbb{R}$ be a Borel measurable function satisfying

$$\int_0^\infty \int_0^\infty |s(x, y)| \exp(-\lambda x - \alpha y) x^{n\alpha-1} dy dx < \infty \quad \text{for all } \alpha > 0, \lambda > 0.$$

Then, the function

$$H(\alpha, \lambda) := \int_0^\infty \int_0^\infty s(x, y) \exp(-\lambda x - \alpha y) x^{n\alpha-1} dy dx$$

is continuous in $(0, \infty)^2$.

Proof: Consider an arbitrary sequence $(\alpha_m, \lambda_m) \rightarrow (\alpha, \lambda) \in (0, \infty)^2$, as $m \rightarrow \infty$. Then, we can find m_0 such that $\frac{\alpha}{2} < \alpha_m < 2\alpha$ and $\frac{\lambda}{2} < \lambda_m < 2\lambda$ for $m \geq m_0$. The function $\delta_1(\lambda) = e^{-\lambda x}$ is obviously decreasing (for each fixed $x > 0$), so that $e^{-\lambda_m x} < e^{-\frac{\lambda}{2}x}$, $m \geq m_0$. Consider also the function $\delta_2(\alpha) = e^{-\alpha y} x^{n\alpha-1}$ (for fixed $x > 0, y > 0$). Then, $\log \delta_2(\alpha) = (n \log x - y)\alpha - \log x$ is a linear function of α , and thus, $\delta_2(\alpha_m) < \max\{\delta_2(\alpha/2), \delta_2(2\alpha)\} < \delta_2(\alpha/2) + \delta_2(2\alpha)$, $m \geq m_0$. Combining the above we obtain the inequality

$$e^{-\lambda_m x - \alpha_m y} x^{n\alpha_m-1} < e^{-\frac{\lambda}{2}x - \frac{\alpha}{2}y} x^{n\frac{\alpha}{2}-1} + e^{-\frac{\lambda}{2}x - 2\alpha y} x^{2n\alpha-1}, \quad (x, y) \in (0, \infty)^2, \quad m \geq m_0.$$

It follows that the sequence

$$w_m(x, y) := s(x, y) e^{-\lambda_m x - \alpha_m y} x^{n\alpha_m-1}, \quad (x, y) \in (0, \infty)^2, \quad m = m_0, m_0 + 1, \dots,$$

is dominated by $W(x, y) := |s(x, y)| e^{-\frac{\lambda}{2}x - \frac{\alpha}{2}y} x^{n\frac{\alpha}{2}-1} + |s(x, y)| e^{-\frac{\lambda}{2}x - 2\alpha y} x^{2n\alpha-1}$, and, by assumption, W is integrable on $(0, \infty)^2$. Clearly, $w_m(x, y) \rightarrow w(x, y) := s(x, y) e^{-\lambda x - \alpha y} x^{n\alpha-1}$ as $m \rightarrow \infty$. Thus, by dominated convergence,

$$H(\alpha_m, \lambda_m) = \int_0^\infty \int_0^\infty w_m(x, y) dy dx \rightarrow \int_0^\infty \int_0^\infty w(x, y) dy dx = H(\alpha, \lambda), \quad \text{as } m \rightarrow \infty,$$

completing the proof.

Proposition 2.2. Let $h(\alpha) : (0, \infty) \rightarrow \mathbb{R}$ be a parametric function of the shape parameter. If there exists an unbiased estimator $U_1 = U_1(\mathbf{X})$ for h , then the UMVUE of $h(\alpha)$ is a function of Y , alone.

Proof: By assumption, $\mathbf{E}_{(\alpha, \lambda)}|U_1(\mathbf{X})| < \infty$ for all $\alpha > 0$, $\lambda > 0$. The function $U_1 : (0, \infty)^n \rightarrow \mathbb{R}$ can always be taken to be Borel measurable. From the RB/LS Theorem, the UMVUE is given by $U(X, Y) = \mathbf{E}(U_1|X, Y)$, where $U : (0, \infty)^2 \rightarrow \mathbb{R}$ can also be taken to be Borel measurable with no loss of generality. Unbiasedness of the estimator $U(X, Y)$ means that (recall that X, Y are independent)

$$\int_0^\infty \int_0^1 \left\{ \frac{\lambda^{n\alpha}}{\Gamma(n\alpha)} x^{n\alpha-1} e^{-\lambda x} \right\} \left\{ K(\alpha) y^{\alpha-1} (1-y)^{(n-3)/2} G(y) \right\} U(x, y) dy dx = h(\alpha),$$

for all $\alpha > 0$, $\lambda > 0$. This relation can be rewritten in the form

$$H(\alpha, \lambda) := \frac{h(\alpha)\Gamma(n\alpha)}{K(\alpha)\lambda^{n\alpha}} = \int_0^\infty \int_0^\infty (1 - e^{-y})^{(n-3)/2} G(e^{-y}) U(x, e^{-y}) \exp(-\lambda x - \alpha y) x^{n\alpha-1} dy dx.$$

Since $\mathbf{E}_{(\alpha, \lambda)}|U(X, Y)| < \infty$ for all $\alpha > 0$ and $\lambda > 0$, we may apply Proposition 2.1 with $s(x, y) = (1 - e^{-y})^{(n-3)/2} G(e^{-y}) U(x, e^{-y})$ to conclude that the function $H(\alpha, \lambda)$ is continuous, and hence, $h(\alpha)$ is a continuous function of $\alpha > 0$; notice that all functions $K(\alpha)$, $\Gamma(n\alpha)$, and $\lambda^{n\alpha}$, are positive and continuous.

Write now the relation $h(\alpha) = \mathbf{E}_{(\alpha, \lambda)} U(X, Y)$ in the form

$$\int_0^\infty e^{-\lambda x} h(\alpha) x^{n\alpha-1} dx = \int_0^\infty e^{-\lambda x} K(\alpha) x^{n\alpha-1} \left\{ \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) U(x, y) dy \right\} dx.$$

It follows that the Laplace transforms of the functions $f_1(x) := h(\alpha) x^{n\alpha-1}$ and $f_2(x) := K(\alpha) x^{n\alpha-1} \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) U(x, y) dy$ are identical, and Lemma 2.1 implies that for every $\alpha > 0$,

$$\frac{h(\alpha)}{K(\alpha)} = \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) U(x, y) dy, \quad \text{for almost all } x \in (0, \infty). \quad (2.6)$$

Write $\mathbb{Q}_+ = \mathbb{Q} \cap (0, \infty)$ and set $A_\alpha = \{x > 0 : (2.6) \text{ holds}\}$, $N_\alpha = (0, \infty) \setminus A_\alpha = \{x > 0 : (2.6) \text{ does not hold}\}$, $A = \bigcap_{\alpha \in \mathbb{Q}_+} A_\alpha$, $N = \bigcup_{\alpha \in \mathbb{Q}_+} N_\alpha = (0, \infty) \setminus A$. By construction, the Lebesgue measure of N is zero. We chose an arbitrary $x_0 \in A$ and define $\tilde{u}(y) := U(x_0, y)$. Notice that $\tilde{u} : (0, \infty) \rightarrow \mathbb{R}$ is not necessarily Borel measurable; it is, however, Lebesgue measurable and, hence, it is equal a.e. to a Borel function $u : (0, \infty) \rightarrow \mathbb{R}$. Obviously, substituting u in place of \tilde{u} , the values of the above integrals are not affected. From (2.6) we have

$$\frac{h(\alpha)}{K(\alpha)} = \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) u(y) dy, \quad \alpha \in \mathbb{Q}_+. \quad (2.7)$$

The right-hand side of (2.7) defines a continuous function of $\alpha > 0$, because

$$\int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) u(y) dy = \int_0^\infty e^{-\alpha y} (1 - e^{-y})^{(n-3)/2} G(e^{-y}) u(e^{-y}) dy$$

is the Laplace transform of the function $y \mapsto (1 - e^{-y})^{(n-3)/2} G(e^{-y}) u(e^{-y})$. Since $h(\alpha)$ and $K(\alpha)$ are both continuous, it follows from (2.7) that $h(\alpha) = \int_0^1 K(\alpha) y^{\alpha-1} (1-y)^{(n-3)/2} G(y) u(y) dy = \int_0^1 f_Y(y; \alpha) u(y) dy$ for all $\alpha > 0$. Hence, $u(Y)$ is an unbiased estimator of $h(\alpha)$, and also, it is a function of the complete, sufficient statistic (X, Y) ; thus, it is the (unique) UMVUE.

Remark 2.1. (A consequence to the Cramér-Rao bound). The Fisher information matrix based on a single observation from $\mathcal{G}(\alpha, \lambda)$ is given by

$$J = J(\alpha, \lambda) = \frac{1}{\lambda^2} \begin{pmatrix} \lambda^2 \psi_1(\alpha) & -\lambda \\ -\lambda & \alpha \end{pmatrix}, \text{ with inverse } J^{-1} = \frac{1}{\alpha \psi_1(\alpha) - 1} \begin{pmatrix} \alpha & \lambda \\ \lambda & \lambda^2 \psi_1(\alpha) \end{pmatrix},$$

where $\psi_1(\alpha) = (\log \Gamma(\alpha))'' > 1/\alpha$. The Cramér-Rao (CR) bound says that for any unbiased estimator $T = T(\mathbf{X})$ of the parametric function $h(\alpha, \lambda)$, $n \text{Var} T \geq (\nabla h(\alpha, \lambda))^T J^{-1} \nabla h(\alpha, \lambda)$. Hence, in the particular case where $h = h(\alpha)$ is a parametric function of the shape parameter alone (i.e., h does not depend on λ), the CR bound takes the form

$$n \text{Var} T(\mathbf{X}) \geq \frac{h'(\alpha)^2}{\psi_1(\alpha) - 1/\alpha}, \text{ whenever } \mathbf{E}_{(\alpha, \lambda)} T(\mathbf{X}) = h(\alpha) \text{ for all } \alpha > 0, \lambda > 0. \quad (2.8)$$

We shall now show that Proposition 2.2 yields a better lower bound. Indeed, since the UMVUE of $h(\alpha)$ is a Borel function of Y , say $u = u(Y)$, we can apply the CR bound, $\text{Var} u(Y) \geq h'(\alpha)^2 / J_Y$, where J_Y is the Fisher information of Y . From (2.1), the density of Y is given by

$$f_Y(y; \alpha) = \exp(a \log y - A(\alpha)) k(y),$$

where $A(\alpha) = -\log K(\alpha)$ and $k(y) = y^{-1} (1-y)^{(n-3)/2} G(y) I_{(0,1)}(y)$, with I_B denoting the indicator function of B . This shows that $\{f_Y(y; \alpha), \alpha > 0\}$ defines a natural exponential family of Lebesgue densities, and thus, the regularity conditions are fulfilled. We calculate $-\frac{\partial^2}{\partial \alpha^2} \log f_Y(y; \alpha) = A''(\alpha)$ and, therefore,

$$\begin{aligned} J_Y &= (-\log K(\alpha))'' = \left(n \log \Gamma(\alpha) + \sum_{i=1}^{n-1} \log \Gamma(i/n) - \sum_{i=1}^{n-1} \log \Gamma(\alpha + i/n) \right)'' \\ &= n \psi_1(\alpha) - \sum_{i=1}^{n-1} \psi_1(\alpha + i/n). \end{aligned}$$

It follows that for any unbiased estimator $T(\mathbf{X})$ of $h(\alpha)$,

$$n \text{Var} T(\mathbf{X}) \geq n \text{Var} u(Y) \geq \frac{h'(\alpha)^2}{\psi_1(\alpha) - n^{-1} \sum_{i=1}^{n-1} \psi_1(\alpha + i/n)}. \quad (2.9)$$

Since the function $\psi_1(\alpha)$ is positive (the Γ function is log-convex) and decreasing, we have $n^{-1} \psi_1(\alpha + i/n) < \int_{(i-1)/n}^{i/n} \psi_1(\alpha + t) dt$, $i = 1, \dots, n$. Therefore,

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^{n-1} \psi_1(\alpha + i/n) &< \frac{1}{n} \sum_{i=1}^n \psi_1(\alpha + i/n) < \int_0^1 \psi_1(\alpha + t) dt = \int_\alpha^{\alpha+1} \psi_1(t) dt \\ &= (\log \Gamma(\alpha + 1))' - (\log \Gamma(\alpha))' = \left(\log \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha)} \right)' = (\log \alpha)' = \frac{1}{\alpha}. \end{aligned}$$

Thus, excluding the trivial case where $h(\alpha)$ is the constant function, the lower bound in (2.9) is strictly larger than the CR bound of (2.8). This means that, for any given non-constant parametric function $h(\alpha)$, no efficient estimator exists, when efficiency is defined in the traditional way, i.e., in terms of attainability of the bound in (2.8). In the contrary, there are functions of the shape parameter that can be efficiently estimated in the sense of (2.9). More precisely, it can be checked that for any constants c_1, c_2 , the parametric function $h(\alpha) = c_1\left(\psi(\alpha) - n^{-1} \sum_{i=1}^{n-1} \psi(\alpha + i/n)\right) + c_2$, where $\psi(\alpha) = \Gamma'(\alpha)/\Gamma(\alpha)$, is efficiently estimated by $T(\mathbf{X}) = u(Y) = (c_1/n) \log Y + c_2$, in the sense that $\mathbf{E}_{(\alpha, \lambda)} T(\mathbf{X}) = h(\alpha)$ for all α, λ , and $n \text{Var} T(\mathbf{X}) = c_1^2\left(\psi_1(\alpha) - n^{-1} \sum_{i=1}^{n-1} \psi_1(\alpha + i/n)\right)$, which is equal to the lower bound given by (2.9). Finally, it can be easily verified that all efficiently estimated functions are of the above form. Notice, however, that asymptotically the lower bounds do coincide, since, by the Riemann integral, $\lim_{n \rightarrow \infty} n^{-1} \sum_{i=1}^{n-1} \psi_1(\alpha + i/n) = \int_{\alpha}^{\alpha+1} \psi_1(t) dt = 1/\alpha$.

We now state the main result of this section. To the best of our knowledge, this result, as well as the existence of u_0 , below, is new.

Theorem 2.1. Let X_1, \dots, X_n , $n \geq 2$, be a random sample from $\mathcal{G}(\alpha, \lambda)$. For every $n \geq 4$, the UMVUE of the shape parameter α is given by

$$u_0(Y) := \frac{1}{2}(n-3) \frac{Y}{1-Y} - \frac{YG'(Y)}{G(Y)}, \quad (2.10)$$

where G is as in (2.2) and $Y^{1/n}$ is the ratio of the geometric to the arithmetic mean of X_1, \dots, X_n . The estimator $u_0(Y)$ has finite variance if and only if $n \geq 6$. Moreover, for $n = 2, 3$, no unbiased estimator of α exists.

Proof: According to Proposition 2.2, if an unbiased estimator (of α) exists, the UMVUE must be a function of Y , say $u(Y)$. Then, the relation $\mathbf{E}_{\alpha} u(Y) = \alpha$ can be written as

$$\frac{1}{K(\alpha)} = \frac{1}{\alpha} \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) u(y) dy, \quad \alpha > 0.$$

Observing that $1/K(\alpha) = \int_0^1 y^{\alpha-1} (1-y)^{(n-3)/2} G(y) dy$ and $1/\alpha = \int_0^{\infty} e^{-\alpha x} dx$, the substitution $y = e^{-x}$ yields

$$\int_0^{\infty} e^{-\alpha x} (1 - e^{-x})^{(n-3)/2} G(e^{-x}) dx = \left\{ \int_0^{\infty} e^{-\alpha x} dx \right\} \left\{ \int_0^{\infty} e^{-\alpha x} (1 - e^{-x})^{(n-3)/2} G(e^{-x}) u(e^{-x}) dx \right\},$$

for all $\alpha > 0$. Since we have assumed that $\mathbf{E}_{\alpha} |u(Y)| < \infty$ for all $\alpha > 0$, we may apply Fubini's theorem to the right-hand side of the above equation, obtaining

$$\int_0^{\infty} e^{-\alpha x} (1 - e^{-x})^{(n-3)/2} G(e^{-x}) dx = \int_0^{\infty} e^{-\alpha x} \left\{ \int_0^x (1 - e^{-t})^{(n-3)/2} G(e^{-t}) u(e^{-t}) dt \right\} dx, \quad \alpha > 0.$$

This relation shows that the Laplace transforms of $f_1(x) := (1 - e^{-x})^{(n-3)/2} G(e^{-x})$ and $f_2(x) := \int_0^x (1 - e^{-t})^{(n-3)/2} G(e^{-t}) u(e^{-t}) dt$ are identical. Hence, $f_1 \equiv f_2$ (notice that both functions are continuous). Setting $x = -\log y$ we get

$$(1-y)^{(n-3)/2} G(y) = \int_0^{-\log y} (1 - e^{-t})^{(n-3)/2} G(e^{-t}) u(e^{-t}) dt = \int_y^1 \frac{1}{x} (1-x)^{(n-3)/2} G(x) u(x) dx, \quad (2.11)$$

$0 < y < 1$, and a differentiation of (2.11) leads to

$$-\frac{n-3}{2}(1-y)^{(n-5)/2}G(y)+(1-y)^{(n-3)/2}G'(y) = \frac{-u(y)}{y}(1-y)^{(n-3)/2}G(y), \quad \text{for almost all } y \in (0, 1).$$

Hence, solving for u we see that $u(y) = ((n-3)/2)y/(1-y) - yG'(y)/G(y) = u_0(y)$, a.e., with u_0 as in (2.10). The preceding argument shows that if an unbiased estimator $T(X_1, \dots, X_n)$ exists then $u_0(Y)$ must be unbiased, but *it does not prove that an unbiased estimator exists*. To see this, let $n = 2$. Then, since $G(y) \equiv 1$ when $n = 2$, (2.11) reads as

$$\frac{1}{\sqrt{1-y}} = \int_y^1 \frac{u(x)}{x\sqrt{1-x}} dx. \quad (2.12)$$

The assumption $\mathbf{E}_\alpha |u(Y)| < \infty$ is equivalent to $\int_0^1 x^{\alpha-1}(1-x)^{-1/2}|u(x)|dx < \infty$ for all $\alpha > 0$. Hence,

$$\left| \int_y^1 \frac{u(x)}{x\sqrt{1-x}} dx \right| \leq \int_y^1 \frac{|u(x)|}{x\sqrt{1-x}} dx = \int_y^1 \frac{x^{\alpha-1}|u(x)|}{\sqrt{1-x}} \frac{1}{x^\alpha} dx \leq \frac{1}{y^\alpha} \int_y^1 \frac{x^{\alpha-1}|u(x)|}{\sqrt{1-x}} dx \rightarrow 0,$$

as $y \rightarrow 1-$, and this contradicts (2.12). When $n = 3$, it is easy to see that the function G of (2.2) is given by

$$G(y) = \sum_{m=0}^{\infty} \frac{(3m)!}{(3^m m!)^3} (1-y)^m, \quad 0 < y < 1.$$

Hence, $G(1-) = 1$. Since $(1-y)^{(n-3)/2} \equiv 1$, assuming $\mathbf{E}_\alpha |u(Y)| < \infty$ we get $\int_0^1 x^{\alpha-1}G(x)|u(x)|dx < \infty$, $\alpha > 0$. Therefore,

$$\left| \int_y^1 \frac{G(x)}{x} u(x) dx \right| \leq \int_y^1 \frac{G(x)}{x} |u(x)| dx = \int_y^1 \frac{x^{\alpha-1}G(x)|u(x)|}{x^\alpha} dx \leq \frac{1}{y^\alpha} \int_y^1 x^{\alpha-1}G(x)|u(x)| dx \rightarrow 0,$$

as $y \rightarrow 1-$. This contradicts (2.11), i.e.,

$$G(y) = \int_y^1 \frac{1}{x} G(x) u(x) dx,$$

since the left-hand side of the above equation approaches 1 as $y \rightarrow 1-$.

The preceding argument shows that there is no unbiased estimator when $n \leq 3$. For $n \geq 4$, however, the situation is completely different: The (positive) estimator $u_0(Y)$ of (2.10) has finite expectation for all $\alpha > 0$ and, indeed, $\mathbf{E}_\alpha u_0(Y) = \alpha$. We now proceed to verify this claim. First observe that $-G'(y) > 0$ so that $u_0(y) > 0$ for $0 < y < 1$. Noting that $G'(y) = -\sum_{m=0}^{\infty} m d_m (1-y)^{m-1}$, we calculate

$$\begin{aligned} \mathbf{E}_\alpha u_0(Y) &= K(\alpha) \int_0^1 y^\alpha (1-y)^{(n-5)/2} \left(\sum_{m=0}^{\infty} \left(\frac{n-3}{2} + m \right) d_m (1-y)^m \right) dy \\ &= K(\alpha) \sum_{m=0}^{\infty} \left(\frac{n-3}{2} + m \right) d_m \int_0^1 y^\alpha (1-y)^{m+(n-5)/2} dy \\ &= \alpha \Gamma(\alpha) K(\alpha) \sum_{m=0}^{\infty} \frac{\Gamma(m+(n-1)/2)}{\Gamma(\alpha+m+(n-1)/2)} d_m, \end{aligned}$$

where, since $d_m > 0$, the interchanging of summation and integration is justified by Beppo Levi's theorem. It remains to verify that

$$S := \sum_{m=0}^{\infty} \frac{\Gamma(m + (n-1)/2)}{\Gamma(\alpha + m + (n-1)/2)} d_m = \frac{1}{\Gamma(\alpha)K(\alpha)}, \quad \alpha > 0. \quad (2.13)$$

From the definition of $d_m = \gamma_{n-1}(m)$ we have (set $i = n - 2$ in (2.3))

$$d_m = \gamma_{n-1}(m) = \frac{\Gamma(m + (n-1)(n-2)/(2n))}{\Gamma(m + (n-1)/2)} \sum_{k=0}^m \frac{\Gamma(m-k + (n-1)/n)}{(m-k)!} \gamma_{n-2}(k).$$

Substituting this expression to the series in (2.13) and changing the order of summation according to Tonelli's theorem, we obtain

$$\begin{aligned} S &= \sum_{k=0}^{\infty} \gamma_{n-2}(k) \sum_{m=k}^{\infty} \frac{\Gamma(m + (n-1)(n-2)/(2n)) \Gamma(m-k + (n-1)/n)}{(m-k)! \Gamma(\alpha + m + (n-1)/2)} \\ &= \sum_{k=0}^{\infty} \gamma_{n-2}(k) \sum_{m=0}^{\infty} \frac{\Gamma(m+k + (n-1)(n-2)/(2n)) \Gamma(m + (n-1)/n)}{m! \Gamma(\alpha + k + (n-1)/2)} \\ &= \frac{\Gamma(\alpha) \Gamma((n-1)/n)}{\Gamma(\alpha + (n-1)/n)} \sum_{k=0}^{\infty} \frac{\Gamma(k + (n-1)(n-2)/(2n))}{\Gamma(\alpha + k + (n-1)(n-2)/(2n))} \gamma_{n-2}(k), \end{aligned} \quad (2.14)$$

where for the last equality we applied the identity

$$\sum_{m=0}^{\infty} \frac{\Gamma(m + \rho_1) \Gamma(m + \rho_2)}{m! \Gamma(m + \rho + \rho_1 + \rho_2)} = \frac{\Gamma(\rho) \Gamma(\rho_1) \Gamma(\rho_2)}{\Gamma(\rho + \rho_1) \Gamma(\rho + \rho_2)}, \quad \rho, \rho_1, \rho_2 > 0,$$

with $\rho = \alpha$, $\rho_1 = k + (n-1)(n-2)/(2n)$, $\rho_2 = (n-1)/n$. Since the function $f_{Y_1}(y; \alpha)$ in (2.4) is a density, we have (see (2.5))

$$\frac{1}{K_1(\alpha)} = \int_0^1 y^{\alpha-1} (1-y)^{(n-1)(n-2)/(2n)-1} G_1(y) dy = \sum_{k=0}^{\infty} \gamma_{n-2}(k) \int_0^1 y^{\alpha-1} (1-y)^{(n-1)(n-2)/(2n)+k-1} dy,$$

and hence,

$$\sum_{k=0}^{\infty} \frac{\Gamma(k + (n-1)(n-2)/(2n))}{\Gamma(\alpha + k + (n-1)(n-2)/(2n))} \gamma_{n-2}(k) = \frac{1}{\Gamma(\alpha)K_1(\alpha)}.$$

Using the fact that $K_1(\alpha) = K(\alpha)\Gamma(\alpha)\Gamma((n-1)/n)/\Gamma(\alpha + (n-1)/n)$ and substituting the above equality in (2.14) we conclude that

$$S = \frac{\Gamma((n-1)/n)}{\Gamma(\alpha + (n-1)/n) K_1(\alpha)} = \frac{\Gamma((n-1)/n) \Gamma(\alpha + (n-1)/n)}{K(\alpha) \Gamma(\alpha) \Gamma((n-1)/n) \Gamma(\alpha + (n-1)/n)} = \frac{1}{K(\alpha)\Gamma(\alpha)};$$

this verifies (2.13) and the claim is proved. Thus, for $n \geq 4$, $u_0(Y)$ is the UMVUE of the parametric function $h(\alpha) = \alpha$.

We shall now investigate the finiteness of the variance of the UMVUE. Assume first that $n \geq 6$. From the inequality $u_0(y)^2 \leq 2c^2 y^2 / (1-y)^2 + 2y^2 |G'(y)|^2 / G(y)^2$, where $c = (n-3)/2$, we see that $\mathbf{E}_{\alpha} u_0(Y)^2 < \infty$ if both integrals

$$\int_0^1 y^{\alpha+1} (1-y)^{(n-7)/2} G(y) dy \quad \text{and} \quad \int_0^1 y^{\alpha+1} (1-y)^{(n-3)/2} \frac{|G'(y)|^2}{G(y)} dy$$

are finite. The integrands are continuous functions in $(0, 1)$, hence, it suffices to verify their integrability in a neighborhood of the endpoints. Obviously,

$$\int_{1-\epsilon}^1 y^{\alpha+1}(1-y)^{(n-7)/2} G(y) dy < \infty \quad \text{and} \quad \int_{1-\epsilon}^1 y^{\alpha+1}(1-y)^{(n-3)/2} \frac{|G'(y)|^2}{G(y)} dy < \infty,$$

since $n \geq 6$, $\lim_{y \rightarrow 1^-} y^{\alpha+1} G(y) = d_0$ and $\lim_{y \rightarrow 1^-} y^{\alpha+1} |G'(y)|^2 / G(y) = d_1^2 / d_0$, so that, close to $y = 1$, the first integrand behaves like $(1-y)^{(n-7)/2}$ and the second one like $(1-y)^{(n-3)/2}$. Regarding the behavior of the first integral close to $y = 0$ it suffices to observe that $G(y)$ is positive and $\int_0^1 y^{\alpha-1}(1-y)^{(n-3)/2} G(y) dy = 1/K(\alpha) < \infty$; see (2.1). Then, we trivially have $\int_0^\delta y^{\alpha+1}(1-y)^{(n-7)/2} G(y) dy < (1-\delta)^{-2} \int_0^\delta y^{\alpha-1}(1-y)^{(n-3)/2} G(y) dy < \infty$. It remains to verify that $\int_0^\delta y^{\alpha+1}(1-y)^{(n-3)/2} |G'(y)|^2 / G(y) dy < \infty$, and since $1/G(y) < 1/d_0$ and $(1-y)^{(n-3)/2} < 1$, it suffices to show that for some $\delta \in (0, 1)$,

$$\int_0^\delta y^{\alpha+1} |G'(y)|^2 dy < \infty. \quad (2.15)$$

Recall that $\mathbf{E}_\alpha u_0(Y) < \infty$; since $u_0(y) > y|G'(y)|/G(y)$, it follows that

$$\int_0^1 y^\alpha (1-y)^{(n-3)/2} |G'(y)| dy < \infty,$$

and thus, $\int_0^{1/2} y^\alpha |G'(y)| dy < \infty$. The function $|G'(y)| = \sum_{m=1}^\infty m d_m (1-y)^{m-1}$ is strictly decreasing in $(0, 1)$ and therefore, with $c_\alpha = (\alpha+1)/(1-2^{-\alpha-1}) > 0$, we get

$$y^{\alpha+1} |G'(y)| = c_\alpha |G'(y)| \int_{y/2}^y x^\alpha dx < c_\alpha \int_{y/2}^y x^\alpha |G'(x)| dx < c_\alpha \int_0^y x^\alpha |G'(x)| dx \rightarrow 0,$$

as $y \rightarrow 0+$, by dominated convergence. Hence, $\lim_{y \rightarrow 0+} y^{\alpha+1} |G'(y)| = 0$ (for all $\alpha > 0$). Thus, for every $\epsilon > 0$ we can find $\delta = \delta(\alpha, \epsilon) > 0$ such that $(y^{\alpha/4+1} |G'(y)|)^2 < \epsilon^2$ for $y \in (0, \delta)$. Then, for this value of δ ,

$$\int_0^\delta y^{\alpha+1} |G'(y)|^2 dy = \int_0^\delta y^{\alpha/2-1} (y^{\alpha/4+1} |G'(y)|)^2 dy < \epsilon^2 \int_0^\delta y^{\alpha/2-1} dy < \infty,$$

and (2.15) is proved. We thus conclude that the variance of $u_0(Y)$ is finite for all $n \geq 6$ and $\alpha > 0$.

Finally, assume that $n = 4$ or 5 . Since $u_0(y) = cy/(1-y) + y|G'(y)|/G(y) > cy/(1-y)$, $c = (n-3)/2 > 0$, and $G(y) > d_0 > 0$, the variance of the UMVUE is infinite:

$$\mathbf{E}_\alpha u_0(Y)^2 > c^2 K(\alpha) \int_0^1 y^{\alpha+1}(1-y)^{(n-7)/2} G(y) dy > c^2 K(\alpha) d_0 \int_0^1 y^{\alpha+1}(1-y)^{(n-7)/2} dy = \infty$$

for all $\alpha > 0$; this completes the proof of the theorem.

Corollary 2.1. For $n \geq 4$, the UMVUE of the rate parameter (reciprocal scale parameter) λ is given by

$$u_1(X, Y) = \frac{u_0(Y) - 1/n}{\bar{X}} = \frac{n u_0(Y) - 1}{X}, \quad n > 1/\alpha,$$

where $u_0(Y)$ is as in (2.10) and $\bar{X} = X/n = n^{-1} \sum_{i=1}^n X_i$. For $n \geq 6$, the estimator $u_1(X, Y)$ has finite variance if and only if $\alpha > 2/n$ and $\lambda > 0$, while for $n = 4$ or 5 its variance is infinite. Finally, if either $n \in \{2, 3\}$ or $n \geq 4$ and $0 < \alpha \leq n^{-1}$, no unbiased estimator for λ exists.

Proof: By definition, u_1 is a function of the complete, sufficient statistic (X, Y) . Since X, Y are independent and $\mathbf{E}_{(\alpha, \lambda)} X^{-1} = \lambda / (n\alpha - 1)$, $n > \alpha^{-1}$, we obtain $\mathbf{E}_{(\alpha, \lambda)} u_1(X, Y) = (n\mathbf{E}_{(\alpha, \lambda)} u_0(Y) - 1) (\mathbf{E}_{(\alpha, \lambda)} X^{-1}) = \lambda$, and $u_1(X, Y)$ is, indeed, the UMVUE, according to Definition 2.1. Let $n = 4$ or 5 . Since $u_1(x, y)^2 = x^{-2}(n^2 u_0(y)^2 - 2nu_0(y) + 1)$ and $\mathbf{E}_{(\alpha, \lambda)} u_0(Y)^2 = \infty$, it follows that $\mathbf{E}_{(\alpha, \lambda)} (n^2 u_0(Y)^2 - 2nu_0(Y) + 1) = \infty$ and, thus, by the independence of X, Y , $\mathbf{E}_{(\alpha, \lambda)} u_1(X, Y)^2 = (\mathbf{E}_{(\alpha, \lambda)} X^{-2}) (\mathbf{E}_{(\alpha, \lambda)} (n^2 u_0(Y)^2 - 2nu_0(Y) + 1)) = \infty$. Next, assume $n \geq 6$. In this case, $\mathbf{E}_{(\alpha, \lambda)} u_1(X, Y)^2 \leq (\mathbf{E}_{(\alpha, \lambda)} X^{-2}) (1 + n^2 \mathbf{E}_{(\alpha, \lambda)} u_0(Y)^2)$, and since $\mathbf{E}_{(\alpha, \lambda)} u_0(Y)^2 < \infty$, we obtain $\mathbf{E}_{(\alpha, \lambda)} u_1(X, Y)^2 < \infty$ if and only if $\mathbf{E}_{(\alpha, \lambda)} X^{-2} < \infty$; equivalently, $\alpha > 2n^{-1}$.

Assume now that $n \geq 2$, $0 < \alpha \leq n^{-1}$, and set $\theta := n\alpha \in (0, 1]$ for fixed α . If for this particular value of α the estimator $w(X, Y)$ were unbiased for $\lambda > 0$, then we could find an interval $I = (\lambda_1, \lambda_2)$, $0 \leq \lambda_1 < \lambda_2 \leq \infty$, such that

$$\lambda = \int_0^\infty \frac{\lambda^\theta}{\Gamma(\theta)} x^{\theta-1} e^{-\lambda x} \delta(x) dx, \quad \lambda \in I, \quad (2.16)$$

where $\delta(x) := K(\theta/n) \int_0^1 y^{\theta/n-1} (1-y)^{(n-3)/2} g(y) w(x, y) dy$. Certainly, the function w is assumed integrable, that is, $\mathbf{E}_{(\theta, \lambda)} |w(X, Y)| < \infty$, $\lambda \in I$. Hence, $\delta(x)$ is a.e. finite, and we can redefine it, if necessary, to be finite for all $x > 0$. Notice that the assumption $\mathbf{E}_{(\theta, \lambda)} |w(X, Y)| < \infty$ for $\lambda \in I$ shows that

$$\int_0^\infty e^{-\lambda x} x^{\theta-1} |\delta(x)| dx \leq \int_0^\infty \int_0^1 e^{-\lambda x} x^{\theta-1} y^{\theta/n-1} (1-y)^{(n-3)/2} g(y) |w(x, y)| dy dx < \infty, \quad \lambda \in I.$$

We now rewrite (2.16) as

$$\frac{\Gamma(\theta)}{\lambda^\theta} = \frac{1}{\lambda} \int_0^\infty e^{-\lambda x} x^{\theta-1} \delta(x) dx, \quad \lambda \in I.$$

The left-hand side of the above equation equals to $\int_0^\infty e^{-\lambda x} x^{\theta-1} dx$, while its right-hand side can be written as

$$\left(\int_0^\infty e^{-\lambda x_2} dx_2 \right) \left(\int_0^\infty e^{-\lambda x_1} x_1^{\theta-1} \delta(x_1) dx_1 \right) = \int_0^\infty e^{-\lambda x} \left\{ \int_0^x t^{\theta-1} \delta(t) dt \right\} dx,$$

from Fubini's theorem. Therefore, we conclude that

$$\int_0^\infty e^{-\lambda x} x^{\theta-1} dx = \int_0^\infty e^{-\lambda x} \left\{ \int_0^x t^{\theta-1} \delta(t) dt \right\} dx, \quad \lambda \in I,$$

and Lemma 2.1 shows that $x^{\theta-1} = \int_0^x t^{\theta-1} \delta(t) dt$, a.e. in $(0, \infty)$. Hence, the only possibility is $\delta(x) = (\theta - 1)/x$, a.e., and since $\delta(x) \leq 0$, this contradicts (2.16). Given $n = 3$ observations from $\mathcal{G}(\alpha, \lambda)$, it can be seen (we omit the details) that the relation $\mathbf{E}_{(\alpha, \lambda)} w(X, Y) = \lambda$ for all $\alpha \in (\alpha_1, \alpha_2)$, $\lambda \in (\lambda_1, \lambda_2)$ [with $\alpha_1 \geq 1/3$, $\lambda_1 \geq 0$] implies $w(x, y) = x^{-1}(y|G'(y)|/G(y) - 1)$, a.e. in $(0, \infty)^2$. Since $\mathbf{E}_{(\alpha, \lambda)} X^{-1} = \lambda / (3\alpha - 1)$, this estimator would be unbiased for λ if and only if $\mathbf{E}_\alpha Y |G'(Y)| / G(Y) = 3\alpha$; however, we showed in Theorem 2.1 that this is impossible. Hence, there is no unbiased estimator for λ when $n = 3$, and therefore, for $n \leq 3$, too.

Several authors use the scale parameter $\beta = 1/\lambda$ in defining the Gamma density, that is,

$$f(x; \alpha, \beta) = \frac{x^{\alpha-1}}{\beta^\alpha \Gamma(\alpha)} e^{-x/\beta}, \quad x > 0.$$

We now proceed to verify that the function G in (2.2) also appears in the form of the UMVUE of both $1/\alpha$ and $1/\lambda$.

Theorem 2.2. For $n \geq 2$, the UMVUE of $1/\alpha$ is given by

$$u_2(Y) := \frac{(1-Y)^{-(n-3)/2}}{G(Y)} \int_Y^1 \frac{1}{x} (1-x)^{(n-3)/2} G(x) dx,$$

with Y and G as in Theorem 2.1; the UMVUE of $1/\lambda$ is given by

$$u_3(X, Y) := \bar{X} u_2(Y) = X u_2(Y)/n.$$

Proof: It suffices to show that u_2 and u_3 are unbiased. For u_2 we have

$$\mathbf{E}_\alpha u_2(Y) = K(\alpha) \int_0^1 y^{\alpha-1} \left\{ \int_y^1 \frac{1}{x} (1-x)^{(n-3)/2} G(x) dx \right\} dy = \frac{1}{\alpha} \int_0^1 f_Y(x; \alpha) dx = \frac{1}{\alpha},$$

where the change in the order of integration is justified by Tonelli's theorem. The result for u_3 is obvious, since $\mathbf{E}_{(\alpha, \lambda)} \bar{X} = \alpha/\lambda$ and the random variables \bar{X} , Y are independent.

Remark 2.2. (Alternative derivation of the closed-form Ye-Chen-type estimators, and their asymptotic efficiency). For $n = 2$, u_2 and u_3 admit closed forms because $G(y) \equiv 1$. More specifically, let X_1, X_2 be independently distributed $\mathcal{G}(\alpha, \lambda)$ random variables, so that $\bar{X} = (X_1 + X_2)/2$ and $Y = 4X_1 X_2 / (X_1 + X_2)^2$. Then,

$$u_2(Y) = \sqrt{1-Y} \int_Y^1 \frac{1}{x \sqrt{1-x}} dx = \sqrt{1-Y} (2 \log(1 + \sqrt{1-Y}) - \log Y).$$

Therefore, after some algebra, one can verify that the unbiased estimators u_2 and u_3 (when $n = 2$) can be written as

$$u_2(Y) = T_2(X_1, X_2) = \frac{(X_2 - X_1)(\log X_2 - \log X_1)}{X_1 + X_2},$$

$$u_3(X, Y) = T_3(X_1, X_2) = \frac{1}{2}(X_2 - X_1)(\log X_2 - \log X_1).$$

From the above form of $u_3 = T_3$ it follows that $\mathbf{E}_{(\alpha, \lambda)} T_3(X_1, X_2) = \text{Cov}_{(\alpha, \lambda)}(X_1, \log X_1)$. The well-known Stein-type identity for the Gamma distribution – see, e.g., Afendras *et al.* (2011) – states that for any absolutely continuous function $w : (0, \infty) \rightarrow \mathbb{R}$,

$$\text{Cov}_{(\alpha, \lambda)}(X_1, w(X_1)) = \frac{1}{\lambda} \mathbf{E}_{(\alpha, \lambda)} X_1 w'(X_1),$$

provided $\mathbf{E}_{(\alpha, \lambda)} X_1 |w'(X_1)| < \infty$. It is now clear that, if we apply Stein's identity to $w(x) = \log x$, we conclude the unbiasedness of u_3 . Consider a random sample X_1, \dots, X_n of size $n \geq 2$ from any distribution; then, as is well-known, under suitable moment conditions, an

unbiased estimator of $\text{Cov}(X_1, w(X_1))$ is given by the corresponding U -statistic with kernel $K(X_1, X_2) = (X_2 - X_1)(w(X_2) - w(X_1))/2$, that is,

$$U_n := \frac{2}{n(n-1)} \sum_{i < j} K(X_i, X_j) = \frac{1}{n-1} \sum_{i=1}^n X_i w(X_i) - \frac{n}{n-1} \bar{X} \bar{w}(X),$$

where $\bar{X} = n^{-1} \sum_{i=1}^n X_i$, $\bar{w}(X) = n^{-1} \sum_{i=1}^n w(X_i)$. With $w(x) = \log x$ we obtain the highly efficient unbiased estimator of the scale parameter $1/\lambda$,

$$\widehat{\lambda^{-1}}(n) = U_n = \frac{1}{n-1} \left\{ \sum_{i=1}^n X_i \log X_i - \bar{X} \sum_{i=1}^n \log X_i \right\},$$

introduced by Ye and Chen (2017) (see their Theorem 4.1), using generalized likelihood equations. From the well-known asymptotic theory of U -statistics, Hoeffding (1948), we have $n \text{Var} U_n \rightarrow 4\zeta_1^2$ and $\sqrt{n}(U_n - \mu) \xrightarrow{d} \mathcal{N}(0, 4\zeta_1^2)$, as $n \rightarrow \infty$, where $\mu = \mathbf{E}K(X_1, X_2)$, $\zeta_1^2 = \text{Var} \mathbf{E}(K(X_1, X_2)|X_2)$. Therefore, with $K = T_3$ (equivalently, $w(x) = \log x$) we obtain

$$\sqrt{n}(\widehat{\lambda^{-1}}(n) - \lambda^{-1}) \xrightarrow{d} \mathcal{N}(0, (1 + \alpha\psi_1(\alpha))/\lambda^2), \quad \text{as } n \rightarrow \infty,$$

in accordance to Theorem 3.2 in Ye and Chen (2017), since $1/\lambda = \beta$. Hence, the asymptotic efficiency of the estimator $\widehat{\lambda^{-1}}(n)$ (compared to the MLE, see Remark 2.1) is given by $\text{ARE}_{\widehat{\lambda^{-1}}(n)}(\alpha, \lambda) = \psi_1(\alpha)/(\alpha^2\psi_1(\alpha)^2 - 1)$, and it is independent of λ . Numerical calculations show that the asymptotic efficiency is greater than 96.4% for all values of $\alpha > 0$ (the minimum of the function $\psi_1(\alpha)/(\alpha^2\psi_1(\alpha)^2 - 1)$ is attained at $\alpha = \alpha^* = .920835\dots$). Applying the delta-method (using the mapping $x \mapsto x^{-1}$) we immediately obtain the corresponding result for the rate parameter λ , namely, $\sqrt{n}(1/\widehat{\lambda^{-1}}(n) - \lambda) \xrightarrow{d} \mathcal{N}(0, (1 + \alpha\psi_1(\alpha))\lambda^2)$, $n \rightarrow \infty$. Certainly, the asymptotic efficiency is the same:

$$\text{ARE}_{1/\widehat{\lambda^{-1}}(n)}(\alpha, \lambda) = \frac{\lambda^2\psi_1(\alpha)/(\alpha\psi_1(\alpha) - 1)}{(1 + \alpha\psi_1(\alpha))\lambda^2} = \frac{\psi_1(\alpha)}{\alpha^2\psi_1(\alpha)^2 - 1} > 0.964.$$

In a similar fashion, a closed-form unbiased estimator for $1/\alpha$ can be constructed as an U -statistic with kernel $K(X_1, X_2) = T_2(X_1, X_2)$. However, it is difficult to evaluate the asymptotic variance of the resulting estimator, because $\text{Var} \mathbf{E}(T_2(X_1, X_2)|X_2)$ is intractable. Ye and Chen proposed a different unbiased estimator for α^{-1} , which is equal to the UMVUE if and only if $n = 2$, namely

$$\widehat{\alpha^{-1}}(n) = \frac{1}{n-1} \frac{\sum_{i=1}^n X_i \log X_i - \bar{X} \sum_{i=1}^n \log X_i}{\bar{X}}.$$

From Theorem 3.2 in Ye and Chen (2017) it follows that $\sqrt{n}(1/\widehat{\alpha^{-1}}(n) - \alpha) \xrightarrow{d} \mathcal{N}(0, \alpha v(\alpha))$ and $\sqrt{n}(\widehat{\alpha^{-1}}(n) - \alpha^{-1}) \xrightarrow{d} \mathcal{N}(0, \alpha^{-3}v(\alpha))$, as $n \rightarrow \infty$, with $v(\alpha) = \alpha^2\psi_1(\alpha) + \alpha - 1$. Therefore, both closed-form estimators $\widehat{\alpha^{-1}}(n)$ (for $1/\alpha$) and $1/\widehat{\alpha^{-1}}(n)$ (for α) have asymptotic efficiency (compared to the MLE – see Remark 2.1) given by $\rho(\alpha) = [(\alpha\psi_1(\alpha) - 1)(\alpha^2\psi_1(\alpha) + \alpha - 1)]^{-1}$. Numerical computations show that $\rho(\alpha) \geq \rho(\alpha^*) = .9281\dots$ for all $\alpha > 0$ (where

$\alpha^* = .41541 \dots$), hence, the asymptotic efficiency is greater than 92.8%. It should be noted, however, that, due to Jensen's inequality, both estimators $1/\widehat{\lambda}^{-1}(n)$ (for λ) and $1/\widehat{\alpha}^{-1}(n)$ (for α) are positively biased.

The present powerful estimation procedure of Ye and Chen was further analyzed by Louzada *et al.* (2019), showing that the bias-corrected estimator

$$\widehat{\alpha}_n = \frac{n-3}{(n-1)\widehat{\alpha}^{-1}(n)} + \frac{2}{5(n+(n-1)\widehat{\alpha}^{-1}(n))} \left(\frac{11}{3} - \frac{2n}{n+(n-1)\widehat{\alpha}^{-1}(n)} \right)$$

improves upon the estimator $1/\widehat{\alpha}^{-1}(n)$ of α for small values of n .

3 Closed form estimators for the two-parameter Beta distribution

Through the present section we consider a random sample X_1, \dots, X_n ($n \geq 2$) from the Beta distribution with density as in (1.1), where the parameter $\theta = (\alpha, \beta) \in (0, \infty)^2$. The Fisher information matrix (based on a single observation) is given by

$$J = J(\alpha, \beta) = \begin{pmatrix} \psi_1(\alpha) - \psi_1(\alpha + \beta) & -\psi_1(\alpha + \beta) \\ -\psi_1(\alpha + \beta) & \psi_1(\beta) - \psi_1(\alpha + \beta) \end{pmatrix}, \text{ with inverse}$$

$$J^{-1} = \frac{1}{\psi_1(\alpha)\psi_1(\beta) - \psi_1(\alpha + \beta)(\psi_1(\alpha) + \psi_1(\beta))} \begin{pmatrix} \psi_1(\beta) - \psi_1(\alpha + \beta) & \psi_1(\alpha + \beta) \\ \psi_1(\alpha + \beta) & \psi_1(\alpha) - \psi_1(\alpha + \beta) \end{pmatrix},$$

with $\psi_1(x) = (\log \Gamma(x))''$, as in Section 2. The complete sufficient statistic has now the form $(X, Y) = (\prod_{i=1}^n X_i, \prod_{i=1}^n (1 - X_i))$, and an extra difficulty, compared to the Gamma-case, is due to the lack of independence of X and Y .

The proposed Ye-Chen-type (biased) estimators for α and β are as follows:

$$\widehat{\alpha}_n = \frac{n\bar{X}_n}{\sum_{i=1}^n X_i \log(X_i/(1 - X_i)) - \bar{X}_n \sum_{i=1}^n \log(X_i/(1 - X_i))}, \quad (3.1)$$

$$\widehat{\beta}_n = \frac{n(1 - \bar{X}_n)}{\sum_{i=1}^n X_i \log(X_i/(1 - X_i)) - \bar{X}_n \sum_{i=1}^n \log(X_i/(1 - X_i))}, \quad (3.2)$$

where $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$. In order to make clear how these estimators have been derived, we recall the Stein-type covariance identity for the Beta distribution,

$$\text{Cov}(X, w(X)) = \frac{1}{\alpha + \beta} \mathbf{E}X(1 - X)w'(X),$$

which holds true for sufficiently smooth functions, namely, when $w : (0, 1) \rightarrow \mathbb{R}$ is absolutely continuous and its derivative, w' , satisfies $\mathbf{E}X(1 - X)|w'(X)| < \infty$; see, e.g., Afendras *et al.* (2011). The special choice $w(x) = \log(x/(1 - x))$ leads to $\text{Cov}(X, w(X)) = 1/(\alpha + \beta)$. Therefore, the U -statistic U_n with kernel

$$K(X_1, X_2) = \frac{1}{2}(X_2 - X_1) \log \frac{X_2(1 - X_1)}{X_1(1 - X_2)}$$

is an unbiased estimator of the parameter $1/(\alpha + \beta)$. One finds

$$U_n = \frac{2 \sum_{i < j} K(X_i, X_j)}{n(n-1)} = \frac{1}{n-1} \left(\sum_{i=1}^n X_i \log(X_i/(1-X_i)) - \bar{X}_n \sum_{i=1}^n \log(X_i/(1-X_i)) \right) = \frac{nV_n}{n-1},$$

say. From the SLLN, $V_n \rightarrow \text{Cov}(X, \log(X/(1-X))) = 1/(\alpha + \beta)$, a.s., as $n \rightarrow \infty$; thus, $1/V_n \rightarrow \alpha + \beta$, a.s. Since, obviously, $\bar{X}_n \rightarrow \alpha/(\alpha + \beta)$ and $1 - \bar{X}_n \rightarrow \beta/(\alpha + \beta)$, we conclude that $\widehat{\alpha}_n = \bar{X}_n/V_n \rightarrow \alpha$ and $\widehat{\beta}_n = (1 - \bar{X}_n)/V_n \rightarrow \beta$, a.s., so both estimators are strongly consistent. In order to investigate their asymptotic efficiency, we have to calculate explicitly the variance of their asymptotic normal distribution. This is the subject of the following

Theorem 3.1. The joint limiting distribution of $\widehat{\alpha}_n, \widehat{\beta}_n$ (see (3.1), (3.2)) is normal. More precisely, as $n \rightarrow \infty$,

$$\sqrt{n} \begin{pmatrix} \widehat{\alpha}_n - \alpha \\ \widehat{\beta}_n - \beta \end{pmatrix} \xrightarrow{d} \mathcal{N}(\mathbf{0}, \Delta), \quad \text{where } \Delta = \frac{1}{\alpha + \beta + 1} \begin{pmatrix} v_{11}(\alpha, \beta) & v_{12}(\alpha, \beta) \\ v_{12}(\alpha, \beta) & v_{22}(\alpha, \beta) \end{pmatrix} \quad (3.3)$$

with

$$\begin{aligned} v_{11}(\alpha, \beta) &= \alpha \left(\alpha(\alpha + \beta + 1) + \alpha^2 \beta \left[\psi_1(\alpha) + \psi_1(\beta) \right] - \beta \right), \\ v_{12}(\alpha, \beta) &= \alpha^2 \beta + \alpha \beta^2 + \alpha^2 \beta^2 \left[\psi_1(\alpha) + \psi_1(\beta) \right] - \alpha^2 - \beta^2, \\ v_{22}(\alpha, \beta) &= \beta \left(\beta(\alpha + \beta + 1) + \alpha \beta^2 \left[\psi_1(\alpha) + \psi_1(\beta) \right] - \alpha \right). \end{aligned}$$

Proof: The result is quite simple, since it is a by-product of the CLT and the delta-method. However, the calculations are somewhat involved. We define the random variables $Y_i = \log(X_i/(1-X_i))$ and $Z_i = X_i \log(X_i/(1-X_i))$. It follows after some algebra that the mean vector and the dispersion matrix of (X_i, Y_i, Z_i) , $i = 1, \dots, n$, are given by

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{pmatrix} = \frac{1}{\alpha + \beta} \begin{pmatrix} \alpha \\ (\alpha + \beta)(\psi(\alpha) - \psi(\beta)) \\ 1 + \alpha(\psi(\alpha) - \psi(\beta)) \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_{11} & \sigma_{12} & \sigma_{13} \\ \sigma_{12} & \sigma_{22} & \sigma_{23} \\ \sigma_{13} & \sigma_{23} & \sigma_{33} \end{pmatrix},$$

where, with $\Psi = \psi(\alpha) - \psi(\beta)$ (recall $\psi(x) = (\log \Gamma(x))'$) and $\Psi_1 = \psi_1(\alpha) + \psi_1(\beta)$,

$$\begin{aligned} \sigma_{11} &= \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)}, \\ \sigma_{12} &= \frac{1}{\alpha + \beta}, \\ \sigma_{13} &= \frac{\beta + \alpha(\alpha + \beta) + \alpha\beta\Psi}{(\alpha + \beta)^2(\alpha + \beta + 1)}, \\ \sigma_{22} &= \Psi_1, \\ \sigma_{23} &= \frac{\Psi + \alpha\Psi_1}{\alpha + \beta}, \\ \sigma_{33} &= \frac{\alpha\beta\Psi^2 + 2(\alpha^2 + \alpha\beta + \beta)\Psi + \alpha(\alpha + 1)(\alpha + \beta)\Psi_1 + \alpha + \beta - 1}{(\alpha + \beta)^2(\alpha + \beta + 1)}. \end{aligned}$$

Hence, from the classical CLT,

$$\sqrt{n} \begin{pmatrix} \bar{X}_n - \mu_1 \\ \bar{Y}_n - \mu_2 \\ \bar{Z}_n - \mu_3 \end{pmatrix} \xrightarrow{d} \mathcal{N}(\mathbf{0}, \Sigma), \quad \text{as } n \rightarrow \infty,$$

where $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$, $\bar{Y}_n = n^{-1} \sum_{i=1}^n Y_i$, $\bar{Z}_n = n^{-1} \sum_{i=1}^n Z_i$. Consider the functions $k_1(x, y, z) = x/(z - xy)$, $k_2(x, y, z) = (1 - x)/(z - xy)$, so that $k_1(\bar{X}_n, \bar{Y}_n, \bar{Z}_n) = \widehat{\alpha}_n$ and $k_2(\bar{X}_n, \bar{Y}_n, \bar{Z}_n) = \widehat{\beta}_n$. The Jacobian matrix is

$$A(x, y, z) := \begin{pmatrix} \frac{\partial}{\partial x} k_1(x, y, z) & \frac{\partial}{\partial y} k_1(x, y, z) & \frac{\partial}{\partial z} k_1(x, y, z) \\ \frac{\partial}{\partial x} k_2(x, y, z) & \frac{\partial}{\partial y} k_2(x, y, z) & \frac{\partial}{\partial z} k_2(x, y, z) \end{pmatrix} = \frac{1}{(xy - z)^2} \begin{pmatrix} z & x^2 & -x \\ y - z & x - x^2 & x - 1 \end{pmatrix}.$$

Hence, with $A_0 := A(\mu_1, \mu_2, \mu_3)$, a straightforward application of the delta-method yields

$$\sqrt{n} \begin{pmatrix} k_1(\bar{X}_n, \bar{Y}_n, \bar{Z}_n) - k_1(\mu_1, \mu_2, \mu_3) \\ k_2(\bar{X}_n, \bar{Y}_n, \bar{Z}_n) - k_2(\mu_1, \mu_2, \mu_3) \end{pmatrix} = \sqrt{n} \begin{pmatrix} \widehat{\alpha}_n - \alpha \\ \widehat{\beta}_n - \beta \end{pmatrix} \xrightarrow{d} \mathcal{N}(\mathbf{0}, A_0 \Sigma A_0^T), \quad \text{as } n \rightarrow \infty,$$

because, obviously, $k_1(\mu_1, \mu_2, \mu_3) = \alpha$ and $k_2(\mu_1, \mu_2, \mu_3) = \beta$. Finally, a lengthy computation reveals that $A_0 \Sigma A_0^T = \Delta$ as in (3.3), completing the proof.

For any unbiased estimator U of $h(\alpha, \beta)$ (provided that such an estimator exists), the CR-bound yields the inequality $n \text{Var } U \geq \nabla h(\alpha, \beta)^T J^{-1} \nabla h(\alpha, \beta)$; in particular, with $h(\alpha, \beta) = \alpha$ we get

$$n \text{Var } U \geq \frac{\psi_1(\beta) - \psi_1(\alpha + \beta)}{\psi_1(\alpha)\psi_1(\beta) - \psi_1(\alpha + \beta)(\psi_1(\alpha) + \psi_1(\beta))},$$

and the right-hand side is the asymptotic variance of the MLE of α , which, as is well known, does not admit a closed form. For $h(\alpha, \beta) = \beta$ we obtain a similar inequality, interchanging the roles of α and β . Hence, from Theorem 3.1, the asymptotic efficiency $\rho_1 = \rho_1(\alpha, \beta)$ of $\widehat{\alpha}_n$ is given by

$$\rho_1 = \frac{(\alpha + \beta + 1)(\psi_1(\beta) - \psi_1(\alpha + \beta))/\alpha}{(\alpha(\alpha + \beta + 1) + \alpha^2\beta [\psi_1(\alpha) + \psi_1(\beta)] - \beta)(\psi_1(\alpha)\psi_1(\beta) - \psi_1(\alpha + \beta)(\psi_1(\alpha) + \psi_1(\beta)))},$$

while the asymptotic efficiency of $\widehat{\beta}_n$ equals to $\rho_2(\alpha, \beta) = \rho_1(\beta, \alpha)$. Despite the fact that the function ρ_1 is too complicated to enable a complete study, we can report some properties supported by numerical evidence. For fixed $\beta > 0$, define $\phi(\alpha) = \rho_1(\alpha, \beta)$. The function ϕ has limits 1, as $\alpha \rightarrow 0+$, and $\psi_1(\beta)/(\beta^2\psi_1(\beta)^2 - 1) > .964$, as $\alpha \rightarrow +\infty$; see Remark 2.2. Hence, either the infimum equals to the limit at $+\infty$, or it is attained at some point $\alpha^* = \alpha^*(\beta)$ such that $\phi'(\alpha^*) = 0$. Several plots showed that the graph of ϕ (for any fixed β) first decreases, then increases, and finally decreases. We calculated numerically the smallest stationary point, α^* , of ϕ (always lying between 0 and 1), resulting to the inequality $\phi(\alpha) \geq \min\{\phi(\alpha^*), .964\}$. Numerical values of $\rho_1^*(\beta) = \rho_1(\alpha^*(\beta), \beta)$, as well as minimization points $\alpha^*(\beta)$, are presented in Table 1. These values indicate that $\rho_1(\alpha^*(\beta), \beta) \rightarrow \min_{x>0} [(x\psi_1(x) - 1)(x^2\psi_1(x) + x - 1)]^{-1} = .9281 \dots$, and $\alpha^*(\beta) \rightarrow \alpha^* = .41541 \dots$ (α^* is the same as in Remark 2.2), as $\beta \rightarrow \infty$; however, we was not able to provide a proof.

The values of Table 1 indicate that for almost the entire parametric range, the asymptotic efficiency of $\widehat{\alpha}_n$ is no less that 90%. The same is clearly true for $\widehat{\beta}_n$, since $\rho_2(\alpha, \beta) = \rho_1(\beta, \alpha)$. This verifies that the proposed estimating procedure is quite powerful and, at the same time, it produces simple, closed-form, estimators.

Table 1.

Least asymptotic efficiency $\rho_1^*(\beta) = \min_{\alpha>0} \rho_1(\alpha, \beta) = \rho_1(\alpha^*(\beta), \beta)$, for $\beta = .05(.05)10$.

β	$\alpha^*(\beta)$	$\rho_1^*(\beta)$	β	$\alpha^*(\beta)$	$\rho_1^*(\beta)$	β	$\alpha^*(\beta)$	$\rho_1^*(\beta)$	β	$\alpha^*(\beta)$	$\rho_1^*(\beta)$
.05	.179398	.712626	.1	.229471	.784172	.15	.258632	.828715	.2	.277437	.85901
.25	.290137	.880541	.3	.298956	.89629	.35	.305205	.908065	.4	.309705	.917024
.45	.312992	.923944	.5	.315422	.929356	.55	.317239	.933636	.6	.318612	.937052
.65	.319661	.939802	.7	.320472	.942029	.75	.321107	.943844	.8	.321614	.945328
.85	.322026	.946546	.9	.322371	.947547	.95	.322669	.94837	1.	.322936	.949046
1.05	.323182	.9496	1.1	.323418	.950051	1.15	.32365	.950416	1.2	.323884	.950709
1.25	.324122	.95094	1.3	.324369	.951118	1.35	.324626	.951251	1.4	.324894	.951345
1.45	.325174	.951405	1.5	.325466	.951437	1.55	.325771	.951445	1.6	.326088	.95143
1.65	.326417	.951397	1.7	.326757	.951348	1.75	.327109	.951284	1.8	.327471	.951209
1.85	.327842	.951122	1.9	.328222	.951027	1.95	.32861	.950924	2.	.329006	.950814
2.05	.329409	.950697	2.1	.329818	.950576	2.15	.330232	.950451	2.2	.330652	.950321
2.25	.331075	.950189	2.3	.331502	.950054	2.35	.331933	.949917	2.4	.332366	.949778
2.45	.332801	.949638	2.5	.333239	.949496	2.55	.333677	.949354	2.6	.334117	.949212
2.65	.334557	.949069	2.7	.334997	.948926	2.75	.335438	.948783	2.8	.335878	.94864
2.85	.336317	.948497	2.9	.336756	.948355	2.95	.337194	.948214	3.	.33763	.948073
3.05	.338065	.947933	3.1	.338498	.947794	3.15	.33893	.947656	3.2	.339359	.947519
3.25	.339786	.947383	3.3	.340212	.947248	3.35	.340635	.947114	3.4	.341055	.946981
3.45	.341473	.946849	3.5	.341888	.946718	3.55	.342301	.946589	3.6	.342711	.946461
3.65	.343118	.946334	3.7	.343522	.946208	3.75	.343924	.946084	3.8	.344322	.945961
3.85	.344718	.945839	3.9	.34511	.945718	3.95	.3455	.945599	4.	.345886	.945481
4.05	.34627	.945364	4.1	.34665	.945248	4.15	.347027	.945134	4.2	.347401	.945021
4.25	.347772	.944909	4.3	.34814	.944798	4.35	.348505	.944689	4.4	.348867	.94458
4.45	.349226	.944473	4.5	.349581	.944367	4.55	.349934	.944262	4.6	.350284	.944159
4.65	.35063	.944056	4.7	.350974	.943955	4.75	.351314	.943855	4.8	.351652	.943756
4.85	.351986	.943658	4.9	.352318	.943561	4.95	.352647	.943465	5.	.352973	.94337
5.05	.353295	.943276	5.1	.353616	.943184	5.15	.353933	.943092	5.2	.354247	.943001
5.25	.354559	.942911	5.3	.354868	.942823	5.35	.355174	.942735	5.4	.355477	.942648
5.45	.355778	.942562	5.5	.356076	.942477	5.55	.356371	.942393	5.6	.356664	.94231
5.65	.356955	.942228	5.7	.357242	.942146	5.75	.357527	.942066	5.8	.35781	.941986
5.85	.35809	.941907	5.9	.358368	.941829	5.95	.358643	.941752	6.	.358916	.941676
6.05	.359186	.9416	6.1	.359454	.941525	6.15	.35972	.941451	6.2	.359983	.941378
6.25	.360244	.941305	6.3	.360503	.941234	6.35	.36076	.941163	6.4	.361014	.941092
6.45	.361266	.941023	6.5	.361516	.940954	6.55	.361764	.940886	6.6	.36201	.940818
6.65	.362254	.940751	6.7	.362495	.940685	6.75	.362735	.94062	6.8	.362972	.940555
6.85	.363208	.940491	6.9	.363441	.940427	6.95	.363673	.940364	7.	.363903	.940302
7.05	.36413	.94024	7.1	.364356	.940179	7.15	.36458	.940118	7.2	.364802	.940058
7.25	.365022	.939999	7.3	.365241	.93994	7.35	.365457	.939882	7.4	.365672	.939824
7.45	.365885	.939767	7.5	.366097	.93971	7.55	.366307	.939654	7.6	.366514	.939599
7.65	.366721	.939544	7.7	.366925	.939489	7.75	.367128	.939435	7.8	.36733	.939381
7.85	.36753	.939328	7.9	.367728	.939276	7.95	.367924	.939224	8.	.368119	.939172
8.05	.368313	.939121	8.1	.368505	.93907	8.15	.368695	.93902	8.2	.368884	.93897
8.25	.369072	.93892	8.3	.369258	.938872	8.35	.369443	.938823	8.4	.369626	.938775
8.45	.369808	.938727	8.5	.369988	.93868	8.55	.370167	.938633	8.6	.370345	.938587
8.65	.370521	.938541	8.7	.370696	.938495	8.75	.37087	.93845	8.8	.371042	.938405
8.85	.371214	.93836	8.9	.371383	.938316	8.95	.371552	.938272	9.	.371719	.938229
9.05	.371885	.938186	9.1	.37205	.938143	9.15	.372214	.938101	9.2	.372376	.938059
9.25	.372538	.938017	9.3	.372698	.937976	9.35	.372857	.937935	9.4	.373014	.937894
9.45	.373171	.937854	9.5	.373327	.937814	9.55	.373481	.937774	9.6	.373634	.937735
9.65	.373786	.937696	9.7	.373938	.937657	9.75	.374088	.937619	9.8	.374237	.937581
9.85	.374385	.937543	9.9	.374531	.937505	9.95	.374677	.937468	10.	.374822	.937431

Remark 3.1. (a) The estimators $\widehat{\alpha}_n$ and $\widehat{\beta}_n$ are positive (w.p. 1) for all $n \geq 2$. This is so because the denominator is a positive multiple of an U -statistic with symmetric kernel $K(X_1, X_2) > 0$, w.p. 1.

(b) The asymptotic covariance of $\widehat{\alpha}_n$ and $\widehat{\beta}_n$ is positive. This follows from the inequality $\psi_1(x) > x^{-2} > x^{-2} - x^{-1}$, $x > 0$, which implies that $p(x) := \psi_1(x) - (x^{-2} - x^{-1}) > 0$. Then, from (3.3), $v_{12}(\alpha, \beta) = \alpha^2 \beta^2 [p(\alpha) + p(\beta)] > 0$. This fact is somewhat counterintuitive, since $\widehat{\alpha}_n = \overline{X}_n/V_n$, $\widehat{\beta}_n = (1 - \overline{X}_n)/V_n$, $V_n > 0$, and the random variables \overline{X}_n and $1 - \overline{X}_n$ are, obviously, negatively correlated.

(c) The limiting distribution of the estimator U_n (which is unbiased for $1/(\alpha + \beta)$) can be derived from Theorem 3.1, since $U_n = (n/(n-1))V_n$ and thus, $\sqrt{n}(U_n - V_n) = o_p(1)$ and $\sqrt{n}(1/U_n - 1/V_n) = o_p(1)$, $n \rightarrow \infty$. Applying the delta-method to (3.3) with $k(x, y) = 1/(x+y)$ we obtain $\sqrt{n}(U_n - (\alpha + \beta)^{-1}) = \sqrt{n}(V_n - (\alpha + \beta)^{-1}) + o_p(1) \xrightarrow{d} \mathcal{N}(0, v^2)$, as $n \rightarrow \infty$, where

$$v^2 = \frac{v_{11}(\alpha, \beta) + v_{22}(\alpha, \beta) + 2v_{12}(\alpha, \beta)}{(\alpha + \beta)^4(\alpha + \beta + 1)} = \frac{\alpha + \beta + \alpha\beta [\psi_1(\alpha) + \psi_1(\beta)] - 1}{(\alpha + \beta)^2(\alpha + \beta + 1)}.$$

Consequently, $\sqrt{n}(1/V_n - (\alpha + \beta)) \xrightarrow{d} \mathcal{N}(0, (\alpha + \beta)^4 v^2)$, as $n \rightarrow \infty$. Using these formulas one can verify that the estimators U_n and $1/V_n$ exhibit high asymptotic efficiency.

(d) For small sample sizes, bias-corrected estimators in the sense of Louzada *et al.* may reduce the mean squared error; this, however, is beyond the scope of the present work.

References

- [1] Afendras, G.; Papadatos, N. (2015). Integrated Pearson family and orthogonality of the Rodrigues polynomials: A review including new results and an alternative classification of the Pearson system. *Applicationes Mathematicae*, **42**(2-3), 231–267.
- [2] Afendras, G.; Papadatos, N.; Papathanasiou, V. (2011). An extended Stein-type covariance identity for the Pearson family, with applications to lower variance bounds. *Bernoulli*, **17**(2), 507–529.
- [3] Glaser, R.E. (1976). The ratio of the geometric mean to the arithmetic mean for a random sample from a gamma distribution. *J. Amer. Statist. Assoc.*, **71**, 480–487.
- [4] Hoeffding, W. (1948). A class of statistics with asymptotically normal distribution. *Ann. Math. Statist.*, **19**(3), 293–325.
- [5] Iliopoulos, G. (2016). Exact confidence intervals for the shape parameter of the gamma distribution. *J. Statist. Comput. Simul.*, **86**(8), 1635–1642.
- [6] Louzada, F.; Ramos P.L.; Ramos, E. (2019). A note on bias of closed-form estimators for the gamma distribution derived from likelihood equations. *The American Statistician*, **73**(2), 195–199.
- [7] Nandi, S.B. (1980). On the exact distribution of the ratio of the weighted geometric mean to the unweighted arithmetic mean in samples from gamma distributions. *J. Amer. Statist. Assoc.*, **75**, 217–220.
- [8] Tang, J.; Gupta, A.K. (1984). On the distribution of the product of independent beta random variables. *Statist. Probab. Lett.*, **2**, 165–168.
- [9] Ye, Z.-S.; Chen, N. (2017). Closed-form estimators for the gamma distribution derived from likelihood equations. *The American Statistician*, **71**(2), 177–181.