

REDUCED TATE–SHAFAREVICH GROUP

HAYATO MORIMURA

ABSTRACT. Given two generic elliptic Calabi–Yau 3-folds over a common base in the sense of Căldăraru, we prove that if their derived categories are equivalent as a triangulated category linear over the base, then they share the common relative Jacobian. It follows that one of them is isomorphic to a relative moduli space of semistable sheaves of rank 1 with an appropriate degree on the fibers of the other. As a consequence, up to linear derived equivalence, generic elliptic Calabi–Yau 3-folds are classified by a certain quotient set of the Brauer group of their relative Jacobian.

1. INTRODUCTION

Let $\pi: X \rightarrow S$ be a *generic elliptic 3-fold* in the sense of [Că00, Definition 6.1.6], which is an elliptic fibration with $X, \dim X = 3$ and $S, \dim S = 2$ smooth over \mathbb{C} satisfying:

- π is flat;
- π does not have multiple fibers;
- π admits a multisection;
- the discriminant locus $\Delta \subset S$ is an integral curve in S with only nodes and cusps as singularities;
- the fiber over a general point of Δ is a rational curve with one node.

It is known that these properties are shared by general members of many families of elliptic 3-folds. When X is Calabi–Yau in the strict sense, we call π a *generic elliptic Calabi–Yau 3-fold*. There exists an étale cover $\{U_i\}_{i \in I}$ of S such that to each U_i the base change $\pi_i: X_i = X \times_S U_i \rightarrow U_i$ admits a section $\sigma_i: U_i \rightarrow X_i$.

Fix a line bundle $\mathcal{O}_{X/S}(1)$ on X relatively ample to S . Let $\varpi: M \rightarrow S$ be the relative coarse moduli space of semistable sheaves of rank 1, degree 0 on the fibers $X_s, s \in S$ of π with respect to the restrictions $\mathcal{O}_{X/S}(1)|_{X_s}$. By universality of M , the morphism ϖ admits a section $\sigma: S \rightarrow M$ whose image is contained in a unique component, called the *relative Jacobian*

$$f: J \rightarrow S.$$

In general, J has singularities all of which are ordinary double points. They arise from the base changes π_i by contracting all components of the fibers which do not intersect $\sigma_i(U_i), i \in I$. One can not always expect to resolve the singularities in the algebraic category. Let

$$\rho: \bar{J} \rightarrow J$$

be any analytic small resolution.

The *smooth part* of π is the base change

$$\pi_U: X_U = X \times_S U \rightarrow U = S \setminus \Delta.$$

Due to Ogg–Shafarevich theory developed by Dolgachev–Gross [DG94], smooth parts of generic elliptic 3-folds with relative Jacobian $f_U: J_U \rightarrow U$ are classified up to U -isomorphism

SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY
E-mail address: hmorimur@sissa.it.

by the Tate–Shafarevich group, which in our setting is isomorphic to the quotient

$$(1.1) \quad \mathrm{Br}(J_U) / \mathrm{Br}(U)$$

of the Brauer group $\mathrm{Br}(J_U)$, where we regard $\mathrm{Br}(U)$ as a subgroup via the pullback along π_U . The goal of this paper is to give a categorical interpretation of the Tate–Shafarevich group.

Let $\alpha_U \in \mathrm{Br}(J_U)$ be a Brauer class representing π_U . It uniquely lifts to $\alpha \in \mathrm{Br}(J)$ along the pullback by the open immersion $J_U \hookrightarrow J$. Note that the Brauer group is a birational invariant. We denote by $\bar{\alpha} \in \mathrm{Br}(\bar{J})$ the image of α under ρ^* , which is injective. In [Că102, Theorem 5.1] Căldăraru constructed an S -linear twisted Fourier–Mukai functor

$$(1.2) \quad \Phi_{\iota_* \mathcal{U}_{\bar{\alpha}}} : D^b(X) \rightarrow D^b(\bar{J}, \bar{\alpha}),$$

where $\mathcal{U}_{\bar{\alpha}}$ is a $\mathrm{pr}_2^* \bar{\alpha}^{-1}$ -twisted pseudo-universal sheaf on $X \times_S \bar{J}$ and $\iota : X \times_S \bar{J} \hookrightarrow X \times \bar{J}$ is the closed immersion. The equivalence (1.2) tells us that any S -linear triangulated category equivalent to $D^b(X)$ is represented by α .

He also constructed an S -linear Fourier–Mukai functor

$$(1.3) \quad \Phi_{\tau_* \mathcal{U}_d} : D^b(X) \rightarrow D^b(X^d),$$

for any d coprime to the order $\mathrm{ord}(\alpha_U)$ of α_U in $\mathrm{Br}(J_U)$, where $\pi_d : X^d \rightarrow S$ is another generic elliptic 3-fold called the d -th twisted power whose smooth part is represented by α_U^d , \mathcal{U}_d is a universal sheaf on $X \times_S X^d$, and $\tau : X \times_S X^d \hookrightarrow X \times X^d$ is the closed immersion. Recall that π_d is the relative moduli space of semistable sheaves of rank 1, degree d on the fibers X_s , $s \in S$ of π with respect to $\mathcal{O}_{X/S}(1)|_{X_s}$. The moduli problem is fine if and only if d is coprime to $\mathrm{ord}(\alpha_U)$ [Că100, Theorem 4.5.2]. In the sequel, we will call such π_d a *coprime d -th twisted power* of π . The equivalence (1.3) tells us that the group (1.1) is too big to classify, up to S -linear triangle equivalence, derived categories of generic elliptic 3-folds with relative Jacobian f .

We prove a reconstruction theorem of generic elliptic 3-folds with a common relative Jacobian modulo coprime twisted powers. Namely, all generic elliptic 3-folds over a fixed base S with relative Jacobian f are determined, modulo coprime twisted powers, by their S -linear derived categories.

Theorem 1.1 (Theorem 3.4). *Let $\pi : X \rightarrow S, \pi' : X' \rightarrow S$ be generic elliptic 3-folds with relative Jacobian $f : J \rightarrow S$. Assume that there exists an S -linear triangle equivalence $\Phi : D^b(X) \rightarrow D^b(X')$. Then π' is isomorphic to a coprime d -th twisted power of π for some $d \in \mathbb{Z}$.*

In particular, two generic elliptic 3-folds are S -linear derived-equivalent if and only if one of them is isomorphic to a coprime d -th twisted power of the other. Define an equivalence relation \sim_U in $\mathrm{Br}(J_U)$ as

$$(1.4) \quad \alpha_U \sim \beta_U \Leftrightarrow \beta_U = \alpha_U^d \text{ for some } d \text{ coprime to } \mathrm{ord}(\alpha_U),$$

which descends to that \sim in the quotient group (1.1). Then the quotient set

$$(\mathrm{Br}(J_U) / \mathrm{Br}(U)) / \sim$$

classifies up to S -linear triangle equivalence derived categories of generic elliptic 3-folds over S with relative Jacobian f .

Our proof works without assuming $\dim X = 3$. However, we are not interested in other cases mainly because this condition was responsible in [Că102, Theorem 5.1] for construction of $\mathcal{U}_{\bar{\alpha}}$. Antieau–Krashen–Ward proved a similar statement exploiting the twisted Brauer space [AKW17, Theorem 1.5, Remark 6.5]. It relies on the existence of a twisted relative Fourier–Mukai functor

$$\Phi_{\mathcal{P}} : D^b(\bar{J}, \bar{\alpha}) \simeq D^b(\bar{J}, \bar{\alpha}')$$

with kernel $\mathcal{P} \in D^b(\bar{J} \times_S \bar{J}')$, which practically would be obtained by concatenating (1.2) with (1.3). Here, we only assume the existence of an S -linear triangle equivalence Φ . Essentially, our contribution is to deduce the derived equivalence of the generic fibers from Φ . After that the argument in the proof of [AKW17, Theorem 1.5] would complete the proof. However, we exploit Ogg–Shafarevich theory to yield a geometric proof, following the original idea by Căldăraru.

The reconstruction problem will be more interesting when X is Calabi–Yau, as the derived category of a Calabi–Yau manifold might have nontrivial autoequivalences. Moreover, most of Calabi–Yau 3-folds admit elliptic fibrations. Some of them are connected with their mirrors admitting dual fibrations via relative Fourier–Mukai functors [HLS09, HM02], which can be seen as S -duality in string theory [Don98, DP08, DP12]. In this case, we strengthen Theorem 1.1 as follows.

Corollary 1.2 (Corollary 4.3). *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic Calabi–Yau 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π' is isomorphic to a coprime d -th twisted power of π for some $d \in \mathbb{Z}$.*

Here, we could remove the assumption from Theorem 1.1 on π, π' to share the common relative Jacobian due to our main result below.

Theorem 1.3 (Theorem 4.2). *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic Calabi–Yau 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π, π' share the relative Jacobian $f: J \rightarrow S$.*

In particular, two generic elliptic Calabi–Yau 3-folds are S -linear derived-equivalent if and only if one of them is isomorphic to a coprime d -th twisted power of the other. Note that if X is Calabi–Yau in the strict sense, so is X' . The key step is reconstruction of fibers based on [LT17]. Also [AKW17, Lemma 2.4] is important to find the generic fibers of X, X' sharing the common Jacobian. We use isomorphisms $\omega_X \cong \mathcal{O}_X, \omega_{X'} \cong \mathcal{O}_{X'}$ to guarantee the coincidence of base changes of relative Jacobians of X, X' to any sufficiently small open subset.

Now, since $\text{Br}(S)$ vanishes [Gro94, Proposition 2.3] and the Brauer group is a birational invariant [Gro68], we obtain an isomorphism

$$\text{Br}(J) = \text{Br}(J)/\text{Br}(S) \rightarrow \text{Br}(J_U)/\text{Br}(U) = \text{Br}(J_U)$$

induced by the pullback along $J_U \hookrightarrow J$. Thus the Tate–Shafarevich group (1.1) gets reduced to the quotient set of $\text{Br}(J)$ by the equivalence relation induced by \sim_U to classify S -linear derived categories of generic elliptic Calabi–Yau 3-folds.

We emphasize that a generic elliptic Calabi–Yau 3-fold π is S -linear derived-equivalent to another one π' if and only if π' is the relative moduli space of semistable sheaves of rank 1 with an appropriate degree. It would be interesting to find such a moduli structure via our result in, for example, the several pairs of Fourier–Mukai partners constructed by Inoue [Ino22], whose derived equivalences follow from homological projective duality for categorical joins developed by Kuznetsov and Perry [KP21].

Notations and conventions. *Throughout the paper, all \mathbb{C} -varieties are integral separated scheme of finite type over \mathbb{C} . Via Serre’s GAGA theorem we will go back and forth from the algebraic to analytic categories for smooth proper \mathbb{C} -varieties, abusing notations without much comments. For any morphism $\pi: X \rightarrow S$ of nonsingular \mathbb{C} -varieties, we consider the canonical S -linear structure on $D^b(X) \simeq \text{perf}(X)$ given by the action*

$$\text{perf}(X) \times \text{perf}(S) \rightarrow \text{perf}(X), (F, G) \mapsto F \otimes_{\mathcal{O}_X} \pi^* G.$$

Then for morphisms $\pi: X \rightarrow S, \pi': X' \rightarrow S$ of nonsingular \mathbb{C} -varieties, a triangle functor $\Phi: D^b(X) \rightarrow D^b(X')$ is S -linear if Φ respects the S -linear structure, i.e., we have functorial isomorphisms

$$\Phi(F \otimes_{\mathcal{O}_X} \pi^* G) \cong \Phi(F) \otimes_{\mathcal{O}_{X'}} \pi'^* G, \quad F \in D^b(X), G \in D^b(S).$$

Since up to isomorphism all Brauer groups we consider are only of smooth quasi-projective \mathbb{C} -varieties, we use a single symbol \mathbf{Br} to denote the Brauer group and the cohomological Brauer group with respect to both étale and analytic topology. See [Jon, Theorem 1.1] and [Că100, Theorem 1.1.4].

Acknowledgements. The author is supported by SISSA PhD scholarships in Mathematics.

2. RECONSTRUCTION OF FIBERS

Lemma 2.1. Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be morphisms of \mathbb{C} -varieties with X', S smooth over \mathbb{C} and $\pi_Z: X_Z \rightarrow Z, \pi'_Z: X'_Z \rightarrow Z$ their base changes to $Z = S \setminus U$ for any open subset $U \subset S$. Then every S -linear triangle functor $\Phi: D^b(X) \rightarrow D^b(X')$ restricts to

$$\Phi_Z: D^b_{\text{coh}_{X_Z}(X)} \rightarrow D^b_{\text{coh}_{X'_Z}(X')},$$

where $D^b_{\text{coh}_{X_Z}(X)} \subset D^b(X), D^b_{\text{coh}_{X'_Z}(X')} \subset D^b(X')$ are the full S -linear triangulated subcategories with cohomology supported on X_Z, X'_Z respectively.

Proof. For each $E \in D^b(X)$ we have

$$\Phi(E) \cong \Phi\left(\bigoplus_i \mathcal{H}^i(E)[-i]\right) \cong \bigoplus_i \Phi(\mathcal{H}^i(E))[-i]$$

as Φ is an S -linear triangle functor. Consider the pullback diagrams

$$\begin{array}{ccc} X_Z & \xrightarrow{\bar{\iota}} & X \\ \bar{\pi} \downarrow & & \downarrow \pi \\ Z & \xrightarrow{\iota} & S \end{array}, \quad \begin{array}{ccc} X'_Z & \xrightarrow{\bar{\iota}'} & X' \\ \bar{\pi}' \downarrow & & \downarrow \pi' \\ Z & \xrightarrow{\iota'} & S \end{array}.$$

If E is supported on X_Z then we have

$$\begin{aligned} \Phi(\mathcal{H}^i(E)) &\cong \Phi(\mathcal{H}^i(E) \otimes_{\mathcal{O}_X} \bar{\iota}_* \mathcal{O}_{X_Z}) \\ &\cong \Phi(\mathcal{H}^i(E) \otimes_{\mathcal{O}_X} \pi^* \iota_* \mathcal{O}_Z) \\ &\cong \Phi(\mathcal{H}^i(E)) \otimes_{\mathcal{O}_{X'}} \pi'^* \iota_* \mathcal{O}_Z \\ &\cong \Phi(\mathcal{H}^i(E)) \otimes_{\mathcal{O}_{X'}} \bar{\iota}'_* \mathcal{O}_{X'_Z}, \end{aligned}$$

where the first, the second, the third, and the fourth isomorphisms follow from E being supported on X_Z , the isomorphism $\bar{\iota}_* \bar{\pi}^* \mathcal{O}_Z \cong \pi^* \iota_* \mathcal{O}_Z$, S -linearity of Φ , and the isomorphism $\bar{\iota}'_* \bar{\pi}'^* \mathcal{O}_Z \cong \pi'^* \iota_* \mathcal{O}_Z$ respectively. Since X' is smooth over \mathbb{C} , replacing it with a quasi-isomorphic object if necessary, we may assume that $\Phi(\mathcal{H}^i(E))$ are perfect. Then the last term becomes $\bar{\iota}'_*(\Phi(\mathcal{H}^i(E))|_{X'_Z})$ by the projection formula. Note that $\bar{\iota}'_*$ is exact as $\bar{\iota}'$ is affine. Hence $\Phi(\mathcal{H}^i(E))$ is supported on X'_Z , which completes the proof. \square

Lemma 2.2. Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be flat morphisms of nonsingular \mathbb{C} -varieties. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then every closed fibers X_s, X'_s over $s \in S$ are derived-equivalent.

Proof. Take an affine open subset $U = \text{Spec } R \subset S$. First, we show that Φ induces an R -linear triangle equivalence

$$\Phi_U: D^b(X_U) \rightarrow D^b(X'_U),$$

where $\pi_U: X_U \rightarrow U, \pi'_U: X'_U \rightarrow U$ are the base changes to U . For their complements $X_Z = X \setminus X_U, X'_Z = X' \setminus X'_U$ we have

$$\text{coh}(X_U) \simeq \text{coh}(X)/\text{coh}_{X_Z}(X), \text{coh}(X'_U) \simeq \text{coh}(X')/\text{coh}_{X'_Z}(X'),$$

where the right hand sides are the quotients by the Serre subcategories

$$\text{coh}_{X_Z}(X) \subset \text{coh}(X), \text{coh}_{X'_Z}(X') \subset \text{coh}(X')$$

of coherent sheaves supported on X_Z, X'_Z respectively. Passing to their derived categories, via [Miy91, Theorem 3.2] we obtain

$$D^b(X_U) \simeq D^b(X)/D^b_{\text{coh}_{X_Z}(X)}(X), D^b(X'_U) \simeq D^b(X')/D^b_{\text{coh}_{X'_Z}(X')}(X').$$

Since X', S are smooth over \mathbb{C} , one can apply Lemma 2.1 to obtain the induced equivalence

$$\Phi_U: D^b(X_U) \simeq D^b(X)/D^b_{\text{coh}_{X_Z}(X)}(X) \rightarrow D^b(X'_U) \simeq D^b(X')/D^b_{\text{coh}_{X'_Z}(X')}(X')$$

by universality of Verdier quotients.

Next, we show that Φ_U induces an equivalence $\Phi_s: \text{perf}(X_s) \rightarrow \text{perf}(X'_s)$ of categories of perfect complexes on every closed fibers X_s, X'_s over $s \in S$. Consider dg enhancements $\text{Perf}(X), \text{Perf}(X')$ of $\text{perf}(X) \simeq D^b(X), \text{perf}(X') \simeq D^b(X')$, which are unique up to quasi-equivalence [CS18, Proposition 6.10]. Here, we use the assumption on X to be smooth over \mathbb{C} . Since π, π' are flat, one can apply [BFN10, Theorem 1.2] and [Töe, Exercise 32] to the pullback diagrams

$$\begin{array}{ccc} X_s & \longrightarrow & X_U \\ \downarrow & & \downarrow \\ \text{Spec } k(s) & \longrightarrow & \text{Spec } R \end{array}, \quad \begin{array}{ccc} X'_s & \longrightarrow & X'_U \\ \downarrow & & \downarrow \\ \text{Spec } k(s) & \longrightarrow & \text{Spec } R \end{array}$$

to obtain a quasi-equivalence $\text{Perf}(X_s) \rightarrow \text{Perf}(X'_s)$. Passing to their homotopy, we obtain the induced equivalence

$$\Phi_s: \text{perf}(X_s) \rightarrow \text{perf}(X'_s).$$

Now, the claim follows from [CS18, Proposition 7.4]. \square

Lemma 2.3 ([Cäl00, Theorem 6.1.9]). *Let $\pi: X \rightarrow S$ be a generic elliptic 3-fold. Then the closed fiber X_s over $s \in S$ is:*

- a smooth elliptic curve when $s \in S \setminus \Delta$;
- a rational curve with one node when $s \in \Delta$ is a smooth point;
- two copies of \mathbb{P}^1 intersecting transversely at two points when $s \in \Delta$ is a node;
- a rational curve with one cusp when $s \in \Delta$ is a cusp.

Proposition 2.4. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then every closed fibers X_s, X'_s over $s \in S$ are isomorphic.*

Proof. By Lemma 2.2 every closed fibers X_s, X'_s over $s \in S$ are derived-equivalent. Since by Lemma 2.3 X_s, X'_s are either a smooth elliptic curve, a rational curve, two copies of \mathbb{P}^1 intersecting transversely at two points, or a rational curve with one cusp, one can apply [LT17, Theorem 7.4(2)] to obtain $X_s \cong X'_s$. \square

3. RECONSTRUCTION OF FIBRATIONS

Lemma 3.1. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be separated morphisms of nonsingular \mathbb{C} -varieties. Then every S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$ induces an $K(S)$ -linear triangle equivalence $\Phi_{K(S)}: D^b(X_\eta) \rightarrow D^b(X'_\eta)$ on the generic fibers X_η, X'_η , where $K(S)$ is the function field of S .*

Proof. By the same argument as in the proof of Lemma 2.2 and [Har77, Corollary III10.7] we may assume that π, π' are smooth over $S = \text{Spec } R$. One can apply [Mor, Theorem 1.1] to obtain $K(S)$ -linear triangle equivalences

$$D^b(X_\eta) \rightarrow D^b(X)/D_0^b(X), \quad D^b(X'_\eta) \rightarrow D^b(X')/D_0^b(X').$$

to the Verdier quotients by the full S -linear triangulated subcategories spanned by complexes with coherent R -torsion cohomology. Since Φ is S -linear, universality of Verdier quotients induces the desired equivalence. \square

Corollary 3.2. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then their generic fibers X_η, X'_η share the common Jacobian.*

Proof. The generic fibers X_η, X'_η are derived-equivalent by Lemma 3.1. Now, the claim follows from [AKW17, Lemma 2.4]. \square

Lemma 3.3. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be smooth elliptic 3-folds with relative Jacobian $f: J \rightarrow S$. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π' is isomorphic to a coprime d -th twisted power of π for some $d \in \mathbb{Z}$.*

Proof. Let $\alpha, \alpha' \in \text{Br}(J)$ be representatives of the Tate–Shafarevich group for π, π' . Take an étale cover $\{U_i\}_{i \in I}$ of S representing both α and α' . Namely, to each U_i the base changes π_i, π'_i admit sections

$$\sigma_i: U_i \rightarrow X_i = X \times_S U_i, \quad \sigma'_i: U_i \rightarrow X'_i = X' \times_S U_i$$

and α, α' give the gluing data for $\{X_i\}_{i \in I}, \{X'_i\}_{i \in I}$ to yield X, X' from $\{J_i\}_{i \in I}$ respectively [Că100, Chapter 4.2, 4.4] via the description of Brauer classes as gerbes [Că100, Chapter 1.1]. Let $\alpha_\eta, \alpha'_\eta \in \text{Br}(J_\eta)$ be the images of α, α' under the injective map induced by the pullback along the base change $J_\eta \rightarrow J$ to the function field $K(S)$ of S . Refining it if necessary, we may assume that the induced étale cover $\{U_i \times_S K(S)\}$ of $\text{Spec } K(S)$ represents both α_η and α'_η .

By assumption and Lemma 3.1 the generic fibers X_η, X'_η are derived-equivalent. From [AKW17, Lemma 2.3] it follows that X'_η is isomorphic to the moduli space of degree d line bundles on X_η for some d coprime to the order $\text{ord}(\alpha_\eta)$ in $\text{Br}(J_\eta)$. Hence both α_η^d and α'_η represent π'_η . Since the map $\text{Br}(J) \rightarrow \text{Br}(J_\eta)$ is injective, both α^d and α' represent π' . As explained in [Că100, Chapter 4.5], the Brauer class α^d represents the d -th twisted power of X . Thus π' is isomorphic to a coprime d -th twisted power of π . \square

Remark 3.1. If the equivalence Φ was given by a relative Fourier–Mukai functor, one could have used [AKW17, Theorem 1.5]. However, still one should deduce the derived equivalence $D^b(J_\eta, \alpha_\eta) \simeq D^b(J_\eta, \alpha'_\eta)$ on the twisted generic fibers via a different way from [AKW17, Proposition 6.2]. We will discuss and remedy this issue in the forthcoming work [MU].

Theorem 3.4. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic 3-folds with relative Jacobian $f: J \rightarrow S$. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π' is isomorphic to a coprime d -th twisted power of π for some $d \in \mathbb{Z}$.*

Proof. Let $\alpha_U, \alpha'_U \in \text{Br}(J_U)$ be the representatives of the Tate–Shafarevich group for the smooth parts π_U, π'_U . By the same argument as in the proof of Lemma 2.2 there is an U -linear triangle equivalence

$$D^b(X_U) \rightarrow D^b(X'_U).$$

One can apply Lemma 3.3 to find some d coprime to $\text{ord}(\alpha_U)$ in $\text{Br}(J_U)$ such that both α_U^d and α'_U represent π'_U . By [Că100, Theorem 6.5.1] or birational invariance of the Brauer group, α_U, α'_U uniquely lift to $\alpha, \alpha' \in \text{Br}(J)$ along the pullback by the open immersion $J_U \hookrightarrow J$.

Let $\tilde{f}: \tilde{J} \rightarrow S$ be the composition of the resolution $\tilde{J} \rightarrow J$ with f . Take an analytic open cover $\{U_i\}_{i \in I}$ of S representing both α and α' such that each U_i contains at most one node. By [Că100, Theorem 6.4.6] there are U_i -isomorphisms

$$X_i \cong \tilde{J}_i = \tilde{J} \times_S U_i \cong X'_i$$

of analytic spaces. As explained in the proof of [Că100, Theorem 6.5.1], both α and α' give the gluing data for $\{X_i\}_{i \in I}, \{X'_i\}_{i \in I}$ to yield X, X' from $\{\tilde{J}_i\}_{i \in I}$. See, also [Că100, Theorem 4.4.1].

Let $\pi'': X'' \rightarrow S$ be the d -th twisted power of π . Refining the analytic open cover $\{U_i\}_{i \in I}$ if necessary, we may assume that to each U_i the base change π''_i admits a section

$$\sigma''_i: U_i \rightarrow X''_i = X'' \times_S U_i.$$

Again, by [Că100, Theorem 6.4.6] there are U_i -isomorphisms

$$X''_i \cong \tilde{J}_i$$

of analytic spaces. Then α^d gives the gluing data for $\{X''_i\}_{i \in I}$ to yield X'' from $\{\tilde{J}_i\}_{i \in I}$. Hence π' and π'' are isomorphic. \square

Remark 3.2. In the above proof we have not used the assumption on X to be 3-dimensional. However, this is crucial for Lemma 2.3 on the classification of the fibers, which in turn is responsible for the construction of local universal sheaves \mathcal{U}_i on $W_i = X_i \times_{U_i} \tilde{J}_i$ [Că100, Theorem 6.4.2]. The collection defines the $\text{pr}_2^* \bar{\alpha}^{-1}$ -twisted pseudo-universal sheaf $\mathcal{U}_{\bar{\alpha}} = (\{\mathcal{U}_i\}, \{\varphi_{ij}\})$ on $W = X \times_S \tilde{J}$, where $\varphi_{ij}: \mathcal{U}_j|_{W_i \times_W W_j} \rightarrow \mathcal{U}_i|_{W_i \times_W W_j}$ is an isomorphism on each double intersection $W_{ij} = W_i \times_W W_j$. Hence without the assumption one would not have the equivalence (1.2).

Remark 3.3. For the convenience of readers we add a little more explanation to the above remark. The Brauer class $\bar{\alpha}$ appeared in (1.2) gives the obstruction against the collection $\mathcal{U}_{\bar{\alpha}}$ to glue to yield a universal sheaf. The discrepancy between the composition $\varphi_{ij} \circ \varphi_{jk} \circ \varphi_{ki}$ and the identity on each triple intersection $W_{ijk} = W_i \times_W W_j \times_W W_k$ defines precisely the $\text{pr}_2^* \bar{\alpha}^{-1}$ -twisting of $\mathcal{U}_{\bar{\alpha}}$. Recall that $\bar{\alpha}$ is the image of α under the injection ρ^* . Consider the image α_X of α under the composition of the isomorphism $\text{Br}(J) \cong \text{Br}(J_U)$ with the projection $\text{Br}(J_U) \rightarrow \text{Br}(J_U)/\text{Br}(U)$. Then [Că100, Theorem 4.4.1] claims that α_X corresponds to π_U via Ogg–Shafarevich theory. Namely, α_X gives the gluing data for J_U cut into pieces to yield X_U via the description of Brauer classes as gerbes. Hence up to elements of $\text{Br}(U)$ the obstruction $\bar{\alpha}$ can be interpreted geometrically, which was tacitly used to obtain the equivalence (1.2)

4. CALABI–YAU CASE

Lemma 4.1. *Let $\pi: X \rightarrow S$ be a projective morphism of nonsingular \mathbb{C} -varieties with $\omega_X \cong \mathcal{O}_X$ such that every closed fiber $X_s, s \in S$ is connected. Then the relative canonical sheaf $\omega_{X/S}$ is invertible and isomorphic to $\pi^* \omega_S^{-1}$. In particular, we have $\pi_* \omega_{X/S} \cong \omega_S^{-1}$.*

Proof. Consider the first fundamental exact sequence

$$(4.1) \quad \pi^* \Omega_S \xrightarrow{u} \Omega_X \rightarrow \Omega_{X/S} \rightarrow 0.$$

If u is injective, then we obtain $\omega_{X/S} \cong \pi^* \omega_S^{-1}$, which is invertible. Since X, S are smooth over \mathbb{C} , for any $x \in X$ there is an open neighborhood $V \subset X$ on which u can be expressed as a homomorphism $\mathcal{O}_V^{\oplus \dim S} \rightarrow \mathcal{O}_V^{\oplus \dim X}$ of free \mathcal{O}_V -modules. As X is integral, it suffices to check the injectivity on the generic point. However, (4.1) restricts to yield the short exact sequence

$$0 \rightarrow \pi_U^* \Omega_U \xrightarrow{u|_{X_U}} \Omega_{X_U} \rightarrow \Omega_{X_U/U} \rightarrow 0.$$

on the smooth part $\pi_U: X_U \rightarrow U$ of π for some open subset $U \subset S$ [Har77, Corollary III10.7].

The second claim follows from the projection formula, since every closed fiber $X_s, s \in S$ is connected, by [Har77, Corollary III 12.9] the coherent sheaf $\pi_* \mathcal{O}_X$ is invertible and hence is isomorphic to \mathcal{O}_S . \square

Theorem 4.2. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic Calabi–Yau 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π, π' share the common relative Jacobian $f: J \rightarrow S$.*

Proof. By Proposition 2.4 every closed fibers X_s, X'_s over $s \in S$ are isomorphic. It follows that π, π' share the common discriminant locus $\Delta \subset S$. Let $f: J \rightarrow S, f': J' \rightarrow S$ be the relative Jacobians of π, π' . For any analytic small resolutions $\rho: \bar{J} \rightarrow J, \rho': \bar{J}' \rightarrow J'$ of singularities, we denote by \bar{f}, \bar{f}' the compositions $f \circ \rho, f' \circ \rho'$ respectively. Take an analytic open cover $\{U_i\}_{i \in I}$ of S such that each U_i contains at most one node and to each U_i the base changes $\pi_i, \pi'_i, \bar{f}_i, \bar{f}'_i$ admit sections

$$\sigma_i: U_i \rightarrow X_i, \sigma'_i: U_i \rightarrow X'_i, \bar{\sigma}_i: U_i \rightarrow \bar{J}_i = \bar{J} \times_S U_i, \bar{\sigma}'_i: U_i \rightarrow \bar{J}'_i = \bar{J}' \times_S U_i.$$

Applying [Nak87, Theorem 2.1] or [DG94, Theorem 2.3], we obtain birational morphisms

$$g_i: \bar{J}_i \rightarrow W(\mathcal{L}_i, a_i, b_i), g'_i: \bar{J}'_i \rightarrow W(\mathcal{L}'_i, a'_i, b'_i),$$

where $W(\mathcal{L}_i, a_i, b_i), W(\mathcal{L}'_i, a'_i, b'_i)$ are the Weierstrass fibrations associated with line bundles $\mathcal{L}_i, \mathcal{L}'_i$ on U_i , global sections a_i, a'_i of $\mathcal{L}_i^{\otimes 4}, \mathcal{L}'_i{}^{\otimes 4}$, and b_i, b'_i of $\mathcal{L}_i^{\otimes 6}, \mathcal{L}'_i{}^{\otimes 6}$ such that $4a_i^3 + 27b_i^2, 4a_i'^3 + 27b_i'^2$ are nonzero global sections of $\mathcal{L}_i^{\otimes 12}, \mathcal{L}'_i{}^{\otimes 12}$. Since g_i, g'_i are the morphism which blow down all components of fibers not intersecting $\bar{\sigma}_i(U_i), \bar{\sigma}'_i(U_i)$ respectively, the fibrations $W(\mathcal{L}_i, a_i, b_i) \rightarrow S, W(\mathcal{L}'_i, a'_i, b'_i) \rightarrow S$ coincide with the base changes of the relative Jacobians f_i, f'_i to U_i . See also [Că102, Theorem 6.4.3].

From Lemma 4.1 it follows

$$\mathcal{L}_i \cong \bar{f}_{i*} \omega_{J_i/U_i} \cong \omega_{U_i}^{-1} \cong \bar{f}'_{i*} \omega_{J'_i/U_i} \cong \mathcal{L}'_i$$

as we have U_i -isomorphisms $X_i \cong \bar{J}_i, X'_i \cong \bar{J}'_i$ of analytic spaces. Note that \bar{f}_i, \bar{f}'_i are locally projective, as they are proper and \bar{J}_i, \bar{J}'_i have relatively ample line bundles [SP, Tag 01WC]. Refining $\{U_i\}_{i \in I}$ if necessary, we may assume that \bar{f}_i, \bar{f}'_i are projective. Since by Corollary 3.2 the generic fibers J_η, J'_η are isomorphic, we obtain U_i -isomorphisms

$$(4.2) \quad W(\mathcal{L}_i, a_i, b_i) \cong W(\mathcal{L}'_i, a'_i, b'_i),$$

as two Weierstrass fibrations $W(\mathcal{L}_i, a_i, b_i), W(\mathcal{L}'_i, a'_i, b'_i)$ must coincide whenever they share the common generic fiber.

Now, f, f' are locally isomorphic elliptic fibrations over the common base S . Since the construction of a Weierstrass fibration is functorial, we have the U_{ij} -isomorphism

$$W(\mathcal{L}_i|_{U_{ij}}, a_i|_{U_{ij}}, b_i|_{U_{ij}}) \cong W(\mathcal{L}'_i, a'_i, b'_i)|_{U_{ij}}$$

for each double intersection $U_{ij} = U_i \times_S U_j$. Then the U_i -isomorphisms (4.2) canonically glue to yield an S -isomorphism $J \rightarrow J'$, which completes the proof. \square

Corollary 4.3. *Let $\pi: X \rightarrow S, \pi': X' \rightarrow S$ be generic elliptic Calabi–Yau 3-folds. Assume that there exists an S -linear triangle equivalence $\Phi: D^b(X) \rightarrow D^b(X')$. Then π' is isomorphic to a coprime d -th twisted power of π for some $d \in \mathbb{Z}$.*

Proof. The claim follows immediately from Theorem 4.2 and Theorem 3.4. □

Remark 4.1. It sufficed to assume either X or X' is Calabi–Yau. Without loss of generality we may assume that X is Calabi–Yau. The triviality of the canonical bundle of X' follows from the uniqueness of Serre functors. The vanishing of $H^1(X', \mathcal{O}_{X'})$ follows from $H^2(X, \mathcal{O}_X) = H^2(X', \mathcal{O}_{X'})$ and Serre duality. Note that the S -linear triangle equivalence is in particular \mathbb{C} -linear triangle equivalence and hence naturally isomorphic to a unique Fourier Mukai functor with kernel belonging to $D^b(X \times X')$ unique up to isomorphism [Orl97, Section 2]. One obtains the induced isometry

$$H^2(X, \mathcal{O}_X) \oplus H^4(X, \mathcal{O}_X) \cong H^2(X', \mathcal{O}_{X'}) \oplus H^4(X', \mathcal{O}_{X'})$$

from [Căl00, Corollary 3.1.13, 3.1.14].

Remark 4.2. If X is Calabi–Yau, then \bar{J} is also Calabi–Yau. Again, the triviality of the canonical bundles of \bar{J} follows from the uniqueness of Serre functors. Refining it if necessary, we may assume that to each U_i the base change $\bar{f}_i = f_i \circ \rho_i$ is projective. Then by [CR11, Corollary 3.2.10] the higher direct image $R^j \rho_* \mathcal{O}_{\bar{J}}$ vanishes for each $j > 0$. Hence we obtain $R\bar{f}_* \mathcal{O}_{\bar{J}} = \mathcal{O}_S$ and Leray spectral sequence implies

$$H^1(\bar{J}, \mathcal{O}_{\bar{J}}) = H^1(S, \mathcal{O}_S).$$

Since X is Calabi–Yau the base S must be either a rational or Enriques surface by [Gro94, Proposition 2.3]. In both cases $H^1(S, \mathcal{O}_S)$ vanishes.

REFERENCES

- [AKW17] B. Antieau, D. Krashen, and M. Ward, *Derived categories of torsors for abelian schemes*, Adv. Math. 306(10), 1-23 (2017).
- [BFN10] D. Ben-zvi, J. Francis, and D. Nadler, *Integral transforms and Drinfeld centers in derived algebraic geometry*, J. Amer. Math. Soc. 23(4), 909-966 (2010).
- [Căl00] A. Căldăraru, *Derived categories of twisted sheaves on Calabi–Yau manifolds*, Ph.D. thesis, Cornell University (2000).
- [Căl02] A. Căldăraru, *Derived categories of twisted sheaves on elliptic threefolds*, J. Reine Angew. Math. 544, 161-179 (2002).
- [CR11] A. Chatzistamatiou and K. Rülling, *Higher direct images of the structure sheaf in positive characteristic*, Algebra Number Theory, 5(6) 693–775 (2011).
- [CS18] A. Canonaco and P. Stellari, *Uniqueness of dg enhancements for the derived category of a Grothendieck category*, J. Eur. Math 20(11), 2607–2641 (2018).
- [DG94] I. Dolgachev and M. Gross, *Elliptic three-folds I: Ogg–Shafarevich theory*, J. Alg. Geom. 3, 39-80 (1994).
- [Don98] R. Donagi, *Taniguchi lectures on principal bundles on elliptic fibrations* in Integral systems and algebraic geometry (Kobe / Kyoto, 1997), World Science publishing, (1998).
- [DP08] R. Donagi and T. Pantev, *Torus fibrations, gerbes, and duality*, Mem. Amer. Math. Soc. 193(901) (2008).
- [DP12] R. Donagi and T. Pantev, *Langrands duality for Hitchin systems*, Invent. Math. 189, 653-735 (2012).
- [Gro94] M. Gross, *Finiteness theorems for elliptic Calabi–Yau threefolds*, Duke. Math. 74(2), 271-299 (1994).
- [Gro68] A. Grothendieck, *Le group de Brauer, I, II, III*, in Dix Exposés sur la Cohomologie des schémas, North–Holland, Amsterdam, 46-188 (1968).
- [Har77] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, 52, Springer-Verlag, (1977).
- [HLS09] D. Hernández Ruipérez, A.C. López Martín and F. Sancho de Salas, *Relative integral functors for singular fibrations and singular partners*, J. Eur. Math. Soc. 11, 597-625 (2009).
- [HM02] D. Hernández Ruipérez and J. M. Muñoz Porras, *stable sheaves on elliptic fibrations*, J. Geom. Phys. 43, 163-183 (2002).

- [Ino22] D. Inoue, *Calabi–Yau 3-folds from projective joins of del Pezzo manifolds*, Adv. Math. 400, <https://doi.org/10.1016/j.aim.2022.108243>
- [Jon] J. Jong, *A result of Gabber*, <https://www.math.columbia.edu/dejong/papers/2-gabber.pdf>.
- [KP21] A. Kuznetsov and A. Perry, *Categorical joins*, J. Amer. Math. Soc. 34(2), 505-564 (2021).
- [LT17] A.C. López Martín and C. Tejero Prieto, *Derived equivalence and Kodaira fibers*, J. Geom. Phys. 122, 69-79 (2017).
- [Nak87] N. Nakayama, *On Weierstrass models*, Algebraic geometry and commutative algebra in honor of Masayoshi Nagata. 405-431 (1991).
- [Miy91] J. Miyachi, *Localization of triangulated categories and derived categories*, J. Algebra. 141(2), 463-483 (1991).
- [Mor] H. Morimura, *Categorical generic fiber*, arXiv:2111.00239v2
- [MU] H. Morimura and T. Ulsnaes *Twisted categorical generic fiber*, in preparation
- [Orl97] D. Orlov, *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. 84, 1361-1381 (1997).
- [SP] The Stacks Project authors, *Stacks project*, <https://stacks.math.columbia.edu>.
- [Töe] B. Töen, *Lectures on DG-categories*, <http://www.math.univ-toulouse.fr/~toen/>