

FUSIONS OF TENSOR POWERS OF JOHNSON SCHEMES

SEAN EBERHARD AND MISHA MUZYCHUK

ABSTRACT. This paper is a follow-up to [E], in which the first author studied primitive association schemes lying between a tensor power \mathcal{T}_m^d of the trivial association scheme and the Hamming scheme $\mathcal{H}(m, d)$. A question which arose naturally in that study was whether all primitive fusions of \mathcal{T}_m^d lie between $\mathcal{T}_{m^e}^{d/e}$ and $\mathcal{H}(m^e, d/e)$ for some $e \mid d$. This note answers this question positively provided that m is large enough. We similarly classify primitive fusions of the d th tensor power of a Johnson scheme on $\binom{m}{k}$ points provided m is large enough in terms of k and d .

1. INTRODUCTION

Association schemes are objects of central importance in algebraic combinatorics. The reader needing an introduction to association schemes could refer to any of [CP, BI, Z, B] or the introduction to [E].

All our association schemes are symmetric. If $\mathfrak{X}, \mathfrak{Y}$ are association schemes on a common vertex set then we write $\mathfrak{X} \leq \mathfrak{Y}$ if \mathfrak{X} refines \mathfrak{Y} as a partition. A fusion is a coarsening which is again an association scheme. We denote the Hamming scheme of order m^d and rank $d + 1$ by $\mathcal{H}(m, d)$ and the Johnson scheme of order $\binom{m}{k}$ and rank $k + 1$ by $\mathcal{J}(m, k)$. The special case $\mathcal{H}(m, 1) \cong \mathcal{J}(m, 1)$ is the *trivial scheme*, denoted \mathcal{T}_m . The d th tensor power of an association scheme \mathfrak{X} is denoted \mathfrak{X}^d . The symmetrized d th tensor power of the Johnson scheme $\mathcal{J}(m, k)$ is called the *Cameron scheme* and denoted $\mathcal{C}(m, k, d)$.

In this paper we classify primitive fusions of $\mathcal{J}(m, k)^d$ assuming m is sufficiently large in terms of k and d .

Theorem 1. *Let \mathfrak{X} be a primitive fusion of $\mathcal{J}(m, k)^d$, where $m \geq m_0(k, d)$. Let $M = \binom{m}{k}$. Then, up to permuting the coordinates, either*

- (1) $\mathcal{J}(m, k)^d \leq \mathfrak{X} \leq \mathcal{C}(m, k, d)$, or
- (2) $\mathcal{T}_{M^e}^{d/e} \leq \mathfrak{X} \leq \mathcal{H}(M^e, d/e)$ for some integer $e \mid d$.

The special cases $k = 1$ and $d = 1$ are worth highlighting individually. In [M2] it was shown that $\mathcal{H}(m, d)$ has no nontrivial fusions for $m > 4$. The case $k = 1$ of Theorem 1 more generally classifies primitive fusions of \mathcal{T}_m^d (for m sufficiently large).

Corollary 2. *Let \mathfrak{X} be a primitive fusion of \mathcal{T}_m^d , where $m \geq m_0(d)$. Then, up to permuting coordinates, $\mathcal{T}_{m^e}^{d/e} \leq \mathfrak{X} \leq \mathcal{H}(m^e, d/e)$ for some integer $e \mid d$.*

In [M3] it was shown that $\mathcal{J}(m, k)$ has no nontrivial fusions for $m \geq 3k + 4$. The case $k = 1$ of Theorem 1 recovers this result, except for the precise lower bound.

Corollary 3. *Let \mathfrak{X} be a fusion of $\mathcal{J}(m, k)$, where $m \geq m_0(k)$. Then either $\mathfrak{X} = \mathcal{J}(m, k)$ or \mathfrak{X} is trivial.*

SE has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 803711).

Association schemes of the type appearing in the conclusion of Theorem 1 are studied in [E], where they are called ‘‘Cameron sandwiches’’ and ‘‘Hamming sandwiches’’, respectively. The main result of [E] is that there are infinite families of nonschurian Hamming sandwiches.

Remark 4. Imprimitive fusions of $\mathcal{J}(m, k)^d$ are not so easily classified. Certainly one must allow arbitrary tensor products of the cases appearing in Theorem 1, but there are still many others. For example, the imprimitive wreath product $\mathcal{T}_m \wr \mathcal{T}_m$ (see [CP, Section 3.4.1]) is an imprimitive fusion of \mathcal{T}_m^2 not fitting this description.

As an application we give an elementary classification of primitive groups containing $(A_m^{(k)})^d$. Here $A_m^{(k)}$ denotes the image of the alternating group A_m in its permutation action on k -sets, and below $S_m^{(k)}$ is defined similarly.

Corollary 5. *Let $G \leq S_n$ be a primitive permutation group containing $(A_m^{(k)})^d$, where $n = \binom{m}{k}^d$ and $m \geq m_0(k, d)$. Let $M = \binom{m}{k}$. Then either*

- (1) $(A_m^{(k)})^d \leq G \leq S_m^{(k)} \wr S_d$ or
- (2) $(A_{M^e})^d \leq G \leq S_{M^e} \wr S_{d/e}$ for some integer $e \mid d$.

This is a special case of Cameron’s theorem [C, L, M], which more generally classifies all large primitive permutation groups. However, while the proof of Cameron’s theorem depends on the classification of finite simple groups, an elementary proof of Corollary 5 is obtained by combining Theorem 1 with [W, Theorem 8.2.1].

2. NOTATION

Let $[0, k]^d = \{0, \dots, k\}^d$. We use the following notation for vectors $a, b, c \in [0, k]^d$:

$$\begin{aligned} a \leq b &\iff a_i \leq b_i \text{ for all } i, \\ |a - b| &= (|a_1 - b_1|, \dots, |a_d - b_d|), \\ \min(a, b) &= (\min(a_1, b_1), \dots, \min(a_d, b_d)), \\ \max(a, b) &= (\max(a_1, b_1), \dots, \max(a_d, b_d)), \\ a! &= a_1! \cdots a_d!, \\ \binom{a}{b} &= \binom{a_1}{b_1} \cdots \binom{a_d}{b_d}, \\ \text{wt}(a) &= a_1 + \cdots + a_d, \\ \text{supp}(a) &= \{i : a_i > 0\}, \\ [a] &= \{b \in [0, k]^d : b \leq a\}, \\ (x)^d &= (x, \dots, x), \\ e_i &= (0, \dots, 0, 1, 0, \dots, 0). \end{aligned}$$

We understand $\binom{a}{b}$ to be zero unless $(0)^d \leq b \leq a$. We call $\text{wt}(a)$ the *weight* of a and $\text{supp}(a)$ the *support* of a .

If p is a polynomial in one variable we write $\deg(p)$ for its degree and $\lambda(p)$ for its leading term.

3. PROOF

The structure constants of $\mathcal{J}(m, k)$ are given by

$$p_{b,c}^a(m) = \sum_i \binom{k-a}{i} \binom{a}{k-b-i} \binom{a}{k-c-i} \binom{m-k-a}{b+c+i-k}$$

(as in [E, KK]). Here $0 \leq a, b, c \leq k$, and i can be restricted to the range

$$\max(0, k-a-b, k-b-c, k-a-c) \leq i \leq \min(k-a, k-b, k-c, m-a-b-c).$$

In the following we always assume $m \geq 3k$. Under this assumption we have $p_{b,c}^a(m) > 0$ if and only if a, b, c satisfy the triangle inequalities [E, Lemma 4.1]. More precisely we have the following.

Lemma 6. *We have $p_{b,c}^a(m) > 0$ if and only if $|b-c| \leq a \leq b+c$. Assuming $0 \leq b-c \leq a \leq b+c$, the leading term of $p_{b,c}^a(m)$ is*

$$\lambda(p_{b,c}^a(m)) = \begin{cases} \binom{a}{b} \binom{a}{c} \frac{1}{(b+c-a)!} m^{b+c-a} & : a \geq b, \\ \binom{k-a}{k-b} \binom{a}{b-c} \frac{1}{c!} m^c & : a \leq b. \end{cases}$$

In particular, $\deg(p_{b,c}^a(m)) \leq \min(b, c)$, and equality holds if and only if $a \leq \max(b, c)$.

Now consider $\mathcal{J}(m, k)^d$. For $a, b, c \in [0, k]^d$, let

$$p_{b,c}^a(m) = \prod_{i=1}^d p_{b_i, c_i}^{a_i}(m).$$

These are the structure constants of $\mathcal{J}(m, k)^d$. The following lemma generalizes the previous one.

Lemma 7. *Let $a, b, c \in [0, k]^d$. Then $p_{b,c}^a(m) > 0$ if and only if $|b-c| \leq a \leq b+c$. Moreover $\deg(p_{b,c}^a(m)) \leq \text{wt}(\min(b, c))$, with equality if and only if $a \leq \max(b, c)$. If $\deg(p_{b,c}^a(m)) = \text{wt}(b) = \text{wt}(c)$ then $a \leq b = c$ and the leading term of $p_{b,c}^a(m)$ is*

$$\lambda(p_{b,c}^a(m)) = \binom{\binom{k}{b}^d - a}{\binom{k}{b}^d - b} \frac{1}{b!} m^{\text{wt}(b)}.$$

Proof. It follows from $p_{b,c}^a(m) = \prod_{i=1}^d p_{b_i, c_i}^{a_i}(m)$ that $p_{b,c}^a(m) > 0$ iff each triple a_i, b_i, c_i satisfies the triangle condition $|b_i - c_i| \leq a_i \leq b_i + c_i$. In this case

$$\deg(p_{b,c}^a(m)) = \sum_{i=1}^d \deg(p_{b_i, c_i}^{a_i}(m)) \leq \sum_{i=1}^d \min(b_i, c_i) = \text{wt}(\min(b, c)).$$

Equality holds if and only if $a_i \leq \max(b_i, c_i)$ for all i .

If $\deg(p_{b,c}^a(m)) = \text{wt}(b) = \text{wt}(c)$ then $\text{wt}(\min(b, c)) = \text{wt}(b) = \text{wt}(c)$, which implies $b = c$, and moreover we have seen that we must have $a \leq b$. Multiplying the leading terms of $p_{b_i, c_i}^{a_i}(m)$ given by the previous lemma, we get the claimed formula. \square

Let \mathfrak{X} be a fusion of $\mathcal{J}(m, k)^d$. Since \mathfrak{X} is a coarsening of $\mathcal{J}(m, k)^d$ there is a partition \mathfrak{S} of $[0, k]^d$ such that $\mathfrak{X} = \{R_\alpha : \alpha \in \mathfrak{S}\}$, where

$$(u, v) \in R_\alpha \iff (|u_1 \setminus v_1|, \dots, |u_d \setminus v_d|) \in \alpha.$$

In this situation we write $\mathfrak{X} = \mathcal{J}(m, k)^\mathfrak{S}$ [E, Section 4]. For $\beta, \gamma \in \mathfrak{S}$ and $a \in [0, k]^d$ define

$$p_{\beta, \gamma}^\alpha(m) = \sum_{b \in \beta, c \in \gamma} p_{b, c}^\alpha(m).$$

For $\mathfrak{X} = \mathcal{J}(m, k)^\mathfrak{S}$ to be an association scheme, \mathfrak{S} must satisfy two conditions:

- (1) $\{(0)^d\} \in \mathfrak{S}$,
- (2) $p_{\beta, \gamma}^\alpha(m) = p_{\beta, \gamma}^{\alpha'}(m)$ for all $\alpha, \beta, \gamma \in \mathfrak{S}$ and $a, a' \in \alpha$.

Assuming $m \geq m_0(k, d)$ (as we do from now on), the latter condition holds only if $p_{\beta, \gamma}^\alpha(m)$ and $p_{\beta, \gamma}^{\alpha'}(m)$ are equal as polynomials. We may denote the common value of $p_{\beta, \gamma}^\alpha(m)$ by $p_{\beta, \gamma}^\alpha$; these are the structure constants of \mathfrak{X} .

We call the sets $\alpha \in \mathfrak{S}$ the *basic \mathfrak{S} -sets*; their unions are called *\mathfrak{S} -sets*. For nonempty $S \subseteq [0, k]^d$ let $\text{wt}(S) = \max\{\text{wt}(b) \mid b \in S\}$. For $\alpha \in \mathfrak{S}$ let α^* be the set of $a \in \alpha$ of maximal weight. Let $D_\alpha = \bigcup_{a \in \alpha^*} [a]$. Note that $\alpha^* \subseteq D_\alpha$.

Proposition 8. *Let $\beta \in \mathfrak{S}$ be a basic \mathfrak{S} -set.*

- (1) D_β is an \mathfrak{S} -set, and $\text{wt}(D_\beta \setminus \beta) < \text{wt}(\beta)$.
- (2) For every basic set $\alpha \in \mathfrak{S}$ there is constant N_α^β such that

$$\sum_{b: a \leq b \in \beta^*} \binom{\binom{k}{d} - a}{\binom{k}{d} - b} = N_\alpha^\beta \quad (a \in \alpha).$$

- (3) Every element of β is dominated by a unique element of β^* .
- (4) Either $\text{wt}(\beta) = \text{wt}(D_\beta \setminus \beta) + 1$ or $\beta^* \subseteq \{0, k\}^d$.

Proof. Let $w = \text{wt}(\beta)$ and $D = D_\beta$. Since the polynomials $p_{b, c}^\alpha(m)$ have positive leading coefficient whenever they are nonzero, the degree of $p_{\beta, \beta}^\alpha(m)$ is, by the previous lemma,

$$\deg(p_{\beta, \beta}^\alpha(m)) = \max_{b, c \in \beta} \deg(p_{b, c}^\alpha(m)) \leq \max_{b, c \in \beta} \text{wt}(\min(b, c)) \leq w,$$

with equality holding if and only if $a \leq b = c \in \beta^*$, i.e., if and only if $a \in D$. Since $p_{\beta, \beta}^\alpha(m)$ should depend only on the cell of \mathfrak{S} containing a , it follows that D is an \mathfrak{S} -set. Since β is basic, $\beta \subseteq D$, and since $\beta \supseteq \beta^*$ we have $\text{wt}(D \setminus \beta) < \text{wt}(\beta)$.

Moreover if $a \in D$ then the leading term of $p_{\beta, \beta}^\alpha$ is

$$\lambda(p_{\beta, \beta}^\alpha(m)) = \sum_{b: a \leq b \in \beta^*} \binom{\binom{k}{d} - a}{\binom{k}{d} - b} \frac{1}{b!} m^w,$$

and again this should depend only on the cell α containing a . Taking $a = b \in \beta^*$ and $\alpha = \beta$, it follows that $b!$ is a constant for $b \in \beta^*$. Hence (2) holds (if α is not contained in D then $N_\alpha^\beta = 0$).

Next apply (2) with $\alpha = \beta$. Taking $a \in \beta^*$ shows $N_\beta^\beta = 1$, so (3) holds.

Finally let $b \in \beta^*$ and suppose $a = b - e_i \geq 0$. If $a \in \beta$ then we get

$$\binom{k - b_i + 1}{k - b_i} = N_\beta^\beta = 1,$$

so $b_i = k$. Hence either $\text{wt}(b) = \text{wt}(\beta) + 1$ or $b_i \in \{0, k\}$ for all i , which implies (4). \square

We can define a partial ordering on \mathfrak{S} by saying $\alpha \preceq \beta$ if every $a \in \alpha$ is dominated by some $b \in \beta$. By part (3) of the proposition, this is equivalent to $N_\alpha^\beta > 0$. It is obvious that \preceq is reflexive and transitive on \mathfrak{S} . To verify antisymmetry, note that if $\alpha \preceq \beta \preceq \alpha$ and $a \in \alpha^*$

then there are $b \in \beta$ and $a' \in \alpha$ such that $a \leq b \leq a'$, which by maximality of a implies $a = b = a'$ and hence $\alpha = \beta$ since \mathfrak{S} is a partition. We say $\alpha \in \mathfrak{S}$ is *minimal* if it is minimal in $(\mathfrak{S} \setminus \{(0)^d\}, \preceq)$.

Corollary 9. *Let $\alpha \in \mathfrak{S}$ be a minimal basic set. Then $D_\alpha = \alpha \cup \{(0)^d\}$. Moreover the elements of α^* have disjoint equal-sized supports, and either $\text{wt}(\alpha) = 1$ or $\alpha^* \subseteq \{0, k\}^d$.*

Proof. Apply the proposition. Note that if β is a basic subset of $D_\alpha \setminus \alpha$ then $\beta \prec \alpha$. By minimality of α this implies $\beta = \{(0)^d\}$. Hence $D_\alpha = \alpha \cup \{(0)^d\}$. Next, for any $a \in \alpha^*$ and $i \in \text{supp}(a)$ we have $e_i \in D_\alpha$. By part (3) of Proposition 8, a is the unique element of α^* dominating e_i . Therefore the elements of α^* have disjoint supports. By part (4), either $\text{wt}(\alpha) = 1$ or $\alpha^* \subseteq \{0, k\}^d$. If $\text{wt}(\alpha) = 1$ then all $a \in \alpha^*$ have singleton support, and otherwise $|\text{supp}(a)| = \text{wt}(a)/k = \text{wt}(\alpha)/k$ for all $a \in \alpha^*$. \square

Until now \mathfrak{X} could be imprimitive. Now we specialize to the primitive case to complete the proof of Theorem 1. Let $\alpha \in \mathfrak{S}$ be minimal. If \mathfrak{X} is primitive then R_α must be connected, which implies that α^* covers $\{1, \dots, d\}$. Hence α^* is an equipartition of $\{1, \dots, d\}$. Let $w = \text{wt}(\alpha)$. If $w = 1$ then α^* must be the set of elements of weight 1, so R_α is the Cameron graph. Since the Weisfeiler–Leman stabilization of the Cameron graph is the Cameron scheme, we find $\mathfrak{X} \leq \mathcal{C}(m, k, d)$. If $w > 1$ then $\alpha^* \subseteq \{0, k\}^d$. If the elements of α^* have support size e then R_α is the Hamming graph $H(M^e, d/e)$, so $\mathfrak{X} \leq \mathcal{H}(M^e, d/e)$. To finish we must show $\mathcal{T}_{M^e}^{d/e} \leq \mathfrak{X}$. For this it suffices to prove that for every $\beta \in \mathfrak{S}$, the elements of β^* are sums of elements of α^* (and in particular $\beta^* \subseteq \{0, k\}^d$).

We apply Proposition 8(2) to α and β . Let $i \in \{1, \dots, d\}$. Taking $a = e_i$, we find that N_α^β is at least the number of $b \in \beta^*$ such that $b_i > 0$. On the other hand, taking a to be the unique element of α^* dominating e_i , since $a \in \{0, k\}^d$ we find that N_α^β is equal to the number of $b \in \beta^*$ such that $a \leq b$. Hence $b_i > 0$ implies $a \leq b$. This implies that b is the sum of those $a \in \alpha^*$ such that $a \leq b$, as required.

REFERENCES

- [B] R. A. Bailey, *Association schemes*, Cambridge Studies in Advanced Mathematics, vol. 84, Cambridge University Press, Cambridge, 2004. Designed experiments, algebra and combinatorics. MR2047311 [↑1](#)
- [BI] E. Bannai and T. Ito, *Algebraic combinatorics. I*, The Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA, 1984. Association schemes. MR882540 [↑1](#)
- [C] P. J. Cameron, *Finite permutation groups and finite simple groups*, Bull. London Math. Soc. **13** (1981), no. 1, 1–22. MR599634 [↑1](#)
- [CP] G. Chen and I. Ponomarenko, *Lectures on coherent configurations* (2019), available at <http://www.pdmi.ras.ru/~inp/ccNOTES.pdf>. Online lecture notes. [↑1](#), [4](#)
- [E] S. Eberhard, *Hamming sandwiches* (2022), [arXiv:2203.03687](https://arxiv.org/abs/2203.03687). [↑\(document\)](#), [1](#), [1](#), [3](#), [3](#)
- [KK] L. A. Kalužnin and M. H. Klin, *Certain maximal subgroups of symmetric and alternating groups*, Mat. Sb. (N.S.) **87(129)** (1972), 91–121. MR0297854 [↑3](#)
- [L] M. W. Liebeck, *On minimal degrees and base sizes of primitive permutation groups*, Arch. Math. (Basel) **43** (1984), no. 1, 11–15. MR758332 [↑1](#)
- [M1] A. Maróti, *On the orders of primitive groups*, J. Algebra **258** (2002), no. 2, 631–640. MR1943938 [↑1](#)
- [M2] M. E. Muzychuk, *Subschemes of Hamming association schemes $H(n, q)$, $q \geq 4$* , 1992, pp. 119–128. Interactions between algebra and combinatorics. MR1192835 [↑1](#)
- [M3] ———, *Subschemes of the Johnson scheme*, European J. Combin. **13** (1992), no. 3, 187–193. MR1164762 [↑1](#)

- [W] J. Wilmes, *Structure, automorphisms, and isomorphisms of regular combinatorial objects*, ProQuest LLC, Ann Arbor, MI, 2016. Ph.D. thesis – The University of Chicago, <https://www.proquest.com/docview/1837431537>. MR3593268 ↑1
- [Z] P.-H. Zieschang, *Theory of association schemes*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2005. MR2184345 ↑1

SEAN EBERHARD, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK

Email address: `eberhard@maths.cam.ac.uk`

MISHA MUZYCHUK, DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY P.O.B. 653, BEER SHEVA 8410501, ISRAEL

Email address: `muzychuk@bgu.ac.il`