

# HODGE-TO-SINGULAR CORRESPONDENCE FOR REDUCED CURVES

MIRKO MAURI AND LUCA MIGLIORINI

ABSTRACT. We study the summands of the decomposition theorem for the Hitchin system for  $GL_n$ , in arbitrary degree, over the locus of reduced spectral curves. A key ingredient is a new correspondence between these summands and the topology of hypertoric quiver varieties. In contrast to the case of meromorphic Higgs fields, the intersection cohomology groups of moduli spaces of regular Higgs bundles depend on the degree. We describe this dependence.

Let  $C$  be a compact Riemann surface of genus  $g$  with canonical bundle  $\omega_C$ . Let  $d$  and  $n$  be integers with  $n > 1$ .

The *Dolbeault moduli space*  $M(n, d)$  is the coarse moduli space which parametrises S-equivalence classes of semistable Higgs bundles on  $C$  of rank  $n$  and degree  $d$ , i.e. polystable pairs  $(\mathcal{E}, \phi)$  consisting of a vector bundle  $\mathcal{E}$  of rank  $n$  and degree  $d$  and a section  $\phi \in H^0(C, \mathcal{E}nd(\mathcal{E}) \otimes \omega_C)$ , called *Higgs field*; see [67]. The Dolbeault moduli space is equipped with a projective fibration called *Hitchin fibration*

$$(1) \quad \chi(n, d): M(n, d) \rightarrow A_n := \bigoplus_{i=1}^n H^0(C, \omega_C^{\otimes i}),$$

$$(\mathcal{E}, \phi) \mapsto \text{char}(\phi) = (\text{tr}(\phi), \text{tr}(\Lambda^2 \phi), \dots, \text{tr}(\Lambda^i \phi), \dots, \det(\phi)),$$

which assigns to  $(\mathcal{E}, \phi)$  the characteristic polynomial  $\text{char}(\phi)$  of the Higgs field  $\phi$ .

The cohomology of  $M(n, d)$  has been extensively studied in the literature, especially under the assumption that  $n$  and  $d$  are coprime, namely when  $M(n, d)$  is smooth; see for instance [40, 33, 36, 35, 38, 54, 31, 16, 54, 19]. Recently studies about the intersection cohomology  $IH^*(M(n, d), \mathbb{Q})$  of singular Dolbeault moduli spaces have started to emerge; see for instance [26, 27, 48, 50, 51, 49, 14, 13, 43, 64].<sup>1</sup> From this viewpoint, the decomposition theorem for the Hitchin map is a key tool to investigate the intersection cohomology of  $M(n, d)$ : it allows to decompose  $IH^*(M(n, d), \mathbb{Q})$  into building blocks, which are cohomology of some perverse sheaves on  $A_n$ .

In this article, we establish a correspondence which we find surprising between the summands of the decomposition theorem for the Hitchin map of  $M(n, d)$ , and the singularities of  $M(n, 0)$ ; see Theorem 1.1. We call it the *Hodge-to-singular correspondence*. This correspondence offers a new perspective on the cohomology of Dolbeault moduli spaces, even in the smooth case. What is surprising is that this relation is not realised via a direct geometric correspondence between  $M(n, d)$

---

<sup>1</sup>The lists of works are not intended to be exhaustive at all. We limit to list a series of selected papers that inspired the current article.

and  $M(n, 0)$ , like a deformation or an alteration, but, as we explain below, by comparing the Hodge theory of  $M(n, d)$  and the singularity theory of  $M(n, 0)$  on the common base of the Hitchin fibration.

The Hodge-to-singular correspondence gives a formula to express the dependence of the intersection cohomology of  $M(n, d)$  on the degree  $d$ . This should be compared with the degree-independence of spaces of meromorphic Higgs bundle [48, Thm 0.1, 0.4]. It also extends the main theorem of [17] to arbitrary degree, and provide a conceptual and elegant explanation for the occurrence of the cographic matroid in [17, §6].

Our correspondence suggests a unifying approach to the study of smooth and singular Dolbeault moduli spaces that may be useful to tackle the topological mirror symmetry conjecture [35, 51] and the P=W conjecture [16, 18] simultaneously in both settings. It also provides new evidence for Davison's conjectures in [13, Conj. 5.6, Rem. 5.10].

In a different context, similar ideas have already been successfully employed to compute the Hodge numbers of the exceptional O'Grady 10 example of compact hyperkähler manifolds, out of the Hodge numbers of the Hilbert scheme of five points on a K3 surface; see [22]. Indeed, both spaces can be interpreted as special moduli spaces of sheaves on K3 surfaces corresponding to different degrees  $d$  (or rather Euler characteristics), and the computation of their Betti numbers can be reduced to determining how their cohomology depends on  $d$ . In fact, the Hodge-to-singular correspondence, formulated here for Dolbeault moduli spaces, can be stated also for Mukai systems on moduli space of sheaves on K3 or abelian surfaces; see Section 10.3.

## 1. MAIN RESULTS

**1.1. Hodge-to-singular correspondence.** In order to state our main results, we need to recall some features of the singularities of  $M(n, d)$ . A singular point of  $M(n, d)$  corresponds to a strictly polystable Higgs bundle

$$(E_1, \phi_1)^{\oplus m_1} \oplus \dots \oplus (E_r, \phi_r)^{\oplus m_r},$$

where  $(E_i, \phi_i)$  are distinct stable Higgs bundles of slope  $\deg E_i / \text{rank } E_i = d/n$ . In particular, the  $r$ -uples of positive integers  $\underline{m} = \{m_i\}$  and  $\underline{n} = \{n_i\} := \{\deg E_i\}$  satisfy the following relations

$$(2) \quad n = \sum_{i=1}^r m_i n_i \quad n_i \cdot \frac{d}{n} = \deg E_i \in \mathbb{Z}.$$

For  $\underline{m}, \underline{n} \in \mathbb{Z}_{>0}^r$  satisfying (2), define  $M_{\underline{m}, \underline{n}}^\circ(d)$  the locus of polystable Higgs bundles in  $M(n, d)$  of multirank  $\underline{n}$  and multiplicity  $\underline{m}$ , and call  $M_{\underline{m}, \underline{n}}(d)$  its closure in  $M(n, d)$ . The loci  $M_{\underline{m}, \underline{n}}^\circ(d)$  are the strata of a complex Whitney stratification of  $M(n, d)$

$$(3) \quad M(n, d) = \bigsqcup_{\underline{m}, \underline{n}} M_{\underline{m}, \underline{n}}^\circ(d),$$

and an analytic normal slice  $W_{\underline{m}, \underline{n}}$  through  $M_{\underline{m}, \underline{n}}^\circ(d)$  is isomorphic to a Nakajima quiver variety; see [50, Proof of Prop. 2.10]. The Whitney stratification (3) induces a filtration of the base  $A_n$  of the

Hitchin fibration by closed subsets  $S_{\underline{m}, \underline{n}} := \chi(n, d)(M_{\underline{m}, \underline{n}}(0))$ , and a stratification

$$(4) \quad A_n = \bigsqcup S_{\underline{m}, \underline{n}}^\circ \quad \text{with } S_{\underline{m}, \underline{n}}^\circ := S_{\underline{m}, \underline{n}} \setminus \bigcup_{\dim S_{\underline{m}', \underline{n}'} < \dim S_{\underline{m}, \underline{n}}} S_{\underline{m}', \underline{n}'}$$

Let  $A_n^{\text{red}} := \bigsqcup_{\underline{m}=(1, \dots, 1)} S_{\underline{m}, \underline{n}}^\circ \subset A_n$  be the open subset of reduced characteristic polynomials. Define

$$M(n, d)^{\text{red}} := \chi(n, d)^{-1}(A_n^{\text{red}}) \quad \text{and} \quad M_{\underline{m}, \underline{n}}(d)^{\text{red}} := M_{\underline{m}, \underline{n}}(d) \cap M(n, d)^{\text{red}}.$$

When  $\underline{m} = \{1, \dots, 1\}$ , we abbreviate the notation by omitting the subscript  $\underline{m}$ , e.g.  $S_{\underline{n}} := S_{\underline{m}, \underline{n}}$ . Remarkably,  $W_{\underline{n}}$  admits diffeomorphic symplectic resolutions  $\widetilde{W}_{\underline{n}}$ , whose top cohomology  $H^{\dim W_{\underline{n}}}(\widetilde{W}_{\underline{n}}, \mathbb{Q})$  is a representation of the fundamental group of  $S_{\underline{n}}^\circ$ ; see Proposition 4.11 and Theorem 7.4.<sup>2</sup> Denote by  $\mathcal{L}_{\underline{n}}$  the corresponding local system on  $S_{\underline{n}}^\circ$  (which carries a pure Hodge structure of weight zero).

The Hodge-to-singular correspondence is a splitting of the mixed Hodge module  $R\chi(n, e)_* \mathbb{Q}_{M(n, e)}$  whose direct summands are controlled by the topology of the singularities  $W_{\underline{n}}$ .

For an integer  $d$  let  $\mathcal{P}_d$  be the set of partitions  $\underline{n} = \{n_i\}$  of  $n$  such that  $n_i d / n \in \mathbb{Z}$  for every  $i$ . Note that this set is in bijection with the partitions of  $q = \gcd(n, d)$ . Given a partition  $\underline{q} = \{a_i\}$  of  $q$ , then  $\underline{q} = \{a_i n / q\} \in \mathcal{P}_d$ .

**Theorem 1.1** (Hodge-to-singular correspondence). *Let  $\langle \bullet \rangle = [-2\bullet](-\bullet)$ . Let  $d$  and  $e$  be integers, with  $e$  coprime to  $n$ . There is an isomorphism in  $D^b \text{MHM}_{\text{alg}}(A_n^{\text{red}})$  or in  $D^b(A_n^{\text{red}})$  (ignoring the Tate shifts):*

$$(5) \quad R\chi(n, e)_* \mathbb{Q}_{M(n, e)} \simeq \bigoplus_{\underline{n} \in \mathcal{P}_d} R\chi(n, d)_* \text{IC}(M_{\underline{n}}(d), \chi(n, d)^* \mathcal{L}_{\underline{n}}) \langle \text{codim } S_{\underline{n}} \rangle.$$

Taking cohomology, we obtain a morphism of mixed Hodge structures

$$(6) \quad H^k(M(n, e)^{\text{red}}, \mathbb{Q}) \simeq \bigoplus_{\underline{n} \in \mathcal{P}_d} IH^{k-2 \text{codim } S_{\underline{n}}}(M_{\underline{n}}(d)^{\text{red}}, \chi(n, d)^* \mathcal{L}_{\underline{n}}) \langle -\text{codim } S_{\underline{n}} \rangle.$$

which respects the perverse filtrations<sup>3</sup> carried by these vector spaces.

Note that the summand of the RHS of (5) corresponding to the trivial partition  $\underline{n} = \{n\}$  is  $R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q})$ . This means that (6) measures the difference between  $H^k(M(n, e)^{\text{red}}, \mathbb{Q})$  and  $IH^k(M(n, d)^{\text{red}}, \mathbb{Q})$ .

Singular Dolbeault moduli space for  $d = 0$  were introduced to study the topology of character varieties, to which they are homomorphic via the non-abelian Hodge correspondence. Historically, to avoid to deal with singular moduli spaces, the moduli problem has been slightly modified or twisted (cf. the "bait-and-switch" in [36, §1]), and the Dolbeault moduli spaces with  $d$  coprime to  $n$  have received more attention in the literature. The Hodge-to-singular correspondence suggests that the historical order, say from smooth to singular spaces, should be reversed at least conceptually:

<sup>2</sup>In the following, we identify  $\widetilde{W}_{\underline{n}}$  with the hypertoric quiver variety  $Y(\Gamma_{\underline{n}}^\pm, \theta)$ . For convenience, we do not introduce hypertoric quiver varieties here, and we postpone their definition to Section 4.

<sup>3</sup>We recall the definition of perverse filtration in (49).

the intersection cohomology of singular Dolbeault moduli spaces should be considered as "primitive" objects into which the cohomology of smooth Dolbeault moduli spaces decomposes.

In [48, Thm 0.1], Maulik and Shen proved an analogue of Theorem 1.1 for twisted Higgs bundles, i.e. pairs  $(\mathcal{E}, \phi)$  where now  $\phi$  is a global section of  $\mathcal{E}nd(\mathcal{E}) \otimes \mathcal{O}_C(D)$  for some effective divisor  $D$  with  $\deg D > 2g - 2$ , the so-called "Fano" condition. Unfortunately, the estimates in [48, §4] are not useful in our setting, namely  $\mathcal{O}_C(D) \simeq \omega_C$  or "Calabi–Yau" condition; see [48, §0.4], [10, §11] and Remark 9.6. This accounts for the existence of proper supports for the complex  $R\chi(n, e)_* \mathrm{IC}(M(n, e), \mathbb{Q})$  in  $A_n$ , which on the contrary do not appear under the "Fano" assumption by [48, Thm. 0.4].

We can suggestively summarise the Hodge-to-singular correspondence in the following table.

HODGE THEORY VS SINGULARITIES OF  $M(n, d)$

$\gcd(n, d) = 1$	$\gcd(n, d) \neq 1, d \neq 0$	$d = 0$
no singularities, more summands of dec. thm	intermediate behaviour	more singular strata, no proper summands of dec. thm

**1.2. Dependence of  $IH^*(M(n, d), \mathbb{Q})$  on degree.** The Hodge-to-singular correspondence asserts that the intersection cohomology  $IH^k(M(n, d)^{\mathrm{red}}, \mathbb{Q})$  for arbitrary  $d \in \mathbb{Z}$  decomposes into the direct sum of isotypic components of tensor products of  $IH^k(M(n', 0)^{\mathrm{red}}, \mathbb{Q})$  for  $n' \leq n$  under the action of a symmetric group; see Remark 10.1.<sup>4</sup> We can refine the result by showing that  $IH^k(M(n', 0)^{\mathrm{red}}, \mathbb{Q})$  is determined only by the smooth locus of the Hitchin fibration. Remarkably, this is in contrast to the smooth case, namely  $d$  coprime to  $n$ , where  $R\chi(n, d)_* \mathbb{Q}_{M(n, d)}$  admits proper supports which cannot be detected on the smooth locus of Hitchin fibration.

To this end, recall that any  $a \in A_n$  can be viewed as a monic polynomial of degree  $n$  with coefficient of the degree  $n - i$  term in  $H^0(C, \omega_C^{\otimes i})$ , or equivalently as the curve  $C_a \subset \mathrm{Tot}(\omega_C)$  of degree  $n$  over  $C$  cut by this monic equation, called *spectral curve*. So an alternative definition of  $S_{\underline{n}}^{\circ}$  is the locus parametrising spectral curves  $C_a = \bigcup_{i=1}^r C_i$  such that the reduced irreducible component  $C_i$  has degree  $n_i$  over  $C$ . We define also the following spaces:

- (i)  $S_{\underline{n}}^{\times} \subseteq S_{\underline{n}}^{\circ}$  is the locus parametrising reducible nodal curves having smooth irreducible components of degree  $n_i$  over  $C$ ;
- (ii)  $\pi_{\underline{n}}: C_{\underline{n}}^{\times} \rightarrow S_{\underline{n}}^{\times}$  is the universal spectral curve over  $S_{\underline{n}}^{\times}$ ;

---

<sup>4</sup>An extension of the Hodge-to-singular correspondence to the whole  $A_n$  would provide a recursive formula to compute the intersection cohomology of  $M(n, d)$  for arbitrary genus  $g$  and degree  $d$ . See also Davison's conjecture [13, Conj. 5.6, Rem. 5.10].

- (iii)  $\text{Pic}^0(\mathcal{C}_{\underline{n}}^\times) \rightarrow S_{\underline{n}}^\times$  is the (relative) Jacobian of the universal spectral curve over  $S_{\underline{n}}^\times$ ;
- (iv)  $g_{\underline{n}} : A_{\underline{n}}^\times \rightarrow S_{\underline{n}}^\times$  is the maximal abelian proper quotient of  $\text{Pic}^0(\mathcal{C}_{\underline{n}}^\times)$ , or equivalently the Jacobian of the normalization of  $\mathcal{C}_{\underline{n}}^\times$ ;
- (v)  $\Lambda_{\underline{n}}^l := R^l(g_{\underline{n}})_* \mathbb{Q}_{A_{\underline{n}}^\times} = \Lambda^l R^1(g_{\underline{n}})_* \mathbb{Q}_{A_{\underline{n}}^\times} = \Lambda^l R^1(\pi_{\underline{n}})_* \mathbb{Q}_{A_{\underline{n}}^\times}$ ; see [16, Lemma 1.3.5].

**Definition 1.2.** *The Ngô strings<sup>5</sup> for the partition  $\underline{n}$  of  $n$  is the mixed Hodge module supported on  $S_{\underline{n}}$  given by*

$$(7) \quad \mathcal{S}_{\underline{n}} := \bigoplus_{l=0}^{2 \dim S_{\underline{n}}} \text{IC}(S_{\underline{n}}, \Lambda_{\underline{n}}^l \otimes \mathcal{L}_{\underline{n}})[-l](\text{codim } S_{\underline{n}}).$$

Note that  $\mathcal{S}_{\underline{n}}$  is independent of  $d$ , since  $\mathcal{L}_{\underline{n}}$  and  $\Lambda^l R^1(g_{\underline{n}})_* \mathbb{Q}_{A_{\underline{n}}^\times}$  are so.

In the coprime case  $\gcd(e, n) = 1$ , the summands of the decomposition theorem of the Hitchin fibration on the locus of reduced spectral curves have been described in [17]. As a first application of the Hodge-to-singular correspondence, we recover [17, Prop. 4.1, Thm 6.12].

**Theorem 1.3** (Ngô strings in degree  $e$  coprime to  $n$ ). *Let  $e$  be an integer coprime to  $n$ . There is an isomorphism in  $D^b \text{MHM}_{\text{alg}}(A_n^{\text{red}})$  or in  $D^b(A_n^{\text{red}})$  (ignoring the Tate shifts):*

$$(8) \quad R\chi(n, e)_* \mathbb{Q}_{M(n, e)}|_{A_n^{\text{red}}} \simeq \bigoplus_{\underline{n} \vdash n} \mathcal{S}_{\underline{n}}|_{A_n^{\text{red}}}.$$

Note that the proof of Theorem 1.3 below is not independent of [17]. What is new is the conceptual interpretation of that result: the rather mysterious occurrence of the cohomology of the cographic matroid in [17, Thm 6.12] can be now interpreted in geometric terms as the local intersection cohomology of the singularities of  $M(n, 0)$ , or the top cohomology of their resolutions, in symbols  $H^{\dim W_{\underline{n}}}(\widetilde{W}_{\underline{n}}, \mathbb{Q})$ . In the upcoming paper [52] we provide a combinatorial and alternative proof of Theorems 1.3 and 1.4, which is independent of [17]. In view of [17], where the locus of reduced spectral curves carries a plethora of proper summands of the decomposition theorem, it could appear bizarre that instead there is none if  $d = 0$ .

**Theorem 1.4** (Ngô strings in degree 0). *The complex  $R\chi(n, 0)_* \text{IC}(M(n, 0), \mathbb{Q})$  has full support on the reduced locus  $A_n^{\text{red}}$ , i.e.*

$$R\chi(n, 0)_* \text{IC}(M(n, 0), \mathbb{Q})|_{A_n^{\text{red}}} \simeq \mathcal{S}_{\{n\}}|_{A_n^{\text{red}}}$$

---

<sup>5</sup>The definition of Ngô strings differs from [22, Def. A.0.4] by a shift and a Tate twist. This convention avoids to write the shifts all the times.

**Question 1.5** (Full support). *Does  $R\chi(n, 0)_* \mathrm{IC}(M(n, 0), \mathbb{Q})$  have full support on the whole Hitchin base  $A_n$ , i.e.*

$$R\chi(n, 0)_* \mathrm{IC}(M(n, 0), \mathbb{Q}) \simeq \mathcal{S}_{\{n\}}?$$

As positive partial answer to Question 1.5, we observe that no summand of  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$  is supported on  $\{0\}$ , which generalises a result of Heinloth [38, Thm 1] to arbitrary degree.

**Proposition 1.6** (Vanishing of intersection form). *The intersection form on  $IH^*(M(n, d), \mathbb{Q})$  is trivial. Equivalently the forgetful map  $IH_c^*(M(n, d), \mathbb{Q}) \rightarrow IH^*(M(n, d), \mathbb{Q})$  is 0.*

In Theorem 1.7 we answer positively Question 1.5 in rank two and illustrate its far-reaching consequences. Note that the intersection cohomology groups of  $M(2, d)$  have been computed in [40, Thm 7.6] and [50, Thm 1.8, 1.9].

**Theorem 1.7** (Ngô strings in rank two). *There exist isomorphisms*

$$R\chi(2, 1)_* \mathbb{Q}_{M(2,1)} \simeq \mathcal{S}_{\{2\}} \oplus \mathcal{S}_{\{1,1\}}, \quad R\chi(2, 0)_* \mathrm{IC}(M(2, 0), \mathbb{Q}) \simeq \mathcal{S}_{\{2\}},$$

*in the bounded derived category  $D^b\mathrm{MMH}(A_n)$  of Hodge modules on  $A_n$ . As a result, there exists a surjective map*

$$H^*(M(2, 1), \mathbb{Q}) \rightarrow IH^*(M(2, 0), \mathbb{Q})$$

*strictly filtered by the perverse filtrations associated to  $\chi(2, d)$ .*

**Remark 1.8** (BPS sheaf). The Hodge-to-singular correspondence gives an explicit formula (Theorem 7.5) for the so-called BPS sheaf [43]. From this viewpoint, Theorem 1.7 should be compared with [43, Ex. 5.4] and [64]. See also [13, Rem. 5.10].

The Hodge-to-singular correspondence provides also a recursive strategy to express the dependence of  $IH^*(M(n, d), \mathbb{Q})$  on the degree  $d$ , when  $\mathrm{gcd}(n, d) \neq 0$  or  $\neq 1$ . We outline it in Section 10.1. The explicit computation of the rank of the Ngô strings for  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})|_{A_n^{\mathrm{red}}}$  will appear in [52].

As a consequence of the Hodge-to-singular correspondence, we obtain an immediate generalisation of [23, Thm. 3.11] and [43, Cor. 5.15, Ex. 5.18] to arbitrary degree on the locus of reduced spectral curves.

**Corollary 1.9.** *Let  $d, d'$  be integers with  $\mathrm{gcd}(d, n) = \mathrm{gcd}(d', n)$ . Then we have*

$$R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})|_{A_n^{\mathrm{red}}} \simeq R\chi(n, d')_* \mathrm{IC}(M(n, d'), \mathbb{Q})|_{A_n^{\mathrm{red}}}.$$

### 1.3. Some applications of the Hodge-to-singular correspondence.

1.3.1. *Stabilisation.* In [12] Coskun and Woolf have proposed a conjecture about the stabilization of the cohomology of moduli space of sheaves on compact complex surface. As a final remark, we observe that  $IH^*(M(n, d), \mathbb{Q})$  does stabilise as the rank  $n$  grows, independently of the degree  $d$ .

**Proposition 1.10** (Stabilisation). *The intersection Betti and Hodge numbers of  $M(n, d)$  stabilize to the stable intersection Betti and Hodge numbers of  $M(n, 0)$  as  $n$  tends to infinity, i.e. the limit*

$$(9) \quad \lim_{n \rightarrow \infty} \dim IH^*(M(n, d))^{p,q}$$

*is finite and independent of  $d$ .*

1.3.2. *Restriction to smooth fibres.* Let  $M$  be a projective irreducible holomorphic symplectic variety of dimension  $2r$  equipped with a Lagrangian fibration  $f: M \rightarrow B$ . In [28] Felisetti, Shen and Yin have proved that the restriction of  $H^*(M, \mathbb{Q})$  to a smooth fiber of  $f$  is isomorphic to  $H^*(\mathbb{P}^r, \mathbb{Q})$ . In general, this is false for non-compact hyperkähler varieties, but it holds true for the Hitchin fibration for  $n > 1$  by [1, Thm 5], up to a copy of  $H^\bullet(C, \mathbb{Q})$ . Here we propose an alternative proof.

**Proposition 1.11.** *The restriction of  $IH^*(M(n, d))$  to any smooth fiber  $M_a \subset M(n, d)$  of the Hitchin fibration  $\chi(n, d)$  is given by*

$$\mathrm{Im}\{IH^*(M(n, d), \mathbb{Q}) \rightarrow H^*(M_a, \mathbb{Q})\} = H^\bullet(C, \mathbb{Q}) \otimes \mathbb{Q}[\alpha|_{M_a}]/(\alpha|_{M_a}^{\dim M(n, d)+1})$$

where  $\alpha$  is a  $\chi(n, d)$ -relative ample class on  $M(n, d)$ .

Let  $\check{M}(n, d)$  be a fibre of the determinant map  $M(n, d) \rightarrow M(1, nd)$ , given by  $(\mathcal{E}, \phi) \mapsto (\det \mathcal{E}, \mathrm{tr} \phi)$ . The restriction of  $IH^*(\check{M}(n, d))$  to any smooth fiber  $\check{M}_a \subset \check{M}(n, d)$  of the Hitchin fibration  $\check{\chi}(n, d)$  is given by

$$\mathrm{Im}\{IH^k(\check{M}(n, d), \mathbb{Q}) \rightarrow H^k(\check{M}_a, \mathbb{Q})\} = \begin{cases} \langle \alpha^{k/2}|_{\check{M}_a} \rangle & \text{if } k \text{ is even,} \\ 0 & \text{if } k \text{ is odd,} \end{cases}$$

1.4. **Strategy.** In this paper we propose a geometric proof of the Hodge-to-singular correspondence. Here we outline the main steps. It is well-known that the Hitchin fibration is a family of compactified Jacobians of spectral curves; see [39, 2, 63]. In Section 5, we enlarge the Hitchin fibration  $\chi(n, d): M(n, d) \rightarrow A_n$  to a family  $\pi: \bar{J}_{d, B} \rightarrow B$  of compactified Jacobians over a versal deformation of spectral curves.<sup>6</sup>

$$\begin{array}{ccc} M(n, d) & \longrightarrow & \bar{J}_{d, B} \\ \chi(n, d) \downarrow & & \downarrow \pi \\ A_n & \longrightarrow & B, \end{array}$$

<sup>6</sup>Actually, to make the diagram work we should slice the Hitchin base, restrict to the locus of nodal spectral curves, and take into consideration the automorphisms of the curves by working over  $\overline{\mathcal{M}}_g$ . For expository reasons, we allow this abuse here, and postpone the precise details to Section 5.

Note that the relative dimension of  $\pi$  is strictly smaller than  $\dim B$ . This is a crucial ingredient to show that  $R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q})$  is independent of  $d$ , while  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$  does not; see Theorem 9.5 and Remark 9.6. In particular, we have

$$(10) \quad R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{e,B}, \mathbb{Q})|_{A_n} \simeq R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q})|_{A_n}.$$

If  $\gcd(n, e) = 1$ , the inclusion  $M(n, e) \hookrightarrow \overline{\mathcal{J}}_{e,B}$  is a regular embedding of smooth varieties, so

$$(11) \quad \mathrm{IC}(\overline{\mathcal{J}}_{e,B}, \mathbb{Q})|_{M(n,e)} \simeq \mathbb{Q}_{\overline{\mathcal{J}}_{e,B}}|_{M(n,e)} \simeq \mathbb{Q}_{M(n,e)},$$

and the LHS of (10) is isomorphic to  $R\chi(n, e)_* \mathbb{Q}_{M(n,e)}$ . If instead  $\gcd(n, d) \neq 1$ , the failure of (11) is measured by the topology of the singularities of  $M(n, d)$ . Following [8], we identify the map  $M(n, d) \hookrightarrow \overline{\mathcal{J}}_{d,B}$  with the inclusion of the hypertoric quiver variety  $Y(\Gamma_{\underline{n}}, 0)$  into the toric Lawrence variety  $X(\Gamma_{\underline{n}}^{\pm}, 0)$

$$\begin{array}{ccc} Y(\Gamma_{\underline{n}}, 0) & \xleftarrow{\iota_M} & X(\Gamma_{\underline{n}}^{\pm}, 0) \\ \chi_{\Gamma_{\underline{n}}} \downarrow & & \downarrow \pi_{\Gamma_{\underline{n}}} \\ \mathbb{C}^{b_1(\Gamma_{\underline{n}})} & \xleftarrow{\iota_A} & \mathbb{C}^s; \end{array}$$

see Theorem 7.4 for the proof, and Section 4 for details about the notation. Now a local version of the Hodge-to-singular correspondence for hypertoric quiver varieties (Theorem 4.18) shapes the RHS of (10), and gives the main result Theorem 1.1; see also Theorem 7.5. The key geometric input is the following:  $X(\Gamma_{\underline{n}}^{\pm}, 0)$  admits small resolutions of singularities that restrict to semismall resolutions of  $Y(\Gamma_{\underline{n}}, 0)$ ; see Section 4.5.

## 1.5. Outline.

- In Section 3 we prove a version of Ngô support theorem for weakly abelian Lagrangian fibrations on singular symplectic varieties; see Theorem 3.6. We specialise the support theorem to the Hitchin fibration in Theorem 3.11.
- In Section 4 we discuss toric Lawrence varieties and hypertoric quiver varieties. The main result is the local Hodge-to-singular correspondence, i.e. Theorem 4.18.
- In Section 5 we explain how (and in which sense) we can embed the Dolbeault moduli space into the universal compactified Jacobian.
- In Section 7 we describe local models for this embedding; see Theorem 7.4. To this purpose we need some auxiliary results: an explicit description of the image of the Kodaira–Spencer map for nodal spectral curves provided in Section 6; the topological trivialization of a tubular neighbourhood of an abelian variety in the universal compactified Jacobian in Section 8.
- In Section 9 we show the degree independence of the intersection cohomology of the universal compactified Jacobian; see Theorem 9.5.
- In Section 10 we collect the proof of the main theorems stated in Section 1.

**1.6. Notation.** The intersection cohomology of a complex variety  $X$  with middle perversity and rational coefficients is denoted by  $IH^*(X, \mathbb{Q}) := H^*(IC(X, \mathbb{Q}))$ , where  $IC(X, \mathbb{Q})$  is the perverse intersection cohomology complex of  $X$  shifted by  $-\dim X$ . Ordinary singular cohomology with rational coefficients is denoted by  $H^*(X, \mathbb{Q})$ . Recall that they all carry mixed Hodge structures. Intersection complexes and perverse sheaves on an algebraic stack are descents of perverse sheaves from a smooth atlas; see [41, 45, 44]. The formalism of six operations works in this context too. We use it in Section 7.

**1.7. Acknowledgements.** We would like to thank Alastair Craw, Mark de Cataldo, Jochen Heinloth, Daniel Huybrechts, Davesh Maulik, Jesse Leo Kass, Tasuki Kinjo, Nicola Pagani, Roberto Pagaria, Alex Perry, Giulia Saccà, Junliang Shen, Filippo Viviani for numerous discussions, emails and helpful advice.

The first author was supported by the University of Michigan and the Max Planck Institute for Mathematics. The second author was supported by PRIN Project 2017 "Moduli and Lie theory".

## 2. STRATIFICATIONS

Recall that a *stratification* of a complex variety  $X$  is a finite collection of locally closed smooth subvarieties  $X_i \subseteq X$ , called *strata*, such that

- $X$  is the disjoint union of  $X_i$  with  $i = 1, \dots, r$ , i.e.  $X = \bigsqcup_{i=1}^r X_i$ ;
- $X_i$  is contained in the closure of  $X_{i+1}$ .

A stratification is called a *Whitney stratification* if all pairs of strata satisfy the Whitney conditions A and B. We omit the precise definition of these conditions (see for instance [30]), since in the following we will only use the stronger property Definition 2.1, that implies Whitney conditions A and B by Lemma 2.2.

**Definition 2.1.** *A stratification  $X = \bigsqcup_i X_i$  is analytically trivial in the normal direction to each stratum, if for any  $x \in X_i$  there exists a normal slice  $N_x$  through  $X_i$  at  $x$ , and a neighbourhood of  $x$  in  $X$  which is locally analytically isomorphic to  $N_x \times T_x X_i$  at  $(x, 0)$ .*

**Lemma 2.2.** *A stratification analytically trivial in the normal direction to each stratum is a Whitney stratification.*

*Proof.* We follow the argument in [53, §4.7]. Whitney conditions A and B are local, so it is enough to check it for  $N_x \times T_x X_i$  at  $(x, 0)$ . Since  $T_x X_i$  is a smooth factor, it is enough to check it for  $N_x$  at  $x$ . Then this follows from [68, Lem. 19.3]. □

### 3. NGÔ THEOREM FOR LAGRANGIAN FIBRATIONS ON SINGULAR SPACES

The goal of this section is to generalise the description of Ngô strings in [22, Thm 7.0.3] to the case of weakly abelian Lagrangian fibrations on singular symplectic varieties.

We will follow closely Maulik and Shen [48]. Let  $B$  be a quasi-projective complex variety. Let  $g: P \rightarrow B$  be a smooth  $B$ -group scheme with connected fibers, and let  $f: X \rightarrow B$  be a projective morphism. Assume that the group scheme  $P$  acts on  $X$  via

$$(12) \quad \text{act}: P \times_B X \rightarrow X.$$

**Definition 3.1** (Weak abelian fibration). *The triple  $(X, P, B)$  is a weak abelian fibration of relative dimension  $c$ , if*

(1) *every fiber of the map  $g$  is pure of dimension  $c$ , and  $X$  has pure dimension*

$$\dim X = c + \dim B,$$

(2) *the action (12) of  $P$  on  $X$  has affine stabilizers, and*

(3) *the Tate module  $T_{\mathbb{Q}_l}(P)$  associated with the group scheme  $P$  is polarizable.*

For a closed point  $b \in B$ , we define  $\delta(b)$  as the dimension of the maximal affine and connected subgroup of  $P_s$ . For any closed subvariety  $Z \subseteq B$ , we denote by  $\delta_Z$  the minimum value of the (upper semi-continuous) function  $\delta$  on  $Z$ .

**Theorem 3.2.** *Let  $(X, P, B)$  be a weak abelian fibration. Assume that*

(1) *( $\delta$ -regularity)  $\text{codim } Z_\delta \geq \delta$  for any  $Z_\delta := \{b \in B \mid \delta(b) = \delta\}$ ;*

(2)  *$\tau_{>2c}(Rf_* \text{IC}(X, \mathbb{Q})) = 0$  for the standard truncation functor  $\tau_{>*}(-)$ .*

*Then every support  $Z$  of  $Rf_* \text{IC}(X, \mathbb{Q})$  satisfies*

$$\text{codim } Z = \delta(Z).$$

*Proof.* The  $\delta$ -regularity and [48, Theorem 1.8] implies that a support  $Z$  of  $Rf_* \text{IC}(X, \mathbb{Q})$  satisfies  $\text{codim } Z = \delta_Z$ . □

**Proposition 3.3** (Relative Dimension Bound). *Let  $f: X \rightarrow B$  be an equidimensional morphism of complex algebraic varieties of relative dimension  $c$ . Suppose that there exists a Whitney stratification  $X = \bigsqcup X_i$  such that for any  $b \in B$  we have*

$$(13) \quad \text{codim}_X X_i \geq 2 \text{codim}_{f^{-1}(b)}(X_i \cap f^{-1}(b)).$$

*Then we have*

- $\tau_{>2c}(Rf_* \text{IC}(X, \mathbb{Q})) = 0$ ;
- $R^{2c}f_* \text{IC}(X, \mathbb{Q}) = R^{2c}f_* \mathbb{Q}_X$ .

*Proof.* Via proper base change we have

$$\mathcal{H}^*(Rf_* \text{IC}(X, \mathbb{Q}))_b = H^*(f^{-1}(b), \text{IC}(X, \mathbb{Q})|_{f^{-1}(b)}) \text{ for any } b \in B.$$

These cohomology groups are the limit of the Grothendieck spectral sequence

$$E_2^{p,q} = H^{p-q}(Z_q \cap f^{-1}(b), \mathcal{H}^q(\text{IC}(X, \mathbb{Q}))|_{Z_q \cap f^{-1}(b)}) \Rightarrow H^p(f^{-1}(b), \text{IC}(X, \mathbb{Q})|_{f^{-1}(b)}),$$

where the closed subset  $Z_q \subset X$  is the support of the constructible sheaf  $\mathcal{H}^q(\text{IC}(X, \mathbb{Q}))$ . In particular, we have

$$E_2^{p,q} = 0 \text{ for } p - q > 2 \dim(Z_q \cap f^{-1}(b)).$$

The irreducible components of  $Z_q$  are the closure of some  $X_i$  with  $q < \text{codim}_X X_i$ , or  $X$  itself, if  $q = 0$ . Indeed, the strong support condition for intersection cohomology implies that  $\mathcal{H}^q(\text{IC}(X, \mathbb{Q}))_x = 0$  for any  $x \in X_i$  and  $q \geq \max\{\text{codim}_X X_i, 1\}$ ; see [20, §2.3]. By (13), we conclude that  $E_2^{p,q} = 0$  for

$$p > q + 2 \dim(Z_q \cap f^{-1}(b)) \geq \max\{\text{codim}_X X_i, 1\} + 2 \dim(Z_q \cap f^{-1}(b)),$$

i.e. for  $p > 2c$  and arbitrary  $q$ , or for  $p = 2c$  and  $q \neq 0$ . This yields

$$\mathcal{H}^{>2c}(Rf_* \text{IC}(X, \mathbb{Q})) = 0, \quad \mathcal{H}^{2c}(Rf_* \text{IC}(X, \mathbb{Q})) = \mathcal{H}^{2c}(Rf_* \mathbb{Q}_X).$$

□

**Proposition 3.4.** *Let  $X$  be a normal complex variety endowed with a holomorphic symplectic form and a proper Lagrangian fibration  $f : X \rightarrow B$ . Then  $X$  admits a complex Whitney stratification*

$$(14) \quad X = \bigsqcup X_i$$

*by locally closed algebraic symplectic submanifolds  $X_i$  such that the restriction map  $f|_{X_i} : X_i \rightarrow B_i := f(X_i)$  is Lagrangian. In particular, for any  $b \in B$  we have*

$$(15) \quad \text{codim}_X X_i = 2 \text{codim}_{f^{-1}(b)}(X_i \cap f^{-1}(b)).$$

*Proof.* By [42, Thm 2.3, Prop. 3.1],  $X$  admits a stratification by locally closed subsets  $X_i$ , with  $i = 1, \dots, r$ , such that:

- $X_i$  are smooth and symplectic;
- $X_{i+1}$  is the singular locus of  $X_i$ ;
- the stratification is analytically trivial in the normal direction to each stratum, so it is a Whitney stratification by Lemma 2.2.

Applying [47, Thm 3.1] iteratively, the restrictions  $f|_{X_i}$  are Lagrangian.

Since both  $f$  and  $f|_{X_i}$  are Lagrangian, we obtain  $\dim X - \dim X_i = 2 \dim f^{-1}(b) - 2 \dim(X_i \cap f^{-1}(b))$ . □

**Example 3.5.** The Whitney stratification (14) for the Dolbeault moduli space  $X = M(n, d)$  is the stratification (3). In particular, Proposition 3.3 gives

$$\tau_{>\dim M(n,d)}(R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})) = 0, \quad R^{\dim M(n,d)} f_* \mathrm{IC}(M(n, d), \mathbb{Q}) = R^{\dim M(n,d)} f_* \mathbb{Q}_{M(n,d)}.$$

Now let  $(X, P, B)$  be a  $\delta$ -regular weak abelian fibration. We denote by  $I$  the set of supports  $Z_\alpha \subseteq B$  for  $Rf_* \mathrm{IC}(X, \mathbb{Q})$ . There exist open dense subsets  $Z_\alpha^\times$  such that the restriction  $P_\alpha$  of the group scheme  $P$  admits a Chevalley decomposition

$$0 \rightarrow R_\alpha \rightarrow P_\alpha \rightarrow A_\alpha \rightarrow 0,$$

where  $R_\alpha \rightarrow Z_\alpha^\times$  is an affine group scheme and  $g_\alpha: A_\alpha \rightarrow Z_\alpha^\times$  is a proper abelian group scheme. Let  $\Lambda_\alpha^l := R^l g_{\alpha*} \mathbb{Q}_{A_\alpha}$ .

**Theorem 3.6** (Ngô strings from singular symplectic varieties). *Let  $(X, P, B)$  be a  $\delta$ -regular weak abelian fibration on a normal holomorphic symplectic variety  $X$  of dimension  $2c$ . Then there exists an isomorphism in  $D^b \mathrm{MHM}_{\mathrm{alg}}(B)$  (resp.  $D^b(B, \mathbb{Q})$ )*

$$(16) \quad Rf_* \mathrm{IC}(X, \mathbb{Q}) \simeq \bigoplus_{\alpha \in I} \bigoplus_{l=0}^{2 \dim Z_\alpha} \mathrm{IC}(Z_\alpha, \Lambda_\alpha^l \otimes \mathcal{L}_\alpha)[-l] \langle \mathrm{codim}(Z_\alpha) \rangle,$$

where  $\mathcal{L}_\alpha$  are local systems on an open subset of  $Z_\alpha$ . Moreover, we have:

(i)  $\mathrm{codim} Z_\alpha = \delta(Z_\alpha)$ ;

(ii)  $R^{2c} f_* \mathbb{Q}_X$  admits the following direct summand  $\bigoplus_{\alpha \in I} i_{\alpha*} \mathcal{L}_\alpha$ , where  $i_\alpha: Z_\alpha^\times \rightarrow B$  is the natural immersion.

*Proof.* The constraint (i) is a combination of Theorem 3.2, Propositions 3.3 and 3.4. In [48, Prop. 1.5] Maulik and Shen provided a generalization of Ngô freeness [58, Prop. 7.4.10] to the singular context. This gives the special form (16) to the decomposition theorem for  $Rf_* \mathrm{IC}(X, \mathbb{Q})$ . Taking cohomology of (16), together with the isomorphism  $R^{2c} f_* \mathrm{IC}(X, \mathbb{Q}) = R^{2c} f_* \mathbb{Q}_X$  in Proposition 3.3, we obtain (ii).  $\square$

**3.1. Ngô strings for Dolbeault moduli spaces.** We specialize Theorem 3.6 to the Hitchin fibration  $\chi(n, d): M(n, d) \rightarrow A_n$ . We rely on the recent key result by Kinjo and Koseki [43], building on the work of Davison [14].

**Proposition 3.7.** *Let  $d$  and  $e$  be integers, with  $e$  coprime to  $n$ . The complex  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$  is a direct summand of  $R\chi(n, e)_* \mathbb{Q}_{M(n,e)}$ .*

*Proof.* [14, Thm 6.6] and [43, Cor. 5.15, Ex. 5.18].  $\square$

Proposition 3.7 allows to extend [17, Thm 4.1] and [38, Thm 1] to Dolbeault moduli spaces of arbitrary degree.

**Corollary 3.8.** *If  $Z$  is a support of  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$  with  $Z \cap A_n^{\mathrm{red}} \neq \emptyset$ , then  $Z = S_{\underline{n}}$  for some partition  $\underline{n}$  of  $n$ . In particular, the generic point of any support  $Z$  of  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$  with  $Z \cap A_n^{\mathrm{red}} \neq \emptyset$  lies in the locus  $A_n^\times$  of nodal spectral curves.*

*Proof.* [17, Thm 4.1] and Proposition 3.7. □

**Corollary 3.9** (Proposition 1.6). *The intersection form on  $IH^*(M(n, d), \mathbb{Q})$  is trivial. Equivalently the forgetful map  $IH_c^*(M(n, d), \mathbb{Q}) \rightarrow IH^*(M(n, d), \mathbb{Q})$  is 0.*

*Proof.* By [37, Prop. 4] the vanishing of the intersection form is equivalent to show that  $\{0\} \subset A_n$  is not a support of  $R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})$ . In the coprime case, this is proved by Heinloth in [38, Thm 1]. Together with Proposition 3.7, we obtain the result in the non-coprime case. □

**Remark 3.10.** Proposition 1.6 holds for moduli space of  $\mathrm{PGL}_n$ - and  $\mathrm{SL}_n$ -Higgs bundles too.

**Theorem 3.11** (Ngô strings from Dolbeault moduli spaces in arbitrary degree). *There exists an isomorphism in  $D^b \mathrm{MHM}_{\mathrm{alg}}(A_n^{\mathrm{red}})$  (resp.  $D^b(A_n^{\mathrm{red}}, \mathbb{Q})$ )*

$$(17) \quad R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})|_{A_n^{\mathrm{red}}} \simeq \bigoplus_{\underline{n} \in I_d} \mathcal{S}(\mathcal{L}_{\underline{n}}(d))|_{A_n^{\mathrm{red}}},$$

where  $I_d$  is a set of partition of  $\underline{n}$  of  $n$ , and the Ngô string  $\mathcal{S}_{\underline{n}}$  is the mixed Hodge module supported on  $S_{\underline{n}}$  given by

$$\mathcal{S}(\mathcal{L}_{\underline{n}}(d)) := \bigoplus_{l=0}^{2 \mathrm{codim} S_{\underline{n}}} \mathrm{IC}(S_{\underline{n}}, \Lambda_{\underline{n}}^l \otimes \mathcal{L}_{\underline{n}}(d))[-l](\mathrm{codim} S_{\underline{n}}),$$

for certain local systems  $\mathcal{L}_{\underline{n}}(d)$  supported on an open set of  $S_{\underline{n}}$ .

*Proof.* It follows from Theorem 3.6 and Corollary 3.8. □

The rest of the paper is devoted to the correspondence between the local systems  $\mathcal{L}_{\underline{n}}(d)$  and the singularities of  $M(n, d)^{\mathrm{red}}$ .

#### 4. HYPERTORIC QUIVER VARIETIES

The singularities of  $M(n, d)^{\mathrm{red}}$  are modelled on hypertoric quiver varieties. In this section we introduce these varieties and establish a local version of the Hodge-to-singular correspondence (Theorem 4.18).

**4.1. Affine toric varieties associated to a quiver.** Let  $Q = (V, E)$  be a connected directed graph (a *quiver*) with  $r$  vertices  $V = \{v_1, \dots, v_r\}$  and  $s$  oriented edges  $E$ . Given  $e \in E$ , let  $s(e), t(e) \in V$  be the source and target of  $e$ , so that  $e : s(e) \rightarrow t(e)$ . The underlying unoriented graph, denoted by  $|Q|$ , has first Betti number  $b_1 := s - r + 1$ .

We consider the group of all  $\mathbb{Z}$ -linear combinations of  $V$  whose coefficients sum to zero. We identify this group with  $\mathbb{Z}^{r-1}$ , by fixing the basis  $\{v_1 - v_2, \dots, v_1 - v_r\}$ . We also identify  $\mathbb{Z}^s$  with the group of  $\mathbb{Z}$ -linear combinations of  $E$ . The boundary map of the quiver  $Q$

$$A : \mathbb{Z}^s \rightarrow \mathbb{Z}^{r-1}, \quad e \mapsto s(e) - t(e)$$

induces an embedding of algebraic tori

$$\mathbb{T}^{r-1} := \text{Spec } \mathbb{C}[\mathbb{Z}^{r-1}] \hookrightarrow \mathbb{T}^s := \text{Spec } \mathbb{C}[\mathbb{Z}^s].$$

The torus  $\mathbb{T}^s$  acts on the affine complex space  $\mathbb{C}^s = \text{Spec } \mathbb{C}[z_e : e \in E]$ , and so does  $\mathbb{T}^{r-1}$  by restriction. Explicitly,  $\lambda = (\lambda_i)_{i=1}^r \in \mathbb{T}^r / \mathbb{T} = \mathbb{T}^{r-1}$  acts as  $\lambda \cdot z_e = \lambda_{s(e)} z_e \lambda_{t(e)}^{-1}$ .

**Definition 4.1.** *The affine toric variety  $X(Q, 0)$  is the affine GIT quotient of  $\mathbb{C}^s$  by  $\mathbb{T}^{r-1}$*

$$X(Q, 0) := \mathbb{C}^s //_0 \mathbb{T}^{r-1} = \text{Spec } \mathbb{C}[z_e : e \in E]^{\mathbb{T}^{r-1}}.$$

The cone of the toric variety  $X(Q, 0)$  is generated by the rows  $\mathcal{B} = \{\beta_1, \dots, \beta_s\} \subset \mathbb{Z}^{b_1}$  of a *Gale dual* of  $A$ , i.e. a  $(s \times b_1)$ -matrix  $B$  whose columns form a base of  $\ker(A : \mathbb{Z}^s \rightarrow \mathbb{Z}^{r-1}) \simeq \mathbb{Z}^{b_1}$ , or equivalently a matrix that sits in the following exact sequence:

$$0 \rightarrow \mathbb{Z}^{b_1} \xrightarrow{B} \mathbb{Z}^s \xrightarrow{A} \mathbb{Z}^{r-1} \rightarrow 0.$$

Recall that a loop in a quiver is an edge whose head and tail coincide.

**Lemma 4.2.** *Let  $Q$  be a quiver with  $m$  loops, and  $Q'$  be the quiver obtained from  $Q$  by deleting all the loops. Then we have*

$$X(Q, 0) \simeq X(Q', 0) \times \mathbb{C}^m.$$

*Proof.* Let  $L \subset E$  be the loops of  $Q$ . We identify the edges  $E'$  of  $Q'$  with  $E \setminus L$ . For any  $l \in L$ , the coordinates  $z_l$  are  $\mathbb{T}^{r-1}$ -invariant by construction, so we have

$$\mathbb{C}[z_e : e \in E]^{\mathbb{T}^{r-1}} \simeq \mathbb{C}[z_e : e \in E']^{\mathbb{T}^{r-1}} \otimes \mathbb{C}[z_l : l \in L].$$

□

**4.2. Semiprojective toric varieties associated to a quiver.** In general,  $X(Q, 0)$  is a singular variety. A toric resolution of singularities can be obtained via projective GIT as follows.

The ring of polynomials  $S := \mathbb{C}[z_e : e \in E]$  is graded by  $\mathbb{Z}^{r-1}$  in such a way that  $\deg(z_e) = Ae$ . For  $\theta \in \mathbb{Z}^{r-1}$ , let  $S_\theta$  be the  $\mathbb{C}$ -vector space of homogenous polynomials of degree  $\theta$ . For instance,  $S_0$  is the coordinate ring of  $X(Q, 0)$ .

**Definition 4.3.** *The semiprojective toric variety  $X(Q, \theta)$  is the projective GIT quotient of  $\mathbb{C}^{2s}$  by the torus  $\mathbb{T}^{r-1}$  with respect to the stability  $\theta \in \mathbb{Z}^{r-1}$ , i.e.*

$$X(Q, \theta) := \mathbb{C}^s //_{\theta} \mathbb{T}^{r-1} := \text{Proj} \left( \bigoplus_{k=0}^{\infty} t^k S_{k\theta} \right).$$

Consider now the coarsest complete fan  $\Gamma(\mathcal{A})$  in  $\mathbb{R}^{r-1} \simeq \mathbb{Z}^{r-1} \otimes \mathbb{R}$  that refines all complete simplicial fans in  $\mathbb{R}^{r-1}$  whose rays lie in  $\mathcal{A} := \{Ae : e \in E\}$ . We say that  $\theta \in \mathbb{Z}^{r-1}$  is generic if it lies in the interior of the maximal cones of  $\Gamma(\mathcal{A})$ .

**Proposition 4.4.** [34, Prop. 8.2] *For generic  $\theta \in \mathbb{Z}^{r-1}$ , the affinization morphism*

$$\pi_X = \pi_X(Q, \theta): X(Q, \theta) \rightarrow \text{Spec } H^0(X(Q, \theta), \mathcal{O}_{X(Q, \theta)}) = X(Q, 0)$$

*is a resolution of singularities.*

**4.3. Lawrence varieties and hypertoric quiver varieties.** We specialize the previous constructions to doubled quivers. Let  $Q$  be a quiver with  $r$  vertices,  $s$  oriented edges  $E$ , and first Betti number  $b_1$ .

**Definition 4.5.** *The double of  $Q$ , denoted by  $Q^\pm$ , is the quiver obtained from  $Q$  by replacing each edge  $e$  of  $Q$  with a pair of edges  $e^+$  and  $e^-$  having the same endpoints as  $e$  but opposite orientations.*

**Definition 4.6.** *The affine Lawrence toric variety associated to  $Q$  is the affine GIT quotient of  $\mathbb{C}^{2s}$  by the torus  $\mathbb{T}^{r-1}$*

$$X(Q^\pm, 0) := \mathbb{C}^{2s} //_0 \mathbb{T}^{r-1}.$$

*The Lawrence toric variety associated to  $Q$  is the projective GIT quotient of  $\mathbb{C}^{2s}$  by the torus  $\mathbb{T}^{r-1}$  with respect to a stability  $\theta \in \mathbb{Z}^{r-1}$ , i.e.*

$$X(Q^\pm, \theta) := \mathbb{C}^{2s} //_{\theta} \mathbb{T}^{r-1} := \text{Proj} \left( \bigoplus_{k=0}^{\infty} t^k S_{k\theta} \right).$$

**Notation 4.7.** For convenience, we relabel the coordinates  $z_{e^+}$  and  $z_{e^-}$  of  $\mathbb{C}^{2s}$  by  $z_e$  and  $w_e$ , indexed by  $e \in E$ . Note that the definition of  $X(Q^\pm, \theta)$  is independent of the choice of an orientation of  $Q$ : a different choice corresponds to exchange some coordinates  $z_e$  with the corresponding  $w_e$ .

Suppose that the matrix  $A = (a_{ie})$  corresponds to the boundary map for  $Q$ . We consider the homogeneous ideal

$$(18) \quad \text{Circ}(\mathcal{B}) := \left\langle \sum_{e \in E} a_{ie} z_e w_e : i = 2, \dots, r \right\rangle \subset S.$$

**Definition 4.8.** *The hypertoric quiver variety  $Y(Q, 0)$  (resp.  $Y(Q, \theta)$ ) is the irreducible subvariety of the affine Lawrence variety  $X(Q^\pm, 0)$  (resp. of the Lawrence variety  $X(Q^\pm, \theta)$ ) cut by the homogeneous ideal  $\text{Circ}(\mathcal{B})$ .*

**Remark 4.9.** Hypertoric quiver variety are both hypertoric varieties in the sense of [5, 32], and Nakajima quiver variety whose dimension vector has all coordinates equals to one; see [57]. In [34],  $Y(Q, 0)$  is called *toric quiver variety*, but note that it is only rarely toric; see [34, §10]. Moreover, Altmann and Hille called toric quiver varieties the fibre over 0 of the morphism  $\pi_Y : Y(Q, \theta) \rightarrow Y(Q, 0)$ . To avoid confusion, we will refer to  $Y(Q, 0)$  and  $Y(Q, \theta)$  as hypertoric quiver variety.

Since the product  $z_e w_e$  is  $\mathbb{T}^{r-1}$ -invariant, the injective morphism of  $\mathbb{C}$ -algebras

$$\mathbb{C}[z_e w_e : e \in E] \hookrightarrow \mathbb{C}[z_e, w_e : e \in E]^{\mathbb{T}^{r-1}}$$

induces the fibre product square

$$(19) \quad \begin{array}{ccc} Y(Q, 0) = \text{Spec} \left( \frac{\mathbb{C}[z_e, w_e : e \in E]^{\mathbb{T}^{r-1}}}{\text{Circ}(\mathcal{B})} \right) & \xleftarrow{\iota_M} & X(Q^\pm, 0) = \text{Spec} (\mathbb{C}[z_e, w_e : e \in E]^{\mathbb{T}^{r-1}}) \\ \chi_Q \downarrow & & \downarrow \pi_Q \\ \mathbb{C}^{b_1} = \text{Spec} \left( \frac{\mathbb{C}[z_e w_e : e \in E]}{\text{Circ}(\mathcal{B})} \right) & \xleftarrow{\iota_A} & \mathbb{C}^s = \text{Spec} (\mathbb{C}[z_e w_e : e \in E]). \end{array}$$

Hence,  $Y(Q, 0)$  is the inverse image under the map  $\pi_Q$  of the linear subspace  $\mathbb{C}^{b_1}$ .

**4.4. Cohomology of hypertoric quiver varieties.** The cohomology of hypertoric quiver varieties can be expressed in terms of the cohomology of the cographic matroid. This permits to compare the Hodge-to-singular correspondence with [17, Thm 6.11].

**Definition 4.10.** *The cographic matroid  $\mathcal{C}_Q$  of the unoriented graph  $|Q|$  is the matroid whose independent subsets are the subsets  $I$  of unoriented edges of  $|Q|$  such that  $|Q| \setminus I$  is connected. The set  $\mathcal{C}_Q$  is partially ordered with respect to inclusions and we denote by  $|\mathcal{C}_Q|$  the associated simplicial complex, i.e. the complex whose  $k$ -dimensional faces are the independent subsets of cardinality  $(k+1)$ .*

We now recall some facts about the affinization morphisms

$$\pi_X : X(Q^\pm, \theta) \rightarrow X(Q^\pm, 0), \quad \pi_Y : Y(Q, \theta) \rightarrow Y(Q, 0).$$

In particular, we observe that the top cohomology of  $Y(Q, \theta)$  can be identified with the top cohomology of  $|\mathcal{C}_Q|$ , which explains the occurrence of the cographic matroid in [17, §6] via the Hodge-to-singular correspondence; see Sections 7 and 10.

In the following all the stability  $\theta, \theta' \in \mathbb{Z}^{r-1}$  are generic.

**Proposition 4.11.** *The following facts hold:*

- (1) [34, Prop. 8.2]  $X(Q^\pm, \theta)$  is a smooth toric variety of dimension  $b_1 + s$ .
- (2) [34, Prop. 8.2]  $Y(Q, \theta)$  is a smooth holomorphic symplectic variety of dimension  $2b_1$ .
- (3) [34, Lem. 6.4]  $\pi_X^{-1}(0) = \pi_Y^{-1}(0) =: C(Q^\pm, \theta)$ .
- (4) The irreducible components of  $C(Q^\pm, \theta)$  are Lagrangian in  $Y(Q, \theta)$ . In particular, they all have dimension  $b_1$ .
- (5) [34, Rmk after Lem. 6.5] The spaces  $C(Q^\pm, \theta) \subset Y(Q, \theta) \subset X(Q^\pm, \theta)$  are deformation retracts in one another.
- (6) [32, Lem. 2.1] [34, Thm 6.3] The symplectic resolutions  $Y(Q, \theta)$  and  $Y(Q, \theta')$  are diffeomorphic.
- (7) By (5) and (6), the cohomology

$$H^*(C(Q^\pm, \theta), \mathbb{Q}) \simeq H^*(Y(Q, \theta), \mathbb{Q}) \simeq H^*(X(Q^\pm, \theta), \mathbb{Q})$$

is independent of  $\theta$ .

- (8) [6, Thm 7.3.3, 7.8.1] There exists an isomorphism

$$H^{2b_1}(Y(Q, \theta), \mathbb{Q}) \simeq \tilde{H}^{2b_1-1}(|\mathcal{C}_Q|, \mathbb{Q}).$$

*Proof.* We just comment on (4). An irreducible component  $F$  of  $C(Q^\pm, \theta)$  is a fibre of  $Y(Q, \theta) \rightarrow Y(Q, 0) \xrightarrow{\chi_Q} \mathbb{C}^{b_1}$ . By semicontinuity,  $\dim F \geq b_1$ . On the other hand, the symplectic form vanishes along  $F$ , because  $F$  is toric and so  $H^0(F, \Omega_F^2) = 0$ . So  $F$  is isotropic, and  $\dim F \leq b_1$ .  $\square$

**4.5. Decomposition theorem for hypertoric quiver varieties.** The goal of this section is to describe the restriction  $\mathrm{IC}(X(Q^\pm, 0), \mathbb{Q})|_{Y(Q, 0)}$ ; see Theorem 4.18. To this end, we briefly recall the statement of the decomposition theorem for semismall maps.

Let  $f: X \rightarrow Y$  be a proper morphism of irreducible varieties. A stratification of  $f$  is a collection of finitely many locally closed subsets  $Y_k$  of dimension  $\dim Y - k$  such that  $f^{-1}(Y_k) \rightarrow Y_k$  are topologically locally trivial fibrations. A stratum  $Y_k$  is *relevant* if  $2 \dim f^{-1}(Y_k) - \dim(Y_k) = \dim X$ .

**Definition 4.12** (Semismall maps). *The map  $f$  is said to be semismall if*

$$2 \dim f^{-1}(Y_k) - \dim(Y_k) \leq \dim X \quad \text{for any } k \geq 0.$$

Further if the stronger inequalities

$$2 \dim f^{-1}(Y_k) - \dim(Y_k) < \dim X \quad \text{for any } k > 0$$

hold, we say that  $f$  is small.

**Theorem 4.13** (Decomposition theorem for semismall maps). *Let  $f: X \rightarrow Y$  be a semismall map from a smooth variety  $X$ . Then there exists a canonical isomorphism in  $D^bMHHM_{alg}(Y)$  (resp.  $D^b(Y, \mathbb{Q})$ )*

$$Rf_*\mathbb{Q}_X \simeq \bigoplus_{Y_k} \text{IC}(\overline{Y}_k, \mathbb{R}^{\text{codim } Y_k} f_*\mathbb{Q}_{f^{-1}(Y_k)})\langle \text{codim } Y_k/2 \rangle,$$

where the summation index runs over all the relevant strata of a stratification of  $f$ .

In particular, if  $f$  is small, then  $Rf_*\mathbb{Q}_X \simeq \text{IC}(Y, \mathbb{Q})$ .

Given a quiver  $Q = (V, E)$ , we describe complex Whitney stratifications of the toric Lawrence variety  $X(Q^\pm, 0)$  and of hypertoric quiver variety  $Y(Q, 0)$ .

**Definition 4.14.** *A partition  $\underline{V}$  of the set  $V$  is a set  $\{V_1, \dots, V_r\}$  of disjoint subsets  $V_i \subset V$  such that  $V = \bigsqcup_{i=1}^r V_i$ . The quiver  $Q_{\underline{V}}$  is obtained from  $Q$  by identifying all vertices in  $V_i$ , for any  $i = 1, \dots, r$ , and by deleting all loops.*

**Proposition 4.15.** *The toric Lawrence variety  $X(Q^\pm, 0)$  admits a complex Whitney stratification*

$$(20) \quad X(Q^\pm, 0) = \bigsqcup_{\underline{V} \dashv V} X_{\underline{V}},$$

by toric-invariant strata  $X_{\underline{V}}$  whose analytic normal slice is isomorphic to  $X(Q_{\underline{V}}^\pm, 0)$ . This induces a complex Whitney stratification of  $Y(Q, 0)$

$$(21) \quad Y(Q, 0) = \bigsqcup_{\underline{V} \dashv V} Y_{\underline{V}} \quad \text{with } Y_{\underline{V}} := X_{\underline{V}} \cap Y(Q, 0),$$

whose normal slice is isomorphic to  $Y(Q_{\underline{V}}, 0)$ .

*Proof.* Let  $N$  be the lattice of one-parameter subgroups of the torus  $\mathbb{T}^{b_1+s} = N \otimes \mathbb{C}^*$ . The affine toric variety  $X(Q^\pm, 0)$  corresponds to the cone  $\sigma \subset N \otimes \mathbb{R}$  generated by the columns of a matrix  $B^\pm$  Gale dual to the boundary map  $A^\pm := (A, -A): \mathbb{Z}^{2s} \rightarrow \mathbb{Z}^{r-1}$  of the quiver  $Q^\pm$ ; see Section 4.1. Let  $\tau$  be a face of  $\sigma$ ,  $N_\tau$  be the sublattice of  $N$  spanned by  $\tau \cap N$ , and  $N(\tau) := N/N_\tau$ . The sets of rays of  $\sigma$ ,  $\tau$ , and those of  $\sigma$  not in  $\tau$  are denoted respectively  $\sigma(1)$ ,  $\tau(1)$  and  $\tau(1)^c$ . The exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^{\tau(1)} & \longrightarrow & \mathbb{Z}^{\sigma(1)} & \longrightarrow & \mathbb{Z}^{\tau(1)^c} \longrightarrow 0 \\ & & \downarrow (B_\tau^\pm)^T & & \downarrow (B^\pm)^T & & \downarrow \\ 0 & \longrightarrow & N_\tau & \longrightarrow & N & \longrightarrow & N(\tau) \longrightarrow 0 \end{array}$$

are dual to

$$\begin{array}{ccccccc}
0 & \longrightarrow & M(\tau) & \longrightarrow & \mathbb{Z}^{\tau(1)^c} & \xrightarrow{A^\pm|_{\tau(1)^c}} & \mathbb{Z}^{r-1} \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & M & \longrightarrow & \mathbb{Z}^{\sigma(1)} & \xrightarrow{A^\pm} & \mathbb{Z}^{r-1} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & M_\tau & \longrightarrow & \mathbb{Z}^{\tau(1)} & \xrightarrow{A_\tau^\pm} & \mathbb{Z}^{r_\tau-1} \longrightarrow 0.
\end{array}$$

In particular, the resulting map  $A_\tau^\pm$  is the boundary map of a quiver  $Q(\tau)$  obtained by contracting the edges corresponding to  $\tau(1)^c$ . Up to loops,  $Q(\tau)$  is isomorphic to  $Q_{\underline{V}}$  for some partition  $\underline{V}$  of  $V$ . Therefore,  $X(Q^\pm, 0)$  is covered by toric charts of the form

$$(22) \quad X(Q(\tau), 0) \times N(\tau) \otimes \mathbb{C}^* \simeq X(Q_{\underline{V}}^\pm, 0) \times \text{smooth factor},$$

where the isomorphism follows from Lemma 4.2. We have obtained the stratification (20) which is Whitney by Lemma 2.2. By inspecting the restriction of the equations (18) to each chart (22), we get the Whitney stratification (21) too; see also [61, §2].  $\square$

**Proposition 4.16.** *The map  $\pi_X : X(Q^\pm, \theta) \rightarrow X(Q^\pm, 0)$  is small.*

*Proof.* The fan of  $X(Q^\pm, \theta)$  is a regular triangulation of the cone  $\sigma \subset N \otimes \mathbb{R}$  of  $X(Q^\pm, 0)$ ; see [34, §2]. Since any regular triangulation of  $\sigma$  restricts to a regular triangulation of its face  $\tau$ , the restriction  $\pi_X$  to the open set  $X(Q_{\underline{V}}^\pm, 0) \times X_{\underline{V}}$  is the resolution

$$X(Q_{\underline{V}}^\pm, [\theta]) \times X_{\underline{V}} \rightarrow X(Q_{\underline{V}}^\pm, 0) \times X_{\underline{V}}.$$

Therefore, given  $y \in X_{\underline{V}} \subset X(Q^\pm, 0) \times X_{\underline{V}}$  we obtain

$$2 \dim \pi_X^{-1}(y) = 2 \dim C(Q_{\underline{V}}^\pm, [\theta]) = 2b_1(Q_{\underline{V}}) < b_1(Q_{\underline{V}}) + s = \dim X(Q_{\underline{V}}^\pm, 0) = \dim X(Q^\pm, 0) - \dim X_{\underline{V}},$$

i.e.  $\pi_X$  is small.  $\square$

**Proposition 4.17.** *The map  $\pi_Y : Y(Q, \theta) \rightarrow Y(Q, 0)$  is semismall.*

*Proof.* By Proposition 4.11.(2),  $Y(Q, \theta)$  is a smooth holomorphic symplectic variety. Any symplectic resolution of singularities is semismall by [42, Lemma 2.11].  $\square$

**Theorem 4.18** (Local Hodge-to-singular correspondence). *The restriction  $\mathrm{IC}(X(Q^\pm, 0), \mathbb{Q})|_{Y(Q, 0)}[2b_1]$  is a semi-simple perverse sheaf. In particular, we have*

$$\mathrm{IC}(X(Q^\pm, 0), \mathbb{Q})|_{Y(Q, 0)}[2b_1] \simeq \bigoplus_{\underline{V} \vdash V} \mathrm{IC}(Y_{\underline{V}}, \mathbb{Q}) \otimes H^{b_1(Q_{\underline{V}})}(Y(Q, \theta), \mathbb{Q})[2b_1](b_1(Q_{\underline{V}})).$$

*Proof.* Consider the cartesian square

$$\begin{array}{ccc} Y(Q, \theta) & \xrightarrow{\iota_{M, \theta}} & X(Q^\pm, \theta) \\ \pi_Y \downarrow & & \downarrow \pi_X \\ Y(Q, 0) & \xrightarrow{\iota_A} & X(Q^\pm, 0). \end{array}$$

The map  $\pi_X$  is small, the map  $\pi_Y$  is semismall and the inclusion  $\iota_{M, \theta}$  is a regular embedding. Therefore, the decomposition theorem for small maps gives  $\mathrm{IC}(X(Q^\pm, 0), \mathbb{Q}) \simeq R\pi_{X,*}\mathbb{Q}_{X(Q^\pm, \theta)}$ . By proper base change, we obtain  $R\pi_{X,*}\mathbb{Q}_{X(Q^\pm, \theta)}|_{Y(Q, 0)} \simeq R\pi_{Y,*}\mathbb{Q}_{Y(Q, \theta)}$ . The statement now follows from the decomposition theorem for semismall maps. In this case, the local systems  $R^{\dim X - \dim Y_k} f_* \mathbb{Q}_{f^{-1}(Y_k)}$  of Theorem 4.13 are trivial, supported on the relevant strata  $Y_{\underline{V}}$  and with fiber  $H^{\mathrm{top}}(\pi_Y^{-1}(y), \mathbb{Q}) \simeq H^{b_1(Q_{\underline{V}})}(Y(Q, \theta), \mathbb{Q})$  for all  $y \in Y_{\underline{V}}$ ; see [61, §5].  $\square$

## 5. UNIVERSAL COMPACTIFIED JACOBIANS

The goal of this section is to enlarge the Hitchin fibration to a universal family of compactified Jacobians. Fix integers  $g > 1$ ,  $n > 1$ ,  $d$  as above.

**Definition 5.1.** *A rank 1 torsion free sheaves  $\mathcal{I}$  on a curve  $X$  is semistable with respect to the canonical polarization  $\omega_X$  if for any proper subcurve  $Y \subset X$  we have that*

$$(23) \quad \frac{\chi(X, \mathcal{I})}{\mathrm{deg}(\omega_X)} \leq \frac{\chi(Y, \mathcal{I}_Y)}{\mathrm{deg}(\omega_X|_Y)},$$

where  $\mathcal{I}_Y$  is the biggest torsion free quotient of the restriction  $\mathcal{I}|_Y$  of  $\mathcal{I}$  to the subcurve  $Y$ . The **compactified Jacobian**  $\overline{\mathcal{J}}_d(X)$  is the projective coarse moduli space of semistable rank 1 torsion free sheaves on  $X$  of degree  $d + (n - n^2)(1 - g)$  with respect to the canonical polarization; see [66].

The BNR correspondence presents the Dolbeault moduli space  $M(n, d)$  as the relative compactified Jacobian of the spectral curve family; see [39, 2, 63]. It says that given a rank 1 torsion free sheaf  $\mathcal{I}$  on the spectral curve  $\pi_a: C_a \rightarrow C$ , the sheaf  $\mathcal{E} \simeq \pi_{a,*}\mathcal{I}$  is a vector bundle of rank  $n$  equipped with a Higgs field  $\phi$ , induced from the  $\mathcal{O}_{C_a}$ -module structure. Viceversa, any Higgs bundle  $(\mathcal{E}, \phi)$  is the pushforward of some rank 1 torsion free sheaf on  $C_a$ .

**Proposition 5.2.** [39, 2, 63, 55] *The fibres of the Hitchin fibrations  $\chi^{-1}(n, d)(a)$  with  $a \in A_n^{\mathrm{red}}$  is isomorphic to the compactified Jacobian  $\overline{\mathcal{J}}_d(C_a)$  of the spectral curve  $C_a$ .*

**Remark 5.3.** The stability condition for rank 1 torsion free sheaves on reduced curves used in [55, (10.5)] and [17] coincides with (23). Suppose that  $X$  is a reduced spectral curve  $C_a = \bigcup_{i=1}^r C_i$  whose component  $C_i$  has degree  $n_i$ , and  $n = \sum_{i=1}^r n_i$ . Let  $\mathcal{I}$  be a rank 1 torsion free sheaf of degree

$d + (n - n^2)(1 - g)$  on  $C_a$ . By Riemann–Roch, (23) reads as follows: for any subcurve  $C_I = \bigcup_{i \in I} C_i$ , with  $I \subset \{1, \dots, r\}$ , we have that

$$\chi(Y, \mathcal{I}_Y) \geq (d + r(1 - g)) \frac{\sum_{i \in I} n_i}{n},$$

which is the stability condition [55, (10.5)].

Following Caporaso, Pandaripande and Esteves [7, 59, 25] (cf also [8, §2.3, Fact 2.12] and [17]) we can enlarge the Hitchin fibration to a relative compactified Jacobian of a versal deformation of  $C_a$ . To this end, let  $\overline{\mathcal{M}}_{g'}$  be the Deligne–Mumford stack of semi-stable curves of genus  $g' := n^2(g - 1) + 1$ , and  $A_n^\times := \bigsqcup_{\underline{n}+n} S_n^\times$ . By [17, Lem. 2.1] the flat universal spectral curve  $\mathcal{C}_n^\times \rightarrow A_n^\times$  induces a morphism

$$f^\times : A_n^\times \rightarrow \overline{\mathcal{M}}_{g'}.$$

**Proposition 5.4** (Universal compactified Jacobian). *There exists an open substack  $\mathcal{B} \subseteq \overline{\mathcal{M}}_{g'}$  containing  $f^\times(A_n^\times)$  and an irreducible (non necessarily smooth) Deligne–Mumford stack  $\pi : \overline{\mathcal{J}}_{d, \mathcal{B}} \rightarrow \mathcal{B}$  that étale locally is a relative compactified Jacobian parametrizing semistable rank 1 torsion free sheaves of degree  $d + (n - n^2)(1 - g)$  with respect to the canonical polarization.*

*Proof.* The first paragraph of the proof of [17, Prop. 5.9] does not require the coprimality of  $n$  and  $d$ , and works verbatim in our context.  $\square$

**Proposition 5.5.** *For every partition  $\underline{n}$  of  $n$ , let  $a \in S_{\underline{n}}^\times$ . There exists a local multisection  $A_a$  of  $f^\times : A_n^\times \rightarrow f^\times(A_n^\times) \subseteq \mathcal{B}$  passing through  $a$ , i.e.  $A_a$  is a locally closed subset of  $A_n^\times$  passing through  $a$ , intersecting  $S_{\underline{n}}^\times$  transversally, and such that the restriction  $f^\times|_{A_a} : A_a \rightarrow f^\times(A_n^\times)$  is étale. Furthermore we have a cartesian diagram*

$$\begin{array}{ccccc} \chi(n, d)^{-1}(A_a) & \xrightarrow{\cong} & \overline{\mathcal{J}}_{d, \mathcal{B}} \times_{\mathcal{B}} A_a & \longrightarrow & \overline{\mathcal{J}}_{d, \mathcal{B}} \\ & \searrow \chi(n, d) & \downarrow & & \downarrow \pi \\ & & A_a & \xrightarrow{f^\times|_{A_a}} & \mathcal{B}. \end{array}$$

*Proof.* [17, Cor. 5.10] works verbatim.  $\square$

**Remark 5.6.** The  $\mathbb{G}_m$ -action on  $A_n$  and the translation action  $H^0(C, \omega_C) \otimes A_n \rightarrow A_n$  lift to the universal spectral curve  $\mathcal{C}_n \rightarrow A_n$  and so to  $M(n, d)$ . Therefore, the group action induces an analytic isomorphism between an open neighbourhood  $U_a \subseteq A_n$  of  $A_a$  and a neighbourhood of the zero-section of trivial vector bundle  $p : A_a \times (H^0(C, \mathcal{O}_C) \oplus H^0(C, \omega_C)) \rightarrow A_a$ , such that there exists an homeomorphism

$$\chi(n, d)^{-1}(U_a) \simeq \chi(n, d)^{-1}(A_a) \times (H^0(C, \mathcal{O}_C) \oplus H^0(C, \omega_C)),$$

and so

$$R\chi(n, d)_* \mathrm{IC}(M(n, d), \mathbb{Q})|_{U_a} \simeq p^*(R\chi(n, d)_* \mathrm{IC}(\chi(n, d)^{-1}(A_a), \mathbb{Q})).$$

## 6. KODAIRA–SPENCER MAP FOR SPECTRAL CURVES

The differential of the map  $f^\times: A_a \rightarrow \mathcal{B}$  in Proposition 5.5 can be identified with the Kodaira–Spencer map; see for instance [17, §5.2]. Its kernel is isomorphic to  $H^0(C, \mathcal{O}_C) \oplus H^0(C, \omega_C)$ ; see again [17, §5.2]. In Proposition 6.3 we study its image.

**6.1. Kodaira–Spencer map.** We start with some recollection about the deformation of nodal spectral curves.

**Definition 6.1.** *The Kodaira–Spencer map is the map*

$$(24) \quad KS_a: T_a A_n \simeq H^0(C_a, N_{C_a/T^*C_a}) \rightarrow \mathrm{Ext}^1(\Omega_{C_a}^1, \mathcal{O}_{C_a})$$

obtained by applying the functor  $\mathrm{Hom}(-, \mathcal{O}_{C_a})$  to the relative cotangent sequence

$$(25) \quad 0 \rightarrow N_{C_a/T^*C}^* \rightarrow \Omega_{T^*C|C_a}^1 \rightarrow \Omega_{C_a}^1 \rightarrow 0.$$

Applying the functor  $\mathcal{H}om(-, \mathcal{O}_{C_a})$  to (25), we obtain the exact sequence of coherent sheaves

$$0 \rightarrow T_{C_a} \rightarrow T_{T^*C|C_a} \rightarrow N_{C_a/T^*C} \rightarrow \mathbb{T}^1 := \mathcal{E}xt^1(\Omega_{C_a}^1, \mathcal{O}_{C_a}) \rightarrow 0.$$

The first tangent sheaf  $\mathbb{T}^1$  is a skyscraper sheaf supported at the nodes  $E(C_a)$  of  $C_a$  and it parametrises smoothings of the nodes. The kernel of  $N_{C_a/T^*C} \rightarrow \mathbb{T}^1$  is easily identified to  $N_{C_a/T^*C} \otimes \mathcal{I}_{E(C_a)}$ , where  $\mathcal{I}_{E(C_a)}$  is the product of the maximal ideals of the nodes  $E(C_a)$ ; see [65, Ex. 4.7.1]. Hence, there exists a short exact sequence

$$0 \rightarrow N_{C_a/T^*C} \otimes \mathcal{I}_{E(C_a)} \rightarrow N_{C_a/T^*C} \rightarrow \mathbb{T}^1 \rightarrow 0$$

inducing the exact sequence in cohomology

$$0 \rightarrow H^0(C_a, N_{C_a/T^*C} \otimes \mathcal{I}_{E(C_a)}) \rightarrow T_a A_n = H^0(C_a, N_{C_a/T^*C}) \rightarrow \mathbb{T}^1(C_a).$$

Note that the last map  $T_a A_n \rightarrow \mathbb{T}^1$  factors as

$$(26) \quad T_a A_n = H^0(C_a, N_{C_a/T^*C}) \xrightarrow{KS_a} \mathrm{Ext}^1(\Omega_{C_a}^1, \mathcal{O}_{C_a}) \rightarrow H^0(C_a, \mathcal{E}xt^1(\Omega_{C_a}^1, \mathcal{O}_{C_a})) = \mathbb{T}^1(C_a),$$

where the latter map comes from the local-to-global Ext spectral sequence.

**6.2. Image of the Kodaira–Spencer map for spectral curves.** Let  $\underline{n}'$  be a partition of  $n$ . Then  $\chi(n, d)^{-1}(S_{\underline{n}'}^\times)$  admits a stratification by type

$$\chi(n, d)^{-1}(S_{\underline{n}'}^\times) = \bigsqcup_{\underline{n} \vdash n} \chi(n, d)^{-1}(S_{\underline{n}'}^\times) \cap M_{\underline{n}}^\circ(d).$$

Given the partition  $\underline{n} = (n_1, \dots, n_r)$ , the condition  $(\mathcal{E}, \phi) \in \chi(n, d)^{-1}(S_{\underline{n}'}^\times) \cap M_{\underline{n}}^\circ(d)$  means that the Higgs bundle  $(\mathcal{E}, \phi)$  splits as

$$(27) \quad (\mathcal{E}, \phi) = \bigoplus_{i=1}^r (\mathcal{E}_i, \phi_i),$$

where  $(\mathcal{E}_i, \phi_i)$  are stable Higgs bundles of rank  $n_i$  and degree  $d_i = dn_i/n$ . Further, the spectral curve  $C_a$  of  $(\mathcal{E}, \phi)$  is nodal and the union of the (possibly reducible) spectral curves  $C_{a_i}$  of  $(\mathcal{E}_i, \phi_i)$

$$(28) \quad C_a = \bigcup_{i=1}^r C_{a_i}.$$

A normal slice  $N_{\underline{n}} \subseteq A_a$  to  $S_{\underline{n}}$  through  $a$  parametrises spectral curves obtained by smoothing the nodes  $E(\Gamma_{\underline{n}}) := \bigcup_{i,j=1}^r C_{a_i} \cap C_{a_j}$  of  $C_a$ .

**Notation 6.2.** For a partition  $\underline{n}$  of  $n$  we will denote by  $\Gamma_{\underline{n}}$  the dual graph of any spectral curve given by a point of  $S_{\underline{n}}^\times$ , i.e.  $\Gamma_{\underline{n}}$  is the graph with vertices  $[r] := \{1, \dots, r\}$  corresponding to the irreducible components of the curve and  $n_i n_j (2g - 2)$  edges between the vertices  $i, j$ , corresponding to the intersection points of the components. By abuse of notation we continue to denote by  $E(\Gamma_{\underline{n}})$  the set of edges of  $\Gamma_{\underline{n}}$ .

**Proposition 6.3.** *The image of the map*

$$T_a N_{\underline{n}} \hookrightarrow T_a A_a \xrightarrow{df^\times} \text{Ext}^1(\Omega_{C_a}^1, \mathcal{O}_{C_a}) \twoheadrightarrow \bigoplus_{e \in E(\Gamma_{\underline{n}})} H^0(C_a, \mathcal{E}xt^1(\Omega_{C_a}^1, \mathcal{O}_{C_a})_e) \simeq \text{Spec } \mathbb{C}[T_e : e \in E(\Gamma_{\underline{n}})]$$

*is cut by the linear equations*

$$\text{Circ}(\mathcal{B}) = \langle \sum_{e \in E} a_{ie} T_e : i = 2, \dots, r \rangle,$$

*where  $A = (a_{ie})$  is the boundary map of a quiver underlying the graph  $\Gamma_{\underline{n}}$  (cf Section 4).*

*Proof.* Let  $\nu_{\underline{n}}: C_{\underline{n},a} := \bigsqcup_{i=1}^r C_{a_i} \rightarrow C_a$  be the partial normalization obtained by blowing-up  $C_a$  at  $E(\Gamma_{\underline{n}})$ . Applying the functor  $\mathcal{H}om(-, \omega_{C_a})$  to the short exact sequence

$$0 \rightarrow \mathcal{O}_{C_a} \rightarrow \nu_{\underline{n},*} \mathcal{O}_{C_{\underline{n},a}} \rightarrow \bigoplus_{e \in E(\Gamma_{\underline{n}})} \mathcal{O}_e \rightarrow 0,$$

we obtain the short exact sequence

$$0 \rightarrow N_{C_a/T^*C} \otimes \mathcal{I}_{E(\Gamma_{\underline{n}})} \rightarrow N_{C_a/T^*C} \rightarrow \mathbb{T}_{\underline{n}}^1 := \bigoplus_{e \in E(\Gamma_{\underline{n}})} \mathcal{E}xt^1(\mathcal{O}_e, \omega_{C_a}) \rightarrow 0,$$

where  $\mathcal{I}_{E(\Gamma_{\underline{n}})}$  is the product of the maximal ideals of the nodes  $E(\Gamma_{\underline{n}})$ , and  $\omega_{C_a}$  is identified with  $N_{C_a/T^*C}$  by adjunction. Taking global cohomology, we get the long exact sequence

$$(29) \quad \begin{aligned} 0 \rightarrow H^0(C_a, N_{C_a/T^*C} \otimes \mathcal{I}_{E(\Gamma_{\underline{n}})}) &\rightarrow T_a A_n = H^0(C_a, N_{C_a/T^*C}) \rightarrow \mathbb{T}_{\underline{n}}^1(C_a) \rightarrow \\ &\rightarrow H^1(C_a, N_{C_a/T^*C} \otimes \mathcal{I}_{E(\Gamma_{\underline{n}})}) \rightarrow H^1(C_a, N_{C_a/T^*C}) \rightarrow 0. \end{aligned}$$

By Serre duality we see

$$(30) \quad \begin{aligned} H^0(C_a, N_{C_a/T^*C} \otimes \mathcal{I}_{E(\Gamma_{\underline{n}})}) &\simeq H^0(C_a, \mathcal{H}om(\nu_{\underline{n},*} \mathcal{O}_{C_{\underline{n},a}}, \omega_{C_a})) \simeq H^1(C_a, \nu_{\underline{n},*} \mathcal{O}_{C_{\underline{n},a}})^\vee \\ &\simeq \bigoplus_{i=1}^r H^1(C_i, \mathcal{O}_{C_i})^\vee \simeq \bigoplus_{i=1}^r H^0(C_i, \omega_{C_i}) \simeq T_a S_{\underline{n}}. \end{aligned}$$

Combining (26), (29) and (30), the image of the map

$$df^\times : T_a N_{\underline{n}} \simeq T_a A_n / T_a S_{\underline{n}} \rightarrow \mathbb{T}_{\underline{n}}^1(C_a)$$

coincides with the kernel of  $\mathbb{T}_{\underline{n}}^1(C_a) \rightarrow \ker\{H^1(C_a, N_{C_a/T^*C} \otimes \mathcal{I}_{E(\Gamma_{\underline{n}})}) \rightarrow H^1(C_a, N_{C_a/T^*C})\}$ . Again via Serre duality, this map is isomorphic to

$$\mathbb{C}^{E(\Gamma_{\underline{n}})} \simeq \bigoplus_{e \in E(\Gamma_{\underline{n}})} H^0(\mathcal{O}_e)^\vee \rightarrow \ker\{H^0(C_a, \nu_{\underline{n},*} \mathcal{O}_{C_{\underline{n},a}})^\vee \rightarrow H^0(C_a, \mathcal{O}_{C_a})^\vee\} \simeq \mathbb{C}^{r-1},$$

which in turn can be identified with the boundary map of a quiver underlying the graph  $\Gamma_{\underline{n}}$ .  $\square$

## 7. SINGULARITIES OF UNIVERSAL COMPACTIFIED JACOBIANS

By the BNR correspondence a Higgs bundle  $(\mathcal{E}, \phi)$  corresponds to a rank 1 torsion free sheaf  $\mathcal{I}$  on the spectral curve  $C_a$ , where  $\mathcal{E}$  is isomorphic to  $\pi_{a,*} \mathcal{I}$  via the projection  $\pi_a : C_a \rightarrow C$ ; see [39, 2, 63]. Suppose that  $C_a$  is nodal, and let  $\Sigma$  be the set of the nodes where  $\mathcal{I}$  fails to be locally free. Define  $M_{\underline{n}}^\times(d)$  as the open subsets of Higgs bundles of  $M_{\underline{n}}^\circ(d)$  with nodal spectral curve.

**Proposition 7.1.** *Let  $\underline{n} = (n_1, \dots, n_r)$  be a partition of  $n$ . Then  $(\mathcal{E}, \phi)$  lies in  $M_{\underline{n}}^\times(d)$  if and only if there exists a partial normalization  $\nu_\Sigma : C_\Sigma \rightarrow C_a$  obtained by blowing-up the nodes  $\Sigma$  and a line bundle  $\mathcal{L}$  on  $C_\Sigma$  such that (1)  $C_\Sigma$  has exactly  $r$  connected components, and (2)  $\mathcal{E} \simeq \nu_{\Sigma,*} \mathcal{L}$ .*

*Proof.* By assumption,  $(\mathcal{E}, \phi)$  splits as  $(\mathcal{E}, \phi) = \bigoplus_{i=1}^r (\mathcal{E}_i, \phi_i)$ , where  $(\mathcal{E}_i, \phi_i)$  are stable Higgs bundles of rank  $n_i$  and degree  $d_i = dn_i/n$ . The spectral curve  $C_a$  of  $(\mathcal{E}, \phi)$  is the union of the spectral curves  $C_{a_i}$

$$C_{a_i} \xrightarrow{j_i} C_a = \bigcup_{i=1}^r C_{a_i} \xrightarrow{\pi_a} C.$$

By the BNR correspondence, there exist rank 1 torsion free sheaves  $\mathcal{I}$  and  $\mathcal{I}_i$  on  $C_a$  and  $C_{a_i}$  respectively such that  $\mathcal{E} = \pi_{a,*} \mathcal{I}$ ,  $\mathcal{E}_i = (\pi_a \circ j_i)_* \mathcal{I}_i$ , and

$$(31) \quad \mathcal{I} = \bigoplus_{i=1}^r j_{i,*} \mathcal{I}_i.$$

Note that  $\mathcal{I}$  fails to be locally free at the node  $C_{a_i} \cap C_{a_j}$ ,  $i \neq j$ . Therefore, the blow-up  $\nu_\Sigma: C_\Sigma \rightarrow C_a$  of  $C_a$  at  $\Sigma$  factors through the disjoint union of  $C_{a_i}$

$$(32) \quad \nu_\Sigma: C_\Sigma \rightarrow \bigsqcup_{i=1}^r C_{a_i} \rightarrow C_a.$$

Since  $C_a$  has only nodal singularities, there exists a line bundle  $\mathcal{L}$  on  $C_\Sigma$  such that  $\mathcal{I} = \nu_{\Sigma,*}\mathcal{L}$ ; see [29, Prop. 3.4]. By (32), the curve  $C_\Sigma$  has at least  $r$  components, and exactly  $r$  by the stability of  $(\mathcal{E}_i, \phi_i)$ . Otherwise, the decomposition (31) could be further refined, and so (27) could be refined too, which is a contradiction.  $\square$

**Proposition 7.2.** *Suppose that  $(\mathcal{E}, \phi)$  lies in  $M_{\underline{n}}^\times(d)$ . After removing all the loops, the dual graph  $\Gamma_{C_a}(\Sigma)$  of the curve obtained by smoothing the nodes not in  $\Sigma$  is the graph  $\Gamma_{\underline{n}}$ .*

*Proof.* As in the proof of Proposition 7.1, we denote the spectral curves of  $(\mathcal{E}, \phi)$  and of its stable summands by  $C_a$  and  $C_{a_i}$  respectively, and we write

$$C_a = \bigcup_{i=1}^r C_{a_i} = \bigcup_{j=1}^{r'} C_j \quad C_{a_i} = \bigcup_{j \in S_i} C_j,$$

where  $C_j$  are the irreducible components of  $C_a$ , and  $[r'] = \bigsqcup_{i=1}^r S_i$  is a partition of the set of  $r'$  elements. Clearly if  $C_j$  has degree  $n'_j$  over  $C$ , the spectral curves  $C_{a_i}$  has degree  $n_i = \sum_{j \in S_i} n'_j$ .

The graph  $\Gamma_{C_a}(\Sigma)$  is obtained from the dual graph  $\Gamma_{C_a}$  of the spectral curve  $C_a$  by contracting all the edges not in  $\Sigma$ .<sup>7</sup> Let  $q: \Gamma_{C_a} \rightarrow \Gamma_{C_a}(\Sigma)$  be the contraction map. If some edge between  $v_j$  and  $v_k$  is not contained in  $\Sigma$ , then  $q(v_j) = q(v_k)$ , and all edges in  $\Sigma$  are sent to loops in  $\Gamma_{C_a}(\Sigma)$ . The graph  $\Gamma_{C_a}(\Sigma)$  has as many vertices as connected components in  $\Gamma_{C_a} \setminus \Sigma \simeq \Gamma_{C_\Sigma}$ , i.e.  $r$  by Proposition 7.1. The edges between two distinct vertices  $w_i$  and  $w_{i'}$  in  $\Gamma_{C_a}(\Sigma)$  are exactly

$$\begin{aligned} \sum_{\substack{v_j \in q^{-1}(w_i), \\ v_k \in q^{-1}(w_{i'})}} (\# \text{ edges between } v_j \text{ and } v_k) &= \sum_{\substack{v_j \in q^{-1}(w_i), \\ v_k \in q^{-1}(w_{i'})}} n'_j n'_k (2g - 2) \\ &= \left( \sum_{j \in S_i} n'_j \right) \cdot \left( \sum_{k \in S_{i'}} n'_k \right) (2g - 2) = n_i n_{i'} (2g - 2). \end{aligned}$$

We conclude that the graphs  $\Gamma_{C_a}(\Sigma)$  and  $\Gamma_{\underline{n}}$  coincide after removing all loops in  $\Gamma_{C_a}(\Sigma)$ .  $\square$

**Remark 7.3.** The dual graph of the spectral curve may depend on the choice of  $(\mathcal{E}, \phi)$  in  $M_{\underline{n}}^\times(d)$ . Proposition 7.2 asserts that up to loops the dual graph  $\Gamma_{C_a}(\Sigma)$  does not!

The singularities of the universal compactified Jacobian  $\bar{J}_{d,B}$  are affine Lawrence varieties  $X(\Gamma_{\underline{n}}^\pm, 0)$ , as explained in [8] and [9]. The corresponding hypertoric quiver variety  $Y(\Gamma_{\underline{n}}, 0)$  provides the local model for the singularities of  $M(n, d)$ .

<sup>7</sup>By abuse of notation, we identify the set of nodes  $\Sigma$  with the corresponding set of edges in the dual graph  $\Gamma_{C_a}$ .

**Theorem 7.4** (Local model of  $\overline{\mathcal{J}}_{d,\mathcal{B}}$  and  $M(n, d)$ ). *Let  $a \in S_{\underline{n}}^\times$ , and  $A_a$  be the local multisection of  $f^\times : A_n^\times \rightarrow f^\times(A_n^\times) \subseteq \mathcal{B}$  defined in Proposition 5.5. Let  $(\mathcal{E}, \phi)$  be a Higgs bundle in  $M_{\underline{n}}^\times(d)$ , and  $\mathcal{I}$  be the corresponding rank 1 torsion free sheaf on the spectral curve  $C_a$ . Then there exists an analytic neighbourhood  $V$  of an étale chart of  $\overline{\mathcal{J}}_{d,\mathcal{B}}$  centered at  $(C_a, \mathcal{I})$  such that the restriction to  $V$  of the fibre product square in Proposition 5.5*

$$\begin{array}{ccc} M(n, d) \supset U & \hookrightarrow & V \subset \overline{\mathcal{J}}_{d,\mathcal{B}} \\ \chi(n, d) \downarrow & & \downarrow \pi \\ \chi(n, d)(U) & \xrightarrow{f^\times} & \pi(V), \end{array}$$

with  $U := V \cap \chi(n, d)^{-1}(A_a)$ , is locally isomorphic to

$$(33) \quad \begin{array}{ccc} Y(\Gamma_{\underline{n}}, 0) \times \mathbb{C}^{\dim M_{\underline{n}} - 1 - g} & \xrightarrow{(\iota_M, l_1)} & X(\Gamma_{\underline{n}}^\pm, 0) \times \mathbb{C}^{c(\underline{n}, g)} \\ (\chi_{\Gamma_{\underline{n}}}, l_2) \downarrow & & \downarrow (\pi_{\Gamma_{\underline{n}}}, l_4) \\ \mathbb{C}^{b_1(\Gamma_{\underline{n}})} \times \mathbb{C}^{d(\underline{n}, g)} & \xrightarrow{(\iota_A, l_3)} & \mathbb{C}^s \times \mathbb{C}^{3n^2(g-1) - s}, \end{array}$$

where the maps  $\chi$ ,  $\pi$ ,  $\iota_M$  and  $\iota_A$  are defined in (19), the maps  $l_i$  are linear,  $s := \#E(\Gamma_{\underline{n}})$ ,  $d(\underline{n}, g) := \frac{1}{2}(\dim M(n, d) - \dim Y(\Gamma_{\underline{n}}, 0)) - g - 1 = (n^2 - 1)(g - 1) - 1 - b_1(\Gamma_{\underline{n}})$  and  $c(\underline{n}, g) := \dim \overline{\mathcal{J}}_{d,\mathcal{B}} - \dim X(\Gamma_{\underline{n}}^\pm, 0) = 4n^2(g - 1) + 1 - b_1(\Gamma_{\underline{n}}) - s$ .

*Proof.* Let  $\Sigma$  be the set of the nodes where  $\mathcal{I}$  fails to be locally free, and  $\Gamma_{C_a}(\Sigma)$  be the dual graph of the curve obtained by smoothing the nodes not in  $\Sigma$ . By [8, Theorem A] there exists an analytic neighbourhoods  $V$  of  $(C_a, \mathcal{I})$  in  $\overline{\mathcal{J}}_{d,\mathcal{B}}$  such that

$$V \simeq_{\text{loc}} X(\Gamma_{C_a}(\Sigma)^\pm, 0) \times \mathbb{C}^{c(\underline{n}, \Sigma)},$$

with  $c(\underline{n}, \Sigma) := 4g(C_a) - 3 - b_1(\Gamma_{C_a}(\Sigma)) - \#E(\Gamma_{C_a}(\Sigma))$ . By [9, Lemma 4.3] and Proposition 7.2 we obtain that

$$V \simeq_{\text{loc}} X(\Gamma_{C_a}(\Sigma)^\pm, 0) \times \mathbb{C}^{c(\underline{n}, \Sigma)} \simeq X(\Gamma_{\underline{n}}^\pm, 0) \times \mathbb{C}^{c(\underline{n}, g)}.$$

The fact that the morphism  $\pi : V \subset \overline{\mathcal{J}}_{d,\mathcal{B}} \rightarrow \pi(V)$  can be identified with  $(\pi_{\Gamma_{\underline{n}}}, l_4)$  is explained in [9, §7.2]. The differential of the map  $f^\times$  is the Kodaira-Spencer map  $KS_a : T_a A_a \rightarrow H^1(C_a, \mathbb{T}_{C_a}^1)$ , and so the identification of the Hitchin fibration  $\chi(n, d)$  with the product of the map  $\chi_{\Gamma_{\underline{n}}} : Y(\Gamma_{\underline{n}}, 0) \rightarrow \mathbb{C}^{b_1(\Gamma_{\underline{n}})}$  with a linear map  $l_2$  follows from Proposition 6.3 and Definition 4.8.  $\square$

**Theorem 7.5.** *Let  $A_a \subset A_n$  as in Proposition 5.5. Then  $\text{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})|_{\chi(n, d)^{-1}(A_a)}[\dim M(n, d)]$  is a semi-simple perverse sheaf. In particular, we have*

$$\text{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})|_{\chi(n, d)^{-1}(A_a)}[\dim M(n, d)] \simeq \bigoplus_{\underline{n} \in \mathcal{P}_d} \text{IC}(M_{\underline{n}}(d), \mathcal{L}'_{\underline{n}})|_{\chi(n, d)^{-1}(A_a)}[\dim M(n, d)] \langle \text{codim } S_{\underline{n}} \rangle,$$

where  $\mathcal{L}'_{\underline{n}} := \mathcal{H}^{\text{codim } M_{\underline{n}}(d)}(\text{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q}))$  is a local system on  $M_{\underline{n}}(d)$  of rank  $h^{b_1(\Gamma_{\underline{n}})}(Y(\Gamma_{\underline{n}}, 0), \mathbb{Q})$ .

We anticipate that the local systems  $\mathcal{L}'_{\underline{n}}$  are pullback of the line bundles  $\mathcal{L}_{\underline{n}}$  in [17, Cor. 6.20]; see Proposition 7.6 and the proof of Theorem 1.3 below. In particular, their monodromy is known and induced by the automorphism of the graph  $\Gamma_{\underline{n}}$ ; see Remark 10.1.

The idea of the proof of Theorem 7.5 is to glue the local Hodge-to-singular correspondence (Theorem 4.18) given the local models in Theorem 7.4.

*Proof of Theorem 7.5.* By Theorem 4.18 and 7.4, the complex  $\mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})|_{\chi(n,d)^{-1}(A_a)}[\dim M(n,d)]$  is self-dual, i.e. it coincides with its own Verdier dual. Since  $\mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})$  is pure and self-dual,

$$\mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})|_{\chi(n,d)^{-1}(A_a)}[\dim M(n,d)] = i^* \mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q}) = i^! \mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})$$

is a pure perverse sheaf, thus semi-simple by [62, Prop. 1.7 and Prop. 1.9].  $\square$

**Proposition 7.6.** *Define  $M_{\underline{n}}^{\circ\circ}(d) := \chi(n,d)^{-1}(S_{\underline{n}}^{\times}) \cap M_{\underline{n}}(d)$ . The local system  $\mathcal{L}'_{\underline{n}}$  on  $M_{\underline{n}}^{\circ\circ}(d)$  descends to a local system  $\mathcal{L}_{\underline{n}}$  on  $S_{\underline{n}}^{\times}$*

$$\mathcal{L}'_{\underline{n}} = \chi(n,d)^* \mathcal{L}_{\underline{n}}.$$

*Proof.* Denote by  $F_b$  the fibre of  $\chi(n,d): M_{\underline{n}}^{\circ\circ}(d) \rightarrow S_{\underline{n}}$ . By Proposition 8.1, the restriction  $\mathrm{IC}(\overline{\mathcal{J}}_{d,\mathcal{B}}, \mathbb{Q})|_{F_b}$  is isomorphic to the complex

$$\mathbb{Q}_{F_b} \otimes \mathrm{IH}^i(X(\Gamma_{\underline{n}}^{\pm}, 0), \mathbb{Q})[-i] \simeq \mathbb{Q}_{F_b} \otimes H^i(Y(\Gamma_{\underline{n}}, 0), \mathbb{Q})[-i]$$

of constant sheaves with trivial differential. Hence,  $\mathcal{L}'_{\underline{n}}|_{F_b} \simeq \mathbb{Q}_{F_b} \otimes H^{b_1(\Gamma_{\underline{n}})}(Y(\Gamma_{\underline{n}}, 0), \mathbb{Q})$  is a component of the complex, so a trivial local system. Therefore,  $\mathcal{L}'_{\underline{n}}$  descends to a local system on  $S_{\underline{n}}^{\times}$ .  $\square$

## 8. TUBULAR NEIGHBOURHOOD OF THE JACOBIAN OF THE NORMALISATION OF A SPECTRAL CURVE

To complete the proof of Proposition 7.6, we must trivialize a tubular neighbourhood of the subvariety  $F_b$  in the universal compactified Jacobian. This is a global statement which asserts that the local trivializations in Theorem 7.4 glue along  $F_b$ .

Let  $f: \mathcal{C} \rightarrow B$  be a flat projective versal family of semistable curves and arithmetic genus  $g'$ , and  $\pi: \overline{\mathcal{J}}_{d,B} \rightarrow B$  be the relative compactified Jacobian with respect to the canonical polarization. For any  $b \in B$ , we denote by  $F_b$  the subvariety in  $\pi^{-1}(b)$  parametrising rank 1 torsion free sheaves on  $C_b := f^{-1}(b)$  of the shape  $\mathcal{I} = \nu_* M$  where  $M$  is a line bundle on the normalization  $\nu: C_b^{\nu} = \bigsqcup_{i=1}^r C_{b,i} \rightarrow C_b$ .

**Proposition 8.1.** *There exists a homeomorphism between a euclidean neighbourhood of  $F_b$  in  $\overline{\mathcal{J}}_{d,B}$  and the product*

$$(34) \quad F_b \times X(\Gamma(C_b)^{\pm}, 0) \times \mathbb{C}^{3g'-3-s},$$

where  $\Gamma(C_b)$  is the dual graph of  $C_b$  with  $s$  edges  $\Sigma$ .

**Remark 8.2.** Let  $B_\Gamma \subseteq B$  the locus of curve with the same dual graph as  $C_b$ , and let  $\mathcal{C}^\nu \rightarrow \mathcal{C} \times_B B_\Gamma$  the simultaneous normalization of  $C_b$ . Proposition 8.1 says in particular that a euclidean neighbourhood of  $F_b$  in the relative Picard scheme  $\text{Pic}^{d'}(\mathcal{C}^\nu) \subseteq \overline{\mathcal{J}}_{d,B}$  is homeomorphic to  $F_b \times \mathbb{C}^{3g'-3-s}$ , but this is not a formal or analytic isomorphism. Otherwise  $T\text{Pic}^{d'}(\mathcal{C}^\nu)|_{F_b} \simeq \mathcal{O}_{F_b}^{\oplus c'}$  with  $c' := \dim(F_b) + 3g' - 3 - s$ , and the boundary map  $H^0(N_{F_b/\text{Pic}^{d'}(\mathcal{C}^\nu)}) \rightarrow H^1(TF_b)$  of the short exact sequence

$$0 \rightarrow TF_b \rightarrow T\text{Pic}^{d'}(\mathcal{C}^\nu)|_{F_b} \rightarrow N_{F_b/\text{Pic}^{d'}(\mathcal{C}^\nu)} \rightarrow 0$$

would be trivial, i.e.  $\mathcal{C}^\nu$  is an isotrivial family, which is a contradiction. This suggests that Proposition 8.1 is false in the analytic category.

In order to show Proposition 8.1, we need to recall the GIT description of  $\overline{\mathcal{J}}_{d,B}$  and the deformation theory of the pair  $(C_b, \mathcal{I})$  as in [8, 9]. The relative compactified Jacobian  $\overline{\mathcal{J}}_{d,B}$  is a GIT-quotient of an open set  $Q^{ss}$  of a relative Quot scheme  $\text{Quot}(\mathcal{O}^{\oplus R})$  over the versal family  $f$ , as in [8, Section 2.3].

Note that  $Q^{ss}$  is smooth along the closed  $\text{SL}_R$ -orbits by [8, Lemma 6.4.(ii)] and by the smoothness of the miniversal deformation of the pair  $(C_b, \mathcal{I})$ ; see for instance [8, §3].

Let  $Z$  be the locus of closed orbits over  $F_b \subset \overline{\mathcal{J}}_{d,B} = Q^{ss} // \text{SL}_R$ . Note that  $F_b$  is isomorphic to the Jacobian  $\text{Pic}^{d'}(C_b^\nu)$  with  $d' := d + (n - n^2)(1 - g)$ , and  $Z$  is a fiber bundle over  $F_b$ .

Now let

$$(35) \quad \mathfrak{n}' : \mathcal{Q} = \text{Bl}_{Z \times 0}(Q^{ss} \times \mathbb{C}) \setminus Q^{ss} \times 0 \rightarrow \mathbb{C}$$

be the deformation of  $Q^{ss}$  to the normal cone of  $Z$ . The central fibre is the total space  $\text{Tot}(N_Z)$  of the normal bundle  $N_Z$  of  $Z$  in  $Q^{ss}$ , and the restriction of  $\mathfrak{n}'$  to  $\mathbb{C}^*$  is the trivial fibration  $Q^{ss} \times \mathbb{C}^* \rightarrow \mathbb{C}^*$ . Since  $Z$  is  $\text{SL}_R$ -invariant,  $\mathcal{Q}$  inherits a  $\text{SL}_R$ -action preserving the fibres of  $\mathfrak{n}'$ . The quotient

$$\mathfrak{n} : \mathcal{Q} // \text{SL}_R \rightarrow \mathbb{C}$$

is a deformation of  $\overline{\mathcal{J}}_{d,B}$  to  $\text{Tot}(N_Z) // \text{SL}_R$ . We can then reduce the proof of Proposition 8.1 to the following statements:

- $\text{Tot}(N_Z) // \text{SL}_R$  is homeomorphic to the product (34);
- $\text{Tot}(N_Z) // \text{SL}_R$  is homeomorphic to a euclidean neighbourhood of  $F_b$  in  $\overline{\mathcal{J}}_{d,B}$ .

We prove the statements respectively in Lemma 8.6 and Section 8.2.

**Remark 8.3.** In general, the deformation  $\mathfrak{n}$  is not the deformation of  $\overline{\mathcal{J}}_{d,B}$  to the normal cone of  $F_b$ . Indeed, a neighbourhood  $U$  of a point in  $F_b$  is locally analytically isomorphic to  $X(\Gamma(C_b)^\pm, 0) \times \mathbb{C}^{c'}$  with  $c' := 4g' - 3 - b_1(\Gamma) - s$  by [8, Theorem A]. Up to a smooth factor, the degeneration to the normal cone deforms  $U$  to the normal cone of  $X(\Gamma(C_b)^\pm, 0)$  at 0. The latter may fail to be isomorphic to  $X(\Gamma(C_b)^\pm, 0)$ : they are isomorphic only if the homogeneous polynomial cutting  $X(\Gamma(C_b)^\pm, 0)$  have

the same degree, equivalently if the number of edges between different vertices of  $\Gamma(C_b)^\pm$  is constant. On the other hand,  $\mathfrak{n}$  is locally trivial, so it deforms  $U$  to  $X(\Gamma(C_b)^\pm, 0) \times \mathbb{C}^{c'}$ .

**8.1. Deformation theory of rank 1 torsion free sheaves.** A slice of  $Q^{ss}$  normal to an  $\mathrm{SL}_R$ -orbit of the pair  $(C_b, \mathcal{I})$  is given by a miniversal deformation  $\mathrm{Def}(C_b, \mathcal{I})$  of the pair; see [8, §6].

The space  $\mathrm{Def}(C_b, \mathcal{I})$  is endowed with a natural action of the torus

$$\mathrm{Aut}(\mathcal{I}) \simeq H^0(C_b^\nu, \mathcal{O}_{C_b^\nu}^*) \simeq \bigoplus_{i=1}^r H^0(C_{b,i}, \mathcal{O}_{C_{b,i}}^*) \simeq \mathbb{T}^r \ni (\lambda_i)_{i=1}^r,$$

see [8, Def. 3.4]. Since  $\mathrm{Aut}(\mathcal{I})$  is reductive, the tangent space  $T\mathrm{Def}(C_b, \mathcal{I})$  splits into the product

$$T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{inv}} \oplus T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{var}},$$

where the invariant part  $T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{inv}}$  is fixed by the action of  $\mathrm{Aut}(\mathcal{I})$ , while the variant part  $T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{var}}$  is the unique  $\mathrm{Aut}(\mathcal{I})$ -invariant complement of  $T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{inv}}$ . In fact, these subspaces have a modular interpretation:

$$T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{inv}} \simeq T\mathrm{Def}^{lt}(C_b, \mathcal{I}) \quad T\mathrm{Def}(C_b, \mathcal{I})_{\mathrm{var}} \simeq \prod_{e \in \Sigma} T\mathrm{Def}(C_{b,e}, \mathcal{I}_e),$$

where  $\mathrm{Def}^{lt}(C_b, \mathcal{I})$  is the miniversal deformation of locally trivial deformations of  $(C_b, \mathcal{I})$ , and  $\mathrm{Def}(C_{b,e}, \mathcal{I}_e)$  is a miniversal deformation of the localised pair  $(C_{b,e}, \mathcal{I}_e)$ ; see [8, §3 and §5]. In particular, it follows from [8, §5] that  $\mathrm{Aut}(\mathcal{I})$  acts on  $T\mathrm{Def}(C_{b,e}, \mathcal{I}_e) \simeq \mathrm{Spec} k[z_e, w_e]$  as

$$(36) \quad z_e \mapsto \lambda_{s(e)} z_e \lambda_{t(e)}^{-1} \quad w_e \mapsto \lambda_{s(e)}^{-1} w_e \lambda_{t(e)}.$$

Now let  $q : Q^{ss} \rightarrow \bar{J}_{d,B}$  be the quotient map. Choose  $\mathcal{U}$  a universal rank 1 torsion-free sheaf over  $Q^{ss} \times_B \mathcal{C}$ , and take  $p : Q^{ss} \times_B \mathcal{C} \rightarrow Q^{ss}$  the natural projection.

**Lemma 8.4.** *Let  $N_Z$  be the normal bundle of  $Z$  in  $Q^{ss}$ . There exists a  $\mathbb{T}^r$ -equivariant splitting*

$$N_Z \simeq N_Z^{\mathbb{T}^r} \oplus R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z),$$

where  $N_Z^{\mathbb{T}^r}$  is the  $\mathbb{T}^r$ -invariant part of  $N_Z$ .

*Proof.* The normal space to  $Z$  in  $Q^{ss}$  at  $z \in Z$  with  $q(z) = (C_b, \mathcal{I})$  admits the  $\mathbb{T}^r$ -equivariant splitting

$$(37) \quad N_{Z,z} \simeq (T_z \mathrm{Def}^{lt}(C_b, \mathcal{I}) / q^* T_{(C_b, \mathcal{I})} F_b) \oplus \bigoplus_{e \in \Sigma} T_z \mathrm{Def}(C_{b,e}, \mathcal{I}_e),$$

and  $\mathbb{T}^r$  acts trivially on the first summand  $(T_z \mathrm{Def}^{lt}(C_b, \mathcal{I}) / q^* T_{(C_b, \mathcal{I})} F_b)$  and as (36) on the second summand. By [24, Lemma 7.13] and the local-to-global Ext spectral sequence,  $\bigoplus_{e \in \Sigma} T_z \mathrm{Def}(C_{b,e}, \mathcal{I}_e)$  can be identified with

$$H^0(C_b, \mathcal{E}xt_{C_b}^1(\mathcal{I}, \mathcal{I})) \simeq \bigoplus_{e \in \Sigma} H^0(C_b, \mathcal{E}xt^1(\mathcal{I}, \mathcal{I})_e) \simeq \mathbb{C}^s,$$

i.e. the fibre of the vector bundle  $R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)$ . □

**Lemma 8.5.**  $R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)$  is a trivial vector bundle on  $Z$ .

*Proof.* The group  $F_b \simeq \text{Pic}^{d'}(C_b^\nu)$  acts on  $Z$ . Its action lifts to the vector bundle  $R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)$  parallelising it, as we show below.

A point in  $Z$  represents a quotient  $[\mathcal{O}_{C_b}^{\otimes R} \rightarrow \nu_* M]$ , where  $M$  is a line bundle on  $C_b^\nu$ . Note that  $\mathcal{E}xt_{C_b}^1(\nu_* M, \nu_* M)$  is independent of the choice of a basis for  $\mathcal{O}_X^{\otimes R}$ , and its rank is independent of  $M$ . Hence,  $R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)$  is the pull-back of a vector bundles  $A$  on  $F_b$ .

The transitive action of  $\text{Pic}^{d'}(C_b^\nu)$  by translation induces identification of the fibers of  $A$ . Indeed, if  $\mathcal{I} := \nu_* M$  and  $\mathcal{I}' := \nu_*(M \otimes L)$  with  $L \in \text{Pic}^{d'}(C_b^\nu)$ , then  $\mathcal{I}$  and  $\mathcal{I}'$  are locally isomorphic, and so there exists an invertible sheaf  $L'$  on  $X$  such that  $\mathcal{I}' = \mathcal{I} \otimes L'$  with  $\nu^* L' = L$ ; see for instance [11, Lemma 2.2.2 or 3.1.2/3.1.3]. Hence, we have

$$\mathcal{E}xt_{C_b}^1(\nu_*(M \otimes L), \nu_*(M \otimes L)) = \mathcal{E}xt_{C_b}^1(\nu_* M \otimes L', \nu_* M \otimes L') = \mathcal{E}xt_{C_b}^1(\nu_* M, \nu_* M).$$

This gives a global trivialization of  $A$ . □

**Lemma 8.6.** Let  $N_{F_b/\overline{\mathcal{J}}_d(\Gamma)}$  be the normal bundle of  $F_b$  in  $\text{Pic}^{d'}(C^\nu) \subseteq \overline{\mathcal{J}}_{d,B}$  as constructed in Remark 8.2. Denote by  $\text{Tot}(N)$  the total space of a vector bundle  $N$ . Then

$$\text{Tot}(N_Z) // \text{SL}_R \simeq \text{Tot}(N_{F_b/\text{Pic}^{d'}(C^\nu)}) \times X(\Gamma(C_b)^\pm, 0).$$

*Proof.* By Lemma 8.4 we have

$$\text{Tot}(N_Z) // \text{SL}_R \simeq (\text{Tot}(N_{\mathbb{Z}}^{\mathbb{T}^r}) \times_Z \text{Tot}(R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)) // \mathbb{T}^{r-1}) // (\text{SL}_r / \mathbb{T}^{r-1}),$$

and Lemma 8.5 gives

$$\text{Tot}(R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)) \simeq Z \times \mathbb{C}^s.$$

The algebraic torus  $\mathbb{T}^{r-1} = \mathbb{T}^r / \mathbb{T}$  acts on  $\mathbb{C}^s$  as prescribed in (36), which coincides with the torus action defining  $X(\Gamma(C_b)^\pm, 0)$ . Therefore, we obtain

$$\begin{aligned} \text{Tot}(R^0 p_* \mathcal{E}xt_{C_b}^1(\mathcal{U}|_Z, \mathcal{U}|_Z)) // \text{SL}_r &\simeq ((Z \times \mathbb{C}^s) // \mathbb{T}^{r-1}) // (\text{SL}_r / \mathbb{T}^{r-1}) \\ &\simeq Z // (\text{SL}_r / \mathbb{T}^{r-1}) \times X(\Gamma(C_b)^\pm, 0) \simeq F_b \times X(\Gamma(C_b)^\pm, 0). \end{aligned}$$

Finally, the vector bundle  $N_{\mathbb{Z}}^{\mathbb{T}^r}$  descends to  $N_{F_b/\text{Pic}^{d'}(C^\nu)}$ , because  $\text{Pic}^{d'}(C^\nu)$  parametrises the locally trivial deformations of the pair  $(C_b, \mathcal{I})$ . □

**8.2. Stratified Ehresmann's theorem.** We use a stratified version of Ehresmann's theorem to prove that  $\mathfrak{n}^{-1}(0)$  is a model of a tubular neighbourhood of  $F_b$  in  $\overline{\mathcal{J}}_{d,B}$ .

**Lemma 8.7.** The degeneration  $\mathfrak{n}: \mathcal{Q} // \text{SL}_R \rightarrow \mathbb{C}$  is locally analytically trivial along  $F_b$ , i.e. for any  $x \in F_b \subset \mathfrak{n}^{-1}(0) \subset \mathcal{Q} // \text{SL}_R$  there exists a trivialising neighbourhood  $U \subset \mathcal{Q} // \text{SL}_R$  of  $x$ , and a local analytic isomorphism

$$(38) \quad \psi: U \rightarrow (\mathbb{C}^c \times X(\Gamma(C_b)^\pm, 0)) \times \mathbb{C},$$

with  $c' := 4g' - 3 - b_1(\Gamma) - s$ , such that  $\mathbf{n}: U \rightarrow \mathbb{C}$  corresponds to the third projection  $(X(\Gamma(C_b)^\pm, 0) \times \mathbb{C}^{c'}) \times \mathbb{C} \rightarrow \mathbb{C}$ .

*Proof.* For any  $z \in Z$ , the degeneration  $\mathbf{n}: Z \times \mathbb{C} \subseteq \mathcal{Q} \rightarrow \mathbb{C}$  admits a section  $\{z\} \times \mathbb{C}$ . By (37), an analytic neighbourhood of  $\{z\} \times \mathbb{C}$  is locally isomorphic to

$$(T_z O_z \times T_z \text{Def}^{lt}(C_b, \mathcal{I}) \times \bigoplus_{e \in \Sigma} T_z \text{Def}(C_{b,e}, \mathcal{I}_e)) \times \mathbb{C},$$

where  $O_z$  is the  $\text{SL}_R$ -orbit of  $z$ . By Luna's slice theorem, an analytic neighbourhood of  $\{[z]\} \times \mathbb{C}$  in  $\mathcal{Q} // \text{SL}_R$  is locally isomorphic to

$$\begin{aligned} & (T_z \text{Def}^{lt}(C_b, \mathcal{I}) \times \bigoplus_{e \in \Sigma} T_z \text{Def}(C_{b,e}, \mathcal{I}_e)) // \mathbb{T}^{r-1} \times \mathbb{C} \\ & \simeq (T_{(C_b, \mathcal{I})} \text{Pic}^{d'}(C^\nu) \times X(\Gamma(C_b)^\pm, 0)) \times \mathbb{C} \simeq (\mathbb{C}^{c'} \times X(\Gamma(C_b)^\pm, 0)) \times \mathbb{C}. \end{aligned}$$

□

*Proof of Proposition 8.1.* Let's first show that a relatively compact euclidean neighbourhood of  $F_b$  in  $\mathbf{n}^{-1}(0)$  is stratified diffeomorphic to a euclidean neighbourhood of  $F_b$  in  $\mathbf{n}^{-1}(0)$  in  $\mathbf{n}^{-1}(t) = \overline{\mathcal{J}}_{d,B}$ , with  $t \neq 0$ .

By Lemma 8.7 the degeneration  $\mathbf{n}: \mathcal{Q} // \text{SL}_R \rightarrow \mathbb{C}$  is locally trivial. This means that we can cover  $F_b \subset \mathbf{n}^{-1}(0)$  by trivialising open sets  $U_k$ , with  $k$  in the index set  $K$ , as in Lemma 8.7. Take now the vector field  $v_k := d\psi_k^{-1}(\partial_t)$  on  $U_k$ , where  $\partial_t$  is a non-vanishing vector field on  $\mathbb{C}$  and  $\psi_k$  is the stratified diffeomorphism (38). Consider a partition of unity  $\{\chi_k\}_{k \in K}$  subordinate to the open cover  $\{U_k\}_{k \in K}$ . Then the flow of the vector field  $v := \sum_{k \in K} \chi_k \cdot v_k$  defines the required stratified diffeomorphism.

We conclude now by noticing that the neighbourhood of  $F_b$  in  $\mathbf{n}^{-1}(0)$  can be chosen homeomorphic to the product

$$\text{Tot}(N_{F_b / \text{Pic}^{d'}(C^\nu)}) \times X(\Gamma(C_b)^\pm, 0) \underset{\text{homeo}}{\simeq} F_b \times \mathbb{C}^{3g'-3-s} \times X(\Gamma(C_b)^\pm, 0)$$

by Lemma 8.6. □

## 9. FULL SUPPORT THEOREM FOR UNIVERSAL COMPACTIFIED JACOBIANS

**Lemma 9.1** (Weak abelian fibration). *Let  $f: \mathcal{C} \rightarrow B$  be a flat projective versal family of semistable curves and arithmetic genus  $g'$ , and  $\pi^P: P := \text{Pic}^0(\mathcal{C}/B) \rightarrow B$  be the relative degree 0 Picard scheme parametrising line bundles on the fibres of  $f$  whose restrictions to each irreducible components are of degree 0. Let  $\pi: \overline{\mathcal{J}}_{d,B} \rightarrow B$  be the relative compactified Jacobian with respect to the canonical polarization. Then the triple  $(\overline{\mathcal{J}}_{d,B}, P, B)$  is a weak abelian fibration.*

*Proof.* We need to check condition (1), (2) and (3) in Definition 3.1. The relative Picard scheme  $\pi^P: P \rightarrow B$  has fibres of pure dimension  $g'$ , and  $\dim \overline{\mathcal{J}}_{d,B} = g' + \dim B$ ; see for instance Theorem 7.4. Hence Condition (1) holds. The proof of [22, Lem. 3.4.5] gives the affineness of the stabilisers, i.e.

Condition (2). The same argument of [58, §4.12] gives Condition (3); see also [15, Thm 3.3.1] where one can drop the assumption that  $\mathcal{C}$  is a family of curves embedded in a surface.  $\square$

**Proposition 9.2** (Whitney stratification of  $\overline{\mathcal{J}}_{d,B}$ ). *In the hypothesis of Lemma 9.1, let  $\mathcal{I}$  be a rank 1 torsion free sheaf on  $\mathcal{C}_b := f^{-1}(b)$  which fails to be locally free at the nodes  $\Sigma$  of  $\mathcal{C}_b$ . Set  $\Gamma_{\mathcal{C}_b}(\Sigma)$  to be the dual graph of any curve obtained from  $\mathcal{C}_b$  by smoothing the nodes not in  $\Sigma$ . Then there exists a complex Whitney stratification*

$$\overline{\mathcal{J}}_{d,B} = \bigsqcup_{\Gamma} \overline{\mathcal{J}}_d^{\circ}(\Gamma),$$

where  $\overline{\mathcal{J}}_d^{\circ}(\Gamma)$  is the locus of rank 1 torsion free sheaves in  $\overline{\mathcal{J}}_{d,B}$  whose  $\Gamma_{\mathcal{C}_b}(\Sigma)$  is isomorphic to  $\Gamma$ , and we have

$$(39) \quad \text{codim}_{\overline{\mathcal{J}}_{d,B}} \overline{\mathcal{J}}_d^{\circ}(\Gamma) \geq 2 \text{codim}_{\pi^{-1}(t)}(\overline{\mathcal{J}}_d^{\circ}(\Gamma) \cap \pi^{-1}(t)).$$

*Proof.* The first statement is a reformulation of [8, Thm A]. For the inequality (39), observe that if  $\Gamma$  has  $r$  vertices and  $s$  edges, then [8, Thm A] gives

$$\begin{aligned} \text{codim}_{\overline{\mathcal{J}}_{d,B}} \overline{\mathcal{J}}_d^{\circ}(\Gamma) &= \dim X(\Gamma^{\pm}, 0) = 2s - r + 1, \\ \text{codim}_{\pi^{-1}(t)}(\overline{\mathcal{J}}_d^{\circ}(\Gamma) \cap \pi^{-1}(t)) &= s - r + 1. \end{aligned}$$

$\square$

**Lemma 9.3.** *In the hypothesis of Lemma 9.1, we have*

$$\tau_{>2R}(R\pi_* \text{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q})) = 0, \quad R := \dim \overline{\mathcal{J}}_{d,B} - \dim B = g'.$$

*Proof.* It follows from Proposition 3.3 and Proposition 9.2.  $\square$

**Lemma 9.4.** *In the hypothesis of Lemma 9.1, any support  $Z$  of  $R\pi_* \text{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q})$  satisfies*

$$\text{codim } Z = \delta_Z$$

*Proof.* The inequality  $\text{codim } Z \geq \delta_Z$  appears for instance in [56, Fact 2.4]. The reverse inequality  $\text{codim } Z \leq \delta_Z$  follows from Lemma 9.1, Lemma 9.3, and [48, Theorem 1.8] or Theorem 3.2.  $\square$

Let  $\pi: \overline{\mathcal{J}}_{d,B} \rightarrow B$  be the relative compactified Jacobian with respect to the canonical polarization of a versal deformation of semistable curves of arithmetic genus  $g'$ . We define the invariant

$$(40) \quad \Psi_{g'} := \dim \overline{\mathcal{J}}_{d,B} - 2 \dim B = 4g' - 3 - 2(3g' - 3) = 3 - 2g'.$$

**Theorem 9.5** (Full support for Jacobians of versal families). *Let  $f: \mathcal{C} \rightarrow B$  be a flat projective versal family of semistable curves and arithmetic genus  $g'$ , and  $U \subset B$  the open subscheme over which the morphism  $f$  is smooth. Let  $\pi: \bar{\mathcal{J}}_{d,B} \rightarrow B$  be the relative compactified Jacobian with respect to the canonical polarization. Then the complex  $R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q})$  has full support on  $B$ , i.e.*

$$(41) \quad R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q}) \simeq \bigoplus_{l=0}^{2g} \mathrm{IC}(B, \Lambda^l(R^1\pi_*\mathbb{Q})|_U)[-l].$$

In particular, for any  $d, d' \in \mathbb{Z}$  there is an isomorphism of mixed Hodge modules

$$(42) \quad R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q}) \simeq R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d',B}, \mathbb{Q}).$$

*Proof.* The argument of [48, §3] can be adapted to work in the present setting. The base  $B$  admits a stratification by type

$$B = \bigsqcup B_{\underline{g}'},$$

which extends (4). For any  $r \geq 1$ , and for any  $r$ -tuples of positive integers  $\underline{g}' = (g'_i)$ , the closed points  $b \in B$  in  $B_{\underline{g}'}$  corresponds to the fibers of  $f$  of the form  $\mathcal{C}_b = \bigcup_i \mathcal{C}_i$ , where  $\mathcal{C}_i$  are irreducible curves of genus  $g'_i = g(\mathcal{C}_i)$ .

Assume that the irreducible subvariety  $Z \subseteq B$  is a support of  $R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q})$ , and that the generic point of  $Z$  is contained in  $B_{\underline{g}'}$ . As in [48, Prop. 4.4], the inequality  $\mathrm{codim} Z \geq \delta_Z$  can be improved to

$$(43) \quad \Psi_{g'} + \mathrm{codim} Z \geq \sum_{i=1}^r \Psi_{g'_i} + \delta_Z.$$

Combining (43), (40) and Lemma 9.4, we obtain

$$3(r-1) + 2(g - \sum_{i=1}^r g_i) = 0.$$

The only possibility is that  $r = 1$ . Hence,  $Z = B$ , which implies (41) and (42).  $\square$

**Remark 9.6.** The proof of Theorem 9.5 does not work for the Hitchin fibration. In this case, the invariant  $\Psi = \dim M(n, d) - 2 \dim A_n$  vanishes, and the inequality (43) is inconclusive.

## 10. PROOF OF THE MAIN THEOREMS

*Proof of Theorem 1.1 (Hodge-to-singular correspondence).* In view of the decomposition (17), it suffices to show that for any partition  $\underline{n}$  of  $n$  the isomorphism (5) holds on an open euclidean neighbourhood of a general  $a \in S_{\underline{n}}^\times$  containing a Zariski-dense open set of  $S_{\underline{n}}^\times$ . By Remark 5.6, it is actually enough to show the isomorphism on the subsets  $A_a$  constructed in the proof of Proposition 5.5.

By Proposition 5.5, proper base change gives

$$R\chi(n, d)_*(\mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q})|_{\chi(n,d)^{-1}(A_a)}) \simeq (R\pi_* \mathrm{IC}(\bar{\mathcal{J}}_{d,B}, \mathbb{Q}))|_{A_a}.$$

For any étale chart  $B \rightarrow \mathcal{B}$ , Theorem 9.5 says that  $R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{e,B}, \mathbb{Q})$  and  $R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q})$  have full support, which implies that

$$(R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{e,B}, \mathbb{Q}))|_{A_a} \simeq (R\pi_* \mathrm{IC}(\overline{\mathcal{J}}_{d,B}, \mathbb{Q}))|_{A_a}.$$

Since  $\chi(n, e)^{-1}(A_a)$  is regularly embedded in  $\overline{\mathcal{J}}_{e,B}$ , we have

$$\mathrm{IC}(\overline{\mathcal{J}}_{e,B}, \mathbb{Q})|_{\chi(n,e)^{-1}(A_a)} \simeq \mathbb{Q}_{\chi(n,e)^{-1}(A_a)}$$

Together with Theorem 7.5 and Proposition 7.6, we obtain Theorem 1.1.  $\square$

*Proof of Theorem 1.3.* By Theorem 7.5, the descent of  $\mathcal{L}'_{\underline{n}}$  on  $S_{\underline{n}}^\times$  in Proposition 7.6 is a sublocal system of the sheaf  $\mathcal{L}_{\underline{n}}(e)$  in the Ngô strings of Theorem 3.11. For dimensional reason, they must coincide by [17, Thm 6.11] and Proposition 4.11.(8).  $\square$

*Proof of Theorem 1.4.* Theorem 1.1 for  $d = 0$  yields

$$(44) \quad R\chi(n, e)_* \mathbb{Q}_{M(n,e)}|_{A_n^{\mathrm{red}}} \simeq \bigoplus_{\underline{n} \vdash n} R\chi(n, 0)_* \mathrm{IC}(M_{\underline{n}}(0), \chi(n, 0)^* \mathcal{L}_{\underline{n}})[2(\dim S_n - c(g, n))]|_{A_n^{\mathrm{red}}}.$$

On the open set of  $S_{\underline{n}}$  over which the Hitchin map  $\chi(n, 0): M_{\underline{n}}(0) \rightarrow S_{\underline{n}}$  is smooth, there are isomorphisms

$$R\chi(n, 0)_* \mathrm{IC}(M_{\underline{n}}(0), \chi(n, 0)^* \mathcal{L}_{\underline{n}}) \simeq \bigoplus_{l=0}^{2 \dim S_{\underline{n}}} \Lambda_{\underline{n}}^l \otimes \mathcal{L}_{\underline{n}}[-l] \simeq \mathcal{S}_{\underline{n}}.$$

Therefore, by the decomposition theorem [3, 62], the string  $\mathcal{S}_{\underline{n}}$  is a direct summand of

$$R\chi(n, 0)_* \mathrm{IC}(M_{\underline{n}}(0), \chi(n, 0)^* \mathcal{L}_{\underline{n}})$$

on the whole  $S_{\underline{n}}$ . Now Theorem 1.3 gives

$$\bigoplus_{\underline{n} \vdash n} \mathcal{S}_{\underline{n}}|_{A_n^{\mathrm{red}}} \langle \mathrm{codim} S_{\underline{n}} \rangle \simeq R\chi(n, e)_* \mathbb{Q}_{M(n,e)}|_{A_n^{\mathrm{red}}} \simeq \mathrm{RHS} \text{ of } (44) \supseteq \bigoplus_{\underline{n} \vdash n} \mathcal{S}_{\underline{n}}|_{A_n^{\mathrm{red}}} \langle \mathrm{codim} S_{\underline{n}} \rangle.$$

This implies that the complexes  $R\chi(n, 0)_* \mathrm{IC}(M_{\underline{n}}(0), \chi(n, 0)^* \mathcal{L}_{\underline{n}})$  have full support. For  $\underline{n} = \{n\}$ , this gives that  $R\chi(n, 0)_* \mathrm{IC}(M(n, 0), \mathbb{Q})$  has full support.  $\square$

*Proof of Theorem 1.7.* By Theorem 1.3 and Theorem 1.4, it suffices to check that no summand of  $R\chi(2, 1)_* \mathbb{Q}_{M(2,1)}$  and  $R\chi(2, 0)_* \mathrm{IC}(M(2, 0), \mathbb{Q})$  is supported over  $A_2 \setminus A_2^{\mathrm{red}} = \{0\}$ . This is indeed the case by Proposition 1.6 (Corollary 3.9).  $\square$

**Remark 10.1.** Let  $\underline{n} = (n_1, \dots, n_r) = 1^{\alpha_1} \dots n^{\alpha_n}$  be a partition of  $n$ , where  $\alpha_i$  is the number of elements in  $n$  equal to  $i$ . Denote by  $\mathfrak{S}_{\underline{n}} := \prod_{i=1}^n \mathfrak{S}_{\alpha_i}$  the subgroup of the symmetric group  $\mathfrak{S}_r$  stabilizing  $\underline{n}$ . Then the horizontal arrows of the following commutative square

$$\begin{array}{ccc} \prod_{i=1}^r M(n_i, d_i) & \longrightarrow & M_{\underline{n}}(d) & & (\mathcal{E}_i, \phi_i)_{i=1}^r & \longmapsto & (\bigoplus_{i=1}^r \mathcal{E}_i, \bigoplus_{i=1}^r \phi_i) \\ \prod_{i=1}^r \chi(n, d) \downarrow & & \downarrow \chi(n, d) & & \downarrow & & \downarrow \\ \prod_{i=1}^r A_{n_i} & \xrightarrow{\mathrm{mult}_{\underline{n}}} & S_{\underline{n}} & & (\mathrm{char}(\phi_i))_{i=1}^r & \longmapsto & \prod_{i=1}^r \mathrm{char}(\phi_i) \end{array}$$

are quotient maps by the action of  $\mathfrak{S}_{\underline{n}}$ . By [17, Cor. 6.20], the monodromy of the local systems  $\mathcal{L}_{\underline{n}}$  is given by the restriction to the subgroup  $\mathfrak{S}_{\underline{n}}$  of the representation of  $\mathfrak{S}_r$  induced by a primitive character of a maximal cyclic subgroup. In particular,  $\text{mult}_{\underline{n}}^* \mathcal{L}_{\underline{n}}$  is a trivial local system. Hence,  $\mathfrak{S}_{\underline{n}}$  acts on the sheaf  $\boxtimes_{i=1}^r R\chi(n_i, d_i)_* \text{IC}(M(n_i, d_i), \mathbb{Q}) \otimes \mathbb{Q}^{\text{rank } \mathcal{L}_{\underline{n}}}$  via the  $\mathfrak{S}_{\underline{n}}$ -representation  $\mathcal{L}_{\underline{n}}$ . This means that the summand of the RHS of (5) are the  $\mathfrak{S}_{\underline{n}}$ -invariant part of

$$(45) \quad \boxtimes_{i=1}^r R\chi(n_i, d_i)_* \text{IC}(M(n_i, d_i), \mathbb{Q}) \otimes \mathbb{Q}^{\text{rank } \mathcal{L}_{\underline{n}}} \langle \text{codim } S_{\underline{n}} \rangle.$$

In particular, the summands of the RHS of (6) are  $\mathfrak{S}_{\underline{n}}$ -subrepresentations of

$$\bigotimes_{i=1}^r IH^{k-2 \text{codim } S_n}(M(n_i, d_i)^{\text{red}}, \mathbb{Q}).$$

*Proof of Corollary 1.9.* The summation indices and the summands in (5) depends only on  $\text{gcd}(n, d)$ .  $\square$

**10.1. Dependence of  $IH^*(M(n, d), \mathbb{Q})$  on degree.** The description of the summands of the decomposition theorem for the Hitchin fibration in the intermediate case,  $\text{gcd}(n, d) \neq 0$  or  $\neq 1$ , is more subtle. It is not clear if one can get a close formula for them, but certainly (5) offers a recursive method. Combining Theorem 1.1 and Theorem 1.4, one obtain the isomorphism in  $D^b M H M_{\text{alg}}(A_n^{\text{red}})$

$$(46) \quad R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q}) \oplus \bigoplus_{\underline{n} \in \mathcal{P}_d, \underline{n} \neq \{n\}} R\chi(n, d)_* \text{IC}(M_{\underline{n}}(d), \chi(n, d)^* \mathcal{L}_{\underline{n}}) \langle \text{codim } S_{\underline{n}} \rangle \simeq \bigoplus_{\underline{n} = \{n_i\} \dashv n} \mathcal{S}_{\underline{n}}.$$

Recursively from this identity, one can determine the Ngô strings for  $R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q})$ , and the subvarieties  $S_{\underline{n}}$  which are actual support of  $R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q})$ . We refer to [52] for the explicit computation of the ranks of the Ngô strings. Here we limit to illustrate the first non-trivial example  $n = 4$ . More care is needed to determine explicitly the monodromy of the associated local systems.

**Example 10.2** ( $n = 4$ ). We determine the rank of the local systems  $\mathcal{L}_{\underline{n}}(d)$  in Theorem 3.11 for  $n = 4$ . To this end, observe that the Hodge-to-singular correspondence gives

$$(47) \quad R\chi(2, 4)_* \text{IC}(M(2, 4), \mathbb{Q}) \oplus R\chi(2, 4)_* \text{IC}(M_{\{2,2\}}(2), \chi(4, 2)^* \mathcal{L}_{\{2,2\}}) \langle 8g - 9 \rangle \simeq \bigoplus_{\underline{n} = \{n_i\} \dashv 4} \mathcal{S}_{\underline{n}}.$$

Consider the commutative square

$$\begin{array}{ccc} M(2, 1)^2 & \longrightarrow & M_{\{2,2\}}(2) \\ \chi(2,1)^2 \downarrow & & \downarrow \chi(4,2) \\ A_2^2 & \xrightarrow{\text{mult}_{\{2,2\}}} & S_{\{2,2\}}. \end{array}$$

Set  $\underline{n} = \{2, 2\}, \{2, 1, 1\}$  or  $\{1, 1, 1, 1\}$ . The map  $\text{mult}_{\{2,2\}}$  is étale at the general points of  $S_{\underline{n}}$ , so  $R\chi(2, 4)_* \text{IC}(M_{\{2,2\}}(2), \chi(4, 2)^* \mathcal{L}_{\{2,2\}})$  is locally isomorphic to  $R\chi(2, 1)_*^2 \text{IC}(M(2, 1)^2, \mathbb{Q}) \otimes \mathbb{Q}^{\text{rank } \mathcal{L}_{\{2,2\}}}$

at those points. We omit the last tensorisation since  $\text{rank } \mathcal{L}_{\{2,2\}} = 1$ . Indeed, recall that  $\text{rank } \mathcal{L}_{\underline{n}} = (|\underline{n}| - 1)!$ ; see [17, cor. 6.20.]. By the Künneth formula and Theorem 1.7, we have

$$R\chi(2, 1)_*^2 \text{IC}(M(2, 1)^2, \mathbb{Q}) \simeq (\mathcal{S}_{\{2\}} \oplus \mathcal{S}_{\{1,1\}}) \boxtimes (\mathcal{S}_{\{2\}} \oplus \mathcal{S}_{\{1,1\}}),$$

so all the Ngô strings of  $R\chi(2, 1)_*^2 \text{IC}(M(2, 1)^2, \mathbb{Q})$  start with a local system of rank one. By (47), this implies that  $\text{rank } \mathcal{L}_{\underline{n}}(2) = \text{rank } \mathcal{L}_{\underline{n}}(1) - 1$ . In particular,  $\mathcal{S}_{\{2,2\}}$  is not a support of  $R\chi(2, 4)_* \text{IC}(M(2, 4), \mathbb{Q})$ . We summarise the results in the following table.

$\text{gcd}(n, d)$	{4}	{3,1}	{2,2}	{2,1,1}	{1,1,1,1}
0	1	0	0	0	0
1	1	1	1	2	6
2	1	1	0	1	5

TABLE 1. The entries of the table are the rank of the local systems  $\mathcal{L}_{\underline{n}}(d)$  for  $n = 4$ . Theorem 3.11 says that the complex  $R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q})$  is a direct sum of Ngô strings

$$\mathcal{S}(\mathcal{L}_{\underline{n}}(d)) := \bigoplus_{l=0}^{2 \text{codim } S_{\underline{n}}} \text{IC}(S_{\underline{n}}, \Lambda_{\underline{n}}^l \otimes \mathcal{L}_{\underline{n}}(d))[-l] \langle \text{codim } S_{\underline{n}} \rangle.$$

By Remark 10.1, the summands  $R\chi(n, d)_* \text{IC}(M_{\underline{n}}(d), \chi(n, d)^* \mathcal{L}_{\underline{n}})$  of the LHS of (46) are the  $\mathfrak{S}_{\underline{n}}$ -invariant part of

$$\bigotimes_{i=1}^r R\chi(n_i, d_i)_* \text{IC}(M(n_i, d_i), \mathbb{Q}) \otimes \mathbb{Q}^{\text{rank } \mathcal{L}_{\underline{n}}},$$

where  $n_i < n$  and  $d_i < d$ . Therefore, using (46), one could rewrite the local system  $\mathcal{L}_{\underline{n}}(d)$  as a combinations of direct sums, linear quotients and  $\mathfrak{S}_{\underline{n}'}$ -quotients of  $\mathcal{L}_{\underline{n}'}(d')$ ,  $\mathcal{L}_{\underline{n}}$  and  $\mathcal{L}_{\underline{n}'}$ , with  $n' < n$  and  $d' < d$ . Recursively, we could remove the dependence of  $\mathcal{L}_{\underline{n}}(d)$  on  $\mathcal{L}_{\underline{n}'}(d')$ , and write  $\mathcal{L}_{\underline{n}}(d)$  as depending only on  $\mathcal{L}_{\underline{k}}$  with  $k \leq n$ . For any partition  $\underline{n} = (n_1, \dots, n_r)$ , Remark 10.1 says that the Ngô string associated to  $\mathcal{L}_{\underline{n}}$ , namely

$$\mathcal{S}_{\underline{n}} := \bigoplus_{l=0}^{2 \text{dim } S_{\underline{n}}} \text{IC}(S_{\underline{n}}, \Lambda_{\underline{n}}^l \otimes \mathcal{L}_{\underline{n}})[-l] \langle \text{codim } S_{\underline{n}} \rangle,$$

is the  $\mathfrak{S}_{\underline{n}}$ -invariant part of  $(\bigotimes_{i=1}^r \mathcal{S}_{n_i})^{\text{rank } \mathcal{L}_{\underline{n}}}$ . This means that  $R\chi(n, d)_* \text{IC}(M(n, d), \mathbb{Q})$  can be expressed in terms of the Ngô strings  $\mathcal{S}_k \simeq R\chi(k, 0)_* \text{IC}(M(k, 0), \mathbb{Q})$  with  $k \leq n$ . In other words, one obtain a recursive (though highly nontrivial) formula for  $IH^*(M^{\text{red}}(n, d), \mathbb{Q})$  in terms of  $IH^*(M^{\text{red}}(k, 0), \mathbb{Q}) = H^*(\mathcal{S}_k)$  with  $k \leq n$ .

## 10.2. Some applications of the Hodge-to-singular correspondence.

*Proof of Proposition 1.10.* Taking global cohomology in (17), we have

$$IH^{\leq k}(M(n, d)) \simeq IH^{\leq k}(M(n, 0))$$

with  $k = \min\{2(\text{dim } \mathcal{A}_n - \text{dim } \mathcal{A}_n^{\text{red}}), \text{dim } M_{\underline{n}} : \underline{n} \neq \{n\}\}$ . By [4, Lem. 2.2], we have

$$(48) \quad k = 2(n^2(g-1) + 1) - 2 \max \left\{ r + (g-1) \sum_{i=1}^r n_i^2 \left| m_i, n_i, \sum_{i=1}^r m_i r_i = n \right. \right\}$$

$$= 4(g-1)(n-1) - 2 = O(n).$$

This gives the independence of the limits (9) by the degree. The finiteness follows from the generation of the cohomology of  $M(n, e)$  with  $\gcd(e, n) = 1$  by finitely many tautological classes in [46, Thm 7].  $\square$

Recall that the perverse filtration on  $IH^*(M(n, d))$  associated to the Hitchin map  $\chi(n, d)$  is given by

$$(49) \quad P_j IH^k(M(n, d), \mathbb{Q}) = \ker\{IH^k(M(n, d), \mathbb{Q}) \rightarrow IH^k(\chi(n, d)^{-1}(\Lambda^{k-j-1}), \mathbb{Q})\},$$

where  $\Lambda^s$  is a general  $s$ -dimensional affine hyperplane of  $A_n$ ; see [21, Thm 4.1.1]. By (48),  $\Lambda^s$  does not intersect the supports of the RHS of (5) which are properly contained in  $A_n$ , provided that  $s < 2(g-1)(n-1) - 1$ . As a result, we obtain the following proposition.

**Proposition 10.3.** *The graded piece of the perverse filtration*

$$\mathrm{Gr}_j^P IH^k(M(n, d), \mathbb{Q}) := P_j IH^k(M(n, d), \mathbb{Q}) / P_{j-1} IH^k(M(n, d), \mathbb{Q})$$

are independent of the degree  $d$  for  $j > k - 2(g-1)(n-1) + 1$ .

*Proof of Proposition 1.11.* By (49) we have

$$\mathrm{Gr}_k^P IH^k(M(n, d), \mathbb{Q}) \simeq \mathrm{Im}\{IH^k(M(n, d), \mathbb{Q}) \rightarrow H^k(M_a, \mathbb{Q})\}.$$

Let  $e$  be an integer coprime with  $n$ . Since

$$\mathrm{Gr}_k^P IH^k(M(n, d), \mathbb{Q}) \simeq \mathrm{Gr}_k^P H^k(M(n, e), \mathbb{Q})$$

by Proposition 10.3, it suffice to determine the latter. Let  $(\mathbb{E}, \Phi)$  be the universal family of Higgs bundles on  $C \times M(n, e)$  normalised such that

$$\mathrm{ch}_1(\mathbb{E})|_{p \times M(n, e)} = 0 \in H^2(M(n, e), \mathbb{Q}), \quad \mathrm{ch}_1(\mathbb{E})|_{C \times q} = 0 \in H^2(C, \mathbb{Q}).$$

For any  $\gamma \in H^j(C, \mathbb{Q})$ , the tautological class  $c(\gamma, k)$  is the integral

$$\int_{\gamma} \mathrm{ch}^k(\mathbb{E}) \in H^{j+2k-2}(M(n, d), \mathbb{Q}).$$

By [46, Thm 7], the cohomology  $H^k(M(n, e), \mathbb{Q})$  is generated by tautological classes as  $\mathbb{Q}$ -algebra. The proof of [16, Cor. 5.1.3] works in arbitrary rank, and shows that the only tautological classes that do not vanish along  $M_a$  are  $c(\gamma, 1)$  with  $\gamma \in H^1(C, \mathbb{Q})$  and the  $\chi(n, e)$ -relative ample class  $\alpha := c([c], 2)$ , where  $[c]$  generates  $H^2(C, \mathbb{Q})$ . We conclude that

$$(50) \quad \mathrm{Gr}_*^P H^*(M(n, e), \mathbb{Q}) = \langle c(\gamma, 1)|_{M_a}, \alpha|_{M_a} \rangle = H^\bullet(C, \mathbb{Q}) \otimes \mathbb{Q}[\alpha|_{M_a}] / (\alpha|_{M_a}^{\dim M(n, d)+1}).$$

The result for  $\check{M}(n, d)$  is an immediate corollary of (50) since

$$\mathrm{Gr}_*^P H^*(M(n, d), \mathbb{Q}) \simeq \mathrm{Gr}_*^P H^*(\check{M}(n, d), \mathbb{Q}) \otimes H^\bullet(C, \mathbb{Q});$$

see for instance [16, §4.4].  $\square$

**10.3. Mukai moduli spaces.** Let  $S$  be a K3 surface. Given an effective Mukai vector<sup>8</sup>  $v \in H_{\text{alg}}^*(S, \mathbb{Z})$ , we denote by  $M(S, v)$  the moduli space of Gieseker semistable sheaves on  $S$  with Mukai vector  $v$  with respect to a polarization; see [66, §1]. Up to deformation, we can suppose that  $S$  contains an ample curve  $C$ , and  $v = (0, nC, -nC^2/2)$ ; see [60, Thm 1.17]. Taking (Fitting) supports defines a Lagrangian fibration

$$M(S, v) \rightarrow |nC|, \quad \mathcal{F} \mapsto \text{Supp}(\mathcal{F}),$$

whose fibers are compactified Jacobians. The Hodge-to-singular correspondence extends to this context with no change, so does Proposition 1.10. Note that the stabilization of the cohomology of  $M(S, v)$  ([12, Conjecture 1.1]) is known in the literature if  $v$  is primitive; see [12, p. 4] and references therein. We can now drop the primitivity assumption or the genericity of the polarization.

**Corollary 10.4.** *Let  $v^2 = mw^2$  and  $w^2 = 2k$ , with  $v, w \in H_{\text{alg}}^*(S, \mathbb{Z})$  and  $m, k \in \mathbb{Z}$ . The intersection Betti and Hodge numbers of  $M(S, v)$  stabilize to the stable Betti and Hodge numbers of the Hilbert scheme of  $(k+1)$ -points as  $k$  tends to infinity, i.e. the limits  $\lim_{k \rightarrow \infty} \dim IH^*(M(S, v))^{p,q}$  are finite and independent of  $m$ .*

## REFERENCES

- [1] D. Baraglia. Monodromy of the  $SL(n)$  and  $GL(n)$  Hitchin fibrations. *Math. Ann.*, 370(3-4):1681–1716, 2018.
- [2] A. Beauville, M.S. Narasimhan, and S. Ramanan. Spectral curves and the generalised theta divisor. *J. Reine Angew. Math.*, 398:169–179, 1989.
- [3] A.A. Beilinson, J.N. Bernstein, and P. Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Astérisque*, pages 5–171. Soc. Math. France, Paris, 1982.
- [4] G. Bellamy and T. Schedler. Symplectic resolutions of character varieties. *arXiv:1909.12545*, 2019.
- [5] R. Bielawski and A.S. Dancer. The geometry and topology of toric hyperkähler manifolds. *Comm. Anal. Geom.*, 8(4):727–760, 2000.
- [6] A. Björner. The homology and shellability of matroids and geometric lattices. In *Matroid applications*, volume 40 of *Encyclopedia Math. Appl.*, pages 226–283. Cambridge Univ. Press, Cambridge, 1992.
- [7] L. Caporaso. A compactification of the universal Picard variety over the moduli space of stable curves. *J. Amer. Math. Soc.*, 7(3):589–660, 1994.
- [8] S. Casalaina-Martin, J.L. Kass, and F. Viviani. The local structure of compactified Jacobians. *Proc. Lond. Math. Soc. (3)*, 110(2):510–542, 2015.
- [9] S. Casalaina-Martin, J.L. Kass, and F. Viviani. The singularities and birational geometry of the compactified universal Jacobian. *Algebr. Geom.*, 4(3):353–393, 2017.
- [10] P.H. Chaudouard and G. Laumon. Un théorème du support pour la fibration de Hitchin. *Ann. Inst. Fourier (Grenoble)*, 66(2):711–727, 2016.
- [11] P. Cook. *Local and Global Aspects of the Module Theory of Singular Curves*. PhD thesis, University of Liverpool, 1993.
- [12] J. Coskun and M. Woolf. The stable cohomology of moduli spaces of sheaves on surfaces. *to appear in J. Differential Geom.*
- [13] B. Davison. Nonabelian Hodge theory for stacks and a stacky P=W conjecture. *arXiv:2112.10830*, 2021.
- [14] B. Davison. Purity and 2-Calabi-Yau categories. *arXiv:2106.07692*, 2021.

---

<sup>8</sup>i.e. there exists a coherent sheaf  $\mathcal{F}$  on  $S$  such that  $v = (rk(\mathcal{F}), c_1(\mathcal{F}), \chi(\mathcal{F}) - rk(\mathcal{F}))$ .

- [15] M.A. de Cataldo. A support theorem for the Hitchin fibration: the case of  $SL_n$ . *Compos. Math.*, 153(6):1316–1347, 2017.
- [16] M.A. de Cataldo, T. Hausel, and L. Migliorini. Topology of Hitchin systems and Hodge theory of character varieties: the case  $A_1$ . *Ann. of Math. (2)*, 175(3):1329–1407, 2012.
- [17] M.A. de Cataldo, J. Heinloth, and L. Migliorini. A support theorem for the Hitchin fibration: the case of  $GL_n$  and  $K_C$ . *J. Reine Angew. Math.*, 780:41–77, 2021.
- [18] M.A. de Cataldo and D. Maulik. The perverse filtration for the Hitchin fibration is locally constant. *Pure Appl. Math. Q.*, 16(5):1441–1464, 2020.
- [19] M.A. de Cataldo, D. Maulik, and J. Shen. Hitchin fibrations, abelian surfaces, and the P=W conjecture. *J. Amer. Math. Soc.*, 2021.
- [20] M.A. de Cataldo and L. Migliorini. The Hodge theory of algebraic maps. *Ann. Sci. École Norm. Sup. (4)*, 38(5):693–750, 2005.
- [21] M.A. de Cataldo and L. Migliorini. The perverse filtration and the Lefschetz hyperplane theorem. *Ann. of Math. (2)*, 171(3):2089–2113, 2010.
- [22] M.A. de Cataldo, A. Rapagnetta, and G. Saccà. The Hodge numbers of O’Grady 10 via Ngô strings. *J. Math. Pures Appl. (9)*, 156:125–178, 2021.
- [23] M.A. de Cataldo and S. Zhang. A Cohomological Non Abelian Hodge Theorem in Positive Characteristic. *arXiv:2104.12970*, 2021.
- [24] J.M. Drézet. Luna’s slice theorem and applications. In *Algebraic group actions and quotients*, pages 39–89. Hindawi Publ. Corp., Cairo, 2004.
- [25] E. Esteves. Compactifying the relative Jacobian over families of reduced curves. *Trans. Amer. Math. Soc.*, 353(8):3045–3095, 2001.
- [26] C. Felisetti. Intersection cohomology of the moduli space of Higgs bundles on a genus 2 curve. *J. Inst. Math. Jussieu*, pages 1–50, 2021.
- [27] C. Felisetti and M. Mauri. P=W conjectures for character varieties with symplectic resolution. *to appear in J. Éc. polytech. Math.*, *arXiv:2006.08752*, 2020.
- [28] C. Felisetti, J. Shen, and Q. Yin. On intersection cohomology and Lagrangian fibrations of irreducible symplectic varieties. *Trans. Amer. Math. Soc.*, 375(4):2987–3001, 2022.
- [29] M. Gagné. Compactified jacobians of integral curves with double points. *Ph.D. Thesis*, 1997.
- [30] M. Goresky and R. MacPherson. *Stratified Morse theory*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1988.
- [31] M. Groechenig, D. Wyss, and P. Ziegler. Mirror symmetry for moduli spaces of Higgs bundles via p-adic integration. *Invent. Math.*, 221(2):505–596, 2020.
- [32] M. Harada and N. Proudfoot. Properties of the residual circle action on a hypertoric variety. *Pacific J. Math.*, 214(2):263–284, 2004.
- [33] T. Hausel and F. Rodriguez-Villegas. Mixed Hodge polynomials of character varieties. *Invent. Math.*, 174(3):555–624, 2008. With an appendix by Nicholas M. Katz.
- [34] T. Hausel and B. Sturmfels. Toric hyperKähler varieties. *Doc. Math.*, 7:495–534, 2002.
- [35] T. Hausel and M. Thaddeus. Mirror symmetry, Langlands duality, and the Hitchin system. *Invent. Math.*, 153(1):197–229, 2003.
- [36] T. Hausel and M. Thaddeus. Generators for the cohomology ring of the moduli space of rank 2 Higgs bundles. *Proc. London Math. Soc. (3)*, 88(3):632–658, 2004.
- [37] J. Heinloth. A conjecture of Hausel on the moduli space of Higgs bundles on a curve. *Astérisque*, (370):157–175, 2015.
- [38] J. Heinloth. The intersection form on moduli spaces of twisted  $PGL_n$ -Higgs bundles vanishes. *Math. Ann.*, 365(3-4):1499–1526, 2016.
- [39] N. Hitchin. Stable bundles and integrable systems. *Duke Math. J.*, 54(1):91–114, 1987.
- [40] N.J. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc. (3)*, 55(1):59–126, 1987.

- [41] R. Joshua. The intersection cohomology and derived category of algebraic stacks. In *Algebraic K-theory and algebraic topology (Lake Louise, AB, 1991)*, volume 407 of *NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci.*, pages 91–145. Kluwer Acad. Publ., Dordrecht, 1993.
- [42] D. Kaledin. Symplectic singularities from the Poisson point of view. *J. Reine Angew. Math.*, 600:135–156, 2006.
- [43] T. Kinjo and N. Koseki. Cohomological  $\chi$ -independence for Higgs bundles and Gopakumar-Vafa invariants. *arXiv:2112.10053*, 2021.
- [44] Y. Laszlo and M. Olsson. Perverse t-structure on artin stacks. *Mathematische Zeitschrift*, 261:737–748, 04 2009.
- [45] G. Laumon and L. Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [46] E. Markman. Generators of the cohomology ring of moduli spaces of sheaves on symplectic surfaces. *J. Reine Angew. Math.*, 544:61–82, 2002.
- [47] D. Matsushita. On base manifolds of Lagrangian fibrations. *Sci. China Math.*, 58(3):531–542, 2015.
- [48] D. Maulik and J. Shen. Cohomological  $\chi$ -independence for moduli of one-dimensional sheaves and moduli of higgs bundles. *arXiv:2012.06627*, 2020.
- [49] D. Maulik and J. Shen. On the intersection cohomology of the moduli of  $SL_n$ -Higgs bundles on a curve. *arXiv:2103.01285*, 2021.
- [50] M. Mauri. Intersection cohomology of rank two character varieties of surface groups. *J. Inst. Math. Jussieu*, 2021.
- [51] M. Mauri. Topological mirror symmetry for rank two character varieties of surface groups. *Abh. Math. Semin. Univ. Hambg.*, 2021.
- [52] M. Mauri, L. Migliorini, and R. Pagaria. The decomposition theorem for Hitchin systems via zonotopes. *In preparation*.
- [53] M. Mayrand. Local structure of singular hyperkahler quotients. *arXiv:1807.05992*, 2018.
- [54] A. Mellit. Cell decompositions of character varieties. *arXiv:1905.10685*, 2019.
- [55] M. Melo, A. Rapagnetta, and F. Viviani. Fourier-Mukai and autoduality for compactified Jacobians. I. *J. Reine Angew. Math.*, 755:1–65, 2019.
- [56] L. Migliorini, V. Shende, and F. Viviani. A support theorem for Hilbert schemes of planar curves, II. *Compos. Math.*, 157(4):835–882, 2021.
- [57] H. Nakajima. Quiver varieties and Kac-Moody algebras. *Duke Math. J.*, 91(3):515–560, 1998.
- [58] B.C. Ngô. Le lemme fondamental pour les algèbres de Lie. *Publ. Math. Inst. Hautes Études Sci.*, (111):1–169, 2010.
- [59] R. Pandharipande. A compactification over  $\overline{M}_g$  of the universal moduli space of slope-semistable vector bundles. *J. Amer. Math. Soc.*, 9(2):425–471, 1996.
- [60] A. Perego and A. Rapagnetta. The moduli spaces of sheaves on K3 surfaces are irreducible symplectic varieties. *arXiv:1802.01182*, 2018.
- [61] N. Proudfoot and B. Webster. Intersection cohomology of hypertoric varieties. *J. Algebraic Geom.*, 16(1):39–63, 2007.
- [62] M. Saito. Introduction to mixed Hodge modules. Number 179-180, pages 10, 145–162. 1989. Actes du Colloque de Théorie de Hodge (Luminy, 1987).
- [63] D. Schaub. Courbes spectrales et compactifications de jacobiniennes. *Math. Z.*, 227(2):295–312, 1998.
- [64] S. Schlegel Mejia. BPS cohomology for rank 2 degree 0 Higgs bundles (and more). *arXiv:2201.09962*, 2022.
- [65] E. Sernesi. *Deformations of algebraic schemes*, volume 334 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin, 2006.
- [66] C.T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. I. *Inst. Hautes Études Sci. Publ. Math.*, (79):47–129, 1994.
- [67] C.T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. II. *Inst. Hautes Études Sci. Publ. Math.*, (80):5–79, 1994.
- [68] H. Whitney. Tangents to an analytic variety. *Ann. of Math. (2)*, 81:496–549, 1965.