

# HOPF-GALOIS EXTENSIONS AND TWISTED HOPF ALGEBROIDS

XIAO HAN AND SHAHN MAJID

ABSTRACT. We show that the Ehresmann-Schauenburg bialgebroid of a quantum principal bundle  $P$  or Hopf Galois extension with structure quantum group  $H$  is in fact a left Hopf algebroid  $\mathcal{L}(P, H)$ . We show further that if  $H$  is coquasitriangular then  $\mathcal{L}(P, H)$  has an antipode map  $S$  obeying certain minimal axioms. Trivial quantum principal bundles or cleft Hopf Galois extensions with base  $B$  are known to be cocycle cross products  $B\#_{\sigma}H$  for a cocycle-action pair  $(\triangleright, \sigma)$  and we look at these of a certain ‘associative type’ where  $\triangleright$  is an actual action. In this case also, we show that the associated left Hopf algebroid has an antipode obeying our minimal axioms. We show that if  $\mathcal{L}$  is any left Hopf algebroid then so is its cotwist  $\mathcal{L}^{\varsigma}$  as an extension of the previous bialgebroid Drinfeld cotwist theory. We show that in the case of associative type,  $\mathcal{L}(B\#_{\sigma}H, H) = \mathcal{L}(B\#H)^{\tilde{\sigma}}$  for a Hopf algebroid cotwist  $\varsigma = \tilde{\sigma}$ . Thus, switching on  $\sigma$  of associative type appears at the Hopf algebroid level as a Drinfeld cotwist. We view the affine quantum group  $\widehat{U_q(sl_2)}$  and the quantum Weyl group of  $u_q(sl_2)$  as examples of associative type.

## 1. INTRODUCTION

A right Hopf-Galois extension or quantum principal bundle is a right comodule algebra  $P$  under a Hopf algebra  $H$  such that the induced map  $P \otimes_B P \rightarrow P \otimes H$  is an isomorphism of vector spaces, where  $B = P^{co(H)}$  is the fixed subalgebra. In noncommutative geometry, one thinks of  $P$  as like the algebra of functions on the total space of a principal bundle,  $H$  as like the algebra of functions on the fibre group and  $B$  as like the functions on the base, except that all algebras may be noncommutative and the group structure of the fibre is now a Hopf algebra or quantum group. This is the starting point for quantum group gauge theory[8] and we refer to [2, Chap. 5] for a recent account of the geometric picture, differential structures and connections. In the classical case, one has an associated Ehresmann groupoid of a principal bundle and similarly for any Hopf Galois extension, one has[23] the ‘Ehresmann-Schauenburg bialgebroid’ which we will show is in fact a left Hopf algebroid and denote  $\mathcal{L}(P, H)$ . This is a main result of the paper and appears in Theorem 4.4.

In general, while the axioms of left bialgebroids [9, 6] are somewhat settled at this point (with a similar notion for right bialgebroids) the correct notion of Hopf algebroid is less clear. By left Hopf algebroid we will mean in the sense of [24] that a certain ‘translation map’ is invertible. This reduces to existence of a usual Hopf algebra antipode  $S$  when the base algebra  $B = k$ , but in general an antipode map appears to be a stronger notion and there are several candidates. One of these is in [6], where one needs both a left and right bialgebroid with  $S$  connecting them and  $S$  antimultiplicative. This notion is, however, too restrictive for our purposes and we introduce in Lemma 3.3 what we consider a minimal formulation for an invertible antipode  $S$ , which drops  $S$  being antimultiplicative but is strong enough to imply a left Hopf algebroid. We expect but do not show that it is a strictly stronger concept than a left Hopf algebroid and we show that  $\mathcal{L}(P, H)$  has such an antipode at least if  $H$  is coquasitriangular. Another general result of the paper is Theorem 3.7 about Drinfeld (co)twists of left Hopf algebroids, extending the bialgebroid

---

*Key words and phrases.* Hopf algebroid, bialgebroid, quantum group, quantum principal bundle, Hopf-Galois extension, cocycle cross product, smash product, bicrossproduct. Version 1.3.

version in [5]. We do not find conditions on the cotwists for the antipode to cotwist, but the example in Corollary 3.14 comes close.

Before these general results, Section 2 starts with the simpler case of a cleft Hopf Galois extension [22] or ‘trivial’ quantum principal bundle [8] defined as a Hopf-Galois extension equipped with a cleaving or ‘trivialisation’ map  $j : H \rightarrow P$  of comodule algebras (where  $H$  coacts on itself by the coproduct) which is convolution-invertible as a map from the coalgebra of  $H$  to the algebra  $P$ . One of the first observations to be made here is that, unlike classical geometry, there is a potentially nontrivial nonAbelian cohomology associated to such seemingly trivial extensions, denoted  $\mathcal{H}^2(H, B)$  in [19][18]. Here it is known that a cleft extension allows for an identification

$$P \cong B \#_{\sigma} H$$

as comodule algebras for certain ‘cocycle data’  $(\triangleright, \sigma)$  where  $\triangleright : H \otimes B \rightarrow B$  is a measuring (but not necessarily an action) and  $\sigma : H \otimes H \rightarrow B$  obeying certain conditions. Gauge transformation by convolution invertible maps  $u : H \rightarrow B$  changes  $(\triangleright, \sigma)$  and the nonAbelian cohomology can be defined as cocycles modulo these. In the special case where  $H$  is cocommutative, it was shown in [10] that there is a bijective correspondence between the equivalence classes of  $H$ -cleft Hopf Galois extensions and the usual cohomology group  $\mathcal{H}^2(H, Z(B))$ , where  $Z(B)$  is the centre of  $B$  and  $\sigma$  is the linear extension of a group 2-cocycle in the case where  $H$  is a group algebra. Another special case is where  $B = k$  is the ground field and in this case  $\sigma$  is a Drinfeld 2-cocycle as in the theory of Hopf algebra cotwists [18, Chap 2]. We introduce and study a joint ‘associative type’ generalisation of these two special cases where  $\triangleright$  is in fact an action so that we also have an ordinary smash product  $B \# H$ .

Cleft extensions are then studied further in Section 3 to illustrate the Hopf algebroid theory more explicitly. First we give a version  $B^e \#_{\sigma} H$  of the Ehresmann-Schaubenburg bialgebroid in the cleft cocycle extension case directly on the vector of  $B^e \otimes H$  where  $B^e = B \otimes B^{op}$ . From our general results, this is necessarily a left Hopf algebroid but we show that it has an antipode at least when  $\sigma$  is of associative type. We also show in the associative type case that there is a Hopf algebroid cotwist  $\tilde{\sigma}$  on the Hopf algebroid  $B^e \# H$  such that  $B^e \#_{\sigma} H = (B^e \# H)^{\tilde{\sigma}}$  is a Hopf algebroid twist. Thus, the introduction of  $\sigma$  at the cocycle level corresponds to a Drinfeld cotwist at the Hopf algebroid level. We also explore how our constructions change both under a Drinfeld cotwist of  $\chi$  on  $H$  and under a gauge transformation  $u$  from our two points of view.

Section 4 contains our results for general quantum principal bundles and the paper then concludes with some examples in Section 5. Of course, many Hopf-Galois extensions are known but the new notion of associative type is less clear and we show how they arise from certain cocycle bicrossproducts of interest in the theory of quantum groups.

## 2. COCYCLE CLEFT EXTENSIONS AND ASSOCIATIVE TYPE

This section starts with recalling basic algebraic preliminaries, then studies cocycle extensions of associative type. We work over a field  $k$ .

**2.1. Basic algebraic preliminaries.** We recall that a Hopf algebra  $H$  means a unital algebra which is also a counital coalgebra with coproduct  $\Delta$  and counit  $\epsilon$ , that these are algebra maps and there is an antipode  $S$  such that  $(Sh_{(1)})h_{(2)} = 1\epsilon(h) = h_{(1)}Sh_{(2)}$ . We use here the sumless Sweedler notation  $\Delta(h) = h_{(1)} \otimes h_{(2)}$  for all  $h \in H$ . We refer to texts such as [25, 18] for more details. If  $C$  is a coalgebra and  $P$  is an algebra, the space  $\text{Hom}_k(C, P)$  is unital associative algebra with the ‘convolution’ product  $\phi \star \psi(c) := \phi(c_{(1)})\psi(c_{(2)})$ , for all  $\phi, \psi \in \text{Hom}_k(C, P)$  and from this point of view  $S$  is the assumed convolution-inverse of the identity map  $\text{id} : H \rightarrow H$ .

Given a Hopf algebra  $H$ , an *left  $H$ -module algebra* is an algebra  $B$  with a left action  $\triangleright$  of  $H$  such that  $h \triangleright (ab) = (h_{(1)} \triangleright a)(h_{(2)} \triangleright b)$  for all  $a, b \in B$  and  $h \in H$ , and  $h \triangleright 1 = \epsilon(h)1$ .

In this case one has a smash or cross product algebra  $B\#H$  built on  $B \otimes H$  with product  $(a\#h)(a'\#g) = a(h_{(1)} \triangleright a')\#h_{(2)}g$  for all  $a, a' \in B$  and  $g, h \in H$ . Similarly, *right  $H$ -comodule algebra* is an algebra  $P$  with a right coaction  $\Delta_R : P \rightarrow P \otimes H$  (the axioms are dual to those of a module) such that  $\Delta_R$  is an algebra map, where  $P \otimes H$  has the tensor product algebra structure. We use the sumless Sweedler notation  $\Delta_R(p) = p_{(0)} \otimes p_{(1)}$ .

The category of left modules  ${}^H\mathcal{M}$  and the category of right comodules  $\mathcal{M}^H$  are both monoidal with finite-dimensional objects rigid. For example, if  $V$  and  $W$  are right comodules the  $V \otimes W$  is right  $H$ -comodule  $\Delta_{R,V \otimes W}(v \otimes w) := v_{(0)} \otimes w_{(0)} \otimes v_{(1)}w_{(1)}$  for all  $v \in V$  and  $w \in W$ . An  $H$ -module algebra  $B$  just means an algebra in  ${}^H\mathcal{M}$  and a comodule algebra  $P$  just means an algebra in  $\mathcal{M}^H$ .

Finally, if  $H$  is a Hopf algebra and  $\chi : H \otimes H \rightarrow k$  a Drinfeld 2-cocycle or *cotwist* in the sense of convolution-invertible and obeying[18]

$$\chi(g_{(1)} \otimes f_{(1)})\chi(h \otimes g_{(2)}f_{(2)}) = \chi(h_{(1)} \otimes g_{(1)})\chi(h_{(1)}g_{(1)} \otimes f), \quad \chi(1 \otimes h) = \chi(h \otimes 1) = \epsilon(h)$$

then one has a new Hopf algebra  $H^\chi$  on the same vector space as  $H$  but with a modified product

$$h \bullet_\chi g = \chi(h_{(1)} \otimes g_{(1)})h_{(2)}g_{(2)}\chi^{-1}(h_{(3)} \otimes g_{(3)})$$

such that the category  $\mathcal{M}^{H^\chi}$  is monoidally equivalent to  $\mathcal{M}^H$ [18]. Moreover, if  $P$  is an  $H$ -comodule algebra then there is a 1-sided cotwisted comodule algebra  $P_\chi$  with new product[18, Sec. 2.3]

$$p \cdot_\chi q := p_{(0)}q_{(0)}\chi^{-1}(p_{(1)} \otimes q_{(1)}), \quad (2.1)$$

for all  $p, q \in P$  and the same coaction on the underlying vector space.

**2.2. Cocycle cross product algebras revisited.** More generally,  $\triangleright : H \otimes B \rightarrow B$  is a measuring of a Hopf algebra  $H$  on an algebra  $B$  if  $h \triangleright (ab) = (h_{(1)} \triangleright a)(h_{(2)} \triangleright b)$  for all  $a, b \in B$  and  $h \in H$ , and  $h \triangleright 1 = \epsilon(h)1$ , i.e. like a module algebra but without requiring  $\triangleright$  to be an action. Given such a measuring  $\triangleright$ , we assume a ‘2-cocycle’ in the sense of a convolution-invertible map  $\sigma : H \otimes H \rightarrow B$  such that

- (1)  $1 \triangleright b = b$ ,
- (2)  $h \triangleright (g \triangleright b) = \sigma(h_{(1)}, g_{(1)})(h_{(2)}g_{(2)} \triangleright b)\sigma^{-1}(h_{(3)}, g_{(3)})$ ,
- (3)  $\sigma(h, 1) = \sigma(1, h) = \epsilon(h)1$ ,
- (4)  $(h_{(1)} \triangleright \sigma(g_{(1)}, f_{(1)}))\sigma(h_{(2)}, g_{(2)}f_{(2)}) = \sigma(h_{(1)}, g_{(1)})\sigma(h_{(2)}g_{(2)}, f)$ ,

for all  $h, g, f \in H$  and  $b \in B$ . Then [11] showed that there is an *cocycle cross product* algebra structure on  $B\#_\sigma H$  built on  $B \otimes H$  as vector space, with the product

$$(b\#_\sigma h)(c\#_\sigma g) = b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)})\#_\sigma h_{(3)}g_{(2)}, \quad (2.2)$$

for all  $h, g \in H$  and  $b, c \in B$ . The converse is also true: given that  $H$  measures  $B$  and  $\sigma : H \otimes H \rightarrow B$  is a convolution invertible linear map, these conditions are needed for the formula given to define an associative product with unit  $1\#_\sigma 1$ . We have the following useful lemma:

**Lemma 2.1.** *cf. [22, Prop. 7.2.7] Given the data for a cocycle cross product  $B\#_\sigma H$ , we have the following properties of  $\sigma$ :*

- (1)  $\sigma^{-1}(h_{(1)}, g_{(1)}f_{(1)})(h_{(2)} \triangleright \sigma^{-1}(g_{(2)}, f_{(2)})) = \sigma^{-1}(h_{(1)}g_{(1)}, f)\sigma^{-1}(h_{(2)}, g_{(2)})$ ,
- (2)  $h \triangleright \sigma(g, f) = \sigma(h_{(1)}, g_{(1)})\sigma(h_{(2)}g_{(2)}, f_{(1)})\sigma^{-1}(h_{(3)}, g_{(3)}f_{(2)})$ ,
- (3)  $h \triangleright \sigma^{-1}(g, f) = \sigma(h_{(1)}, g_{(1)}f_{(1)})\sigma^{-1}(h_{(2)}g_{(2)}, f_{(2)})\sigma^{-1}(h_{(3)}, g_{(3)})$ ,
- (4)  $(h_{(1)} \triangleright \sigma^{-1}(S(h_{(4)}), h_{(5)}))\sigma(h_{(2)}, S(h_{(3)})) = \epsilon(h)1$ .
- (5)  $\sigma^{-1}(S(h_{(1)}), S(g_{(1)}))\sigma(h_{(2)}g_{(2)}) = \sigma^{-1}(S(h_{(3)}), h_{(4)})(S(h_{(2)} \triangleright \sigma^{-1}(S(g_{(2)}), g_{(3)}h_{(5)}))\sigma(S(h_{(1)}), S(g_{(1)})))$ .
- (6)  $\sigma^{-1}(S(h_{(2)}), h_{(3)})S(h_{(1)} \triangleright \sigma(h_{(4)}, S(h_{(5)}))) = \epsilon(h)1$ .

*Proof.* (1) and (2) can be derived by applying the convolution inverse on both side of the 2-cocycle conditions, and (3) can be derived from (1). For (4),

$$\begin{aligned}
& (h_{(1)} \triangleright \sigma^{-1}(S(h_{(4)}), h_{(5)})) \sigma(h_{(2)}, h_{(3)}) \\
&= \sigma(h_{(1)}, S(h_{(8)})h_{(9)}) \sigma^{-1}(h_{(2)}, S(h_{(7)}), h_{(10)}) \sigma^{-1}(h_{(3)}, S(h_{(6)})) \sigma(h_{(4)}, S(h_{(5)})) \\
&= \sigma(h_{(1)}, S(h_{(4)})h_{(5)}) \sigma^{-1}(h_{(2)}, S(h_{(3)}), h_{(6)}) \\
&= \epsilon(h)1,
\end{aligned}$$

for any  $h \in H$ , and similarly for (6). For (5)

$$\begin{aligned}
& \sigma^{-1}(S(h_{(3)}), h_{(4)})(S(h_{(2)}) \triangleright \sigma^{-1}(S(g_{(2)}), g_{(3)}h_{(5)})) \sigma(S(h_{(1)}), S(g_{(1)})) \\
&= \sigma^{-1}(S(h_{(5)}), h_{(6)}) \sigma(S(h_{(4)}), S(g_{(4)})g_{(5)}h_{(7)}) \sigma^{-1}(S(h_{(3)})S(g_{(3)}), g_{(6)}h_{(8)}) \\
&\quad \sigma^{-1}(S(h_{(2)}), S(g_{(2)})) \sigma(S(h_{(1)}), S(g_{(1)})) \\
&= \sigma^{-1}(S(h_{(1)})S(g_{(1)}), h_{(2)}g_{(2)}),
\end{aligned}$$

where the first step use (3).  $\square$

From an extension theory point of view, a cleft extension of an algebra  $B$  by a Hopf algebra  $H$  is a comodule algebra  $P$  and a convolution-invertible unital comodule map  $j : H \rightarrow P$  and an algebra inclusion such that

$$B \hookrightarrow P \leftarrow H$$

such that the tensor product of these maps combined with the product of  $P$  gives an isomorphism  $B \otimes H \rightarrow P$  of right comodule and left  $B$ -module, with canonical left- $B$  module and right  $H$  comodule structure on  $B \otimes H$  (this entails that  $j$  is injective). It is easy to see that  $B \#_{\sigma} H$  meets these conditions and conversely, given the extension data, one can show that  $B = P^{co(H)}$  is the invariant subalgebra and construct  $(\triangleright, \sigma)$  by [19][18, Prop. 6.3.6]

$$h \triangleright b = j(h_{(1)})bj^{-1}(h_{(2)}), \quad \sigma(h \otimes g) = j(h_{(1)})j(g_{(1)})j^{-1}(h_{(2)}g_{(2)}) \quad (2.3)$$

with images in  $B$  due to the assumed equivariance of  $j$ . A map between extensions is a  $H$ -comodule map which is the identity on  $B$ . Moreover, if  $(\triangleright, \sigma)$  is cocycle data then

$$h \triangleright^u b = u^{-1}(h_{(1)})(h_{(2)} \triangleright b)u(h_{(3)}), \quad (2.4)$$

$$\sigma^u(h, g) = u^{-1}(h_{(1)})(h_{(2)} \triangleright u^{-1}(g_{(1)})) \sigma(h_{(3)}, g_{(2)})u(h_{(4)}g_{(3)}). \quad (2.5)$$

for a convolution-invertible unital map  $u : H \rightarrow B$  then  $(\triangleright^u, \sigma^u)$  gives an isomorphic cocycle extension by

$$\Theta_u : B \#_{\sigma^u} H \rightarrow B \#_{\sigma} H, \quad \Theta(b \#_{\sigma^u} h) = bu^{-1}(h_{(1)}) \#_{\sigma} h_{(2)},$$

and conversely isomorphic extensions are related in this way [19][18, Props. 6.3.4, 6.3.5]. This therefore defines a kind of ‘nonAbelian cohomology’ theory  $\mathcal{H}^2(H, B)$  realised as cocycles modulo such ‘gauge equivalence’. We will recall in Section 4 how such cleft extensions are Hopf-Galois and hence can be viewed as trivial quantum principal bundles.

Cleft cocycle extensions and their dual have also been applied in a theory of cocycle bicrossproduct extensions [17][18] where  $B$  and  $P = B_{\sigma} \blacktriangleright^{\psi} H$  are Hopf algebras and the extension is both cleft and ‘cocleft’. This will be a source of examples in Section 5. Finally, a different operation one can perform on cocycle cross products is a Drinfeld cotwist.

**Proposition 2.2.** *If  $(\triangleright, \sigma)$  is a cocycle on for  $B \#_{\sigma} H$  then  $\triangleright_{\chi} = \triangleright$  and  $\sigma_{\chi} = \sigma * \chi^{-1}$  are cocycle data for  $B \#_{\sigma_{\chi}} H^{\chi}$ . Moreover,  $\mathcal{H}^2(H^{\chi}, B) \cong \mathcal{H}^2(H, B)$ .*

*Proof.* One can simply check that the twisted data meets the conditions for a cocycle. Or, from the point of view of cleft extensions, assume  $j : H \rightarrow P$  is the cleaving map of the cleft extension. We define  $j_\chi = j$  on the same underlying vector space of  $H$  now viewed as a  $H^\chi$ -comodule map. Now we claim that its inverse is given by

$$j_\chi^{-1}(h) := j^{-1}(h_{(3)})\chi(h_{(1)} \otimes S(h_{(2)})),$$

where  $j^{-1}$  is the inverse of the original cleaving map. On the one hand,

$$\begin{aligned} j_\chi^{-1} * j_\chi(h) &= \chi(h_{(1)}, S(h_{(2)}))j^{-1}(h_{(3)}) \cdot_\chi j(h_{(4)}) \\ &= \chi(h_{(1)}, S(h_{(2)}))j^{-1}(h_{(4)})j(h_{(5)})\chi^{-1}(S(h_{(3)}), h_{(6)}) \\ &= \chi(h_{(1)}, S(h_{(2)}))\chi^{-1}(S(h_{(3)}), h_{(4)}) = \epsilon(h), \end{aligned}$$

where the second step uses the fact that  $j^{-1}(h_{(0)}) \otimes j^{-1}(h_{(1)}) = j^{-1}(h_{(2)}) \otimes S(h_{(1)})$  for any  $h \in H$ , the last step uses the fact that  $\chi$  is a 2-cocycle on the Hopf algebra  $H$ . On the other hand,

$$\begin{aligned} j_\chi * j_\chi^{-1}(h) &= j(h_{(1)}) \cdot_\chi j^{-1}(h_{(4)})\chi(h_{(2)}, S(h_{(3)})) \\ &= j(h_{(1)})j^{-1}(h_{(6)})\chi(h_{(2)}, S(h_{(5)}))\chi(h_{(3)}, S(h_{(4)})) = \epsilon(h), \end{aligned}$$

According to (2.3), the deformed 2-cocycle action  $\triangleright_\chi$  is given by

$$\begin{aligned} h \triangleright_\chi b &= j_\chi(h_{(1)}) \cdot_\chi b \cdot_\chi j_\chi^{-1}(h_{(2)}) \\ &= (1 \#_\sigma h_{(1)}) \cdot_\chi (b\chi(h_{(2)}, S(h_{(3)}))j^{-1}(h_{(4)})) \\ &= (1 \#_\sigma h_{(1)}) \cdot_\chi (b\chi(h_{(2)}, S(h_{(3)}))\sigma^{-1}(S(h_{(5)}), h_{(6)})\#_\sigma S(h_{(4)})) \\ &= (1 \#_\sigma h_{(1)})(b\chi(h_{(3)}, S(h_{(4)}))\sigma^{-1}(S(h_{(7)}), h_{(8)})\#_\sigma S(h_{(6)}))\chi^{-1}(h_{(2)}, S(h_{(5)})) \\ &= (1 \#_\sigma h_{(1)})(b\sigma^{-1}(S(h_{(3)}), h_{(4)})\#_\sigma S(h_{(2)})) \\ &= (h_{(1)} \triangleright b)(h_{(2)} \triangleright \sigma^{-1}(S(h_{(5)}), h_{(6)}))\sigma(h_{(3)}, S(h_{(4)}))\#_\sigma 1 \\ &= h \triangleright b \#_\sigma 1, \end{aligned}$$

where the last step uses Proposition 2.1. So we can see the the twisted action is unchanged. And the deformed 2-cocycle  $\sigma_\chi$  is given by

$$\begin{aligned} \sigma_\chi(h, g) &= j_\chi(h_{(1)}) \cdot_\chi j_\chi(g_{(1)}) \cdot_\chi j_\chi^{-1}(h_{(2)}) \cdot_\chi g_{(2)} \\ &= (j(h_{(1)})j(g_{(1)})\chi^{-1}(h_{(2)}, g_{(2)})) \cdot_\chi j_\chi^{-1}(\chi(h_{(3)}, g_{(3)})h_{(4)}g_{(4)}\chi^{-1}(h_{(5)}, g_{(5)})) \\ &= (j(h_{(1)})j(g_{(1)})) \cdot_\chi j_\chi^{-1}(h_{(2)}g_{(2)})\chi^{-1}(h_{(3)}, g_{(3)}) \\ &= (j(h_{(1)})j(g_{(1)})) \cdot_\chi (\chi(h_{(2)}g_{(2)}, S(h_{(3)}g_{(3)}))j^{-1}(h_{(4)}g_{(4)}))\chi^{-1}(h_{(5)}, g_{(5)}) \\ &= j(h_{(1)})j(g_{(1)})j^{-1}(h_{(6)}g_{(6)})\chi^{-1}(h_{(2)}g_{(2)}, S(h_{(5)}g_{(5)}))\chi(h_{(3)}g_{(3)}, S(h_{(4)}g_{(4)}))\chi^{-1}(h_{(7)}, g_{(7)}) \\ &= j(h_{(1)})j(g_{(1)})j^{-1}(h_{(2)}g_{(2)})\chi^{-1}(h_{(3)}, g_{(3)}) \\ &= \sigma(h_{(1)}, g_{(1)})\chi^{-1}(h_{(2)}, g_{(2)}) \end{aligned}$$

for any  $h, g \in H$  and  $b \in B$ .

For the last part, if we we gauge transform by  $u : H \rightarrow B$  then  $(\sigma^u)_\chi = (\sigma_\chi)^u$ . Indeed,

$$\begin{aligned} (\sigma^u)_\chi(h, g) &= u^{-1}(h_{(1)})(h_{(2)} \triangleright u^{-1}(g_{(1)}))\sigma(h_{(3)}, g_{(2)})u(h_{(4)}g_{(3)})\chi^{-1}(h_{(5)}, g_{(4)}) \\ &= u^{-1}(h_{(1)})(h_{(2)} \triangleright u^{-1}(g_{(1)}))\sigma(h_{(3)}, g_{(2)})\chi^{-1}(h_{(4)}, g_{(5)})\chi(h_{(6)}, g_{(5)})u(h_{(7)}g_{(6)})\chi^{-1}(h_{(8)}, g_{(7)}) \\ &= u^{-1}(h_{(1)})(h_{(2)} \triangleright u^{-1}(g_{(1)}))\sigma(h_{(3)}, g_{(2)})\chi^{-1}(h_{(4)}, g_{(5)})u(h_{(5)} \cdot_\chi g_{(4)}) \\ &= (\sigma_\chi)^u(h, g), \end{aligned}$$

so that the equivalence classes are in bijective correspondence under the cotwist.  $\square$

### 2.3. Cocycles of associative type.

**Definition 2.3.** We call a crossed product  $B\#_\sigma H$  associative type or we say  $(\triangleright, \sigma)$  is a cocycle of associative type, if  $B$  is an  $H$ -module algebra under  $\triangleright$ .

It is easy to see that this happens iff

$$\sigma(h_{(1)}, g_{(1)})((h_{(2)}g_{(2)}) \triangleright b) = ((h_{(1)}g_{(1)}) \triangleright b)\sigma(h_{(2)}, g_{(2)}), \quad (2.6)$$

for any  $b \in B$ ,  $h, g \in H$ . Clearly, in the associative type case,  $B$  is an  $H$ -module algebra under the same left action, and  $B\#H$  is well defined. A lemma needed later is

**Lemma 2.4.** Let  $B\#_\sigma H$  be of associative type. Then

$$(S(h_{(3)}) \triangleright b)\sigma^{-1}(S(h_{(1)}), h_{(2)}) = \sigma^{-1}(S(h_{(2)}), h_{(3)})(S(h_{(1)}) \triangleright b)$$

*Proof.* Since  $B\#_\sigma H$  is a associative type, we have

$$b\sigma^{-1}(S(h_{(1)}), h_{(2)}) = ((S(h_{(2)})h_{(3)}) \triangleright b)\sigma^{-1}(S(h_{(1)}), h_{(4)}) = \sigma^{-1}(S(h_{(2)}), h_{(3)})(S(h_{(1)})h_{(4)}) \triangleright b).$$

Then

$$\begin{aligned} (S(h_{(3)}) \triangleright b)\sigma^{-1}(S(h_{(1)}), h_{(2)}) &= \sigma^{-1}(S(h_{(2)}), h_{(3)})((S(h_{(1)})h_{(4)}) \triangleright (S(h_{(5)}) \triangleright b)) \\ &= \sigma^{-1}(S(h_{(2)}), h_{(3)})(S(h_{(1)}) \triangleright b). \end{aligned}$$

□

On the other hand, if  $B\#_\sigma H$  is of associative type then  $B\#_{\sigma^u} H$  need not be. We say an cleft extension is of associative type if its underlying  $(\triangleright, \sigma)$  as in Section 2.2 obey Definition 2.3. Let  $Z_{as}^2(H, B)$  be the set of such action-cocycle pairs for a fixed Hopf algebra  $H$  and algebra  $B$ .

**Proposition 2.5.** Cleft extensions of associative type are classified by  $\mathcal{H}_{as}(H, B)$  defined as  $Z_{as}^2(H, B)$  modulo transformation  $(\triangleright, \sigma)$  to  $(\triangleright^u, \sigma^u)$  where  $u : H \rightarrow B$  is convolution invertible and obeys

$$u^{-1}(h_{(1)})(h_{(2)} \triangleright (u^{-1}(g_{(1)})(g_{(2)} \triangleright b)u(g_{(3)})))u(h_{(3)}) = u^{-1}(h_{(1)}g_{(1)})((h_{(2)}g_{(2)}) \triangleright b)u(h_{(3)}g_{(3)}),$$

for any  $h, g \in H$  and  $b \in B$ .

*Proof.* Any two equivalent cleft extensions are related by some convolution-invertible  $u$  as recalled in Section 2.2. Not all of them will preserve associative type, the condition for this being  $h\triangleright^u(g\triangleright^u b) = (hg)\triangleright^u b$  which is equivalent to the condition stated using the relation (2.4). It follows, but can be checked explicitly, that if  $(\triangleright, \sigma) \sim (\triangleright^u, \sigma^u) \sim ((\triangleright^u)^v, (\sigma^u)^v)$  so that  $v$  obeys the stated condition with respect to  $\triangleright^u$ , then the convolution product  $u * v$  obeys the stated conditions and  $((\triangleright^u)^v, (\sigma^u)^v) = (\triangleright^{u*v}, \sigma^{u*v})$ . Similarly for  $u^{-1}$  to go back from  $(\triangleright^u, \sigma^u)$ . □

*Example 2.6.* Let  $H$  be a Hopf algebra,  $B$  an  $H$ -module algebra of an inner form  $h\triangleright b = u(h_{(1)})bu^{-1}(h_{(2)})$  where  $u : H \rightarrow B$  is convolution invertible and such that

$$u^{-1}(g_{(1)})u^{-1}(h_{(1)})u(h_{(2)}g_{(2)}) \in Z(B)$$

for all  $h, g \in H$  (this is necessary and sufficient to have an action). Then  $(\triangleright, 1\epsilon \otimes \epsilon)$  (i.e. with trivial cocycle) is of associative type and cohomologous in  $\mathcal{H}_{as}(H, B)$  to  $(\text{id}\epsilon, \sigma^u)$  (i.e. with trivial action) where  $\sigma^u(h, g) = u^{-1}(g_{(1)})u^{-1}(h_{(1)})u(h_{(2)}g_{(2)})$ . Conversely, if  $B$  is an  $H$ -module algebra  $(\triangleright, \sigma = 1\epsilon \otimes \epsilon)$  is cohomologous to a cocycle of the form  $(\text{id}\epsilon, \sigma)$  (i.e. for trivial action) then  $\triangleright$  is of this inner form. For the first part, the condition in Proposition 2.5 reduces to the assumed property of  $u$  and for this combination to lie in the centre  $Z(B)$  is exactly the condition (2.6) for  $\sigma^u$  to be of associative type when the action is trivial. In the other direction for  $\triangleright^u$  to end up trivial,  $\triangleright$  has to be of the inner form in view of (2.4). For a very concrete example one could let  $H = \mathbb{C}\mathbb{Z}_2$  and  $B$  an algebra with an automorphism  $\alpha$ . If  $\alpha^2 = \text{id}$  then we make  $B$  into an  $H$ -module algebra

with the generator of  $\mathbb{Z}_2$  acting by  $\alpha$ . This action is of the inner form if and only if  $\alpha$  is an inner automorphism.

Note that one could further restrict our gauge transformations to obey

$$u(h_{(1)})(h_{(2)}\triangleright b) = (h_{(1)}\triangleright b)u(h_{(2)}) \quad (2.7)$$

which is stronger than the condition in Proposition 2.5 and enforces that  $\triangleright^u = \triangleright$ . This would define a cohomology  $\mathcal{H}_{\triangleright}^2(H, B)$  for cocycle cross products for a fixed  $H$ -module algebra  $B$ . This condition is, however, quite restrictive. Also note that our associative type construction should not be confused with lazy Sweedler cohomology in the terminology of [4] where  $\sigma$  is  $k$ -valued.

Similarly, not every cotwist preserves associative type:

**Proposition 2.7.** *A Drinfeld co-twist  $\chi$  as in Proposition 2.2 preserves associative type iff  $(\triangleright, \chi)$  viewed as a cocycle with values in  $k$ .  $1 \in B$  is of associative type.*

*Proof.* Indeed,

$$(h \cdot_{\chi} g) \triangleright_{\chi} b = \chi(h_{(1)}, g_{(1)})((h_{(2)}g_{(2)}) \triangleright b)\chi^{-1}(h_{(3)}, g_{(3)}) \neq h \triangleright_{\chi} (g \triangleright_{\chi} b) = h \triangleright (g \triangleright b) = (hg) \triangleright b.$$

would need  $(h \cdot_{\chi} g) \triangleright = (hg) \triangleright$  i.e. act the same on  $B$  and we see that this happens if and only if

$$\chi(h_{(1)}, g_{(1)})((h_{(2)}g_{(2)}) \triangleright b)\chi^{-1}(h_{(3)}, g_{(3)}) = (hg) \triangleright b$$

which we interpret as stated.  $\square$

### 3. ANTIPODE AND TWISTING OF COCYCLE HOPF ALGEBROIDS

We start with preliminaries on bialgebroids before proceeding to the our new results for certain ‘cocycle Hopf algebroids’  $B^e \#_{\sigma} H$ . These are a special case of Ehresmann-Schauburg in Section 4 but of interest in their own right.

**3.1. Preliminaries on bialgebroids and Hopf algebroids.** Here we recall the basic definitions (cf. [9], [5]). Let  $B$  be a unital algebra over a field  $k$ . A  $B$ -ring just means a unital algebra in the category  ${}_B\mathcal{M}_B$  of  $B$ -bimodules. Likewise, a  $B$ -coring is a coalgebra in  ${}_B\mathcal{M}_B$ . Morphisms of  $B$  (co)rings map the (co)algebra structures but now in the category  ${}_B\mathcal{M}_B$ .

In practice, specifying a unital  $B$ -ring  $\mathcal{L}$  is equivalent to specifying a unital algebra  $\mathcal{L}$  (over  $k$ ) and an algebra map  $\eta : B \rightarrow \mathcal{L}$ . Left and right multiplication in  $\mathcal{L}$  pull back to left and right  $B$ -actions as a bimodule (so  $b.p.c = \eta(b)p\eta(c)$  for all  $b, c \in B$  and  $p \in \mathcal{L}$ ) and the product descends to the product  $\mu_B : \mathcal{L} \otimes_B \mathcal{L} \rightarrow \mathcal{L}$  with  $\eta$  the unit map. Conversely, given  $\mu_B$  we can pull back to an associative product on  $\mathcal{L}$  with unit  $\eta(1)$ .

Now suppose that  $s : B \rightarrow \mathcal{L}$  and  $t : B^{op} \rightarrow \mathcal{L}$  are algebra maps with images that commute. Then  $\eta(b \otimes c) = s(b)t(c)$  is an algebra map  $\eta : B^e \rightarrow \mathcal{L}$  where  $B^e = B \otimes B^{op}$  and is equivalent to making  $\mathcal{L}$  a  $B^e$ -ring. The left  $B^e$ -action part of this is equivalent to a  $B$ -bimodule structure

$$b.p.c = s(b)t(c)p \quad (3.1)$$

for all  $b, c \in B$  and  $p \in \mathcal{L}$ .

**Definition 3.1.** *Let  $B$  be a unital algebra. A left  $B$ -bialgebroid is an algebra  $\mathcal{L}$  and (‘source’ and ‘target’) commute algebra maps  $s : B \rightarrow \mathcal{L}$  and  $t : B^{op} \rightarrow \mathcal{L}$  (and hence a  $B^e$ -ring) and a  $B$ -coring for the bimodule structure (3.1) which is compatible in the sense*

(i) *The coproduct  $\Delta$  corestricts to an algebra map  $\mathcal{L} \rightarrow \mathcal{L} \times_B \mathcal{L}$  where*

$$\mathcal{L} \times_B \mathcal{L} := \left\{ \sum_i p_i \otimes_B q_i \mid \sum_i p_i t(b) \otimes_B q_i = \sum_i p_i \otimes_B q_i s(b), \quad \forall b \in B \right\} \subseteq \mathcal{L} \otimes_B \mathcal{L},$$

*is an algebra via factorwise multiplication.*



(ii) The counit  $\epsilon$  is a left character in the following sense:

$$\epsilon(1_{\mathcal{L}}) = 1_B, \quad \epsilon(s(b)p) = b\epsilon(p), \quad \epsilon(ps(\epsilon(q))) = \epsilon(pq) = \epsilon(pt(\epsilon(q)))$$

for all  $p, q \in \mathcal{L}$  and  $b \in B$ .

Morphisms between left  $B$ -bialgebroids are  $B$ -coring maps which are also algebra maps.

**Definition 3.2.** A left bialgebroid  $\mathcal{L}$  is a left Hopf algebroid ([24], Thm and Def 3.5.) if

$$\lambda : \mathcal{L} \otimes_{B^{op}} \mathcal{L} \rightarrow \mathcal{L} \otimes_B \mathcal{L}, \quad \lambda(p \otimes_{B^{op}} q) = p_{(1)} \otimes_B p_{(2)}q$$

is invertible, where  $\otimes_{B^{op}}$  is induced by  $t$  (so  $pt(b) \otimes_{B^{op}} q = p \otimes_{B^{op}} t(b)q$ , for all  $p, q \in \mathcal{L}$  and  $b \in B$ ) while the  $\otimes_B$  is the standard one (3.1) (so  $t(b)p \otimes_B q = p \otimes_B s(b)q$ ).

In the following we will always call Hopf algebroid instead of left Hopf algebroid. If  $B = k$  then this reduces to the map  $\mathcal{L} \otimes \mathcal{L} \rightarrow \mathcal{L} \otimes \mathcal{L}$  given by  $p \otimes q \mapsto p_{(1)} \otimes p_{(2)}q$  which for a usual Hopf algebra has inverse  $p \otimes q \mapsto p_{(1)} \otimes (Sp_{(2)})q$  if there is an antipode.

**Lemma 3.3.** If there is an invertible linear map  $S : \mathcal{L} \rightarrow \mathcal{L}$ , such that:

- (1)  $S(t(b)X) = S(X)s(b)$ ,  $S^{-1}(s(b)X) = S^{-1}(X)t(b)$
- (2)  $S^{-1}(X_{(2)})_{(1)} \otimes_B S^{-1}(X_{(2)})_{(2)}X_{(1)} = S^{-1}(X) \otimes_B 1$ ,
- (3)  $S(X_{(1)})_{(1)}X_{(2)} \otimes_B S(X_{(1)})_{(2)} = 1 \otimes_B S(X)$ ,

for any  $X, Y \in \mathcal{L}$  and  $b \in B$ . Then  $\mathcal{L}$  is a left Hopf algebroid with  $\lambda^{-1}$  given by

$$\lambda^{-1}(X \otimes_B Y) := S^{-1}(S(X)_{(2)}) \otimes_{B^{op}} S(X)_{(1)}Y.$$

We call  $S$  a left antipode for  $\mathcal{L}$ .

*Proof.* First we check condition (2), (3) and the stated  $\lambda^{-1}$  are well defined. For (3),

$$S(t(b)X_{(1)})_{(1)}X_{(2)} \otimes_B S(t(b)X_{(1)})_{(2)} = S(X_{(1)})_{(1)}s(b)X_{(2)} \otimes_B S(X_{(1)})_{(2)}.$$

Similar for (2). For  $\lambda^{-1}$ ,

$$\begin{aligned} S^{-1}(s(b)S(X)_{(2)}) \otimes_{B^{op}} S(X)_{(1)}Y &= S^{-1}(S(X)_{(2)})t(b) \otimes_{B^{op}} S(X)_{(1)}Y \\ &= S^{-1}(S(X)_{(2)}) \otimes_{B^{op}} t(b)S(X)_{(1)}Y. \end{aligned}$$

Now we are going to show that  $\lambda^{-1}$  is the inverse of  $\lambda$ :

$$\begin{aligned} \lambda(\lambda^{-1}(X \otimes_B Y)) &= \lambda(S^{-1}(S(X)_{(2)}) \otimes_{B^{op}} S(X)_{(1)}Y) \\ &= S^{-1}(S(X)_{(2)})_{(1)} \otimes_B S^{-1}(S(X)_{(2)})_{(2)}S(X)_{(1)}Y \\ &= X \otimes_B Y. \end{aligned}$$

$$\begin{aligned} \lambda^{-1}(\lambda(X \otimes_{B^{op}} Y)) &= \lambda^{-1}(X_{(1)} \otimes_B X_{(2)}Y) \\ &= S^{-1}(S(X_{(1)})_{(2)}) \otimes_{B^{op}} S(X_{(1)})_{(1)}X_{(2)}Y \\ &= X \otimes_{B^{op}} Y. \end{aligned}$$

□

Note that we are not claiming that  $S$  here is unique. We adopt the shorthand

$$X_+ \otimes_{B^{op}} X_- := \lambda^{-1}(X \otimes_B 1) \tag{3.2}$$

*Remark 3.4.* (1) If  $B = k$  then Definition 3.2 reduces to the usual definition of a Hopf algebra. We first observe that

- (i)  $h_{(1)+} \otimes h_{(1)-}h_{(2)} = h \otimes 1$ .
- (ii)  $h_{+(1)} \otimes h_{+(2)}h_- = h \otimes 1$
- (iii)  $h_{+(1)} \otimes h_{+(2)} \otimes h_- = h_{(1)} \otimes h_{(2)+} \otimes h_{(2)-}$
- (iv)  $h_+h_- = \epsilon(h)$



where (i) and (ii) are from applying  $\lambda^{-1} \circ \lambda$  and  $\lambda \circ \lambda^{-1}$  on  $1 \otimes h$ . Then (iv) is the result by applying  $\epsilon \otimes \text{id}$  to (ii). For (iii), apply  $\text{id} \otimes \lambda$  on the right hand side of (iii) to get  $h_{(1)} \otimes h_{(2)} \otimes 1$ . But applying  $\text{id} \otimes \lambda$  on the left hand side of (iii) we get

$$h_{+(1)} \otimes h_{+(2)} \otimes h_{+(3)} h_- = h_{+(1)(1)} \otimes h_{+(1)(2)} \otimes h_{+(2)} h_- = h_{(1)} \otimes h_{(2)} \otimes 1$$

also, where the second steps uses (ii). Since  $\text{id} \otimes \lambda$  is a bijective map, we have (iii). Now set  $S = \epsilon(h_+) h_-$  and check

$$S(h_{(1)}) h_{(2)} = \epsilon(h_{(1)+}) h_{(1)-} a_{(2)} = \epsilon(h),$$

here we use (i) in the second step. On the other hand,

$$h_{(1)} S(h_{(2)}) = a_{(1)} \epsilon(h_{(2)+}) h_{(2)-} = h_{+(1)} \epsilon(h_{+(2)}) h_- = h_+ h_- = \epsilon(h),$$

where the second step uses (iii) and the last step uses (iv).

(2) If  $B = k$  then the conditions in Lemma 3.3 is equivalent to a usual Hopf algebra with invertible antipode. Here (2) and (3) in Lemma 3.3 are the same and say  $S$  is an anticoalgebra maps. By applying  $\epsilon$  one can show that these say that  $S$  obeys half of the antipode axiom and  $S^{-1}$  obeys half of its usual which is equivalent to the other half axiom of  $S$ .

We will need some identities, and we recall from [24, Prop. 3.7] that for a left Hopf algebroid,

$$X_{+(1)} \otimes_B X_{+(2)} X_- = X \otimes_B 1; \quad (3.3)$$

$$X_{(1)+} \otimes_{B^{op}} X_{(1)-} X_{(2)} = X \otimes_{B^{op}} 1; \quad (3.4)$$

$$(XY)_+ \otimes_{B^{op}} (XY)_- = X_+ Y_+ \otimes_{B^{op}} Y_- X_-; \quad (3.5)$$

$$1_+ \otimes_{B^{op}} 1_- = 1 \otimes_{B^{op}} 1; \quad (3.6)$$

$$X_{+(1)} \otimes_B X_{+(2)} \otimes_{B^{op}} X_- = X_{(1)} \otimes_B X_{(2)+} \otimes_{B^{op}} X_{(2)-}; \quad (3.7)$$

$$X_+ \otimes_{B^{op}} X_{-(1)} \otimes_B X_{-(2)} = X_{++} \otimes_{B^{op}} X_- \otimes_B X_{+-}; \quad (3.8)$$

$$X = X_+ t(\epsilon(X_-)); \quad (3.9)$$

$$X_+ X_- = s(\epsilon(X)). \quad (3.10)$$

Since  $\lambda(s(b) \otimes_{B^{op}} 1) = s(b) \otimes_B 1$  and  $\lambda(1 \otimes_{B^{op}} t(b)) = t(b) \otimes_B 1$ , we have

$$s(b)_+ \otimes_{B^{op}} s(b)_- = s(b) \otimes_{B^{op}} 1$$

$$t(b)_+ \otimes_{B^{op}} t(b)_- = 1 \otimes_{B^{op}} t(b) = t(b) \otimes_{B^{op}} 1$$

As a result from (3.5), we get

$$(Xs(b))_+ \otimes_{B^{op}} (Xs(b))_- = X_+ s(b) \otimes_{B^{op}} X_- \quad (3.11)$$

$$(Xt(b))_+ \otimes_{B^{op}} (Xt(b))_- = X_+ \otimes_{B^{op}} t(b) X_- = X_+ t(b) \otimes_{B^{op}} X_- \quad (3.12)$$

$$(s(b)X)_+ \otimes_{B^{op}} (s(b)X)_- = s(b) X_+ \otimes_{B^{op}} X_- \quad (3.13)$$

$$(t(b)X)_+ \otimes_{B^{op}} (t(b)X)_- = X_+ \otimes_{B^{op}} X_- t(b) = t(b) X_+ \otimes_{B^{op}} X_- \quad (3.14)$$

Moreover, since  $X_+ \otimes_{B^{op}} X_- t(b) = \lambda^{-1}(t(b) X \otimes_B 1) = \lambda^{-1}(X \otimes_B s(b)) = X_+ \otimes_{B^{op}} X_- s(b)$ , we have

$$X_+ \otimes_{B^{op}} X_- t(b) = X_+ \otimes_{B^{op}} X_- s(b). \quad (3.15)$$

**3.2. Twisted Hopf algebroids.** Given a left  $B$ -bialgebroid  $\mathcal{L}$ , there is an algebra structure on  ${}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$  with the (convolution) product and unit are given by

$$f \star g(X, Y) := f(X_{(1)}, Y_{(1)}) g(X_{(2)}, Y_{(2)}), \quad \tilde{\epsilon}(X \otimes Y) = \epsilon(XY) \quad (3.16)$$

for all  $X, Y \in \mathcal{L}$  and  $f, g \in {}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$ . The  $B$ -bimodule structure on  $\mathcal{L} \otimes_{B^e} \mathcal{L}$  is just the left  $B^e$  action (so  $b.(X \otimes Y).c = s(b)t(c)X \otimes Y$  for all  $b, c \in B$ ) as induced by the algebra map  $\eta: B^e \rightarrow \mathcal{L}$ .

**Definition 3.5.** [5] Let  $\mathcal{L}$  be a left  $B$ -bialgebroid, a normalised left 2-cocycle on  $\mathcal{L}$  is a convolution invertible element  $\varsigma \in {}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$ , such that

$$\varsigma(X, s(\varsigma(Y_{(1)}, Z_{(1)}))Y_{(2)}Z_{(2)}) = \varsigma(s(\varsigma(X_{(1)}, Y_{(1)}))X_{(2)}Y_{(2)}, Z), \quad \varsigma(1_{\mathcal{L}}, X) = \epsilon(X) = \varsigma(X, 1_{\mathcal{L}})$$

for all  $X, Y, Z \in \mathcal{L}$ .

Similarly, a normalised right 2-cocycle on  $\mathcal{L}$  is a convolution invertible element  $\xi \in {}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$ , such that

$$\xi(X, t(\xi(Y_{(2)}, Z_{(2)}))Y_{(1)}Z_{(1)}) = \xi(t(\xi(X_{(2)}, Y_{(2)}))X_{(1)}Y_{(1)}, Z), \quad \xi(1_{\mathcal{L}}, X) = \epsilon(X) = \xi(X, 1_{\mathcal{L}})$$

for all  $X, Y, Z \in \mathcal{L}$ . A normalised left 2-cocycle is called an invertible normalised 2-cocycle, if its convolution inverse is a normalised right 2-cocycle.

If  $B = k$  the field then a bialgebroid reduces to a usual bialgebra and  $\varsigma$  reduces to a usual Drinfeld cotwist cocycle in the sense of [18]. However, unlike a 2-cocycle on bialgebra, for any normalised left 2-cocycle, its convolution inverse is not automatically a normalised right 2-cocycle. The corresponding ‘Drinfeld cotwist’ for bialgebroids is as follows.

**Lemma 3.6.** Let  $\mathcal{L}$  be a left  $B$ -bialgebroid and  $\varsigma \in {}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$  be an invertible normalised 2-cocycle, with inverse  $\varsigma^{-1}$ , then

$$\varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}Y_{(2)}, Z_{(2)}) = \varsigma(Xs(\varsigma^{-1}(Y_{(1)}, Z)), Y_{(2)}) \quad (3.17)$$

$$\varsigma(X_{(1)}Y_{(1)}, Z_{(1)})\varsigma^{-1}(X_{(2)}, Y_{(2)}Z_{(2)}) = \varsigma^{-1}(X, s(\varsigma(Y_{(2)}, Z))Y_{(1)}) \quad (3.18)$$

*Proof.* Since the inverse of  $\varsigma$  satisfies

$$\varsigma^{-1}(X, t(\varsigma^{-1}(Y_{(2)}, Z_{(2)}))Y_{(1)}Z_{(1)}) = \varsigma^{-1}(t(\varsigma^{-1}(X_{(2)}, Y_{(2)}))X_{(1)}Y_{(1)}, Z).$$

We will have a well defined equation:

$$\begin{aligned} & \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}, t(\varsigma^{-1}(Y_{(3)}, Z_{(3)}))Y_{(2)}Z_{(2)})\varsigma(X_{(3)}, Y_{(4)}) \\ & = \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(t(\varsigma^{-1}(X_{(3)}, Y_{(3)}))X_{(2)}Y_{(2)}, Z)\varsigma(X_{(4)}, Y_{(4)}) \end{aligned}$$

The left hand side is equal to

$$\begin{aligned} & \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}, t(\varsigma^{-1}(Y_{(3)}, Z_{(3)}))Y_{(2)}Z_{(2)})\varsigma(X_{(3)}, Y_{(4)}) \\ & = \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}, Y_{(2)}Z_{(2)})\varsigma(X_{(3)}s(\varsigma^{-1}(Y_{(3)}, Z_{(3)})), Y_{(4)}) \\ & = \epsilon(X_{(1)}Y_{(1)}Z_{(1)})\varsigma(X_{(2)}s(\varsigma^{-1}(Y_{(2)}, Z_{(2)})), Y_{(3)}) \\ & = \epsilon(X_{(1)}Y_{(1)}t(\epsilon(Z_{(1)})))\varsigma(X_{(2)}s(\varsigma^{-1}(Y_{(2)}, Z_{(2)})), Y_{(3)}) \\ & = \epsilon(X_{(1)}Y_{(1)})\varsigma(X_{(2)}s(\varsigma^{-1}(Y_{(2)}s(\epsilon(Z_{(1)}))), Z_{(2)})), Y_{(3)}) \\ & = \epsilon(X_{(1)}t(\epsilon(Y_{(1)})))\varsigma(X_{(2)}s(\varsigma^{-1}(Y_{(2)}, Z)), Y_{(3)}) \\ & = \epsilon(X_{(1)})\varsigma(X_{(2)}s(\epsilon(Y_{(1)}))s(\varsigma^{-1}(Y_{(2)}, Z))), Y_{(3)}) \\ & = \epsilon(X_{(1)})\varsigma(X_{(2)}s(\varsigma^{-1}(\epsilon(Y_{(1)})Y_{(2)}, Z))), Y_{(3)}) \\ & = \epsilon(X_{(1)})\varsigma(X_{(2)}s(\varsigma^{-1}(Y_{(1)}, Z))), Y_{(2)}) \\ & = \varsigma(Xs(\varsigma^{-1}(Y_{(1)}, Z))), Y_{(2)}. \end{aligned}$$

The right hand side is equal to

$$\begin{aligned} & \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(t(\varsigma^{-1}(X_{(3)}, Y_{(3)}))X_{(2)}Y_{(2)}, Z_{(2)})\varsigma(X_{(4)}, Y_{(4)}) \\ & = \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}Y_{(2)}, Z_{(2)})\varsigma^{-1}(X_{(3)}, Y_{(3)})\varsigma(X_{(4)}, Y_{(4)}) \\ & = \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}Y_{(2)}, Z_{(2)})\epsilon(X_{(3)}Y_{(3)}) \\ & = \varsigma(X_{(1)}, Y_{(1)}Z_{(1)})\varsigma^{-1}(X_{(2)}Y_{(2)}, Z_{(2)}) \end{aligned}$$

So we have (3.17). Similarly, we have (3.18).  $\square$

Using such cotwists, it shown in [5] that  $B$ -coring  $\mathcal{L}$  with the twisted product [5]

$$X \cdot_{\varsigma} Y := s(\varsigma(X_{(1)}, Y_{(1)}))t(\varsigma^{-1}(X_{(3)}, Y_{(3)}))X_{(2)}Y_{(2)}, \quad (3.19)$$

for all  $X, Y \in \mathcal{L}$ , and the original coproduct, counit,  $s, t$  constitute a left bilgebroid  $\mathcal{L}^{\varsigma}$ .

**Theorem 3.7.** *Let  $\mathcal{L}$  be a left Hopf algebroid and  $\varsigma \in {}_B \text{Hom}_B(\mathcal{L} \otimes_{B^e} \mathcal{L}, B)$  be an invertible normalised 2-cocycle, with inverse  $\varsigma^{-1}$ , then the left bialgebroid  $\mathcal{L}^{\varsigma}$  with twisted product (3.19) is a left Hopf algebroid, with*

$$(\lambda^{\varsigma})^{-1}(X \otimes_B Y) = t(\varsigma(X_{(1)+(2)}, X_{(1)-(3)}))X_{(1)+(1)} \otimes_{B^{op}} s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}Y.$$

*Proof.* We need to show that  $\lambda^{\varsigma} : X \otimes_{B^{op}} Y \mapsto X_{(1)} \otimes_B X_{(2)} \cdot_{\varsigma} Y$  is an invertible map. We claim the inverse of  $\lambda^{\varsigma}$  is as stated. We show that this is a well defined over the balanced tensor product of  $X_{(1)} \otimes_B X_{(2)}$ , with other details left similarly to the reader. Thus,

$$\begin{aligned} & t(\varsigma((t(b)X_{(1)})_{+(2)}, (t(b)X_{(1)})_{-(3)}))(t(b)X_{(1)})_{+(1)} \otimes_{B^{op}} s(\varsigma^{-1}((t(b)X_{(1)})_{-(1)}, X_{(2)}))(t(b)X_{(1)})_{-(2)} \\ &= t(\varsigma(X_{(1)+(2)}, (X_{(1)}-t(b))_{(3)}))X_{(1)+(1)} \otimes_{B^{op}} s(\varsigma^{-1}((X_{(1)}-t(b))_{(1)}, X_{(2)}))(X_{(1)}-t(b))_{(2)} \\ &= t(\varsigma(X_{(1)+(2)}, (X_{(1)}-s(b))_{(3)}))X_{(1)+(1)} \otimes_{B^{op}} s(\varsigma^{-1}((X_{(1)}-s(b))_{(1)}, X_{(2)}))(X_{(1)}-s(b))_{(2)} \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(3)}))X_{(1)+(1)} \otimes_{B^{op}} s(\varsigma^{-1}(X_{(1)-(1)}, s(b)X_{(2)}))X_{(1)-(2)}, \end{aligned}$$

where the 1st step uses (3.14), the 2nd step uses (3.15). Next, it is sufficient to check  $(\lambda^{\varsigma})^{-1} \circ \lambda^{\varsigma}$  and  $\lambda^{\varsigma} \circ (\lambda^{\varsigma})^{-1}$  on  $X \otimes_{B^{op}} 1$  and  $X \otimes_B 1$ . On the one side we have:

$$\begin{aligned} & (\lambda^{\varsigma})^{-1} \circ \lambda^{\varsigma}(X \otimes_{B^{op}} 1) \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(3)}))X_{(1)+(1)} \otimes_{B^{op}} (s(\varsigma(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}) \cdot_{\varsigma} X_{(3)} \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(5)}))X_{(1)+(1)} \\ & \quad \otimes_{B^{op}} s(\varsigma(s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}, X_{(3)}))t(\varsigma^{-1}(X_{(1)-(4)}, X_{(5)}))X_{(1)-(3)}X_{(4)} \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(5)}))X_{(1)+(1)} \\ & \quad \otimes_{B^{op}} s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)})\varsigma(X_{(1)-(2)}, X_{(3)}))t(\varsigma^{-1}(X_{(1)-(4)}, X_{(5)}))X_{(1)-(3)}X_{(4)} \\ &= t(\varsigma(X_{(1)++(2)}, X_{(1)+-(2)}))X_{(1)++(1)} \otimes_{B^{op}} t(\varsigma^{-1}(X_{(1)+-(1)}, X_{(3)}))X_{(1)-}X_{(2)} \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(2)}))X_{(1)+(1)} \otimes_{B^{op}} t(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)})) \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(2)}))X_{(1)+(1)}t(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)})) \otimes_{B^{op}} 1 \\ &= t(\varsigma(X_{(1)+(2)}s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)})), X_{(1)-(2)}))X_{(1)+(1)} \otimes_{B^{op}} 1 \\ &= t(\varsigma(X_{(1)+(2)}, X_{(1)-(1)}X_{(2)})\varsigma^{-1}(X_{(1)+(3)}X_{(1)-(2)}, X_{(3)}))X_{(1)+(1)} \otimes_{B^{op}} 1 \\ &= t(\varsigma(X_{(1)++(2)}, X_{(1)-}X_{(2)})\varsigma^{-1}(X_{(1)++(3)}X_{(1)+-}, X_{(3)}))X_{(1)++(1)} \otimes_{B^{op}} 1 \\ &= t(\epsilon(X_{(1)+(2)})\varsigma^{-1}(X_{(1)+(3)}X_{(1)-}, X_{(2)}))X_{(1)+(1)} \otimes_{B^{op}} 1 \\ &= t(\varsigma^{-1}(X_{(1)+(2)}X_{(1)-}, X_{(2)}))X_{(1)+(1)} \otimes_{B^{op}} 1 \\ &= t(\epsilon(X_{(2)}))X_{(1)} \otimes_{B^{op}} 1 \\ &= X \otimes_{B^{op}} 1, \end{aligned}$$

where the 4th and 9th steps use (3.8), the 5th and 10th steps use (3.4), the 8th step use (3.17), the 12th step uses (3.3). On another side we have

$$\begin{aligned} & \lambda^{\varsigma} \circ (\lambda^{\varsigma})^{-1}(X \otimes_B 1) \\ &= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)})) \\ & \quad t(\varsigma^{-1}(t(\varsigma(X_{(1)+(5)}, X_{(1)-(5)}))X_{(1)+(4)}, X_{(1)-(4)}))X_{(1)+(3)}X_{(1)-(3)} \\ &= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}))X_{(1)+(3)}X_{(1)-(3)} \\ &= X_{(1)++(1)} \otimes_B s(\varsigma(X_{(1)++(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}))X_{(1)++(3)}X_{(1)+-} \\ &= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}))X_{(1)+(3)+}X_{(1)+(3)-} \end{aligned}$$

$$\begin{aligned}
&= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)}))s(\epsilon(X_{(1)+(3)})) \\
&= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, s(\varsigma^{-1}(X_{(1)-(1)}, X_{(2)}))X_{(1)-(2)})) \\
&= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, X_{(1)-(1)}X_{(2)})\varsigma^{-1}(X_{(1)+(3)}X_{(1)-(2)}, X_{(3)})) \\
&= X_{(1)+(1)} \otimes_B s(\varsigma(X_{(1)+(2)}, X_{(1)-}X_{(2)})\varsigma^{-1}(X_{(1)+(3)}X_{(1)+}, X_{(3)})) \\
&= X_{(1)+(1)} \otimes_B s(\epsilon(X_{(1)+(2)}))\varsigma^{-1}(X_{(1)+(3)}X_{(1)+}, X_{(2)}) \\
&= X_{(1)+(1)} \otimes_B s(\varsigma^{-1}(X_{(1)+(2)}X_{(1)-}, X_{(2)})) \\
&= X_{(1)} \otimes_B s(\epsilon(X_{(2)})) \\
&= t(\epsilon(X_{(2)}))X_{(1)} \otimes_B 1 \\
&= X \otimes_B 1
\end{aligned}$$

where the 3rd and 8th steps use (3.8), the 4th step uses (3.7), the 5th step uses (3.10), the 7th step use (3.17), the 11th step use (3.3).  $\square$

*Remark 3.8.* The right handed version of the theorem also holds. Thus, for an invertible normalised 2-cocycle  $\varsigma$  on a left bialgebroid  $\mathcal{L}$ , if the map  $\mu : \mathcal{L} \otimes_B \mathcal{L} \rightarrow \mathcal{L} \otimes_B \mathcal{L}$  given by

$$\mu(X \otimes_B Y) := Y_{(1)}X \otimes_B Y_{(2)}$$

is invertible, then  $\mu^\varsigma : \mathcal{L}^\varsigma \otimes_B \mathcal{L}^\varsigma \rightarrow \mathcal{L}^\varsigma \otimes_B \mathcal{L}^\varsigma$  given by  $\mu(X \otimes_B Y) := Y_{(1)} \cdot_\varsigma X \otimes_B Y_{(2)}$  is invertible as well. The balanced tensor product on the left side is given by  $X \otimes_B s(b)Y = Xs(b) \otimes_B Y$ , The balanced tensor product on the right side is given by  $X \otimes_B s(b)Y = t(b)X \otimes_B Y$ . More explicitly, if the image of  $\mu^{-1}(1 \otimes_B X)$  is denoted by  $X_{(-)} \otimes_B X_{(+)}$ , then

$$(\mu^\varsigma)^{-1}(1 \otimes_B X) = t(\varsigma(X_{(2)(-)(3)}, X_{(1)}))X_{(2)(-)(2)} \otimes_B s(\varsigma^{-1}(X_{(2)(+)(1)}, X_{(2)(-)(1)}))X_{(2)(+)(2)}.$$

**3.3. Cocycle Hopf algebroids.** Associated to any cleft cocycle extension we have a Hopf algebroid as follows.

**Proposition 3.9.** *Associated to a cocycle cross product  $B\#_\sigma H$  we have a cocycle Hopf algebroid  $B^e\#_\sigma H$  built on  $B^e \otimes H$  with with product*

$$(b \otimes b' \#_\sigma h)(c \otimes c' \#_\sigma g) = b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)}) \otimes c'(Sg_{(4)} \triangleright b')\sigma(Sg_{(3)}, Sh_{(4)})\#_\sigma h_{(3)}g_{(2)}$$

and source and target maps

$$s(b) = b \otimes 1\#_\sigma 1; \quad t(b) = 1 \otimes b\#_\sigma 1; \quad \eta(b \otimes b') = s(b)t(b') = b \otimes b'\#_\sigma 1$$

The coproduct is

$$\Delta(b \otimes b' \#_\sigma h) = (b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)})\#_\sigma h_{(1)}) \otimes_B (1 \otimes b' \#_\sigma h_{(4)})$$

where the  $B$ -bimodule structure is

$$c \triangleright (b \otimes b' \#_\sigma h) \triangleleft c' = cb \otimes b'(S(h_{(2)}) \triangleright c')\#_\sigma h_{(1)}$$

The counit is

$$\epsilon(b \otimes b' \#_\sigma h) = b(h_{(1)} \triangleright b')\sigma(h_{(2)}, S(h_{(3)})).$$

The inverse of  $\lambda$  is given by

$$\lambda^{-1}(X \otimes_B Y) = (b \otimes 1 \otimes h_{(1)}) \otimes_{B^{op}} (b' \otimes \sigma^{-1}(S^2(h_{(4)}), S(h_{(3)}))\#_\sigma S(h_{(5)}))Y$$

*Proof.* First, we can show  $B^e\#_\sigma H$  is a  $B$ -coring and  $B^e$ -ring.

It is not hard to see both the coproduct and the counit are  $B$ -bimodule map.

$$(\epsilon \otimes_B \text{id})(\Delta(b \otimes b' \#_\sigma h)) = b(h_{(1)} \triangleright \sigma^{-1}(S(h_{(4)}), h_{(5)}))\sigma(h_{(2)}, S(h_{(3)})) \otimes b' \#_\sigma h_{(6)} = b \otimes b' \#_\sigma h,$$

and

$$(\text{id} \otimes_B \epsilon)(\Delta(b \otimes b' \#_\sigma h)) = b \otimes \sigma^{-1}(S(h_{(3)}), h_{(4)})(S(h_{(2)}) \triangleright ((h_{(5)} \triangleright b')\sigma(h_{(6)}, S(h_{(7)})))\#_\sigma h_{(1)}$$

$$\begin{aligned}
 &= b \otimes (S(h_{(4)})h_{(5)} \triangleright b')\sigma^{-1}(S(h_{(3)}), h_{(6)})(S(h_{(2)}) \triangleright \sigma(h_{(7)}, S(h_{(8)})))\#_{\sigma} h_{(1)} \\
 &= b \otimes b'\sigma^{-1}(S(h_{(5)}), h_{(6)})\sigma(S(h_{(4)}), h_{(7)})\sigma(S(h_{(3)})h_{(8)}, S(h_{(11)})) \\
 &\quad \sigma^{-1}(S(h_{(2)}), h_{(9)})\sigma(S(h_{(10)}))\#_{\sigma} h_{(1)} \\
 &= b \otimes b'\#_{\sigma} h,
 \end{aligned}$$

where we use Lemma 2.1 repeatedly.

$$\begin{aligned}
 &(\text{id} \otimes_B \Delta)(\Delta(b \otimes b'\#_{\sigma} h)) \\
 &= (b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)}))\#_{\sigma} h_{(1)} \otimes_B (1 \otimes \sigma^{-1}(S(h_{(5)}), h_{(6)}))\#_{\sigma} h_{(4)} \otimes_B (1 \otimes b'\#_{\sigma} h_{(7)}) \\
 &= (\Delta \otimes_B \text{id})(\Delta(b \otimes b'\#_{\sigma} h)).
 \end{aligned}$$

For the  $B^e$  structure, we can see  $\eta$  is an algebra map. It is a direct computation that the product is associative, since  $\sigma$  is a 2-cocycle. Before we show the coproduct is an algebra map, we first check the image of coproduct belongs to the Takeuchi product. Let  $X = b \otimes b' \otimes h$  and  $d \in B$ ,

$$\begin{aligned}
 X_{(1)} \otimes_B X_{(2)}s(d) &= (b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)}))\#_{\sigma} h_{(1)} \otimes_B (h_{(4)} \triangleright d \otimes b'\#_{\sigma} h_{(5)}) \\
 &= t(h_{(4)} \triangleright d)(b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)}))\#_{\sigma} h_{(1)} \otimes_B (1 \otimes b'\#_{\sigma} h_{(5)}) \\
 &= b \otimes \sigma^{-1}(S(h_{(3)}), h_{(4)})S(h_{(2)}) \triangleright (h_{(5)} \triangleright d)\#_{\sigma} h_{(1)} \otimes_B (1 \otimes b'\#_{\sigma} h_{(6)}) \\
 &= (b \otimes d\sigma^{-1}(S(h_{(2)}), h_{(3)}))\#_{\sigma} h_{(1)} \otimes_B 1 \otimes b'\#_{\sigma} h_{(4)} \\
 &= X_{(1)}t(d) \otimes_B X_{(2)}.
 \end{aligned}$$

Finally, by direct computation one can show that the coproduct is an algebra map and  $\lambda^{-1}$  is the inverse of  $\lambda$ . We omit details since this follows from a more general result Theorem 4.4 later.  $\square$

Next we show that  $B^e \#_{\sigma} H$  has an antipode in the sense of Lemma 3.3 at least in the associative type case.

**Proposition 3.10.** *Let  $B \#_{\sigma} H$  be an associative type, then*

$$\begin{aligned}
 S(b \otimes b'\#_{\sigma} h) &:= b' \otimes \sigma^{-1}(S^2(h_{(2)}), S(h_{(1)}))S^2(h_{(3)}) \triangleright b\#_{\sigma} S(h_{(4)}) \\
 S^{-1}(b \otimes b'\#_{\sigma} h) &:= (h_{(2)} \triangleright b')\sigma(h_{(3)}, S(h_{(4)})) \otimes b\#_{\sigma} S^{-1}(h_{(1)}).
 \end{aligned}$$

is an antipode for the Hopf algebroid  $B^e \#_{\sigma} H$ . Here  $S$  on the right is the antipode of  $H$ .

*Proof.* Let  $X = b \otimes b'\#_{\sigma} h$ , then we can define a linear map  $S : B^e \#_{\sigma} H \rightarrow B^e \#_{\sigma} H$  as stated with the inverse given by the formula stated. It is not hard to see the images of  $S^{-1}$  and  $S^{-1}$  are contained in  $B^e \#_{\sigma} H$ . We also check

$$\begin{aligned}
 S^{-1}(S(X)) &= S^{-1}(b' \otimes \sigma^{-1}(S^2(h_{(2)}), S(h_{(1)}))S^2(h_{(3)}) \triangleright b\#_{\sigma} S(h_{(4)})) \\
 &= g_{(2)} \triangleright (\sigma^{-1}(S(g_{(6)}), g_{(7)})S(g_{(5)}) \triangleright b)\sigma(g_{(3)}, S(g_{(4)})) \otimes_b \#_{\sigma} S^{-1}(g_{(1)}) \\
 &= g_{(2)} \triangleright ((S(g_{(8)}) \triangleright b)g_{(3)} \triangleright (\sigma^{-1}(S(g_{(6)}), g_{(7)}))\sigma(g_{(4)}, S(g_{(5)}))) \otimes_b \#_{\sigma} S^{-1}(g_{(1)}) \\
 &= b \otimes b'\#_{\sigma} S^{-1}(g) = X.
 \end{aligned}$$

In the 2nd step, we define  $g := S(h)$ . In the 3rd step we use Lemma 2.4. In the 4th step we use (4) of Lemma 2.1. On the other side,

$$\begin{aligned}
 S(S^{-1}(X)) &= S((h_{(2)} \triangleright b')\sigma(h_{(3)}, S(h_{(4)})) \otimes b\#_{\sigma} S^{-1}(h_{(1)})) \\
 &= b \otimes \sigma^{-1}(S(h_{(3)}), h_{(4)})S(h_{(2)}) \triangleright ((h_{(5)} \triangleright b')\sigma(h_{(6)}, S(h_{(7)})))\#_{\sigma} h_{(1)} \\
 &= b \otimes S(h_{(4)}) \triangleright ((h_{(5)} \triangleright b')\sigma(h_{(6)}, S(h_{(7)})))\sigma^{-1}(S(h_{(2)}), h_{(3)})\#_{\sigma} h_{(1)} \\
 &= b \otimes b'\sigma^{-1}(S(h_{(3)}), h_{(4)})S(h_{(2)}) \triangleright \sigma(h_{(5)}, S(h_{(6)}))\#_{\sigma} h_{(1)}
 \end{aligned}$$

$$=b \otimes b' \#_{\sigma} h.$$

In the 3rd and 4th steps we use Lemma 2.4. In the 5th step we use (6) of Lemma 2.1. Now we are going to check  $S$  and its inverse satisfy the conditions of Lemma 3.3. By direct computation

$$S(t(d)X) = b'(S(h_{(4)}) \triangleright d) \otimes \sigma^{-1}(S^2(h_{(2)}), S(h_{(1)}))S^2(h_{(3)}) \triangleright b \#_{\sigma} S^2(h_{(2)}) = S(X)s(d),$$

Similarly, we have  $S^{-1}(s(b)X) = S(X)t(b)$ . For condition (2) of Lemma 3.3, we have:

$$\begin{aligned} & S^{-1}(X_{(2)})_{(1)} \otimes_B S^{-1}(X_{(2)})_{(1)} X_{(1)} \\ &= (h_{(8)} \triangleright b') \sigma(h_{(9)}, S(h_{(10)})) \otimes \sigma^{-1}(h_{(6)}, S^{-1}(h_{(5)})) \#_{\sigma} S^{-1}(h_{(7)}) \\ & \quad \otimes_B (1 \otimes 1 \#_{\sigma} S^{-1}(h_{(4)})) (b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)})) \#_{\sigma} h_{(1)} \\ &= (h_{(7)} \triangleright b') \sigma(h_{(8)}, S(h_{(9)})) \otimes \sigma^{-1}(h_{(5)}, S^{-1}(h_{(4)})) \#_{\sigma} S^{-1}(h_{(6)}) \\ & \quad \otimes_B s((S^{-1}(h_{(3)}) \triangleright b) \sigma(S^{-1}(h_{(2)}), h_{(1)})) \\ &= (h_{(8)} \triangleright b') \sigma(h_{(9)}, S(h_{(10)})) \otimes \sigma^{-1}(h_{(5)}, S^{-1}(h_{(4)})) h_{(6)} \triangleright ((S^{-1}(h_{(3)}) \triangleright b) \sigma(S^{-1}(h_{(2)}), h_{(1)})) \\ & \quad \#_{\sigma} S^{-1}(h_{(7)}) \otimes_B 1_{\mathcal{L}} \\ &= (h_{(8)} \triangleright b') \sigma(h_{(9)}, S(h_{(10)})) \otimes h_{(4)} \triangleright ((S^{-1}(h_{(3)}) \triangleright b) \sigma(S^{-1}(h_{(2)}), h_{(1)})) \sigma^{-1}(h_{(6)}, S^{-1}(h_{(5)})) \\ & \quad \#_{\sigma} S^{-1}(h_{(7)}) \otimes_B 1_{\mathcal{L}} \\ &= (h_{(7)} \triangleright b') \sigma(h_{(8)}, S(h_{(9)})) \otimes b(h_{(3)} \triangleright \sigma(S^{-1}(h_{(2)}), h_{(1)})) \sigma^{-1}(h_{(5)}, S^{-1}(h_{(4)})) \#_{\sigma} S^{-1}(h_{(6)}) \otimes_B 1_{\mathcal{L}} \\ &= (h_{(7)} \triangleright b') \sigma(h_{(8)}, S(h_{(9)})) \otimes b \sigma^{-1}(h_{(4)}, S^{-1}(h_{(3)})) (h_{(5)} \triangleright \sigma(S^{-1}(h_{(2)}), h_{(1)})) \#_{\sigma} S^{-1}(h_{(6)}) \otimes_B 1_{\mathcal{L}} \\ &= (h_{(2)} \triangleright b') \sigma(h_{(3)}, S(h_{(4)})) \otimes b \#_{\sigma} S^{-1}(h_{(1)}) \otimes_B 1_{\mathcal{L}} \\ &= S^{-1}(X) \otimes_B 1_{\mathcal{L}} \end{aligned}$$

where the 4th and 6th steps use Lemma 2.4, the 7th step use (6) of Lemma 2.1. For condition (3),

$$\begin{aligned} & S(X_{(1)})_{(1)} X_{(2)} \otimes_B S(X_{(1)})_{(2)} \\ &= (\sigma^{-1}(S(h_{(8)}), h_{(9)})) \otimes \sigma^{-1}(S^2(h_{(6)}), S(h_{(5)})) \#_{\sigma} S(h_{(7)}) (1 \otimes b' \#_{\sigma} h_{(10)}) \\ & \quad \otimes_B 1 \otimes \sigma^{-1}(S^2(h_{(2)}), h_{(1)}) S^2(h_{(3)}) \triangleright b \#_{\sigma} S(h_{(4)}) \\ &= \sigma^{-1}(S(h_{(10)}), h_{(11)}) \sigma(S(h_{(9)}), h_{(12)}) \otimes b'(S(h_{(15)}) \triangleright \sigma^{-1}(S^2(h_{(6)}), h_{(5)})) \sigma(S(h_{(14)}), S^2(h_{(7)})) \\ & \quad \#_{\sigma} S(h_{(8)}) h_{(13)} \otimes_B 1 \otimes \sigma^{-1}(S^2(h_{(2)}), h_{(1)}) S^2(h_{(3)}) \triangleright b \#_{\sigma} S(h_{(4)}) \\ &= 1 \otimes b'(S(h_{(9)}) \triangleright \sigma^{-1}(S^2(h_{(6)}), h_{(5)})) \sigma(S(h_{(8)}), S^2(h_{(7)})) \#_{\sigma} 1 \\ & \quad \otimes_B 1 \otimes \sigma^{-1}(S^2(h_{(2)}), h_{(1)}) S^2(h_{(3)}) \triangleright b \#_{\sigma} S(h_{(4)}) \\ &= t(b') \otimes_B 1 \otimes \sigma^{-1}(S^2(h_{(2)}), h_{(1)}) S^2(h_{(3)}) \triangleright b \#_{\sigma} S(h_{(4)}) \\ &= 1_{\mathcal{L}} \otimes_B b' \otimes \sigma^{-1}(S^2(h_{(2)}), h_{(1)}) S^2(h_{(3)}) \triangleright b \#_{\sigma} S(h_{(4)}) \\ &= 1_{\mathcal{L}} \otimes_B S(X), \end{aligned}$$

where the 4th step uses Lemma 2.4.  $\square$

**3.4. Cocycle Hopf algebroid in the associative type case.** We now this class of cocycle Hopf algebroids to illustrate our twisting theory.

**Lemma 3.11.** *If  $B \#_{\sigma} H$  is a associative type, then the linear map  $\tilde{\sigma} \in {}_B \text{Hom}_B(B^e \#_{\sigma} H \otimes_{B^e} B^e \#_{\sigma} H)$  given by*

$$\tilde{\sigma}(b \otimes b' \#_{\sigma} h, c \otimes c' \#_{\sigma} g) := b(h_{(1)} \triangleright c) \sigma(h_{(2)}, g_{(1)}) ((h_{(3)} g_{(2)}) \triangleright c') (h_{(4)} \triangleright b'),$$

for all  $b \otimes b' \#_{\sigma} h, c \otimes c' \#_{\sigma} g \in B^e \#_{\sigma} H$  is an invertible normalised 2-cocycle.

*Proof.* Since  $B\#_{\sigma}H$  is a associative type, we know  $B$  is a left  $H$ -module algebra, which ensures the smash product  $B\#H$  is well defined. Let  $X = b \otimes b'\#_{\sigma}h$ ,  $Y = c \otimes c'\#_{\sigma}g$  and  $Z = d \otimes d'\#_{\sigma}f$  be three elements in  $B^e\#_{\sigma}H$ . First we can show this cocycle is well defined over the balanced tensor over  $B^e$ : On the one hand we have

$$\begin{aligned}\tilde{\sigma}(X\eta(d \otimes d'), Y) &= \tilde{\sigma}((b \otimes b'\#_{\sigma}h)\eta(d \otimes d'), c \otimes c'\#_{\sigma}g) \\ &= b(h_{(1)} \triangleright d)(h_{(2)} \triangleright c)\sigma(h_{(3)}, g_{(1)})((h_{(4)}g_{(2)} \triangleright c')(h_{(5)} \triangleright (d'b'))),\end{aligned}$$

for all  $d, d' \in B$ ; On the other hand

$$\begin{aligned}\tilde{\sigma}(X, \eta(d \otimes d')Y) &= \tilde{\sigma}((b \otimes b'\#_{\sigma}h), \eta(d \otimes d')(c \otimes c'\#_{\sigma}g)) \\ &= \tilde{\sigma}(b \otimes b'\#_{\sigma}h, dc \otimes c'(S(g_{(2)}) \triangleright d')\#_{\sigma}g_{(1)}) \\ &= b(h_{(1)} \triangleright (dc))\sigma(h_{(2)}, g_{(1)})((h_{(3)}g_{(2)} \triangleright (c'(S(g_{(3)}) \triangleright d')))(h_{(4)} \triangleright b')) \\ &= b(h_{(1)} \triangleright d)(h_{(2)} \triangleright c)\sigma(h_{(3)}, g_{(1)})((h_{(4)}g_{(2)} \triangleright c')(h_{(5)} \triangleright (d'b')).\end{aligned}$$

The inverse  $\tilde{\sigma}^{-1}$  is given by

$$\tilde{\sigma}^{-1}(X, Y) := b(h_{(1)} \triangleright c)\sigma^{-1}(h_{(2)}, g_{(1)})((h_{(3)}g_{(2)} \triangleright c')(h_{(4)} \triangleright b')).$$

We can see that

$$\begin{aligned}\tilde{\sigma}^{-1} \star \tilde{\sigma}(X, Y) &= \tilde{\sigma}^{-1} \star \tilde{\sigma}(b \otimes b'\#_{\sigma}h, c \otimes c'\#_{\sigma}g) \\ &= b(h_{(1)} \triangleright c)\sigma^{-1}(h_{(2)}, g_{(1)})\sigma(h_{(3)}, g_{(2)})((h_{(4)}g_{(3)} \triangleright c')(h_{(5)} \triangleright b')) \\ &= \epsilon((b \otimes b'\#_{\sigma}h)(c \otimes c'\#_{\sigma}g)) \\ &= \tilde{\epsilon}(X, Y),\end{aligned}$$

where  $\tilde{\epsilon}$  is the unit in the algebra  ${}_B\text{Hom}_B(B^e\#_{\sigma}H \otimes_{B^e} B^e\#_{\sigma}H, B)$ . Similarly, we can also see  $\tilde{\sigma} \star \tilde{\sigma}^{-1} = \tilde{\epsilon}$ .

It is clear that  $\tilde{\sigma}$  is left  $B$ -linear, we can also show that  $\tilde{\sigma}$  is also right  $B$ -linear:

$$\begin{aligned}\tilde{\sigma}(t(b'')X, Y) &= \tilde{\sigma}(t(b'')(b \otimes b'\#_{\sigma}h, c \otimes c'\#_{\sigma}g)) \\ &= \tilde{\sigma}(b \otimes b'(S(h_{(2)}) \triangleright b'')\#_{\sigma}h_{(1)}, c \otimes c'\#_{\sigma}g) \\ &= b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)})((h_{(3)}g_{(2)} \triangleright c')(h_{(4)} \triangleright (b'(S(h_{(5)}) \triangleright b''))) \\ &= b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)})((h_{(3)}g_{(2)} \triangleright c')(h_{(4)} \triangleright b')b'') \\ &= \tilde{\sigma}(X, Y)b'',\end{aligned}$$

for all  $b'' \in B$ . Now let's show the cocycle condition of  $\tilde{\sigma}$ . On the one hand we have:

$$\begin{aligned}\tilde{\sigma}(X, s(\tilde{\sigma}(Y_{(1)}, Z_{(1)}))Y_{(2)}Z_{(2)}) &= \tilde{\sigma}(b \otimes b'\#_{\sigma}h, c(g_{(1)} \triangleright d)\sigma(g_{(2)}, f_{(1)}) \otimes d'(S(f_{(3)}) \triangleright c')\#_{\sigma}g_{(3)}f_{(2)}) \\ &= b(h_{(1)} \triangleright (c(g_{(1)} \triangleright d)\sigma(g_{(2)}, f_{(1)})))\sigma(h_{(2)}, g_{(3)}f_{(2)})((h_{(3)}g_{(4)}f_{(3)} \triangleright (d'(S(f_{(4)}) \triangleright c')))(h_{(4)} \triangleright b')).\end{aligned}$$

On the other hand,

$$\begin{aligned}\tilde{\sigma}(s(\tilde{\sigma}(X_{(1)}, Y_{(1)}))X_{(2)}Y_{(2)}, Z) &= \tilde{\sigma}(b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)}) \otimes c'(S(g_{(3)}) \triangleright b')\#_{\sigma}h_{(3)}g_{(2)}, d \otimes d'\#_{\sigma}f) \\ &= b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)})((h_{(3)}g_{(2)} \triangleright d)\sigma(h_{(4)}g_{(3)}, f_{(1)})(h_{(5)}g_{(4)}f_{(2)} \triangleright d') \\ &\quad ((h_{(6)}g_{(5)} \triangleright (c'(S(g_{(6)}) \triangleright b')))).\end{aligned}$$

Comparing the results on both hand sides, it is sufficient to show

$$h_{(1)} \triangleright ((g_{(1)} \triangleright d)\sigma(g_{(2)}, f_{(1)}))\sigma(h_{(2)}, g_{(3)}f_{(2)}) = \sigma(h_{(1)}, g_{(1)})((h_{(2)}g_{(2)} \triangleright d)\sigma(h_{(3)}g_{(3)}, f)).$$

We can see the left hand side is

$$\begin{aligned}h_{(1)} \triangleright ((g_{(1)} \triangleright d)\sigma(g_{(2)}, f_{(1)}))\sigma(h_{(2)}, g_{(3)}f_{(2)}) &= (h_{(1)} \triangleright (g_{(1)} \triangleright d))(h_{(2)} \triangleright \sigma(g_{(2)}, f_{(1)}))\sigma(h_{(3)}, g_{(3)}f_{(2)})\end{aligned}$$



$$\begin{aligned}
&= (h_{(1)} \triangleright (g_{(1)} \triangleright d))\sigma(h_{(2)}, g_{(2)})\sigma(h_{(3)}g_{(3)}, f) \\
&= \sigma(h_{(1)}, g_{(1)})((h_{(2)}g_{(2)}) \triangleright d)\sigma(h_{(3)}g_{(3)}, f).
\end{aligned}$$

Finally, we can show the normalisation condition:

$$\tilde{\sigma}(X, 1) = b(h_{(1)} \triangleright b') = \epsilon(X), \quad \tilde{\sigma}(1, X) = b(h_{(1)} \triangleright b') = \epsilon(X).$$

By using the same method, we can check  $\tilde{\sigma}^{-1}$  satisfies the definition of normalised right 2-cocycle. This is another long calculation but sufficiently similar that we omit the details. As a result,  $\tilde{\sigma}$  is an invertible normalised 2-cocycle.  $\square$

**Lemma 3.12.** *If  $B\#_{\sigma}H$  is of associative type, then the map  $\phi : B^e\#H \rightarrow B^e\#_{\sigma}H$  is an invertible  $B$ -coring map, which is given by*

$$\phi(b \otimes b' \# h) := b \otimes b' \sigma^{-1}(S(h_{(2)}), h_{(3)}) \#_{\sigma} h_{(1)}$$

for any  $b \otimes b' \# h \in B^e\#H$ .

*Proof.* First we check  $\epsilon = \epsilon^{\sigma} \circ \phi$ , where  $\epsilon^{\sigma}$  is the counit of  $B^e\#_{\sigma}H$ . Let  $X = b \otimes b' \# h \in B^e\#H$ , then

$$\begin{aligned}
\epsilon^{\sigma}(\phi(X)) &= (b\#_{\sigma}h_{(1)})(b'\sigma^{-1}(S(h_{(3)}), h_{(4)})\#_{\sigma}S(h_{(2)})) \\
&= b(h_{(1)} \triangleright b')(h_{(2)} \triangleright \sigma^{-1}(S(h_{(5)}), h_{(6)}))\sigma(h_{(3)}, S(h_{(4)}))\#_{\sigma}1 \\
&= b(h \triangleright b')\#_{\sigma}1 \\
&= \epsilon(b \otimes b' \# h),
\end{aligned}$$

where in the third step we use Proposition 2.1. Here we always identify  $B$  with its image in  $B\#H$  and  $B\#_{\sigma}H$  by  $b \mapsto b\#1$  and  $b \mapsto b\#_{\sigma}1$  respectively. We can see  $\phi$  is left  $B$ -module map. We can also check it is right  $B$ -linear:

$$\begin{aligned}
\phi(X \triangleleft b'') &= \phi(b \otimes b'(S(h_{(2)}) \triangleright b'')\#h_{(1)}) \\
&= b \otimes b'(S(h_{(4)}) \triangleright b'')\sigma^{-1}(S(h_{(2)}), h_{(3)})\#_{\sigma}h_{(1)} \\
&= b \otimes b'\sigma^{-1}(S(h_{(3)}), h_{(4)})(S(h_{(2)}) \triangleright b'')\#_{\sigma}h_{(1)} \\
&= \phi(X) \triangleleft b'',
\end{aligned}$$

for all  $b'' \in B$ , where the third step uses Lemma 2.4. Recall that the translation map of the Hopf Galois extension  $B = (B\#_{\sigma}H)^{coH} \subseteq B\#_{\sigma}H$  is given by

$$\tau(h) = \sigma^{-1}(S(h_{(2)}), h_{(3)})\#_{\sigma}S(h_{(1)}) \otimes_B 1\#_{\sigma}h_{(4)},$$

for all  $h \in H$ . So we have

$$\begin{aligned}
(\phi \otimes_B \phi)(\Delta(X)) &= (\phi \otimes_B \phi)(b \otimes 1\#h_{(1)} \otimes_B 1 \otimes b'\#h_{(2)}) \\
&= (b \otimes \sigma^{-1}(S(h_{(2)}), h_{(3)})\#_{\sigma}h_{(1)}) \otimes_B (1 \otimes b'\sigma^{-1}(S(h_{(5)}), h_{(6)})\#_{\sigma}h_{(4)}) \\
&= \Delta^{\sigma}(b \otimes b'\sigma^{-1}(S(h_{(2)}), h_{(3)})\#_{\sigma}h_{(1)}) \\
&= \Delta^{\sigma}(\phi(X)),
\end{aligned}$$

where  $\Delta^{\sigma}$  is the coproduct of  $B^e\#_{\sigma}H$ . We can see  $\phi$  is invertible with the inverse  $\phi$  given by:

$$\phi^{-1}(b \otimes b' \#_{\sigma}h) = b \otimes b'(S(h_{(2)}) \triangleright \sigma(h_{(3)}, S(h_{(4)}))\#h_{(1)}).$$

Here

$$\phi\phi^{-1}(b \otimes b' \#_{\sigma}h) = b \otimes b'\sigma^{-1}(S(h_{(3)}), h_{(4)})S(h_{(2)})\triangleright\sigma(h_{(5)}, S(h_{(6)}))\#_{\sigma}h_{(1)}$$

and using Lemma 2.1(2), we have

$$\begin{aligned}
&\sigma^{-1}(S(h_{(2)}), h_{(3)})S(h_{(1)})\triangleright\sigma(h_{(4)}, S(h_{(5)})) \\
&= \sigma^{-1}(S(h_{(4)}), h_{(5)})\sigma(S(h_{(3)}), h_{(6)})\sigma(S(h_{(2)})h_{(7)}, S(h_{(10)}))\sigma^{-1}(S(h_{(1)}), h_{(8)}S(h_{(9)})) = \epsilon(h).
\end{aligned}$$

In the other side, we similarly have

$$\phi^{-1}\phi(b \otimes b' \# h) = b \otimes b' \sigma^{-1}(S(h_{(5)}), h_{(6)})S(h_{(2)}) \triangleright \sigma(h_{(3)}, S(h_{(4)})) \# h_{(1)}$$

and using Lemma 2.1(4), we have

$$\begin{aligned} \sigma^{-1}(S(h_{(4)}), h_{(5)})S(h_{(1)}) \triangleright \sigma(h_{(2)}, S(h_{(3)})) &= (S(h_{(1)})) \triangleright ((h_{(2)}) \triangleright \sigma^{-1}(S(h_{(5)}), h_{(6)})) \sigma(h_{(3)}, S(h_{(4)})) \\ &= S(h_{(1)}) \triangleright \epsilon(h_{(2)}) 1 = \epsilon(h) \end{aligned}$$

where, in the associative type case we use that  $\triangleright$  is a module algebra.  $\square$

**Theorem 3.13.** *If  $B \#_{\sigma} H$  is a associative type, then there is an invertible normalised 2-cocycle  $\tilde{\sigma}$  on  $B^e \# H$ , such that  $\phi : (B^e \# H)^{\tilde{\sigma}} \rightarrow B^e \#_{\sigma} H$  is an isomorphism of left  $B$ -bialgebroids, where  $\tilde{\sigma}$  is given by Lemma 3.11 and  $\phi$  is given by Lemma 3.12.*

*Proof.* Since  $\phi$  is a coring map, so we only need to show  $\phi$  is an algebra map. Let  $X = b \otimes b' \# h$ ,  $Y = c \otimes c' \# g \in B^e \# H$ .

On the one hand,

$$\begin{aligned} \phi(X \cdot_{\tilde{\sigma}} Y) &= \phi\left(\tilde{\sigma}(b \otimes 1 \# h_{(1)}, c \otimes 1 \# g_{(1)}) \otimes ((S(g_{(3)})S(h_{(3)})) \triangleright \tilde{\sigma}^{-1}(1 \otimes b' \# h_{(4)}, 1 \otimes c' \# g_{(4)})) \# h_{(2)} g_{(2)}\right) \\ &= \phi\left(b(h_{(1)} \triangleright c) \sigma(h_{(2)}, g_{(1)}) \otimes (S(g_{(3)})S(h_{(4)})) \triangleright (\sigma^{-1}(h_{(5)}, g_{(4)})(h_{(6)} g_{(5)} \triangleright c')(h_{(7)} \triangleright b'))\right) \# h_{(3)} g_{(2)} \\ &= b(h_{(1)} \triangleright c) \sigma(h_{(2)}, g_{(1)}) \otimes (S(g_{(5)})S(h_{(6)})) \triangleright (\sigma^{-1}(h_{(7)}, g_{(6)})(h_{(8)} g_{(7)} \triangleright c')(h_{(9)} \triangleright b')) \\ &\quad \sigma^{-1}(S(g_{(3)})S(h_{(4)}), h_{(5)} g_{(4)}) \#_{\sigma} h_{(3)} g_{(2)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \phi(X)\phi(Y) &= (b \otimes b' \sigma^{-1}(S(h_{(2)}), h_{(3)}) \#_{\sigma} h_{(1)})(c \otimes c' \sigma^{-1}(S(g_{(2)}), g_{(3)}) \#_{\sigma} g_{(1)}) \\ &= b(h_{(1)} \triangleright c) \sigma(h_{(2)}, g_{(1)}) \otimes \\ &\quad c' \sigma^{-1}(S(g_{(5)}), g_{(6)})(S(g_{(4)}) \triangleright (b' \sigma^{-1}(S(h_{(5)}), h_{(6)}))) \sigma(S(g_{(3)}), S(h_{(4)})) \#_{\sigma} h_{(3)} g_{(2)}, \end{aligned}$$

Comparing with both hand sides, it is enough to show

$$\begin{aligned} (S(g_{(3)})S(h_{(3)})) \triangleright (\sigma^{-1}(h_{(4)}, g_{(4)})(h_{(5)} g_{(5)} \triangleright c')(h_{(6)} \triangleright b')) \sigma^{-1}(S(g_{(1)})S(h_{(1)}), h_{(2)} g_{(2)}) \\ = c' \sigma^{-1}(S(g_{(3)}), g_{(4)})(S(g_{(2)}) \triangleright (b' \sigma^{-1}(S(h_{(2)}), h_{(3)}))) \sigma(S(g_{(1)}), S(h_{(1)})). \end{aligned}$$

Indeed,

$$\begin{aligned} (S(g_{(3)})S(h_{(3)})) \triangleright (\sigma^{-1}(h_{(4)}, g_{(4)})(h_{(5)} g_{(5)} \triangleright c')(h_{(6)} \triangleright b')) \sigma^{-1}(S(g_{(1)})S(h_{(1)}), h_{(2)} g_{(2)}) \\ = (S(g_{(3)})S(h_{(3)})) \triangleright (((h_{(4)} g_{(4)}) \triangleright c') \sigma^{-1}(h_{(5)}, g_{(5)})(h_{(6)} \triangleright b')) \sigma^{-1}(S(g_{(1)})S(h_{(1)}), h_{(2)} g_{(2)}) \\ = c' (S(g_{(3)})S(h_{(3)})) \triangleright (\sigma^{-1}(h_{(4)}, g_{(4)})(h_{(5)} \triangleright b')) \sigma^{-1}(S(g_{(1)})S(h_{(1)}), h_{(2)} g_{(2)}) \\ = c' \sigma^{-1}(S(g_{(2)})S(h_{(2)}), h_{(3)} g_{(3)})(S(g_{(1)})S(h_{(1)})) \triangleright (\sigma^{-1}(h_{(4)}, g_{(4)})(h_{(5)} \triangleright b')) \\ = c' \sigma^{-1}(S(g_{(4)}), g_{(5)})(S(g_{(3)}) \triangleright \sigma^{-1}(S(h_{(3)}), h_{(4)} g_{(6)})) \\ \quad \sigma(S(g_{(2)}), S(h_{(2)}))(S(g_{(1)})S(h_{(1)})) \triangleright (\sigma^{-1}(h_{(5)}, g_{(7)})(h_{(6)} \triangleright b')) \\ = c' \sigma^{-1}(S(g_{(4)}), g_{(5)})(S(g_{(3)}) \triangleright \sigma^{-1}(S(h_{(3)}), h_{(4)} g_{(6)})) \\ \quad (S(g_{(2)})S(h_{(2)})) \triangleright (\sigma^{-1}(h_{(5)}, g_{(7)})(h_{(6)} \triangleright b')) \sigma(S(g_{(1)}), S(h_{(1)})) \\ = c' \sigma^{-1}(S(g_{(3)}), g_{(4)})S(g_{(2)}) \triangleright \left( \sigma^{-1}(S(h_{(3)}), h_{(4)} g_{(5)})S(h_{(2)}) \triangleright (\sigma^{-1}(h_{(5)}, g_{(6)})(h_{(6)} \triangleright b')) \right) \\ \quad \sigma(S(g_{(1)}), S(h_{(1)})) \\ = c' \sigma^{-1}(S(g_{(3)}), g_{(4)})S(g_{(2)}) \triangleright \left( \sigma^{-1}(S(h_{(6)}), h_{(7)} g_{(5)}) \sigma(S(h_{(5)}), h_{(8)} g_{(6)}) \sigma^{-1}(S(h_{(4)})h_{(9)}, g_{(7)}) \right) \\ \quad \sigma^{-1}(S(h_{(3)}), h_{(10)})(S(h_{(2)})h_{(11)} \triangleright b')) \sigma(S(g_{(1)}), S(h_{(1)})) \end{aligned}$$

$$\begin{aligned}
&= c' \sigma^{-1}(S(g_{(3)}), g_{(4)}) S(g_{(2)}) \triangleright \left( \sigma^{-1}(S(h_{(3)}), h_{(4)}) S(h_{(2)}) \triangleright (h_{(5)} \triangleright b') \right) \sigma(S(g_{(1)}), S(h_{(1)})) \\
&= c' \sigma^{-1}(S(g_{(3)}), g_{(4)}) S(g_{(2)}) \triangleright \left( S(h_{(4)}) \triangleright (h_{(5)} \triangleright b') \sigma^{-1}(S(h_{(2)}), h_{(3)}) \right) \sigma(S(g_{(1)}), S(h_{(1)})) \\
&= c' \sigma^{-1}(S(g_{(3)}), g_{(4)}) S(g_{(2)}) \triangleright \left( b' \sigma^{-1}(S(h_{(2)}), h_{(3)}) \right) \sigma(S(g_{(1)}), S(h_{(1)})),
\end{aligned}$$

where the 1st, 5th steps use the property of associative type. The 3rd, 9th steps use Lemma 2.4. The 4th step uses Proposition 2.1. So we are done,  $\phi(X \cdot_{\tilde{\sigma}} Y) = \phi(X)\phi(Y)$ .  $\square$

The simplest case of a Galois object (see Section 5) was already shown in [23]. By Proposition 3.9 and Theorem 3.13 we have the following corollary

**Corollary 3.14.** *If  $B \#_{\sigma} H$  is of associative type, then  $(B^e \# H)^{\tilde{\sigma}}$  is a left Hopf algebroid with antipode and its inverse given by*

$$\begin{aligned}
S^{\tilde{\sigma}}(X) &:= \phi \circ S \circ \phi^{-1}(X) = b' \sigma^{-1}(S(h_{(3)}), h_{(4)}) \otimes S^2(h_{(1)}) \triangleright b \# S(h_{(2)}) \\
(S^{\tilde{\sigma}})^{-1}(X) &:= \phi \circ S^{-1} \circ \phi^{-1}(X) = h_{(5)} \triangleright b' \otimes b h_{(3)} \triangleright \sigma(S^{-1}(h_{(2)}), h_{(1)}) \# S^{-1}(h_{(4)})
\end{aligned}$$

as corresponding via  $\phi$  to  $S$  on  $B^e \#_{\sigma} H$  in Proposition 3.9.

Moreover, by substituting

$$\begin{aligned}
X_{+}^{\tilde{\sigma}} \otimes_{B^{\circ p}} X_{-}^{\tilde{\sigma}} &= t(\tilde{\sigma}(S^{-1}(S(X_{(1)}_{(4)}_{(2)}), S(X_{(1)}_{(3)}))) S^{-1}(S(X_{(1)}_{(4)}_{(1)})) \\
&\quad \otimes s((\tilde{\sigma})^{-1}(S(X_{(1)}_{(1)}), X_{(2)})) S(X_{(1)}_{(2)}),
\end{aligned}$$

one can check that

$$S^{\tilde{\sigma}}(X) = t(\epsilon(S^{\tilde{\sigma}}(X_{+}^{\tilde{\sigma}}))) X_{-}^{\tilde{\sigma}} \quad (3.20)$$

which suggests that there should be a general theory of twistings of antipodes with  $\epsilon_R = \epsilon \circ S^{\tilde{\sigma}}$  constructed independently by twisting and in the spirit of the left-right antipode theory in [5]. This will be looked at elsewhere.

*Example 3.15.* Following on from Example 2.6, we start with a smash product with inner action given by  $u$  and no cocycle and we saw that this is cohomologous to a trivial action and  $\sigma^u$ . By Lemma 3.11 the corresponding 2-cocycle (which cotwists the left Hopf algebroid  $B^e \# H$ ) is

$$\begin{aligned}
\tilde{\sigma}^u(b \otimes b' \#_{\sigma^u} h, c \otimes c' \#_{\sigma^u} g) &= bc \sigma^u(h, g) c' b' = bc u^{-1}(g_{(1)}) u^{-1}(h_{(1)}) u(h_{(2)} g_{(2)}) c' b' \\
(\tilde{\sigma}^u)^{-1}(b \otimes b' \#_{\sigma^u} h, c \otimes c' \#_{\sigma^u} g) &= bc (\sigma^u)^{-1}(h, g) c' b' = bc u^{-1}(h_{(1)} g_{(1)}) u(h_{(2)}) h(g_{(2)}) c' b'
\end{aligned}$$

and the resulting Hopf algebroid  $(B^e \# H)^{\tilde{\sigma}^u}$  has, by Corollary 3.14,

$$\begin{aligned}
S^{\tilde{\sigma}^u}(b \otimes b' \# h) &= b' u(h_{(2)}) u(S(h_{(3)})) \otimes b \# S(h_{(1)}) \\
(S^{\tilde{\sigma}^u})^{-1}(b \otimes b' \# h) &= b' \otimes u^{-1}(S^{-1}(h_{(2)})) u^{-1}(h_{(1)}) b \# S^{-1}(h_{(3)}).
\end{aligned}$$

Moreover, this Hopf algebroid is isomorphic to  $B^e \#_{\sigma^u} H$ , which by Proposition 3.10, has antipode and its inverse

$$\begin{aligned}
S(b \otimes b' \#_{\sigma^u} h) &= b' \otimes (\sigma^u)^{-1}(S^2(h_{(2)}), S(h_{(1)})) b \#_{\sigma^u} S(h_{(3)}) \\
&= b' \otimes u(S^2(h_{(2)})) u(S(h_{(1)})) b \#_{\sigma^u} S(h_{(3)}) \\
S^{-1}(b \otimes b' \#_{\sigma^u} h) &= b' \sigma^u(h_{(2)}, S(h_{(3)})) \otimes b \#_{\sigma^u} S^{-1}(h_{(1)}) \\
&= b' u^{-1}(h_{(2)}) u^{-1}(S(h_{(3)})) \otimes b \#_{\sigma^u} S^{-1}(h_{(1)}).
\end{aligned}$$

If  $u$  is an algebra map then  $\sigma^u(h, g) = \epsilon(h)\epsilon(g)$  and then the antipode (and its inverse) are just the flipping of  $b$  and  $b'$  together  $S$  (and its inverse) on  $H$ .

## 4. RESULTS FOR GENERAL HOPF GALOIS EXTENSIONS

Here we cast our results on cocycle extensions into a more general context of Hopf Galois extensions.

**4.1. Preliminaries on Hopf Galois extensions.** A Hopf-Galois extension or quantum principal bundle with universal calculus means an  $H$ -comodule algebra  $P$  with coinvariant subalgebra  $B := \{b \in P \mid \Delta_R(b) = b \otimes 1_H\} \subseteq P$  such that the canonical map

$$\text{can} : P \otimes_B P \rightarrow P \otimes H, \quad p \otimes_B q \mapsto pq_{(0)} \otimes q_{(1)}$$

is bijective, where  $\otimes_B$  is the balanced tensor product by  $B \subseteq P$  as a subalgebra. We also require for convenience that  $P$  is a faithful flat left  $B$  module. In fact the inverse of can is determined by its restriction, the *translation map*,

$$\text{can}^{-1}|_{1 \otimes H} : H \rightarrow P \otimes_B P, \quad h \mapsto h^{<1>} \otimes_B h^{<2>}.$$

and it is known, e.g. [7, Prop. 3.6][9, Lemma 34.4] that it obeys

$$h^{<1>} \otimes_B h^{<2>}_{(0)} \otimes h^{<2>}_{(1)} = h_{(1)}^{<1>} \otimes_B h_{(1)}^{<2>} \otimes h_{(2)}, \quad (4.1)$$

$$h_{(2)}^{<1>} \otimes_B h_{(2)}^{<2>} \otimes S(h_{(1)}) = h^{<1>}_{(0)} \otimes_B h^{<2>} \otimes h^{<1>}_{(1)}, \quad (4.2)$$

$$h^{<1>} h^{<2>}_{(0)} \otimes h^{<2>}_{(1)} = 1_P \otimes h, \quad (4.3)$$

$$p_{(0)} p_{(1)}^{<1>} \otimes_B p_{(1)}^{<2>} = 1_P \otimes_B p, \quad (4.4)$$

for all  $h \in H$  and  $p \in P$ .

Finally, a Hopf-Galois extension is *cleft* if there is a *cleaving map* defined as convolution-invertible and  $H$ -equivariant. This is known cf [22, Theorem 8.2.4] to be equivalent to requiring that  $P \simeq B \otimes H$  as left  $B$ -modules and right  $H$ -comodules (i.e. that it has the ‘normal basis property’). This means that a cleft Hopf Galois extension is equivalent to a cocycle cross product. The converse is also true.

**Lemma 4.1.**  *$B \#_\sigma H$  for a cocycle  $(\triangleright, \sigma)$  is a cleft Hopf-Galois extension. Hence the two notions are equivalent.*

*Proof.* We have already seen how the data of an extension with the normal basis property implies a cocycle cross product. Conversely, if we are given a cocycle cross product  $B \#_\sigma H$  by data  $(\triangleright, \sigma)$  then we can view it as a cleft Hopf-Galois extension with  $\Delta_R = \text{id} \otimes \Delta$  and  $B \otimes 1 = B^{\text{co}(H)}$ ,  $j(h) = 1 \otimes h$  as previously discussed. This is clear but we explain it explicitly. Indeed, the inverse to the canonical map is provided by [22]

$$\text{can}^{-1}(b \#_\sigma g \otimes h) = (b \#_\sigma g)(\sigma^{-1}(S(h_{(2)}), h_{(3)}) \#_\sigma S(h_{(1)})) \otimes_B 1 \#_\sigma h_{(4)}, \quad (4.5)$$

for all  $b \in B$  and  $g, h \in H$ . One can check that

$$\begin{aligned} & \text{can}((b \#_\sigma g)(\sigma^{-1}(S(h_{(2)}), h_{(3)}) \#_\sigma S(h_{(1)})) \otimes_B 1 \#_\sigma h_{(4)}) \\ &= (b \#_\sigma g)(\sigma^{-1}(S(h_{(3)}), h_{(4)}) \sigma(S(h_{(2)}), h_{(5)}) \#_\sigma S(h_{(1)}) h_{(6)}) \otimes h_{(7)} = b \#_\sigma g \otimes h. \end{aligned}$$

$\text{can}^{-1}(\text{can}(b \#_\sigma g \otimes_B b' \#_\sigma h))$

$$\begin{aligned} &= (b \#_\sigma g)(b' \#_\sigma h_{(1)})(\sigma^{-1}(S(h_{(3)}), h_{(4)}) \#_\sigma S(h_{(2)})) \otimes_B 1 \#_\sigma h_{(5)} \\ &= (b \#_\sigma g)(b'(h_{(1)} \triangleright \sigma^{-1}(S(h_{(6)}), h_{(7)})) \sigma(h_{(2)}, S(h_{(5)})) \#_\sigma h_{(3)} S(h_{(4)})) \otimes_B 1 \#_\sigma h_{(8)} \\ &= (b \#_\sigma g)(b' \#_\sigma 1) \otimes_B 1 \#_\sigma h = b \#_\sigma g \otimes_B b' \#_\sigma h, \end{aligned}$$

for all  $b, b' \in B$  and  $g, h \in H$ , where the third step uses Proposition 2.1.  $\square$

Given the lemma, our various results in Section 2 translate to the point of view of cleft Hopf-Galois extensions as follows, with the more elementary proofs left to the reader. First of all, our notion of associative type from the point of view of cleft Hopf-Galois extensions is equivalent to

$$j^{-1}(g_{(1)}) j^{-1}(h_{(1)}) j(h_{(2)} g_{(2)}) \in Z(B). \quad (4.6)$$

Next, ‘gauge transform’ from  $(\triangleright, \sigma)$  to  $(\triangleright^u, \sigma^u)$  by convolution-invertible  $u : H \rightarrow B$  is equivalent from the point of view of cleft Hopf-Galois extensions to a change of the cleaving map  $j$  to  $j^u = u^{-1} * j$ , and preserves associative type as in Proposition 2.5 if and only if

$$j^{-1}(g_{(1)})u(g_{(2)})j^{-1}(h_{(1)})u(h_{(2)})u^{-1}(h_{(3)}g_{(3)})j(h_{(4)}g_{(4)}) \in Z(B). \quad (4.7)$$

Lastly, it was recently observed [1] that if  $\chi : H \otimes H \rightarrow k$  is an invertible normalised Drinfeld cotwist on  $H$  then a Hopf-Galois extension  $B = P^{coH} \subseteq P$  cotwists to another one,  $B = P_{\chi}^{coH^X} \subseteq P_{\chi}$  with product (2.1). We can make this a little more explicit:

**Lemma 4.2.** *If  $P, H$  is a Hopf-Galois extension with translation map as above and  $\chi$  a Drinfeld cotwist then  $P_{\chi}, H^X$  has translation map*

$$\tau_{\chi}(h) := h_{(3)}^{<1>} \otimes_B h_{(3)}^{<2>} \chi(h_{(1)} \otimes S(h_{(2)})),$$

for all  $h \in H$ .

*Proof.* We give an elementary direct proof. On the one hand,

$$\begin{aligned} (\text{can}_{\chi} \circ \tau_{\chi})(h) &= h_{(3)}^{<1>} h_{(3)}^{<2>} \otimes \chi^{-1}(h_{(3)}^{<1>} h_{(3)}^{<2>}) h_{(3)}^{<2>} \chi(h_{(1)} \otimes S(h_{(2)})) \\ &= h_{(4)}^{<1>} h_{(4)}^{<2>} \otimes \chi^{-1}(S(h_{(3)}) \otimes h_{(4)}^{<2>}) h_{(4)}^{<2>} \chi(h_{(1)} \otimes S(h_{(2)})) \\ &= 1 \otimes \chi^{-1}(S(h_{(3)}) \otimes h_{(4)}) h_{(5)} \chi(h_{(1)} \otimes S(h_{(2)})) = 1 \otimes h, \end{aligned}$$

for all  $h \in H$ , where the 2nd step uses (4.2), the 3rd step uses (4.3), and the last step uses the fact that  $\chi$  is a cotwist on the Hopf algebra  $H$ . (This is sufficient since  $\text{can}_{\chi}$  is a left  $P_{\chi}$  module map.) On the other side,

$$\begin{aligned} (\text{can}_{\chi}^{-1} \circ \text{can}_{\chi})(p \otimes_B q) &= p \cdot_{\chi} q_{(0)} \cdot_{\chi} q_{(3)}^{<1>} \otimes_B q_{(3)}^{<2>} \chi(q_{(1)} \otimes S(q_{(2)})) \\ &= p \cdot_{\chi} (q_{(0)} q_{(4)}^{<1>}) \otimes_B q_{(4)}^{<2>} \chi^{-1}(q_{(1)} \otimes q_{(4)}^{<1>}) \chi(q_{(2)}, S(q_{(3)})) \\ &= p \cdot_{\chi} (q_{(0)} q_{(5)}^{<1>}) \otimes_B q_{(5)}^{<2>} \chi^{-1}(q_{(1)} \otimes S(q_{(4)})) \chi(q_{(2)} \otimes S(q_{(3)})) \\ &= p \cdot_{\chi} (q_{(0)} q_{(1)}^{<1>}) \otimes_B q_{(1)}^{<2>} = p \otimes_B q, \end{aligned}$$

for all  $p, q \in P_{\chi}$ , where the 3rd step uses (4.2), and the last step uses (4.4). Now, in the cleft case, as the isomorphism  $P \cong B \otimes H$  is left  $B$ -linear (since the coaction is the same on the underlying vector space) and right  $H^X$ -colinear (since  $b \cdot_{\chi} p = bp$  as  $b$  is coinvariant) the twisting reduces to Proposition 2.2.  $\square$

In the cleft case, this agrees with the twisting of the cocycle extension in Proposition 2.2. From the point of view of Hopf-Galois extensions, the condition that twisting by  $\chi$  preserves associative type as in Proposition 2.7 is

$$j^{-1}(h_{(1)}g_{(1)})\chi(h_{(2)}, g_{(2)})j(h_{(3)}g_{(3)}) \in Z(B) \quad (4.8)$$

on using the form of the action  $\triangleright$  as given by  $j$ .

**4.2. Ehresmann-Schauenburg Hopf algebroid.** The classical construction here is the Ehresmann groupoid  $\mathcal{G}$  associated to a classical principal bundle  $X \rightarrow M$  where  $M$  is classical manifold, with structure group  $G$ . Then  $\mathcal{G} = (X \times X)/G$  where we identify  $(x, y) \sim (x.u, y.u)$  for all  $u \in G$  and  $x, y \in X$ . The base is  $M = X/G$  similarly identified. The composition is  $(x, y) \circ (y, z) = (x, z)$  and the source and target maps are represented by  $s(x, y) = x$  and  $t(x, y) = y$  which one can check are all well-defined on the quotient spaces (so the results here are  $\pi(x)$  and  $\pi(y)$ ).

**Definition 4.3.** [9, §34.13]. *Let  $B = P^{coH} \subseteq P$  be a Hopf Galois extension such that  $P$  is a faithful flat left  $B$ -module. The Ehresmann Schauenburg bialgebroid is*

$$\mathcal{B} := \{p \otimes q \in P \otimes P \mid p_{(0)} \otimes \tau(p_{(1)})q = p \otimes q \otimes_B 1\} \subset P \otimes P$$

with  $B$ -bimodule inherited from  $P$  and  $B$ -coring coproduct and counit

$$\Delta(p \otimes q) = p_{(0)} \otimes \tau(p_{(1)}) \otimes q, \quad \epsilon(p \otimes q) = pq.$$

Moreover,  $\mathcal{B}$  is a  $B^e$ -ring with the product and unit

$$(p \otimes q) \bullet_c (r \otimes u) = pr \otimes uq, \quad \eta(b \otimes c) = b \otimes c$$

for all  $p \otimes q, r \otimes u \in \mathcal{B}$  and  $b \otimes c \in B^e$ . Here  $s(b) = b \otimes 1$  and  $t(b) = 1 \otimes b$ .

It was shown in [9], [13] that the  $B$ -bimodule  $\mathcal{B}$  is isomorphic to the  $B$ -bimodule of coinvariant elements,

$$\mathcal{B} \simeq (P \otimes P)^{coH} := \{p \otimes q \in P \otimes P \mid p_{(0)} \otimes q_{(0)} \otimes p_{(1)} q_{(1)} = p \otimes q \otimes 1_H\}. \quad (4.9)$$

The last result and the fact that it is true classically suggests that  $\mathcal{B}$  should generally be a left Hopf algebroid. We prove this now.

**Theorem 4.4.** *If  $B \subseteq P$  is a  $H$ -Hopf Galois extension, then  $\mathcal{B}$  is a Hopf algebroid which we denote  $\mathcal{L}(P, H)$ .*

*Proof.* Let  $X = a \otimes a', Y = d \otimes d'$ . We define  $\rho : \mathcal{L}(P, H) \otimes_B \mathcal{L}(P, H) \rightarrow \mathcal{L}(P, H) \otimes_{B^{op}} \mathcal{L}(P, H)$  by:

$$\rho(X \otimes_B Y) := a \otimes a'_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y,$$

this is well defined, since we have

$$\begin{aligned} \rho(t(b)X \otimes_B Y) &= a \otimes a'_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a'_{(0)} b \otimes a'_{(1)} \langle 1 \rangle) Y \\ &= \rho(X \otimes_B s(b)Y). \end{aligned}$$

And

$$\begin{aligned} a \otimes ba'_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y &= (a \otimes a'_{(1)} \langle 2 \rangle) t(b) \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y \\ &= a \otimes a'_{(1)} \langle 2 \rangle \otimes_{B^{op}} t(b) (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y \\ &= a \otimes a'_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle b) Y \end{aligned}$$

Moreover,  $(a \otimes a'_{(1)} \langle 2 \rangle) \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) \in \mathcal{L}(P, H) \otimes_{B^{op}} \mathcal{L}(P, H)$ , since

$$\begin{aligned} \Delta_R^{\text{diag}}(a \otimes a'_{(1)} \langle 2 \rangle) \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) &= (a_{(0)} \otimes a'_{(1)} \langle 2 \rangle_{(0)}) \otimes a_{(1)} a'_{(1)} \langle 2 \rangle_{(1)} \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) \\ &= (a_{(0)} \otimes a'_{(1)} \langle 2 \rangle) \otimes a_{(1)} a'_{(2)} \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) \\ &= (a \otimes a'_{(1)} \langle 2 \rangle) \otimes 1 \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) \end{aligned}$$

We can see  $\lambda$  is the inverse of  $\rho$ . On the one hand:

$$\begin{aligned} \lambda(\rho(X \otimes_B Y)) &= \lambda(a \otimes a'_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y) \\ &= a_{(0)} \otimes a_{(0)} \langle 1 \rangle \otimes_B (a_{(1)} \langle 2 \rangle \otimes a'_{(1)} \langle 2 \rangle) (a'_{(0)} \otimes a'_{(1)} \langle 1 \rangle) Y \\ &= a_{(0)} \otimes a_{(0)} \langle 1 \rangle \otimes_B (a_{(1)} \langle 2 \rangle a' \otimes 1) Y \\ &= X \otimes_B Y. \end{aligned}$$

On the other hand,

$$\begin{aligned} \rho(\lambda(X \otimes_{B^{op}} Y)) &= \rho(a_{(0)} \otimes a_{(1)} \langle 1 \rangle \otimes_B (a_{(1)} \langle 2 \rangle \otimes a') Y) \\ &= a_{(0)} \otimes a_{(1)} \langle 1 \rangle_{(1)} \langle 2 \rangle \otimes_{B^{op}} (a_{(1)} \langle 1 \rangle_{(0)} \otimes a_{(1)} \langle 1 \rangle_{(1)} \langle 1 \rangle) (a_{(1)} \langle 2 \rangle \otimes a') Y \\ &= a_{(0)} \otimes S(a_{(1)}) \langle 2 \rangle \otimes_{B^{op}} (a_{(2)} \langle 1 \rangle \otimes S(a_{(1)}) \langle 1 \rangle) (a_{(2)} \langle 2 \rangle \otimes a') Y \\ &= a_{(0)} \otimes S(a_{(1)}) \langle 2 \rangle \otimes_{B^{op}} (1 \otimes a' S(a_{(1)}) \langle 1 \rangle) Y \\ &= a_{(0)} \otimes a' S(a_{(1)}) \langle 1 \rangle S(a_{(1)}) \langle 2 \rangle \otimes_{B^{op}} Y \\ &= X \otimes_{B^{op}} Y. \end{aligned}$$

□

This generalises [24] for the case of a Galois object (see Section 5). We also connect to Section 3 as follows.

**Lemma 4.5.** *In the cleft case, we have  $\mathcal{L}(B\#_\sigma H, H) \cong B^e\#_\sigma H$  as left Hopf algebroids*

*Proof.* Using (4.9), invariant elements of  $P \otimes P$  are then of the form

$$\mathcal{L}(B\#_\sigma H, H) = \{b\#_\sigma h_{(1)} \otimes b'\#_\sigma S(h_{(2)}) \mid b, b' \in B, h \in H\}.$$

We can also see the coproduct is given by

$$\Delta(b\#_\sigma h_{(1)} \otimes b'\#_\sigma S(h_{(2)})) = b\#_\sigma h_{(1)} \otimes \sigma^{-1}(S(h_{(3)}), h_{(4)})\#_\sigma S(h_{(2)}) \otimes_B 1\#_\sigma h_{(5)} \otimes b'\#_\sigma S(h_{(6)}),$$

since from (4.5) we know the translation map  $\tau : H \rightarrow B\#H \otimes_B B\#H$  is

$$\tau(h) = \sigma^{-1}(S(h_{(2)}), h_{(3)})\#_\sigma S(h_{(1)}) \otimes_B 1\#_\sigma h_{(4)},$$

for all  $h \in H$ . Next,

$$H \rightarrow H \otimes H, \quad h \mapsto h_{(1)} \otimes S(h_{(2)})$$

is a vector space inclusion with image exactly the elements as shown. So we have an induced isomorphism of vector spaces

$$\Theta : B^e\#_\sigma H \rightarrow \mathcal{L}(B\#_\sigma H, H), \quad \Theta(b \otimes b'\#_\sigma h) := b\#_\sigma h_{(1)} \otimes b'\#_\sigma S(h_{(2)}).$$

It remains to show that this preserves the bialgebroid structures.

First, it is not hard to see this is a  $B$ -bimodule map

$$\begin{aligned} \Theta(c \triangleright (b \otimes b'\#_\sigma h) \triangleleft c') &= \Theta(cb \otimes b'(S(h_{(2)})) \triangleright c')\#_\sigma h_{(1)} \\ &= cb\#_\sigma h_{(1)} \otimes b'(S(h_{(3)})) \triangleright c'\#_\sigma S(h_{(2)}). \end{aligned}$$

We can see  $\Theta$  is a coring map:

$$\begin{aligned} (\Theta \otimes_B \Theta)(\Delta(b \otimes b'\#_\sigma h)) &= b\#_\sigma h_{(1)} \otimes \sigma^{-1}(S(h_{(3)}), h_{(4)})\#_\sigma S(h_{(2)}) \otimes_B 1 \otimes_\sigma h_{(5)} \otimes b'\#_\sigma S(h_{(6)}) \\ &= \Delta(b\#_\sigma h_{(1)} \otimes b'\#_\sigma S(h_{(2)})). \end{aligned}$$

We also have  $\epsilon(\Theta(X)) = \epsilon(X)$ :

$$\epsilon(b\#_\sigma h_{(1)} \otimes b'\#_\sigma) = b(h_{(1)} \triangleright b')\sigma(h_{(2)}, S(h_{(3)})) = \epsilon(b \otimes b'\#_\sigma h).$$

Next, we show  $\Theta(XY) = \Theta(X)\Theta(Y)$  for all  $X, Y \in B^e\#_\sigma H$ . Indeed,

$$\begin{aligned} &\Theta((b \otimes b'\#_\sigma h).(c \otimes c'\#_\sigma g)) \\ &= \Theta(b(h_{(1)} \triangleright c)\sigma(h_{(2)}, g_{(1)}) \otimes c'(S(g_{(4)})) \triangleright b')\sigma(S(g_{(3)}), S(h_{(4)}))\#_\sigma h_{(3)}g_{(2)}) \\ &= (b\#_\sigma h_{(1)} \otimes b'\#_\sigma S(h_{(2)}))(c\#_\sigma g_{(1)} \otimes c'\#_\sigma S(g_{(2)})) \end{aligned}$$

□

We next want to know if  $\mathcal{L}(P, H)$  has an antipode. In view of the lemma, we already know this if  $\sigma$  is of associative type. Classically, we know the inverse of a gauge groupoid is just the flipping of the two entries. However, when we flip a element  $a \otimes a' \in \mathcal{L}(P, H)$ , the result  $a' \otimes a$  doesn't belong to  $\mathcal{L}(P, H)$ , since  $H$  is not commutative. Rather, we need some kind of braiding to generalise the flip and we provide this now if  $H$  is coquasitriangular.

We recall that a Hopf algebra  $H$  is a coquasitriangular if there is a convolution invertible linear map  $\mathcal{R} : H \otimes H \rightarrow k$  satisfies

$$\begin{aligned} g_{(1)}h_{(1)}\mathcal{R}(h_{(2)} \otimes g_{(2)}) &= \mathcal{R}(h_{(1)} \otimes g_{(1)})h_{(2)}g_{(2)} \\ \mathcal{R}(h \otimes gf) &= \mathcal{R}(h_{(1)} \otimes g)\mathcal{R}(h_{(2)} \otimes f), \quad \mathcal{R}(hg \otimes f) = \mathcal{R}(h \otimes f_{(1)})\mathcal{R}(g \otimes f_{(2)}) \end{aligned}$$

for all  $h, g, f \in H$ . It is known (for example [18], Prop 2.2.4) that the antipode of a coquasitriangular Hopf algebra is invertible. Now, since  $P$  is a right  $H$ -comodule, we know[18] that  $\mathcal{R}$  induces a braiding

$$\Psi(p \otimes q) = q_{(0)} \otimes p_{(0)}\mathcal{R}(p_{(1)} \otimes q_{(1)})$$



which is an  $H$ -comodule map and invertible, with inverse

$$\Psi^{-1}(p \otimes q) = q_{(0)} \otimes p_{(0)} \mathcal{R}^{-1}(q_{(1)} \otimes p_{(1)}).$$

We use this for the notion of ‘flip’ need for the antipode.

**Proposition 4.6.** *If  $H$  is coquasitriangular then  $\mathcal{L}(P, H)$  has an antipode.*

*Proof.* Since  $\Psi$  is a morphism in the category of  $H$ -comodules, it and its inverse restrict to the invariant subspace of  $P \otimes P$ , hence we can set  $S = \Psi^{-1}$ . We know this is invertible and now we show that the remaining conditions of Lemma 3.3 apply.

First, we can see  $S(t(b)X) = S(X)s(b)$  and  $S^{-1}(s(b)X) = S^{-1}(X)t(b)$ . For condition (2) of Lemma 3.3, let  $X = p \otimes q$  we have:

$$\begin{aligned} S^{-1}(X_{(2)})_{(1)} \otimes_B \otimes_B S^{-1}(X_{(2)})_{(1)} X_{(1)} &= (q_{(0)} \otimes \mathcal{R}(p_{(1)}^{<2>}_{(1)} \otimes q_{(1)}) p_{(1)}^{<2>}_{(0)})_{(1)} \\ &\quad \otimes_B (q_{(0)} \otimes \mathcal{R}(p_{(1)}^{<2>}_{(1)} \otimes q_{(1)}) p_{(1)}^{<2>}_{(0)})_{(2)} (p_{(0)} \otimes p_{(1)}^{<1>}) \\ &= (\mathcal{R}(p_{(2)} \otimes S(p_{(3)})) q_{(0)} \otimes q_{(1)}^{<1>}) \otimes_B (q_{(1)}^{<2>} \otimes p_{(1)}^{<2>}) (p_{(0)} \otimes p_{(1)}^{<1>}) \\ &= (\mathcal{R}(p_{(1)} \otimes S(p_{(2)})) q_{(0)} \otimes q_{(1)}^{<1>}) \otimes_B (q_{(1)}^{<2>} p_{(0)} \otimes 1) \\ &= (\mathcal{R}(p_{(1)} \otimes S(p_{(2)})) q_{(0)} \otimes q_{(1)}^{<1>} q_{(1)}^{<2>} p_{(0)}) \otimes_B 1_{\mathcal{C}} \\ &= (\mathcal{R}(p_{(1)} \otimes S(p_{(2)})) q \otimes p_{(0)}) \otimes_B 1_{\mathcal{C}} \\ &= (\mathcal{R}(p_{(1)} \otimes q_{(1)}) q_{(0)} \otimes p_{(0)}) \otimes_B 1_{\mathcal{C}} \\ &= S^{-1}(X) \otimes_B 1_{\mathcal{C}}. \end{aligned}$$

For condition (3),

$$\begin{aligned} S(X_{(1)})_{(1)} X_{(2)} \otimes_B S(X_{(1)})_{(2)} &= \mathcal{R}^{-1}(S(p_{(2)} \otimes p_{(1)}) (p_{(3)}^{<1>}_{(0)} \otimes p_{(3)}^{<1>}_{(1)}^{<1>})) (p_{(3)}^{<2>} \otimes q) \otimes_B (a_{(3)}^{<1>}_{(1)}^{<2>} \otimes p_{(0)}) \\ &= \mathcal{R}^{-1}(S(p_{(2)} \otimes p_{(1)}) (p_{(4)}^{<1>} \otimes S(p_{(3)}^{<1>})) (p_{(4)}^{<2>} \otimes q) \otimes_B (S(p_{(3)}^{<2>} \otimes p_{(0)})) \\ &= (\mathcal{R}^{-1}(S(p_{(2)} \otimes p_{(1)}) \otimes q S(p_{(3)}^{<1>})) \otimes_B (S(p_{(3)}^{<2>} \otimes p_{(0)})) \\ &= 1_{\mathcal{C}} \otimes_B (\mathcal{R}^{-1}(S(p_{(2)} \otimes p_{(1)}) q \otimes p_{(0)})) \\ &= 1_{\mathcal{C}} \otimes_B (\mathcal{R}^{-1}(q_{(1)} \otimes p_{(1)}) q_{(0)} \otimes p_{(0)}) \\ &= 1_{\mathcal{C}} \otimes_B S(X). \end{aligned}$$

□

In the case of a cleft Hopf-Galois extension we obtain an antipode on the corresponding cocycle Hopf algebroid.

**Corollary 4.7.** *If  $H$  is coquasitriangular then  $B^e \#_{\sigma} H$  has a left antipode*

$S(b \otimes b' \#_{\sigma} h) = u^{-1}(h_{(1)}) b' \otimes b \#_{\sigma} S(h_{(2)})$ ,  $S^{-1}(b \otimes b' \#_{\sigma} h) = v(h_{(2)}) b' \otimes b \#_{\sigma} S^{-1}(h_{(1)})$  obeying the conditions in Lemma 3.3, where  $v(h) = \mathcal{R}(h_{(1)} \otimes S(h_{(2)}))$  and  $u^{-1}(h) = \mathcal{R}(S^2(h_{(2)}) \otimes h_{(1)})$  are standard convolution-invertible maps related to the square of the antipode on  $H$ .

*Proof.* We use the isomorphism in Lemma 4.5 to transfer back  $S, S^{-1}$  on  $\mathcal{L}(B \#_{\sigma} H, H)$  computed from Proposition 4.6 as

$$\begin{aligned} S(b \#_{\sigma} h_{(1)} \otimes b' \#_{\sigma} S(h_{(2)})) &= b' \#_{\sigma} S(h_{(4)}) \otimes b \#_{\sigma} h_{(1)} \mathcal{R}^{-1}(S(h_{(3)}) \otimes h_{(2)}) \\ &= b' \#_{\sigma} S(h_{(4)}) \otimes b \#_{\sigma} h_{(1)} \mathcal{R}(S^2(h_{(3)}) \otimes h_{(2)}) \\ &= b' \#_{\sigma} S(h_{(4)}) \otimes b \#_{\sigma} S^2(h_{(3)}) \mathcal{R}(S^2(h_{(2)}) \otimes h_{(1)}), \end{aligned}$$

where the last step uses [18, Lem. 2.2.2, Prop. 2.2.3]. For the inverse, writing  $g = S^{-1}(h)$ ,

$$S^{-1}(b \#_{\sigma} h_{(1)} \otimes b' \#_{\sigma} S(h_{(2)})) = b' \#_{\sigma} S(h_{(4)}) \otimes b \#_{\sigma} h_{(1)} \mathcal{R}(h_{(2)} \otimes S(h_{(3)}))$$

$$\begin{aligned}
&= b' \#_{\sigma} S^2(g_{(1)}) \otimes b \#_{\sigma} S(g_{(4)}) \mathcal{R}(S(g_{(3)}) \otimes S^2(g_{(2)})) \\
&= b' \#_{\sigma} S^2(g_{(1)}) \mathcal{R}(g_{(3)} \otimes S(g_{(2)})) \otimes b \#_{\sigma} S(g_{(4)}) \\
&= b' \#_{\sigma} g_{(3)} \mathcal{R}(g_{(2)} \otimes S(g_{(1)})) \otimes b \#_{\sigma} S(g_{(4)}) \\
&= b' \#_{\sigma} \mathcal{R}(S^{-1}(h_{(3)}) \otimes h_{(4)}) S^{-1}(h_{(2)}) \otimes b \#_{\sigma} h_{(1)}
\end{aligned}$$

where the 2nd step use of [18, Lem. 2.2.2] and the 3rd step uses [18, Prop. 2.2.4]. We use invariance of  $\mathcal{R}$  under  $S \otimes S$  to recognise the answer in terms of the Drinfeld functions  $u^{-1}, v$  in [18, Prop. 2.2.4].  $\square$

Note that this construction is very different from the one for a cocycle Hopf algebroid of associative type in Proposition 3.10, hence it is clear that  $S$  obeying the conditions in Lemma 3.3 need not be unique. Indeed, for the two antipodes to coincide, we would need

$$\sigma^{-1}(S^2(h_{(2)}), S(h_{(1)}))(S^2(h_{(3)}) \triangleright b) = u^{-1}(h)b, \quad (h_{(1)} \triangleright b)\sigma(h_{(2)}, S(h_{(3)})) = v(h)b,$$

for all  $h \in H$  and  $b \in B$ . Setting  $b = 1$ , this requires  $\sigma^{-1}(S^2(h_{(2)}), S(h_{(1)})) = u^{-1}(h)$  and  $\sigma(h_{(1)}, S h_{(2)}) = v(h)$  for all  $h$ , which requires  $\sigma$  to be  $k$ -valued and  $h \triangleright b = \epsilon(h)b$  the trivial action. For an example, we can let  $\sigma = \mathcal{R}$  be viewed as a cocycle with values in  $k$  which, together with the trivial action means  $B \#_{\sigma} H = B \otimes_{(\sigma} H)$ , where  ${}_{\sigma} H$  is a version of (2.1) one-sided cotwisted on the left with the left coaction of  $H$  on itself, i.e.

$$h \cdot g = \sigma(h_{(1)} \otimes g_{(1)}) h_{(2)} g_{(2)}. \quad (4.10)$$

with  $\sigma = \mathcal{R}$  in the present case. Thus, the two antipodes can agree in very special cases, but in general are different.

## 5. MORE EXAMPLES OF ASSOCIATIVE TYPE

There are several ways to arrive at a cleft extension of associative type. The most obvious is  $H$  cocommutative and  $\sigma$  has its image in  $Z(B)$ . In this case, the condition in Definition 2.3 is clearly automatic.

**5.1. Galois objects.** The next most obvious case is  $\triangleright$  trivial  $h \triangleright b = \epsilon(h)b$  and in this case (2.6) tells us that the extension is of associative type iff the image of  $\sigma$  is in  $Z(B)$ . We saw how this arises from gauge transform by an inner action in Example 2.6 but clearly one could have more general form of  $\sigma$ .

The simplest such example is that of a *Galois object*. By definition, this as a Hopf-Galois extension where  $B = k$ , the field. In the cleft case, it means  $P \cong H$  as a vector space and clearly in his case  $\triangleright$  is trivial and the conditions for a cocycle  $\sigma$  reduce to those of a Drinfeld 2-cocycle. Then every such Galois object is a 1-sided cotwist

$$B \#_{\sigma} H = {}_{\sigma} H$$

as in (4.10).

By contrast, the associated Hopf algebroid is now just a Hopf algebra and we clearly have from Proposition 3.9 and Theorem 3.13 that

$$B^e \#_{\sigma} H = (B^e \# H)^{\bar{\sigma}} = H^{\sigma}$$

where the second step uses  $B^e \# H = H$  as Hopf algebra and  $\bar{\sigma} = \sigma$  by Lemma 3.11 as a Drinfeld cotwist in agreement with both [23] where Galois objects were looked at, and with our twisting Theorem 3.13 with  $\bar{\sigma} = \sigma$  as the base is trivial.

This also agrees with our observations about Drinfeld cotwists. Thus, if we start with a cleft Galois object above with cocycle  $\sigma$  and structure quantum group  $H$ , and then twist by  $\chi$  then

$$B \#_{\sigma * \chi^{-1}} H^{\chi} = {}_{\sigma * \chi^{-1}}(H^{\chi}) = {}_{\sigma} H_{\chi}, \quad B^e \#_{\sigma * \chi^{-1}} H^{\chi} = (H^{\chi})^{\sigma * \chi^{-1}} = H^{\sigma}$$

where  $H_\chi$  is one-sided right cotwist using (2.1) and the right coaction by the coproduct. We used Proposition 2.2 for the twisted cocycle extension and then our previous observations. The condition in Proposition 2.7 is automatic as  $\triangleright$  is trivial.

**5.2. Affine quantum group.** The abstract structure here [18, Lem. 6.3.15] starts with  $B \hookrightarrow P \twoheadrightarrow H$  algebra maps between Hopf algebras with  $P = B \blacktriangleright H$  as a coalgebra (so with respect to a coaction  $\Delta_R(h) = h^{(1)} \otimes h^{(2)} \in H \otimes B$ ), the inclusion given by  $b \mapsto 1 \otimes b$  and the surjection by  $b \otimes h \mapsto \epsilon(b)h$ . Suppose that  $j : H \rightarrow P$  given by  $j(h) = 1 \otimes h$  obeys  $bj(h) = b \otimes h = j(h)b$ . Then  $j^{-1}(h) = (Sj(h^{(1)}))h^{(2)}$  is the convolution-inverse of  $j$  and makes  $P$  a cleft cocycle extension with

$$\begin{aligned}\sigma(h, g) &= j(h_{(1)})j(g_{(1)})(Sj(h_{(2)}^{(1)}g_{(2)}^{(1)}))h_{(2)}^{(2)}g_{(2)}^{(2)} \\ \sigma^{-1}(h, g) &= j(h_{(1)}g_{(1)})(Sj(g_{(2)}^{(1)}g_{(2)}^{(2)}))(Sj(h_{(2)}^{(1)}h_{(2)}^{(2)})\end{aligned}$$

and trivial action. Moreover, the coalgebra is cocleft and the extension is a cocycle bicrossproduct  $P = B_\sigma \blacktriangleright H$ . The concrete example in [18, Ex. 6.3.14]  $B = \mathbb{C}\mathbb{Z} = \mathbb{C}[c, c^{-1}]$  and  $H = U_q(Lsl_2)$  the quantum loop group version of  $U_q(sl_2)$ . By definition, this can be taken as generated by  $k, k^{-1}, e_0, f_0, e_1, f_1$  with

$$\begin{aligned}ke_0k^{-1} &= q^{-2}e_0, & ke_1k^{-1} &= q^2e_1, & kfk^{-1} &= q^2f_0, & kf_1k^{-1} &= q^{-2}f_1 \\ q^2e_0f_0 - f_0e_0 &= \frac{k^{-2} - 1}{q - q^{-1}}, & q^2e_1f_1 - f_1e_1 &= \frac{k^2 - 1}{q - q^{-1}}\end{aligned}$$

and certain  $q$ -Serre relations involving powers of  $e_0, f_0$ . The coproduct and antipode are  $k$  grouplike,

$$\Delta e_0 = e_0 \otimes k^{-1} + 1 \otimes e_0, \quad \Delta e_1 = e_1 \otimes k + 1 \otimes e_1, \quad S e_0 = -e_0 k, \quad S e_1 = -e_0 k^{-1}$$

and similarly for  $f_i$ . The algebra has a grading  $||$  by the total power of  $e_0, f_0$  when expressions are ordered with all  $e_i$  to the left of all  $f_j$ , which we view as a right coaction of  $B$  by  $\Delta_R h = h \otimes c^{|h|}$  for any  $h \in U_q(Lsl_2)$  of homogeneous degree, and we make the associated cross product coalgebra. The 2-cocycle has, for example

$$\sigma(f_0, e_0) = \frac{1 - c^2}{q - q^{-1}}$$

and  $\mathbb{C}\mathbb{Z}_\sigma \blacktriangleright U_q(Lsl_2)$  is isomorphic to  $\widehat{U_q(sl_2)}$ . (The latter has similar relations but with some central elements  $c$  appearing from the cocycle.)

Here, we observe that by the same methods one can compute that

$$\begin{aligned}\sigma(e_1, k^{\pm 1}) &= \sigma(k^{\pm 1}, e_0) = \sigma(e_0, k) = 0, & \sigma(k^{-1}, k) &= \sigma(k, k^{-1}) = 1 \\ \sigma^{-1}(k^{\pm 1}, S e_0) &= \sigma^{-1}(S e_0, k^{\pm 1}) = 0, & \sigma^{-1}(k^{-1}, k) &= \sigma^{-1}(k, k^{-1}) = 1.\end{aligned}$$

Hence the antipode of the cocycle Hopf algebroid  $B^e \#_\sigma H$  by Proposition 3.10, simplifies to

$$\begin{aligned}S(b \otimes b' \#_\sigma e_0) &= b' \otimes \sigma^{-1}(S^2(k^{-1}) \otimes S(e_0))b \#_\sigma S(k^{-1}) + b' \otimes \sigma^{-1}(S^2(e_0) \otimes S(1))b \#_\sigma S(e_0) \\ &\quad + b' \otimes \sigma^{-1}(S^2(1) \otimes S(1))b \#_\sigma S(e_0) \\ &= b' \otimes \sigma^{-1}(S^2(1) \otimes S(1))b \#_\sigma S(e_0) = b' \otimes b \#_\sigma S(e_0)\end{aligned}$$

and similarly

$$S^{-1}(b \otimes b' \#_\sigma e_0) = b' \otimes b \#_\sigma S^{-1}(e_0)$$

i.e. just given by flip and  $S^{\pm 1}$  on  $H$ . The same applies with  $e_0$  can be replaced by  $e_1, f_i$  and  $k^\pm$ . We also know from our twisting theory that this Hopf algebroid is isomorphic to a cotwist of  $B^e \otimes H$  (as the action is trivial) by a cotwist  $\tilde{\sigma}$  as in Lemma 3.11, for example

$$\tilde{\sigma}(b \otimes b' \#_\sigma e_0, d \otimes d' \#_\sigma f_0) = \frac{bdd'b(1 - c^2)}{q - q^{-1}}.$$

**5.3. Quantum Weyl group.** The abstract structure here (a left-right swap of [18, Prop. 6.3.12]) starts with a Hopf algebra  $B$  with Drinfeld twist cocycle so that  ${}^\psi B$  (say) is the twisted Hopf algebra with conjugated coproduct. We also need  $T : B \rightarrow {}^\psi B$  a Hopf algebra map such that

$$x^{-1}ax = T^2(a), \quad T(x) = x, \quad \Delta x = (x \otimes x)((T \otimes T)(\psi))\psi, \quad \epsilon x = 1$$

for some invertible  $x \in B$ . Then there is a cocycle cross product algebra  $B \#_\sigma \mathbb{C}\mathbb{Z}_2$  where if  $w \in \mathbb{C}\mathbb{Z}_2$  is the generator of  $\mathbb{Z}_2$ , the cocycle and action are

$$w \triangleright b = T(b), \quad \sigma(w, w) = x^{-1} \quad \sigma(w, 1) = \sigma(1, w) = \sigma(1, 1) = 1.$$

The result is then a cocycle bicrossproduct Hopf algebra  $B_{\sigma \blacktriangleleft \psi} \mathbb{C}\mathbb{Z}_2$  with coproduct

$$\Delta(b \otimes 1) = b_{(1)} \otimes 1 \otimes b_{(2)} \otimes 1, \quad \Delta(b \otimes w) = b_{(1)} \psi^1 \otimes w \otimes b_{(2)} \psi^2 \otimes w$$

where  $\psi = \psi^1 \otimes \psi^2$  (sum understood). The concrete example in [18, Ex 6.3.13] is to take  $B$  a ribbon Hopf algebra with ribbon element  $\nu$  and  $T$  an algebra automorphism and anticoalgebra map such that  $T^2 = \text{id}$  and  $T(\nu) = \nu$ . The ribbon element is central and hence so is the image of  $\sigma$ . We take  $\psi = \mathcal{R}$  the quasitriangular structure regarded as a cocycle. The result of the construction is an extension of  $B$  such that  $w^2 = \nu$  in the extended algebra. The required ‘Lusztig automorphism’  $T$  exists for the standard  $q$ -deformation enveloping algebras[21] as well as for reduced quantum groups at roots of unity. The ribbon element in the latter case is given explicitly in [15].

Here, we compute the antipode for the associated cocycle Hopf algebroid  $B^e \#_\sigma \mathbb{C}\mathbb{Z}_2$  using Proposition 3.10 as

$$\begin{aligned} S^{\pm 1}(b \otimes b' \#_\sigma 1) &= b' \otimes b \#_\sigma 1 \\ S(b \otimes b' \#_\sigma w) &= b' \otimes \sigma^{-1}(w \otimes w) T(b) \#_\sigma w^{-1} = b' \otimes \nu^{-1} T(b) \#_\sigma w, \\ S^{-1}(b \otimes b' \#_\sigma w) &= T(b') \sigma(w \otimes w) \otimes b \#_\sigma w^{-1} = T(b') \nu \otimes b \#_\sigma w. \end{aligned}$$

Also, we know that this Hopf algebroid is a cotwist of  $B^e \# \mathbb{C}\mathbb{Z}_2$  by a cocycle in Lemma 3.11,

$$\begin{aligned} \tilde{\sigma}(b \otimes b' \# 1, c \otimes c' \# 1) &= b c c' b', \quad \tilde{\sigma}(b \otimes b' \# 1, c \otimes c' \# w) = b c (w \triangleright c') b' = b c T(c') b', \\ \tilde{\sigma}(b \otimes b' \# w, c \otimes c' \# 1) &= b (w \triangleright (c c' b')) = b T(c c' b'), \\ \tilde{\sigma}(b \otimes b' \# w, c \otimes c' \# w) &= b (w \triangleright c) \sigma(w \otimes w) ((w)^2 \triangleright c') (w \triangleright b') = b T(c) \nu c' T(b'). \end{aligned}$$

with antipode

$$\begin{aligned} (S^{\tilde{\sigma}})^{\pm 1}(b \otimes b' \# 1) &= b' \otimes b \# 1 \\ (S^{\tilde{\sigma}})(b \otimes b' \# w) &= b' \sigma^{-1}(w \otimes w) \otimes T(b) \# w^{-1} = b' \nu^{-1} \otimes T(b) \# w, \\ (S^{\tilde{\sigma}})^{-1}(b \otimes b' \# w) &= T(b') \otimes b (w \triangleright \sigma(w \otimes w)) \# w = T(b') \otimes b \nu \# w. \end{aligned}$$

where  $w^{-1} = w$  and in the last step we use  $w \triangleright \nu = \nu$ .

**Acknowledgment:** XH would like to thank Prof. Dr. Giovanni Landi, Dr. Song Cheng and A. Ghobadi for discussions. XH was supported by Marie Curie Fellowship HADG - 101027463 agreed between QMUL and the European Commission.

## REFERENCES

- [1] P. Aschieri, P. Bieliavsky, C. Pagani, A. Schenkel, *Noncommutative principal bundles through twist deformation*. Commun. Math. Phys. 352 (2017) 287–344.
- [2] E.J. Beggs and S. Majid, *Quantum Riemannian Geometry*, Grundlehren der mathematischen Wissenschaften, vol. 355, Springer (2020) 809pp.
- [3] J. Bichon, *Hopf-Galois objects and cogroupoids*. Rev. Un. Mat. Argentina, 5 (2014) 11–69.
- [4] J. Bichon and G. Carnovale *Lazy cohomology: an analogue of the Schur multiplier for arbitrary Hopf algebras* J. Pure and Appl. Algebra, 204 (2006) 627–665
- [5] G. Böhm, *Hopf algebroids*. Handbook of Algebra, Vol. 6, North-Holland, 2009, Pages 173–235.
- [6] G. Böhm, K. Szlachányi *Hopf algebroids with bijective antipodes: axioms, integrals, and duals*. Journal of Algebra, 274(2):708-750, 2004.

- [7] T. Brzeziński, *Translation map in quantum principal bundles*. J. Geom. Phys. 20 (1996) 349–370.
- [8] T. Brzezinski and S. Majid, *Quantum group gauge theory on quantum spaces*, Comm. Math. Phys. 157 (1993) 591–638.
- [9] T. Brzeziński, R. Wisbauer, *Corings and comodules*. London Mathematical Society Lecture Notes 309, CUP 2003.
- [10] Y. Doi, *equivalent crossed products for a Hopf algebra*. Comm. Alg. 17 (1989), 3053-3085.
- [11] Y. Doi, M. Takauchi, *Cleft comodule algebras for a bialgebra*. Comm. Alg. 14 (1986), 801-818.
- [12] A. Ghobadi, *Hopf algebroids, bimodule connections and noncommutative geometry*, arXiv:2001.08673 [math.qa].
- [13] X. Han, G. Landi, *Gauge groups and bialgebroids*. Letters in Mathematical Physics volume 111, Article number: 140 (2021).
- [14] C. Kassel, *Principal fiber bundles in non-commutative geometry*. In: Quantization, geometry and noncommutative structures in mathematics and physics, Mathematical Physics Studies, Springer 2017, pp. 75–133.
- [15] V. Lyubashenko and S. Majid, *Braided groups and quantum Fourier transform*, J. Algebra. 166 (1994) 506–528
- [16] K. Mackenzie, *General theory of Lie groupoids and Lie algebroids*. London Mathematical Society Lecture Notes Series 213, CUP 2005.
- [17] S. Majid, *More examples of bicrossproduct and double cross product Hopf algebras*, Isr. J. Math. 72 (1990) 133–148
- [18] S. Majid, *Foundations of quantum group theory*. CUP 1995 and 2000.
- [19] S. Majid, *Cross product quantisation, nonabelian cohomology and twisting of Hopf algebras*. In H.-D. Doebner et al., eds., Generalized Symmetries in Physics. World Sci. (1994) 13-41 (arXiv:hep-th/9311184v1)
- [20] A. Masuoka, *Cleft extensions for a Hopf algebra generated by a nearly primitive element*. Comm. Algebra 22 (1994) 4537–4559.
- [21] S. Majid and Ya. S. Soibelman, *Bicrossproduct structure of the quantum Weyl group*, J. Algebra 163 (1994) 68–87
- [22] S. Montgomery, *Hopf algebras and their actions on rings*. AMS 1993.
- [23] P. Schauenburg, *Hopf bi-Galois extensions*. Commun. Algebra 24 (1996) 3797–3825.
- [24] P. Schauenburg, *Duals and doubles of quantum groupoids ( $\times_R$ -Hopf algebras)*. in ‘New trends in Hopf algebra theory’, (Proc. of the colloquium on quantum groups and Hopf algebras, La Falda, Sierras de Cordoba, Argentina, 1999), AMS Contemporary Mathematics 267 (2000), 273–299
- [25] M. Sweedler, *Hopf Algebras*, Benjamin, 1969.
- [26] K. Szlachanyi, *Monoidal Morita equivalence*. arXiv:math/0410407 [math.QA].

QUEEN MARY UNIVERSITY OF LONDON, SCHOOL OF MATHEMATICAL SCIENCES, MILE END RD, LONDON E1 4NS, UK

*Email address:* x.h.han@qmul.ac.uk, s.majid@qmul.ac.uk