

# Entanglement dynamics: Generalized master equation for uniformly accelerated two-level systems

M. S. Soares\*, N. F. Svaiter<sup>†</sup>

Centro Brasileiro de Pesquisas Físicas

Rua Xavier Sigaud, 150 - Urca, Rio de Janeiro - RJ, 22290-180, Brazil

G. Menezes<sup>‡</sup>

Universidade Federal Rural do Rio de Janeiro,

Rodovia BR 465, Km 07,

Seropédica-RJ, - Brazil

## Abstract

We propose a new form for the quantum master equation in the theory of open quantum systems. This new formalism allows one to describe the dynamics of two-level systems moving along different hyperbolic trajectories with distinct proper times. In the Born-Markov approximation, we consider a quantum massless scalar field coupled with two-level systems. Starting from a separable state we show the emergence of entanglement harvesting. For different proper accelerations we verify also the entanglement sudden death.

## 1 Introduction

One of the consequences of describing quantum systems by infinite number of degrees of freedom is the non-uniqueness of the vacuum state. This situation appears in a generic curved space-time, where there is not a natural decomposition of the Fourier modes in terms of negative and positive frequency parts [1, 2]. Even for observers in Minkowski space-time, there are situations where the vacuum state is not the usual one [3–5]. For instance, a uniformly accelerated observer in flat space-time perceives the Minkowski vacuum as a thermal bath. This is known as the Unruh-Davies effect and has a close connection to the Hawking effect [6, 7]. The discussion of quantum fields in non-inertial frames and related subjects have been a topic under vigorous investigation throughout the years [8–26].

An apparatus device in such studies is the Unruh-deWitt particle detector [27, 28]. It is defined as an idealized two-level system coupled with a scalar field through a monopole interaction. Other descriptions for particle detectors are also possible as, for example, is given by the Glauber theory [29]. A recent discussion on particle-detector models can be found in Refs. [30]. In this work, the authors have highlighted the importance of the Glauber model in the clarification on the different interpretations of the measurements performed by an accelerated detector. See also Refs. [31–35].

An essential tool in quantum computation and quantum information theory is the quantum entanglement [36]. The literature has examined sources of entangled quantum systems, recurrently found in solid-state physics and atoms in cavity electrodynamics. In fact, several

proposals to generate entangled states in systems of two-level systems coupled to bosonic fields were established [37–41].

In recent years, quantum entanglement has been analysed through the lens of the relativistic quantum information theory, which studies the behaviour of atoms interacting with relativistic quantum fields [42–56]. A well known phenomenon in this scenario is the so-called entanglement degradation, when correlated states become uncorrelated by the interaction with a quantum field [57–61]. Another effect is the entanglement harvesting. Atoms initially prepared in a separated state can extract entanglement from the quantum vacuum [62–68]. Vacuum fluctuations are able to act as a source of entanglement for atoms coupled with quantum fields. This is realised when atoms are uniformly accelerated, since the Minkowski vacuum state can be conceived as an entangled state between the left and right Rindler wedges when stated on the Rindler vacuum [5]. In any case, it is important to bear in mind that entanglement harvesting protocol can be considered as legitimate only when the detectors are not able to communicate [69].

Relativistic quantum entanglement has been investigated in different setups. Many of these works reveal how entanglement is an observer-dependent quantity, in addition to unraveling how the effects of entanglement dynamics can depend on the space-time separation between atoms, as well as the space-time dimension [70–82]. Here we wish to report progress on the discussion of entanglement dynamics for two-level systems in non inertial reference frames. We treat this system, namely  $\mathcal{A}$ , coupled with a larger system  $\mathcal{B}$  which is the quantum field and make use of the theory of open quantum systems to describe its dynamics [83, 84]. We wish to generalize the construction of the *master equation*, which is the object that describes the time evolution of the density matrix of the subsystem  $\mathcal{A}$  [85]. Benatti and Floreanini have derived this master equation for non-inertial atoms weakly interacting with a quantum scalar field [86]. See also Ref. [87–89]. Recently their results were generalized to the case of atoms coupled with an electromagnetic field [58]. Both of these works describe the entanglement dynamics for atoms with the same proper acceleration. Here we will extend the formalism to a more general scenario, since we consider that the atoms travel along different world lines which are parametrized by different proper times.

The generalization presented in this work allows one to investigate other aspects that can influence the dynamics of the system in addition to those already studied in previous works, such as the orthogonal distance to the motion between atoms and the polarization of the atoms in the electromagnetic field case. We investigate, within this method, the implications of different accelerations for entanglement degradation and entanglement harvesting. In order to quantify the content of entanglement between the two-level system, we calculate the concurrence introduced by Wootters [90].

The organization of this paper is as follows. In Sec. 2 we present the derivation of the master equation for two-level systems travelling in different world lines. In Sec. 3 we apply this equation for two two-level systems interacting with a scalar field. In Sec. 4 we discuss entanglement dynamics. Conclusions are given in Sec. 5. We use units such that  $\hbar = c = k_B = 1$ .

## 2 Generalized Master Equation

In this section we derive a generalized form of the master equation [91]. The term *generalized* is been used since we are working with two-level systems travelling along different world lines with different proper times. The Hamiltonian of the system that we are interested in can be

written as

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_{\text{int}} = \mathcal{H}_A + \mathcal{H}_F + \mathcal{H}_{\text{int}}. \quad (1)$$

Usually, the interaction Hamiltonian ( $\mathcal{H}_{\text{int}}$ ) has the form

$$\mathcal{H}_{\text{int}} = \sum_i \sum_{\alpha} A_i^{(\alpha)} \otimes B_i^{(\alpha)}, \quad (2)$$

where  $A$  denotes the two-level systems operators and  $B$  are the quantum field operators. The  $i$ -sum is associated with the Hilbert space dimensionality and the  $\alpha$ -sum counts the number of two-level systems. In order to describe the evolution of the subsystem  $\mathcal{A}$  we choose a parameter  $\eta$ , which will be duly specified in due course. At the moment we only make the reasonable assumption that the proper times of the two-level systems are  $\tau_{\alpha} = \tau_{\alpha}(\eta)$ . The interaction Hamiltonian in the interaction picture is then

$$\mathcal{H}_{\text{int}}(\eta) = \sum_{\omega} \sum_i \sum_{\alpha} e^{-i\omega\tau_{\alpha}(\eta)} A_i^{(\alpha)}(\omega) \otimes B_i^{(\alpha)}[x(\tau_{\alpha}(\eta))] \frac{d\tau_{\alpha}(\eta)}{d\eta}. \quad (3)$$

In order to describe the dynamics of the subsystem  $\mathcal{A}$  we make use of the density matrix theory. The density matrix of the whole system ( $\rho_{AB}$ ) in the interaction picture is described by the Liouville-von Neumann equation

$$\frac{d}{d\eta} \tilde{\rho}_{AB}(\eta) = -i [\mathcal{H}_{\text{int}}(\eta), \tilde{\rho}_{AB}(\eta)]. \quad (4)$$

Although this equation gives the dynamics of the whole system, we are interested only in the behaviour of the subsystem  $A$ . Our aim is to construct an equation for the reduced density matrix  $\rho_A(\eta)$  of the two-level systems which preserves all the properties of the density matrices. By making use of perturbation theory and considering the weak-coupling approximation, we find that

$$\Delta \tilde{\rho}_{AB}(\eta) = -i \int_{\eta}^{\eta+\Delta\eta} d\eta_1 [\mathcal{H}_{\text{int}}(\eta_1), \tilde{\rho}_{AB}(\eta_1)] - \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 [\mathcal{H}_{\text{int}}(\eta_1), [\mathcal{H}_{\text{int}}(\eta_2), \tilde{\rho}_{AB}(\eta)]] \quad (5)$$

where we have kept only terms up to the second order in interaction Hamiltonian. As we wish to study the evolution of the reduced density matrix  $\rho_A$  of the subsystem  $\mathcal{A}$ , we trace over the field states,  $\tilde{\rho}_A(\eta) = \text{Tr}_B [\tilde{\rho}_{AB}(\eta)]$ , to obtain that

$$\begin{aligned} \Delta \tilde{\rho}_A(\eta) = & -i \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \text{Tr}_B [\mathcal{H}_{\text{int}}(\eta_1), \tilde{\rho}_{AB}(\eta_1)] \\ & - \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 \text{Tr}_B [\mathcal{H}_{\text{int}}(\eta_1), [\mathcal{H}_{\text{int}}(\eta_2), \tilde{\rho}_{AB}(\eta)]] . \end{aligned} \quad (6)$$

Before proceeding, we must resort to some additional approximations. We assume that there are two distinct time scales. One is specified by  $\tau_B$ , the characteristic time associated with the internal correlations in the system  $\mathcal{B}$ . The other time scale is  $\tau_A$  which describes the time evolution of the density matrix in the presence of the interaction between the two systems. We assume that

$$\tau_B \ll \Delta\eta \ll \tau_A, \quad (7)$$

which implies that

$$\frac{\Delta \tilde{\rho}_A(\eta)}{\Delta \eta} \approx \frac{\tilde{\rho}_A(\eta)}{\tau_A}. \quad (8)$$

This means that the time scale relaxation of the system  $\mathcal{B}$  is smaller than the time scale relaxation of the two-level systems. Therefore, when the correlation between the two systems starts, the field has enough time to reach the thermodynamic equilibrium. This implies that we are within the regime of the *Born approximation*, which enables one to neglect initial correlations induced by the interactions between the subsystems:

$$\tilde{\rho}_{AB}(\eta) \approx \tilde{\rho}_A(\eta) \otimes \tilde{\rho}_B, \quad (9)$$

where  $\tilde{\rho}_B$  is the reduced density operator associated with subsystem  $\mathcal{B}$ . Furthermore, such approximations also allow us to conclude that the operator  $\tilde{\rho}_B$  can be approximately regarded as being stationary:

$$\tilde{\rho}_B(\eta) \approx \sigma_B \rightarrow [\sigma_B, \mathcal{H}_B] = 0. \quad (10)$$

As mentioned above, the subsystem  $\mathcal{B}$  is associated with the quantum field to which the two-level systems couple. Therefore, one expects that

$$\text{Tr}_B \left\{ B_i^{(\alpha)} [x(\tau_\alpha(\eta))] \sigma_B \right\} = 0. \quad (11)$$

So we can rewrite Eq. (6) as

$$\begin{aligned} \Delta \tilde{\rho}_A(\eta) = & - \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 \\ & \times \text{Tr}_B [\mathcal{H}_{\text{int}}(\eta_1), [\mathcal{H}_{\text{int}}(\eta_2), \tilde{\rho}_{AB}(\eta)]], \end{aligned} \quad (12)$$

Expanding the commutators and employing the aforementioned discussion of time intervals, we find that

$$\begin{aligned} \frac{\Delta \tilde{\rho}_A(\eta)}{\Delta \eta} = & \frac{1}{\Delta \eta} \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 \\ & \times \text{Tr}_B [\mathcal{H}_{\text{int}}(\eta_2) \tilde{\rho}_A(\eta) \otimes \sigma_B \mathcal{H}_{\text{int}}(\eta_1) \\ & - \mathcal{H}_{\text{int}}(\eta_1) \mathcal{H}_{\text{int}}(\eta_2) \tilde{\rho}_A(\eta) \otimes \sigma_B] + \text{h.c.} \end{aligned} \quad (13)$$

Using Eq. (3):

$$\begin{aligned} \frac{\Delta \tilde{\rho}_A(\eta)}{\Delta \eta} = & \frac{1}{\Delta \eta} \sum_{\omega, \omega'} \sum_{i, j} \sum_{\alpha, \beta} \int_{\eta}^{\eta+\Delta\eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 e^{-i\omega\tau_\beta(\eta_2)} e^{i\omega'\tau_\alpha(\eta_1)} \\ & \times \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega') \text{Tr}_B \left[ B_j^{(\beta)}(\eta_2) \sigma_B B_i^{\dagger(\alpha)}(\eta_1) \right] \right. \\ & \left. - A_i^{\dagger(\alpha)}(\omega') A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) \text{Tr}_B \left[ B_i^{\dagger(\alpha)}(\eta_1) B_j^{(\beta)}(\eta_2) \sigma_B \right] \right) \frac{d\tau_\beta(\eta_1)}{d\eta_1} \frac{d\tau_\alpha(\eta_2)}{d\eta_2} + \text{h.c.}, \end{aligned} \quad (14)$$

where for brevity we wrote  $B_i^{(\alpha)} [x(\tau_\alpha(\eta))] = B_i^{(\alpha)}(\eta)$ . We define the field correlation function as

$$\text{Tr}_B \left[ B_i^{\dagger(\alpha)}(\eta_1) B_j^{(\beta)}(\eta_2) \sigma_B \right] = G_{ij}^{(\alpha\beta)}(\tau_\beta(\eta_1), \tau_\alpha(\eta_2)). \quad (15)$$

In many cases that we are interested, this function is the positive Wightman function, which depends only on the difference of the proper times of the two-level systems  $\tau_\beta(\eta_1) - \tau_\alpha(\eta_2)$ . Here this function only depends on the difference of the parameters  $\eta_1 - \eta_2$ . Therefore

$$G_{ij}^{(\alpha\beta)}(\tau_\beta(\eta_1), \tau_\alpha(\eta_2)) = G_{ij}^{(\alpha\beta)}(\eta_1 - \eta_2). \quad (16)$$

Using the cyclic property of the trace, and after a simple algebraic manipulation, we find that

$$\begin{aligned} \frac{\Delta \tilde{\rho}_A(\eta)}{\Delta \eta} &= \frac{1}{\Delta \eta} \sum_{\omega, \omega'} \sum_{i, j} \sum_{\alpha, \beta} \int_{\eta}^{\eta + \Delta \eta} d\eta_1 \int_{\eta}^{\eta_1} d\eta_2 e^{i(\omega' - \omega)\tau_\alpha(\eta_1)} e^{i\omega(\tau_\alpha(\eta_1) - \tau_\beta(\eta_2))} G_{ij}^{(\alpha\beta)}(\eta_1 - \eta_2) \\ &\times \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega') - A_i^{\dagger(\alpha)}(\omega') A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) \right) \frac{d\tau_\beta(\eta_1)}{d\eta_1} \frac{d\tau_\alpha(\eta_2)}{d\eta_2} + h.c. \end{aligned} \quad (17)$$

Up to now we have not specified the behaviour of the functions  $\tau_\alpha(\eta)$ . For simplicity, let us assume a linear relation of the form

$$\tau_\alpha(\eta_i) = a_\alpha \eta_i, \quad (18)$$

for some constant  $a_\alpha$ . This is not a purely academic exercise; indeed this relation is obeyed in several interesting physical situations, as, for example, for some classes of observers in Rindler and Schwarzschild space-times – in the former case,  $a_\alpha$  can be associated with the proper acceleration of the two-level systems.

Defining  $s = \eta_1 - \eta_2$  and performing a simple change of coordinates, one obtains that

$$\begin{aligned} \frac{\Delta \tilde{\rho}_A(\eta)}{\Delta \eta} &= \sum_{\omega, \omega'} \sum_{i, j} \sum_{\alpha, \beta} J(\omega' a_\alpha - \omega a_\beta) W_{ij}^{(\alpha\beta)}(a_\beta \omega) \\ &\times \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega') - A_i^{\dagger(\alpha)}(\omega') A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) \right) + h.c., \end{aligned} \quad (19)$$

where we have defined the following functions

$$J(\omega' a_\alpha - \omega a_\beta) = \int_{\eta}^{\eta + \Delta \eta} d\eta_1 \frac{e^{i(\omega' a_\alpha - \omega a_\beta) \eta_1}}{\Delta \eta}, \quad (20)$$

and

$$W_{ij}^{(\alpha\beta)}(a_\beta \omega) = a_\alpha a_\beta \int_0^\infty ds e^{i\omega a_\beta s} G_{ij}^{(\alpha\beta)}(s). \quad (21)$$

Notice that we have moved the upper limit of integration over  $ds$  to infinity as  $\tau_B \ll \Delta \eta$  and the lower limit of the integral over  $d\eta_1$  was simply taken to be  $\eta$  since only small values of  $s$  contribute significantly to the integral.

In the following we will study the form of the function  $J(x)$ . We can solve explicitly the integral:

$$\begin{aligned} J(\omega' a_\alpha - \omega a_\beta) &= \int_{\eta}^{\eta + \Delta \eta} d\eta_1 \frac{e^{i(\omega' a_\alpha - \omega a_\beta) \eta_1}}{\Delta \eta} \\ &= e^{i(\omega' a_\alpha - \omega a_\beta) \eta} F(\omega' a_\alpha - \omega a_\beta), \end{aligned} \quad (22)$$

with

$$F(\omega' a_\alpha - \omega a_\beta) = e^{i(\omega' a_\alpha - \omega a_\beta) \frac{\eta}{2}} \frac{\sin \left[ \left( \frac{\omega' a_\alpha - \omega a_\beta}{2} \right) \Delta \eta \right]}{\left[ \left( \frac{\omega' a_\alpha - \omega a_\beta}{2} \right) \Delta \eta \right]}. \quad (23)$$

We observe that, for  $|\omega' a_\alpha - \omega a_\beta| \ll (\Delta\eta)^{-1}$ , the function  $|F|$  is very close to one, and, on the other hand, for  $|\omega' a_\alpha - \omega a_\beta| \gg (\Delta\eta)^{-1}$ ,  $|F|$  is very close to zero. In other words,  $|F|$  is sharply peaked around the value  $|\omega' a_\alpha - \omega a_\beta| = 0$ . Therefore, within our assumptions, we can make the approximation

$$J(\omega' a_\alpha - \omega a_\beta) \approx e^{i(\omega' a_\alpha - \omega a_\beta)\eta} \delta(\omega' a_\alpha - \omega a_\beta). \quad (24)$$

Let us now obtain the standard form of the master equation. By *standard form* we mean the derivation known by the Lindblad-Kossakowski equation [92, 93]. We start by inserting Eq. (24) into equation (19). We obtain

$$\begin{aligned} \frac{\Delta \tilde{\rho}_A(\eta)}{\Delta\eta} = & \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} W_{ij}^{(\alpha\beta)}(a_\beta\omega) \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)} \left( \frac{a_\beta}{a_\alpha} \omega \right) - A_i^{\dagger(\alpha)} \left( \frac{a_\beta}{a_\alpha} \omega \right) A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) \right) \\ & + \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} W_{ij}^{*(\alpha\beta)}(a_\alpha\omega) \left( A_i^{(\alpha)}(\omega) \tilde{\rho}_A(\eta) A_j^{\dagger(\beta)} \left( \frac{a_\alpha\omega}{a_\beta} \right) - \tilde{\rho}_A(\eta) A_j^{\dagger(\beta)} \left( \frac{a_\alpha\omega}{a_\beta} \right) A_i^{(\alpha)}(\omega) \right). \end{aligned} \quad (25)$$

This equations describes the change of  $\tilde{\rho}_A(\eta)$  within the interval  $(\eta, \eta + \Delta\eta)$ . The ratio  $\Delta \tilde{\rho}_A / \Delta\eta$  can be thought as an operation of averaging which smooths out rapid variations of  $\tilde{\rho}_A(\eta)$ . The investigation of the time evolution on a time scale  $\tau_A \ll \Delta\eta$  allows us to neglect these small and rapid fluctuations. This coarse-graining approximation is also known as the *Markovian approximation*. Moreover, in our case the operators  $A(\omega)$  will take the form of linear combinations of the Pauli matrices. Then  $\omega$  is to be associated with the eigenvalues of the atomic Hamiltonian ( $\mathcal{H}_A$ ). Therefore, we can write  $A(\lambda\omega) = A(\omega)$  for some positive constant  $\lambda$ . Hence

$$\begin{aligned} \frac{d}{d\eta} \tilde{\rho}_A(\eta) = & \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} W_{ij}^{(\alpha\beta)}(a_\beta\omega) \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega) - A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) \right) \\ & + \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} W_{ji}^{*(\beta\alpha)}(a_\beta\omega) \left( A_j^{(\beta)}(\omega) \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega) - \tilde{\rho}_A(\eta) A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega) \right). \end{aligned} \quad (26)$$

where we have changed  $i \longleftrightarrow j$  and  $\alpha \longleftrightarrow \beta$  in the second term of the right-hand side. Moving to the Schrödinger representation,  $\tilde{\rho}_A(\eta) = e^{i\mathcal{H}_A\eta} \rho_A(\eta) e^{-i\mathcal{H}_A\eta}$ , and also

$$e^{i\mathcal{H}_A\eta} A_i^{(\alpha)}(\omega) e^{-i\mathcal{H}_A\eta} = e^{-i\omega\eta} A_i^{(\alpha)}(\omega), \quad (27)$$

and defining the following functions

$$\Gamma_{ij}^{(\alpha\beta)}(a_\beta, \omega) = W_{ij}^{(\alpha\beta)}(a_\beta\omega) + W_{ji}^{*(\beta\alpha)}(a_\beta\omega), \quad (28)$$

$$\Delta_{ij}^{(\alpha\beta)}(a_\beta, \omega) = \frac{W_{ij}^{(\alpha\beta)}(a_\beta\omega) - W_{ji}^{*(\beta\alpha)}(a_\beta\omega)}{2i}, \quad (29)$$

we obtain that

$$\begin{aligned} \dot{\rho}_A(\eta) = & -i [\mathcal{H}_A, \rho_A(\eta)] - i \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} \Delta_{ij}^{(\alpha\beta)}(a_\beta, \omega) \left[ A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega), \rho_A(\eta) \right] \\ & + \frac{1}{2} \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} \Gamma_{ij}^{(\alpha\beta)}(a_\beta, \omega) \left( 2A_j^{(\beta)}(\omega) \rho_A(\eta) A_i^{\dagger(\alpha)}(\omega) - \left\{ A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega), \rho_A(\eta) \right\} \right), \end{aligned} \quad (30)$$

where  $\dot{F} = dF/d\eta$  and the commutator and the anticommutator of operators  $C, D$  are defined in the usual way. If we define the following effective Hamiltonian

$$\mathcal{H}_{\text{eff}} \equiv \mathcal{H}_A + \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} \Delta_{ij}^{(\alpha\beta)}(a_{\beta}, \omega) A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega), \quad (31)$$

which describes the unitary evolution of the subsystem, we finally obtain the standard form of the generalized master equation describing the time evolution of atoms travelling along different world lines:

$$\dot{\rho}_A(\eta) = -i [\mathcal{H}_{\text{eff}}, \rho_A(\eta)] + \mathcal{L} \{ \rho_A(\eta) \}, \quad (32)$$

where the term  $\mathcal{L} \{ \rho_A(\eta) \}$ , the dissipative contribution to generalized master equation, is given by

$$\mathcal{L} \{ \rho_A(\eta) \} = \frac{1}{2} \sum_{\omega} \sum_{i,j} \sum_{\alpha,\beta} \Gamma_{ij}^{(\alpha\beta)}(a_{\beta}, \omega) \left( 2A_j^{(\beta)}(\omega) \rho_A(\eta) A_i^{\dagger(\alpha)}(\omega) - \left\{ A_i^{\dagger(\alpha)}(\omega) A_j^{(\beta)}(\omega), \rho_A(\eta) \right\} \right). \quad (33)$$

As we shall see, it is the non-unitary term  $\mathcal{L} \{ \rho_A(\eta) \}$  that is responsible for decoherence effects as well as entanglement generation. Observe the generality of our derivation of the generalized master equation, that can be applied to a scalar or a vector field.

### 3 Generalized master equation for a couple of two-level systems

Our task now is to specify the generalized master equation for one particular case. The generalization of the results presented in Ref. [86] will be obtained. We consider the situation of two identical two-level systems interacting with a quantum massless scalar field  $\phi(x)$ . The parameter we choose to describe the time evolution of the system is the proper time of one of the two-level systems. The different contributions to the Hamiltonian of the system can be cast in the following form

$$H = H_A + H_F + H_{\text{int}} \quad (34)$$

where

$$H_A = \frac{\omega_0}{2} \left[ S_1^z \otimes \mathbb{1}_2 \frac{d\tau_1(\tau)}{d\tau} + \mathbb{1}_1 \otimes S_2^z \frac{d\tau_2(\tau)}{d\tau} \right], \quad (35)$$

$$H_F = \frac{1}{2} \int d^3\mathbf{x} \left[ (\dot{\phi}(x))^2 + (\nabla\phi(x))^2 \right], \quad (36)$$

$$H_{\text{int}} = \sum_{j=1}^2 \mu_j \chi_j[\tau_j(\tau)] m^{(j)}(\tau_j(\tau)) \phi[x_j(\tau_j(\tau))] \frac{d\tau_j(\tau)}{d\tau}. \quad (37)$$

In the above equation,  $H_A$  describes the Hamiltonian of the two-level system,  $H_F$  is the scalar field Hamiltonian and  $H_{\text{int}}$  describes the interaction Hamiltonian. In addition,  $\mu_j$  is the coupling constant of the  $j$ -th two-level system,  $\chi_j(x)$  is the switching function and  $m^{(j)}$  is the monopole moment of the two-level systems, given by

$$\begin{aligned} m^{(j)}(\tau_j(\tau)) &= |e^j\rangle \langle g^j| e^{i\omega_0\tau_j(\tau)} + |g^j\rangle \langle e^j| e^{-i\omega_0\tau_j(\tau)} \\ &= S_+^{(j)} e^{i\omega_0\tau_j(\tau)} + S_-^{(j)} e^{-i\omega_0\tau_j(\tau)}. \end{aligned} \quad (38)$$

For simplicity, henceforth we take  $\mu_1 = \mu_2 = \mu$ . We would like to study the situation where both two-level systems are in uniformly accelerated world lines. It is known that the coordinates of these two-level systems are given by the so called *Rindler coordinates*. These coordinates for uniform acceleration occurring only in the  $x$ -direction are expressed as

$$t = \frac{e^{a\xi}}{a} \sinh(a\eta), \quad x = \frac{e^{a\xi}}{a} \cosh(a\eta). \quad (39)$$

The proper acceleration is defined by  $\alpha^{-1} = ae^{-a\xi}$  and the proper time is associated with  $\eta$  and  $\xi$  via the relation  $\tau = e^{a\xi}\eta$ . We consider that our two-level systems are travelling along different hyperbolic trajectories  $x^\mu(\tau_1) = (t(\tau_1), x(\tau_1), y_1, z_1)$  and  $x^\mu(\tau_2) = (t(\tau_2), x(\tau_2), y_2, z_2)$ , whose functional dependence  $x_i(\tau_i), x_i(\tau_i)$  is given by equations similar to Eq. (39).

We now identify the time-evolution parameter of the generalized master equation derived in the previous section with the Rindler coordinate  $\eta$ . This allows us to perform a simple change of variables in the generalized master equation so that we obtain a description in terms of the proper time. However, recall that we have two different proper times at our disposal. Nevertheless, considering that the two different hyperbolas are in the same Rindler wedge, they are cut by a single line of constant  $\eta$ , which means that we have a simple relation between the proper times of the two-level systems. In what follows we find it convenient to choose the proper time  $\tau_1$  as the parameter that describes the time evolution of the system. Using standard methods, we can then relate the other proper time  $\tau_2$  to  $\tau_1$ . This relation is given by

$$\tau_2(\tau_1) = \tau_1 e^{a(\xi_2 - \xi_1)} = \frac{\alpha_2}{\alpha_1} \tau_1, \quad (40)$$

and, therefore, the  $a_\beta$  constant that we have used in Eq. (18) can be written as  $a_1 = 1$  and  $a_2 = \alpha_2/\alpha_1$ . As we are dealing with a scalar field, the  $i, j$ -summation is removed. In addition, the  $A$  operators are to be identified with the ladder operators  $S_\pm$ . We then write the generalized master equation as

$$\dot{\rho}_A(\eta) = -i [\mathcal{H}_{\text{eff}}, \rho_A(\eta)] + \mathcal{L} \{ \rho_A(\eta) \}, \quad (41)$$

where the dissipative contribution is given by

$$\begin{aligned} \mathcal{L} \{ \rho_A(\eta) \} = & \frac{1}{2} \sum_{\omega=-\omega_0}^{\omega_0} \sum_{\alpha, \beta=1}^2 \Gamma^{(\alpha\beta)}(a_\beta \omega) \\ & \left( 2S_-^{(\beta)} \rho_A(\eta) S_+^{(\alpha)} - \left\{ S_+^{(\alpha)} S_-^{(\beta)}, \rho_A(\eta) \right\} \right), \end{aligned} \quad (42)$$

with  $H_{\text{eff}}$  being given by

$$\mathcal{H}_{\text{eff}} = \mathcal{H}_A + \sum_{\omega} \sum_{\alpha, \beta} \Delta^{(\alpha\beta)}(a_\beta \omega) S_+^{(\alpha)} S_-^{(\beta)}. \quad (43)$$

The  $\Gamma(x)$  and  $\Delta(x)$  functions are given by Eqs. (28) and (29), removing the dependence on the  $i$  and  $j$  indices. We find that

$$W^{(\alpha\beta)}(a_\beta \omega) = a_\alpha a_\beta \int_0^\infty ds e^{i\omega a_\beta s} G^{+(\alpha\beta)}(s), \quad (44)$$

where  $G^{+(\alpha\beta)}(s)$  is the positive-frequency Wightman function whose form will be presented in the following. The reason that there is no switching function's products in Eq. (21) is that in



the derivation of the generalized master equation we already have discussed how the interaction is turned on and off between the two-level systems and the field. We have chosen the switching functions as simple theta functions which does not change the limits of integration.

In order to compute  $\Gamma^{(\alpha\beta)}(a_\beta\omega)$ , we must compute first  $G^{+(\alpha\beta)}(s)$  and  $W^{(\alpha\beta)}(a_\beta\omega)$ . The positive-frequency Wightman function for a massless real scalar field in Minkowski space-time is given by

$$G^+(x, x') = -\frac{1}{4\pi^2} \frac{1}{[(t - t' - i\epsilon)^2 - |\mathbf{x} - \mathbf{x}'|^2]}. \quad (45)$$

As we require that the atoms are in a uniformly accelerated motion coupled with a scalar field prepared in the Minkowski vacuum  $|0, M\rangle$ , the procedure is to change the inertial coordinates  $(x, x')$  in Eq. (45) to the Rindler coordinates using the expressions defined above. By doing so we have, for the usual term represented by  $\alpha = \beta$

$$G^{+(\beta\beta)}(s) = -\frac{1}{16\pi^2\alpha_\beta^2} \frac{1}{\sinh^2\left(\frac{1}{2\alpha_1}(s - 2i\epsilon)\right)}, \quad (46)$$

and for the crossed terms  $\alpha \neq \beta$  we have  $G^{+(12)}(s) = G^{+(21)}(s) = G_c^+(s)$ , where

$$G_c^+(s) = -\frac{1}{16\pi^2\alpha_1\alpha_2} \left[ \sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} + \frac{\phi}{2}\right) \times \sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} - \frac{\phi}{2}\right) \right]^{-1}, \quad (47)$$

with the following definition

$$\cosh \phi = 1 + \frac{(\alpha_1 - \alpha_2)^2 + |\Delta\mathbf{y}|^2}{2\alpha_1\alpha_2}, \quad (48)$$

where  $|\Delta\mathbf{x}|^2 = (y_2 - y_1)^2 + (z_1 - z_2)^2$  is the orthogonal (to the  $(t, x)$ -plane) distance between the two-level systems which we will assume to be constant.

Let us calculate the different contributions to the dynamics separately. Observe that

$$\begin{aligned} \Gamma^{(\alpha\beta)}(a_\beta\omega) &= W^{(\alpha\beta)}(a_\beta\omega) + W^{*(\beta\alpha)}(a_\beta\omega) \\ &= a_\alpha a_\beta \int_0^\infty ds e^{i\omega a_\beta s} G^{+(\alpha\beta)}(s) \\ &\quad + a_\beta a_\alpha \int_0^\infty ds e^{-i\omega a_\beta s} G^{+(\beta\alpha)}(s). \end{aligned} \quad (49)$$

By changing  $s \rightarrow -s$  in the second integral and observing that  $G^{+(\beta\alpha)}(-s) = G^{+(\alpha\beta)}(s)$  we have that

$$\Gamma^{(\alpha\beta)}(a_\beta\omega) = a_\alpha a_\beta \int_{-\infty}^\infty ds e^{i\omega a_\beta s} G^{+(\alpha\beta)}(s). \quad (50)$$

The explicit computation of Eq. (50) is given in the Appendix A. We find that

$$\begin{aligned} \Gamma^{(\beta\beta)}(a_\beta\omega) &= \frac{|\omega|\alpha_\beta}{2\pi\alpha_1} \left[ \theta(\omega) \left( 1 + \frac{1}{e^{2\pi\alpha_\beta\omega} - 1} \right) \right. \\ &\quad \left. + \theta(-\omega) \left( \frac{1}{e^{2\pi\alpha_\beta|\omega|} - 1} \right) \right], \end{aligned} \quad (51)$$

and for the crossed contribution

$$\begin{aligned}
\Gamma^{(\alpha\beta)}(a_\beta\omega) &= W^{(\alpha\beta)}(a_\beta\omega) + W^{*(\alpha\beta)}(a_\beta\omega) \\
&= \frac{1}{4\pi\alpha_1 \sinh \phi} \\
&\times \left\{ \theta(\omega) \left[ 2 \operatorname{Re}(\mathcal{U}(\alpha_\beta\omega)) \left( 1 + \frac{1}{e^{2\pi\alpha_\beta\omega} - 1} \right) \right] \right. \\
&\left. + \theta(-\omega) \left[ \frac{2 \operatorname{Re}(\mathcal{U}(\alpha_\beta|\omega|))}{e^{2\pi\alpha_\beta|\omega|} - 1} \right] \right\}. \tag{52}
\end{aligned}$$

It is useful to write Eq. (41) in terms of the Pauli matrices using that  $S_+^{(\alpha)} = S_1^{(\alpha)} + iS_2^{(\alpha)}$  and  $S_-^{(\alpha)} = S_1^{(\alpha)} - iS_2^{(\alpha)}$ , where  $S_i^{(1)} = \sigma^i \otimes \sigma^0$  and  $S_i^{(2)} = \sigma^0 \otimes \sigma^i$  with  $\sigma^i$  the  $i$ -th Pauli matrix. See, for example, Ref. [94]. The dissipative contribution can be written as (hereafter we omit the subscript  $A$  in the reduced density matrix  $\rho_A$ )

$$\begin{aligned}
\mathcal{L}\{\rho(\tau)\} &= \frac{1}{2} \sum_{i,j=1}^3 \sum_{\alpha,\beta=1}^2 \mathcal{C}_{ij}^{(\alpha\beta)} \left[ 2S_j^{(\beta)} \rho(\tau) S_i^{(\alpha)} \right. \\
&\quad \left. - \left\{ S_i^{(\alpha)} S_j^{(\beta)}, \rho(\tau) \right\} \right], \tag{53}
\end{aligned}$$

with the Kossakowski matrix  $\mathcal{C}_{ij}^{(\alpha\beta)}$  defined as

$$\mathcal{C}_{ij}^{(\alpha\beta)} = A^{(\alpha\beta)} \delta_{ij} - iB^{(\alpha\beta)} \epsilon_{ijk} \delta_{3k} - A^{(\alpha\beta)} \delta_{3i} \delta_{3j}. \tag{54}$$

We have also defined the following functions

$$A^{(\alpha\beta)} = \frac{\mu^2}{4} (\Gamma^{(\alpha\beta)}(a_\beta\omega) + \Gamma^{(\alpha\beta)}(-a_\beta\omega)), \tag{55}$$

$$B^{(\alpha\beta)} = \frac{\mu^2}{4} (\Gamma^{(\alpha\beta)}(a_\beta\omega) - \Gamma^{(\alpha\beta)}(-a_\beta\omega)). \tag{56}$$

Using Eqs. (51) and (52) we can compute explicitly the functions given by Eqs. (55) and (56). For  $\alpha = \beta$  we have

$$A^{(\beta)} = A^{(\beta\beta)} = \frac{\mu^2 \omega \alpha_\beta}{8\pi\alpha_1} \left( 1 + \frac{2}{e^{2\pi\alpha_\beta\omega} - 1} \right), \tag{57}$$

$$B^{(\beta)} = B^{(\beta\beta)} = \frac{\mu^2 \omega \alpha_\beta}{8\pi\alpha_1}, \tag{58}$$

and for  $\alpha \neq \beta$ ,

$$A^{(\alpha\beta)} = \frac{\mu^2 \sin(\omega\alpha_\beta\phi)}{8\pi\alpha_1 \sinh \phi} \left( 1 + \frac{2}{e^{2\pi\alpha_\beta\omega} - 1} \right), \tag{59}$$

$$B^{(\alpha\beta)} = \frac{\mu^2 \sin(\omega\alpha_\beta\phi)}{8\pi\alpha_1 \sinh \phi}. \tag{60}$$

## 4 Entanglement dynamics

To describe the entanglement dynamics we may ignore the unitary contribution given by Eq. (43). We write the dissipative part given by Eq. (53) in the basis  $\{|1\rangle = |g_1\rangle \otimes |g_2\rangle, |2\rangle = |e_1\rangle \otimes |e_2\rangle, |3\rangle = |g_1\rangle \otimes |e_2\rangle, |4\rangle = |e_1\rangle \otimes |g_2\rangle\}$ . We assume that the density matrix of the system has the block form

$$\rho(\tau) = \begin{pmatrix} \rho_{11} & \rho_{12} & 0 & 0 \\ \rho_{21} & \rho_{22} & 0 & 0 \\ 0 & 0 & \rho_{33} & \rho_{34} \\ 0 & 0 & \rho_{43} & \rho_{44} \end{pmatrix}. \quad (61)$$

The problem simplifies by working in the basis of the collective states of the system. For identical two-level systems, the collective states are

$$|G\rangle = |g_1\rangle \otimes |g_2\rangle, \quad (62)$$

$$|E\rangle = |e_1\rangle \otimes |e_2\rangle, \quad (63)$$

$$|S\rangle = \frac{1}{\sqrt{2}}(|g_1\rangle \otimes |e_2\rangle + |e_1\rangle \otimes |g_2\rangle), \quad (64)$$

$$|A\rangle = \frac{1}{\sqrt{2}}(|e_1\rangle \otimes |g_2\rangle - |g_1\rangle \otimes |e_2\rangle), \quad (65)$$

and we obtain that in this basis, the density matrix still has a block diagonal form as

$$\rho(\tau) = \begin{pmatrix} \rho_{GG} & \rho_{GE} & 0 & 0 \\ \rho_{EG} & \rho_{EE} & 0 & 0 \\ 0 & 0 & \rho_{SS} & \rho_{SA} \\ 0 & 0 & \rho_{AS} & \rho_{AA} \end{pmatrix}, \quad (66)$$

with  $\rho_{ij} = \langle i | \rho | j \rangle$ . We compute the matrix elements of the generalized master equation given by Eq. (41) in the aforementioned basis. We obtain eight coupled linear differential equations involving all the matrix components of Eq. (66). As we are dealing with a more general situation than the ones studied in Refs. [58, 86], our set of equations is more involved and are presented in the appendix B. We observe from Eqs. (92-99) that if we take  $\alpha_1 = \alpha_2$ , *i.e.*, set both two-level systems at the same world line, we then have  $A^{(11)} = A^{(22)}$ ,  $B^{(11)} = B^{(22)}$ ,  $A^{(12)} = A^{(21)}$  and  $B^{(12)} = B^{(21)}$ . This simplifies the set of equations and we recover the previous result of the literature [58, 86, 88]. Therefore, as expected, the case where the two-level systems are with the same proper acceleration is a special case of the master equation derived in this work.

The problem follows by treating the two two-level system as a single four-level system. Next, we must study how much of entanglement is stored in this quantum system. There are many alternatives to proceed this study. We here choose to compute the *concurrence* of this system. The concurrence  $C(\rho)$  is a function of the eigenvalues of the matrix  $R = \rho \tilde{\rho}$ , with  $\tilde{\rho}$  given by

$$\tilde{\rho} = \sigma_y \otimes \sigma_y \rho^T \sigma_y \otimes \sigma_y, \quad (67)$$

which has value  $C = 1$  for maximally entangled state and  $C = 0$  for separated states. In the basis given by Eqs. (62), (63), (64) and (65) we have the following form of the concurrence [40, 58]

$$C[\rho(\tau)] = \max \{0, \mathcal{C}_1(\tau), \mathcal{C}_2(\tau)\}, \quad (68)$$

with

$$\mathcal{C}_1(\tau) = \sqrt{[\rho_{AA}(\tau) - \rho_{SS}(\tau)]^2 - [\rho_{AS}(\tau) - \rho_{SA}(\tau)]^2 - 2\sqrt{\rho_{GG}(\tau)\rho_{EE}(\tau)}}, \quad (69)$$

$$\mathcal{C}_2(\tau) = -\sqrt{[\rho_{AA}(\tau) + \rho_{SS}(\tau)]^2 - [\rho_{AS}(\tau) + \rho_{SA}(\tau)]^2} + 2|\rho_{GE}(\tau)|. \quad (70)$$

In the following, we numerically solve the dissipative part of the generalized master equation for different initial configuration of the two-level systems and use this solution to evaluate the concurrence of Eq. (68) to study the entanglement dynamics of this system.

## 4.1 Entanglement degradation

Let us first consider the system to be initially in a maximally entangled state and compute its evolution. It is well known that this setup provides the scenario to study a phenomenon called *entanglement degradation*. The interest in studying this phenomenon is that the entanglement degradation for accelerated atoms will behave differently from inertial atoms due to the Unruh effect [95]. This phenomenon can be investigated by using as the initial state the symmetric state  $|S\rangle$  or the antisymmetric state  $|A\rangle$ . For the the system prepared in the symmetric state the results are depicted in Fig. 1, where we have fixed the value of acceleration of the second two-level system ( $\alpha_2^{-1}$ ) and varied the acceleration of the first two-level system ( $\alpha_1^{-1}$ ). In Fig. 2 we used the antisymmetric state as the initial state of the system where  $\alpha_2$  is fixed and we varied  $\alpha_1$ . We observe that the entanglement degradation occurs faster for the system prepared in the state  $|S\rangle$ . Moreover, notice the early-stage disentanglement developed by the two-level system – the profile seen in the degradation of the correlations of the two-level systems by environmental noise is compatible with the known effect of entanglement sudden death [96]. Such results are in agreement with the outcomes presented in Ref. [88].

The master equation derived in this work allows us to investigate how the entanglement occurs when the two-level systems are in distinct world lines, *i.e.*, for different values of  $\alpha_i$ . Here we consider  $\alpha_1 \neq \alpha_2$  but observe that for  $\alpha_1 = \alpha_2$  the master equation is the same as the one used in Ref. [86] and, therefore, we obtain the same results. By construction we have  $\alpha_2 \geq \alpha_1$  and we discuss how entanglement behaves when we vary both proper accelerations. In Fig. 3 we have for a fixed value of time parameter  $\tau$  the values that the concurrence  $C(\tau)$  assumes for different values of  $\alpha_1$  and  $\alpha_2$ . We observe that as both accelerations decrease (as  $\alpha_i$  increases) the lifetime of the entangled state increases.




Figure 1: Comparison of entanglement degradation for different values of  $\alpha_1$ . The initial state were prepared to be  $|S\rangle$ . We use  $\omega = 0.01$  and  $\alpha_2 = 10$ .




Figure 2: Comparison of entanglement degradation for different values of  $\alpha_1$ . The initial state were prepared to be  $|A\rangle$ . We use  $\omega = 0.01$  and  $\alpha_2 = 10$ .




Figure 3: Comparison of entanglement degradation for different values of  $\alpha_1$  and  $\alpha_2$ . The initial state were prepared to be  $|S\rangle$ . We use  $\omega = 0.01$ .

## 4.2 Entanglement Harvesting

Another interesting phenomenon is the so called *entanglement harvesting* [62]. This phenomenon occurs when a system is prepared in a separated state and, due to the interaction of the atoms with the field, an atomic entangled state is created. As discussed above, for a massless scalar field, the system is truly harvesting entanglement from the field only if the two-level systems are spacelike separated [69]. As we are dealing with two-level systems in different hyperbolas we can definitely assert that the entanglement that is being generated has to come from entanglement harvesting. To study this phenomena within our formalism we prepare the two-level system in the  $|G\rangle$  state or in the  $|E\rangle$  state. We observe in Fig. 4 that, for the initial state being  $|G\rangle$ , an entangled state can be created and the concurrence increases with the value of  $\alpha_1$ . One interesting result is that entanglement is only created by larger values of  $\alpha_2$ . If we employ the same configuration used in the discussion of the entanglement degradation, we cannot observe any entangled state being created, so  $C[\rho(\tau)] = 0$  in this configuration.




Figure 4: Comparison of entanglement harvesting for different values of  $\alpha_1$ . The initial state were prepared to be  $|G\rangle$ . We use  $\omega = 0.01$  and  $\alpha_2 = 70$ .

When setting the initial state to be  $|E\rangle$ , we obtain the results illustrated in Fig. 5. Again, we observe that the entangled state is only created by significantly decreasing the acceleration of the second two-level system  $\alpha_2^{-1}$ . Moreover, the phenomenon of entanglement harvesting only occurs for small values of this acceleration. In this situation, we observe that we only have entangled states for  $\alpha_2 \gg \alpha_1$  – we observe that this entangled state has short lifetime. Furthermore, when we increase the value of  $\alpha_1$ , the associated entanglement decreases. It is important to emphasize that although  $\alpha_2$  is the parameter that most assists the entanglement harvesting, this phenomenon only occurs for relatively high values for  $\alpha_1$  as well. This difficulty in creating entangled state is in agreement with the results discussed in Ref. [88].

As claimed above, our generalized master equation allows us to study how the entanglement of creation evolves when we scan through many different values for both proper accelerations. In Fig. 6 we consider a fixed time when the concurrence reaches its maximum value. To have a non-trivial result, we set values for  $\alpha_2$  much greater than  $\alpha_1$ . An interesting work that studies how the acceleration can assist entanglement harvesting is presented in Ref. [80]. The authors investigate three different acceleration scenarios using the Unruh-DeWitt particle-detector model – this is the main difference between the procedure that we have used here. This system configuration can now be revisited using the theory of quantum open systems and the master equation derived in this paper.




Figure 5: Comparison of entanglement harvesting for different values of  $\alpha_1$ . The initial state were prepared to be  $|E\rangle$ . We use  $\omega = 0.01$  and  $\alpha_2 = 145$ .




Figure 6: Comparison of entanglement harvesting for different values of  $\alpha_1$  and  $\alpha_2$ . The initial state were prepared to be  $|G\rangle$  and we fixed the time parameter to  $\tau = 150$ . We use  $\omega = 0.01$ .

## 5 Conclusions

Using the theory of open quantum system we were able to derive a generalized master equation for a system of two two-level system travelling along different world lines. We have also established a suitable generalized Lindblad-Kossakowski form and made use of the Kossakowski matrix formalism. We observe that this equation has a similar form of the one used for two-level systems in the same world lines. The crucial distinction here relies on the function given by Eq. (28), which allows us to investigate how differences in the proper accelerations of the two-level systems affect entanglement dynamics.

We studied the phenomenon of entanglement degradation and observed that as the proper acceleration of the two-level systems decreases, the lifetime of the entangled state increases, in agreement with the Unruh-Davies effect. In particular, the antisymmetric entangled state presents a slower decoherence in comparison with the symmetric entangled state. In any case, we were able to observe the occurrence of entanglement sudden death in all situations under studied. On the other hand, we also observed that the entanglement harvesting is also possible in this scenario and we have investigated its dependence on the values of the proper acceleration of both two-level systems. We have demonstrated that a quantum system initially prepared in the ground state supports entanglement for a longer time in comparison with one initially prepared in the excited separable state. Furthermore, entanglement creation is favoured when the system undergoes small values of accelerations.

The formalism developed in this work provides us with a new way to understand entanglement dynamics in non-inertial reference frames. The procedure used here can be used to study a more physical situation, for example a pair of dipoles interacting with the electromagnetic field, and even more complex situations, as with two-level systems in different space-time regions in a Kerr black-hole background. These subjects are under investigation by the authors.



## Acknowledgements

This work was partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq, under grants 303436/2015-8 (N.F.S.) and 317548/2021-2 (G.M.), and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro – FAPERJ under grants E-26/202.725/2018 and E-26/201.142/2022 (G.M.). We would like to thanks C. A. D. Zarro and F. de Melo for useful discussions about the mathematical and physical implications in whole the construction of this paper.

## A Computation of the $\Gamma^{(\alpha\beta)}$ function

In order to compute the function  $\Gamma^{(\alpha\beta)}$  from Eq. (49) we have two alternative ways to follow. One is to compute it by the Fourier transform presented in Eq. (50). This procedure will be straightforward when dealing with the  $\alpha = \beta$  contribution. Instead, for the crossed terms we find it more convenient to first compute the function  $W^{(\alpha\beta)}(a_\beta\omega)$  and then use Eq. (49).

Let us start with the  $\alpha = \beta$  contribution. For simplicity we write  $\Gamma^{(\alpha\beta)}(a_\beta\omega) = \Gamma^{(\beta)}(\omega)$  We use the positive wightman function given by (46) into Eq. (50) and then

$$\Gamma^{(\beta)}(\omega) = -\frac{a_\beta^2}{16\pi^2\alpha_\beta^2} \int_{-\infty}^{\infty} ds \frac{e^{i\omega a_\beta s}}{\sinh^2\left(\frac{1}{2\alpha_1}(s - 2i\epsilon)\right)}. \quad (71)$$

To solve this integral we use the following identity [1]

$$\csc^2(\pi x) = \frac{1}{\pi^2} \sum_{k=-\infty}^{\infty} \frac{1}{(x - k)^2} \quad \text{and} \quad (72)$$

$$\operatorname{csch} x = i \csc(ix). \quad (73)$$

Therefore Eq. (71) can be written as

$$\Gamma^{(\beta)}(\omega) = -\frac{a_\beta^2\alpha_1^2}{4\pi^2\alpha_\beta^2} \sum_{k=-\infty}^{\infty} \int_{-\infty}^{\infty} ds \frac{e^{i\omega a_\beta s}}{(s - 2i\epsilon + 2\pi i k \alpha_1)^2}. \quad (74)$$

By performing a change of variables  $u = a_\beta s$  in Eq. (74) we find that

$$\Gamma^{(\beta)}(\omega) = -\frac{a_\beta^3\alpha_1^2}{4\pi^2\alpha_\beta^2} \sum_{k=-\infty}^{\infty} \int_{-\infty}^{\infty} du \frac{e^{i\omega u}}{(u - 2ia_\beta\epsilon + 2\pi i k \alpha_1 a_\beta)^2}. \quad (75)$$

The integral presented in Eq. (75) can be solved with the help of a contour integration methods. In this integral we observe the existence of second order poles given by  $u = 2ia_\beta\epsilon - 2\pi i k \alpha_1 a_\beta$ , where  $k$  is an integer. For different values of  $k$  ( $k \geq 0$  and  $k < 0$ ) we have poles over and above the complex plane. For  $\omega > 0$  we use a semicircle of radius  $R$  that we close on the upper-half ( $\operatorname{Im}[u] \geq 0$ ) plane. This contour encloses the poles for  $k \leq 0$  and runs in anticlockwise direction. For  $\omega < 0$  we close the contour with the lower-half semicircle ( $\operatorname{Im}[u] \leq 0$  of radius  $R$ . In this contour we have  $k > 0$  and it runs in clockwise direction. We consider the limit  $R \rightarrow \infty$  such that the contribution from the arcs will vanish by the Jordan's lemma. From this discussion we can have

$$\Gamma^{(\beta)}(\omega) = -\frac{a_\beta^3 \alpha_1^2}{4\pi^2 \alpha_\beta^2} \left\{ 2\pi i \theta(\omega) \sum_{k=0}^{\infty} \text{Res} \left[ \frac{e^{i\omega u}}{(u - 2ia_\beta \epsilon - 2\pi i k \alpha_1 a_\beta)^2}; u = 2ia_\beta \epsilon + 2\pi i k \alpha_1 a_\beta \right] \right. \\ \left. - 2\pi i \theta(-\omega) \sum_{k=1}^{\infty} \text{Res} \left[ \frac{e^{-i|\omega|u}}{(u + 2\pi i k \alpha_1 a_\beta)^2}; u = -2\pi i k \alpha_1 a_\beta \right] \right\}, \quad (76)$$

where in the first summation we performed the change  $k \rightarrow -k$ . By finding the residues of Eq. (76) and considering that  $a_\beta = \alpha_\beta/\alpha_1$ , we finally obtain

$$\Gamma^{(\beta)}(\omega) = \frac{|\omega| \alpha_\beta}{2\pi \alpha_1} \left[ \theta(\omega) \left( \sum_{k=0}^{\infty} e^{-2\pi \omega \alpha_1 a_\beta k} \right) + \theta(-\omega) \left( \sum_{k=1}^{\infty} e^{-2\pi |\omega| \alpha_1 a_\beta k} \right) \right] \\ = \frac{|\omega| \alpha_\beta}{2\pi \alpha_1} \left[ \theta(\omega) \left( 1 + \frac{1}{e^{2\pi \omega \alpha_1 a_\beta} - 1} \right) + \theta(-\omega) \left( \frac{1}{e^{2\pi |\omega| \alpha_1 a_\beta} - 1} \right) \right]. \quad (77)$$

As discussed above, to compute  $\Gamma^{(\alpha\beta)}(a_\beta \omega)$  for  $\alpha \neq \beta$  we first compute the functions  $W^{(\alpha\beta)}(a_\beta \omega)$ . Using Eq. (47) into Eq. (44) we have

$$W^{(\alpha\beta)}(a_\beta \omega) = -\frac{a_\alpha a_\beta}{16\pi^2 \alpha_1 \alpha_2} \int_0^\infty ds \frac{e^{i\omega a_\beta s}}{\sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} + \frac{\phi}{2}\right) \sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} - \frac{\phi}{2}\right)}. \quad (78)$$

Let us start with  $W^{(21)}(\omega)$ . By Eq. (78) we have

$$W^{(21)}(\omega) = -\frac{1}{16\pi^2 \alpha_1^2} \int_0^\infty ds \frac{e^{i\omega s}}{\sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} + \frac{\phi}{2}\right) \sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1 + \alpha_2} - \frac{\phi}{2}\right)} \\ = -\frac{1}{16\pi^2 \alpha_1^2} \int_0^\infty ds f(s). \quad (79)$$




Figure 7: Contours in the complex plane for the evaluation of the integrals for  $W^{(\alpha\beta)}(a_\beta \omega)$ . In (a) we have the contour for  $\omega > 0$ . We observe that in this case the poles inside the contour for  $R \rightarrow \infty$  have  $n \geq 0$ . Figure (b) corresponds to  $\omega < 0$  and we have  $n \geq 1$ .

To solve this integral we use the contours illustrated in Fig. (7). The sign of  $\omega$  will define if we close the contour in the upper-half of the semicircle of radius  $R$  or in its lower-half. For  $\omega > 0$ , the poles inside the upper-half semicircle contour, see Fig. (7), are given by

$$s_n^{(1)} = \alpha_1 \phi + \frac{8\alpha_1 i \epsilon}{\alpha_1 + \alpha_2} + 2\alpha_1 \pi i n, \quad (80)$$

with  $n \geq 0$ . The associated contour integral can be split up to

$$\begin{aligned} \oint_{\omega > 0} ds f(s) &= \int_0^R ds f(s) + \int_{C_R} ds f(s) + \int_{iR}^0 ds f(s) \\ &= 2\pi i \sum_{n=0}^{\infty} \text{Res}(f(s); s = s_n^{(1)}) \\ &= \frac{4\pi i \alpha_1}{\sinh \phi} e^{i\omega \alpha_1 \phi} \sum_{n=0}^{\infty} e^{-2\pi \omega \alpha_1 n} \end{aligned} \quad (81)$$

with  $C_R$  the quarter of circle presented in Fig. (7) and we have taken  $\epsilon \rightarrow 0$ . For  $\omega < 0$ , the poles inside the lower-half semicircle contour, see Fig. (7), are given by

$$s_n^{(2)} = \alpha_1 \phi - 2\alpha_1 \pi i n, \quad (82)$$

with  $n > 0$ . The associated contour integral can be split up to

$$\begin{aligned} \oint_{\omega < 0} ds f(s) &= \int_0^R ds f(s) + \int_{C'_R} ds f(s) + \int_{-iR}^0 ds f(s) \\ &= -2\pi i \sum_{n=1}^{\infty} \text{Res}(f(s); s = s_n^{(2)}) \\ &= -\frac{4\pi i \alpha_1}{\sinh \phi} e^{-i|\omega| \alpha_1 \phi} \sum_{n=1}^{\infty} e^{-2\pi \alpha_1 |\omega| n}, \end{aligned} \quad (83)$$

with  $C'_R$  the quarter of circle presented in Fig. (7). By taking the limit  $R \rightarrow \infty$  the integrals along  $C_R$  and  $C'_R$  vanishes by the Jordan lemma. After some algebraic manipulations, one gets

$$\begin{aligned} W^{(21)}(\omega) &= \frac{\text{csch} \phi}{4\pi \alpha_1} \left\{ \theta(\omega) \left[ \mathcal{U}(\alpha_1 \omega) \left( 1 + \frac{1}{e^{2\pi \alpha_1 \omega} - 1} \right) + \mathcal{V}(\alpha_1 \omega) \right] \right. \\ &\quad \left. + \theta(-\omega) \left( \frac{\mathcal{U}^*(\alpha_1 |\omega|)}{e^{2\pi \alpha_1 |\omega|} - 1} + \mathcal{V}^*(\alpha_1 |\omega|) \right) \right\}, \end{aligned} \quad (84)$$

where we have defined the following functions

$$\mathcal{U}(\alpha_1 \omega) = -i e^{i\omega \alpha_1 \phi}, \quad (85)$$

$$\mathcal{V}(\alpha_1 \omega) = \frac{i \sinh \phi}{2} \int_0^{2\pi} du \frac{e^{u \alpha_1 \omega}}{\cos u - \cosh \phi}. \quad (86)$$

Now we look for the solution for  $W^{(12)}(a_2\omega)$ . From Eq. (78) we have

$$W^{(12)}(a_2\omega) = W^{(12)}\left(\frac{\alpha_2\omega}{\alpha_1}\right) = -\frac{1}{16\pi^2\alpha_1^2} \int_0^\infty ds \frac{e^{i\omega\frac{\alpha_2}{\alpha_1}s}}{\sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1+\alpha_2} + \frac{\phi}{2}\right) \sinh\left(\frac{s}{2\alpha_1} - \frac{4i\epsilon}{\alpha_1+\alpha_2} - \frac{\phi}{2}\right)}. \quad (87)$$

We observe that the integral presented in Eq. (87) can be solved similarly to the one from Eq. (79) by changing  $w \rightarrow w\alpha_2/\alpha_1$ . Therefore, we can write

$$W^{(12)}(a_2\omega) = \frac{\text{csch}\phi}{4\pi\alpha_1} \left\{ \theta(\omega) \left[ \mathcal{U}(\alpha_2\omega) \left(1 + \frac{1}{e^{2\pi\alpha_2\omega} - 1}\right) + \mathcal{V}(\alpha_2\omega) \right] + \theta(-\omega) \left( \frac{\mathcal{U}^*(\alpha_2|\omega|)}{e^{2\pi\alpha_2|\omega|} - 1} + \mathcal{V}^*(\alpha_2|\omega|) \right) \right\}, \quad (88)$$

or in a general form

$$W^{(\alpha\beta)}(a_\beta\omega) = \frac{\text{csch}\phi}{4\pi\alpha_1} \left\{ \theta(\omega) \left[ \mathcal{U}(\alpha_\beta\omega) \left(1 + \frac{1}{e^{2\pi\alpha_\beta\omega} - 1}\right) + \mathcal{V}(\alpha_\beta\omega) \right] + \theta(-\omega) \left( \frac{\mathcal{U}^*(\alpha_\beta|\omega|)}{e^{2\pi\alpha_\beta|\omega|} - 1} + \mathcal{V}^*(\alpha_\beta|\omega|) \right) \right\}. \quad (89)$$

With Eq. (89) in hands we are able to compute  $\Gamma^{(\alpha\beta)}(a_\beta\omega)$  from Eq. (49). We first observe that  $\phi$  is invariant under the change  $\alpha_1 \rightarrow \alpha_2$ . This lead us to write

$$W^{(\alpha\beta)}(a_\beta\omega) = W^{(\beta\alpha)}(a_\beta\omega). \quad (90)$$

Therefore, by using Eq. (89) into Eq. (49) we finally have

$$\Gamma^{(\alpha\beta)}(a_\beta\omega) = \frac{1}{4\pi\alpha_1 \sinh \phi} \left\{ \theta(\omega) \left[ 2 \text{Re}(\mathcal{U}(\alpha_\beta\omega)) \left(1 + \frac{1}{e^{2\pi\alpha_\beta\omega} - 1}\right) \right] + \theta(-\omega) \left[ \frac{2 \text{Re}(\mathcal{U}(\alpha_\beta|\omega|))}{e^{2\pi\alpha_\beta|\omega|} - 1} \right] \right\}. \quad (91)$$

## B Explicit form of the generalized master equation

In this appendix we present the explicit form of the master equation (41) where we consider the contribution only of the dissipative part of it as discussed in Sec. 4. From Eq. (66) we only have to compute eight components. These are given by the following set of differential

equations:

$$\begin{aligned}
\dot{\rho}_{GG}(\tau) = & -2(A^{(11)} + A^{(22)} - B^{(11)} - B^{(22)})\rho_{GG}(\tau) \\
& + (A^{(11)} + A^{(22)} + B^{(11)} + B^{(22)} \\
& - A^{(12)} - A^{(21)} - B^{(12)} - B^{(21)})\rho_{AA}(\tau) \\
& + (A^{(11)} - A^{(22)} + B^{(11)} - B^{(22)} \\
& - A^{(12)} + A^{(21)} + B^{(12)} - B^{(21)})\rho_{AS}(\tau) \\
& + (A^{(11)} - A^{(22)} + B^{(11)} - B^{(22)} \\
& + A^{(12)} - A^{(21)} + B^{(12)} - B^{(21)})\rho_{SA}(\tau) \\
& + (A^{(11)} + A^{(22)} + B^{(11)} + B^{(22)} \\
& + A^{(12)} + A^{(21)} + B^{(12)} + B^{(21)})\rho_{SS}(\tau),
\end{aligned} \tag{92}$$

$$\begin{aligned}
\dot{\rho}_{SS}(\tau) = & -2(A^{(11)} + A^{(22)} + A^{(12)} + A^{(21)})\rho_{SS}(\tau) \\
& + (B^{(22)} - B^{(11)} + B^{(12)} - B^{(21)})\rho_{AS}(\tau) \\
& + (A^{(11)} + A^{(22)} + B^{(11)} + B^{(22)} \\
& + A^{(12)} + A^{(21)} + B^{(12)} + B^{(21)})\rho_{EE}(\tau) \\
& + (A^{(11)} + A^{(22)} - B^{(11)} - B^{(22)} \\
& + A^{(12)} + A^{(21)} - B^{(12)} - B^{(21)})\rho_{GG}(\tau) \\
& + (B^{(22)} - B^{(11)} - B^{(12)} + B^{(21)})\rho_{SA}(\tau),
\end{aligned} \tag{93}$$

$$\begin{aligned}
\dot{\rho}_{AA}(\tau) = & -2(A^{(11)} + A^{(22)} - A^{(12)} - A^{(21)})\rho_{AA}(\tau) \\
& + (B^{(22)} - B^{(11)} + B^{(12)} - B^{(21)})\rho_{AS}(\tau) \\
& + (A^{(11)} + A^{(22)} + B^{(11)} + B^{(22)} \\
& + A^{(12)} + A^{(21)} + B^{(12)} + B^{(21)})\rho_{EE}(\tau) \\
& + (A^{(11)} + A^{(22)} - B^{(11)} - B^{(22)} \\
& + A^{(12)} + A^{(21)} - B^{(12)} - B^{(21)})\rho_{GG}(\tau) \\
& + (B^{(22)} - B^{(11)} - B^{(12)} + B^{(21)})\rho_{SA}(\tau),
\end{aligned} \tag{94}$$

$$\begin{aligned}
\dot{\rho}_{EE}(\tau) = & -2(A^{(11)} + A^{(22)} + B^{(11)} + B^{(22)})\rho_{EE}(\tau) \\
& + (A^{(22)} - A^{(11)} + B^{(11)} - B^{(22)} \\
& - A^{(12)} + A^{(21)} + B^{(12)} - B^{(21)})\rho_{SA}(\tau) \\
& + (A^{(11)} + A^{(22)} - B^{(11)} - B^{(22)} \\
& - A^{(12)} - A^{(21)} + B^{(12)} + B^{(21)})\rho_{AA}(\tau) \\
& + (A^{(22)} - A^{(11)} + B^{(11)} - B^{(22)} \\
& + A^{(12)} - A^{(21)} - B^{(12)} + B^{(21)})\rho_{AS}(\tau) \\
& + (A^{(22)} + A^{(11)} - B^{(11)} - B^{(22)} \\
& + A^{(12)} + A^{(21)} - B^{(12)} - B^{(21)})\rho_{SS}(\tau),
\end{aligned} \tag{95}$$

$$\begin{aligned}
\dot{\rho}_{AS}(\tau) = & -2(A^{(11)} + A^{(22)})\rho_{AS}(\tau) \\
& + (B^{(22)} - B^{(11)} - B^{(12)} + B^{(21)})\rho_{AA}(\tau) \\
& + (A^{(22)} - A^{(11)} - B^{(11)} + B^{(22)} \\
& + A^{(12)} - A^{(21)} + B^{(12)} - B^{(21)})\rho_{EE}(\tau) \\
& + (A^{(11)} - A^{(22)} - B^{(11)} + B^{(22)} \\
& - A^{(12)} + A^{(21)} + B^{(12)} - B^{(21)})\rho_{GG}(\tau) \\
& + (B^{(22)} - B^{(11)} - B^{(12)} + B^{(21)})\rho_{SS}(\tau),
\end{aligned} \tag{96}$$

$$\begin{aligned}
\dot{\rho}_{SA}(\tau) = & -2(A^{(11)} + A^{(22)})\rho_{SA}(\tau) \\
& + (B^{(22)} - B^{(11)} + B^{(12)} - B^{(21)})\rho_{AA}(\tau) \\
& + (A^{(22)} - A^{(11)} - B^{(11)} + B^{(22)} \\
& - A^{(12)} + A^{(21)} - B^{(12)} + B^{(21)})\rho_{EE}(\tau) \\
& + (A^{(11)} - A^{(22)} - B^{(11)} + B^{(22)} \\
& + A^{(12)} - A^{(21)} - B^{(12)} + B^{(21)})\rho_{GG}(\tau) \\
& + (B^{(22)} - B^{(11)} + B^{(12)} - B^{(21)})\rho_{SS}(\tau),
\end{aligned} \tag{97}$$

$$\dot{\rho}_{GE}(\tau) = -2(A^{(11)} + A^{(22)})\rho_{GE}(\tau), \tag{98}$$

$$\dot{\rho}_{EG}(\tau) = -2(A^{(11)} + A^{(22)})\rho_{EG}(\tau). \tag{99}$$

## References

- [1] N. D. Birell and P. C. W. Davies. *Quantum Fields in Curved Space*. Cambridge University Press, Cambridge, 1982.
- [2] Leonard E. Parker and D. Toms. *Quantum Field Theory in Curved Spacetime*. Cambridge University Press, 8 2009.
- [3] P. C. W. Davies. Scalar particle production in Schwarzschild and Rindler metrics. *J. Phys. A*, 8:609–616, 1975.
- [4] W. G. Unruh. Notes on black hole evaporation. *Phys. Rev. D*, 14:870, 1976.
- [5] William G Unruh and Robert M Wald. What happens when an accelerating observer detects a Rindler particle. *Phys. Rev. D*, 29(6):1047, 1984.
- [6] S. W. Hawking. Particle creation by black holes. *Commun. Math. Phys.*, 43:199–220, Aug 1975.
- [7] Hongwei Yu and Jialin Zhang. Understanding hawking radiation in the framework of open quantum systems. *Phys. Rev. D*, 77:024031, Jan 2008.
- [8] Stephen A Fulling. Nonuniqueness of canonical field quantization in Riemannian spacetime. *Phys. Rev. D*, 7(10):2850, 1973.

- [9] P. Candelas and D. J. Raine. Quantum Field Theory in Incomplete Manifolds. *J. Math. Phys.*, 17:2101–2112, 1976.
- [10] Bernar F Svaiter and Nami F Svaiter. Inertial and noninertial particle detectors and vacuum fluctuations. *Phys. Rev. D*, 46(12):5267, 1992. [Erratum: *Phys. Rev. D* **47**, 4802(E) (1993)].
- [11] L. H. Ford, N. F. Svaiter, and Marcelo L. Lyra. Radiative properties of a two-level system in the presence of mirrors. *Phys. Rev. A*, 49:1378–1386, 1994.
- [12] Shih-Yuin Lin and B. L. Hu. Accelerated Detector - Quantum Field Correlations: From Vacuum Fluctuations to Radiation Flux. *Phys. Rev. D*, 73:124018, 2006.
- [13] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas. The Unruh effect and its applications. *Rev. Mod. Phys.*, 80:787, 2008.
- [14] F. Lenz, K. Ohta, and K. Yazaki. Static interactions and stability of matter in Rindler space. *Phys. Rev. D*, 83:064037, 2011.
- [15] E. Arias, G. Krein, G. Menezes, and N. F. Svaiter. Thermal Radiation from a Fluctuating Event Horizon. *Int. J. Mod. Phys. A*, 27:1250129, 2012.
- [16] G. Menezes and N. F. Svaiter. Radiative processes of uniformly accelerated entangled atoms. *Phys. Rev. A*, 93(5):052117, 2016.
- [17] J. S. Ben-Benjamin et al. Unruh Acceleration Radiation Revisited. *Int. J. Mod. Phys. A*, 34(28):1941005, 2019.
- [18] C. D. Rodríguez-Camargo, N. F. Svaiter, and G. Menezes. Finite-time response function of uniformly accelerated entangled atoms. *Annals Phys.*, 396:266–291, 2018.
- [19] E. Arias, J. G. Dueñas, G. Menezes, and N. F. Svaiter. Boundary effects on radiative processes of two entangled atoms. *JHEP*, 07:147, 2016.
- [20] Christopher J. Fewster, Benito A. Juárez-Aubry, and Jorma Louko. Waiting for Unruh. *Class. Quant. Grav.*, 33(16):165003, 2016.
- [21] G. Menezes. Spontaneous excitation of an atom in a Kerr spacetime. *Phys. Rev. D*, 95(6):065015, 2017. [Erratum: *Phys.Rev.D* 97, 029901(E) (2018)].
- [22] G. Menezes, N. F. Svaiter, and C. A. D. Zarro. Entanglement dynamics in random media. *Phys. Rev. A*, 96(6):062119, 2017.
- [23] Gabriel Picanço, Nami F. Svaiter, and Carlos A. D. Zarro. Radiative processes of entangled detectors in rotating frames. *JHEP*, 08:025, 2020.
- [24] Wenting Zhou and Hongwei Yu. Radiation-reaction-induced transitions of two maximally entangled atoms in noninertial motion. *Phys. Rev. D*, 101:025009, Jan 2020.
- [25] Erickson Tjoa and Robert B. Mann. Unruh-DeWitt detector in dimensionally-reduced static spherically symmetric spacetimes. *JHEP*, 03:014, 2022.

- [26] Hui Hu, Jialin Zhang, and Hongwei Yu. Harvesting Entanglement by non-identical detectors with different energy gaps. 4 2022.
- [27] Bryce S DeWitt. Quantum Field Theory in Curved Spacetime. *Phys. Rep.*, 19(6):295–357, 1975.
- [28] S. W. Hawking and W. Israel. *General Relativity: An Einstein Centenary Survey*. Univ. Pr., Cambridge, UK, 1979.
- [29] Roy J Glauber. The quantum theory of optical coherence. *Phys. Rev.*, 130(6):2529, 1963.
- [30] M. S. Soares, N. F. Svaiter, C. A. D. Zarro, and G. Menezes. Uniformly accelerated quantum counting detector in minkowski and fulling vacuum states. *Phys. Rev. A*, 103:042225, Apr 2021.
- [31] Eduardo Martin-Martinez and Jorma Louko. Particle detectors and the zero mode of a quantum field. *Phys. Rev. D*, 90(2):024015, 2014.
- [32] Eduardo Martin-Martinez. Causality issues of particle detector models in QFT and Quantum Optics. *Phys. Rev. D*, 92(10):104019, 2015.
- [33] Daniel Hümmer, Eduardo Martin-Martinez, and Achim Kempf. Renormalized Unruh-DeWitt Particle Detector Models for Boson and Fermion Fields. *Phys. Rev. D*, 93(2):024019, 2016.
- [34] Erickson Tjoa, Irene Lopez-Gutiérrez, Allison Sachs, and Eduardo Martín-Martínez. What makes a particle detector click. *Phys. Rev. D*, 103(12):125021, 2021.
- [35] José de Ramón, Maria Papageorgiou, and Eduardo Martín-Martínez. Relativistic causality in particle detector models: Faster-than-light signaling and impossible measurements. *Phys. Rev. D*, 103(8):085002, 2021.
- [36] Michael A Nielsen and Isaac Chuang. *Quantum computation and quantum information*. Cambridge University Press, 2010.
- [37] M. B. Plenio, S. F. Huelga, A. Beige, and P. L. Knight. Cavity loss induced generation of entangled atoms. *Phys. Rev. A*, 59:2468–2475, 1999.
- [38] A. M. Basharov. Entanglement of atomic states upon collective radiative decay. *JETP Letters*, 75(3):123–126, 2002.
- [39] Z. Ficek and R. Tanas. Entanglement induced by spontaneous emission in spatially extended two-atom systems. *J. Mod. Opt.*, 50(18):2765–2779, 2003.
- [40] R Tanas and Z Ficek. Entangling two atoms via spontaneous emission. *J. opt., B Quantum semiclass. opt.*, 6(3):S90–S97, mar 2004.
- [41] Luigi Amico, Rosario Fazio, Andreas Osterloh, and Vlatko Vedral. Entanglement in many-body systems. *Rev. Mod. Phys.*, 80:517–576, 2008.
- [42] Asher Peres, Petra F. Scudo, and Daniel R. Terno. Quantum entropy and special relativity. *Phys. Rev. Lett.*, 88:230402, 2002.



- [43] Robert M. Gingrich and Christoph Adami. Quantum Entanglement of Moving Bodies. *Phys. Rev. Lett.*, 89(27):270402, 2002.
- [44] Robert M. Gingrich, Attila J. Bergou, and Christoph Adami. Entangled light in moving frames. *Phys. Rev. A*, 68:042102, 2003.
- [45] Yu Shi. Entanglement in relativistic quantum field theory. *Phys. Rev. D*, 70:105001, 2004.
- [46] Marek Czachor. Comment on “Quantum Entropy and Special Relativity”. *Phys. Rev. Lett.*, 94:078901, 2005.
- [47] Thomas F. Jordan, Anil Shaji, and E. C. G. Sudarshan. Lorentz transformations that entangle spins and entangle momenta. *Phys. Rev. A*, 75:022101, 2007.
- [48] Gerardo Adesso, Ivette Fuentes-Schuller, and Marie Ericsson. Continuous variable entanglement sharing in non-inertial frames. *Phys. Rev. A*, 76:062112, 2007.
- [49] Animesh Datta. Quantum discord between relatively accelerated observers. *Phys. Rev. A*, 80:052304, 2009.
- [50] Asher Peres and Daniel R. Terno. Quantum information and relativity theory. *Rev. Mod. Phys.*, 76:93–123, 2004.
- [51] Shih-Yuin Lin and B. L. Hu. Entanglement creation between two causally disconnected objects. *Phys. Rev. D*, 81:045019, 2010.
- [52] David E. Bruschi, Jorma Louko, Eduardo Martin-Martinez, Andrzej Dragan, and Ivette Fuentes. The Unruh effect in quantum information beyond the single-mode approximation. *Phys. Rev. A*, 82:042332, 2010.
- [53] David C. M. Ostapchuk, Shih-Yuin Lin, Robert B. Mann, and B. L. Hu. Entanglement Dynamics between Inertial and Non-uniformly Accelerated Detectors. *JHEP*, 07:072, 2012.
- [54] B. L. Hu, Shih-Yuin Lin, and Jorma Louko. Relativistic Quantum Information in Detectors-Field Interactions. *Class. Quant. Grav.*, 29:224005, 2012.
- [55] Relativistic quantum information, special issue of Classical and Quantum Gravity. *Class. Quantum Grav.*, 29:(11), 2012.
- [56] Shih-Yuin Lin, Chung-Hsien Chou, and Bei-Lok Hu. Entanglement Dynamics of Detectors in an Einstein Cylinder. *JHEP*, 03:047, 2016.
- [57] Eduardo Martin-Martinez, Luis J. Garay, and Juan Leon. Unveiling quantum entanglement degradation near a Schwarzschild black hole. *Phys. Rev. D*, 82:064006, 2010.
- [58] Yiquan Yang, Jiawei Hu, and Hongwei Yu. Entanglement dynamics for uniformly accelerated two-level atoms coupled with electromagnetic vacuum fluctuations. *Phys. Rev. A*, 94:032337, Sep 2016.
- [59] Jiaozhen She, Jiawei Hu, and Hongwei Yu. Entanglement dynamics for circularly accelerated two-level atoms coupled with electromagnetic vacuum fluctuations. *Phys. Rev. D*, 99(10):105009, 2019.

- [60] Pingyang He, Hongwei Yu, and Jiawei Hu. Entanglement dynamics for static two-level atoms in cosmic string spacetime. *Eur. Phys. J. C*, 80(2):134, 2020.
- [61] Sourav Bhattacharya and Nitin Joshi. Entanglement degradation in multi-event horizon spacetimes. *Phys. Rev. D*, 105(6):065007, 2022.
- [62] Alejandro Pozas-Kerstjens and Eduardo Martín-Martínez. Harvesting correlations from the quantum vacuum. *Phys. Rev. D*, 92:064042, Sep 2015.
- [63] Allison Sachs, Robert B. Mann, and Eduardo Martín-Martínez. Entanglement harvesting and divergences in quadratic Unruh-DeWitt detector pairs. *Phys. Rev. D*, 96(8):085012, 2017.
- [64] Alejandro Pozas-Kerstjens, Jorma Louko, and Eduardo Martín-Martínez. Degenerate detectors are unable to harvest spacelike entanglement. *Phys. Rev. D*, 95(10):105009, 2017.
- [65] Alejandro Pozas-Kerstjens and Eduardo Martín-Martínez. Entanglement harvesting from the electromagnetic vacuum with hydrogenlike atoms. *Phys. Rev. D*, 94:064074, Sep 2016.
- [66] Zhihong Liu, Jialin Zhang, and Hongwei Yu. Entanglement harvesting in the presence of a reflecting boundary. *JHEP*, 08:020, 2021.
- [67] Erickson Tjoa and Eduardo Martín-Martínez. Vacuum entanglement harvesting with a zero mode. *Phys. Rev. D*, 101(12):125020, 2020.
- [68] T. Rick Perche, Caroline Lima, and Eduardo Martín-Martínez. Harvesting entanglement from complex scalar and fermionic fields with linearly coupled particle detectors. *Phys. Rev. D*, 105(6):065016, 2022.
- [69] Erickson Tjoa and Eduardo Martín-Martínez. When entanglement harvesting is not really harvesting. *Phys. Rev. D*, 104:125005, Dec 2021.
- [70] Ivette Fuentes-Schuller and Robert B. Mann. Alice falls into a black hole: Entanglement in non-inertial frames. *Phys. Rev. Lett.*, 95:120404, 2005.
- [71] RB Mann and I Fuentes-Schuller. Quantum entanglement in noninertial frames. *Physics Essays*, 21(1):1, 2008.
- [72] Paul M Alsing and Ivette Fuentes. Observer-dependent entanglement. *Class. Quantum Grav.*, 29(22):224001, oct 2012.
- [73] Grant Salton, Robert B Mann, and Nicolas C Menicucci. Acceleration-assisted entanglement harvesting and rangefinding. *New J. Phys.*, 17(3):035001, mar 2015.
- [74] Kensuke Gallock-Yoshimura, Erickson Tjoa, and Robert B. Mann. Harvesting entanglement with detectors freely falling into a black hole. *Phys. Rev. D*, 104:025001, Jul 2021.
- [75] Alejandro Pozas-Kerstjens and Eduardo Martín-Martínez. Harvesting correlations from the quantum vacuum. *Phys. Rev. D*, 92:064042, Sep 2015.
- [76] Erickson Tjoa and Eduardo Martín-Martínez. When entanglement harvesting is not really harvesting. *Phys. Rev. D*, 104:125005, Dec 2021.

- [77] Matheus H. Zambianco, André G. S. Landulfo, and George E. A. Matsas. Observer dependence of entanglement in nonrelativistic quantum mechanics. *Phys. Rev. A*, 100:062126, Dec 2019.
- [78] Jialin Zhang and Hongwei Yu. Entanglement harvesting for unruh-dewitt detectors in circular motion. *Phys. Rev. D*, 102:065013, Sep 2020.
- [79] Wan Cong, Chen Qian, Michael RR Good, and Robert B Mann. Effects of horizons on entanglement harvesting. *JHEP*, 2020(10):1–19, 2020.
- [80] Zhihong Liu, Jialin Zhang, Robert B Mann, and Hongwei Yu. Does acceleration assist entanglement harvesting? *arXiv preprint arXiv:2111.04392*, 2021.
- [81] G. Menezes and N. F. Svaiter. Vacuum fluctuations and radiation reaction in radiative processes of entangled states. *Phys. Rev. A*, 92(6):062131, 2015.
- [82] G. Menezes. Radiative processes of two entangled atoms outside a Schwarzschild black hole. *Phys. Rev. D*, 94(10):105008, 2016.
- [83] Heinz-Peter Breuer, Francesco Petruccione, et al. *The theory of open quantum systems*. Oxford University Press on Demand, 2002.
- [84] Stanislaw Kryszewski and Justyna Czechowska-Kryszk. Master equation-tutorial approach. *arXiv preprint arXiv:0801.1757*, 2008.
- [85] Claude Cohen-Tannoudji, Bernard Diu, and Frank Laloe. Quantum mechanics, volume 2. *Quantum Mechanics*, 2:626, 1986.
- [86] F. Benatti and R. Floreanini. Entanglement generation in uniformly accelerating atoms: Reexamination of the unruh effect. *Phys. Rev. A*, 70:012112, Jul 2004.
- [87] F. Benatti, R. Floreanini, and U. Marzolino. Entangling two unequal atoms through a common bath. *Phys. Rev. A*, 81:012105, Jan 2010.
- [88] Jiawei Hu and Hongwei Yu. Entanglement dynamics for uniformly accelerated two-level atoms. *Phys. Rev. A*, 91:012327, Jan 2015.
- [89] Boris Sokolov, Jorma Louko, Sabrina Maniscalco, and Iiro Vilja. Unruh effect and information flow. *Phys. Rev. D*, 101:024047, Jan 2020.
- [90] William K. Wootters. Entanglement of formation of an arbitrary state of two qubits. *Phys. Rev. Lett.*, 80:2245–2248, Mar 1998.
- [91] Stanislaw Kryszewski and Justyna Czechowska-Kryszk. Master equation - tutorial approach. 2008.
- [92] Goran Lindblad. On the generators of quantum dynamical semigroups. *Communications in Mathematical Physics*, 48(2):119–130, 1976.
- [93] Vittorio Gorini, Andrzej Kossakowski, and Ennackal Chandy George Sudarshan. Completely positive dynamical semigroups of n-level systems. *Journal of Mathematical Physics*, 17(5):821–825, 1976.

- [94] Zbigniew Ficek and Ryszard Tanas. Entangled states and collective nonclassical effects in two-atom systems. *Phys. Rep.*, 372(5):369–443, 2002.
- [95] Kensuke Gallock-Yoshimura and Robert B. Mann. Entangled detectors nonperturbatively harvest mutual information. *Phys. Rev. D*, 104:125017, Dec 2021.
- [96] Ting Yu and J. H. Eberly. Sudden death of entanglement. *Science*, 323(5914):598–601, 2009.