

Randomly Initialized One-Layer Neural Networks Make Data Linearly Separable

Promit Ghosal
MIT
promit@mit.edu

Srinath Mahankali
MIT
srinathm@mit.edu

Yihang Sun
MIT
kimisun@mit.edu

Abstract

Recently, neural networks have been shown to perform exceptionally well in transforming two arbitrary sets into two linearly separable sets. Doing this with a randomly initialized neural network is of immense interest because the associated computation is cheaper than using fully trained networks. In this paper, we show that, with sufficient width, a randomly initialized one-layer neural network transforms two sets into two linearly separable sets with high probability. Furthermore, we provide explicit bounds on the required width of the neural network for this to occur. Our first bound is exponential in the input dimension and polynomial in all other parameters, while our second bound is independent of the input dimension, thereby overcoming the curse of dimensionality. We also perform an experimental study comparing the separation capacity of randomly initialized one-layer and two-layer neural networks. With correctly chosen biases, our study shows for low-dimensional data, the two-layer neural network outperforms the one-layer network. However, the opposite is observed for higher-dimensional data.

1 Introduction

In the last decade, neural networks (NNs) have shown to be enormously successful in countless applications [27, 37, 14]. In particular, random initialization enhances the ability of NNs and makes computation cheaper than using fully trained NNs [17, 14, 3]. In data classification, a pre-processing step with a randomly initialized neural network (RINN) combined with any standard linear classifier has been observed to perform exceedingly well [21, 35, 46]. However, a theoretical justification for this is still mostly unexplored, until very recently [1, 11]. In this paper, we address the question of linear separability of datasets via RINNs.

Problem 1.1. Consider two bounded, possibly infinite sets $X^+, X^- \subset \mathbb{R}^d$ that are δ -separated, i.e., $\|x^+ - x^-\| \geq \delta$ for all $x^+ \in X^+$ and $x^- \in X^-$. Let $F : \mathbb{R}^d \rightarrow \mathbb{R}^n$ denote the function defined by a feedforward neural network. The fundamental problem of data separability asks if $F(X^+)$ and $F(X^-)$ are *linearly separable*, i.e., if there exists a nonzero vector $a \in \mathbb{R}^n$ and constant c such that

$$a^T x + c = \begin{cases} > 0 & \text{if } x \in X^+ \\ < 0 & \text{if } x \in X^- \end{cases}.$$

The *margin* of separation of this hyperplane is defined as $\inf\{|a^T x + c|/\|a\| : x \in X^+ \cup X^-\}$.

This problem probes the geometric underpinning of NNs by asking about its *separation capacity*. We show in this paper that a sufficiently wide one-layer RINN solves this problem, i.e., it transforms δ -separated sets into linearly separable sets with high probability. The width bound depends polynomially on the parameters of the random distributions of the RINN but exponentially in the input dimension in our first main result. The second result removes this exponential dependence.

1.1 Our Contributions

The focus of our study is twofold: shedding light on the geometry of the linear separation capacity of one-layer RINNs and doing extensive experimentation to compare the performance of one-layer and two-layer RINNs by varying the associated parameters, widths of the networks, and the input dimension of the given sets. Our contributions can be summarized as follows:

- (1) Our first main result, Theorem 1.3, shows that a ReLU-activated one-layer RINN transforms any two δ -separated sets into two linearly separable sets with high probability, where the required width is polynomial in the parameters of the random weight distributions and δ^{-1} . This result is an improvement on the bound on the width in [11, Theorem 2.3] where with depth two, the width grows exponentially in these variables. However, the bound on the width in Theorem 1.3 is exponential in the input dimension, which was not the case in [11].
- (2) Our second main result, Theorem 1.4, provides a different bound as Theorem 1.3 for the same problem. It shows that the width of the one-layer RINN does not need to grow exponentially in dimension, thereby overcoming the curse of dimensionality. However, it grows exponentially in certain parameters relating to the weights and input datasets.
- (3) In showing the separation capacity of RINNs, we present two algorithms in Section 2.2 that together, on finite input datasets, construct a deterministic one-layer NN that makes them separable and also constructs an explicit hyperplane separating them. We hope to study this approach more in a future work.
- (4) We conduct an empirical study to compare the separation capacity of one-layer and two-layer RINNs and report the results in Section 3. We address the effects the random weight parameters, the width, and the depth of the network have on the separation capacity. We demonstrate how the two-layer RINN can outperform the one-layer RINN by a data-driven choice of parameters for any width. However, as dimension increases, our study shows the opposite can hold, indicating the importance of a dimension-dependent choice of depth.

1.2 Setup and Notations

Let $\|\cdot\|$ be the Euclidean norm. Let $\mathbb{B}_2^d(c, R)$ be the Euclidean ball of radius R in \mathbb{R}^d centered around c . We often use the shorthand $R\mathbb{B}_2^d := \mathbb{B}_2^d(0, R)$. Let \mathbb{S}^{d-1} be the surface of \mathbb{B}_2^d . Also, define the distance between any point $x \in \mathbb{R}^d$ and a set $S \subset \mathbb{R}^d$ as $\text{dist}(x, S) := \inf\{\|x - y\| : y \in S\}$.

In this paper, the *rectified linear activation function* (ReLU) defined by $\text{ReLU}(x) = x\mathbb{1}_{\{x>0\}}$ and extended to \mathbb{R}^n coordinate-wise will be our standalone choice of activation function. We denote the standard Gaussian matrix and vector where each entry is i.i.d. $N(0, 1)$ by $N(0, I_{n \times d})$ and $N(0, I_d)$ with appropriate dimensions. We sometimes refer to NNs by the function it defines. For a one-layer NN, it is $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}^n$ where $\Phi(x) = \text{ReLU}(Wx + b)$ for weights $W \in \mathbb{R}^{n \times d}$ and bias $b \in \mathbb{R}^n$.

The *mutual complexity* of any two finite sets X^+ and X^- is measured by the total number of points $N = |X^+| + |X^-|$. For two arbitrary sets $X^+, X^- \subset R\mathbb{B}_2^d$, we use the notion of mutual complexity indexed by parameters (δ, μ) where δ measures their separation and μ indexes the mutual metric entropy of those two sets [41, Section 7]. Our definition below is influenced by the one found in [11].

Definition 1.2. For any $n \in \mathbb{N}$, let $[n] := \{1, 2, \dots, n\}$. Let $X^-, X^+ \subset R\mathbb{B}_2^d$ be two δ -separated sets. Suppose there exist two finite sets of *centers* $C^+ = \{c_i^+\}_{i=1}^{N^+} \subset \mathbb{R}_2^d$ and $C^- = \{c_j^-\}_{j=1}^{N^-} \subset \mathbb{R}^d$, and two sets of constants $r_1^+, r_2^+, \dots, r_{N^+}^+, r_1^-, r_2^-, \dots, r_{N^-}^- \geq 0$ of *radii* such that

- (1) $X^+ \subset \bigcup_{i=1}^{N^+} X_i^+$ and $X^- \subset \bigcup_{j=1}^{N^-} X_j^-$ where $X_i^+ := X^+ \cap \mathbb{B}_2^d(c_i^+, r_i^+)$ and $X_j^- := X^- \cap \mathbb{B}_2^d(c_j^-, r_j^-)$ satisfies $c_i^+ \in \text{conv}(X_i^+)$ and $c_j^- \in \text{conv}(X_j^-)$ for all $i \in [N^+], j \in [N^-]$.
- (2) $r_i^+ \leq \mu^{-1} \text{dist}^2(c_i^+, C^-)$ for $i \in [N^+]$, and $r_j^- \leq \mu^{-1} \text{dist}^2(c_j^-, C^+)$ for $j \in [N^-]$,
- (3) C^+ and C^- are δ -separated.

Then, we say X^- and X^+ have (δ, μ) -mutual complexity N , where $N = N^- + N^+$.

1.3 Main Results

In this section, we state our two main results. We discuss their proof sketch in Section 2. For more details, we refer to Appendix A.6. Our first main result is twofold. The first part shows that any two finite δ -separated sets can be made linearly separable by applying a one-layer RINN; the second part shows the same result for two arbitrary δ -separated sets with a given mutual complexity N . Both parts of our first main result are proven for two different choices of weight distributions for the random initialization of the neural network: one result samples i.i.d. copies from \mathbb{S}^{d-1} while the other samples a standard Gaussian matrix.

Theorem 1.3. *Consider any two δ -separated sets $X^+, X^- \subset R\mathbb{B}_2^d$ with $d \geq 2$. Define the mutual complexity parameter N and the separation parameter γ by $N := |X^+| + |X^-|$ and $\gamma := \delta^2/8Rd$ if X^+, X^- are finite. Otherwise N is the $(\delta, 64\delta^{-2}R^3d)$ -mutual complexity and $\gamma := \delta^2/16Rd$. Let*

$$p := \frac{R}{8(d-1)\lambda} \left(\frac{\delta^2}{8R^2} \right)^d \quad \text{and} \quad \mathcal{M}(\gamma, N) := \sqrt{\frac{4R(R+\gamma)}{N(1+2R/\gamma)^{2N}-N}} \quad (1.1)$$

For any $\eta \in (0, 1)$, consider a one-layer ReLU-activated RINN $\Phi(x) = \text{ReLU}(Wx + b)$ with bias $b \sim \text{Unif}([- \lambda, \lambda]^n)$ where width n and maximal bias λ satisfy one of the following conditions:

- (i) $n \geq \frac{\log(N/\eta)}{p}$ and $\lambda \geq R$ if weights $W \in \mathbb{R}^{n \times d}$ satisfy that each row $W_i \stackrel{iid}{\sim} \text{Unif}(\mathbb{S}^{d-1})$,
- (ii) $n \geq \frac{10 \log(N/\eta)}{p}$ and $\lambda \geq 3R\sqrt{d}$ if weights $W \sim \text{N}(0, I_{n \times d})$ are standard Gaussian.

Then, the sets $\Phi(X^+), \Phi(X^-) \subset \mathbb{R}^n$ are linearly separable with probability at least $1 - \eta$ by a margin at least $\mathcal{M}(\gamma, N)$.

Remark. One can extend Theorem 1.3 to show the separation capacity of one-layer RINNs for ℓ -many δ -separated sets by running the argument between each set and all the other sets put together. Let N be the total number of points in the finite case or $(\delta, 64\delta^{-2}R^3d)$ -mutual complexity of all sets. Let γ be the same as in Theorem 1.3. With ℓ times the width lower bound, Φ makes each set linearly separable from the rest with the same margin and probability at least $1 - \ell\gamma$ by a union bound.

The following result extends the scope of Theorem 1.3 by improving the bound on the width of the one-layer RINN. Unlike the exponential dependence of width on dimension in Theorem 1.3, the proposed width in our next result will only grow polynomially with the dimension of the data. To state the result, we define the *Gaussian width* $w(T)$ of a subset $T \subset \mathbb{R}^n$ ([41, Definition 7.5.1]) as

$$w(T) := \mathbb{E} \left[\sup_{x \in T} g^T x \right], \quad \text{where } g \sim \text{N}(0, I_{n \times n}). \quad (1.2)$$

Theorem 1.4. *Let $X^+, X^- \subset R\mathbb{B}_2^d$ be δ -separated sets with $d \geq 2$. There exists a constant $C > 0$ depending only on δ and R such that the following holds. Define the mutual complexity parameter $N := |X^+| + |X^-|$ and the separation parameter $\gamma := \delta^2/18R\sqrt{k}$ if X^+ and X^- are finite where*

$$k := C \left(\frac{32R}{\delta^2} \right)^2 \left(w^2((X^+ - X^-) \cup X^+ \cup X^-) + R^2 \right). \quad (1.3)$$

Otherwise N is the $(\delta, 144\delta^{-2}R^3\sqrt{k})$ -mutual complexity and $\gamma := \delta^2/36R\sqrt{k}$.

Consider a one-layer ReLU-activated RINN $\Phi(x) = \text{ReLU}(Wx + b)$ with standard Gaussian weights $W \sim \text{N}(0, I_{n \times d})$ and bias $b \sim \text{Unif}([- \lambda, \lambda]^n)$ for maximal bias $\lambda \geq 9R\sqrt{k}/8$. Then, for any $\eta \in (0, 1)$, the sets $\Phi(X^+), \Phi(X^-) \subset \mathbb{R}^n$ are linearly separable with probability at least $1 - \eta$ by a margin at least $\mathcal{M}(\gamma, N)$ defined in (1.1) provided that width n satisfies the following condition

$$n \geq \frac{\log(N/\eta)}{q}, \quad \text{where } q := \frac{R}{4\lambda\sqrt{k}} \left(\frac{2\delta^2}{81R^2} \right)^k. \quad (1.4)$$

1.4 Related Works

Expressivity of NNs has been the focus of the literature on NNs since the 1960s. Earlier works along this direction study the memorization capacity of linear threshold networks [7, 9, 24, 38]. Later,

their counterparts with sigmoids, ReLU, and other bounded activation functions have been studied [22, 20, 19, 43]. More works show the empirical evidence of training and performance advantages [29, 25, 28, 13, 10, 42]. While the amount of empirical evidence is highly impressive, only few provided theoretical justification for such performance. Most theoretical work focuses on topics such as universal approximation power [39, 2, 44, 30, 18], finite sample expressivity, or memorization capacity of NNs [46, 16, 31, 45], none of which guarantees that NNs perform well in data separation.

The separation capacity of NNs has recently been explored in [1], where the authors put forth a systematic geometric framework for showing that any two disjoint subsets $X^+, X^- \subset \mathbb{R}^d$ can be linearly separated using a deterministic two-layer NN. Their construction has very recently been adopted for RINNs by [11] where a probabilistic approach is undertaken to study the separation capacity of two-layer RINNs, with asymptotic bounds on the output dimension, unlike the existential results in [1]. Our results show that linear separability of the data can actually be achieved with a one-layer RINN without the imposition of the curse of dimensionality.

Kernel-based methods are a major tool for linear separation of the data. Introduced by [34], random feature methods provide a cheap alternative to the kernel evaluation of the data via the construction of random feature maps in high dimensions. The generalization error of random feature maps is further studied in [34, 35, 4, 36] and are shown to be comparable to the kernel-based methods. The idea of using RINNs [1, 11] to separate data shares many similarities with the use of random feature methods. However, our results have no direct implication for the theory of random feature maps.

One of the key points in our analysis is the preservation of Euclidean distance between the points by RINNs with ReLU activation. This feature, which was first sketched in [1] and shown rigorously in [11], is reminiscent of nonlinear random embedding in compressed sensing [23, 33, 32, 12].

We end this section by discussing similarities between the separation capacity of one-layer RINNs in our main results and the rare eclipse problem. For any two linearly separable sets, the rare eclipse problem searches for a random lower-dimensional embedding that makes them disjoint with high probability. It was shown in [6] using Gordon’s escape through the mesh theorem [15] that multiplication by a random Gaussian matrix achieves such an embedding with a sharp lower bound on the dimension. This bound, which was stated (see [6, Corollary 3.2]) in terms of the Gaussian width of the difference between two sets, shares close ties with the expression of k in (1.3) of Theorem 1.3.

2 Proof Strategy

In this section, we discuss the main ideas behind the proofs of our main results. We first consider the geometry of linear separation in Section 2.1. Then, we show analogs of the main results for deterministic one-layer NN in Section 2.2 via Algorithm 1 and Algorithm 2 which search for the hyperplanes separating each data point of one set from the other set. Leveraging the algorithms, we explain the separation capacity of one-layer RINN for arbitrary sets in Section 2.3.

2.1 Key Insight: One-Layer ReLU Networks and Separation

Recall that i -th node of a neural network Φ encodes the i -th row W_i of the weight matrix W and the i -th entry b_i of b , i.e., $(\Phi(x))_i = \text{ReLU}(W_i^T x + b_i)$. For any $x \in \mathbb{R}^n$, assuming that $W_i \in \mathbb{S}^{d-1}$ is normalized, $h_i(x) = W_i^T x + b_i$ equals the signed distance between x and the hyperplane $\{h_i = 0\}$. Thus, $(\Phi(x))_i$ measures the distance of x and h_i if x is on the positive side of h_i and is zero otherwise.

To show $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable, it suffices to find a vector a such that $a^T \Phi(x^+) > 0$ for every $x^+ \in X^+$ and $a^T \Phi(x^-) < 0$ for every $x^- \in X^-$. Constructing such a boils down to pinning down which of the coordinates of $\Phi(x^+)$ for $x^+ \in X^+$ and $\Phi(x^-)$ for $x^- \in X^-$ are strictly positive and then assigning some weight vector a so that above property will be satisfied. We will use the following lemma that shows that the construction of the weight vector on a subset of the coordinates that satisfies the above property suffices to show the linear separability of $\Phi(X^+)$ and $\Phi(X^-)$. The proof of this lemma is rather simple and will be deferred to Appendix A.2 for brevity.

Lemma 2.1. *For $S^+, S^- \subset \mathbb{R}^k$ and any projection π onto a subspace indexed by coordinates $\Sigma \subset \{1, \dots, k\}$, if $\pi(S^+)$ and $\pi(S^-)$ are linearly separable with margin μ , then so are S^+ and S^- .*

2.2 A One-Layer Deterministic Neural Network Separates Finite Sets

Here, we sketch the deterministic NN analogs of the finite set cases of Theorem 1.3. We use Algorithm 1 below to construct the one-layer deterministic NN by drawing the hyperplanes separating one point at a time in descending order of norm. Before giving Algorithm 1, we introduce some notations. Recall k from (1.3). Let $N := |X^+| + |X^-|$ and define *sign* $\sigma : X^+ \cup X^- \rightarrow \{+1, -1\}$ by

$$\sigma(x) = +1 \quad \text{if } x \in X^+, \quad \text{and} \quad \sigma(x) = -1 \quad \text{if } x \in X^-. \quad (2.1)$$

Algorithm 1 To construct a one-layer neural network separating two finite sets with a margin

Input: Two δ -separated finite sets $X^+, X^- \subset R\mathbb{B}_2^d$ and $\gamma \in [0, \max(\delta^2/8Rd, \delta^2/18R\sqrt{k})]$.

Output: A one-layer neural network Φ such that $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable.

```

1: for  $i = 1, 2, \dots, N$  do
2:    $x_i \leftarrow x$  where  $x$  has the  $i$ -th largest norm in  $X^+ \cup X^-$ .
3: end for
4: for  $i = 1, 2, \dots, N$  do
5:   Find  $w \in \mathbb{S}^{d-1}$  and  $b \in \mathbb{R}$  such that  $w^T x_i + b \geq \gamma$  and  $w^T x_j + b \leq -\gamma$  for all  $j$  with  $j > i$ 
   and  $\sigma(x_i) \neq \sigma(x_j)$ . If no such  $j$  exists, ask  $w^T x_i + b \geq \gamma$  and  $w^T x + b \leq 0$  for some  $x \in R\mathbb{B}_2^d$ .
6:    $w_i \leftarrow w, b_i \leftarrow b, h_i \leftarrow$  the function  $w_i^T x + b_i$ 
7: end for
8:  $W \leftarrow$  the matrix with row vectors  $w_i^T$ 
9:  $b \leftarrow$  the vector with components  $b_i$ 
10: return the function  $\Phi(x) = \text{ReLU}(Wx + b)$ 

```

For convenience of visualization, we will often refer to h_i (see line 6 of Algorithm 1) as the hyperplane $\{h_i = 0\}$. The goal of Algorithm 1 is to construct the hyperplane h_i which separates x_i by a margin γ from all points $x_j \in X^+ \cup X^-$ which have smaller norm than x_i and $\sigma(x_j) \neq \sigma(x_i)$. It is easy to see that each hyperplane h_i exists in the case where no such j exists on line 5 (see Appendix A.6). However, it is not obvious why h_i exists when j exists. In the next result, we will show this via the probabilistic method that h_i exists when γ is at most either term in the maxima in the input condition.

Theorem 2.2. *For any two δ -separated finite sets $X^+, X^- \subset R\mathbb{B}_2^d$, define x_i and σ as in Algorithm 1 and (2.1). Consider the random affine function $h(z) = v^T z + t$ where $t \in \mathbb{R}, v \in \mathbb{R}^d$ are independent random variables, $t \sim \text{Unif}([- \lambda, \lambda])$, and the distribution of v is specified below. For any $\gamma > 0$, let*

$$B_i := \{h(x_i) \geq \gamma\} \cap \bigcap_{j: j > i, \sigma(x_i) \neq \sigma(x_j)} \{h(x_j) \leq -\gamma\}.$$

for each $i \in \{1, 2, \dots, N\}$. Recall p from (1.1) and q from (1.4). There exists a constant C depend only on δ and R in the definition (1.3) of k such that the following statements hold.

- (i) For all i and $\gamma \leq \delta^2/8Rd$, $\mathbb{P}(B_i) \geq p$ if $v \sim \text{Unif}(\mathbb{S}^{d-1})$ and $\lambda \geq R$.
- (ii) For all i and $\gamma \leq \delta^2/8Rd$, $\mathbb{P}(B_i) \geq p/10$ if $v \sim \text{N}(0, I_d)$ and $\lambda \geq 3R\sqrt{d}$.
- (iii) For all i and $\gamma \leq \delta^2/18R\sqrt{k}$, $\mathbb{P}(B_i) \geq q$ if $v \sim \text{N}(0, I_d)$ and $\lambda \geq 9R\sqrt{k}/8$.

Assuming the existence of hyperplanes h_i , we now argue the correctness of the algorithm and bound the margin of separation, thereby showing that one deterministic layer separates finite sets.

Theorem 2.3. *For output Φ of Algorithm 1 on any sets X^+, X^- and parameter γ satisfying the input conditions, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable with margin at least $\sqrt{N}\mathcal{M}(\gamma, N)$ in (1.1).*

Proof Sketch of Theorem 2.2. Since $\mathbb{P}(B_i)$ increases as γ decreases, it suffices to show lower bounds for $\mathbb{P}(B_i)$ when γ is equal to the respective upper bounds in (i), (ii), and (iii).

For each i , we draw two balls $\mathbb{B}_2^d(x_i, \delta)$ and $\|x_i\|\mathbb{B}_2^d$ as in the left picture in Figure 1. Since x_i is separated by δ from all x such that $\sigma(x) \neq \sigma(x_i)$, the points x_j with higher index than x_i (i.e., smaller norm) and different *sign* (e.g. the blue points x_{i+1} and x_{i+2}) must lie in $\|x_i\|\mathbb{B}_2^d \setminus \mathbb{B}_2^d(x_i, \delta)$.

In particular, they lie below the orange hyperplane formed by the intersection of the surfaces of the two balls. Roughly, we see that the distance between the orange hyperplane and x_i is large enough compared to the choices of γ , so we can find a hyperplane h through the region between x_i and the orange hyperplane while maintaining distance at least γ to both of them. We see that any such h satisfies B_i as all x_j 's lie below the orange hyperplane, so their distance to h is at least that of the orange hyperplane, namely γ . Now, we discuss how to bound probability of this occurring when $h(z) = v^T z + t$ is chosen randomly as in (i), (ii), and (iii). Detailed calculations are in Appendix A.3.

For Case (i) where $v \sim \text{Unif}(\mathbb{S}^{d-1})$, we condition on v being close to $x_i/\|x_i\|$. Then, h will be roughly parallel to the orange hyperplane with normal vector x_i , so we can lower bound the probability that t translates h to go through the region while maintaining a γ distance. Case (ii) follows by relating the Gaussian v to the uniform v in Case (i). We defer it to Appendix A.3.

For Case (iii) where $v \sim \text{N}(0, I_d)$, by [11, Proof of Theorem 1.6], v is distributed identically as $G^T u$ with independent $G \sim \text{N}(0, I_{k \times d})$ and $u \sim \text{Unif}(\mathbb{S}^{k-1})$ for all $k \in \mathbb{N}$. Let $A := G/\sqrt{k}$ and $H(z) := u^T z + t/\sqrt{k}$. Note that $h(z) = H(Az)/\sqrt{k}$ for all z , so the condition B_i on h is equivalent to $H(Ax_i) \geq \gamma/\sqrt{k}$ and $H(Ax_j) \leq -\gamma/\sqrt{k}$ for all $j > i$ and $\sigma(x_i) \neq \sigma(x_j)$. We want to apply Case (i) to points $Ax_i \in \mathbb{R}^k$, set $\{Ax_j : j > i, \sigma(x_j) \neq \sigma(x_i)\} \subset \mathbb{R}^k$, and hyperplane H . To do so, we need to show δ -separation and our ordering of x_i is preserved when we multiply everything by A . This follows the Matrix Deviation Inequality in [41, Exercise 9.1.8]: there exists a constant C in the definition (1.3) of k such that with at least constant probability, $\|Ax\| - \|x\| \leq \theta := \delta^2/32R$ for all $x \in X^+ \cup X^- \cup (X^+ - X^-)$. Conditioned on it, a similar proof as Case (i) suffices. \square




Figure 1: **Illustration of Algorithm 1 and Theorem 2.3.** Given input sets X^+ and X^- , let x_i be the point with the i -th largest norm. The left picture illustrates how Algorithm 1 picks each h_i . See the proof sketch of Theorem 2.3. The right picture is a concrete example of this on a input with 6 points.

Proof Sketch of Theorem 2.3. Define x_i and h_j as in Algorithm 1 and σ as in (2.1). Each h_j exists by Theorem 2.2. We can see that $(\Phi(x_i))_j = \text{ReLU}(h_j(x_i))$ is at least γ for $j = i$, exactly zero for all $j < i$ such that $\sigma(x_j) \neq \sigma(x_i)$, and at least zero for all other coordinates. We state an algorithm that uses this to find a hyperplane separating $\Phi(X^+)$ and $\Phi(X^-)$ with the desired margin.

Algorithm 2 To separate the image of the two sets under the deterministic network from Algorithm 1

Input: Two δ -separated finite sets $X^+, X^- \subset \mathbb{R}\mathbb{B}_2^d$ and parameter γ satisfying the input conditions of Algorithm 1, and the output Φ of Algorithm 1 on X^+, X^- , and γ .

Output: A hyperplane $H \subset \mathbb{R}^N$ separating $\Phi(X^+)$ and $\Phi(X^-)$ with margin at least $\sqrt{N}\mathcal{M}(\gamma, N)$.

1: **for** $i = N, N-1, \dots, 1$ **do**

2: $a_i \leftarrow \sigma(x_i) \left(1 + \left| \sum_{j=i+1}^N a_j (\Phi(x_i))_j \right| \right) / \gamma$.

3: **end for**

4: **return** the hyperplane $H(z) = a^T z$ where $a = (a_i)_i \in \mathbb{R}^N$.

See Appendix A.4 for the correctness of above including a bound on the margin of separation. \square

2.3 Generalizations to Arbitrary Sets and Random Initialization

We first generalize Theorem 2.3 to arbitrary sets X^+ and X^- which may not be finite. With carefully chosen margin parameter γ and mutual complexity parameters, if the output of Algorithm 1 separates the centers in Definition 1.2, the next theorem shows that it must separate the entire sets X^+ and X^- .

Theorem 2.4. *Let $X^+, X^- \subset R\mathbb{B}_2^d$ be any two δ -separated sets and k as defined in (1.3). For any parameter $\gamma \in [0, \max(\delta^2/8Rd, \delta^2/18R\sqrt{k})]$, let N be the $(\delta, 8R^2/\gamma)$ -mutual complexity of X^+ and X^- and let centers C^+ and C^- be as guaranteed by Definition 1.2. Then*

- (i) *The centers C^+ and C^- with parameter γ satisfy the input conditions of Algorithm 1.*
- (ii) *Let Φ be the output of Algorithm 1 on C^+ and C^- with parameter γ . Then, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable with margin at least $\sqrt{N}\mathcal{M}(\gamma/2, N)$ where \mathcal{M} is as in (1.1).*

Proof Sketch. (i) follows from Definition 1.2 and the assumptions on X^+, X^- , and γ . We defer the proof to Appendix A.5. For (ii), the radii used in Definition 1.2 for $(\delta, 8R^2/\gamma)$ -mutual complexity can be bounded above by $\gamma/2$. By Theorem 2.3 on the finite sets C^+ and C^- , for each center of C^+ , we get a hyperplane that is γ away from that center and any center from C^- with smaller norm, and vice versa. Then, those hyperplanes are at least $\gamma/2$ away from any point in the balls guaranteed by Definition 1.2 around those centers. The result now follows from the proof of Theorem 2.3. \square

To generalize Theorem 2.3 to one-layer RINN Φ , we show that given sufficient width, for every i , with high probability there is a node in Φ that defines a hyperplane satisfying the conditions for h_i in Algorithm 1. Then, those nodes collected over all i will make X^+ and X^- separable, which will be enough for Φ to also make them separable. This amounts to the following two conditions:

- (1) The condition for each h_i is satisfied by a random hyperplane with non-trivial probability.
- (2) The other hyperplanes that do not fulfill the conditions for any h_i do not hurt separation.

Notice that (1) is precisely Theorem 2.2 and (2) is precisely Lemma 2.1. Finally, we sketch our main results (Theorem 1.3 and Theorem 1.4) that a one-layer RINN separates sets.

Proof Sketches of Theorem 1.3 and Theorem 1.4. We first show both Theorem 1.3 and Theorem 1.4 in the case where X^+ and X^- are finite. Then, we generalize to the case of arbitrary sets.

For finite sets X^+ and X^- , let B_i^ℓ for $1 \leq i \leq N$ and $1 \leq \ell \leq n$ be the event that the ℓ -th random hyperplane H_ℓ of Φ satisfies the condition for h_i on line 5 of Algorithm 1.

We claim that $\Phi(X^+)$ and $\Phi(X^-)$ are separable with the desired margin if, for every i , B_i^ℓ holds for some ℓ . If this holds, then we can find some hyperplanes in Φ that form an output of Algorithm 1 on X^+, X^- , and γ . We proceed as Theorem 2.3 to show these hyperplanes make X^+ and X^- separable via Algorithm 2. However, it is possible that the same H_ℓ may satisfy the conditions of multiple h_i 's, so we slightly modify Algorithm 2 to fix this issue (see Appendix A.6). This modification shrinks the separation margin by a factor of \sqrt{N} compared to Theorems 2.3 and 2.4. Then, by Lemma 2.1, Φ makes X^+ and X^- separable with the desired margin, thus proving the claim.

Now, it suffices to bound $\mathbb{P}((\forall i)(\exists \ell)B_i^\ell)$. We show that for any i and ℓ , $\mathbb{P}(B_i^\ell) \geq p$ or $p/10$ or q depending on the distribution of Φ . This holds by Theorem 2.2 if j exists on line 5 of Algorithm 1. Otherwise, it follows from a calculation in Appendix A.6. We now show the finite set case for Theorem 1.3(i). By a union bound over i and independence of ℓ

$$\mathbb{P}(\Phi(X^+), \Phi(X^-) \text{ are separable}) \geq \mathbb{P}((\forall i)(\exists \ell)B_i^\ell) \geq 1 - \sum_{i=1}^N \mathbb{P}((\forall \ell)(B_i^\ell)^C) \geq 1 - N(1-p)^n$$

which is at least $1 - \eta$ if $n \geq \log(N/\eta)/p$. The finite set cases of Theorem 1.3(ii) and Theorem 1.4 follow similarly by replacing p with $p/10$ and q , respectively.

For arbitrary sets X^+ and X^- , let B_i^ℓ be the same except based on the centers C^+ and C^- as guaranteed by Definition 1.2. Then, the argument above combined with Theorem 2.4 shows that $\Phi(X^+)$ and $\Phi(X^-)$ are separable with the desired margin if, for every i , B_i^ℓ holds for some ℓ . Since C^+ and C^- are δ -separated by Definition 1.2, the arbitrary set cases of Theorem 1.3 and Theorem 1.4 now follow from the proof of the finite set cases applied to C^+ and C^- . \square

3 Experimental Results

In this section, we empirically compare the performance of one-layer and two-layer RINNs with the same width on simulated datasets (all experiments are done using a single personal computer). Given a point $x \in \mathbb{R}^d$, we slightly modify the definition of the one-layer RINN Φ to

$$\Phi(x) = \sqrt{\frac{2}{n}} \text{ReLU}(Wx + b), \quad (3.1)$$

where n is the width of the layer, W is a $d \times n$ matrix with standard Gaussian entries, and $b \in \mathbb{R}^n$ has entries uniformly sampled from $[-\lambda, \lambda]$ for a fixed λ . We normalize the neural network Φ by $\sqrt{2/n}$ so that for each point $x \in X$, $\|\Phi(x)\|_2^2 \approx \|x\|_2^2 + \lambda^2/3$ with high probability, as shown in [11, Theorem 4.1]. Our two-layer network adds an extra layer $\hat{\Phi}$ so that the network $\hat{\Phi} \circ \Phi$ satisfies

$$\hat{\Phi}(\Phi(x)) = \sqrt{\frac{2}{n}} \text{ReLU} \left(\hat{W} \left(\sqrt{\frac{2}{n}} \text{ReLU}(Wx + b) \right) + \hat{b} \right), \quad (3.2)$$

where \hat{W} is an $n \times n$ matrix with standard Gaussian entries, and $\hat{b} \in \mathbb{R}^n$ has entries uniformly sampled from $[-\hat{\lambda}, \hat{\lambda}]$, where $\hat{\lambda}$ does not need to be equal to λ .

3.1 Choosing λ and $\hat{\lambda}$

We choose $\hat{\lambda}$ so that with high probability, $\hat{\lambda}$ is approximately equal to the radius of the set of points $\Phi(X)$. Combining this motivation with the estimate $\|\Phi(x)\|_2^2 \approx \|x\|_2^2 + \lambda^2/3$, we choose

$$\hat{\lambda} = \sqrt{R^2 + \frac{\lambda^2}{3}}. \quad (3.3)$$

We first consider a dataset $X \subset \mathbb{R}^2$ of three concentric rings, as shown in Figure 2. The radius of X is approximately 360. We consider randomly initialized one-layer and two-layer networks of the form (3.1) and (3.2) with width $n = 30$. We compute the output $\Phi(x)$ for each point $x \in X$, and consider whether the classes of points are linearly separable after applying this transformation by using a linear SVM [8]. To estimate the probability that the classes are linearly separable, we repeat this 1000 times and compute the proportion of times that the SVM achieves 100% accuracy. We repeat this for different choices of λ . We also repeat this for a two-layer network $\hat{\Phi} \circ \Phi$, where $\hat{\lambda}$ is chosen according to (3.3). Finally, we repeat this calculation for the case where we choose $\hat{\lambda} = \lambda$.




Figure 2: **Left:** Original dataset of three concentric rings with 100 points each. **Middle:** Estimated separation probability of one-layer and two-layer networks with width 30 and different choices of maximal bias λ . The separation probability of a two-layer network where $\hat{\lambda} = \lambda$ is in green. **Right:** Estimated separation probability of one-layer and two-layer networks for different widths.

We note that Theorem 1.3 applies for $\lambda \geq 3R\sqrt{d}$, which is approximately 1500 since $d = 2$. However, as shown in Figure 2, the probability that the dataset is linearly separable after plugging it into the neural network peaks around $\lambda = 200$ for both one-layer and two-layer networks (shown in blue and orange, respectively). Furthermore, if we instead choose $\hat{\lambda} = \lambda$, the performance of the two-layer network (shown in green) is much worse compared to choosing $\hat{\lambda}$ according to (3.3). In fact, for certain choices of λ , the two-layer network performs worse than the one-layer network. However, the two-layer network performs much better than the one-layer network when $\hat{\lambda}$ is chosen using (3.3). This suggests that, in some special cases, it is possible to prove sharper bounds of the probability that a randomly initialized two-layer network separates two sets.

Finally, we compare the performance of one-layer and two-layer networks on the dataset shown in Figure 2. Motivated by the plots in the middle of Figure 2, where the optimal choice of λ is much less than $3R\sqrt{2}$, we set $\lambda = R$ and $\hat{\lambda} = R\sqrt{4/3}$ according to (3.3). The plot on the right of Figure 2 shows the separation probability of one-layer and two-layer networks with varying widths, and the two-layer network performs better.

3.2 Comparing Performance on High-Dimensional Data

Next, we consider a second dataset $X \subset \mathbb{R}^{100}$ of three concentric hyperspheres with radius approximately 360 and 100 points in each ring, similar to Section 3.1. We choose X of this form to observe the effect of high dimensionality of data on the required width for an RINN to make two sets linearly separable, and we visualize the data in the left of Figure 3 using t-SNE [40]. In the right of Figure 3, we compare the performance of one-layer and two-layer networks. Compared to the previous dataset, which is 2-dimensional, this dataset is 100-dimensional. However, the necessary width to consistently separate the two sets of points increases from over 60 to approximately 150, multiplying by a factor of less than 3. Thus, the effect of dimensionality on the required width does not appear to be very major. We also note that the one-layer network performs better than the two-layer network in Figure 3, contrasting with the rightmost figure in Figure 2.




Figure 3: **Left:** Visualization of the 100-dimensional dataset using t-SNE. **Right:** Estimated separation probability of one-layer and two-layer neural network for different widths.

4 Conclusion

We showed that a sufficiently wide one-layer RINN transforms two δ -separated sets in \mathbb{R}^d into two linearly separable sets with high probability. We derived explicit bounds on the width of the network for this to happen in two distinct scenarios. One of our bounds depends exponentially on d , but polynomially on all the parameters of the weight distribution of the RINN, thereby improving on the existing bound on the width when the two sets are d -dimensional. In comparison, our second bound depends polynomially on d , overcoming the curse of dimensionality. We performed an empirical study comparing the linear separation capacity of one-layer and two-layer RINNs. Our findings underline the importance of appropriate choices of parameters of weight distribution and demonstrate that the separation capacity of one-layer RINNs can outperform that of two-layer RINNs when d is

large. These observations warrant a detailed study on the effect of increasing depth on the separation capacity of RINNs, which we hope to pursue in future work.

Acknowledgments

YS was supported through the Simons Investigator Award grant (622132) of Elchanan Mossel and the UROP program of MIT. PG, SM and YS thankfully acknowledge the productive research environment of the mathematics department at MIT.

References

- [1] S. An, F. Boussaid, and M. Bannour. How can deep rectifier networks achieve linear separability and preserve distances? In *International Conference on Machine Learning*, pages 514–523. PMLR, 2015.
- [2] A. Andoni, R. Panigrahy, G. Valiant, and L. Zhang. Learning sparse polynomial functions. In *Proceedings of the twenty-fifth annual ACM-SIAM symposium on Discrete algorithms*, pages 500–510. SIAM, 2014.
- [3] D. Arpit and Y. Bengio. The benefits of over-parameterization at initialization in deep relu networks. *arXiv preprint arXiv:1901.03611*, 2019.
- [4] F. Bach. On the equivalence between kernel quadrature rules and random feature expansions. *The Journal of Machine Learning Research*, 18(1):714–751, 2017.
- [5] K. Ball et al. An elementary introduction to modern convex geometry. *Flavors of geometry*, 31(1-58):26, 1997.
- [6] A. S. Bandeira, D. G. Mixon, and B. Recht. Compressive classification and the rare eclipse problem. In *Compressed Sensing and its Applications*, pages 197–220. Springer, 2017.
- [7] E. B. Baum. On the capabilities of multilayer perceptrons. *Journal of complexity*, 4(3):193–215, 1988.
- [8] C. Cortes and V. Vapnik. Support-vector networks. *Machine learning*, 20(3):273–297, 1995.
- [9] T. M. Cover. Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition. *IEEE transactions on electronic computers*, (3):326–334, 1965.
- [10] G. E Dahl, T. N. Sainath, and G. E. Hinton. Improving deep neural networks for lvcsr using rectified linear units and dropout. *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 8609–8613, 2013.
- [11] S. Dirksen, M. Genzel, L. Jacques, and A. Stollenwerk. The separation capacity of random neural networks. *arXiv preprint arXiv:2108.00207*, 2021.
- [12] S. Dirksen and S. Mendelson. Non-gaussian hyperplane tessellations and robust one-bit compressed sensing. *Journal of the European Mathematical Society*, 23(9):2913–2947, 2021.
- [13] X. Glorot, A. Bordes, and Y. Bengio. Deep sparse rectifier networks. *Proceedings of the 14th International Conference on Artificial Intelligence and Statistics*, 15:315–323, 2011.
- [14] I. Goodfellow, Y. Bengio, and A. Courville. *Deep learning*. MIT press, 2016.
- [15] Y. Gordon. On milman’s inequality and random subspaces which escape through a mesh in \mathbb{R}^n . In *Geometric aspects of functional analysis*, pages 84–106. Springer, 1988.
- [16] M. Hardt and T. Ma. Identity matters in deep learning. *arXiv preprint arXiv:1611.04231*, 2016.
- [17] K. He, X. Zhang, S. Ren, and J. Sun. Delving deep into rectifiers: Surpassing human-level performance on imagenet classification. In *Proceedings of the IEEE international conference on computer vision*, pages 1026–1034, 2015.
- [18] D. Hsu, C. H. Sanford, R. Servedio, and Emmanouil Vasileios Vlatakis-Gkaragkounis. On the approximation power of two-layer networks of random relus. In *Conference on Learning Theory*, pages 2423–2461. PMLR, 2021.

- [19] G. B. Huang. Learning capability and storage capacity of two-hidden-layer feedforward networks. *IEEE transactions on neural networks*, 14(2):274–281, 2003.
- [20] G. B. Huang and H. A. Babri. Upper bounds on the number of hidden neurons in feedforward networks with arbitrary bounded nonlinear activation functions. *IEEE transactions on neural networks*, 9(1):224–229, 1998.
- [21] G. B. Huang, Q. Y. Zhu, and C. K. Siew. Extreme learning machine: theory and applications. *Neurocomputing*, 70(1-3):489–501, 2006.
- [22] S. C. Huang and Y. F. Huang. Bounds on the number of hidden neurons in multilayer perceptrons. *IEEE transactions on neural networks*, 2(1):47–55, 1991.
- [23] L. Jacques, J. N. Laska, P. T. Boufounos, and R. G. Baraniuk. Robust 1-bit compressive sensing via binary stable embeddings of sparse vectors. *IEEE Transactions on Information Theory*, 59(4):2082–2102, 2013.
- [24] A. Kowalczyk. Estimates of storage capacity of multilayer perceptron with threshold logic hidden units. *Neural networks*, 10(8):1417–1433, 1997.
- [25] A. Krizhevsky, I. Sutskever, and G. E. Hinton. Imagenet classification with deep convolutional neural networks. *Advances in Neural Information Processing Systems*, pages 1097–1105, 2012.
- [26] B. Laurent and P. Massart. Adaptive estimation of a quadratic functional by model selection. *Annals of Statistics*, pages 1302–1338, 2000.
- [27] Y. LeCun, Y. Bengio, and G. Hinton. Deep learning. *nature*, 521(7553):436–444, 2015.
- [28] A. L. Maas, A. Y. Hannun, and A. Y. Ng. Rectifier nonlinearities improve neural network acoustic models. *ICML Workshop on Deep Learning for Audio, Speech, and Language Processing*, 2013.
- [29] V. Nair and G. E. Hinton. Rectified linear units improve restricted boltzmann machines. *Proceedings of the 27th International Conference on Machine Learning*, pages 807–814, 2010.
- [30] D. Needell, A. A. Nelson, R. Saab, and P. Salanevich. Random vector functional link networks for function approximation on manifolds. *arXiv preprint arXiv:2007.15776*, 2020.
- [31] Q. Nguyen and M. Hein. Optimization landscape and expressivity of deep cnns. In *International conference on machine learning*, pages 3730–3739. PMLR, 2018.
- [32] S. Oymak and B. Recht. Near-optimal bounds for binary embeddings of arbitrary sets. *arXiv preprint arXiv:1512.04433*, 2015.
- [33] Y. Plan and R. Vershynin. Dimension reduction by random hyperplane tessellations. *Discrete & Computational Geometry*, 51(2):438–461, 2014.
- [34] A. Rahimi and B. Recht. Random features for large-scale kernel machines. *Advances in neural information processing systems*, 20, 2007.
- [35] A. Rahimi and B. Recht. Weighted sums of random kitchen sinks: Replacing minimization with randomization in learning. *Advances in neural information processing systems*, 21, 2008.
- [36] A. Rudi and L. Rosasco. Generalization properties of learning with random features. *Advances in neural information processing systems*, 30, 2017.
- [37] J. Schmidhuber. Deep learning in neural networks: An overview. *Neural networks*, 61:85–117, 2015.
- [38] E. D. Sontag. Shattering all sets of ‘k’ points in “general position” requires $(k-1)/2$ parameters. *Neural Computation*, 9(2):337–348, 1997.
- [39] Y. Sun, A. Gilbert, and A. Tewari. On the approximation properties of random relu features. *arXiv preprint arXiv:1810.04374*, 2018.
- [40] L. Van der Maaten and G. Hinton. Visualizing data using t-sne. *Journal of machine learning research*, 9(11), 2008.
- [41] R. Vershynin. *High-dimensional probability*, volume 47 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2018. An introduction with applications in data science, With a foreword by Sara van de Geer.
- [42] Sun Y., Chen Y., Wang X., and Tang X. Deep learning face representation by joint identification-verification. *Advances in Neural Information Processing Systems*, pages 1988–1996, 2014.

- [43] G. B. Yamasaki. The lower bound of the capacity for a neural network with multiple hidden layers. *ICANN*, 14(2):546–549, 1993.
- [44] G. Yehudai and O. Shamir. On the power and limitations of random features for understanding neural networks. *Advances in Neural Information Processing Systems*, 32, 2019.
- [45] C. Yun, S. Sra, and A. Jadbabaie. Small relu networks are powerful memorizers: a tight analysis of memorization capacity. *Advances in Neural Information Processing Systems*, 32, 2019.
- [46] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals. Understanding deep learning requires rethinking generalization. arxiv preprint (in iclr 2017). *arXiv preprint arXiv:1611.03530*, 2016.

A Appendix

A.1 Some Auxiliary Lemmas

In this section, we will state and prove a few results in convex geometry and probability.

Lemma A.1. Fix $d \in \mathbb{Z}_{\geq 2}$. Denote the convex hull of X by $\text{conv}(X)$.

(1) For any $0 \leq r \leq 2$, $u \in \mathbb{S}^{d-1}$, and $v \sim \text{Unif}(\mathbb{S}^{d-1})$, we have

$$\mathbb{P}_v(\|u - v\| \leq r) \geq \frac{1}{2} \left(\frac{r}{2}\right)^{d-1}.$$

(2) For any set $X \subset \mathbb{R}^d$ and $c \in \text{conv}(X)$, there exists $x \in X$ with $\|x\| \geq \|c\|$. Moreover, for any vector $g \in \mathbb{R}^d$, then there exists some $x \in X$ such that $g^T x \geq g^T c$.

Proof. (1) follows from [5, Lemma 2.3]. For (2), note that since $c \in \text{conv}(X)$, there are weights $\{w_x : x \in X\} \subset [0, 1]$ such that

$$c = \sum_{x \in X} w_x x \quad \text{and} \quad \sum_{x \in X} w_x = 1$$

By the Triangle Inequality and linearity of inner product with g

$$\|c\| \leq \sum_{x \in X} w_x \|x\| \quad \text{and} \quad g^T c = \sum_{x \in X} w_x g^T x.$$

Since w_x are non-negative and $\sum_{x \in X} w_x = 1$, (2) follows directly from the above display. \square

Lemma A.2. Let χ_d^2 be the chi-squared distributions with parameter d .

(1) For $v \sim N(0, I_d)$, $\rho := \|v\|$ satisfy $\rho^2 \sim \chi_d^2$, and $\hat{v} := v/\rho \sim \text{Unif}(\mathbb{S}^{d-1})$ are independent.

(2) $\mathbb{P}(z - d \geq 2\sqrt{dx} + 2x) \leq e^{-x}$ and $\mathbb{P}(d - z \geq 2\sqrt{dx}) \leq e^{-x}$ for all $x > 0$ and $z \sim \chi_d^2$.

Proof. (1) is well-known. (2) follows from [26, Lemma 1]. \square

Proposition A.3. Suppose $\rho^2 \sim \chi_d^2$, then $\mathbb{P}(1 \leq \rho \leq 3\sqrt{d}) \geq 1/10$.

Proof. We apply the second inequality of Lemma A.2(2) with $x = (d-1)^2/4d$ to obtain

$$\mathbb{P}(\rho \leq 1) = \mathbb{P}(d - \rho^2 \geq d - 1) \leq e^{-(d-1)^2/4d}.$$

The derivative of $\exp(-(d-1)^2/4d)$ with respect to d is $-(d^2-1)\exp(-(d-1)^2/4d)/4d^2$ which is negative for all $d \geq 2$. Hence, on $d \geq 2$, $\exp(-(d-1)^2/4d)$ attains its maximum at $d = 2$, so

$$\mathbb{P}(\rho \leq 1) \leq e^{-(d-1)^2/4d} \leq e^{-1/8}.$$

Now, let $x = 9/2$. We see that $9d \geq d + \sqrt{2xd} + dx$ for $d \geq 2$, so

$$\begin{aligned} \mathbb{P}(\rho \geq 3\sqrt{d}) &= \mathbb{P}(\rho^2 \geq 9d) \leq \mathbb{P}(\rho^2 \geq d + \sqrt{2xd} + dx) \\ &\leq \mathbb{P}(\rho^2 \geq d + 2\sqrt{dx} + 2x) \leq \mathbb{P}(\rho^2 - d \geq 2\sqrt{dx} + 2x) \leq e^{-9/2} \end{aligned}$$

Finally, by direct computation $\mathbb{P}(1 \leq \rho \leq 3\sqrt{d}) \geq 1 - e^{-1/8} - e^{-9/2} \geq 1/10$. \square

A.2 Proof of Lemma 2.1

Let $\Sigma = \{\sigma_1, \dots, \sigma_\ell\} \subset \{1, \dots, k\}$. If $\pi(S^+)$ and $\pi(S^-)$ are linearly separable, then there exists $a = (a_{\sigma(x_i)})_i \in \mathbb{R}^\ell$ with $\|a\| = 1$ and $b \in \mathbb{R}, \mu \in \mathbb{R}_{\geq 0}$ such that

$$a^T \pi(s) + b = \begin{cases} \geq \mu & \text{if } s \in S^+ \\ \leq -\mu & \text{if } s \in S^- \end{cases}$$

Let $\tilde{a} \in \mathbb{R}^k$ be equal to a on the ℓ coordinates indexed by Σ and 0 everywhere else, then

$$\tilde{a}^T s + b = \sum_{j=1}^k a_j s_j + b = \sum_{i=1}^{\ell} a_{\sigma(x_i)} s_{\sigma(x_i)} + b = a^T \pi(s) + b = \begin{cases} \geq \mu & \text{if } s \in S^+ \\ \leq -\mu & \text{if } s \in S^- \end{cases}$$

As $\|\tilde{a}\| = \|a\| = 1$, we have that S^+ and S^- are linearly separable with margin at least μ .

A.3 Proof of Theorem 2.2

In this section, we prove the lower bound on $\mathbb{P}(B_i)$ in Theorem 2.2 for the three different cases (i), (ii), and (iii) separately. We will show Case (i), reduce Case (ii) to Case (i), and separate Case (iii) further into two cases and will be shown using ideas from [11, Proof of Theorem 1.6].

Proof of Case (i). Here, $v \sim \text{Unif}(\mathbb{S}^{d-1})$. The idea is to condition on v being close to $x_i/\|x_i\|$ and find which t makes $h(z) = v^T z + t$ satisfy B_i . We break it down into steps.

Note, for any $x_j \in X^+ \cup X^-$ with $j > i$ and $\sigma(x_i) \neq \sigma(x_j)$, we know $\|x_j\| \leq \|x_i\|$ by our ordering and $\|x - x_i\| \geq \delta$ by δ -separation. First, for any j as in the definition of B_i , by the above

$$\delta^2 \leq (x_i - x_j)^T (x_i - x_j) = \|x_i\|^2 + \|x_j\|^2 - 2x_i^T x_j \leq 2\|x_i\|^2 - 2x_i^T x_j.$$

Hence, $x_i^T x_j \leq \|x_i\|^2 - \delta^2/2$. Now, define $\hat{x}_i = x_i/\|x_i\|$. For every $v \in \mathbb{S}^{d-1}$, we note $\hat{x}_i^T x_i = \|x_i\| \leq R$ and bound $(v - \hat{x}_i)^T x_i \geq -\|v - \hat{x}_i\| \cdot \|x_i\|$ by Cauchy-Schwarz to obtain

$$v^T x_i = \hat{x}_i^T x_i + (v - \hat{x}_i)^T x_i \geq \|x_i\| - \|v - \hat{x}_i\| \cdot \|x_i\| \geq \|x_i\| - R\|v - \hat{x}_i\| \quad (\text{A.1})$$

Now, for any j as in the definition of B_i , we had $x_i^T x_j \leq \|x_i\|^2 - \delta^2/2$, so $\hat{x}_i^T x_j \leq \|x_i\| - \delta^2/2\|x_i\|$. Since $\|x_j\| \leq \|x_i\| \leq R$ and $(v - \hat{x}_i)^T x_j \leq \|v - \hat{x}_i\| \cdot \|x_j\|$ by Cauchy-Schwarz, we obtain

$$v^T x_j = \hat{x}_i^T x_j + (v - \hat{x}_i)^T x_j \leq \|x_i\| - \frac{\delta^2}{2\|x_i\|} + \|v - \hat{x}_i\| \cdot \|x_j\| \leq \|x_i\| - \frac{\delta^2}{2R} + R\|v - \hat{x}_i\| \quad (\text{A.2})$$

Recall that $h(z) = v^T z + t$ and B_i occurs if and only if $v^T x_i + t \geq \gamma$ and $v^T x_j + t \leq -\gamma$ for all j where $j > i$ and $\sigma(x_j) \neq \sigma(x_i)$. This is equivalent to $t \geq \gamma - v^T x_i$ and $t \leq -\gamma - v^T x_j$ for all such j . We bound $|v^T x_i| \leq \|v\| \cdot \|x_i\| \leq R$ and similarly $|v^T x_j| \leq R$. Hence, for a fixed v , B_i occurs if

$$t \in I := \left[\gamma - v^T x_i, -\gamma - \max_{j:j>i, \sigma(x_j) \neq \sigma(x_i)} v^T x_j \right] \subset [-R, R] \subset [-\lambda, \lambda]$$

as $\lambda \geq R$, so may now bound $\mathbb{P}(B_i | v)$ below from by the length of I over 2λ , i.e.

$$\mathbb{P}_t(B_i | v) \geq \frac{1}{2\lambda} \left(v^T x_i - \max_{j:j>i, \sigma(x_j) \neq \sigma(x_i)} (v^T x_j) - 2\gamma \right) \geq \frac{1}{2\lambda} \left(\frac{\delta^2}{2R} - 2R\|v - \hat{x}_i\| - 2\gamma \right)$$

where the second inequality follows from (A.1) and (A.2). Let $r := (\delta^2/4R^2)(1 - 1/d) \in [0, 2]$ as $\delta \leq 2R$. As $v \sim \text{Unif}(\mathbb{S}^{d-1})$, we can apply Lemma A.1(1) to bound

$$\mathbb{P}(B_i) \geq \mathbb{P}(B_i | \|v - \hat{x}_i\| \leq r) \cdot \mathbb{P}_t(\|v - \hat{x}_i\| \leq r) \geq \frac{1}{2\lambda} \left(\frac{\delta^2}{2R} - 2Rr - 2\gamma \right) \left(\frac{1}{2} \right) \left(\frac{r}{2} \right)^{d-1}$$

Plug in r and bound $\gamma \leq \delta^2/8Rd$ to obtain

$$\begin{aligned}
\mathbb{P}(B_i) &\geq \frac{1}{4\lambda} \left[\frac{\delta^2}{2R} - 2R \cdot \frac{\delta^2}{4R^2} \left(1 - \frac{1}{d}\right) - 2 \cdot \frac{\delta^2}{8Rd} \right] \left[\frac{\delta^2}{8R^2} \left(1 - \frac{1}{d}\right) \right]^{d-1} \\
&= \frac{1}{4\lambda} \left(\frac{\delta^2}{4Rd} \right) \left(\frac{\delta^2}{8R^2} \right)^{d-1} \left(1 - \frac{1}{d}\right)^{d-1} \\
&\geq \frac{R}{8(d-1)\lambda} \left(\frac{\delta^2}{8R^2} \right)^d \\
&= p
\end{aligned}$$

where the second inequality follows from the fact that $(1 - 1/d)^d$ is increasing in d for $d \geq 2$, so it can be lower bounded by its value $1/4$ at $d = 2$. This establishes Case (i). \square

Proof of Case (ii). Here, $v \sim N(0, I_d)$. By Lemma A.2(1), we can change variables to $\rho = \|v\|$ with $\rho^2 \sim \chi_d^2$ and $\hat{v} = v/\rho \sim \text{Unif}(\mathbb{S}^{d-1})$, which are independent. For any $1 \leq \rho \leq 3\sqrt{d}$, we compute

$$\begin{aligned}
\mathbb{P}(B_i | \rho) &= \mathbb{P}(\hat{v}^T x + t/\rho \geq \gamma/\rho \forall x \in X^+ \text{ and } \hat{v}^T x + t/\rho \leq -\gamma/\rho \forall x \in X^-) \\
&\geq \mathbb{P}(\hat{v}^T x + t/\rho \geq \gamma \forall x \in X^+ \text{ and } \hat{v}^T x + t/\rho \leq -\gamma \forall x \in X^-) \\
&\geq \mathbb{P}(\hat{v}^T x + t/\rho \text{ satisfies event } B_i)
\end{aligned}$$

Now, $\hat{v} \sim \text{Unif}(\mathbb{S}^{d-1})$ and, for a fixed ρ , $t/\rho \sim \text{Unif}([-\lambda/\rho, \lambda/\rho])$. Since $\rho \leq 3\sqrt{d}$ and $\lambda \geq 3\sqrt{d}R$, $\lambda/\rho \geq R$. Hence, Case (i) applies to show

$$\mathbb{P}(B_i | \rho) \geq \mathbb{P}(\hat{v}^T x + t/\rho \text{ satisfies event } B_i) \geq p$$

This holds for every ρ where $1 \leq \rho \leq 3\sqrt{d}$. By conditioning on it, we obtain

$$\mathbb{P}(B_i) \geq \mathbb{P}(B_i | 1 \leq \rho \leq 3\sqrt{d}) \cdot \mathbb{P}(1 \leq \rho \leq 3\sqrt{d}) \geq \frac{1}{10} \inf \left\{ \mathbb{P}(B_i | \rho) : 1 \leq \rho \leq 3\sqrt{d} \right\} \geq \frac{p}{10}.$$

where $\mathbb{P}(1 \leq \rho \leq 3\sqrt{d}) \geq 1/10$ by Proposition A.3, showing the desired bound for Case (ii). \square

Proof of Case (iii). We prove this in several steps which are outlined here. First, we will relate this case to Case (i) but with ambient space \mathbb{R}^k instead of \mathbb{R}^d where k is defined in (1.3). The idea is to write the Gaussian vector v in \mathbb{R}^d as a product of a Gaussian random matrix and a Gaussian vector u in \mathbb{R}^k , and apply a Matrix Deviation Inequality [41, Exercise 9.1.8]. Then, we will separate into two sub-cases according to the norm of x_i . If $\|x_i\|$ is large, we will do a similar analysis as Case (i) in \mathbb{R}^k to get bound $\mathbb{P}(B_i) \geq q$; if $\|x_i\|$ is small, we will calculate $\mathbb{P}(B_i)$ and see it is much greater than q .

Choosing Dimension k For every positive integer k , $v \sim N(0, I_d)$ is distributed identically as $G^T u$ with independent $G \sim N(0, I_{k \times d})$ and $u \sim \text{Unif}(\mathbb{S}^{k-1})$. We can require C to be such that k defined in (1.3) satisfy $k \in \mathbb{Z}_{\geq 2}$. We will specify k now below.

Let $A := G/\sqrt{k}$. By [41, Exercise 9.1.8], for any constants $\theta > 0$ and $\omega \in [0, 1]$, there exists C in the definition of k in (1.3) such that with probability at least ω , for all $x \in X^+ \cup X^- \cup (X^+ - X^-)$

$$\| \|Ax\| - \|x\| \| \leq \theta. \quad (\text{A.3})$$

Fix $\theta := \delta^2/32R$ and $\omega := 8/9$, and denote the event of the above display as E . We can choose C in (1.3) such that $k \in \mathbb{Z}_{\geq 2}$ and $\mathbb{P}(E) \geq 8/9$. Note, C depends on ω and θ , which depends on R and δ .

Relating to Case (i) Let $H(z) := u^T z + t/\sqrt{k}$. For every $1 \leq j \leq N$, let $y_j := Ax_j \in \mathbb{R}^k$ and $\sigma(y_j) := \sigma(x_j)$. Note that for all z , $h(z) = \sqrt{k}H(Az)$, so B_i holds if and only if $H(y_i) \geq \gamma/\sqrt{k}$ and $H(y_j) \leq -\gamma/\sqrt{k}$ for all $j > i$ and $\sigma(y_i) \neq \sigma(y_j)$. Let B'_i be B_i conditioned on E . Note

$$\mathbb{P}(B_i) \geq \mathbb{P}(B_i | E) \cdot \mathbb{P}(E) \geq \frac{8}{9} \cdot \mathbb{P}(B_i | E)$$

For the rest of the proof, we lower bound $\mathbb{P}(B_i | E)$, i.e. we bound the probability that $H(y_i) \geq \gamma/\sqrt{k}$ and $H(y_j) \leq -\gamma/\sqrt{k}$ for all $j > i$ and $\sigma(y_i) \neq \sigma(y_j)$ given that the event E occurs. By the ordering of the points by their norms, we first notice that for all $j > i$,

$$\|x_i\| + \theta \geq \|y_i\| = \|Ax_i\| \geq \|x_i\| - \theta \geq \|x_j\| - \theta \geq \|Ax_j\| - 2\theta = \|y_j\| - 2\theta \quad (\text{A.4})$$

On the other hand, since AX^+ and AX^- are $(\delta - \theta)$ -separable, so $\|y_i - y_j\| \geq \delta - \theta$ whenever $\sigma(y_i) \neq \sigma(y_j)$. These conditions are similar to the δ -separation assumption and ordering of x_i in Case (i). We show Case (iii) in two sub-cases Case (iiia) and Case (iiib) according to whether $\|x_i\| \geq \delta/2$ or $\|x_i\| \leq \delta/2$.

Case (iiia) In this case, assume $\ell := \|x_i\| \geq \delta/2$. We follow Case (i) applied to the y_i 's. On the event E defined in (A.3), for every y_j with $j > i$ and $\sigma(y_j) \neq \sigma(y_i)$, by (A.4) and $(\delta - \theta)$ -separation

$$(\delta - \theta)^2 \leq (y_i - y_j)^T (y_i - y_j) = \|y_i\|^2 + \|y_j\|^2 - 2y_i^T y_j \leq 2(\ell + \theta)^2 - 2y_i^T y_j$$

so we have $y_i^T y_j \leq (\ell + \theta)^2 - (\delta - \theta)^2/2$. Let $\hat{y}_i = y_i/\|y_i\|$. For every $u \in \mathbb{S}^{k-1}$, we bound $(u - \hat{y}_i)^T y_i \geq -\|u - \hat{y}_i\| \cdot \|y_i\|$ by Cauchy-Schwarz and $\ell - \theta \leq \|y_i\| \leq \ell + \theta$ from (A.4) to obtain

$$u^T y_i = \hat{y}_i^T y_i + (u - \hat{y}_i)^T y_i \geq \|y_i\| - \|u - \hat{y}_i\| \cdot \|y_i\| \geq (\ell - \theta) - (\ell + \theta)\|u - \hat{y}_i\| \quad (\text{A.5})$$

Similarly, for any j where $j > i$ and $\sigma(y_j) \neq \sigma(y_i)$, we had $y_i^T y_j \leq (\ell + \theta)^2 - (\delta - \theta)^2/2$, so $\hat{y}_i^T y_j \leq (\ell + \theta)^2/\|y_i\| - (\delta - \theta)^2/2\|y_i\|$. Combined with the bound $(u - \hat{y}_i)^T y_j \leq \|u - \hat{y}_i\| \cdot \|y_j\|$ by Cauchy-Schwarz inequality, we obtain the first inequality in the following

$$\begin{aligned} u^T y_j &= \hat{y}_i^T y_j + (u - \hat{y}_i)^T y_j \\ &\leq \frac{(\ell + \theta)^2}{\|y_i\|} - \frac{(\delta - \theta)^2}{2\|y_i\|} + \|u - \hat{y}_i\| \cdot \|y_j\| \\ &\leq \frac{(\ell + \theta)^2}{\ell - \theta} - \frac{(\delta - \theta)^2}{2(\ell + \theta)} + (\ell + \theta)\|u - \hat{y}_i\| \end{aligned} \quad (\text{A.6})$$

where we use $\ell - \theta \leq \|y_i\| \leq \ell + \theta$ and $\|y_j\| \leq \ell + \theta$ from (A.4) to obtain the second inequality.

Recall that $H(z) = u^T z + t/\sqrt{k}$ and B_i occurs if and only if $u^T x_i + t/\sqrt{k} \geq \gamma/\sqrt{k}$ and $u^T x_j + t/\sqrt{k} \leq -\gamma/\sqrt{k}$ for all $j > i$ and $\sigma(y_j) \neq \sigma(y_i)$. This is equivalent to $t \geq \gamma - \sqrt{k}u^T y_i$ and $t \leq -\gamma - \sqrt{k}u^T y_j$ for all such j . Then, for a fixed u , the event B_i occurs if

$$t \in I := \left[\gamma - \sqrt{k}u^T y_i, -\gamma - \sqrt{k} \max_{j:j>i, \sigma(y_j) \neq \sigma(y_i)} u^T y_j \right].$$

We bound $|u^T y_i| \leq \|u\| \cdot \|y_i\| \leq \|x_i\| + \theta \leq R + \theta$ and similarly $|u^T y_j| \leq R + \theta$. Since $\theta = \delta^2/32R \leq (2R)^2/32R = R/8$ and $\lambda \geq 9R\sqrt{k}/8$, we have

$$I \subset [-\sqrt{k}(R + \theta), \sqrt{k}(R + \theta)] \subset [-9\sqrt{k}R/8, 9\sqrt{k}R/8] \subset [-\lambda, \lambda].$$

So we may now bound $\mathbb{P}(B_i | E, u)$ from below by the length of I over 2λ , i.e.

$$\begin{aligned} \mathbb{P}_t(B_i | E, u) &\geq \frac{\sqrt{k}}{2\lambda} \left(u^T y_i - \max_{j:j>i, \sigma(y_j) \neq \sigma(y_i)} (u^T y_j) - \frac{2\gamma}{\sqrt{k}} \right) \\ &\geq \frac{\sqrt{k}}{2\lambda} \left((\ell - \theta) - \frac{(\ell + \theta)^2}{\ell - \theta} + \frac{(\delta - \theta)^2}{2(\ell + \theta)} - 2(\ell + \theta)\|u - \hat{y}_i\| - \frac{2\gamma}{\sqrt{k}} \right) \\ &= \frac{\sqrt{k}}{\lambda} \left(\frac{(\delta - \theta)^2}{4(\ell + \theta)} - \frac{2\ell\theta}{\ell - \theta} - (\ell + \theta)\|u - \hat{y}_i\| - \frac{\gamma}{\sqrt{k}} \right) \\ &= \frac{\sqrt{k}}{\lambda} \left(\frac{(\delta - \theta)^2}{4(\ell + \theta)} - \frac{2(\delta^2/32R)}{1 - \theta/\ell} - (\ell + \theta)\|u - \hat{y}_i\| - \frac{\gamma}{\sqrt{k}} \right) \end{aligned}$$

where we plug in (A.5) and (A.6) for the second inequality and the last equality is due to $\theta = \delta^2/32R$. We know $\delta/2 \leq \ell \leq R$, so we have $\theta \leq \delta/16$. Note that the expression inside the parenthesis of the

last line of the above display decreases when we replace every θ by its upper bound $\delta/16$, replace ℓ in the first and third terms by its upper bound R , and replace ℓ in the second term by its lower bound $\delta/2$. By these substitutions, we obtain the following inequality

$$\begin{aligned}
\mathbb{P}(B_i | E, u) &\geq \frac{\sqrt{k}}{\lambda} \left(\frac{(15\delta/16)^2}{4(R + \delta/16)} - \frac{2(\delta^2/32R)}{1 - (\delta/16)/(\delta/2)} - (R + \delta/16)\|u - \hat{y}_i\| - \frac{\gamma}{\sqrt{k}} \right) \\
&\geq \frac{\sqrt{k}}{\lambda} \left(\frac{(15\delta/16)^2}{4(R + 2R/16)} - \frac{2(\delta^2/32R)}{1 - (\delta/16)/(\delta/2)} - (R + 2R/16)\|u - \hat{y}_i\| - \frac{\gamma}{\sqrt{k}} \right) \\
&= \frac{\sqrt{k}}{\lambda} \left(\frac{111}{896} \cdot \frac{\delta^2}{R} - (9R/8) \cdot \|u - \hat{y}_i\| - \frac{\gamma}{\sqrt{k}} \right) \\
&\geq \frac{\sqrt{k}}{\lambda} \left(\frac{\delta^2}{18R} - (9R/8) \cdot \|u - \hat{y}_i\| \right)
\end{aligned}$$

where the second inequality follows by replacing δ by its upper bound $2R$, and the last inequality follows $\gamma \leq \delta^2\sqrt{k}/18R$ by bounding $111/896 \geq 1/9$.

We condition on $\|u - y_i\| \leq r := (4\delta^2/81R^2)(1 - 1/k) \in [0, 2]$ to obtain the first inequality below. As $u \sim \text{Unif}(\mathbb{S}^{k-1})$, Lemma A.1(1) implies the second inequality below, i.e.

$$\begin{aligned}
\mathbb{P}(B_i | E) &\geq \mathbb{P}(B_i | E, \|u - \hat{y}_i\| \leq r) \cdot \mathbb{P}_t(\|u - \hat{y}_i\| \leq r) \\
&\geq \frac{\sqrt{k}}{\lambda} \left(\frac{\delta^2}{18R} - \frac{9Rr}{8} \right) \cdot \left(\frac{1}{2} \right) \left(\frac{r}{2} \right)^{k-1} \\
&= \frac{\sqrt{k}}{\lambda} \left(\frac{\delta^2}{18R} - \frac{9R}{8} \cdot \frac{4\delta^2}{81R^2} \left(1 - \frac{1}{k} \right) \right) \left(\frac{1}{2} \right) \left(\frac{2\delta^2}{81R^2} \left(1 - \frac{1}{k} \right) \right)^{k-1} \\
&= \frac{\sqrt{k}}{2\lambda} \left(\frac{\delta^2}{18Rk} \right) \left(\frac{2\delta^2}{81R^2} \right)^{k-1} \left(1 - \frac{1}{k} \right)^{k-1} \\
&\geq \frac{\sqrt{k}}{2\lambda} \left(\frac{9R}{4k} \right) \left(\frac{2\delta^2}{81R^2} \right)^k \left(1 - \frac{1}{k} \right)^{-1} \left(\frac{1}{4} \right) \\
&\geq \frac{9}{32} \left(\frac{R}{\lambda\sqrt{k}} \right) \left(\frac{2\delta^2}{81R^2} \right)^k
\end{aligned}$$

where the third inequality follows from the fact that $(1 - 1/k)^k$ is increasing in k for $k \geq 2$, so it can be lower bounded by its value $1/4$ at $k = 2$, and the last inequality is due to $(1 - 1/k)^{-1} \geq 1$. Hence

$$\mathbb{P}(B_i) \geq \mathbb{P}(E) \cdot \mathbb{P}(B_i | E) \geq \frac{8}{9} \cdot \frac{9}{32} \left(\frac{R}{\lambda\sqrt{k}} \right) \left(\frac{2\delta^2}{81R^2} \right)^k = \frac{R}{4\lambda\sqrt{k}} \left(\frac{2\delta^2}{81R^2} \right)^k =: q$$

establishing Case (iiia), as desired.

Case (iiib) In this case, we have $\ell := \|x_i\| \leq \delta/2$. By the Triangle Inequality, $\mathbb{B}_2^d(x_i, \delta) \supset (\delta/2)\mathbb{B}_2^d \supset \|x_i\|\mathbb{B}_2^d$, so there cannot be any x_j with smaller norm in the other set as x_i . Then, $B_i = \{h(x_i) \geq \gamma\}$, which intuitively happens with much higher probability than q . Continuing with the same notation as in Case (iiia), we have that conditioned on the event E (recall from (A.3))

$$B_i \iff h(x_i) \geq \gamma \iff H(y_i) = u^T y_i + t/\sqrt{k} \geq \gamma/\sqrt{k} \iff t \geq \gamma - \sqrt{k}u^T y_i$$

Since $u \sim \text{Unif}(\mathbb{S}^{k-1})$, for any $y_i \in \mathbb{R}^k$, we see $\mathbb{P}(u^T y_i \geq 0) = 1/2$. With this, we get

$$\mathbb{P}(B_i | E) \geq \mathbb{P}\left(t \geq \gamma - \sqrt{k}u^T y_i \mid E, u^T y_i \geq 0\right) \cdot \mathbb{P}(u^T y_i \geq 0) \geq \frac{1}{2} \cdot \mathbb{P}(t \geq \gamma \mid E) \quad (\text{A.7})$$

Since the event E is independent of the random variable t , $\mathbb{P}(t \geq \gamma \mid E) = \mathbb{P}(t \geq \gamma)$. Since $t \sim \text{Unif}([-\lambda, \lambda])$, $\lambda \geq 9R\sqrt{k}/8$, and $\gamma \leq \delta^2/18R\sqrt{k}$, we obtain the first inequality below

$$\mathbb{P}(t \geq \gamma) = \frac{\lambda - \gamma}{2\lambda} \geq \frac{1}{2} - \frac{\delta^2/18R\sqrt{k}}{2(9R\sqrt{k})/8} = \frac{1}{2} - \frac{2\delta^2}{81kR^2} \geq \frac{1}{2} - \frac{2(2R)^2}{162R^2} > \frac{1}{3}$$

where the second inequality follows from $k \geq 2$ and $\delta \leq 2R$. Plugging this back into (A.7) gives $\mathbb{P}(B_i | E) \geq 1/6$. Finally, as before, we condition on E to bound

$$\mathbb{P}(B_i) \geq \mathbb{P}(B_i | E) \cdot \mathbb{P}(E) \geq \frac{1}{6} \cdot \frac{8}{9} = \frac{4}{27}$$

On the other hand, using $\lambda \geq 9R\sqrt{k}/8 \geq R\sqrt{k}$ and $\delta \leq 2R$, we obtain the first inequality below

$$q = \frac{R}{4\lambda\sqrt{k}} \left(\frac{2\delta^2}{81R^2} \right)^k \leq \frac{R}{4Rk} \left(\frac{2(2R)^2}{81R^2} \right)^2 \leq \frac{1}{8} \left(\frac{8}{81} \right)^2 \leq \frac{4}{27} \leq \mathbb{P}(B_i)$$

where the second inequality follows since $k \geq 2$. This proves Case (iiib), establishing Theorem 2.2. \square

A.4 Proof of Theorem 2.3

For every $i, j \in \{1, 2, \dots, N\}$, define x_i and h_j as in Algorithm 1, which exists under its input conditions by Theorem 2.2. By definition of h_j , we have that

$$h_j(x_i) = \begin{cases} \leq -\gamma & \text{if } j < i \text{ and } \sigma(x_j) \neq \sigma(x_i) \\ \geq \gamma & \text{if } j = i \\ * & \text{otherwise} \end{cases} \quad (\text{A.8})$$

where $*$ denotes unknown reals. Hence, after activation,

$$(\Phi(x_i))_j = \text{ReLU}(h_j(x_i)) = \begin{cases} 0 & \text{if } j < i \text{ and } \sigma(x_j) \neq \sigma(x_i) \\ \geq \gamma & \text{if } j = i \\ \geq 0 & \text{otherwise} \end{cases} \quad (\text{A.9})$$

We choose the weight vector $a \in \mathbb{R}^N$ by picking a_i recursively for $i = N, N-1, \dots, 1$ as follows:

$$a_i = \frac{\sigma(x_i)}{\gamma} \left(1 + \left| \sum_{j=i+1}^N a_j (\Phi(x_i))_j \right| \right) \quad (\text{A.10})$$

Note that a_i has same sign as $\sigma(x_i)$. Let $H(z) = a^T z$ be an affine function associated with the hyperplane $\{z : H(z) = 0\}$. We first claim and prove that H separates the points of $\Phi(X^+)$ and $\Phi(X^-)$, and then derive the lower bound on the margin of separation.

Claim A.4. H separates $\Phi(X^+)$ and $\Phi(X^-)$ with margin at least $1/\|a\|$.

Proof of Claim A.4. It suffices to show $H(\Phi(x_i)) \geq 1$ if $x_i \in X^+$ and $H(\Phi(x_i)) \leq -1$ if $x_i \in X^-$. For every $1 \leq i \leq N$, we have

$$\begin{aligned} \sigma(x_i)H(\Phi(x_i)) &= \sigma(x_i) \sum_{j=1}^N a_j (\Phi(x_i))_j \\ &= \sigma(x_i) \sum_{j < i, \sigma(x_j) = \sigma(x_i)} a_j (\Phi(x_i))_j + \sigma(x_i) \sum_{j < i, \sigma(x_j) \neq \sigma(x_i)} a_j (\Phi(x_i))_j \\ &\quad + \sigma(x_i) a_i (\Phi(x_i))_i + \sigma(x_i) \sum_{j > i} a_j (\Phi(x_i))_j \end{aligned} \quad (\text{A.11})$$

where the second equality is obtained by splitting the summation into four groups depending on whether $j < i$, $j = i$, or $j > i$, and depending on whether $\sigma(x_j) = \sigma(x_i)$. We consider each quantity.

For each term in the first quantity, a_j has the same sign as $\sigma(x_j) = \sigma(x_i)$, so we get

$$\sigma(x_i) \sum_{j < i, \sigma(x_j) = \sigma(x_i)} a_j (\Phi(x_i))_j = \sum_{j < i, \sigma(x_j) = \sigma(x_i)} \sigma(x_j) a_j (\Phi(x_i))_j \geq 0. \quad (\text{A.12})$$

For each term in the first quantity, $(\Phi(x_i))_j = 0$ if $j < i$ and $\sigma(x_j) \neq \sigma(x_i)$, we get

$$\sigma(x_i) \sum_{j < i, \sigma(x_j) \neq \sigma(x_i)} a_j(\Phi(x_i))_j = 0. \quad (\text{A.13})$$

For the third quantity, from (A.9) we have $(\Phi(x_i))_i \geq \gamma$, and from (A.10) we get

$$\sigma(x_i) a_i(\Phi(x_i))_i \geq \frac{\sigma(x_i)^2}{\gamma} \left(1 + \left| \sum_{j > i} a_j(\Phi(x_i))_j \right| \right) \gamma = 1 + \left| \sum_{j > i} a_j(\Phi(x_i))_j \right| \quad (\text{A.14})$$

Thus, plugging (A.12), (A.13), and (A.14) into (A.11) gives the first inequality below

$$\sigma(x_i) H(\Phi(x_i)) \geq 0 + 0 + 1 + \left| \sum_{j > i} a_j(\Phi(x_i))_j \right| + \sigma(x_i) \sum_{j > i} a_j(\Phi(x_i))_j \geq 1,$$

where the second inequality follows by the triangle inequality. This shows $H(\Phi(x_i)) \geq 1$ if $x_i \in X^+$ and $H(\Phi(x_i)) \leq -1$ if $x_i \in X^-$ and hence, proves the claim. \square

We now further lower bound the margin of separation $1/\|a\|$.

Claim A.5. For a defined in (A.10) and \mathcal{M} defined in (1.1), $1/\|a\| \geq \sqrt{N} \mathcal{M}(\gamma, N)$.

Proof of Claim A.5. We first show that $(\Phi(x_i))_j \leq 2R$ for all $j > i$ such that $\sigma(x_i) \neq \sigma(x_j)$. We start by showing for all i , $h_i(x) = 0$ for some $x \in R\mathbb{B}_2^d$. This just follows from the observations that each of our hyperplanes $\{h_i = 0\}$ intersects $R\mathbb{B}_2^d$ by assumption. More rigorously, consider the two cases on line 5 of Algorithm 1, either way we have $h_i(x_i) > 0$ and $h_i(x_j) \leq 0$ for some $x_i, x_j \in R\mathbb{B}_2^d$. Then, by the intermediate value theorem, there is some x on the segment $x_i x_j$ such that $h_i(x) = 0$. Since $R\mathbb{B}_2^d$ is convex, $x \in R\mathbb{B}_2^d$. Then, for every $y \in R\mathbb{B}_2^d$, since $w_i \in \mathbb{S}^{d-1}$ we have

$$|h_i(y)| \leq |h_i(x)| + |h_i(y) - h_i(x)| \leq |w_i^T(y - x)| \leq \|w_i\| \cdot \|y - x\| \leq 1 \cdot (R + R) = 2R. \quad (\text{A.15})$$

Here, we used the Triangle Inequality to obtain the first inequality, used that $h_i(x) = 0$ to get the second inequality, applied the Cauchy-Schwarz Inequality in the third inequality, and used the fact that $x, y \in R\mathbb{B}_2^d$ to get the final inequality. By using the upper bound in (A.15),

$$|a_i| = \frac{1}{\gamma} \left(1 + \left| \sum_{j=i+1}^N a_j(\Phi(x_i))_j \right| \right) \leq \frac{1}{\gamma} \left(1 + 2R \sum_{j=i+1}^N |a_j| \right),$$

where we used the triangle Inequality to obtain the last inequality. Thus, $|a_i| \leq \tilde{a}_i$ where \tilde{a}_i is a sequence defined recursively by $\tilde{a}_N = 1/\gamma = |a_N|$ and

$$\tilde{a}_i = \frac{1}{\gamma} \left(1 + 2R \sum_{j=i+1}^N \tilde{a}_j \right).$$

We compute \tilde{a}_i explicitly. Taking the successive differences

$$\tilde{a}_{i-1} - \tilde{a}_i = \frac{1}{\gamma} \left(1 + 2R \sum_{j=i}^N \tilde{a}_j \right) - \frac{1}{\gamma} \left(1 + 2R \sum_{j=i+1}^N \tilde{a}_j \right) = \frac{2R\tilde{a}_i}{\gamma},$$

and using the fact that $\tilde{a}_N = 1/\gamma$, we get

$$\tilde{a}_i = \frac{1}{\gamma} \left(1 + \frac{2R}{\gamma}\right)^{N-i}.$$

Hence,

$$\|a\|^2 \leq \sum_{i=1}^N \frac{1}{\gamma^2} \left(1 + \frac{2R}{\gamma}\right)^{2(N-i)} = \frac{1}{\gamma^2} \cdot \frac{(1 + 2R/\gamma)^{2N} - 1}{(1 + 2R/\gamma)^2 - 1} = \frac{(1 + 2R/\gamma)^{2N} - 1}{4R(R + \gamma)}.$$

which gives the desired bound by rearranging. \square

Therefore, Claim A.4 and Claim A.5 together implies that H separates $\Phi(X^+)$ and $\Phi(X^-)$ by a margin at least $\sqrt{N} \mathcal{M}(\gamma, N)$, as desired.

A.5 Proof of Theorem 2.4

Consider Definition 1.2 for the fact that X^+ and X^- have $(\delta, 8R^2/\gamma)$ -mutual complexity N . Since $C^+, C^- \subset R\mathbb{B}_2^d$ are δ -separated, for any center c_i^\pm

$$\delta \leq \text{dist}(c_i^\pm, C^\mp) \leq 2R$$

Then, for any radius r_i^\pm , we have that

$$r_i^\pm \leq \frac{\gamma}{8R^2} \text{dist}(c_i^\pm, C^\mp)^2 \leq \frac{\gamma}{8R^2} (2R)^2 = \frac{\gamma}{2} \quad (\text{A.16})$$

We re-order the points in $C^+ \cup C^-$ as in Algorithm 1, i.e., for every i , let c_i be the point in $C^+ \cup C^-$ with the i -th largest norm. Similarly, let r_i now be the radius associated to c_i . Hence, (A.16) implies $r_i \leq \gamma/2$ for all i . Also, let $\sigma(c_i) = +1$ if $c_i \in C^+$ and $\sigma(c_i) = -1$ if $c_i \in C^-$ as in (2.1). Hence, the first condition of Definition 1.2 says that $c_i \in \text{conv}(X_i)$ where $X_i := X^{\sigma(c_i)} \cap \mathbb{B}_2^d(c_i, r_i)$.

Part (i) Here we show that if the input conditions of Algorithm 1 are satisfied for X^+ and X^- , then those must be satisfied for the sets C^+ and C^- . For convenience, let k_X be k in (1.3) for X^+ and X^- and let k_C be k in (1.3) where we replace X^+ and X^- by C^+ and C^- . We check the following:

- The dimension d doesn't change since $C^+, C^- \subset \mathbb{R}^d$, so d is preserved.
- By Definition 1.2, C^+ and C^- are δ -separated finite sets, so δ is preserved.
- By Definition 1.2, $X_i \neq \emptyset$ and $c_i \in \text{conv}(X_i)$ for every i . Hence, Lemma A.1(2) implies $\|c_i\| \leq \|x\| \leq R$ for some $x \in X_i$. Thus, $C^+, C^- \subset R\mathbb{B}_2^d$, which means R is preserved.

Finally, we check γ satisfies the input condition now on C^+ and C^- if it satisfies that for X^+ and X^- . Since $c_i \in \text{conv}(X_i)$ for every i , Lemma A.1(2) implies for every vector g and indices i with $\sigma(c_i) = +1$ and $\sigma(c_j) = -1$, there exists $x_i \in X_i$ and $x_j \in X_j$ such that $g^T x_i \geq g^T c_i$ and $(-g)^T x_j \geq (-g)^T c_j$. Hence

$$w(C^+) = \mathbb{E}_g \sup\{g^T(c_i) : \sigma(c_i) = +1\} \leq \mathbb{E}_g \sup\{g^T(x_i)\} \leq \mathbb{E}_g \sup\{g^T x : x \in X^+\} = w(X^+)$$

and similarly $w(C^-) \leq w(X^-)$. Moreover, by a similar calculation

$$\begin{aligned} w(C^+ - C^-) &= \mathbb{E}_g \sup\{g^T c : c \in C^+ - C^-\} \\ &= \mathbb{E}_g \sup\{g^T(c_i - c_j) : \sigma(c_i) = +1, \sigma(c_j) = -1\} \\ &\leq \mathbb{E}_g \sup\{g^T(x_i - x_j) : x_i \in X_i, x_j \in X_j\} \\ &\leq \mathbb{E}_g \sup\{g^T x : x \in X^+ - X^-\} \\ &= w(X^+ - X^-) \end{aligned}$$

Since we showed R, δ, d are preserved, this implies $k_X \geq k_C$, we obtain that

$$\gamma \leq \max(\delta^2/8Rd, \delta^2/18R\sqrt{k_X}) \leq \max(\delta^2/8Rd, \delta^2/18R\sqrt{k_C})$$

Therefore, as γ satisfies its input condition for X^+ and X^- , it does for C^+ and C^- .

Part (ii) Here we show that the output of Algorithm 1 on C^+ and C^- linearly separates the points from X^+ and X^- with a margin $\sqrt{N}\mathcal{M}(\gamma/2, N)$. By Part (i), we can run Algorithm 1 on C^+ and C^- with parameter γ to define h_j for each $1 \leq j \leq N$ as in line 6. By definition of h_j , we obtain the exact same statement as (A.8) but for C^+ and C^- , i.e.

$$h_j(c_i) = \begin{cases} \leq -\gamma & \text{if } j < i \text{ and } \sigma(c_j) \neq \sigma(c_i) \\ \geq \gamma & \text{if } j = i \\ * & \text{otherwise} \end{cases} \quad (\text{A.17})$$

where $*$ denotes unknown reals. As $r_i \leq \gamma/2$ by (A.16), for every i, j and $x_i \in \mathbb{B}_2^d(c_i, r_i)$, we have

$$|h_j(x_i) - h_j(c_i)| = |w_j^T(x_i - c_i)| \leq \|x_i - c_i\| \leq r_i \leq \frac{\gamma}{2}.$$

By the triangle inequality on (A.17), we obtain an analog of (A.17) that for every $x_i \in X_i$

$$h_j(x_i) = \begin{cases} \leq -\gamma/2 & \text{if } j < i \text{ and } \sigma(x_j) \neq \sigma(x_i) \\ \geq \gamma/2 & \text{if } j = i \\ * & \text{otherwise} \end{cases} \quad (\text{A.18})$$

Hence, after activation, for every $x_i \in X_i$

$$(\Phi(x_i))_j = \text{ReLU}(h_j(x_i)) = \begin{cases} 0 & \text{if } j < i \text{ and } \sigma(x_j) \neq \sigma(x_i) \\ \geq \gamma/2 & \text{if } j = i \\ \geq 0 & \text{otherwise} \end{cases} \quad (\text{A.19})$$

We now proceed exactly as the proof of Theorem 2.4, except since x_i can be any point in cluster X_i , we use supremum over X_i instead of just x_i . Then, the proof will show the clusters X_i are linearly separable according to $\sigma(x_i)$, which suffices since the clusters partition $X^+ \cup X^-$.

We choose the weight vector $a \in \mathbb{R}^N$ by picking a_i recursively for $i = N, N-1, \dots, 1$ as follows:

$$a_i = \frac{2\sigma(c_i)}{\gamma} \sup_{x \in X_i} \left(1 + \left| \sum_{j=i+1}^N a_j (\Phi(x_i))_j \right| \right)$$

and so for hyperplane or affine function $H(z) = a^T z$, every $x_i \in X_i$, and every $1 \leq i \leq N$,

$$\begin{aligned} \sigma(x_i)H(\Phi(x_i)) &= \sigma(x_i) \sum_{j=1}^N a_j (\Phi(x_i))_j \\ &= \sigma(x_i) \sum_{j < i, \sigma(x_j) = \sigma(x_i)} a_j (\Phi(x_i))_j + \sigma(x_i) \sum_{j < i, \sigma(x_j) \neq \sigma(x_i)} a_j (\Phi(x_i))_j \\ &\quad + \sigma(x_i) a_i (\Phi(x_i))_i + \sigma(x_i) \sum_{j > i} a_j (\Phi(x_i))_j \end{aligned}$$

by breaking into four cases similar to the proof of Theorem 2.3. The first term is non-negative since a_j has the same sign as $\sigma(x_j) = \sigma(x_i)$ and coordinates of $\Phi(x_i)$ are non-negative due to ReLU-activation. The second term is zero since $(\Phi(x_i))_j = 0$ for such j by (A.19). Hence

$$\begin{aligned} \sigma(x_i)H(\Phi(x_i)) &\geq 0 + 0 + \frac{2\sigma(x_i)^2}{\gamma} \sup_{x \in X_i} \left(1 + \left| \sum_{j > i} a_j (\Phi(x_i))_j \right| \right) \left(\frac{\gamma}{2} \right) + \sigma(x_i) \sum_{j > i} a_j (\Phi(x_i))_j \\ &= \sup_{x \in X_i} \left(1 + \left| \sum_{j > i} a_j (\Phi(x_i))_j \right| \right) + \sigma(x_i) \sum_{j > i} a_j (\Phi(x_i))_j \\ &\geq 1. \end{aligned}$$

by the Triangle Inequality. Thus, $H(\Phi(x_i)) \geq 1$ if $x_i \in X^+$ and $H(\Phi(x_i)) \leq -1$ if $x_i \in X^-$, so $H(z) = a^T z$ separates $\Phi(X^+)$ and $\Phi(X^-)$. The margin of separation is at least $1/\|a\|$. We bound this as follows. Exactly as (A.15), $|h_i(y)| \leq 2R$ for every $y \in R\mathbb{B}_2^d$. Now, we bound each weight a_i

$$|a_i| = \frac{2}{\gamma} \sup_{x \in X_i} \left(1 + \left| \sum_{j=i+1}^N a_j (\Phi(x_i))_j \right| \right) \leq \frac{2}{\gamma} \left(1 + 2R \sum_{j=i+1}^N |a_j| \right)$$

Thus, $|a_i| \leq \tilde{a}_i$ where \tilde{a}_i is a sequence defined recursively by $\tilde{a}_N = 2/\gamma = |a_N|$ and

$$\tilde{a}_i = \frac{2}{\gamma} \left(1 + 2R \sum_{j=i+1}^N \tilde{a}_j \right).$$

We compute \tilde{a}_i explicitly. Taking the successive differences

$$\tilde{a}_{i-1} - \tilde{a}_i = \frac{2}{\gamma} \left(1 + 2R \sum_{j=i}^N \tilde{a}_j \right) - \frac{2}{\gamma} \left(1 + 2R \sum_{j=i+1}^N \tilde{a}_j \right) = \left(\frac{4R}{\gamma} \right) \tilde{a}_i,$$

and using the fact that $\tilde{a}_N = 2/\gamma$, we get

$$\tilde{a}_i = \frac{2}{\gamma} \left(1 + \frac{4R}{\gamma}\right)^{N-i}.$$

Hence,

$$\|a\|^2 \leq \sum_{i=1}^N \frac{4}{\gamma^2} \left(1 + \frac{4R}{\gamma}\right)^{2(N-i)} = \frac{4}{\gamma^2} \cdot \frac{(1 + 4R/\gamma)^{2N} - 1}{(1 + 4R/\gamma)^2 - 1} = \frac{(1 + 2R/(\gamma/2))^{2N} - 1}{4R(R + \gamma/2)}.$$

Since the margin is at least $1/\|a\|$, we obtain the desired margin bound $\sqrt{N}\mathcal{M}(\gamma/2, N)$.

A.6 Proof of Theorem 1.3 and Theorem 1.4

We modify the notation slightly for convenience of proving both finite set cases and arbitrary set cases together. Fix γ to be either $\delta^2/8Rd$ or $\delta^2/18R\sqrt{k}$ depending on the distribution of Φ , as in the finite set cases. For the arbitrary set cases, the parameter will be $\gamma/2$. We show in all three cases that provided the respective width bounds are met, with probability $1 - \eta$, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable. If $|X^+| + |X^-| = N < \infty$, then $\Phi(X^+)$ and $\Phi(X^-)$ will be linearly separable with margin at least $\mathcal{M}(\gamma, N)$. Otherwise, in the case of arbitrary sets, $\Phi(X^+)$ and $\Phi(X^-)$ will be linearly separable with margin at least $\mathcal{M}(\gamma/2, N)$ where N is now the $(\delta, 8R^2/\gamma)$ -mutual complexity. We can check it is so for $\gamma \in \{\delta^2/8Rd, \delta^2/18R\sqrt{k}\}$.

For finite sets X^+ and X^- , recall x_i and σ from Algorithm 1 and (2.1). For arbitrary sets X^+ and X^- , let C^+ and C^- be the finite sets of centers guaranteed by Definition 1.2 for $(\delta, 8R^2/\gamma)$ -mutual complexity. Let c_i be the point in $C^+ \cup C^-$ with the i -th largest norm and let $\sigma : C^+ \cup C^- \rightarrow \{-1, +1\}$ such that $\sigma(c) = +1$ if $c \in C^+$ and $\sigma(c) = -1$ if $c \in C^-$.

For a random layer $\Phi(x) = \text{ReLU}(Wx + b)$ with width n , we let B_i^j for $1 \leq i \leq N$ and $1 \leq \ell \leq n$ be the event that the ℓ -th random hyperplane $H_\ell(z) = W_\ell^T z + b_\ell$ of Φ satisfies the condition for h_i in Algorithm 1 applied to X^+ and X^- when they are finite or on C^+ and C^- otherwise, and parameter γ . More precisely, if X^+ and X^- are finite sets, let

$$B_i^\ell(X) := \begin{cases} \{H_\ell(x_i) \geq \gamma\} \cap \bigcap_{j:j>i, \sigma(x_i) \neq \sigma(x_j)} \{H_\ell(x_j) \leq -\gamma\} & \text{if such } j \text{ exists} \\ \{H_\ell(x_i) \geq \gamma\} \cap \{\exists x \in R\mathbb{B}_2^d : H_\ell(x) \leq 0\} & \text{otherwise} \end{cases}. \quad (\text{A.20})$$

For arbitrary sets X^+ and X^- , analogously define

$$B_i^\ell(C) := \begin{cases} \{H_\ell(c_i) \geq \gamma\} \cap \bigcap_{j:j>i, \sigma(c_i) \neq \sigma(c_j)} \{H_\ell(c_j) \leq -\gamma\} & \text{if such } j \text{ exists} \\ \{H_\ell(c_i) \geq \gamma\} \cap \{\exists c \in R\mathbb{B}_2^d : H_\ell(c) \leq 0\} & \text{otherwise} \end{cases}. \quad (\text{A.21})$$

with γ chosen to be either $\delta^2/8Rd$ or $\delta^2/18R\sqrt{k}$ depending on the distribution of Φ . This is similar to B_i in Theorem 2.2, except we add the case where j does not exist on line 5 of Algorithm 1. We now resolve the duplication issue mentioned in the proof sketch.

Claim A.6. Fix γ to be $\delta^2/8Rd$ or $\delta^2/18R\sqrt{k}$. Recall \mathcal{M} from (1.1). Then, the following holds:

- (1) Suppose that X^+ and X^- are finite and that for every i , $B_i^\ell(X)$ with parameter γ holds for some ℓ . Then, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable with margin $\mathcal{M}(\gamma, N)$.
- (2) For arbitrary sets X^+ and X^- , suppose that for every i , $B_i^\ell(C)$ with parameter γ holds for some ℓ . Then, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable with margin $\mathcal{M}(\gamma/2, N)$.

Proof of Claim A.6. We break this proof down into four steps which we outline here. First, we set up some notation to clarify why we cannot directly apply Algorithm 2. Then, we use Theorem 2.3 and Theorem 2.4 to show that if we allow a single random hyperplane $H_\ell(z)$ to satisfy the condition for two hyperplanes h_{i_1} and h_{i_2} , it makes the sets separable. Next, we show how to remove one of such hyperplanes while still making the sets separable. Finally, we use Lemma 2.1 to show Φ also makes the sets separable. The separation margin will also be tracked throughout.

SETUP: By assumption, we can define some map $f : [N] \rightarrow [n]$ of the indices by letting $f(i) = \ell$ if B_i^ℓ holds, where B_i^ℓ is either $B_i^\ell(X)$ or $B_i^\ell(C)$ depending on whether X^+ and X^- are finite. For a given i , if there are multiple ℓ such that B_i^ℓ holds, define $f(i)$ to be any such ℓ . Define the range $\Sigma := f([N]) \subset [n]$ and let $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^N$ be the projection onto the coordinates indexed by Σ . When f is not injective, we cannot apply Theorem 2.3 directly to $\{H_j : j \in \Sigma\}$ since there are less than N hyperplanes. Let $h_i := H_{f(i)}$ with another *duplicated* hyperplane $h_{i'}$, i.e. h_i and $h_{i'}$ are two copies of H_j if $f(i) = f(i') = j$. Then, h_i satisfies the condition on line 5 of Algorithm 1 for this i . Consider the one-layer neural network $\Psi(x) = \text{ReLU}(h_1(x), \dots, h_N(x))$.

SHOWING Ψ SEPARATES: To show Φ makes X^+ and X^- separable, we first need to show Ψ does. For Claim A.6(1), we first note that Ψ is a valid output of Algorithm 1 on X^+, X^- , and the chosen γ . Then, by Theorem 2.3, we can run Algorithm 2 to find vector $a \in \mathbb{R}^N$ such that the hyperplane $H(z) := a^T z$ separates $\Psi(X^+)$ and $\Psi(X^-)$ with the margin at least $\sqrt{N}\mathcal{M}(\gamma, N)$, i.e.

$$H(\Psi(x)) = \sum_{i=1}^N a_i \text{ReLU}(h_i(x)) = \begin{cases} \geq \|a\| \sqrt{N}\mathcal{M}(\gamma, N) & \text{if } x \in X^+ \\ \leq -\|a\| \sqrt{N}\mathcal{M}(\gamma, N) & \text{if } x \in X^- \end{cases} \quad (\text{A.22})$$

For Claim A.6(2), we first note that Ψ is a valid output of Algorithm 1 on C^+, C^- , and the chosen γ . By Theorem 2.4, $\Psi(X^+)$ and $\Psi(X^-)$ are linearly separable with margin at least $\sqrt{N}\mathcal{M}(\gamma/2, N)$. In fact, by the proof of Theorem 2.4, the separating hyperplane is of the form $H(z) = a^T z$, i.e.

$$H(\Psi(x)) = \sum_{i=1}^N a_i \text{ReLU}(h_i(x)) = \begin{cases} \geq \|a\| \sqrt{N}\mathcal{M}(\gamma/2, N) & \text{if } x \in X^+ \\ \leq -\|a\| \sqrt{N}\mathcal{M}(\gamma/2, N) & \text{if } x \in X^- \end{cases} \quad (\text{A.23})$$

SHOWING $\pi \circ \Phi$ SEPARATES: Define real vector $\tilde{a} \in \mathbb{R}^\Sigma$ as follows: for all $j \in \Sigma$

$$\tilde{a}_j := \sum_{i: f(i)=j} a_i.$$

In words, \tilde{a}_j is the sum of the weights a_i over all i where h_i is a duplicate of H_j . Now, we show that the hyperplane $\tilde{H}(z) := \tilde{a}^T z$ separates $\pi(\Phi(X^+))$ and $\pi(\Phi(X^-))$ with margin at least $\mathcal{M}(\gamma, N)$. Note, for $j \in \Sigma$, $(\pi(\Phi(x)))_j = (\Phi(x))_j = \text{ReLU}(H_j(x))$, and $H_j = h_i$ if $f(i) = j$. Hence

$$\tilde{H}(\pi(\Phi(x))) = \tilde{a}^T \pi(\Phi(x)) = \sum_{j \in \Sigma} \tilde{a}_j (\pi(\Phi(x)))_j = \sum_{j \in \Sigma} \left(\sum_{i: f(i)=j} a_i \right) \text{ReLU}(H_j(x))$$

by the definition of \tilde{a}_j . Since $H_j = h_i$ if $f(i) = j$, we get

$$\tilde{H}(\pi(\Phi(x))) = \sum_{j \in \Sigma} \sum_{i: f(i)=j} a_i \text{ReLU}(h_i(x)) = \sum_{i=1}^N a_i \text{ReLU}(h_i(x)) = H(\Psi(x))$$

by (A.23). By (A.22) and (A.23), $(\pi \circ \Phi)(X^+)$ and $(\pi \circ \Phi)(X^-)$ are linearly separable with margin at least $\|a\| \sqrt{N}\mathcal{M}(\gamma, N) / \|\tilde{a}\|$ if X^+ and X^- are finite, and margin at least $\|a\| \sqrt{N}\mathcal{M}(\gamma/2, N) / \|\tilde{a}\|$ otherwise. For each j , by Cauchy-Schwarz on each vector $(a_i)_i$ indexed by $i \in f^{-1}(j) \subset [N]$

$$\sum_{j \in \Sigma} \tilde{a}_j^2 = \sum_{j \in \Sigma} \left(\sum_{i: f(i)=j} a_i \right)^2 \leq \sum_{j \in \Sigma} |f^{-1}(j)| \sum_{i: f(i)=j} a_i^2 \leq N \sum_{j \in \Sigma} \sum_{i: f(i)=j} a_i^2 = N \sum_{i=1}^N a_i^2,$$

where for a given j , $|f^{-1}(j)|$ is the number of indices i such that $f(i) = j$. From this inequality, we get $\|\tilde{a}\| \leq \sqrt{N}\|a\|$. Thus, the separation margins are at least $\mathcal{M}(\gamma, N)$ and $\mathcal{M}(\gamma/2, N)$, respectively.

SHOWING Φ SEPARATES: By Lemma 2.1, $\Phi(X^+)$ and $\Phi(X^-)$ are linearly separable with margin at least $\mathcal{M}(\gamma, N)$ or $\mathcal{M}(\gamma/2, N)$, depending on if X^+ and X^- are finite. This proves Claim A.6. \square

Now, to show Theorem 1.3 and Theorem 1.4, it suffices to bound the probability that for every i , $B_i^\ell(X)$ and $B_i^\ell(C)$ occurs for some ℓ . We do so in the following claim.

Claim A.7. For any δ -separated sets $X^+, X^- \subset R\mathbb{B}_2^d$ with $N = |X^+| + |X^-| < \infty$, define Φ as in the three cases in Theorem 1.3 and 1.4, and define $B_i^\ell(X)$ as in (A.20). Then, $\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X)) \geq 1 - \eta$ provided the width bounds on n from Theorem 1.3 and 1.4 are met in each case.

Proof of Claim A.7. We break this proof down into two steps which we outline here. First, we will obtain bounds of $\mathbb{P}(B_i^\ell(X))$ similar to Theorem 2.2, except we include the case where no such j exists as in (A.20) and (A.21). Then, we use these bounds to obtain a final bound on $\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X))$.

BOUNDING EACH $\mathbb{P}(B_i^\ell(X))$: If j exists in (A.20), Theorem 2.2 gives bounds p , $p/10$, and q of $\mathbb{P}(B_i^\ell(X))$ for any i and ℓ , depending on Φ as in Theorem 1.3 and Theorem 1.4. We show the same result for the easier case where no such j exists by reducing it to Theorem 2.2. If no such j exists

$$B_i^\ell(X) = \{H_\ell(x_i) \geq \gamma\} \cap \{\exists x \in R\mathbb{B}_2^d : H_\ell(x) \leq 0\}.$$

Since $\delta \leq 2R$, we can pick any $x_j \in R\mathbb{B}_2^d$ such that $\|x_j - x_i\| \geq \delta$. Then

$$\mathbb{P}(\{H_\ell(x_i) \geq \gamma\} \cap \{\exists x \in R\mathbb{B}_2^d : H_\ell(x) \leq 0\}) \geq \mathbb{P}(\{H_\ell(x_i) \geq \gamma\} \cap \{H_\ell(x_j) \leq -\gamma\}).$$

If $\|x_i\| \geq \|x_j\|$, then this is precisely event B_i in Theorem 2.2 where we want the random hyperplane H_ℓ to separate x_i from the singleton set $\{x_j\}$ with parameter γ . If $\|x_i\| \leq \|x_j\|$, then this is precisely event B_j in Theorem 2.2 where we want the random hyperplane $-H_\ell$ to separate x_j from the singleton set $\{x_i\}$ with parameter γ . Then, by Theorem 2.2, $\mathbb{P}(B_i^\ell(X)) \geq p$ or $p/10$ or q for every i and ℓ , depending on the distribution of weights as in Theorem 1.3 and Theorem 1.4.

BOUNDING $\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X))$: We show Claim A.7 when the distribution of Φ is from Theorem 1.3(i). The cases for Theorem 1.3(ii) and Theorem 1.4 follow by replacing p with $p/10$ and q , respectively. For convenience, let \overline{B} be the complement of event B . By a union bound over i , we get

$$\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X)) = 1 - \mathbb{P}((\exists i)(\forall \ell)\overline{B_i^\ell(X)}) \geq 1 - \sum_{i=1}^N \mathbb{P}((\forall \ell)\overline{B_i^\ell(X)}).$$

Using the independence of $\ell \in [n]$ and the fact that $\mathbb{P}(B_i^\ell(X)) \geq p$, we get

$$\mathbb{P}((\forall \ell)\overline{B_i^\ell(X)}) \leq (1 - p)^n,$$

and thus,

$$\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X)) \geq 1 - N(1 - p)^n.$$

Note, $\log(1/(1 - p)) \geq p$ as $p \geq 0$. Then, our width bound implies

$$n \geq \frac{\log(N/\eta)}{p} \geq \frac{\log(N/\eta)}{\log(1/(1 - p))} = \log_{1-p}(\eta/N).$$

Hence, if the width bound is met, we get

$$\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(X)) \geq 1 - N(1 - p)^n \geq 1 - N(1 - p)^{\log_{1-p}(\eta/N)} = 1 - \eta$$

as desired. This concludes the proof of Claim A.7. \square

Finally, Claim A.6 and Claim A.7 together implies the finite set cases of Theorem 1.3 and Theorem 1.4. For the arbitrary set cases, apply Claim A.7 to the δ -separated finite sets $C^+, C^- \subset R\mathbb{B}_2^d$ to get that $\mathbb{P}((\forall i)(\exists \ell)B_i^\ell(C)) \geq 1 - \eta$ provided the width bounds on n in Theorem 1.3 and Theorem 1.4 are met. Then, Claim A.6 implies the arbitrary set cases of Theorem 1.3 and Theorem 1.4.