

Optimal temperature estimation in polariton Bose-Einstein Condensate

Dong Xie^{1,*}

¹College of Science, Guilin University of Aerospace Technology,
Guilin, Guangxi 541004, People's Republic of China

Improving the measurement precision of temperature is very important and challenging, especially in the low temperature range. Based on the existence of invariant subspaces during the polariton thermalization, we propose a new way to enhance the measurement precision of the low temperature and obtain Landau bound to avoid that the measurement uncertainty of the temperature diverges as the temperature approaches zero. The measurement precision of the low temperature increases significantly with the number of polariton states. In order to resist the dissipation, the incoherent pumping is necessary for obtaining the information of the temperature encoded in the steady state. It should be noted that too strong incoherent pumping is wasteful due to that the quantum Fisher information of the temperature becomes less and less dependent on the total number of the polaritons.

Introduction.-Precise estimation of temperature is significant and crucial for the fundamental natural science and the changing quantum technology[1–11]. Since quantum devices generally work at low temperature[12–14] and the development of the field of quantum thermodynamics[15–19] also need the low temperature to reserve the quantum properties, precisely control and measurement of the low temperature is becoming an important subject in quantum metrology[11] and quantum sensing.

Enhancing the estimation precision of temperature with quantum resources and investigating the fundamental limitations on temperature estimation have attracted a lot of attention[20–31]. Temperature measurements can generally be divided into two categories: one is to measure the temperature encoded in the thermal equilibrium state of the probe system, the other is in the non-equilibrium state. Either way, the measurement of the low temperature has always been a challenging and arduous task. The uncertainty of the temperature diverges as the temperature approaches zero[32]. Recently, some works attempted to alleviate the divergence. Correa *et al.*[33] showed that the thermometric precision at low temperature could be significantly enhanced by the strong probe-sampling coupling. Mukherjee *et al.*[34] utilized the periodic modulation to obtain the low-temperature thermometry with temperature-independent relative uncertainty. More importantly, Zhang *et al.*[35] obtained the Landau bound[36] by using a continuous-variable system to detect the temperature of a non-Markovian reservoir.

In this work, we propose a new way to obtain the optimal low temperature estimation, and the Landau bound can also be achieved. The exciton-polariton Bose-Einstein Condensates (BECs) are used as the thermometer to measure the temperature of the phonons in the semiconductors or the temperature of intermolecular oscillations of the organic dyes. Different with the Bose

polaron model[12] in BEC, there are invariant subspaces due to the polariton thermalization, in which the final thermal state is not unique. For the infinite states of the polaritons, the Landau bound can also be achieved at the low temperature. In addition, we find that incoherent pumping can resist the dissipation. But too strong incoherent pumping is useless in enhancing the estimation precision of the temperature due to that the temperature estimation precision will not increase with the increase of the total number of the polaritons for enough polaritons.

The physical model of polaritons.- When cavity photons strongly interact with an optical transition of active material, new eigenstates can be generated, i.e., lower and upper exciton-polariton branches[37, 38]. We only consider the lower branch in which the BEC occurs. Due to pair particle scattering, the exciton polaritons of the lower polariton branch can be treated as harmonic oscillators by neglecting the nonlinearity[39]. The Hamiltonian of the polaritons can be described as

$$H_{LP} = \sum_{j=0}^M \hbar\omega_j a_j^\dagger a_j. \quad (1)$$

where the ground state (mode) is the state with $j = 0$ and a_j (a_j^\dagger) denotes the bosonic annihilation (creation) operator. The master equation for the the density matrix of the polaritons ρ , subject only to the polariton thermalization, can be expressed as

$$\frac{d}{dt}\rho(t) = \mathcal{L}\rho = -i[H, \rho] + L_{\text{thermal}}(\rho), \quad (2)$$

where the polariton thermalization is described by the Lindblad superoperator

$$L_{\text{thermal}}(\rho) = \sum_{j=0}^M \sum_{k=0}^M \Gamma_{jk} (a_j a_k^\dagger \rho a_k a_j^\dagger - \frac{1}{2} \rho a_k a_j^\dagger a_k a_j^\dagger - \frac{1}{2} a_k a_j^\dagger a_k a_j^\dagger \rho), \quad (3)$$

where Γ_{jk} is the transition rate from the j th polariton state to the k th state. The thermalization rates

* xiedong@mail.ustc.edu.cn

obey the Kubo-Martin-Schwinger relation[40, 41] $\Gamma_{jk} = \exp[\frac{\hbar(\omega_j - \omega_k)}{\kappa_B T}] \Gamma_{kj}$, where κ_B is the Boltzmann constant and T is the temperature of intermolecular oscillations of the organic dyes or the temperature of the phonons in the semiconductors that we want to estimate. There are different underlying mechanisms of the polariton thermalization, which is dependent on the detail system. For example, the polariton thermalization comes from the nonlinear interaction with low frequency vibrations in organic polariton systems[42, 43].

The operator of the total polariton number $\sum_{j=0}^M a_j^\dagger a_j$ is the constant of motion during the thermalization process. The constant of motion implies that there are invariant subspaces $|n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|$ with the total number of polaritons equal to $\sum_{j=0}^M n_j = N$. When there are invariant subspaces, the stationary solution is not unique[44]. The Gibbs distribution over the states of a given invariant subspace is also a stationary solution, which is given by[39]

$$\rho_s = \sum_{N=0}^{\infty} P_N(0) \frac{1}{Z_N} \sum_{n_0 + \dots + n_M = N} \nu_0^{n_0} \dots \nu_M^{n_M} \times |n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|, \quad (4)$$

where $P_N(0)$ denotes the probability that there are N polaritons in total in the low polariton branch at the initial time, which is given by $P_N(0) = \sum_{n_0 + \dots + n_M = N} \text{Tr}[\rho(0) |n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|]$ with $\sum_{j=0}^M n_j = N$, $\nu_j = \exp[\frac{\hbar(\omega_0 - \omega_j)}{\kappa_B T}]$, and Z_N is the partition function discribed by

$$Z_N = \sum_{n_0 + \dots + n_M = N} \nu_0^{n_0} \dots \nu_M^{n_M}. \quad (5)$$

The estimation uncertainty of the unbiased estimator $\delta^2 T$ is bounded by the quantum Cramér-Rao lower bound as $\delta^2 T \geq 1/F_T$ [45, 46], where F_T is the quantum Fisher information (QFI)[47] of temperature T in the steady state ρ_s , which is given by

$$F_T = \sum_{N=0}^{\infty} \sum_{n_0 + \dots + n_M = N} \frac{(\partial_T P_{N, n_0, \dots, n_M})^2}{P_{N, n_0, \dots, n_M}}, \quad (6)$$

where $P_{N, n_0, \dots, n_M} = \frac{P_N(0)}{Z_N} \nu_0^{n_0} \dots \nu_M^{n_M}$ denotes the probability of projection into the state $|n_0, n_1, \dots, n_M\rangle$ with $\sum_{j=0}^M n_j = N$, and the shorthand notation $\partial_T = \frac{\partial}{\partial T}$. Hereafter, we set $\hbar = \kappa_B = 1$ for convenience.

Without loss of generality, we first consider that the system is consisted of $M + 1$ states equidistant in frequency, i.e., $\omega_j = \omega_0 + \omega \times j/M$ with $\omega = \omega_M - \omega_0$.

In the case of two polariton states.-For $M = 1$, we can obtain the probability of the state $|n_0, n_1\rangle$ with $n_0 + n_1 = N$

$$P_{N, n_0, n_1} = \langle n_0, n_1 | \rho_s(M=1) | n_0, n_1 \rangle = \frac{\lambda^{-n_1} (\lambda - 1)}{\lambda - \lambda^{-N}}, \quad (7)$$

where the factor λ is defined as $\lambda = \exp(\omega/T)$, and only one invariant subspace is considered, i.e., $P_N(0) = 1$. By utilizing the formula in Eq. (6), the QFI with $M = 1$ is analytically expressed as

$$F_T = \frac{\omega^2}{T^4} \left[\frac{1}{\lambda - 1} + \frac{1}{(\lambda - 1)^2} - \frac{(1 + N)^2}{\lambda^{1+N} - 1} - \frac{(1 + N)^2}{(\lambda^{1+N} - 1)^2} \right]. \quad (8)$$

For $N \gg 1$, we can obtain the simplified form $F_T \simeq \frac{\lambda \omega^2}{(\lambda - 1)^2 T^4}$. This result is equivalent to measuring the temperature T in the thermal equilibrium state of harmonic oscillator with the Hamiltonian $H_e = \omega a^\dagger a$ [28]. This does not reflect the advantage of having invariant subspaces. This is mainly due to the low number of states. Next, we investigate the QFI of the temperature with a large number of states, i.e., $M > 1$.

In the case of single polarization: $N = 1$.- We then consider another simple case, which can be analytically calculated for $N = 1$. In this case, the probability of projection into the state $|0, 0, \dots, n_j = 1, \dots, 0\rangle$ is $P_{N=1, n_j=1} = \frac{\lambda^{-j/M}}{\sum_{j=0}^M \lambda^{-j/M}} = \frac{\lambda^{1-j/M} (\lambda^{1/M} - 1)}{\lambda^{(1+1/M)} - 1}$. The corresponding QFI is given by

$$F_T = 2\omega^2 \lambda^{1+2/M} \times \frac{M(2 + M) - (1 + M)^2 \cosh(\frac{\omega}{MT}) + \cosh[(1 + M)\frac{\omega}{MT}]}{(\lambda^{1/M} - 1)^2 [\lambda^{(1+M)/M} - 1]^2 M^2 T^4}. \quad (9)$$

For the infinite M , the QFI of the temperature T tends to be

$$F_T|_{M \rightarrow \infty} = \frac{1}{T^2} - \frac{\lambda \omega^2}{(\lambda - 1)^2 T^4}. \quad (10)$$

For $\omega \gg T$, we can obtain $F_T|_{M \rightarrow \infty} \simeq 1/T^2$, which is the maximal QFI of the temperature. According to the quantum Cramér-Rao lower bound, we obtain the Landau bound[36], i.e., $\delta T \geq T$. It means that we can obtain the infinite QFI for the temperature $T \rightarrow 0$, leading to that the optimal estimation precision $\delta T \approx 0$ is achieved. The result shows that we can perform very accurate low temperature measurement in the case of the invariant subspace with infinite states. As shown in Fig. 1, the QFI increases with the value of M for $\omega \gg T$. On the contrary, the QFI decreases with the value of M for high temperature. For M is larger than a certain characteristic value M_c , the relationship between QFI and M becomes increasingly independent. From Fig. 1, we can see that the characteristic value M_c decreases as the temperature T increases. In the low temperature region, the increasing number of polariton states M promotes the estimation precision of temperature more obviously.

For the general case, i.e., $N > 1$ and $M > 1$, there are no analytical solutions. It can be calculated numerically to investigate the effects of the total number of polaritons N and the number of polariton states $M + 1$ on the QFI of the temperature T . As shown in Fig. 2,




Figure 1. The change relation diagram of the ratio of the QFI corresponding to finite M value and the maximal QFI $1/T^2$ corresponding to infinite M , i.e., $T^2 F_T$, versus M for three different temperatures in arbitrary units: $T = 0.5, 3.0, 10$. Here the dimensionless parameters are given by $\omega = 10$ and $N = 1$.




Figure 2. The QFI of the temperature changes with the temperature T for the four numbers of polaritons: $N = 1, 2, 3, 4$. Here the dimensionless parameters are given by $\omega = 10$ and $M = 10$.

with the increase of the polaritons N , the QFI of the temperature T increases when the temperature T above a certain value. However, the enhancement effect becomes less and less obvious as N increases. This result is consistent with the previous analytical result of $M = 1$, which is independent of the total number of polaritons N for large number $N \gg 1$. By contrast, increasing M can significantly improve the QFI with $N = 4$, especially in the low-temperature areas, as shown in Fig. 3. This conclusion also supports the previous analytical results.

As a summary, one of our main results is that increasing the number of polariton states can greatly enhance the measurement precision of the low temperature. However, the total number of polaritons N plays a smaller and smaller role in enhancing the measurement precision of the temperature as N increases.

The polariton dissipation and incoherent pumping.- We consider that there is a polariton dissipation in the polari-




Figure 3. The QFI of the temperature changes with the temperature T for the three numbers of polariton states: $M = 2, 4, 6$. Here the dimensionless parameters are given by $\omega = 10$, and $N = 4$.

ton system, which is generally unavoidable in real quantum systems. We assume that the coupling between the polariton system and the environment is weak, leading to that the Born-Markov approximation[48, 49] can be utilized. Therefore, the polariton dissipation can be described by the Lindblad superoperator

$$L_{\text{diss}}(\rho) = \sum_{j=0}^M \gamma_j (a_j \rho a_j^\dagger - \frac{1}{2} \rho a_j^\dagger a_j - \frac{1}{2} a_j^\dagger a_j \rho), \quad (11)$$

where γ_j is the dissipation rate of the j th polariton state. Without extra pumping, the total number of the polaritons in the steady state will be 0 due to the dissipation, i.e., the steady system is in the vacuum state $|0, 0, \dots, 0\rangle$. The information of temperature can not be obtained by the steady state due to that the dissipation erases all information. In order to get the information of the temperature, it is necessary to take the measurements in advance before the system reaches the steady state. In this work, we assume that the thermalization process is much faster than the dissipation process, i.e., $\Gamma_{0j}(1 + \langle a_0^\dagger a_0 \rangle) \gg \gamma_j$. The density matrix in Eq. (4) can be obtained approximately at time $\gamma_j^{-1} \gg t \gg \Gamma_{0j}^{-1}(1 + \langle a_0^\dagger a_0 \rangle)^{-1}$. In thus case, the optimal interrogation time should be much smaller than the characteristic time of the dissipation process and larger than that of the thermalization process.

In order not to control the interrogation time, extra pumping is required to obtain a non-vacuum steady state. We consider that there is a incoherent pumping. The energy transfers from the upper branch and uncoupled excitons towards the lower branch can be treated as an effective incoherent pumping, which can be described by

the Lindblad master equation

$$L_{\text{pump}}(\rho) = \sum_{j=0}^M \kappa_j (a_j \rho a_j^\dagger - \frac{1}{2} \rho a_j^\dagger a_j - \frac{1}{2} a_j^\dagger a_j \rho) + \sum_{j=0}^M \kappa_j (a_j^\dagger \rho a_j - \frac{1}{2} \rho a_j a_j^\dagger - \frac{1}{2} a_j a_j^\dagger \rho), \quad (12)$$

where κ_j is the pumping rate of the j th polariton state. Including the dissipation and the incoherent pumping, the density matrix of the polaritons ρ is dominated by the master equation

$$\frac{d}{dt} \rho = -i[H, \rho] + L_{\text{thermal}}(\rho) + L_{\text{diss}}(\rho) + L_{\text{pump}}(\rho). \quad (13)$$

In general, the above equation is difficult to be solved numerically and analytically. It can be approximately solved by assuming that the thermalization is much faster than the dissipation and the incoherent pumping, i.e., $\Gamma_{0j}(1 + \langle a_0^\dagger a_0 \rangle) \gg \gamma_j, \kappa_j$ [50, 51]. The general expression for the density matrix in the steady state ρ_s is also described by

$$\rho_s = \sum_{N=0}^{\infty} P_N \frac{1}{Z_N} \sum_{n_0 + \dots + n_M = N} \nu_0^{n_0} \dots \nu_M^{n_M} \times |n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|, \quad (14)$$

where P_N denotes the probability that there are N polaritons in total in the steady state. In this case, P_N is independent of the initial value $P_N(0)$. Without loss of generality, we consider $\kappa_j = \kappa$ and $\gamma_j = \gamma$. For only two polariton states ($M = 1$), we can obtain the general form of the probabilities (see Appendix A for details)

$$P_N = \left(\frac{\kappa}{\kappa + \gamma} \right)^N (N + 1) P_0, \quad (15)$$

$$P_0 = \frac{1}{1 + \frac{\kappa(\kappa + 2\gamma)}{\gamma^2}}. \quad (16)$$

In the steady state, the probability of the state $|n_0, n_1\rangle$ with $n_0 + n_1 = N$ is given by

$$P_{N, n_0, n_1} = P_N \lambda^{-n_1} (\lambda - 1) / (\lambda - \lambda^{-N}), \quad (17)$$

where n_1 ranges from 0 to N . The corresponding QFI can be calculated by combing Eq. (6) and Eq. (17), and the results are directly shown in Fig. 4. The QFI can increase with the pumping rate κ for different temperature, but not all the time. When the pumping rate κ is larger than a certain value, the QFI is independent of the pumping rate. It is because that the number of polaritons N is significantly increased by the strong enough pumping, i.e., $\sum_{N \gg 1} P_N \sim 1$. As shown in Eq. (8), the QFI is independent of the number of polaritons N in the case of $N \gg 1$. Therefore, too strong pumping is wasteful




Figure 4. Diagram of the QFI F_T as a function of pumping rate κ with four different temperature: $T = 1, 3, 5, 7$. Here, the dimensionless parameters are given by $\omega = 10$, and $\gamma = 10$.

in improving the measurement precision of the temperature T . Only the appropriate strength of the incoherent pumping is required to resist the effects of dissipation.

M degenerate states.- In order to deal with the case of $N \gg 1$ and $M \gg 1$, we consider that there are M degenerate states (modes), i.e., $\omega_j = \omega_0 + \omega$ for $j \geq 1$. The general form of the partition function Z_N^d in the case of M degenerate states are obtained (see Appendix B for details)

$$Z_N^d = \frac{\lambda^M}{(\lambda - 1)^M} - \frac{(M + N)! {}_2F_1[1, 1 + M + N, 2 + N; \frac{1}{\lambda}]}{(M - 1)!(N + 1)!\lambda^{(1+N)}}, \quad (18)$$

where ${}_2F_1(a, b, c; d) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k d^k}{(c)_k k!}$ denotes the generalized Hypergeometric function with $(r)_k = \Gamma(r + k)/\Gamma(r)$. The general form of the probability of the state $|n_0, \dots, n_M\rangle$ with $n_0 + \dots + n_M = N$ is given by

$$P_{N, n_0, \dots, n_M}^d = \frac{\lambda^{n_0 - N} P_N^d}{Z_N}, \quad (19)$$

where P_N^d denotes the probability that there are N polaritons in total in the steady state: $P_N^d = P_N(0)$ in the absence of the dissipation and the incoherent pumping; $P_N^d = P_N$ with the dissipation and the incoherent pumping. We note that there are $(N - n_0 + M - 1)! / [(N - n_0)!(M - 1)!]$ different states $|n_0, \dots, n_M\rangle$ with the same probability P_{N, n_0, \dots, n_M}^d due to the M degenerate modes. This simplifies the calculation somewhat. We numerically calculate the corresponding QFI F_T^d with the large N and M .

As shown in Fig. 5, the QFI increases with the number of states M with the fixed number of the polaritons $N = 100$, especially in the low temperature. It means that the number of polariton states M is an important resource for enhancing the estimation precision of the temperature, which is consistent with the previous results by using the small N and M in the case of states equidistant in frequency.




Figure 5. Diagram of the QFI F_T^d as the function of the temperature T with three different values of M : $M = 100, 200, 300$. The number of the polaritons is fixed, i.e., $P_N^d = 1$. Here, the dimensionless parameters are given by $\omega = 10$, and $N = 100$.




Figure 6. Diagram of the QFI F_T as the function of the temperature T with three different numbers of the polaritons: $N = 10, 200, 300$. Here, the dimensionless parameters are given by $\omega = 10$, $P_N^d = 1$, and $M = 100$.

As shown in Fig. 6, in the case of large M , the line $N = 100$ is coincident with the line $N = 200$. It implies that the QFI of the temperature is independent of the number of the polaritons for large N in the case of large M . The value of M can not change the relation between the QFI and the number of the polaritons N . *Discussion and Conclusion.*- We have proposed a new mechanism to improve the measurement precision of the low temperature and obtained Landau bound $\delta T \simeq T$. It is based on the existence of invariant subspaces during the polariton thermalization. The measurement precision of the low temperature increases significantly with the number of polariton states. It is interesting to apply our mechanism to different systems beyond polariton BECs by looking for invariant subspaces with many modes during the thermalization.

When the polaritons suffer from the dissipation, the incoherent pumping can be used to enhancing the estimation precision of temperature. Due to that the QFI becomes less and less dependent on the total number of the polaritons, too strong incoherent pumping is a waste of energy. Whether the periodic coherent modulation and nonlinearity can be used to enhancing the estimation precision of low temperature deserves further study.

Acknowledgements.-This research was supported by the National Natural Science Foundation of China under Grant No. 62001134, Guangxi Natural Science Foundation under Grant No. 2020GXNSFAA159047 and National Key R&D Program of China under Grant No. 2018YFB1601402-2.

-
- [1] Y. Gao, and Y. Bando, Nanotechnology: carbon nanothermometer containing gallium, Nature 415, 599 (2002).
 [2] D. M. Weld, P. Medley, H. Miyake, D. Hucul, D. E. Pritchard, and W. Ketterle, Spin gradient thermometry

- for ultracold atoms in optical lattices, Phys. Rev. Lett. 103, 245301 (2009).
 [3] P. Neumann, I. Jakobi, F. Dolde, C. Burk, R. Reuter, G. Waldherr, J. Honert, T. Wolf, A. Brunner, and J. H.

- Shim, High-precision nanoscale temperature sensing using single defects in diamond. *Nano Lett.* 13, 2738 (2013).
- [4] F. Giazotto, T. T. Heikkilä, A. Luukanen, A. M. Savin, and J. P. Pekola, Opportunities for mesoscopics in thermometry and refrigeration: Physics and applications, *Rev. Mod. Phys.* 78, 217 (2006).
- [5] N. Navon, S. Nascimbène, F. Chevy, and C. Salomon, The Equation of State of a Low-Temperature Fermi Gas with Tunable Interactions, *Science* 328, 729 (2010).
- [6] M. J. H. Ku, A. T. Sommer, L. W. Cheuk, and M. W. Zwierlein, Revealing the Superfluid Lambda Transition in the Universal Thermodynamics of a Unitary Fermi Gas, *Science* 335, 563 (2012).
- [7] L. D. Carlos, F. Palacio, eds., *Thermometry at the Nanoscale* (The Royal Society of Chemistry, Cambridge, 2016).
- [8] W. Hofstetter and T. Qin, Quantum simulation of strongly correlated condensed matter systems, *J. Phys. B* 51, 082001 (2018).
- [9] L. Tarruell and L. Sanchez-Palencia, Quantum simulation of the Hubbard model with ultracold fermions in optical lattices, *C. R. Phys.* 19, 365 (2018).
- [10] A. De Pasquale, and T. M. Stace, in *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions*, edited by F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso (Springer International Publishing, Cham, 2018), p. 503.
- [11] M. Mehboudi, A. Sanpera, and L. A. Correa, Thermometry in the quantum regime: Recent theoretical progress, *J. Phys. A: Math. Theor.* 52, 303001 (2019).
- [12] M. Mehboudi, A. Lampo, C. Charalambous, L. A. Correa, M. A. Garca-March, and M. Lewenstein, Using Polarons for sub-NK Quantum Nondemolition Thermometry in a Bose-Einstein Condensate, *Phys. Rev. Lett.* 122, 030403 (2019).
- [13] Q. Bouton, J. Nettersheim, D. Adam, F. Schmidt, D. Mayer, T. Lausch, E. Tiemann, and A. Widera, Single-Atom Quantum Probes for Ultracold Gases Boosted by Nonequilibrium Spin Dynamics, *Phys. Rev. X* 10, 011018 (2020).
- [14] M. T. Mitchison, T. Fogarty, G. Guarnieri, S. Campbell, T. Busch, and J. Goold, In Situ Thermometry of a Cold Fermi gas via Dephasing Impurities, *Phys. Rev. Lett.* 125, 080402 (2020).
- [15] M. Campisi, P. Hänggi, and P. Talkner, Colloquium: Quantum fluctuation relations: Foundations and applications, *Rev. Mod. Phys.* 83, 771 (2011).
- [16] F. Brandão, M. Horodecki, N. Ng, J. Oppenheim, and S. Wehner, The second laws of quantum thermodynamics, *Proc. Natl. Acad. Sci. USA* 112, 3275 (2015).
- [17] S. Vinjanampathy and J. Anders, Quantum thermodynamics, *Contemp. Phys.* 57, 545 (2016).
- [18] S. Deffner and S. Campbell, *Quantum Thermodynamics* (Morgan & Claypool Publishers, 2019).
- [19] U. Marzolino and D. Braun, Precision measurements of temperature and chemical potential of quantum gases, *Phys. Rev. A* 88, 063609 (2013).
- [20] D. Xie, C. Xu, and A. Wang, Optimal quantum thermometry by dephasing, *Quantum Inf Process* 16, 155 (2017).
- [21] J. Yang, C. Elouard, J. Splettstoesser, B. Sothmann, R. Sánchez, and A. N. Jordan, Thermal transistor and thermometer based on Coulomb-coupled conductors, *Phys. Rev. B* 100, 045418 (2019).
- [22] P. P. Hofer, J. B. Brask, M. Perarnau-Llobet, and N. Brunner, Quantum Thermal Machine as a Thermometer, *Phys. Rev. Lett.* 119, 090603 (2017).
- [23] L. Spietz, K. W. Lehnert, I. Siddiqi, and R. J. Schoelkopf, Primary Electronic Thermometry Using the Shot Noise of a Tunnel Junction, *Science* 300, 1929 (2003).
- [24] L. Spietz, R. J. Schoelkopf, and P. Pari, Shot noise thermometry down to 10mK, *Appl. Phys. Lett.* 89, 183123 (2006).
- [25] S. Gasparinetti, F. Deon, G. Biasiol, L. Sorba, F. Beltram, and F. Giazotto, Probing the local temperature of a two-dimensional electron gas microdomain with a quantum dot: Measurement of electron-phonon interaction, *Phys. Rev. B* 83, 201306(R) (2011).
- [26] M. Brunelli, S. Olivares, and M. G. A. Paris, Qubit thermometry for micromechanical resonators, *Phys. Rev. A* 84, 032105 (2011).
- [27] S. Jevtic, D. Newman, T. Rudolph, and T. M. Stace, Single qubit thermometry, *Phys. Rev. A* 91, 012331 (2015).
- [28] L. A. Correa, M. Mehboudi, G. Adesso, and A. Sanpera, Individual Quantum Probes for Optimal Thermometry, *Phys. Rev. Lett.* 114, 220405 (2015).
- [29] S. Campbell, M. Mehboudi, G. D. Chiara, and M. Paternostro, Global and local thermometry schemes in coupled quantum systems, *New J. Phys.* 19, 103003 (2017).
- [30] A. H. Kiilerich, A. De Pasquale, and V. Giovannetti, Dynamical approach to ancilla-assisted quantum thermometry, *Phys. Rev. A* 98, 042124 (2018).
- [31] D. Xie, F. Sun, and C. Xu, Quantum thermometry based on a cavity-QED setup, *Phys. Rev. A* 101, 063844 (2020).
- [32] G. De Palma, A. De Pasquale, and V. Giovannetti, Universal locality of quantum thermal susceptibility, *Phys. Rev. A* 95, 052115 (2017).
- [33] Luis A. Correa, Mart Perarnau-Llobet, Karen V. Hovhannisyanyan, Senaida Hernandez-Santana, Mohammad Mehboudi, and Anna Sanpera, Enhancement of low-temperature thermometry by strong coupling, *Phys. Rev. A* 96, 062103 (2017).
- [34] V. Mukherjee, A. Zwick, A. Ghosh, X. Chen, and G. Kurizki, Enhanced precision bound of low-temperature quantum thermometry via dynamical control, *Commun. Phys.* 2, 162 (2019).
- [35] Ning Zhang, Chong Chen, Si-Yuan Bai, Wei Wu, and Jun-Hong An, Non-Markovian Quantum Thermometry, *Phys. Rev. Applied* 17, 034073 (2022).
- [36] M. G. A. Paris, Achieving the Landau bound to precision of quantum thermometry in systems with vanishing gap, *J. Phys. A: Math. Theor.* 49, 03LT02 (2015).
- [37] T. Byrnes, N. Y. Kim, and Y. Yamamoto, Exciton-polariton condensates, *Nat. Phys.* 10, 803 (2014).
- [38] A. V. Zasedatelev, A. V. Baranikov, D. Urbonas, F. Scafrimuto, U. Scherf, T. Stöferle, R. F. Mahrt, and P. G. Lagoudakis, A room-temperature organic polariton transistor, *Nat. Photonics* 13, 378 (2019).
- [39] Vladislav Yu. Shishkov and Evgeny S. Andrianov, Exact Analytical Solution for the Density Matrix of a Nonequilibrium Polariton Bose-Einstein Condensate, *Phys. Rev. Lett.* 128, 065301 (2022).
- [40] R. Kubo, Statistical-Mechanical Theory of Irreversible Processes. I. General Theory and Simple Applications to Magnetic and Conduction Problems, *J. Phys. Soc. Jpn.* 12, 570 (1957).
- [41] P. C. Martin and J. Schwinger, Theory of Many-Particle Systems. I, *Phys. Rev.* 115, 1342 (1959).

- [42] Artem Strashko, Peter Kirton, and Jonathan Keeling, Organic Polariton Lasing and the Weak to Strong Coupling Crossover, *Phys. Rev. Lett.* 121, 193601 (2018).
- [43] Mohammad Ramezani, Quynh Le-Van, Alexei Halpin, and Jaime G3mez Rivas, Nonlinear Emission of Molecular Ensembles Strongly Coupled to Plasmonic Lattices with Structural Imperfections, *Phys. Rev. Lett.* 121, 243904 (2018).
- [44] V. Yu. Shishkov, E. S. Andrianov, A. A. Pukhov, A. P. Vinogradov, and A. A. Lisiansky, Zeroth law of thermodynamics for thermalized open quantum systems having constants of motion, *Phys. Rev. E* 98, 022132 (2018).
- [45] Samuel L. Braunstein, Carlton M. Caves G. J. Milburn, Generalized Uncertainty Relations: Theory, Examples, and Lorentz Invariance, *Ann. Phys. (NY)* 247, 135 (1996).
- [46] Samuel L. Braunstein and Carlton M. Caves, Statistical distance and the geometry of quantum states, *Phys. Rev. Lett.* 72, 3439 (1994).
- [47] V. Giovannetti, S. Lloyd, and L. Maccone, Advances in quantum metrology, *Nat. Photonics* 5, 222 (2011).
- [48] Vittorio Gorini and Andrzej Kossakowski, Completely positive dynamical semigroups of N-level systems, *J. Math. Phys.* 17, 821 (1976).
- [49] H.-P. Breuer, and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press on Demand, New York, 2002).
- [50] T. K. Hakala, A. J. Moilanen, A. I. V4ikevainen, R. Guo, J.-P. Martikainen, K. S. Daskalakis, H. T. Rekola, A. Julku, and P. Torma, BoseEinstein condensation in a plasmonic lattice, *Nature Physics* 14, 739 (2018).
- [51] A. I. V4ikev4inen, A. J. Moilanen, M. Ne3ada, T. K. Hakala, K. S. Daskalakis, and P. T3rm4, Sub-picosecond thermalization dynamics in condensation of strongly coupled lattice plasmons, *Nature communications* 11, 1 (2020).

APPENDIX A

We consider that there are polariton dissipation and incoherent pumping of the lower polariton states, which are described by the Lindblad superoperators

$$L_{\text{diss}}(\rho) = \sum_{j=0}^M \gamma_j (a_j \rho a_j^\dagger - \frac{1}{2} \rho a_j^\dagger a_j - \frac{1}{2} a_j^\dagger a_j \rho) \quad (\text{S1})$$

$$L_{\text{pump}}(\rho) = \sum_{j=0}^M \kappa_j (a_j \rho a_j^\dagger - \frac{1}{2} \rho a_j^\dagger a_j - \frac{1}{2} a_j^\dagger a_j \rho) +$$

$$\sum_{j=0}^M \kappa_j (a_j^\dagger \rho a_j - \frac{1}{2} \rho a_j a_j^\dagger - \frac{1}{2} a_j a_j^\dagger \rho), \quad (\text{S2})$$

where γ_j (κ_j) is the dissipation (the incoherent pumping) rate of the j th state. The general solution is difficult and we also consider that the thermalization process is very rapid, i.e., $\Gamma_{0j}(1 + \langle a_0^\dagger a_0 \rangle) \gg \kappa_j, \gamma_j$, which has been proved to be valid in ref.[39]. In the first stage, i.e., $\kappa_j^{-1}, \gamma_j^{-1} \gg t \gg \Gamma_{0j}^{-1}(1 + \langle a_0^\dagger a_0 \rangle)^{-1}$, the density matrix $\rho(t)$ obeys the approximate differential equation by

ignoring the effect of the dissipation and the incoherent pumping,

$$\frac{d}{dt} \rho \approx -i[H, \rho] + L_{\text{thermal}}(\rho), \quad (\text{S3})$$

which leads to the general expression for the density matrix ρ [39]

$$\rho = \sum_{N=0}^{\infty} P_N(t) \frac{1}{Z_N} \sum_{n_0+\dots+n_M=N} \nu_0^{n_0} \dots \nu_M^{n_M} \times |n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|, \quad (\text{S4})$$

where $P_N(t) \approx P_N(0)$ for $\kappa_j^{-1}, \gamma_j^{-1} \gg t \gg \Gamma_{0j}^{-1}(1 + \langle a_0^\dagger a_0 \rangle)^{-1}$.

In the second stage, one can substitute the above equation into the whole evolution equation of the density matrix ρ , which is described as

$$\frac{d}{dt} \rho = -i[H, \rho] + L_{\text{thermal}}(\rho) + L_{\text{diss}}(\rho) + L_{\text{pump}}(\rho). \quad (\text{S5})$$

Substituting Eq. (S4) into Eq. (S5) and using the steady conditions $\frac{d}{dt} P_N(t \rightarrow \infty) = 0$, the steady-state solutions of the probabilities $P_N = P_N(t \rightarrow \infty)$ are achieved[39]

$$P_{N+1} = \frac{(d_{N-1} + \beta_N) Z_{N+1}}{d_N Z_N} P_N - \frac{\beta_{N-1} Z_{N+1}}{d_N Z_{N-1}} P_{N-1}, \quad (\text{S6})$$

$$P_1 = \frac{\beta_0 Z_1}{d_0 Z_0} P_0, \quad (\text{S7})$$

where $d_N = \sum_{n=0}^N \sum_{j=0}^M (\gamma_j + \kappa_j) \nu_j^{n+1} Z_{N-n}$ and $\beta_N = \sum_{n=0}^N \sum_{j=0}^M \kappa_j \nu_j^n Z_{N-n}$. Utilizing the above equations, we further obtain that

$$P_2 = \frac{\beta_1 Z_2}{d_1 Z_1} P_1, \quad (\text{S8})$$

$$P_3 = \frac{\beta_2 Z_3}{d_2 Z_2} P_2, \quad (\text{S9})$$

$$P_4 = \frac{\beta_3 Z_4}{d_3 Z_3} P_3, \quad (\text{S10})$$

$$P_5 = \frac{\beta_4 Z_5}{d_4 Z_4} P_4, \quad (\text{S11})$$

...

By induction, we can obtain the general form of the probability P_N with $N \geq 1$

$$P_N = \frac{\beta_{N-1} Z_N}{d_{N-1} Z_{N-1}} P_{N-1} \quad (\text{S12})$$

$$= \frac{\prod_{n=0}^{N-1} \beta_n Z_N}{\prod_{n=0}^{N-1} d_n} P_0. \quad (\text{S13})$$

Without loss of generality, we consider $\kappa_j = \kappa$ and $\gamma_j = \gamma$. In the case of two states ($M = 1$), we can obtain the analytical results about the distribution

$$d_N = \frac{\lambda^{-(1+N)}(\lambda^{2+N} - 1)(1 + N)(\kappa + \gamma)}{(\lambda - 1)}, \quad (\text{S14})$$

$$\beta_N = \frac{\lambda^{-N}(\lambda^{1+N} - 1)(2 + N)\kappa}{(\lambda - 1)}, \quad (\text{S15})$$

$$Z_N = \sum_{n=0}^N \exp[-n\omega/T] = \frac{\lambda - \lambda^{-N}}{\lambda - 1}. \quad (\text{S16})$$

Substituting the above equations into Eq. (S13), we can derive a simplified form of the probability P_N , which is described as

$$P_N = \left(\frac{\kappa}{\kappa + \gamma}\right)^N (N + 1)P_0. \quad (\text{S17})$$

Due to the normalization condition $\sum_{N=0}^{\infty} P_N = 1$, we can obtain that

$$P_N = \left(\frac{\kappa}{\kappa + \gamma}\right)^N (N + 1)P_0, \quad (\text{S18})$$

$$P_0 = \frac{1}{1 + \frac{\kappa(\kappa + 2\gamma)}{\gamma^2}}. \quad (\text{S19})$$

APPENDIX B

In the case of M degenerate states, $\omega_j = \omega_0 + \omega$ for $j \geq 1$, the corresponding density matrix in the steady state $\rho_s^d = \rho^d(t \rightarrow \infty)$ is also written as

$$\rho_s^d = \sum_{N=0}^{\infty} P_N^d \frac{1}{Z_N^d} \sum_{n_0 + \dots + n_M = N} \nu_0^{n_0} \dots \nu_M^{n_M} \times |n_0, n_1, \dots, n_M\rangle \langle n_0, n_1, \dots, n_M|, \quad (\text{S20})$$

The partition function can be derived

$$Z_N^d = \sum_{n_0 + \dots + n_M = N} \lambda^{-(n_1 + \dots + n_M)} = \sum_{n_0=0}^N \mu(n_0) \lambda^{n_0 - N} \quad (\text{S21})$$

$$= \frac{\lambda^M}{(\lambda - 1)^M} - \frac{(M + N)! {}_2F_1[1, 1 + M + N, 2 + N; \frac{1}{\lambda}]}{(M - 1)!(N + 1)!\lambda^{(1+N)}}, \quad (\text{S22})$$

where the degeneration coefficient $\mu(n_0) = (N - n_0 + M - 1)!/[N!(M - 1)!]$ comes from the M degenerate modes, and ${}_2F_1(a, b, c; d) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k d^k}{(c)_k k!}$ denotes the generalized Hypergeometric function with $(r)_k = \Gamma(r + k)/\Gamma(r)$. The general form of the probability of the state $|n_0, n_1, \dots, n_M\rangle$ with the total number $n_0 + n_1 + \dots + n_M = N$ is given by

$$P_{N, n_0, \dots, n_M}^d = \frac{\lambda^{n_0 - N} P_N^d}{Z_N^d}, \quad (\text{S23})$$

where P_N^d denotes the probability that there are N polaritons in total in the steady state: $P_N^d = P_N(0)$ in the absence of the dissipation and the incoherent pumping; $P_N^d = P_N$ with the dissipation and the incoherent pumping. In the case of M degenerate states, the general formula of QFI F_T^d can be described as

$$F_T^d = \sum_{N=0}^{\infty} \sum_{n_0 + \dots + n_M = N} \frac{(\partial_T P_{N, n_0, \dots, n_M}^d)^2}{P_{N, n_0, \dots, n_M}^d} \quad (\text{S24})$$

$$= \sum_{N=0}^{\infty} \sum_{n_0=0}^N \mu(n_0) \times \frac{(\partial_T \frac{\lambda^{n_0 - N} P_N^d}{Z_N^d})^2}{\frac{\lambda^{n_0 - N} P_N^d}{Z_N^d}} \quad (\text{S25})$$

$$= \sum_{N=0}^{\infty} \sum_{n_0=0}^N \mu(N - n_0) \times \frac{(\partial_T \frac{\lambda^{-n_0} P_N^d}{Z_N^d})^2}{\frac{\lambda^{-n_0} P_N^d}{Z_N^d}}. \quad (\text{S26})$$

For the fixed number of the polaritons, i.e., $P_N^d = 1$, the QFI can be calculated by

$$F_T^d = \sum_{n_0=0}^N \mu(N - n_0) \times \frac{(\partial_T \frac{\lambda^{-n_0}}{Z_N^d})^2}{\frac{\lambda^{-n_0}}{Z_N^d}}. \quad (\text{S27})$$