

FINITE QUOTIENTS OF SINGULAR ARTIN MONOIDS AND CATEGORIFICATION OF THE DESINGULARIZATION MAP

HELENA JONSSON, VOLODYMYR MAZORCHUK, ELIN PERSSON WESTIN,
SHRADDHA SRIVASTAVA, MATEUSZ STROIŃSKI AND XIAOYU ZHU

ABSTRACT. We study various aspects of the structure and representation theory of singular Artin monoids. This includes a number of generalizations of the desingularization map and explicit presentations for certain finite quotient monoids of diagrammatic nature. The main result is a categorification of the classical desingularization map for singular Artin monoids associated to finite Weyl groups using BGG category \mathcal{O} .

1. INTRODUCTION AND DESCRIPTION OF THE RESULTS

1.1. Brief introduction. Singular braid monoids are certain natural generalizations of braid groups that play an important role in the theory of finite type invariant of knots and links (a.k.a. Vassiliev invariants), see [Bi93]. A singular braid allows for two strands to cross each other, creating a so-called *singular crossing*. A singular link is the closure of a singular braid and to study invariants of singular links is a natural and interesting problem. A singular braid monoid admits a presentation which generalizes Artin's presentation for the corresponding braid group, see [Bi93, Lemma 3]. Extrapolating to the setup of an arbitrary Coxeter matrix allows one to define a singular Artin monoid, for each Artin braid group.

There is an interesting relation between a singular Artin monoid and the corresponding (regular) Artin braid group. Interpreting a singular crossing as the difference between the left and right crossings defines a homomorphism from a singular Artin monoid to the integral group algebra of the corresponding Artin braid group. In [Pa04], it is shown that this map is injective for all singular braid monoids. A similar result is known for some singular Artin monoids, see [GP05, Ea06a], however, to the best of our knowledge, the general case is still open. Various structure properties of singular Artin monoids and their applications to low dimensional topology were studied by many authors during the last 30 years, see e.g. [An04, An05, Bae92, Bi93, Co00, Co05, Ea06a, Ea15, GP05, Ja17, Pa04, Ve10] and references therein.

Each braid group admits an interesting upgrade to a 2-category, see, for example, [Ro06]. One of the classical 2-representations of this object, usually referred to as a 2-braid group, is given by the action of derived shuffling functors on the blocks of the Bernstein-Gelfand-Gelfand (BGG) category \mathcal{O} for the Lie algebra \mathfrak{sl}_n , see [MaStr05]. It is even known that this action is faithful, see [KS02]. In [MaStr07, Theorem 1] (see also [CM17, Theorem 5.1]), this was extended to a categorical action of the singular braid monoid. Unfortunately, this construction does not have any obvious connection to the desingularization map, even on the level of the Grothendieck group.

1.2. Story of the paper: how and why. During the academic year 2020-2021, the second author gave a PhD level course about category \mathcal{O} at Uppsala University (all lectures are available on YouTube via the Uppsala Algebra Channel). One of these lectures was exactly about the categorical action of the singular braid monoid on category

\mathcal{O} . The research presented in this paper started from some questions asked by some participants of the course after that lecture. One of the main questions was whether one could define another categorical action of the singular braid monoid on category \mathcal{O} that would be better related to the classical desingularization map. The discussions that started during the spring of 2021 were organized as a study group on singular Artin monoids during the academic year 2021-2022 in Uppsala. The present paper is the report, by all participants, of the discussions during the meetings of this study group.

Now we will briefly describe what we discussed. This will also help to summarize the results and the structure of the present paper.

We looked into three main directions:

- Generalizations of the desingularization map.
- Finite diagrammatic quotients of singular Artin monoids.
- Categorification of the desingularization map.

One could argue that some of the results of the paper could be cleaned up, strengthened and pushed further. However, the time constraints on the organization of this study group put strict bounds on what we could include in this paper.

1.3. Results: generalized desingularization map. The first of the research directions, on various generalizations of the desingularization map, was originally motivated by the construction in [MaStr07, Theorem 1]. If the latter result is not related to the classical desingularization map, then what kind of map is it related to? In Section 3 we investigate various generalizations of the desingularization map. In particular, in Subsection 3.2 we consider the odd skeleton of a Coxeter graph, that is, the graph obtained from the Coxeter graph by removing all even edges. We show that one can define an analogue of the desingularization map for each assignment of a Laurent polynomial to a connected component of the odd skeleton, see Proposition 2. Moreover, for all cases for which the classical desingularization map is known to be injective, we establish, in Proposition 3, the injectivity of a generic choice for the generalized desingularization map constructed in Proposition 2. In Subsection 3.2, we also discuss some further generalizations of the classical desingularization map in the setup when one replaces the integral group algebra of the Artin braid group by the semialgebra of the Artin braid group over the Boolean semiring.

The situation encountered in [MaStr07, Theorem 1] is explained in Section 4. The group algebra of the braid group has an interesting quotient, called the Hecke algebra. The latter controls the combinatorics of category \mathcal{O} . The categorification of the singular braid monoid proposed in [MaStr07, Theorem 1] corresponds to a generalized desingularization map for this quotient. This latter map is given by assigning to the connected components of the odd skeleton of the corresponding Coxeter graph the Laurent polynomial describing the Kazhdan-Lusztig basis element of the Hecke algebra associated to a simple reflection.

1.4. Results: finite diagrammatic quotients of singular Artin monoids. The second major direction which we explore studies various finite diagrammatic quotients of singular Artin monoids. Diagram algebras are intensively studied in modern representation theory due to their numerous applications in various areas of contemporary mathematics and theoretical physics. Such algebras usually have a basis given by some kind of combinatorially defined diagrams, and multiplication is based on concatenation

of these diagrams. The most general such object (which contains all other as subobjects) is the algebra of partitioned binary relations, studied in [MM13]. Other examples include the Brauer algebra, see [Br37], and the partition algebra, see [Mar94].

Diagram algebras are usually defined as deformations of the corresponding diagram monoid. In the latter, multiplication is given exactly by concatenation of diagrams, which usually involves some straightening procedure. A diagram monoid usually has the symmetric group as the group of invertible elements. The symmetric group is, of course, a quotient of the braid group. It is therefore natural to ask whether the canonical projection from the braid group to the symmetric group can be extended to a homomorphism from the singular braid monoid to the diagram monoid in question. In case such an extension is possible, one could try to characterize its kernel, leading to a presentation of the image by generators and relations. As a next step, based on this presentation, one could then try to define analogues of this diagram monoid for other Coxeter types. That was our main idea for this second direction addressed in the paper.

All results related to finite quotients of singular Artin monoids are collected in Section 5. As it turned out, most of them can be organized so that one starts with a maximal possible quotient and then, step by step, projects further.

Our starting point is a homomorphism from the singular braid monoid to the monoid of all binary relations, see Proposition 6. The homomorphism itself is inspired by our generalized desingularization maps. In this case, instead of the usual integral group algebra of the braid group, we consider, as the target of our homomorphism, the semialgebra of the braid group over the Boolean semiring. The image of the singular generator τ_s under this map should be thought of as the set $\{e, \sigma_s\}$ consisting of the identity e and the corresponding regular element σ_s . This map appeared previously in the context of 0-Hecke monoids, see [MaSte12]. The monoid generated by the images of the singular elements (in type A) was studied in [MaSte12] under the name of double Catalan monoid. The image of the singular braid monoid under this map is also related to the factor power of the symmetric group as defined and studied in [GM95a, GM95b]. In Subsections 5.2.3 and 5.2.4 we explore type B analogues of these results.

The next target for a homomorphism is the dual symmetric inverse monoid \mathbf{I}_n^* , that is, the inverse monoid of all bijections between quotients of the finite set $\underline{n} = \{1, 2, \dots, n\}$. In Proposition 8 we show that this monoid is a natural quotient of the monoid of essential binary relations on \underline{n} . The latter contains the image of the singular braid monoid. Composing with the projection map onto the quotient, we get the induced homomorphism from the singular braid monoid to \mathbf{I}_n^* . The image of this homomorphism is the maximal factorizable submonoid \mathbf{F}_n^* of \mathbf{I}_n^* . The monoid \mathbf{F}_n^* was studied in detail by several authors. In particular, there are various presentations of this monoid which can be found in the literature, see [Ea06b, EEF05, Fi03].

In Subsections 5.3.4 and 5.3.5 we look at certain type B analogues of the monoids \mathbf{F}_n^* and \mathbf{I}_n^* . In particular, in Proposition 14, by examining the kernel of the natural projection from the type B singular Artin monoid, we obtain a presentation for the type B analogue of the monoid \mathbf{F}_n^* .

We also note that the congruence describing the kernel of the map from the essential binary relations to \mathbf{I}_n^* can be extended to all binary relations. The corresponding quotient is the monoid of all bijections between quotients of subsets of \underline{n} , studied in [KMU15].

The next step is the classical symmetric inverse monoid \mathbf{IS}_n of all bijections between subsets of \underline{n} . It turns out, see Proposition 15, that restricting elements of \mathbf{F}_n^* to

singletons defines a homomorphism to \mathbf{IS}_n . This leads to a natural homomorphism from the singular braid monoid to \mathbf{IS}_n . This homomorphism is not surjective, its image “misses” all elements with singleton defect. Again, there is the obvious analogue of the story in type B , explored in Subsections 5.4.4 – 5.4.7. This includes a presentation for the signed rook monoid in Proposition 20, see also [EF13] and [CE22] for further generalizations.

Finally, in Subsection 5.5 we study natural surjections from singular braid monoids to Brauer monoids. The type A story is fairly expected, see Proposition 22. We also consider the natural type B analogue of this story, which turns out to be different from another type B analogue of Brauer algebras considered in [CL15]. As explained in Subsection 5.5.6, our type B Brauer monoid is related to the partial Brauer monoid introduced in [Ma98]. In fact, our type B Brauer monoid is a partially \mathbb{Z}_2 -colored version of the partial Brauer monoid. In Theorem 26 we extend the presentation of singular braid monoids and type A Brauer monoids to a presentation of type B Brauer monoids.

1.5. Results: categorification of the desingularization map. Our final main direction describes a new categorical action of singular Artin monoids on category \mathcal{O} . All necessary preliminaries on category \mathcal{O} are collected in Section 6.

The classical realization of the generators of the braid group as endofunctors of category \mathcal{O} uses derived (co)shuffling functors introduced in [Ca86]. In [MaStr07, Theorem 1] it is proposed to realize the action of the singular crossings using indecomposable projective functors (associated to the corresponding simple root). All this is recalled in Section 7. The most difficult part of the proof of this results is to check the mixed braid relations between the singular and the regular generators. In type A , the corresponding argument in the proof of [MaStr07, Theorem 1] is incomplete, a complete proof is given in [CM17, Theorem 5.1] based on an alternative approach outlined in [MaStr07]. In Theorem 32 we give a fairly short general proof of these mixed braid relations in any type. In Proposition 30 we also give a general proof of braid relations for (co)shuffling functors in all types.

Our final main results are in Section 8. To make a connection with the desingularization map, we observe that there are two obvious natural transformations between the derived shuffling and coshuffling functors. The usual philosophy of categorification suggests that the singular crossing should act via the cone of one of these. And we, indeed, show that one of the cones works, see Theorem 35, while the other one fails, see Proposition 36. The proof of Theorem 35 is split into verifying the defining relations for the generators of the singular braid monoid, one at a time.

1.6. Structure: preliminaries. Additionally to the above, in Section 2 below we collected all necessary preliminaries, including the definitions for all main protagonists of this paper which are illustrated by a detailed example in type A .

Acknowledgments. This research is partially supported by the Swedish Research Council. We thank Hankyung Ko for very helpful discussions. We thank James East for helpful comments.

2. SINGULAR ARTIN MONOIDS

2.1. Coxeter groups. Let S be a finite set with $n > 0$ elements. Consider an $n \times n$ matrix $M = (m_{s,t})_{s,t \in S}$ with entries in $\mathbb{Z}_{>0} \cup \{\infty\}$ satisfying the following conditions:

- $m_{s,s} = 1$, for all $s \in S$,

- $m_{s,t} = m_{t,s}$, for all $s, t \in S$,
- $m_{s,t} \neq 1$ provided that $s \neq t$.

Such a matrix M is called a *Coxeter matrix*. Associated to M , we have the corresponding *Coxeter group* $W = W(M)$ generated by S subject to the relations

$$(st)^{m_{s,t}} = e, \quad \text{for all } s, t \in S \text{ such that } m_{s,t} \neq \infty.$$

Note that, for $s = t$, we have the relation $s^2 = e$. Taking this into account, we have the following alternative set of relations defining W :

$$\begin{aligned} s^2 &= e, & \text{for all } s \in S; \\ \underbrace{ststs\dots}_{m_{s,t} \text{ factors}} &= \underbrace{tstst\dots}_{m_{t,s} \text{ factors}}, & \text{for all } s \neq t \in S \text{ such that } m_{s,t} \neq \infty. \end{aligned}$$

The same information as given by matrix M can be described by the corresponding Coxeter graph Γ_M defined as follows:

- the set of vertices of Γ_M is S ;
- Γ_M has no loops;
- there is an (unoriented) edge between two different vertices s and t in Γ_M if and only if $m_{s,t} > 2$;
- in case $m_{s,t} > 3$, this edge is marked by $m_{s,t}$.

2.2. Artin groups. The Artin group (a.k.a. Artin-Tits group) associated to W is the group $\mathbf{B}(W)$ generated by σ_s , where $s \in S$, subject to the relations

$$\underbrace{\sigma_s \sigma_t \sigma_s \dots}_{m_{s,t} \text{ factors}} = \underbrace{\sigma_t \sigma_s \sigma_t \dots}_{m_{t,s} \text{ factors}}, \quad \text{for all } s \neq t \in S \text{ such that } m_{s,t} \neq \infty.$$

2.3. Singular Artin monoids. The singular Artin monoid associated to W is the monoid $\mathbf{SB}(W)$ generated by σ_s and τ_s , where $s \in S$, subject to the relations

- (1) $\underbrace{\sigma_s \sigma_t \sigma_s \dots}_{m_{s,t} \text{ factors}} = \underbrace{\sigma_t \sigma_s \sigma_t \dots}_{m_{t,s} \text{ factors}}, \quad \text{for all } s \neq t \in S \text{ s. t. } m_{s,t} \neq \infty;$
- (2) $\underbrace{\tau_s \sigma_t \sigma_s \dots \sigma_t \sigma_s}_{m_{s,t} \text{ factors}} = \underbrace{\sigma_t \sigma_s \sigma_t \dots \sigma_s \tau_t}_{m_{t,s} \text{ factors}}, \quad \text{for all } s \neq t \in S \text{ s. t. } m_{s,t} \neq \infty \text{ is odd};$
- (3) $\underbrace{\tau_s \sigma_t \sigma_s \dots \sigma_s \sigma_t}_{m_{s,t} \text{ factors}} = \underbrace{\sigma_t \sigma_s \sigma_t \dots \sigma_t \tau_s}_{m_{t,s} \text{ factors}}, \quad \text{for all } s \neq t \in S \text{ s. t. } m_{s,t} \neq \infty \text{ is even};$
- (4) $\tau_s \tau_t = \tau_t \tau_s, \quad m_{s,t} = 2;$
- (5) $\tau_s \sigma_s = \sigma_s \tau_s.$

2.4. Special case: singular braid monoid. In the special case when we take $S = \{s_1, s_2, \dots, s_{k-1}\}$, for some $k \geq 2$, and the matrix

$$M = \begin{pmatrix} 1 & 3 & 2 & 2 & \dots & 2 & 2 & 2 \\ 3 & 1 & 3 & 2 & \dots & 2 & 2 & 2 \\ 2 & 3 & 1 & 3 & \dots & 2 & 2 & 2 \\ 2 & 2 & 3 & 1 & \dots & 2 & 2 & 2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 2 & 2 & 2 & 2 & \dots & 1 & 3 & 2 \\ 2 & 2 & 2 & 2 & \dots & 3 & 1 & 3 \\ 2 & 2 & 2 & 2 & \dots & 2 & 3 & 1 \end{pmatrix},$$

which corresponds to the following Coxeter graph:

$$(6) \quad s_1 \text{ --- } s_2 \text{ --- } s_3 \text{ --- } \dots \text{ --- } s_{k-1},$$

we obtain the following classical objects.

The associated Coxeter group is isomorphic to the symmetric group S_k , via the isomorphism which sends s_i to the transposition $(i, i+1)$.

The associated Artin group is isomorphic to the braid group on k strands via the isomorphism which sends $\sigma_i := \sigma_{s_i}$ to the following braid:

$$\begin{array}{cccccccc} 1 & 2 & \dots & i-1 & i & i+1 & i+2 & \dots & k \\ | & | & & | & \diagdown & \diagup & | & & | \\ 1 & 2 & \dots & i-1 & i & i+1 & i+2 & \dots & k \end{array}$$

The associated singular Artin monoid is isomorphic to the singular braid monoid on k strands via the isomorphism which sends $\sigma_i := \sigma_{s_i}$ to the above braid and which sends the generator $\tau_i := \tau_{s_i}$ to the following singular braid:

$$\begin{array}{cccccccc} 1 & 2 & \dots & i-1 & i & i+1 & i+2 & \dots & k \\ | & | & & | & \diagdown & \diagup & | & & | \\ 1 & 2 & \dots & i-1 & i & i+1 & i+2 & \dots & k \end{array}$$

3. DESINGULARIZATION MAP AND ITS GENERALIZATIONS

3.1. Desingularization map. Let M be a Coxeter matrix with the associated Coxeter group W . Consider the integral group algebra $\mathbb{Z}(\mathbf{B}(W))$. This algebra is, in particular, a monoid with respect to multiplication.

Lemma 1. *There is a monoid homomorphism $\Delta : \mathbf{SB}(W) \rightarrow \mathbb{Z}(\mathbf{B}(W))$ which is the identity on all σ_s and sends each τ_s to $\sigma_s - \sigma_s^{-1}$.*

Proof. It is straightforward to verify that the images in $\mathbb{Z}(\mathbf{B}(W))$ of the generators of $\mathbf{SB}(W)$ under Δ satisfy the defining relations of $\mathbf{SB}(W)$. \square

The map Δ was originally considered in [Bae92]. It is called the *desingularization map*, see [Bi93]. For the classical singular braid monoid, the map Δ has the following topological interpretation:

$$\begin{array}{c} \diagup \quad \diagdown \\ \bullet \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ \diagdown \quad \diagup \end{array}$$

In [Pa04], it is shown that, in the case of the classical singular braid monoid, the map Δ is injective. Similar results are also known in a number of other cases, see [GP05, Ea06a].

3.2. The odd skeleton. We want to generalize the desingularization map. In order to give the full generalization, we need to introduce the notion of the *odd skeleton* of a Coxeter graph.

Let Γ be a Coxeter graph. The *odd skeleton* $\mathbf{O}(\Gamma)$ of Γ is the graph obtained from Γ by removing all edges that are marked by even numbers or by ∞ . First, to mention a

trivial example, if Γ is the graph given by (6), then $\mathbf{O}(\Gamma) = \Gamma$. At the same time, in type B_4 , we have:

$$\Gamma : \bullet \overset{4}{\text{---}} \bullet \text{---} \bullet \text{---} \bullet, \quad \mathbf{O}(\Gamma) : \bullet \quad \bullet \text{---} \bullet \text{---} \bullet.$$

3.3. Generalized desingularization map. Let M be a Coxeter matrix with the corresponding Coxeter group W . Let Γ_M be the corresponding Coxeter graph and K the set of connected components of $\mathbf{O}(\Gamma)$. For $s \in S$, we denote by K_s the connected component of $\mathbf{O}(\Gamma)$ containing s .

For a fixed map $\Phi : K \rightarrow \mathbb{Z}[x, x^{-1}]$, define

$$\Delta_\Phi(\sigma_s) = \sigma_s \text{ and } \Delta_\Phi(\tau_s) = \Phi(K_s)[\sigma_s], \text{ for } s \in S.$$

Proposition 2. *The above assignment extends uniquely to a monoid homomorphism $\Delta_\Phi : \mathbf{SB}(W) \rightarrow \mathbb{Z}(\mathbf{B}(W))$.*

Proof. We need to check that the images of the generators of $\mathbf{SB}(W)$ in $\mathbb{Z}(\mathbf{B}(W))$ under Δ_Φ satisfy the defining relations of $\mathbf{SB}(W)$. The relations in (1) are obvious. The relations in (4) follow from the relations $\sigma_s \sigma_t = \sigma_t \sigma_s$ provided that $m_{s,t} = 2$, which, in particular, implies that $\sigma_s^i \sigma_t^j = \sigma_t^j \sigma_s^i$ provided that $m_{s,t} = 2$, for all $i, j \in \mathbb{Z}$.

The relations in (3) follow similarly from the corresponding relations for $\mathbf{B}(W)$. The latter, basically, say that σ_s commutes with $\sigma_t \sigma_s \dots \sigma_t$ (the latter word is of length $m_{s,t} - 1$) and hence σ_s^i commutes with $\sigma_t \sigma_s \dots \sigma_t$, for all $i \in \mathbb{Z}$.

Next, to check the relations in (2), we will use that the map $\Phi(K_-) : S \rightarrow \mathbb{Z}[x, x^{-1}]$ is constant on the connected components of the odd skeleton. The relations in $\mathbf{B}(W)$ say that moving σ_s past $\sigma_t \sigma_s \dots \sigma_s$ (of length $m_{s,t} - 1$) to the right, we get σ_t . This implies a similar claim for σ_s^i and σ_t^i , for any $i \in \mathbb{Z}$. Now, taking into account that the Laurent polynomials assigned to s and t by Φ are the same (since $m_{s,t}$ is odd), the necessary relation follows.

Finally, the relations in (5) follow from the fact that σ_s commutes with all its integer powers. \square

It is natural to call Δ_Φ a *generalized desingularization map*.

3.4. Generic generalized desingularization map. For a fixed $k \in \mathbb{Z}_{\geq 0}$, denote by \mathcal{A}_k the set of all possible choices of $\Phi : K \rightarrow \mathbb{Z}[x, x^{-1}]$ such that $\Phi(K_s)$ is of the form

$$a_{-k}x^{-k} + a_{-(k-1)}x^{-(k-1)} + \dots + a_{k-1}x^{k-1} + a_kx^k, \text{ for some } a_{-k}, \dots, a_k \in \mathbb{Z},$$

for any simple reflection s . The set \mathcal{A}_k has the obvious structure of a free \mathbb{Z} -module of rank $(2k+1)^{|K|}$.

As usual, we will say that a subset of \mathcal{A}_k is *generic* provided that this subset is dense in $\mathbb{C} \otimes_{\mathbb{Z}} \mathcal{A}_k$ with respect to Zariski topology. The next proposition shows that, in many cases, "almost all" choices for the generalized desingularization map lead to injective monoid homomorphisms.

Proposition 3. *Assume that the Coxeter group W belongs to one of the following cases:*

- (a) *the symmetric group (i.e. type A),*
- (b) *the dihedral group (i.e. type $I_2(n)$),*

(c) *the right-angled group (i.e. each $m_{s,t} \in \{2, \infty\}$, for all $s \neq t$).*

Then, for each $k \in \mathbb{Z}_{\geq 1}$, the set of all $\Phi \in \mathcal{A}_k$, for which Δ_Φ is injective, is generic.

Proof. We start by extending the scalars from \mathbb{Z} to \mathbb{C} . We consider the group algebra $\mathbb{C}(\mathbf{B}(W))$ and, for any choice of $\Phi : K \rightarrow \mathbb{C}[x, x^{-1}]$, the corresponding monoid homomorphism $\Delta_\Phi : \mathbf{SB}(W) \rightarrow \mathbb{C}(\mathbf{B}(W))$. Then we also have the corresponding set $\mathcal{A}_k^\mathbb{C}$ which is naturally isomorphic to $\mathbb{C} \otimes_{\mathbb{Z}} \mathcal{A}_k$ as a vector space.

Let $\Phi \in \mathcal{A}_k^\mathbb{C}$ and u and v be two different elements in $\mathbf{SB}(W)$. Then the fact that $\Delta_\Phi(u) = \Delta_\Phi(v)$ means that the coefficients of $\Delta_\Phi(u)$ with respect to the standard basis of $\mathbb{C}(\mathbf{B}(W))$ coincide with the corresponding coefficients of $\Delta_\Phi(v)$. Each such coefficient is, by construction, a polynomial in the coefficients of all $\Phi(K_s)$. This means that the condition $\Delta_\Phi(u) = \Delta_\Phi(v)$ defines a closed subset of $\mathcal{A}_k^\mathbb{C}$. Taking the intersection over all u and v , we obtain that the set \mathcal{B} of all $\Phi \in \mathcal{A}_k^\mathbb{C}$, for which Δ_Φ is not injective, is closed.

Under our assumptions on the type of W , the fact that the set \mathcal{B} does not coincide with $\mathcal{A}_k^\mathbb{C}$ follows from the results of [Pa04, GP05, Ea06a] because the classical desingularization map corresponds to the choice of a particular element in $\mathcal{A}_1 \subset \mathcal{A}_k \subset \mathcal{A}_k^\mathbb{C}$. Since \mathcal{A}_k is dense in $\mathcal{A}_k^\mathbb{C}$, it follows that $\mathcal{A}_k \setminus \mathcal{B}$ is dense in $\mathcal{A}_k^\mathbb{C}$ as well. This completes the proof. \square

3.5. Further generalizations. The content of this subsection is inspired by the results of [MaSte12, Subsection 2.2]. Let $R = (R, +, \cdot, 0, 1)$ be a commutative semiring (e.g. in the sense of [C-Z18, Subsection 2.1]). Then we can consider the corresponding semialgebra $R(\mathbf{B}(W))$ and, for any choice of $\Phi : K \rightarrow R[x, x^{-1}]$, from the obvious analogue of Proposition 2, we obtain the corresponding monoid homomorphism

$$\Delta_\Phi : \mathbf{SB}(W) \rightarrow R(\mathbf{B}(W)).$$

One particular choice of R leads to a very natural construction and interpretation. Consider as R the Boolean semiring $\mathbf{B} = \{0 := \text{false}, 1 := \text{true}\}$ with respect to the usual operations $+$ $:= \vee$ and \cdot $:= \wedge$. Then the semialgebra $\mathbf{B}(\mathbf{B}(W))$ can be naturally identified with the set of all finite subsets of $\mathbf{B}(W)$. Similarly, we can view $\mathbf{B}(\mathbb{Z})$ as the set of all finite subsets of \mathbb{Z} . In this case, the obvious analogue of Proposition 2 can be reformulated as follows:

Proposition 4. *For a fixed map $\Phi : K \rightarrow \mathbf{B}(\mathbb{Z})$, define*

$$\Delta_\Phi(\sigma_s) = \sigma_s \text{ and } \Delta_\Phi(\tau_s) = \{\sigma_s^i : i \in \Phi(K_s)\}, \text{ for } s \in S.$$

This assignment extends uniquely to a monoid homomorphism

$$\Delta_\Phi : \mathbf{SB}(W) \rightarrow \mathbf{B}(\mathbf{B}(W)).$$

We do not know whether any of these Δ_Φ 's is injective or not.

4. SINGULAR ARTIN MONOIDS AND HECKE ALGEBRAS

4.1. Hecke algebra. Let M be a Coxeter matrix and (W, S) the corresponding Coxeter system.

Consider the Laurent polynomial ring $\mathbb{A} := \mathbb{Z}[v, v^{-1}]$ and the corresponding group algebra $\mathbb{A}(\mathbf{B}(W))$. Denote by $\mathbf{H} = \mathbf{H}(W)$ the quotient of $\mathbb{A}(\mathbf{B}(W))$ modulo the ideal generated by the following relations:

$$(\sigma_s - v)(\sigma_s + v^{-1}) = 0, \quad \text{for } s \in S.$$

Traditionally, the image of σ_s in \mathbf{H} is denoted by H_s .

For $w \in W$ with a fixed reduced expression $w = s_1 s_2 \dots s_k$, set $H_w := H_{s_1} H_{s_2} \dots H_{s_k}$. Then $\{H_w : w \in W\}$ is an \mathbb{A} -basis of \mathbf{H} , called the *standard basis*.

Note that each H_s is invertible in \mathbf{H} with inverse $H_s^{-1} = H_s - (v - v^{-1})H_e$. The algebra \mathbf{H} admits a unique involution $\overline{}$, called the *bar-involution*, satisfying $\overline{H_s} = H_s^{-1}$ and $\overline{\overline{v}} = v^{-1}$. Furthermore, \mathbf{H} has a unique basis $\{\underline{H}_w : w \in W\}$ such that, for $w \in W$, we have:

- $\underline{H}_w \in H_w + \sum_{x \in W} v\mathbb{Z}[v]H_x$;
- $\overline{\underline{H}_w} = \underline{H}_w$.

This basis is called the *Kazhdan-Lusztig (KL) basis*, see [KL79] and note that we are using the normalization of [So07].

4.2. Monoid homomorphisms from $\mathbf{SB}(W)$ to \mathbf{H} . Fix a map $\Phi : K \rightarrow \mathbb{A}[x, x^{-1}]$. Define

$$\Upsilon_{\Phi}(\sigma_s) = H_s \text{ and } \Upsilon_{\Phi}(\tau_s) = \Phi(K_s)[H_s], \text{ for } s \in S.$$

Corollary 5. *The above assignment extends uniquely to a monoid homomorphism*

$$\Delta_{\Phi} : \mathbf{SB}(W) \rightarrow \mathbf{H}(W).$$

A very special case of the above construction is the case when Φ sends each K_s to $v + x$. In this case $\Upsilon_{\Phi}(\tau_s) = \underline{H}_s$.

We note that the restriction of Δ_{Φ} to $\mathbf{B}(W)$ is not faithful, in general.

5. DIAGRAM ALGEBRAS AND SINGULAR ARTIN MONOIDS

In this section we investigate relations between singular Artin monoids and various families of diagram algebras.

5.1. The group algebra of the Coxeter group.

5.1.1. Connection via a generalized desingularization map. Let M be a Coxeter matrix with the associated Coxeter group W , the Artin braid group $\mathbf{B}(W)$ and the singular Artin monoid $\mathbf{SB}(W)$. For a fixed map $\Phi : K \rightarrow \mathbb{Z}[x, x^{-1}]$, we have the monoid homomorphism

$$\mathbf{SB}(W) \rightarrow \mathbb{Z}(\mathbf{B}(W)) \rightarrow \mathbb{Z}(W),$$

where the first map is Δ_{Φ} and the second map is the natural epimorphism. This induces an algebra epimorphism $\overline{\Delta}_{\Phi} : \mathbb{Z}(\mathbf{SB}(W)) \rightarrow \mathbb{Z}(W)$.

Since $\overline{\Delta}_{\Phi}(\sigma_s) = \overline{\Delta}_{\Phi}(\sigma_s^{-1})$, the number of independent parameters in this construction is $2 \cdot |K|$ as, for each element of K , without loss of generality, we may assume that its image under Φ is of the form $a + bx$.

5.1.2. Pulling back Coxeter group modules. Pulling back via $\overline{\Delta}_{\Phi}$ defines a functor from W -mod to $\mathbf{SB}(W)$ -mod. Generically, this gives a $2 \cdot |K|$ -parametric family of lifts to $\mathbf{SB}(W)$ of simple W -modules.

We note that, for some modules, the number of essential parameters might be lower. For example, this happens if the action of s and e on such a module are linearly dependent. This is true, for example, for the trivial and the sign W -modules. In these cases we only have a $|K|$ -parametric family of different lifts.

5.2. Binary relations.

the matrices which are invariant under the 180° rotation with respect to the center of the matrix.

In terms of Γ_ρ , if both parts of Γ_ρ are given by $\overline{n}, \overline{n-1}, \dots, n$ left-to-right, then $\rho \in \mathbf{Bin}_n^B$ if and only if Γ_ρ is invariant under the flip which swaps i and \bar{i} in both parts.

The group of invertible elements in \mathbf{Bin}_n^B is naturally isomorphic to the group $\{\pm 1\} \wr S_n$ of signed permutations. The latter is a standard realization of the Coxeter group of type B_n .

5.2.4. *Map from a type B singular Artin monoid.* Let W be the Coxeter group corresponding to the following (type B) Coxeter graph:

$$s_0 \xrightarrow{4} s_1 \text{ --- } s_2 \text{ --- } \dots \text{ --- } s_{n-1}$$

Denote by \tilde{s}_0 the element of \mathbf{Bin}_n^B which is obtained from the equality relation by removing $(1, 1)$ and $(\bar{1}, \bar{1})$ and adding $(1, \bar{1})$, $(\bar{1}, 1)$. In other words, this is the transposition of 1 and $\bar{1}$. Further, for all $i = 1, 2, \dots, n-1$, we denote by \tilde{s}_i the transposition of i and $i+1$ and, simultaneously, of \bar{i} and $\bar{i}+1$. Then $s_i \mapsto \tilde{s}_i$, for $i = 1, 2, \dots, n-1$, gives rise to a natural monomorphism from S_n to $\{\pm 1\} \wr S_n$.

For $i = 1, 2, \dots, n-1$, we denote by $\tilde{\tilde{s}}_i$ the element of \mathbf{Bin}_n^B which is obtained from the equality relation by adding $(i, i+1)$, $(i+1, i)$, $(\bar{i}, \bar{i}+1)$ and $(\bar{i}+1, \bar{i})$. We also denote by $\tilde{\tilde{s}}_0$ the element of \mathbf{Bin}_n^B which is obtained from the equality relation by adding $(1, \bar{1})$ and $(\bar{1}, 1)$. For example, here are the graphs of the elements \tilde{s}_0 and $\tilde{\tilde{s}}_1$ in \mathbf{Bin}_2^B :

$$\begin{array}{cccc} \bar{2} & \bar{1} & 1 & 2 \\ \bar{1} & | \times | & | & \bar{1} \\ \bar{2} & \bar{1} & 1 & 2 \end{array} \qquad \begin{array}{cccc} \bar{2} & \bar{1} & 1 & 2 \\ \bar{1} \times \bar{1} & | \times | & & \\ \bar{2} & \bar{1} & 1 & 2 \end{array}$$

Proposition 7. *There is a unique homomorphism $\eta : \mathbf{SB}(W) \rightarrow \mathbf{Bin}_n^B$ such that*

$$\eta(\sigma_{s_i}) = \tilde{s}_i \text{ and } \eta(\tau_{s_i}) = \tilde{\tilde{s}}_i, \text{ for } i = 0, 1, \dots, n-1.$$

Proof. This follows from the straightforward verification of the analogues of the relations (1), (2), (3), (4) and (5) for the elements \tilde{s}_i and $\tilde{\tilde{s}}_i$. \square

The submonoid of \mathbf{Bin}_n^B generated by the elements \tilde{s}_i is a natural type B analogue of the *double Catalan monoid*, see also [MaSte12, Subsection 6.2].

5.3. Dual symmetric inverse monoid.

5.3.1. *Dual symmetric inverse monoid and its factorizable submonoid.* For a positive integer n , consider the symmetric inverse monoid \mathbf{IS}_n , defined as the set of all bijections between subsets of \underline{n} . We will talk about \mathbf{IS}_n in more detail in Subsection 5.4. In this subsection we will focus on the dual object, called the *dual symmetric inverse monoid* \mathbf{I}_n^* which is defined as the set of all bijections between quotients of $\underline{n} := \{1, 2, \dots, n\}$, see [FL06]. The reason why we start with \mathbf{I}_n^* will be explained in Lemma 15.

We can view each element of \mathbf{I}_n^* as a binary relation on \underline{n} in the natural way. That is, given two set partitions

$$A_1 \amalg A_2 \amalg \dots \amalg A_k = \underline{n} = B_1 \amalg B_2 \amalg \dots \amalg B_k$$

and $\sigma \in S_k$, the (unique) element ξ of \mathbf{I}_n^* which sends A_i to $B_{\sigma(i)}$, for $i = 1, 2, \dots, k$, corresponds to the binary relation

$$\rho_\xi := \bigcup_{i=1}^k (B_{\sigma(i)} \times A_i).$$

Note that ρ_ξ is essential. The bipartite graph Γ_{ρ_ξ} has the property that each connected component of Γ_{ρ_ξ} is a complete bipartite graph. Conversely, given an essential binary relation ρ such that each connected component of Γ_ρ is a complete bipartite graph, there exists $\xi \in \mathbf{I}_n^*$ such that $\rho = \rho_\xi$. In fact, in Proposition 8 below we show that \mathbf{I}_n^* is a quotient of $\mathbf{Bin}_n^{\text{ess}}$.

For $\rho \in \mathbf{Bin}_n$, denote by $\bar{\rho}$ the unique element in \mathbf{Bin}_n such that $\Gamma_{\bar{\rho}}$ is obtained from Γ_ρ by completing each connected component of Γ_ρ to a complete bipartite graph.

Proposition 8.

- (a) Call ρ and ρ' in \mathbf{Bin}_n equivalent provided that $\bar{\rho} = \bar{\rho}'$. This equivalence relation is a congruence on \mathbf{Bin}_n .
- (b) The subsemigroup $\mathbf{Bin}_n^{\text{ess}}$ is a union of congruence classes.
- (c) The quotient of $\mathbf{Bin}_n^{\text{ess}}$ modulo this congruence is isomorphic to \mathbf{I}_n^* via the map which sends $\xi \in \mathbf{I}_n^*$ to the equivalence class of ρ_ξ .

Proof. Equivalence of ρ and ρ' can be alternatively described as follows: two vertices of Γ_ρ belong to the same connected component if and only if the corresponding vertices of $\Gamma_{\rho'}$ belong to the same connected component.

Let now ρ_1 and ρ_2 be two elements in \mathbf{Bin}_n . Then the product $\rho_1\rho_2$ can be described as follows in terms of Γ_{ρ_1} and Γ_{ρ_2} : the graph $\Gamma_{\rho_1\rho_2}$ is obtained by first taking the union of Γ_{ρ_1} and Γ_{ρ_2} under the assumption that the lower vertices of Γ_{ρ_2} are identified with the corresponding upper vertices of Γ_{ρ_1} and then removing these common vertices making the following adjustment of edges whenever possible (here the red vertex is removed):



Taking the union of connected graphs with a common vertex produces a connected graph. This implies the following: assume that ρ_1 and ρ'_1 are equivalent and ρ_2 and ρ'_2 are equivalent. Then, for any fixed vertex, while the connected components of this vertex in $\Gamma_{\rho_1\rho_2}$ and $\Gamma_{\rho'_1\rho'_2}$ might be non-isomorphic as graphs, the sets of vertices in these two components coincide. This means exactly that $\rho_1\rho_2$ and $\rho'_1\rho'_2$ are equivalent. This shows that our equivalence relation is a congruence, proving Claim (a).

Claim (b) follows directly from the definitions. To prove Claim (c), we note that the map which sends $\xi \in \mathbf{I}_n^*$ to the equivalence class of ρ_ξ in \mathbf{Bin}_n is, clearly, injective. Moreover, it is bijective if we restrict to $\mathbf{Bin}_n^{\text{ess}}$ modulo the congruence in Claim (a). The fact that it is a homomorphism of semigroups follows by comparing the definitions of multiplications in \mathbf{I}_n^* and $\mathbf{Bin}_n^{\text{ess}}$. \square

Remark 9. The quotient of \mathbf{Bin}_n modulo the congruence given by Proposition 8(a) is the inverse semigroup of all bijections between quotients of subsets of \underline{n} (alternatively, between subsets of quotients of \underline{n}). This semigroup was studied in [KM11, KMU15].

We denote by $\pi : \mathbf{Bin}_n^{\text{ess}} \rightarrow \mathbf{I}_n^*$ the surjective map given by Proposition 8.

The monoid \mathbf{I}_n^* is an inverse semigroup. The semigroup \mathbf{I}_n^* contains a subsemigroup, denoted \mathbf{F}_n^* , defined by the condition that $|B_{\sigma(i)}| = |A_i|$, for all i , in the above notation. The subsemigroup \mathbf{F}_n^* is usually called the *maximal factorizable submonoid* of \mathbf{I}_n^* . The meaning of this is the following: The group of units of \mathbf{I}_n^* is the symmetric group S_n . As usual, we denote by s_i the elementary transposition $(i, i+1) \in S_n$, for $i = 1, 2, \dots, n-1$. The idempotents of \mathbf{I}_n^* are in a natural bijection with the equivalence relations on (alternatively, the set partitions of) \underline{n} . In the above notation, to get an idempotent, we take $A_i = B_i$, for all i , and $\sigma = e \in S_k$. The set $E(\mathbf{I}_n^*)$ of all idempotents of \mathbf{I}_n^* is a commutative semigroup whose operation can be interpreted as “taking the minimal equivalence relation which contains the two given equivalence relations”. Note that $E(\mathbf{I}_n^*)$ is closed under the conjugation by elements in S_n . For an equivalence relation ρ on \underline{n} , we denote by ξ_ρ the corresponding idempotent in \mathbf{I}_n^* .

The subsemigroup \mathbf{F}_n^* consists of all elements of the form $\sigma\xi$, where $\sigma \in S_n$ and $\xi \in E(\mathbf{I}_n^*)$. That is the *factorization* mentioned in the name. We note that $\sigma\xi = \sigma'\xi'$ does not imply $\sigma = \sigma'$, in general (but it does imply $\xi = \xi'$). The product of two elements in \mathbf{F}_n^* can be computed as follows:

$$(\sigma'\xi')(\sigma\xi) = (\sigma'\sigma)((\sigma^{-1}\xi'\sigma)\xi).$$

The subsemigroup \mathbf{F}_n^* is an inverse semigroup as well.

For $i \neq j$ in $\{1, 2, \dots, n\}$, denote by $\xi_{\{i,j\}}$ the idempotent in \mathbf{F}_n^* corresponding to the equivalence relation with equivalence classes $\{i, j\}$ and $\{s\}$, where $s \neq i, j$. It is easy to check that \mathbf{F}_n^* is generated by S_n and any $\xi_{\{i,j\}}$. For $i = 1, 2, \dots, n-1$, set $\xi_i := \xi_{\{i, i+1\}}$. Note that $\xi_i = \pi(s_i)$.

5.3.2. Map from singular braid monoid.

Proposition 10. *There is a unique homomorphism $\lambda : \mathbf{SB}(S_n) \rightarrow \mathbf{F}_n^*$ such that*

$$\lambda(\sigma_{s_i}) = s_i \text{ and } \lambda(\tau_{s_i}) = \xi_i, \text{ for } i = 1, 2, \dots, n-1.$$

This homomorphism is surjective.

Proof. The existence claim follows from the straightforward verification of the analogues of the relations (1), (2), (3), (4) and (5) for the elements s_i and ξ_i . The uniqueness claim follows from the fact that σ_{s_i} and τ_{s_i} generate $\mathbf{SB}(S_n)$. Surjectivity follows by combining the fact that the s_i 'th generate S_n with the fact that S_n and any ξ_i generate \mathbf{F}_n^* . \square

We note that $\lambda = \pi \circ \bar{\eta}$.

5.3.3. Presentation for \mathbf{F}_n^* .

Proposition 11. *The monoid \mathbf{F}_n^* is generated by the elements s_i and ξ_i , where we have $i = 1, 2, \dots, n-1$, subject to the relations (1), (2), (3), (4) and (5) (for s_i instead of σ_{s_i} and ξ_i instead of τ_{s_i}) and, additionally, the relations*

$$(7) \quad s_i^2 = e, \quad \text{for } i = 1, 2, \dots, n-1;$$

$$(8) \quad \xi_i^2 = \xi_i, \quad \text{for } i = 1, 2, \dots, n-1;$$

$$(9) \quad \xi_i \xi_{i+1} = \xi_{i+1} \xi_i, \quad \text{for } i = 1, 2, \dots, n-2;$$

$$(10) \quad \xi_i s_i = \xi_i, \quad \text{for } i = 1, 2, \dots, n-1.$$

The associative algebra defined by our presentation was studied in [Ko06] (however, the relation to the diagrammatic realization for this algebra is not explained with all details in this paper). A slight variation of this presentation is given in [Fi03], see also [EEF05] for a general approach to presentation of factorizable inverse monoids. Our proof is based on a direct reduction to the presentation in [Fi03, Theorem 3]. Our presentation can also be deduced from [Ea06b, Theorem 6.4 and Remark 6.5].

Proof. We show that our relations imply the relations in [Fi03, Theorem 3]. We note that, in [Fi03, Theorem 3], the generating set for \mathbf{F}_n^* is smaller than ours, namely, it consists of the Coxeter generators for S_n and ξ_1 . It follows from the relations (2) and (3) that all other ξ_i are conjugates of ξ_1 under the action of the symmetric group (and hence are redundant as generators).

Relations (S) in [Fi03, Theorem 3] are just the usual Coxeter relations for the symmetric group. Relation (F2) in [Fi03, Theorem 3] is our relation (8), Relation (F3) in [Fi03, Theorem 3] is a special case of our relation (10) combined with (5), Relation (F4) in [Fi03, Theorem 3] is a special case of our relation (3), Relation (F5) in [Fi03, Theorem 3] is a special case of our relation (9), and, finally, Relation (F6) in [Fi03, Theorem 3] is a special case of our relation (4). Therefore our claim follows from [Fi03, Theorem 3]. \square

5.3.4. *Type B dual symmetric inverse monoid and its factorizable submonoid.* The congruence on \mathbf{Bin}_n described in Proposition 8 restricts to the monoid \mathbf{Bin}_n^B . It is natural to call the quotient of the submonoid

$$\mathbf{Bin}_n^{B,\text{ess}} := \mathbf{Bin}_n^B \cap \mathbf{Bin}_n^{(2),\text{ess}}$$

of \mathbf{Bin}_n^B modulo this restricted congruence the *type B dual symmetric inverse monoid*. We denote it by \mathbf{IB}_n^* . The maximal factorizable submonoid of \mathbf{IB}_n^* is denoted \mathbf{FB}_n^* .

Note that the idempotents of \mathbf{IB}_n^* are the equivalence relations on $\overline{\underline{n}} \cup \underline{n}$ which are invariant the map $i \mapsto \bar{i}$. For an equivalence class X of such an equivalence relation, there are two possibilities:

- $|X \cap \{i, \bar{i}\}| \leq 1$, for all $i \in \underline{n}$, and, in this case, \overline{X} is another equivalence class;
- $|X \cap \{i, \bar{i}\}| = 2$, for some $i \in \underline{n}$, and, in this case, $\overline{X} = X$.

Such an equivalence relation can be described by a tuple (ρ, Y, f) , where

- ρ is an equivalence relation on \underline{n} ,
- Y is the union of some of the equivalence classes of ρ ,
- $f : Y \rightarrow \{\pm\}$ is a function,

modulo the equivalence relation \sim defined by $(\rho, Y, f) \sim (\rho, Y, f')$ provided that f differs from f' by changing the sign on the union of some of the equivalence classes of ρ inside Y . Here Y consists of the absolute values of all elements appearing in those classes $X \subset (\overline{\underline{n}} \cup \underline{n})$ for which $|X \cap \{i, \bar{i}\}| \leq 1$, for all $i \in \underline{n}$. For such an X , the restrictions of the functions f and $-f$ to X determine X and \overline{X} , up to swapping these two sets. The equivalence relation \sim compensates for this swapping.

For $i = 1, 2, \dots, n-1$, we denote by $\tilde{\xi}_i$ the idempotent of \mathbf{IB}_n^* corresponding to the tuple (ρ, \underline{n}, f) , where ρ is the equivalence relation on \underline{n} which has only one non-singleton part, namely $\{i, i+1\}$, and $f(i) = +$, for all $i \in \underline{n}$. We denote by $\tilde{\xi}_0$ the idempotent of \mathbf{IB}_n^* corresponding to the tuple $(\rho, \underline{n} \setminus \{1\}, f)$, where ρ is the equality relation and $f(i) = +$, for all $i \in \underline{n} \setminus \{1\}$.

5.3.5. *Map from a type B singular braid monoid.* Let W be the Coxeter group corresponding to the following (type B) Coxeter graph:

$$s_0 \xrightarrow{4} s_1 \text{ --- } s_2 \text{ --- } \dots \text{ --- } s_{n-1}$$

The obvious analogue of the map λ from Proposition 10 gives rise to a natural epimorphism from $\mathbf{SB}(W)$, where W is of type B_n onto \mathbf{FB}_n^* .

Proposition 12. *There is a unique homomorphism $\lambda : \mathbf{SB}(W) \rightarrow \mathbf{FB}_n^*$, such that*

$$\lambda(\sigma_{s_i}) = \tilde{s}_i \text{ and } \lambda(\tau_{s_i}) = \tilde{\xi}_i, \text{ for } i = 0, 1, \dots, n-1.$$

This homomorphism is surjective.

Proof. Existence follows from the straightforward verification of the analogues of the relations (1), (2), (3), (4) and (5) for the elements \tilde{s}_i and $\tilde{\xi}_i$. Uniqueness is due to the fact that the homomorphism is defined on the generators of $\mathbf{SB}(W)$. Surjectivity follows from the fact that the images of the generators of $\mathbf{SB}(W)$ generate \mathbf{FB}_n^* . \square

Remark 13. We note that the homomorphism λ is the composition of η followed by the natural projection from $\mathbf{Bin}_n^{B, \text{ess}}$ onto \mathbf{FB}_n^* .

5.3.6. *Presentation for \mathbf{FB}_n^* .*

Proposition 14. *The monoid \mathbf{FB}_n^* is generated by the elements \tilde{s}_i and $\tilde{\xi}_i$, where we have $i = 0, 1, \dots, n-1$, subject to*

- the (analogues of the) relations (1), (2), (3), (4) and (5);
- the (analogues of the) relations (7), (8), (9) and (10), for all i including 0;
- the additional relations

$$(11) \quad \tilde{s}_1 \tilde{\xi}_0 \tilde{s}_1 \tilde{\xi}_0 = \tilde{\xi}_0 \tilde{s}_1 \tilde{\xi}_0 \tilde{s}_1,$$

$$(12) \quad \tilde{s}_0 \tilde{\xi}_1 \tilde{s}_0 \tilde{\xi}_1 = \tilde{\xi}_1 \tilde{s}_0 \tilde{\xi}_1 \tilde{s}_0,$$

$$(13) \quad \tilde{\xi}_0 \tilde{\xi}_1 = \tilde{s}_0 \tilde{\xi}_1 \tilde{s}_0 \tilde{\xi}_1.$$

Proof. Below we give a very detailed sketch of the idea of the proof, leaving it for the reader to verify most of the technical details.

It is easy to check that the generators of \mathbf{FB}_n^* satisfy all the relations in the formulation. Denote by Q the monoid with the presentation described in the formulation generated by x_i (instead of \tilde{s}_i) and \mathbf{x}_i (instead of $\tilde{\xi}_i$). We have the obvious surjection $\psi : Q \twoheadrightarrow \mathbf{FB}_n^*$ which we want to prove is an isomorphism. For this we need to prove that $|Q| \leq |\mathbf{FB}_n^*|$.

Due to (1) and (7), the group G of invertible elements in Q is isomorphic to the group of signed permutations on \underline{n} (i.e. the Coxeter group of type B_n). Denote by T the set of all G -conjugates of all \mathbf{x}_i , where $i = 0, 1, 2, \dots, n-1$.

By (10), (2), (3) and (5), the G -conjugate stabilizer of \mathbf{x}_0 contains both x_0 and all signed permutations on $\{2, 3, \dots, n\}$. Hence the number of the G -conjugates of \mathbf{x}_0 is at most $\frac{2^n \cdot n!}{2 \cdot 2^{n-1} (n-1)!} = n$. Since the number of the G -conjugates of $\tilde{\xi}_0$ is exactly n , we obtain that the number of the G -conjugates of \mathbf{x}_0 is exactly n . For $i = 1, \dots, n$, we denote by \mathbf{t}_i the conjugate of \mathbf{x}_0 by the transposition $(1, i)$. Note that $\mathbf{t}_1 = \mathbf{x}_0$.

Similarly, the G -conjugate stabilizer of \mathbf{x}_1 contains both the elements x_1 and $x_0 x_1 x_0$, their product $x_1 x_0 x_1 x_0$, and, furthermore, all signed permutations on $\{3, 4, \dots, n\}$.

Hence the number of the G -conjugates of x_1 is at most $\frac{2^n \cdot n!}{4 \cdot 2^{n-2} (n-2)!} = n(n-1)$. Since the number of the G -conjugates of ξ_1 is exactly $n(n-1)$, we obtain that the number of the G -conjugates of x_1 is exactly $n(n-1)$. For different $i, j \in \{1, \dots, n\}$, we denote by $\mathfrak{t}_{i,j}$ the conjugate of x_1 by $(1, i)(2, j)$. We also denote by $\bar{\mathfrak{t}}_{i,j}$ the conjugate of x_1 by $(1, \bar{i})(2, j)$. From (10), it follows that $\mathfrak{t}_{i,j} = \mathfrak{t}_{j,i}$ and $\bar{\mathfrak{t}}_{i,j} = \bar{\mathfrak{t}}_{j,i}$. Also, we have $x_i = \mathfrak{t}_{i,i+1}$, for all $i > 1$, using (2) and (3).

From (11), (12), (3), (4) and (9), it follows that all elements in T commute with each other. From (8), it also follows that all these elements are idempotent.

We note the following property (involving (11)):

$$(14) \quad x_1 \mathfrak{t}_1 x_1 \mathfrak{t}_{1,2} = \mathfrak{t}_1 \mathfrak{t}_{1,2}.$$

Indeed, using the already established commutativity, we have

$$\begin{aligned} x_1 \mathfrak{t}_1 x_1 \mathfrak{t}_{1,2} &\stackrel{(10)}{=} x_1 \mathfrak{t}_1 \mathfrak{t}_{1,2} \\ &= x_1 \mathfrak{t}_{1,2} \mathfrak{t}_1 \\ &\stackrel{(10)}{=} \mathfrak{t}_{1,2} \mathfrak{t}_1 \\ &= \mathfrak{t}_1 \mathfrak{t}_{1,2}. \end{aligned}$$

Also, note that $x_1 \mathfrak{t}_1 x_1 = \mathfrak{t}_2$. This implies the following extension of (14):

$$(15) \quad \mathfrak{t}_1 \mathfrak{t}_{1,2} = \mathfrak{t}_2 \mathfrak{t}_{1,2} = \mathfrak{t}_{1,2} \mathfrak{t}_1 = \mathfrak{t}_{1,2} \mathfrak{t}_2.$$

Now we can classify the elements in the monoid generated by T . Let ω be a product of elements in T . Due to commutativity and idempotency for elements of T , we may assume that each factor appearing in ω appears there only once. We associate to ω an unoriented graph Λ_ω with the set of vertices $\bar{n} \cup \underline{n}$ and the edges defined as follows:

- For each factor \mathfrak{t}_i appearing in ω , we connect i and \bar{i} by an edge in Λ_ω .
- For each factor $\mathfrak{t}_{i,j}$, where $i \neq j$, appearing in ω , we connect i and j by an edge in Λ_ω and we also connect \bar{i} with \bar{j} by an edge in Λ_ω .
- For each factor $\bar{\mathfrak{t}}_{i,j}$, where $i \neq j$, appearing in ω , we connect i and \bar{j} by an edge in Λ_ω and we also connect j with \bar{i} by an edge in Λ_ω .

Let $\tilde{\Lambda}_\omega$ be the graph obtained from Λ_ω by extending each connected component to a complete graph on the set of vertices of this connected component.

From the relations (11), (12), (13) and (15), it follows that $\omega = \omega'$ if and only if $\tilde{\Lambda}_\omega = \tilde{\Lambda}_{\omega'}$. Here is an illustration of how this works by an example. In terms of the graphs, the relation (13) says that the following two graphs define the same element of Q :

$$\begin{array}{c} \bar{1} - 1 \\ | \quad | \\ \bar{2} \quad 2 \end{array} \quad \text{and} \quad \begin{array}{c} \bar{1} \quad 1 \\ | \quad | \\ \bar{2} \quad 2 \end{array} \times$$

Using (9) and (10), it then follows that the following two graphs define the same element of Q :

$$\begin{array}{c} \bar{1} \quad 1 \\ | \quad | \\ \bar{2} - 2 \end{array} \quad \text{and} \quad \begin{array}{c} \bar{1} \quad 1 \\ | \quad | \\ \bar{2} \quad 2 \end{array} \times$$

Now, using the idempotency of all factors, it follows that the element of Q defined by any of the above graphs is equal to the element of Q defined by the full graph on $\{1, 2, \bar{1}, \bar{2}\}$. The general case follows from this local example inductively.

Combining the fact that $\omega = \omega'$ if and only if $\tilde{\Lambda}_\omega = \tilde{\Lambda}_{\omega'}$ with the fact that full symmetric (under $i \mapsto \bar{i}$) graphs on $\bar{n} \cup \underline{n}$ whose connected components are complete graphs classify all idempotents in \mathbf{FB}_n^* , we obtain that the restriction of ψ to the submonoid $\langle T \rangle$ generated by T in Q is injective.

Finally, let ω be a product of elements in T with the associated graph $\tilde{\Lambda}_\omega$. Each edge of $\tilde{\Lambda}_\omega$ corresponds, using (2), (3), (5) and (10), to a reflection in G which stabilizes ω under the left multiplication. It is easy to check that the $\psi(G)$ -stabilizer of $\psi(\omega)$ with respect to the left multiplication is generated by such reflections. This implies that the cardinality of the stabilizer of ω in G with respect to the left multiplication is at least as large as that of the corresponding stabilizer of $\psi(\omega)$ in $\psi(G)$. Combining the facts that both Q and \mathbf{FB}_n^* are factorizable and ψ is bijective on both G and $\langle T \rangle$, it follows that $|Q| \leq |\mathbf{FB}_n^*|$, completing the proof. \square

5.4. Symmetric inverse semigroup (a.k.a. the rook monoid).

5.4.1. *Symmetric inverse semigroup.* For $n \in \mathbb{Z}_{\geq 1}$, consider the corresponding symmetric inverse semigroup \mathbf{IS}_n on \underline{n} . The elements of \mathbf{IS}_n are all bijections between subsets of \underline{n} . The semigroup operation is given by the usual composition of partial maps, see [GM09, Section 2.1] for details. The symmetric group S_n is the group of invertible elements in \mathbf{IS}_n . The semigroup \mathbf{IS}_n is also known as the rook monoid.

Let $\alpha \in \mathbf{IS}_n$ be a bijection from $X \subset \underline{n}$ to $Y \subset \underline{n}$. The cardinality of X is called the rank of α , denoted $\text{rank}(\alpha)$. The subset $\widetilde{\mathbf{IS}}_n$ of \mathbf{IS}_n consisting of all elements whose rank is different from $n - 1$ is a subsemigroup (containing S_n).

For a subset $X \subset \underline{n}$, the semigroup \mathbf{IS}_n contains the identity map Id_X on X . We will use the notation $\varepsilon_X = \text{Id}_{\underline{n} \setminus X}$. The element ε_X is an idempotent and each idempotent of \mathbf{IS}_n is of this form.

The following lemma describes a close relation between \mathbf{F}_n^* and \mathbf{IS}_n .

Lemma 15. *Restricting an element of \mathbf{F}_n^* to singletons defines a homomorphism from \mathbf{F}_n^* to \mathbf{IS}_n which we denote by v . The image of v coincides with $\widetilde{\mathbf{IS}}_n$.*

Proof. Let α and β be two elements of \mathbf{F}_n^* . If β sends some singleton x to a singleton y and α sends y to a singleton z , then $\alpha\beta$ sends x to z .

If β sends some singleton x to a singleton y , however y is not a singleton for α but is, rather, contained in a non-singleton class Y , then α sends Y to a non-singleton Z (as $|\alpha(Y)| = |Y|$) and therefore $\alpha\beta$ sends some non-singleton class containing x to a non-singleton class containing Y . Similar arguments, applied to other possibilities of classes for α and β , show that the map in the formulation is a homomorphism of monoids.

The image of this map, clearly, contains S_n and $\varepsilon_{\{1,2\}}$ and these generate $\widetilde{\mathbf{IS}}_n$. Therefore the image of our homomorphism coincides with $\widetilde{\mathbf{IS}}_n$. \square

5.4.2. Map from singular braid monoid.

Proposition 16. *There is a unique homomorphism $\varphi : \mathbf{SB}(S_n) \rightarrow \mathbf{IS}_n$ such that*

$$\varphi(\sigma_{s_i}) = s_i \text{ and } \varphi(\tau_{s_i}) = \varepsilon_{\{i,i+1\}}, \text{ for } i = 1, 2, \dots, n-1.$$

The image of φ coincides with $\widetilde{\mathbf{IS}}_n$.

Proof. We note that $\varphi = \nu \circ \lambda$ and hence the claim follows from Proposition 10 and Lemma 15. \square

The statement of Proposition 16 admits a slight generalization when working with algebras. For example, for a fixed commutative ring \mathbb{k} and $a \in \mathbb{k}$, there is a unique homomorphism $\varphi_a : \mathbf{SB}(S_n) \rightarrow \mathbb{k}[\mathbf{IS}_n]$ such that

$$\varphi_a(\sigma_{s_i}) = s_i \text{ and } \varphi_a(\tau_{s_i}) = a\varepsilon_{\{i,i+1\}}, \text{ for } i = 1, 2, \dots, n-1.$$

5.4.3. *Presentation in type A.* We have the following presentation for the monoid $\widetilde{\mathbf{IS}}_n$.

Proposition 17. *The monoid $\widetilde{\mathbf{IS}}_n$ is generated by the elements s_i and $\varepsilon_{\{i,i+1\}}$, where $i = 1, 2, \dots, n-1$, subject to the relations (1), (2), (3), (4) and (5) (for s_i instead of σ_{s_i} and $\varepsilon_{\{i,i+1\}}$ instead of τ_{s_i}), the relations (7), (8), (9) and (10) (for $\varepsilon_{\{i,i+1\}}$ instead of $\xi_{\{i,i+1\}}$) and, additionally, the relation*

$$(16) \quad \varepsilon_{\{1,2\}}\varepsilon_{\{3,4\}} = \varepsilon_{\{1,2\}}\varepsilon_{\{2,3\}}\varepsilon_{\{3,4\}}.$$

Proof. It is clear that s_i and $\varepsilon_{\{i,i+1\}}$ satisfy all these relations, so we only need to check that the monoid defined by this presentation is exactly $\widetilde{\mathbf{IS}}_n$. Since $\varphi = \nu \circ \lambda$, from Proposition 11 we only need to prove that the kernel of ν is generated by the relation given by (16). Denote by Q the quotient of \mathbf{F}_n^* modulo the congruence generated by (16). Then we have the quotient map $Q \rightarrow \widetilde{\mathbf{IS}}_n$ which is surjective by Lemma 15. We need to prove that $|Q| \leq |\widetilde{\mathbf{IS}}_n|$.

If ρ is an equivalence relation on \underline{n} , then the corresponding idempotent $\xi_\rho \in \mathbf{I}_n^*$ (see Subsection 5.3.1) is the product of all $\xi_{\{i,j\}}$, where i and j belong to the same class of ρ . Using the S_n -conjugates of the (16), allows us to merge non-singleton classes of equivalence relations. In particular, we see that the images of two equivalence relations ρ and ρ' in Q coincide if ρ and ρ' have the same singletons. Comparing the outcomes with $E(\widetilde{\mathbf{IS}}_n)$, we get that the image of $E(\mathbf{F}_n^*)$ in Q is in a natural bijection with $E(\widetilde{\mathbf{IS}}_n)$.

Each element of Q can be written as $\sigma\xi$, where $\sigma \in S_n$ and ξ is in the image of $E(\mathbf{F}_n^*)$. From the relation (10) it follows that, for fixed ξ with k singletons, the number of different elements of the form $\sigma\xi$ is at most $\frac{n!}{(n-k)!}$. Adding this up over all ξ , implies $|Q| \leq |\widetilde{\mathbf{IS}}_n|$. This completes the proof. \square

5.4.4. *Signed symmetric inverse semigroup.* For a positive integer n , consider the symmetric inverse semigroup \mathbf{IS}_X , where $X = \{-n, -n+1, \dots, -1, 1, \dots, n-1, n\}$. There is the obvious automorphism $\bar{\omega}$ of \mathbf{IS}_X induced by the endomorphism ω of X which changes the sign. The set of all elements in \mathbf{IS}_X that are invariant under $\bar{\omega}$ is a subsemigroup denoted \mathbf{SIS}_n , the *signed symmetric inverse semigroup*.

The semigroup \mathbf{IS}_n can be realized as the monoid of all $n \times n$ matrices which have the property that each row and each column of the matrix contains at most one non-zero element, and this non-zero element equals 1. Such a matrix can be thought of as the indicator matrix for a placement of rooks which do not attack each other on an $n \times n$ square board. This is the justification for the alias *rook monoid*.

Similarly, the semigroup \mathbf{SIS}_n can be realized as the monoid of all $n \times n$ matrices which have the property that each row and each column of the matrix contains at

most one non-zero element, and this non-zero element equals ± 1 . In this realization, \mathbf{SIS}_n is usually called the *signed rook monoid* or a *generalized rook monoid*, see [MaSr21].

For $i = 1, 2, \dots, n$, we denote by $\epsilon_i \in \mathbf{SIS}_n$ the transposition $(i, -i)$.

5.4.5. \mathbf{SIS}_n as a quotient of \mathbf{FB}_n^* . We have a natural type B analogue of Lemma 15. Consider the monoid \mathbf{FB}_n^* and an element of this monoid written in the form $\sigma\xi$, where σ is an invertible element and ξ is an idempotent. Recall from Subsection 5.3.4 that ξ corresponds to a certain triple (ρ, Y, f) , up to some equivalence. A singleton equivalence class of ρ contained in Y will be called a *pure singleton* for ξ .

Note that $\sigma\xi = (\sigma\xi\sigma^{-1})\sigma$, where $\sigma\xi\sigma^{-1}$ is an idempotent. The element σ maps each pure singleton for ξ to a pure singleton for $\sigma\xi\sigma^{-1}$. In other words, each $\sigma\xi = (\sigma\xi\sigma^{-1})\sigma$ induces a map from pure singletons for ξ to pure singletons for $\sigma\xi\sigma^{-1}$.

Lemma 18. *Restricting an element of \mathbf{FB}_n^* to pure singletons defines a homomorphism from \mathbf{FB}_n^* to \mathbf{SIS}_n which we denote by v . The homomorphism v is surjective.*

Proof. Similar to the proof of Lemma 15. □

5.4.6. Map from a type B singular Artin monoid. Let W be the Coxeter group corresponding to the following (type B) Coxeter graph:

$$s_0 \xrightarrow{4} s_1 \text{ --- } s_2 \text{ --- } \dots \text{ --- } s_{n-1}$$

For $i = 1, 2, \dots, n-1$, we denote by $\tilde{\epsilon}_{\{i, i+1\}}$ the idempotent of \mathbf{SIS}_n given by the identity on the complement to $\{i, i+1, \bar{i}, \bar{i}+1\}$. Similarly, for $i = 1, 2, \dots, n$, we denote by $\tilde{\epsilon}_{\{i\}}$ the idempotent of \mathbf{SIS}_n given by the identity on the complement to $\{i, \bar{i}\}$.

Proposition 19. *There is a unique homomorphism $\varphi : \mathbf{SB}(W) \rightarrow \mathbf{SIS}_n$ such that*

$$\begin{aligned} \varphi(\sigma_{s_i}) &= \tilde{s}_i, & \text{for } i = 1, 2, \dots, n-1; \\ \varphi(\sigma_{s_0}) &= \epsilon_1 \\ \varphi(\tau_{s_i}) &= \tilde{\epsilon}_{\{i, i+1\}}, & \text{for } i = 1, 2, \dots, n-1; \\ \varphi(\tau_{s_0}) &= \tilde{\epsilon}_{\{1\}}. \end{aligned}$$

The homomorphism φ is surjective.

Note that $\varphi = v \circ \lambda$.

Proof. The first statement follows by checking the defining relations of $\mathbf{SB}(W)$ for the images of the generators under φ as prescribed in the formulation. The second statement follows from the easy fact that \mathbf{SIS}_n is generated by \mathbf{IS}_n and ϵ_1 . □

As in type A , there is some space for generalizations of this map in the linear setting by using some scalars on the right hand side.

5.4.7. *Presentation in type B.* Set $\tilde{s}_0 := \epsilon_1$, $t_0 := \varepsilon_{\{1\}}$ and also $t_i := \varepsilon_{\{i, i+1\}}$, for $i = 1, 2, \dots, n-1$. We have the following presentation for \mathbf{SIS}_n .

Proposition 20. *The monoid \mathbf{SIS}_n is generated by the elements \tilde{s}_i and t_i , where we have $i = 0, 1, \dots, n-1$, subject to*

- the relations (1), (2), (3), (4) and (5) (for \tilde{s}_i instead of σ_{s_i} and t_i instead of τ_{s_i});
- the relations (7), (8), (9) and (10) (including for $i = 0$ and for \tilde{s}_i instead of s_i and for t_i instead of ξ_i);
- the relation (16) and, additionally,
- the relation

$$(17) \quad t_0 \tilde{s}_1 t_0 = t_1.$$

Proof. It is easy to check that the generators of \mathbf{SIS}_n satisfy all the prescribed relations. Let Q denote the semigroup generated by s_i and t_i , for $i = 0, 1, \dots, n-1$, subject to the analogues of the relations (1)–(5), the relations (7)–(10), the relation (16) and the relation (17). We have a canonical surjective map $\pi : Q \rightarrow \mathbf{SIS}_n$ sending s_i to \tilde{s}_i and t_i to t_i , for $i = 0, 1, \dots, n-1$. We need to show that π is an isomorphism.

The group W of units of Q is generated by the elements s_i , where $i = 0, 1, \dots, n-1$, and hence is isomorphic to the group of signed permutations of n elements (which is a Coxeter group of type B_n). Consider the set

$$T := \{\sigma t_0 \sigma^{-1} : \sigma \in W\} \subset Q.$$

The centralizer C of t_0 in W under conjugation contains all s_i , for $i \neq 1$, due to (3) and (5) as well as $s_1 s_0 s_1$ (again, due to (3)). The subgroup N of W generated by all these elements has index n . Since $|\pi(T)| = n$, it follows that $C = N$. Consequently, $|T| = n$ and π is injective, when restricted to T . Set

$$(18) \quad r_1 := t_0, \quad r_2 := s_1 t_0 s_1, \quad r_3 := s_2 s_1 t_0 s_1 s_2, \text{ and so on.}$$

Then $T = \{r_i : i = 1, 2, \dots, n\}$ and $\pi(r_i) = \tilde{\varepsilon}_{\{i\}}$, for $i = 1, 2, \dots, n$.

Lemma 21. *The elements in T commute.*

Proof. Using conjugation by elements in W , it is enough to show that r_1 commutes with all other r_i . Note that, for $i > 1$, the element r_i is a product of r_2 and some s_j , for $j > 1$. Therefore, using (3) and (18), it is enough to show that $r_1 r_2 = r_2 r_1$.

Using (17) and (10), we have

$$r_1 r_2 = t_0 s_1 t_0 s_1 = t_1 s_1 = t_1.$$

Similarly, $r_2 r_1 = t_1$ and we are done. \square

Each element of T is an idempotent by Proposition 11(8) and now we know that all these elements commute. Therefore the submonoid $\langle T \rangle$ of Q generated by T has at most 2^n elements. Since $\pi(\langle T \rangle)$ has exactly 2^n elements, we conclude that $|\langle T \rangle| = 2^n$.

Using conjugation by W , from $r_2 r_1 = t_1$, we obtain that all t_i belong to $\langle T \rangle$. For $k = 1, 2, \dots, n$, using (10) we obtain that

$$q r_1 r_2 \cdots r_k = r_1 r_2 \cdots r_k$$

for any q in the subgroup G of W generated by s_0, s_1, \dots, s_{k-1} . Note that G is a Coxeter group of type B_k .

We can write each element in Q in the form ab , where $a \in W$ and $b \in \langle T \rangle$. If b is a product of k different r_i , then, by the previous paragraph, a can be chosen modulo a type B_k Coxeter subgroup of W . Now the claim of the proposition follows by comparing the cardinality of Q with that of \mathbf{SIS}_n . \square

The signed rook monoid and its presentation appears also in [EF13]. In the recent paper [CE22], one can find an independent general treatment of presentations for the wreath product of an abelian group with \mathbf{IS}_n .

5.4.8. *Possible analogues in other types.* Propositions 17 and 20 can be used to extrapolate a candidate for an analogue of the symmetric inverse monoid for an arbitrary Coxeter group. One might note that the additional relation given by Proposition 20 involves an asymmetry that is not a part of the Coxeter datum. Looking carefully into the proofs of Propositions 17 and 20 in order to find some common ground, it is natural to suggest the following definition.

Let M be a Coxeter matrix with the associated Coxeter group W , the Artin braid group $\mathbf{B}(W)$ and the singular Artin monoid $\mathbf{SB}(W)$. We define the monoid $\mathbf{FB}(W)$ as the quotient of $\mathbf{SB}(W)$ modulo the relations (7)-(10) in Proposition 11 and, additionally, the relations that all conjugates of all τ_s by all elements in W commute. This set of relations is, clearly, not minimal and can be cleaned up.

Let T be the set of all conjugates of all τ_s by all elements in W . This generates a commutative subsemigroup $\langle T \rangle$ of $\mathbf{FB}(W)$ consisting of idempotents. From the proofs of Propositions 17 and 20, we see that each element of $\mathbf{FB}(W)$ can be written as σx , where $\sigma \in W$ and $x \in \langle T \rangle$. In particular, this implies that $\mathbf{FB}(W)$ is finite if and only if W is.


The semigroup $\mathbf{FB}(W)$ is regular as $x\sigma^{-1}$ is an inverse of σx . Furthermore, it is easy to see that $\langle T \rangle$ coincides with the set of all idempotents of $\mathbf{FB}(W)$. In particular, all idempotents of $\mathbf{FB}(W)$ commute and hence $\mathbf{FB}(W)$ is an inverse semigroup. Note that, in type A , the semigroup $\mathbf{FB}(W)$ is isomorphic to \mathbf{F}_n^* (and, in particular, is bigger than \mathbf{IS}_n). One could probably add the relation (16) for type A parabolic subgroups of W to get a smaller inverse semigroup.

In type B , the semigroup $\mathbf{FB}(W)$ is, in general, bigger than \mathbf{SIS}_n , as the relation (17), being asymmetric (for example, in the dihedral types), cannot follow from the symmetric relations defining $\mathbf{FB}(W)$, in general.


To understand the structure and properties of $\mathbf{FB}(W)$ for general Coxeter groups seems to be a natural and interesting problem.

5.5. Brauer algebra.

5.5.1. *The classical Brauer algebra.* Let $n \in \mathbb{Z}_{>0}$ and δ be an indeterminate. The classical Brauer algebra $\mathbf{Br}_n(\mathbb{Z}, \delta)$, defined in [Br37], is an algebra over $\mathbb{Z}[\delta]$ with a basis given by all Brauer diagrams on $2n$ vertices $\underline{n} \cup \underline{n}'$. Such a Brauer diagram is a partition of $\underline{n} \cup \underline{n}'$ into two-element disjoint subsets, depicted as an unoriented graph with the vertices from \underline{n} in the top row, the vertices from \underline{n}' in the bottom row and edges between the elements belonging to the same subsets. Here is an example of a Brauer diagram for $n = 4$:




Multiplication is given by concatenation of two such diagrams by identifying the primed vertices of the first with the ordinary vertices of the second, which induces a partition of the remaining vertices into two-element subsets, and then multiplying with δ^k , where k is the number of connected components supported only on the identified (and subsequently removed) vertices. Here is an example:



The diagrams in which each subset intersects both \underline{n} and \underline{n}' form a group isomorphic to the symmetric group S_n . In particular, $\mathbb{Z}[S_n] \subset \mathbf{Br}_n(\mathbb{Z}, \delta)$. One can extend scalars from \mathbb{Z} to any commutative ring \mathbb{k} and evaluate δ at any element d of the latter ring. This gives the algebra $\mathbf{Br}_n(\mathbb{k}, d)$.

5.5.2. Map from singular braid monoid. As usual, for $i = 1, 2, \dots, n-1$, we denote by s_i the transposition $(i, i+1)$ considered as an element of $\mathbf{Br}_n(\mathbb{Z}, \delta)$. We also denote by s_i the Brauer diagram



Proposition 22. *There is a unique homomorphism $\chi : \mathbf{SB}(S_n) \rightarrow \mathbf{Br}_n(\mathbb{Z}, \delta)$ such that*

$$\chi(\sigma_{s_i}) = s_i \text{ and } \chi(\tau_{s_i}) = s_i, \text{ for } i = 1, 2, \dots, n-1.$$

Proof. The existence claim follows from the straightforward verification of the analogues of the relations (1), (2), (3), (4) and (5) for the elements s_i and s_i . The uniqueness claim follows from the fact that σ_{s_i} and τ_{s_i} generate $\mathbf{SB}(S_n)$. \square

5.5.3. Presentation for the Brauer algebra. In [KMaz06, Theorem 3.1], one can find a presentation of the Brauer monoid \mathbf{Br}_n (this corresponds to the case $\delta = 1$) which is obtained from the presentation of $\mathbf{SB}(S_n)$ by adding the additional relations (for the images of the generators under χ):

- (19) $s_i^2 = e,$
- (20) $\mathbf{s}_i^2 = \mathbf{s}_i,$
- (21) $\mathbf{s}_i \mathbf{s}_i = s_i \mathbf{s}_i = \mathbf{s}_i,$
- (22) $\mathbf{s}_i \mathbf{s}_{i\pm 1} \mathbf{s}_i = \mathbf{s}_i,$
- (23) $\mathbf{s}_i \mathbf{s}_{i\pm 1} \mathbf{s}_i = \mathbf{s}_i \mathbf{s}_{i\pm 1},$
- (24) $s_i \mathbf{s}_{i\pm 1} \mathbf{s}_i = s_{i\pm 1} \mathbf{s}_i.$

In other words, the kernel of χ is generated by the above relations. We note the following:

Lemma 23. *The relation (22) is redundant.*

Proof. We have:

$$\begin{aligned}
 \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i &\stackrel{(19)}{=} \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i^2 \mathbf{s}_i \\
 &\stackrel{(21)}{=} \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i \mathbf{s}_i \\
 &\stackrel{(23)}{=} \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i \\
 &\stackrel{(19)}{=} \mathbf{s}_{i+1}^2 \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i^2 \mathbf{s}_i \\
 &\stackrel{(19),(21)}{=} \mathbf{s}_{i+1} \mathbf{s}_i^2 \mathbf{s}_{i+1} \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i \mathbf{s}_i \\
 &\stackrel{(2)}{=} \mathbf{s}_{i+1} \mathbf{s}_i \mathbf{s}_{i+1} \mathbf{s}_i \\
 &\stackrel{(24)}{=} \mathbf{s}_{i+1} \mathbf{s}_{i+1} \mathbf{s}_i \\
 &\stackrel{(19)}{=} \mathbf{s}_i
 \end{aligned}$$

□

If \mathbb{k} is a commutative ring and $\delta \in \mathbb{k}$ is invertible, then the above implies a presentation for $\mathbf{Br}_n(\mathbb{k}, \delta)$ by correcting $\mathbf{s}_i^2 = \mathbf{s}_i$ to $\mathbf{s}_i^2 = \delta \mathbf{s}_i$.

5.5.4. *Type B Brauer algebra.* Similarly to all previous situations (i.e. binary relations, dual symmetric inverse semigroups and symmetric inverse semigroups), consider the Brauer algebra $\mathbf{Br}_n^{(2)}(\mathbb{Z}, \delta)$ on $\underline{n} \cup \overline{n}$. It has an obvious automorphism induced by the swapping $i \leftrightarrow \bar{i}$. We denote by $\mathbf{Br}_n^B(\mathbb{Z}, \delta)$ the subalgebra of $\mathbf{Br}_n^{(2)}(\mathbb{Z}, \delta)$ generated by all Brauer diagrams on $\underline{n} \cup \overline{n}$ that are invariant under this automorphism. We emphasize that this is smaller than the subalgebra of all elements in $\mathbf{Br}_n^{(2)}(\mathbb{Z}, \delta)$ invariant under the above automorphism.

For $i = 1, 2, \dots, n-1$, we denote by \tilde{s}_i the unique element of $\mathbf{Br}_n^B(\mathbb{Z}, \delta)$ whose restriction to \underline{n} coincides with \mathbf{s}_i . We also denote by \tilde{s}_0 the element

$$\begin{array}{ccccccc}
 \bar{n} & \dots & \bar{2} & \bar{1} & 1 & 2 & \dots & n \\
 | & & | & \frown & & | & & | \\
 \bar{n}' & \dots & \bar{2}' & \bar{1}' & 1' & 2' & \dots & n'
 \end{array}$$

Let $\tilde{s}_0, \tilde{s}_1, \dots, \tilde{s}_{n-1}$ be the usual generators of the type B Coxeter group.

A different candidate for a type B Brauer algebra appeared in [CL15]. As far as we can judge, this candidate does not coincide with our algebra. For example, the relation [CL15, 2.12], that is $\tilde{s}_0 \tilde{s}_1 \tilde{s}_0 = \tilde{s}_1 \tilde{s}_0$, does not hold for our generators.

5.5.5. *Map from a type B singular Artin monoid.* Let W be the Coxeter group corresponding to the following (type B) Coxeter graph:


$$s_0 \xrightarrow{4} s_1 \text{ --- } s_2 \text{ --- } \dots \text{ --- } s_{n-1}$$

Proposition 24. *There is a unique homomorphism $\chi : \mathbf{SB}(W) \rightarrow \mathbf{Br}_n^B(\mathbb{Z}, \delta)$ such that*

$$\chi(\sigma_{s_i}) = \tilde{s}_i \text{ and } \chi(\tau_{s_i}) = \tilde{s}_i, \text{ for } i = 0, 1, \dots, n-1.$$

Proof. The existence claim follows from the straightforward verification of the analogues of the relations (1), (2), (3), (4) and (5) for the elements \tilde{s}_i and \tilde{s}_i . The uniqueness claim follows from the fact that σ_{s_i} and τ_{s_i} generate $\mathbf{SB}(S_n)$. □

5.5.6. *Connection to the partial Brauer algebra.* A partial analogue of Brauer monoid was introduced in [Ma98] and the representation theory of the corresponding partial Brauer algebra was studied in [MM14]. This algebra, denoted $\mathbf{PB}_n(\delta, \delta')$, has a basis consisting of *partial Brauer diagrams*. Such a diagram is a partition of $\underline{n} \cup \underline{n}'$ into singletons and two-element disjoint subsets, depicted as an unoriented graph with the vertices from \underline{n} in the top row, the vertices from \underline{n}' in the bottom row and edges between the elements belonging to the same subsets. Here is an example of a partial Brauer diagram for $n = 4$:



Multiplication is given by concatenation of two such diagrams by identifying the primed vertices of the first with the ordinary vertices of the second, which induces a partition of the remaining vertices into singletons and two-element subsets, and then multiplying with $\delta^k(\delta')^{k'}$, where k is the number of closed connected components supported only on the identified (and subsequently removed) vertices and k' is the number of not closed connected components supported only on the identified (and subsequently removed) vertices.

Let G be an abelian semigroup and π a partial Brauer diagram. An *almost G -coloring* of π is a map f from the two-element components of π to G . Let $\delta = (\delta_i)_{i \in G}$ be a collection of parameters. Given two almost G -colored partial Brauer diagrams (π, f) and (π', f') , we can define their product as $(\delta')^{k'} \prod_{i \in G} \delta_i^{k_i} (\pi\pi', \mathbf{f})$, where $\pi\pi'$ and k' are as in the previous paragraph, the value of \mathbf{f} at some two-element component C is defined by taking the product of the values of f , respectively, f' over all two-element components of π and π' that contributed to C , and k_i denotes the number of closed connected components of total value i (computed similarly as for C) that are removed during the straightening procedure. This defined on the linear span of all almost G -colored partial Brauer diagrams the structure of an associative algebra, denoted by $\mathbf{APB}_n(G, \delta, \delta')$. It is informative to compare this construction with the construction of colored partition algebras studied in [MaSr22].

By considering only the diagrams and forgetting δ and δ' , we obtain the corresponding partial Brauer monoid \mathbf{PB}_n and the almost colored partial Brauer monoid $\mathbf{APB}_n(G)$.


Proposition 25. *The monoid \mathbf{Br}_n^B is isomorphic to the monoid $\mathbf{APB}_n(\mathbb{Z}_2)$.*

Proof. Given a diagram $\pi \in \mathbf{Br}_n^B$, we define the corresponding diagram $\sigma \in \mathbf{APB}_n(\mathbb{Z}_2)$ as follows (here all i and j are positive):

- If π connects i with \bar{i} , then this i is a singleton in σ .
- If π connects i' with \bar{i}' , then this i' is a singleton in σ .
- If π connects i with j , then i and j are connected in σ and colored by 0.
- If π connects i' with j' , then i' and j' are connected in σ and colored by 0.
- If π connects i with \bar{j} , where $i \neq j$, then i and j are connected in σ and colored by 1.
- If π connects i' with \bar{j}' , where $i \neq j$, then i' and j' are connected in σ and colored by 1.
- If π connects i with j' , then i and j' are connected in σ and colored by 0.

- If π connects i with j' , then i and j' are connected in σ and colored by 1.

Here is an example:



It is easy to check that this defines an isomorphism between \mathbf{Br}_n^B and $\mathbf{APB}_n(\mathbb{Z}_2)$. \square

5.5.7. *Presentation of $\mathbf{Br}_n^B(\mathbb{Z}, \delta)$.* We now give a presentation for the type B Brauer monoid \mathbf{Br}_n^B , that is the semigroup formed by all type B Brauer diagrams. This corresponds to the case $\delta = 1$. Note that χ restricts to a homomorphism from $\mathbf{SB}(S_n)$ to \mathbf{Br}_n^B .

Theorem 26. *The monoid \mathbf{Br}_n^B is generated by \tilde{s}_i and \tilde{s}_i , where $i = 0, 1, \dots, n-1$, subject to*

- the (analogues of the) relations (1), (2), (3), (4) and (5);
- the (analogues of the) relations (19), (20), (21), (23), (24), where all indices are greater than 0;
- the additional relations

$$\begin{aligned}
 (25) \quad & \tilde{s}_0^2 = \tilde{s}_0, \\
 (26) \quad & \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 = \tilde{s}_1, \\
 (27) \quad & \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 = \tilde{s}_1, \\
 (28) \quad & \tilde{s}_0 \tilde{s}_1 \tilde{s}_0 = \tilde{s}_0 \tilde{s}_1 \tilde{s}_0, \\
 (29) \quad & \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 \tilde{s}_0 = \tilde{s}_0 \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 = \tilde{s}_0 \tilde{s}_1 \tilde{s}_0, \\
 (30) \quad & \tilde{s}_1 \tilde{s}_0 = \tilde{s}_1 \tilde{s}_1 \tilde{s}_0 \tilde{s}_1, \\
 (31) \quad & \tilde{s}_0 \tilde{s}_1 = \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 \tilde{s}_1, \\
 (32) \quad & \tilde{s}_0 \tilde{s}_0 = \tilde{s}_0 \tilde{s}_0 = \tilde{s}_0.
 \end{aligned}$$

Proof. Denote by Q the monoid with the presentation described in the formulation generated by \tilde{x}_i (instead of \tilde{s}_i) and \tilde{x}_i (instead of \tilde{s}_i). We have the obvious surjection $\tilde{\pi} : Q \rightarrow \mathbf{Br}_n^B$ which we want to prove is an isomorphism. For this we need to prove that $|Q| \leq |\mathbf{Br}_n^B|$.

Due to (1) and (20), the group G of invertible element in Q is isomorphic to the signed permutation group on \underline{n} . Denote by T the set of all G -conjugates of all \tilde{x}_i , where $i = 0, 1, 2, \dots, n-1$. From (25) and (20) it follows that all elements in T are idempotents.

By (21), (32), (2), (3) and (5), the G -conjugate stabilizer of \tilde{x}_0 contains both \tilde{x}_0 and all signed permutations on $\{2, 3, \dots, n\}$. Hence the number of the G -conjugates of \tilde{x}_0 is at most $\frac{2^n \cdot n!}{2 \cdot 2^{n-1} (n-1)!} = n$. Since the number of the G -conjugates of \tilde{s}_0 is exactly n , we obtain that the number of the G -conjugates of \tilde{x}_0 is exactly n . For $i = 1, \dots, n$, we denote by t_i the conjugate of \tilde{x}_0 by the transposition $(1, i)$.

Similarly, the G -conjugate stabilizer of \tilde{x}_1 contains both \tilde{x}_1 , $\tilde{x}_0 \tilde{x}_1 \tilde{x}_0$, $\tilde{x}_1 \tilde{x}_0 \tilde{x}_1 \tilde{x}_0$ and all signed permutations on $\{3, 4, \dots, n\}$. Hence the number of the G -conjugates of \tilde{x}_1 is

at most $\frac{2^n \cdot n!}{4 \cdot 2^{n-2} (n-2)!} = n(n-1)$. Since the number of the G -conjugates of \tilde{s}_1 is exactly $n(n-1)$, we obtain that the number of the G -conjugates of \tilde{x}_1 is exactly $n(n-1)$. For different $i, j \in \{1, \dots, n\}$, we denote by $\mathfrak{t}_{i,j}$ the conjugate of \tilde{x}_1 by $(1, i)(2, j)$. From (21), it follows that $\mathfrak{t}_{i,j} = \mathfrak{t}_{j,i}$. Also, for different $i, j \in \{1, \dots, n\}$, we denote by $\bar{\mathfrak{t}}_{i,j}$ the conjugate of \tilde{x}_1 by $(1, \bar{i})(2, j)$. We similarly have $\bar{\mathfrak{t}}_{i,j} = \bar{\mathfrak{t}}_{j,i}$.

From (29) and the relations for the singular Artin monoids, it follows that all \mathfrak{t}_i commute with each other. From the relations for the singular Artin monoids, it follows also that \mathfrak{t}_i commutes with both $\bar{\mathfrak{t}}_{k,j}$ and $\mathfrak{t}_{k,j}$ provided that $i \notin \{k, j\}$. Similarly, both $\bar{\mathfrak{t}}_{i,j}$ and $\mathfrak{t}_{i,j}$ commute with both $\bar{\mathfrak{t}}_{k,l}$ and $\mathfrak{t}_{k,l}$ provided that $\{i, j\} \cap \{k, l\} = \emptyset$. This implies that different elements of T commute with each other in Q if and only if their images in \mathbf{Br}_n^B commute with each other. We will call $\{i\}$ the support of \mathfrak{t}_i . Similarly, we will call $\{i, j\}$ the support of both $\bar{\mathfrak{t}}_{i,j}$ and $\mathfrak{t}_{i,j}$.

Let $\omega \in Q$. Denote by ω_l the product of all $\mathfrak{t} \in T$ such that $\mathfrak{t}\omega = \omega$. Define the left defect of ω as the union of the supports of all the factors of ω_l . Note that, if \mathfrak{t} is a factor of ω_l , then $\tilde{\pi}(\mathfrak{t})\tilde{\pi}(\omega) = \tilde{\pi}(\omega)$. In the monoid \mathbf{Br}_n^B , different $\tilde{\pi}(\mathfrak{t})$ with the above property commute with each other. This implies that all factors of ω_l commute with each other. In particular, the left defect of ω is the disjoint union of the supports of the factors of ω_l . Since all elements in T are idempotents, we have $\omega = \omega_l\omega$.

Denote by ω_r the product of all $\mathfrak{t} \in T$ such that $\omega\mathfrak{t} = \omega$. Define the right defect of ω as the union of the supports of all the factors of ω_r . Similarly to the previous paragraph, all factor of ω_r commute with each other and the right defect of ω is the disjoint union of the supports of the factors of ω_r . Since all elements in T are idempotents, we have $\omega = \omega\omega_r$.

The following lemma will be useful.

Lemma 27. *Let ω be a product of pairwise commuting elements of T with disjoint supports. Assume that the left defect of ω is \underline{n} . Then $\omega = \omega u \omega$, for any $u \in Q$.*

Proof. For the classical Brauer and partial Brauer algebras, the corresponding claim is clear. In our generalization, we note that the presentation of the Brauer algebra is included into our presentation by definition. The presentation of the partial Brauer algebra from [KMaz06, Section 5] follows easily taking into account our additional relations (25)–(32). Indeed, [KMaz06, (15)] follows from (25) and the commutativity of the \mathfrak{t}_i 's established above; [KMaz06, (16)] follows from our construction of the \mathfrak{t}_i 's and (29); [KMaz06, (17)] is a singular braid relation; [KMaz06, (18)] follows from (30) and (31) and, finally, [KMaz06, (19)] follows from (26), (28) and (29).

In the terminology of Proposition 25, we observe that we have the similar cancellation relations (26) and (27) for the “partial” element \tilde{s}_0 and the “coloring” element \tilde{s}_0 , as well as the additional connection between these two elements given by the relation (32) (this relation is, in particular, responsible for the fact that singletons cannot be colored). Therefore cancellation of the factors of u that are conjugates of the “coloring” element \tilde{s}_0 can be done similarly to the arguments used in [KMaz06, Section 5] for the cancellation of the conjugates of the “partial” element \tilde{s}_0 . We leave the details to the reader. \square

We claim that every $\omega \in Q$ can be written as $\omega = \omega_l \omega_m \omega_r$, where ω_l and ω_r are defined above and ω_m is invertible and such that $\tilde{\pi}(\omega_m)$ induces a map from the right to the left defect of ω . We will call this a normal form of ω . We note that ω_l and ω_r are uniquely defined. At the same time, from (21) and (32) it follows that ω_m is not uniquely defined in general. In fact, from (an appropriate analogue of) Lemma 27

it follows that, to the very least, it is defined up to a signed bijection between the elements in the left and the right defects of ω .

In \mathbf{Br}_n^B , it is easy to see that each element can be written in the above normal form where the middle term is defined exactly up to the signed bijection between the elements in the left and the right defects of ω . In particular, the sizes of the left and the right defects of ω coincide. Another consequence is that $\tilde{\pi}$ induces a bijection between the elements of Q which have a normal form and the elements of \mathbf{Br}_n^B . In other words, to complete the proof, we just need to show the above claim that each element in Q has a normal form.

We prove the claim by induction on the number of factors from T in a shortest expression for $\omega \in Q$. If this shortest expression of ω has no factors from T , the ω is invertible and the claim is clear. This is the basis of our induction.

To prove the induction step, consider a normal form $\omega_l \omega_m \omega_r$, for some $\omega \in Q$. For any invertible u , it is easy to see that $(u\omega_l u^{-1})(u\omega_m)\omega_r$ is a normal form for $u\omega_l \omega_m \omega_r$. Therefore, it remains to prove that, for any $t \in T$, the element $t\omega_l \omega_m \omega_r$ has a normal form.

If the support of t is disjoint from the left defect of ω , then, taking into account that t is idempotent, we can move one copy of t past ω_l . Note that, because of our assumption on ω_m , the support of $\omega_m^{-1}t\omega_m \in T$ is disjoint from the support of ω_r . Using some of the commutativity relations that we already established, it follows that

$$(t\omega_l)\omega_m(\omega_m^{-1}t\omega_m\omega_r)$$

is a normal form for $t\omega_l \omega_m \omega_r$.

Next, let $t = t_i$, for some i . Because of the previous paragraph, we only need to consider the case when i belongs to the left defect of ω . If t_i is a factor of ω_l , then $t\omega_l \omega_m \omega_r = \omega$ due to the idempotency of t_i and we have nothing to prove. In the other case, some $t_{i,j}$ or $\bar{t}_{i,j}$ is a factor of ω . We consider the former case, with the latter being similar. Using (31) and the idempotency of t_i , we have $t_i t_{i,j} = t_i t_j t_{i,j}$. Denote by ω'_l the element obtained from ω_l by replacing the factor $t_{i,j}$ by the factor $t_i t_j$. We claim that $\omega'_l \omega_m \omega_r$ is a normal form for $t\omega_l \omega_m \omega_r$. It follows from the definitions that $\omega'_l \omega_m \omega_r$ is a normal form, so the only thing to show is that $\omega'_l \omega_m \omega_r = t\omega_l \omega_m \omega_r$, that is, that $\omega'_l \omega_m \omega_r = \omega'_l(t_{i,j}\omega_m)\omega_r$.

From (26), it follows that

$$(33) \quad t_{s,t} t_s t_{s,t} = t_{s,t},$$

for all $s \neq t$. Using (28) and (29), we have $\tilde{s}_0 \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 \tilde{s}_0 \tilde{s}_1 \tilde{s}_0 = \tilde{s}_0 \tilde{s}_1 \tilde{s}_0$. This implies

$$(34) \quad t_s t_{s,t} t_s t_{s,t} t_s = t_s t_{s,t} t_s,$$

for all $s \neq t$. Of course, we also have similar identities with $t_{s,t}$ replaced by $\bar{t}_{s,t}$. Using these observations, the fact that $\omega'_l \omega_m \omega_r = \omega'_l(t_{i,j}\omega_m)\omega_r$ follows from (an appropriate analogue) of Lemma 27.

Next consider the case $t = t_{i,j}$ with $\{i, j\}$ inside the left defect of ω (for $t = \bar{t}_{i,j}$, the arguments are similar). If both, t_i and t_j are factors of ω_l , then one can use the same argument as in the previous case. The remaining case is when either $t_{i,s}$ or $t_{j,s}$ or $\bar{t}_{i,s}$ or $\bar{t}_{j,s}$ is a factor of ω_l , for some $s \neq i, j$. In this case, using (24), we reduce the situation to the case of multiplication of ω with an invertible element considered above.

Finally, consider the case $t = t_{i,j}$ with only i belonging to the left defect of ω (again, for $t = \bar{t}_{i,j}$, the arguments are similar). The case when either $t_{i,s}$ or $\bar{t}_{i,s}$ is a factor of

ω_l is dealt with as in the previous paragraph. It remains to consider the case when τ_i is a factor of ω_l . Define ω'_l as ω_l with τ_i replaced by $\tau_{i,j}$.

Assume that $\tilde{\pi}(\omega_m)$ matches j with some k . Define ω'_r as the product $\omega_r \tau_k$. We claim that $\omega'_l \omega_m \omega'_r$ is a normal form for $\tau_{i,j} \omega$. It follows directly from the construction that $\omega'_l \omega_m \omega'_r$ is a normal form. It remains to note that $\omega'_l \omega_m \omega'_r = \tau_{i,j} \omega$ follows from the definitions of ω'_l and ω'_m and (an appropriate analogue) of Lemma 27. \square

If \mathbb{k} is a commutative ring and $\delta \in \mathbb{k}$ is invertible, then the above implies a presentation for $\mathbf{Br}_n^B(\mathbb{k}, \delta)$ by correcting $\mathfrak{s}_i^2 = \mathfrak{s}_i$ to $\mathfrak{s}_0^2 = \delta \mathfrak{s}_0$ and $\mathfrak{s}_i^2 = \delta^2 \mathfrak{s}_i$, for $i > 0$.

6. RECAP OF CATEGORY \mathcal{O}

6.1. Setup and definitions. From now on we work over the field \mathbb{C} of complex numbers. We use [Hu72, Hu08] as standard references.

Let \mathfrak{g} be a finite dimensional, semi-simple complex Lie algebra with a fixed triangular decomposition

$$(35) \quad \mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+.$$

Here \mathfrak{h} is a Cartan subalgebra and \mathfrak{n}_\pm are the positive and the negative parts, respectively. We denote by $\mathbf{R} \subset \mathfrak{h}^*$ the root system of the pair $(\mathfrak{g}, \mathfrak{h})$ with the induced decomposition $\mathbf{R} = \mathbf{R}_- \cup \mathbf{R}_+$ into negative and positive roots. Let W be the Weyl group of \mathbf{R} and S the set of simple reflections associated to our choice of positive roots. Then (W, S) is a Coxeter system. We denote by w_0 the longest element of W .

Let \mathcal{O} be the Bernstein-Gelfand-Gelfand (BGG) category associated with the triangular decomposition in (35), see [BGG75, BGG76]. For $\lambda \in \mathfrak{h}^*$, we denote by

- $\Delta(\lambda)$ the Verma module with highest weight λ ,
- $L(\lambda)$ the unique simple quotient of $\Delta(\lambda)$,
- $P(\lambda)$ the indecomposable projective cover of $L(\lambda)$ in \mathcal{O} ,
- $I(\lambda)$ the indecomposable injective envelope of $L(\lambda)$ in \mathcal{O} ,
- $T(\lambda)$ the indecomposable tilting envelope of $\Delta(\lambda)$ in \mathcal{O} .

Let \mathcal{O}_0 be the principal block of \mathcal{O} , that is the indecomposable direct summand containing the trivial \mathfrak{g} -module $L(0)$. Let ρ be the half of the sum of all positive roots. The dot-action of W on \mathfrak{h}^* is defined via $w \cdot \lambda := w(\lambda + \rho) - \rho$. The simple objects in \mathcal{O}_0 are in bijection with the elements in W via $w \mapsto L_w := L(w \cdot 0)$. We use similar notation for other structural modules in \mathcal{O}_0 , for example $\Delta_w := \Delta(w \cdot 0)$, $P_w := P(w \cdot 0)$ etc.

Denote by A a basic algebra of \mathcal{O}_0 , that is the (opposite of the) endomorphism algebra of a multiplicity-free projective generator of \mathcal{O}_0 . Then we have the usual equivalence $\mathcal{O}_0 \cong A\text{-mod}$.

Denote by \star the usual simple preserving duality on \mathcal{O} . For any $w \in W$, we then have $L_w^\star \cong L_w$, $P_w^\star \cong I_w$ and $T_w^\star \cong T_w$. We also have the dual Verma modules $\nabla_w := \Delta_w^\star$.

6.2. Graded lift of \mathcal{O}_0 . Denote by \mathbb{C} the coinvariant algebra of the W -module \mathfrak{h}^* . It is defined as the quotient of the \mathbb{Z} -graded polynomial algebra of \mathfrak{h} , under the convention $\deg \mathfrak{h} = 2$, modulo the ideal generated by all homogeneous W -invariant polynomials of positive degree. Directly from the definition, we have that \mathbb{C} is naturally \mathbb{Z} -graded. In what follows, by *graded* we mean \mathbb{Z} -graded.

Soergel's combinatorial description of \mathcal{O}_0 , see [So90], asserts, among other things, the following:

- $\text{End}_{\mathcal{O}}(P_{w_0}) \cong \mathbb{C}$;
- the functor $\text{Hom}_{\mathcal{O}}(P_{w_0}, -) : \mathcal{O}_0 \rightarrow \text{mod-}\mathbb{C}$ is full and faithful on projectives and, for $w \in W$, the \mathbb{C} -module $\text{Hom}_{\mathcal{O}}(P_{w_0}, P_w)$ is naturally graded.

This gives the algebra A the natural structure of a graded algebra

$$A = \bigoplus_{i \in \mathbb{Z}} A_i.$$

Moreover, we have $A_i = 0$ for $i < 0$ and A_0 is semisimple. In other words, A is positively graded, see [So90, St03] for details.

Denote by $\mathcal{O}_0^{\mathbb{Z}}$ the category of finite dimensional graded A -modules (where morphisms are homogeneous maps of degree zero). There is the obvious functor from $\mathcal{O}_0^{\mathbb{Z}}$ to \mathcal{O}_0 which forgets the grading.

We denote by $\langle 1 \rangle$ the endofunctor of $\mathcal{O}_0^{\mathbb{Z}}$ which shifts the grading, under the normalization that it sends degree 0 to degree -1 . We fix the standard graded lifts of all structural modules:

- L_w is concentrated in degree 0;
- Δ_w with the top concentrated in degree 0;
- P_w with the top concentrated in degree 0;
- ∇_w with the socle concentrated in degree 0;
- I_w with the socle concentrated in degree 0;
- T_w such that $\Delta_w \hookrightarrow T_w$ has degree 0.

The duality \star admits a graded lift, denoted by the same symbol. It has the usual property that $(M^*)_{-i} \cong \text{Hom}_{\mathbb{C}}(M_i, \mathbb{C})$, for $M \cong \bigoplus_{i \in \mathbb{Z}} M_i \in \mathcal{O}_0^{\mathbb{Z}}$.

6.3. Combinatorics of \mathcal{O}_0 . Consider the Grothendieck group $[\mathcal{O}_0^{\mathbb{Z}}]$ of $\mathcal{O}_0^{\mathbb{Z}}$ and the Hecke algebra \mathbf{H} of the Coxeter system (W, S) . There exists a unique isomorphism of abelian groups as follows:

$$\Xi : [\mathcal{O}_0^{\mathbb{Z}}] \rightarrow \mathbf{H} \text{ such that } \Xi([\Delta_w]) = H_w, \text{ for } w \in W, \text{ and } \Xi \circ \langle 1 \rangle = v^{-1} \Xi.$$

As a consequence of the Kazhdan-Lusztig conjecture (see [KL79], proved in [BB81, BK81]), we have $\Xi([P_w]) = \underline{H}_w$, for $w \in W$.

6.4. Endofunctors of \mathcal{O}_0 . For $w \in W$, denote by θ_w the projective endofunctor on \mathcal{O}_0 or $\mathcal{O}_0^{\mathbb{Z}}$ such that $\theta_w P_e = P_w$, see [BG80]. These are exactly the indecomposable objects (up to shift of grading) in the corresponding monoidal categories \mathcal{P} and $\mathcal{P}^{\mathbb{Z}}$ of projective endofunctors on \mathcal{O}_0 or $\mathcal{O}_0^{\mathbb{Z}}$, respectively. The split Grothendieck ring $[\mathcal{P}^{\mathbb{Z}}]_{\oplus}$ is isomorphic to \mathbf{H} under $[\theta_w] \mapsto \underline{H}_w$, for $w \in W$, with the usual convention that $\langle 1 \rangle$ corresponds to v^{-1} . Homomorphisms between projective functors are described in [Bac92]. Soergel's combinatorial description of \mathcal{O}_0 implies that $\mathcal{P}(\theta_e, \theta_e) \cong \mathbb{C}$. Furthermore, each $\mathcal{P}(\theta_x, \theta_y)$ is a free \mathbb{C} -module as a basis of which one can take an arbitrary lift of any basis in $\text{Hom}_{\mathcal{O}}(P_x, P_y)$. Consequently, the endomorphism algebra B of $\bigoplus_{w \in W} \theta_w$ is positively graded. More generally, Soergel's combinatorial description of \mathcal{O}_0 describes \mathcal{P} via the biequivalent monoidal category of *Soergel bimodules* for (W, S) . The latter is the minimal isomorphism closed monoidal subcategory of $\mathbb{C}\text{-mod-}\mathbb{C}$

containing the bimodules $\mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C}$, where, for a simple reflection s , we denote by \mathbb{C}^s the algebra of all s -invariants in \mathbb{C} . This biequivalence sends θ_s to $\mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C}$, for $s \in S$.

For $s \in S$, we have the adjunction morphisms $\theta_e \langle -1 \rangle \xrightarrow{\text{adj}_s} \theta_s$ and $\theta_s \xrightarrow{\text{adj}_s} \theta_e \langle 1 \rangle$. The cokernel of adj_s is called the shuffling functor and denoted C_s . The kernel of adj_s is called the coshuffling functor and denoted K_s .

The functor C_s is right exact and the derived functor $\mathcal{L}C_s$ is given by tensoring with the complex $0 \rightarrow \theta_e \langle -1 \rangle \xrightarrow{\text{adj}_s} \theta_s \rightarrow 0$, where θ_s is in the homological position 0, followed by taking the total complex.

Similarly, the functor K_s is left exact and the derived functor $\mathcal{R}K_s$ is given by tensoring with the complex $0 \rightarrow \theta_s \xrightarrow{\text{adj}_s} \theta_e \langle 1 \rangle \rightarrow 0$, where θ_s is in the homological position 0, followed by taking the total complex. The functors $\mathcal{L}C_s$ and $\mathcal{R}K_s$ are mutually inverse equivalences. We refer to [Ca86, MaStr05] for details.

For $w \in W$ and $s \in S$ such that $ws > w$, we have the exact sequences

$$\Delta_w \langle -1 \rangle \xrightarrow{\text{adj}_s(\Delta_w)} \theta_s \Delta_w \rightarrow \Delta_{ws} \quad \text{and} \quad \Delta_w \hookrightarrow \theta_s \Delta_{ws} \xrightarrow{\text{adj}_s(\Delta_{ws})} \Delta_{ws} \langle 1 \rangle.$$

In particular, we have

$$(36) \quad C_s \Delta_w \cong \Delta_{ws} \quad \text{and} \quad K_s \Delta_{ws} \cong \Delta_w.$$

Corollary 28. *Let $w \in W$ and $w = s_1 s_2 \dots s_k$ be a reduced expression. Then we have an isomorphism $C_{s_k} \dots C_{s_2} C_{s_1} \Delta_e \cong \Delta_w$.*

Proof. Apply (36) inductively along the reduced expression. \square

6.5. Homotopy category of complexes of projective functors. Consider the homotopy category $\mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$ of complexes of projective functors. This category has the natural structure of a tensor triangulated category since it is the homotopy category of an additive finitary monoidal category. The category $\mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$ acts naturally on $\mathcal{D}^b(\mathcal{O}_0^{\mathbb{Z}})$ via the component-wise action followed by taking the total complex. The following elementary observation will be very handy:

Lemma 29. *Let $F_{\bullet} \in \mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$ be minimal (in the sense that it does not have non-trivial null-homotopic direct summands). Then $F_{\bullet} \Delta_e \in \mathcal{K}^b(\text{proj}(\mathcal{O}_0^{\mathbb{Z}}))$ is also minimal.*

Proof. A complex of projective modules is not minimal provided that it contains a (null-homotopic) direct summand of the form $0 \rightarrow P \xrightarrow{\cong} P \rightarrow 0$, for some indecomposable P .

Let $\theta_w \langle i \rangle \xrightarrow{\alpha} \theta_w \langle i \rangle$ be a piece of F_{\bullet} which, after the evaluation at Δ_e , produces a null-homotopic direct summand $P_w \langle i \rangle \xrightarrow{\alpha(\Delta_e)_i \cong} P_w \langle i \rangle$ of $F_{\bullet} \Delta_e$. Then α must be non-zero and thus an isomorphism due to the positivity of the grading on B . This contradicts our assumption that F_{\bullet} is minimal. \square

7. ORIGINAL CATEGORIFICATION OF SINGULAR BRAID MONOID USING PROJECTIVE FUNCTORS

7.1. Braid group action. Let G be a (semi)group and \mathcal{C} a category. Recall that a *weak action* of G on \mathcal{C} is an assignment of an endofunctor $F_g : \mathcal{C} \rightarrow \mathcal{C}$, to each $g \in G$, such that $F_g F_h \cong F_{gh}$, for any $g, h \in G$. The following statement is well-known (see [MaStr05, MaStr07]). However, there does not seem to be an available proof in

the literature in full generality, so we give a proof that appears to be based on a new approach that we will also extensively use later on.

Proposition 30. *The assignment $\sigma_s \mapsto \mathcal{L}C_s$ and $\sigma_s^{-1} \mapsto \mathcal{R}K_s$, for $s \in S$, extends uniquely (up to isomorphism) to a weak action of the Artin group $\mathbf{B}(W)$ on $\mathcal{D}^b(\mathcal{O}_0^{\mathbb{Z}})$.*

Proof. We need to check that every defining relation for $\mathbf{B}(W)$ gives rise to an isomorphism of functors between the two sides of the relation. The relations $\sigma_s \sigma_s^{-1} = \sigma_s^{-1} \sigma_s = e$ are checked in [MaStr05, Theorem 5.9], this means exactly that the functors $\mathcal{L}C_s$ and $\mathcal{R}K_s$ are mutually inverse equivalences. Hence we only need to check the braid relations.


Let $s \neq t \in S$ and $w = \underbrace{sts \dots}_{m_{s,t} \text{ factors}}$. Consider the functor

$$F = \dots \underbrace{\mathcal{L}C_s \mathcal{L}C_t \mathcal{L}C_s}_{m_{s,t} \text{ factors}} \dots$$

Evaluating F at Δ_e and noting that all shuffling functors are acyclic on Verma modules, we obtain $F\Delta_e \cong \Delta_w$ using Corollary 28.

Let $G_{\bullet} \in \mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$ be a minimal complex representing F . Then, by Lemma 29, the evaluation $G_{\bullet}\Delta_e$ gives a minimal projective resolution of Δ_w . To complete the proof, we need to show that this property describes G_{\bullet} uniquely, up to isomorphism.

For $i > 0$, denote by s_i the word $stst \dots$ of length i and by t_i the word $tsts \dots$ of length i . Since the parabolic subgroup of W generated by s and t has rank two, all KL-polynomials for this parabolic subgroup are trivial and hence the minimal projective resolution of Δ_w is easy to describe. The Bruhat interval $[e, w]$ in W has the following form:



A minimal projective resolution \mathcal{P}_w^{\bullet} of Δ_w then has the following form:

$$0 \rightarrow P_e\langle -\ell(w) \rangle \rightarrow P_s\langle -\ell(w) + 1 \rangle \oplus P_t\langle -\ell(w) + 1 \rangle \rightarrow \dots \rightarrow P_{s_{\ell(w)-1}}\langle -1 \rangle \oplus P_{t_{\ell(w)-1}}\langle -1 \rangle \rightarrow P_w \rightarrow 0.$$

For any two summands $P_x\langle -i \rangle$ and $P_y\langle -i + 1 \rangle$ in the neighboring positions of this resolution, there is a unique, up to scalar, map from $P_x\langle -i \rangle$ to $P_y\langle -i + 1 \rangle$. Since, in this case, $x \neq y$, there are no non-zero homomorphisms from $P_x\langle -i \rangle$ to $P_y\langle -i \rangle$. Using Backelin's description of homomorphisms between projective functors, this implies that there is a unique, up to scalar, map from $\theta_x\langle -i \rangle$ to $\theta_y\langle -i + 1 \rangle$ as well. Similarly, for any two summands $P_x\langle -i \rangle$ and $P_y\langle -i + 2 \rangle$ in the positions of this resolution with one gap, there is a unique, up to scalar, map from $P_x\langle -i \rangle$ to $P_y\langle -i + 2 \rangle$ and also from $\theta_x\langle -i \rangle$ to $\theta_y\langle -i + 2 \rangle$.

Finally, note that the evaluation map gives rise to a morphism of complexes $G_\bullet \rightarrow \mathcal{P}_w^\bullet$. If we now fix \mathcal{P}_w^\bullet and look at $G_\bullet \rightarrow \mathcal{P}_w^\bullet$, the properties described in previous paragraph identify G_\bullet uniquely, up to isomorphism. This completes the proof. \square

Remark 31. It is interesting to see how the argument used in the above proof relates to naïve computations of compositions of derived shuffling functors. For $w = sts$, the naïve composition followed by taking the total complex evaluates the composition $\mathcal{L}C_s \mathcal{L}C_t \mathcal{L}C_s$ to

$$0 \rightarrow \theta_e \langle -3 \rangle \rightarrow \theta_s \langle -2 \rangle \oplus \theta_t \langle -2 \rangle \oplus \theta_s \langle -2 \rangle \rightarrow \theta_s \theta_t \langle -1 \rangle \oplus \theta_s \theta_s \langle -1 \rangle \oplus \theta_t \theta_s \langle -1 \rangle \rightarrow \theta_s \theta_t \theta_s \rightarrow 0$$

Further, the computation of all individual compositions gives:

$$0 \rightarrow \theta_e \langle -3 \rangle \rightarrow \theta_s \langle -2 \rangle \oplus \theta_t \langle -2 \rangle \oplus \theta_s \langle -2 \rangle \rightarrow \theta_{ts} \langle -1 \rangle \oplus \theta_s \langle -2 \rangle \oplus \theta_s \oplus \theta_{st} \langle -1 \rangle \rightarrow \theta_{sts} \oplus \theta_s \rightarrow 0$$

Since the projective resolution of Δ_w is linear (see, e.g., [Ma09-1, Corollary 2.8]), the indecomposables appearing at homological position i of a minimal representative of the above complex must be shifted by i . In other words, the summands $\theta_s \langle -2 \rangle \oplus \theta_s$ must belong to null-homotopic summands and should be removed. This removal forces the removal of the summands θ_s and $\theta_s \langle -2 \rangle$ resulting in the minimal complex

$$0 \rightarrow \theta_e \langle -3 \rangle \rightarrow \theta_s \langle -2 \rangle \oplus \theta_t \langle -2 \rangle \rightarrow \theta_{ts} \langle -1 \rangle \oplus \theta_{st} \langle -1 \rangle \rightarrow \theta_{sts} \rightarrow 0$$

which is exactly the one that appears in the proof above.

7.2. Singular crosses via projective functors. In type A , the following result can be found in [MaStr07, Theorem 1] and [CM17, Theorem 5.1]. We give a general proof.

Theorem 32. *The assignment $\sigma_s \mapsto \mathcal{L}C_s$ and $\tau_s \mapsto \theta_s$, for $s \in S$, extends uniquely (up to isomorphism) to a weak action of the singular Artin monoid $\mathbf{SB}(W)$ on $\mathcal{D}^b(\mathcal{O}_0^{\mathbb{Z}})$.*

Proof. After Proposition 30, we only need to check the relations involving singular generators. That, under the assumption $st = ts$, we have $\theta_s \theta_t \cong \theta_t \theta_s \cong \theta_{st}$ follows directly from the classification of projective functors.

Now let us check the braid relations for $m_{s,t} > 2$, so we assume that. We do the case of even $m_{s,t}$ and the odd case is similar. Let $w = \mathbf{t}_{m_{s,t}-1}$ in the notation of the proof of Proposition 30. Then our braid relation reads $sw = ws$. We will use the same approach as in the proof of Proposition 30. We show that the functors on the two sides of this relation send Δ_e to the same module, up to isomorphism, and that the projective resolution of this module has similar properties as the projective resolutions of Verma modules used in the proof of Proposition 30.

Consider the functors

$$F = \theta_s \circ \mathcal{L}C_w \quad \text{and} \quad G = \mathcal{L}C_w \circ \theta_s.$$

Lemma 33. *We have $F\Delta_e \cong G\Delta_e \cong \theta_s \Delta_w$.*

Proof. That $F\Delta_e \cong \theta_s \Delta_w$ follows directly from Subsection 6.4. Also from Subsection 6.4 it follows that there is a short exact sequence

$$(37) \quad \Delta_w \langle -1 \rangle \hookrightarrow G\Delta_e \twoheadrightarrow \Delta_{ws}.$$

The module $\theta_s \Delta_e \cong P_s$ is indecomposable. Since $\mathcal{L}C_w$ is an equivalence, we have that $G\Delta_e$ is indecomposable as well. Therefore the sequence in (37) is nonsplit. Now, the fact that $F\Delta_e \cong G\Delta_e$ follows from $\dim \text{Ext}_{\mathcal{O}}^1(\Delta_{ws}, \Delta_w \langle -1 \rangle) = 1$ (see, e.g., [Ma07, Corollary 23]). \square

Lemma 34. *The minimal projective resolution of $\theta_s \Delta_w$ has the following form:*

$$0 \rightarrow P_{tws}\langle -1 \rangle \rightarrow P_{ws} \rightarrow 0.$$

Proof. We start by noting that $\ell(tws) = \ell(ws) - 1$. There is a unique, up to scalar, map from $P_{tws}\langle -1 \rangle$ to P_{ws} .

Note that both w and tws belong to a rank two parabolic subgroup of W . Since the Kazhdan-Lusztig theory in the rank two case is trivial, the multiplicities of Verma modules in $P_{tws}\langle -1 \rangle$ and P_{ws} are either 0 or 1 and are determined by the Bruhat order. This implies that both $P_{tws}\langle -1 \rangle$ and P_{ws} have simple socles (by [St03, Theorem 8.1]) and, consequently, any non-zero map from $P_{tws}\langle -1 \rangle$ to P_{ws} is injective.

By comparing the Verma flags of $P_{tws}\langle -1 \rangle$ and P_{ws} , we see that the images of $\theta_s \Delta_w$ and $P_{ws}/P_{tws}\langle -1 \rangle$ in the Grothendieck group coincide. The module $P_{ws}/P_{tws}\langle -1 \rangle$ has, by construction, simple top. In particular, it is indecomposable. Now, that fact that we have an isomorphism $P_{ws}/P_{tws}\langle -1 \rangle \cong \theta_s \Delta_w$ follows from

$$\dim \text{ext}_{\mathcal{O}}^1(\Delta_{ws}, \Delta_w\langle -1 \rangle) = 1,$$

see, e.g., [Ma07, Corollary 23]. \square

Taking Lemmata 33 and 34 into account, the same reasoning as in the proof of Proposition 30 shows that both F and G have the following minimal representative (where the middle map is unique up to scalar and non-zero) which is unique, up to isomorphism:

$$(38) \quad 0 \rightarrow \theta_{tws}\langle -1 \rangle \rightarrow \theta_{ws} \rightarrow 0.$$

It remains to check the relation $\tau_s \sigma_s = \sigma_s \tau_s$. We have

$$\theta_s \circ \mathcal{L}C_s \Delta_e \cong \mathcal{L}C_s \circ \theta_s \Delta_e \cong P_s\langle 1 \rangle.$$

The argument from the proof of Proposition 30 implies $\theta_s \circ \mathcal{L}C_s \cong \mathcal{L}C_s \circ \theta_s$. This completes the proof. \square

8. CATEGORIFICATION OF THE DESINGULARIZATION MAP

8.1. New candidate for the singular crossing. For a simple reflection s , consider the following morphism ξ in $\mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$, where θ_s is in the homological position 0:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & 0 & \longrightarrow & \theta_e\langle 1 \rangle & \xrightarrow{\text{adj}_s} & \theta_s\langle 2 \rangle & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \theta_s & \xrightarrow{\text{adj}_s^s} & \theta_e\langle 1 \rangle & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

Setting $\alpha_s = \text{adj}_s \text{adj}_s^s$, we obtain that the complex

$$(39) \quad 0 \longrightarrow \theta_s \xrightarrow{\alpha_s} \theta_s\langle 2 \rangle \longrightarrow 0$$

is a minimal representative for the cone of ξ . Note that the identity map on $\theta_e\langle 1 \rangle$ splits off as a direct summand of the cone and hence disappears in this minimal representative. We denote the complex in (39) by $\hat{\theta}_s$.

We note that α_s is non-zero since, when evaluated at the simple module L_s , the adjunction morphism adj_s^s is surjective while the adjunction morphism adj_s is injective.

8.2. Main theorem.

Theorem 35. *The assignment $\sigma_s \mapsto \mathcal{L}C_s$ and $\tau_s \mapsto \hat{\theta}_s$, for $s \in S$, extends uniquely (up to isomorphism) to a weak action of the singular Artin monoid $\mathbf{SB}(W)$ on $\mathcal{D}^b(\mathcal{O}_0^{\mathbb{Z}})$.*

Since $\hat{\theta}_s$ is defined as a cone of a morphism from $\mathcal{L}C_s[-2]\langle 2 \rangle$ to $\mathcal{L}C_s$, one can interpret Theorem 35 as a categorification of the desingularization map.

Note that Relation 1 is proved in Theorem 30, so, to prove Theorem 35, we only need to check Relations 2, 3, 4 and 5. We will do this, one at a time, below.

8.3. Proof: Relation 4. We need to prove that, for $s, t \in S$, we have

$$(40) \quad \hat{\theta}_s \circ \hat{\theta}_t \cong \hat{\theta}_t \circ \hat{\theta}_s$$

provided that $st = ts$. The LHS of (40) is given by taking the total complex of the following commutative diagram:

$$(41) \quad \begin{array}{ccc} \theta_s \theta_t & \xrightarrow{\text{id}_s \circ_h \alpha_t} & \theta_s \theta_t \langle 2 \rangle \\ \alpha_s \circ_h \text{id}_t \downarrow & & \downarrow \alpha_s \circ_h \text{id}_t \langle 2 \rangle \\ \theta_s \theta_t \langle 2 \rangle & \xrightarrow{\text{id}_s \circ_h \alpha_t \langle 2 \rangle} & \theta_s \theta_t \langle 4 \rangle \end{array}$$

Using Soergel's combinatorial description, we can represent this diagram as the following complexes of bimodules over the coinvariant algebra \mathbb{C} :

$$(42) \quad \begin{array}{ccc} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} & \xrightarrow{\text{id}_s \circ_h \alpha_t} & \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \langle 2 \rangle \\ \alpha_s \circ_h \text{id}_t \downarrow & & \downarrow \alpha_s \circ_h \text{id}_t \langle 2 \rangle \\ \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \langle 2 \rangle & \xrightarrow{\text{id}_s \circ_h \alpha_t \langle 2 \rangle} & \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \langle 4 \rangle \end{array}$$

here \mathbb{C}^s and \mathbb{C}^t are the subalgebras of s - and t -invariants in \mathbb{C} , respectively. Let x_s and x_t denote the images in \mathbb{C} of the coroots corresponding to s and t , respectively. Then the map α_s maps the generator $1 \otimes 1$ of $\mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C}$ to $1 \otimes x_s + x_s \otimes 1$. Similarly, the map α_t maps the generator $1 \otimes 1$ of $\mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C}$ to $1 \otimes x_t + x_t \otimes 1$.

Since $st = ts$, we have the unique isomorphism from the indecomposable \mathbb{C} - \mathbb{C} -bimodule $\mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C}$ to the indecomposable \mathbb{C} - \mathbb{C} -bimodule $\mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C}$ which sends $1 \otimes 1 \otimes 1$ to $1 \otimes 1 \otimes 1$. Applying this isomorphism component-wise to (42) and taking the formulae of the previous paragraph into account, we get

$$(43) \quad \begin{array}{ccc} \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} & \xrightarrow{\alpha_t \circ_h \text{id}_s} & \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \langle 2 \rangle \\ \text{id}_t \circ_h \alpha_s \downarrow & & \downarrow \text{id}_t \circ_h \alpha_s \langle 2 \rangle \\ \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \langle 2 \rangle & \xrightarrow{\alpha_t \circ_h \text{id}_s \langle 2 \rangle} & \mathbb{C} \otimes_{\mathbb{C}^t} \mathbb{C} \otimes_{\mathbb{C}^s} \mathbb{C} \langle 4 \rangle \end{array}$$

which represents the RHS of (40) (note that the two middle bimodules also got swapped compared to (42)).

8.4. Proof: Relation 5. We need to prove that

$$(44) \quad \hat{\theta}_s \circ \mathcal{L}C_s \cong \mathcal{L}C_s \circ \hat{\theta}_s.$$

The LHS of (44) is given by taking the total complex of the following commutative diagram:

$$(45) \quad \begin{array}{ccc} \theta_s \langle -1 \rangle & \xrightarrow{\text{id}_s \circ_h \text{adj}_s} & \theta_s \theta_s \\ \alpha_s \langle -1 \rangle \downarrow & & \downarrow \alpha_s \circ_h \text{id}_s \\ \theta_s \langle 1 \rangle & \xrightarrow{\text{id}_s \circ_h \text{adj}_s \langle 2 \rangle} & \theta_s \theta_s \langle 2 \rangle \end{array}$$

Using Soergel's combinatorial description, we can look at the image of (45) in the homotopy category of bimodules over the coinvariant algebra.

Let us first consider the case of the Lie algebra \mathfrak{sl}_2 . In this case, the coinvariant algebra is isomorphic to the algebra $D = \mathbb{C}[x]/(x^2)$ of dual numbers, graded such that $\deg(x) = 2$. The identity functor θ_e is represented by the regular D - D -bimodule D graded naturally. The functor θ_s is represented by the D - D -bimodule $D \otimes_{\mathbb{C}} D$, graded such that $\deg(1 \otimes 1) = -1$. The adjunction morphism $\text{adj}_s^s : \theta_s \rightarrow \theta_e \langle 1 \rangle$ is given by the multiplication map while the adjunction morphism $\text{adj}_s^s : \theta_e \langle -1 \rangle \rightarrow \theta_s$ sends 1 to $1 \otimes x + x \otimes 1$. This means that the map α_s is the endomorphism of $D \otimes_{\mathbb{C}} D$ sending $1 \otimes 1$ to $1 \otimes x + x \otimes 1$. The two ways around the diagram in (45) send $1 \otimes 1$ to $x \otimes x \otimes 1 + x \otimes 1 \otimes x + 1 \otimes x \otimes x$.

The upper map of the diagram sends $1 \otimes 1$ to $1 \otimes 1 \otimes x + 1 \otimes x \otimes 1$. The latter element generates a summand of $\theta_s \theta_s$, call it X . We also have the corresponding summand of $\theta_s \theta_s \langle 2 \rangle$, call it Y . These two summands, together with the left part of the diagram, form a subdiagram (that is they are closed with respect to all maps on the diagram). By construction, the total complex of this subdiagram is homotopic to zero.

The image of the element $1 \otimes 1 \otimes 1$ under the right vertical map is $x \otimes 1 \otimes 1 + 1 \otimes x \otimes 1$. When factoring out the (homotopic to zero) subcomplex from the previous paragraph, we send $1 \otimes 1 \otimes x + 1 \otimes x \otimes 1$ to zero, implying that

$$x \otimes 1 \otimes 1 + 1 \otimes x \otimes 1 = x \otimes 1 \otimes 1 - 1 \otimes 1 \otimes x.$$

This means that the total complex of (45) is represented in the homotopy category of complexes by the following complex of D - D -bimodules:

$$(46) \quad 0 \longrightarrow D \otimes_{\mathbb{C}} D \xrightarrow{\beta_s} D \otimes_{\mathbb{C}} D \longrightarrow 0,$$

where the map β_s sends $1 \otimes 1$ to $x \otimes 1 - 1 \otimes x$.

The same computation for the RHS of (45) results in the complex

$$(47) \quad 0 \longrightarrow D \otimes_{\mathbb{C}} D \xrightarrow{\gamma_s} D \otimes_{\mathbb{C}} D \longrightarrow 0,$$

where the map γ_s sends $1 \otimes 1$ to $1 \otimes x - x \otimes 1$. Since $\beta_s = -\gamma_s$, the complexes in (46) and (47) are isomorphic, for example, by choosing the identity map between the left non-zero components and minus the identity map between the right non-zero components. This completes the proof in the case of the algebra \mathfrak{sl}_2 .

The general case is similar. One uses that the coinvariant algebra \mathbb{C} is a free module over the subalgebra \mathbb{C}^s of s -invariants with the basis given by 1 and the coroot corresponding to s (which we can denote by x). Then all the above arguments go through, mutatis mutandis.

8.5. Proof: Relations 2 and 3. We prove the relation given by Equation 3. The proof of the relation given by Equation 2 is similar. We assume that $m_{s,t}$ is finite and even and need to prove that

$$(48) \quad \hat{\theta}_s \circ \underbrace{\mathcal{L}C_t \circ \mathcal{L}C_s \circ \cdots \circ \mathcal{L}C_t}_{m_{s,t}-1 \text{ factors}} \cong \underbrace{\mathcal{L}C_t \circ \mathcal{L}C_s \circ \cdots \circ \mathcal{L}C_t}_{m_{s,t}-1 \text{ factors}} \circ \hat{\theta}_s.$$

Let us take a look at the LHS of (48). Set

$$w = \underbrace{sts\dots t}_{m_{s,t} \text{ factors}} = \underbrace{tst\dots s}_{m_{s,t} \text{ factors}}.$$

From (38), it follows that the LHS of (48) is given by the total complex of the following complex:

$$(49) \quad \begin{array}{ccc} \theta_{sw} & \xrightarrow{\gamma_1} & \theta_w\langle 1 \rangle, \\ \gamma_2 \downarrow & & \downarrow \gamma_4 \\ \theta_{sw}\langle 2 \rangle & \xrightarrow{\gamma_3} & \theta_w\langle 3 \rangle \end{array}$$

for some natural transformations γ_i , with γ_1 and γ_3 non-zero. Furthermore, from [Bac92], see Subsection 6.4, it follows that $\dim \mathcal{P}^{\mathbb{Z}}(\theta_{sw}, \theta_w\langle 1 \rangle) = 1$. The projective module P_{sw} is a submodule of P_w , which implies that the maps γ_1 and γ_3 are injections.

By construction of α_s , the map γ_4 factors through θ_{ws} . Again, [Bac92], see Subsection 6.4, it follows that $\dim \mathcal{P}^{\mathbb{Z}}(\theta_w, \theta_{ws}\langle 1 \rangle) = 1$ and $\dim \mathcal{P}^{\mathbb{Z}}(\theta_{ws}, \theta_w\langle 1 \rangle) = 1$. This means that γ_4 is unique up to scalar. Since γ_3 is an injection, it follows that commutativity of (49) defines the map γ_2 uniquely (as soon as the scalar for γ_4 is fixed).

It remains to note that the RHS of (48) is also given by the total complex of a commutative diagram of the form (49) (possibly for different choices of the γ_i 's). However, we have just established that the only real choices in (49) are scalars. This implies (48).

8.6. Another candidate that fails. There is another obvious candidate for categorification of the desingularization map. For a simple reflection s , consider the following morphism ξ' in $\mathcal{K}^b(\mathcal{P}^{\mathbb{Z}})$, where θ_s is in the homological position 0:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & 0 & \longrightarrow & \theta_s & \xrightarrow{\text{adj}_s^s} & \theta_e\langle 1 \rangle & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \theta_e\langle -1 \rangle & \xrightarrow{\text{adj}_s} & \theta_s & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

Setting $\beta_s = \text{adj}_s^s \text{adj}_s$, we obtain that the complex

$$(50) \quad 0 \longrightarrow \theta_e\langle -1 \rangle \xrightarrow{\beta_s} \theta_e\langle 1 \rangle \longrightarrow 0$$

is a minimal representative for the cone of ξ' . Note that the identity map on θ_s splits off as a direct summand of the cone and hence disappears in this minimal representative. We denote the complex in (50) by $\check{\theta}_s$. The following negative result shows that sending τ_s to $\check{\theta}_s$ does not define a weak action of the singular braid monoid.

Proposition 36. *For $s \in S$, we have $\check{\theta}_s \circ \mathcal{L}C_s \not\cong \mathcal{L}C_s \circ \check{\theta}_s$.*

Proof. It is enough to consider the case of the Lie algebra \mathfrak{sl}_2 . Again, we interpret all objects using bimodules over the dual numbers D . Then the map β_s is given by the bimodule homomorphism $D \rightarrow D$ sending 1 to $2x$.

The composition $\check{\theta}_s \circ \mathcal{LC}_s$ is then given by the total complex of the following commutative diagram:

$$\begin{array}{ccc} D\langle -2 \rangle & \xrightarrow{\gamma_1} & D \otimes_{\mathbb{C}} D\langle -1 \rangle \\ \gamma_2 \downarrow & & \downarrow \gamma_4 \\ D & \xrightarrow{\gamma_3} & D \otimes_{\mathbb{C}} D\langle 1 \rangle \end{array}$$

Here

- γ_1 and γ_3 send 1 to $1 \otimes x + x \otimes 1$,
- γ_2 sends 1 to $2x$,
- γ_4 sends $1 \otimes 1$ to $2x \otimes 1$.

In turn, the composition $\mathcal{LC}_s \circ \check{\theta}_s$ is given by the total complex of the following commutative diagram:

$$\begin{array}{ccc} D\langle -2 \rangle & \xrightarrow{\gamma_1} & D \otimes_{\mathbb{C}} D\langle -1 \rangle \\ \gamma_2 \downarrow & & \downarrow \gamma'_4 \\ D & \xrightarrow{\gamma_3} & D \otimes_{\mathbb{C}} D\langle 1 \rangle \end{array}$$

Here γ_1 , γ_2 and γ_3 are as in the previous paragraph, while γ'_4 sends $1 \otimes 1$ to $1 \otimes 2x$.

Since γ_4 and γ'_4 are linearly independent, it follows that the two total complexes are not isomorphic. \square

REFERENCES

- [An04] Antony, N. On the injectivity of the Vassiliev homomorphism of singular Artin monoids. *Bull. Austral. Math. Soc.* **70** (2004), no. 3, 401–422.
- [An05] Antony, N. On singular Artin monoids and contributions to Birman's conjecture. *Comm. Algebra* **33** (2005), no. 11, 4043–4056.
- [Bac92] Backelin, E. The Hom-spaces between projective functors. *Represent. Theory* **5** (2001), 267–283.
- [Bae92] Baez, J. Link invariants of finite type and perturbation theory. *Lett. Math. Phys.* **26** (1992), no. 1, 43–51.
- [BB81] Beilinson, A.; Bernstein, J. Localisation de \mathfrak{g} -modules. *C. R. Acad. Sci. Paris Ser. I Math.* **292** (1981), no. 1, 15–18.
- [BG80] Bernstein, J.; Gelfand, S. Tensor products of finite- and infinite-dimensional representations of semisimple Lie algebras. *Compositio Math.* **41** (1980), no. 2, 245–285.
- [BGG75] Bernstein, I.; Gelfand, I.; Gelfand, S. Differential operators on the base affine space and a study of \mathfrak{g} -modules. *Lie groups and their representations (Proc. Summer School, Bolyai Janos Math. Soc., Budapest, 1971)*, pp. 21–64. Halsted, New York, 1975.
- [BGG76] Bernstein, I.; Gelfand, I.; Gelfand, S. A certain category of \mathfrak{g} -modules. (Russian) *Funkcional. Anal. i Priložen.* **10** (1976), no. 2, 1–8.
- [Bi93] Birman, J. New points of view in knot theory. *Bull. Amer. Math. Soc. (N.S.)* **28** (1993), no. 2, 253–287.
- [Br37] Brauer, R. On algebras which are connected with the semisimple continuous groups. *Ann. of Math. (2)* **38** (1937), no. 4, 857–872.
- [BK81] Brylinski, J.-L.; Kashiwara, M. Kazhdan-Lusztig conjecture and holonomic systems. *Invent. Math.* **64** (1981), no. 3, 387–410.
- [Ca86] Carlin, K. Extensions of Verma modules. *Trans. Amer. Math. Soc.* **294** (1986), no. 1, 29–43.
- [CE22] Clark, C.; East, J. Presentations for wreath products involving symmetric inverse monoids and categories. Preprint arXiv:2204.06992.
- [C-Z18] Chen, C.-W.; Frisk Dubsky, B.; Jonsson, H.; Mazorchuk, V.; Persson Westin, E.; Zhang, X.; Zimmermann, J. Extreme representations of semirings. *Serdica Math. J.* **44** (2018), no. 3-4, 365–412.
- [CL15] Cohen, A.; Liu, S. Brauer algebras of type B. *Forum Math.* **27** (2015), no. 2, 1163–1202.

- [Co00] Corran, R. A normal form for a class of monoids including the singular braid monoids. *J. Algebra* **223** (2000), no. 1, 256–282.
- [Co05] Corran, R. Conjugacy in singular Artin monoids. *J. Aust. Math. Soc.* **79** (2005), no. 2, 183–212.
- [CM17] Coulembier, K.; Mazorchuk, V. Dualities and derived equivalences for category \mathcal{O} . *Israel J. Math.* **219** (2017), no. 2, 661–706.
- [EEF05] Easdown, D.; East, J.; FitzGerald, D. Presentations of factorizable inverse monoids. *Acta Sci. Math. (Szeged)* **71** (2005), no. 3-4, 509–520.
- [Ea06a] East, J. Birman’s conjecture is true for $I_2(p)$. *J. Knot Theory Ramifications* **15** (2006), no. 2, 167–177.
- [Ea06b] East, J. The factorizable braid monoid. *Proc. Edinb. Math. Soc. (2)* **49** (2006), no. 3, 609–636.
- [Ea15] East, J. Singular braids and partial permutations. *Acta Sci. Math. (Szeged)* **81** (2015), no. 1-2, 55–77.
- [EF13] Everitt, B.; Fountain, J. Partial mirror symmetry, lattice presentations and algebraic monoids. *Proc. Lond. Math. Soc. (3)* **107** (2013), no. 2, 414–450.
- [Fi03] FitzGerald, D. G. A presentation for the monoid of uniform block permutations. *Bull. Austral. Math. Soc.* **68** (2003), no. 2, 317–324.
- [FL06] FitzGerald, D.; Leech, J. Dual symmetric inverse monoids and representation theory. *J. Austral. Math. Soc. Ser. A* **64** (1998), no. 3, 345–367.
- [GM93] Ganyushkin, O.; Mazorchuk, V. Factor powers of semigroups of transformations. (Ukrainian) *Dopov./Dokl. Akad. Nauk Ukraini* **1993**, no. 12, 5–9 (1994).
- [GM95a] Ganyushkin, O.; Mazorchuk, V. Factor powers of finite symmetric groups. (Russian) *Mat. Zametki* **58** (1995), no. 2, 176–188; translation in *Math. Notes* **58** (1995), no. 1-2, 794–802 (1996)
- [GM95b] Ganyushkin, O.; Mazorchuk, V. The structure of subsemigroups of factor powers of finite symmetric groups. (Russian) *Mat. Zametki* **58** (1995), no. 3, 341–354, 478; translation in *Math. Notes* **58** (1995), no. 3-4, 910–920 (1996)
- [GM03] Ganyushkin, O.; Mazorchuk, V. On the radical of $\mathcal{FP}^+(S_n)$. *Mat. Stud.* **20** (2003), no. 1, 17–26.
- [GM09] Ganyushkin, O.; Mazorchuk, V. *Classical finite transformation semigroups. An introduction. Algebra and Applications*, **9**. Springer-Verlag London, Ltd., London, 2009.
- [GP05] Goddelle, E.; Paris, L. On singular Artin monoids. *Geometric methods in group theory*, 43–57, *Contemp. Math.*, **372**, Amer. Math. Soc., Providence, RI, 2005.
- [Hu10] Hudzenko, S. V. Automorphisms of a finitary factor power of an infinite symmetric group. *Ukrainian Math. J.* **62** (2010), no. 7, 1158–1162; translated from *Ukrain. Mat. Zh.* **62** (2010), no. 7, 997–1001.
- [Hu72] Humphreys, J. *Introduction to Lie algebras and representation theory. Graduate Texts in Mathematics, Vol. 9*. Springer-Verlag, New York-Berlin, 1972. xii+169 pp.
- [Hu08] Humphreys, J. *Representations of semisimple Lie algebras in the BGG category \mathcal{O} . Graduate Studies in Mathematics*, **94**. American Mathematical Society, Providence, RI, 2008. xvi+289 pp.
- [Ja17] Jabłonowski, M. Presentations and representations of surface singular braid monoids. *J. Korean Math. Soc.* **54** (2017), no. 3, 749–762.
- [KL79] Kazhdan, D.; Lusztig, G. Representations of Coxeter groups and Hecke algebras. *Invent. Math.* **53** (1979), no. 2, 165–184.
- [KS02] Khovanov, M.; Seidel, P. Quivers, Floer cohomology, and braid group actions. *J. Amer. Math. Soc.* **15** (2002), no. 1, 203–271.
- [Ko06] Kosuda, M. Irreducible representations of the party algebra. *Osaka J. Math.* **43** (2006), no. 2, 431–474.
- [KM11] Kudryavtseva, G.; Maltcev, V. Two generalisations of the symmetric inverse semigroups. *Publ. Math. Debrecen* **78** (2011), no. 2, 253–282.
- [KMU15] Kudryavtseva, G.; Maltcev, V.; Umar, A. Presentation for the partial dual symmetric inverse monoid. *Comm. Algebra* **43** (2015), no. 4, 1621–1639.
- [KMaz06] Kudryavtseva, G.; Mazorchuk, V. On presentations of Brauer-type monoids. *Cent. Eur. J. Math.* **4** (2006), no. 3, 413–434.
- [Mar94] Martin, P., Temperley-Lieb algebras for nonplanar statistical mechanics – the partition algebra construction, *J. Knot Theory Ramifications* **3** (1994), 51–82.
- [MM13] Martin, P.; Mazorchuk, V. Partitioned binary relations. *Math. Scand.* **113** (2013), no. 1, 30–52.
- [MM14] Martin, P.; Mazorchuk, V. On the representation theory of partial Brauer algebras. *Q. J. Math.* **65** (2014), no. 1, 225–247.

- [Ma98] Mazorchuk, V. On the structure of Brauer semigroup and its partial analogue, *Problems in Algebra*, **13** (1998), 29–45.
- [Ma00] Mazorchuk, V. All automorphisms of $\mathcal{FP}^+(S_n)$ are inner. *Semigroup Forum* **60** (2000), no. 3, 486–490.
- [Ma07] Mazorchuk, V. Some homological properties of the category \mathcal{O} . *Pacific J. Math.* **232** (2007), no. 2, 313–341.
- [Ma09-1] Mazorchuk, V. Applications of the category of linear complexes of tilting modules associated with the category \mathcal{O} . *Algebr. Represent. Theory* **12** (2009), no. 6, 489–512.
- [Ma09-2] Mazorchuk, V. Simple modules over factorpowers. *Acta Sci. Math. (Szeged)* **75** (2009), no. 3-4, 467–485.
- [MaSr21] Mazorchuk, V.; Srivastava, S. Jucys-Murphy elements and Grothendieck groups for generalized rook monoids. Preprint arXiv:2104.13632. To appear in the *Journal of Combinatorial Algebra*.
- [MaSr22] Mazorchuk, V.; Srivastava, S. Multiparameter colored partition category and the product of reduced Kronecker coefficients. Preprint arXiv:2204.13564.
- [MaSte12] Mazorchuk, V.; Steinberg, B. Double Catalan monoids. *J. Algebraic Combin.* **36** (2012), no. 3, 333–354.
- [MaStr05] Mazorchuk, V.; Stroppel, C. Translation and shuffling of projectively presentable modules and a categorification of a parabolic Hecke module. *Trans. Amer. Math. Soc.* **357** (2005), no. 7, 2939–2973.
- [MaStr07] Mazorchuk, V.; Stroppel, C. On functors associated to a simple root. *J. Algebra* **314** (2007), no. 1, 97–128.
- [OSSZ21] Orellana, R.; Saliola, F.; Schilling, A.; Zabrocki, M. Plethysm and the algebra of uniform block permutations. Preprint arXiv:2112.13909.
- [Pa04] Paris, L. The proof of Birman’s conjecture on singular braid monoids. *Geom. Topol.* **8** (2004), 1281–1300.
- [PW70] Plemmons, R.; West, M. On the semigroup of binary relations. *Pacific J. Math.* **35** (1970), 743–753.
- [Ro06] Rouquier, R. Categorification of \mathfrak{sl}_2 and braid groups. *Trends in representation theory of algebras and related topics*, 137–167, *Contemp. Math.*, **406**, Amer. Math. Soc., Providence, RI, 2006.
- [So90] Soergel, W. Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. *J. Amer. Math. Soc.* **3** (1990), no. 2, 421–445.
- [So07] Soergel, W. Kazhdan-Lusztig-Polynome und unzerlegbare Bimoduln über Polynomringen. *J. Inst. Math. Jussieu* **6** (2007), no. 3, 501–525.
- [St03] Stroppel, C. Category \mathcal{O} : gradings and translation functors. *J. Algebra* **268** (2003), no. 1, 301–326.
- [Ve10] Vershinin, V. On the singular braid monoid. *St. Petersburg Math. J.* **21** (2010), no. 5, 693–704.

H. J.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: helena.jonsson@math.uu.se

V. M.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: mazor@math.uu.se

E. P. W.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: elin.persson.westin@math.uu.se

S. S.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: shraddha.srivastava@math.uu.se

M. S.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: mateusz.stroinski@math.uu.se

X. Z.: School of Mathematical Sciences, Tongji University, Shanghai 200092, P. R. CHINA, email: 1810079@tongji.edu.cn