

Nonnegative scalar curvature on manifolds with at least two ends

Simone Cecchini^{*1}, Daniel Råde^{†2}, and Rudolf Zeidler^{‡3}

¹University of Göttingen, Mathematisches Institut, Germany,
simone.cecchini@mathematik.uni-goettingen.de, simonececchini.org

²Augsburg University, Institut für Mathematik, Germany, daniel.raede@math.uni-augsburg.de

³University of Münster, Mathematisches Institut, Germany, math@rzeidler.eu, www.rzeidler.eu

Abstract

Let M be an orientable connected n -dimensional manifold with $n \in \{6, 7\}$ and let $Y \subset M$ be a two-sided closed connected incompressible hypersurface which does not admit a metric of positive scalar curvature (abbreviated by psc). Moreover, suppose that the universal covers of M and Y are either both spin or both non-spin. Using Gromov’s μ -bubbles, we show that M does not admit a complete metric of psc. We provide an example showing that the spin/non-spin hypothesis cannot be dropped from the statement of this result. This answers, up to dimension 7, a question by Gromov for a large class of cases. Furthermore, we prove a related result for submanifolds of codimension two. We deduce as special cases that, if Y does not admit a metric of psc and $\dim(Y) \neq 4$, then $M := Y \times \mathbb{R}$ does not carry a complete metric of psc and $N := Y \times \mathbb{R}^2$ does not carry a complete metric of uniformly psc provided that $\dim(M) \leq 7$ and $\dim(N) \leq 7$, respectively. This solves, up to dimension 7, a conjecture due to Rosenberg and Stolz in the case of orientable manifolds.

1 Introduction

In the 1994 survey article [28, Section 7], Rosenberg and Stolz proposed the following conjectures concerning the (non-)existence of *positive scalar curvature* (abbreviated by *psc*) metrics on certain non-compact manifolds:

Conjecture 1.1. Let Y be a closed manifold of dimension $(n - 1) \neq 4$ which does not admit a metric of positive scalar curvature. Then $Y \times \mathbb{R}$ does not admit a complete metric of positive scalar curvature.

Conjecture 1.2. Let Y be a closed manifold of dimension $(n - 2) \neq 4$ which does not admit a metric of positive scalar curvature. Then $Y \times \mathbb{R}^2$ does not admit a complete metric with uniformly positive scalar curvature.

Recently Gromov phrased a related conjecture [14, 11.12, Conjecture C] which is a cornerstone of his program regarding the study of metric inequalities with scalar curvature:

Conjecture 1.3. Let Y be a closed manifold of dimension $(n - 1) \neq 4$ which does not admit a metric of positive scalar curvature and $X = Y \times [-1, 1]$. If g is a Riemannian metric on X with $\text{scal}(X, g) \geq n(n - 1)$, then

$$\text{width}(X, g) := \text{dist}_g(Y \times \{-1\}, Y \times \{1\}) \leq \frac{2\pi}{n}.$$

^{*}Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Priority Programme “Geometry at Infinity” (SPP 2026, CE 393/1-1).

[†]Funded by a Doctoral Scholarship of the ‘Studienstiftung des deutschen Volkes’.

[‡]Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics–Geometry–Structure, and through the Priority Programme “Geometry at Infinity” (SPP 2026, ZE 1123/2-2).

Remark 1.4. The condition $\dim(Y) \neq 4$ (which we have added in our formulation of Conjectures 1.1 and 1.2) is necessary since in this case there are well known counterexamples using Seiberg–Witten obstructions to positive scalar curvature. It is possible to show that there exists a closed simply connected 4-manifold Y which does not admit a metric of positive scalar curvature while $Y \times S^1$ does (see [29, Counterexample 4.16]). But then $Y \times \mathbb{R}$ (and consequently $Y \times \mathbb{R}^2$) admits a complete metric with uniformly positive scalar curvature which would violate Conjectures 1.1 to 1.3.

As is common in the study of scalar curvature, there are two broad families of methods to approach these conjectures: One is based on the spinor Dirac operator on spin manifolds originating in the work of Lichnerowicz [21]. Indeed, using different variants of index theory on non-compact manifolds, it is now well established that Conjectures 1.1 and 1.2 can be proved whenever Y admits an index-theoretic obstruction to positive scalar curvature such as the Rosenberg index [27], see [1; 17; 35]. The other family of methods is based on geometric measure theory and originates in the minimal hypersurface method of Schoen and Yau [31]. The first established cases of Conjecture 1.3 by Gromov [14, Section 2] used the classical minimal hypersurface method. Subsequently also a Dirac operator approach to Conjecture 1.3 was developed by Cecchini and Zeidler [2; 3; 36; 37].

In [15, Sections 3.6, 5], Gromov proposes a different approach towards Conjecture 1.3 using a modified version of the minimal hypersurface method involving so called μ -bubbles. Following this idea, Råde [25] proved Conjecture 1.3 and generalizations thereof in case that Y is orientable and $n \leq 7$. Conjectures 1.1 and 1.2 have not been directly approached via minimal hypersurface techniques so far, in particular due to the non-compactness inherent to the problem. However, in recent work of various authors (e.g. [6; 19; 20; 39]), μ -bubbles have turned out to be a useful tool to deal with non-compact situations. Even though there is no direct formal implication between Conjecture 1.1 and Conjecture 1.3, it was observed in [37] that the Dirac operator methods used in [2; 36] to attack Conjecture 1.3 can be refined to prove a more general statement (compare [37, Conjecture 7.1]) which imply both Conjectures 1.1 and 1.3 for closed spin manifolds with non-vanishing Rosenberg index.

The main objective of this article is to combine the ideas from [37] with μ -bubble methods, in particular from [25], to prove a generalization of Conjecture 1.1 in case Y is orientable and of dimension ≤ 6 . By a reduction argument to a codimension one situation (compare [13, Theorem 7.5; 17]), we also establish Conjecture 1.2 in case Y is orientable and of dimension ≤ 5 . We note that ideas related to Conjecture 1.1 have also appeared recently in the work of Chen, Liu, Shi, and Zhu [5] in connection with the positive mass theorem.

More generally than Conjecture 1.1, one may ask under which circumstances the existence of a hypersurface $Y \subset M$ which does not admit psc already is an obstruction to the existence of a complete psc metric on the ambient manifold M . This question has been discussed by Gromov in [14, §11.6], where in particular he asked if it would be enough to assume that Y is a two-sided *incompressible hypersurface*, that is, the map $\pi_1 Y \rightarrow \pi_1 M$ induced by the inclusion is injective. In the case that the ambient manifold is spin of dimension $n \in \{6, 7\}$ and under further geometric conditions, a proof confirming this was sketched in [14, pp. 708 sqq.] and it was asked if the spin hypothesis can be dropped. We answer this question in our first theorem below together with the following Example 1.6. Here we say that a connected manifold M is *almost spin*, if its universal covering \tilde{M} is spin, and we say it is *totally non-spin* if \tilde{M} is non-spin. Since spin structures lift to coverings, being almost spin is equivalent to the existence of some covering which is spin. Unless explicitly stated otherwise, we consider manifolds without boundary.

Theorem 1.5. *Let M be an orientable connected n -dimensional manifold with $n \in \{6, 7\}$ and let $Y \subset M$ be a two-sided closed connected incompressible hypersurface which does not admit a metric of positive scalar curvature. Suppose that one of the following two conditions holds:*

- (a) M is almost spin.

(b) Y is totally non-spin.

Then M does not admit a complete metric of positive scalar curvature. More precisely, if g is a complete metric of non-negative scalar curvature on M , then (M, g) admits a connected Riemannian covering isometric to $(N \times \mathbb{R}, g_N + dx^2)$, where (N, g_N) is a closed Ricci flat manifold.

Note that if M is almost spin, then any two-sided hypersurface $Y \subset M$ is almost spin itself. Conversely, if a two-sided hypersurface is totally non-spin, then so is the ambient manifold. Thus the alternative hypotheses (a) and (b) of Theorem 1.5 simply say that either M and Y are both almost spin or neither is. The following example shows that this restriction cannot be dropped.

Example 1.6. Fix $n \geq 6$. Let L be the K3 surface, that is, a 4-dimensional simply-connected spin manifold such that $\widehat{A}(L) \neq 0$. Consider the closed manifold $M := (L \times \mathbb{T}^{n-4}) \# (\mathbb{C}P^2 \times S^{n-4})$. Then, since L is oriented cobordant to a psc manifold by [12, §3], the cobordism class represented by M in $\Omega_n^{\text{SO}}(\text{B}\mathbb{Z}^{n-4})$ has a psc representative. Thus the totally non-spin manifold M itself admits a psc metric, e.g. by [26, Theorem 2.13] or [8, Theorem 1.5].¹ It contains $Y = L \times \mathbb{T}^{n-5}$ as an incompressible hypersurface which does not admit psc by [13, Corollary 5.22]. By passing to the covering $\widehat{M} \rightarrow M$ with $\pi_1 \widehat{M} = \mathbb{Z}^{n-5}$ corresponding to $\pi_1 Y$, we even find an example of a complete manifold which admits a complete uniform psc metric although it contains a closed incompressible *separating* hypersurface which does not.

This shows that the almost spin condition is fundamentally relevant for this kind of problem even though the proof Theorem 1.5 does not use the Dirac operator at all. Moreover, even in the spin case, Theorem 1.5 is stronger (in the dimension range where it applies) than what can be proved using index-theoretic techniques because there are examples of closed spin manifolds which do not admit psc even though all known index invariants vanish [30].

The upper dimension bound $n \leq 7$ comes from the usual problem pertaining to singularities of minimal hypersurfaces and μ -bubbles. Thus, if this issue was resolved, the upper dimensional bound in Theorem 1.5 (and in all other results of this paper) should conjecturally be removable. The case $n = 8$ likely is more easily accessible via an adaptation of the work of N. Smale [32].

Remark 1.7. In [14, §11.6], Gromov conjectured that in general, given a complete Riemannian manifold M , the existence of a two-sided incompressible closed embedded hypersurface which does not carry a psc metric obstructs the existence of a function h on M such that

$$\frac{n}{n-1}h^2 - 2|dh| + \text{scal}_g \geq 0.$$

This condition is motivated by the potential function used to construct μ -bubbles. From this point of view, Theorem 1.5 gives a complete description of the case when $h = 0$, which includes many geometrically interesting cases. We point out that our method can be adapted to treat cases with non-trivial h . In the direction of Gromov’s motivating example [14, pp. 708 sqq.], compare Corollary 3.7. We also point out that, compared to the method sketched in [14, §11.6], our technique does not require any extra assumption on the geometry of the manifold.

For $n = 5$, Theorem 1.5 fails as mentioned in Remark 1.4. On the other hand, for $2 \leq n \leq 4$, it holds even independently of the hypotheses (a) and (b), but for a different reason than in high dimensions. To formulate this, we consider the following notion:

Definition 1.8 (compare [25, Definition 2.20; 5, “ \mathcal{C}_{deg} ”). A closed connected oriented manifold Y is called NPSC⁺ if it satisfies the following property: No closed oriented manifold Z which admits a continuous map of non-zero degree $Z \rightarrow Y$ admits a metric of positive scalar curvature.

¹Alternatively, the existence of a psc metric on M can also be deduced from [23, Theorem 5.6].

For instance, it has recently been shown [6; 7; 16] that closed oriented aspherical manifolds of dimension ≤ 5 are NPSC⁺. Moreover, a closed oriented manifold of dimension ≤ 3 which does not admit psc necessarily admits a non-zero degree map to an aspherical NPSC⁺ manifold.² Thus the low-dimensional counterpart to Theorem 1.5 is contained in the following theorem which already appeared recently in [5].

Theorem 1.9 (compare [5, Theorem 1.1]). *Let M be an orientable connected n -dimensional manifold with $n \leq 7$ and let $\iota: Y \hookrightarrow M$ be a two-sided closed hypersurface that admits a map of non-zero degree $\phi: Y \rightarrow Y_0$ to an aspherical NPSC⁺ manifold Y_0 and such that $\ker(\pi_1 Y \xrightarrow{\iota_*} \pi_1 M) \subseteq \ker(\pi_1 Y \xrightarrow{\phi_*} \pi_1 Y_0)$. Then M does not admit a complete metric of positive scalar curvature. More precisely, if g is a complete metric of non-negative scalar curvature on M , then (M, g) admits a connected Riemannian covering isometric to $(N \times \mathbb{R}, g_N + dx^2)$, where (N, g_N) is a closed Ricci flat manifold.*

In particular, applying Theorems 1.5 and 1.9 to $M = Y \times \mathbb{R}$, we deduce:

Corollary 1.10. *Conjecture 1.1 holds for orientable manifolds in dimensions $5 \neq n \leq 7$.*

However, we note that the low-dimensional cases $n \leq 4$ of Corollary 1.10 were already known because an oriented manifold of dimension ≤ 3 which does not admit positive scalar curvature is necessarily spin and has rationally non-vanishing Rosenberg index, and so the situation is within the scope of index-theoretic results such as [1, Theorem A].

We now turn to our results corresponding to Conjecture 1.2:

Theorem 1.11. *Let M be an orientable connected 7-dimensional manifold and let $Y \subset M$ be a closed connected 5-dimensional submanifold with trivial normal bundle such that the inclusion induces an injection $\pi_1 Y \hookrightarrow \pi_1 M$ and a surjection $\pi_2 Y \twoheadrightarrow \pi_2 M$. Suppose that Y does not admit a metric of positive scalar curvature. Then M does not admit a complete metric of uniformly positive scalar curvature.*

Note that, unlike Theorem 1.5, we do not need to impose any conditions involving spin structures. The intuitive reason for this is that the hypotheses already imply that the induced map $\tilde{Y} \rightarrow \tilde{M}$ between universal covers is 2-connected and so M is almost spin if and only if Y is. On the other hand, the surjectivity condition $\pi_2 Y \twoheadrightarrow \pi_2 M$ cannot be omitted as the example $M = Y \times S^2$ shows.

We also have a codimension 2 counterpart to Theorem 1.9 exploiting the NPSC⁺ property.

Theorem 1.12. *Let M be an orientable connected n -dimensional manifold with $n \leq 7$ and let $Y \subset M$ be a closed connected $(n - 2)$ -dimensional submanifold with trivial normal bundle such that the inclusion induces an injection $\pi_1 Y \hookrightarrow \pi_1 M$ and a surjection $\pi_2 Y \twoheadrightarrow \pi_2 M$. Suppose that Y admits a map of non-zero degree $Y \rightarrow Y_0$ to an aspherical NPSC⁺ manifold Y_0 . Then M does not admit a complete metric of uniformly positive scalar curvature.*

Similarly as before, this includes a version of Theorem 1.11 in dimensions $n \leq 5$:

Corollary 1.13. *Let $6 \neq n \leq 7$. Let M be an orientable connected n -dimensional manifold and let $Y \subset M$ be a closed connected $(n - 2)$ -dimensional submanifold with trivial normal bundle such that the inclusion induces an injection $\pi_1 Y \hookrightarrow \pi_1 M$ and a surjection $\pi_2 Y \twoheadrightarrow \pi_2 M$. Suppose that Y does not admit a metric of positive scalar curvature. Then M does not admit a complete metric of uniformly positive scalar curvature.*

²In dimension 2 this is a consequence of the classification of surfaces and the Gauß–Bonnet theorem, whereas in dimension 3 it is a consequence of the classification of 3-manifolds which admit psc following from Perelman’s work, see e.g. the discussion in [22].

Proof. For $n = 7$, this is a restatement of Theorem 1.11. If $n \leq 5$, then $\dim(Y) \leq 3$ and so it admits a non-zero degree map $Y \rightarrow Y_0$, where Y_0 is aspherical and NPSC⁺. Thus Theorem 1.12 is applicable in this case. \square

Specializing to $M = Y \times \mathbb{R}^2$, we finally obtain:

Corollary 1.14. *Conjecture 1.2 holds for orientable manifolds in dimensions $6 \neq n \leq 7$.*

Again we like to point out that the low-dimensional cases $n \leq 5$ of Corollary 1.14 were already known due to index-theoretic results [17; 36, Theorem 1.10].

This article is organized as follows: In Section 2, we prepare an abstract setup for the study of manifolds with at least two ends which underpins our work. In Section 3, we state quantitative comparison results in the spirit of Conjecture 1.3 which are then used together with topological arguments in Section 4 to prove our codimension one results. The codimension two results are deduced in Section 5. Finally, in Section 6 we provide the analytic proofs of the comparison statements from Section 3.

Acknowledgement

The authors thank Georg Frenck for helpful discussions and suggestions.

2 Bands

A natural class of manifolds generalizing the situation of Conjecture 1.1 are connected non-compact manifolds, where we partition the set of ends into two parts and we consider hypersurfaces separating these parts from each other. To make this precise, we will make use of the Freudenthal end compactification [9] of a connected manifold M , denoted by $\mathcal{F}M = M \cup \mathcal{E}M$, where $\mathcal{E}M$ is the space of ends.

Definition 2.1.

- (i) An *open band* is a connected non-compact manifold M without boundary and a decomposition

$$\mathcal{E}M = \mathcal{E}_-M \sqcup \mathcal{E}_+M,$$

where $\mathcal{E}_\pm M$ are non-empty closed³ subsets $\mathcal{E}_\pm M \subset \mathcal{E}M$; in particular M has at least two ends.

- (ii) Given an open band M , a *separating hypersurface* $\Sigma \subset M$ is a compact hypersurface which separates each end in \mathcal{E}_-M from every end in \mathcal{E}_+M . Moreover, we say a separating hypersurface $\Sigma \subset M$ is *properly separating* if every component of Σ can be connected to both \mathcal{E}_+M and \mathcal{E}_-M inside $M \setminus \Sigma$.
- (iii) Let $\Sigma_-, \Sigma_+ \subset M$ be two properly separating hypersurfaces in an open band M . Then we write $\Sigma_- \prec \Sigma_+$ if the hypersurface Σ_- is contained in the union of those components of $M \setminus \Sigma_+$ that contain the ends in \mathcal{E}_-M (or equivalently, Σ_+ is contained in the union of those components of $M \setminus \Sigma_-$ containing \mathcal{E}_+M).

The condition of being *properly* separating simply means that there are no superfluous components as the observation recorded in the following lemma illustrates.

Lemma 2.2. *Let $\Sigma \subset M$ be a separating hypersurface in an open band M . Then there exists a union of components of Σ which is a properly separating hypersurface in M .*

³While the space of ends is totally disconnected, it is in general not necessarily discrete. In this case it is important to assume $\mathcal{E}_\pm M$ to be closed (and thus clopen) subsets of $\mathcal{E}M$.

Proof. Suppose that Σ is a separating hypersurface that contains a component not connected to both \mathcal{E}_-M and \mathcal{E}_+M inside $M \setminus \Sigma$. Then the hypersurface Σ' obtained from Σ by deleting this component is still a separating hypersurface. This shows that a minimal collection of components of Σ such that its union is still separating yields the desired properly separating hypersurface. \square

The next elementary lemma shows that (properly) separating hypersurfaces always exist and we can find them arbitrarily far out.

Lemma 2.3. *Let M be an open band and $K \subset M$ be an arbitrary compact subset. Then there exists a properly separating hypersurface $\Sigma \subset M$ which also separates K from \mathcal{E}_+M (or \mathcal{E}_-M , respectively).*

Proof. Note that the end compactification $\mathcal{F}M$ is a compact Hausdorff space which in our case of a connected manifold is also second countable. Thus, since $\mathcal{E}_\pm M$ are two disjoint closed subsets of $\mathcal{F}M$, Urysohn's lemma implies the existence of a continuous function $f: \mathcal{F}M \rightarrow [-1, 1]$ such that $\mathcal{E}_\pm M = f^{-1}(\pm 1)$. Since $K \subseteq M$ is compact, there exists $0 < r < 1$ such that $K \subseteq f^{-1}([-r, r])$. Choose $s \in (r, 1)$. Then $f^{-1}(s) \subseteq M$ is a compact subset which separates \mathcal{E}_-M from \mathcal{E}_+M . Now choose a connected compact n -dimensional submanifold $V \subset M$ with boundary, where $n = \dim(M)$, such that $f^{-1}(s) \subseteq \mathring{V} \subseteq f^{-1}([s - \varepsilon, s + \varepsilon])$ for some $\varepsilon > 0$ with $r < s - \varepsilon$. Then ∂V is a separating hypersurface and it contains a properly separating hypersurface $\Sigma \subseteq \partial V$ by Lemma 2.2. Since by construction $f(x) \leq r < s - \varepsilon \leq f(y) \leq s + \varepsilon < 1$ for each $x \in K$ and $y \in \Sigma$, it follows that Σ must separate K from \mathcal{E}_+M . A completely analogous argument also provides a properly separating hypersurface that separates K from \mathcal{E}_-M . \square

In the spirit of Conjecture 1.1 we will be interested in open bands with:

Property A. No separating hypersurface admits a metric of positive scalar curvature.

We will also work with compact bands, which may be viewed as a special case of open bands in the following sense:

Definition 2.4. A *compact band*, to which we will often simply refer to as a “band”, is a connected compact manifold X together with the structure of an open band on its interior \mathring{X} . In other words, this amounts to a decomposition $\partial X = \partial_- X \sqcup \partial_+ X$, where $\partial_\pm X$ are (non-empty) unions of boundary components.

The notions of (properly) separating hypersurfaces and Property A make sense for compact bands by considering them for the interior.

We also note that, if M is an open band and Σ_\pm are two properly separating hypersurfaces such that $\Sigma_- \prec \Sigma_+$, then the hypersurfaces Σ_\pm bound a compact band $X \subset M$. In this case, if M has Property A, then so has X .

3 The partitioned comparison principle

In this section, we establish the main analytic results on which our main theorems rely. The central concept we study here is the $\text{width}(X, g)$ of a compact band X endowed with a Riemannian metric g , that is, the distance between $\partial_- X$ and $\partial_+ X$ with respect to g . As is explained in [15, Section 3.6] the width of an n -dimensional Riemannian band (X, g) with $\text{scal} \geq n(n - 1)$ is bounded from above by $\frac{2\pi}{n}$ if X has Property A and $n \leq 7$. We will work in the setting where (X, g) has Property A and is partitioned into multiple segments with possibly different lower scalar curvature bounds. It turns out that in this case positivity of the scalar curvature in a single segment can often have global effects on the geometry of (X, g) .

Definition 3.1. Let X be a compact band and let Σ_i , for $i \in \{1, \dots, k\}$, be properly separating hypersurfaces such that $\Sigma_{i-1} \prec \Sigma_i$ for $1 < i \leq k$. We call the triple (X, Σ_i, k) a *partitioned band* and denote by V_j , for $j \in \{1, \dots, k + 1\}$, the segment of X bounded by Σ_{j-1} and Σ_j , where $\Sigma_0 = \partial_- X$ and $\Sigma_{k+1} = \partial_+ X$.

Definition 3.2. A smooth function $\varphi: [a, b] \rightarrow \mathbb{R}_+$ is called *log-concave* if

$$\frac{d^2}{dt^2} \log(\varphi(t)) = \left(\frac{\varphi'(t)}{\varphi(t)} \right)' \leq 0$$

for all $t \in [a, b]$. If the inequality is strict, then φ is called *strictly log-concave*.

Definition 3.3. Let (N, g_N) be a closed scalar flat Riemannian manifold. A warped product

$$(M, g_\varphi) = (N \times [a, b], \varphi^2(t)g_N + dt^2)$$

is called a *model space* if $\text{scal}(M, g_\varphi)$ is constant and φ is strictly log-concave.

Remark 3.4. This notion of a model space arises in the scalar and mean curvature comparison theory surrounding Conjecture 1.3. In [3; 25] a compact Riemannian band (X, g) is compared, in scalar curvature, mean curvature and width, to a warped product (M, g_φ) over an arbitrary scalar flat manifold (N, g_N) with strictly log-concave warping function. It turns out that, if X has Property A, $\text{scal}(X, g) \geq \text{scal}(M, g_\varphi)$ and $H(\partial X, g) \geq H(\partial M, g_\varphi)$, then $\text{width}(X, g) \leq \text{width}(M, g_\varphi)$. Furthermore, Cecchini and Zeidler [3, Theorem 8.3, Theorem 9.1] showed that, under further topological assumptions on X , equality of widths can only be achieved if (X, g) itself is isometric to such a warped product. In this section we adapt these ideas and compare a compact partitioned Riemannian band with Property A to a sequence of model spaces, whose scalar curvatures may differ, but whose mean curvatures fit together pairwise.

For a compact Riemannian manifold with boundary (N, h) and a connected component of the boundary $\partial_j N$, we denote by $H(\partial_j N, h)$ the mean curvature of $\partial_j N$ with respect to the interior unit normal field. According to our convention, the boundary of the unit disk in \mathbb{R}^3 has mean curvature 2. In the following theorem, we give estimates on the widths of segments inside a partitioned band based on scalar and mean curvature.

Theorem 3.5. *Let $n \leq 7$ and let (X, Σ_i, k) be an orientable partitioned n -dimensional band with Property A. For $j \in \{1, \dots, k+1\}$, let (M_j, g_{φ_j}) be model spaces over a fixed closed scalar flat Riemannian manifold (N, g) . If*

- ▷ $\text{scal}(V_j, g) \geq \text{scal}(M_j, g_{\varphi_j})$ for all $j \in \{1, \dots, k+1\}$,
- ▷ $H(\partial_- X, g) \geq H(\partial_- M_1, g_{\varphi_1})$ and $H(\partial_+ X, g) \geq H(\partial_+ M_{k+1}, g_{\varphi_{k+1}})$,
- ▷ $H(\partial_+ M_j, g_{\varphi_j}) = -H(\partial_- M_{j+1}, g_{\varphi_{j+1}})$ for all $j \in \{1, \dots, k\}$,

then $\text{width}(V_j, g) \leq \text{width}(M_j, g_{\varphi_j})$ for at least one $j \in \{1, \dots, k+1\}$.

The following result is more or less a direct application of Theorem 3.5:

Corollary 3.6. *Let $n \leq 7$ and let $(X, \Sigma_i, 2)$ be an orientable partitioned n -dimensional band with Property A. Let g be a metric on X and let κ be a positive constant. If*

- ▷ $\text{scal}(V_2, g) \geq \kappa n(n-1)$,
- ▷ $\text{scal}(X, g) \geq 0$,

then $\min\{\text{width}(V_1, g), \text{width}(V_3, g)\} < \ell = \frac{2}{\sqrt{\kappa n}} \cot\left(\frac{\sqrt{\kappa n d}}{4}\right)$, where $d := \text{width}(V_2, g) < \frac{2\pi}{\sqrt{\kappa n}}$.

If, instead of $\text{scal}(X, g) \geq 0$, one assumes that the scalar curvature of the partitioned band is bounded from below by a negative constant, Theorem 3.5 provides the following estimate, which is very much in the same spirit of Corollary 3.6 and should be compared with Zeidler's result [37, Theorem 1.4] in the spin setting.

Corollary 3.7. *Let $n \leq 7$ and let $(X, \Sigma_i, 2)$ be an orientable partitioned n -dimensional band with Property A. Let g be a metric on X and let κ be a positive constant. If*

$$\triangleright \text{scal}(V_2, g) \geq \kappa n(n-1),$$

$$\triangleright \text{scal}(X, g) \geq -\sigma > -\kappa n(n-1) \tan\left(\frac{\sqrt{\kappa}nd}{4}\right)^2, \text{ where } d := \text{width}(V_2, g) < \frac{2\pi}{\sqrt{\kappa n}},$$

then $\min\{\text{width}(V_1, g), \text{width}(V_3, g)\} < \ell$, where ℓ is such that

$$\sqrt{\kappa}(n-1) \tan\left(\frac{\sqrt{\kappa}nd}{4}\right) = \sqrt{\frac{\sigma(n-1)}{n}} \coth\left(\frac{\sqrt{\sigma n} \ell}{2\sqrt{n-1}}\right).$$

We postpone the proofs of Theorem 3.5 and Corollaries 3.6 and 3.7 to Section 6.

4 Obstructions on open bands

We want to use Corollary 3.6 or Corollary 3.7 to attack Conjecture 1.1. If $X = Y \times \mathbb{R}$ and g is a complete metric on X , we consider the compact segment $Y \times [-C, C]$ for any $C > 0$, which is partitioned into the bands $Y \times [-C, -1]$, $Y \times [-1, 1]$ and $Y \times [1, C]$. If the scalar curvature of (X, g) is assumed to be positive and $Y \times [-C, C]$ has Property A, the minimum of the widths of $(Y \times [-C, -1], g)$ and $(Y \times [1, C], g)$ is bounded from above in terms of $\text{width}(Y \times [-1, 1], g)$ and the infimum of $\text{scal}(Y \times [-1, 1], g)$. For $C > 0$ large enough, this produces a contradiction.

In this section, we formulate the most general result that one can prove in this manner based on our notion of an open band from Definition 2.1. Indeed, Corollary 3.6 or Corollary 3.7 will yield obstructions to the existence of a complete metric of positive scalar curvature on an open band with Property A. Moreover, since any open band M has at least two ends by definition, we can also establish a rigidity result using the Cheeger–Gromoll splitting theorem.

Theorem 4.1. *Let $n \leq 7$ and let M be an open n -dimensional band with Property A. If g is a complete metric on M with nonnegative scalar curvature, then (M, g) is isometric to $(Y \times \mathbb{R}, g_Y + dt^2)$, where (Y, g_Y) is a closed Ricci flat manifold.*

Proof. If (M, g) is not Ricci flat, then M admits a complete metric \hat{g} of positive scalar curvature by [18, Theorem B]. Let $\Sigma \subset M$ be a properly separating hypersurface which exists by Lemma 2.3. There is a $\kappa > 0$ such that $\text{scal}(M, \hat{g}) \geq \kappa n(n-1)$ in a neighborhood of width $d < \frac{2\pi}{\sqrt{\kappa n}}$, which is bounded by two properly separating hypersurfaces Σ_1 and Σ_2 such that $\Sigma_1 \prec \Sigma \prec \Sigma_2$. For every $C > 0$, we can use Lemma 2.3 to find further properly separating hypersurfaces Σ_{\pm}^C such that $\Sigma_-^C \prec \Sigma_1 \prec \Sigma_2 \prec \Sigma_+^C$ and $\text{dist}_{\hat{g}}(\Sigma_-^C, \Sigma_1) \geq C \leq \text{dist}_{\hat{g}}(\Sigma_2, \Sigma_+^C)$. Let X^C be the compact band bounded by Σ_{\pm}^C . Then for C large enough, Corollary 3.6 or Corollary 3.7 applied to X^C yields a contradiction. Hence (X, g) is Ricci flat.

Since M is an open band it is disconnected at infinity and so admits a geodesic line (see e.g. [24, Lemma 7.3.1]). Thus by the Cheeger–Gromoll splitting theorem (see e.g. [24, Theorem 7.3.5]), (M, g) is isometric to $(Y \times \mathbb{R}, g_Y + dt^2)$, where (Y, g_Y) is a Ricci flat manifold. Furthermore, M has more than one end by the definition of an open band, and so Y must be compact. \square

Using a similar (and somewhat simpler argument), we also obtain the following statement:

Proposition 4.2. *Let $n \leq 7$ and let M be an open n -dimensional band with Property A. Then M does not admit a complete metric which has uniformly positive scalar curvature outside a compact subset.*

Proof. Assume, by contradiction, that M admits a complete metric g which has scalar curvature $\text{scal}_g \geq \kappa n(n-1) > 0$ on $M \setminus K$ for some compact subset $K \subset M$ and some $\kappa > 0$. Then, applying Lemma 2.3 twice, we can find a compact band $X \subset M \setminus K$ which is bounded by

two properly separating hypersurfaces $\partial_- X \prec \partial_+ X \subset M$ such that $\text{width}(X, g) > \frac{2\pi}{\sqrt{\kappa n}}$. By construction, we also have $\text{scal}_g \geq \kappa n(n-1) > 0$ on X . But this contradicts Corollary 3.7 or even the usual band width estimate on compact bands [15, Section 3.6; 25] of dimension ≤ 7 . \square

In the rest of this section, we investigate topological conditions which imply the existence of open bands with Property A and use them together with Theorem 4.1 to prove the main Theorems 1.5 and 1.9. Let us start with the notion of an incompressible hypersurface.

Definition 4.3. Let X be a connected manifold and let $\Sigma \subset X$ be a connected hypersurface. We say that Σ is incompressible if the induced map $\iota_* : \pi_1(\Sigma) \rightarrow \pi_1(X)$ is injective.

Our first goal is to provide the most general conditions under which the existence of an incompressible hypersurface which does not admit psc shows that the manifold is covered by an open band with Property A. The arguments will be based on positive scalar curvature bordism and surgery techniques which usually require a case distinction depending on the presence of a spin structure. To treat this in a unified way, we use the language of tangential structures, for details see e.g. [10, §5]. We briefly recall that an n -dimensional tangential structure is given by a fibration $\theta: B \rightarrow \text{BO}(n)$ and a θ -structure on an n -manifold M is a lift $M \rightarrow B$ of the classifying map of the tangent bundle $M \rightarrow \text{BO}(n)$ along $\theta: B \rightarrow \text{BO}(n)$. Then, given an n -dimensional tangential structure $\theta: B \rightarrow \text{BO}(n)$, one may define a notion of θ -cobordism for $(n-1)$ -dimensional θ -manifolds⁴ and form the corresponding cobordism group which we denote by Ω_{n-1}^θ . Now the positive scalar curvature surgery principle can be phrased as follows: Let $\theta: B \rightarrow \text{BO}(n)$ be an n -dimensional tangential structure and fix an $(n-1)$ -dimensional θ -manifold N such that the map $N \rightarrow B$ is 2-connected. If N is θ -cobordant to a θ -manifold which admits a metric of psc, then N itself already admits psc, see [8, Theorem 1.5].

We now connect this back to the study of open bands and first observe that an open band endowed with a θ -structure induces a well-defined θ -cobordism class of $(n-1)$ -manifolds.

Lemma 4.4. *Let $\theta: B \rightarrow \text{BO}(n)$ be a tangential structure and let M be an open band endowed with a θ -structure. Then any two properly separating hypersurfaces $\Sigma_1, \Sigma_2 \subset M$ are θ -cobordant.*

Proof. First consider the case that $\Sigma_1 \prec \Sigma_2$. Then the band in M bounded in between Σ_1 and Σ_2 is a cobordism witnessing that Σ_1 and Σ_2 are θ -cobordant. In general, Lemma 2.3 implies the existence of a properly separating hypersurface $\Sigma' \subset M$ such that $\Sigma_1 \prec \Sigma'$ and $\Sigma_2 \prec \Sigma'$. Thus, by the first case, both Σ_1 and Σ_2 are θ -cobordant to Σ' which proves the desired statement. \square

Next we use the psc surgery principle as stated above to derive a sufficient condition for the existence of open bands with Property A.

Lemma 4.5. *Let $n \geq 6$ and let $\theta: B \rightarrow \text{BO}(n)$ be a tangential structure. Let M be a connected θ -manifold without boundary and let $Y \subset M$ be a closed connected two-sided hypersurface. Assume that Y does not support a psc metric and that the composition $Y \subset M \rightarrow B$ induces an injection $\pi_1 Y \hookrightarrow \pi_1 B$ and a surjection $\pi_2 Y \twoheadrightarrow \pi_2 B$. Then the connected covering \hat{M} of M with $\pi_1 \hat{M} = \pi_1 Y$ is an open band with Property A.*

Proof. First let $\hat{B} \rightarrow B$ be the covering with $\pi_1 \hat{B} = \pi_1 Y$. Then $\hat{M} \rightarrow M$ is the pullback of the covering $\hat{B} \rightarrow B$ along $\theta: M \rightarrow B$. The embedding $Y \subset M$ lifts to an embedding $Y \subset \hat{M}$ as a hypersurface which separates \hat{M} into two components (otherwise there existed a loop in \hat{M} intersecting Y transversally in precisely one point which would contradict $\pi_1 Y \cong \pi_1 \hat{M}$).

⁴To be precise, here one implicitly considers the $(n-1)$ -dimensional tangential structure obtained from θ via the pullback along $\text{BO}(n-1) \rightarrow \text{BO}(n)$.

Moreover, we obtain a $\hat{\theta}$ -structure on \hat{M} , where $\hat{\theta}: \hat{B} \rightarrow B \xrightarrow{\theta} \text{BO}(n)$.

$$\begin{array}{ccccc}
 & & \hat{M} & \longrightarrow & \hat{B} \\
 & \nearrow & \downarrow & & \searrow \hat{\theta} \\
 Y & \hookrightarrow & M & \longrightarrow & B \xrightarrow{\theta} \text{BO}(n)
 \end{array}$$

By assumption, the composition $Y \hookrightarrow \hat{M} \rightarrow \hat{B}$ is 2-connected. Let U_{\pm} be the two components of $\hat{M} \setminus Y$. We observe that U_{\pm} are both non-compact because otherwise Y would be $\hat{\theta}$ -nullbordant and thus support a psc metric by [8, Theorem 1.5]. We can now define the ends of U_{-} to be $\mathcal{E}_{-}\hat{M}$ and the ends of U_{+} to be $\mathcal{E}_{+}\hat{M}$ in order to turn \hat{M} into an open band. Then, by construction, the open band \hat{M} contains Y as a properly separating hypersurface.

Finally, we need to verify Property A. To this end, assume by contradiction that $\Sigma \subset \hat{M}$ is a separating hypersurface that admits a psc metric. By Lemma 2.2, we may assume without loss of generality that Σ is properly separating. Then it follows from Lemma 4.4 that Σ and Y are $\hat{\theta}$ -cobordant. But since the map $Y \rightarrow \hat{B}$ is 2-connected and Σ admits a psc metric, [8, Theorem 1.5] implies that Y also admits a psc metric, a contradiction. \square

To describe the possible concrete applications of Lemma 4.5, it turns out to be enough to go through the possible tangential 2-types of M . Recall that for an arbitrary n -manifold, its tangential 2-type is the (unique up to homotopy) tangential structure θ_M which factors the tangent bundle as $M \rightarrow B_M \xrightarrow{\theta_M} \text{BO}(n)$, where $M \rightarrow B_M$ is 2-connected and $\theta_M: B_M \rightarrow \text{BO}(n)$ is 2-co-connected. In other words, B_M is the second stage of the Moore–Postnikov tower for the map $M \rightarrow \text{BO}(n)$. Now a simple diagram chase shows that if the hypotheses of Lemma 4.5 are satisfied for some tangential structure θ on M , then they are already satisfied for $\theta = \theta_M$. Moreover, the tangential 2-type can be described algebraically in terms of the fundamental group $\pi = \pi_1 M$, the first Stiefel–Whitney class $w: \pi \rightarrow \mathbb{Z}/2$, and an extension $\hat{\pi} \rightarrow \pi$ determined by the second Stiefel–Whitney class whose kernel has order at most 2, see [33, §2] for details. However, for our purposes we do not need such a full description, and we only need to distinguish two cases depending on whether $\pi_2 B_M \cong \mathbb{Z}/2$ or $\pi_2 B_M = 0$ as the following proposition demonstrates.

Proposition 4.6. *Let $n \geq 6$. Let M be a connected n -dimensional manifold without boundary and $Y \subset M$ a closed two-sided incompressible hypersurface that does not admit a psc metric. Suppose that one of the following conditions holds:*

- (a) M is almost spin.
- (b) Y is totally non-spin.

Then there exists a covering $\hat{M} \rightarrow M$ which is an open band with Property A.

Proof. In light of the discussion in the previous paragraph, we need to check that in either case we can apply Lemma 4.5 for the tangential structure $\theta = \theta_M$ given by its 2-type.

- (a) If M is almost spin, that is, the universal covering \tilde{M} of M is spin, then the map $M \rightarrow \text{BO}(n)$ classifying the tangent bundle induces the zero map $\pi_2 M \rightarrow \pi_2 \text{BO}(n) = \mathbb{Z}/2$ because this map can be identified with the second Stiefel–Whitney class of \tilde{M} via the Hurewicz isomorphism $H_2(\tilde{M}) \cong \pi_2(\tilde{M}) \cong \pi_2(M)$. Thus it follows that $\pi_2(B_M) = 0$ and so Lemma 4.5 applies to every incompressible hypersurface $Y \subset M$ which does not admit psc.
- (b) On the other hand, if M is totally non-spin, that is \tilde{M} is non-spin, then by an analogous consideration involving the second Stiefel–Whitney class of \tilde{M} we necessarily have $\pi_2 B_M \cong \mathbb{Z}/2$ and $\pi_2 M \rightarrow \pi_2 B_M = \mathbb{Z}/2$ is surjective. Now in this situation the condition on the hypersurface $Y \subset M$ in Lemma 4.5 is that it is incompressible, itself totally non-spin, and does not admit psc. \square

We are now ready to deduce Theorem 1.5:

Proof of Theorem 1.5. By Proposition 4.6, there exists a covering \hat{M} which is an open band with Property A. Now suppose that M admits a complete metric of non-negative scalar curvature g . Let \hat{g} denote its lift to \hat{M} . Then Theorem 4.1 implies that (\hat{M}, \hat{g}) must be isometric to $(N \times \mathbb{R}, g_N + dt^2)$ for a closed Ricci flat manifold (N, g_N) . \square

We now turn to Theorem 1.9. Note that a proof of Theorem 1.9 has already appeared recently in [5, Theorem 1.1], but we give a separate argument here because it also fits directly into our topological setup. In fact, it is simpler than Theorem 1.5 as it does not need surgery arguments and instead only relies on Lemma 4.4 together with some homological considerations:

Lemma 4.7. *Let M be an oriented open band and $Y \subseteq M$ a properly separating hypersurface together with a map $\phi: M \rightarrow Y_0$ to an NPSC⁺ manifold Y_0 such that the restriction $\phi|_Y: Y \rightarrow Y_0$ has non-zero degree. Then M has Property A.*

Proof. We define a tangential structure $M \xrightarrow{l} B \xrightarrow{\theta} \text{BO}(n)$, where we set $B = Y_0 \times \text{BSO}(n)$ and l is induced by the map $M \rightarrow Y_0$ together with the orientation of M . Note that a θ -structure on an $(n-1)$ -manifold N is the same as an orientation on N together with a map $N \rightarrow Y_0$, and thus $\Omega_{n-1}^\theta = \Omega_{n-1}^{\text{SO}}(Y_0)$. In this picture, the degree of the map $N \rightarrow Y_0$ can be read off from the transformation $\Omega_{n-1}^{\text{SO}}(Y_0) \rightarrow \text{H}_{n-1}(Y_0; \mathbb{Z}) \cong \mathbb{Z}$. Suppose that $\Sigma \subset M$ is a separating hypersurface, and assume without loss of generality that it is properly separating. Then, by Lemma 4.4, $\Sigma \rightarrow Y_0$ and $Y \rightarrow Y_0$ represent the same class of $\Omega_{n-1}^{\text{SO}}(Y_0)$. In particular, $\deg(\Sigma \rightarrow Y_0) = \deg(Y \rightarrow Y_0) \neq 0$. Since Y_0 is NPSC⁺, this proves that Σ does not admit a psc metric and so M must have Property A. \square

Proposition 4.8. *Let M be an oriented connected n -dimensional manifold and let $\iota: Y \hookrightarrow M$ be a two-sided closed connected hypersurface that admits a map of non-zero degree $\phi: Y \rightarrow Y_0$ to an aspherical NPSC⁺ manifold Y_0 and such that $\ker(\pi_1 Y \xrightarrow{\iota_*} \pi_1 M) \subseteq \ker(\pi_1 Y \xrightarrow{\phi_*} \pi_1 Y_0)$. Then there exists a connected covering $\hat{M} \rightarrow M$ that is an open band with Property A.*

Proof. Let $\Lambda := \iota_*(\pi_1 Y) \subseteq \pi_1 M$ and let $\hat{M} \rightarrow M$ be the covering with $\pi_1 \hat{M} = \Lambda$. Then $Y \hookrightarrow M$ lifts to an embedding $Y \hookrightarrow \hat{M}$ which induces a surjection $\pi_1 Y \twoheadrightarrow \Lambda$. By the assumption on the kernels of the induced maps on fundamental groups, it follows that the homomorphism $\phi_*: \pi_1 Y \rightarrow \pi_1 Y_0$ factors as a composition $\pi_1 Y \twoheadrightarrow \Lambda \rightarrow \pi_1 Y_0$. Since Y_0 is aspherical and $\pi_1 \hat{M} = \Lambda$, this implies that the map $\phi: Y \rightarrow Y_0$ extends to a map $\hat{M} \rightarrow Y_0$. As in the proof of Lemma 4.5, let U_\pm be the two components of $\hat{M} \setminus Y$. Then U_\pm must be non-compact because otherwise $[Y \rightarrow Y_0] = 0 \in \Omega_{n-1}^{\text{SO}}(Y_0)$ which would contradict the hypothesis $\deg(Y \rightarrow Y_0) \neq 0$ (compare the proof of Lemma 4.7 above). Thus \hat{M} can be turned into an open band such that Y is a properly separating hypersurface. Thus the proposition follows from Lemma 4.7. \square

Proof of Theorem 1.9. Combine Proposition 4.8 and Theorem 4.1 analogously as in the proof of Theorem 1.5 above. \square

5 The codimension two obstruction

In this section, we prove our codimension two obstruction results. These are based on a reduction to a codimension one problem essentially following original ideas of Gromov and Lawson [13, Theorem 7.5] and their adaptation by Hanke, Pape, and Schick [17].

Lemma 5.1 (cf. [17, Theorem 4.3; 34, Lemma 4.1.4]). *Let X be a connected manifold without boundary. Let $Y \subset X$ be a submanifold without boundary of codimension two whose normal bundle is trivial and suppose that the pair (X, Y) is 2-connected. Consider the manifold $W := X \setminus \mathcal{U}$*

obtained by deleting a small open tubular neighborhood $Y \times B^2 \cong \mathcal{U} \subseteq X$. Then the map $\partial W \cong Y \times S^1 \xrightarrow{\text{pr}_2} S^1$ given by projection onto the second factor extends to a continuous map $W \rightarrow S^1$. In particular, the map $\pi_1 Y \times \mathbb{Z} = \pi_1(\partial W) \rightarrow \pi_1 W$ induced by the inclusion $\partial W \hookrightarrow W$ is split-injective.

Proof. It suffices to show that the induced homomorphism $(\text{pr}_2)_*: \pi_1(\partial W) \rightarrow \pi_1(S^1) = \mathbb{Z}$ extends to a homomorphism $\pi_1(W) \rightarrow \mathbb{Z}$. The hypotheses imply that the pair (X, \mathcal{U}) is also 2-connected and so excision and the Hurewicz theorem show that $H_k(W, \partial W) \cong H_k(X, \mathcal{U}) = 0$ for $0 \leq k \leq 2$. In particular, the map $H_1(\partial W) \xrightarrow{\cong} H_1(W)$ induced by the inclusion $\partial W \hookrightarrow W$ is an isomorphism. The existence of the desired extension now follows from the following diagram because the map $(\text{pr}_2)_*: \pi_1(\partial W) \rightarrow \pi_1(S^1) = \mathbb{Z}$ factors through the Hurewicz homomorphism.

$$\begin{array}{ccccc} \pi_1(\partial W) & \xrightarrow{\text{hur}} & H_1(\partial W) & \xrightarrow{(\text{pr}_2)_*} & H_1(S^1) & \xrightarrow{\text{hur}} & \pi_1(S^1) = \mathbb{Z} \\ \downarrow & & \downarrow \cong & \nearrow \text{---} & & & \\ \pi_1(W) & \xrightarrow{\text{hur}} & H_1(W) & & & & \end{array}$$

This also implies that the map $\pi_1 Y \times \mathbb{Z} = \pi_1(\partial W) \rightarrow \pi_1 W$ is split injective because a retraction can be constructed using the map $\pi_1 W \rightarrow \mathbb{Z}$ from the previous paragraph together with the map $\pi_1 W \rightarrow \pi_1 X \cong \pi_1 Y$ induced by the inclusion $W \hookrightarrow X$. \square

Proposition 5.2. *Let $n \geq 6$. Let X be a connected n -dimensional manifold and let $Y \subset X$ be a closed connected submanifold of codimension two with trivial normal bundle such that the pair (X, Y) is 2-connected. Consider the manifold $W := X \setminus \mathcal{U}$ obtained by deleting a small open tubular neighborhood $Y \times B^2 \cong \mathcal{U} \subseteq X$. If $Y \times S^1$ does not admit a metric of positive scalar curvature, then the double dW of W is an open band with Property A.*

Proof. We start with considering the tangential 2-type $X \rightarrow B_X \rightarrow \text{BO}(n)$ of X . By restriction to $W \subset X$ this induces a tangential structure $l': W \rightarrow B_X$. Let $p: W \rightarrow S^1$ be a map extending the projection $\partial W \cong Y \times S^1 \rightarrow S^1$ which exists by Lemma 5.1. We obtain a new tangential structure $W \xrightarrow{l} B := (B_X \times S^1) \xrightarrow{\theta} \text{BO}(n)$, where $l = (l', p)$, and θ is defined as projection to B_X followed by $B_X \rightarrow \text{BO}(n)$. Furthermore, this tangential structure l extends to the double dW by reflection. Its restriction to the hypersurface $dW \supset \partial W \cong Y \times S^1$ is a 2-connected map $Y \times S^1 \rightarrow B = B_X \times S^1$ because by construction it is homotopic to $l_Y \times \text{id}_{S^1}$, where l_Y denotes the restriction of $X \rightarrow B_X$ to $Y \subset X$ and l_Y is 2-connected. Thus Lemma 4.5 (applied to the trivial covering) proves the desired conclusion. \square

Proof of Theorem 1.11. Note that since Y does not admit positive scalar curvature and $\dim(Y) = 5$, Theorem 1.5 implies that $Y \times S^1$ does not admit positive scalar curvature either. Then we let X be the connected covering of M with $\pi_1 X = \pi_1 Y$. It follows $Y \hookrightarrow X$ and the pair (X, Y) satisfies the hypotheses of Proposition 5.2. Thus, if we let W be as in the statement of Proposition 5.2, then its double dW is an open band with Property A. Now assume by contradiction that M admits a complete metric of uniformly positive scalar curvature. Then by first lifting it to X and restricting it to W , we obtain a complete metric of uniformly positive scalar curvature on W . After changing this metric in a compact neighborhood of ∂W , we obtain a smooth complete metric on dW which has uniformly positive scalar curvature outside a compact subset, a contradiction to Proposition 4.2. \square

Proof of Theorem 1.12. We first verify that $Y_0 \times S^1$ is also NPSC⁺. To this end, let N be an oriented $(n-1)$ -manifold and $\phi: N \rightarrow Y_0 \times S^1$ a map of non-zero degree, where we assume N to be connected without loss of generality. Then let $Z = \phi^{-1}(Y_0 \times \{*\})$ be a transversal pre-image. It follows that $\iota: Z \hookrightarrow N$ and $\phi: Z \rightarrow Y_0 \times \{*\} = Y_0$ satisfies the hypotheses of Theorem 1.9 and so N does not admit a psc metric. This proves that $Y_0 \times S^1$ is NPSC⁺.

To prove the theorem, we again consider the connected covering of $X \rightarrow M$ with $\pi_1 X = \pi_1 Y$, and we let W be as in the statement of Proposition 5.2. By Lemma 5.1, the map $\pi_1 Y \times \mathbb{Z} = \pi_1(\partial W) \rightarrow \pi_1 W$ induced by the inclusion $Y \times S^1 = \partial W \hookrightarrow W$ is injective and admits a retraction $r: \pi_1 W \rightarrow \pi_1 Y \times \mathbb{Z}$. Since $Y_0 \times S^1$ is aspherical, this implies that the map $Y \times S^1 \rightarrow Y_0 \times S^1$ extends to a map $W \rightarrow Y_0 \times S^1$ and subsequently to a map $dW \rightarrow Y_0 \times S^1$ on the double. In summary, dW is an open band that contains $Y \times S^1$ as a properly separating hypersurface and it satisfies the hypotheses of Lemma 4.7 because $Y_0 \times S^1$ is NPSC⁺. Thus dW has Property A and the theorem now follows again from Proposition 4.2 as in the proof of Theorem 1.11 above. \square

6 Proof of the Partitioned Comparison Principle

We will prove Theorem 3.5 by contradiction. We stress that all bands considered in this section are compact. Under the assumption that $\text{width}(V_j, g) > \text{width}(M_j, g_{\varphi_j})$ for all $j \in \{1, \dots, k+1\}$, we will produce a closed embedded hypersurface $\Sigma \subset X$ which separates $\partial_- X$ and $\partial_+ X$ and admits a metric of positive scalar curvature.

This hypersurface Σ will appear as the boundary of a μ -bubble. The key ingredient for the corresponding functional is the potential function $h: X \rightarrow \mathbb{R}$. We use the ideas from [25, Section 3] for each band (V_j, g) and model space (M_j, g_{φ_j}) separately to produce $h_j: V_j \rightarrow \mathbb{R}$ as the concatenation of a strictly 1-Lipschitz band map $(V_j, g) \rightarrow (M_j, g_{\varphi_j})$ and the function $h_{\varphi_j}: M_j \rightarrow \mathbb{R}$. Subsequently, we use a gluing construction to paste all of the h_j together to obtain a smooth function $h: X \rightarrow \mathbb{R}$ which is suitable for our purposes.

The idea to combine potential functions in this way, was already used in [4; 37]. The gluing construction is based on the following result:

Lemma 6.1. *Let $h: [a, b] \rightarrow \mathbb{R}$ be a smooth strictly monotonously decreasing function such that*

$$-\frac{n}{n-1}h^2 - 2h' = \sigma,$$

for some constant $\sigma \in \mathbb{R}$. Then, for every sufficiently small $\varepsilon > 0$, there exists a function $\hat{h}: [a - \varepsilon, b + \varepsilon] \rightarrow \mathbb{R}$ such that:

- ▷ $\hat{h}(t) = h(t)$ for $t \in [a + \varepsilon, b - \varepsilon]$,
- ▷ $\hat{h}(t) = h(a)$ in a neighborhood of $a - \varepsilon$ and $\hat{h}(t) = h(b)$ in a neighborhood of $b + \varepsilon$,
- ▷ $\hat{h}' \leq 0$,
- ▷ $-\frac{n}{n-1}\hat{h}^2 - 2\hat{h}' \leq \sigma$ and $-\frac{n}{n-1}\hat{h}^2(t) - 2\hat{h}'(t) < \sigma$ if $\hat{h}'(t) = 0$.

Proof. Let $\rho: \mathbb{R} \rightarrow [a, b]$ be a smooth function with:

- ▷ $\rho(t) = a$ for $t \in (-\infty, a - \frac{\varepsilon}{2}]$, $\rho(t) = t$ for $t \in [a + \frac{\varepsilon}{2}, b - \frac{\varepsilon}{2}]$ and $\rho(t) = b$ for $t \in [b + \frac{\varepsilon}{2}, \infty)$.
- ▷ $0 < \rho'(t) < 1$ for $t \in (a - \frac{\varepsilon}{2}, a + \frac{\varepsilon}{2})$ and $t \in (b - \frac{\varepsilon}{2}, b + \frac{\varepsilon}{2})$.

Then the function $\hat{h}: [a - \varepsilon, b + \varepsilon] \rightarrow \mathbb{R}$ defined by $\hat{h} = h \circ \rho$ has all of the desired properties. The first two are immediate from the definition. The third one holds since $\hat{h}'(t) = h'(\rho(t))\rho'(t)$ and $h' < 0$ while $\rho' \geq 0$. To check the last property we point out that

$$-\frac{n}{n-1}\hat{h}^2(t) - 2\hat{h}'(t) = -\frac{n}{n-1}h(\rho(t))^2 - 2h'(\rho(t))\rho'(t) = \sigma + 2h'(\rho(t))(1 - \rho'(t)).$$

Since $h'(\rho(t)) < 0$ and $0 \leq \rho' \leq 1$, the above is always $\leq \sigma$ and it is $< \sigma$ if $\rho'(t) < 1$. This holds true in particular when $\hat{h}'(t) = 0$, that is, $\rho'(t) = 0$. \square

In order to construct our functions $h_j: V_j \rightarrow \mathbb{R}$, we make use of the following basic existence result of band maps; for the proof, we refer to [38, Lemma 4.1; 3, Lemma 7.2].

Lemma 6.2. *Let (V, g) be a Riemannian band and let $a < b$ be two real numbers. If $\text{width}(V, g) > b - a$, then there exists a smooth function $\beta: V \rightarrow [a, b]$ with $\beta(\partial_- V) = a$, $\beta(\partial_+ V) = b$ and $\text{Lip}(\beta) < 1$.*

Next, for a Riemannian band (X, g) , we give conditions on the scalar curvature, the mean curvature of ∂X and on the potential function h so that the μ -bubble associated to h produces a separating closed hypersurface admitting a metric of positive scalar curvature. For more details on μ -bubbles, we refer the reader to [15, Section 5].

Proposition 6.3. *Let $n \leq 7$ and let (X, g) be an n -dimensional oriented Riemannian band. Let $h: X \rightarrow \mathbb{R}$ be a smooth function with the property that*

$$\text{scal}(X, g) + \frac{n}{n-1}h^2 - 2|\nabla h| > 0. \quad (6.1)$$

Furthermore, suppose that the mean curvature satisfies

$$\mathbb{H}(\partial_\pm X, g) > \pm h|_{\partial_\pm X}. \quad (6.2)$$

Then there exist a closed embedded hypersurface Σ which separates $\partial_- X$ and $\partial_+ X$ and a constant $b > 0$ such that

$$\int_\Sigma \left(|\nabla_\Sigma \psi|^2 + \frac{1}{2} \text{scal}(\Sigma, g) \psi^2 \right) d\text{vol}_\Sigma \geq b \int_\Sigma \psi^2 d\text{vol}_\Sigma, \quad \forall \psi \in C^\infty(\Sigma). \quad (6.3)$$

Proof. Denote by $\mathcal{C}(X)$ the set of all Caccioppoli sets in X which contain an open neighborhood of $\partial_- X$ and are disjoint from $\partial_+ X$. For $\hat{\Omega} \in \mathcal{C}(X)$ consider the functional

$$\mathcal{A}_h(\hat{\Omega}) = \mathcal{H}^{n-1}(\partial^* \hat{\Omega} \cap \mathring{X}) - \int_{\hat{\Omega}} h d\mathcal{H}^n,$$

where $\partial^* \hat{\Omega}$ is the reduced boundary [11, Chapters 3, 4] of $\hat{\Omega}$. By Condition (6.2) and [25, Lemma 4.2], there exists a smooth μ -bubble $\Omega \in \mathcal{C}(X)$, that is, a smooth Caccioppoli set with

$$\mathcal{A}_h(\Omega) = \mathcal{I} := \inf \{ \mathcal{A}_h(\hat{\Omega}) \mid \hat{\Omega} \in \mathcal{C}(X) \}.$$

Then $\Sigma := \partial\Omega \cap \mathring{X}$ is a closed embedded hypersurface that separates $\partial_- X$ and $\partial_+ X$. Let ν be the outward pointing unit normal vector field to Σ . By the first variation formula for \mathcal{A}_h (see [25, Lemma 4.3]) the mean curvature of Σ (computed with respect to $-\nu$) is equal to $h|_\Sigma$. By stability, from the second variation formula (see [25, Lemma 4.4]) we deduce

$$\begin{aligned} \int_\Sigma \left(2|\nabla_\Sigma \psi|^2 + \text{scal}(\Sigma, g) \psi^2 \right) d\text{vol}_\Sigma &\geq \int_\Sigma \left(\text{scal}(X, g) + \frac{n}{n-1}h^2 + 2g(\nabla_X h, \nu) \right) \psi^2 d\text{vol}_\Sigma \\ &\geq \min_\Sigma \left\{ \text{scal}(X, g) + \frac{n}{n-1}h^2 - 2|\nabla h| \right\} \int_\Sigma \psi^2 d\text{vol}_\Sigma \end{aligned}$$

for all $\psi \in C^\infty(\Sigma)$. By Condition (6.1), the previous inequality yields a constant $b > 0$ such that Inequality (6.3) holds. \square

We have gathered all the ingredients we need to prove Theorem 3.5.

Proof of Theorem 3.5. Assume, by contradiction, that $\text{width}(V_j, g) > \text{width}(M_j, g_{\varphi_j})$ for all $j \in \{1, \dots, k+1\}$. Consider the functions

$$h_{\varphi_j}(t) = (n-1) \frac{\varphi_j'(t)}{\varphi_j(t)}: [a_j, b_j] \rightarrow \mathbb{R}.$$

Since the φ_j are strictly log-concave, the functions h_{φ_j} are strictly monotonously decreasing and the scalar curvature of (M_j, g_{φ_j}) is given by

$$\sigma_j = \text{scal}(M_j, g_{\varphi_j}) = -\frac{n}{n-1}h_{\varphi_j}^2 - 2h'_{\varphi_j}. \quad (6.4)$$

For $j \in \{2, \dots, k\}$, we apply Lemma 6.1 to h_{φ_j} and obtain smooth functions

$$\hat{h}_{\varphi_j}: [a_j - \varepsilon, b_j + \varepsilon] \rightarrow \mathbb{R}$$

with the aforementioned properties.

For $j = 1$ we extend the domain of h_{φ_j} to $[a_1 - \varepsilon, b_1]$ and apply the interpolation procedure of Lemma 6.1 only on the right hand side of the interval to produce $\hat{h}_{\varphi_1}: [a_1 - \varepsilon, b_1 + \varepsilon] \rightarrow \mathbb{R}$. For $j = k+1$ we extend the domain of $h_{\varphi_{k+1}}$ to $[a_{k+1}, b_{k+1} + \varepsilon]$ and apply the interpolation procedure of Lemma 6.1 only on the left hand side of the interval to produce $\hat{h}_{\varphi_{k+1}}: [a_{k+1} - \varepsilon, b_{k+1} + \varepsilon] \rightarrow \mathbb{R}$.

By Lemma 6.2, there are smooth maps

$$\beta_j: V_j \rightarrow [a_j - \varepsilon, b_j + \varepsilon]$$

such that $\beta_j(\partial_- V_j) = a_j - \varepsilon$, $\beta_j(\partial_+ V_j) = b_j + \varepsilon$ and $\text{Lip}(\beta_j) < 1$. Define $h: X \rightarrow \mathbb{R}$ by $h(x) = \hat{h}_{\varphi_j} \circ \beta_j(x)$ if $x \in V_j$. The function h is continuous since

$$\hat{h}_{\varphi_j}(b_j + \varepsilon) = \text{H}(\partial_+ M_j, g_{\varphi_j}) = -\text{H}(\partial_- M_{j+1}, g_{\varphi_{j+1}}) = \hat{h}_{\varphi_{j+1}}(a_{j+1} - \varepsilon)$$

for all $j \in \{1, \dots, k\}$. It is smooth since the $\hat{h}_{\varphi_j} \circ \beta_j$ are constant in a neighborhood of the separating hypersurfaces Σ_i , which partition the band.

Note that h satisfies Condition (6.1) by the chain rule, the fourth property of the \hat{h}_{φ_j} from Lemma 6.1, and since $\text{Lip}(\beta_j) < 1$. For the mean curvature of the boundary, the following holds true:

$$\text{H}(\partial_- X, g) \geq \text{H}(\partial_- M_1, g_{\varphi_1}) = -\hat{h}_{\varphi_1}(a_1) > -\hat{h}_{\varphi_1}(a_1 - \varepsilon) = -h|_{\partial_- X}$$

and

$$\text{H}(\partial_+ X, g) \geq \text{H}(\partial_+ M_{k+1}, g_{\varphi_{k+1}}) = \hat{h}_{\varphi_{k+1}}(b_{k+1}) > \hat{h}_{\varphi_{k+1}}(b_{k+1} + \varepsilon) = h|_{\partial_+ X}.$$

Hence, Condition (6.2) is satisfied as well. By Proposition 6.3, there exist a closed embedded hypersurface Σ which separates $\partial_- X$ and $\partial_+ X$ and a constant $b > 0$ such that Inequality (6.3) holds.

If $n = 2$, this yields an immediate contradiction by choosing $\psi = 1$ in (6.3). If $n = 3$, we again choose $\psi = 1$ and use Gauß-Bonnet to see that Σ admits a psc metric. If $n \geq 4$, then Σ admits a metric of positive scalar curvature by the conformal change argument of Schoen and Yau [31]. This contradicts the fact that X has Property A. \square

Proof of Corollary 3.6. Consider the function

$$\varphi_2: \left(-\frac{\pi}{\sqrt{\kappa n}}, \frac{\pi}{\sqrt{\kappa n}}\right) \rightarrow \mathbb{R}_+ \quad t \mapsto \cos\left(\frac{\sqrt{\kappa n} t}{2}\right)^{\frac{2}{n}},$$

which is strictly log-concave and has

$$h_{\varphi_2}(t) = -\sqrt{\kappa}(n-1) \tan\left(\frac{\sqrt{\kappa n} t}{2}\right).$$

Consider the function

$$\varphi_1: \mathbb{R}_+ \rightarrow \mathbb{R}_+ \quad t \mapsto t^{\frac{2}{n}},$$

which is strictly log-concave and has

$$h_{\varphi_1}(t) = \frac{2(n-1)}{nt}.$$

Since $h_{\varphi_1}(t) \rightarrow \infty$ as $t \rightarrow 0$, there is a value $t_- > 0$ such that $H(\partial_- X, g) \geq -h_{\varphi_1}(t_-)$. By continuity, there are $\delta_1, \delta_2 > 0$ small enough such that

$$h_{\varphi_2}\left(\frac{-d + \delta_1}{2}\right) = h_{\varphi_1}(\ell + \delta_2),$$

while $\delta_2 < t_-$ and hence $\ell + \delta_2 - t_- < \ell$. Let (N, g_N) be a closed scalar flat Riemannian manifold. We fix the model space

$$(M_1, g_{\varphi_1}) = (N \times [t_-, \ell + \delta_2], \varphi_1^2(t)g_N + dt^2)$$

with scalar curvature equal to zero and width $< \ell$.

Let $\varphi_3: \mathbb{R}_- \rightarrow \mathbb{R}_+$ be defined by $\varphi_3(t) = \varphi_1(-t)$. This function is strictly log-concave and $h_{\varphi_3}(t) = -h_{\varphi_1}(-t)$. Since $h_{\varphi_3}(t) \rightarrow -\infty$ as $t \rightarrow 0$, there is a value $t_+ < 0$ such that $H(\partial_X, g) \geq h_{\varphi_3}(t_+)$. Similarly as before, we find $\delta_3, \delta_4 > 0$ such that

$$h_{\varphi_2}\left(\frac{d - \delta_3}{2}\right) = h_{\varphi_3}(-\ell - \delta_4),$$

while $\delta_4 < -t_+$ and hence $\ell + \delta_4 + t_+ < \ell$. We fix the model space

$$(M_3, g_{\varphi_3}) = (N \times [t_-, \ell + \delta_4], \varphi_3^2(t)g_N + dt^2)$$

with scalar curvature equal to zero and width $< \ell$.

Finally, we fix the model space

$$(M_2, g_{\varphi_2}) = \left(N \times \left[\frac{-d + \delta_1}{2}, \frac{d - \delta_3}{2}\right], \varphi_2^2(t)g_N + dt^2\right)$$

with scalar curvature equal to $\kappa n(n-1)$ and width $< d$.

It follows from Theorem 3.5 that $\text{width}(V_j, g) \leq \text{width}(M_j, g_{\varphi_j})$ for at least one $i \in \{1, 2, 3\}$. Since $d = \text{width}(V_2, g) > \text{width}(M_2, g_{\varphi_2})$, we conclude that

$$\min\{\text{width}(V_1, g), \text{width}(V_3, g)\} < \ell,$$

which is what we wanted to prove. □

Proof of Corollary 3.7. Consider the function

$$\varphi_2: \left(-\frac{\pi}{\sqrt{\kappa n}}, \frac{\pi}{\sqrt{\kappa n}}\right) \rightarrow \mathbb{R}_+ \quad t \mapsto \cos\left(\frac{\sqrt{\kappa n} t}{2}\right)^{\frac{2}{n}},$$

which is strictly log-concave and has

$$h_{\varphi_2}(t) = -\sqrt{\kappa}(n-1) \tan\left(\frac{\sqrt{\kappa n} t}{2}\right).$$

Consider the function

$$\varphi_1: \mathbb{R}_+ \rightarrow \mathbb{R}_+ \quad t \mapsto \sinh\left(\frac{\sqrt{\sigma n} t}{2\sqrt{n-1}}\right)^{\frac{2}{n}},$$

which is strictly log-concave and has

$$h_{\varphi_1}(t) := \sqrt{\frac{\sigma(n-1)}{n}} \coth\left(\frac{\sqrt{\sigma nt}}{2\sqrt{n-1}}\right).$$

Since $h_{\varphi_1}(t) \rightarrow \infty$ as $t \rightarrow 0$, there is a value $t_- > 0$ such that the mean curvature $H(\partial_- X, g)$ is greater or equal to $-h_{\varphi_1}(t_-)$. By continuity, there are $\delta_1, \delta_2 > 0$ small enough such that $-\sigma > -\kappa n(n-1) \tan\left(\frac{\sqrt{\kappa n(d-2\delta_1)}}{4}\right)^2$ and

$$h_{\varphi_2}\left(\frac{-d + \delta_1}{2}\right) = h_{\varphi_1}(\ell + \delta_2),$$

while $\delta_2 < t_-$ and hence $\ell + \delta_2 - t_- < \ell$. Let (N, g_N) be a closed scalar flat Riemannian manifold. We fix the model space

$$(M_1, g_{\varphi_1}) = (N \times [t_-, \ell + \delta_2], \varphi_1^2(t)g_N + dt^2)$$

with scalar curvature equal to $-\sigma$ and width $< \ell$.

Let $\varphi_3: \mathbb{R}_- \rightarrow \mathbb{R}_+$ be defined by $\varphi_3(t) = \varphi_1(-t)$. This function is strictly log-concave and $h_{\varphi_3}(t) = -h_{\varphi_1}(-t)$. Since $h_{\varphi_3}(t) \rightarrow -\infty$ as $t \rightarrow 0$, there is a value $t_+ < 0$ such that $H(\partial_X, g) \geq h_{\varphi_3}(t_+)$. Similarly as before, we find $\delta_3, \delta_4 > 0$ such that $-\sigma > -\kappa n(n-1) \tan\left(\frac{\sqrt{\kappa n(d-2\delta_3)}}{4}\right)^2$ and

$$h_{\varphi_2}\left(\frac{d - \delta_3}{2}\right) = h_{\varphi_3}(-\ell - \delta_4),$$

while $\delta_4 < -t_+$ and hence $\ell + \delta_4 + t_+ < \ell$. We fix the model space

$$(M_3, g_{\varphi_3}) = (N \times [-\ell - \delta_4, t_+], \varphi_3^2(t)g_N + dt^2)$$

with scalar curvature equal to $-\sigma$ and width $< \ell$.

Finally we fix the model space

$$(M_2, g_{\varphi_2}) = \left(N \times \left[\frac{-d + \delta_1}{2}, \frac{d - \delta_3}{2}\right], \varphi_2^2(t)g_N + dt^2\right)$$

with scalar curvature equal to $\kappa n(n-1)$ and width $< d$.

It follows from Theorem 3.5 that $\text{width}(V_j, g) \leq \text{width}(M_j, g_{\varphi_j})$ for at least one $i \in \{1, 2, 3\}$. Since $d = \text{width}(V_2, g) > \text{width}(M_2, g_{\varphi_2})$, we conclude that

$$\min\{\text{width}(V_1, g), \text{width}(V_3, g)\} < \ell,$$

which is what we wanted to prove. \square

References

- [1] S. Cecchini. “Callias-type operators in C^* -algebras and positive scalar curvature on noncompact manifolds”. In: *Journal of Topology and Analysis* (2018). DOI: 10.1142/S1793525319500687.
- [2] S. Cecchini. “A long neck principle for Riemannian spin manifolds with positive scalar curvature”. In: *Geom. Funct. Anal.* 30.5 (2020), pp. 1183–1223. DOI: 10.1007/s00039-020-00545-1.
- [3] S. Cecchini and R. Zeidler. “Scalar and mean curvature comparison via the Dirac operator”. In: *Geom. Topol.* (to appear). arXiv: 2103.06833 [math.DG].
- [4] S. Cecchini and R. Zeidler. *The positive mass theorem and distance estimates in the spin setting*. 2021. arXiv: 2108.11972 [math.DG].
- [5] J. Chen, P. Liu, Y. Shi, and J. Zhu. *Incompressible hypersurface, positive scalar curvature and positive mass theorem*. 2021. arXiv: 2112.14442v1 [math.DG].

- [6] O. Chodosh and C. Li. *Generalized soap bubbles and the topology of manifolds with positive scalar curvature*. arXiv: 2008.11888 [math.DG].
- [7] O. Chodosh, C. Li, and Y. Liokumovich. “Classifying sufficiently connected PSC manifolds in 4 and 5 dimensions”. In: *Geom. Topol.* (to appear). arXiv: 2105.07306 [math.DG].
- [8] J. Ebert and G. Frenck. “The Gromov-Lawson-Chernysh surgery theorem”. In: *Bol. Soc. Mat. Mex. (3)* 27.2 (2021), Paper No. 37, 43. DOI: 10.1007/s40590-021-00310-w.
- [9] H. Freudenthal. “Über die Enden topologischer Räume und Gruppen”. In: *Math. Z.* 33.1 (1931), pp. 692–713. DOI: 10.1007/BF01174375.
- [10] S. Galatius, U. Tillmann, I. Madsen, and M. Weiss. “The homotopy type of the cobordism category”. In: *Acta Math.* 202.2 (2009), pp. 195–239. DOI: 10.1007/s11511-009-0036-9.
- [11] E. Giusti. *Minimal surfaces and functions of bounded variation*. Vol. 80. Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984, pp. xii+240. DOI: 10.1007/978-1-4684-9486-0.
- [12] M. Gromov and H. B. Lawson Jr. “The classification of simply connected manifolds of positive scalar curvature”. In: *Ann. of Math. (2)* 111.3 (1980), pp. 423–434. DOI: 10.2307/1971103.
- [13] M. Gromov and H. B. Lawson Jr. “Positive scalar curvature and the Dirac operator on complete Riemannian manifolds”. In: *Inst. Hautes Études Sci. Publ. Math.* 58 (1983), 83–196 (1984).
- [14] M. Gromov. “Metric inequalities with scalar curvature”. In: *Geom. Funct. Anal.* 28.3 (2018), pp. 645–726. DOI: 10.1007/s00039-018-0453-z.
- [15] M. Gromov. *Four Lectures on Scalar Curvature*. arXiv Version 6. 2019. arXiv: 1908.10612v6 [math.DG].
- [16] M. Gromov. *No metrics with Positive Scalar Curvatures on Aspherical 5-Manifolds*. 2020. arXiv: 2009.05332 [math.DG].
- [17] B. Hanke, D. Pape, and T. Schick. “Codimension two index obstructions to positive scalar curvature”. In: *Ann. Inst. Fourier (Grenoble)* 65.6 (2015), pp. 2681–2710. DOI: 10.5802/aif.3000.
- [18] J. L. Kazdan. “Deformation to positive scalar curvature on complete manifolds”. In: *Math. Ann.* 261.2 (1982), pp. 227–234. DOI: 10.1007/BF01456220.
- [19] M. Lesourd, R. Unger, and S.-T. Yau. *Positive Scalar Curvature on Noncompact Manifolds and the Liouville Theorem*. 2020. arXiv: 2009.12618 [math.DG].
- [20] M. Lesourd, R. Unger, and S.-T. Yau. “The Positive Mass Theorem with Arbitrary Ends”. In: *J. Differential Geom.* (to appear). arXiv: 2103.02744 [math.DG].
- [21] A. Lichnerowicz. “Spineurs harmoniques”. In: *C. R. Acad. Sci. Paris* 257 (1963), pp. 7–9.
- [22] F. C. Marques. “Deforming three-manifolds with positive scalar curvature”. In: *Ann. of Math. (2)* 176.2 (2012), pp. 815–863. DOI: 10.4007/annals.2012.176.2.3.
- [23] T. Miyazaki. “On the existence of positive scalar curvature metrics on non-simply-connected manifolds”. In: *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* 30.3 (1984), pp. 549–561.
- [24] P. Petersen. *Riemannian geometry*. Third. Vol. 171. Graduate Texts in Mathematics. Springer, Cham, 2016, pp. xviii+499. DOI: 10.1007/978-3-319-26654-1.
- [25] D. Råde. *Scalar and mean curvature comparison via μ -bubbles*. 2021. arXiv: 2104.10120 [math.DG].
- [26] J. Rosenberg. “ C^* -algebras, positive scalar curvature and the Novikov conjecture. II”. In: *Geometric methods in operator algebras (Kyoto, 1983)*. Vol. 123. Pitman Res. Notes Math. Ser. Longman Sci. Tech., Harlow, 1986, pp. 341–374.
- [27] J. Rosenberg. “ C^* -algebras, positive scalar curvature, and the Novikov conjecture”. In: *Inst. Hautes Études Sci. Publ. Math.* 58 (1983), 197–212 (1984).
- [28] J. Rosenberg and S. Stolz. “Manifolds of positive scalar curvature”. In: *Algebraic topology and its applications*. Vol. 27. Math. Sci. Res. Inst. Publ. Springer, New York, 1994, pp. 241–267. DOI: 10.1007/978-1-4613-9526-3_8.
- [29] J. Rosenberg and S. Stolz. “Metrics of positive scalar curvature and connections with surgery”. In: *Surveys on surgery theory, Vol. 2*. Vol. 149. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 2001, pp. 353–386.

- [30] T. Schick. “A counterexample to the (unstable) Gromov-Lawson-Rosenberg conjecture”. In: *Topology* 37.6 (1998), pp. 1165–1168. DOI: 10.1016/S0040-9383(97)00082-7.
- [31] R. Schoen and S. T. Yau. “On the structure of manifolds with positive scalar curvature”. In: *Manuscripta Math.* 28.1-3 (1979), pp. 159–183. DOI: 10.1007/BF01647970.
- [32] N. Smale. “Generic regularity of homologically area minimizing hypersurfaces in eight-dimensional manifolds”. In: *Comm. Anal. Geom.* 1.2 (1993), pp. 217–228. DOI: 10.4310/CAG.1993.v1.n2.a2.
- [33] S. Stolz. *Concordance classes of positive scalar curvature metrics*. Preprint. 1998. URL: <http://www3.nd.edu/~stolz/concordance.ps>.
- [34] R. Zeidler. “Secondary large-scale index theory and positive scalar curvature”. Doctoral dissertation. University of Göttingen, 2016.
- [35] R. Zeidler. “An index obstruction to positive scalar curvature on fiber bundles over aspherical manifolds”. In: *Algebr. Geom. Topol.* 17.5 (2017), pp. 3081–3094. DOI: 10.2140/agt.2017.17.3081.
- [36] R. Zeidler. “Band width estimates via the Dirac operator”. In: *J. Differential Geom.* (to appear). arXiv: 1905.08520 [math.DG].
- [37] R. Zeidler. “Width, largeness and index theory”. In: *SIGMA Symmetry Integrability Geom. Methods Appl.* 16 (2020), Paper No. 127, 15. DOI: 10.3842/SIGMA.2020.127.
- [38] J. Zhu. “Width estimate and doubly warped product”. In: *Trans. Amer. Math. Soc.* 374.2 (2021), pp. 1497–1511. DOI: 10.1090/tran/8263.
- [39] J. Zhu. “Rigidity results for complete manifolds with nonnegative scalar curvature”. In: *J. Differential Geom.* (to appear). arXiv: 2008.07028 [math.DG].