

Differential Expansion for antiparallel triple pretzels: the way the factorization is deformed

A. Morozov, N. Tselousov

MIPT, ITEP & IITP, Moscow, Russia

ABSTRACT

For a peculiar family of double braid knots there is a remarkable factorization formula for the coefficients of the differential (cyclotomic) expansion (DE), which nowadays is widely used to construct the exclusive Racah matrices S and \bar{S} in arbitrary representations. The origins of the factorization remain obscure and the special role of double braids remains a mystery. In an attempt to broaden the perspective, we extend the family of double braids to antiparallel triple pretzels, which are obtained by the defect-preserving deformation from the trefoil and all have defect zero. It turns out that factorization of DE coefficients is violated quite strongly, still remains described by an elegant formula, at least for all symmetric representations.

1 Introduction

An important task of modern theoretical physics is to study the hidden symmetries. They are well known to play the crucial role in selecting of physically relevant theoretical models. The oldest example is provided by gauge symmetries [1], which do not affect any physical observables, but allow to give physical theories a simple and well distinguished formulation. Another example is hidden integrability of exact non-perturbative (functional) integrals (considered as functions of the coupling constants), which are invariant under any change of integration variables, which preserve the boundary conditions [2]. The newly emerging concept, nicknamed *superintegrability* [3–6], is the often-unexpected factorization of the answers for cleverly chosen observables – which turns out to reflect additional hidden properties of the theory. Such factorization is registered in many different circumstances, and is not yet clearly explained in most cases – often treated as no more than an amusing accident. Still, the evidence is mounting that the phenomenon is quite general [7, 8] and should have a universal explanation – once again distinguishing between the physically relevant models and artificial creations of the human mind.

In this paper we address a particular question about particular factorization phenomenon – that of the coefficients of the differential expansion for HOMFLY-PT polynomials in knot theory. While looking rather special, this is quite a typical example – it concerns the obviously-relevant $3d$ Chern-Simons theory, and appropriate variables – the coefficients of the *differential (cyclotomic) expansion* [9–44] – are far from obvious in the original formulation. Still, it is slowly being recognized that these are (at least rather close to) the right variables [11] and their properties are the ones that should be studied by the future knot theory. The first, still not truly impressive, attempts were made at the level of \mathcal{C} -polynomials [45]. A much more profound is the factorization property for the particular case of double braid knots, discovered in [21] (see [14–18] for numerous further developments).

This paper is an attempt to understand what is so special about double braids, and what is the deformation of factorization property beyond them. This can be an important step in the search for the truly relevant variables beyond the double-braid example. What we do, we consider the most straightforward generalization of double braids – to antiparallel triple pretzels. This is the most general antiparallel evolution of the trefoil and – in accordance with the recent conjecture [46] about defect-invariance under antiparallel evolution – all these pretzels have defect zero. Thus this is indeed a kind of a *minimal* deformation of double braids and the coefficients of DE remain defined in just the same way. Still, as mentioned already in [21], factorization is dramatically violated. We wonder what is the way to describe this violation. The answer is an unexpectedly elegant formula – much simpler than one could anticipate and not immediately continued beyond the distinguished triple-pretzel family. We do not know yet, what implications it will have, but clearly it will help us to further advance the story of “the proper variables” and factorization properties.

As an additional test of our formulas for triple-pretzels we check that they possess still another factorization property – of colored HOFMLY at the roots of unite [7], which is a far-going generalization of the celebrated factorization of special polynomials [47, 48]. For even more amusing facts about HOMFLY at roots of unity see [49] and references therein.

2 Differential expansion (DE)

HOMFLY-PT polynomials are the Wilson-line averages in 3d Chern-Simons theory [50]:

$$H_R^K = \left\langle \text{Tr}_R \text{Pexp} \left(\oint_{\mathcal{K}} \mathcal{A} \right) \right\rangle \quad (1)$$

which depend on integration contour (knot) \mathcal{K} , the coupling constant g , the size N of the \mathfrak{sl}_N gauge algebra and its representation R . Dependence on g and N is actually through the non-perturbative variables $q = \exp\left(\frac{2\pi i}{g+N}\right)$ and $A = q^N$, of which H^K is – mysteriously – just a polynomial. For a recent progress in perturbative approach see [51, 52].

Dependence on \mathcal{K} and R is more tricky, and the better (less redundant) variables are the coefficients \mathcal{F}_Q^K of the *differential (cyclotomic) expansion* [9–44], which we write down restricted in two ways – to symmetric representations and to defect [23] zero – according to the limited consideration in this paper. This allows to avoid lengthier formulas and extra notations, unnecessary for what follows:


$$H_{[r]}^{K_0}(A, q) = 1 + \sum_{s=1}^r \frac{[r]_q!}{[s]_q! [r-s]_q!} \cdot \underbrace{\mathcal{F}_{[s]}^{K_0}(A, q)}_{\text{depends on knot}} \cdot \prod_{i=0}^{s-1} \{Aq^{r+i}\} \{Aq^{i-1}\} \quad (2)$$

where $\{x\} := x - x^{-1}$ and $[n]_q := \frac{\{q^n\}}{\{q\}}$. We do not discuss origins of this formula and refer to other papers on the subject [13, 46]. The main advantage of DE formulas is that it extracts functions, such as \mathcal{F}_Q^K , which truly distinguish knots from the other constituents of HOMFLY that are common for all knots. Equation (2) is literally true for defect-0 knots [23], thus \mathcal{K} carries additional label 0. For other defects factorization is slightly less pronounced. In this paper we deal with defect zero and do not need to go into more details on this difference.

The first example of zero defect knot is the trefoil knot 3_1 . In other words, zero defect means that the corresponding colored HOMFLY have the form (2) with the following DE coefficients:

$$\mathcal{F}_{[s]}^{3_1} = (-)^s A^{2s} q^{s(s-1)} \quad (3)$$

It is observed [46] that the form of DE (i.e. the defect of DE) does not change under the antiparallel evolution of knots. By antiparallel evolution we mean replacement some of the crossings of knot by the antiparallel two-strand braid of odd length. In particular, this defect evolution hypothesis means that the whole three parametric family of triple antiparallel pretzels $\text{Pr}(2n-1, 2m-1, 2l-1)$ have the same form of DE (2) as the trefoil knot, since they could be considered as descendants of the trefoil under antiparallel evolution (4).



(4)

The double braid knots are included in this family as $\text{Pr}(2n-1, 2m-1, 1)$ and posses surprisingly simple behaviour of DE coefficients:

$$\boxed{\mathcal{F}_{[s]}^{(n,m,1)} = \frac{F_{[s]}^{(n)} \cdot F_{[s]}^{(m)}}{F_{[s]}^{(1)}}} \quad (5)$$

Here (n, m, l) is a shortcut notation for $\text{Pr}(2n-1, 2m-1, 2l-1)$ and $F_{[s]}^{(n)}$ are the DE coefficients of twist knots:

$$F_{[s]}^{(n)} := \mathcal{F}_{[s]}^{(n,1,1)} \quad (6)$$

because twist knots are included in triple pretzels as $\text{Pr}(2n-1, 1, 1)$. This mysterious factorization property of double braid DE coefficients (5) holds beyond symmetric and even rectangular representations [15, 16]. In order to understand the origins and nature of this factorization property we consider the double braid family as the part of a bigger family of triple pretzels and investigate how the factorization is deformed – see the resulting formula for triple pretzels (33).

The figure-eight knot 4_1 ($n = -1$), trefoil 3_1 ($n = 1$) and unknot 0_1 ($n = 0$) have simplest DE coefficients:

$$F_{[s]}^{(-1)} = 1, \quad F_{[s]}^{(0)} = 0, \quad F_{[s]}^{(1)} = (-)^s A^{2s} q^{s(s-1)} \quad (7)$$

for other twist knots there is a general formula [12]:

$$F_{[s]}^{(n)} = q^{\frac{s(s-1)}{2}} A^s \cdot \sum_{j=0}^s (-)^j \cdot \frac{[s]!}{[j]![s-j]!} \cdot \frac{(Aq^{j-1})^{2jn} \{Aq^{2j-1}\}}{\prod_{i=j-1}^{s+j-1} \{Aq^i\}} \quad (8)$$

All these DE coefficients are encoded into the *one* semi-infinite lower triangular matrix \mathcal{B} which we call the KNTZ matrix [14, 16, 17]:

$$F_{[s]}^{(n)} = \sum_{j=0}^s (\mathcal{B}^{n+1})_{s,j} \quad (9)$$

where explicit form of matrix elements for $i, j = 0, 1, 2, \dots$:

$$\mathcal{B}_{i,j} = (-)^{i-j} A^{2i} q^{(i+j)(i-1)} \frac{[i]_q!}{[j]_q! [i-j]_q!}, \quad j \leq i \quad (10)$$

$$\mathcal{B}_{i,j} = 0, \quad j > i \quad (11)$$

$$\mathcal{B} = \begin{pmatrix} 1 & 0 & 0 & 0 & \dots \\ -A^2 & A^2 & 0 & 0 & \dots \\ A^4 q^2 & -[2]_q A^4 q^3 & A^4 q^4 & 0 & \dots \\ -A^6 q^6 & [3]_q A^6 q^8 & -[3]_q A^6 q^{10} & A^6 q^{12} & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix} \quad (12)$$

The approach to the twist knot DE coefficients through the KNTZ matrix turn out to be extremely productive: this structures generalizes to higher representations [15] and superpolynomial T-deformation [16].

3 Technicalities of HOMFLY-PT calculus for pretzel knots

At present there is yet no direct way to calculate DE coefficients $\mathcal{F}_Q^{\mathcal{K}}$ for arbitrary knots and they are extracted from the answers for colored HOMFLY that are derived with the help of various techniques. All pretzels are very simple examples of arborescent knots and all the calculations are very straightforward due to the well-developed methods of arborescent calculus [53–56]. In this section we provide explicit formulas that allow to compute any symmetric HOMFLY for a 3-pretzel family that we discuss in this paper.

The calculations involve only 4 matrices: S, \bar{S}, T, \bar{T} . The \mathcal{R} matrix eigenvalues in symmetric representations $R = [r]$ in the two channels $[r] \otimes [r]$ and $[r] \otimes [\bar{r}]$ are very simple:

$$T = \text{diag}\left((-)^{r+i-1} q^{-r^2+i(i-1)} A^{-r}\right), \quad \bar{T} = \text{diag}\left((-)^{r+i-1} q^{(i-1)(i-2)} A^{i-1}\right), \quad (13)$$

for $i = 1, \dots, r+1$. For symmetric representations the matrices S , which switch between $[r] \otimes \overline{[r]}$ and $\overline{[r]} \otimes [r]$ and \bar{S} acting within the space $\overline{[r]} \otimes \overline{[r]}$, are also well known [57, 58]: for $i, j = 1, \dots, r+1$

$$S_{ij} = \sqrt{\frac{\bar{d}_i}{d_j}} \cdot \alpha_{i-1, j-1} \quad \bar{S}_{ij} = \sqrt{\frac{\bar{d}_i}{\bar{d}_j}} \cdot \bar{\alpha}_{i-1, j-1} \quad (14)$$

where d_X and \bar{d}_X are dimensions of representations from $[r] \otimes [r]$ and $[r] \otimes \overline{[r]}$ respectively,

$$[r] \otimes [r] = \bigoplus_{j=1}^{r+1} [r+j-1, r-j+1] \quad (15)$$

$$d_j = \dim_{[r+j-1, r-j+1]} := \frac{[2j-1]_q}{\prod_{i=1}^{r+j} [i]_q \prod_{i=1}^{r+1-j} [i]_q} \cdot \prod_{i=1}^{r+j-2} \{Aq^i\} \prod_{i=-1}^{r-1-j} \{Aq^i\}$$

$$\otimes \overline{[r]} = \bigoplus_{i=1}^{r+1} [2(i-1), (i-1)^{N-2}] \quad (16)$$

$$\bar{d}_i = \dim_{[2(i-1), (i-1)^{N-2}]} := \{Aq^{2i-3}\} \{Aq^{-1}\} \prod_{j=0}^{i-3} \left(\frac{\{Aq^j\}}{[j+2]} \right)^2$$

The matrices S, \bar{S} are build from 6-j symbols []:

$$\begin{pmatrix} \alpha_{km}(r) \\ \bar{\alpha}_{km}(r) \end{pmatrix} = \frac{(-)^{r+k+m} [2m+1] \left([k!] [m!] \right)^2 [r-k]! [r-m]!}{[r+k+1]! [r+m+1]!} \cdot \sum_{j=\max(r+m, r+k)}^{\min(r+k+m, 2r)} \frac{(-)^j [j+1]!}{[2r-j]! \left([j-r-k]! [j-r-m]! [r+k+m-j]! \right)^2} \cdot \begin{pmatrix} \frac{\mathcal{D}_{r-m} \mathcal{D}_{j+1}}{\mathcal{D}_{r+k+1} \mathcal{D}_{j-r-m}} \\ \frac{\mathcal{D}_m^2 \mathcal{D}_{j+1}}{\mathcal{D}_{r+k+1} \mathcal{D}_{r+m+1} \mathcal{D}_{r+k+m-j}} \end{pmatrix} \quad (17)$$

where $\mathcal{D}_n := \frac{1}{[n]!} \prod_{j=-1}^{n-2} \{Aq^j\}$ are responsible for the deviation from the \mathfrak{sl}_2 case when $A = q^2$ and $\mathcal{D}_n = 1$. The resulting answer for antiparallel 3-pretzel family:

$$H_{[r]}^{(n, m, l)} = d_{[r]}^2 \cdot \sum_{j=1}^{r+1} d_j^{-1/2} (\bar{S} \bar{T}^{2n-1} S)_{1, j} (\bar{S} \bar{T}^{2m-1} S)_{1, j} (\bar{S} \bar{T}^{2l-1} S)_{1, j} \quad (18)$$

where (n, m, l) is the shortcut notation for odd antiparallel triple pretzels $\text{Pr}(2n-1, 2m-1, 2l-1)$. For example, $3_1 = (1, 1, 1)$ and $4_1 = (-1, 1, 1)$:

$$H_{[1]}^{(1, 1, 1)} = -A^4 + A^2 q^2 + \frac{A^2}{q^2}, \quad H_{[1]}^{(-1, 1, 1)} = A^2 + \frac{1}{A^2} - q^2 - \frac{1}{q^2} + 1. \quad (19)$$

4 Triple antiparallel pretzels

In this section we analyze DE coefficients for triple antiparallel pretzels $\text{Pr}(2n-1, 2m-1, 2l-1)$. All these knots have defect zero, thus the differential expansion is

$$\begin{aligned} H_{[1]}^{(n, m, l)} &= 1 + \mathcal{F}_{[1]}^{(n, m, l)}(A) \cdot \{Aq\} \{A/q\} \\ H_{[2]}^{(n, m, l)} &= 1 + [2] \cdot \mathcal{F}_{[1]}^{(n, m, l)}(A) \cdot \{Aq^2\} \{A/q\} + \mathcal{F}_{[2]}^{(n, m, l)}(A, q) \cdot \{Aq^3\} \{Aq^2\} \{A\} \{A/q\} \\ &\dots \\ H_{[r]}^{(n, m, l)} &= 1 + \sum_{s=1}^r \frac{[r]!}{[s]! [r-s]!} \cdot \mathcal{F}_{[s]}^{(n, m, l)} \cdot \prod_{j=0}^{s-1} \{Aq^{r+j}\} \{Aq^{j-1}\} \end{aligned} \quad (20)$$

Dealing with triple pretzel DE coefficients we follow the strategy: we consider twist knot DE coefficients $F_{[s]}^{(n)}$ as "building blocks" and express the resulting answers through them, in the direct analogy with double braid family (5).

4.1 The lowest coefficient $\mathcal{F}_{[1]}^{(n,m,l)}$

The first DE coefficient is

$$\mathcal{F}_{[1]}^{(2n-1,2m-1,2l-1)} = -\frac{A^{2(n+m+l-1)} + A^{2(n+m+l-2)} - A^{2(n+m-1)} - A^{2(n+l-1)} - A^{2(m+l-1)} + 1}{\{A\}^2} \quad (21)$$

so that the Alexander polynomial ($A = 1$ limit of HOMFLY polynomial) has the following form:

$$\text{Al}_{[1]}^{(2n-1,2m-1,2l-1)} = 1 - \underline{(nm + nl + ml - m - n - l + 1)} \cdot \{q\}^2 \quad (22)$$

and the underlined expression is independent of q , as required for defect-zero knots [23, 46].

Remarkably the first coefficient of the differential expansion is equal to

$$\boxed{\mathcal{F}_{[1]}^{(n,m,l)} = \frac{F_{[1]}^{(n)} F_{[1]}^{(m)} F_{[1]}^{(l)}}{\left(F_{[1]}^{(1)}\right)^2} + \frac{F_{[1]}^{(n-1)} F_{[1]}^{(m-1)} F_{[1]}^{(l-1)}}{F_{[1]}^{(1)}}} \quad (23)$$

i.e. has a simple expression through twist knot DE coefficients (8):

$$F_{[1]}^{(n)} = -\frac{A^2(A^{2n} - 1)}{A^2 - 1} = A \cdot \frac{1 - A^{2n}}{\{A\}} \quad (24)$$

Since $F_{[s]}^{(0)} = 0$, this formula reproduces factorization property (5) for all double braids with $(n, m, l) = (n, m, 1)$. Also (23) reproduces purely topological statement $(n, m, 0) = (n-1, m-1, 1)$, due to $F_{[s]}^{(-1)} = 1$.

4.2 The second coefficient $\mathcal{F}_{[2]}^{(n,m,l)}$

The second DE coefficient $\mathcal{F}_{[2]}^{(n,m,l)}$ has analogous first and second term as (23) with an additional "correction":

$$\boxed{F_{[2]}^{(n,m,l)} = \frac{F_{[2]}^{(n)} F_{[2]}^{(m)} F_{[2]}^{(l)}}{\left(F_{[2]}^{(1)}\right)^2} + \frac{F_{[2]}^{(n-1)} F_{[2]}^{(m-1)} F_{[2]}^{(l-1)}}{F_{[2]}^{(1)}} - q^3 [2] \left(F_{[2]}^{(n-1)} - F_{[1]}^{(n-1)}\right) \left(F_{[2]}^{(m-1)} - F_{[1]}^{(m-1)}\right) \left(F_{[2]}^{(l-1)} - F_{[1]}^{(l-1)}\right)} \quad (25)$$

with

$$F_{[2]}^{(n)} = qA^2 \cdot \left(\frac{1}{\{Aq\}\{A\}} - \frac{[2]A^{2n}}{\{Aq^2\}\{A\}} + \frac{q^{4n} A^{4n}}{\{Aq^2\}\{Aq\}} \right) \quad (26)$$

4.3 Generic symmetric representation $\mathcal{F}_{[s]}^{(n,m,l)}$

Going through several examples we managed to write the following conjectural formula:

$$\boxed{\mathcal{F}_{[r]}^{(n,m,k)} = \frac{F_{[r]}^{(n)} F_{[r]}^{(m)} F_{[r]}^{(k)}}{\left(F_{[r]}^{(1)}\right)^2} + \frac{F_{[r]}^{(n-1)} F_{[r]}^{(m-1)} F_{[r]}^{(l-1)}}{F_{[r]}^{(1)}} + \sum_{i=1}^{r-1} (-1)^i (A^2 q^{r-1})^{2i} \cdot \frac{[r]!}{[i]![r-i]!} \cdot \frac{G_{[r],i}^{(n)} G_{[r],i}^{(m)} G_{[r],i}^{(k)}}{F_{[r]}^{(1)}}} \quad (27)$$

where we define new functions $G_{[r],i}^{(n)}$ describing additional "correction" terms:

$$G_{[r],i}^{(n)} := q^{i(r-1)} \cdot \sum_{j=0}^i (-1)^j q^{-j(i-1)} \cdot \frac{[i]_q!}{[j]_q![i-j]_q!} \cdot F_{[r-j]}^{(n-1)} \quad (28)$$

Surprisingly the first and the second terms in (27) correspond to the "correction" summands for $i = 0$ and $i = r$:

$$\frac{F_{[r]}^{(n-1)} F_{[r]}^{(m-1)} F_{[r]}^{(l-1)}}{F_{[r]}^{(1)}} = \frac{G_{[r],0}^{(n)} G_{[r],0}^{(m)} G_{[r],0}^{(k)}}{F_{[r]}^{(1)}} \quad (29)$$

$$\frac{F_{[r]}^{(n)} F_{[r]}^{(m)} F_{[r]}^{(k)}}{\left(F_{[r]}^{(1)}\right)^2} = (-)^r \left(A^{2r} q^{r(r-1)}\right)^2 \cdot \frac{G_{[r],r}^{(n)} G_{[r],r}^{(m)} G_{[r],r}^{(k)}}{F_{[r]}^{(1)}} \quad (30)$$

due to $F_{[r]}^{(1)} = (-)^r A^{2r} q^{r(r-1)}$ and the following relations:

$$G_{[r],0}^{(n-1)} = F_{[r]}^{(n-1)} \quad (31)$$

$$(-)^r G_{[r],r}^{(n)} = \frac{F_{[r]}^{(n)}}{F_{[r]}^{(1)}} \quad (32)$$

The first relation follows directly from the definition (28), while the second is less trivial and we discuss it in Sec. 6. Finally, the resulting answer could be represented as extended tridiagonal sum:

$$\boxed{\mathcal{F}_{[r]}^{(n,m,k)} = \sum_{i=0}^r (-)^{i-r} \left(A^2 q^{r-1}\right)^{2i-r} \cdot \frac{[r]!}{[i]! [r-i]!} \cdot G_{[r],i}^{(n)} \cdot G_{[r],i}^{(m)} \cdot G_{[r],i}^{(k)}} \quad (33)$$

5 Factorization at the roots of unity

5.1 Generalities

It may seem that at $q = 1$ the Chern-Simons theory becomes trivial, since quantization and all \mathcal{R} -matrices disappear. However, this is not quite true for *reduced* HOMFLY polynomials, because they are sensitive to the next order of the $1/N$ expansion, according to the rule $q = \exp\left(\frac{2\pi i}{N+g}\right) = 1 + \frac{2\pi i}{N} + \dots$ and $A = q^N$. What happens, however, is that all the Wilson loop correlators in this approximation are factorized and *colored* HOMFLY are just powers of the fundamental ones (which are sometimes called *special* polynomials [47]) at $q = 1$:

$$H_R^{\mathcal{K}} \Big|_{q=1} = \left(H_{[1]}^{\mathcal{K}}\right)^{|R|} \Big|_{q=1} \quad (34)$$

This implies some factorization property of the DE coefficients for symmetric HOMFLY. Namely, substituting $q = 1$ to differential expansion for defect-zero knots (2) and taking into account $[s]_{q=1} = s$

$$H_R^{\mathcal{K}} \Big|_{q=1} = 1 + \sum_{s=1}^r \frac{r!}{s!(r-s)!} \cdot \{A\}^{2s} \cdot \mathcal{F}_{[s]}^{\mathcal{K}} \Big|_{q=1} = \left(1 + \{A\}^2 \cdot \mathcal{F}_{[1]}^{\mathcal{K}} \Big|_{q=1}\right)^r = \left(H_{[1]}^{\mathcal{K}}\right)^r \Big|_{q=1} \quad (35)$$

one concludes power law for DE coefficients of defect zero knots:

$$\mathcal{F}_{[s]}^{\mathcal{K}_0} \Big|_{q=1} = \left(\mathcal{F}_{[1]}^{\mathcal{K}_0}\right)^s \Big|_{q=1} \quad (36)$$

Less trivially, similar factorizations occur at other unimodular values of $q = e^{\frac{\pi\sqrt{-1}}{m}}$ [7]:

$$H_{[r]}^{\mathcal{K}} \Big|_{q=\sqrt{-1}} = H_{[\text{Rem}(r/m)]}^{\mathcal{K}} \cdot \left(H_{[m]}^{\mathcal{K}}\right)^{\text{IntPart}(r/m)} \Big|_{q=\sqrt{-1}} \quad (37)$$

where $r = m \cdot \text{IntPart}(r/m) + \text{Rem}(r/m)$. IntPart, Rem are respectively the integer part and the remainder correspondingly. In this paper we analyze first nontrivial root of unity $q = \sqrt{-1}$ ($m = 2$).

We can now check that these generic factorization properties are indeed respected by our expressions for the triple pretzels (33).

5.3.2 Simple examples

The factorization property can be seen from the simple examples:

$$H_{[1]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = 1 + \mathcal{F}_{[1]}^{\mathcal{K}_0} \cdot \{A\}_+^2 \Big|_{q=\sqrt{-1}} \quad (45)$$

$$H_{[2]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = 1 + \mathcal{F}_{[2]}^{\mathcal{K}_0} \cdot \{A\}^2 \{A\}_+^2 \Big|_{q=\sqrt{-1}} \quad (46)$$

For representation $[r] = [3]$:

$$\begin{aligned} H_{[3]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} &= 1 + \mathcal{F}_{[1]}^{\mathcal{K}_0} \cdot \{A\}_+^2 + \mathcal{F}_{[2]}^{\mathcal{K}_0} \cdot \{A\}^2 \{A\}_+^2 + \mathcal{F}_{[3]}^{\mathcal{K}_0} \cdot \{A\}^2 \{A\}_+^4 \Big|_{q=\sqrt{-1}} \\ H_{[3]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} &= H_{[1]}^{\mathcal{K}_0} \cdot H_{[2]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} \implies \mathcal{F}_{[3]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \mathcal{F}_{[1]}^{\mathcal{K}_0} \cdot \mathcal{F}_{[2]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} \end{aligned} \quad (47)$$

For representation $[r] = [4]$:

$$\begin{aligned} H_{[4]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} &= 1 + \mathcal{F}_{[2]}^{\mathcal{K}_0} \cdot \{A\}^2 \{A\}_+^2 + \mathcal{F}_{[4]}^{\mathcal{K}_0} \cdot \{A\}^4 \{A\}_+^4 \Big|_{q=\sqrt{-1}} \\ H_{[4]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} &= H_{[2]}^{\mathcal{K}_0} \cdot H_{[2]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} \implies \mathcal{F}_{[4]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \mathcal{F}_{[2]}^{\mathcal{K}_0} \cdot \mathcal{F}_{[2]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} \end{aligned} \quad (48)$$

5.3.3 Arbitrary representations

For arbitrary representation of even length $[r] = [2n]$:

$$H_{[2n]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \sum_{s=0}^n \frac{n!}{s!(n-s)!} \cdot \{A\}^{2s} \{A\}_+^{2s} \cdot \mathcal{F}_{[2s]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} \quad (49)$$

where all the q -binomial coefficient are evaluated at the point $q = \sqrt{-1}$ following (42). From the factorization property (37) and the last formula follows the corresponding factorization of the DE coefficients:

$$H_{[2n]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \left(H_{[2]}^{\mathcal{K}_0} \right)^n \Big|_{q=\sqrt{-1}} \implies \mathcal{F}_{[2n]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \left(\mathcal{F}_{[2]}^{\mathcal{K}_0} \right)^n \Big|_{q=\sqrt{-1}} \quad (50)$$

For representations of odd length $[r] = [2n+1]$ the situation is a little more involved:

$$H_{[2n+1]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \sum_{s=0}^n \frac{n!}{s!(n-s)!} \cdot \{A\}^{2s} \{A\}_+^{2s} \cdot \left(\mathcal{F}_{[2s]}^{\mathcal{K}_0} + \{A\}_+^2 \mathcal{F}_{[2s+1]}^{\mathcal{K}_0} \right) \Big|_{q=\sqrt{-1}} \quad (51)$$

when evaluation we used the following relations that appear from (42): $(-)^{\frac{1}{2}\text{Rem}(\frac{2n+2s-1}{4}) \cdot \text{Rem}(\frac{2s}{4})} = (-)^s$ and $(-)^{\frac{1}{2}\text{Rem}(\frac{2n+2s}{4}) \cdot \text{Rem}(\frac{2s+1}{4})} = (-)^{s+n}$. Therefore we conclude the factorization property for DE coefficients for zero defect knots:

$$H_{[2n+1]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = H_{[1]}^{\mathcal{K}_0} \cdot \left(H_{[2]}^{\mathcal{K}_0} \right)^n \Big|_{q=\sqrt{-1}} \implies \mathcal{F}_{[2n+1]}^{\mathcal{K}_0} \Big|_{q=\sqrt{-1}} = \mathcal{F}_{[1]}^{\mathcal{K}_0} \cdot \left(\mathcal{F}_{[2]}^{\mathcal{K}_0} \right)^n \Big|_{q=\sqrt{-1}} \quad (52)$$

5.3.4 Direct check of (33)

Now we evaluate main "building blocks" $F_{[r]}^{(n)} \Big|_{q=\sqrt{-1}}$ and $G_{[r],i}^{(n)} \Big|_{q=\sqrt{-1}}$ at the point $q = \sqrt{-1}$. From the explicit form (8) and (42):

$$F_{[2s]}^{(n)} \Big|_{q=\sqrt{-1}} = A^{2s} \cdot \sum_{j=0}^s (-)^j \cdot \frac{s!}{j!(s-j)!} \cdot \frac{A^{4jn}}{\{A\}^s \{A\}_+^s} = \left(\frac{A^2}{\{A\} \{A\}_+} \right)^s (1 - A^{4n})^s = \left(F_{[2]}^{(n)} \right)^s \Big|_{q=\sqrt{-1}} \quad (53)$$

For the odd length of the representations $[2s+1]$ the calculation involve splitting the sum for even and odd indices:

$$\begin{aligned} F_{[2s+1]}^{(n)} \Big|_{q=\sqrt{-1}} &= A^{2s+1} \cdot \sum_{j=0}^s (-)^j \cdot \frac{s!}{j!(s-j)!} \cdot \left(\frac{A^{4jn}}{\{A\}^{s+1} \{A\}_+^s} - \frac{A^{4jn+2n}}{\{A\}^{s+1} \{A\}_+^s} \right) = \\ &= \frac{A}{\{A\}} (1 - A^{2n}) \cdot \left(\frac{A^2}{\{A\} \{A\}_+} \right)^s (1 - A^{4n})^s = F_{[1]}^{(n)} \cdot \left(F_{[2]}^{(n)} \right)^s \Big|_{q=\sqrt{-1}} \end{aligned} \quad (54)$$

$$\begin{aligned}
G_{[2r],2i}^{(n)} \Big|_{q=\sqrt{-1}} &= \left(G_{[2],0}^{(n)}\right)^{r-i} \cdot \left(G_{[2],2}^{(n)}\right)^i \Big|_{q=\sqrt{-1}} \\
G_{[2r+1],2i}^{(n)} \Big|_{q=\sqrt{-1}} &= (-)^i \cdot G_{[1],0}^{(n)} \cdot G_{[2r],2i}^{(n)} \Big|_{q=\sqrt{-1}} \\
G_{[2r+2],2i+1}^{(n)} \Big|_{q=\sqrt{-1}} &= (-)^r \cdot G_{[2],1}^{(n)} \cdot G_{[2r],2i}^{(n)} \Big|_{q=\sqrt{-1}} \\
G_{[2r+1],2i+1}^{(n)} \Big|_{q=\sqrt{-1}} &= (-)^{i+r} \cdot G_{[1],1}^{(n)} \cdot G_{[2r],2i}^{(n)} \Big|_{q=\sqrt{-1}}
\end{aligned} \tag{55}$$

We check (37) for 3-pretzel DE coefficients (33) for even length of representations $[2r]$:

$$\begin{aligned}
\mathcal{F}_{[2r]}^{(n,m,l)} \Big|_{q=\sqrt{-1}} &= \left(-A^{-4} \cdot G_{[2],0}^{(n,m,l)}\right)^r \cdot \sum_{i=0}^r \frac{r!}{i!(r-i)!} \cdot A^{8i} \cdot \left(\frac{G_{[2],2}^{(n,m,l)}}{G_{[2],0}^{(n,m,l)}}\right)^i \Big|_{q=\sqrt{-1}} = \\
&= \left(-A^{-4} \cdot G_{[2],0}^{(n,m,l)} - A^4 \cdot G_{[2],2}^{(n,m,l)}\right)^r \Big|_{q=\sqrt{-1}} = \left(\mathcal{F}_{[2]}^{(n,m,l)}\right)^r \Big|_{q=\sqrt{-1}}
\end{aligned} \tag{56}$$

and odd length of representations $[2r+1]$:

$$\begin{aligned}
\mathcal{F}_{[2r+1]}^{(n,m,l)} \Big|_{q=\sqrt{-1}} &= \left(-A^{-2} G_{[1],0}^{(n,m,l)} + A^2 G_{[1],1}^{(n,m,l)}\right) \left(-A^{-4} G_{[2],0}^{(n,m,l)}\right)^r \sum_{i=0}^r \frac{r!}{i!(r-i)!} \cdot A^{8i} \cdot \left(\frac{G_{[2],2}^{(n,m,l)}}{G_{[2],0}^{(n,m,l)}}\right)^i \Big|_{q=\sqrt{-1}} = \\
&= \left(-A^{-2} \cdot G_{[1],0}^{(n,m,l)} + A^2 \cdot G_{[1],1}^{(n,m,l)}\right) \left(-A^{-4} \cdot G_{[2],0}^{(n,m,l)} - A^4 \cdot G_{[2],2}^{(n,m,l)}\right)^r \Big|_{q=\sqrt{-1}} = \\
&= \left(\mathcal{F}_{[1]}^{(n,m,l)}\right) \cdot \left(\mathcal{F}_{[2]}^{(n,m,l)}\right)^r \Big|_{q=\sqrt{-1}}
\end{aligned} \tag{57}$$

where we define $G_{[r],i}^{(n,m,l)} := G_{[r],i}^{(n)} \cdot G_{[r],i}^{(m)} \cdot G_{[r],i}^{(l)}$.

6 New \mathcal{C} -polynomial for the twist knots

The colored HOMFLY polynomials are observed to satisfy linear recurrence relations with respect to representation [60–63] known as quantum A-polynomials. For example the simplest relation hold for symmetric representations for the trefoil knot:

$$H_{[r]}^{3_1} + w_1(r) \cdot H_{[r-1]}^{3_1} + w_2(r) \cdot H_{[r-2]}^{3_1} = 0 \tag{58}$$

$$\begin{aligned}
w_1(r) &= -\frac{A(Aq^{2r}-q^2)(Aq^{2r}+q^2)(A^4q^{8r}-A^2q^{4r+2}-A^2q^{4r+6}-A^2q^{6r+4}-q^{2r+8}+q^{4r+6}+q^{4r+8}+q^8)}{q^7(q^r-1)(q^r+1)(q^3-Aq^{2r})(Aq^{2r}+q^3)} \\
w_2(r) &= \frac{A^2q^{6r-6}(Aq^r-q^2)(Aq^r+q^2)(Aq^{2r}-q)(Aq^{2r}+q)}{(q^r-1)(q^r+1)(q^3-Aq^{2r})(Aq^{2r}+q^3)}
\end{aligned} \tag{59}$$

with "initial conditions" $H_{[0]}^{3_1} = 1$ and $H_{[1]}^{3_1} = -A^4 + A^2q^2 + \frac{A^2}{q^2}$.

Within the framework of differential expansion (2) one can study recurrence relations on DE coefficients that known as \mathcal{C} -polynomials [45, 64]. For the trefoil knot this relation has the following form:

$$\mathcal{F}_{[s]}^{3_1} + A^2q^{2(s-1)} \cdot \mathcal{F}_{[s-1]}^{3_1} = 0 \tag{60}$$

that is much simpler than the original quantum A-polynomial. The simplicity is not understood completely. One of the possible explanations is that the quantum A-polynomial describes highly nontrivial knot-independent factors of DE (2), while \mathcal{C} polynomial deals directly with rather simple DE coefficients.

In recent work [45] was proposed to consider evolution equations on DE coefficients in another direction – so called A -evolution:

$$\sum_{i=0} t_i(A, q) \cdot \mathcal{F}_{[s]}^{\mathcal{K}}(Aq^i, q) = 0 \tag{61}$$

where different terms in the sum undergo a substitution $A \rightarrow Aq^i$ in the DE coefficients.

We propose to consider one more evolution direction – in the knot space . We argue that there is a simple \mathcal{C} -polynomial of mixed "representation-knot" evolution for twist knot family. Recall the relation (32) from sec.4.3:

$$(-)^r G_{[r],r}^{(n)} = \frac{F_{[r]}^{(n)}}{F_{[r]}^{(1)}} \quad (62)$$

that provide a simple form of the resulting formula (33). Using explicit form of $G_{[r],r}^{(n)}$ functions (28) it takes a form

$$F_{[r]}^{(n)} = A^{2r} q^{2r(r-1)} \sum_{j=0}^r (-1)^{j+1} q^{-j(r-1)} \cdot \frac{[r]!}{[j]![r-j]!} \cdot F_{[r-j]}^{(n-1)} \quad (63)$$

of nontrivial relation that connects DE coefficients for different representations of two twist knots. Remarkably, this relation is nothing but a consequence of formula (9) involving KNTZ matrix $\mathcal{B}_{i,j}$ (10):

$$F_{[r]}^{(n)} = \sum_{j=0}^i \mathcal{B}_{r,j} \cdot F_{[j]}^{(n-1)} \quad (64)$$

We consider this relation as a \mathcal{C} -polynomial of *mixed* evolution along representation $[r]$ and twist knot direction (n) . This mixed \mathcal{C} -polynomial has much simpler form comparing with pure evolution along representation $[r]$ (see (59) in [45]). Moreover, this equation has a remarkable property: if one knows all DE coefficients $F_{[r]}$ for a particular twist knot, then one can find DE coefficients for *all* other twist knot. In particular,

$$F_{[r]}^{(-1)} = 1 \quad \implies \quad F_{[r]}^{(n)} = \sum_{j=0}^r (\mathcal{B}^{n+1})_{r,j} \quad (65)$$

that reproduces (9).

7 Conclusion

In this paper we report a remarkably simple formula for the coefficients of the differential expansion of defect-zero family of antiparallel pretzel knots. It would be interesting to extend this result in several directions:

- **Non-symmetric representations.**

The generalization of the results (33) to higher representations looks straightforward, still technically involved and interesting. Based on the form of DE coefficients (33) one could try the following ansatz:

$$G_{R,Q}^{(n)} := \sum_{\lambda: Q \subset \lambda \subset R} V_{R,Q}^\lambda \cdot F_\lambda^{(n-1)} \quad (66)$$

$$\mathcal{F}_R^{(n,m,l)} = \sum_{Q \in R} W_{R,Q} \cdot G_{R,Q}^{(n)} \cdot G_{R,Q}^{(m)} \cdot G_{R,Q}^{(l)} \quad (67)$$

with appropriate functions $W_{R,Q}$ and $V_{R,Q}^\lambda$. We expect triple factorization for higher representations for triple pretzels due to the fact that double factorization holds in arbitrary representation for double-braid family. Still it is unclear if this optimistic suggestion will work.

- **Other knot families and higher defects.**

The formulas of DE coefficients for other pretzels and more general families of knots, including non-arborescent ones, are less simple – especially when defect is non-vanishing. It is interesting to see to what extent the problem will reduce to KNTZ matrix and twist DE coefficients, like it happened for the zero-defect triple pretzel family.

Even more interesting is to find appropriate interpretation of these remarkable formulas for triple pretzels per se — and the way they deviate from factorization in the double-braid case. It can happen that the corollaries and applications of such understanding will be no less impressive than those for the double braids.

Acknowledgements

We are indebted to E. Lanina, And.Morozov and A.Sleptsov for conversations and help. This work is supported by the Russian Science Foundation (Grant No.20-71-10073).

References

- [1] L.B. Okun. *Leptons and Quarks*. 1980.
- [2] A. Morozov. “Integrability and matrix models”. In: *Phys. Usp.* 37 (1994), pp. 1–55. DOI: [10.1070/PU1994v037n01ABEH0000](https://doi.org/10.1070/PU1994v037n01ABEH0000) arXiv: [hep-th/9303139](https://arxiv.org/abs/hep-th/9303139).
- [3] A. Mironov and A. Morozov. “Superintegrability summary”. In: (Jan. 2022). arXiv: [2201.12917](https://arxiv.org/abs/2201.12917) [[hep-th](#)].
- [4] A. D. Mironov and A. Morozov. “Generalized Q-functions for GKM”. In: *Phys. Lett. B* 819 (2021), p. 136474. DOI: [10.1016/j.physletb.2021.136474](https://doi.org/10.1016/j.physletb.2021.136474). arXiv: [2101.08759](https://arxiv.org/abs/2101.08759) [[hep-th](#)].
- [5] Andrei Mironov and Alexei Morozov. “Superintegrability of Kontsevich matrix model”. In: *Eur. Phys. J. C* 81.3 (2021), p. 270. DOI: [10.1140/epjc/s10052-021-09030-x](https://doi.org/10.1140/epjc/s10052-021-09030-x). arXiv: [2011.12917](https://arxiv.org/abs/2011.12917) [[hep-th](#)].
- [6] H. Itoyama, A. Mironov, and A. Morozov. “Tensorial generalization of characters”. In: *JHEP* 12 (2019), p. 127. DOI: [10.1007/JHEP12\(2019\)127](https://doi.org/10.1007/JHEP12(2019)127). arXiv: [1909.06921](https://arxiv.org/abs/1909.06921) [[hep-th](#)].
- [7] Ya. Kononov and A. Morozov. “Factorization of colored knot polynomials at roots of unity”. In: *Phys. Lett. B* 747 (2015), pp. 500–510. DOI: [10.1016/j.physletb.2015.06.043](https://doi.org/10.1016/j.physletb.2015.06.043). arXiv: [1505.06170](https://arxiv.org/abs/1505.06170) [[hep-th](#)].
- [8] Ya. Kononov and A. Morozov. “On Factorization of Generalized Macdonald Polynomials”. In: *Eur. Phys. J. C* 76.8 (2016), p. 424. DOI: [10.1140/epjc/s10052-016-4276-5](https://doi.org/10.1140/epjc/s10052-016-4276-5). arXiv: [1607.00615](https://arxiv.org/abs/1607.00615) [[hep-th](#)].
- [9] H. Itoyama et al. “HOMFLY and superpolynomials for figure eight knot in all symmetric and antisymmetric representations”. In: *JHEP* 07 (2012), p. 131. DOI: [10.1007/JHEP07\(2012\)131](https://doi.org/10.1007/JHEP07(2012)131). arXiv: [1203.5978](https://arxiv.org/abs/1203.5978) [[hep-th](#)].
- [10] Andrei Mironov, Alexei Morozov, and Andrey Morozov. “On colored HOMFLY polynomials for twist knots”. In: *Mod. Phys. Lett. A* 29.34 (2014), p. 1450183. DOI: [10.1142/S0217732314501831](https://doi.org/10.1142/S0217732314501831). arXiv: [1408.3076](https://arxiv.org/abs/1408.3076) [[hep-th](#)].
- [11] S. B. Arthamonov, A. Mironov, and A. Morozov. “Differential hierarchy and additional grading of knot polynomials”. In: *Theor. Math. Phys.* 179 (2014), pp. 509–542. DOI: [10.1007/s11232-014-0159-9](https://doi.org/10.1007/s11232-014-0159-9). arXiv: [1306.5682](https://arxiv.org/abs/1306.5682) [[hep-th](#)].
- [12] A. Mironov, A. Morozov, and An. Morozov. “Evolution method and “differential hierarchy” of colored knot polynomials”. In: *AIP Conf. Proc.* 1562.1 (2013). Ed. by Wen-Xiu Ma and David Kaup, pp. 123–155. DOI: [10.1063/1.4828688](https://doi.org/10.1063/1.4828688). arXiv: [1306.3197](https://arxiv.org/abs/1306.3197) [[hep-th](#)].
- [13] L. Bishler and A. Morozov. “Perspectives of differential expansion”. In: *Phys. Lett. B* 808 (2020), p. 135639. DOI: [10.1016/j.physletb.2020.135639](https://doi.org/10.1016/j.physletb.2020.135639). arXiv: [2006.01190](https://arxiv.org/abs/2006.01190) [[hep-th](#)].
- [14] A. Morozov. “KNTZF trick from arborescent calculus and the structure of differential expansion”. In: *Theor. Math. Phys.* 204 (2020), pp. 863–889. DOI: [10.1134/S0040577920080036](https://doi.org/10.1134/S0040577920080036). arXiv: [2001.10254](https://arxiv.org/abs/2001.10254) [[hep-th](#)].
- [15] A. Morozov. “Extension of KNTZ trick to non-rectangular representations”. In: *Phys. Lett. B* 793 (2019), pp. 464–468. DOI: [10.1016/j.physletb.2019.05.016](https://doi.org/10.1016/j.physletb.2019.05.016). arXiv: [1903.00259](https://arxiv.org/abs/1903.00259) [[hep-th](#)].
- [16] A. Morozov. “On exclusive Racah matrices \bar{S} for rectangular representations”. In: *Phys. Lett. B* 793 (2019), pp. 116–125. DOI: [10.1016/j.physletb.2019.04.034](https://doi.org/10.1016/j.physletb.2019.04.034). arXiv: [1902.04140](https://arxiv.org/abs/1902.04140) [[hep-th](#)].
- [17] R.Tao M. Kameyama S.Nawata and H. D. Zhang. “Cyclotomic expansions of HOMFLY-PT colored by rectangular Young diagrams”. In: *Lett. Math. Phys.* 110 (2020), pp. 2573–2583. DOI: [10.1007/s11005-020-01318-5](https://doi.org/10.1007/s11005-020-01318-5). arXiv: [1902.02275](https://arxiv.org/abs/1902.02275) [[math.GT](#)].
- [18] Shengmao Zhu Qingtao Chen Kefeng Liu. “Cyclotomic expansions for the colored HOMFLY-PT invariants of double twist knots”. In: (2021). arXiv: [2110.03616](https://arxiv.org/abs/2110.03616) [[math.GT](#)].
- [19] A. Morozov. “Pentad and triangular structures behind the Racah matrices”. In: *Eur. Phys. J. Plus* 135.2 (2020), p. 196. DOI: [10.1140/epjp/s13360-020-00234-w](https://doi.org/10.1140/epjp/s13360-020-00234-w). arXiv: [1906.09971](https://arxiv.org/abs/1906.09971) [[hep-th](#)].
- [20] A. Morozov. “Differential expansion and rectangular HOMFLY for the figure eight knot”. In: *Nucl. Phys. B* 911 (2016), pp. 582–605. DOI: [10.1016/j.nuclphysb.2016.08.027](https://doi.org/10.1016/j.nuclphysb.2016.08.027). arXiv: [1605.09728](https://arxiv.org/abs/1605.09728) [[hep-th](#)].
- [21] A. Morozov. “Factorization of differential expansion for antiparallel double-braid knots”. In: *JHEP* 09 (2016), p. 135. DOI: [10.1007/JHEP09\(2016\)135](https://doi.org/10.1007/JHEP09(2016)135). arXiv: [1606.06015](https://arxiv.org/abs/1606.06015) [[hep-th](#)].
- [22] C. Bai et al. “Differential expansion for link polynomials”. In: *Phys. Lett. B* 778 (2018), pp. 197–206. DOI: [10.1016/j.physletb.2018.01.026](https://doi.org/10.1016/j.physletb.2018.01.026). arXiv: [1709.09228](https://arxiv.org/abs/1709.09228) [[hep-th](#)].
- [23] Ya. Kononov and A. Morozov. “On the defect and stability of differential expansion”. In: *JETP Lett.* 101.12 (2015), pp. 831–834. DOI: [10.1134/S0021364015120127](https://doi.org/10.1134/S0021364015120127). arXiv: [1504.07146](https://arxiv.org/abs/1504.07146) [[hep-th](#)].

- [24] Ya. Kononov and A. Morozov. “Rectangular superpolynomials for the figure-eight knot 4_1 ”. In: *Theor. Math. Phys.* 193.2 (2017), pp. 1630–1646. DOI: [10.1134/S0040577917110058](https://doi.org/10.1134/S0040577917110058). arXiv: [1609.00143](https://arxiv.org/abs/1609.00143) [hep-th].
- [25] Ya. Kononov and A. Morozov. “On rectangular HOMFLY for twist knots”. In: *Mod. Phys. Lett. A* 31.38 (2016), p. 1650223. DOI: [10.1142/S0217732316502230](https://doi.org/10.1142/S0217732316502230). arXiv: [1610.04778](https://arxiv.org/abs/1610.04778) [hep-th].
- [26] H. Itoyama et al. “Character expansion for HOMFLY polynomials. III. All 3-Strand braids in the first symmetric representation”. In: *Int. J. Mod. Phys. A* 27 (2012), p. 1250099. DOI: [10.1142/S0217751X12500996](https://doi.org/10.1142/S0217751X12500996). arXiv: [1204.4785](https://arxiv.org/abs/1204.4785) [hep-th].
- [27] H. Itoyama et al. “Eigenvalue hypothesis for Racah matrices and HOMFLY polynomials for 3-strand knots in any symmetric and antisymmetric representations”. In: *Int. J. Mod. Phys. A* 28 (2013), p. 1340009. DOI: [10.1142/S0217751X13400095](https://doi.org/10.1142/S0217751X13400095). arXiv: [1209.6304](https://arxiv.org/abs/1209.6304) [math-ph].
- [28] Kazuo Habiro. “A unified Witten-Reshetikhin-Turaev invariant for integral homology spheres”. In: *Inventiones mathematicae* 171.1 (2007), pp. 1–81. DOI: [10.1007/s00222-007-0071-0](https://doi.org/10.1007/s00222-007-0071-0).
- [29] Satoshi Nawata and Alexei Oblomkov. “Lectures on knot homology”. In: *Contemp. Math.* 680 (2016). Ed. by Sergei Gukov, Mikhail Khovanov, and Johannes Walcher, p. 137. DOI: [10.1090/conm/680/13702](https://doi.org/10.1090/conm/680/13702). arXiv: [1510.01795](https://arxiv.org/abs/1510.01795) [math-ph].
- [30] Qintao Chen et al. “Congruent skein relations for colored HOMFLY-PT invariants and colored Jones polynomials”. In: (Feb. 2014). arXiv: [1402.3571](https://arxiv.org/abs/1402.3571) [math.GT].
- [31] Qingtao Chen, Kefeng Liu, and Shengmao Zhu. “Volume conjecture for $SU(n)$ -invariants”. In: (Nov. 2015). arXiv: [1511.00658](https://arxiv.org/abs/1511.00658) [math.QA].
- [32] Qingtao Chen. “Cyclotomic expansion and volume conjecture for superpolynomials of colored HOMFLY-PT homology and colored Kauffman homology”. In: (Dec. 2015). arXiv: [1512.07906](https://arxiv.org/abs/1512.07906) [math.QA].
- [33] Kenichi Kawagoe. “On the formulae for the colored HOMFLY polynomials”. In: *J. Geom. Phys.* 106 (2016), pp. 143–154. DOI: [10.1016/j.geomphys.2016.02.012](https://doi.org/10.1016/j.geomphys.2016.02.012). arXiv: [1210.7574](https://arxiv.org/abs/1210.7574) [math.GT].
- [34] Kenichi Kawagoe. “The colored HOMFLY-PT polynomials of the trefoil knot, the figure-eight knot and twist knots”. In: (July 2021). arXiv: [2107.08678](https://arxiv.org/abs/2107.08678) [math.GT].
- [35] Anna Beliakova and Eugene Gorsky. *Cyclotomic expansions for \mathfrak{gl}_N knot invariants via interpolation Macdonald polynomials*. 2021. arXiv: [2101.08243](https://arxiv.org/abs/2101.08243).
- [36] Eugene Gorsky, Sergei Gukov, and Marko Stosic. “Quadruply-graded colored homology of knots”. In: (Apr. 2013). arXiv: [1304.3481](https://arxiv.org/abs/1304.3481) [math.QA].
- [37] Sergei Gukov and Marko Stošić. “Homological Algebra of Knots and BPS States”. In: *Proc. Symp. Pure Math.* 85 (2012). Ed. by Jonathan Block et al., pp. 125–172. DOI: [10.1090/pspum/085/1377](https://doi.org/10.1090/pspum/085/1377). arXiv: [1112.0030](https://arxiv.org/abs/1112.0030) [hep-th].
- [38] Nathan M. Dunfield, Sergei Gukov, and Jacob Rasmussen. “The Superpolynomial for knot homologies”. In: (May 2005). arXiv: [math/0505662](https://arxiv.org/abs/math/0505662).
- [39] Yuri Berest, Joseph Gallagher, and Peter Samuelson. “Cyclotomic expansion of generalized Jones polynomials”. In: *Letters in Mathematical Physics* 111.2 (2021), pp. 1–32. arXiv: [1908.04415](https://arxiv.org/abs/1908.04415).
- [40] Jeremy Lovejoy and Robert Osburn. “The colored Jones polynomial and Kontsevich-Zagier series for double twist knots, II”. In: (2019). arXiv: [1903.05060](https://arxiv.org/abs/1903.05060).
- [41] Jeremy Lovejoy and Robert Osburn. “The colored Jones polynomial and Kontsevich-Zagier series for double twist knots”. In: (2017). arXiv: [1710.04865](https://arxiv.org/abs/1710.04865).
- [42] Kazuhiro Hikami and Jeremy Lovejoy. “Torus knots and quantum modular forms”. In: *Research in the Mathematical Sciences* 2.1 (2015), pp. 1–15. arXiv: [1409.6243](https://arxiv.org/abs/1409.6243).
- [43] Stavros Garoufalidis and Thang TQ Le. “An analytic version of the melvin-morton-rozansky conjecture”. In: (2005). arXiv: [math/0503641](https://arxiv.org/abs/math/0503641).
- [44] Stavros Garoufalidis and Thang T Q Lê. “Asymptotics of the colored Jones function of a knot”. In: *Geometry & Topology* 15.4 (2011), pp. 2135–2180. arXiv: [math/0508100](https://arxiv.org/abs/math/0508100).
- [45] Andrei Mironov and A. Morozov. “Algebra of quantum \mathcal{C} -polynomials”. In: *JHEP* 02 (2021), p. 142. DOI: [10.1007/JHEP02\(2021\)142](https://doi.org/10.1007/JHEP02(2021)142). arXiv: [2009.11641](https://arxiv.org/abs/2009.11641) [hep-th].
- [46] A. Morozov and N. Tselousov. “Evolution properties of the knot’s defect”. In: (Apr. 2022). arXiv: [2204.05977](https://arxiv.org/abs/2204.05977) [hep-th].

- [47] P. Dunin-Barkowski et al. “Superpolynomials for toric knots from evolution induced by cut-and-join operators”. In: *JHEP* 03 (2013), p. 021. DOI: [10.1007/JHEP03\(2013\)021](https://doi.org/10.1007/JHEP03(2013)021). arXiv: [1106.4305](https://arxiv.org/abs/1106.4305) [[hep-th](#)].
- [48] Shengmao Zhu. “Colored HOMFLY polynomials via skein theory”. In: *JHEP* 10 (2013), p. 229. DOI: [10.1007/JHEP10\(2013\)229](https://doi.org/10.1007/JHEP10(2013)229). arXiv: [1206.5886](https://arxiv.org/abs/1206.5886) [[math.GT](#)].
- [49] Liudmila Bishler, Andrei Mironov, and Andrey Morozov. “Invariants of knots and links at roots of unity”. In: (May 2022). arXiv: [2205.05650](https://arxiv.org/abs/2205.05650) [[hep-th](#)].
- [50] Shiin-Shen Chern and James Simons. “Characteristic forms and geometric invariants”. In: *Annals Math.* 99 (1974), pp. 48–69. DOI: [10.2307/1971013](https://doi.org/10.2307/1971013).
- [51] E. Lanina, A. Sleptsov, and N. Tselousov. “Implications for colored HOMFLY polynomials from explicit formulas for group-theoretical structure”. In: *Nucl. Phys. B* 974 (2022), p. 115644. DOI: [10.1016/j.nuclphysb.2021.115644](https://doi.org/10.1016/j.nuclphysb.2021.115644). arXiv: [2111.11751](https://arxiv.org/abs/2111.11751) [[hep-th](#)].
- [52] E. Lanina, A. Sleptsov, and N. Tselousov. “Chern-Simons perturbative series revisited”. In: *Phys. Lett. B* 823 (2021), p. 136727. DOI: [10.1016/j.physletb.2021.136727](https://doi.org/10.1016/j.physletb.2021.136727). arXiv: [2105.11565](https://arxiv.org/abs/2105.11565) [[hep-th](#)].
- [53] A. Mironov et al. “Colored HOMFLY polynomials of knots presented as double fat diagrams”. In: *JHEP* 07 (2015), p. 109. DOI: [10.1007/JHEP07\(2015\)109](https://doi.org/10.1007/JHEP07(2015)109). arXiv: [1504.00371](https://arxiv.org/abs/1504.00371) [[hep-th](#)].
- [54] Nafaa Chbili and Vivek Kumar Singh. “Colored HOMFLY-PT polynomials of quasi-alternating 3-braid knots”. In: (Feb. 2022). arXiv: [2202.09169](https://arxiv.org/abs/2202.09169) [[hep-th](#)].
- [55] Saswati Dhara et al. “Eigenvalue hypothesis for multistrand braids”. In: *Phys. Rev. D* 97.12 (2018), p. 126015. DOI: [10.1103/PhysRevD.97.126015](https://doi.org/10.1103/PhysRevD.97.126015). arXiv: [1711.10952](https://arxiv.org/abs/1711.10952) [[hep-th](#)].
- [56] A. Mironov et al. “Tabulating knot polynomials for arborescent knots”. In: *J. Phys. A* 50.8 (2017), p. 085201. DOI: [10.1088/1751-8121/aa5574](https://doi.org/10.1088/1751-8121/aa5574). arXiv: [1601.04199](https://arxiv.org/abs/1601.04199) [[hep-th](#)].
- [57] L.D. Landau and E.M. Lifshitz, eds. *Quantum Mechanics*. Third Edition. Pergamon, 1977, p. ii. ISBN: 978-0-08-020940-1. DOI: doi.org/10.1016/B978-0-08-020940-1.50001-3.
- [58] A. Mironov, A. Morozov, and A. Sleptsov. “Colored HOMFLY polynomials for the pretzel knots and links”. In: *JHEP* 07 (2015), p. 069. DOI: [10.1007/JHEP07\(2015\)069](https://doi.org/10.1007/JHEP07(2015)069). arXiv: [1412.8432](https://arxiv.org/abs/1412.8432) [[hep-th](#)].
- [59] Yuxiao Wang Quanlin Chen Tianze Jiang. “The generational behavior of Gaussian binomial coefficients at roots of unity”. In: ().
- [60] Stavros Garoufalidis, Aaron D. Lauda, and Thang T. Q. Lê. “The colored HOMFLYPT function is q -holonomic”. In: *Duke Math. J.* 167.3 (2018), pp. 397–447. DOI: [10.1215/00127094-2017-0030](https://doi.org/10.1215/00127094-2017-0030). arXiv: [1604.08502](https://arxiv.org/abs/1604.08502) [[math.GT](#)].
- [61] Stavros Garoufalidis. “The colored HOMFLY polynomial is q -holonomic”. In: (Nov. 2012). arXiv: [1211.6388](https://arxiv.org/abs/1211.6388) [[math.GT](#)].
- [62] Stavros Garoufalidis and Thang T Q Le. “The colored Jones function is q -holonomic”. In: *Geometry and Topology* 9.3 (July 2005), pp. 1253–1293. DOI: [10.2140/gt.2005.9.1253](https://doi.org/10.2140/gt.2005.9.1253).
- [63] Stavros Garoufalidis and Xinyu Sun. *The non-commutative A-polynomial of twist knots*. 2008. DOI: [10.48550/ARXIV.0802.1153](https://doi.org/10.48550/ARXIV.0802.1153).
- [64] Stavros Garoufalidis and Xinyu Sun. “The C-polynomial of a knot”. In: *Algebraic Geometric Topology* 6.4 (Oct. 2006), pp. 1623–1653. DOI: [10.2140/agt.2006.6.1623](https://doi.org/10.2140/agt.2006.6.1623).