

A simple upper bound for the perimeter of an ellipse

Jean-Christophe PAIN

CEA, DAM, DIF, F-91297 Arpajon, France
 Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
 91680 Bruyères-le-Châtel, France

May 26, 2022

Abstract

We propose a simple derivation of an upper bound for the perimeter of an ellipse. The procedure, which relies on the use of elliptic integrals, consists in introducing, *via* inequalities and convexity properties, specific integrals which can be calculated analytically.

1 Introduction

The perimeter of an ellipse is the length of the continuous line forming its boundary. Computing accurate approximations to the perimeter of an ellipse has been a subject of interest for mathematicians for a long time [1–3]. Many approaches have been used to estimate its value, such as approximation formulas (two formulas derived by Ramanujan are particularly famous [4, 5] and we provide a non-exhaustive list in Appendix B) [6–8], infinite series (Maclaurin, Gauss-Kummer, Euler, etc.), hypergeometric functions [9, 10] (see Appendix C), and of course (elliptic) integrals [11–15].

It is well known that the perimeter of an ellipse can be expressed exactly as a complete elliptic integral of the second kind. Approximate formulas can, of course, be obtained by truncating the series representations of exact formulas. The difficulty of calculating exactly the perimeter has aroused a particular interest in the search for bounds. A long time ago, Kepler found the geometric mean of semi-major and semi-minor axis, as a lower bound for the perimeter, and many sophisticated approaches were developed along the years (see for instance [16–18]). The subject is still an active field of research [19, 20], important for astrophysical applications (trajectories of planets, satellites, *etc.*).

An ellipse with semi-major axis a and semi-minor axis b is parametrized by $x = a \cos \phi$ and $y = b \sin \phi$. The infinitesimal variation of curvilinear abscissa s is $ds = \sqrt{dx^2 + dy^2}$. The length of a quarter of the ellipse is provided by the elliptic integral

$$L = \int_0^{\pi/2} ds = \int_0^{\pi/2} \sqrt{a^2 \sin^2 \phi + b^2 \cos^2 \phi} d\phi. \quad (1)$$

We will use the latter formula to derive the upper bound. The total perimeter of the ellipse will therefore be

$$L_e = 4L. \quad (2)$$

The well-known simple lower bound $2\pi\sqrt{ab}$ and upper bound $4a + \pi b$ are recalled in sections 2 and 3 respectively, and the main result of the present work, which is an upper bound more accurate than the one of section 3, but still rather simple, is presented in section 4.

2 The geometric-mean lower bound

Let us consider two parametrizations of an ellipse

$$\begin{aligned} x_1 &= a \cos \phi, \\ y_1 &= b \sin \phi \end{aligned} \quad (3)$$

and

$$\begin{aligned} x_2 &= b \cos \phi, \\ y_2 &= a \sin \phi. \end{aligned} \quad (4)$$

The two associated lengths of the quarter of the ellipse are

$$L_1 = \int_0^{\pi/2} \sqrt{a^2 \cos^2 \phi + b^2 \sin^2 \phi} \, d\phi \quad (5)$$

and

$$L_2 = \int_0^{\pi/2} \sqrt{b^2 \cos^2 \phi + a^2 \sin^2 \phi} \, d\phi. \quad (6)$$

Since $t \rightarrow \sqrt{t}$ is a concave function, one has

$$\sqrt{\lambda t_1 + (1 - \lambda)t_2} \geq \lambda\sqrt{t_1} + (1 - \lambda)\sqrt{t_2}, \quad (7)$$

yielding ($\lambda = \cos^2 \phi$, $t_1 = a^2$ and $t_2 = b^2$):

$$L_1 \geq \int_0^{\pi/2} (a \cos^2 \phi + b \sin^2 \phi) \, d\phi \quad (8)$$

and

$$L_2 \geq \int_0^{\pi/2} (b \cos^2 \phi + a \sin^2 \phi) \, d\phi \quad (9)$$

and therefore

$$L = L_1 = L_2 = \frac{1}{2}(L_1 + L_2) \geq (a + b)\frac{\pi}{4}. \quad (10)$$

Setting $ab = R^2$, the minimum of $a + b = a + R/a^2 = f(a)$ is obtained for $R = \sqrt{ab}$ and is equal to $2R$. In other words, the ellipse which has the minimum perimeter for a given area is the circle of radius R . This yields the well-known result

$$\frac{\pi}{2}\sqrt{ab} \leq L \quad (11)$$

and for the total perimeter of the ellipse

$$2\pi\sqrt{ab} \leq L_e. \quad (12)$$




Figure 1: quarters of ellipse with semi-major and semi-minor axis $(a = 2, b = \sqrt{3})$ and $(a = 2, b = \sqrt{3})$ respectively.

3 First upper bound

Equation (1) is equivalent to

$$L = a \int_0^{\pi/2} \sqrt{1 - \frac{(a^2 - b^2)}{a^2} \cos^2 \phi} d\phi \quad (13)$$

which can be put in the form

$$L = a \frac{\pi}{2} \mathcal{F} \left(\frac{b^2 - a^2}{a^2} \right) \cos^2 \phi d\phi = a \frac{\pi}{2} \mathcal{F} (-e^2) \quad (14)$$

where we have introduced the function

$$\mathcal{F}(t) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{1 + t \cos^2 \phi} d\phi \quad (15)$$

where e represents the eccentricity of the ellipse defined by

$$\frac{b^2}{a^2} = 1 - e^2. \quad (16)$$

The asymptotic behaviour of the function \mathcal{F} is studied in Appendix A. If $-1 < t < 0$, then $-t/(1+t) > 0$ and

$$\mathcal{F}(t) - \mathcal{F}(-1) = \frac{2}{\pi} \int_0^{\pi/2} \left(\sqrt{1 + t \cos^2 \phi} - \sqrt{1 - \cos^2 \phi} \right) d\phi. \quad (17)$$

Since

$$\sqrt{u} - \sqrt{v} = \frac{u - v}{\sqrt{u} + \sqrt{v}}, \quad (18)$$

we get

$$\mathcal{F}(t) - \mathcal{F}(-1) = \frac{2}{\pi} \int_0^{\pi/2} \frac{(1+t) \cos^2 \phi}{\sqrt{1+t \cos^2 \phi} + \sqrt{1-\cos^2 \phi}} d\phi. \quad (19)$$

and

$$\mathcal{F}(t) - \mathcal{F}(-1) \leq \frac{2}{\pi}(1+t) \int_0^{\pi/2} \frac{\cos^2 \phi}{\sqrt{1+t}} d\phi, \quad (20)$$

i.e.

$$\mathcal{F}(t) - \mathcal{F}(-1) \leq \frac{2}{\pi} \sqrt{1+t} \frac{\pi}{4}. \quad (21)$$

because

$$\int_0^{\pi/2} \cos^2 \phi \, d\phi = \frac{\pi}{4}. \quad (22)$$

This enables us to write

$$\mathcal{F}(t) - \mathcal{F}(-1) \leq \frac{\sqrt{1+t}}{2}. \quad (23)$$

Since $-1 < -e^2 < 0$, we get

$$\mathcal{F}(-e^2) - \mathcal{F}(-1) \leq \frac{\sqrt{1-e^2}}{2}, \quad (24)$$

leading to

$$L \leq a + \frac{\pi}{4}b. \quad (25)$$

and for the total perimeter of the ellipse

$$\boxed{L_e \leq 4a + \pi b.} \quad (26)$$

4 Second upper bound

Let us go back to Eq. (19):

$$\mathcal{F}(t) - \mathcal{F}(-1) = \frac{2}{\pi} \int_0^{\pi/2} \frac{(1+t) \cos^2 \phi}{\sqrt{1+t \cos^2 \phi} + \sqrt{1-\cos^2 \phi}} d\phi. \quad (27)$$

We have

$$\mathcal{F}(t) - \mathcal{F}(-1) \leq \frac{2}{\pi}(1+t) \int_0^{\pi/2} \frac{\cos^2 \phi}{\sqrt{1+t} + \sin \phi} d\phi, \quad (28)$$

where we introduce, for $0 < \xi < 1$, the following integral

$$\mathcal{H}(\xi) = \int_0^{\pi/2} \frac{\cos^2 \phi}{\xi + \sin \phi} d\phi. \quad (29)$$




Figure 2: Comparison between the first and second upper bounds of the perimeter of a quarter of an ellipse with semi-major axis $a = 2$. On has $f_1(a, b) = a + \frac{\pi}{4}b$ (Eq. (25)) and $f_2(a, b) = a + \frac{b^2}{a^2} \left[\sqrt{a^2 - b^2} \ln \left(\frac{a + \sqrt{a^2 - b^2}}{b} \right) + \frac{\pi}{2}b - a \right]$ (Eq. (44)).

We obtain

$$\begin{aligned}
 \mathcal{H}(\xi) &= \int_0^{\pi/2} \frac{1 - \sin^2 \phi}{\xi + \sin \phi} d\phi = \int_0^{\pi/2} \frac{(\xi^2 - \sin^2 \phi) + (1 - \xi^2)}{1 + \sin \phi} d\phi \\
 &= \int_0^{\pi/2} (\xi - \sin \phi) d\phi + (1 - \xi^2) \int_0^{\pi/2} \frac{d\phi}{\xi + \sin \phi} \\
 &= \xi \frac{\pi}{2} - 1 + (1 - \xi^2) \mathcal{J}(\xi)
 \end{aligned} \tag{30}$$

with

$$\mathcal{J}(\xi) = \int_0^{\pi/2} \frac{d\phi}{\xi + \sin \phi}. \tag{31}$$

Setting $t = \tan(\phi/2)$, we have

$$dt = \frac{1}{2} \left[1 + \tan^2 \left(\frac{\phi}{2} \right) \right] d\phi, \tag{32}$$

i.e. $d\phi = 2dt/(1 + t^2)$. Then, since

$$\sin \phi = \frac{2t}{(1 + t^2)}, \tag{33}$$

we get

$$\frac{d\phi}{\xi + \sin \phi} = \frac{2dt}{\xi t^2 + 2t + \xi}. \tag{34}$$

Writing

$$\frac{2}{\xi t^2 + 2t + \xi} = \frac{2}{\xi(t - \chi_1)(t - \chi_2)} \quad (35)$$

with

$$\chi_1 = \frac{-1 - \sqrt{1 - \xi^2}}{\lambda} \quad (36)$$

and

$$\chi_2 = \frac{-1 + \sqrt{1 - \xi^2}}{\lambda}, \quad (37)$$

enables us to write

$$\begin{aligned} \mathcal{J}(\xi) &= \frac{2}{\xi} \int_0^1 \frac{dt}{(t - \chi_1)(t - \chi_2)} = \frac{2}{\xi(\chi_1 - \chi_2)} \left[\frac{1}{t - \chi_1} - \frac{1}{t - \chi_2} \right] dt \\ &= \frac{2}{\xi(\chi_1 - \chi_2)} \left| \ln \left(\frac{t - \chi_1}{t - \chi_2} \right) \right|_0^1 = \frac{2}{\xi(\chi_1 - \chi_2)} \ln \left[\frac{(1 - \chi_1)\chi_2}{(1 - \chi_2)\chi_1} \right]. \end{aligned} \quad (38)$$

We have also

$$\chi_1 \chi_2 = 1 \quad (39)$$

and

$$\chi_1 - \chi_2 = -\frac{2}{\xi} \sqrt{1 - \xi^2} \quad (40)$$

and thus

$$\begin{aligned} \mathcal{J}(\xi) &= -\frac{1}{\sqrt{1 - \xi^2}} \ln \left(\frac{\chi_2 - 1}{\chi_1 - 1} \right) = \frac{1}{\sqrt{1 - \xi^2}} \ln \left(\frac{\xi + 1 + \sqrt{1 - \xi^2}}{\xi + 1 - \sqrt{1 - \xi^2}} \right) \\ &= \frac{1}{\sqrt{1 - \xi^2}} \ln \left(\frac{\sqrt{\xi + 1} + \sqrt{1 - \xi}}{\sqrt{\xi + 1} - \sqrt{1 - \xi}} \right) \\ &= \frac{1}{\sqrt{1 - \xi^2}} \ln \left(\frac{1 + \sqrt{1 - \xi^2}}{\xi} \right), \end{aligned} \quad (41)$$

yielding

$$\mathcal{K}(\xi) = \frac{\xi\pi}{2} - 1 + \sqrt{1 - \xi^2} \ln \left(\frac{1 + \sqrt{1 - \xi^2}}{\xi} \right). \quad (42)$$

Therefore

$$L = \frac{\pi}{2} a \mathcal{F} \left(\frac{b^2}{a^2} - 1 \right) \leq \frac{\pi}{2} a \left[\mathcal{F}(-1) + \frac{2}{\pi} \left(\frac{b^2}{a^2} \right) \mathcal{K} \left(\frac{b}{a} \right) \right], \quad (43)$$

or finally

$$L \leq a + \frac{b^2}{a^2} \left[\sqrt{a^2 - b^2} \ln \left(\frac{a + \sqrt{a^2 - b^2}}{b} \right) + \frac{\pi}{2} b - a \right] \quad (44)$$

and for the total perimeter of the ellipse

$$\boxed{L_e \leq 4a + 4 \frac{b^2}{a^2} \left[\sqrt{a^2 - b^2} \ln \left(\frac{a + \sqrt{a^2 - b^2}}{b} \right) + \frac{\pi}{2} b - a \right]}. \quad (45)$$

This upper bound is better than the first one in Eq. (26) (see figure 2).

5 Conclusion

In this article, we obtained an upper bound for the perimeter of an ellipse. The calculation consists in simplifying elliptic integrals by applying inequalities and convexity properties. It is hoped that the method may stimulate the derivation of further bounds, improving the compromise between simplicity and accuracy.

References

- [1] S. Adlaj, *An eloquent formula for the perimeter of an ellipse*, Notices of the AMS **59**, 1094-1099 (2012).
- [2] J. M. Borwein and P. B. Borwein, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity* (New York: Wiley, 1987).
- [3] D. H. Bailey, J. M. Borwein, *Pi: The Next Generation : A Sourcebook on the Recent History of Pi and Its Computation* (Springer International Publishing AG, 2016).
- [4] G. Almkvist and B. Berndt, *Gauss, Landen, Ramanujan, the arithmetic-geometric mean, ellipses, π , and the Ladies diary*, Amer. Math. Monthly **95**,5 585-608 (1988).
- [5] M. B. Villarino, *Ramanujan's perimeter of an ellipse*, <https://arxiv.org/abs/math/0506384>
- [6] Jean-Baptiste Hiriart-Urruty, *Propos elliptiques...*, <https://www.math.univ-toulouse.fr/~jbhu/Ellipses.pdf>
- [7] <http://www.ebyte.it/library/docs/math05a/EllipsePerimeterApprox05.html>
- [8] R. W .Barnard, K. Pearce, and L. Schovanec, *Inequalities for the Perimeter of an Ellipse*, J. Math. Anal. Appl. **260**, 295–306 (2001).
- [9] P. C. Abbott, *On the Perimeter of an Ellipse*, The Mathematica journal **11**, 172-185 (2009).
- [10] T. R. Chandrupatla and T. J. Osler, *The perimeter of an ellipse*, Math. Sci. **35**, 122-131 (2010).
- [11] N. H. Abel, *Recherches sur les fonctions elliptiques*, J. für die Reine und Angew. Math. **2**, 101–181 (1827).
- [12] M. Abramowitz and I. A. Stegun, *Elliptic Integrals*, pp. 587-607, Ch. 17 in Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing (New York: Dover, 1972).
- [13] G. Arfken, *Elliptic Integrals*, section 5.8, pp. 321-327 in Mathematical Methods for Physicists, 3rd ed. (Orlando, FL: Academic Press, 1985).

- [14] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, *Integrals and Series, Vol. 1: Elementary Functions* (New York: Gordon and Breach, 1986).
- [15] E. T. Whittaker and G. N. Watson, *A Course in Modern Analysis*, 4th ed. (Cambridge, England: Cambridge University Press, 1990).
- [16] R. E. Pfeifer, *Bounds on the Perimeter of an Ellipse via Minkowski Sums*, Coll. Math. J. **19**, 348-350 (1988).
- [17] M.-K. Wang, Y.-M. Chu, S.-L. Qiu and Y.-P. Jiang, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164**, 928–937 (2012).
- [18] I. Gusić, *On the bounds for the perimeter of an ellipse*, Math. Gaz. **99**, 540-541 (2015).
- [19] Z.-Y. He, M.-K. Wang, Y.-P. Jiang and Y.-M. Chu, *Bounds for the perimeter of an ellipse in terms of power means*, J. Math. Inequalities **14**, 887-899 (2020).
- [20] T. Zhao, M. Wang and Y. Chu *On the Bounds of the Perimeter of an Ellipse*, Acta Math. Sci. **42**, 491-501 (2022).

Appendix A: Asymptotic behavior

Let us consider $t > -1$, i.e. $-t/(1+t) > -1$. We have

$$\begin{aligned} \mathcal{F}\left(\frac{-t}{1+t}\right) &= \frac{2}{\pi} \int_0^{\pi/2} \frac{\sqrt{1+t-t\cos^2\phi}}{\sqrt{1+t}} d\phi \\ &= \frac{2}{\pi} \frac{1}{\sqrt{1+t}} \int_0^{\pi/2} \sqrt{1+t\sin^2\phi} d\phi \end{aligned} \quad (46)$$

Making the change of variables $\psi = \pi/2 - \phi$, we get

$$\int_0^{\pi/2} \sqrt{1+t\sin^2\phi} d\phi = \int_0^{\pi/2} \sqrt{1+t\cos^2\phi} d\phi \quad (47)$$

and thus

$$\sqrt{1+t} \mathcal{F}\left(-\frac{t}{1+t}\right) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{1+t\cos^2\phi} d\phi \quad (48)$$

and

$$\mathcal{F}(t) = \sqrt{1+t} \mathcal{F}\left(-\frac{t}{1+t}\right) \quad (49)$$

$$\frac{\mathcal{F}(t)}{\sqrt{t}} = \frac{\mathcal{F}(t)}{\sqrt{1+t}} \frac{\sqrt{1+t}}{\sqrt{t}} = \mathcal{F}\left(-\frac{t}{1+t}\right) \sqrt{1+\frac{1}{t}} \quad (50)$$

If $t \rightarrow \infty$, $-t/(1+t) \rightarrow -1$ and $\mathcal{F}(t)/\sqrt{t} \rightarrow \mathcal{F}(-1) = 2/\pi$. Therefore

$$\mathcal{F}(t) \approx \frac{2}{\pi} \sqrt{t}. \quad (51)$$

Appendix B: A few approximate formulas

In this appendix, we provide a non-exhaustive of approximate formulas for the perimeter of an ellipse:

Kepler (1609) [lower bound of section 2]:

$$L_e \approx 2\pi\sqrt{ab} \quad (52)$$

$$L_e \approx \pi(a + b) \quad (53)$$

Euler

$$L_e \approx 2\pi\sqrt{\frac{a^2 + b^2}{2}} \quad (54)$$

Sipos

$$L_e \approx 2\pi\sqrt{\frac{(a + b)^2}{(\sqrt{a} + \sqrt{b})^2}} \quad (55)$$

Cesaro

$$L_e \approx \pi(a + b) + \frac{\pi}{4} \frac{(a - b)^2}{a + b} \quad (56)$$

Muir

$$L_e \approx 2\pi \left[\frac{a^{3/2} + b^{3/2}}{2} \right]^{2/3} \quad (57)$$

Peano, Boussinesq

$$L_e \approx \pi \left[\frac{3}{2}(a + b) - \sqrt{ab} \right] \quad (58)$$

Almkvist

$$L_e \approx 2\pi \frac{2(a + b)^2 - (\sqrt{a} - \sqrt{b})^4}{(\sqrt{a} + \sqrt{b})^2 + 2\sqrt{2(a + b)} (ab)^{1/4}} \quad (59)$$

Lindner

$$L_e \approx \pi(a + b) \left[1 + \frac{1}{8} \left(\frac{a - b}{a + b} \right)^2 \right]^2 \quad (60)$$

Ramanujan: first formula

$$L_e \approx \pi \left[3(a + b) - \sqrt{(3a + b)(a + 3b)} \right] \quad (61)$$

Ramanujan: second formula

$$L_e \approx \pi \left[(a + b) - \frac{3(a - b)^2}{10(a + b) + \sqrt{a^2 + 14ab + b^2}} \right]. \quad (62)$$

Appendix C: Expressions in terms of special functions or infinite sums

The integral $\mathcal{F}(t)$ can be expanded as

$$\begin{aligned}\mathcal{F}(t) &= \frac{2}{\pi} \int_0^{\pi/2} \sqrt{1+t \cos^2 \phi} \, d\phi \\ &= 1 + \sum_{k=1}^{\infty} (-1)^{k+1} t^k \frac{(2k-3)!!}{2^k k!} \frac{2}{\pi} \int_0^{\pi/2} \cos^{2k} \phi \, d\phi\end{aligned}\quad (63)$$

with $(2k-3)!! = (2k-3) \cdots 5 \times 3 \times 1$ and $(-1)!! = 1$. Setting

$$\begin{aligned}\mathcal{A}_k &= \int_0^{\pi/2} \cos^{2k} \phi \, d\phi = \int_0^{\pi/2} \cos^{2k-1} \phi \cos \phi \, d\phi \\ &= \left| \cos^{2k-1} \phi \sin \phi \right|_0^{\pi/2} + \int_0^{\pi/2} (2k-1) \cos^{2k-2} \phi \sin^2 \phi \, d\phi \\ &= (2k-1) \mathcal{A}_{k-1} - (2k-1) \mathcal{A}_k,\end{aligned}\quad (64)$$

we have

$$\mathcal{A}_k = \frac{(2k-1)}{2k} \mathcal{A}_{k-1} = \frac{(2k-1)!!}{2^k k!} \mathcal{A}_0, \quad (65)$$

where $\mathcal{A}_0 = \pi/2$ and finally

$$\mathcal{F}(t) = 1 + \sum_{k=1}^{\infty} (-1)^{k+1} \left[\frac{(2k-1)!!}{2^k k!} \right]^2 \frac{t^k}{2k-1} \quad (66)$$

giving the final Euler-Mac-Laurin series

$$L_e = 2a\pi \left\{ 1 + \sum_{k=1}^{\infty} (-1)^{k+1} \left[\frac{(2k-1)!!}{2^k k!} \right]^2 \frac{\left(\frac{b^2}{a^2} - 1 \right)^k}{2k-1} \right\}. \quad (67)$$

We mention below a few other formulas:

McLaurin

$$L_e = 2a\pi \, {}_2F_1 \left[\begin{matrix} -\frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix}; 1 - \frac{b^2}{a^2} \right] \quad (68)$$

Gauss-Kummer

$$L_e = \pi(a+b) \, {}_2F_1 \left[\begin{matrix} -\frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix}; \left(\frac{a-b}{a+b} \right)^2 \right] \quad (69)$$

Cayley

$$\begin{aligned}L_e &= 4a \left[1 + \frac{1}{2} \left(\ln \left(\frac{4a}{b} \right) - \frac{1}{1 \times 2} \right) \left(\frac{b}{a} \right)^2 + \frac{1^2 \times 3}{2^2 \times 4} \left(\ln \left(\frac{4a}{b} \right) - \frac{2}{1 \times 2} - \frac{1}{3 \times 4} \right) \left(\frac{b}{a} \right)^4 \right. \\ &\quad \left. + \frac{1^2 \times 3^2 \times 5}{2^2 \times 4^2 \times 6} \left(\ln \left(\frac{4a}{b} \right) - \frac{2}{1.2} - \frac{2}{3.4} - \frac{1}{5.6} \right) \left(\frac{b}{a} \right)^6 + \cdots \right] \quad (70)\end{aligned}$$

Euler

$$L_e = \pi \sqrt{2(a^2 + b^2)} {}_2F_1 \left[\begin{matrix} \frac{1}{4}, -\frac{1}{4} \\ 1 \end{matrix} ; \left(\frac{a^2 - b^2}{a^2 + b^2} \right)^2 \right] \quad (71)$$

Abbott

$$L_e = 2\pi \sqrt{ab} \mathcal{P}_{1/2} \left(\frac{a^2 + b^2}{2ab} \right), \quad (72)$$

where $\mathcal{P}_{1/2}$ is a Legendre function.